

| Adolescent Behaviors and Experiences Survey | | | | | | | | | | | | | | |
|--|-----------------------------------|---------------------|------------|---------------------|----------------------------|---------------------|---------------------|---------------------|--|---------------------|------------|---------------------|----------------|---------------------|
| Sexual Behaviors | American Indian or Alaska Native* | | Asian* | | Black or African American* | | Hispanic or Latino* | | Native Hawaiian or Other Pacific Islander* | | White* | | Multiple Race* | |
| | Percentage | Confidence Interval | Percentage | Confidence Interval | Percentage | Confidence Interval | Percentage | Confidence Interval | Percentage | Confidence Interval | Percentage | Confidence Interval | Percentage | Confidence Interval |
| Ever had sexual intercourse | 43.4 | 25.2 - 63.6 | 13.7 | 9.6 - 19.2 | 28.9 | 24.1 - 34.2 | 26.7 | 21.9 - 32.1 | N/A | N/A | 29.9 | 26.9 - 33.0 | 36.5 | 27.2 - 46.8 |
| Had sexual intercourse for the first time before age 13 years | 8.2 | 3.0 - 20.4 | 0.6 | 0.2 - 2.3 | 4.3 | 3.0 - 6.0 | 2.6 | 1.9 - 3.6 | N/A | N/A | 1.2 | 0.8 - 2.0 | 5.1 | 2.8 - 9.1 |
| Had sexual intercourse with four or more persons during their life | 11.9 | 6.0 - 22.3 | 0.8 | 0.2 - 3.3 | 7.8 | 5.5 - 10.8 | 3.8 | 2.7 - 5.3 | N/A | N/A | 6.2 | 4.6 - 8.2 | 10.6 | 7.1 - 15.6 |
| Were currently sexually active (had sexual intercourse with at least one person, during the 3 months before the survey) | 27.5 | 17.6 - 40.2 | 10.1 | 7.2 - 14.1 | 18.3 | 14.9 - 22.2 | 14.7 | 11.9 - 18.0 | N/A | N/A | 20.9 | 18.4 - 23.6 | 26.2 | 18.1 - 36.4 |
| Drank alcohol or used drugs before last sexual intercourse (among students who were currently sexually active) | N/A | N/A | 1.6 | 0.2 - 10.1 | 17.8 | 13.0 - 23.9 | 16.5 | 11.6 - 22.9 | N/A | N/A | 17.1 | 13.4 - 21.5 | 18.4 | 9.7 - 32.3 |
| Used a condom during last sexual intercourse (among students who were currently sexually active) | N/A | N/A | 49.1 | 32.2 - 66.2 | 54.3 | 45.1 - 63.2 | 53.3 | 45.7 - 60.8 | N/A | N/A | 55.0 | 50.9 - 59.0 | 50.4 | 37.0 - 63.6 |
| Used birth control pills before last sexual intercourse with opposite-sex partner (to prevent pregnancy, not counting emergency contraception such as Plan B or the "morning after" pill, among students who were currently sexually active) | N/A | N/A | N/A | N/A | 11.5 | 6.6 - 19.2 | 13.7 | 9.2 - 19.9 | N/A | N/A | 26.5 | 21.9 - 31.7 | 25.6 | 17.4 - 36.0 |
| Were ever tested for human immunodeficiency virus (HIV) (not counting tests done if they donated blood) | 7.4 | 3.2 - 16.0 | 5.2 | 2.9 - 9.0 | 7.2 | 5.8 - 8.9 | 4.6 | 3.3 - 6.2 | N/A | N/A | 6.3 | 5.1 - 7.9 | 5.6 | 3.2 - 9.5 |
| Were tested for a sexually transmitted disease (STD) other than HIV, such as chlamydia or gonorrhea (during the 12 months before the survey) | 8.5 | 2.9 - 22.4 | 4.2 | 1.6 - 10.1 | 7.1 | 4.8 - 10.3 | 2.9 | 2.1 - 4.0 | N/A | N/A | 4.8 | 3.8 - 6.2 | 5.6 | 2.7 - 11.1 |
| Used an IUD (such as Mirena or ParaGard) or implant (such as Implanon or Nexplanon) before last sexual intercourse with an opposite-sex partner (to prevent pregnancy, among students who were currently sexually active) | N/A | N/A | N/A | N/A | 3.4 | 1.1 - 10.1 | 6.7 | 3.7 - 11.7 | N/A | N/A | 6.4 | 4.5 - 9.1 | 4.7 | 2.4 - 8.8 |
| Used birth control pills; an IUD (such as Mirena or ParaGard) or implant (such as Implanon or Nexplanon); or a shot (such as Depo-Provera), patch (such as OrthoEvra), or birth control ring (such as NuvaRing) before last sexual intercourse with an opposite-sex partner (to prevent pregnancy, among students who were currently sexually active) | N/A | N/A | N/A | N/A | 19.9 | 13.9 - 27.7 | 26.2 | 20.3 - 33.2 | N/A | N/A | 38.6 | 32.9 - 44.6 | 39.1 | 27.3 - 52.5 |
| Used both a condom during last sexual intercourse and birth control pills; an IUD (such as Mirena or ParaGard) or implant (such as Implanon or Nexplanon); or a shot (such as Depo-Provera), patch (such as OrthoEvra), or birth control ring (such as NuvaRing) before last sexual intercourse with an opposite-sex partner (to prevent pregnancy, among students who were currently sexually active) | N/A | N/A | 3.3 | 0.4 - 21.2 | 5.8 | 2.5 - 12.6 | 7.9 | 5.0 - 12.1 | N/A | N/A | 13.9 | 11.2 - 17.2 | 15.9 | 8.8 - 27.0 |
| Did not use any method to prevent pregnancy during last sexual intercourse with an opposite-sex partner (among students who were currently sexually active) | N/A | N/A | N/A | N/A | 23.7 | 18.4 - 30.0 | 14.1 | 9.0 - 21.4 | N/A | N/A | 11.1 | 8.3 - 14.5 | 18.0 | 11.0 - 28.0 |

*Non-Hispanic

N/A: Less than 30 respondents for the subgroup