

Adolescent Behaviors and Experiences Survey

	Female		Male	
	Percentage	Confidence Interval	Percentage	Confidence Interval
Alcohol and Other Drug Use				
Had their first drink of alcohol before age 13 years (other than a few sips)	11.0	9.4 - 12.8	11.0	9.3 - 13.0
Currently drank alcohol (at least one drink of alcohol, on at least 1 day during the 30 days before the survey)	22.4	18.5 - 27.0	16.4	14.3 - 18.7
Currently were binge drinking (had four or more drinks of alcohol in a row if they were female or five or more drinks of alcohol in a row if they were male, within a couple of hours, on at least 1 day during the 30 days before the survey)	9.5	6.6 - 13.5	5.9	4.6 - 7.6
Reported that the largest number of drinks they had in a row was 10 or more (within a couple of hours, during the 30 days before the survey)	1.5	0.7 - 3.0	2.9	2.0 - 4.2
Usually got the alcohol they drank by someone giving it to them (during the 30 days before the survey, among students who currently drank alcohol)	41.9	36.4 - 47.5	33.1	27.4 - 39.4
Ever used marijuana (one or more times during their life)	25.6	22.4 - 29.0	24.3	22.2 - 26.6
Tried marijuana for the first time before age 13 years	3.0	2.4 - 3.9	3.8	2.8 - 5.0
Currently used marijuana (one or more times during the 30 days before the survey)	12.7	10.2 - 15.6	12.9	11.2 - 14.8
Ever used synthetic marijuana (one or more times during their life)	6.2	4.9 - 7.7	5.3	4.3 - 6.6
Ever took prescription pain medicine without a doctor's prescription or differently than how a doctor told them to use it (counting drugs such as codeine, Vicodin, OxyContin, Hydrocodone, and Percocet, one or more times during their life)	11.4	10.0 - 12.9	7.6	6.2 - 9.3
Currently took prescription pain medicine without a doctor's prescription or differently than how a doctor told them to use it (one or more times during the 30 days before the survey)	5.4	4.4 - 6.5	3.2	2.3 - 4.4
Ever used cocaine (any form of cocaine, including powder, crack, or freebase, one or more times during their life)	0.8	0.6 - 1.2	2.0	1.4 - 3.0
Ever used inhalants (sniffed glue, breathed the contents of aerosol spray cans, or inhaled any paints or sprays to get high, one or more times during their life)	7.2	5.8 - 8.9	4.2	3.4 - 5.3
Ever used heroin (also called "smack," "junk," or "China White," one or more times during their life)	0.2	0.1 - 0.5	1.1	0.7 - 1.7
Ever used methamphetamines (also called "speed," "crystal meth," "crank," "ice," or "meth," one or more times during their life)	0.6	0.3 - 1.0	1.1	0.8 - 1.6
Ever used ecstasy (also called "MDMA" or "Molly," one or more times during their life)	1.5	1.0 - 2.3	2.3	1.7 - 3.2
Ever used hallucinogenic drugs (such as LSD, acid, PCP, angel dust, mescaline, or mushrooms, one or more times during their life)	3.0	2.2 - 4.1	4.5	3.4 - 5.8
Ever used select illicit drugs (counting cocaine, inhalants, heroin, methamphetamines, ecstasy, or hallucinogens, one or more times during their life)	9.6	7.9 - 11.6	8.5	7.0 - 10.3
Ever injected any illegal drug (used a needle to inject any illegal drug into their body, one or more times during their life)	0.3	0.1 - 0.5	0.8	0.5 - 1.3
Were offered, sold, or given an illegal drug on school property (during the 12 months before the survey, among students who attended school in-person during the 12 months before the survey)	8.5	6.7 - 10.8	10.1	7.5 - 13.4