

Adolescent Behaviors and Experiences Survey				
Dietary Behaviors	Female		Male	
	Percentage	Confidence Interval	Percentage	Confidence Interval
Did not drink fruit juice (100% fruit juices one or more times during the 7 days before the survey)	42.1	39.7 - 44.5	37.2	34.5 - 40.0
Did not eat fruit (one or more times during the 7 days before the survey)	13.1	10.8 - 15.7	16.1	13.2 - 19.4
Did not eat green salad (one or more times during the 7 days before the survey)	44.9	41.8 - 48.1	51.9	48.4 - 55.4
Did not eat potatoes (one or more times during the 7 days before the survey)	39.7	37.1 - 42.3	34.9	33.0 - 36.7
Did not eat carrots (one or more times during the 7 days before the survey)	60.7	57.9 - 63.6	56.0	53.1 - 58.8
Did not eat other vegetables (one or more times during the 7 days before the survey)	17.8	15.4 - 20.5	19.4	16.7 - 22.3
Did not eat fruit or drink 100% fruit juices (such as orange juice, apple juice, or grape juice, during the 7 days before the survey)	8.2	6.6 - 10.2	9.8	7.6 - 12.5
Ate fruit or drank 100% fruit juices one or more times per day (such as orange juice, apple juice, or grape juice, during the 7 days before the survey)	45.4	42.6 - 48.2	49.5	46.1 - 52.8
Ate fruit or drank 100% fruit juices two or more times per day (such as orange juice, apple juice, or grape juice, during the 7 days before the survey)	20.7	18.5 - 23.2	23.2	21.0 - 25.5
Did not eat vegetables (green salad, potatoes [excluding french fries, fried potatoes, or potato chips], carrots, or other vegetables, during the 7 days before the survey)	7.0	5.9 - 8.3	9.2	7.8 - 10.8
Ate vegetables one or more times per day (green salad, potatoes [excluding french fries, fried potatoes, or potato chips], carrots, or other vegetables, during the 7 days before the survey)	53.1	49.7 - 56.5	56.7	54.1 - 59.3
Ate vegetables two or more times per day (green salad, potatoes [excluding french fries, fried potatoes, or potato chips], carrots, or other vegetables, during the 7 days before the survey)	20.6	18.7 - 22.6	22.9	20.6 - 25.3
Ate vegetables three or more times per day (green salad, potatoes [excluding french fries, fried potatoes, or potato chips], carrots, or other vegetables, during the 7 days before the survey)	10.2	8.9 - 11.5	12.6	10.9 - 14.5
Drank a can, bottle, or glass of soda or pop one or more times per day (such as Coke, Pepsi, or Sprite, not counting diet soda or diet pop, during the 7 days before the survey)	13.2	9.9 - 17.4	18.7	14.9 - 23.1
Did not drink a can, bottle, or glass of soda or pop (such as Coke, Pepsi, or Sprite, not counting diet soda or diet pop, one or more times during the 7 days before the survey)	37.7	34.3 - 41.3	31.9	29.1 - 34.8
Drank a can, bottle, or glass of soda or pop two or more times per day (such as Coke, Pepsi, or Sprite, not counting diet soda or diet pop, during the 7 days before the survey)	8.3	5.9 - 11.4	11.0	8.2 - 14.6
Did not drink milk (during the 7 days before the survey)	43.2	40.6 - 45.9	25.9	23.8 - 28.0
Drank one or more glasses per day of milk (counting the milk they drank in a glass or cup, from a carton, or with cereal and counting the half pint of milk served at school as equal to one glass, during the 7 days before the survey)	15.8	14.0 - 17.8	31.7	29.6 - 33.8
Drank three or more glasses per day of milk (counting the milk they drank in a glass or cup, from a carton, or with cereal and counting the half pint of milk served at school as equal to one glass, during the 7 days before the survey)	2.6	2.0 - 3.4	7.1	5.9 - 8.7
Did not eat breakfast (during the 7 days before the survey)	15.5	13.9 - 17.2	14.5	12.5 - 16.7
Ate breakfast on all 7 days (during the 7 days before the survey)	25.4	23.0 - 28.1	32.5	30.0 - 35.1
Did not drink a can, bottle, or glass of a sports drink (such as Gatorade or PowerAde, not counting low-calorie sports drinks such as Propel or G2, during the 7 days before the survey)	62.0	57.9 - 66.0	47.1	43.5 - 50.7
Drank a can, bottle, or glass of a sports drink one or more times per day (such as Gatorade or PowerAde, not counting low-calorie sports drinks such as Propel or G2, during the 7 days before the survey)	6.5	5.1 - 8.3	13.7	11.6 - 16.1
Drank a can, bottle, or glass of a sports drink two or more times per day (such as Gatorade or PowerAde, not counting low-calorie sports drinks such as Propel or G2, during the 7 days before the survey)	3.3	2.4 - 4.4	7.8	6.2 - 9.7
Did not drink a bottle or glass of plain water (counting tap, bottled, and unflavored sparkling water, during the 7 days before the survey)	2.9	2.2 - 3.7	5.3	4.1 - 6.7
Drank a bottle or glass of plain water one or more times per day (counting tap, bottled, and unflavored sparkling water, during the 7 days before the survey)	75.9	73.1 - 78.5	75.6	72.9 - 78.2
Drank a bottle or glass of plain water two or more times per day (counting tap, bottled, and unflavored sparkling water, during the 7 days before the survey)	69.1	65.9 - 72.2	67.8	64.7 - 70.7
Drank a bottle or glass of plain water three or more times per day (counting tap, bottled, and unflavored sparkling water, during the 7 days before the survey)	52.6	49.3 - 55.9	54.9	51.5 - 58.2