

Adolescent Behaviors and Experiences Survey				
Alcohol and Other Drug Use	Opposite Sex Only		Same Sex Only or Both Sexes	
	Percentage	Confidence Interval	Percentage	Confidence Interval
Had their first drink of alcohol before age 13 years (other than a few sips)	16.4	14.2 - 18.8	27.9	22.5 - 34.1
Currently drank alcohol (at least one drink of alcohol, on at least 1 day during the 30 days before the survey)	35.8	30.9 - 41.0	40.0	33.0 - 47.4
Currently were binge drinking (had four or more drinks of alcohol in a row if they were female or five or more drinks of alcohol in a row if they were male, within a couple of hours, on at least 1 day during the 30 days before the survey)	16.1	12.4 - 20.6	17.3	13.2 - 22.4
Reported that the largest number of drinks they had in a row was 10 or more (within a couple of hours, during the 30 days before the survey)	5.2	3.8 - 7.2	3.2	1.6 - 6.3
Usually got the alcohol they drank by someone giving it to them (during the 30 days before the survey, among students who currently drank alcohol)	36.7	31.4 - 42.4	31.1	23.6 - 39.8
Ever used marijuana (one or more times during their life)	46.2	42.6 - 49.8	55.7	48.3 - 62.9
Tried marijuana for the first time before age 13 years	5.9	4.4 - 7.7	12.6	9.3 - 16.9
Currently used marijuana (one or more times during the 30 days before the survey)	25.2	22.7 - 27.8	31.7	24.6 - 39.8
Ever used synthetic marijuana (one or more times during their life)	10.3	8.4 - 12.5	15.6	11.3 - 21.1
Ever took prescription pain medicine without a doctor's prescription or differently than how a doctor told them to use it (counting drugs such as codeine, Vicodin, OxyContin, Hydrocodone, and Percocet, one or more times during their life)	12.3	10.5 - 14.4	26.1	21.1 - 31.8
Currently took prescription pain medicine without a doctor's prescription or differently than how a doctor told them to use it (one or more times during the 30 days before the survey)	6.0	4.9 - 7.3	10.5	7.0 - 15.5
Ever used cocaine (any form of cocaine, including powder, crack, or freebase, one or more times during their life)	2.3	1.4 - 3.7	5.0	3.0 - 8.2
Ever used inhalants (sniffed glue, breathed the contents of aerosol spray cans, or inhaled any paints or sprays to get high, one or more times during their life)	6.8	5.4 - 8.5	19.6	14.1 - 26.6
Ever used heroin (also called "smack," "junk," or "China White," one or more times during their life)	0.7	0.4 - 1.2	2.4	1.0 - 5.6
Ever used methamphetamines (also called "speed," "crystal meth," "crank," "ice," or "meth," one or more times during their life)	1.1	0.6 - 2.0	2.7	1.2 - 6.0
Ever used ecstasy (also called "MDMA" or "Molly," one or more times during their life)	3.6	2.5 - 5.3	6.4	3.5 - 11.3
Ever used hallucinogenic drugs (such as LSD, acid, PCP, angel dust, mescaline, or mushrooms, one or more times during their life)	7.1	5.4 - 9.5	13.3	8.7 - 19.8
Ever used select illicit drugs (counting cocaine, inhalants, heroin, methamphetamines, ecstasy, or hallucinogens, one or more times)	14.1	11.7 - 16.8	26.4	20.2 - 33.7
Ever injected any illegal drug (used a needle to inject any illegal drug into their body, one or more times during their life)	0.7	0.4 - 1.2	2.3	0.9 - 5.5
Were offered, sold, or given an illegal drug on school property (during the 12 months before the survey, among students who attended)	13.8	10.6 - 17.8	23.2	16.5 - 31.5