

<b>Adolescent Behaviors and Experiences Survey</b>				
	<b>Opposite Sex Only</b>		<b>Same Sex Only or Both Sexes</b>	
<b>Mental Health</b>	<b>Percentage</b>	<b>Confidence Interval</b>	<b>Percentage</b>	<b>Confidence Interval</b>
Who felt sad or hopeless (almost every day for two or more weeks in a row so that they stopped doing some usual activities, ever during the 12 months before the survey)	52.7	49.4 - 55.9	81.9	75.8 - 86.8
Seriously considered attempting suicide (during the 12 months before the survey)	24.9	21.8 - 28.4	56.6	49.1 - 63.9
Made a plan about how they would attempt suicide (during the 12 months before the survey)	19.0	16.1 - 22.3	48.1	42.0 - 54.3
Actually attempted suicide (one or more times during the 12 months before the survey)	11.7	9.4 - 14.4	35.0	28.7 - 41.9
Had a suicide attempt that resulted in an injury, poisoning, or overdose that had to be treated by a doctor or nurse (during the 12 months before the survey)	2.3	1.7 - 3.3	8.9	5.6 - 13.7
Reported that their mental health was most of the time or always not good (including stress, anxiety, and depression, during the 30 days before the survey)	35.3	31.7 - 39.1	55.8	49.1 - 62.3