

| Adolescent Behaviors and Experiences Survey  |              |                     |                           |                     |                   |                     |
|--|--------------|---------------------|---------------------------|---------------------|-------------------|---------------------|
| Alcohol and Other Drug Use   | Heterosexual |                     | Gay, Lesbian, or Bisexual |                     | Other/Questioning |                     |
|  | Percentage   | Confidence Interval | Percentage                | Confidence Interval | Percentage        | Confidence Interval |
| Had their first drink of alcohol before age 13 years (other than a few sips)   | 9.5          | 8.2 - 11.1          | 19.4                      | 15.7 - 23.7         | 14.8              | 11.2 - 19.4         |
| Currently drank alcohol (at least one drink of alcohol, on at least 1 day during the 30 days before the survey)  | 18.8         | 15.6 - 22.5         | 26.4                      | 22.1 - 31.3         | 20.1              | 15.0 - 26.4         |
| Currently were binge drinking (had four or more drinks of alcohol in a row if they were female or five or more drinks of alcohol in a row if they were male, within a couple of hours, on at least 1 day during the 30 days before the survey) | 7.8          | 5.6 - 10.8          | 9.0                       | 6.5 - 12.5          | 7.5               | 4.8 - 11.7          |
| Reported that the largest number of drinks they had in a row was 10 or more (within a couple of hours, during the 30 days before the survey)   | 2.4          | 1.6 - 3.5           | 2.1                       | 1.1 - 3.9           | 0.6               | 0.2 - 1.7           |
| Usually got the alcohol they drank by someone giving it to them (during the 30 days before the survey, among students who currently drank alcohol)   | 37.3         | 32.8 - 42.0         | 44.4                      | 36.1 - 53.0         | 36.8              | 25.8 - 49.4         |
| Ever used marijuana (one or more times during their life)  | 23.5         | 21.1 - 26.0         | 35.7                      | 31.4 - 40.3         | 25.8              | 22.0 - 29.9         |
| Tried marijuana for the first time before age 13 years   | 2.9          | 2.2 - 3.9           | 6.1                       | 4.4 - 8.4           | 4.4               | 2.6 - 7.2           |
| Currently used marijuana (one or more times during the 30 days before the survey)  | 12.0         | 10.3 - 13.8         | 18.5                      | 14.7 - 22.9         | 12.1              | 9.1 - 15.8          |
| Ever used synthetic marijuana (one or more times during their life)  | 5.1          | 4.0 - 6.5           | 8.1                       | 5.9 - 10.9          | 6.7               | 4.5 - 9.7           |
| Ever took prescription pain medicine without a doctor's prescription or differently than how a doctor told them to use it (counting drugs such as codeine, Vicodin, OxyContin, Hydrocodone, and Percocet, one or more times during their life) | 8.0          | 6.9 - 9.2           | 16.0                      | 13.1 - 19.4         | 15.5              | 12.2 - 19.5         |
| Currently took prescription pain medicine without a doctor's prescription or differently than how a doctor told them to use it (one or more times during the 30 days before the survey)  | 3.8          | 3.0 - 4.7           | 6.8                       | 5.2 - 9.0           | 5.4               | 3.8 - 7.8           |
| Ever used cocaine (any form of cocaine, including powder, crack, or freebase, one or more times during their life)   | 1.2          | 0.8 - 1.7           | 1.9                       | 1.1 - 3.3           | 1.7               | 0.8 - 3.8           |
| Ever used inhalants (sniffed glue, breathed the contents of aerosol spray cans, or inhaled any paints or sprays to get high, one or more times during their life)  | 4.0          | 3.2 - 5.0           | 13.7                      | 10.6 - 17.5         | 10.1              | 6.9 - 14.5          |
| Ever used heroin (also called "smack," "junk," or "China White," one or more times during their life)  | 0.4          | 0.2 - 0.8           | 1.1                       | 0.5 - 2.3           | 0.9               | 0.4 - 2.0           |
| Ever used methamphetamines (also called "speed," "crystal meth," "crank," "ice," or "meth," one or more times during their life)   | 0.5          | 0.3 - 0.9           | 1.5                       | 0.8 - 2.9           | 1.4               | 0.5 - 3.4           |
| Ever used ecstasy (also called "MDMA" or "Molly," one or more times during their life)   | 1.8          | 1.2 - 2.6           | 1.9                       | 1.0 - 3.7           | 3.1               | 1.5 - 6.1           |
| Ever used hallucinogenic drugs (such as LSD, acid, PCP, angel dust, mescaline, or mushrooms, one or more times during their life)  | 3.1          | 2.3 - 4.3           | 6.0                       | 4.2 - 8.6           | 5.8               | 3.9 - 8.6           |
| Ever used select illicit drugs (counting cocaine, inhalants, heroin, methamphetamines, ecstasy, or hallucinogens, one or more times during their life)   | 7.4          | 6.1 - 9.0           | 17.5                      | 14.3 - 21.3         | 12.5              | 9.0 - 17.2          |
| Ever injected any illegal drug (used a needle to inject any illegal drug into their body, one or more times during their life)   | 0.3          | 0.2 - 0.6           | 0.9                       | 0.4 - 2.0           | 0.8               | 0.3 - 2.1           |
| Were offered, sold, or given an illegal drug on school property (during the 12 months before the survey, among students who attended school in-person during the 12 months before the survey)  | 8.5          | 6.4 - 11.3          | 11.3                      | 8.2 - 15.2          | 14.4              | 10.4 - 19.5         |