

## Adolescent Behaviors and Experiences Survey

Dietary Behaviors	Percentage	Confidence Interval
Did not drink fruit juice (100% fruit juices one or more times during the 7 days before the survey)	39.7	37.8 - 41.7
Did not eat fruit (one or more times during the 7 days before the survey)	14.6	12.2 - 17.2
Did not eat green salad (one or more times during the 7 days before the survey)	48.3	45.5 - 51.2
Did not eat potatoes (one or more times during the 7 days before the survey)	37.3	35.5 - 39.0
Did not eat carrots (one or more times during the 7 days before the survey)	58.3	55.8 - 60.8
Did not eat other vegetables (one or more times during the 7 days before the survey)	18.6	16.3 - 21.0
Did not eat fruit or drink 100% fruit juices (such as orange juice, apple juice, or grape juice, during the 7 days before the survey)	9.0	7.3 - 11.0
Ate fruit or drank 100% fruit juices one or more times per day (such as orange juice, apple juice, or grape juice, during the 7 days before the survey)	47.4	44.8 - 50.0
Ate fruit or drank 100% fruit juices two or more times per day (such as orange juice, apple juice, or grape juice, during the 7 days before the survey)	21.9	20.1 - 23.8
Did not eat vegetables (green salad, potatoes [excluding french fries, fried potatoes, or potato chips], carrots, or other vegetables, during the 7 days before the survey)	8.1	7.0 - 9.4
Ate vegetables one or more times per day (green salad, potatoes [excluding french fries, fried potatoes, or potato chips], carrots, or other vegetables, during the 7 days before the survey)	54.9	52.4 - 57.4
Ate vegetables two or more times per day (green salad, potatoes [excluding french fries, fried potatoes, or potato chips], carrots, or other vegetables, during the 7 days before the survey)	21.7	20.0 - 23.6
Ate vegetables three or more times per day (green salad, potatoes [excluding french fries, fried potatoes, or potato chips], carrots, or other vegetables, during the 7 days before the survey)	11.3	10.3 - 12.5
Drank a can, bottle, or glass of soda or pop one or more times per day (such as Coke, Pepsi, or Sprite, not counting diet soda or diet pop, during the 7 days before the survey)	15.9	12.5 - 19.9
Did not drink a can, bottle, or glass of soda or pop (such as Coke, Pepsi, or Sprite, not counting diet soda or diet pop, one or more times during the 7 days before the survey)	34.9	32.1 - 37.8
Drank a can, bottle, or glass of soda or pop two or more times per day (such as Coke, Pepsi, or Sprite, not counting diet soda or diet pop, during the 7 days before the survey)	9.6	7.1 - 12.8
Did not drink milk (during the 7 days before the survey)	34.8	32.6 - 37.0
Drank one or more glasses per day of milk (counting the milk they drank in a glass or cup, from a carton, or with cereal and counting the half pint of milk served at school as equal to one glass, during the 7 days before the survey)	23.5	21.9 - 25.1
Drank three or more glasses per day of milk (counting the milk they drank in a glass or cup, from a carton, or with cereal and counting the half pint of milk served at school as equal to one glass, during the 7 days before the survey)	4.8	4.1 - 5.7
Did not eat breakfast (during the 7 days before the survey)	14.9	13.5 - 16.6
Ate breakfast on all 7 days (during the 7 days before the survey)	28.9	26.6 - 31.2
Did not drink a can, bottle, or glass of a sports drink (such as Gatorade or PowerAde, not counting low-calorie sports drinks such as Propel or G2, during the 7 days before the survey)	54.8	51.6 - 58.0
Drank a can, bottle, or glass of a sports drink one or more times per day (such as Gatorade or PowerAde, not counting low calorie sports drinks such as Propel or G2, during the 7 days before the survey)	10.0	8.4 - 11.8
Drank a can, bottle, or glass of a sports drink two or more times per day (such as Gatorade or PowerAde, not counting low calorie sports drinks such as Propel or G2, during the 7 days before the survey)	5.5	4.4 - 6.8
Did not drink a bottle or glass of plain water (counting tap, bottled, and unflavored sparkling water, during the 7 days before the survey)	4.0	3.3 - 4.9
Drank a bottle or glass of plain water one or more times per day (counting tap, bottled, and unflavored sparkling water, during the 7 days before the survey)	75.7	73.4 - 78.0
Drank a bottle or glass of plain water two or more times per day (counting tap, bottled, and unflavored sparkling water, during the 7 days before the survey)	68.4	65.6 - 71.1
Drank a bottle or glass of plain water three or more times per day (counting tap, bottled, and unflavored sparkling water, during the 7 days before the survey)	53.7	50.8 - 56.6