

Adolescent Behaviors and Experiences Survey

Tobacco Use	Percentage	Confidence Interval
Ever tried cigarette smoking (even one or two puffs)	16.1	13.2 - 19.6
First tried cigarette smoking before age 13 years (even one or two puffs)	6.0	4.5 - 7.8
Currently smoked cigarettes (on at least 1 day during the 30 days before the survey)	3.3	2.5 - 4.4
Smoked more than 10 cigarettes per day (on the days they smoked during the 30 days before the survey, among students who currently smoked cigarettes)	6.9	3.3 - 13.9
Ever used an electronic vapor product (including e-cigarettes, vapes, vape pens, e-cigars, e-hookahs, hookah pens, and mods [such as JUUL, SMOK, Suorin, Vuse, and blu])	36.1	33.0 - 39.2
Currently used an electronic vapor product (including e-cigarettes, vapes, vape pens, e-cigars, e-hookahs, hookah pens, and mods [such as JUUL, SMOK, Suorin, Vuse, and blu]), on at least 1 day during the 30 days before the survey)	15.4	13.0 - 18.1
Usually got their electronic vapor products by buying them themselves in a convenience store, supermarket, discount store, or gas station (including e-cigarettes, vapes, vape pens, e-cigars, e-hookahs, hookah pens, and mods [such as JUUL, SMOK, Suorin, Vuse, and blu]), during the 30 days before the survey, among students who currently used electronic vapor products)	8.4	6.1 - 11.4
Currently used smokeless tobacco (chewing tobacco, snuff, dip, snus, or dissolvable tobacco products [such as Copenhagen, Grizzly, Skoal, or Camel Snus], not counting any electronic vapor products, on at least 1 day during the 30 days before the survey)	1.9	1.3 - 2.8
Currently smoked cigars (cigars, cigarillos, or little cigars, on at least 1 day during the 30 days before the survey)	2.3	1.8 - 2.9
Tried to quit using all tobacco products (including cigarettes, electronic vapor products, smokeless tobacco, cigars, shisha or hookah tobacco, or pipe tobacco, during the 12 months before the survey, among students who used any tobacco products during the 12 months before the survey)	55.7	52.6 - 58.9
Currently smoked cigarettes frequently (on 20 or more days during the 30 days before the survey)	0.5	0.3 - 0.9
Currently smoked cigarettes daily (on all 30 days during the 30 days before the survey)	0.4	0.3 - 0.7
Currently used electronic vapor products frequently (on 20 or more days during the 30 days before the survey)	5.8	4.5 - 7.5
Currently used electronic vapor products daily (on all 30 days during the 30 days before the survey)	4.8	3.7 - 6.2
Currently used smokeless tobacco frequently (chewing tobacco, snuff, dip, snus, or dissolvable tobacco products [such as Copenhagen, Grizzly, Skoal, or Camel Snus], not counting any electronic vapor products, on 20 or more days during the 30 days before the survey)	0.5	0.3 - 0.9
Currently used smokeless tobacco daily (chewing tobacco, snuff, dip, snus, or dissolvable tobacco products [such as Copenhagen, Grizzly, Skoal, or Camel Snus], not counting any electronic vapor products, on all 30 days during the 30 days before the survey)	0.4	0.3 - 0.7
Currently smoked cigars frequently (cigars, cigarillos, or little cigars, on 20 or more days during the 30 days before the survey)	0.5	0.3 - 0.9
Currently smoked cigars daily (cigars, cigarillos, or little cigars, on all 30 days during the 30 days before the survey)	0.4	0.3 - 0.8
Currently smoked cigarettes or cigars (on at least 1 day during the 30 days before the survey)	4.4	3.4 - 5.7
Currently smoked cigarettes or cigars or used smokeless tobacco (on at least 1 day during the 30 days before the survey)	5.0	3.9 - 6.4
Currently smoked cigarettes or cigars or used smokeless tobacco or electronic vapor products (on at least 1 day during the 30 days before the survey)	15.6	13.3 - 18.2
Currently smoked cigarettes or used electronic vapor products (on at least 1 day during the 30 days before the survey)	15.2	12.9 - 17.7