

TABLE 10. Estimated percent using inhaled short acting beta agonists among children¹ with current² or active³ asthma status by state/territory - BRFSS Asthma Call-back Survey, United States, 2006-2010

State/Territory	Current				Active			
	Sample size [±]	%	SE [§]	95% CI [†]	Sample size	%	SE	95% CI
Arizona	125	46.8	7.3	(32.5 - 61.1)	149	40.1	6.4	(27.6 - 52.6)
California	170	57.0	5.1	(47.0 - 67.0)	206	51.2	4.8	(41.8 - 60.6)
Connecticut	347	53.6	3.6	(46.5 - 60.7)	413	47.4	3.3	(40.9 - 53.9)
District of Columbia	114	48.3	6.0	(36.5 - 60.1)	140	42.3	5.2	(32.1 - 52.5)
Georgia	341	43.3	3.7	(36.0 - 50.6)	399	38.9	3.3	(32.4 - 45.4)
Hawaii	417	34.6	2.9	(28.9 - 40.3)	444	33.5	2.8	(28.0 - 39.0)
Illinois	168	48.4	5.0	(38.6 - 58.2)	196	43.0	4.6	(34.0 - 52.0)
Indiana	348	46.9	3.6	(39.8 - 54.0)	391	40.3	3.3	(33.8 - 46.8)
Iowa	249	44.9	3.6	(37.8 - 52.0)	274	40.7	3.4	(34.0 - 47.4)
Kansas	588	53.8	2.4	(49.1 - 58.5)	657	48.8	2.3	(44.3 - 53.3)
Louisiana	107	32.2	5.4	(21.6 - 42.8)	122	29.5	5.0	(19.7 - 39.3)
Maine	284	54.8	3.4	(48.1 - 61.5)	332	47.2	3.1	(41.1 - 53.3)
Maryland	385	48.3	3.4	(41.6 - 55.0)	467	39.5	3.0	(33.6 - 45.4)
Massachusetts	269	46.6	4.5	(37.8 - 55.4)	315	41.8	4.0	(34.0 - 49.6)
Michigan	480	45.9	3.0	(40.0 - 51.8)	533	43.6	2.8	(38.1 - 49.1)
Mississippi	80	31.9	6.1	(19.9 - 43.9)	94	28.5	5.3	(18.1 - 38.9)
Missouri	230	51.2	4.7	(42.0 - 60.4)	255	46.1	4.5	(37.3 - 54.9)
Montana	210	46.7	4.4	(38.1 - 55.3)	253	42.2	4.0	(34.4 - 50.0)
Nebraska	467	47.2	3.9	(39.6 - 54.8)	528	43.4	3.7	(36.1 - 50.7)
New Hampshire	150	48.9	5.1	(38.9 - 58.9)	173	44.4	4.8	(35.0 - 53.8)
New Jersey	296	38.6	4.0	(30.8 - 46.4)	337	35.3	3.7	(28.0 - 42.6)
New Mexico	234	45.9	4.5	(37.1 - 54.7)	262	41.9	4.2	(33.7 - 50.1)
New York	276	42.7	4.2	(34.5 - 50.9)	336	35.4	3.7	(28.1 - 42.7)
North Dakota	98	46.7	5.7	(35.5 - 57.9)	120	36.7	5.1	(26.7 - 46.7)
Ohio	222	43.2	4.8	(33.8 - 52.6)	258	36.6	4.4	(28.0 - 45.2)
Oklahoma	250	53.0	3.9	(45.4 - 60.6)	256	52.4	3.9	(44.8 - 60.0)
Oregon	193	44.1	4.4	(35.5 - 52.7)	224	44.7	4.2	(36.5 - 52.9)
Pennsylvania	134	49.5	5.9	(37.9 - 61.1)	156	42.4	5.3	(32.0 - 52.8)
Rhode Island	153	51.5	4.8	(42.1 - 60.9)	180	47.8	4.4	(39.2 - 56.4)
Texas	479	41.5	3.8	(34.1 - 48.9)	558	34.9	3.3	(28.4 - 41.4)
Utah	368	49.4	3.7	(42.1 - 56.7)	414	46.2	3.5	(39.3 - 53.1)
Vermont	389	52.4	3.0	(46.5 - 58.3)	443	48.3	2.8	(42.8 - 53.8)
Virginia	88	44.7	6.6	(31.8 - 57.6)	100	42.6	6.2	(30.4 - 54.8)
Washington	341	49.0	3.5	(42.1 - 55.9)	389	44.4	3.3	(37.9 - 50.9)
West Virginia	193	48.6	4.3	(40.2 - 57.0)	213	45.3	4.0	(37.5 - 53.1)
Wisconsin	147	51.6	5.4	(41.0 - 62.2)	206	43.4	4.4	(34.8 - 52.0)
36 state total	9,390	46.8	1.1	(44.6 - 49.0)	10,793	41.5	1.1	(39.3 - 43.7)
Median		47.1				42.5		
Range		31.9-57.0				28.5-52.4		

¹ Aged < 18 years

² "Yes" response to "Do you still have asthma?"

³ Reported any of the following during the past year: asthma medication use, asthma symptoms, or MD visits for asthma

[±] Sample size is for multiple years of participation. Responses of "Don't Know" and "Refused" are excluded.

[§] Standard error

[†] Confidence interval