

Table C2_1
Child Current Asthma Prevalence Rate (Percent)
by Sex and State or Territory: BRFSS 2009

State	Male Sample Size	Male Prevalence	Male Standard Error	Male 95% CI *		Female Sample Size	Female Prevalence	Female Standard Error	Female 95% CI *
Total **	37,878	9.6	0.30	(9.0 - 10.2)		35,592	7.6	0.26	(7.1 - 8.1)
CA	2,379	7.4	0.75	(6.0 - 9.0)		2,179	6.0	0.68	(4.8 - 7.4)
CT	824	12.7	1.47	(10.1 - 15.9)		789	11.3	1.48	(8.7 - 14.5)
DE	592	15.9	2.01	(12.4 - 20.3)		548	11.1	1.62	(8.3 - 14.7)
DC	369	17.2	2.53	(12.8 - 22.8)		335	10.9	2.05	(7.4 - 15.6)
GA	783	11.2	1.55	(8.4 - 14.6)		753	8.7	1.37	(6.4 - 11.8)
HI	940	13.4	1.45	(10.8 - 16.5)		921	9.0	1.14	(7.0 - 11.5)
ID	706	4.5	0.89	(3.1 - 6.6)		701	4.7	0.94	(3.2 - 6.9)
IL	844	9.4	1.30	(7.2 - 12.3)		829	9.2	1.28	(7.0 - 12.0)
IN	1,142	11.1	1.29	(8.8 - 13.9)		1,056	8.6	1.08	(6.7 - 10.9)
IA	782	5.6	0.95	(4.0 - 7.8)		797	3.9	0.74	(2.7 - 5.6)
KS	2,616	9.1	0.64	(8.0 - 10.5)		2,350	7.4	0.57	(6.4 - 8.6)
LA	1,223	10.4	1.15	(8.4 - 12.9)		1,054	8.4	1.08	(6.5 - 10.8)
ME	515	10.0	1.41	(7.6 - 13.1)		483	9.4	1.49	(6.8 - 12.7)
MD	1,298	15.4	1.57	(12.6 - 18.8)		1,175	8.5	1.14	(6.5 - 11.1)
MA	735	9.2	1.52	(6.7 - 12.7)		666	9.2	1.45	(6.7 - 12.5)
MI	1,167	8.5	1.08	(6.6 - 10.9)		1,128	7.8	0.98	(6.1 - 10.0)
MS	1,373	11.3	1.04	(9.5 - 13.5)		1,254	9.5	1.07	(7.6 - 11.8)
MT	965	7.8	1.07	(5.9 - 10.2)		866	4.9	0.94	(3.4 - 7.1)
NE	1,929	5.2	0.75	(3.9 - 6.9)		1,821	5.2	0.88	(3.8 - 7.3)
NV	441	6.2	1.63	(3.7 - 10.3)		440	7.6	2.30	(4.2 - 13.5)
NJ	1,847	9.6	0.96	(7.9 - 11.6)		1,800	8.6	0.89	(7.0 - 10.5)
NM	782	9.3	1.47	(6.8 - 12.6)		722	7.4	1.20	(5.3 - 10.1)
NY	809	11.2	1.44	(8.7 - 14.4)		816	8.9	1.37	(6.6 - 12.0)
ND	617	5.4	0.86	(3.9 - 7.3)		598	6.2	1.17	(4.2 - 8.9)
OH	497	9.4	1.50	(6.8 - 12.8)		490	7.3	1.33	(5.1 - 10.4)
OK	497	11.2	1.63	(8.4 - 14.8)		440	10.5	1.70	(7.6 - 14.3)
OR	406	12.6	1.97	(9.2 - 17.0)		385	6.5	1.35	(4.3 - 9.7)
PA	1,099	12.4	1.62	(9.5 - 15.9)		953	8.3	1.29	(6.1 - 11.2)
RI	855	11.7	1.25	(9.5 - 14.4)		812	10.4	1.27	(8.1 - 13.1)

Notes:

* CI denotes confidence interval.

** Total includes data from 35 states plus the District of Columbia and excludes the territory, Puerto Rico.

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the sample size is less than 50, estimates are not precise and should be interpreted with caution.

Table C2_1
Child Current Asthma Prevalence Rate (Percent)
by Sex and State or Territory: BRFSS 2009

State	Male Sample Size	Male Prevalence	Male Standard Error	Male 95% CI *		Female Sample Size	Female Prevalence	Female Standard Error	Female 95% CI *
TX	1,719	9.4	1.15	(7.4 - 11.9)		1,648	6.9	0.96	(5.3 - 9.1)
UT	1,984	9.1	0.85	(7.6 - 10.9)		1,870	5.4	0.64	(4.3 - 6.8)
VT	837	11.6	1.30	(9.3 - 14.4)		797	8.7	1.19	(6.7 - 11.3)
VA	697	10.6	1.90	(7.4 - 15.0)		616	5.9	1.13	(4.0 - 8.5)
WA	2,542	7.8	0.63	(6.6 - 9.1)		2,478	5.9	0.57	(4.9 - 7.2)
WV	510	8.0	1.30	(5.8 - 11.0)		546	9.0	1.43	(6.6 - 12.2)
WI	557	6.4	1.65	(3.9 - 10.5)		476	7.4	1.72	(4.7 - 11.6)
Territories									
PR	568	14.8	1.74	(11.7 - 18.5)		482	9.1	1.62	(6.4 - 12.8)

Notes:

* CI denotes confidence interval.

** Total includes data from 35 states plus the District of Columbia and excludes the territory, Puerto Rico.

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the sample size is less than 50, estimates are not precise and should be interpreted with caution.