

Table L2_1
Child Lifetime Asthma Prevalence Rate (Percent)
by Sex and State or Territory: BRFSS 2009

State	Male Sample Size	Male Prevalence	Male Standard Error	Male 95% CI *		Female Sample Size	Female Prevalence	Female Standard Error	Female 95% CI *
Total **	38,051	15.4	0.36	(14.7 - 16.1)		35,716	11.1	0.30	(10.5 - 11.7)
CA	2,382	13.6	0.97	(11.8 - 15.6)		2,183	9.2	0.77	(7.8 - 10.8)
CT	826	19.8	1.77	(16.6 - 23.5)		793	14.7	1.63	(11.8 - 18.2)
DE	594	20.5	2.21	(16.5 - 25.2)		549	15.9	1.90	(12.5 - 20.0)
DC	373	26.2	2.82	(21.1 - 32.1)		337	14.5	2.37	(10.5 - 19.8)
GA	788	18.6	1.96	(15.0 - 22.7)		754	11.9	1.53	(9.2 - 15.3)
HI	945	21.6	1.77	(18.3 - 25.3)		927	14.0	1.38	(11.5 - 17.0)
ID	712	8.7	1.17	(6.7 - 11.3)		707	8.9	1.30	(6.7 - 11.8)
IL	845	14.6	1.57	(11.8 - 18.0)		830	11.5	1.40	(9.0 - 14.5)
IN	1,150	16.7	1.50	(14.0 - 19.9)		1,067	12.6	1.26	(10.3 - 15.3)
IA	784	8.3	1.17	(6.3 - 10.9)		799	5.2	0.84	(3.8 - 7.1)
KS	2,623	13.7	0.78	(12.2 - 15.3)		2,359	10.4	0.70	(9.1 - 11.9)
LA	1,228	17.0	1.40	(14.4 - 19.9)		1,055	11.8	1.23	(9.6 - 14.4)
ME	517	16.5	1.82	(13.3 - 20.4)		484	12.5	1.72	(9.5 - 16.3)
MD	1,309	21.4	1.72	(18.2 - 25.0)		1,183	12.8	1.32	(10.5 - 15.7)
MA	740	13.2	1.70	(10.2 - 16.9)		666	14.3	1.82	(11.1 - 18.2)
MI	1,175	12.5	1.21	(10.3 - 15.0)		1,131	10.3	1.10	(8.3 - 12.7)
MS	1,382	16.6	1.20	(14.4 - 19.1)		1,259	12.7	1.20	(10.5 - 15.2)
MT	967	11.0	1.27	(8.7 - 13.7)		869	7.3	1.07	(5.5 - 9.7)
NE	1,946	9.6	1.10	(7.7 - 12.0)		1,823	8.1	1.08	(6.2 - 10.5)
NV	442	10.8	2.21	(7.1 - 15.9)		441	10.6	2.45	(6.7 - 16.5)
NJ	1,850	15.6	1.13	(13.5 - 17.9)		1,806	12.9	1.06	(11.0 - 15.2)
NM	782	14.5	1.67	(11.5 - 18.1)		724	11.6	1.44	(9.0 - 14.7)
NY	814	18.7	1.74	(15.5 - 22.3)		820	14.4	1.64	(11.4 - 17.9)
ND	620	9.3	1.18	(7.2 - 11.9)		601	9.2	1.37	(6.9 - 12.3)
OH	498	15.9	1.96	(12.4 - 20.1)		491	9.4	1.46	(6.9 - 12.7)
OK	499	15.8	1.91	(12.4 - 19.9)		440	13.2	1.85	(10.0 - 17.3)
OR	411	19.8	2.22	(15.8 - 24.5)		386	9.7	1.68	(6.8 - 13.5)
PA	1,105	17.0	1.72	(13.9 - 20.7)		960	12.6	1.46	(10.0 - 15.8)
RI	863	16.6	1.44	(14.0 - 19.6)		815	15.0	1.47	(12.3 - 18.1)

Notes:
* CI denotes confidence interval.
** Total includes data from 35 states plus the District of Columbia and excludes the territory, Puerto Rico.

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the sample size is less than 50, estimates are not precise and should be interpreted with caution.

Table L2_1
Child Lifetime Asthma Prevalence Rate (Percent)
by Sex and State or Territory: BRFSS 2009

State	Male Sample Size	Male Prevalence	Male Standard Error	Male 95% CI *		Female Sample Size	Female Prevalence	Female Standard Error	Female 95% CI *
TX	1,730	15.9	1.38	(13.4 - 18.8)		1,651	10.6	1.12	(8.6 - 13.0)
UT	1,992	13.5	1.00	(11.6 - 15.6)		1,874	8.3	0.78	(6.9 - 9.9)
VT	840	17.0	1.49	(14.3 - 20.1)		803	14.3	1.46	(11.7 - 17.4)
VA	697	15.8	2.06	(12.2 - 20.3)		619	8.5	1.31	(6.2 - 11.4)
WA	2,553	13.0	0.80	(11.5 - 14.7)		2,485	9.4	0.71	(8.1 - 10.9)
WV	512	12.9	1.62	(10.1 - 16.5)		549	12.5	1.61	(9.7 - 16.0)
WI	557	10.5	1.90	(7.3 - 14.8)		476	9.2	1.84	(6.1 - 13.5)
Territories									
PR	569	27.6	2.14	(23.6 - 32.0)		482	19.3	2.10	(15.5 - 23.7)

Notes:
* CI denotes confidence interval.
** Total includes data from 35 states plus the District of Columbia and excludes the territory, Puerto Rico.

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the sample size is less than 50, estimates are not precise and should be interpreted with caution.