

**Table C2\_1**  
**Child Current Asthma Prevalence Rate (Percent)**  
**by Sex and State or Territory: BRFSS 2010**

State	Male Sample Size	Male Prevalence	Male Standard Error	Male 95% CI *		Female Sample Size	Female Prevalence	Female Standard Error	Female 95% CI *
Total **	36,498	9.8	0.33	(9.1 - 10.4)		34,217	7.0	0.29	(6.5 - 7.6)
AL	893	12.0	1.51	(9.4 - 15.3)		860	11.1	1.46	(8.6 - 14.3)
AZ	621	11.6	1.94	(8.3 - 16.0)		588	7.3	1.47	(4.9 - 10.8)
CA	834	7.5	1.01	(5.8 - 9.8)		798	4.4	0.85	(3.0 - 6.4)
CT	860	12.9	1.55	(10.1 - 16.3)		859	9.4	1.31	(7.2 - 12.3)
DC	373	22.2	3.13	(16.7 - 28.9)		344	13.8	2.57	(9.5 - 19.7)
GA	786	9.7	1.29	(7.4 - 12.5)		723	8.5	1.34	(6.3 - 11.6)
HI	853	13.0	1.46	(10.4 - 16.2)		794	9.2	1.33	(6.9 - 12.2)
IL	701	10.8	1.77	(7.8 - 14.8)		640	8.8	1.63	(6.1 - 12.5)
IN	1,185	8.3	0.98	(6.5 - 10.4)		1,125	9.3	1.24	(7.2 - 12.1)
IA	791	6.6	1.15	(4.7 - 9.2)		720	5.7	1.07	(3.9 - 8.2)
KS	1,171	7.8	0.88	(6.2 - 9.7)		1,044	7.3	0.93	(5.6 - 9.3)
KY	967	12.0	1.42	(9.5 - 15.1)		906	9.3	1.49	(6.8 - 12.7)
LA	983	8.8	1.10	(6.9 - 11.2)		879	7.9	1.09	(6.0 - 10.3)
ME	452	8.9	1.49	(6.3 - 12.3)		431	8.0	1.54	(5.4 - 11.6)
MD	1,219	13.9	1.43	(11.3 - 17.0)		1,194	9.9	1.25	(7.7 - 12.7)
MA	700	12.6	1.77	(9.5 - 16.4)		673	6.6	1.33	(4.4 - 9.7)
MI	1,013	12.3	1.43	(9.7 - 15.3)		979	9.9	1.26	(7.7 - 12.7)
MS	988	9.5	1.16	(7.5 - 12.0)		951	7.7	1.10	(5.8 - 10.1)
MO	650	12.0	2.01	(8.6 - 16.5)		600	10.0	1.70	(7.1 - 13.8)
MT	904	7.0	1.12	(5.1 - 9.6)		832	6.9	1.29	(4.8 - 9.9)
NE	1,940	6.7	1.12	(4.8 - 9.3)		1,920	5.4	0.92	(3.9 - 7.5)
NV	484	9.7	2.26	(6.1 - 15.1)		409	7.5	1.99	(4.4 - 12.4)
NJ	1,793	11.2	1.05	(9.3 - 13.4)		1,543	6.9	0.78	(5.5 - 8.6)
NM	859	9.2	1.24	(7.0 - 11.9)		826	6.8	1.03	(5.1 - 9.1)
NY	373	9.4	2.03	(6.1 - 14.2)		312	5.5	1.65	(3.0 - 9.8)
ND	602	8.6	1.44	(6.2 - 11.9)		547	4.3	0.90	(2.8 - 6.4)
OH	411	9.6	2.07	(6.3 - 14.5)		370	8.5	2.19	(5.1 - 13.9)
OK	1,022	11.3	1.19	(9.2 - 13.9)		979	9.2	1.15	(7.1 - 11.7)
OR	482	8.7	1.39	(6.3 - 11.8)		456	6.6	1.33	(4.4 - 9.7)

Notes:

\* CI denotes confidence interval.

\*\* Total includes data from 38 states plus the District of Columbia and excludes the territory, Puerto Rico.

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the sample size is less than 50, estimates are not precise and should be interpreted with caution.

**Table C2\_1**  
**Child Current Asthma Prevalence Rate (Percent)**  
**by Sex and State or Territory: BRFSS 2010**

State	Male Sample Size	Male Prevalence	Male Standard Error	Male 95% CI *		Female Sample Size	Female Prevalence	Female Standard Error	Female 95% CI *
PA	1,300	12.4	1.39	(10.0 - 15.4)		1,138	6.8	0.89	(5.3 - 8.8)
RI	779	11.9	1.38	(9.4 - 14.9)		758	11.8	1.46	(9.3 - 15.0)
TN	201	7.6	2.17	(4.3 - 13.1)		194	5.1	1.94	(2.4 - 10.5)
TX	2,559	9.1	1.08	(7.2 - 11.4)		2,433	6.1	0.99	(4.4 - 8.4)
UT	1,909	8.0	0.80	(6.6 - 9.7)		1,856	5.7	0.71	(4.5 - 7.2)
VT	782	10.1	1.45	(7.6 - 13.3)		779	9.7	1.19	(7.6 - 12.3)
WA	2,347	7.1	0.67	(5.9 - 8.5)		2,185	4.9	0.50	(4.0 - 6.0)
WV	486	6.8	1.28	(4.7 - 9.8)		415	6.0	1.17	(4.1 - 8.8)
WI	590	10.0	1.63	(7.2 - 13.6)		528	7.6	1.52	(5.1 - 11.2)
WY	635	7.5	1.10	(5.6 - 10.0)		629	5.7	1.04	(4.0 - 8.2)
Territories									
PR	398	17.4	2.30	(13.3 - 22.4)		380	7.2	1.59	(4.6 - 11.0)

Notes:

\* CI denotes confidence interval.

\*\* Total includes data from 38 states plus the District of Columbia and excludes the territory, Puerto Rico.

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the sample size is less than 50, estimates are not precise and should be interpreted with caution.