

**Table L2\_1**  
**Child Lifetime Asthma Prevalence Rate (Percent)**  
**by Sex and State or Territory: BRFSS 2010**

State	Male Sample Size	Male Prevalence	Male Standard Error	Male 95% CI *		Female Sample Size	Female Prevalence	Female Standard Error	Female 95% CI *
Total **	36,642	14.7	0.40	(13.9 - 15.5)		34,321	10.7	0.37	(10.0 - 11.4)
AL	895	14.6	1.58	(11.8 - 18.0)		861	14.5	1.63	(11.6 - 18.0)
AZ	624	14.5	2.16	(10.7 - 19.2)		589	10.4	1.93	(7.2 - 14.8)
CA	835	13.6	1.43	(11.1 - 16.7)		801	8.6	1.15	(6.6 - 11.1)
CT	862	17.7	1.72	(14.6 - 21.3)		863	12.8	1.49	(10.2 - 16.0)
DC	373	27.1	3.25	(21.2 - 33.9)		344	17.9	2.84	(13.0 - 24.1)
GA	788	16.3	1.63	(13.3 - 19.7)		723	12.9	1.56	(10.1 - 16.3)
HI	856	19.8	1.77	(16.6 - 23.5)		798	14.1	1.56	(11.3 - 17.4)
IL	701	14.6	1.92	(11.3 - 18.8)		642	12.4	1.76	(9.4 - 16.3)
IN	1,188	14.1	1.35	(11.7 - 17.0)		1,127	12.8	1.38	(10.3 - 15.8)
IA	793	10.7	1.37	(8.3 - 13.7)		720	6.7	1.14	(4.8 - 9.3)
KS	1,175	11.5	1.05	(9.6 - 13.7)		1,046	9.2	1.02	(7.4 - 11.4)
KY	971	16.2	1.55	(13.3 - 19.4)		911	13.2	1.88	(10.0 - 17.4)
LA	990	12.9	1.27	(10.6 - 15.6)		880	10.3	1.20	(8.2 - 12.9)
ME	457	15.5	1.87	(12.2 - 19.5)		434	10.9	1.69	(8.0 - 14.7)
MD	1,227	19.0	1.57	(16.1 - 22.2)		1,197	14.0	1.39	(11.5 - 17.0)
MA	702	17.1	2.07	(13.4 - 21.5)		675	9.9	1.55	(7.2 - 13.3)
MI	1,017	16.0	1.55	(13.2 - 19.3)		981	12.8	1.47	(10.2 - 15.9)
MS	990	14.7	1.45	(12.1 - 17.8)		954	11.7	1.34	(9.3 - 14.5)
MO	653	17.3	2.34	(13.1 - 22.4)		602	11.9	1.79	(8.8 - 15.9)
MT	909	11.0	1.34	(8.7 - 13.9)		834	8.3	1.44	(5.9 - 11.6)
NE	1,942	10.1	1.23	(7.9 - 12.8)		1,920	7.0	1.02	(5.3 - 9.3)
NV	484	11.5	2.33	(7.7 - 16.9)		409	9.7	2.12	(6.3 - 14.8)
NJ	1,802	17.5	1.25	(15.2 - 20.1)		1,546	11.0	1.05	(9.1 - 13.2)
NM	862	13.5	1.46	(10.9 - 16.6)		830	11.0	1.25	(8.8 - 13.7)
NY	374	16.2	2.49	(11.9 - 21.7)		314	10.5	2.46	(6.6 - 16.4)
ND	606	13.6	1.68	(10.6 - 17.2)		548	6.1	1.03	(4.4 - 8.5)
OH	412	13.7	2.32	(9.8 - 18.9)		372	12.4	2.63	(8.1 - 18.5)
OK	1,025	15.3	1.34	(12.8 - 18.1)		983	12.9	1.29	(10.5 - 15.6)
OR	486	13.7	1.67	(10.8 - 17.3)		458	10.0	1.55	(7.3 - 13.5)

Notes:  
\* CI denotes confidence interval.  
\*\* Total includes data from 38 states plus the District of Columbia and excludes the territory, Puerto Rico.

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the sample size is less than 50, estimates are not precise and should be interpreted with caution.

**Table L2\_1**  
**Child Lifetime Asthma Prevalence Rate (Percent)**  
**by Sex and State or Territory: BRFSS 2010**

State	Male Sample Size	Male Prevalence	Male Standard Error	Male 95% CI *		Female Sample Size	Female Prevalence	Female Standard Error	Female 95% CI *
PA	1,309	18.6	1.54	(15.8 - 21.8)		1,149	10.1	1.05	(8.2 - 12.4)
RI	783	15.8	1.53	(13.0 - 19.0)		766	14.7	1.56	(11.9 - 18.0)
TN	201	9.9	2.34	(6.2 - 15.5)		195	11.1	2.97	(6.4 - 18.3)
TX	2,574	13.8	1.30	(11.5 - 16.6)		2,439	9.4	1.11	(7.4 - 11.8)
UT	1,916	10.8	0.88	(9.2 - 12.7)		1,862	8.0	0.81	(6.5 - 9.7)
VT	784	15.4	1.62	(12.4 - 18.8)		781	12.9	1.32	(10.5 - 15.7)
WA	2,362	11.8	0.82	(10.3 - 13.6)		2,194	8.8	0.75	(7.5 - 10.4)
WV	486	11.3	1.66	(8.4 - 15.0)		415	9.8	1.76	(6.8 - 13.8)
WI	590	13.1	1.80	(10.0 - 17.1)		529	10.2	1.65	(7.4 - 13.9)
WY	638	10.6	1.30	(8.3 - 13.4)		629	8.0	1.16	(6.0 - 10.6)
Territories									
PR	398	27.8	2.65	(22.9 - 33.3)		380	17.2	2.18	(13.3 - 21.9)

Notes:  
\* CI denotes confidence interval.  
\*\* Total includes data from 38 states plus the District of Columbia and excludes the territory, Puerto Rico.

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the sample size is less than 50, estimates are not precise and should be interpreted with caution.