

Table C1
Child Current Asthma Prevalence
and Weighted Number by State or Territory: BRFSS 2016

State	Unweighted Number	Prevalence (Percent)	Standard Error	95% CI * (Percent)		Weighted Number	95% CI * (Number)
Total **	62,623	8.1	0.20	(7.7–8.4)		3,861,228	(3,673,998–4,048,458)
AZ	1,831	8.1	0.93	(6.4–10.1)		129,244	(99,588–158,900)
CA	2,584	7.7	0.64	(6.5–9.0)		588,421	(491,145–685,698)
CT	2,156	11.0	0.86	(9.4–12.8)		78,916	(66,607–91,226)
FL	6,709	7.8	0.61	(6.7–9.1)		304,053	(257,053–351,052)
GA	927	8.8	1.14	(6.9–11.3)		207,109	(154,413–259,806)
HI	1,642	10.2	1.00	(8.4–12.4)		30,630	(24,620–36,640)
IL	987	7.3	0.90	(5.7–9.3)		207,644	(157,688–257,599)
IN	2,122	8.2	0.72	(6.9–9.7)		124,718	(103,026–146,410)
KS	2,686	9.0	0.66	(7.8–10.3)		61,729	(52,770–70,689)
KY	2,169	7.8	0.77	(6.5–9.5)		74,500	(60,013–88,987)
ME	1,915	7.4	0.87	(5.8–9.3)		18,009	(13,784–22,235)
MA	709	9.6	1.39	(7.2–12.7)		125,763	(89,516–162,010)
MI	2,686	8.9	0.65	(7.7–10.3)		185,141	(158,580–211,703)
MN	3,798	6.9	0.50	(6.0–7.9)		85,117	(72,985–97,248)
MS	1,160	6.5	0.79	(5.1–8.3)		44,741	(34,162–55,321)
MO	1,321	8.7	1.01	(6.9–10.9)		115,545	(88,916–142,173)
MT	1,305	5.8	0.96	(4.2–8.0)		12,728	(8,569–16,888)
NE	3,672	6.4	0.63	(5.3–7.7)		29,825	(23,969–35,681)
NV	932	6.5	1.20	(4.5–9.3)		41,264	(26,240–56,287)
NJ	1,601	8.1	0.96	(6.4–10.2)		154,201	(118,085–190,317)
NM	1,346	9.9	1.17	(7.8–12.4)		47,112	(35,958–58,266)
NY	2,290	9.0	1.00	(7.2–11.2)		364,877	(284,905–444,850)
OH	2,421	6.9	0.69	(5.6–8.3)		171,051	(137,338–204,764)
OK	1,200	9.8	1.04	(7.9–12.0)		88,720	(69,932–107,508)
OR	1,076	5.6	0.76	(4.3–7.3)		46,438	(34,103–58,774)
PA	1,469	8.8	0.89	(7.2–10.7)		224,979	(180,196–269,762)
RI	1,129	9.4	1.25	(7.2–12.1)		18,846	(13,799–23,893)

Notes:

* CI denotes confidence interval.

** Total includes data from 32 states based on cellphone and landline data combined excludes the territory, Puerto Rico.

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the sample size is less than 50, estimates are not precise and should be interpreted with caution.

Table C1
Child Current Asthma Prevalence
and Weighted Number by State or Territory: BRFSS 2016

State	Unweighted Number	Prevalence (Percent)	Standard Error	95% CI * (Percent)		Weighted Number	95% CI * (Number)
UT	3,511	5.8	0.48	(5.0–6.9)		52,108	(43,615–60,600)
VT	1,266	8.3	1.24	(6.2–11.1)		9,397	(6,553–12,242)
WA	2,964	7.3	0.65	(6.1–8.7)		115,898	(95,101–136,696)
WI	1,039	8.3	1.27	(6.1–11.2)		102,501	(71,022–133,980)
Territory							
PR	1,176	15.5	1.58	(12.6–18.8)		123,102	(97,153–149,050)

Notes:

* CI denotes confidence interval.

** Total includes data from 32 states based on cellphone and landline data combined excludes the territory, Puerto Rico.

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the sample size is less than 50, estimates are not precise and should be interpreted with caution.