

Table C2_1
Child Current Asthma Prevalence
by Sex and State or Territory: BRFSS 2016

State	Male Unweighted Number	Male Prevalence	Male Standard Error	Male 95% CI *		Female Unweighted Number	Female Prevalence	Female Standard Error	Female 95% CI *
Total **	31,797	9.6	0.30	(9.1–10.3)		29,938	6.6	0.26	(6.1–7.1)
AZ	970	9.9	1.38	(7.5–12.9)		839	6.2	1.25	(4.2–9.2)
CA	1,339	9.5	0.97	(7.8–11.6)		1,240	6.2	0.84	(4.7–8.0)
CT	1,091	12.4	1.25	(10.2–15.1)		1,025	9.8	1.20	(7.7–12.4)
FL	3,410	8.8	0.89	(7.2–10.7)		3,160	7.1	0.86	(5.6–9.0)
GA	474	10.9	1.79	(7.9–14.9)		435	7.1	1.44	(4.8–10.5)
HI	837	11.3	1.42	(8.8–14.4)		787	9.5	1.44	(7.1–12.8)
IL	489	9.7	1.52	(7.1–13.1)		489	5.0	0.96	(3.4–7.2)
IN	1,080	9.1	1.07	(7.3–11.5)		1,020	7.3	0.98	(5.6–9.5)
KS	1,366	11.2	1.04	(9.4–13.4)		1,290	6.8	0.81	(5.4–8.6)
KY	1,058	9.1	1.15	(7.1–11.6)		1,091	6.7	1.06	(4.9–9.1)
ME	975	8.4	1.27	(6.2–11.3)		930	6.1	1.16	(4.1–8.8)
MA	365	12.9	2.29	(9.0–18.1)		324	6.2	1.55	(3.8–10.0)
MI	1,350	11.4	1.07	(9.5–13.7)		1,310	6.4	0.71	(5.2–7.9)
MN	1,943	7.6	0.71	(6.3–9.1)		1,780	6.3	0.70	(5.0–7.8)
MS	590	8.6	1.30	(6.4–11.5)		552	4.6	0.91	(3.1–6.7)
MO	651	9.0	1.43	(6.6–12.2)		653	8.4	1.48	(5.9–11.7)
MT	666	5.5	1.17	(3.7–8.3)		624	6.1	1.53	(3.7–9.9)
NE	1,841	8.0	1.04	(6.2–10.3)		1,783	4.9	0.72	(3.6–6.5)
NV	459	8.6	2.04	(5.4–13.6)		459	4.4	1.19	(2.6–7.4)
NJ	806	9.6	1.41	(7.1–12.7)		760	6.9	1.37	(4.7–10.1)
NM	703	11.1	1.75	(8.1–15.0)		616	9.0	1.60	(6.3–12.7)
NY	1,170	10.9	1.60	(8.1–14.4)		1,101	7.3	1.24	(5.2–10.2)
OH	1,202	7.8	1.06	(6.0–10.2)		1,177	5.9	0.89	(4.3–7.9)
OK	587	12.3	1.65	(9.5–16.0)		602	7.3	1.26	(5.2–10.2)
OR	552	6.1	1.10	(4.2–8.6)		505	5.3	1.10	(3.6–7.9)
PA	736	10.2	1.39	(7.8–13.3)		714	7.5	1.13	(5.6–10.1)
RI	558	9.7	1.77	(6.7–13.7)		550	9.2	1.81	(6.2–13.4)
UT	1,830	6.9	0.75	(5.6–8.5)		1,647	4.8	0.62	(3.7–6.1)
VT	673	8.9	1.43	(6.5–12.1)		573	6.9	1.98	(3.9–11.9)
WA	1,497	8.5	1.02	(6.7–10.8)		1,408	5.9	0.78	(4.6–7.6)

Notes:

* CI denotes confidence interval.

** Total includes data from 32 states based on cellphone and landline data combined excludes the territory, Puerto Rico.

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the sample size is less than 50, estimates are not precise and should be interpreted with caution.

Table C2_1
Child Current Asthma Prevalence
by Sex and State or Territory: BRFSS 2016

State	Male Unweighted Number	Male Prevalence	Male Standard Error	Male 95% CI *	Female Unweighted Number	Female Prevalence	Female Standard Error	Female 95% CI *
WI	529	9.3	1.96	(6.1–13.9)	494	7.5	1.69	(4.7–11.5)
Territory								
PR	616	17.1	2.29	(13.0–22.0)	550	14.2	2.20	(10.4–19.1)

Notes:

* CI denotes confidence interval.

** Total includes data from 32 states based on cellphone and landline data combined excludes the territory, Puerto Rico.

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the sample size is less than 50, estimates are not precise and should be interpreted with caution.