

Table 2-2
Standard Errors* for Lifetime Prevalence Percents
by Age, United States: National Health Interview Survey, 2005

| | All ages | Children | Adults | Age (years) | | | | | | | |
|---|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|
| | | | | 15 - 34 | | | | | | | |
| | Total | Age <18 | Age 18+ | 0 - 4 | 5 - 14 | 15 - 34 | 15 - 19 | 20 - 24 | 25 - 34 | 35 - 64 | 65 + |
| Total: | 0.18 | 0.36 | 0.22 | 0.57 | 0.48 | 0.40 | 0.96 | 0.85 | 0.48 | 0.27 | 0.44 |
| Male | 0.26 | 0.53 | 0.31 | 0.96 | 0.76 | 0.56 | 1.37 | 1.16 | 0.70 | 0.37 | 0.64 |
| Female | 0.25 | 0.50 | 0.30 | 0.70 | 0.63 | 0.56 | 1.40 | 1.22 | 0.66 | 0.40 | 0.61 |
| White Non-Hispanic: | 0.23 | 0.47 | 0.27 | 0.61 | 0.64 | 0.52 | 1.24 | 1.10 | 0.68 | 0.33 | 0.51 |
| Male | 0.32 | 0.70 | 0.38 | 1.00 | 0.96 | 0.74 | 1.74 | 1.51 | 1.00 | 0.46 | 0.72 |
| Female | 0.32 | 0.66 | 0.37 | 0.79 | 0.82 | 0.73 | 1.81 | 1.56 | 0.95 | 0.47 | 0.70 |
| Black Non-Hispanic: | 0.53 | 1.05 | 0.54 | 2.08 | 1.28 | 1.05 | 2.49 | 2.27 | 1.00 | 0.72 | 1.41 |
| Male | 0.82 | 1.70 | 0.81 | 3.62 | 2.07 | 1.63 | 3.50 | 2.98 | 1.54 | 1.01 | 1.89 |
| Female | 0.65 | 1.34 | 0.76 | 2.29 | 1.87 | 1.34 | 3.14 | 3.45 | 1.34 | 1.05 | 1.98 |
| Other Races Non-Hispanic: | 0.88 | 1.59 | 1.15 | 2.69 | 2.20 | 2.03 | 4.79 | 5.22 | 1.86 | 1.09 | 2.29 |
| Male | 1.26 | 2.54 | 1.80 | 4.40 | 3.55 | 3.05 | 8.66 | 5.18 | 1.78 | 1.50 | 3.61 |
| Female | 1.33 | 2.20 | 1.67 | 3.17 | 3.16 | 2.94 | 6.69 | 8.38 | 3.08 | 1.69 | 3.11 |
| Hispanic: | 0.38 | 0.70 | 0.45 | 1.12 | 0.94 | 0.68 | 1.49 | 1.26 | 0.78 | 0.62 | 1.28 |
| Male | 0.52 | 1.00 | 0.55 | 1.62 | 1.41 | 0.77 | 1.71 | 1.71 | 0.99 | 0.67 | 2.22 |
| Female | 0.59 | 0.99 | 0.68 | 1.68 | 1.14 | 1.03 | 2.59 | 1.93 | 1.13 | 1.03 | 1.58 |
| Puerto Rican: | 1.49 | 3.21 | 1.72 | 4.02 | 4.36 | 3.14 | 7.06 | 5.95 | 4.40 | 3.33 | 3.81 |
| Male | 2.80 | 5.01 | 2.88 | 6.89 | 6.76 | 4.69 | 9.63 | 8.40 | 6.76 | 4.44 | 6.35 |
| Female | 1.99 | 3.19 | 2.44 | 5.18 | 5.01 | 3.78 | 7.75 | 8.11 | 5.52 | 4.57 | 4.15 |
| Mexican/Mexican-American: | 0.46 | 0.87 | 0.51 | 1.37 | 1.19 | 0.81 | 1.51 | 1.57 | 0.85 | 0.59 | 1.95 |
| Male | 0.62 | 1.09 | 0.70 | 1.87 | 1.65 | 0.97 | 1.70 | 2.02 | 1.10 | 0.70 | 3.50 |
| Female | 0.67 | 1.31 | 0.71 | 2.13 | 1.51 | 1.22 | 2.91 | 2.43 | 1.27 | 0.99 | 2.13 |
| Region: | | | | | | | | | | | |
| Northeast | 0.40 | 0.81 | 0.52 | 1.40 | 1.07 | 0.94 | 2.50 | 2.54 | 1.15 | 0.62 | 0.81 |
| Midwest | 0.38 | 0.79 | 0.42 | 1.11 | 1.08 | 0.78 | 2.01 | 1.58 | 0.94 | 0.48 | 0.89 |
| South | 0.32 | 0.62 | 0.39 | 1.02 | 0.82 | 0.68 | 1.52 | 1.38 | 0.87 | 0.45 | 0.77 |
| West | 0.36 | 0.69 | 0.44 | 1.04 | 0.90 | 0.81 | 1.84 | 1.68 | 0.86 | 0.65 | 1.10 |
| Ratio of Family Income to Poverty Threshold: | | | | | | | | | | | |
| 0 - 0.99 | 0.56 | 0.85 | 0.68 | 1.42 | 1.22 | 1.00 | 2.42 | 1.77 | 0.97 | 0.97 | 1.26 |
| 1.00 - 2.49 | 0.35 | 0.68 | 0.40 | 1.15 | 0.92 | 0.75 | 1.95 | 1.30 | 0.92 | 0.61 | 0.68 |
| 2.50 - 4.49 | 0.39 | 0.78 | 0.43 | 1.15 | 1.00 | 0.89 | 1.88 | 2.19 | 1.13 | 0.56 | 0.82 |
| 4.50 and above | 0.36 | 0.74 | 0.41 | 0.86 | 1.04 | 0.83 | 1.84 | 2.47 | 0.97 | 0.44 | 1.06 |

Source: National Health Interview Survey, National Center for Health Statistics, Centers for Disease Control and Prevention

compiled 5/22/2007

* See the "Methods" section of the documentation found at "<http://www.cdc.gov/asthma/NHIS/default.htm>" for information regarding standard errors.

Air Pollution and Respiratory Health Branch, National Center for Environmental Health
Centers for Disease Control and Prevention