Table 3. Recommended Oseltamivir and Peramivir Dose Adjustments for Treatment or Chemoprophylaxis of Influenza in Adult Patients with Renal Impairment or End Stage Renal Disease (ESRD) on Dialysis*

	Creatinine Clearance	Recommended Treatment Regimen	Recommended Chemoprophylaxis Regimen
Oral Oseltamivir ¹	Creatinine clearance 61 to 90 mL/min	75 mg twice a day	75 mg once daily
	Creatinine clearance 31 to 60 mL/min	30 mg twice a day	30 mg once daily
	Creatinine clearance 11 to 30 mL/min	30 mg once daily	30 mg every other day
	ESRD Patients on Hemodialysis Creatinine clearance ≤10 mL/min	30 mg after every hemodialysis cycle. Treatment duration not to exceed 5 days ²	30 mg after alternate hemodialysis cycles ³
	ESRD Patients on Continuous Ambulatory Peritoneal Dialysis⁴ Creatinine clearance ≤10 mL/min	A single 30 mg dose administered immediately after a dialysis exchange	30 mg once weekly immediately after dialysis exchange
Intravenous Peramivir (single dose) ⁵	Creatinine clearance ≥50 mL/min	600 mg	N/A
	Creatinine clearance 30 to 49 mL/min	200 mg	N/A
	Creatinine clearance 10 to 29 mL/min	100 mg	N/A
	ESRD Patients on Hemodialysis	Dose administered after dialysis at a dose adjusted based on creatinine clearance	

^{*} From package inserts for oseltamivir and peramivir; see FDA Influenza (Flu) Antiviral Drugs and Related Information.

Abbreviations: N/A = approved, not recommended

- 1. Renal dosing of oseltamivir is not available in the <u>package insert</u> for pediatric patients. However, these tables may be useful for children who qualify for adult doses based on weight >40 kg.
- 2. Assuming 3 hemodialysis sessions are performed in the 5- day period. Treatment can be initiated immediately if influenza symptoms develop during the 48 hours between hemodialysis sessions; however, the post-hemodialysis dose should still be administered independently of time of administration of the initial dose.
- 3. An initial dose can be administered prior to the start of dialysis.
- 4. Data derived from studies in continuous ambulatory peritoneal dialysis (CAPD) patients.
- 5. Renal dosing from peramivir <u>package insert</u> is available for pediatric patients: Creatinine clearance ≥50 mL/min: 12 mg/kg (up to maximum dose of 600 mg); Creatinine clearance 30 to 49 mL/min: 4 mg/kg; Creatinine clearance 10 to 29 mL/min: 2 mg/kg.