

Suicide Rates by Major Occupational Group — 17 States, 2012 and 2015

Cora Peterson, PhD¹; Deborah M. Stone, ScD²; Suzanne M. Marsh, MPA³; Pamela K. Schumacher⁴; Hope M. Tiesman, PhD³; Wendy LiKamWa McIntosh, MPH²; Colby N. Lokey, MS²; Aimée-Rika T. Trudeau, MPH²; Brad Bartholow, PhD²; Feijun Luo, PhD¹

During 2000–2016, the suicide rate among the U.S. working age population (persons aged 16–64 years) increased 34%, from 12.9 per 100,000 population to 17.3 (<https://www.cdc.gov/injury/wisqars>). To better understand suicide among different occupational groups and inform suicide prevention efforts, CDC analyzed suicide deaths by Standard Occupational Classification (SOC) major groups for decedents aged 16–64 years from the 17 states participating in both the 2012 and 2015 National Violent Death Reporting System (NVDRS) (<https://www.cdc.gov/violenceprevention/nvdrs>). The occupational group with the highest male suicide rate in 2012 and 2015 was Construction and Extraction (43.6 and 53.2 per 100,000 civilian noninstitutionalized working persons, respectively), whereas the group with the highest female suicide rate was Arts, Design, Entertainment, Sports, and Media (11.7 [2012] and 15.6 [2015]). The largest suicide rate increase among males from 2012 to 2015 (47%) occurred in the Arts, Design, Entertainment, Sports, and Media occupational group (26.9 to 39.7) and among females, in the Food Preparation and Serving Related group, from 6.1 to 9.4 (54%). CDC's technical package of strategies to prevent suicide is a resource for communities, including workplace settings (1).

NVDRS combines data on all violent deaths (defined as those resulting from the intentional use of physical force or power, threatened or actual, against oneself, another person, or a group or community), including suicide, based on death certificates, coroner/medical examiner reports, and law enforcement reports. Data on usual lifetime occupation among 22,053 suicide decedents aged 16–64 years from the 17 states* that participated in NVDRS in 2012 and 2015 were analyzed. CDC's National Institute for Occupational Safety and Health Industry and Occupation Computerized Coding System (NIOCCS 3.0)

(<https://wwwn.cdc.gov/nioccs3>) was used to assign 2010 U.S. Census civilian occupation and industry codes to NVDRS decedent records based on decedents' usual lifetime occupation and industry as reported on the death certificate. Results are reported by 2010 SOC major groups, converted from U.S. Census codes by NIOCCS. Records that could not be coded by NIOCCS were manually coded using the NIOCCS computer-assisted feature. All coding assignments were reviewed by industry and occupation coding experts for accuracy.

Suicide counts are presented by year, sex, and usual lifetime occupational group. Suicide rates were calculated using annual

INSIDE

- 1261 Global Routine Vaccination Coverage — 2017
- 1265 Progress Toward Global Eradication of Dracunculiasis — January 2017–June 2018
- 1271 Updated Framework for Development of Evidence-Based Recommendations by the Advisory Committee on Immunization Practices
- 1273 Increase in Acute Flaccid Myelitis — United States, 2018
- 1276 Notes from the Field: Use of Electronic Cigarettes and Any Tobacco Product Among Middle and High School Students — United States, 2011–2018
- 1278 Retraction Notice: Suicide Rates by Occupational Group — 17 States, 2012
- 1279 Correction and Republication: Prevalence and Characteristics of Autism Spectrum Disorder Among Children Aged 8 Years — Autism and Developmental Disabilities Monitoring Network, 11 Sites, United States, 2012
- 1281 QuickStats

Continuing Education examination available at https://www.cdc.gov/mmwr/cme/conted_info.html#weekly.

*Alaska, Colorado, Georgia, Kentucky, Maryland, Massachusetts, New Jersey, New Mexico, North Carolina, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, Utah, Virginia, and Wisconsin.



civilian noninstitutionalized working population counts by occupational group (based on longest job held during the previous calendar year) from the Current Population Survey Annual Social and Economic Supplement (2) as the denominator; 95% confidence intervals (CIs) were calculated using the National Center for Health Statistics method for death rates (3). The rate change from 2012 to 2015 is presented for each occupational group by sex, as is each group's rank for rate change (i.e., where rank position 1 signifies the greatest suicide rate increase). Decedents whose NVDRS data from coroner/medical examiner reports or law enforcement reports indicated that the decedent was not employed at the time of death (unemployed, disabled, incarcerated, homemaker, or student) were excluded from rate calculations, as were decedents with military or unpaid occupations, and those with insufficient information to classify occupation. Separate analyses of suicide deaths among males in agriculture-related SOC detailed groups were conducted; such rates were not calculated for female decedents because of small numbers.

NIOCCS classified 83% (8,858 in 2012 and 9,508 in 2015) of decedent records (Table 1); this count includes those that the NIOCCS program determined to have insufficient information to classify occupation. After expert review of NIOCCS automated code assignments, 231 (3%) of 2012 records and 290 (3%) of 2015 records were recoded. The remaining 1,799 (2012) and 1,888 (2015) (17% for both years) records were coded using the NIOCCS computer-assisted feature. For 2012 and 2015 combined, 5,089 (23%) decedents were not included

in suicide rate calculations because they were in the military, had unpaid occupations (e.g., did not work, homemaker, or student), or had insufficient information to classify lifetime occupation. Another 2,236 (10%) were excluded because they were not employed at the time of death.

In both 2012 and 2015, the largest percentage of male suicides (19%–20% of decedents) occurred among those in the Construction and Extraction group (SOC 47) (Table 2); the largest percentage of female suicides in both years occurred among decedents with unpaid occupations (29%). The largest percentage of female suicides among classifiable occupations occurred in the Office and Administrative Support group (SOC 43) in both years (15%). In both years, the highest suicide rates among males were in the Construction and Extraction group (43.6 in 2012 and 53.2 in 2015 per 100,000 civilian noninstitutionalized working persons) (Table 3). Among females, the highest suicide rates in both years were in the Arts, Design, Entertainment, Sports, and Media group (SOC 27) (11.7 in 2012 and 15.6 in 2015). Among males, the largest suicide rate increase from 2012 to 2015 (47%) occurred in the Arts, Design, Entertainment, Sports, and Media group (from 26.9 to 39.7), and among females (54%) in the Food Preparation and Serving Related group (SOC 35) (from 6.1 to 9.4). Rate changes among females in six SOC major groups were not reported because of small numbers (≤ 20 decedents in one or both years).

The 2012 and 2015 male suicide rates among Farmers, Ranchers, and Other Agricultural Managers (SOC 11–9013,

The *MMWR* series of publications is published by the Center for Surveillance, Epidemiology, and Laboratory Services, Centers for Disease Control and Prevention (CDC), U.S. Department of Health and Human Services, Atlanta, GA 30329-4027.

Suggested citation: [Author names; first three, then et al., if more than six.] [Report title]. *MMWR Morb Mortal Wkly Rep* 2018;67:[inclusive page numbers].

Centers for Disease Control and Prevention

Robert R. Redfield, MD, *Director*
 Anne Schuchat, MD, *Principal Deputy Director*
 Leslie Dauphin, PhD, *Acting Associate Director for Science*
 Barbara Ellis, PhD, MS, *Acting Director, Office of Science Quality*
 Chesley L. Richards, MD, MPH, *Deputy Director for Public Health Scientific Services*
 William R. MacKenzie, MD, *Acting Director, Center for Surveillance, Epidemiology, and Laboratory Services*

MMWR Editorial and Production Staff (Weekly)

Charlotte K. Kent, PhD, MPH, *Acting Editor in Chief, Executive Editor*
 Jacqueline Gindler, MD, *Editor*
 Mary Dott, MD, MPH, *Online Editor*
 Teresa F. Rutledge, *Managing Editor*
 Douglas W. Weatherwax, *Lead Technical Writer-Editor*
 Glenn Damon, Soumya Dunworth, PhD, Teresa M. Hood, MS,
Technical Writer-Editors

Martha F. Boyd, *Lead Visual Information Specialist*
 Maureen A. Leahy, Julia C. Martinroe,
 Stephen R. Spriggs, Tong Yang,
Visual Information Specialists
 Quang M. Doan, MBA, Phyllis H. King,
 Terraye M. Starr, Moua Yang,
Information Technology Specialists

MMWR Editorial Board

Timothy F. Jones, MD, *Chairman*
 Robin Ikeda, MD, MPH
 Phyllis Meadows, PhD, MSN, RN
 Jewel Mullen, MD, MPH, MPA
 Jeff Niederdeppe, PhD
 Patricia Quinlisk, MD, MPH
 Matthew L. Boulton, MD, MPH
 Virginia A. Caine, MD
 Katherine Lyon Daniel, PhD
 Jonathan E. Fielding, MD, MPH, MBA
 David W. Fleming, MD
 William E. Halperin, MD, DrPH, MPH

Stephen C. Redd, MD,
 Patrick L. Remington, MD, MPH
 Carlos Roig, MS, MA
 William Schaffner, MD
 Morgan Bobb Swanson, BS

TABLE 1. Procedure for Identification of analysis cohort of suicide decedents, by occupation — National Violent Death Reporting System (NVDRS) — 17 U.S. states* 2012 and 2015

Analytic procedure	2012 no.	2015 no.
Suicide decedents obtained from NVDRS data set [†]	12,811	13,967
Excluded before assignment of occupation code		
Aged <16 years or >64 years or missing sex	2,154	2,571
Assignment of occupation code		
Assigned based on decedent usual lifetime occupation	10,657	11,396
Autocoded by NIOCCS	8,858	9,508
Manually reassigned using the NIOCCS computer-assisted feature	231	290
Manually assigned using the NIOCCS computer-assisted feature	1,799	1,888
Rate analysis		
Decedents presumed to be in the labor force at time of death [§]	6,881	7,847

Abbreviation: NIOCCS = National Institute for Occupational Safety and Health Industry and Occupation Computerized Coding System.

* Alaska, Colorado, Georgia, Kentucky, Maryland, Massachusetts, New Jersey, New Mexico, North Carolina, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, Utah, Virginia, and Wisconsin.

[†] The total number of deaths (including suicides) reported in the NVDRS data set (June 2018) for the 17 U.S. states analyzed included 19,885 decedents in 2012 and 21,884 decedents in 2015.

[§] Because annual U.S. population count data by occupational group reflects the longest held job in the previous year among civilian, noninstitutionalized persons, suicide decedents whose NVDRS data from coroner/medical examiner or law enforcement reports indicated no employment (e.g., retired, unemployed, disabled, incarcerated, home maker, or student) at the time of death were excluded from rate analysis. Decedents were excluded if NVDRS current occupation information contained any of the following: “student, unemp*, not empl*, laid off, retir*, disab*, incarcer*, inmate, prisoner, homemaker, home maker, housewife, house wife, never worked, or not working.” Manual review of records excluded based on these criteria resulted in five records being reinstated (e.g., “student teacher” and two occupations listed for a decedent with just one occupation explicitly identified as retired).

a subgroup of the SOC 11 Management major group) were 44.9 (CI = 34.2–57.9) and 32.2 (CI = 24.2–42.0) per 100,000, based on 59 and 54 suicides in 2012 and 2015, respectively. The 2012 and 2015 male suicide rates for Agricultural Workers (SOC 45–2000, a subgroup of the SOC 45 Farming, Fishing, and Forestry major group) were 20.4 (CI = 13.8–29.1) and 17.3 (CI = 12.1–23.9), based on 30 and 36 suicides in 2012 and 2015, respectively.

Discussion

Suicide rates varied widely across occupational groups in both 2012 and 2015, and rates among males and females increased in many occupational groups. The etiology of suicide is multifactorial, and identifying the specific role that occupational factors might play in suicide risk is complicated; both work (e.g., little job control or job insecurity) and nonwork (e.g., relationship conflict) factors are associated with psychological distress and suicide (4). The relationship between occupation and suicide might be confounded by access to lethal means on the job and socioeconomic factors such as lower income

TABLE 2. Number and percentage of suicide decedents* in Standard Occupational Classification (SOC) major group, by year and sex — National Violent Death Reporting System, 17 states,† 2012 and 2015

SOC code	Occupational group	Male		Female	
		2012 no. (%)	2015 no. (%)	2012 no. (%)	2015 no. (%)
11	Management	534 (8)	611 (9)	117 (7)	118 (7)
13	Business and Financial Operations	155 (2)	145 (2)	81 (5)	84 (5)
15	Computer and Mathematical	208 (3)	237 (3)	22 (1)	32 (2)
17	Architecture and Engineering	172 (3)	167 (2)	10 (1)	15 (1)
19	Life, Physical, and Social Science	56 (1)	52 (1)	15 (1)	21 (1)
21	Community and Social Service	41 (1)	48 (1)	39 (2)	40 (2)
23	Legal	54 (1)	49 (1)	34 (2)	29 (2)
25	Education, Training, and Library	91 (1)	87 (1)	82 (5)	84 (5)
27	Arts, Design, Entertainment, Sports, and Media	140 (2)	186 (3)	54 (3)	76 (4)
29	Health Care Practitioners and Technical occupations	145 (2)	169 (2)	220 (14)	225 (12)
31	Health Care Support	35 (1)	34 (<1)	97 (6)	124 (7)
33	Protective Service	232 (4)	226 (3)	29 (2)	32 (2)
35	Food Preparation and Serving Related	214 (3)	301 (4)	112 (7)	154 (9)
37	Building and Grounds Cleaning and Maintenance	316 (5)	315 (4)	36 (2)	46 (3)
39	Personal Care and Service	81 (1)	85 (1)	98 (6)	102 (6)
41	Sales and Related	555 (9)	553 (8)	170 (11)	212 (12)
43	Office and Administrative Support	244 (4)	260 (4)	234 (15)	268 (15)
45	Farming, Fishing, and Forestry	68 (1)	71 (1)	7 (<1)	5 (<1)
47	Construction and Extraction	1,216 (19)	1,404 (20)	12 (1)	17 (1)
49	Installation, Maintenance, and Repair	549 (9)	621 (9)	8 (1)	NR
51	Production	605 (9)	679 (10)	64 (4)	81 (4)
53	Transportation and Material Moving	736 (11)	817 (11)	52 (3)	39 (2)
NA	Military	228 (3)	203 (2)	15 (1)	13 (<1)
NA	Unpaid	822 (10)	913 (11)	724 (29)	795 (29)
NA	Insufficient Information to Classify Occupation	651 (8)	425 (5)	177 (9)	123 (4)

Abbreviation: NA = not assigned; NR = not reported due to cell size <5.

* Aged 16–64 years.

[†] Alaska, Colorado, Georgia, Kentucky, Maryland, Massachusetts, New Jersey, New Mexico, North Carolina, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, Utah, Virginia, and Wisconsin.

and education (5,6). Previous studies have employed a range of methodologies to study the proposed association between suicide and occupation and, at times, have arrived at different conclusions. For example, although this analysis aligns with another that found high suicide rates among construction workers in Colorado (7), a meta-analysis using an international occupational classification system found persons in other less-skilled occupations, such as laborers and cleaners, to be at higher risk (6).

A better understanding of how suicides are distributed by occupational group might help inform prevention programs and policies. Because many adults spend a substantial amount of their time at work, the workplace is an important but underutilized location for suicide prevention (8). Workplaces could potentially benefit from suicide prevention activities.

TABLE 3. Suicide rate per 100,000 civilian, noninstitutionalized working persons aged 16–64 years, by sex, based on suicide decedents (N = 14,728) presumed in the labor force at time of death using Standard Occupational Classification (SOC) major groups — National Violent Death Reporting System, 17 states,* 2012 and 2015

Males					Females						
SOC code	Occupational group	2012	2015	Rate change		SOC code	Occupational group	2012	2015	Rate change	
				%	Rank [†]					%	Rank [†]
47	Construction and Extraction					27	Arts, Design, Entertainment, Sports, and Media				
	Rate rank [§]	1	1	+22%	5		Rate rank [§]	1	1	+34%	2
	Rate per 100,000	43.6	53.2				Rate per 100,000	11.7	15.6		
	95% CI [¶]	40.9–46.3	50.2–56.1				95% CI [¶]	8.6–15.5	12.1–19.8		
	Suicide decedents, no.	1,009	1,248				Suicide decedents, no.	47	67		
	Population, no.	2,313,934	2,345,952				Population, no.	403,305	429,424		
27	Arts, Design, Entertainment, Sports, and Media					33	Protective Service				
	Rate rank	7	2	+47%	1		Rate rank	2	2	+5%	9
	Rate per 100,000	26.9	39.7				Rate per 100,000	11.6	12.2		
	95% CI	22.1–31.8	33.6–45.8				95% CI	7.5–17.1	8.1–17.7		
	Suicide decedents, no.	117	162				Suicide decedents, no.	25	28		
	Population, no.	434,177	408,113				Population, no.	215,345	228,862		
49	Installation, Maintenance, and Repair					31	Health Care Support				
	Rate rank	2	3	+24%	3		Rate rank	5	3	+31%	3
	Rate per 100,000	31.6	39.1				Rate per 100,000	8.4	11.0		
	95% CI	28.7–34.4	35.8–42.3				95% CI	6.7–10.4	8.9–13.0		
	Suicide decedents, no.	473	542				Suicide decedents, no.	83	108		
	Population, no.	1,498,263	1,387,681				Population, no.	993,407	984,369		
53	Transportation and Material Moving					35	Food Preparation and Serving Related				
	Rate rank	4	4	+9%	8		Rate rank	11	4	+54%	1
	Rate per 100,000	28.4	30.9				Rate per 100,000	6.1	9.4		
	95% CI	26.2–30.7	28.6–33.1				95% CI	4.9–7.5	7.8–11.0		
	Suicide decedents, no.	615	721				Suicide decedents, no.	94	139		
	Population, no.	2,164,530	2,336,133				Population, no.	1,539,199	1,479,822		
51	Production					23	Legal				
	Rate rank	3	5	+7%	10		Rate rank	3	5	-17%	15
	Rate per 100,000	28.4	30.5				Rate per 100,000	11.1	9.2		
	95% CI	26.0–30.9	28.1–33.0				95% CI	7.5–15.9	5.8–13.9		
	Suicide decedents, no.	524	607				Suicide decedents, no.	30	22		
	Population, no.	1,843,879	1,987,864				Population, no.	269,243	238,870		
33	Protective Service					29	Health Care Practitioners and Technical				
	Rate rank	6	6	+4%	11		Rate rank	4	6	-13%	13
	Rate per 100,000	27.1	28.2				Rate per 100,000	10.3	9.0		
	95% CI	23.3–30.9	24.2–32.1				95% CI	8.9–11.8	7.7–10.3		
	Suicide decedents, no.	198	194				Suicide decedents, no.	195	193		
	Population, no.	730,044	689,034				Population, no.	1,890,885	2,140,217		
37	Building and Grounds Cleaning and Maintenance					51	Production				
	Rate rank	5	7	-2%	14		Rate rank	7	7	+18%	6
	Rate per 100,000	27.3	26.8				Rate per 100,000	7.6	9.0		
	95% CI	24.1–30.5	23.6–30.0				95% CI	5.8–10.0	7.0–11.3		
	Suicide decedents, no.	281	276				Suicide decedents, no.	55	72		
	Population, no.	1,028,779	1,029,385				Population, no.	719,183	800,640		
29	Health Care Practitioners and Technical					39	Personal Care and Service				
	Rate rank	14	8	+23%	4		Rate rank	9	8	+14%	7
	Rate per 100,000	20.8	25.6				Rate per 100,000	6.8	7.7		
	95% CI	17.1–24.6	21.5–29.8				95% CI	5.5–8.4	6.2–9.5		
	Suicide decedents, no.	119	145				Suicide decedents, no.	89	92		
	Population, no.	571,387	565,768				Population, no.	1,308,535	1,187,811		

See table footnotes on page 1258.

TABLE 3. (Continued) Suicide rate per 100,000 civilian, noninstitutionalized working persons aged 16–64 years, by sex, based on suicide decedents (N = 14,728) presumed in the labor force at time of death using Standard Occupational Classification (SOC) major groups — National Violent Death Reporting System, 17 states,* 2012 and 2015

Males					Females						
SOC code	Occupational group	2012	2015	Rate change		SOC code	Occupational group	2012	2015	Rate change	
				%	Rank†					%	Rank†
45	Farming, Fishing, and Forestry					41	Sales and Related				
	Rate rank	8	9	-13%	21		Rate rank	10	9	+20%	5
	Rate per 100,000	26.3	22.8				Rate per 100,000	6.4	7.7		
	95% CI	20.0–34.0	17.7–29.0				95% CI	5.3–7.4	6.6–8.7		
	Suicide decedents, no.	58	67				Suicide decedents, no.	148	192		
	Population, no.	220,364	293,746				Population, no.	2,325,223	2,505,186		
41	Sales and Related					15	Computer and Mathematical				
	Rate rank	11	10	+1%	12		Rate rank	NR	10	NR	NR
	Rate per 100,000	21.3	21.5				Rate per 100,000	NR	7.3		
	95% CI	19.4–23.2	19.6–23.4				95% CI	NR	5.0–10.5		
	Suicide decedents, no.	487	489				Suicide decedents, no.	20	30		
	Population, no.	2,282,361	2,276,666				Population, no.	390,260	408,410		
35	Food Preparation and Serving Related					53	Transportation and Material Moving				
	Rate rank	19	11	+43%	2		Rate rank	6	11	-17%	14
	Rate per 100,000	14.6	20.9				Rate per 100,000	8.3	6.9		
	95% CI	12.5–16.7	18.4–23.3				95% CI	6.0–11.2	4.8–9.7		
	Suicide decedents, no.	180	276				Suicide decedents, no.	43	33		
	Population, no.	1,234,381	1,321,800				Population, no.	517,082	477,143		
31	Health Care Support					21	Community and Social Service				
	Rate rank	9	12	-12%	18		Rate rank	8	12	-17%	16
	Rate per 100,000	22.1	19.5				Rate per 100,000	7.3	6.0		
	95% CI	14.8–31.7	12.5–29.0				95% CI	5.1–10.2	4.2–8.4		
	Suicide decedents, no.	29	24				Suicide decedents, no.	34	36		
	Population, no.	131,497	123,003				Population, no.	464,942	595,582		
17	Architecture and Engineering					43	Office and Administrative Support				
	Rate rank	10	13	-10%	15		Rate rank	14	13	+27%	4
	Rate per 100,000	21.6	19.4				Rate per 100,000	4.7	6.0		
	95% CI	18.1–25.1	16.3–22.6				95% CI	4.1–5.4	5.2–6.8		
	Suicide decedents, no.	145	147				Suicide decedents, no.	201	239		
	Population, no.	670,938	756,515				Population, no.	4,267,892	3,985,105		
23	Legal					13	Business and Financial Operations				
	Rate rank	12	14	-12%	19		Rate rank	12	14	-5%	11
	Rate per 100,000	21.3	18.7				Rate per 100,000	5.7	5.4		
	95% CI	15.7–28.2	13.4–25.4				95% CI	4.4–7.2	4.2–6.8		
	Suicide decedents, no.	48	41				Suicide decedents, no.	70	71		
	Population, no.	225,681	219,171				Population, no.	1,235,880	1,321,724		
11	Management					37	Building and Grounds Cleaning and Maintenance				
	Rate rank	17	15	+8%	9		Rate rank	15	15	+14%	8
	Rate per 100,000	16.4	17.8				Rate per 100,000	4.6	5.2		
	95% CI	14.9–17.9	16.3–19.3				95% CI	3.1–6.5	3.7–7.2		
	Suicide decedents, no.	477	530				Suicide decedents, no.	31	36		
	Population, no.	2,906,468	2,981,498				Population, no.	673,483	688,809		
39	Personal Care and Service					11	Management				
	Rate rank	13	16	-21%	22		Rate rank	13	16	-12%	12
	Rate per 100,000	20.9	16.5				Rate per 100,000	5.6	4.9		
	95% CI	16.2–26.4	12.9–20.7				95% CI	4.5–6.7	4.0–5.9		
	Suicide decedents, no.	68	73				Suicide decedents, no.	104	103		
	Population, no.	326,037	443,543				Population, no.	1,855,055	2,083,968		

See table footnotes on page 1258.

TABLE 3. (Continued) Suicide rate per 100,000 civilian, noninstitutionalized working persons aged 16–64 years, by sex, based on suicide decedents (N = 14,728) presumed in the labor force at time of death using Standard Occupational Classification (SOC) major groups — National Violent Death Reporting System, 17 states,* 2012 and 2015

Males					Females						
SOC code	Occupational group	2012	2015	Rate change		SOC code	Occupational group	2012	2015	Rate change	
				%	Rank [†]					%	Rank [†]
15	Computer and Mathematical					25	Education, Training, and Library				
	Rate rank	15	17	-11%	16		Rate rank	16	17	+3%	10
	Rate per 100,000	18.1	16.1				Rate per 100,000	3.3	3.4		
	95% CI	15.5–20.8	13.9–18.4				95% CI	2.6–4.2	2.7–4.2		
	Suicide decedents, no.	179	202				Suicide decedents, no.	69	74		
	Population, no.	986,994	1,252,275				Population, no.	2,091,706	2,186,483		
43	Office and Administrative Support					17	Architecture and Engineering				
	Rate rank	20	18	+12%	7		Rate rank	NR	NR	NR	NR
	Rate per 100,000	14.1	15.8				Rate per 100,000	NR	NR		
	95% CI	12.2–16.1	13.7–17.9				95% CI	NR	NR		
	Suicide decedents, no.	206	223				Suicide decedents, no.	10	12		
	Population, no.	1,456,242	1,411,453				Population, no.	135,632	144,852		
19	Life, Physical, and Social Science					19	Life, Physical, and Social Science				
	Rate rank	16	19	-13%	20		Rate rank	NR	NR	NR	NR
	Rate per 100,000	17.3	15.0				Rate per 100,000	NR	NR		
	95% CI	12.7–23.0	11.0–20.0				95% CI	NR	NR		
	Suicide decedents, no.	47	47				Suicide decedents, no.	13	19		
	Population, no.	271,690	312,925				Population, no.	225,992	204,566		
21	Community and Social Service					45	Farming, Fishing, and Forestry				
	Rate rank	21	20	+15%	6		Rate rank	NR	NR	NR	NR
	Rate per 100,000	12.8	14.6				Rate per 100,000	NR	NR		
	95% CI	8.8–17.9	10.7–19.6				95% CI	NR	NR		
	Suicide decedents, no.	33	45				Suicide decedents, no.	7	5		
	Population, no.	258,744	307,829				Population, no.	54,068	91,967		
13	Business and Financial Operations					47	Construction and Extraction				
	Rate rank	18	21	-11%	17		Rate rank	NR	NR	NR	NR
	Rate per 100,000	14.6	13.0				Rate per 100,000	NR	NR		
	95% CI	12.1–17.2	10.7–15.3				95% CI	NR	NR		
	Suicide decedents, no.	125	122				Suicide decedents, no.	9	14		
	Population, no.	855,329	941,806				Population, no.	55,164	76,173		
25	Education, Training, and Library					49	Installation, Maintenance, and Repair				
	Rate rank	22	22	-1%	13		Rate rank	NR	NR	NR	NR
	Rate per 100,000	10.9	10.9				Rate per 100,000	NR	NR		
	95% CI	8.6–13.6	8.6–13.5				95% CI	NR	NR		
	Suicide decedents, no.	78	79				Suicide decedents, no.	8	NR		
	Population, no.	713,321	727,167				Population, no.	73,231	46,136		

Abbreviations: CI = confidence interval; NR = not reported; number of decedents not reported <5, and rates were not calculated for occupational groups with ≤20 decedents; SOC = Standard Occupational Classification.

[†] Rate change rank refers to each occupational group's rank order for rate change from 2012 to 2015, where rank position 1 signifies the greatest suicide rate increase.

[§] Occupational groups were ranked by 2015 suicide rate, separately for males and females. Because of rounding, some rate and rate change results are not precisely calculable from the data presented.

[¶] 95% CIs were calculated using CDC's National Center for Health Statistics methods, including confidence limit factors for mortality rates based on <100 decedents.

Additional and tailored prevention approaches might be necessary to support workers at higher risk. Workplace suicide prevention efforts to date have focused primarily on early detection and tertiary intervention through the training of persons (i.e. gatekeepers) to identify those at risk for suicide and refer them to supporting services. However, more research on the role of the workplace in primary suicide prevention is

needed, including improving working conditions and reducing stress (8).

The findings in this report are subject to at least four limitations. First, because of the nature of the data that were available, and consistent with previous research methods, this report compared decedents' usual lifetime occupation as recorded on the death certificate with occupations of the employed population to calculate suicide rates. Additional

Summary**What is already known about this topic?**

From 2000 to 2016, the U.S. suicide rate among working aged (16–64 years) adults increased 34% from 12.9 per 100,000 population to 17.3.

What is added by this report?

2012 and 2015 National Violent Death Reporting System data from 17 states indicated the major occupational group with the highest male suicide rate was Construction and Extraction (43.6 [2012] and 53.2 [2015]). The Arts, Design, Entertainment, Sports, and Media major occupation group had the highest female suicide rate in 2012 (11.7) and 2015 (15.6).

What are the implications for public health practice?

A comprehensive approach to suicide prevention, including workplace-based approaches, is needed. CDC's technical package of strategies to prevent suicide is a resource for communities and workplaces to identify prevention strategies with the best available evidence.

data from coroner/medical examiner and law enforcement reports were used to exclude decedents identified as not in the labor force at time of death. Separate analyses indicated that if no such exclusion were applied, suicide rates would have been higher for all groups, although the top and bottom eight ranked occupational groups in 2015 by male suicide rate would maintain the same rank position, as would the top three and bottom four occupational groups by female suicide rate. Second, this report did not address confounding factors that might account for higher or lower rates of suicide between and within occupational groups, including education and income (9,10). Within SOC major occupational groups, employee education and income might vary widely. For example, the Management SOC major group includes farmers, ranchers, and chief executives of large companies, and the Construction and Extraction group includes both employees who might be salaried (e.g., supervisors) and those who might be paid hourly wages (e.g., roofer helpers). Future research might benefit from using more narrowly defined occupational groups and controlling for education and income to refine understanding of the relationship between occupation and suicide. Third, industry and occupation data obtained from death certificates rely on the accuracy and completeness of employment information provided by decedents' family members. It is also possible that completeness and accuracy of that information might be associated with decedents' job history. For example, categorization based on single lifetime industry and occupation might not accurately reflect employment for those persons with multiple lifetime occupations and those who worked across industries. Finally, this report is based on data from 17 U.S. states that

participated in NVDRS in 2012 and 2015, and the data are not nationally representative.

To address the multifactorial etiology of suicide, CDC recommends a comprehensive approach to prevention (1). Strategies might include enhancing social connectedness and expanding access to relevant resources, strengthening state or local economic supports, implementing practices that encourage help-seeking and decrease stigma, and providing referrals to mental health and other services (1). Strategies can be implemented to assure support and reduce access to lethal means among persons at risk. Decision makers, including employers, can create a response plan, should a suicide affect their organization. Surviving family and friends can be supported to reduce their own suicide risk. The media can follow reporting recommendations to avoid sensationalized reporting and can refrain from providing details on suicide methods (1). Further workplace prevention resources are available at <https://theactionalliance.org/>, and help is available at 1-800-273-TALK.

Acknowledgments

Jeff Purdin, Matt Hirst, Susan Burton, Jenny Huddleston, Elizabeth Smith, Division of Surveillance, Hazard Evaluations, and Field Studies, National Institute for Occupational Safety and Health, CDC.

Corresponding author: Deborah M. Stone, dstone3@cdc.gov, 770-488-3942.

¹Division of Analysis, Research, and Practice Integration, National Center for Injury Prevention and Control, CDC; ²Division of Violence Prevention, National Center for Injury Prevention and Control, CDC; ³Division of Safety Research, National Institute for Occupational Safety and Health, CDC; ⁴Division of Surveillance, Hazard Evaluations, and Field Studies, National Institute for Occupational Safety and Health, CDC.

All authors have completed and submitted the ICMJE form for disclosure of potential conflicts of interest. No potential conflicts of interest were disclosed.

References

1. Stone DM, Holland K, Bartholow B, Crosby A, Davis S, Wilkins N. Preventing suicide: a technical package of policies, programs, and practices. Atlanta, GA: US Department of Health and Human Services, CDC; 2017. <https://www.cdc.gov/violenceprevention/pdf/suicideTechnicalPackage.pdf>
2. US Census Bureau. Current population survey annual social and economic supplement 2013 and 2016. Washington, DC: US Census Bureau; 2018.
3. National Center for Health Statistics. Vital statistics of the United States: mortality, 1999 technical appendix. Hyattsville, MD: US Department of Health and Human Services, CDC, National Center for Health Statistics; 2004. <https://www.cdc.gov/nchs/data/statab/techap99.pdf>
4. Milner A, Witt K, LaMontagne AD, Niedhammer I. Psychosocial job stressors and suicidality: a meta-analysis and systematic review. *Occup Environ Med* 2018;75:245–53. <https://doi.org/10.1136/oemed-2017-104531>

5. Milner A, Witt K, Maheen H, LaMontagne AD. Access to means of suicide, occupation and the risk of suicide: a national study over 12 years of coronial data. *BMC Psychiatry* 2017;17:125. <https://doi.org/10.1186/s12888-017-1288-0>
6. Milner A, Spittal MJ, Pirkis J, LaMontagne AD. Suicide by occupation: systematic review and meta-analysis. *Br J Psychiatry* 2013;203:409–16. <https://doi.org/10.1192/bjp.bp.113.128405>
7. Stallones L, Doenges T, Dik BJ, Valley MA. Occupation and suicide: Colorado, 2004-2006. *Am J Ind Med* 2013;56:1290–5.
8. Milner A, Page K, Spencer-Thomas S, Lamotagne AD. Workplace suicide prevention: a systematic review of published and unpublished activities. *Health Promot Int* 2015;30:29–37. <https://doi.org/10.1093/heapro/dau085>
9. Daly MC, Wilson DJ, Johnson NJ. Relative status and well-being: evidence from U.S. suicide deaths. *Rev Econ Stat* 2013;95:1480–500. https://doi.org/10.1162/REST_a_00355
10. Phillips JA, Hempstead K. Differences in U.S. suicide rates by educational attainment, 2000–2014. *Am J Prev Med* 2017;53:e123–30. <https://doi.org/10.1016/j.amepre.2017.04.010>