



Surveillance Summaries / Vol. 73 / No. 2

May 2, 2024

Preventable Premature Deaths from the Five Leading Causes of Death in Nonmetropolitan and Metropolitan Counties, United States, 2010–2022



CONTENTS

troduction	2
ethods	3
esults	3
iscussion	7
mitations	8
ıture Directions	9
onclusion	9
eferences	9

The MMWR series of publications is published by the Office of Science, Centers for Disease Control and Prevention (CDC), U.S. Department of Health and Human Services, Atlanta, GA 30329-4027.

Suggested citation: [Author names; first three, then et al., if more than six.] [Title]. MMWR Surveill Summ 2024;73(SS-#):[inclusive page numbers].

Centers for Disease Control and Prevention

Mandy K. Cohen, MD, MPH, Director
Debra Houry, MD, MPH, Chief Medical Officer and Deputy Director for Program and Science
Samuel F. Posner, PhD, Director, Office of Sciencee

MMWR Editorial and Production Staff (Serials) Phyllis H. King,

Charlotte K. Kent, PhD, MPH, Editor in Chief Rachel Gorwitz, MD, MPH, Acting Executive Editor Christine G. Casey, MD, Editor Mary Dott, MD, MPH, Online Editor Terisa F. Rutledge, Managing Editor David C. Johnson, Lead Technical Writer-Editor Marella Meadows, Project Editor

Acting Lead Health Communication Specialist
Alexander J. Gottardy, Maureen A. Leahy,
Stephen R. Spriggs, Armina Velarde, Tong Yang,
Visual Information Specialists
Quang M. Doan, MBA,
Terraye M. Starr, Moua Yang,
Information Technology Specialists

Symone Hairston, MPH,
Acting Lead Health Communication Specialist
Kiana Cohen, MPH,
Leslie Hamlin, Lowery Johnson,
Health Communication Specialists
Dewin Jimenez, Will Yang, MA,
Visual Information Specialists

Matthew L. Boulton, MD, MPH Carolyn Brooks, ScD, MA Virginia A. Caine, MD Jonathan E. Fielding, MD, MPH, MBA Timothy F. Jones, MD, *Chairman*David W. Fleming, MD
William E. Halperin, MD, DrPH, MPH
Jewel Mullen, MD, MPH, MPA
Jeff Niederdeppe, PhD
Patricia Quinlisk, MD, MPH

MMWR Editorial Board

Patrick L. Remington, MD, MPH Carlos Roig, MS, MA William Schaffner, MD Morgan Bobb Swanson, MD, PhD

Preventable Premature Deaths from the Five Leading Causes of Death in Nonmetropolitan and Metropolitan Counties, United States, 2010–2022

Macarena C. García, DrPH¹; Lauren M. Rossen, PhD²; Kevin Matthews, PhD³; Gery Guy, PhD⁴; Katrina F. Trivers, PhD⁵; Cheryll C. Thomas, MSPH⁵; Linda Schieb, MSPH⁵; Michael F. Iademarco, MD¹

Office of Science and Medicine, Office of the Assistant Secretary of Health, U.S. Department of Health and Human Services, Washington, DC;
²National Center for Health Statistics, CDC, Hyattsville, Maryland; ³National Center for State, Tribal, Local, and Territorial Public Health Infrastructure and Workforce, CDC, Atlanta, Georgia; ⁴National Center for Injury Prevention and Control, CDC, Atlanta, Georgia;
⁵National Center for Chronic Disease Prevention and Health Promotion, CDC, Atlanta, Georgia

Abstract

Problem/Condition: A 2019 report quantified the higher percentage of potentially excess (preventable) deaths in U.S. nonmetropolitan areas compared with metropolitan areas during 2010–2017. In that report, CDC compared national, regional, and state estimates of preventable premature deaths from the five leading causes of death in nonmetropolitan and metropolitan counties during 2010–2017. This report provides estimates of preventable premature deaths for additional years (2010–2022).

Period Covered: 2010–2022.

Description of System: Mortality data for U.S. residents from the National Vital Statistics System were used to calculate preventable premature deaths from the five leading causes of death among persons aged <80 years. CDC's National Center for Health Statistics urban-rural classification scheme for counties was used to categorize the deaths according to the urban-rural county classification level of the decedent's county of residence (1: large central metropolitan [most urban], 2: large fringe metropolitan, 3: medium metropolitan, 4: small metropolitan, 5: micropolitan, and 6: noncore [most rural]). Preventable premature deaths were defined as deaths among persons aged <80 years that exceeded the number expected if the death rates for each cause in all states were equivalent to those in the benchmark states (i.e., the three states with the lowest rates). Preventable premature deaths were calculated separately for the six urban-rural county categories nationally, the 10 U.S. Department of Health and Human Services public health regions, and the 50 states and the District of Columbia.

Results: During 2010–2022, the percentage of preventable premature deaths among persons aged <80 years in the United States increased for unintentional injury (e.g., unintentional poisoning including drug overdose, unintentional motor vehicle traffic crash, unintentional drowning, and unintentional fall) and stroke, decreased for cancer and chronic lower respiratory disease (CLRD), and remained stable for heart disease. The percentages of preventable premature deaths from the five leading causes of death were higher in rural counties in all years during 2010–2022. When assessed by the six urban-rural county classifications, percentages of preventable premature deaths in the most rural counties (noncore) were consistently higher than in the most urban counties (large central metropolitan and fringe metropolitan) for the five leading causes of death during the study period.

During 2010–2022, preventable premature deaths from heart disease increased most in noncore (+9.5%) and micropolitan counties (+9.1%) and decreased most in large central metropolitan counties (-10.2%). Preventable premature deaths from cancer decreased in all county categories, with the largest decreases in large central metropolitan and large fringe metropolitan counties (-100.0%; benchmark achieved in both county categories in 2019). In all county categories, preventable premature deaths from unintentional injury increased, with the largest increases occurring in large central metropolitan (+147.5%) and large fringe metropolitan (+97.5%) counties. Preventable premature deaths from CLRD decreased most in large central metropolitan counties where the benchmark was achieved in 2019 and increased slightly in noncore counties (+0.8%). In all county categories, preventable premature deaths from stroke decreased from 2010 to 2013, remained constant from 2013 to 2019, and then increased in 2020 at the start of the COVID-19 pandemic. Percentages of preventable premature deaths varied across states by urban-rural county classification during 2010–2022.

Corresponding author: Macarena C. García, Office of Science and Medicine, Office of the Assistant Secretary of Health, U.S. Department of Health and Human Services, Washington, DC. Telephone: 678-770-6220; Email: mcgarcia@cdc.gov.

Interpretation: During 2010–2022, nonmetropolitan counties had higher percentages of preventable premature deaths from the five leading causes of death than did metropolitan counties nationwide, across public health regions, and in most states.

The gap between the most rural and most urban counties for preventable premature deaths increased during 2010–2022 for four causes of death (cancer, heart disease, CLRD, and stroke) and decreased for unintentional injury. Urban and suburban counties (large central metropolitan, large fringe metropolitan, medium metropolitan, and small metropolitan) experienced increases in preventable premature deaths from unintentional injury during 2010–2022, leading to a narrower gap between the already high (approximately 69% in 2022) percentage of preventable premature deaths in noncore and micropolitan counties. Sharp increases in preventable premature deaths from unintentional injury, heart disease, and stroke were observed in 2020, whereas preventable premature deaths from CLRD and cancer continued to decline. CLRD deaths decreased during 2017–2020 but increased in 2022. An increase in the percentage of preventable premature deaths for multiple leading causes of death was observed in 2020 and was likely associated with COVID-19–related conditions that contributed to increased mortality from heart disease and stroke.

Public Health Action: Routine tracking of preventable premature deaths based on urban-rural county classification might enable public health departments to identify and monitor geographic disparities in health outcomes. These disparities might be related to different levels of access to health care, social determinants of health, and other risk factors. Identifying areas with a high prevalence of potentially preventable mortality might be informative for interventions.

Introduction

Premature deaths, all-cause mortality, and poor health outcomes are greater among residents of rural counties than of urban counties in the United States (1). In 2021, the all-cause age-adjusted death rate in the United States was 841.6 per 100,000 population. The gap in all-cause mortality between rural (nonmetropolitan) and urban (metropolitan) areas of the United States continues to widen. In 1999, the death rate in rural areas was 7% higher than in urban areas; by 2019, it was 20% higher (2). Describing premature mortality rates from the five leading causes of death (cancer, unintentional injury [e.g., unintentional poisoning including drug overdose, unintentional motor vehicle traffic crash, unintentional drowning, and unintentional fall], heart disease, stroke, and chronic lower respiratory disease [CLRD]) and related rural disparities might help guide public health messaging and interventions.

The risk for premature death is associated with modifiable factors that vary by disease (3). Four of the five leading risk factors for premature death are more prevalent in rural areas of the United States: using tobacco, obesity, physical inactivity, and drinking alcohol or drinking in excess (4,5). Extensive literature on social determinants of health has established the importance of community context in shaping all aspects of health (6). Structural factors (e.g., lower socioeconomic status, limited access to health care professionals, and limited job opportunities) increase the risk for premature death among rural residents (7).

Multiple factors influence the rural-urban gap in preventable premature deaths. Because each of the five leading causes of death is age related, these conditions are more prevalent in rural areas of the United States where residents typically are older than their urban counterparts. Working-age adults might leave rural areas to seek better economic opportunities elsewhere (8), and older persons might be more likely to retire in rural areas (9). However, the population's age structure alone does not explain the disparity in mortality. Instead, differences in social circumstances, socioeconomic characteristics, health-related behaviors, and access to health care services affect mortality and potentially contribute to approximately half of all preventable premature deaths (10). County-level disparities in all-cause premature deaths by rurality, race, and ethnicity have been documented (11). Data on cause-specific preventable premature deaths from the leading causes of death by rurality, sex, race, and ethnicity are limited, and direct comparisons accounted for by these factors will be reported in subsequent analyses.

Rural public health needs and sociodemographic characteristics of rural populations are changing (12). Although the proportion of the U.S. population that lives in rural areas is gradually declining, any rural population growth can be attributed to in-migration, which might require sensitivity to cultural differences (13,14). With gradual declines in population, the wealth and tax bases of rural counties also are decreasing, resulting in reduced funding for social and health services (15).

In this analysis, mortality data were used to estimate the number and percentage of deaths from each of the five leading causes of death that could have been prevented if all states had similarly low death rates. Disparities in premature mortality from the five leading causes of death in rural areas in the United States during 2010–2022 also were estimated. The results of this analysis are intended to serve as a critical resource for policymakers, public health officials, and researchers striving to understand and address the root causes of preventable premature deaths.

Methods

This analysis used mortality data for U.S. residents from the National Vital Statistics System (https://www.cdc.gov/nchs/index.htm) to calculate preventable premature deaths by urban-rural county classification from the five leading causes of death during 2010–2022 (heart disease, cancer, unintentional injury, CLRD, and stroke). Deaths from COVID-19 were excluded to maintain consistency and facilitate the assessment of trends over time. Data for 2022 are provisional counts from January through June and were annualized for comparability with previous years.

The number of preventable premature deaths for a specific cause (also described as potentially preventable premature or excess deaths) is equal to the difference in the number of observed deaths among persons aged <80 years and the number of deaths expected if the mortality rate in all states were equivalent to the average rate of the three states with the lowest mortality. Rates in the three states define the benchmarks. The benchmark for each cause of death is derived from a unique set of three states.

Rural and urban categories were identified using the National Center for Health Statistics 2013 urban-rural classification scheme for counties (16). County of residence of the decedent was used to determine urban-rural county classification. The categories are 1: large central metropolitan (most urban), 2: large fringe metropolitan, 3: medium metropolitan, 4: small metropolitan, 5: micropolitan, and 6: noncore (most rural).

Preventable premature deaths were calculated individually for the two nonmetropolitan categories (micropolitan and noncore) and the four metropolitan categories (large central metropolitan, large fringe metropolitan, medium metropolitan, and small metropolitan) as well as for the broader categories of metropolitan and nonmetropolitan. Analyses were restricted to deaths with an underlying cause of death from the five leading causes of death based on the International Classification of Diseases, 10th Revision (ICD-10): heart disease (I00-I09, I11, I13, and I20-I51), cancer (C00-C97), unintentional injury (V01-X59 and Y85-Y86), CLRD (J40-J47), and stroke (I60–I69). The analysis of preventable premature deaths during 2010-2022 was restricted to persons aged <80 years at the time of death. The age restriction is consistent with the average life expectancy for the U.S. population in 2010, which was approximately 79 years (17).

Age-specific mortality rates for each of the five leading causes of death were used to derive the number of preventable premature deaths using methods described elsewhere (18). Age groupings varied by cause of death. (Most were 10-year age groups; however, the size of the youngest age group ranged


from 0 to 9 years for unintentional injury to 0 to 49 years for CLRD and cerebrovascular disease because deaths from those causes are rare among younger persons.) For each age group and cause of death, the death rates of the three states with the lowest rates during 2008-2010 (benchmark states) were averaged to produce benchmark rates (18) (https://stacks.cdc.gov/view/ cdc/42342). These benchmarks were chosen to represent the lowest death rates achievable by states at the beginning of the study period and did not vary by year to allow for the examination of trends over time. Although using time-varying benchmarks would better account for potential improvements over time in the benchmark rates, time-varying benchmarks also would make temporal and geographic comparisons more difficult. The same benchmarks were applied to both nonmetropolitan and metropolitan counties, and benchmarks were not adjusted for other characteristics that might affect death rates (e.g., race, ethnicity, socioeconomic status, and urbanicity). Deaths attributed to COVID-19 from 2020 through June 2022 were excluded from this study.


The numbers of preventable premature deaths for each cause of death were assumed to follow a Poisson distribution. SEs were calculated using standard formulas that incorporated the variance around both the observed and the expected counts (18), and pairwise z-tests were performed to determine whether the differences during 2010–2022 were statistically significant (p<0.05). All differences during 2010–2022 are statistically significant unless otherwise noted. The percentage of preventable premature deaths was calculated by dividing the number of preventable premature deaths by the total observed number of premature deaths.

Results

The percentage of preventable premature deaths from cancer decreased from 2010 through June 2022 (from 21% to 0.3%) (Figure 1). Regardless of urban-rural classification, all county categories experienced decreases (Figure 2). However, the decreases in urban counties were larger than those in rural counties, which widened the rural-urban disparities in preventable premature deaths from cancer (Figure 2) (Table). The percentage of preventable premature deaths from cancer in noncore counties in 2022 (18.1%) was similar to the percentage in large central metropolitan counties in 2010 (17.9%).

The percentage of preventable premature deaths from heart disease decreased from 2010 through 2019 (from 33.5% to 28.8%), followed by a steep increase to 33.6% from 2020 through June 2022 (Figure 1). Increases from 2020 through June 2022 occurred across all rural-urban





^{*} Preventable premature deaths are defined as deaths among persons aged <80 years in excess of the number that would be expected if the death rates for each cause in all states were equivalent to those in the benchmark states (i.e., the three states with the lowest rates).

 † Data for 2022 are provisional counts from January through June and were annualized for comparability with previous years.


categories except for large central metropolitan counties, which experienced a decrease from 32.9% in 2020 to 30.1% in 2021 (Figure 2) (Table). Rural counties had the highest percentage of preventable premature deaths from heart disease in 2022 (45.8% in micropolitan counties and 49.4% in noncore counties) (Figure 2) (Table). Most states experienced an increase in preventable early deaths from heart disease and stroke (96% and 88% of states, respectively) from 2019 through June 2022 (Supplementary Table, https://stacks.cdc.gov/view/cdc/147842).

The percentage of preventable premature deaths from unintentional injury increased from 2010 to 2019 (from 38.8% to 53.8%), followed by a steep increase from 2019 to 2021 and a slight decrease through June 2022 to 63.5% (Figure 1). Increases in preventable premature deaths from unintentional injury during 2010–2022 were statistically

significant for all metropolitan categories except micropolitan. Rural percentages were higher than in urban areas, but the gap narrowed (Figure 2). The percentages increased in all states except Wyoming, but the increase varied widely at the state level (Supplementary Table, https://stacks.cdc.gov/view/cdc/147842).

The percentage of preventable premature deaths from CLRD decreased from 2010 through 2022 (from 38.6% to 25.5%) (Figure 1). The percentage of preventable premature deaths varied widely when stratified by rural-urban county category, but all county categories except for noncore counties experienced decreases. Rural-urban disparities widened when large central metropolitan percentages decreased from 23.4% in 2010 to 0% in 2022, whereas the rural percentages hovered between 50.7% and 54.8% in 2022 (Figure 2) (Table).

FIGURE 2. Percentages of preventable premature deaths* among persons aged <80 years from the five leading causes of death by rural-urban county classification and year — National Vital Statistics System, United States, 2010–2022 †



^{*} Preventable premature deaths are defined as deaths among persons aged <80 years in excess of the number that would be expected if the death rates for each cause in all states were equivalent to those in the benchmark states (i.e., the three states with the lowest rates).

[†] Data for 2022 are provisional counts from January through June and were annualized for comparability with previous years.

TABLE. Numbers and percentages of preventable premature deaths* among persons aged <80 years from the five leading causes of death, by rural-urban county classification — National Vital Statistics System, United States, 2010–2022

Cause	2010 No. (%)	2011 No. (%)	2012 No. (%)	2013 No. (%)	2014 No. (%)	2015 No. (%)	2016 No. (%)	2017 No. (%)	2018 No. (%)	2019 No. (%)	2020 No. (%)	2021 No. (%)	2022 [†] No. (%)	Percent change from 2010 to 2022
Heart disease														
Large central metropolitan	24,859 (33.5)	23,836 (31.9)	23,173 (30.8)	23,344 (30.3)	22,655 (29.0)	23,660 (29.4)	24,515 (29.7)	23,496 (28.0)	24,158 (28.1)	22,929 (26.6)	31,779 (32.9)	28,410 (30.0)	14,212 (30.1)	-10.2
Large fringe metropolitan		12,360 (21.4)	11,574 (19.9)	11,358 (19.0)	11,261 (18.4)	11,762 (18.6)	12,159 (18.7)	12,399 (18.5)	11,887 (17.4)	11,184 (16.2)	16,012 (21.2)	16,593 (21.3)	8,635 (22.0)	-8.8
Medium metropolitan	17,781 (31.7)	17,089 (30.3)	17,130 (29.8)	17,992 (30.2)	17,999 (29.6)	18,951 (30.1)	19,583 (30.3)	20,029 (30.1)	19,963 (29.4)	20,148 (29.0)	24,813 (32.9)	26,038 (33.5)	13,143 (33.7)	6.4
Small metropolitan	10,420 (36.9)	10,021 (35.5)	9,924 (34.7)	10,147 (34.7)	10,295 (34.4)	10,686 (34.7)	11,089 (35.1)	11,282 (34.8)	11,822 (35.2)	11,378 (33.8)	13,943 (37.9)	15,089 (39.4)	7,579 (39.5)	7.0
Micropolitan	12,913 (42.0)	12,320 (40.4)	12,585 (40.5)	12,861 (40.5)	13,139 (40.5)	14,326 (42.2)	14,445 (42.1)	14,611 (41.8)	14,686 (41.4)	14,711 (40.9)	17,167 (44.3)	18,723 (46.1)	9,285 (45.8)	9.1
Noncore	11,509 (45.1)	11,111 (43.9)	11,132 (43.5)	11,556 (44.0)	12,015 (44.6)	12,384 (45.1)	12,892 (45.8)	12,644 (44.9)	12,399 (43.9)	12,577 (43.9)	14,482 (47.1)	15,910 (49.4)	7,981 (49.4)	9.5
Cancer														
Large central metropolitan	18,754 (17.9)	16,709 (15.8)	15,945 (14.9)	12,844 (12.0)	11,830 (10.9)	9,546 (8.7)	7,693 (7.0)	3,443 (3.2)	736 (0.7)	0 (0)	0 (0)	0 (0)	0 (0)	-100.0
Large fringe metropolitan	16,566 (17.7)	14,208 (15.2)	12,930 (13.6)	11,229 (11.7)	9,870 (10.1)	8,102 (8.3)	6,948 (7.0)	4,000 (4.0)	1,112 (1.1)	0 (0)	0 (0)	0 (0)	0 (0)	-100.0
Medium metropolitan	18,252 (21.4)	16,330 (19.2)	16,337 (18.8)	14,635 (16.7)	14,282 (16.0)	13,232 (14.7)	11,366 (12.6)	9,328 (10.3)	7,057 (7.8)	4,761 (5.3)	4,512 (4.9)	3,117 (3.4)	1,302 (2.8)	-86.9
Small metropolitan	9,606 (23.6)	8,883 (21.8)	8,836 (21.3)	8,263 (19.8)	7,773 (18.4)	7,500 (17.6)	6,798 (15.9)	5,766 (13.5)	5,322 (12.3)	4,497 (10.4)	3,908 (9.0)	3,422 (7.8)	1,383 (6.4)	-72.8
Micropolitan	11,313 (26.7)	11,254 (26.1)	10,912 (25.2)	10,837 (24.7)	10,060 (23.0)	10,050 (22.6)	9,416 (21.3)	8,819 (19.9)	7,758 (17.6)	7,438 (16.7)	7,399 (16.4)	6,619 (14.8)	3,115 (14.0)	-47.4
Noncore	9,806 (28.7)	9,503 (27.7)	9,192 (26.7)	8,831 (25.6)	8,723 (25.1)	8,563 (24.5)	8,321 (23.8)	7,538 (21.7)	7,291 (20.9)	6,456 (18.8)	6,434 (18.5)	7,139 (20.1)	3,130 (18.1)	-37.0
Unintentional injury§														
Large central metropolitan	5,924 (25.4)	6,941 (28.2)	7,072 (28.2)	8,024 (30.6)	9,021 (32.8)	11,168 (37.3)	15,900 (45.7)	17,580 (47.8)	17,140 (47.0)	19,258 (49.8)	28,868 (59.6)	33,518 (63.0)	16,603 (62.8)	147.5
Large fringe metropolitan	5,765 (28.8)	6,645 (31.5)	7,059 (32.5)	7,229 (32.7)	8,049 (34.8)	10,236 (40.1)	14,422 (48.2)	16,118 (50.6)	14,814 (48.2)	15,219 (48.6)	20,303 (55.5)	22,544 (57.6)	10,953 (56.9)	97.5
Medium metropolitan	9,129 (43.1)	9,589 (43.9)	9,242 (42.8)	9,804 (43.9)	10,808 (46.0)	12,849 (50.1)	15,930 (55.2)	17,774 (57.6)	17,023 (56.2)	17,824 (57.1)	24,187 (64.2)	27,794 (67.0)	13,439 (66.2)	53.7
Small metropolitan	4,342 (44.5)	4,775 (46.6)	4,546 (45.1)	4,715 (45.9)	5,197 (48.0)	5,892 (51.0)	6,587 (53.6)	7,171 (55.5)	7,129 (55.1)	7,213 (55.2)	9,511 (61.7)	11,339 (65.5)	5,595 (65.2)	46.6
Micropolitan	5,865 (53.1)	6,293 (54.7)	6,263 (54.5)	6,107 (53.7)	6,375 (54.7)	6,973 (56.8)	7,786 (59.5)	8,376 (61.1)	8,007 (59.9)	8,418 (61.0)	10,200 (65.4)	12,042 (68.9)	5,755 (67.9)	27.8
Noncore	5,781 (60.9)	5,972 (61.6)	5,924 (61.4)	5,656 (60.2)	5,758 (60.6)	6,304 (62.8)	6,683 (64.2)	6,673 (64.1)	6,379 (63.0)	6,571 (63.6)	8,028 (68.1)	9,289 (71.2)	4,384 (70.0)	15.1
Chronic lower respiratory	disease													
Large central metropolitan	3,756 (23.4)	3,587 (22.1)	3,250 (20.0)	3,581 (21.0)	2,627 (15.9)	2,986 (17.2)	2,906 (16.5)	2,322 (13.0)	1,902 (10.7)	977 (5.6)	524 (3.0)	0 (0)	0 (0)	-100.0
Large fringe metropolitan	4,546 (29.4)	4,717 (29.5)	4,244 (26.6)	4,357 (26.4)	4,136 (24.6)	4,472 (25.5)	4,237 (23.9)	4,314 (23.4)	3,787 (20.5)	3,305 (17.9)	2,152 (12.1)	855 (5.0)	564 (6.5)	-77.8
Medium metropolitan	6,794 (41.1)	6,815 (40.4)	6,928 (40.1)	7,426 (41.0)	7,275 (39.6)	7,714 (40.3)	7,970 (40.4)	8,104 (39.8)	7,573 (37.3)	6,873 (34.3)	6,432 (32.2)	5,195 (27.3)	2,955 (29.9)	-27.2
Small metropolitan	4,079 (47.1)	4,225 (47.3)	4,243 (46.8)	4,613 (48.1)	4,377 (46.0)	4,928 (48.3)	4,916 (47.6)	5,024 (47.2)	5,244 (47.4)	4,777 (44.4)	4,564 (42.5)	4,230 (40.3)	2,281 (42.1)	-10.7
Micropolitan	5,203 (53.1)	5,284 (52.9)	5,413 (53.0)	5,906 (54.5)	5,843 (53.6)	6,355 (55.2)	6,441 (55.0)	6,538 (54.6)	6,394 (53.4)	6,491 (53.2)	6,210 (51.5)	5,767 (49.3)	3,050 (50.7)	-4.6
Noncore	4,342 (54.3)	4,434 (54.3)	4,621 (54.9)	4,807 (55.3)	4,801 (54.7)	5,307 (56.8)	5,153 (55.8)	5,586 (57.1)	5,469 (56.1)	5,524 (55.8)	5,307 (54.4)	5,116 (53.4)	2,699 (54.8)	0.8

See table footnotes on the next page.

The percentage of preventable premature deaths from stroke decreased slightly from 2010 through 2019 (32.4% to 26.4%), followed by an increase to 33.9% through June 2022 (Figure 1). Each rural-urban category experienced steep increases from 2019 to June 2022, except for noncore counties that experienced a slight decrease from 2021 to

June 2022; rural counties had the highest percentages from January to June 2022 (42.0% in micropolitan counties and 40.9% in noncore counties) (Figure 2) (Table). The highest percentages of preventable premature deaths from stroke in 2022 were in southern states (Supplementary Table, https://stacks.cdc.gov/view/cdc/147842).

TABLE. (Continued) Numbers and percentages of preventable premature deaths* among persons aged <80 years from the five leading causes of death, by rural-urban county classification — National Vital Statistics System, United States, 2010–2022

Cause	2010 No. (%)	2011 No. (%)	2012 No. (%)	2013 No. (%)	2014 No. (%)	2015 No. (%)	2016 No. (%)	2017 No. (%)	2018 No. (%)	2019 No. (%)	2020 No. (%)	2021 No. (%)	2022 [†] No. (%)	Percent change from 2010 to 2022
Stroke														
Large central metropolitan	4,465	4,263	3,838	3,806	3,779	4,132	4,391	4,506	4,280	4,412	5,713	6,082	3,209	3.4
	(31.7)	(30.1)	(27.5)	(26.7)	(25.9)	(27.1)	(27.9)	(27.4)	(25.9)	(26.0)	(30.7)	(31.6)	(32.8)	
Large fringe metropolitan	2,494	2,227	2,209	1,732	2,009	2,333	2,345	2,209	2,406	2,494	3,369	3,720	2,160	14.8
	(22.7)	(20.3)	(19.7)	(15.6)	(17.1)	(18.9)	(18.5)	(17.0)	(17.7)	(17.8)	(22.1)	(23.3)	(26.1)	
Medium metropolitan	3,762	3,527	3,220	3,258	3,492	3,665	3,667	3,784	3,905	3,736	4,732	5,527	2,904	7.2
	(33.3)	(31.3)	(28.9)	(28.4)	(29.2)	(29.6)	(29.1)	(28.9)	(28.8)	(27.3)	(31.6)	(34.6)	(35.7)	
Small metropolitan	1,886	1,795	1,623	1,625	1,651	1,817	1,882	1,941	1,760	1,866	2,422	2,667	1,345	4.0
	(34.9)	(33.3)	(30.5)	(29.9)	(29.7)	(31.2)	(31.4)	(31.4)	(28.6)	(29.3)	(34.3)	(36.1)	(36.3)	
Micropolitan	2,393	2,255	2,229	2,211	2,183	2,251	2,212	2,201	2,287	2,155	2,751	3,087	1,610	3.6
	(40.5)	(38.6)	(37.9)	(37.1)	(36.3)	(36.5)	(35.8)	(35.0)	(35.3)	(33.4)	(38.6)	(41.0)	(42.0)	
Noncore	1,970	1,939	1,723	1,745	1,854	1,808	1,956	1,906	1,944	1,914	2,341	2,371	1,147	-1.7
	(41.6)	(40.8)	(37.5)	(37.3)	(38.3)	(37.4)	(39.0)	(37.8)	(37.8)	(37.0)	(41.4)	(41.7)	(40.9)	

^{*} Preventable premature deaths are defined as deaths among persons aged <80 years in excess of the number that would be expected if the death rates for each cause in all states were equivalent to those in the benchmark states (i.e., the three states with the lowest rates). Estimates of potentially excess deaths that were negative were set to zero. These negative excess estimates occurred in cases where deaths were fewer than expected (i.e., mortality was lower than benchmark rates).

Discussion

Rural residents, particularly those in noncore counties, experienced high percentages of preventable premature deaths during the study period. The rural-urban disparities in premature deaths varied by cause of death. However, disparities were not limited to place of residence. Disparities in all-cause premature deaths also were associated with other demographic factors (e.g., sex, race, and ethnicity) (11). For example, the highest rates of premature deaths were observed in rural counties where a majority of the population was Black, African American, American Indian, or Alaska Native (11). To address disparities in preventable premature deaths across rural and urban counties, data on disparities in cause-specific premature deaths from the five leading causes by rural-urban county category, race, and ethnicity are needed to inform interventions and health care policies for specific racial and ethnic groups. A follow-up of this analysis stratified by race and ethnicity will be published in subsequent reports, further contributing evidence to guide existing and new programs and policies.

Cancer

Overall, the decrease in preventable premature deaths from cancer was substantial and was greatest in urban counties where access to preventive services, treatment, survivor care, and specialty care is much higher than in rural counties (19). Large central metropolitan and fringe metropolitan areas achieved the benchmark rates in 2019. This is consistent with

overall declines in cancer mortality, which decreased 27% between 2001 and 2020 (20). The decrease in preventable premature deaths likely reflects multiple factors. Increases in recommended screening for the leading causes of deaths from cancer (e.g., lung, colon, cervical, and female breast) have led to earlier detection, when treatment is more effective, and prevention by detecting cellular changes before they turn into cancer, as in the case of colorectal cancer (21). Increases in vaccination rates for cancer-causing viruses and decreases in prevalence of risk factors (e.g., combustible tobacco use) also have driven cancer mortality downward (22). Access to these cancer prevention and early detection strategies was increased with the expansion of Medicaid (23). New cancer treatments and therapies, specifically for lung cancer and melanoma, also have led to longer survival for those with a cancer diagnosis (24). CDC conducted a demonstration project on how to best provide care for persons living in rural areas who had cancer diagnosed (25). Although cancer is categorized as a single disease group in this analysis, each cancer site has different risk factors, has varying treatment methods, and can manifest itself in different ways among groups by sex, age, race, and ethnicity. Preventable premature deaths might vary depending on the cancer site and might not have decreased for cancers with increasing prevalence of risk factors (e.g., obesity), no recommended screening modalities, or therapies that have not changed. Lung cancer, the leading cause of cancer mortality, accounted for 23% of all cancer deaths in 2020 (20). Geographic differences in combustible tobacco use and use of lung cancer screening likely partially drive differences in lung cancer mortality. Access to lung cancer screening facilities is

[†] Data for 2022 are provisional counts from January through June and were annualized for comparability with previous years.

[§] Includes unintentional poisoning (e.g., drug overdose), unintentional motor vehicle traffic crash, unintentional drowning, and unintentional fall (https://www.cdc.gov/injury/wisqars/animated-leading-causes.html).

more limited in rural counties than in urban counties (26). Despite overall reductions in preventable premature deaths from cancer, premature deaths surpass the national average in micropolitan and noncore counties, highlighting the need in rural areas to reduce cancer-related premature deaths. Because more urban areas surpassed the 2010 benchmarks for cancer death rates in 2019, future updates to the cancer-specific benchmarks using more recent years of data might better reflect the lowest achievable death rates.

Unintentional Injury

The worsening and expanding drug overdose epidemic, increases in motor vehicle traffic fatalities, and falls drive the growth in preventable premature deaths from unintentional injury (27). Narrowing rural-urban disparities in the percentage of preventable premature deaths from unintentional injury were driven by worsening rates of preventable mortality in more urban areas, with the percentage more than doubling in large central metropolitan areas over the study period. For drug overdoses, access to medications for opioid use disorder continues to be more limited in rural counties, as evidenced by low buprenorphine dispensing rates and reduced treatment capacity (28). For motor vehicle traffic crashes, rural residents have an increased risk for death and are less likely than urban residents to wear seat belts (29). Evidence-based interventions reduce rural-urban disparities in seat belt use and motor vehicle death rates (30). Many fall risk factors are modifiable, implying that many falls can be prevented (31).

Heart Disease and Stroke

Disparities in preventable premature deaths from heart disease and stroke between rural and urban areas existed across the study period. These gaps increased from 2019 to June 2022, except in large central metropolitan counties where a decrease of three percentage points was observed from 2020 to 2021. Increases in preventable premature deaths from heart disease and stroke in 2020 and 2021 were likely associated with COVID-19-related conditions that contributed to riskassociated increased mortality from heart disease and stroke (32). Increases in systolic and diastolic blood pressure, a leading risk factor for heart disease and stroke, were observed among all age groups when comparing 2020 with 2019 (33). Inequities in control of hypertension (i.e., systolic blood pressure values of ≥130 mm Hg, diastolic blood pressure of >80 mm Hg, or both) were observed during the COVID-19 pandemic and are related to insufficient health care access, medication adherence, and monitoring (34). Patients might have delayed or avoided seeking emergency care when experiencing a life-threatening event during the height of the COVID-19 pandemic (35). Emergency department visits for heart attack and stroke decreased by 20% during the weeks after the declaration of COVID-19 as a national emergency on March 13, 2020, and hospital admissions for heart attack and stroke decreased during the pandemic (35). In addition, COVID-19 was associated with an increased risk for stroke and heart disease (36,37).

Chronic Lower Respiratory Disease

Despite the overall decrease during 2010–2020 (because of decreases observed in larger urban areas), the percentage of preventable premature deaths from CLRD was relatively stable in medium and small urban counties and rural counties during 2010–2015. During 2010–2022, the sharpest decline in preventable premature death from CLRD in urban areas occurred from 2019 through 2021 and could be the result of deaths from COVID-19 that otherwise would have been attributable to CLRD. Persons with CLRD (e.g., chronic obstructive pulmonary disease) are at increased risk for death from COVID-19 (38).

Limitations

The findings in this report are subject to at least six limitations. First, applying benchmarks (e.g., the three states with the lowest rates) to all urban-rural county categories facilitates comparisons but might not represent the lowest death rates achievable by certain subgroups. For example, large metropolitan areas met the benchmarks for cancer in 2019 and, therefore, had negative estimates of preventable premature deaths during 2019-2022. In these instances, negative estimates were truncated at zero to indicate that the 2010 benchmarks had been achieved. Using urban-ruralspecific benchmarks would likely result in larger numbers of preventable premature deaths in certain categories and larger rural-urban disparities in certain cases. However, using benchmarks specific to state, age, cause, and subgroup (e.g., urban-rural, sex, race, and ethnicity) also could lead to less stable estimates because of smaller numbers of deaths when stratifying by multiple demographic and geographic dimensions. Second, the differences cannot be attributed solely to population size and geographic location because risk factors do not occur randomly in populations and are related to wellknown social, demographic, environmental, economic, and geographic attributes of the neighborhoods in which persons live and work (39). Third, estimates of preventable premature

deaths using historical benchmarks (e.g., 2008-2010) might not reflect improvements in mortality that could have occurred in a later year. For example, the 2010 benchmarks for cancer are higher than if benchmarks were based on 2022 data because of decreases in cancer deaths during 2010-2022. Fourth, the numbers of preventable premature deaths by cause are not necessarily independent, and the numbers of potentially excess deaths from the five causes cannot be combined to generate a total. For example, the number of preventable premature deaths from cancer might be lower because persons with cancer died from another cause (e.g., heart disease). Fifth, deaths from certain causes might have increased partly because of COVID-19 and the pandemic. Specifically, misclassification of deaths from COVID-19 that were attributed to other causes (because of lack of testing or reporting on the death certificate) could have contributed to a greater number of deaths across other causes and by rural and urban areas. In addition, certain causes of death might have increased because of indirect pandemic-related effects (e.g., reduced access to emergency care or life-saving treatments). Finally, data for 2022 were based on provisional data from the first half of the year, and results might differ when the final data for the full year are available.

Future Directions

The findings in this report demonstrate the value of analyzing preventable premature deaths according to the six National Center for Health Statistics 2013 urban-rural county classifications. Reporting trends in preventable premature deaths over a 12-year period highlights differences over time and might aid in understanding underlying structural, environmental, and social risk factors. Because of increasing percentages of preventable premature deaths in recent years for specific causes of death and certain demographic groups, these data might augment traditional rate comparisons and help guide focused public health interventions. Comparing the findings in this report with data from tools such as the CDC Interactive Atlas of Heart Disease and Stroke (https://www. cdc.gov/dhdsp/maps/atlas/index.htm) might help identify social determinants of health, health care infrastructure, and public policies that could be related to increases or decreases in preventable premature deaths in specific nonmetropolitan areas. Detailed community-based evaluations might clarify how various risk factors and social determinants of health relate to premature mortality and related rural-urban disparities. In addition, other methods for developing benchmark rates might be helpful, including using benchmarks based on the

nonmetropolitan areas with the lowest death rates, or updating benchmarks based on more recent data, especially for causes of death (e.g., cancer) that have decreased substantially over time. In addition, more detailed analyses by race, ethnicity, and age, along with examining preventable deaths among persons aged >80 years, and preventable premature deaths from other causes (especially causes that are more prevalent in rural counties) might be informative.

Conclusion

Defining preventable premature deaths within the context of the five leading causes of death does not capture the full spectrum of preventable mortality. The degree to which these deaths could be prevented is related to various factors, including distinct prevention strategies, varying risk factors, and the availability of effective interventions, all of which vary by cause. Not all premature deaths are equally preventable among the leading causes or even within each specific leading cause category, as exemplified by certain types of cancer (40).

Routine tracking of preventable premature deaths by urbanrural county classification might facilitate the identification of areas with high prevalence of preventable premature mortality along with related geographic and urban-rural disparities in health outcomes. These disparities might be related to different levels of access to health care, social determinants of health, and other risk factors. Findings might help guide more focused interventions to reduce premature death from the five leading causes of death and reduce disparities by rural-urban residence and geographic region.

Acknowledgments

Health care professionals, medical examiners, coroners, vital registrars, and vital statistics offices across the United States.

Conflicts of Interest

All authors have completed and submitted the International Committee of Medical Journal Editors form for disclosure of potential conflicts of interest. No potential conflicts of interest were disclosed.

References

- 1. James CV, Moonesinghe R, Wilson-Frederick SM, Hall JE, Penman-Aguilar A, Bouye K. Racial/ethnic health disparities among rural adults—United States, 2012–2015. MMWR Surveill Summ 2017;66(No. SS-23):1–9. PMID:29145359 https://doi.org/10.15585/mmwr.ss6623a1
- Curtin SC, Spencer MR. Trends in death rates in urban and rural areas: United States, 1999–2019. NCHS data brief no. 417, September 2021. Hyattsville, MD: US Department of Health and Human Services, CDC, National Center for Health Statistics; 2021. https://www.cdc.gov/nchs/data/databriefs/db417.pdf

- Committee on Population; Division of Behavioral and Social Sciences and Education; Board on Health Care Services; National Research Council; Institute of Medicine. Measuring the risks and causes of premature death: summary of workshops. Washington, DC: The National Academies Press; 2015.
- Matthews KA, Croft JB, Liu Y, et al. Health-related behaviors by urbanrural county classification—United States, 2013. MMWR Surveill Summ 2017;66(No. SS-5):1–8. PMID:28151923 https://doi. org/10.15585/mmwr.ss6605a1
- Yoon PW, Bastian B, Anderson RN, Collins JL, Jaffe HW; CDC. Potentially preventable deaths from the five leading causes of death— United States, 2008–2010. MMWR Morb Mortal Wkly Rep 2014;63:369–74. PMID:24785982
- 6. Office of the Assistant Secretary of Health, Office of Disease Prevention and Health Promotion. Healthy People 2030. Social determinants of health [Internet]. Washington, DC: US Department of Health and Human Services, Office of the Secretary, Office of the Assistant Secretary of Health, Office of Disease Prevention and Health Promotion. https:// health.gov/healthypeople/priority-areas/social-determinants-health
- McMaughan DJ, Oloruntoba O, Smith ML. Socioeconomic status and access to healthcare: interrelated drivers for healthy aging. Front Public Health 2020;8:231. PMID:32626678 https://doi.org/10.3389/ fpubh.2020.00231
- Jensen L, Monnat SM, Green JJ, Hunter LM, Sliwinski MJ. Rural population health and aging: toward a multilevel and multidimensional research agenda for the 2020s. Am J Public Health 2020;110:1328–31. PMID:32673118 https://doi.org/10.2105/AJPH.2020.305782
- Hayward MD, Majmundar MK, eds. Future directions for the demography of aging: proceedings of a workshop. Washington, DC: The National Academies Press, 2018.
- 10. Committee on Population; Division of Behavioral and Social Sciences and Education; Board on Health Care Services; National Research Council; Institute of Medicine. Measuring the risks and causes of premature death: summary of workshops, Washington, DC: The National Academies Press; 2015.
- Henning-Smith C, Hernandez A, Ramirez M, et al. Dying too soon: county-level disparities in premature death by rurality, race, and ethnicity. Minneapolis, MN: University of Minnesota Rural Health Research Center; 2019. 1552267547UMNpolicybriefPrematureDeath.pdf
- 12. Gilbert PA, Laroche HH, Wallace RB, Parker EA, Curry SJ. Extending work on rural health disparities: a commentary on Matthews and colleagues' report. J Rural Health 2018;34:119–21. PMID:28397970 https://doi.org/10.1111/jrh.12241
- Lichter DT, Johnson KM. A demographic lifeline? Immigration and Hispanic population growth in rural America. Popul Res Policy Rev 2020;39:785–803. https://doi.org/10.1007/s11113-020-09605-8
- Khullar D, Chokshi DA. Challenges for immigrant health in the USA the road to crisis. Lancet 2019;393:2168–74. PMID:30981536 https://doi.org/10.1016/S0140-6736(19)30035-2
- Leider JP, Meit M, McCullough JM, et al. The state of rural public health: enduring needs in a new decade. Am J Public Health 2020;110:1283–90. PMID:32673103 https://doi.org/10.2105/ AJPH.2020.305728
- Ingram DD, Franco SJ. 2013 NCHS urban-rural classification scheme for counties [Internet]. Hyattsville, MD: US Department of Health and Human Services, CDC, National Center for Health Statistics; 2014. https://www.cdc.gov/nchs/data_access/urban_rural.htm
- Murphy SL, Xu J, Kochanek KD. Deaths: final data for 2010. Natl Vital Stat Rep 2013;61:1–117. PMID:24979972
- García MC, Rossen LM, Bastian B, et al. Potentially excess deaths from the five leading causes of death in metropolitan and nonmetropolitan counties—United States, 2010–2017. MMWR Surveill Summ 2019;68(No. SS-10):1–11. PMID:31697657 https://doi.org/10.15585/ mmwr.ss6810a1

- Hung P, Shi K, Probst JC, et al. Trends in cancer treatment service availability across critical access hospitals and prospective payment system hospitals. Med Care 2022;60:196–205. PMID:34432764 https://doi. org/10.1097/MLR.0000000000001635
- CDC. An update on cancer deaths in the United States. Atlanta, GA: US Department of Health and Human Services, CDC; 2022. https://stacks.cdc.gov/view/cdc/119728
- Zheng S, Schrijvers JJA, Greuter MJW, Kats-Ugurlu G, Lu W, de Bock GH. Effectiveness of colorectal cancer (CRC) screening on all-cause and CRC-specific mortality reduction: a systematic review and meta-analysis. Cancers (Basel) 2023;15:1948. PMID:37046609 https://doi. org/10.3390/cancers15071948
- Williams PA, Zaidi SK, Sengupta R. AACR cancer progress report 2023: advancing the frontiers of cancer science and medicine. Clin Cancer Res 2023;29:3850–1. PMID:37702621 https://doi.org/10.1158/1078-0432.CCR-23-2591
- 23. Barnes JM, Johnson KJ, Adjei Boakye E, et al. Early Medicaid expansion and cancer mortality. J Natl Cancer Inst 2021;113:1714–22. PMID:34259321 https://doi.org/10.1093/jnci/djab135
- 24. Cronin KA, Scott S, Firth AU, et al. Annual report to the nation on the status of cancer, part 1: national cancer statistics. Cancer 2022;128:4251–84. PMID:36301149 https://doi.org/10.1002/ cncr.34479
- 25. CDC. Using project ECHO and patient navigation to improve the health and wellness of cancer survivors in rural communities. Atlanta, GA: US Department of Health and Human Services, CDC; 2023. https://www.cdc.gov/cancer/ncccp/success-stories/echo-navigation.htm
- 26. Sahar L, Douangchai Wills VL, Liu KKA, et al. Geographic access to lung cancer screening among eligible adults living in rural and urban environments in the United States. Cancer 2022;128:1584–94. PMID:35167123 https://doi.org/10.1002/cncr.33996
- CDC. WISQARS fatal injury data visualization. Atlanta, GA: US Department of Health and Human Services, CDC; 2020. https://wisqars.cdc.gov/data/lcd/home
- 28. Roehler DR, Guy GP Jr, Jones CM. Buprenorphine prescription dispensing rates and characteristics following federal changes in prescribing policy, 2017–2018: a cross-sectional study. Drug Alcohol Depend 2020;213:108083. PMID:32505044 https://doi.org/10.1016/j. drugalcdep.2020.108083
- National Highway Traffic Safety Administration. US Census Bureau American Community Survey. Fatality Analysis Reporting System (FARS), 2016–2020. Washington, DC: National Highway Traffic Safety Administration. https://www-fars.nhtsa.dot.gov/Main/index.aspx
- 30. Shaw KM, West B, Kendi S, Zonfrillo MR, Sauber-Schatz E. Urban and rural child deaths from motor vehicle crashes: United States, 2015–2019. J Pediatr 2022;250:93–9. PMID:35809653 https://doi.org/10.1016/j.jpeds.2022.07.001
- 31. Stevens JA, Lee R. The potential to reduce falls and avert costs by clinically managing fall risk. Am J Prev Med 2018;55:290–7. PMID:30122212 https://doi.org/10.1016/j.amepre.2018.04.035
- 32. Sidney S, Lee C, Liu J, Khan SS, Lloyd-Jones DM, Rana JS. Age-adjusted mortality rates and age and risk-associated contributions to change in heart disease and stroke mortality, 2011–2019 and 2019–2020. JAMA Netw Open 2022;5:e223872. PMID:35319764 https://doi.org/10.1001/jamanetworkopen.2022.3872
- Laffin LJ, Kaufman HW, Chen Z, et al. Rise in blood pressure observed among US adults during the COVID-19 pandemic. Circulation 2022;145:235–7. PMID:34865499 https://doi.org/10.1161/ CIRCULATIONAHA.121.057075
- 34. Bress AP, Cohen JB, Anstey DE, et al. Inequities in hypertension control in the United States exposed and exacerbated by COVID-19 and the role of home blood pressure and virtual health care during and after the COVID-19 pandemic. J Am Heart Assoc 2021;10:e020997. PMID:34006116 https://doi.org/10.1161/JAHA.121.020997

Surveillance Summaries

- 35. Lange SJ, Ritchey MD, Goodman AB, et al. Potential indirect effects of the COVID-19 pandemic on use of emergency departments for acute life-threatening conditions—United States, January—May 2020. Am J Transplant 2020;20:2612–7. PMID:32862556 https://doi.org/10.1111/ajt.16239
- 36. Yang Q, Tong X, George MG, Chang A, Merritt RK. COVID-19 and risk of acute ischemic stroke among Medicare beneficiaries aged 65 years or older: self-controlled case series study. Neurology 2022;98:e778–89. PMID:35115387 https://doi.org/10.1212/WNL.0000000000013184
- 37. Italia L, Tomasoni D, Bisegna Š, et al. COVID-19 and heart failure: from epidemiology during the pandemic to myocardial injury, myocarditis, and heart failure sequelae. Front Cardiovasc Med 2021;8:713560. PMID:34447795 https://doi.org/10.3389/fcvm.2021.713560
- 38. Schultze A, Walker AJ, MacKenna B, et al.; OpenSAFELY Collaborative. Risk of COVID-19-related death among patients with chronic obstructive pulmonary disease or asthma prescribed inhaled corticosteroids: an observational cohort study using the OpenSAFELY platform. Lancet Respir Med 2020;8:1106–20. PMID:32979987 https://doi.org/10.1016/S2213-2600(20)30415-X
- 39. Frieden TR. Foreword. In: CDC health disparities and inequalities report—United States, 2013. MMWR Suppl 2013;62:(No. Suppl-3):1–2. PMID:24264482
- Kilic H, Arguder E, Karalezli A, et al. Effect of chronic lung diseases on mortality of prevariant COVID-19 pneumonia patients. Front Med (Lausanne) 2022;9:957598. PMID:36314036 https://doi.org/10.3389/ fmed.2022.957598

The *Morbidity and Mortality Weekly Report (MMWR)* Series is prepared by the Centers for Disease Control and Prevention (CDC) and is available free of charge in electronic format. To receive an electronic copy each week, visit *MMWR* at https://www.cdc.gov/mmwr/index.html.

Readers who have difficulty accessing this PDF file may access the HTML file at https://www.cdc.gov/mmwr/volumes/73/ss/ss7302a1.htm?s_cid=ss7302a1_w. Address all inquiries about the MMWR Series to Editor-in-Chief, MMWR Series, Mailstop V25-5, CDC, 1600 Clifton Rd., N.E., Atlanta, GA 30329-4027 or to mmurq@cdc.gov.

All material in the *MMWR* Series is in the public domain and may be used and reprinted without permission; citation as to source, however, is appreciated. *MMWR* and *Morbidity and Mortality Weekly Report* are service marks of the U.S. Department of Health and Human Services.

Use of trade names and commercial sources is for identification only and does not imply endorsement by the U.S. Department of Health and Human Services.

References to non-CDC sites on the Internet are provided as a service to MMWR readers and do not constitute or imply endorsement of these organizations or their programs by CDC or the U.S. Department of Health and Human Services. CDC is not responsible for the content of these sites. URL addresses listed in MMWR were current as of the date of publication.

ISSN: 1546-0738 (Print)