

Data Brief 396. Secondhand Smoke Exposure Among Nonsmoking Adults: United States, 2015–2018

Data table for Figure 1. Percentage of secondhand smoke exposure among nonsmoking adults, by sex and age: United States, 2015–2018

Sex and age	Percent	Standard error
All		
18 and over	20.8	1.2
18–39	25.6	1.4
40–59	19.1	1.7
60 and over	17.6	1.4
Men		
18 and over	20.8	1.7
18–39	25.0	2.4
40–59	18.8	2.4
60 and over	18.7	2.3
Women		
18 and over	20.7	1.0
18–39	26.0	1.5
40–59	19.3	1.5
60 and over	16.7	1.3

NOTE: Secondhand smoke is defined as serum cotinine levels of 0.05–10.00 ng/mL.

SOURCE: National Center for Health Statistics, National Health and Nutrition Examination Survey, 2015–2018.

Data Brief 396. Secondhand Smoke Exposure Among Nonsmoking Adults: United States, 2015–2018

Data table for Figure 2. Percentage of secondhand smoke exposure among nonsmoking adults, by sex and race and Hispanic origin: United States, 2015–2018

Sex and race and Hispanic origin	Percent	Standard error
All		
Non-Hispanic white	18.4	1.5
Non-Hispanic black	39.7	1.9
Non-Hispanic Asian	20.9	1.8
Hispanic	17.2	1.1
Men		
Non-Hispanic white	18.8	2.2
Non-Hispanic black	38.4	2.9
Non-Hispanic Asian	20.2	2.1
Hispanic	18.1	1.7
Women		
Non-Hispanic white	18.1	1.3
Non-Hispanic black	40.5	2.1
Non-Hispanic Asian	21.4	2.3
Hispanic	16.4	1.5

NOTES: Secondhand smoke is defined as serum cotinine levels of 0.05–10.00 ng/mL. Only persons reporting a single race are included.

SOURCE: National Center for Health Statistics, National Health and Nutrition Examination Survey, 2015–2018.

Data Brief 396. Secondhand Smoke Exposure Among Nonsmoking Adults: United States, 2015–2018

Data table for Figure 3. Percentage of secondhand smoke exposure among nonsmoking adults, by education and income: United States, 2015–2018

Characteristic	Percent	Standard error
Education		
Less than high school	27.3	2.4
High school graduate or GED	29.5	1.8
Some college or more	16.8	1.1
Family income		
Less than 100% FPL	35.1	2.9
100% to less than 200% FPL	30.2	1.9
200% to less than 400% FPL	21.1	1.7
At or above 400% FPL	11.8	1.4

NOTES: Secondhand smoke is defined as serum cotinine levels of 0.05–10.00 ng/mL. FPL is federal poverty level.

SOURCE: National Center for Health Statistics, National Health and Nutrition Examination Survey, 2015–2018.

Data Brief 396. Secondhand Smoke Exposure Among Nonsmoking Adults: United States, 2015–2018

Data table for Figure 4. Trends in the percentage of secondhand smoke exposure among nonsmoking adults: United States, 2009–2018

Survey years	Percent	Standard error
2009–2010	27.7	1.4
2011–2012	22.1	1.3
2013–2014	22.4	1.9
2015–2016	20.8	1.8
2017–2018	20.7	1.5

NOTE: Secondhand smoke is defined as serum cotinine levels of 0.05–10.00 ng/mL.

SOURCE: National Center for Health Statistics, National Health and Nutrition Examination Survey, 2009–2018.