

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, all ages				
...	All causes	2,448,017	100.0	825.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	652,091	26.6	220.0
2	Malignant neoplasms (C00-C97)	559,312	22.8	188.7
3	Cerebrovascular diseases (I60-I69)	143,579	5.9	48.4
4	Chronic lower respiratory diseases (J40-J47)	130,933	5.3	44.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	117,809	4.8	39.7
6	Diabetes mellitus (E10-E14)	75,119	3.1	25.3
7	Alzheimer's disease (G30)	71,599	2.9	24.2
8	Influenza and pneumonia (J10-J18)	63,001	2.6	21.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	43,901	1.8	14.8
10	Septicemia (A40-A41)	34,136	1.4	11.5
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	32,637	1.3	11.0
12	Chronic liver disease and cirrhosis (K70,K73-K74)	27,530	1.1	9.3
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	24,902	1.0	8.4
14	Parkinson's disease (G20-G21)	19,544	0.8	6.6
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	18,124	0.7	6.1
...	All other causes (Residual)	433,800	17.7	146.4
All races, both sexes, 1-4 years				
...	All causes	4,756	100.0	29.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,664	35.0	10.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	522	11.0	3.2
3	Malignant neoplasms (C00-C97)	377	7.9	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	375	7.9	2.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	151	3.2	0.9
6	Influenza and pneumonia (J10-J18)	110	2.3	0.7
7	Septicemia (A40-A41)	85	1.8	0.5
8	Cerebrovascular diseases (I60-I69)	62	1.3	0.4
9	Certain conditions originating in the perinatal period (P00-P96)	58	1.2	0.4
10	Chronic lower respiratory diseases (J40-J47)	56	1.2	0.3
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	52	1.1	0.3
12	Meningitis (G00,G03)	32	0.7	0.2
13	Anemias (D50-D64)	23	0.5	0.1
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	16	0.3	*
...	All other causes (Residual)	1,157	24.3	7.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 5-9 years				
...	All causes	2,837	100.0	14.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,072	37.8	5.5
2	Malignant neoplasms (C00-C97)	485	17.1	2.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	196	6.9	1.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	121	4.3	0.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	106	3.7	0.5
6	Cerebrovascular diseases (I60-I69)	52	1.8	0.3
7	Influenza and pneumonia (J10-J18)	51	1.8	0.3
8	Chronic lower respiratory diseases (J40-J47)	49	1.7	0.3
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	40	1.4	0.2
10	Septicemia (A40-A41)	36	1.3	0.2
11	Certain conditions originating in the perinatal period (P00-P96)	14	0.5	*
12	Meningitis (G00,G03)	13	0.5	*
13	Anemias (D50-D64)	11	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	10	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.3	*
...	All other causes (Residual)	572	20.2	2.9
All races, both sexes, 10-14 years				
...	All causes	3,765	100.0	18.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,343	35.7	6.4
2	Malignant neoplasms (C00-C97)	515	13.7	2.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	270	7.2	1.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	220	5.8	1.1
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	200	5.3	1.0
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	146	3.9	0.7
7	Influenza and pneumonia (J10-J18)	55	1.5	0.3
7	Chronic lower respiratory diseases (J40-J47)	55	1.5	0.3
9	Septicemia (A40-A41)	45	1.2	0.2
10	Cerebrovascular diseases (I60-I69)	43	1.1	0.2
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	36	1.0	0.2
12	Diabetes mellitus (E10-E14)	34	0.9	0.2
13	Anemias (D50-D64)	15	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	0.4	*
15	Meningitis (G00,G03)	13	0.3	*
...	All other causes (Residual)	761	20.2	3.6

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 15-19 years				
...	All causes	13,703	100.0	65.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,616	48.3	31.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,076	15.1	9.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,613	11.8	7.7
4	Malignant neoplasms (C00-C97)	731	5.3	3.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	389	2.8	1.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	253	1.8	1.2
7	Cerebrovascular diseases (I60-I69)	76	0.6	0.4
8	Influenza and pneumonia (J10-J18)	68	0.5	0.3
9	Diabetes mellitus (E10-E14)	67	0.5	0.3
10	Septicemia (A40-A41)	61	0.4	0.3
11	Chronic lower respiratory diseases (J40-J47)	60	0.4	0.3
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	50	0.4	0.2
13	Pregnancy, childbirth and the puerperium (O00-099)	42	0.3	0.2
14	Anemias (D50-D64)	40	0.3	0.2
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	34	0.2	0.2
...	All other causes (Residual)	1,527	11.1	7.3
All races, both sexes, 20-24 years				
...	All causes	20,531	100.0	97.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,137	44.5	43.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,390	16.5	16.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,599	12.7	12.4
4	Malignant neoplasms (C00-C97)	986	4.8	4.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	730	3.6	3.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	251	1.2	1.2
7	Pregnancy, childbirth and the puerperium (O00-099)	141	0.7	0.7
8	Diabetes mellitus (E10-E14)	135	0.7	0.6
9	Human immunodeficiency virus (HIV) disease (B20-B24)	131	0.6	0.6
10	Cerebrovascular diseases (I60-I69)	120	0.6	0.6
11	Influenza and pneumonia (J10-J18)	104	0.5	0.5
12	Chronic lower respiratory diseases (J40-J47)	88	0.4	0.4
13	Septicemia (A40-A41)	87	0.4	0.4
14	Anemias (D50-D64)	63	0.3	0.3
15	Legal intervention (Y35,Y89.0)	61	0.3	0.3
...	All other causes (Residual)	2,508	12.2	11.9

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 25-29 years				
...	All causes	19,568	100.0	97.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,215	36.9	36.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,735	14.0	13.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,377	12.1	11.8
4	Malignant neoplasms (C00-C97)	1,337	6.8	6.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,180	6.0	5.9
6	Human immunodeficiency virus (HIV) disease (B20-B24)	412	2.1	2.1
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	232	1.2	1.2
8	Diabetes mellitus (E10-E14)	210	1.1	1.0
9	Cerebrovascular diseases (I60-I69)	192	1.0	1.0
10	Pregnancy, childbirth and the puerperium (O00-099)	155	0.8	0.8
11	Influenza and pneumonia (J10-J18)	147	0.8	0.7
12	Septicemia (A40-A41)	129	0.7	0.6
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	112	0.6	0.6
14	Chronic lower respiratory diseases (J40-J47)	100	0.5	0.5
15	Anemias (D50-D64)	82	0.4	0.4
...	All other causes (Residual)	2,953	15.1	14.7
All races, both sexes, 30-34 years				
...	All causes	22,357	100.0	111.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,782	30.3	33.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,613	11.7	13.0
3	Malignant neoplasms (C00-C97)	2,264	10.1	11.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,069	9.3	10.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,017	9.0	10.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	906	4.1	4.5
7	Diabetes mellitus (E10-E14)	407	1.8	2.0
8	Cerebrovascular diseases (I60-I69)	354	1.6	1.8
9	Chronic liver disease and cirrhosis (K70,K73-K74)	236	1.1	1.2
10	Influenza and pneumonia (J10-J18)	207	0.9	1.0
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	204	0.9	1.0
12	Septicemia (A40-A41)	182	0.8	0.9
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	173	0.8	0.9
14	Chronic lower respiratory diseases (J40-J47)	158	0.7	0.8
15	Pregnancy, childbirth and the puerperium (O00-099)	157	0.7	0.8
...	All other causes (Residual)	3,628	16.2	18.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 35-39 years				
...	All causes	31,420	100.0	149.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,419	23.6	35.3
2	Malignant neoplasms (C00-C97)	4,378	13.9	20.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,076	13.0	19.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,893	9.2	13.8
5	Human immunodeficiency virus (HIV) disease (B20-B24)	1,693	5.4	8.1
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,582	5.0	7.5
7	Diabetes mellitus (E10-E14)	762	2.4	3.6
8	Chronic liver disease and cirrhosis (K70,K73-K74)	719	2.3	3.4
9	Cerebrovascular diseases (I60-I69)	693	2.2	3.3
10	Influenza and pneumonia (J10-J18)	318	1.0	1.5
11	Septicemia (A40-A41)	310	1.0	1.5
12	Chronic lower respiratory diseases (J40-J47)	261	0.8	1.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	260	0.8	1.2
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	233	0.7	1.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	140	0.4	0.7
...	All other causes (Residual)	5,683	18.1	27.1
All races, both sexes, 40-44 years				
...	All causes	53,365	100.0	233.4
1	Malignant neoplasms (C00-C97)	10,188	19.1	44.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,500	17.8	41.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,612	16.1	37.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,657	6.9	16.0
5	Human immunodeficiency virus (HIV) disease (B20-B24)	2,670	5.0	11.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1,969	3.7	8.6
7	Cerebrovascular diseases (I60-I69)	1,567	2.9	6.9
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,527	2.9	6.7
9	Diabetes mellitus (E10-E14)	1,283	2.4	5.6
10	Chronic lower respiratory diseases (J40-J47)	629	1.2	2.8
11	Influenza and pneumonia (J10-J18)	616	1.2	2.7
12	Septicemia (A40-A41)	530	1.0	2.3
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	482	0.9	2.1
14	Viral hepatitis (B15-B19)	385	0.7	1.7
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	288	0.5	1.3
...	All other causes (Residual)	9,462	17.7	41.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 45-49 years				
...	All causes	79,383	100.0	353.1
1	Malignant neoplasms (C00-C97)	19,452	24.5	86.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,295	19.3	68.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,019	12.6	44.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,764	4.7	16.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	3,376	4.3	15.0
6	Cerebrovascular diseases (I60-I69)	2,629	3.3	11.7
7	Human immunodeficiency virus (HIV) disease (B20-B24)	2,546	3.2	11.3
8	Diabetes mellitus (E10-E14)	2,233	2.8	9.9
9	Chronic lower respiratory diseases (J40-J47)	1,455	1.8	6.5
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,223	1.5	5.4
11	Influenza and pneumonia (J10-J18)	955	1.2	4.2
12	Viral hepatitis (B15-B19)	949	1.2	4.2
13	Septicemia (A40-A41)	898	1.1	4.0
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	864	1.1	3.8
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	504	0.6	2.2
...	All other causes (Residual)	13,221	16.7	58.8
All races, both sexes, 50-54 years				
...	All causes	104,147	100.0	520.8
1	Malignant neoplasms (C00-C97)	30,953	29.7	154.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	22,808	21.9	114.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,320	8.0	41.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	4,141	4.0	20.7
5	Cerebrovascular diseases (I60-I69)	3,752	3.6	18.8
6	Diabetes mellitus (E10-E14)	3,458	3.3	17.3
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,227	3.1	16.1
8	Chronic lower respiratory diseases (J40-J47)	2,522	2.4	12.6
9	Human immunodeficiency virus (HIV) disease (B20-B24)	1,970	1.9	9.9
10	Viral hepatitis (B15-B19)	1,365	1.3	6.8
11	Septicemia (A40-A41)	1,313	1.3	6.6
12	Influenza and pneumonia (J10-J18)	1,228	1.2	6.1
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,164	1.1	5.8
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	837	0.8	4.2
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	662	0.6	3.3
...	All other causes (Residual)	16,427	15.8	82.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 55-59 years				
...	All causes	127,478	100.0	734.6
1	Malignant neoplasms (C00-C97)	44,358	34.8	255.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	29,681	23.3	171.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,168	4.8	35.5
4	Diabetes mellitus (E10-E14)	5,150	4.0	29.7
5	Chronic lower respiratory diseases (J40-J47)	4,672	3.7	26.9
6	Cerebrovascular diseases (I60-I69)	4,480	3.5	25.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	3,995	3.1	23.0
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,477	1.9	14.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,835	1.4	10.6
10	Septicemia (A40-A41)	1,822	1.4	10.5
11	Influenza and pneumonia (J10-J18)	1,522	1.2	8.8
12	Human immunodeficiency virus (HIV) disease (B20-B24)	1,094	0.9	6.3
13	Viral hepatitis (B15-B19)	928	0.7	5.3
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	873	0.7	5.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	579	0.5	3.3
...	All other causes (Residual)	17,844	14.0	102.8
All races, both sexes, 60-64 years				
...	All causes	147,823	100.0	1,136.9
1	Malignant neoplasms (C00-C97)	54,882	37.1	422.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	35,527	24.0	273.2
3	Chronic lower respiratory diseases (J40-J47)	8,075	5.5	62.1
4	Diabetes mellitus (E10-E14)	6,151	4.2	47.3
5	Cerebrovascular diseases (I60-I69)	5,548	3.8	42.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,685	3.2	36.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	3,131	2.1	24.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,306	1.6	17.7
9	Septicemia (A40-A41)	2,090	1.4	16.1
10	Influenza and pneumonia (J10-J18)	1,900	1.3	14.6
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,733	1.2	13.3
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,081	0.7	8.3
13	Aortic aneurysm and dissection (I71)	878	0.6	6.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	739	0.5	5.7
15	Human immunodeficiency virus (HIV) disease (B20-B24)	518	0.4	4.0
...	All other causes (Residual)	18,579	12.6	142.9

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 65-69 years				
...	All causes	172,236	100.0	1,700.0
1	Malignant neoplasms (C00-C97)	63,036	36.6	622.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	41,084	23.9	405.5
3	Chronic lower respiratory diseases (J40-J47)	12,016	7.0	118.6
4	Cerebrovascular diseases (I60-I69)	7,473	4.3	73.8
5	Diabetes mellitus (E10-E14)	7,260	4.2	71.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,060	2.4	40.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,958	1.7	29.2
8	Septicemia (A40-A41)	2,602	1.5	25.7
9	Chronic liver disease and cirrhosis (K70,K73-K74)	2,569	1.5	25.4
10	Influenza and pneumonia (J10-J18)	2,486	1.4	24.5
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,413	0.8	13.9
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,198	0.7	11.8
13	Aortic aneurysm and dissection (I71)	1,185	0.7	11.7
14	Alzheimer's disease (G30)	983	0.6	9.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	966	0.6	9.5
...	All other causes (Residual)	20,947	12.2	206.8
All races, both sexes, 70-74 years				
...	All causes	226,119	100.0	2,657.6
1	Malignant neoplasms (C00-C97)	75,410	33.3	886.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	55,645	24.6	654.0
3	Chronic lower respiratory diseases (J40-J47)	17,894	7.9	210.3
4	Cerebrovascular diseases (I60-I69)	11,366	5.0	133.6
5	Diabetes mellitus (E10-E14)	8,923	3.9	104.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,572	2.0	53.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,362	1.9	51.3
8	Influenza and pneumonia (J10-J18)	4,137	1.8	48.6
9	Septicemia (A40-A41)	3,471	1.5	40.8
10	Alzheimer's disease (G30)	2,830	1.3	33.3
11	Chronic liver disease and cirrhosis (K70,K73-K74)	2,497	1.1	29.3
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,893	0.8	22.2
13	Parkinson's disease (G20-G21)	1,729	0.8	20.3
14	Aortic aneurysm and dissection (I71)	1,685	0.7	19.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,384	0.6	16.3
...	All other causes (Residual)	28,321	12.5	332.9

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 75-79 years				
...	All causes	307,888	100.0	4,154.0
1	Malignant neoplasms (C00-C97)	85,861	27.9	1,158.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	80,843	26.3	1,090.7
3	Chronic lower respiratory diseases (J40-J47)	24,501	8.0	330.6
4	Cerebrovascular diseases (I60-I69)	18,800	6.1	253.6
5	Diabetes mellitus (E10-E14)	11,218	3.6	151.4
6	Alzheimer's disease (G30)	7,596	2.5	102.5
7	Influenza and pneumonia (J10-J18)	7,123	2.3	96.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,348	2.1	85.6
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,126	2.0	82.7
10	Septicemia (A40-A41)	4,715	1.5	63.6
11	Parkinson's disease (G20-G21)	3,724	1.2	50.2
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	3,000	1.0	40.5
13	Aortic aneurysm and dissection (I71)	2,340	0.8	31.6
14	Chronic liver disease and cirrhosis (K70,K73-K74)	2,208	0.7	29.8
15	Pneumonitis due to solids and liquids (J69)	2,109	0.7	28.5
...	All other causes (Residual)	41,376	13.4	558.2
All races, both sexes, 80-84 years				
...	All causes	378,777	100.0	6,712.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	109,850	29.0	1,946.8
2	Malignant neoplasms (C00-C97)	80,560	21.3	1,427.7
3	Cerebrovascular diseases (I60-I69)	28,059	7.4	497.3
4	Chronic lower respiratory diseases (J40-J47)	25,832	6.8	457.8
5	Alzheimer's disease (G30)	15,543	4.1	275.5
6	Diabetes mellitus (E10-E14)	11,918	3.1	211.2
7	Influenza and pneumonia (J10-J18)	11,440	3.0	202.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,055	2.1	142.8
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,728	2.0	137.0
10	Septicemia (A40-A41)	5,911	1.6	104.8
11	Parkinson's disease (G20-G21)	5,570	1.5	98.7
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	4,256	1.1	75.4
13	Pneumonitis due to solids and liquids (J69)	3,449	0.9	61.1
14	Aortic aneurysm and dissection (I71)	2,631	0.7	46.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,526	0.7	44.8
...	All other causes (Residual)	55,449	14.6	982.7

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 85-89 years				
...	All causes	351,398	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	113,840	32.4	---
2	Malignant neoplasms (C00-C97)	52,871	15.0	---
3	Cerebrovascular diseases (I60-I69)	28,909	8.2	---
4	Alzheimer's disease (G30)	19,870	5.7	---
5	Chronic lower respiratory diseases (J40-J47)	19,338	5.5	---
6	Influenza and pneumonia (J10-J18)	13,109	3.7	---
7	Diabetes mellitus (E10-E14)	9,313	2.7	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,662	2.2	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,303	2.1	---
10	Septicemia (A40-A41)	5,125	1.5	---
11	Parkinson's disease (G20-G21)	4,624	1.3	---
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	4,601	1.3	---
13	Pneumonitis due to solids and liquids (J69)	3,987	1.1	---
14	Atherosclerosis (I70)	2,710	0.8	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,231	0.6	---
...	All other causes (Residual)	55,905	15.9	---
All races, both sexes, 90-94 years				
...	All causes	239,706	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	86,322	36.0	---
2	Malignant neoplasms (C00-C97)	23,759	9.9	---
3	Cerebrovascular diseases (I60-I69)	20,446	8.5	---
4	Alzheimer's disease (G30)	16,058	6.7	---
5	Influenza and pneumonia (J10-J18)	10,979	4.6	---
6	Chronic lower respiratory diseases (J40-J47)	9,809	4.1	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,937	2.1	---
8	Diabetes mellitus (E10-E14)	4,910	2.0	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,791	2.0	---
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	3,891	1.6	---
11	Septicemia (A40-A41)	3,138	1.3	---
12	Pneumonitis due to solids and liquids (J69)	2,993	1.2	---
13	Atherosclerosis (I70)	2,485	1.0	---
14	Parkinson's disease (G20-G21)	2,163	0.9	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,262	0.5	---
...	All other causes (Residual)	41,763	17.4	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 95-99 years				
...	All causes	92,761	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	35,457	38.2	---
2	Cerebrovascular diseases (I60-I69)	7,406	8.0	---
3	Alzheimer's disease (G30)	6,770	7.3	---
4	Malignant neoplasms (C00-C97)	6,068	6.5	---
5	Influenza and pneumonia (J10-J18)	4,937	5.3	---
6	Chronic lower respiratory diseases (J40-J47)	2,923	3.2	---
7	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,811	2.0	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,778	1.9	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,769	1.9	---
10	Diabetes mellitus (E10-E14)	1,455	1.6	---
11	Atherosclerosis (I70)	1,238	1.3	---
12	Pneumonitis due to solids and liquids (J69)	1,153	1.2	---
13	Septicemia (A40-A41)	1,098	1.2	---
14	Parkinson's disease (G20-G21)	486	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	363	0.4	---
...	All other causes (Residual)	18,049	19.5	---
All races, both sexes, 100 years and over				
...	All causes	19,304	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,885	40.8	---
2	Cerebrovascular diseases (I60-I69)	1,422	7.4	---
3	Influenza and pneumonia (J10-J18)	1,242	6.4	---
4	Alzheimer's disease (G30)	1,208	6.3	---
5	Malignant neoplasms (C00-C97)	757	3.9	---
6	Chronic lower respiratory diseases (J40-J47)	403	2.1	---
7	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	400	2.1	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	371	1.9	---
9	Atherosclerosis (I70)	340	1.8	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	325	1.7	---
11	Diabetes mellitus (E10-E14)	225	1.2	---
12	Pneumonitis due to solids and liquids (J69)	219	1.1	---
13	Septicemia (A40-A41)	183	0.9	---
14	Anemias (D50-D64)	83	0.4	---
15	Nutritional deficiencies (E40-E64)	71	0.4	---
...	All other causes (Residual)	4,170	21.6	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, all ages				
...	All causes	1,207,675	100.0	827.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	322,841	26.7	221.1
2	Malignant neoplasms (C00-C97)	290,422	24.0	198.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	76,375	6.3	52.3
4	Chronic lower respiratory diseases (J40-J47)	62,435	5.2	42.8
5	Cerebrovascular diseases (I60-I69)	56,586	4.7	38.8
6	Diabetes mellitus (E10-E14)	36,538	3.0	25.0
7	Influenza and pneumonia (J10-J18)	28,052	2.3	19.2
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	25,907	2.1	17.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21,268	1.8	14.6
10	Alzheimer's disease (G30)	20,559	1.7	14.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	17,937	1.5	12.3
12	Septicemia (A40-A41)	15,322	1.3	10.5
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	14,376	1.2	9.8
14	Parkinson's disease (G20-G21)	11,247	0.9	7.7
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	9,458	0.8	6.5
...	All other causes (Residual)	198,352	16.4	135.9
All races, male, 1-4 years				
...	All causes	2,765	100.0	33.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,027	37.1	12.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	264	9.5	3.2
3	Malignant neoplasms (C00-C97)	217	7.8	2.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	214	7.7	2.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	81	2.9	1.0
6	Influenza and pneumonia (J10-J18)	56	2.0	0.7
7	Septicemia (A40-A41)	47	1.7	0.6
8	Cerebrovascular diseases (I60-I69)	41	1.5	0.5
9	Certain conditions originating in the perinatal period (P00-P96)	36	1.3	0.4
10	Chronic lower respiratory diseases (J40-J47)	33	1.2	0.4
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	31	1.1	0.4
12	Meningitis (G00,G03)	21	0.8	0.3
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	10	0.4	*
15	Anemias (D50-D64)	9	0.3	*
...	All other causes (Residual)	666	24.1	8.0

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 5-9 years				
...	All causes	1,556	100.0	15.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	627	40.3	6.3
2	Malignant neoplasms (C00-C97)	263	16.9	2.6
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	86	5.5	0.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	68	4.4	0.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	50	3.2	0.5
6	Chronic lower respiratory diseases (J40-J47)	28	1.8	0.3
7	Cerebrovascular diseases (I60-I69)	27	1.7	0.3
8	Influenza and pneumonia (J10-J18)	25	1.6	0.3
9	Septicemia (A40-A41)	24	1.5	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	1.2	*
11	Certain conditions originating in the perinatal period (P00-P96)	7	0.4	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.4	*
13	Anemias (D50-D64)	6	0.4	*
13	Meningitis (G00,G03)	6	0.4	*
15	Diseases of appendix (K35-K38)	5	0.3	*
...	All other causes (Residual)	308	19.8	3.1
All races, male, 10-14 years				
...	All causes	2,297	100.0	21.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	866	37.7	8.1
2	Malignant neoplasms (C00-C97)	294	12.8	2.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	202	8.8	1.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	136	5.9	1.3
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	109	4.7	1.0
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	81	3.5	0.8
7	Chronic lower respiratory diseases (J40-J47)	35	1.5	0.3
8	Cerebrovascular diseases (I60-I69)	28	1.2	0.3
8	Influenza and pneumonia (J10-J18)	28	1.2	0.3
10	Septicemia (A40-A41)	25	1.1	0.2
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	0.7	*
12	Diabetes mellitus (E10-E14)	13	0.6	*
13	Meningitis (G00,G03)	9	0.4	*
14	Anemias (D50-D64)	8	0.3	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	7	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.3	*
...	All other causes (Residual)	432	18.8	4.0

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 15-19 years				
...	All causes	9,886	100.0	91.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,645	47.0	43.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,815	18.4	16.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,303	13.2	12.1
4	Malignant neoplasms (C00-C97)	442	4.5	4.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	268	2.7	2.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	172	1.7	1.6
7	Influenza and pneumonia (J10-J18)	49	0.5	0.5
8	Chronic lower respiratory diseases (J40-J47)	43	0.4	0.4
9	Diabetes mellitus (E10-E14)	40	0.4	0.4
10	Cerebrovascular diseases (I60-I69)	34	0.3	0.3
11	Septicemia (A40-A41)	33	0.3	0.3
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	28	0.3	0.3
13	Legal intervention (Y35,Y89.0)	22	0.2	0.2
14	Anemias (D50-D64)	21	0.2	0.2
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	0.1	*
...	All other causes (Residual)	957	9.7	8.9
All races, male, 20-24 years				
...	All causes	15,623	100.0	143.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,182	46.0	66.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,950	18.9	27.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,195	14.0	20.2
4	Malignant neoplasms (C00-C97)	593	3.8	5.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	502	3.2	4.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	154	1.0	1.4
7	Human immunodeficiency virus (HIV) disease (B20-B24)	78	0.5	0.7
8	Diabetes mellitus (E10-E14)	75	0.5	0.7
9	Influenza and pneumonia (J10-J18)	68	0.4	0.6
10	Cerebrovascular diseases (I60-I69)	60	0.4	0.6
10	Legal intervention (Y35,Y89.0)	60	0.4	0.6
12	Chronic lower respiratory diseases (J40-J47)	52	0.3	0.5
13	Septicemia (A40-A41)	49	0.3	0.5
14	Anemias (D50-D64)	36	0.2	0.3
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	31	0.2	0.3
...	All other causes (Residual)	1,538	9.8	14.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 25-29 years				
...	All causes	14,242	100.0	138.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,622	39.5	54.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,342	16.4	22.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,975	13.9	19.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	814	5.7	7.9
5	Malignant neoplasms (C00-C97)	749	5.3	7.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	249	1.7	2.4
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	132	0.9	1.3
8	Diabetes mellitus (E10-E14)	118	0.8	1.1
9	Cerebrovascular diseases (I60-I69)	109	0.8	1.1
10	Influenza and pneumonia (J10-J18)	79	0.6	0.8
11	Septicemia (A40-A41)	72	0.5	0.7
12	Chronic lower respiratory diseases (J40-J47)	58	0.4	0.6
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	56	0.4	0.5
14	Chronic liver disease and cirrhosis (K70,K73-K74)	50	0.4	0.5
15	Anemias (D50-D64)	44	0.3	0.4
15	Legal intervention (Y35,Y89.0)	44	0.3	0.4
...	All other causes (Residual)	1,729	12.1	16.8
All races, male, 30-34 years				
...	All causes	15,041	100.0	148.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,043	33.5	49.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,091	13.9	20.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,666	11.1	16.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,392	9.3	13.7
5	Malignant neoplasms (C00-C97)	1,056	7.0	10.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	560	3.7	5.5
7	Diabetes mellitus (E10-E14)	254	1.7	2.5
8	Cerebrovascular diseases (I60-I69)	192	1.3	1.9
9	Chronic liver disease and cirrhosis (K70,K73-K74)	146	1.0	1.4
10	Influenza and pneumonia (J10-J18)	117	0.8	1.2
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	112	0.7	1.1
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	97	0.6	1.0
13	Septicemia (A40-A41)	80	0.5	0.8
14	Chronic lower respiratory diseases (J40-J47)	72	0.5	0.7
15	Legal intervention (Y35,Y89.0)	65	0.4	0.6
...	All other causes (Residual)	2,098	13.9	20.7

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 35-39 years				
...	All causes	20,011	100.0	189.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,347	26.7	50.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,828	14.1	26.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,259	11.3	21.4
4	Malignant neoplasms (C00-C97)	1,842	9.2	17.4
5	Human immunodeficiency virus (HIV) disease (B20-B24)	1,186	5.9	11.2
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,179	5.9	11.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	476	2.4	4.5
8	Diabetes mellitus (E10-E14)	456	2.3	4.3
9	Cerebrovascular diseases (I60-I69)	355	1.8	3.4
10	Influenza and pneumonia (J10-J18)	192	1.0	1.8
11	Septicemia (A40-A41)	143	0.7	1.4
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	141	0.7	1.3
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	129	0.6	1.2
14	Chronic lower respiratory diseases (J40-J47)	124	0.6	1.2
15	Aortic aneurysm and dissection (I71)	94	0.5	0.9
...	All other causes (Residual)	3,260	16.3	30.9
All races, male, 40-44 years				
...	All causes	33,298	100.0	292.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,679	20.1	58.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,103	18.3	53.6
3	Malignant neoplasms (C00-C97)	4,496	13.5	39.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,804	8.4	24.6
5	Human immunodeficiency virus (HIV) disease (B20-B24)	1,960	5.9	17.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1,310	3.9	11.5
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,138	3.4	10.0
8	Diabetes mellitus (E10-E14)	814	2.4	7.2
9	Cerebrovascular diseases (I60-I69)	793	2.4	7.0
10	Influenza and pneumonia (J10-J18)	348	1.0	3.1
11	Chronic lower respiratory diseases (J40-J47)	302	0.9	2.7
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	282	0.8	2.5
13	Septicemia (A40-A41)	271	0.8	2.4
14	Viral hepatitis (B15-B19)	262	0.8	2.3
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	174	0.5	1.5
...	All other causes (Residual)	5,562	16.7	48.9

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 45-49 years				
...	All causes	49,279	100.0	443.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,967	22.3	98.7
2	Malignant neoplasms (C00-C97)	9,263	18.8	83.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,057	14.3	63.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,826	5.7	25.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,361	4.8	21.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	1,913	3.9	17.2
7	Diabetes mellitus (E10-E14)	1,413	2.9	12.7
8	Cerebrovascular diseases (I60-I69)	1,345	2.7	12.1
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	927	1.9	8.3
10	Chronic lower respiratory diseases (J40-J47)	707	1.4	6.4
11	Viral hepatitis (B15-B19)	696	1.4	6.3
12	Influenza and pneumonia (J10-J18)	567	1.2	5.1
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	532	1.1	4.8
14	Septicemia (A40-A41)	484	1.0	4.4
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	307	0.6	2.8
...	All other causes (Residual)	7,914	16.1	71.3
All races, male, 50-54 years				
...	All causes	65,193	100.0	666.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	16,519	25.3	168.8
2	Malignant neoplasms (C00-C97)	16,140	24.8	164.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,945	9.1	60.7
4	Chronic liver disease and cirrhosis (K70,K73-K74)	3,062	4.7	31.3
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,443	3.7	25.0
6	Cerebrovascular diseases (I60-I69)	2,101	3.2	21.5
7	Diabetes mellitus (E10-E14)	2,054	3.2	21.0
8	Human immunodeficiency virus (HIV) disease (B20-B24)	1,504	2.3	15.4
9	Chronic lower respiratory diseases (J40-J47)	1,327	2.0	13.6
10	Viral hepatitis (B15-B19)	1,001	1.5	10.2
11	Influenza and pneumonia (J10-J18)	732	1.1	7.5
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	705	1.1	7.2
13	Septicemia (A40-A41)	691	1.1	7.1
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	652	1.0	6.7
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	426	0.7	4.4
...	All other causes (Residual)	9,891	15.2	101.0

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 55-59 years				
...	All causes	77,988	100.0	925.7
1	Malignant neoplasms (C00-C97)	23,993	30.8	284.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	21,012	26.9	249.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,318	5.5	51.3
4	Diabetes mellitus (E10-E14)	3,062	3.9	36.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,905	3.7	34.5
6	Cerebrovascular diseases (I60-I69)	2,535	3.3	30.1
7	Chronic lower respiratory diseases (J40-J47)	2,397	3.1	28.5
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,888	2.4	22.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,046	1.3	12.4
10	Septicemia (A40-A41)	970	1.2	11.5
11	Influenza and pneumonia (J10-J18)	903	1.2	10.7
12	Human immunodeficiency virus (HIV) disease (B20-B24)	883	1.1	10.5
13	Viral hepatitis (B15-B19)	687	0.9	8.2
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	535	0.7	6.4
15	Aortic aneurysm and dissection (I71)	445	0.6	5.3
...	All other causes (Residual)	10,409	13.3	123.5
All races, male, 60-64 years				
...	All causes	87,441	100.0	1,410.0
1	Malignant neoplasms (C00-C97)	30,046	34.4	484.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	23,883	27.3	385.1
3	Chronic lower respiratory diseases (J40-J47)	4,192	4.8	67.6
4	Diabetes mellitus (E10-E14)	3,508	4.0	56.6
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,189	3.6	51.4
6	Cerebrovascular diseases (I60-I69)	3,101	3.5	50.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,189	2.5	35.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,362	1.6	22.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,306	1.5	21.1
10	Septicemia (A40-A41)	1,104	1.3	17.8
11	Influenza and pneumonia (J10-J18)	1,085	1.2	17.5
12	Aortic aneurysm and dissection (I71)	618	0.7	10.0
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	616	0.7	9.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	448	0.5	7.2
15	Human immunodeficiency virus (HIV) disease (B20-B24)	409	0.5	6.6
...	All other causes (Residual)	10,385	11.9	167.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 65-69 years				
...	All causes	98,412	100.0	2,084.2
1	Malignant neoplasms (C00-C97)	35,112	35.7	743.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	25,958	26.4	549.7
3	Chronic lower respiratory diseases (J40-J47)	6,274	6.4	132.9
4	Cerebrovascular diseases (I60-I69)	3,996	4.1	84.6
5	Diabetes mellitus (E10-E14)	3,890	4.0	82.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,579	2.6	54.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,600	1.6	33.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,483	1.5	31.4
9	Influenza and pneumonia (J10-J18)	1,400	1.4	29.6
10	Septicemia (A40-A41)	1,349	1.4	28.6
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	973	1.0	20.6
12	Aortic aneurysm and dissection (I71)	807	0.8	17.1
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	751	0.8	15.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	569	0.6	12.1
15	Parkinson's disease (G20-G21)	454	0.5	9.6
...	All other causes (Residual)	11,217	11.4	237.6
All races, male, 70-74 years				
...	All causes	124,395	100.0	3,267.0
1	Malignant neoplasms (C00-C97)	41,573	33.4	1,091.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	33,092	26.6	869.1
3	Chronic lower respiratory diseases (J40-J47)	9,140	7.3	240.0
4	Cerebrovascular diseases (I60-I69)	5,692	4.6	149.5
5	Diabetes mellitus (E10-E14)	4,615	3.7	121.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,764	2.2	72.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,250	1.8	59.1
8	Influenza and pneumonia (J10-J18)	2,248	1.8	59.0
9	Septicemia (A40-A41)	1,746	1.4	45.9
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,452	1.2	38.1
11	Alzheimer's disease (G30)	1,195	1.0	31.4
12	Parkinson's disease (G20-G21)	1,142	0.9	30.0
13	Aortic aneurysm and dissection (I71)	1,104	0.9	29.0
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	967	0.8	25.4
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	888	0.7	23.3
...	All other causes (Residual)	14,527	11.7	381.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 75-79 years				
...	All causes	159,114	100.0	5,103.4
1	Malignant neoplasms (C00-C97)	45,989	28.9	1,475.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	44,088	27.7	1,414.1
3	Chronic lower respiratory diseases (J40-J47)	12,273	7.7	393.6
4	Cerebrovascular diseases (I60-I69)	8,598	5.4	275.8
5	Diabetes mellitus (E10-E14)	5,425	3.4	174.0
6	Influenza and pneumonia (J10-J18)	3,719	2.3	119.3
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,429	2.2	110.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,203	2.0	102.7
9	Alzheimer's disease (G30)	2,989	1.9	95.9
10	Parkinson's disease (G20-G21)	2,337	1.5	75.0
11	Septicemia (A40-A41)	2,234	1.4	71.7
12	Aortic aneurysm and dissection (I71)	1,365	0.9	43.8
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,307	0.8	41.9
14	Pneumonitis due to solids and liquids (J69)	1,193	0.7	38.3
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,161	0.7	37.2
...	All other causes (Residual)	19,804	12.4	635.2
All races, male, 80-84 years				
...	All causes	176,121	100.0	8,147.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	52,492	29.8	2,428.3
2	Malignant neoplasms (C00-C97)	41,107	23.3	1,901.6
3	Chronic lower respiratory diseases (J40-J47)	12,206	6.9	564.7
4	Cerebrovascular diseases (I60-I69)	11,090	6.3	513.0
5	Influenza and pneumonia (J10-J18)	5,484	3.1	253.7
6	Alzheimer's disease (G30)	5,337	3.0	246.9
7	Diabetes mellitus (E10-E14)	5,276	3.0	244.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,974	2.3	183.8
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,908	2.2	180.8
10	Parkinson's disease (G20-G21)	3,321	1.9	153.6
11	Septicemia (A40-A41)	2,589	1.5	119.8
12	Pneumonitis due to solids and liquids (J69)	1,864	1.1	86.2
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,610	0.9	74.5
14	Aortic aneurysm and dissection (I71)	1,413	0.8	65.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,286	0.7	59.5
...	All other causes (Residual)	23,164	13.2	1,071.6

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 85-89 years				
...	All causes	139,812	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	46,068	32.9	---
2	Malignant neoplasms (C00-C97)	24,960	17.9	---
3	Cerebrovascular diseases (I60-I69)	9,662	6.9	---
4	Chronic lower respiratory diseases (J40-J47)	8,332	6.0	---
5	Alzheimer's disease (G30)	5,778	4.1	---
6	Influenza and pneumonia (J10-J18)	5,419	3.9	---
7	Diabetes mellitus (E10-E14)	3,615	2.6	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,514	2.5	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,311	2.4	---
10	Parkinson's disease (G20-G21)	2,469	1.8	---
11	Septicemia (A40-A41)	2,001	1.4	---
12	Pneumonitis due to solids and liquids (J69)	1,989	1.4	---
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,476	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	968	0.7	---
15	Aortic aneurysm and dissection (I71)	950	0.7	---
...	All other causes (Residual)	19,300	13.8	---
All races, male, 90-94 years				
...	All causes	75,026	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	27,191	36.2	---
2	Malignant neoplasms (C00-C97)	9,940	13.2	---
3	Cerebrovascular diseases (I60-I69)	5,237	7.0	---
4	Influenza and pneumonia (J10-J18)	3,844	5.1	---
5	Chronic lower respiratory diseases (J40-J47)	3,792	5.1	---
6	Alzheimer's disease (G30)	3,431	4.6	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,891	2.5	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,650	2.2	---
9	Diabetes mellitus (E10-E14)	1,547	2.1	---
10	Pneumonitis due to solids and liquids (J69)	1,264	1.7	---
11	Parkinson's disease (G20-G21)	1,004	1.3	---
12	Septicemia (A40-A41)	950	1.3	---
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	893	1.2	---
14	Atherosclerosis (I70)	709	0.9	---
15	Aortic aneurysm and dissection (I71)	422	0.6	---
...	All other causes (Residual)	11,261	15.0	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 95-99 years				
...	All causes	20,972	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,012	38.2	---
2	Malignant neoplasms (C00-C97)	2,108	10.1	---
3	Cerebrovascular diseases (I60-I69)	1,350	6.4	---
4	Influenza and pneumonia (J10-J18)	1,305	6.2	---
5	Alzheimer's disease (G30)	954	4.5	---
6	Chronic lower respiratory diseases (J40-J47)	931	4.4	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	550	2.6	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	497	2.4	---
9	Pneumonitis due to solids and liquids (J69)	365	1.7	---
10	Diabetes mellitus (E10-E14)	334	1.6	---
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	290	1.4	---
12	Septicemia (A40-A41)	257	1.2	---
13	Atherosclerosis (I70)	246	1.2	---
14	Parkinson's disease (G20-G21)	180	0.9	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	87	0.4	---
...	All other causes (Residual)	3,506	16.7	---
All races, male, 100 years and over				
...	All causes	2,986	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,218	40.8	---
2	Influenza and pneumonia (J10-J18)	234	7.8	---
3	Malignant neoplasms (C00-C97)	189	6.3	---
4	Cerebrovascular diseases (I60-I69)	163	5.5	---
5	Alzheimer's disease (G30)	99	3.3	---
6	Chronic lower respiratory diseases (J40-J47)	95	3.2	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	83	2.8	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	63	2.1	---
9	Pneumonitis due to solids and liquids (J69)	59	2.0	---
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	57	1.9	---
11	Atherosclerosis (I70)	51	1.7	---
12	Septicemia (A40-A41)	30	1.0	---
13	Diabetes mellitus (E10-E14)	27	0.9	---
14	Parkinson's disease (G20-G21)	13	0.4	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.4	---
15	Nutritional deficiencies (E40-E64)	11	0.4	---
...	All other causes (Residual)	583	19.5	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, all ages				
...	All causes	1,240,342	100.0	824.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	329,250	26.5	218.9
2	Malignant neoplasms (C00-C97)	268,890	21.7	178.8
3	Cerebrovascular diseases (I60-I69)	86,993	7.0	57.8
4	Chronic lower respiratory diseases (J40-J47)	68,498	5.5	45.5
5	Alzheimer's disease (G30)	51,040	4.1	33.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	41,434	3.3	27.5
7	Diabetes mellitus (E10-E14)	38,581	3.1	25.7
8	Influenza and pneumonia (J10-J18)	34,949	2.8	23.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22,633	1.8	15.0
10	Septicemia (A40-A41)	18,814	1.5	12.5
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	15,444	1.2	10.3
12	Chronic liver disease and cirrhosis (K70,K73-K74)	9,593	0.8	6.4
13	Pneumonitis due to solids and liquids (J69)	8,519	0.7	5.7
14	Parkinson's disease (G20-G21)	8,297	0.7	5.5
15	Atherosclerosis (I70)	7,369	0.6	4.9
...	All other causes (Residual)	230,038	18.5	152.9
All races, female, 1-4 years				
...	All causes	1,991	100.0	25.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	637	32.0	8.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	258	13.0	3.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	161	8.1	2.0
4	Malignant neoplasms (C00-C97)	160	8.0	2.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	70	3.5	0.9
6	Influenza and pneumonia (J10-J18)	54	2.7	0.7
7	Septicemia (A40-A41)	38	1.9	0.5
8	Chronic lower respiratory diseases (J40-J47)	23	1.2	0.3
9	Certain conditions originating in the perinatal period (P00-P96)	22	1.1	0.3
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	21	1.1	0.3
10	Cerebrovascular diseases (I60-I69)	21	1.1	0.3
12	Anemias (D50-D64)	14	0.7	*
13	Meningitis (G00,G03)	11	0.6	*
14	Acute bronchitis and bronchiolitis (J20-J21)	6	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.3	*
...	All other causes (Residual)	489	24.6	6.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 5-9 years				
...	All causes	1,281	100.0	13.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	445	34.7	4.7
2	Malignant neoplasms (C00-C97)	222	17.3	2.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	110	8.6	1.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	56	4.4	0.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	53	4.1	0.6
6	Influenza and pneumonia (J10-J18)	26	2.0	0.3
7	Cerebrovascular diseases (I60-I69)	25	2.0	0.3
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	21	1.6	0.2
8	Chronic lower respiratory diseases (J40-J47)	21	1.6	0.2
10	Septicemia (A40-A41)	12	0.9	*
11	Meningitis (G00,G03)	7	0.5	*
11	Certain conditions originating in the perinatal period (P00-P96)	7	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.5	*
14	Anemias (D50-D64)	5	0.4	*
15	Pneumonitis due to solids and liquids (J69)	4	0.3	*
...	All other causes (Residual)	261	20.4	2.7
All races, female, 10-14 years				
...	All causes	1,468	100.0	14.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	477	32.5	4.7
2	Malignant neoplasms (C00-C97)	221	15.1	2.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	91	6.2	0.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	84	5.7	0.8
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	68	4.6	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	65	4.4	0.6
7	Influenza and pneumonia (J10-J18)	27	1.8	0.3
8	Diabetes mellitus (E10-E14)	21	1.4	0.2
9	Septicemia (A40-A41)	20	1.4	0.2
9	Chronic lower respiratory diseases (J40-J47)	20	1.4	0.2
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	1.3	*
12	Cerebrovascular diseases (I60-I69)	15	1.0	*
13	Certain conditions originating in the perinatal period (P00-P96)	8	0.5	*
14	Anemias (D50-D64)	7	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.5	*
...	All other causes (Residual)	318	21.7	3.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 15-19 years				
...	All causes	3,817	100.0	37.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,971	51.6	19.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	310	8.1	3.0
3	Malignant neoplasms (C00-C97)	289	7.6	2.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	261	6.8	2.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	121	3.2	1.2
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	81	2.1	0.8
7	Cerebrovascular diseases (I60-I69)	42	1.1	0.4
7	Pregnancy, childbirth and the puerperium (O00-099)	42	1.1	0.4
9	Septicemia (A40-A41)	28	0.7	0.3
10	Diabetes mellitus (E10-E14)	27	0.7	0.3
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	22	0.6	0.2
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20	0.5	0.2
13	Anemias (D50-D64)	19	0.5	*
13	Influenza and pneumonia (J10-J18)	19	0.5	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	17	0.4	*
15	Chronic lower respiratory diseases (J40-J47)	17	0.4	*
...	All other causes (Residual)	531	13.9	5.2
All races, female, 20-24 years				
...	All causes	4,908	100.0	48.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,955	39.8	19.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	440	9.0	4.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	404	8.2	4.0
4	Malignant neoplasms (C00-C97)	393	8.0	3.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	228	4.6	2.2
6	Pregnancy, childbirth and the puerperium (O00-099)	141	2.9	1.4
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	97	2.0	1.0
8	Diabetes mellitus (E10-E14)	60	1.2	0.6
8	Cerebrovascular diseases (I60-I69)	60	1.2	0.6
10	Human immunodeficiency virus (HIV) disease (B20-B24)	53	1.1	0.5
11	Septicemia (A40-A41)	38	0.8	0.4
12	Influenza and pneumonia (J10-J18)	36	0.7	0.4
12	Chronic lower respiratory diseases (J40-J47)	36	0.7	0.4
14	Anemias (D50-D64)	27	0.6	0.3
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	26	0.5	0.3
...	All other causes (Residual)	914	18.6	9.0

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 25-29 years				
...	All causes	5,326	100.0	54.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,593	29.9	16.3
2	Malignant neoplasms (C00-C97)	588	11.0	6.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	402	7.5	4.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	393	7.4	4.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	366	6.9	3.7
6	Human immunodeficiency virus (HIV) disease (B20-B24)	163	3.1	1.7
7	Pregnancy, childbirth and the puerperium (O00-099)	155	2.9	1.6
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	100	1.9	1.0
9	Diabetes mellitus (E10-E14)	92	1.7	0.9
10	Cerebrovascular diseases (I60-I69)	83	1.6	0.8
11	Influenza and pneumonia (J10-J18)	68	1.3	0.7
12	Septicemia (A40-A41)	57	1.1	0.6
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	56	1.1	0.6
14	Chronic lower respiratory diseases (J40-J47)	42	0.8	0.4
15	Anemias (D50-D64)	38	0.7	0.4
...	All other causes (Residual)	1,130	21.2	11.5
All races, female, 30-34 years				
...	All causes	7,316	100.0	73.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,739	23.8	17.5
2	Malignant neoplasms (C00-C97)	1,208	16.5	12.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	677	9.3	6.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	522	7.1	5.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	351	4.8	3.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	346	4.7	3.5
7	Cerebrovascular diseases (I60-I69)	162	2.2	1.6
8	Pregnancy, childbirth and the puerperium (O00-099)	157	2.1	1.6
9	Diabetes mellitus (E10-E14)	153	2.1	1.5
10	Septicemia (A40-A41)	102	1.4	1.0
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	92	1.3	0.9
12	Influenza and pneumonia (J10-J18)	90	1.2	0.9
12	Chronic liver disease and cirrhosis (K70,K73-K74)	90	1.2	0.9
14	Chronic lower respiratory diseases (J40-J47)	86	1.2	0.9
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	76	1.0	0.8
...	All other causes (Residual)	1,465	20.0	14.8

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 35-39 years				
...	All causes	11,409	100.0	109.3
1	Malignant neoplasms (C00-C97)	2,536	22.2	24.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,072	18.2	19.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,248	10.9	12.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	634	5.6	6.1
5	Human immunodeficiency virus (HIV) disease (B20-B24)	507	4.4	4.9
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	403	3.5	3.9
7	Cerebrovascular diseases (I60-I69)	338	3.0	3.2
8	Diabetes mellitus (E10-E14)	306	2.7	2.9
9	Chronic liver disease and cirrhosis (K70,K73-K74)	243	2.1	2.3
10	Septicemia (A40-A41)	167	1.5	1.6
11	Chronic lower respiratory diseases (J40-J47)	137	1.2	1.3
12	Pregnancy, childbirth and the puerperium (O00-099)	133	1.2	1.3
13	Influenza and pneumonia (J10-J18)	126	1.1	1.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	119	1.0	1.1
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	104	0.9	1.0
...	All other causes (Residual)	2,336	20.5	22.4
All races, female, 40-44 years				
...	All causes	20,067	100.0	174.7
1	Malignant neoplasms (C00-C97)	5,692	28.4	49.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,821	14.1	24.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,509	12.5	21.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	853	4.3	7.4
5	Cerebrovascular diseases (I60-I69)	774	3.9	6.7
6	Human immunodeficiency virus (HIV) disease (B20-B24)	710	3.5	6.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	659	3.3	5.7
8	Diabetes mellitus (E10-E14)	469	2.3	4.1
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	389	1.9	3.4
10	Chronic lower respiratory diseases (J40-J47)	327	1.6	2.8
11	Influenza and pneumonia (J10-J18)	268	1.3	2.3
12	Septicemia (A40-A41)	259	1.3	2.3
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	200	1.0	1.7
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	141	0.7	1.2
15	Viral hepatitis (B15-B19)	123	0.6	1.1
...	All other causes (Residual)	3,873	19.3	33.7

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 45-49 years				
...	All causes	30,104	100.0	264.6
1	Malignant neoplasms (C00-C97)	10,189	33.8	89.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,328	14.4	38.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,962	9.8	26.0
4	Cerebrovascular diseases (I60-I69)	1,284	4.3	11.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,015	3.4	8.9
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	938	3.1	8.2
7	Diabetes mellitus (E10-E14)	820	2.7	7.2
8	Chronic lower respiratory diseases (J40-J47)	748	2.5	6.6
9	Human immunodeficiency virus (HIV) disease (B20-B24)	633	2.1	5.6
10	Septicemia (A40-A41)	414	1.4	3.6
11	Influenza and pneumonia (J10-J18)	388	1.3	3.4
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	332	1.1	2.9
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	296	1.0	2.6
14	Viral hepatitis (B15-B19)	253	0.8	2.2
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	197	0.7	1.7
...	All other causes (Residual)	5,307	17.6	46.6
All races, female, 50-54 years				
...	All causes	38,954	100.0	381.6
1	Malignant neoplasms (C00-C97)	14,813	38.0	145.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,289	16.1	61.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,375	6.1	23.3
4	Cerebrovascular diseases (I60-I69)	1,651	4.2	16.2
5	Diabetes mellitus (E10-E14)	1,404	3.6	13.8
6	Chronic lower respiratory diseases (J40-J47)	1,195	3.1	11.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,079	2.8	10.6
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	784	2.0	7.7
9	Septicemia (A40-A41)	622	1.6	6.1
10	Influenza and pneumonia (J10-J18)	496	1.3	4.9
11	Human immunodeficiency virus (HIV) disease (B20-B24)	466	1.2	4.6
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	459	1.2	4.5
13	Viral hepatitis (B15-B19)	364	0.9	3.6
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	236	0.6	2.3
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	185	0.5	1.8
...	All other causes (Residual)	6,536	16.8	64.0

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 55-59 years				
...	All causes	49,490	100.0	554.3
1	Malignant neoplasms (C00-C97)	20,365	41.1	228.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,669	17.5	97.1
3	Chronic lower respiratory diseases (J40-J47)	2,275	4.6	25.5
4	Diabetes mellitus (E10-E14)	2,088	4.2	23.4
5	Cerebrovascular diseases (I60-I69)	1,945	3.9	21.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,850	3.7	20.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,090	2.2	12.2
8	Septicemia (A40-A41)	852	1.7	9.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	789	1.6	8.8
10	Influenza and pneumonia (J10-J18)	619	1.3	6.9
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	589	1.2	6.6
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	338	0.7	3.8
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	273	0.6	3.1
14	Viral hepatitis (B15-B19)	241	0.5	2.7
15	Human immunodeficiency virus (HIV) disease (B20-B24)	211	0.4	2.4
...	All other causes (Residual)	7,296	14.7	81.7
All races, female, 60-64 years				
...	All causes	60,382	100.0	887.9
1	Malignant neoplasms (C00-C97)	24,836	41.1	365.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,644	19.3	171.2
3	Chronic lower respiratory diseases (J40-J47)	3,883	6.4	57.1
4	Diabetes mellitus (E10-E14)	2,643	4.4	38.9
5	Cerebrovascular diseases (I60-I69)	2,447	4.1	36.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,496	2.5	22.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,000	1.7	14.7
8	Septicemia (A40-A41)	986	1.6	14.5
9	Chronic liver disease and cirrhosis (K70,K73-K74)	942	1.6	13.9
10	Influenza and pneumonia (J10-J18)	815	1.3	12.0
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	465	0.8	6.8
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	371	0.6	5.5
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	291	0.5	4.3
14	Aortic aneurysm and dissection (I71)	260	0.4	3.8
15	Alzheimer's disease (G30)	253	0.4	3.7
...	All other causes (Residual)	8,050	13.3	118.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 65-69 years				
...	All causes	73,824	100.0	1,364.7
1	Malignant neoplasms (C00-C97)	27,924	37.8	516.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,126	20.5	279.6
3	Chronic lower respiratory diseases (J40-J47)	5,742	7.8	106.1
4	Cerebrovascular diseases (I60-I69)	3,477	4.7	64.3
5	Diabetes mellitus (E10-E14)	3,370	4.6	62.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,481	2.0	27.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,475	2.0	27.3
8	Septicemia (A40-A41)	1,253	1.7	23.2
9	Influenza and pneumonia (J10-J18)	1,086	1.5	20.1
10	Chronic liver disease and cirrhosis (K70,K73-K74)	969	1.3	17.9
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	662	0.9	12.2
12	Alzheimer's disease (G30)	530	0.7	9.8
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	397	0.5	7.3
14	Aortic aneurysm and dissection (I71)	378	0.5	7.0
15	Pneumonitis due to solids and liquids (J69)	248	0.3	4.6
...	All other causes (Residual)	9,706	13.1	179.4
All races, female, 70-74 years				
...	All causes	101,724	100.0	2,164.0
1	Malignant neoplasms (C00-C97)	33,837	33.3	719.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	22,553	22.2	479.8
3	Chronic lower respiratory diseases (J40-J47)	8,754	8.6	186.2
4	Cerebrovascular diseases (I60-I69)	5,674	5.6	120.7
5	Diabetes mellitus (E10-E14)	4,308	4.2	91.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,112	2.1	44.9
7	Influenza and pneumonia (J10-J18)	1,889	1.9	40.2
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,808	1.8	38.5
9	Septicemia (A40-A41)	1,725	1.7	36.7
10	Alzheimer's disease (G30)	1,635	1.6	34.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1,045	1.0	22.2
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,005	1.0	21.4
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	611	0.6	13.0
14	Parkinson's disease (G20-G21)	587	0.6	12.5
15	Aortic aneurysm and dissection (I71)	581	0.6	12.4
...	All other causes (Residual)	13,600	13.4	289.3

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 75-79 years				
...	All causes	148,774	100.0	3,464.7
1	Malignant neoplasms (C00-C97)	39,872	26.8	928.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	36,755	24.7	856.0
3	Chronic lower respiratory diseases (J40-J47)	12,228	8.2	284.8
4	Cerebrovascular diseases (I60-I69)	10,202	6.9	237.6
5	Diabetes mellitus (E10-E14)	5,793	3.9	134.9
6	Alzheimer's disease (G30)	4,607	3.1	107.3
7	Influenza and pneumonia (J10-J18)	3,404	2.3	79.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,145	2.1	73.2
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,697	1.8	62.8
10	Septicemia (A40-A41)	2,481	1.7	57.8
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,693	1.1	39.4
12	Parkinson's disease (G20-G21)	1,387	0.9	32.3
13	Chronic liver disease and cirrhosis (K70,K73-K74)	1,047	0.7	24.4
14	Aortic aneurysm and dissection (I71)	975	0.7	22.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	935	0.6	21.8
...	All other causes (Residual)	21,553	14.5	501.9
All races, female, 80-84 years				
...	All causes	202,656	100.0	5,822.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	57,358	28.3	1,647.8
2	Malignant neoplasms (C00-C97)	39,453	19.5	1,133.4
3	Cerebrovascular diseases (I60-I69)	16,969	8.4	487.5
4	Chronic lower respiratory diseases (J40-J47)	13,626	6.7	391.5
5	Alzheimer's disease (G30)	10,206	5.0	293.2
6	Diabetes mellitus (E10-E14)	6,642	3.3	190.8
7	Influenza and pneumonia (J10-J18)	5,956	2.9	171.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,081	2.0	117.2
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,820	1.9	109.7
10	Septicemia (A40-A41)	3,322	1.6	95.4
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,646	1.3	76.0
12	Parkinson's disease (G20-G21)	2,249	1.1	64.6
13	Pneumonitis due to solids and liquids (J69)	1,585	0.8	45.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,240	0.6	35.6
15	Atherosclerosis (I70)	1,224	0.6	35.2
...	All other causes (Residual)	32,279	15.9	927.3

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 85-89 years				
...	All causes	211,586	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	67,772	32.0	---
2	Malignant neoplasms (C00-C97)	27,911	13.2	---
3	Cerebrovascular diseases (I60-I69)	19,247	9.1	---
4	Alzheimer's disease (G30)	14,092	6.7	---
5	Chronic lower respiratory diseases (J40-J47)	11,006	5.2	---
6	Influenza and pneumonia (J10-J18)	7,690	3.6	---
7	Diabetes mellitus (E10-E14)	5,698	2.7	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,148	2.0	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,992	1.9	---
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	3,125	1.5	---
11	Septicemia (A40-A41)	3,124	1.5	---
12	Parkinson's disease (G20-G21)	2,155	1.0	---
13	Pneumonitis due to solids and liquids (J69)	1,998	0.9	---
14	Atherosclerosis (I70)	1,764	0.8	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,263	0.6	---
...	All other causes (Residual)	36,601	17.3	---
All races, female, 90-94 years				
...	All causes	164,680	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	59,131	35.9	---
2	Cerebrovascular diseases (I60-I69)	15,209	9.2	---
3	Malignant neoplasms (C00-C97)	13,819	8.4	---
4	Alzheimer's disease (G30)	12,627	7.7	---
5	Influenza and pneumonia (J10-J18)	7,135	4.3	---
6	Chronic lower respiratory diseases (J40-J47)	6,017	3.7	---
7	Diabetes mellitus (E10-E14)	3,363	2.0	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,141	1.9	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,046	1.8	---
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,998	1.8	---
11	Septicemia (A40-A41)	2,188	1.3	---
12	Atherosclerosis (I70)	1,776	1.1	---
13	Pneumonitis due to solids and liquids (J69)	1,729	1.0	---
14	Parkinson's disease (G20-G21)	1,159	0.7	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	842	0.5	---
...	All other causes (Residual)	30,500	18.5	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 95-99 years				
...	All causes	71,789	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	27,445	38.2	---
2	Cerebrovascular diseases (I60-I69)	6,056	8.4	---
3	Alzheimer's disease (G30)	5,816	8.1	---
4	Malignant neoplasms (C00-C97)	3,960	5.5	---
5	Influenza and pneumonia (J10-J18)	3,632	5.1	---
6	Chronic lower respiratory diseases (J40-J47)	1,992	2.8	---
7	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,521	2.1	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,281	1.8	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,219	1.7	---
10	Diabetes mellitus (E10-E14)	1,121	1.6	---
11	Atherosclerosis (I70)	992	1.4	---
12	Septicemia (A40-A41)	841	1.2	---
13	Pneumonitis due to solids and liquids (J69)	788	1.1	---
14	Parkinson's disease (G20-G21)	306	0.4	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	276	0.4	---
...	All other causes (Residual)	14,543	20.3	---
All races, female, 100 years and over				
...	All causes	16,318	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,667	40.9	---
2	Cerebrovascular diseases (I60-I69)	1,259	7.7	---
3	Alzheimer's disease (G30)	1,109	6.8	---
4	Influenza and pneumonia (J10-J18)	1,008	6.2	---
5	Malignant neoplasms (C00-C97)	568	3.5	---
6	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	343	2.1	---
7	Chronic lower respiratory diseases (J40-J47)	308	1.9	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	308	1.9	---
9	Atherosclerosis (I70)	289	1.8	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	242	1.5	---
11	Diabetes mellitus (E10-E14)	198	1.2	---
12	Pneumonitis due to solids and liquids (J69)	160	1.0	---
13	Septicemia (A40-A41)	153	0.9	---
14	Anemias (D50-D64)	76	0.5	---
15	Nutritional deficiencies (E40-E64)	60	0.4	---
...	All other causes (Residual)	3,570	21.9	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, all ages				
...	All causes	2,098,097	100.0	873.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	564,796	26.9	235.2
2	Malignant neoplasms (C00-C97)	482,132	23.0	200.8
3	Cerebrovascular diseases (I60-I69)	121,868	5.8	50.7
4	Chronic lower respiratory diseases (J40-J47)	120,884	5.8	50.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	100,406	4.8	41.8
6	Alzheimer's disease (G30)	66,191	3.2	27.6
7	Diabetes mellitus (E10-E14)	59,755	2.8	24.9
8	Influenza and pneumonia (J10-J18)	55,540	2.6	23.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	34,806	1.7	14.5
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	29,527	1.4	12.3
11	Septicemia (A40-A41)	27,194	1.3	11.3
12	Chronic liver disease and cirrhosis (K70,K73-K74)	24,003	1.1	10.0
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	19,254	0.9	8.0
14	Parkinson's disease (G20-G21)	18,496	0.9	7.7
15	Pneumonitis due to solids and liquids (J69)	15,338	0.7	6.4
...	All other causes (Residual)	357,907	17.1	149.0
White, both sexes, 1-4 years				
...	All causes	3,408	100.0	27.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,262	37.0	10.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	385	11.3	3.1
3	Malignant neoplasms (C00-C97)	293	8.6	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	217	6.4	1.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	93	2.7	0.7
6	Influenza and pneumonia (J10-J18)	76	2.2	0.6
7	Septicemia (A40-A41)	59	1.7	0.5
8	Cerebrovascular diseases (I60-I69)	45	1.3	0.4
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	42	1.2	0.3
10	Certain conditions originating in the perinatal period (P00-P96)	35	1.0	0.3
11	Chronic lower respiratory diseases (J40-J47)	24	0.7	0.2
12	Meningitis (G00,G03)	20	0.6	0.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	13	0.4	*
15	Acute bronchitis and bronchiolitis (J20-J21)	12	0.4	*
...	All other causes (Residual)	819	24.0	6.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 5-9 years				
...	All causes	2,018	100.0	13.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	775	38.4	5.1
2	Malignant neoplasms (C00-C97)	380	18.8	2.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	140	6.9	0.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	66	3.3	0.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	63	3.1	0.4
6	Influenza and pneumonia (J10-J18)	37	1.8	0.2
7	Cerebrovascular diseases (I60-I69)	35	1.7	0.2
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	30	1.5	0.2
9	Chronic lower respiratory diseases (J40-J47)	25	1.2	0.2
10	Septicemia (A40-A41)	24	1.2	0.2
11	Certain conditions originating in the perinatal period (P00-P96)	12	0.6	*
12	Meningitis (G00,G03)	10	0.5	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	9	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.3	*
15	Pneumonitis due to solids and liquids (J69)	5	0.2	*
15	Diseases of appendix (K35-K38)	5	0.2	*
...	All other causes (Residual)	395	19.6	2.6
White, both sexes, 10-14 years				
...	All causes	2,692	100.0	16.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,010	37.5	6.3
2	Malignant neoplasms (C00-C97)	407	15.1	2.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	208	7.7	1.3
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	145	5.4	0.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	102	3.8	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	96	3.6	0.6
7	Influenza and pneumonia (J10-J18)	40	1.5	0.2
8	Septicemia (A40-A41)	33	1.2	0.2
9	Cerebrovascular diseases (I60-I69)	29	1.1	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	27	1.0	0.2
11	Diabetes mellitus (E10-E14)	21	0.8	0.1
12	Chronic lower respiratory diseases (J40-J47)	20	0.7	0.1
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	0.4	*
14	Certain conditions originating in the perinatal period (P00-P96)	10	0.4	*
15	Anemias (D50-D64)	8	0.3	*
15	Meningitis (G00,G03)	8	0.3	*
...	All other causes (Residual)	517	19.2	3.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 15-19 years				
...	All causes	10,268	100.0	62.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,562	54.2	33.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,369	13.3	8.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	866	8.4	5.3
4	Malignant neoplasms (C00-C97)	590	5.7	3.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	262	2.6	1.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	188	1.8	1.1
7	Cerebrovascular diseases (I60-I69)	51	0.5	0.3
7	Influenza and pneumonia (J10-J18)	51	0.5	0.3
9	Diabetes mellitus (E10-E14)	41	0.4	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	38	0.4	0.2
11	Septicemia (A40-A41)	35	0.3	0.2
12	Chronic lower respiratory diseases (J40-J47)	34	0.3	0.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21	0.2	0.1
13	Pregnancy, childbirth and the puerperium (O00-099)	21	0.2	0.1
15	Pneumonitis due to solids and liquids (J69)	16	0.2	*
...	All other causes (Residual)	1,123	10.9	6.8
White, both sexes, 20-24 years				
...	All causes	15,142	100.0	91.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,763	51.3	46.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,171	14.3	13.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,306	8.6	7.9
4	Malignant neoplasms (C00-C97)	775	5.1	4.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	474	3.1	2.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	198	1.3	1.2
7	Pregnancy, childbirth and the puerperium (O00-099)	89	0.6	0.5
8	Cerebrovascular diseases (I60-I69)	85	0.6	0.5
9	Diabetes mellitus (E10-E14)	83	0.5	0.5
9	Influenza and pneumonia (J10-J18)	83	0.5	0.5
11	Septicemia (A40-A41)	60	0.4	0.4
12	Legal intervention (Y35,Y89.0)	48	0.3	0.3
13	Human immunodeficiency virus (HIV) disease (B20-B24)	44	0.3	0.3
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	39	0.3	0.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	38	0.3	0.2
...	All other causes (Residual)	1,886	12.5	11.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 25-29 years				
...	All causes	14,079	100.0	89.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,066	43.1	38.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,976	14.0	12.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,133	8.0	7.2
4	Malignant neoplasms (C00-C97)	991	7.0	6.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	759	5.4	4.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	180	1.3	1.1
7	Human immunodeficiency virus (HIV) disease (B20-B24)	127	0.9	0.8
8	Diabetes mellitus (E10-E14)	123	0.9	0.8
9	Cerebrovascular diseases (I60-I69)	118	0.8	0.7
10	Influenza and pneumonia (J10-J18)	111	0.8	0.7
11	Septicemia (A40-A41)	80	0.6	0.5
12	Pregnancy, childbirth and the puerperium (O00-099)	78	0.6	0.5
13	Chronic lower respiratory diseases (J40-J47)	64	0.5	0.4
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	60	0.4	0.4
15	Chronic liver disease and cirrhosis (K70,K73-K74)	59	0.4	0.4
...	All other causes (Residual)	2,154	15.3	13.7
White, both sexes, 30-34 years				
...	All causes	16,024	100.0	101.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,562	34.7	35.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,253	14.1	14.3
3	Malignant neoplasms (C00-C97)	1,703	10.6	10.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,325	8.3	8.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	889	5.5	5.6
6	Human immunodeficiency virus (HIV) disease (B20-B24)	337	2.1	2.1
7	Diabetes mellitus (E10-E14)	249	1.6	1.6
8	Cerebrovascular diseases (I60-I69)	238	1.5	1.5
9	Chronic liver disease and cirrhosis (K70,K73-K74)	172	1.1	1.1
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	159	1.0	1.0
11	Influenza and pneumonia (J10-J18)	151	0.9	1.0
12	Septicemia (A40-A41)	111	0.7	0.7
13	Pregnancy, childbirth and the puerperium (O00-099)	101	0.6	0.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	92	0.6	0.6
15	Chronic lower respiratory diseases (J40-J47)	88	0.5	0.6
...	All other causes (Residual)	2,594	16.2	16.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 35-39 years				
...	All causes	23,177	100.0	138.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,164	26.6	36.8
2	Malignant neoplasms (C00-C97)	3,376	14.6	20.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,793	12.1	16.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,586	11.2	15.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	794	3.4	4.7
6	Human immunodeficiency virus (HIV) disease (B20-B24)	710	3.1	4.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	609	2.6	3.6
8	Diabetes mellitus (E10-E14)	510	2.2	3.0
9	Cerebrovascular diseases (I60-I69)	461	2.0	2.7
10	Influenza and pneumonia (J10-J18)	231	1.0	1.4
11	Septicemia (A40-A41)	199	0.9	1.2
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	192	0.8	1.1
13	Chronic lower respiratory diseases (J40-J47)	181	0.8	1.1
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	147	0.6	0.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	108	0.5	0.6
...	All other causes (Residual)	4,116	17.8	24.6
White, both sexes, 40-44 years				
...	All causes	40,407	100.0	217.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,008	19.8	43.2
2	Malignant neoplasms (C00-C97)	7,976	19.7	43.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,308	15.6	34.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,331	8.2	18.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,675	4.1	9.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	1,248	3.1	6.7
7	Cerebrovascular diseases (I60-I69)	989	2.4	5.3
8	Diabetes mellitus (E10-E14)	918	2.3	4.9
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	821	2.0	4.4
10	Chronic lower respiratory diseases (J40-J47)	466	1.2	2.5
11	Influenza and pneumonia (J10-J18)	452	1.1	2.4
12	Septicemia (A40-A41)	353	0.9	1.9
13	Viral hepatitis (B15-B19)	320	0.8	1.7
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	254	0.6	1.4
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	235	0.6	1.3
...	All other causes (Residual)	7,053	17.5	38.0

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 45-49 years				
...	All causes	60,286	100.0	326.6
1	Malignant neoplasms (C00-C97)	15,135	25.1	82.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,324	18.8	61.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,372	13.9	45.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,497	5.8	18.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,850	4.7	15.4
6	Cerebrovascular diseases (I60-I69)	1,674	2.8	9.1
7	Diabetes mellitus (E10-E14)	1,588	2.6	8.6
8	Chronic lower respiratory diseases (J40-J47)	1,142	1.9	6.2
9	Human immunodeficiency virus (HIV) disease (B20-B24)	1,097	1.8	5.9
10	Viral hepatitis (B15-B19)	788	1.3	4.3
11	Influenza and pneumonia (J10-J18)	709	1.2	3.8
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	700	1.2	3.8
13	Septicemia (A40-A41)	602	1.0	3.3
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	500	0.8	2.7
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	281	0.5	1.5
...	All other causes (Residual)	10,027	16.6	54.3
White, both sexes, 50-54 years				
...	All causes	79,938	100.0	480.7
1	Malignant neoplasms (C00-C97)	24,582	30.8	147.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	17,050	21.3	102.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,763	8.5	40.7
4	Chronic liver disease and cirrhosis (K70,K73-K74)	3,492	4.4	21.0
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,008	3.8	18.1
6	Diabetes mellitus (E10-E14)	2,491	3.1	15.0
7	Cerebrovascular diseases (I60-I69)	2,409	3.0	14.5
8	Chronic lower respiratory diseases (J40-J47)	2,066	2.6	12.4
9	Viral hepatitis (B15-B19)	1,107	1.4	6.7
10	Influenza and pneumonia (J10-J18)	914	1.1	5.5
11	Septicemia (A40-A41)	897	1.1	5.4
12	Human immunodeficiency virus (HIV) disease (B20-B24)	850	1.1	5.1
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	721	0.9	4.3
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	482	0.6	2.9
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	347	0.4	2.1
...	All other causes (Residual)	12,759	16.0	76.7

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 55-59 years				
...	All causes	101,212	100.0	689.7
1	Malignant neoplasms (C00-C97)	36,277	35.8	247.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	22,984	22.7	156.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,097	5.0	34.7
4	Chronic lower respiratory diseases (J40-J47)	4,060	4.0	27.7
5	Diabetes mellitus (E10-E14)	3,798	3.8	25.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	3,446	3.4	23.5
7	Cerebrovascular diseases (I60-I69)	3,101	3.1	21.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,338	2.3	15.9
9	Septicemia (A40-A41)	1,314	1.3	9.0
10	Influenza and pneumonia (J10-J18)	1,175	1.2	8.0
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,127	1.1	7.7
12	Viral hepatitis (B15-B19)	725	0.7	4.9
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	518	0.5	3.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	490	0.5	3.3
15	Human immunodeficiency virus (HIV) disease (B20-B24)	472	0.5	3.2
...	All other causes (Residual)	14,290	14.1	97.4
White, both sexes, 60-64 years				
...	All causes	121,006	100.0	1,085.7
1	Malignant neoplasms (C00-C97)	46,162	38.1	414.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	28,423	23.5	255.0
3	Chronic lower respiratory diseases (J40-J47)	7,312	6.0	65.6
4	Diabetes mellitus (E10-E14)	4,660	3.9	41.8
5	Cerebrovascular diseases (I60-I69)	4,039	3.3	36.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,935	3.3	35.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,692	2.2	24.2
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,641	1.4	14.7
9	Septicemia (A40-A41)	1,563	1.3	14.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,536	1.3	13.8
11	Influenza and pneumonia (J10-J18)	1,525	1.3	13.7
12	Aortic aneurysm and dissection (I71)	742	0.6	6.7
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	693	0.6	6.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	619	0.5	5.6
15	Alzheimer's disease (G30)	413	0.3	3.7
...	All other causes (Residual)	15,051	12.4	135.0

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 65-69 years				
...	All causes	143,202	100.0	1,649.5
1	Malignant neoplasms (C00-C97)	53,698	37.5	618.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	33,445	23.4	385.2
3	Chronic lower respiratory diseases (J40-J47)	10,955	7.7	126.2
4	Cerebrovascular diseases (I60-I69)	5,695	4.0	65.6
5	Diabetes mellitus (E10-E14)	5,495	3.8	63.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,457	2.4	39.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,302	1.6	26.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,072	1.4	23.9
9	Influenza and pneumonia (J10-J18)	2,023	1.4	23.3
10	Septicemia (A40-A41)	1,985	1.4	22.9
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,128	0.8	13.0
12	Aortic aneurysm and dissection (I71)	1,019	0.7	11.7
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	920	0.6	10.6
14	Alzheimer's disease (G30)	869	0.6	10.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	842	0.6	9.7
...	All other causes (Residual)	17,297	12.1	199.2
White, both sexes, 70-74 years				
...	All causes	193,160	100.0	2,618.1
1	Malignant neoplasms (C00-C97)	65,582	34.0	888.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	46,828	24.2	634.7
3	Chronic lower respiratory diseases (J40-J47)	16,562	8.6	224.5
4	Cerebrovascular diseases (I60-I69)	9,134	4.7	123.8
5	Diabetes mellitus (E10-E14)	6,981	3.6	94.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,967	2.1	53.8
7	Influenza and pneumonia (J10-J18)	3,467	1.8	47.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,271	1.7	44.3
9	Septicemia (A40-A41)	2,766	1.4	37.5
10	Alzheimer's disease (G30)	2,514	1.3	34.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	2,279	1.2	30.9
12	Parkinson's disease (G20-G21)	1,606	0.8	21.8
13	Aortic aneurysm and dissection (I71)	1,520	0.8	20.6
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,302	0.7	17.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,225	0.6	16.6
...	All other causes (Residual)	24,156	12.5	327.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 75-79 years				
...	All causes	270,366	100.0	4,128.9
1	Malignant neoplasms (C00-C97)	76,278	28.2	1,164.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	70,272	26.0	1,073.2
3	Chronic lower respiratory diseases (J40-J47)	22,879	8.5	349.4
4	Cerebrovascular diseases (I60-I69)	15,990	5.9	244.2
5	Diabetes mellitus (E10-E14)	9,164	3.4	139.9
6	Alzheimer's disease (G30)	6,895	2.6	105.3
7	Influenza and pneumonia (J10-J18)	6,229	2.3	95.1
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,444	2.0	83.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,152	1.9	78.7
10	Septicemia (A40-A41)	3,856	1.4	58.9
11	Parkinson's disease (G20-G21)	3,517	1.3	53.7
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,291	0.8	35.0
13	Aortic aneurysm and dissection (I71)	2,126	0.8	32.5
14	Chronic liver disease and cirrhosis (K70,K73-K74)	2,018	0.7	30.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,878	0.7	28.7
...	All other causes (Residual)	36,377	13.5	555.5
White, both sexes, 80-84 years				
...	All causes	340,454	100.0	6,731.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	98,439	28.9	1,946.5
2	Malignant neoplasms (C00-C97)	72,454	21.3	1,432.7
3	Cerebrovascular diseases (I60-I69)	24,842	7.3	491.2
4	Chronic lower respiratory diseases (J40-J47)	24,353	7.2	481.5
5	Alzheimer's disease (G30)	14,377	4.2	284.3
6	Influenza and pneumonia (J10-J18)	10,318	3.0	204.0
7	Diabetes mellitus (E10-E14)	10,048	3.0	198.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,080	2.1	140.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,843	2.0	135.3
10	Parkinson's disease (G20-G21)	5,280	1.6	104.4
11	Septicemia (A40-A41)	4,973	1.5	98.3
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	3,514	1.0	69.5
13	Pneumonitis due to solids and liquids (J69)	3,103	0.9	61.4
14	Aortic aneurysm and dissection (I71)	2,406	0.7	47.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,342	0.7	46.3
...	All other causes (Residual)	50,082	14.7	990.3

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 85-89 years				
...	All causes	321,335	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	104,488	32.5	---
2	Malignant neoplasms (C00-C97)	47,874	14.9	---
3	Cerebrovascular diseases (I60-I69)	26,229	8.2	---
4	Alzheimer's disease (G30)	18,555	5.8	---
5	Chronic lower respiratory diseases (J40-J47)	18,278	5.7	---
6	Influenza and pneumonia (J10-J18)	12,008	3.7	---
7	Diabetes mellitus (E10-E14)	8,001	2.5	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,842	2.1	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,703	2.1	---
10	Parkinson's disease (G20-G21)	4,433	1.4	---
11	Septicemia (A40-A41)	4,402	1.4	---
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	3,928	1.2	---
13	Pneumonitis due to solids and liquids (J69)	3,650	1.1	---
14	Atherosclerosis (I70)	2,544	0.8	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,085	0.6	---
...	All other causes (Residual)	51,315	16.0	---
White, both sexes, 90-94 years				
...	All causes	219,853	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	79,681	36.2	---
2	Malignant neoplasms (C00-C97)	21,491	9.8	---
3	Cerebrovascular diseases (I60-I69)	18,645	8.5	---
4	Alzheimer's disease (G30)	14,977	6.8	---
5	Influenza and pneumonia (J10-J18)	10,127	4.6	---
6	Chronic lower respiratory diseases (J40-J47)	9,241	4.2	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,494	2.0	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,337	2.0	---
9	Diabetes mellitus (E10-E14)	4,160	1.9	---
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	3,359	1.5	---
11	Pneumonitis due to solids and liquids (J69)	2,750	1.3	---
12	Septicemia (A40-A41)	2,682	1.2	---
13	Atherosclerosis (I70)	2,340	1.1	---
14	Parkinson's disease (G20-G21)	2,074	0.9	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,185	0.5	---
...	All other causes (Residual)	38,310	17.4	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 95-99 years				
...	All causes	84,513	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	32,484	38.4	---
2	Cerebrovascular diseases (I60-I69)	6,750	8.0	---
3	Alzheimer's disease (G30)	6,264	7.4	---
4	Malignant neoplasms (C00-C97)	5,395	6.4	---
5	Influenza and pneumonia (J10-J18)	4,546	5.4	---
6	Chronic lower respiratory diseases (J40-J47)	2,714	3.2	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,655	2.0	---
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,573	1.9	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,534	1.8	---
10	Diabetes mellitus (E10-E14)	1,246	1.5	---
11	Atherosclerosis (I70)	1,171	1.4	---
12	Pneumonitis due to solids and liquids (J69)	1,040	1.2	---
13	Septicemia (A40-A41)	907	1.1	---
14	Parkinson's disease (G20-G21)	459	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	330	0.4	---
...	All other causes (Residual)	16,445	19.5	---
White, both sexes, 100 years and over				
...	All causes	16,838	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,950	41.3	---
2	Cerebrovascular diseases (I60-I69)	1,230	7.3	---
3	Influenza and pneumonia (J10-J18)	1,107	6.6	---
4	Alzheimer's disease (G30)	1,064	6.3	---
5	Malignant neoplasms (C00-C97)	644	3.8	---
6	Chronic lower respiratory diseases (J40-J47)	360	2.1	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	342	2.0	---
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	318	1.9	---
9	Atherosclerosis (I70)	301	1.8	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	270	1.6	---
11	Pneumonitis due to solids and liquids (J69)	190	1.1	---
12	Diabetes mellitus (E10-E14)	172	1.0	---
13	Septicemia (A40-A41)	132	0.8	---
14	Anemias (D50-D64)	78	0.5	---
15	Nutritional deficiencies (E40-E64)	58	0.3	---
...	All other causes (Residual)	3,622	21.5	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, male, all ages				
...	All causes	1,028,152	100.0	864.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	279,324	27.2	234.9
2	Malignant neoplasms (C00-C97)	250,478	24.4	210.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	64,600	6.3	54.3
4	Chronic lower respiratory diseases (J40-J47)	56,911	5.5	47.9
5	Cerebrovascular diseases (I60-I69)	47,194	4.6	39.7
6	Diabetes mellitus (E10-E14)	29,628	2.9	24.9
7	Influenza and pneumonia (J10-J18)	24,425	2.4	20.5
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	23,478	2.3	19.7
9	Alzheimer's disease (G30)	18,990	1.8	16.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17,137	1.7	14.4
11	Chronic liver disease and cirrhosis (K70,K73-K74)	15,679	1.5	13.2
12	Septicemia (A40-A41)	12,172	1.2	10.2
13	Parkinson's disease (G20-G21)	10,649	1.0	9.0
14	Pneumonitis due to solids and liquids (J69)	7,759	0.8	6.5
15	Aortic aneurysm and dissection (I71)	7,203	0.7	6.1
...	All other causes (Residual)	162,525	15.8	136.7
White, male, 1-4 years				
...	All causes	1,996	100.0	30.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	793	39.7	12.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	201	10.1	3.1
3	Malignant neoplasms (C00-C97)	168	8.4	2.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	119	6.0	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	55	2.8	0.9
6	Influenza and pneumonia (J10-J18)	36	1.8	0.6
7	Septicemia (A40-A41)	33	1.7	0.5
8	Cerebrovascular diseases (I60-I69)	28	1.4	0.4
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	26	1.3	0.4
10	Certain conditions originating in the perinatal period (P00-P96)	20	1.0	0.3
11	Meningitis (G00,G03)	13	0.7	*
12	Chronic lower respiratory diseases (J40-J47)	11	0.6	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.5	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	9	0.5	*
15	Acute bronchitis and bronchiolitis (J20-J21)	8	0.4	*
...	All other causes (Residual)	467	23.4	7.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 5-9 years				
...	All causes	1,109	100.0	14.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	457	41.2	5.9
2	Malignant neoplasms (C00-C97)	201	18.1	2.6
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	64	5.8	0.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	40	3.6	0.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	33	3.0	0.4
6	Cerebrovascular diseases (I60-I69)	17	1.5	*
7	Influenza and pneumonia (J10-J18)	16	1.4	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	15	1.4	*
8	Chronic lower respiratory diseases (J40-J47)	15	1.4	*
10	Septicemia (A40-A41)	14	1.3	*
11	Certain conditions originating in the perinatal period (P00-P96)	7	0.6	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.5	*
13	Diseases of appendix (K35-K38)	5	0.5	*
14	Pneumonitis due to solids and liquids (J69)	4	0.4	*
15	Meningitis (G00,G03)	3	0.3	*
15	Acute bronchitis and bronchiolitis (J20-J21)	3	0.3	*
...	All other causes (Residual)	209	18.8	2.7
White, male, 10-14 years				
...	All causes	1,640	100.0	19.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	644	39.3	7.8
2	Malignant neoplasms (C00-C97)	228	13.9	2.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	159	9.7	1.9
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	81	4.9	1.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	57	3.5	0.7
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	56	3.4	0.7
7	Influenza and pneumonia (J10-J18)	20	1.2	0.2
8	Septicemia (A40-A41)	19	1.2	*
9	Cerebrovascular diseases (I60-I69)	18	1.1	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	0.9	*
11	Chronic lower respiratory diseases (J40-J47)	11	0.7	*
12	Diabetes mellitus (E10-E14)	9	0.5	*
13	Meningitis (G00,G03)	6	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.3	*
15	Meningococcal infection (A39)	4	0.2	*
15	Anemias (D50-D64)	4	0.2	*
15	Certain conditions originating in the perinatal period (P00-P96)	4	0.2	*
...	All other causes (Residual)	301	18.4	3.6

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 15-19 years				
...	All causes	7,299	100.0	86.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,889	53.3	46.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,119	15.3	13.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	726	9.9	8.6
4	Malignant neoplasms (C00-C97)	358	4.9	4.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	176	2.4	2.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	129	1.8	1.5
7	Influenza and pneumonia (J10-J18)	37	0.5	0.4
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	24	0.3	0.3
9	Diabetes mellitus (E10-E14)	22	0.3	0.3
9	Chronic lower respiratory diseases (J40-J47)	22	0.3	0.3
11	Cerebrovascular diseases (I60-I69)	21	0.3	0.2
12	Septicemia (A40-A41)	19	0.3	*
13	Legal intervention (Y35,Y89.0)	11	0.2	*
14	Aortic aneurysm and dissection (I71)	9	0.1	*
14	Pneumonitis due to solids and liquids (J69)	9	0.1	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.1	*
...	All other causes (Residual)	719	9.9	8.5
White, male, 20-24 years				
...	All causes	11,493	100.0	133.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,146	53.5	71.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,826	15.9	21.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,074	9.3	12.5
4	Malignant neoplasms (C00-C97)	468	4.1	5.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	333	2.9	3.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	123	1.1	1.4
7	Influenza and pneumonia (J10-J18)	53	0.5	0.6
8	Diabetes mellitus (E10-E14)	51	0.4	0.6
9	Legal intervention (Y35,Y89.0)	47	0.4	0.5
10	Cerebrovascular diseases (I60-I69)	44	0.4	0.5
11	Septicemia (A40-A41)	37	0.3	0.4
12	Human immunodeficiency virus (HIV) disease (B20-B24)	31	0.3	0.4
13	Chronic lower respiratory diseases (J40-J47)	24	0.2	0.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	0.2	0.3
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	0.2	0.3
...	All other causes (Residual)	1,190	10.4	13.9

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 25-29 years				
...	All causes	10,323	100.0	126.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,740	45.9	58.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,653	16.0	20.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	907	8.8	11.1
4	Malignant neoplasms (C00-C97)	572	5.5	7.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	552	5.3	6.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	101	1.0	1.2
7	Human immunodeficiency virus (HIV) disease (B20-B24)	91	0.9	1.1
8	Diabetes mellitus (E10-E14)	71	0.7	0.9
9	Cerebrovascular diseases (I60-I69)	68	0.7	0.8
10	Influenza and pneumonia (J10-J18)	58	0.6	0.7
11	Septicemia (A40-A41)	46	0.4	0.6
12	Chronic liver disease and cirrhosis (K70,K73-K74)	42	0.4	0.5
13	Chronic lower respiratory diseases (J40-J47)	36	0.3	0.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	29	0.3	0.4
14	Legal intervention (Y35,Y89.0)	29	0.3	0.4
...	All other causes (Residual)	1,328	12.9	16.3
White, male, 30-34 years				
...	All causes	10,909	100.0	135.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,136	37.9	51.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,794	16.4	22.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	917	8.4	11.4
4	Malignant neoplasms (C00-C97)	806	7.4	10.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	680	6.2	8.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	246	2.3	3.0
7	Diabetes mellitus (E10-E14)	156	1.4	1.9
8	Cerebrovascular diseases (I60-I69)	129	1.2	1.6
9	Chronic liver disease and cirrhosis (K70,K73-K74)	110	1.0	1.4
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	92	0.8	1.1
11	Influenza and pneumonia (J10-J18)	84	0.8	1.0
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	55	0.5	0.7
13	Septicemia (A40-A41)	51	0.5	0.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	40	0.4	0.5
15	Legal intervention (Y35,Y89.0)	38	0.3	0.5
...	All other causes (Residual)	1,575	14.4	19.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 35-39 years				
...	All causes	15,031	100.0	176.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,445	29.6	52.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,023	13.5	23.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,998	13.3	23.4
4	Malignant neoplasms (C00-C97)	1,457	9.7	17.1
5	Human immunodeficiency virus (HIV) disease (B20-B24)	579	3.9	6.8
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	553	3.7	6.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	412	2.7	4.8
8	Diabetes mellitus (E10-E14)	301	2.0	3.5
9	Cerebrovascular diseases (I60-I69)	241	1.6	2.8
10	Influenza and pneumonia (J10-J18)	137	0.9	1.6
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	102	0.7	1.2
12	Septicemia (A40-A41)	90	0.6	1.1
13	Chronic lower respiratory diseases (J40-J47)	86	0.6	1.0
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	83	0.6	1.0
15	Viral hepatitis (B15-B19)	73	0.5	0.9
...	All other causes (Residual)	2,451	16.3	28.7
White, male, 40-44 years				
...	All causes	25,793	100.0	276.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,651	21.9	60.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,646	18.0	49.8
3	Malignant neoplasms (C00-C97)	3,615	14.0	38.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,549	9.9	27.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,127	4.4	12.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	1,029	4.0	11.0
7	Diabetes mellitus (E10-E14)	611	2.4	6.5
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	581	2.3	6.2
9	Cerebrovascular diseases (I60-I69)	513	2.0	5.5
10	Influenza and pneumonia (J10-J18)	254	1.0	2.7
11	Chronic lower respiratory diseases (J40-J47)	225	0.9	2.4
12	Viral hepatitis (B15-B19)	222	0.9	2.4
13	Septicemia (A40-A41)	186	0.7	2.0
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	155	0.6	1.7
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	118	0.5	1.3
...	All other causes (Residual)	4,311	16.7	46.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 45-49 years				
...	All causes	38,117	100.0	413.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,389	22.0	91.0
2	Malignant neoplasms (C00-C97)	7,302	19.2	79.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,886	15.4	63.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,627	6.9	28.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,009	5.3	21.8
6	Diabetes mellitus (E10-E14)	1,039	2.7	11.3
7	Human immunodeficiency virus (HIV) disease (B20-B24)	921	2.4	10.0
8	Cerebrovascular diseases (I60-I69)	880	2.3	9.5
9	Viral hepatitis (B15-B19)	588	1.5	6.4
10	Chronic lower respiratory diseases (J40-J47)	556	1.5	6.0
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	513	1.3	5.6
12	Influenza and pneumonia (J10-J18)	416	1.1	4.5
13	Septicemia (A40-A41)	332	0.9	3.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	316	0.8	3.4
15	Aortic aneurysm and dissection (I71)	176	0.5	1.9
...	All other causes (Residual)	6,167	16.2	66.9
White, male, 50-54 years				
...	All causes	50,765	100.0	616.6
1	Malignant neoplasms (C00-C97)	12,892	25.4	156.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,784	25.2	155.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,828	9.5	58.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	2,624	5.2	31.9
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,279	4.5	27.7
6	Diabetes mellitus (E10-E14)	1,516	3.0	18.4
7	Cerebrovascular diseases (I60-I69)	1,359	2.7	16.5
8	Chronic lower respiratory diseases (J40-J47)	1,085	2.1	13.2
9	Viral hepatitis (B15-B19)	821	1.6	10.0
10	Human immunodeficiency virus (HIV) disease (B20-B24)	692	1.4	8.4
11	Influenza and pneumonia (J10-J18)	543	1.1	6.6
12	Septicemia (A40-A41)	468	0.9	5.7
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	452	0.9	5.5
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	368	0.7	4.5
15	Aortic aneurysm and dissection (I71)	229	0.5	2.8
...	All other causes (Residual)	7,825	15.4	95.0

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 55-59 years				
...	All causes	62,213	100.0	864.0
1	Malignant neoplasms (C00-C97)	19,566	31.5	271.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	16,675	26.8	231.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,544	5.7	49.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	2,517	4.0	35.0
5	Diabetes mellitus (E10-E14)	2,305	3.7	32.0
6	Chronic lower respiratory diseases (J40-J47)	2,052	3.3	28.5
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,784	2.9	24.8
8	Cerebrovascular diseases (I60-I69)	1,742	2.8	24.2
9	Septicemia (A40-A41)	696	1.1	9.7
10	Influenza and pneumonia (J10-J18)	679	1.1	9.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	642	1.0	8.9
12	Viral hepatitis (B15-B19)	544	0.9	7.6
13	Human immunodeficiency virus (HIV) disease (B20-B24)	402	0.6	5.6
14	Aortic aneurysm and dissection (I71)	366	0.6	5.1
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	326	0.5	4.5
...	All other causes (Residual)	8,373	13.5	116.3
White, male, 60-64 years				
...	All causes	72,059	100.0	1,341.8
1	Malignant neoplasms (C00-C97)	25,250	35.0	470.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	19,560	27.1	364.2
3	Chronic lower respiratory diseases (J40-J47)	3,769	5.2	70.2
4	Diabetes mellitus (E10-E14)	2,740	3.8	51.0
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,641	3.7	49.2
6	Cerebrovascular diseases (I60-I69)	2,244	3.1	41.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,890	2.6	35.2
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,296	1.8	24.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	903	1.3	16.8
10	Influenza and pneumonia (J10-J18)	866	1.2	16.1
11	Septicemia (A40-A41)	848	1.2	15.8
12	Aortic aneurysm and dissection (I71)	537	0.7	10.0
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	401	0.6	7.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	381	0.5	7.1
15	Pneumonitis due to solids and liquids (J69)	242	0.3	4.5
...	All other causes (Residual)	8,491	11.8	158.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 65-69 years				
...	All causes	82,421	100.0	2,016.7
1	Malignant neoplasms (C00-C97)	29,911	36.3	731.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	21,564	26.2	527.6
3	Chronic lower respiratory diseases (J40-J47)	5,665	6.9	138.6
4	Diabetes mellitus (E10-E14)	3,056	3.7	74.8
5	Cerebrovascular diseases (I60-I69)	3,038	3.7	74.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,193	2.7	53.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,431	1.7	35.0
8	Influenza and pneumonia (J10-J18)	1,129	1.4	27.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,107	1.3	27.1
10	Septicemia (A40-A41)	1,028	1.2	25.2
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	925	1.1	22.6
12	Aortic aneurysm and dissection (I71)	701	0.9	17.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	505	0.6	12.4
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	498	0.6	12.2
15	Parkinson's disease (G20-G21)	407	0.5	10.0
...	All other causes (Residual)	9,263	11.2	226.7
White, male, 70-74 years				
...	All causes	107,129	100.0	3,208.8
1	Malignant neoplasms (C00-C97)	36,169	33.8	1,083.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	28,331	26.4	848.6
3	Chronic lower respiratory diseases (J40-J47)	8,385	7.8	251.2
4	Cerebrovascular diseases (I60-I69)	4,607	4.3	138.0
5	Diabetes mellitus (E10-E14)	3,722	3.5	111.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,393	2.2	71.7
7	Influenza and pneumonia (J10-J18)	1,885	1.8	56.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,760	1.6	52.7
9	Septicemia (A40-A41)	1,395	1.3	41.8
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,330	1.2	39.8
11	Alzheimer's disease (G30)	1,066	1.0	31.9
12	Parkinson's disease (G20-G21)	1,065	1.0	31.9
13	Aortic aneurysm and dissection (I71)	1,022	1.0	30.6
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	923	0.9	27.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	701	0.7	21.0
...	All other causes (Residual)	12,375	11.6	370.7

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 75-79 years				
...	All causes	141,070	100.0	5,065.8
1	Malignant neoplasms (C00-C97)	40,996	29.1	1,472.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	39,051	27.7	1,402.3
3	Chronic lower respiratory diseases (J40-J47)	11,324	8.0	406.6
4	Cerebrovascular diseases (I60-I69)	7,419	5.3	266.4
5	Diabetes mellitus (E10-E14)	4,583	3.2	164.6
6	Influenza and pneumonia (J10-J18)	3,279	2.3	117.7
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,064	2.2	110.0
8	Alzheimer's disease (G30)	2,715	1.9	97.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,670	1.9	95.9
10	Parkinson's disease (G20-G21)	2,214	1.6	79.5
11	Septicemia (A40-A41)	1,857	1.3	66.7
12	Aortic aneurysm and dissection (I71)	1,255	0.9	45.1
13	Chronic liver disease and cirrhosis (K70,K73-K74)	1,060	0.8	38.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,045	0.7	37.5
15	Pneumonitis due to solids and liquids (J69)	1,034	0.7	37.1
...	All other causes (Residual)	17,504	12.4	628.6
White, male, 80-84 years				
...	All causes	159,158	100.0	8,169.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	47,649	29.9	2,445.7
2	Malignant neoplasms (C00-C97)	36,919	23.2	1,894.9
3	Chronic lower respiratory diseases (J40-J47)	11,369	7.1	583.5
4	Cerebrovascular diseases (I60-I69)	9,845	6.2	505.3
5	Alzheimer's disease (G30)	4,954	3.1	254.3
6	Influenza and pneumonia (J10-J18)	4,946	3.1	253.9
7	Diabetes mellitus (E10-E14)	4,587	2.9	235.4
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,563	2.2	182.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,469	2.2	178.1
10	Parkinson's disease (G20-G21)	3,160	2.0	162.2
11	Septicemia (A40-A41)	2,201	1.4	113.0
12	Pneumonitis due to solids and liquids (J69)	1,677	1.1	86.1
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,342	0.8	68.9
14	Aortic aneurysm and dissection (I71)	1,308	0.8	67.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,206	0.8	61.9
...	All other causes (Residual)	20,963	13.2	1,076.0

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 85-89 years				
...	All causes	128,448	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	42,732	33.3	---
2	Malignant neoplasms (C00-C97)	22,593	17.6	---
3	Cerebrovascular diseases (I60-I69)	8,794	6.8	---
4	Chronic lower respiratory diseases (J40-J47)	7,771	6.0	---
5	Alzheimer's disease (G30)	5,400	4.2	---
6	Influenza and pneumonia (J10-J18)	4,969	3.9	---
7	Diabetes mellitus (E10-E14)	3,197	2.5	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,155	2.5	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,098	2.4	---
10	Parkinson's disease (G20-G21)	2,367	1.8	---
11	Pneumonitis due to solids and liquids (J69)	1,829	1.4	---
12	Septicemia (A40-A41)	1,706	1.3	---
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,267	1.0	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	918	0.7	---
15	Atherosclerosis (I70)	902	0.7	---
...	All other causes (Residual)	17,750	13.8	---
White, male, 90-94 years				
...	All causes	69,060	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	25,283	36.6	---
2	Malignant neoplasms (C00-C97)	8,972	13.0	---
3	Cerebrovascular diseases (I60-I69)	4,781	6.9	---
4	Influenza and pneumonia (J10-J18)	3,541	5.1	---
5	Chronic lower respiratory diseases (J40-J47)	3,524	5.1	---
6	Alzheimer's disease (G30)	3,196	4.6	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,695	2.5	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,545	2.2	---
9	Diabetes mellitus (E10-E14)	1,352	2.0	---
10	Pneumonitis due to solids and liquids (J69)	1,177	1.7	---
11	Parkinson's disease (G20-G21)	959	1.4	---
12	Septicemia (A40-A41)	814	1.2	---
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	769	1.1	---
14	Atherosclerosis (I70)	673	1.0	---
15	Aortic aneurysm and dissection (I71)	397	0.6	---
...	All other causes (Residual)	10,382	15.0	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 95-99 years				
...	All causes	18,939	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,341	38.8	---
2	Malignant neoplasms (C00-C97)	1,843	9.7	---
3	Cerebrovascular diseases (I60-I69)	1,220	6.4	---
4	Influenza and pneumonia (J10-J18)	1,180	6.2	---
5	Alzheimer's disease (G30)	877	4.6	---
6	Chronic lower respiratory diseases (J40-J47)	846	4.5	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	480	2.5	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	454	2.4	---
9	Pneumonitis due to solids and liquids (J69)	325	1.7	---
10	Diabetes mellitus (E10-E14)	287	1.5	---
11	Atherosclerosis (I70)	235	1.2	---
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	234	1.2	---
13	Septicemia (A40-A41)	214	1.1	---
14	Parkinson's disease (G20-G21)	171	0.9	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	77	0.4	---
15	Aortic aneurysm and dissection (I71)	77	0.4	---
...	All other causes (Residual)	3,078	16.3	---
White, male, 100 years and over				
...	All causes	2,543	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,054	41.4	---
2	Influenza and pneumonia (J10-J18)	199	7.8	---
3	Malignant neoplasms (C00-C97)	151	5.9	---
4	Cerebrovascular diseases (I60-I69)	140	5.5	---
5	Alzheimer's disease (G30)	86	3.4	---
6	Chronic lower respiratory diseases (J40-J47)	84	3.3	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	69	2.7	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	56	2.2	---
9	Pneumonitis due to solids and liquids (J69)	53	2.1	---
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	44	1.7	---
10	Atherosclerosis (I70)	44	1.7	---
12	Septicemia (A40-A41)	22	0.9	---
13	Diabetes mellitus (E10-E14)	21	0.8	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	0.4	---
15	Nutritional deficiencies (E40-E64)	9	0.4	---
15	Parkinson's disease (G20-G21)	9	0.4	---
15	Hyperplasia of prostate (N40)	9	0.4	---
...	All other causes (Residual)	483	19.0	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, female, all ages				
...	All causes	1,069,945	100.0	882.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	285,472	26.7	235.5
2	Malignant neoplasms (C00-C97)	231,654	21.7	191.1
3	Cerebrovascular diseases (I60-I69)	74,674	7.0	61.6
4	Chronic lower respiratory diseases (J40-J47)	63,973	6.0	52.8
5	Alzheimer's disease (G30)	47,201	4.4	38.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	35,806	3.3	29.5
7	Influenza and pneumonia (J10-J18)	31,115	2.9	25.7
8	Diabetes mellitus (E10-E14)	30,127	2.8	24.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17,669	1.7	14.6
10	Septicemia (A40-A41)	15,022	1.4	12.4
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	12,236	1.1	10.1
12	Chronic liver disease and cirrhosis (K70,K73-K74)	8,324	0.8	6.9
13	Parkinson's disease (G20-G21)	7,847	0.7	6.5
14	Pneumonitis due to solids and liquids (J69)	7,579	0.7	6.3
15	Atherosclerosis (I70)	6,752	0.6	5.6
...	All other causes (Residual)	194,494	18.2	160.5
White, female, 1-4 years				
...	All causes	1,412	100.0	22.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	469	33.2	7.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	184	13.0	3.0
3	Malignant neoplasms (C00-C97)	125	8.9	2.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	98	6.9	1.6
5	Influenza and pneumonia (J10-J18)	40	2.8	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	38	2.7	0.6
7	Septicemia (A40-A41)	26	1.8	0.4
8	Cerebrovascular diseases (I60-I69)	17	1.2	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	1.1	*
10	Certain conditions originating in the perinatal period (P00-P96)	15	1.1	*
11	Chronic lower respiratory diseases (J40-J47)	13	0.9	*
12	Anemias (D50-D64)	7	0.5	*
12	Meningitis (G00,G03)	7	0.5	*
14	Meningococcal infection (A39)	5	0.4	*
15	Acute bronchitis and bronchiolitis (J20-J21)	4	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.3	*
...	All other causes (Residual)	340	24.1	5.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 5-9 years				
...	All causes	909	100.0	12.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	318	35.0	4.3
2	Malignant neoplasms (C00-C97)	179	19.7	2.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	76	8.4	1.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	30	3.3	0.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	26	2.9	0.4
6	Influenza and pneumonia (J10-J18)	21	2.3	0.3
7	Cerebrovascular diseases (I60-I69)	18	2.0	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	15	1.7	*
9	Septicemia (A40-A41)	10	1.1	*
9	Chronic lower respiratory diseases (J40-J47)	10	1.1	*
11	Meningitis (G00,G03)	7	0.8	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.6	*
12	Certain conditions originating in the perinatal period (P00-P96)	5	0.6	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.3	*
15	Anemias (D50-D64)	2	0.2	*
15	Diabetes mellitus (E10-E14)	2	0.2	*
15	Nutritional deficiencies (E40-E64)	2	0.2	*
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2	0.2	*
...	All other causes (Residual)	178	19.6	2.4
White, female, 10-14 years				
...	All causes	1,052	100.0	13.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	366	34.8	4.7
2	Malignant neoplasms (C00-C97)	179	17.0	2.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	64	6.1	0.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	49	4.7	0.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	46	4.4	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	39	3.7	0.5
7	Influenza and pneumonia (J10-J18)	20	1.9	0.3
8	Septicemia (A40-A41)	14	1.3	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	1.2	*
10	Diabetes mellitus (E10-E14)	12	1.1	*
11	Cerebrovascular diseases (I60-I69)	11	1.0	*
12	Chronic lower respiratory diseases (J40-J47)	9	0.9	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.6	*
13	Certain conditions originating in the perinatal period (P00-P96)	6	0.6	*
15	Anemias (D50-D64)	4	0.4	*
...	All other causes (Residual)	214	20.3	2.7

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 15-19 years				
...	All causes	2,969	100.0	37.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,673	56.3	21.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	250	8.4	3.1
3	Malignant neoplasms (C00-C97)	232	7.8	2.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	140	4.7	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	86	2.9	1.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	59	2.0	0.7
7	Cerebrovascular diseases (I60-I69)	30	1.0	0.4
8	Pregnancy, childbirth and the puerperium (O00-099)	21	0.7	0.3
9	Diabetes mellitus (E10-E14)	19	0.6	*
10	Septicemia (A40-A41)	16	0.5	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	0.5	*
11	Influenza and pneumonia (J10-J18)	14	0.5	*
13	Chronic lower respiratory diseases (J40-J47)	12	0.4	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	0.4	*
15	Meningococcal infection (A39)	7	0.2	*
15	Pneumonitis due to solids and liquids (J69)	7	0.2	*
...	All other causes (Residual)	377	12.7	4.7
White, female, 20-24 years				
...	All causes	3,649	100.0	45.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,617	44.3	20.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	345	9.5	4.3
3	Malignant neoplasms (C00-C97)	307	8.4	3.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	232	6.4	2.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	141	3.9	1.8
6	Pregnancy, childbirth and the puerperium (O00-099)	89	2.4	1.1
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	75	2.1	0.9
8	Cerebrovascular diseases (I60-I69)	41	1.1	0.5
9	Diabetes mellitus (E10-E14)	32	0.9	0.4
10	Influenza and pneumonia (J10-J18)	30	0.8	0.4
11	Septicemia (A40-A41)	23	0.6	0.3
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	0.4	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	15	0.4	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	13	0.4	*
14	Chronic lower respiratory diseases (J40-J47)	13	0.4	*
...	All other causes (Residual)	660	18.1	8.3

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 25-29 years				
...	All causes	3,756	100.0	49.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,326	35.3	17.4
2	Malignant neoplasms (C00-C97)	419	11.2	5.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	323	8.6	4.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	226	6.0	3.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	207	5.5	2.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	79	2.1	1.0
7	Pregnancy, childbirth and the puerperium (O00-099)	78	2.1	1.0
8	Influenza and pneumonia (J10-J18)	53	1.4	0.7
9	Diabetes mellitus (E10-E14)	52	1.4	0.7
10	Cerebrovascular diseases (I60-I69)	50	1.3	0.7
11	Human immunodeficiency virus (HIV) disease (B20-B24)	36	1.0	0.5
12	Septicemia (A40-A41)	34	0.9	0.4
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	33	0.9	0.4
14	Chronic lower respiratory diseases (J40-J47)	28	0.7	0.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	24	0.6	0.3
...	All other causes (Residual)	788	21.0	10.4
White, female, 30-34 years				
...	All causes	5,115	100.0	66.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,426	27.9	18.6
2	Malignant neoplasms (C00-C97)	897	17.5	11.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	459	9.0	6.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	408	8.0	5.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	209	4.1	2.7
6	Cerebrovascular diseases (I60-I69)	109	2.1	1.4
7	Pregnancy, childbirth and the puerperium (O00-099)	101	2.0	1.3
8	Diabetes mellitus (E10-E14)	93	1.8	1.2
9	Human immunodeficiency virus (HIV) disease (B20-B24)	91	1.8	1.2
10	Influenza and pneumonia (J10-J18)	67	1.3	0.9
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	67	1.3	0.9
12	Chronic liver disease and cirrhosis (K70,K73-K74)	62	1.2	0.8
13	Septicemia (A40-A41)	60	1.2	0.8
14	Chronic lower respiratory diseases (J40-J47)	52	1.0	0.7
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	37	0.7	0.5
...	All other causes (Residual)	977	19.1	12.7

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 35-39 years				
...	All causes	8,146	100.0	99.0
1	Malignant neoplasms (C00-C97)	1,919	23.6	23.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,719	21.1	20.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	795	9.8	9.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	563	6.9	6.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	241	3.0	2.9
6	Cerebrovascular diseases (I60-I69)	220	2.7	2.7
7	Diabetes mellitus (E10-E14)	209	2.6	2.5
8	Chronic liver disease and cirrhosis (K70,K73-K74)	197	2.4	2.4
9	Human immunodeficiency virus (HIV) disease (B20-B24)	131	1.6	1.6
10	Septicemia (A40-A41)	109	1.3	1.3
11	Chronic lower respiratory diseases (J40-J47)	95	1.2	1.2
12	Influenza and pneumonia (J10-J18)	94	1.2	1.1
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	90	1.1	1.1
14	Pregnancy, childbirth and the puerperium (O00-099)	80	1.0	1.0
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	64	0.8	0.8
...	All other causes (Residual)	1,620	19.9	19.7
White, female, 40-44 years				
...	All causes	14,614	100.0	158.6
1	Malignant neoplasms (C00-C97)	4,361	29.8	47.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,357	16.1	25.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,662	11.4	18.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	782	5.4	8.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	548	3.7	5.9
6	Cerebrovascular diseases (I60-I69)	476	3.3	5.2
7	Diabetes mellitus (E10-E14)	307	2.1	3.3
8	Chronic lower respiratory diseases (J40-J47)	241	1.6	2.6
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	240	1.6	2.6
10	Human immunodeficiency virus (HIV) disease (B20-B24)	219	1.5	2.4
11	Influenza and pneumonia (J10-J18)	198	1.4	2.1
12	Septicemia (A40-A41)	167	1.1	1.8
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	117	0.8	1.3
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	99	0.7	1.1
15	Viral hepatitis (B15-B19)	98	0.7	1.1
...	All other causes (Residual)	2,742	18.8	29.7

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 45-49 years				
...	All causes	22,169	100.0	239.9
1	Malignant neoplasms (C00-C97)	7,833	35.3	84.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,935	13.2	31.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,486	11.2	26.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	870	3.9	9.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	841	3.8	9.1
6	Cerebrovascular diseases (I60-I69)	794	3.6	8.6
7	Chronic lower respiratory diseases (J40-J47)	586	2.6	6.3
8	Diabetes mellitus (E10-E14)	549	2.5	5.9
9	Influenza and pneumonia (J10-J18)	293	1.3	3.2
10	Septicemia (A40-A41)	270	1.2	2.9
11	Viral hepatitis (B15-B19)	200	0.9	2.2
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	187	0.8	2.0
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	184	0.8	2.0
14	Human immunodeficiency virus (HIV) disease (B20-B24)	176	0.8	1.9
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	132	0.6	1.4
...	All other causes (Residual)	3,833	17.3	41.5
White, female, 50-54 years				
...	All causes	29,173	100.0	347.5
1	Malignant neoplasms (C00-C97)	11,690	40.1	139.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,266	14.6	50.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,935	6.6	23.0
4	Cerebrovascular diseases (I60-I69)	1,050	3.6	12.5
5	Chronic lower respiratory diseases (J40-J47)	981	3.4	11.7
6	Diabetes mellitus (E10-E14)	975	3.3	11.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	868	3.0	10.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	729	2.5	8.7
9	Septicemia (A40-A41)	429	1.5	5.1
10	Influenza and pneumonia (J10-J18)	371	1.3	4.4
11	Viral hepatitis (B15-B19)	286	1.0	3.4
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	269	0.9	3.2
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	162	0.6	1.9
14	Human immunodeficiency virus (HIV) disease (B20-B24)	158	0.5	1.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	147	0.5	1.8
...	All other causes (Residual)	4,857	16.6	57.9

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 55-59 years				
...	All causes	38,999	100.0	521.9
1	Malignant neoplasms (C00-C97)	16,711	42.8	223.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,309	16.2	84.4
3	Chronic lower respiratory diseases (J40-J47)	2,008	5.1	26.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,553	4.0	20.8
5	Diabetes mellitus (E10-E14)	1,493	3.8	20.0
6	Cerebrovascular diseases (I60-I69)	1,359	3.5	18.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	929	2.4	12.4
8	Septicemia (A40-A41)	618	1.6	8.3
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	554	1.4	7.4
10	Influenza and pneumonia (J10-J18)	496	1.3	6.6
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	485	1.2	6.5
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	227	0.6	3.0
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	192	0.5	2.6
14	Viral hepatitis (B15-B19)	181	0.5	2.4
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	168	0.4	2.2
...	All other causes (Residual)	5,716	14.7	76.5
White, female, 60-64 years				
...	All causes	48,947	100.0	847.6
1	Malignant neoplasms (C00-C97)	20,912	42.7	362.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,863	18.1	153.5
3	Chronic lower respiratory diseases (J40-J47)	3,543	7.2	61.4
4	Diabetes mellitus (E10-E14)	1,920	3.9	33.2
5	Cerebrovascular diseases (I60-I69)	1,795	3.7	31.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,294	2.6	22.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	802	1.6	13.9
8	Septicemia (A40-A41)	715	1.5	12.4
9	Influenza and pneumonia (J10-J18)	659	1.3	11.4
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	633	1.3	11.0
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	345	0.7	6.0
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	292	0.6	5.1
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	238	0.5	4.1
14	Alzheimer's disease (G30)	225	0.5	3.9
15	Aortic aneurysm and dissection (I71)	205	0.4	3.5
...	All other causes (Residual)	6,506	13.3	112.7

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 65-69 years				
...	All causes	60,781	100.0	1,322.9
1	Malignant neoplasms (C00-C97)	23,787	39.1	517.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,881	19.5	258.6
3	Chronic lower respiratory diseases (J40-J47)	5,290	8.7	115.1
4	Cerebrovascular diseases (I60-I69)	2,657	4.4	57.8
5	Diabetes mellitus (E10-E14)	2,439	4.0	53.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,264	2.1	27.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	965	1.6	21.0
8	Septicemia (A40-A41)	957	1.6	20.8
9	Influenza and pneumonia (J10-J18)	894	1.5	19.5
10	Chronic liver disease and cirrhosis (K70,K73-K74)	871	1.4	19.0
11	Alzheimer's disease (G30)	470	0.8	10.2
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	422	0.7	9.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	337	0.6	7.3
14	Aortic aneurysm and dissection (I71)	318	0.5	6.9
15	Parkinson's disease (G20-G21)	211	0.3	4.6
...	All other causes (Residual)	8,018	13.2	174.5
White, female, 70-74 years				
...	All causes	86,031	100.0	2,129.8
1	Malignant neoplasms (C00-C97)	29,413	34.2	728.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	18,497	21.5	457.9
3	Chronic lower respiratory diseases (J40-J47)	8,177	9.5	202.4
4	Cerebrovascular diseases (I60-I69)	4,527	5.3	112.1
5	Diabetes mellitus (E10-E14)	3,259	3.8	80.7
6	Influenza and pneumonia (J10-J18)	1,582	1.8	39.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,574	1.8	39.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,511	1.8	37.4
9	Alzheimer's disease (G30)	1,448	1.7	35.8
10	Septicemia (A40-A41)	1,371	1.6	33.9
11	Chronic liver disease and cirrhosis (K70,K73-K74)	949	1.1	23.5
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	699	0.8	17.3
13	Parkinson's disease (G20-G21)	541	0.6	13.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	524	0.6	13.0
15	Aortic aneurysm and dissection (I71)	498	0.6	12.3
...	All other causes (Residual)	11,461	13.3	283.7

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 75-79 years				
...	All causes	129,296	100.0	3,435.6
1	Malignant neoplasms (C00-C97)	35,282	27.3	937.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	31,221	24.1	829.6
3	Chronic lower respiratory diseases (J40-J47)	11,555	8.9	307.0
4	Cerebrovascular diseases (I60-I69)	8,571	6.6	227.7
5	Diabetes mellitus (E10-E14)	4,581	3.5	121.7
6	Alzheimer's disease (G30)	4,180	3.2	111.1
7	Influenza and pneumonia (J10-J18)	2,950	2.3	78.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,482	1.9	66.0
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,380	1.8	63.2
10	Septicemia (A40-A41)	1,999	1.5	53.1
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,307	1.0	34.7
12	Parkinson's disease (G20-G21)	1,303	1.0	34.6
13	Chronic liver disease and cirrhosis (K70,K73-K74)	958	0.7	25.5
14	Aortic aneurysm and dissection (I71)	871	0.7	23.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	833	0.6	22.1
...	All other causes (Residual)	18,823	14.6	500.2
White, female, 80-84 years				
...	All causes	181,296	100.0	5,831.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	50,790	28.0	1,633.6
2	Malignant neoplasms (C00-C97)	35,535	19.6	1,143.0
3	Cerebrovascular diseases (I60-I69)	14,997	8.3	482.4
4	Chronic lower respiratory diseases (J40-J47)	12,984	7.2	417.6
5	Alzheimer's disease (G30)	9,423	5.2	303.1
6	Diabetes mellitus (E10-E14)	5,461	3.0	175.7
7	Influenza and pneumonia (J10-J18)	5,372	3.0	172.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,517	1.9	113.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,374	1.9	108.5
10	Septicemia (A40-A41)	2,772	1.5	89.2
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,172	1.2	69.9
12	Parkinson's disease (G20-G21)	2,120	1.2	68.2
13	Pneumonitis due to solids and liquids (J69)	1,426	0.8	45.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,136	0.6	36.5
15	Atherosclerosis (I70)	1,126	0.6	36.2
...	All other causes (Residual)	29,091	16.0	935.7

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 85-89 years				
...	All causes	192,887	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	61,756	32.0	---
2	Malignant neoplasms (C00-C97)	25,281	13.1	---
3	Cerebrovascular diseases (I60-I69)	17,435	9.0	---
4	Alzheimer's disease (G30)	13,155	6.8	---
5	Chronic lower respiratory diseases (J40-J47)	10,507	5.4	---
6	Influenza and pneumonia (J10-J18)	7,039	3.6	---
7	Diabetes mellitus (E10-E14)	4,804	2.5	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,744	1.9	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,548	1.8	---
10	Septicemia (A40-A41)	2,696	1.4	---
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,661	1.4	---
12	Parkinson's disease (G20-G21)	2,066	1.1	---
13	Pneumonitis due to solids and liquids (J69)	1,821	0.9	---
14	Atherosclerosis (I70)	1,642	0.9	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,167	0.6	---
...	All other causes (Residual)	33,565	17.4	---
White, female, 90-94 years				
...	All causes	150,793	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	54,398	36.1	---
2	Cerebrovascular diseases (I60-I69)	13,864	9.2	---
3	Malignant neoplasms (C00-C97)	12,519	8.3	---
4	Alzheimer's disease (G30)	11,781	7.8	---
5	Influenza and pneumonia (J10-J18)	6,586	4.4	---
6	Chronic lower respiratory diseases (J40-J47)	5,717	3.8	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,949	2.0	---
8	Diabetes mellitus (E10-E14)	2,808	1.9	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,642	1.8	---
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,590	1.7	---
11	Septicemia (A40-A41)	1,868	1.2	---
12	Atherosclerosis (I70)	1,667	1.1	---
13	Pneumonitis due to solids and liquids (J69)	1,573	1.0	---
14	Parkinson's disease (G20-G21)	1,115	0.7	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	789	0.5	---
...	All other causes (Residual)	27,927	18.5	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 95-99 years				
...	All causes	65,574	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	25,143	38.3	---
2	Cerebrovascular diseases (I60-I69)	5,530	8.4	---
3	Alzheimer's disease (G30)	5,387	8.2	---
4	Malignant neoplasms (C00-C97)	3,552	5.4	---
5	Influenza and pneumonia (J10-J18)	3,366	5.1	---
6	Chronic lower respiratory diseases (J40-J47)	1,868	2.8	---
7	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,339	2.0	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,201	1.8	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,054	1.6	---
10	Diabetes mellitus (E10-E14)	959	1.5	---
11	Atherosclerosis (I70)	936	1.4	---
12	Pneumonitis due to solids and liquids (J69)	715	1.1	---
13	Septicemia (A40-A41)	693	1.1	---
14	Parkinson's disease (G20-G21)	288	0.4	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	253	0.4	---
...	All other causes (Residual)	13,290	20.3	---
White, female, 100 years and over				
...	All causes	14,295	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,896	41.2	---
2	Cerebrovascular diseases (I60-I69)	1,090	7.6	---
3	Alzheimer's disease (G30)	978	6.8	---
4	Influenza and pneumonia (J10-J18)	908	6.4	---
5	Malignant neoplasms (C00-C97)	493	3.4	---
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	286	2.0	---
7	Chronic lower respiratory diseases (J40-J47)	276	1.9	---
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	274	1.9	---
9	Atherosclerosis (I70)	257	1.8	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	201	1.4	---
11	Diabetes mellitus (E10-E14)	151	1.1	---
12	Pneumonitis due to solids and liquids (J69)	137	1.0	---
13	Septicemia (A40-A41)	110	0.8	---
14	Anemias (D50-D64)	71	0.5	---
15	Nutritional deficiencies (E40-E64)	49	0.3	---
...	All other causes (Residual)	3,118	21.8	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, all ages				
...	All causes	292,808	100.0	749.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	74,159	25.3	189.8
2	Malignant neoplasms (C00-C97)	63,165	21.6	161.7
3	Cerebrovascular diseases (I60-I69)	17,541	6.0	44.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,652	4.7	34.9
5	Diabetes mellitus (E10-E14)	12,970	4.4	33.2
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8,669	3.0	22.2
7	Chronic lower respiratory diseases (J40-J47)	8,229	2.8	21.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,075	2.8	20.7
9	Human immunodeficiency virus (HIV) disease (B20-B24)	7,022	2.4	18.0
10	Septicemia (A40-A41)	6,221	2.1	15.9
11	Influenza and pneumonia (J10-J18)	5,780	2.0	14.8
12	Certain conditions originating in the perinatal period (P00-P96)	5,050	1.7	12.9
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	4,953	1.7	12.7
14	Alzheimer's disease (G30)	4,620	1.6	11.8
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,530	0.9	6.5
...	All other causes (Residual)	50,172	17.1	128.4
Black, both sexes, 1-4 years				
...	All causes	1,097	100.0	41.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	327	29.8	12.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	140	12.8	5.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	106	9.7	4.0
4	Malignant neoplasms (C00-C97)	64	5.8	2.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	45	4.1	1.7
6	Chronic lower respiratory diseases (J40-J47)	31	2.8	1.2
7	Influenza and pneumonia (J10-J18)	29	2.6	1.1
8	Certain conditions originating in the perinatal period (P00-P96)	23	2.1	0.9
9	Septicemia (A40-A41)	20	1.8	0.8
10	Cerebrovascular diseases (I60-I69)	14	1.3	*
11	Anemias (D50-D64)	11	1.0	*
12	Meningitis (G00,G03)	6	0.5	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.5	*
13	Pneumonitis due to solids and liquids (J69)	5	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.3	*
...	All other causes (Residual)	268	24.4	10.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 5-9 years				
...	All causes	663	100.0	21.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	236	35.6	7.5
2	Malignant neoplasms (C00-C97)	82	12.4	2.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	45	6.8	1.4
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	41	6.2	1.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	35	5.3	1.1
6	Chronic lower respiratory diseases (J40-J47)	23	3.5	0.7
7	Cerebrovascular diseases (I60-I69)	14	2.1	*
8	Septicemia (A40-A41)	11	1.7	*
8	Influenza and pneumonia (J10-J18)	11	1.7	*
10	Anemias (D50-D64)	8	1.2	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	1.1	*
12	Meningitis (G00,G03)	3	0.5	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.3	*
13	Pneumonitis due to solids and liquids (J69)	2	0.3	*
13	Certain conditions originating in the perinatal period (P00-P96)	2	0.3	*
...	All other causes (Residual)	141	21.3	4.5
Black, both sexes, 10-14 years				
...	All causes	880	100.0	25.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	265	30.1	7.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	105	11.9	3.0
3	Malignant neoplasms (C00-C97)	90	10.2	2.6
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	47	5.3	1.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	42	4.8	1.2
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	40	4.5	1.1
7	Chronic lower respiratory diseases (J40-J47)	30	3.4	0.9
8	Diabetes mellitus (E10-E14)	12	1.4	*
9	Influenza and pneumonia (J10-J18)	11	1.3	*
10	Septicemia (A40-A41)	10	1.1	*
10	Cerebrovascular diseases (I60-I69)	10	1.1	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	7	0.8	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.8	*
12	Anemias (D50-D64)	7	0.8	*
15	Meningitis (G00,G03)	4	0.5	*
15	Diseases of appendix (K35-K38)	4	0.5	*
...	All other causes (Residual)	189	21.5	5.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 15-19 years				
...	All causes	2,839	100.0	83.6
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,138	40.1	33.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	773	27.2	22.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	148	5.2	4.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	116	4.1	3.4
5	Malignant neoplasms (C00-C97)	103	3.6	3.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	55	1.9	1.6
7	Anemias (D50-D64)	30	1.1	0.9
8	Chronic lower respiratory diseases (J40-J47)	25	0.9	0.7
9	Diabetes mellitus (E10-E14)	24	0.8	0.7
10	Human immunodeficiency virus (HIV) disease (B20-B24)	22	0.8	0.6
11	Septicemia (A40-A41)	20	0.7	0.6
11	Pregnancy, childbirth and the puerperium (O00-099)	20	0.7	0.6
13	Cerebrovascular diseases (I60-I69)	19	0.7	*
14	Influenza and pneumonia (J10-J18)	13	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	0.4	*
...	All other causes (Residual)	321	11.3	9.5
Black, both sexes, 20-24 years				
...	All causes	4,532	100.0	143.1
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,939	42.8	61.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,007	22.2	31.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	289	6.4	9.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	218	4.8	6.9
5	Malignant neoplasms (C00-C97)	161	3.6	5.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	87	1.9	2.7
7	Chronic lower respiratory diseases (J40-J47)	50	1.1	1.6
8	Diabetes mellitus (E10-E14)	49	1.1	1.5
8	Pregnancy, childbirth and the puerperium (O00-099)	49	1.1	1.5
10	Anemias (D50-D64)	47	1.0	1.5
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	45	1.0	1.4
12	Cerebrovascular diseases (I60-I69)	32	0.7	1.0
13	Septicemia (A40-A41)	24	0.5	0.8
14	Influenza and pneumonia (J10-J18)	18	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	0.4	*
...	All other causes (Residual)	500	11.0	15.8

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 25-29 years				
...	All causes	4,647	100.0	162.0
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,521	32.7	53.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	844	18.2	29.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	375	8.1	13.1
4	Human immunodeficiency virus (HIV) disease (B20-B24)	278	6.0	9.7
5	Malignant neoplasms (C00-C97)	263	5.7	9.2
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	252	5.4	8.8
7	Diabetes mellitus (E10-E14)	80	1.7	2.8
8	Pregnancy, childbirth and the puerperium (O00-099)	66	1.4	2.3
9	Anemias (D50-D64)	65	1.4	2.3
10	Cerebrovascular diseases (I60-I69)	64	1.4	2.2
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	47	1.0	1.6
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	46	1.0	1.6
13	Septicemia (A40-A41)	44	0.9	1.5
14	Chronic lower respiratory diseases (J40-J47)	34	0.7	1.2
15	Influenza and pneumonia (J10-J18)	28	0.6	1.0
...	All other causes (Residual)	640	13.8	22.3
Black, both sexes, 30-34 years				
...	All causes	5,389	100.0	197.3
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,070	19.9	39.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	952	17.7	34.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	647	12.0	23.7
4	Human immunodeficiency virus (HIV) disease (B20-B24)	558	10.4	20.4
5	Malignant neoplasms (C00-C97)	425	7.9	15.6
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	233	4.3	8.5
7	Diabetes mellitus (E10-E14)	139	2.6	5.1
8	Cerebrovascular diseases (I60-I69)	99	1.8	3.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	74	1.4	2.7
10	Anemias (D50-D64)	66	1.2	2.4
11	Septicemia (A40-A41)	64	1.2	2.3
12	Chronic lower respiratory diseases (J40-J47)	63	1.2	2.3
13	Pregnancy, childbirth and the puerperium (O00-099)	51	0.9	1.9
14	Influenza and pneumonia (J10-J18)	50	0.9	1.8
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	36	0.7	1.3
...	All other causes (Residual)	862	16.0	31.6

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 35-39 years				
...	All causes	7,048	100.0	254.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,152	16.3	41.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,001	14.2	36.1
3	Human immunodeficiency virus (HIV) disease (B20-B24)	951	13.5	34.3
4	Malignant neoplasms (C00-C97)	779	11.1	28.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	734	10.4	26.5
6	Diabetes mellitus (E10-E14)	224	3.2	8.1
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	199	2.8	7.2
8	Cerebrovascular diseases (I60-I69)	188	2.7	6.8
9	Septicemia (A40-A41)	101	1.4	3.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	98	1.4	3.5
11	Influenza and pneumonia (J10-J18)	72	1.0	2.6
12	Chronic lower respiratory diseases (J40-J47)	71	1.0	2.6
13	Anemias (D50-D64)	62	0.9	2.2
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	62	0.9	2.2
15	Chronic liver disease and cirrhosis (K70,K73-K74)	57	0.8	2.1
...	All other causes (Residual)	1,297	18.4	46.8
Black, both sexes, 40-44 years				
...	All causes	11,138	100.0	379.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,037	18.3	69.4
2	Malignant neoplasms (C00-C97)	1,781	16.0	60.6
3	Human immunodeficiency virus (HIV) disease (B20-B24)	1,386	12.4	47.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,207	10.8	41.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	635	5.7	21.6
6	Cerebrovascular diseases (I60-I69)	501	4.5	17.1
7	Diabetes mellitus (E10-E14)	317	2.8	10.8
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	220	2.0	7.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	215	1.9	7.3
10	Chronic liver disease and cirrhosis (K70,K73-K74)	182	1.6	6.2
11	Septicemia (A40-A41)	161	1.4	5.5
12	Chronic lower respiratory diseases (J40-J47)	148	1.3	5.0
13	Influenza and pneumonia (J10-J18)	145	1.3	4.9
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	135	1.2	4.6
15	Anemias (D50-D64)	85	0.8	2.9
...	All other causes (Residual)	1,983	17.8	67.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 45-49 years				
...	All causes	16,714	100.0	602.7
1	Malignant neoplasms (C00-C97)	3,581	21.4	129.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,563	21.3	128.5
3	Human immunodeficiency virus (HIV) disease (B20-B24)	1,420	8.5	51.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,374	8.2	49.5
5	Cerebrovascular diseases (I60-I69)	847	5.1	30.5
6	Diabetes mellitus (E10-E14)	575	3.4	20.7
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	478	2.9	17.2
8	Chronic liver disease and cirrhosis (K70,K73-K74)	395	2.4	14.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	335	2.0	12.1
10	Chronic lower respiratory diseases (J40-J47)	291	1.7	10.5
11	Septicemia (A40-A41)	277	1.7	10.0
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	237	1.4	8.5
13	Influenza and pneumonia (J10-J18)	220	1.3	7.9
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	169	1.0	6.1
15	Viral hepatitis (B15-B19)	131	0.8	4.7
...	All other causes (Residual)	2,821	16.9	101.7
Black, both sexes, 50-54 years				
...	All causes	21,035	100.0	914.9
1	Malignant neoplasms (C00-C97)	5,320	25.3	231.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,153	24.5	224.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,315	6.3	57.2
4	Cerebrovascular diseases (I60-I69)	1,157	5.5	50.3
5	Human immunodeficiency virus (HIV) disease (B20-B24)	1,100	5.2	47.8
6	Diabetes mellitus (E10-E14)	835	4.0	36.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	497	2.4	21.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	419	2.0	18.2
9	Chronic lower respiratory diseases (J40-J47)	412	2.0	17.9
10	Septicemia (A40-A41)	381	1.8	16.6
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	319	1.5	13.9
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	295	1.4	12.8
13	Influenza and pneumonia (J10-J18)	279	1.3	12.1
14	Viral hepatitis (B15-B19)	224	1.1	9.7
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	145	0.7	6.3
...	All other causes (Residual)	3,184	15.1	138.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 55-59 years				
...	All causes	22,656	100.0	1,254.6
1	Malignant neoplasms (C00-C97)	6,760	29.8	374.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,902	26.1	326.8
3	Cerebrovascular diseases (I60-I69)	1,181	5.2	65.4
4	Diabetes mellitus (E10-E14)	1,148	5.1	63.6
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	898	4.0	49.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	646	2.9	35.8
7	Human immunodeficiency virus (HIV) disease (B20-B24)	610	2.7	33.8
8	Chronic lower respiratory diseases (J40-J47)	549	2.4	30.4
9	Septicemia (A40-A41)	460	2.0	25.5
10	Chronic liver disease and cirrhosis (K70,K73-K74)	444	2.0	24.6
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	332	1.5	18.4
12	Influenza and pneumonia (J10-J18)	303	1.3	16.8
13	Viral hepatitis (B15-B19)	166	0.7	9.2
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	150	0.7	8.3
15	Aortic aneurysm and dissection (I71)	102	0.5	5.6
...	All other causes (Residual)	3,005	13.3	166.4
Black, both sexes, 60-64 years				
...	All causes	22,760	100.0	1,812.2
1	Malignant neoplasms (C00-C97)	7,200	31.6	573.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,213	27.3	494.7
3	Diabetes mellitus (E10-E14)	1,292	5.7	102.9
4	Cerebrovascular diseases (I60-I69)	1,249	5.5	99.5
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	700	3.1	55.7
6	Chronic lower respiratory diseases (J40-J47)	658	2.9	52.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	573	2.5	45.6
8	Septicemia (A40-A41)	482	2.1	38.4
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	359	1.6	28.6
10	Chronic liver disease and cirrhosis (K70,K73-K74)	319	1.4	25.4
11	Influenza and pneumonia (J10-J18)	308	1.4	24.5
12	Human immunodeficiency virus (HIV) disease (B20-B24)	301	1.3	24.0
13	Aortic aneurysm and dissection (I71)	111	0.5	8.8
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	98	0.4	7.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	94	0.4	7.5
...	All other causes (Residual)	2,803	12.3	223.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 65-69 years				
...	All causes	24,368	100.0	2,466.3
1	Malignant neoplasms (C00-C97)	7,733	31.7	782.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,543	26.9	662.2
3	Diabetes mellitus (E10-E14)	1,484	6.1	150.2
4	Cerebrovascular diseases (I60-I69)	1,434	5.9	145.1
5	Chronic lower respiratory diseases (J40-J47)	909	3.7	92.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	785	3.2	79.4
7	Septicemia (A40-A41)	551	2.3	55.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	469	1.9	47.5
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	441	1.8	44.6
10	Influenza and pneumonia (J10-J18)	387	1.6	39.2
11	Chronic liver disease and cirrhosis (K70,K73-K74)	198	0.8	20.0
12	Human immunodeficiency virus (HIV) disease (B20-B24)	164	0.7	16.6
13	Aortic aneurysm and dissection (I71)	131	0.5	13.3
14	Pneumonitis due to solids and liquids (J69)	103	0.4	10.4
15	Alzheimer's disease (G30)	102	0.4	10.3
...	All other causes (Residual)	2,934	12.0	296.9
Black, both sexes, 70-74 years				
...	All causes	27,350	100.0	3,514.3
1	Malignant neoplasms (C00-C97)	7,986	29.2	1,026.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,462	27.3	958.8
3	Cerebrovascular diseases (I60-I69)	1,801	6.6	231.4
4	Diabetes mellitus (E10-E14)	1,644	6.0	211.2
5	Chronic lower respiratory diseases (J40-J47)	1,111	4.1	142.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	971	3.6	124.8
7	Septicemia (A40-A41)	632	2.3	81.2
8	Influenza and pneumonia (J10-J18)	531	1.9	68.2
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	517	1.9	66.4
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	458	1.7	58.9
11	Alzheimer's disease (G30)	288	1.1	37.0
12	Chronic liver disease and cirrhosis (K70,K73-K74)	159	0.6	20.4
13	Pneumonitis due to solids and liquids (J69)	132	0.5	17.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	128	0.5	16.4
15	Aortic aneurysm and dissection (I71)	124	0.5	15.9
...	All other causes (Residual)	3,406	12.5	437.7

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 75-79 years				
...	All causes	30,903	100.0	5,161.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,949	29.0	1,494.7
2	Malignant neoplasms (C00-C97)	7,763	25.1	1,296.6
3	Cerebrovascular diseases (I60-I69)	2,229	7.2	372.3
4	Diabetes mellitus (E10-E14)	1,698	5.5	283.6
5	Chronic lower respiratory diseases (J40-J47)	1,322	4.3	220.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,053	3.4	175.9
7	Septicemia (A40-A41)	751	2.4	125.4
8	Influenza and pneumonia (J10-J18)	693	2.2	115.8
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	603	2.0	100.7
10	Alzheimer's disease (G30)	601	1.9	100.4
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	523	1.7	87.4
12	Pneumonitis due to solids and liquids (J69)	245	0.8	40.9
13	Aortic aneurysm and dissection (I71)	153	0.5	25.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	144	0.5	24.1
15	Parkinson's disease (G20-G21)	136	0.4	22.7
...	All other causes (Residual)	4,040	13.1	674.8
Black, both sexes, 80-84 years				
...	All causes	31,095	100.0	7,547.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,402	30.2	2,282.2
2	Malignant neoplasms (C00-C97)	6,532	21.0	1,585.5
3	Cerebrovascular diseases (I60-I69)	2,517	8.1	611.0
4	Diabetes mellitus (E10-E14)	1,553	5.0	377.0
5	Chronic lower respiratory diseases (J40-J47)	1,117	3.6	271.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,050	3.4	254.9
7	Alzheimer's disease (G30)	995	3.2	241.5
8	Septicemia (A40-A41)	839	2.7	203.7
9	Influenza and pneumonia (J10-J18)	821	2.6	199.3
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	631	2.0	153.2
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	457	1.5	110.9
12	Pneumonitis due to solids and liquids (J69)	285	0.9	69.2
13	Parkinson's disease (G20-G21)	206	0.7	50.0
14	Aortic aneurysm and dissection (I71)	159	0.5	38.6
15	Atherosclerosis (I70)	139	0.4	33.7
...	All other causes (Residual)	4,392	14.1	1,066.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 85-89 years				
...	All causes	24,013	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,573	31.5	---
2	Malignant neoplasms (C00-C97)	3,991	16.6	---
3	Cerebrovascular diseases (I60-I69)	2,054	8.6	---
4	Alzheimer's disease (G30)	1,104	4.6	---
5	Diabetes mellitus (E10-E14)	1,050	4.4	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	832	3.5	---
7	Influenza and pneumonia (J10-J18)	783	3.3	---
8	Chronic lower respiratory diseases (J40-J47)	767	3.2	---
9	Septicemia (A40-A41)	631	2.6	---
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	567	2.4	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	315	1.3	---
12	Pneumonitis due to solids and liquids (J69)	266	1.1	---
13	Atherosclerosis (I70)	148	0.6	---
14	Parkinson's disease (G20-G21)	129	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	116	0.5	---
...	All other causes (Residual)	3,687	15.4	---
Black, both sexes, 90-94 years				
...	All causes	16,002	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,365	33.5	---
2	Malignant neoplasms (C00-C97)	1,855	11.6	---
3	Cerebrovascular diseases (I60-I69)	1,408	8.8	---
4	Alzheimer's disease (G30)	922	5.8	---
5	Diabetes mellitus (E10-E14)	612	3.8	---
6	Influenza and pneumonia (J10-J18)	597	3.7	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	504	3.1	---
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	437	2.7	---
9	Chronic lower respiratory diseases (J40-J47)	412	2.6	---
10	Septicemia (A40-A41)	408	2.5	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	219	1.4	---
12	Pneumonitis due to solids and liquids (J69)	192	1.2	---
13	Atherosclerosis (I70)	122	0.8	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	58	0.4	---
14	Aortic aneurysm and dissection (I71)	58	0.4	---
...	All other causes (Residual)	2,833	17.7	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 95-99 years				
...	All causes	6,795	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,438	35.9	---
2	Malignant neoplasms (C00-C97)	583	8.6	---
3	Cerebrovascular diseases (I60-I69)	513	7.5	---
4	Alzheimer's disease (G30)	418	6.2	---
5	Influenza and pneumonia (J10-J18)	272	4.0	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	203	3.0	---
7	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	200	2.9	---
8	Diabetes mellitus (E10-E14)	186	2.7	---
9	Septicemia (A40-A41)	173	2.5	---
10	Chronic lower respiratory diseases (J40-J47)	159	2.3	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	97	1.4	---
12	Pneumonitis due to solids and liquids (J69)	91	1.3	---
13	Atherosclerosis (I70)	63	0.9	---
14	Nutritional deficiencies (E40-E64)	36	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	26	0.4	---
...	All other causes (Residual)	1,337	19.7	---
Black, both sexes, 100 years and over				
...	All causes	2,142	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	813	38.0	---
2	Cerebrovascular diseases (I60-I69)	166	7.7	---
3	Alzheimer's disease (G30)	136	6.3	---
4	Influenza and pneumonia (J10-J18)	111	5.2	---
5	Malignant neoplasms (C00-C97)	99	4.6	---
6	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	72	3.4	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	49	2.3	---
8	Septicemia (A40-A41)	48	2.2	---
9	Diabetes mellitus (E10-E14)	47	2.2	---
10	Chronic lower respiratory diseases (J40-J47)	34	1.6	---
11	Atherosclerosis (I70)	33	1.5	---
12	Pneumonitis due to solids and liquids (J69)	24	1.1	---
13	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	23	1.1	---
14	Nutritional deficiencies (E40-E64)	11	0.5	---
15	Parkinson's disease (G20-G21)	5	0.2	---
...	All other causes (Residual)	471	22.0	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, all ages				
...	All causes	149,108	100.0	799.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	36,343	24.4	194.8
2	Malignant neoplasms (C00-C97)	32,726	21.9	175.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,329	6.3	50.0
4	Cerebrovascular diseases (I60-I69)	7,519	5.0	40.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7,412	5.0	39.7
6	Diabetes mellitus (E10-E14)	5,730	3.8	30.7
7	Human immunodeficiency virus (HIV) disease (B20-B24)	4,684	3.1	25.1
8	Chronic lower respiratory diseases (J40-J47)	4,464	3.0	23.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,645	2.4	19.5
10	Certain conditions originating in the perinatal period (P00-P96)	2,828	1.9	15.2
11	Septicemia (A40-A41)	2,786	1.9	14.9
12	Influenza and pneumonia (J10-J18)	2,729	1.8	14.6
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,128	1.4	11.4
14	Chronic liver disease and cirrhosis (K70,K73-K74)	1,656	1.1	8.9
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,621	1.1	8.7
...	All other causes (Residual)	23,508	15.8	126.0
Black, male, 1-4 years				
...	All causes	623	100.0	46.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	184	29.5	13.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	88	14.1	6.6
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	48	7.7	3.6
4	Malignant neoplasms (C00-C97)	37	5.9	2.8
5	Chronic lower respiratory diseases (J40-J47)	22	3.5	1.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	19	3.0	*
7	Influenza and pneumonia (J10-J18)	17	2.7	*
8	Certain conditions originating in the perinatal period (P00-P96)	16	2.6	*
9	Septicemia (A40-A41)	13	2.1	*
10	Cerebrovascular diseases (I60-I69)	10	1.6	*
11	Anemias (D50-D64)	5	0.8	*
12	Meningitis (G00,G03)	3	0.5	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.5	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.3	*
14	Pneumonitis due to solids and liquids (J69)	2	0.3	*
...	All other causes (Residual)	154	24.7	11.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 5-9 years				
...	All causes	361	100.0	22.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	132	36.6	8.3
2	Malignant neoplasms (C00-C97)	47	13.0	2.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	25	6.9	1.6
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	18	5.0	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	14	3.9	*
6	Chronic lower respiratory diseases (J40-J47)	12	3.3	*
7	Septicemia (A40-A41)	9	2.5	*
7	Cerebrovascular diseases (I60-I69)	9	2.5	*
9	Influenza and pneumonia (J10-J18)	7	1.9	*
10	Anemias (D50-D64)	5	1.4	*
11	Meningitis (G00,G03)	3	0.8	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.6	*
13	Nutritional deficiencies (E40-E64)	1	0.3	*
13	Acute bronchitis and bronchiolitis (J20-J21)	1	0.3	*
13	Hernia (K40-K46)	1	0.3	*
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1	0.3	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.3	*
...	All other causes (Residual)	73	20.2	4.6
Black, male, 10-14 years				
...	All causes	548	100.0	30.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	182	33.2	10.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	72	13.1	4.1
3	Malignant neoplasms (C00-C97)	56	10.2	3.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	26	4.7	1.5
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	23	4.2	1.3
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	22	4.0	1.2
6	Chronic lower respiratory diseases (J40-J47)	22	4.0	1.2
8	Human immunodeficiency virus (HIV) disease (B20-B24)	6	1.1	*
8	Cerebrovascular diseases (I60-I69)	6	1.1	*
8	Influenza and pneumonia (J10-J18)	6	1.1	*
11	Septicemia (A40-A41)	5	0.9	*
12	Anemias (D50-D64)	4	0.7	*
13	Diabetes mellitus (E10-E14)	3	0.5	*
13	Meningitis (G00,G03)	3	0.5	*
13	Diseases of appendix (K35-K38)	3	0.5	*
...	All other causes (Residual)	109	19.9	6.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 15-19 years				
...	All causes	2,189	100.0	127.2
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,032	47.1	60.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	560	25.6	32.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	124	5.7	7.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	85	3.9	4.9
5	Malignant neoplasms (C00-C97)	58	2.6	3.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	36	1.6	2.1
7	Chronic lower respiratory diseases (J40-J47)	20	0.9	1.2
8	Anemias (D50-D64)	18	0.8	*
9	Diabetes mellitus (E10-E14)	17	0.8	*
10	Septicemia (A40-A41)	13	0.6	*
11	Legal intervention (Y35,Y89.0)	11	0.5	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	10	0.5	*
13	Influenza and pneumonia (J10-J18)	9	0.4	*
14	Cerebrovascular diseases (I60-I69)	8	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.2	*
...	All other causes (Residual)	184	8.4	10.7
Black, male, 20-24 years				
...	All causes	3,520	100.0	220.4
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,758	49.9	110.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	774	22.0	48.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	258	7.3	16.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	144	4.1	9.0
5	Malignant neoplasms (C00-C97)	97	2.8	6.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	47	1.3	2.9
7	Chronic lower respiratory diseases (J40-J47)	28	0.8	1.8
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	28	0.8	1.8
9	Anemias (D50-D64)	27	0.8	1.7
10	Diabetes mellitus (E10-E14)	22	0.6	1.4
11	Cerebrovascular diseases (I60-I69)	16	0.5	*
12	Influenza and pneumonia (J10-J18)	14	0.4	*
13	Legal intervention (Y35,Y89.0)	13	0.4	*
14	Septicemia (A40-A41)	11	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.2	*
...	All other causes (Residual)	276	7.8	17.3

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 25-29 years				
...	All causes	3,334	100.0	239.1
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,375	41.2	98.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	658	19.7	47.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	230	6.9	16.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	208	6.2	14.9
5	Human immunodeficiency virus (HIV) disease (B20-B24)	153	4.6	11.0
6	Malignant neoplasms (C00-C97)	130	3.9	9.3
7	Diabetes mellitus (E10-E14)	41	1.2	2.9
8	Cerebrovascular diseases (I60-I69)	36	1.1	2.6
9	Anemias (D50-D64)	34	1.0	2.4
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	29	0.9	2.1
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	24	0.7	1.7
12	Septicemia (A40-A41)	23	0.7	1.6
13	Chronic lower respiratory diseases (J40-J47)	21	0.6	1.5
14	Influenza and pneumonia (J10-J18)	16	0.5	*
15	Legal intervention (Y35,Y89.0)	12	0.4	*
...	All other causes (Residual)	344	10.3	24.7
Black, male, 30-34 years				
...	All causes	3,509	100.0	270.7
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	944	26.9	72.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	711	20.3	54.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	403	11.5	31.1
4	Human immunodeficiency virus (HIV) disease (B20-B24)	305	8.7	23.5
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	194	5.5	15.0
6	Malignant neoplasms (C00-C97)	191	5.4	14.7
7	Diabetes mellitus (E10-E14)	87	2.5	6.7
8	Cerebrovascular diseases (I60-I69)	53	1.5	4.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	37	1.1	2.9
10	Anemias (D50-D64)	35	1.0	2.7
11	Chronic lower respiratory diseases (J40-J47)	32	0.9	2.5
12	Influenza and pneumonia (J10-J18)	29	0.8	2.2
13	Septicemia (A40-A41)	27	0.8	2.1
13	Legal intervention (Y35,Y89.0)	27	0.8	2.1
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	21	0.6	1.6
...	All other causes (Residual)	413	11.8	31.9

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 35-39 years				
...	All causes	4,247	100.0	323.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	736	17.3	56.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	730	17.2	55.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	586	13.8	44.7
4	Human immunodeficiency virus (HIV) disease (B20-B24)	584	13.8	44.5
5	Malignant neoplasms (C00-C97)	285	6.7	21.7
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	158	3.7	12.0
7	Diabetes mellitus (E10-E14)	141	3.3	10.8
8	Cerebrovascular diseases (I60-I69)	93	2.2	7.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	48	1.1	3.7
10	Septicemia (A40-A41)	47	1.1	3.6
11	Influenza and pneumonia (J10-J18)	45	1.1	3.4
12	Aortic aneurysm and dissection (I71)	38	0.9	2.9
13	Anemias (D50-D64)	36	0.8	2.7
14	Chronic lower respiratory diseases (J40-J47)	35	0.8	2.7
15	Chronic liver disease and cirrhosis (K70,K73-K74)	33	0.8	2.5
...	All other causes (Residual)	652	15.4	49.7
Black, male, 40-44 years				
...	All causes	6,379	100.0	463.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,253	19.6	91.1
2	Human immunodeficiency virus (HIV) disease (B20-B24)	905	14.2	65.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	827	13.0	60.1
4	Malignant neoplasms (C00-C97)	688	10.8	50.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	505	7.9	36.7
6	Cerebrovascular diseases (I60-I69)	239	3.7	17.4
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	179	2.8	13.0
8	Diabetes mellitus (E10-E14)	169	2.6	12.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	117	1.8	8.5
10	Chronic liver disease and cirrhosis (K70,K73-K74)	111	1.7	8.1
11	Influenza and pneumonia (J10-J18)	84	1.3	6.1
12	Septicemia (A40-A41)	75	1.2	5.5
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	72	1.1	5.2
14	Chronic lower respiratory diseases (J40-J47)	68	1.1	4.9
15	Anemias (D50-D64)	43	0.7	3.1
15	Aortic aneurysm and dissection (I71)	43	0.7	3.1
...	All other causes (Residual)	1,001	15.7	72.8

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 45-49 years				
...	All causes	9,765	100.0	755.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,272	23.3	175.9
2	Malignant neoplasms (C00-C97)	1,640	16.8	127.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	978	10.0	75.7
4	Human immunodeficiency virus (HIV) disease (B20-B24)	966	9.9	74.8
5	Cerebrovascular diseases (I60-I69)	413	4.2	32.0
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	379	3.9	29.3
7	Diabetes mellitus (E10-E14)	335	3.4	25.9
8	Chronic liver disease and cirrhosis (K70,K73-K74)	269	2.8	20.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	205	2.1	15.9
10	Chronic lower respiratory diseases (J40-J47)	142	1.5	11.0
11	Septicemia (A40-A41)	141	1.4	10.9
12	Influenza and pneumonia (J10-J18)	135	1.4	10.5
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	131	1.3	10.1
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	127	1.3	9.8
15	Viral hepatitis (B15-B19)	87	0.9	6.7
...	All other causes (Residual)	1,545	15.8	119.6
Black, male, 50-54 years				
...	All causes	12,483	100.0	1,184.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,297	26.4	312.9
2	Malignant neoplasms (C00-C97)	2,725	21.8	258.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	958	7.7	90.9
4	Human immunodeficiency virus (HIV) disease (B20-B24)	794	6.4	75.4
5	Cerebrovascular diseases (I60-I69)	637	5.1	60.5
6	Diabetes mellitus (E10-E14)	445	3.6	42.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	334	2.7	31.7
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	254	2.0	24.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	238	1.9	22.6
10	Chronic lower respiratory diseases (J40-J47)	212	1.7	20.1
11	Septicemia (A40-A41)	205	1.6	19.5
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	183	1.5	17.4
13	Influenza and pneumonia (J10-J18)	167	1.3	15.9
14	Viral hepatitis (B15-B19)	153	1.2	14.5
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	109	0.9	10.3
...	All other causes (Residual)	1,772	14.2	168.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 55-59 years				
...	All causes	13,614	100.0	1,667.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,765	27.7	461.0
2	Malignant neoplasms (C00-C97)	3,754	27.6	459.7
3	Cerebrovascular diseases (I60-I69)	683	5.0	83.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	656	4.8	80.3
5	Diabetes mellitus (E10-E14)	637	4.7	78.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	472	3.5	57.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	374	2.7	45.8
8	Chronic liver disease and cirrhosis (K70,K73-K74)	313	2.3	38.3
9	Chronic lower respiratory diseases (J40-J47)	306	2.2	37.5
10	Septicemia (A40-A41)	249	1.8	30.5
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	199	1.5	24.4
12	Influenza and pneumonia (J10-J18)	194	1.4	23.8
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	125	0.9	15.3
14	Viral hepatitis (B15-B19)	117	0.9	14.3
15	Aortic aneurysm and dissection (I71)	72	0.5	8.8
...	All other causes (Residual)	1,698	12.5	207.9
Black, male, 60-64 years				
...	All causes	13,097	100.0	2,380.7
1	Malignant neoplasms (C00-C97)	4,013	30.6	729.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,740	28.6	679.8
3	Cerebrovascular diseases (I60-I69)	734	5.6	133.4
4	Diabetes mellitus (E10-E14)	650	5.0	118.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	438	3.3	79.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	376	2.9	68.3
7	Chronic lower respiratory diseases (J40-J47)	363	2.8	66.0
8	Septicemia (A40-A41)	239	1.8	43.4
9	Human immunodeficiency virus (HIV) disease (B20-B24)	224	1.7	40.7
10	Chronic liver disease and cirrhosis (K70,K73-K74)	223	1.7	40.5
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	201	1.5	36.5
12	Influenza and pneumonia (J10-J18)	183	1.4	33.3
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	71	0.5	12.9
14	Aortic aneurysm and dissection (I71)	66	0.5	12.0
15	Viral hepatitis (B15-B19)	58	0.4	10.5
...	All other causes (Residual)	1,518	11.6	275.9

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 65-69 years				
...	All causes	13,358	100.0	3,174.7
1	Malignant neoplasms (C00-C97)	4,312	32.3	1,024.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,693	27.6	877.7
3	Cerebrovascular diseases (I60-I69)	778	5.8	184.9
4	Diabetes mellitus (E10-E14)	681	5.1	161.8
5	Chronic lower respiratory diseases (J40-J47)	530	4.0	126.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	334	2.5	79.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	308	2.3	73.2
8	Septicemia (A40-A41)	287	2.1	68.2
9	Influenza and pneumonia (J10-J18)	226	1.7	53.7
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	222	1.7	52.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	130	1.0	30.9
12	Human immunodeficiency virus (HIV) disease (B20-B24)	125	0.9	29.7
13	Aortic aneurysm and dissection (I71)	79	0.6	18.8
14	Pneumonitis due to solids and liquids (J69)	65	0.5	15.4
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	53	0.4	12.6
...	All other causes (Residual)	1,535	11.5	364.8
Black, male, 70-74 years				
...	All causes	14,258	100.0	4,509.2
1	Malignant neoplasms (C00-C97)	4,412	30.9	1,395.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,983	27.9	1,259.7
3	Cerebrovascular diseases (I60-I69)	875	6.1	276.7
4	Diabetes mellitus (E10-E14)	749	5.3	236.9
5	Chronic lower respiratory diseases (J40-J47)	629	4.4	198.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	422	3.0	133.5
7	Septicemia (A40-A41)	307	2.2	97.1
8	Influenza and pneumonia (J10-J18)	286	2.0	90.5
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	286	2.0	90.5
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	250	1.8	79.1
11	Alzheimer's disease (G30)	114	0.8	36.1
12	Chronic liver disease and cirrhosis (K70,K73-K74)	100	0.7	31.6
13	Pneumonitis due to solids and liquids (J69)	81	0.6	25.6
14	Human immunodeficiency virus (HIV) disease (B20-B24)	59	0.4	18.7
14	Parkinson's disease (G20-G21)	59	0.4	18.7
...	All other causes (Residual)	1,646	11.5	520.6

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 75-79 years				
...	All causes	14,718	100.0	6,579.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,209	28.6	1,881.5
2	Malignant neoplasms (C00-C97)	4,034	27.4	1,803.3
3	Cerebrovascular diseases (I60-I69)	911	6.2	407.2
4	Chronic lower respiratory diseases (J40-J47)	773	5.3	345.5
5	Diabetes mellitus (E10-E14)	695	4.7	310.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	467	3.2	208.8
7	Influenza and pneumonia (J10-J18)	323	2.2	144.4
8	Septicemia (A40-A41)	321	2.2	143.5
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	292	2.0	130.5
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	277	1.9	123.8
11	Alzheimer's disease (G30)	232	1.6	103.7
12	Pneumonitis due to solids and liquids (J69)	139	0.9	62.1
13	Parkinson's disease (G20-G21)	81	0.6	36.2
14	Aortic aneurysm and dissection (I71)	76	0.5	34.0
15	Chronic liver disease and cirrhosis (K70,K73-K74)	73	0.5	32.6
...	All other causes (Residual)	1,815	12.3	811.3
Black, male, 80-84 years				
...	All causes	13,351	100.0	9,376.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,860	28.9	2,710.8
2	Malignant neoplasms (C00-C97)	3,352	25.1	2,354.0
3	Cerebrovascular diseases (I60-I69)	933	7.0	655.2
4	Chronic lower respiratory diseases (J40-J47)	602	4.5	422.8
5	Diabetes mellitus (E10-E14)	558	4.2	391.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	432	3.2	303.4
7	Influenza and pneumonia (J10-J18)	369	2.8	259.1
8	Septicemia (A40-A41)	340	2.5	238.8
9	Alzheimer's disease (G30)	317	2.4	222.6
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	238	1.8	167.1
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	220	1.6	154.5
12	Pneumonitis due to solids and liquids (J69)	151	1.1	106.0
13	Parkinson's disease (G20-G21)	117	0.9	82.2
14	Aortic aneurysm and dissection (I71)	67	0.5	47.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	61	0.5	42.8
...	All other causes (Residual)	1,734	13.0	1,217.8

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 85-89 years				
...	All causes	8,615	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,549	29.6	---
2	Malignant neoplasms (C00-C97)	1,864	21.6	---
3	Cerebrovascular diseases (I60-I69)	632	7.3	---
4	Chronic lower respiratory diseases (J40-J47)	391	4.5	---
5	Diabetes mellitus (E10-E14)	312	3.6	---
6	Alzheimer's disease (G30)	307	3.6	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	295	3.4	---
8	Influenza and pneumonia (J10-J18)	291	3.4	---
9	Septicemia (A40-A41)	252	2.9	---
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	169	2.0	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	134	1.6	---
12	Pneumonitis due to solids and liquids (J69)	118	1.4	---
13	Parkinson's disease (G20-G21)	69	0.8	---
14	Aortic aneurysm and dissection (I71)	39	0.5	---
15	Atherosclerosis (I70)	38	0.4	---
...	All other causes (Residual)	1,155	13.4	---
Black, male, 90-94 years				
...	All causes	4,417	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,403	31.8	---
2	Malignant neoplasms (C00-C97)	771	17.5	---
3	Cerebrovascular diseases (I60-I69)	321	7.3	---
4	Alzheimer's disease (G30)	188	4.3	---
5	Chronic lower respiratory diseases (J40-J47)	187	4.2	---
6	Influenza and pneumonia (J10-J18)	181	4.1	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	156	3.5	---
8	Diabetes mellitus (E10-E14)	144	3.3	---
9	Septicemia (A40-A41)	107	2.4	---
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	92	2.1	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	73	1.7	---
12	Pneumonitis due to solids and liquids (J69)	66	1.5	---
13	Atherosclerosis (I70)	29	0.7	---
14	Parkinson's disease (G20-G21)	26	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	0.3	---
15	Aortic aneurysm and dissection (I71)	14	0.3	---
...	All other causes (Residual)	645	14.6	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 95-99 years				
...	All causes	1,487	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	475	31.9	---
2	Malignant neoplasms (C00-C97)	221	14.9	---
3	Cerebrovascular diseases (I60-I69)	86	5.8	---
4	Influenza and pneumonia (J10-J18)	74	5.0	---
5	Chronic lower respiratory diseases (J40-J47)	56	3.8	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	55	3.7	---
7	Alzheimer's disease (G30)	50	3.4	---
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	41	2.8	---
9	Diabetes mellitus (E10-E14)	39	2.6	---
10	Septicemia (A40-A41)	37	2.5	---
11	Pneumonitis due to solids and liquids (J69)	29	2.0	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	29	2.0	---
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.6	---
13	Atherosclerosis (I70)	9	0.6	---
15	Parkinson's disease (G20-G21)	8	0.5	---
...	All other causes (Residual)	269	18.1	---
Black, male, 100 years and over				
...	All causes	337	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	124	36.8	---
2	Malignant neoplasms (C00-C97)	31	9.2	---
3	Influenza and pneumonia (J10-J18)	26	7.7	---
4	Cerebrovascular diseases (I60-I69)	19	5.6	---
5	Alzheimer's disease (G30)	13	3.9	---
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	3.9	---
7	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	9	2.7	---
8	Septicemia (A40-A41)	7	2.1	---
9	Chronic lower respiratory diseases (J40-J47)	6	1.8	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6	1.8	---
11	Diabetes mellitus (E10-E14)	5	1.5	---
11	Atherosclerosis (I70)	5	1.5	---
13	Pneumonitis due to solids and liquids (J69)	4	1.2	---
14	Hernia (K40-K46)	3	0.9	---
15	Nutritional deficiencies (E40-E64)	2	0.6	---
15	Parkinson's disease (G20-G21)	2	0.6	---
...	All other causes (Residual)	62	18.4	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, all ages				
...	All causes	143,700	100.0	703.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	37,816	26.3	185.2
2	Malignant neoplasms (C00-C97)	30,439	21.2	149.1
3	Cerebrovascular diseases (I60-I69)	10,022	7.0	49.1
4	Diabetes mellitus (E10-E14)	7,240	5.0	35.5
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,430	3.1	21.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,323	3.0	21.2
7	Chronic lower respiratory diseases (J40-J47)	3,765	2.6	18.4
8	Septicemia (A40-A41)	3,435	2.4	16.8
9	Alzheimer's disease (G30)	3,328	2.3	16.3
10	Influenza and pneumonia (J10-J18)	3,051	2.1	14.9
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,825	2.0	13.8
12	Human immunodeficiency virus (HIV) disease (B20-B24)	2,338	1.6	11.5
13	Certain conditions originating in the perinatal period (P00-P96)	2,222	1.5	10.9
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,257	0.9	6.2
15	Chronic liver disease and cirrhosis (K70,K73-K74)	874	0.6	4.3
...	All other causes (Residual)	26,335	18.3	129.0
Black, female, 1-4 years				
...	All causes	474	100.0	36.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	143	30.2	11.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	58	12.2	4.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	52	11.0	4.0
4	Malignant neoplasms (C00-C97)	27	5.7	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	26	5.5	2.0
6	Influenza and pneumonia (J10-J18)	12	2.5	*
7	Chronic lower respiratory diseases (J40-J47)	9	1.9	*
8	Septicemia (A40-A41)	7	1.5	*
8	Certain conditions originating in the perinatal period (P00-P96)	7	1.5	*
10	Anemias (D50-D64)	6	1.3	*
11	Cerebrovascular diseases (I60-I69)	4	0.8	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.6	*
12	Meningitis (G00,G03)	3	0.6	*
12	Pneumonitis due to solids and liquids (J69)	3	0.6	*
15	Acute bronchitis and bronchiolitis (J20-J21)	2	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
...	All other causes (Residual)	110	23.2	8.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 5-9 years				
...	All causes	302	100.0	19.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	104	34.4	6.7
2	Malignant neoplasms (C00-C97)	35	11.6	2.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	23	7.6	1.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	21	7.0	1.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	20	6.6	1.3
6	Chronic lower respiratory diseases (J40-J47)	11	3.6	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	1.7	*
7	Cerebrovascular diseases (I60-I69)	5	1.7	*
9	Influenza and pneumonia (J10-J18)	4	1.3	*
10	Anemias (D50-D64)	3	1.0	*
11	Septicemia (A40-A41)	2	0.7	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.7	*
11	Pneumonitis due to solids and liquids (J69)	2	0.7	*
11	Certain conditions originating in the perinatal period (P00-P96)	2	0.7	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
...	All other causes (Residual)	62	20.5	4.0
Black, female, 10-14 years				
...	All causes	332	100.0	19.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	83	25.0	4.8
2	Malignant neoplasms (C00-C97)	34	10.2	2.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	33	9.9	1.9
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	24	7.2	1.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	20	6.0	1.2
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	14	4.2	*
7	Diabetes mellitus (E10-E14)	9	2.7	*
8	Chronic lower respiratory diseases (J40-J47)	8	2.4	*
9	Septicemia (A40-A41)	5	1.5	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	1.5	*
9	Influenza and pneumonia (J10-J18)	5	1.5	*
12	Cerebrovascular diseases (I60-I69)	4	1.2	*
13	Anemias (D50-D64)	3	0.9	*
14	Syphilis (A50-A53)	1	0.3	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.3	*
14	Meningitis (G00,G03)	1	0.3	*
14	Diseases of appendix (K35-K38)	1	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
14	Certain conditions originating in the perinatal period (P00-P96)	1	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.3	*
...	All other causes (Residual)	78	23.5	4.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 15-19 years				
...	All causes	650	100.0	38.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	213	32.8	12.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	106	16.3	6.3
3	Malignant neoplasms (C00-C97)	45	6.9	2.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	31	4.8	1.9
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	24	3.7	1.4
6	Pregnancy, childbirth and the puerperium (O00-099)	20	3.1	1.2
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	19	2.9	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	12	1.8	*
8	Anemias (D50-D64)	12	1.8	*
10	Cerebrovascular diseases (I60-I69)	11	1.7	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	1.2	*
12	Septicemia (A40-A41)	7	1.1	*
12	Diabetes mellitus (E10-E14)	7	1.1	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.8	*
14	Chronic lower respiratory diseases (J40-J47)	5	0.8	*
...	All other causes (Residual)	125	19.2	7.5
Black, female, 20-24 years				
...	All causes	1,012	100.0	64.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	233	23.0	14.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	181	17.9	11.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	74	7.3	4.7
4	Malignant neoplasms (C00-C97)	64	6.3	4.1
5	Pregnancy, childbirth and the puerperium (O00-099)	49	4.8	3.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	40	4.0	2.5
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	31	3.1	2.0
8	Diabetes mellitus (E10-E14)	27	2.7	1.7
9	Chronic lower respiratory diseases (J40-J47)	22	2.2	1.4
10	Anemias (D50-D64)	20	2.0	1.3
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	17	1.7	*
12	Cerebrovascular diseases (I60-I69)	16	1.6	*
13	Septicemia (A40-A41)	13	1.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	1.0	*
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	4	0.4	*
15	Influenza and pneumonia (J10-J18)	4	0.4	*
...	All other causes (Residual)	207	20.5	13.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 25-29 years				
...	All causes	1,313	100.0	89.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	186	14.2	12.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	146	11.1	9.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	145	11.0	9.8
4	Malignant neoplasms (C00-C97)	133	10.1	9.0
5	Human immunodeficiency virus (HIV) disease (B20-B24)	125	9.5	8.5
6	Pregnancy, childbirth and the puerperium (O00-099)	66	5.0	4.5
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	44	3.4	3.0
8	Diabetes mellitus (E10-E14)	39	3.0	2.6
9	Anemias (D50-D64)	31	2.4	2.1
10	Cerebrovascular diseases (I60-I69)	28	2.1	1.9
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	1.8	1.6
12	Septicemia (A40-A41)	21	1.6	1.4
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	17	1.3	*
14	Chronic lower respiratory diseases (J40-J47)	13	1.0	*
15	Influenza and pneumonia (J10-J18)	12	0.9	*
...	All other causes (Residual)	284	21.6	19.3
Black, female, 30-34 years				
...	All causes	1,880	100.0	131.0
1	Human immunodeficiency virus (HIV) disease (B20-B24)	253	13.5	17.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	244	13.0	17.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	241	12.8	16.8
4	Malignant neoplasms (C00-C97)	234	12.4	16.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	126	6.7	8.8
6	Diabetes mellitus (E10-E14)	52	2.8	3.6
7	Pregnancy, childbirth and the puerperium (O00-099)	51	2.7	3.6
8	Cerebrovascular diseases (I60-I69)	46	2.4	3.2
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	39	2.1	2.7
10	Septicemia (A40-A41)	37	2.0	2.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	37	2.0	2.6
12	Anemias (D50-D64)	31	1.6	2.2
12	Chronic lower respiratory diseases (J40-J47)	31	1.6	2.2
14	Influenza and pneumonia (J10-J18)	21	1.1	1.5
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	21	1.1	1.5
...	All other causes (Residual)	416	22.1	29.0

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 35-39 years				
...	All causes	2,801	100.0	191.5
1	Malignant neoplasms (C00-C97)	494	17.6	33.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	416	14.9	28.4
3	Human immunodeficiency virus (HIV) disease (B20-B24)	367	13.1	25.1
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	271	9.7	18.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	148	5.3	10.1
6	Cerebrovascular diseases (I60-I69)	95	3.4	6.5
7	Diabetes mellitus (E10-E14)	83	3.0	5.7
8	Septicemia (A40-A41)	54	1.9	3.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	50	1.8	3.4
10	Pregnancy, childbirth and the puerperium (O00-099)	41	1.5	2.8
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	41	1.5	2.8
12	Chronic lower respiratory diseases (J40-J47)	36	1.3	2.5
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	30	1.1	2.1
14	Influenza and pneumonia (J10-J18)	27	1.0	1.8
15	Anemias (D50-D64)	26	0.9	1.8
...	All other causes (Residual)	622	22.2	42.5
Black, female, 40-44 years				
...	All causes	4,759	100.0	304.8
1	Malignant neoplasms (C00-C97)	1,093	23.0	70.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	784	16.5	50.2
3	Human immunodeficiency virus (HIV) disease (B20-B24)	481	10.1	30.8
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	380	8.0	24.3
5	Cerebrovascular diseases (I60-I69)	262	5.5	16.8
6	Diabetes mellitus (E10-E14)	148	3.1	9.5
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	130	2.7	8.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	98	2.1	6.3
9	Septicemia (A40-A41)	86	1.8	5.5
10	Chronic lower respiratory diseases (J40-J47)	80	1.7	5.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	71	1.5	4.5
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	63	1.3	4.0
13	Influenza and pneumonia (J10-J18)	61	1.3	3.9
14	Anemias (D50-D64)	42	0.9	2.7
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	41	0.9	2.6
...	All other causes (Residual)	939	19.7	60.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 45-49 years				
...	All causes	6,949	100.0	469.0
1	Malignant neoplasms (C00-C97)	1,941	27.9	131.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,291	18.6	87.1
3	Human immunodeficiency virus (HIV) disease (B20-B24)	454	6.5	30.6
4	Cerebrovascular diseases (I60-I69)	434	6.2	29.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	396	5.7	26.7
6	Diabetes mellitus (E10-E14)	240	3.5	16.2
7	Chronic lower respiratory diseases (J40-J47)	149	2.1	10.1
8	Septicemia (A40-A41)	136	2.0	9.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	130	1.9	8.8
10	Chronic liver disease and cirrhosis (K70,K73-K74)	126	1.8	8.5
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	110	1.6	7.4
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	99	1.4	6.7
13	Influenza and pneumonia (J10-J18)	85	1.2	5.7
14	Viral hepatitis (B15-B19)	44	0.6	3.0
15	Anemias (D50-D64)	38	0.5	2.6
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	38	0.5	2.6
...	All other causes (Residual)	1,238	17.8	83.6
Black, female, 50-54 years				
...	All causes	8,552	100.0	686.6
1	Malignant neoplasms (C00-C97)	2,595	30.3	208.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,856	21.7	149.0
3	Cerebrovascular diseases (I60-I69)	520	6.1	41.8
4	Diabetes mellitus (E10-E14)	390	4.6	31.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	357	4.2	28.7
6	Human immunodeficiency virus (HIV) disease (B20-B24)	306	3.6	24.6
7	Chronic lower respiratory diseases (J40-J47)	200	2.3	16.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	181	2.1	14.5
9	Septicemia (A40-A41)	176	2.1	14.1
10	Chronic liver disease and cirrhosis (K70,K73-K74)	163	1.9	13.1
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	112	1.3	9.0
11	Influenza and pneumonia (J10-J18)	112	1.3	9.0
13	Viral hepatitis (B15-B19)	71	0.8	5.7
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	65	0.8	5.2
15	Anemias (D50-D64)	40	0.5	3.2
...	All other causes (Residual)	1,408	16.5	113.0

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 55-59 years				
...	All causes	9,042	100.0	914.1
1	Malignant neoplasms (C00-C97)	3,006	33.2	303.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,137	23.6	216.0
3	Diabetes mellitus (E10-E14)	511	5.7	51.7
4	Cerebrovascular diseases (I60-I69)	498	5.5	50.3
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	272	3.0	27.5
6	Chronic lower respiratory diseases (J40-J47)	243	2.7	24.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	242	2.7	24.5
8	Septicemia (A40-A41)	211	2.3	21.3
9	Human immunodeficiency virus (HIV) disease (B20-B24)	138	1.5	14.0
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	133	1.5	13.4
11	Chronic liver disease and cirrhosis (K70,K73-K74)	131	1.4	13.2
12	Influenza and pneumonia (J10-J18)	109	1.2	11.0
13	Viral hepatitis (B15-B19)	49	0.5	5.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	36	0.4	3.6
15	Aortic aneurysm and dissection (I71)	30	0.3	3.0
...	All other causes (Residual)	1,296	14.3	131.0
Black, female, 60-64 years				
...	All causes	9,663	100.0	1,369.2
1	Malignant neoplasms (C00-C97)	3,187	33.0	451.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,473	25.6	350.4
3	Diabetes mellitus (E10-E14)	642	6.6	91.0
4	Cerebrovascular diseases (I60-I69)	515	5.3	73.0
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	324	3.4	45.9
6	Chronic lower respiratory diseases (J40-J47)	295	3.1	41.8
7	Septicemia (A40-A41)	243	2.5	34.4
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	158	1.6	22.4
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	135	1.4	19.1
10	Influenza and pneumonia (J10-J18)	125	1.3	17.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	96	1.0	13.6
12	Human immunodeficiency virus (HIV) disease (B20-B24)	77	0.8	10.9
13	Aortic aneurysm and dissection (I71)	45	0.5	6.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	44	0.5	6.2
15	Viral hepatitis (B15-B19)	33	0.3	4.7
...	All other causes (Residual)	1,271	13.2	180.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 65-69 years				
...	All causes	11,010	100.0	1,940.8
1	Malignant neoplasms (C00-C97)	3,421	31.1	603.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,850	25.9	502.4
3	Diabetes mellitus (E10-E14)	803	7.3	141.5
4	Cerebrovascular diseases (I60-I69)	656	6.0	115.6
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	451	4.1	79.5
6	Chronic lower respiratory diseases (J40-J47)	379	3.4	66.8
7	Septicemia (A40-A41)	264	2.4	46.5
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	219	2.0	38.6
9	Influenza and pneumonia (J10-J18)	161	1.5	28.4
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	161	1.5	28.4
11	Chronic liver disease and cirrhosis (K70,K73-K74)	68	0.6	12.0
12	Alzheimer's disease (G30)	53	0.5	9.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	52	0.5	9.2
13	Aortic aneurysm and dissection (I71)	52	0.5	9.2
15	Human immunodeficiency virus (HIV) disease (B20-B24)	39	0.4	6.9
...	All other causes (Residual)	1,381	12.5	243.4
Black, female, 70-74 years				
...	All causes	13,092	100.0	2,833.5
1	Malignant neoplasms (C00-C97)	3,574	27.3	773.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,479	26.6	753.0
3	Cerebrovascular diseases (I60-I69)	926	7.1	200.4
4	Diabetes mellitus (E10-E14)	895	6.8	193.7
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	549	4.2	118.8
6	Chronic lower respiratory diseases (J40-J47)	482	3.7	104.3
7	Septicemia (A40-A41)	325	2.5	70.3
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	267	2.0	57.8
9	Influenza and pneumonia (J10-J18)	245	1.9	53.0
10	Alzheimer's disease (G30)	174	1.3	37.7
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	172	1.3	37.2
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	71	0.5	15.4
13	Aortic aneurysm and dissection (I71)	70	0.5	15.2
14	Chronic liver disease and cirrhosis (K70,K73-K74)	59	0.5	12.8
15	Pneumonitis due to solids and liquids (J69)	51	0.4	11.0
...	All other causes (Residual)	1,753	13.4	379.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 75-79 years				
...	All causes	16,185	100.0	4,316.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,740	29.3	1,264.0
2	Malignant neoplasms (C00-C97)	3,729	23.0	994.4
3	Cerebrovascular diseases (I60-I69)	1,318	8.1	351.5
4	Diabetes mellitus (E10-E14)	1,003	6.2	267.5
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	586	3.6	156.3
6	Chronic lower respiratory diseases (J40-J47)	549	3.4	146.4
7	Septicemia (A40-A41)	430	2.7	114.7
8	Influenza and pneumonia (J10-J18)	370	2.3	98.7
9	Alzheimer's disease (G30)	369	2.3	98.4
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	326	2.0	86.9
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	231	1.4	61.6
12	Pneumonitis due to solids and liquids (J69)	106	0.7	28.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	81	0.5	21.6
14	Aortic aneurysm and dissection (I71)	77	0.5	20.5
15	Atherosclerosis (I70)	61	0.4	16.3
...	All other causes (Residual)	2,209	13.6	589.1
Black, female, 80-84 years				
...	All causes	17,744	100.0	6,581.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,542	31.2	2,055.7
2	Malignant neoplasms (C00-C97)	3,180	17.9	1,179.6
3	Cerebrovascular diseases (I60-I69)	1,584	8.9	587.6
4	Diabetes mellitus (E10-E14)	995	5.6	369.1
5	Alzheimer's disease (G30)	678	3.8	251.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	618	3.5	229.2
7	Chronic lower respiratory diseases (J40-J47)	515	2.9	191.0
8	Septicemia (A40-A41)	499	2.8	185.1
9	Influenza and pneumonia (J10-J18)	452	2.5	167.7
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	411	2.3	152.5
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	219	1.2	81.2
12	Pneumonitis due to solids and liquids (J69)	134	0.8	49.7
13	Aortic aneurysm and dissection (I71)	92	0.5	34.1
14	Parkinson's disease (G20-G21)	89	0.5	33.0
15	Atherosclerosis (I70)	82	0.5	30.4
...	All other causes (Residual)	2,654	15.0	984.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 85-89 years				
...	All causes	15,398	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,024	32.6	---
2	Malignant neoplasms (C00-C97)	2,127	13.8	---
3	Cerebrovascular diseases (I60-I69)	1,422	9.2	---
4	Alzheimer's disease (G30)	797	5.2	---
5	Diabetes mellitus (E10-E14)	738	4.8	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	537	3.5	---
7	Influenza and pneumonia (J10-J18)	492	3.2	---
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	398	2.6	---
9	Septicemia (A40-A41)	379	2.5	---
10	Chronic lower respiratory diseases (J40-J47)	376	2.4	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	181	1.2	---
12	Pneumonitis due to solids and liquids (J69)	148	1.0	---
13	Atherosclerosis (I70)	110	0.7	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	80	0.5	---
15	Aortic aneurysm and dissection (I71)	71	0.5	---
...	All other causes (Residual)	2,518	16.4	---
Black, female, 90-94 years				
...	All causes	11,585	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,962	34.2	---
2	Cerebrovascular diseases (I60-I69)	1,087	9.4	---
3	Malignant neoplasms (C00-C97)	1,084	9.4	---
4	Alzheimer's disease (G30)	734	6.3	---
5	Diabetes mellitus (E10-E14)	468	4.0	---
6	Influenza and pneumonia (J10-J18)	416	3.6	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	348	3.0	---
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	345	3.0	---
9	Septicemia (A40-A41)	301	2.6	---
10	Chronic lower respiratory diseases (J40-J47)	225	1.9	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	146	1.3	---
12	Pneumonitis due to solids and liquids (J69)	126	1.1	---
13	Atherosclerosis (I70)	93	0.8	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	44	0.4	---
14	Aortic aneurysm and dissection (I71)	44	0.4	---
...	All other causes (Residual)	2,162	18.7	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 95-99 years				
...	All causes	5,308	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,963	37.0	---
2	Cerebrovascular diseases (I60-I69)	427	8.0	---
3	Alzheimer's disease (G30)	368	6.9	---
4	Malignant neoplasms (C00-C97)	362	6.8	---
5	Influenza and pneumonia (J10-J18)	198	3.7	---
6	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	159	3.0	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	148	2.8	---
8	Diabetes mellitus (E10-E14)	147	2.8	---
9	Septicemia (A40-A41)	136	2.6	---
10	Chronic lower respiratory diseases (J40-J47)	103	1.9	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	68	1.3	---
12	Pneumonitis due to solids and liquids (J69)	62	1.2	---
13	Atherosclerosis (I70)	54	1.0	---
14	Nutritional deficiencies (E40-E64)	29	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	0.3	---
15	Anemias (D50-D64)	17	0.3	---
15	Aortic aneurysm and dissection (I71)	17	0.3	---
...	All other causes (Residual)	1,033	19.5	---
Black, female, 100 years and over				
...	All causes	1,805	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	689	38.2	---
2	Cerebrovascular diseases (I60-I69)	147	8.1	---
3	Alzheimer's disease (G30)	123	6.8	---
4	Influenza and pneumonia (J10-J18)	85	4.7	---
5	Malignant neoplasms (C00-C97)	68	3.8	---
6	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	63	3.5	---
7	Diabetes mellitus (E10-E14)	42	2.3	---
8	Septicemia (A40-A41)	41	2.3	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	36	2.0	---
10	Atherosclerosis (I70)	28	1.6	---
10	Chronic lower respiratory diseases (J40-J47)	28	1.6	---
12	Pneumonitis due to solids and liquids (J69)	20	1.1	---
13	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17	0.9	---
14	Nutritional deficiencies (E40-E64)	9	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.2	---
15	Anemias (D50-D64)	4	0.2	---
...	All other causes (Residual)	401	22.2	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, all ages				
...	All causes	13,918	100.0	440.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,659	19.1	84.1
2	Malignant neoplasms (C00-C97)	2,465	17.7	78.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,626	11.7	51.4
4	Diabetes mellitus (E10-E14)	818	5.9	25.9
5	Cerebrovascular diseases (I60-I69)	627	4.5	19.8
6	Chronic liver disease and cirrhosis (K70,K73-K74)	596	4.3	18.9
7	Chronic lower respiratory diseases (J40-J47)	518	3.7	16.4
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	392	2.8	12.4
9	Influenza and pneumonia (J10-J18)	354	2.5	11.2
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	265	1.9	8.4
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	257	1.8	8.1
12	Septicemia (A40-A41)	217	1.6	6.9
13	Alzheimer's disease (G30)	173	1.2	5.5
14	Certain conditions originating in the perinatal period (P00-P96)	130	0.9	4.1
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	117	0.8	3.7
...	All other causes (Residual)	2,704	19.4	85.5
American Indian, both sexes, 1-4 years				
...	All causes	100	100.0	59.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	44	44.0	26.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	11	11.0	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	10	10.0	*
4	Malignant neoplasms (C00-C97)	4	4.0	*
5	Meningitis (G00,G03)	3	3.0	*
6	Septicemia (A40-A41)	1	1.0	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	1.0	*
6	Influenza and pneumonia (J10-J18)	1	1.0	*
6	Chronic lower respiratory diseases (J40-J47)	1	1.0	*
...	All other causes (Residual)	24	24.0	14.2
American Indian, both sexes, 5-9 years				
...	All causes	46	100.0	17.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	27	58.7	10.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	10.9	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	8.7	*
4	Malignant neoplasms (C00-C97)	3	6.5	*
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	2.2	*
...	All other causes (Residual)	6	13.0	*

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 10-14 years				
...	All causes	64	100.0	22.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	28	43.8	9.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	14	21.9	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	4.7	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	3.1	*
4	Chronic lower respiratory diseases (J40-J47)	2	3.1	*
6	Human immunodeficiency virus (HIV) disease (B20-B24)	1	1.6	*
6	Malignant neoplasms (C00-C97)	1	1.6	*
6	Meningitis (G00,G03)	1	1.6	*
6	Influenza and pneumonia (J10-J18)	1	1.6	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	1.6	*
...	All other causes (Residual)	10	15.6	*
American Indian, both sexes, 15-19 years				
...	All causes	282	100.0	93.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	142	50.4	47.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	59	20.9	19.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	29	10.3	9.6
4	Malignant neoplasms (C00-C97)	10	3.5	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	1.4	*
6	Septicemia (A40-A41)	2	0.7	*
6	Diabetes mellitus (E10-E14)	2	0.7	*
8	Salmonella infections (A01-A02)	1	0.4	*
8	Cerebrovascular diseases (I60-I69)	1	0.4	*
8	Influenza and pneumonia (J10-J18)	1	0.4	*
8	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.4	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.4	*
8	Pregnancy, childbirth and the puerperium (O00-099)	1	0.4	*
...	All other causes (Residual)	28	9.9	9.3
American Indian, both sexes, 20-24 years				
...	All causes	353	100.0	121.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	179	50.7	61.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	69	19.5	23.8
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	53	15.0	18.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	9	2.5	*
5	Malignant neoplasms (C00-C97)	5	1.4	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.6	*
6	Diabetes mellitus (E10-E14)	2	0.6	*
6	Chronic liver disease and cirrhosis (K70,K73-K74)	2	0.6	*
9	Septicemia (A40-A41)	1	0.3	*
9	Anemias (D50-D64)	1	0.3	*
9	Cerebrovascular diseases (I60-I69)	1	0.3	*
9	Pregnancy, childbirth and the puerperium (O00-099)	1	0.3	*
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.3	*
...	All other causes (Residual)	27	7.6	9.3

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 25-29 years				
...	All causes	364	100.0	145.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	175	48.1	69.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	49	13.5	19.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	37	10.2	14.8
4	Malignant neoplasms (C00-C97)	17	4.7	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	13	3.6	*
6	Chronic liver disease and cirrhosis (K70,K73-K74)	11	3.0	*
7	Human immunodeficiency virus (HIV) disease (B20-B24)	5	1.4	*
8	Diabetes mellitus (E10-E14)	4	1.1	*
9	Influenza and pneumonia (J10-J18)	3	0.8	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.8	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	0.5	*
11	Legal intervention (Y35,Y89.0)	2	0.5	*
13	Septicemia (A40-A41)	1	0.3	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.3	*
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1	0.3	*
13	Cerebrovascular diseases (I60-I69)	1	0.3	*
13	Aortic aneurysm and dissection (I71)	1	0.3	*
13	Chronic lower respiratory diseases (J40-J47)	1	0.3	*
13	Pregnancy, childbirth and the puerperium (O00-099)	1	0.3	*
...	All other causes (Residual)	36	9.9	14.4
American Indian, both sexes, 30-34 years				
...	All causes	356	100.0	156.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	149	41.9	65.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	42	11.8	18.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	33	9.3	14.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	25	7.0	11.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	23	6.5	10.1
6	Diabetes mellitus (E10-E14)	13	3.7	*
7	Malignant neoplasms (C00-C97)	12	3.4	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	5	1.4	*
9	Septicemia (A40-A41)	3	0.8	*
9	Cerebrovascular diseases (I60-I69)	3	0.8	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.8	*
12	Influenza and pneumonia (J10-J18)	2	0.6	*
12	Chronic lower respiratory diseases (J40-J47)	2	0.6	*
14	Tuberculosis (A16-A19)	1	0.3	*
14	Pneumonitis due to solids and liquids (J69)	1	0.3	*
14	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.3	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.3	*
...	All other causes (Residual)	37	10.4	16.3

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 35-39 years				
...	All causes	497	100.0	222.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	147	29.6	65.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	43	8.7	19.3
3	Chronic liver disease and cirrhosis (K70,K73-K74)	41	8.2	18.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	39	7.8	17.5
5	Malignant neoplasms (C00-C97)	35	7.0	15.7
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	21	4.2	9.4
7	Human immunodeficiency virus (HIV) disease (B20-B24)	15	3.0	*
7	Diabetes mellitus (E10-E14)	15	3.0	*
9	Cerebrovascular diseases (I60-I69)	13	2.6	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	2.0	*
11	Influenza and pneumonia (J10-J18)	7	1.4	*
12	Septicemia (A40-A41)	6	1.2	*
13	Chronic lower respiratory diseases (J40-J47)	5	1.0	*
14	Viral hepatitis (B15-B19)	4	0.8	*
15	Pregnancy, childbirth and the puerperium (O00-099)	3	0.6	*
15	Legal intervention (Y35,Y89.0)	3	0.6	*
...	All other causes (Residual)	90	18.1	40.3
American Indian, both sexes, 40-44 years				
...	All causes	726	100.0	305.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	158	21.8	66.5
2	Chronic liver disease and cirrhosis (K70,K73-K74)	84	11.6	35.4
3	Malignant neoplasms (C00-C97)	81	11.2	34.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	77	10.6	32.4
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	45	6.2	18.9
6	Diabetes mellitus (E10-E14)	26	3.6	10.9
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	24	3.3	10.1
8	Human immunodeficiency virus (HIV) disease (B20-B24)	20	2.8	8.4
9	Cerebrovascular diseases (I60-I69)	19	2.6	*
10	Influenza and pneumonia (J10-J18)	11	1.5	*
11	Septicemia (A40-A41)	10	1.4	*
12	Chronic lower respiratory diseases (J40-J47)	7	1.0	*
13	Viral hepatitis (B15-B19)	6	0.8	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	0.8	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.7	*
...	All other causes (Residual)	147	20.2	61.9

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 45-49 years				
...	All causes	874	100.0	394.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	150	17.2	67.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	147	16.8	66.3
3	Malignant neoplasms (C00-C97)	137	15.7	61.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	97	11.1	43.7
5	Diabetes mellitus (E10-E14)	36	4.1	16.2
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	31	3.5	14.0
7	Cerebrovascular diseases (I60-I69)	21	2.4	9.5
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	17	1.9	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	15	1.7	*
10	Influenza and pneumonia (J10-J18)	14	1.6	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	1.6	*
12	Septicemia (A40-A41)	10	1.1	*
	Chronic lower respiratory diseases (J40-J47)	10	1.1	*
14	Viral hepatitis (B15-B19)	9	1.0	*
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	3	0.3	*
15	Pneumonitis due to solids and liquids (J69)	3	0.3	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.3	*
...	All other causes (Residual)	157	18.0	70.8
American Indian, both sexes, 50-54 years				
...	All causes	1,053	100.0	570.2
1	Malignant neoplasms (C00-C97)	203	19.3	109.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	183	17.4	99.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	108	10.3	58.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	99	9.4	53.6
5	Diabetes mellitus (E10-E14)	62	5.9	33.6
6	Cerebrovascular diseases (I60-I69)	43	4.1	23.3
7	Chronic lower respiratory diseases (J40-J47)	23	2.2	12.5
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	22	2.1	11.9
9	Septicemia (A40-A41)	14	1.3	*
9	Viral hepatitis (B15-B19)	14	1.3	*
11	Influenza and pneumonia (J10-J18)	13	1.2	*
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	12	1.1	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	11	1.0	*
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	7	0.7	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.6	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.6	*
...	All other causes (Residual)	227	21.6	122.9

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 55-59 years				
...	All causes	1,075	100.0	729.8
1	Malignant neoplasms (C00-C97)	256	23.8	173.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	237	22.0	160.9
3	Diabetes mellitus (E10-E14)	95	8.8	64.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	72	6.7	48.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	72	6.7	48.9
6	Cerebrovascular diseases (I60-I69)	41	3.8	27.8
7	Chronic lower respiratory diseases (J40-J47)	32	3.0	21.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	2.1	15.6
9	Septicemia (A40-A41)	22	2.0	14.9
10	Influenza and pneumonia (J10-J18)	19	1.8	*
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	10	0.9	*
12	Viral hepatitis (B15-B19)	9	0.8	*
13	Atherosclerosis (I70)	5	0.5	*
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	0.5	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.4	*
...	All other causes (Residual)	173	16.1	117.4
American Indian, both sexes, 60-64 years				
...	All causes	1,165	100.0	1,158.8
1	Malignant neoplasms (C00-C97)	324	27.8	322.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	248	21.3	246.7
3	Diabetes mellitus (E10-E14)	82	7.0	81.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	71	6.1	70.6
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	63	5.4	62.7
6	Chronic lower respiratory diseases (J40-J47)	59	5.1	58.7
7	Cerebrovascular diseases (I60-I69)	48	4.1	47.7
8	Influenza and pneumonia (J10-J18)	24	2.1	23.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	24	2.1	23.9
10	Septicemia (A40-A41)	23	2.0	22.9
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	6	0.5	*
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	0.5	*
13	Viral hepatitis (B15-B19)	5	0.4	*
13	Atherosclerosis (I70)	5	0.4	*
13	Pneumonitis due to solids and liquids (J69)	5	0.4	*
...	All other causes (Residual)	172	14.8	171.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 65-69 years				
...	All causes	1,202	100.0	1,662.7
1	Malignant neoplasms (C00-C97)	334	27.8	462.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	260	21.6	359.7
3	Diabetes mellitus (E10-E14)	106	8.8	146.6
4	Cerebrovascular diseases (I60-I69)	64	5.3	88.5
5	Chronic lower respiratory diseases (J40-J47)	63	5.2	87.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	37	3.1	51.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	31	2.6	42.9
8	Chronic liver disease and cirrhosis (K70,K73-K74)	30	2.5	41.5
9	Septicemia (A40-A41)	26	2.2	36.0
10	Influenza and pneumonia (J10-J18)	23	1.9	31.8
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	11	0.9	*
12	Aortic aneurysm and dissection (I71)	8	0.7	*
13	Pneumonitis due to solids and liquids (J69)	6	0.5	*
14	Viral hepatitis (B15-B19)	4	0.3	*
14	Parkinson's disease (G20-G21)	4	0.3	*
14	Alzheimer's disease (G30)	4	0.3	*
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4	0.3	*
...	All other causes (Residual)	187	15.6	258.7
American Indian, both sexes, 70-74 years				
...	All causes	1,285	100.0	2,467.2
1	Malignant neoplasms (C00-C97)	342	26.6	656.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	292	22.7	560.6
3	Diabetes mellitus (E10-E14)	128	10.0	245.8
4	Chronic lower respiratory diseases (J40-J47)	79	6.1	151.7
5	Cerebrovascular diseases (I60-I69)	78	6.1	149.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	38	3.0	73.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	34	2.6	65.3
8	Chronic liver disease and cirrhosis (K70,K73-K74)	31	2.4	59.5
9	Influenza and pneumonia (J10-J18)	27	2.1	51.8
10	Septicemia (A40-A41)	21	1.6	40.3
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	17	1.3	*
12	Atherosclerosis (I70)	9	0.7	*
12	Pneumonitis due to solids and liquids (J69)	9	0.7	*
14	Parkinson's disease (G20-G21)	6	0.5	*
14	Alzheimer's disease (G30)	6	0.5	*
...	All other causes (Residual)	168	13.1	322.6

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 75-79 years				
...	All causes	1,263	100.0	3,372.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	307	24.3	819.8
2	Malignant neoplasms (C00-C97)	293	23.2	782.4
3	Diabetes mellitus (E10-E14)	92	7.3	245.7
4	Chronic lower respiratory diseases (J40-J47)	82	6.5	219.0
5	Cerebrovascular diseases (I60-I69)	81	6.4	216.3
6	Influenza and pneumonia (J10-J18)	39	3.1	104.1
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	35	2.8	93.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	34	2.7	90.8
9	Alzheimer's disease (G30)	32	2.5	85.4
10	Septicemia (A40-A41)	21	1.7	56.1
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	21	1.7	56.1
12	Chronic liver disease and cirrhosis (K70,K73-K74)	17	1.3	*
13	Parkinson's disease (G20-G21)	11	0.9	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.7	*
14	Atherosclerosis (I70)	9	0.7	*
...	All other causes (Residual)	180	14.3	480.7
American Indian, both sexes, 80-84 years				
...	All causes	1,170	100.0	4,777.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	290	24.8	1,184.2
2	Malignant neoplasms (C00-C97)	225	19.2	918.7
3	Cerebrovascular diseases (I60-I69)	82	7.0	334.8
3	Chronic lower respiratory diseases (J40-J47)	82	7.0	334.8
5	Diabetes mellitus (E10-E14)	67	5.7	273.6
6	Influenza and pneumonia (J10-J18)	59	5.0	240.9
7	Alzheimer's disease (G30)	37	3.2	151.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	28	2.4	114.3
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	25	2.1	102.1
10	Septicemia (A40-A41)	24	2.1	98.0
11	Parkinson's disease (G20-G21)	20	1.7	81.7
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	17	1.5	*
13	Chronic liver disease and cirrhosis (K70,K73-K74)	10	0.9	*
14	Pneumonitis due to solids and liquids (J69)	9	0.8	*
15	Atherosclerosis (I70)	7	0.6	*
15	Aortic aneurysm and dissection (I71)	7	0.6	*
...	All other causes (Residual)	181	15.5	739.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 85-89 years				
...	All causes	871	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	247	28.4	---
2	Malignant neoplasms (C00-C97)	127	14.6	---
3	Cerebrovascular diseases (I60-I69)	65	7.5	---
4	Diabetes mellitus (E10-E14)	57	6.5	---
5	Influenza and pneumonia (J10-J18)	55	6.3	---
6	Alzheimer's disease (G30)	42	4.8	---
7	Chronic lower respiratory diseases (J40-J47)	40	4.6	---
8	Septicemia (A40-A41)	17	2.0	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	2.0	---
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	14	1.6	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14	1.6	---
12	Parkinson's disease (G20-G21)	10	1.1	---
13	Pneumonitis due to solids and liquids (J69)	9	1.0	---
14	Atherosclerosis (I70)	5	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.5	---
15	Chronic liver disease and cirrhosis (K70,K73-K74)	4	0.5	---
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	4	0.5	---
...	All other causes (Residual)	140	16.1	---
American Indian, both sexes, 90-94 years				
...	All causes	539	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	169	31.4	---
2	Cerebrovascular diseases (I60-I69)	45	8.3	---
3	Malignant neoplasms (C00-C97)	43	8.0	---
4	Alzheimer's disease (G30)	33	6.1	---
5	Influenza and pneumonia (J10-J18)	27	5.0	---
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	27	5.0	---
7	Diabetes mellitus (E10-E14)	24	4.5	---
8	Chronic lower respiratory diseases (J40-J47)	21	3.9	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10	1.9	---
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	9	1.7	---
11	Septicemia (A40-A41)	7	1.3	---
11	Pneumonitis due to solids and liquids (J69)	7	1.3	---
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.9	---
13	Parkinson's disease (G20-G21)	5	0.9	---
13	Atherosclerosis (I70)	5	0.9	---
...	All other causes (Residual)	102	18.9	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 95-99 years				
...	All causes	208	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	68	32.7	---
2	Alzheimer's disease (G30)	16	7.7	---
2	Influenza and pneumonia (J10-J18)	16	7.7	---
4	Malignant neoplasms (C00-C97)	13	6.3	---
4	Cerebrovascular diseases (I60-I69)	13	6.3	---
6	Chronic lower respiratory diseases (J40-J47)	7	3.4	---
7	Septicemia (A40-A41)	6	2.9	---
7	Diabetes mellitus (E10-E14)	6	2.9	---
9	Pneumonitis due to solids and liquids (J69)	5	2.4	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5	2.4	---
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	1.9	---
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	3	1.4	---
13	Anemias (D50-D64)	2	1.0	---
13	Aortic aneurysm and dissection (I71)	2	1.0	---
13	Cholelithiasis and other disorders of gallbladder (K80-K82)	2	1.0	---
...	All other causes (Residual)	40	19.2	---
American Indian, both sexes, 100 years and over				
...	All causes	66	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	24	36.4	---
2	Influenza and pneumonia (J10-J18)	7	10.6	---
3	Cerebrovascular diseases (I60-I69)	6	9.1	---
4	Alzheimer's disease (G30)	3	4.5	---
5	Parkinson's disease (G20-G21)	2	3.0	---
6	Anemias (D50-D64)	1	1.5	---
6	Diabetes mellitus (E10-E14)	1	1.5	---
6	Nutritional deficiencies (E40-E64)	1	1.5	---
6	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1	1.5	---
6	Acute bronchitis and bronchiolitis (J20-J21)	1	1.5	---
6	Chronic lower respiratory diseases (J40-J47)	1	1.5	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	1.5	---
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1	1.5	---
...	All other causes (Residual)	16	24.2	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, all ages				
...	All causes	7,607	100.0	481.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,471	19.3	93.2
2	Malignant neoplasms (C00-C97)	1,297	17.1	82.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,118	14.7	70.8
4	Diabetes mellitus (E10-E14)	412	5.4	26.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	348	4.6	22.0
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	313	4.1	19.8
7	Chronic lower respiratory diseases (J40-J47)	257	3.4	16.3
8	Cerebrovascular diseases (I60-I69)	249	3.3	15.8
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	193	2.5	12.2
10	Influenza and pneumonia (J10-J18)	180	2.4	11.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	105	1.4	6.7
12	Septicemia (A40-A41)	104	1.4	6.6
13	Certain conditions originating in the perinatal period (P00-P96)	75	1.0	4.8
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	61	0.8	3.9
15	Human immunodeficiency virus (HIV) disease (B20-B24)	59	0.8	3.7
...	All other causes (Residual)	1,365	17.9	86.5
American Indian, male, 1-4 years				
...	All causes	62	100.0	72.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	31	50.0	36.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	6.5	*
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	6.5	*
4	Malignant neoplasms (C00-C97)	3	4.8	*
5	Meningitis (G00,G03)	2	3.2	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	1.6	*
...	All other causes (Residual)	17	27.4	*
American Indian, male, 5-9 years				
...	All causes	27	100.0	20.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18	66.7	*
2	Malignant neoplasms (C00-C97)	3	11.1	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	7.4	*
4	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	3.7	*
...	All other causes (Residual)	3	11.1	*

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 10-14 years				
...	All causes	37	100.0	25.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18	48.6	*
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11	29.7	*
3	Human immunodeficiency virus (HIV) disease (B20-B24)	1	2.7	*
3	Malignant neoplasms (C00-C97)	1	2.7	*
3	Influenza and pneumonia (J10-J18)	1	2.7	*
3	Chronic lower respiratory diseases (J40-J47)	1	2.7	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	2.7	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1	2.7	*
...	All other causes (Residual)	2	5.4	*
American Indian, male, 15-19 years				
...	All causes	180	100.0	117.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	97	53.9	63.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	37	20.6	24.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	20	11.1	13.1
4	Malignant neoplasms (C00-C97)	7	3.9	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	1.7	*
6	Diabetes mellitus (E10-E14)	1	0.6	*
6	Cerebrovascular diseases (I60-I69)	1	0.6	*
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.6	*
...	All other causes (Residual)	13	7.2	*
American Indian, male, 20-24 years				
...	All causes	259	100.0	173.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	126	48.6	84.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	62	23.9	41.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	47	18.1	31.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	1.5	*
5	Malignant neoplasms (C00-C97)	3	1.2	*
6	Chronic liver disease and cirrhosis (K70,K73-K74)	2	0.8	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.4	*
7	Diabetes mellitus (E10-E14)	1	0.4	*
...	All other causes (Residual)	13	5.0	*

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 25-29 years				
...	All causes	264	100.0	201.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	131	49.6	100.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	38	14.4	29.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	27	10.2	20.6
4	Malignant neoplasms (C00-C97)	9	3.4	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	9	3.4	*
6	Chronic liver disease and cirrhosis (K70,K73-K74)	6	2.3	*
7	Diabetes mellitus (E10-E14)	4	1.5	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	3	1.1	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	1.1	*
10	Influenza and pneumonia (J10-J18)	2	0.8	*
10	Legal intervention (Y35,Y89.0)	2	0.8	*
12	Septicemia (A40-A41)	1	0.4	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.4	*
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1	0.4	*
12	Cerebrovascular diseases (I60-I69)	1	0.4	*
12	Aortic aneurysm and dissection (I71)	1	0.4	*
...	All other causes (Residual)	25	9.5	19.1
American Indian, male, 30-34 years				
...	All causes	248	100.0	211.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	109	44.0	93.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	35	14.1	29.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	24	9.7	20.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	18	7.3	*
5	Chronic liver disease and cirrhosis (K70,K73-K74)	16	6.5	*
6	Malignant neoplasms (C00-C97)	7	2.8	*
6	Diabetes mellitus (E10-E14)	7	2.8	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	4	1.6	*
9	Influenza and pneumonia (J10-J18)	2	0.8	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.8	*
11	Tuberculosis (A16-A19)	1	0.4	*
11	Cerebrovascular diseases (I60-I69)	1	0.4	*
11	Chronic lower respiratory diseases (J40-J47)	1	0.4	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.4	*
...	All other causes (Residual)	20	8.1	17.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 35-39 years				
...	All causes	306	100.0	270.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	100	32.7	88.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	31	10.1	27.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	28	9.2	24.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	21	6.9	18.6
5	Malignant neoplasms (C00-C97)	16	5.2	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	15	4.9	*
7	Diabetes mellitus (E10-E14)	10	3.3	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	9	2.9	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	2.0	*
10	Influenza and pneumonia (J10-J18)	5	1.6	*
11	Septicemia (A40-A41)	3	1.0	*
11	Viral hepatitis (B15-B19)	3	1.0	*
11	Cerebrovascular diseases (I60-I69)	3	1.0	*
11	Legal intervention (Y35,Y89.0)	3	1.0	*
15	Peptic ulcer (K25-K28)	1	0.3	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.3	*
...	All other causes (Residual)	51	16.7	45.1
American Indian, male, 40-44 years				
...	All causes	471	100.0	400.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	113	24.0	96.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	56	11.9	47.6
3	Chronic liver disease and cirrhosis (K70,K73-K74)	49	10.4	41.6
4	Malignant neoplasms (C00-C97)	46	9.8	39.1
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	37	7.9	31.4
6	Diabetes mellitus (E10-E14)	17	3.6	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	17	3.6	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	13	2.8	*
8	Cerebrovascular diseases (I60-I69)	13	2.8	*
10	Septicemia (A40-A41)	7	1.5	*
10	Influenza and pneumonia (J10-J18)	7	1.5	*
12	Viral hepatitis (B15-B19)	4	0.8	*
13	Chronic lower respiratory diseases (J40-J47)	3	0.6	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.6	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.6	*
...	All other causes (Residual)	83	17.6	70.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 45-49 years				
...	All causes	524	100.0	486.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	108	20.6	100.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	100	19.1	92.8
3	Malignant neoplasms (C00-C97)	62	11.8	57.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	53	10.1	49.2
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	25	4.8	23.2
6	Diabetes mellitus (E10-E14)	18	3.4	*
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	15	2.9	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	14	2.7	*
9	Viral hepatitis (B15-B19)	8	1.5	*
9	Cerebrovascular diseases (I60-I69)	8	1.5	*
9	Influenza and pneumonia (J10-J18)	8	1.5	*
12	Septicemia (A40-A41)	7	1.3	*
13	Chronic lower respiratory diseases (J40-J47)	4	0.8	*
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	3	0.6	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.6	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.6	*
...	All other causes (Residual)	85	16.2	78.9
American Indian, male, 50-54 years				
...	All causes	636	100.0	712.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	121	19.0	135.6
2	Malignant neoplasms (C00-C97)	99	15.6	110.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	73	11.5	81.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	63	9.9	70.6
5	Diabetes mellitus (E10-E14)	45	7.1	50.4
6	Cerebrovascular diseases (I60-I69)	19	3.0	*
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	17	2.7	*
8	Chronic lower respiratory diseases (J40-J47)	14	2.2	*
9	Viral hepatitis (B15-B19)	10	1.6	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	10	1.6	*
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	10	1.6	*
12	Influenza and pneumonia (J10-J18)	8	1.3	*
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	6	0.9	*
14	Septicemia (A40-A41)	5	0.8	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.6	*
...	All other causes (Residual)	132	20.8	147.9

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 55-59 years				
...	All causes	678	100.0	950.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	161	23.7	225.6
2	Malignant neoplasms (C00-C97)	149	22.0	208.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	57	8.4	79.9
4	Diabetes mellitus (E10-E14)	53	7.8	74.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	49	7.2	68.7
6	Cerebrovascular diseases (I60-I69)	25	3.7	35.0
7	Chronic lower respiratory diseases (J40-J47)	15	2.2	*
8	Septicemia (A40-A41)	14	2.1	*
9	Influenza and pneumonia (J10-J18)	11	1.6	*
10	Viral hepatitis (B15-B19)	9	1.3	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	1.3	*
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	9	1.3	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	3	0.4	*
13	Atherosclerosis (I70)	3	0.4	*
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	0.4	*
...	All other causes (Residual)	108	15.9	151.4
American Indian, male, 60-64 years				
...	All causes	663	100.0	1,383.0
1	Malignant neoplasms (C00-C97)	172	25.9	358.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	167	25.2	348.4
3	Chronic liver disease and cirrhosis (K70,K73-K74)	46	6.9	96.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	42	6.3	87.6
5	Diabetes mellitus (E10-E14)	41	6.2	85.5
6	Chronic lower respiratory diseases (J40-J47)	30	4.5	62.6
7	Cerebrovascular diseases (I60-I69)	18	2.7	*
8	Influenza and pneumonia (J10-J18)	13	2.0	*
9	Septicemia (A40-A41)	8	1.2	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	1.2	*
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	0.9	*
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	4	0.6	*
12	Pneumonitis due to solids and liquids (J69)	4	0.6	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.5	*
14	Cholelithiasis and other disorders of gallbladder (K80-K82)	3	0.5	*
...	All other causes (Residual)	98	14.8	204.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 65-69 years				
...	All causes	624	100.0	1,830.6
1	Malignant neoplasms (C00-C97)	175	28.0	513.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	151	24.2	443.0
3	Diabetes mellitus (E10-E14)	54	8.7	158.4
4	Cerebrovascular diseases (I60-I69)	29	4.6	85.1
5	Chronic lower respiratory diseases (J40-J47)	24	3.8	70.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	22	3.5	64.5
7	Septicemia (A40-A41)	16	2.6	*
7	Chronic liver disease and cirrhosis (K70,K73-K74)	16	2.6	*
9	Influenza and pneumonia (J10-J18)	15	2.4	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	1.3	*
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	6	1.0	*
11	Aortic aneurysm and dissection (I71)	6	1.0	*
13	Alzheimer's disease (G30)	3	0.5	*
13	Pneumonitis due to solids and liquids (J69)	3	0.5	*
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3	0.5	*
...	All other causes (Residual)	93	14.9	272.8
American Indian, male, 70-74 years				
...	All causes	673	100.0	2,869.6
1	Malignant neoplasms (C00-C97)	183	27.2	780.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	168	25.0	716.3
3	Diabetes mellitus (E10-E14)	61	9.1	260.1
4	Chronic lower respiratory diseases (J40-J47)	41	6.1	174.8
5	Cerebrovascular diseases (I60-I69)	36	5.3	153.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	23	3.4	98.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20	3.0	85.3
8	Septicemia (A40-A41)	13	1.9	*
8	Chronic liver disease and cirrhosis (K70,K73-K74)	13	1.9	*
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	10	1.5	*
11	Influenza and pneumonia (J10-J18)	9	1.3	*
12	Atherosclerosis (I70)	6	0.9	*
13	Pneumonitis due to solids and liquids (J69)	5	0.7	*
14	Aortic aneurysm and dissection (I71)	4	0.6	*
15	Anemias (D50-D64)	3	0.4	*
15	Nutritional deficiencies (E40-E64)	3	0.4	*
15	Alzheimer's disease (G30)	3	0.4	*
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.4	*
...	All other causes (Residual)	66	9.8	281.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 75-79 years				
...	All causes	611	100.0	3,750.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	156	25.5	957.5
2	Malignant neoplasms (C00-C97)	146	23.9	896.1
3	Diabetes mellitus (E10-E14)	43	7.0	263.9
3	Chronic lower respiratory diseases (J40-J47)	43	7.0	263.9
5	Cerebrovascular diseases (I60-I69)	36	5.9	221.0
6	Influenza and pneumonia (J10-J18)	25	4.1	153.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15	2.5	*
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	11	1.8	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	1.8	*
10	Septicemia (A40-A41)	10	1.6	*
11	Alzheimer's disease (G30)	9	1.5	*
12	Parkinson's disease (G20-G21)	7	1.1	*
13	Pneumonitis due to solids and liquids (J69)	6	1.0	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.8	*
14	Chronic liver disease and cirrhosis (K70,K73-K74)	5	0.8	*
...	All other causes (Residual)	83	13.6	509.5
American Indian, male, 80-84 years				
...	All causes	533	100.0	5,401.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	138	25.9	1,398.6
2	Malignant neoplasms (C00-C97)	121	22.7	1,226.3
3	Chronic lower respiratory diseases (J40-J47)	49	9.2	496.6
4	Influenza and pneumonia (J10-J18)	27	5.1	273.6
5	Diabetes mellitus (E10-E14)	26	4.9	263.5
6	Cerebrovascular diseases (I60-I69)	24	4.5	243.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13	2.4	*
8	Parkinson's disease (G20-G21)	11	2.1	*
9	Alzheimer's disease (G30)	10	1.9	*
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	7	1.3	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	1.3	*
12	Septicemia (A40-A41)	6	1.1	*
12	Pneumonitis due to solids and liquids (J69)	6	1.1	*
12	Chronic liver disease and cirrhosis (K70,K73-K74)	6	1.1	*
15	Aortic aneurysm and dissection (I71)	4	0.8	*
...	All other causes (Residual)	78	14.6	790.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 85-89 years				
...	All causes	367	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	104	28.3	---
2	Malignant neoplasms (C00-C97)	67	18.3	---
3	Cerebrovascular diseases (I60-I69)	23	6.3	---
4	Influenza and pneumonia (J10-J18)	22	6.0	---
5	Diabetes mellitus (E10-E14)	21	5.7	---
6	Chronic lower respiratory diseases (J40-J47)	19	5.2	---
7	Alzheimer's disease (G30)	10	2.7	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	2.7	---
9	Septicemia (A40-A41)	9	2.5	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9	2.5	---
11	Parkinson's disease (G20-G21)	7	1.9	---
12	Pneumonitis due to solids and liquids (J69)	6	1.6	---
13	Chronic liver disease and cirrhosis (K70,K73-K74)	3	0.8	---
14	Tuberculosis (A16-A19)	2	0.5	---
14	Anemias (D50-D64)	2	0.5	---
14	Nutritional deficiencies (E40-E64)	2	0.5	---
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2	0.5	---
14	Atherosclerosis (I70)	2	0.5	---
14	Cholelithiasis and other disorders of gallbladder (K80-K82)	2	0.5	---
...	All other causes (Residual)	45	12.3	---
American Indian, male, 90-94 years				
...	All causes	179	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	56	31.3	---
2	Malignant neoplasms (C00-C97)	20	11.2	---
3	Influenza and pneumonia (J10-J18)	15	8.4	---
4	Chronic lower respiratory diseases (J40-J47)	12	6.7	---
5	Alzheimer's disease (G30)	11	6.1	---
6	Cerebrovascular diseases (I60-I69)	9	5.0	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	5.0	---
8	Diabetes mellitus (E10-E14)	8	4.5	---
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	5	2.8	---
10	Septicemia (A40-A41)	4	2.2	---
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	1.1	---
11	Anemias (D50-D64)	2	1.1	---
11	Atherosclerosis (I70)	2	1.1	---
11	Pneumonitis due to solids and liquids (J69)	2	1.1	---
15	Tuberculosis (A16-A19)	1	0.6	---
15	Parkinson's disease (G20-G21)	1	0.6	---
15	Diseases of appendix (K35-K38)	1	0.6	---
15	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1	0.6	---
...	All other causes (Residual)	18	10.1	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 95-99 years				
...	All causes	55	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	17	30.9	---
2	Malignant neoplasms (C00-C97)	8	14.5	---
3	Influenza and pneumonia (J10-J18)	6	10.9	---
4	Diabetes mellitus (E10-E14)	2	3.6	---
4	Alzheimer's disease (G30)	2	3.6	---
4	Cerebrovascular diseases (I60-I69)	2	3.6	---
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2	3.6	---
8	Septicemia (A40-A41)	1	1.8	---
8	Nutritional deficiencies (E40-E64)	1	1.8	---
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1	1.8	---
8	Chronic lower respiratory diseases (J40-J47)	1	1.8	---
8	Pneumonitis due to solids and liquids (J69)	1	1.8	---
...	All other causes (Residual)	11	20.0	---
American Indian, male, 100 years and over				
...	All causes	13	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	6	46.2	---
2	Influenza and pneumonia (J10-J18)	2	15.4	---
3	Parkinson's disease (G20-G21)	1	7.7	---
...	All other causes (Residual)	4	30.8	---
American Indian, female, all ages				
...	All causes	6,311	100.0	398.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,188	18.8	75.1
2	Malignant neoplasms (C00-C97)	1,168	18.5	73.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	508	8.0	32.1
4	Diabetes mellitus (E10-E14)	406	6.4	25.7
5	Cerebrovascular diseases (I60-I69)	378	6.0	23.9
6	Chronic lower respiratory diseases (J40-J47)	261	4.1	16.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	248	3.9	15.7
8	Influenza and pneumonia (J10-J18)	174	2.8	11.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	160	2.5	10.1
10	Alzheimer's disease (G30)	125	2.0	7.9
11	Septicemia (A40-A41)	113	1.8	7.1
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	79	1.3	5.0
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	64	1.0	4.0
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	57	0.9	3.6
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	56	0.9	3.5
...	All other causes (Residual)	1,326	21.0	83.8

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 1-4 years				
...	All causes	38	100.0	45.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13	34.2	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	18.4	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	15.8	*
4	Septicemia (A40-A41)	1	2.6	*
4	Malignant neoplasms (C00-C97)	1	2.6	*
4	Meningitis (G00,G03)	1	2.6	*
4	Influenza and pneumonia (J10-J18)	1	2.6	*
4	Chronic lower respiratory diseases (J40-J47)	1	2.6	*
...	All other causes (Residual)	7	18.4	*
American Indian, female, 5-9 years				
...	All causes	19	100.0	*
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9	47.4	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	26.3	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	10.5	*
...	All other causes (Residual)	3	15.8	*
American Indian, female, 10-14 years				
...	All causes	27	100.0	18.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10	37.0	*
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3	11.1	*
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	7.4	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	7.4	*
5	Meningitis (G00,G03)	1	3.7	*
5	Chronic lower respiratory diseases (J40-J47)	1	3.7	*
...	All other causes (Residual)	8	29.6	*
American Indian, female, 15-19 years				
...	All causes	102	100.0	68.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	45	44.1	30.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	22	21.6	14.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	9	8.8	*
4	Malignant neoplasms (C00-C97)	3	2.9	*
5	Septicemia (A40-A41)	2	2.0	*
6	Salmonella infections (A01-A02)	1	1.0	*
6	Diabetes mellitus (E10-E14)	1	1.0	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	1.0	*
6	Influenza and pneumonia (J10-J18)	1	1.0	*
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1	1.0	*
6	Pregnancy, childbirth and the puerperium (O00-O99)	1	1.0	*
...	All other causes (Residual)	15	14.7	*

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 20-24 years				
...	All causes	94	100.0	66.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	53	56.4	37.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7	7.4	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	6.4	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	5.3	*
5	Malignant neoplasms (C00-C97)	2	2.1	*
6	Septicemia (A40-A41)	1	1.1	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.1	*
6	Anemias (D50-D64)	1	1.1	*
6	Diabetes mellitus (E10-E14)	1	1.1	*
6	Cerebrovascular diseases (I60-I69)	1	1.1	*
6	Pregnancy, childbirth and the puerperium (O00-099)	1	1.1	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	1.1	*
...	All other causes (Residual)	14	14.9	*
American Indian, female, 25-29 years				
...	All causes	100	100.0	83.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	44	44.0	36.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11	11.0	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	10	10.0	*
4	Malignant neoplasms (C00-C97)	8	8.0	*
5	Chronic liver disease and cirrhosis (K70,K73-K74)	5	5.0	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	4.0	*
7	Human immunodeficiency virus (HIV) disease (B20-B24)	2	2.0	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	2.0	*
9	Influenza and pneumonia (J10-J18)	1	1.0	*
9	Chronic lower respiratory diseases (J40-J47)	1	1.0	*
9	Pregnancy, childbirth and the puerperium (O00-099)	1	1.0	*
...	All other causes (Residual)	11	11.0	*
American Indian, female, 30-34 years				
...	All causes	108	100.0	98.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	40	37.0	36.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	9	8.3	*
2	Chronic liver disease and cirrhosis (K70,K73-K74)	9	8.3	*
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7	6.5	*
5	Diabetes mellitus (E10-E14)	6	5.6	*
6	Malignant neoplasms (C00-C97)	5	4.6	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	4.6	*
8	Septicemia (A40-A41)	3	2.8	*
9	Cerebrovascular diseases (I60-I69)	2	1.9	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.9	*
10	Chronic lower respiratory diseases (J40-J47)	1	0.9	*
10	Pneumonitis due to solids and liquids (J69)	1	0.9	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.9	*
10	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.9	*
...	All other causes (Residual)	17	15.7	*

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 35-39 years				
...	All causes	191	100.0	173.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	47	24.6	42.7
2	Chronic liver disease and cirrhosis (K70,K73-K74)	20	10.5	18.2
3	Malignant neoplasms (C00-C97)	19	9.9	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	15	7.9	*
5	Cerebrovascular diseases (I60-I69)	10	5.2	*
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8	4.2	*
7	Human immunodeficiency virus (HIV) disease (B20-B24)	6	3.1	*
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	3.1	*
9	Diabetes mellitus (E10-E14)	5	2.6	*
9	Chronic lower respiratory diseases (J40-J47)	5	2.6	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	2.1	*
12	Septicemia (A40-A41)	3	1.6	*
12	Pregnancy, childbirth and the puerperium (O00-099)	3	1.6	*
14	Influenza and pneumonia (J10-J18)	2	1.0	*
15	Viral hepatitis (B15-B19)	1	0.5	*
15	Meningitis (G00,G03)	1	0.5	*
15	Hernia (K40-K46)	1	0.5	*
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.5	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.5	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.5	*
...	All other causes (Residual)	32	16.8	29.1
American Indian, female, 40-44 years				
...	All causes	255	100.0	212.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	45	17.6	37.6
2	Malignant neoplasms (C00-C97)	35	13.7	29.2
2	Chronic liver disease and cirrhosis (K70,K73-K74)	35	13.7	29.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	21	8.2	17.5
5	Diabetes mellitus (E10-E14)	9	3.5	*
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8	3.1	*
7	Human immunodeficiency virus (HIV) disease (B20-B24)	7	2.7	*
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7	2.7	*
9	Cerebrovascular diseases (I60-I69)	6	2.4	*
10	Influenza and pneumonia (J10-J18)	4	1.6	*
10	Chronic lower respiratory diseases (J40-J47)	4	1.6	*
12	Septicemia (A40-A41)	3	1.2	*
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	1.2	*
14	Viral hepatitis (B15-B19)	2	0.8	*
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2	0.8	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.8	*
...	All other causes (Residual)	62	24.3	51.8

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 45-49 years				
...	All causes	350	100.0	307.0
1	Malignant neoplasms (C00-C97)	75	21.4	65.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	47	13.4	41.2
3	Chronic liver disease and cirrhosis (K70,K73-K74)	44	12.6	38.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	42	12.0	36.8
5	Diabetes mellitus (E10-E14)	18	5.1	*
6	Cerebrovascular diseases (I60-I69)	13	3.7	*
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	3.1	*
8	Influenza and pneumonia (J10-J18)	6	1.7	*
8	Chronic lower respiratory diseases (J40-J47)	6	1.7	*
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6	1.7	*
11	Septicemia (A40-A41)	3	0.9	*
12	Pneumonitis due to solids and liquids (J69)	2	0.6	*
12	Cholelithiasis and other disorders of gallbladder (K80-K82)	2	0.6	*
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	0.6	*
15	Viral hepatitis (B15-B19)	1	0.3	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.3	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.3	*
15	Aortic aneurysm and dissection (I71)	1	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.3	*
...	All other causes (Residual)	68	19.4	59.6
American Indian, female, 50-54 years				
...	All causes	417	100.0	437.0
1	Malignant neoplasms (C00-C97)	104	24.9	109.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	62	14.9	65.0
3	Chronic liver disease and cirrhosis (K70,K73-K74)	36	8.6	37.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	35	8.4	36.7
5	Cerebrovascular diseases (I60-I69)	24	5.8	25.2
6	Diabetes mellitus (E10-E14)	17	4.1	*
7	Septicemia (A40-A41)	9	2.2	*
7	Chronic lower respiratory diseases (J40-J47)	9	2.2	*
9	Influenza and pneumonia (J10-J18)	5	1.2	*
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5	1.2	*
11	Viral hepatitis (B15-B19)	4	1.0	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.7	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.7	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.5	*
14	Nutritional deficiencies (E40-E64)	2	0.5	*
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	0.5	*
...	All other causes (Residual)	95	22.8	99.6

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 55-59 years				
...	All causes	397	100.0	522.7
1	Malignant neoplasms (C00-C97)	107	27.0	140.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	76	19.1	100.1
3	Diabetes mellitus (E10-E14)	42	10.6	55.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	23	5.8	30.3
5	Chronic lower respiratory diseases (J40-J47)	17	4.3	*
6	Cerebrovascular diseases (I60-I69)	16	4.0	*
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15	3.8	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	3.5	*
9	Septicemia (A40-A41)	8	2.0	*
9	Influenza and pneumonia (J10-J18)	8	2.0	*
11	Atherosclerosis (I70)	2	0.5	*
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	0.5	*
13	Tuberculosis (A16-A19)	1	0.3	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.3	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.3	*
13	Nutritional deficiencies (E40-E64)	1	0.3	*
13	Parkinson's disease (G20-G21)	1	0.3	*
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1	0.3	*
13	Peptic ulcer (K25-K28)	1	0.3	*
13	Hernia (K40-K46)	1	0.3	*
13	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.3	*
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1	0.3	*
...	All other causes (Residual)	57	14.4	75.1
American Indian, female, 60-64 years				
...	All causes	502	100.0	954.4
1	Malignant neoplasms (C00-C97)	152	30.3	289.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	81	16.1	154.0
3	Diabetes mellitus (E10-E14)	41	8.2	77.9
4	Cerebrovascular diseases (I60-I69)	30	6.0	57.0
5	Chronic lower respiratory diseases (J40-J47)	29	5.8	55.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	25	5.0	47.5
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	21	4.2	39.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	3.2	*
9	Septicemia (A40-A41)	15	3.0	*
10	Influenza and pneumonia (J10-J18)	11	2.2	*
11	Viral hepatitis (B15-B19)	4	0.8	*
12	Atherosclerosis (I70)	3	0.6	*
13	Parkinson's disease (G20-G21)	2	0.4	*
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2	0.4	*
13	Aortic aneurysm and dissection (I71)	2	0.4	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	0.4	*
...	All other causes (Residual)	66	13.1	125.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 65-69 years				
...	All causes	578	100.0	1,513.0
1	Malignant neoplasms (C00-C97)	159	27.5	416.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	109	18.9	285.3
3	Diabetes mellitus (E10-E14)	52	9.0	136.1
4	Chronic lower respiratory diseases (J40-J47)	39	6.7	102.1
5	Cerebrovascular diseases (I60-I69)	35	6.1	91.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	4.0	60.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15	2.6	*
8	Chronic liver disease and cirrhosis (K70,K73-K74)	14	2.4	*
9	Septicemia (A40-A41)	10	1.7	*
10	Influenza and pneumonia (J10-J18)	8	1.4	*
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	5	0.9	*
12	Atherosclerosis (I70)	3	0.5	*
12	Pneumonitis due to solids and liquids (J69)	3	0.5	*
12	Hernia (K40-K46)	3	0.5	*
15	Viral hepatitis (B15-B19)	2	0.3	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.3	*
15	Anemias (D50-D64)	2	0.3	*
15	Parkinson's disease (G20-G21)	2	0.3	*
15	Aortic aneurysm and dissection (I71)	2	0.3	*
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	0.3	*
...	All other causes (Residual)	88	15.2	230.3
American Indian, female, 70-74 years				
...	All causes	612	100.0	2,137.5
1	Malignant neoplasms (C00-C97)	159	26.0	555.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	124	20.3	433.1
3	Diabetes mellitus (E10-E14)	67	10.9	234.0
4	Cerebrovascular diseases (I60-I69)	42	6.9	146.7
5	Chronic lower respiratory diseases (J40-J47)	38	6.2	132.7
6	Influenza and pneumonia (J10-J18)	18	2.9	*
6	Chronic liver disease and cirrhosis (K70,K73-K74)	18	2.9	*
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15	2.5	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	2.3	*
10	Septicemia (A40-A41)	8	1.3	*
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	7	1.1	*
12	Parkinson's disease (G20-G21)	5	0.8	*
13	Pneumonitis due to solids and liquids (J69)	4	0.7	*
14	Alzheimer's disease (G30)	3	0.5	*
14	Atherosclerosis (I70)	3	0.5	*
...	All other causes (Residual)	87	14.2	303.9

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 75-79 years				
...	All causes	652	100.0	3,081.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	151	23.2	713.7
2	Malignant neoplasms (C00-C97)	147	22.5	694.8
3	Diabetes mellitus (E10-E14)	49	7.5	231.6
4	Cerebrovascular diseases (I60-I69)	45	6.9	212.7
5	Chronic lower respiratory diseases (J40-J47)	39	6.0	184.3
6	Alzheimer's disease (G30)	23	3.5	108.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	3.5	108.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	20	3.1	94.5
9	Influenza and pneumonia (J10-J18)	14	2.1	*
10	Chronic liver disease and cirrhosis (K70,K73-K74)	12	1.8	*
11	Septicemia (A40-A41)	11	1.7	*
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	10	1.5	*
13	Atherosclerosis (I70)	6	0.9	*
14	Aortic aneurysm and dissection (I71)	5	0.8	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.6	*
15	Parkinson's disease (G20-G21)	4	0.6	*
...	All other causes (Residual)	89	13.7	420.7
American Indian, female, 80-84 years				
...	All causes	637	100.0	4,356.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	152	23.9	1,039.5
2	Malignant neoplasms (C00-C97)	104	16.3	711.2
3	Cerebrovascular diseases (I60-I69)	58	9.1	396.6
4	Diabetes mellitus (E10-E14)	41	6.4	280.4
5	Chronic lower respiratory diseases (J40-J47)	33	5.2	225.7
6	Influenza and pneumonia (J10-J18)	32	5.0	218.8
7	Alzheimer's disease (G30)	27	4.2	184.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21	3.3	143.6
9	Septicemia (A40-A41)	18	2.8	*
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12	1.9	*
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	10	1.6	*
12	Parkinson's disease (G20-G21)	9	1.4	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.6	*
13	Atherosclerosis (I70)	4	0.6	*
13	Chronic liver disease and cirrhosis (K70,K73-K74)	4	0.6	*
13	Cholelithiasis and other disorders of gallbladder (K80-K82)	4	0.6	*
...	All other causes (Residual)	104	16.3	711.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 85-89 years				
...	All causes	504	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	143	28.4	---
2	Malignant neoplasms (C00-C97)	60	11.9	---
3	Cerebrovascular diseases (I60-I69)	42	8.3	---
4	Diabetes mellitus (E10-E14)	36	7.1	---
5	Influenza and pneumonia (J10-J18)	33	6.5	---
6	Alzheimer's disease (G30)	32	6.3	---
7	Chronic lower respiratory diseases (J40-J47)	21	4.2	---
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	12	2.4	---
9	Septicemia (A40-A41)	8	1.6	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	1.4	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5	1.0	---
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.6	---
12	Parkinson's disease (G20-G21)	3	0.6	---
12	Atherosclerosis (I70)	3	0.6	---
12	Aortic aneurysm and dissection (I71)	3	0.6	---
12	Pneumonitis due to solids and liquids (J69)	3	0.6	---
...	All other causes (Residual)	90	17.9	---
American Indian, female, 90-94 years				
...	All causes	360	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	113	31.4	---
2	Cerebrovascular diseases (I60-I69)	36	10.0	---
3	Malignant neoplasms (C00-C97)	23	6.4	---
4	Alzheimer's disease (G30)	22	6.1	---
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18	5.0	---
6	Diabetes mellitus (E10-E14)	16	4.4	---
7	Influenza and pneumonia (J10-J18)	12	3.3	---
8	Chronic lower respiratory diseases (J40-J47)	9	2.5	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9	2.5	---
10	Pneumonitis due to solids and liquids (J69)	5	1.4	---
11	Parkinson's disease (G20-G21)	4	1.1	---
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	4	1.1	---
13	Septicemia (A40-A41)	3	0.8	---
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.8	---
13	Atherosclerosis (I70)	3	0.8	---
...	All other causes (Residual)	80	22.2	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 95-99 years				
...	All causes	153	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	51	33.3	---
2	Alzheimer's disease (G30)	14	9.2	---
3	Cerebrovascular diseases (I60-I69)	11	7.2	---
4	Influenza and pneumonia (J10-J18)	10	6.5	---
5	Chronic lower respiratory diseases (J40-J47)	6	3.9	---
6	Septicemia (A40-A41)	5	3.3	---
6	Malignant neoplasms (C00-C97)	5	3.3	---
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	2.6	---
8	Diabetes mellitus (E10-E14)	4	2.6	---
8	Pneumonitis due to solids and liquids (J69)	4	2.6	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3	2.0	---
12	Anemias (D50-D64)	2	1.3	---
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2	1.3	---
12	Aortic aneurysm and dissection (I71)	2	1.3	---
12	Cholelithiasis and other disorders of gallbladder (K80-K82)	2	1.3	---
...	All other causes (Residual)	28	18.3	---
American Indian, female, 100 years and over				
...	All causes	53	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	18	34.0	---
2	Cerebrovascular diseases (I60-I69)	6	11.3	---
3	Influenza and pneumonia (J10-J18)	5	9.4	---
4	Alzheimer's disease (G30)	3	5.7	---
5	Anemias (D50-D64)	1	1.9	---
5	Diabetes mellitus (E10-E14)	1	1.9	---
5	Nutritional deficiencies (E40-E64)	1	1.9	---
5	Parkinson's disease (G20-G21)	1	1.9	---
5	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1	1.9	---
5	Acute bronchitis and bronchiolitis (J20-J21)	1	1.9	---
5	Chronic lower respiratory diseases (J40-J47)	1	1.9	---
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	1.9	---
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1	1.9	---
...	All other causes (Residual)	12	22.6	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, all ages				
...	All causes	43,194	100.0	307.7
1	Malignant neoplasms (C00-C97)	11,550	26.7	82.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,477	24.3	74.6
3	Cerebrovascular diseases (I60-I69)	3,543	8.2	25.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,125	4.9	15.1
5	Diabetes mellitus (E10-E14)	1,576	3.6	11.2
6	Influenza and pneumonia (J10-J18)	1,327	3.1	9.5
7	Chronic lower respiratory diseases (J40-J47)	1,302	3.0	9.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	755	1.7	5.4
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	726	1.7	5.2
10	Alzheimer's disease (G30)	615	1.4	4.4
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	582	1.3	4.1
12	Septicemia (A40-A41)	504	1.2	3.6
13	Certain conditions originating in the perinatal period (P00-P96)	463	1.1	3.3
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	428	1.0	3.0
15	Chronic liver disease and cirrhosis (K70,K73-K74)	401	0.9	2.9
...	All other causes (Residual)	6,820	15.8	48.6
Asian or Pacific Islander, both sexes, 1-4 years				
...	All causes	151	100.0	19.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	31	20.5	3.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	20	13.2	2.5
3	Malignant neoplasms (C00-C97)	16	10.6	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	12	7.9	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8	5.3	*
6	Septicemia (A40-A41)	5	3.3	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	3.3	*
8	Influenza and pneumonia (J10-J18)	4	2.6	*
9	Meningitis (G00,G03)	3	2.0	*
9	Cerebrovascular diseases (I60-I69)	3	2.0	*
11	Anemias (D50-D64)	1	0.7	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.7	*
...	All other causes (Residual)	42	27.8	5.3

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 5-9 years				
...	All causes	110	100.0	12.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	34	30.9	3.7
2	Malignant neoplasms (C00-C97)	20	18.2	2.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	10	9.1	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	8	7.3	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	5.5	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	2.7	*
6	Cerebrovascular diseases (I60-I69)	3	2.7	*
6	Influenza and pneumonia (J10-J18)	3	2.7	*
9	Meningococcal infection (A39)	1	0.9	*
9	Septicemia (A40-A41)	1	0.9	*
9	Chronic lower respiratory diseases (J40-J47)	1	0.9	*
9	Pneumonitis due to solids and liquids (J69)	1	0.9	*
...	All other causes (Residual)	19	17.3	*
Asian or Pacific Islander, both sexes, 10-14 years				
...	All causes	129	100.0	13.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	40	31.0	4.3
2	Malignant neoplasms (C00-C97)	17	13.2	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	10	7.8	*
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8	6.2	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	5.4	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	6	4.7	*
7	Cerebrovascular diseases (I60-I69)	4	3.1	*
8	Influenza and pneumonia (J10-J18)	3	2.3	*
8	Chronic lower respiratory diseases (J40-J47)	3	2.3	*
10	Septicemia (A40-A41)	2	1.6	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	1.6	*
12	Diabetes mellitus (E10-E14)	1	0.8	*
12	Diseases of appendix (K35-K38)	1	0.8	*
12	Certain conditions originating in the perinatal period (P00-P96)	1	0.8	*
...	All other causes (Residual)	24	18.6	2.6

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 15-19 years				
...	All causes	314	100.0	33.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	139	44.3	15.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	43	13.7	4.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	37	11.8	4.0
4	Malignant neoplasms (C00-C97)	28	8.9	3.0
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	10	3.2	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	7	2.2	*
7	Cerebrovascular diseases (I60-I69)	5	1.6	*
8	Septicemia (A40-A41)	4	1.3	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	1.3	*
10	Influenza and pneumonia (J10-J18)	3	1.0	*
11	Tuberculosis (A16-A19)	1	0.3	*
11	Anemias (D50-D64)	1	0.3	*
11	Nutritional deficiencies (E40-E64)	1	0.3	*
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1	0.3	*
11	Chronic lower respiratory diseases (J40-J47)	1	0.3	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.3	*
...	All other causes (Residual)	28	8.9	3.0
Asian or Pacific Islander, both sexes, 20-24 years				
...	All causes	504	100.0	49.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	188	37.3	18.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	92	18.3	8.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	70	13.9	6.8
4	Malignant neoplasms (C00-C97)	45	8.9	4.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	29	5.8	2.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	1.4	*
7	Influenza and pneumonia (J10-J18)	3	0.6	*
8	Tuberculosis (A16-A19)	2	0.4	*
8	Septicemia (A40-A41)	2	0.4	*
8	Viral hepatitis (B15-B19)	2	0.4	*
8	Anemias (D50-D64)	2	0.4	*
8	Meningitis (G00,G03)	2	0.4	*
8	Cerebrovascular diseases (I60-I69)	2	0.4	*
8	Pregnancy, childbirth and the puerperium (O00-099)	2	0.4	*
15	Diabetes mellitus (E10-E14)	1	0.2	*
15	Chronic lower respiratory diseases (J40-J47)	1	0.2	*
15	Pneumonitis due to solids and liquids (J69)	1	0.2	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
...	All other causes (Residual)	51	10.1	5.0

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 25-29 years				
...	All causes	478	100.0	40.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	130	27.2	11.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	100	20.9	8.4
3	Malignant neoplasms (C00-C97)	66	13.8	5.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	44	9.2	3.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	33	6.9	2.8
6	Pregnancy, childbirth and the puerperium (O00-099)	10	2.1	*
7	Cerebrovascular diseases (I60-I69)	9	1.9	*
8	Influenza and pneumonia (J10-J18)	5	1.0	*
9	Septicemia (A40-A41)	4	0.8	*
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	0.8	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.6	*
11	Diabetes mellitus (E10-E14)	3	0.6	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.4	*
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2	0.4	*
13	Aortic aneurysm and dissection (I71)	2	0.4	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
...	All other causes (Residual)	59	12.3	5.0
Asian or Pacific Islander, both sexes, 30-34 years				
...	All causes	588	100.0	43.2
1	Malignant neoplasms (C00-C97)	124	21.1	9.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	119	20.2	8.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	85	14.5	6.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	64	10.9	4.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	35	6.0	2.6
6	Cerebrovascular diseases (I60-I69)	14	2.4	*
7	Chronic liver disease and cirrhosis (K70,K73-K74)	10	1.7	*
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	1.5	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	6	1.0	*
9	Diabetes mellitus (E10-E14)	6	1.0	*
9	Aortic aneurysm and dissection (I71)	6	1.0	*
12	Chronic lower respiratory diseases (J40-J47)	5	0.9	*
12	Pregnancy, childbirth and the puerperium (O00-099)	5	0.9	*
14	Septicemia (A40-A41)	4	0.7	*
14	Influenza and pneumonia (J10-J18)	4	0.7	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.7	*
...	All other causes (Residual)	88	15.0	6.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 35-39 years				
...	All causes	698	100.0	56.3
1	Malignant neoplasms (C00-C97)	188	26.9	15.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	107	15.3	8.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	88	12.6	7.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	69	9.9	5.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	33	4.7	2.7
6	Cerebrovascular diseases (I60-I69)	31	4.4	2.5
7	Human immunodeficiency virus (HIV) disease (B20-B24)	17	2.4	*
8	Diabetes mellitus (E10-E14)	13	1.9	*
9	Chronic liver disease and cirrhosis (K70,K73-K74)	12	1.7	*
10	Pregnancy, childbirth and the puerperium (O00-099)	9	1.3	*
11	Influenza and pneumonia (J10-J18)	8	1.1	*
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	5	0.7	*
12	Aortic aneurysm and dissection (I71)	5	0.7	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.7	*
15	Septicemia (A40-A41)	4	0.6	*
15	Chronic lower respiratory diseases (J40-J47)	4	0.6	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	0.6	*
15	Legal intervention (Y35,Y89.0)	4	0.6	*
...	All other causes (Residual)	92	13.2	7.4
Asian or Pacific Islander, both sexes, 40-44 years				
...	All causes	1,094	100.0	96.2
1	Malignant neoplasms (C00-C97)	350	32.0	30.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	190	17.4	16.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	127	11.6	11.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	61	5.6	5.4
5	Cerebrovascular diseases (I60-I69)	58	5.3	5.1
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	47	4.3	4.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	28	2.6	2.5
8	Diabetes mellitus (E10-E14)	22	2.0	1.9
9	Human immunodeficiency virus (HIV) disease (B20-B24)	16	1.5	*
10	Aortic aneurysm and dissection (I71)	12	1.1	*
11	Viral hepatitis (B15-B19)	10	0.9	*
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	8	0.7	*
12	Influenza and pneumonia (J10-J18)	8	0.7	*
12	Chronic lower respiratory diseases (J40-J47)	8	0.7	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.7	*
...	All other causes (Residual)	141	12.9	12.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 45-49 years				
...	All causes	1,509	100.0	146.7
1	Malignant neoplasms (C00-C97)	599	39.7	58.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	261	17.3	25.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	123	8.2	12.0
4	Cerebrovascular diseases (I60-I69)	87	5.8	8.5
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	67	4.4	6.5
6	Diabetes mellitus (E10-E14)	34	2.3	3.3
6	Chronic liver disease and cirrhosis (K70,K73-K74)	34	2.3	3.3
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	28	1.9	2.7
9	Viral hepatitis (B15-B19)	21	1.4	2.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	1.0	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	14	0.9	*
11	Aortic aneurysm and dissection (I71)	14	0.9	*
13	Influenza and pneumonia (J10-J18)	12	0.8	*
13	Chronic lower respiratory diseases (J40-J47)	12	0.8	*
15	Septicemia (A40-A41)	9	0.6	*
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	9	0.6	*
...	All other causes (Residual)	170	11.3	16.5
Asian or Pacific Islander, both sexes, 50-54 years				
...	All causes	2,121	100.0	239.6
1	Malignant neoplasms (C00-C97)	848	40.0	95.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	422	19.9	47.7
3	Cerebrovascular diseases (I60-I69)	143	6.7	16.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	134	6.3	15.1
5	Diabetes mellitus (E10-E14)	70	3.3	7.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	53	2.5	6.0
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	52	2.5	5.9
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	24	1.1	2.7
9	Influenza and pneumonia (J10-J18)	22	1.0	2.5
10	Septicemia (A40-A41)	21	1.0	2.4
10	Chronic lower respiratory diseases (J40-J47)	21	1.0	2.4
12	Viral hepatitis (B15-B19)	20	0.9	2.3
12	Aortic aneurysm and dissection (I71)	20	0.9	2.3
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18	0.8	*
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	13	0.6	*
...	All other causes (Residual)	240	11.3	27.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 55-59 years				
...	All causes	2,535	100.0	348.9
1	Malignant neoplasms (C00-C97)	1,065	42.0	146.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	558	22.0	76.8
3	Cerebrovascular diseases (I60-I69)	157	6.2	21.6
4	Diabetes mellitus (E10-E14)	109	4.3	15.0
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	101	4.0	13.9
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	44	1.7	6.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	39	1.5	5.4
8	Chronic liver disease and cirrhosis (K70,K73-K74)	33	1.3	4.5
9	Chronic lower respiratory diseases (J40-J47)	31	1.2	4.3
10	Viral hepatitis (B15-B19)	28	1.1	3.9
11	Septicemia (A40-A41)	26	1.0	3.6
12	Influenza and pneumonia (J10-J18)	25	1.0	3.4
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	22	0.9	3.0
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	20	0.8	2.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	0.6	*
...	All other causes (Residual)	261	10.3	35.9
Asian or Pacific Islander, both sexes, 60-64 years				
...	All causes	2,892	100.0	578.2
1	Malignant neoplasms (C00-C97)	1,196	41.4	239.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	643	22.2	128.6
3	Cerebrovascular diseases (I60-I69)	212	7.3	42.4
4	Diabetes mellitus (E10-E14)	117	4.0	23.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	114	3.9	22.8
6	Chronic liver disease and cirrhosis (K70,K73-K74)	49	1.7	9.8
7	Chronic lower respiratory diseases (J40-J47)	46	1.6	9.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	46	1.6	9.2
9	Influenza and pneumonia (J10-J18)	43	1.5	8.6
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	31	1.1	6.2
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	23	0.8	4.6
12	Septicemia (A40-A41)	22	0.8	4.4
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	22	0.8	4.4
12	Aortic aneurysm and dissection (I71)	22	0.8	4.4
15	Viral hepatitis (B15-B19)	19	0.7	*
...	All other causes (Residual)	287	9.9	57.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 65-69 years				
...	All causes	3,464	100.0	889.1
1	Malignant neoplasms (C00-C97)	1,271	36.7	326.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	836	24.1	214.6
3	Cerebrovascular diseases (I60-I69)	280	8.1	71.9
4	Diabetes mellitus (E10-E14)	175	5.1	44.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	97	2.8	24.9
6	Chronic lower respiratory diseases (J40-J47)	89	2.6	22.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	70	2.0	18.0
8	Influenza and pneumonia (J10-J18)	53	1.5	13.6
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	41	1.2	10.5
10	Septicemia (A40-A41)	40	1.2	10.3
11	Chronic liver disease and cirrhosis (K70,K73-K74)	39	1.1	10.0
12	Aortic aneurysm and dissection (I71)	27	0.8	6.9
13	Viral hepatitis (B15-B19)	26	0.8	6.7
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	25	0.7	6.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	0.7	5.9
...	All other causes (Residual)	372	10.7	95.5
Asian or Pacific Islander, both sexes, 70-74 years				
...	All causes	4,324	100.0	1,440.6
1	Malignant neoplasms (C00-C97)	1,500	34.7	499.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,063	24.6	354.2
3	Cerebrovascular diseases (I60-I69)	353	8.2	117.6
4	Diabetes mellitus (E10-E14)	170	3.9	56.6
5	Chronic lower respiratory diseases (J40-J47)	142	3.3	47.3
6	Influenza and pneumonia (J10-J18)	112	2.6	37.3
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	109	2.5	36.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	86	2.0	28.7
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	57	1.3	19.0
10	Septicemia (A40-A41)	52	1.2	17.3
11	Viral hepatitis (B15-B19)	38	0.9	12.7
12	Aortic aneurysm and dissection (I71)	36	0.8	12.0
13	Parkinson's disease (G20-G21)	32	0.7	10.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	29	0.7	9.7
15	Chronic liver disease and cirrhosis (K70,K73-K74)	28	0.6	9.3
...	All other causes (Residual)	517	12.0	172.3

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 75-79 years				
...	All causes	5,356	100.0	2,354.3
1	Malignant neoplasms (C00-C97)	1,527	28.5	671.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,315	24.6	578.0
3	Cerebrovascular diseases (I60-I69)	500	9.3	219.8
4	Diabetes mellitus (E10-E14)	264	4.9	116.0
5	Chronic lower respiratory diseases (J40-J47)	218	4.1	95.8
6	Influenza and pneumonia (J10-J18)	162	3.0	71.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	124	2.3	54.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	109	2.0	47.9
9	Septicemia (A40-A41)	87	1.6	38.2
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	85	1.6	37.4
11	Alzheimer's disease (G30)	68	1.3	29.9
12	Parkinson's disease (G20-G21)	60	1.1	26.4
13	Chronic liver disease and cirrhosis (K70,K73-K74)	54	1.0	23.7
14	Aortic aneurysm and dissection (I71)	53	1.0	23.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	37	0.7	16.3
...	All other causes (Residual)	693	12.9	304.6
Asian or Pacific Islander, both sexes, 80-84 years				
...	All causes	6,058	100.0	4,071.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,719	28.4	1,155.4
2	Malignant neoplasms (C00-C97)	1,349	22.3	906.7
3	Cerebrovascular diseases (I60-I69)	618	10.2	415.4
4	Chronic lower respiratory diseases (J40-J47)	280	4.6	188.2
5	Diabetes mellitus (E10-E14)	250	4.1	168.0
6	Influenza and pneumonia (J10-J18)	242	4.0	162.7
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	166	2.7	111.6
8	Alzheimer's disease (G30)	134	2.2	90.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	134	2.2	90.1
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	94	1.6	63.2
11	Septicemia (A40-A41)	75	1.2	50.4
12	Parkinson's disease (G20-G21)	64	1.1	43.0
13	Aortic aneurysm and dissection (I71)	59	1.0	39.7
14	Pneumonitis due to solids and liquids (J69)	52	0.9	35.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	41	0.7	27.6
...	All other causes (Residual)	781	12.9	524.9

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 85-89 years				
...	All causes	5,179	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,532	29.6	---
2	Malignant neoplasms (C00-C97)	879	17.0	---
3	Cerebrovascular diseases (I60-I69)	561	10.8	---
4	Influenza and pneumonia (J10-J18)	263	5.1	---
5	Chronic lower respiratory diseases (J40-J47)	253	4.9	---
6	Diabetes mellitus (E10-E14)	205	4.0	---
7	Alzheimer's disease (G30)	169	3.3	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	132	2.5	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	110	2.1	---
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	92	1.8	---
11	Septicemia (A40-A41)	75	1.4	---
12	Pneumonitis due to solids and liquids (J69)	62	1.2	---
13	Parkinson's disease (G20-G21)	52	1.0	---
14	Aortic aneurysm and dissection (I71)	42	0.8	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	26	0.5	---
...	All other causes (Residual)	726	14.0	---
Asian or Pacific Islander, both sexes, 90-94 years				
...	All causes	3,312	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,107	33.4	---
2	Malignant neoplasms (C00-C97)	370	11.2	---
3	Cerebrovascular diseases (I60-I69)	348	10.5	---
4	Influenza and pneumonia (J10-J18)	228	6.9	---
5	Chronic lower respiratory diseases (J40-J47)	135	4.1	---
6	Alzheimer's disease (G30)	126	3.8	---
7	Diabetes mellitus (E10-E14)	114	3.4	---
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	86	2.6	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	69	2.1	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	68	2.1	---
11	Pneumonitis due to solids and liquids (J69)	44	1.3	---
12	Septicemia (A40-A41)	41	1.2	---
13	Parkinson's disease (G20-G21)	29	0.9	---
14	Aortic aneurysm and dissection (I71)	26	0.8	---
15	Atherosclerosis (I70)	18	0.5	---
...	All other causes (Residual)	503	15.2	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 95-99 years				
...	All causes	1,245	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	467	37.5	---
2	Cerebrovascular diseases (I60-I69)	130	10.4	---
3	Influenza and pneumonia (J10-J18)	103	8.3	---
4	Malignant neoplasms (C00-C97)	77	6.2	---
5	Alzheimer's disease (G30)	72	5.8	---
6	Chronic lower respiratory diseases (J40-J47)	43	3.5	---
7	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	35	2.8	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	31	2.5	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	21	1.7	---
10	Diabetes mellitus (E10-E14)	17	1.4	---
10	Pneumonitis due to solids and liquids (J69)	17	1.4	---
12	Septicemia (A40-A41)	12	1.0	---
13	Aortic aneurysm and dissection (I71)	8	0.6	---
13	Peptic ulcer (K25-K28)	8	0.6	---
15	Anemias (D50-D64)	4	0.3	---
15	Atherosclerosis (I70)	4	0.3	---
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	4	0.3	---
...	All other causes (Residual)	192	15.4	---
Asian or Pacific Islander, both sexes, 100 years and over				
...	All causes	258	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	98	38.0	---
2	Cerebrovascular diseases (I60-I69)	20	7.8	---
3	Influenza and pneumonia (J10-J18)	17	6.6	---
4	Malignant neoplasms (C00-C97)	14	5.4	---
5	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	9	3.5	---
6	Chronic lower respiratory diseases (J40-J47)	8	3.1	---
7	Atherosclerosis (I70)	6	2.3	---
8	Diabetes mellitus (E10-E14)	5	1.9	---
8	Alzheimer's disease (G30)	5	1.9	---
8	Pneumonitis due to solids and liquids (J69)	5	1.9	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	1.9	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5	1.9	---
13	Septicemia (A40-A41)	3	1.2	---
14	Parkinson's disease (G20-G21)	2	0.8	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.4	---
15	Nutritional deficiencies (E40-E64)	1	0.4	---
15	Peptic ulcer (K25-K28)	1	0.4	---
15	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.4	---
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.4	---
...	All other causes (Residual)	51	19.8	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, all ages				
...	All causes	22,808	100.0	333.9
1	Malignant neoplasms (C00-C97)	5,921	26.0	86.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,703	25.0	83.5
3	Cerebrovascular diseases (I60-I69)	1,624	7.1	23.8
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,328	5.8	19.4
5	Chronic lower respiratory diseases (J40-J47)	803	3.5	11.8
6	Diabetes mellitus (E10-E14)	768	3.4	11.2
7	Influenza and pneumonia (J10-J18)	718	3.1	10.5
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	495	2.2	7.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	381	1.7	5.6
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	314	1.4	4.6
11	Septicemia (A40-A41)	260	1.1	3.8
12	Certain conditions originating in the perinatal period (P00-P96)	258	1.1	3.8
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	256	1.1	3.7
14	Chronic liver disease and cirrhosis (K70,K73-K74)	254	1.1	3.7
15	Alzheimer's disease (G30)	229	1.0	3.4
...	All other causes (Residual)	3,496	15.3	51.2
Asian or Pacific Islander, male, 1-4 years				
...	All causes	84	100.0	20.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19	22.6	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	11	13.1	*
3	Malignant neoplasms (C00-C97)	9	10.7	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	6	7.1	*
5	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	3.6	*
5	Meningitis (G00,G03)	3	3.6	*
5	Cerebrovascular diseases (I60-I69)	3	3.6	*
5	Influenza and pneumonia (J10-J18)	3	3.6	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	3.6	*
10	Septicemia (A40-A41)	1	1.2	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	1	1.2	*
...	All other causes (Residual)	22	26.2	5.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 5-9 years				
...	All causes	59	100.0	12.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	20	33.9	4.3
2	Malignant neoplasms (C00-C97)	12	20.3	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	6.8	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	5.1	*
5	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	3.4	*
5	Influenza and pneumonia (J10-J18)	2	3.4	*
7	Meningococcal infection (A39)	1	1.7	*
7	Septicemia (A40-A41)	1	1.7	*
7	Cerebrovascular diseases (I60-I69)	1	1.7	*
7	Chronic lower respiratory diseases (J40-J47)	1	1.7	*
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1	1.7	*
...	All other causes (Residual)	11	18.6	*
Asian or Pacific Islander, male, 10-14 years				
...	All causes	72	100.0	15.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	22	30.6	4.6
2	Malignant neoplasms (C00-C97)	9	12.5	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7	9.7	*
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6	8.3	*
5	Cerebrovascular diseases (I60-I69)	4	5.6	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	5.6	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	2.8	*
8	Septicemia (A40-A41)	1	1.4	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.4	*
8	Diabetes mellitus (E10-E14)	1	1.4	*
8	Influenza and pneumonia (J10-J18)	1	1.4	*
8	Chronic lower respiratory diseases (J40-J47)	1	1.4	*
...	All other causes (Residual)	13	18.1	*

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 15-19 years				
...	All causes	218	100.0	45.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	99	45.4	20.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	37	17.0	7.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	23	10.6	4.8
4	Malignant neoplasms (C00-C97)	19	8.7	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	3.2	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	1.8	*
6	Cerebrovascular diseases (I60-I69)	4	1.8	*
8	Influenza and pneumonia (J10-J18)	3	1.4	*
9	Tuberculosis (A16-A19)	1	0.5	*
9	Septicemia (A40-A41)	1	0.5	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.5	*
9	Chronic lower respiratory diseases (J40-J47)	1	0.5	*
9	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.5	*
...	All other causes (Residual)	17	7.8	*
Asian or Pacific Islander, male, 20-24 years				
...	All causes	351	100.0	67.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	136	38.7	26.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	71	20.2	13.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	49	14.0	9.4
4	Malignant neoplasms (C00-C97)	25	7.1	4.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	21	6.0	4.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.9	*
7	Anemias (D50-D64)	2	0.6	*
8	Tuberculosis (A16-A19)	1	0.3	*
8	Septicemia (A40-A41)	1	0.3	*
8	Viral hepatitis (B15-B19)	1	0.3	*
8	Diabetes mellitus (E10-E14)	1	0.3	*
8	Meningitis (G00,G03)	1	0.3	*
8	Influenza and pneumonia (J10-J18)	1	0.3	*
8	Pneumonitis due to solids and liquids (J69)	1	0.3	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
...	All other causes (Residual)	36	10.3	6.9

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 25-29 years				
...	All causes	321	100.0	54.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	93	29.0	15.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	76	23.7	13.0
3	Malignant neoplasms (C00-C97)	38	11.8	6.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	33	10.3	5.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	23	7.2	3.9
6	Cerebrovascular diseases (I60-I69)	4	1.2	*
7	Influenza and pneumonia (J10-J18)	3	0.9	*
8	Septicemia (A40-A41)	2	0.6	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.6	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.6	*
8	Diabetes mellitus (E10-E14)	2	0.6	*
8	Aortic aneurysm and dissection (I71)	2	0.6	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.6	*
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	0.6	*
15	Tuberculosis (A16-A19)	1	0.3	*
15	Viral hepatitis (B15-B19)	1	0.3	*
15	Nutritional deficiencies (E40-E64)	1	0.3	*
15	Meningitis (G00,G03)	1	0.3	*
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1	0.3	*
15	Chronic lower respiratory diseases (J40-J47)	1	0.3	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.3	*
15	Legal intervention (Y35,Y89.0)	1	0.3	*
...	All other causes (Residual)	29	9.0	5.0
Asian or Pacific Islander, male, 30-34 years				
...	All causes	375	100.0	56.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	87	23.2	13.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	68	18.1	10.2
3	Malignant neoplasms (C00-C97)	52	13.9	7.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	48	12.8	7.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	24	6.4	3.6
6	Cerebrovascular diseases (I60-I69)	9	2.4	*
7	Chronic liver disease and cirrhosis (K70,K73-K74)	7	1.9	*
8	Aortic aneurysm and dissection (I71)	6	1.6	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	5	1.3	*
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	1.3	*
11	Diabetes mellitus (E10-E14)	4	1.1	*
12	Chronic lower respiratory diseases (J40-J47)	3	0.8	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.8	*
14	Septicemia (A40-A41)	2	0.5	*
14	Viral hepatitis (B15-B19)	2	0.5	*
14	Influenza and pneumonia (J10-J18)	2	0.5	*
...	All other causes (Residual)	48	12.8	7.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 35-39 years				
...	All causes	427	100.0	70.5
1	Malignant neoplasms (C00-C97)	84	19.7	13.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	72	16.9	11.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	66	15.5	10.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	47	11.0	7.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	25	5.9	4.1
6	Cerebrovascular diseases (I60-I69)	18	4.2	*
7	Human immunodeficiency virus (HIV) disease (B20-B24)	14	3.3	*
8	Chronic liver disease and cirrhosis (K70,K73-K74)	10	2.3	*
9	Influenza and pneumonia (J10-J18)	5	1.2	*
10	Diabetes mellitus (E10-E14)	4	0.9	*
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	4	0.9	*
10	Aortic aneurysm and dissection (I71)	4	0.9	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.9	*
10	Legal intervention (Y35,Y89.0)	4	0.9	*
15	Septicemia (A40-A41)	3	0.7	*
15	Chronic lower respiratory diseases (J40-J47)	3	0.7	*
...	All other causes (Residual)	60	14.1	9.9
Asian or Pacific Islander, male, 40-44 years				
...	All causes	655	100.0	118.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	148	22.6	26.9
2	Malignant neoplasms (C00-C97)	147	22.4	26.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	88	13.4	16.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	39	6.0	7.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	35	5.3	6.4
6	Cerebrovascular diseases (I60-I69)	28	4.3	5.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	23	3.5	4.2
8	Diabetes mellitus (E10-E14)	17	2.6	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	13	2.0	*
10	Aortic aneurysm and dissection (I71)	11	1.7	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	1.1	*
12	Viral hepatitis (B15-B19)	6	0.9	*
12	Chronic lower respiratory diseases (J40-J47)	6	0.9	*
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	5	0.8	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.6	*
...	All other causes (Residual)	78	11.9	14.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 45-49 years				
...	All causes	873	100.0	179.0
1	Malignant neoplasms (C00-C97)	259	29.7	53.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	206	23.6	42.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	85	9.7	17.4
4	Cerebrovascular diseases (I60-I69)	44	5.0	9.0
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	43	4.9	8.8
6	Chronic liver disease and cirrhosis (K70,K73-K74)	30	3.4	6.2
7	Diabetes mellitus (E10-E14)	21	2.4	4.3
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	20	2.3	4.1
9	Viral hepatitis (B15-B19)	13	1.5	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	12	1.4	*
11	Aortic aneurysm and dissection (I71)	11	1.3	*
12	Influenza and pneumonia (J10-J18)	8	0.9	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.9	*
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	7	0.8	*
15	Chronic lower respiratory diseases (J40-J47)	5	0.6	*
...	All other causes (Residual)	101	11.6	20.7
Asian or Pacific Islander, male, 50-54 years				
...	All causes	1,309	100.0	317.5
1	Malignant neoplasms (C00-C97)	424	32.4	102.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	317	24.2	76.9
3	Cerebrovascular diseases (I60-I69)	86	6.6	20.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	86	6.6	20.9
5	Diabetes mellitus (E10-E14)	48	3.7	11.6
6	Chronic liver disease and cirrhosis (K70,K73-K74)	41	3.1	9.9
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	38	2.9	9.2
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	20	1.5	4.9
9	Viral hepatitis (B15-B19)	17	1.3	*
9	Aortic aneurysm and dissection (I71)	17	1.3	*
11	Chronic lower respiratory diseases (J40-J47)	16	1.2	*
12	Influenza and pneumonia (J10-J18)	14	1.1	*
13	Septicemia (A40-A41)	13	1.0	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	0.9	*
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	10	0.8	*
...	All other causes (Residual)	150	11.5	36.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 55-59 years				
...	All causes	1,483	100.0	441.1
1	Malignant neoplasms (C00-C97)	524	35.3	155.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	411	27.7	122.2
3	Cerebrovascular diseases (I60-I69)	85	5.7	25.3
4	Diabetes mellitus (E10-E14)	67	4.5	19.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	61	4.1	18.1
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	30	2.0	8.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	26	1.8	7.7
8	Chronic lower respiratory diseases (J40-J47)	24	1.6	7.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21	1.4	6.2
10	Influenza and pneumonia (J10-J18)	19	1.3	*
11	Viral hepatitis (B15-B19)	17	1.1	*
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	15	1.0	*
13	Septicemia (A40-A41)	11	0.7	*
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	10	0.7	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.5	*
...	All other causes (Residual)	155	10.5	46.1
Asian or Pacific Islander, male, 60-64 years				
...	All causes	1,622	100.0	695.8
1	Malignant neoplasms (C00-C97)	611	37.7	262.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	416	25.6	178.5
3	Cerebrovascular diseases (I60-I69)	105	6.5	45.0
4	Diabetes mellitus (E10-E14)	77	4.7	33.0
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	68	4.2	29.2
6	Chronic lower respiratory diseases (J40-J47)	30	1.8	12.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	30	1.8	12.9
8	Influenza and pneumonia (J10-J18)	23	1.4	9.9
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	20	1.2	8.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19	1.2	*
11	Viral hepatitis (B15-B19)	15	0.9	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	0.9	*
12	Aortic aneurysm and dissection (I71)	14	0.9	*
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	10	0.6	*
15	Septicemia (A40-A41)	9	0.6	*
...	All other causes (Residual)	161	9.9	69.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 65-69 years				
...	All causes	2,009	100.0	1,115.6
1	Malignant neoplasms (C00-C97)	714	35.5	396.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	550	27.4	305.4
3	Cerebrovascular diseases (I60-I69)	151	7.5	83.8
4	Diabetes mellitus (E10-E14)	99	4.9	55.0
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	56	2.8	31.1
6	Chronic lower respiratory diseases (J40-J47)	55	2.7	30.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	34	1.7	18.9
8	Influenza and pneumonia (J10-J18)	30	1.5	16.7
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	25	1.2	13.9
10	Chronic liver disease and cirrhosis (K70,K73-K74)	23	1.1	12.8
11	Aortic aneurysm and dissection (I71)	21	1.0	11.7
12	Septicemia (A40-A41)	18	0.9	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	0.8	*
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	16	0.8	*
15	Parkinson's disease (G20-G21)	14	0.7	*
...	All other causes (Residual)	186	9.3	103.3
Asian or Pacific Islander, male, 70-74 years				
...	All causes	2,335	100.0	1,805.1
1	Malignant neoplasms (C00-C97)	809	34.6	625.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	610	26.1	471.6
3	Cerebrovascular diseases (I60-I69)	174	7.5	134.5
4	Chronic lower respiratory diseases (J40-J47)	85	3.6	65.7
5	Diabetes mellitus (E10-E14)	83	3.6	64.2
6	Influenza and pneumonia (J10-J18)	68	2.9	52.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	62	2.7	47.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	48	2.1	37.1
9	Septicemia (A40-A41)	31	1.3	24.0
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	25	1.1	19.3
11	Aortic aneurysm and dissection (I71)	24	1.0	18.6
12	Parkinson's disease (G20-G21)	17	0.7	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	0.6	*
14	Viral hepatitis (B15-B19)	13	0.6	*
15	Alzheimer's disease (G30)	12	0.5	*
...	All other causes (Residual)	260	11.1	201.0

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 75-79 years				
...	All causes	2,715	100.0	2,918.7
1	Malignant neoplasms (C00-C97)	813	29.9	874.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	672	24.8	722.4
3	Cerebrovascular diseases (I60-I69)	232	8.5	249.4
4	Chronic lower respiratory diseases (J40-J47)	133	4.9	143.0
5	Diabetes mellitus (E10-E14)	104	3.8	111.8
6	Influenza and pneumonia (J10-J18)	92	3.4	98.9
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	58	2.1	62.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	55	2.0	59.1
9	Septicemia (A40-A41)	46	1.7	49.5
10	Parkinson's disease (G20-G21)	35	1.3	37.6
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	35	1.3	37.6
12	Alzheimer's disease (G30)	33	1.2	35.5
13	Aortic aneurysm and dissection (I71)	31	1.1	33.3
14	Chronic liver disease and cirrhosis (K70,K73-K74)	23	0.8	24.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	0.7	21.5
...	All other causes (Residual)	333	12.3	358.0
Asian or Pacific Islander, male, 80-84 years				
...	All causes	3,079	100.0	5,038.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	845	27.4	1,382.8
2	Malignant neoplasms (C00-C97)	715	23.2	1,170.1
3	Cerebrovascular diseases (I60-I69)	288	9.4	471.3
4	Chronic lower respiratory diseases (J40-J47)	186	6.0	304.4
5	Influenza and pneumonia (J10-J18)	142	4.6	232.4
6	Diabetes mellitus (E10-E14)	105	3.4	171.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	94	3.1	153.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	66	2.1	108.0
9	Alzheimer's disease (G30)	56	1.8	91.6
10	Septicemia (A40-A41)	42	1.4	68.7
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	41	1.3	67.1
12	Aortic aneurysm and dissection (I71)	34	1.1	55.6
13	Parkinson's disease (G20-G21)	33	1.1	54.0
14	Pneumonitis due to solids and liquids (J69)	30	1.0	49.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	21	0.7	34.4
...	All other causes (Residual)	381	12.4	623.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 85-89 years				
...	All causes	2,382	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	683	28.7	---
2	Malignant neoplasms (C00-C97)	436	18.3	---
3	Cerebrovascular diseases (I60-I69)	213	8.9	---
4	Chronic lower respiratory diseases (J40-J47)	151	6.3	---
5	Influenza and pneumonia (J10-J18)	137	5.8	---
6	Diabetes mellitus (E10-E14)	85	3.6	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	70	2.9	---
8	Alzheimer's disease (G30)	61	2.6	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	54	2.3	---
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	38	1.6	---
11	Pneumonitis due to solids and liquids (J69)	36	1.5	---
12	Septicemia (A40-A41)	34	1.4	---
13	Aortic aneurysm and dissection (I71)	27	1.1	---
14	Parkinson's disease (G20-G21)	26	1.1	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	0.5	---
...	All other causes (Residual)	318	13.4	---
Asian or Pacific Islander, male, 90-94 years				
...	All causes	1,370	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	449	32.8	---
2	Malignant neoplasms (C00-C97)	177	12.9	---
3	Cerebrovascular diseases (I60-I69)	126	9.2	---
4	Influenza and pneumonia (J10-J18)	107	7.8	---
5	Chronic lower respiratory diseases (J40-J47)	69	5.0	---
6	Diabetes mellitus (E10-E14)	43	3.1	---
7	Alzheimer's disease (G30)	36	2.6	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	31	2.3	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	31	2.3	---
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	27	2.0	---
11	Septicemia (A40-A41)	25	1.8	---
12	Pneumonitis due to solids and liquids (J69)	19	1.4	---
13	Parkinson's disease (G20-G21)	18	1.3	---
14	Aortic aneurysm and dissection (I71)	11	0.8	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.6	---
...	All other causes (Residual)	193	14.1	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 95-99 years				
...	All causes	491	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	179	36.5	---
2	Influenza and pneumonia (J10-J18)	45	9.2	---
3	Cerebrovascular diseases (I60-I69)	42	8.6	---
4	Malignant neoplasms (C00-C97)	36	7.3	---
5	Chronic lower respiratory diseases (J40-J47)	28	5.7	---
6	Alzheimer's disease (G30)	25	5.1	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	3.1	---
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	14	2.9	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12	2.4	---
10	Pneumonitis due to solids and liquids (J69)	10	2.0	---
11	Diabetes mellitus (E10-E14)	6	1.2	---
11	Aortic aneurysm and dissection (I71)	6	1.2	---
13	Septicemia (A40-A41)	5	1.0	---
14	Peptic ulcer (K25-K28)	3	0.6	---
15	Atherosclerosis (I70)	2	0.4	---
15	Hyperplasia of prostate (N40)	2	0.4	---
...	All other causes (Residual)	61	12.4	---
Asian or Pacific Islander, male, 100 years and over				
...	All causes	93	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	34	36.6	---
2	Malignant neoplasms (C00-C97)	7	7.5	---
2	Influenza and pneumonia (J10-J18)	7	7.5	---
4	Chronic lower respiratory diseases (J40-J47)	5	5.4	---
5	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	4	4.3	---
5	Cerebrovascular diseases (I60-I69)	4	4.3	---
7	Atherosclerosis (I70)	2	2.2	---
7	Pneumonitis due to solids and liquids (J69)	2	2.2	---
9	Septicemia (A40-A41)	1	1.1	---
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.1	---
9	Diabetes mellitus (E10-E14)	1	1.1	---
9	Parkinson's disease (G20-G21)	1	1.1	---
9	Peptic ulcer (K25-K28)	1	1.1	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	1.1	---
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	1.1	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1	1.1	---
...	All other causes (Residual)	20	21.5	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, all ages				
...	All causes	20,386	100.0	282.8
1	Malignant neoplasms (C00-C97)	5,629	27.6	78.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,774	23.4	66.2
3	Cerebrovascular diseases (I60-I69)	1,919	9.4	26.6
4	Diabetes mellitus (E10-E14)	808	4.0	11.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	797	3.9	11.1
6	Influenza and pneumonia (J10-J18)	609	3.0	8.4
7	Chronic lower respiratory diseases (J40-J47)	499	2.4	6.9
8	Alzheimer's disease (G30)	386	1.9	5.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	374	1.8	5.2
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	326	1.6	4.5
11	Septicemia (A40-A41)	244	1.2	3.4
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	231	1.1	3.2
13	Certain conditions originating in the perinatal period (P00-P96)	205	1.0	2.8
14	Chronic liver disease and cirrhosis (K70,K73-K74)	147	0.7	2.0
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	146	0.7	2.0
...	All other causes (Residual)	3,292	16.1	45.7
Asian or Pacific Islander, female, 1-4 years				
...	All causes	67	100.0	17.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12	17.9	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	13.4	*
3	Malignant neoplasms (C00-C97)	7	10.4	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	6	9.0	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	7.5	*
6	Septicemia (A40-A41)	4	6.0	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	3.0	*
8	Anemias (D50-D64)	1	1.5	*
8	Influenza and pneumonia (J10-J18)	1	1.5	*
...	All other causes (Residual)	20	29.9	5.2
Asian or Pacific Islander, female, 5-9 years				
...	All causes	51	100.0	11.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14	27.5	*
2	Malignant neoplasms (C00-C97)	8	15.7	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	11.8	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	9.8	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	9.8	*
6	Cerebrovascular diseases (I60-I69)	2	3.9	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	2.0	*
7	Influenza and pneumonia (J10-J18)	1	2.0	*
7	Pneumonitis due to solids and liquids (J69)	1	2.0	*
...	All other causes (Residual)	8	15.7	*

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 10-14 years				
...	All causes	57	100.0	12.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18	31.6	*
2	Malignant neoplasms (C00-C97)	8	14.0	*
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	7.0	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	5.3	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	5.3	*
6	Influenza and pneumonia (J10-J18)	2	3.5	*
6	Chronic lower respiratory diseases (J40-J47)	2	3.5	*
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2	3.5	*
9	Septicemia (A40-A41)	1	1.8	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.8	*
9	Diseases of appendix (K35-K38)	1	1.8	*
9	Certain conditions originating in the perinatal period (P00-P96)	1	1.8	*
...	All other causes (Residual)	11	19.3	*
Asian or Pacific Islander, female, 15-19 years				
...	All causes	96	100.0	21.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	40	41.7	8.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	14	14.6	*
3	Malignant neoplasms (C00-C97)	9	9.4	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	6.3	*
5	Septicemia (A40-A41)	3	3.1	*
5	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	3.1	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	3.1	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	3.1	*
9	Anemias (D50-D64)	1	1.0	*
9	Nutritional deficiencies (E40-E64)	1	1.0	*
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1	1.0	*
9	Cerebrovascular diseases (I60-I69)	1	1.0	*
...	All other causes (Residual)	11	11.5	*
Asian or Pacific Islander, female, 20-24 years				
...	All causes	153	100.0	30.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	52	34.0	10.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	21	13.7	4.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	21	13.7	4.2
4	Malignant neoplasms (C00-C97)	20	13.1	4.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	8	5.2	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	2.6	*
7	Cerebrovascular diseases (I60-I69)	2	1.3	*
7	Influenza and pneumonia (J10-J18)	2	1.3	*
7	Pregnancy, childbirth and the puerperium (O00-099)	2	1.3	*
10	Tuberculosis (A16-A19)	1	0.7	*
10	Septicemia (A40-A41)	1	0.7	*
10	Viral hepatitis (B15-B19)	1	0.7	*
10	Meningitis (G00,G03)	1	0.7	*
10	Chronic lower respiratory diseases (J40-J47)	1	0.7	*
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.7	*
...	All other causes (Residual)	15	9.8	*

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 25-29 years				
...	All causes	157	100.0	26.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	37	23.6	6.2
2	Malignant neoplasms (C00-C97)	28	17.8	4.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	24	15.3	4.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	11	7.0	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	10	6.4	*
5	Pregnancy, childbirth and the puerperium (O00-099)	10	6.4	*
7	Cerebrovascular diseases (I60-I69)	5	3.2	*
8	Septicemia (A40-A41)	2	1.3	*
8	Influenza and pneumonia (J10-J18)	2	1.3	*
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	1.3	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.6	*
11	Anemias (D50-D64)	1	0.6	*
11	Diabetes mellitus (E10-E14)	1	0.6	*
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1	0.6	*
11	Certain conditions originating in the perinatal period (P00-P96)	1	0.6	*
...	All other causes (Residual)	21	13.4	3.5
Asian or Pacific Islander, female, 30-34 years				
...	All causes	213	100.0	30.7
1	Malignant neoplasms (C00-C97)	72	33.8	10.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	32	15.0	4.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	17	8.0	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	16	7.5	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	11	5.2	*
6	Cerebrovascular diseases (I60-I69)	5	2.3	*
6	Pregnancy, childbirth and the puerperium (O00-099)	5	2.3	*
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	1.9	*
9	Chronic liver disease and cirrhosis (K70,K73-K74)	3	1.4	*
10	Septicemia (A40-A41)	2	0.9	*
10	Diabetes mellitus (E10-E14)	2	0.9	*
10	Influenza and pneumonia (J10-J18)	2	0.9	*
10	Chronic lower respiratory diseases (J40-J47)	2	0.9	*
14	Tuberculosis (A16-A19)	1	0.5	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.5	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.5	*
14	Inflammatory diseases of female pelvic organs (N70-N76)	1	0.5	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.5	*
...	All other causes (Residual)	34	16.0	4.9

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 35-39 years				
...	All causes	271	100.0	42.7
1	Malignant neoplasms (C00-C97)	104	38.4	16.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	35	12.9	5.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	22	8.1	3.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	22	8.1	3.5
5	Cerebrovascular diseases (I60-I69)	13	4.8	*
6	Diabetes mellitus (E10-E14)	9	3.3	*
6	Pregnancy, childbirth and the puerperium (O00-099)	9	3.3	*
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8	3.0	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	3	1.1	*
9	Influenza and pneumonia (J10-J18)	3	1.1	*
11	Chronic liver disease and cirrhosis (K70,K73-K74)	2	0.7	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	0.7	*
13	Septicemia (A40-A41)	1	0.4	*
13	Viral hepatitis (B15-B19)	1	0.4	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.4	*
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1	0.4	*
13	Aortic aneurysm and dissection (I71)	1	0.4	*
13	Chronic lower respiratory diseases (J40-J47)	1	0.4	*
13	Pneumonitis due to solids and liquids (J69)	1	0.4	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.4	*
...	All other causes (Residual)	31	11.4	4.9
Asian or Pacific Islander, female, 40-44 years				
...	All causes	439	100.0	74.9
1	Malignant neoplasms (C00-C97)	203	46.2	34.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	42	9.6	7.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	39	8.9	6.7
4	Cerebrovascular diseases (I60-I69)	30	6.8	5.1
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	22	5.0	3.8
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	12	2.7	*
7	Diabetes mellitus (E10-E14)	5	1.1	*
7	Influenza and pneumonia (J10-J18)	5	1.1	*
7	Chronic liver disease and cirrhosis (K70,K73-K74)	5	1.1	*
10	Viral hepatitis (B15-B19)	4	0.9	*
11	Septicemia (A40-A41)	3	0.7	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	3	0.7	*
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	3	0.7	*
11	Pregnancy, childbirth and the puerperium (O00-099)	3	0.7	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.5	*
15	Anemias (D50-D64)	2	0.5	*
15	Chronic lower respiratory diseases (J40-J47)	2	0.5	*
...	All other causes (Residual)	54	12.3	9.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 45-49 years				
...	All causes	636	100.0	117.6
1	Malignant neoplasms (C00-C97)	340	53.5	62.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	55	8.6	10.2
3	Cerebrovascular diseases (I60-I69)	43	6.8	8.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	38	6.0	7.0
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	24	3.8	4.4
6	Diabetes mellitus (E10-E14)	13	2.0	*
7	Viral hepatitis (B15-B19)	8	1.3	*
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8	1.3	*
9	Chronic lower respiratory diseases (J40-J47)	7	1.1	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	1.1	*
11	Septicemia (A40-A41)	5	0.8	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.8	*
13	Influenza and pneumonia (J10-J18)	4	0.6	*
13	Chronic liver disease and cirrhosis (K70,K73-K74)	4	0.6	*
15	Aortic aneurysm and dissection (I71)	3	0.5	*
15	Pregnancy, childbirth and the puerperium (O00-099)	3	0.5	*
...	All other causes (Residual)	69	10.8	12.8
Asian or Pacific Islander, female, 50-54 years				
...	All causes	812	100.0	171.7
1	Malignant neoplasms (C00-C97)	424	52.2	89.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	105	12.9	22.2
3	Cerebrovascular diseases (I60-I69)	57	7.0	12.1
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	48	5.9	10.2
5	Diabetes mellitus (E10-E14)	22	2.7	4.7
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	14	1.7	*
7	Chronic liver disease and cirrhosis (K70,K73-K74)	12	1.5	*
8	Septicemia (A40-A41)	8	1.0	*
8	Influenza and pneumonia (J10-J18)	8	1.0	*
10	Tuberculosis (A16-A19)	6	0.7	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.7	*
12	Chronic lower respiratory diseases (J40-J47)	5	0.6	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.5	*
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	0.5	*
15	Viral hepatitis (B15-B19)	3	0.4	*
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	3	0.4	*
15	Aortic aneurysm and dissection (I71)	3	0.4	*
...	All other causes (Residual)	80	9.9	16.9

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 55-59 years				
...	All causes	1,052	100.0	269.5
1	Malignant neoplasms (C00-C97)	541	51.4	138.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	147	14.0	37.7
3	Cerebrovascular diseases (I60-I69)	72	6.8	18.4
4	Diabetes mellitus (E10-E14)	42	4.0	10.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	40	3.8	10.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18	1.7	*
7	Septicemia (A40-A41)	15	1.4	*
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	14	1.3	*
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	12	1.1	*
10	Viral hepatitis (B15-B19)	11	1.0	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.9	*
12	Chronic lower respiratory diseases (J40-J47)	7	0.7	*
12	Chronic liver disease and cirrhosis (K70,K73-K74)	7	0.7	*
14	Influenza and pneumonia (J10-J18)	6	0.6	*
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	0.5	*
...	All other causes (Residual)	106	10.1	27.2
Asian or Pacific Islander, female, 60-64 years				
...	All causes	1,270	100.0	475.6
1	Malignant neoplasms (C00-C97)	585	46.1	219.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	227	17.9	85.0
3	Cerebrovascular diseases (I60-I69)	107	8.4	40.1
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	46	3.6	17.2
5	Diabetes mellitus (E10-E14)	40	3.1	15.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	27	2.1	10.1
7	Influenza and pneumonia (J10-J18)	20	1.6	7.5
8	Chronic liver disease and cirrhosis (K70,K73-K74)	19	1.5	*
9	Chronic lower respiratory diseases (J40-J47)	16	1.3	*
10	Septicemia (A40-A41)	13	1.0	*
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	13	1.0	*
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11	0.9	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.6	*
13	Aortic aneurysm and dissection (I71)	8	0.6	*
15	Peptic ulcer (K25-K28)	5	0.4	*
...	All other causes (Residual)	125	9.8	46.8

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 65-69 years				
...	All causes	1,455	100.0	694.4
1	Malignant neoplasms (C00-C97)	557	38.3	265.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	286	19.7	136.5
3	Cerebrovascular diseases (I60-I69)	129	8.9	61.6
4	Diabetes mellitus (E10-E14)	76	5.2	36.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	41	2.8	19.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	36	2.5	17.2
7	Chronic lower respiratory diseases (J40-J47)	34	2.3	16.2
8	Influenza and pneumonia (J10-J18)	23	1.6	11.0
9	Septicemia (A40-A41)	22	1.5	10.5
10	Viral hepatitis (B15-B19)	17	1.2	*
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	16	1.1	*
11	Chronic liver disease and cirrhosis (K70,K73-K74)	16	1.1	*
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	9	0.6	*
14	Parkinson's disease (G20-G21)	7	0.5	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.4	*
15	Alzheimer's disease (G30)	6	0.4	*
15	Aortic aneurysm and dissection (I71)	6	0.4	*
...	All other causes (Residual)	168	11.5	80.2
Asian or Pacific Islander, female, 70-74 years				
...	All causes	1,989	100.0	1,164.6
1	Malignant neoplasms (C00-C97)	691	34.7	404.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	453	22.8	265.2
3	Cerebrovascular diseases (I60-I69)	179	9.0	104.8
4	Diabetes mellitus (E10-E14)	87	4.4	50.9
5	Chronic lower respiratory diseases (J40-J47)	57	2.9	33.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	47	2.4	27.5
7	Influenza and pneumonia (J10-J18)	44	2.2	25.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	38	1.9	22.3
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	32	1.6	18.7
10	Viral hepatitis (B15-B19)	25	1.3	14.6
11	Septicemia (A40-A41)	21	1.1	12.3
12	Chronic liver disease and cirrhosis (K70,K73-K74)	19	1.0	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	15	0.8	*
13	Parkinson's disease (G20-G21)	15	0.8	*
15	Aortic aneurysm and dissection (I71)	12	0.6	*
...	All other causes (Residual)	254	12.8	148.7

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 75-79 years				
...	All causes	2,641	100.0	1,963.8
1	Malignant neoplasms (C00-C97)	714	27.0	530.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	643	24.3	478.1
3	Cerebrovascular diseases (I60-I69)	268	10.1	199.3
4	Diabetes mellitus (E10-E14)	160	6.1	119.0
5	Chronic lower respiratory diseases (J40-J47)	85	3.2	63.2
6	Influenza and pneumonia (J10-J18)	70	2.7	52.1
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	66	2.5	49.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	54	2.0	40.2
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	50	1.9	37.2
10	Septicemia (A40-A41)	41	1.6	30.5
11	Alzheimer's disease (G30)	35	1.3	26.0
12	Chronic liver disease and cirrhosis (K70,K73-K74)	31	1.2	23.1
13	Parkinson's disease (G20-G21)	25	0.9	18.6
14	Aortic aneurysm and dissection (I71)	22	0.8	16.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	0.6	*
...	All other causes (Residual)	360	13.6	267.7
Asian or Pacific Islander, female, 80-84 years				
...	All causes	2,979	100.0	3,397.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	874	29.3	996.9
2	Malignant neoplasms (C00-C97)	634	21.3	723.2
3	Cerebrovascular diseases (I60-I69)	330	11.1	376.4
4	Diabetes mellitus (E10-E14)	145	4.9	165.4
5	Influenza and pneumonia (J10-J18)	100	3.4	114.1
6	Chronic lower respiratory diseases (J40-J47)	94	3.2	107.2
7	Alzheimer's disease (G30)	78	2.6	89.0
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	72	2.4	82.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	68	2.3	77.6
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	53	1.8	60.5
11	Septicemia (A40-A41)	33	1.1	37.6
12	Parkinson's disease (G20-G21)	31	1.0	35.4
13	Aortic aneurysm and dissection (I71)	25	0.8	28.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	0.8	26.2
15	Pneumonitis due to solids and liquids (J69)	22	0.7	25.1
...	All other causes (Residual)	397	13.3	452.8

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 85-89 years				
...	All causes	2,797	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	849	30.4	---
2	Malignant neoplasms (C00-C97)	443	15.8	---
3	Cerebrovascular diseases (I60-I69)	348	12.4	---
4	Influenza and pneumonia (J10-J18)	126	4.5	---
5	Diabetes mellitus (E10-E14)	120	4.3	---
6	Alzheimer's disease (G30)	108	3.9	---
7	Chronic lower respiratory diseases (J40-J47)	102	3.6	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	62	2.2	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	56	2.0	---
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	54	1.9	---
11	Septicemia (A40-A41)	41	1.5	---
12	Parkinson's disease (G20-G21)	26	0.9	---
12	Pneumonitis due to solids and liquids (J69)	26	0.9	---
14	Aortic aneurysm and dissection (I71)	15	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	0.5	---
15	Peptic ulcer (K25-K28)	13	0.5	---
...	All other causes (Residual)	395	14.1	---
Asian or Pacific Islander, female, 90-94 years				
...	All causes	1,942	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	658	33.9	---
2	Cerebrovascular diseases (I60-I69)	222	11.4	---
3	Malignant neoplasms (C00-C97)	193	9.9	---
4	Influenza and pneumonia (J10-J18)	121	6.2	---
5	Alzheimer's disease (G30)	90	4.6	---
6	Diabetes mellitus (E10-E14)	71	3.7	---
7	Chronic lower respiratory diseases (J40-J47)	66	3.4	---
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	59	3.0	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	38	2.0	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	37	1.9	---
11	Pneumonitis due to solids and liquids (J69)	25	1.3	---
12	Septicemia (A40-A41)	16	0.8	---
13	Aortic aneurysm and dissection (I71)	15	0.8	---
14	Atherosclerosis (I70)	13	0.7	---
15	Parkinson's disease (G20-G21)	11	0.6	---
...	All other causes (Residual)	307	15.8	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 95-99 years				
...	All causes	754	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	288	38.2	---
2	Cerebrovascular diseases (I60-I69)	88	11.7	---
3	Influenza and pneumonia (J10-J18)	58	7.7	---
4	Alzheimer's disease (G30)	47	6.2	---
5	Malignant neoplasms (C00-C97)	41	5.4	---
6	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	21	2.8	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	2.1	---
8	Chronic lower respiratory diseases (J40-J47)	15	2.0	---
9	Diabetes mellitus (E10-E14)	11	1.5	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9	1.2	---
11	Septicemia (A40-A41)	7	0.9	---
11	Pneumonitis due to solids and liquids (J69)	7	0.9	---
13	Peptic ulcer (K25-K28)	5	0.7	---
14	Anemias (D50-D64)	3	0.4	---
14	Cholelithiasis and other disorders of gallbladder (K80-K82)	3	0.4	---
...	All other causes (Residual)	135	17.9	---
Asian or Pacific Islander, female, 100 years and over				
...	All causes	165	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	64	38.8	---
2	Cerebrovascular diseases (I60-I69)	16	9.7	---
3	Influenza and pneumonia (J10-J18)	10	6.1	---
4	Malignant neoplasms (C00-C97)	7	4.2	---
5	Alzheimer's disease (G30)	5	3.0	---
5	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	5	3.0	---
7	Diabetes mellitus (E10-E14)	4	2.4	---
7	Atherosclerosis (I70)	4	2.4	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	2.4	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4	2.4	---
11	Chronic lower respiratory diseases (J40-J47)	3	1.8	---
11	Pneumonitis due to solids and liquids (J69)	3	1.8	---
13	Septicemia (A40-A41)	2	1.2	---
14	Nutritional deficiencies (E40-E64)	1	0.6	---
14	Parkinson's disease (G20-G21)	1	0.6	---
14	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.6	---
...	All other causes (Residual)	31	18.8	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available