

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, all ages				
...	All causes	2,423,712	100.0	803.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	616,067	25.4	204.3
2	Malignant neoplasms (C00-C97)	562,875	23.2	186.6
3	Cerebrovascular diseases (I60-I69)	135,952	5.6	45.1
4	Chronic lower respiratory diseases (J40-J47)	127,924	5.3	42.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	123,706	5.1	41.0
6	Alzheimer's disease (G30)	74,632	3.1	24.7
7	Diabetes mellitus (E10-E14)	71,382	2.9	23.7
8	Influenza and pneumonia (J09-J18)	52,717	2.2	17.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	46,448	1.9	15.4
10	Septicemia (A40-A41)	34,828	1.4	11.5
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	34,598	1.4	11.5
12	Chronic liver disease and cirrhosis (K70,K73-K74)	29,165	1.2	9.7
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	23,965	1.0	7.9
14	Parkinson's disease (G20-G21)	20,058	0.8	6.7
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	18,361	0.8	6.1
...	All other causes (Residual)	451,034	18.6	149.5
All races, both sexes, 1-4 years				
...	All causes	4,703	100.0	28.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,588	33.8	9.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	546	11.6	3.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	398	8.5	2.4
4	Malignant neoplasms (C00-C97)	364	7.7	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	173	3.7	1.1
6	Influenza and pneumonia (J09-J18)	109	2.3	0.7
7	Septicemia (A40-A41)	78	1.7	0.5
8	Certain conditions originating in the perinatal period (P00-P96)	70	1.5	0.4
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	59	1.3	0.4
10	Chronic lower respiratory diseases (J40-J47)	57	1.2	0.3
11	Cerebrovascular diseases (I60-I69)	52	1.1	0.3
12	Complications of medical and surgical care (Y40-Y84,Y88)	23	0.5	0.1
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22	0.5	0.1
14	Anemias (D50-D64)	17	0.4	*
15	Meningitis (G00,G03)	16	0.3	*
15	Acute bronchitis and bronchiolitis (J20-J21)	16	0.3	*
...	All other causes (Residual)	1,115	23.7	6.8

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 5-9 years				
...	All causes	2,711	100.0	13.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	965	35.6	4.9
2	Malignant neoplasms (C00-C97)	480	17.7	2.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	196	7.2	1.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	133	4.9	0.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	110	4.1	0.6
6	Chronic lower respiratory diseases (J40-J47)	54	2.0	0.3
7	Influenza and pneumonia (J09-J18)	48	1.8	0.2
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	41	1.5	0.2
9	Cerebrovascular diseases (I60-I69)	38	1.4	0.2
10	Septicemia (A40-A41)	36	1.3	0.2
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	0.5	*
12	Anemias (D50-D64)	13	0.5	*
13	Diabetes mellitus (E10-E14)	10	0.4	*
13	Certain conditions originating in the perinatal period (P00-P96)	10	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	10	0.4	*
...	All other causes (Residual)	553	20.4	2.8
All races, both sexes, 10-14 years				
...	All causes	3,436	100.0	16.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,229	35.8	6.0
2	Malignant neoplasms (C00-C97)	479	13.9	2.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	213	6.2	1.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	180	5.2	0.9
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	178	5.2	0.9
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	131	3.8	0.6
7	Chronic lower respiratory diseases (J40-J47)	64	1.9	0.3
8	Influenza and pneumonia (J09-J18)	55	1.6	0.3
9	Cerebrovascular diseases (I60-I69)	45	1.3	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	43	1.3	0.2
11	Septicemia (A40-A41)	38	1.1	0.2
12	Anemias (D50-D64)	19	0.6	*
13	Meningitis (G00,G03)	16	0.5	*
14	Certain conditions originating in the perinatal period (P00-P96)	12	0.3	*
15	Diabetes mellitus (E10-E14)	11	0.3	*
15	Pneumonitis due to solids and liquids (J69)	11	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	11	0.3	*
...	All other causes (Residual)	701	20.4	3.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 15-19 years				
...	All causes	13,299	100.0	61.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,493	48.8	30.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,224	16.7	10.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,481	11.1	6.9
4	Malignant neoplasms (C00-C97)	673	5.1	3.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	346	2.6	1.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	191	1.4	0.9
7	Cerebrovascular diseases (I60-I69)	71	0.5	0.3
7	Chronic lower respiratory diseases (J40-J47)	71	0.5	0.3
9	Diabetes mellitus (E10-E14)	55	0.4	0.3
10	Septicemia (A40-A41)	53	0.4	0.2
11	Influenza and pneumonia (J09-J18)	52	0.4	0.2
12	Pregnancy, childbirth and the puerperium (O00-O99)	44	0.3	0.2
13	Anemias (D50-D64)	35	0.3	0.2
14	Human immunodeficiency virus (HIV) disease (B20-B24)	33	0.2	0.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	32	0.2	0.1
...	All other causes (Residual)	1,445	10.9	6.7
All races, both sexes, 20-24 years				
...	All causes	20,683	100.0	98.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,404	45.5	44.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,327	16.1	15.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,659	12.9	12.6
4	Malignant neoplasms (C00-C97)	980	4.7	4.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	738	3.6	3.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	211	1.0	1.0
7	Human immunodeficiency virus (HIV) disease (B20-B24)	127	0.6	0.6
8	Cerebrovascular diseases (I60-I69)	124	0.6	0.6
9	Pregnancy, childbirth and the puerperium (O00-O99)	116	0.6	0.6
10	Diabetes mellitus (E10-E14)	113	0.5	0.5
11	Influenza and pneumonia (J09-J18)	111	0.5	0.5
12	Septicemia (A40-A41)	107	0.5	0.5
13	Chronic lower respiratory diseases (J40-J47)	78	0.4	0.4
14	Legal intervention (Y35,Y89.0)	73	0.4	0.3
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	65	0.3	0.3
...	All other causes (Residual)	2,450	11.8	11.6

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 25-29 years				
...	All causes	20,931	100.0	99.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,134	38.9	38.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,717	13.0	12.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,673	12.8	12.7
4	Malignant neoplasms (C00-C97)	1,318	6.3	6.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,269	6.1	6.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	412	2.0	2.0
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	226	1.1	1.1
8	Diabetes mellitus (E10-E14)	215	1.0	1.0
9	Cerebrovascular diseases (I60-I69)	180	0.9	0.9
10	Pregnancy, childbirth and the puerperium (O00-O99)	158	0.8	0.8
11	Influenza and pneumonia (J09-J18)	155	0.7	0.7
12	Septicemia (A40-A41)	127	0.6	0.6
12	Chronic lower respiratory diseases (J40-J47)	127	0.6	0.6
14	Chronic liver disease and cirrhosis (K70,K73-K74)	111	0.5	0.5
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	104	0.5	0.5
...	All other causes (Residual)	3,005	14.4	14.3
All races, both sexes, 30-34 years				
...	All causes	21,641	100.0	110.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,843	31.6	35.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,605	12.0	13.3
3	Malignant neoplasms (C00-C97)	2,145	9.9	11.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,041	9.4	10.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,954	9.0	10.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	679	3.1	3.5
7	Diabetes mellitus (E10-E14)	395	1.8	2.0
8	Cerebrovascular diseases (I60-I69)	325	1.5	1.7
9	Chronic liver disease and cirrhosis (K70,K73-K74)	273	1.3	1.4
10	Congenital malformations, deformations and chromosomal	191	0.9	1.0
11	Influenza and pneumonia (J09-J18)	176	0.8	0.9
12	Septicemia (A40-A41)	170	0.8	0.9
13	Pregnancy, childbirth and the puerperium (O00-O99)	168	0.8	0.9
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	157	0.7	0.8
15	Chronic lower respiratory diseases (J40-J47)	136	0.6	0.7
...	All other causes (Residual)	3,383	15.6	17.3

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 35-39 years				
...	All causes	30,881	100.0	145.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,737	25.1	36.5
2	Malignant neoplasms (C00-C97)	4,237	13.7	20.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,898	12.6	18.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,022	9.8	14.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,619	5.2	7.6
6	Human immunodeficiency virus (HIV) disease (B20-B24)	1,382	4.5	6.5
7	Diabetes mellitus (E10-E14)	749	2.4	3.5
8	Chronic liver disease and cirrhosis (K70,K73-K74)	742	2.4	3.5
9	Cerebrovascular diseases (I60-I69)	733	2.4	3.5
10	Septicemia (A40-A41)	328	1.1	1.5
11	Influenza and pneumonia (J09-J18)	287	0.9	1.4
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	281	0.9	1.3
13	Congenital malformations, deformations and chromosomal	233	0.8	1.1
14	Chronic lower respiratory diseases (J40-J47)	228	0.7	1.1
15	Pregnancy, childbirth and the puerperium (O00-O99)	156	0.5	0.7
...	All other causes (Residual)	5,249	17.0	24.8
All races, both sexes, 40-44 years				
...	All causes	48,725	100.0	221.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,194	18.9	41.8
2	Malignant neoplasms (C00-C97)	9,051	18.6	41.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,941	16.3	36.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,700	7.6	16.8
5	Human immunodeficiency virus (HIV) disease (B20-B24)	2,190	4.5	10.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1,828	3.8	8.3
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,433	2.9	6.5
8	Cerebrovascular diseases (I60-I69)	1,400	2.9	6.4
9	Diabetes mellitus (E10-E14)	1,235	2.5	5.6
10	Septicemia (A40-A41)	582	1.2	2.6
11	Chronic lower respiratory diseases (J40-J47)	568	1.2	2.6
12	Influenza and pneumonia (J09-J18)	497	1.0	2.3
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	473	1.0	2.2
14	Viral hepatitis (B15-B19)	373	0.8	1.7
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	280	0.6	1.3
...	All other causes (Residual)	7,980	16.4	36.3

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 45-49 years				
...	All causes	77,738	100.0	340.0
1	Malignant neoplasms (C00-C97)	18,778	24.2	82.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,865	19.1	65.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,717	13.8	46.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,051	5.2	17.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	3,540	4.6	15.5
6	Cerebrovascular diseases (I60-I69)	2,616	3.4	11.4
7	Human immunodeficiency virus (HIV) disease (B20-B24)	2,267	2.9	9.9
8	Diabetes mellitus (E10-E14)	2,161	2.8	9.5
9	Chronic lower respiratory diseases (J40-J47)	1,397	1.8	6.1
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,234	1.6	5.4
11	Viral hepatitis (B15-B19)	1,019	1.3	4.5
12	Septicemia (A40-A41)	1,002	1.3	4.4
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	883	1.1	3.9
14	Influenza and pneumonia (J09-J18)	800	1.0	3.5
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	435	0.6	1.9
...	All other causes (Residual)	11,973	15.4	52.4
All races, both sexes, 50-54 years				
...	All causes	106,948	100.0	509.0
1	Malignant neoplasms (C00-C97)	31,389	29.3	149.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	22,569	21.1	107.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,598	9.0	45.7
4	Chronic liver disease and cirrhosis (K70,K73-K74)	4,672	4.4	22.2
5	Cerebrovascular diseases (I60-I69)	3,769	3.5	17.9
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,727	3.5	17.7
7	Diabetes mellitus (E10-E14)	3,592	3.4	17.1
8	Chronic lower respiratory diseases (J40-J47)	2,756	2.6	13.1
9	Human immunodeficiency virus (HIV) disease (B20-B24)	1,889	1.8	9.0
10	Viral hepatitis (B15-B19)	1,796	1.7	8.5
11	Septicemia (A40-A41)	1,429	1.3	6.8
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,350	1.3	6.4
13	Influenza and pneumonia (J09-J18)	1,109	1.0	5.3
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	906	0.8	4.3
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	800	0.7	3.8
...	All other causes (Residual)	15,597	14.6	74.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 55-59 years				
...	All causes	132,458	100.0	726.3
1	Malignant neoplasms (C00-C97)	45,476	34.3	249.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	29,793	22.5	163.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,100	5.4	38.9
4	Diabetes mellitus (E10-E14)	5,085	3.8	27.9
5	Cerebrovascular diseases (I60-I69)	4,745	3.6	26.0
6	Chronic lower respiratory diseases (J40-J47)	4,641	3.5	25.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	4,485	3.4	24.6
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,962	2.2	16.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,977	1.5	10.8
10	Septicemia (A40-A41)	1,958	1.5	10.7
11	Viral hepatitis (B15-B19)	1,641	1.2	9.0
12	Influenza and pneumonia (J09-J18)	1,396	1.1	7.7
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,100	0.8	6.0
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	951	0.7	5.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	634	0.5	3.5
...	All other causes (Residual)	18,514	14.0	101.5
All races, both sexes, 60-64 years				
...	All causes	154,652	100.0	1,068.3
1	Malignant neoplasms (C00-C97)	57,695	37.3	398.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	35,734	23.1	246.9
3	Chronic lower respiratory diseases (J40-J47)	8,136	5.3	56.2
4	Diabetes mellitus (E10-E14)	6,219	4.0	43.0
5	Cerebrovascular diseases (I60-I69)	5,755	3.7	39.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,093	3.3	35.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	3,519	2.3	24.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,463	1.6	17.0
9	Septicemia (A40-A41)	2,273	1.5	15.7
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,107	1.4	14.6
11	Influenza and pneumonia (J09-J18)	1,756	1.1	12.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,173	0.8	8.1
13	Aortic aneurysm and dissection (I71)	873	0.6	6.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	782	0.5	5.4
15	Viral hepatitis (B15-B19)	772	0.5	5.3
...	All other causes (Residual)	20,302	13.1	140.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 65-69 years				
...	All causes	174,991	100.0	1,627.5
1	Malignant neoplasms (C00-C97)	65,624	37.5	610.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	39,690	22.7	369.1
3	Chronic lower respiratory diseases (J40-J47)	11,975	6.8	111.4
4	Cerebrovascular diseases (I60-I69)	7,461	4.3	69.4
5	Diabetes mellitus (E10-E14)	6,879	3.9	64.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,187	2.4	38.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,248	1.9	30.2
8	Septicemia (A40-A41)	2,796	1.6	26.0
9	Chronic liver disease and cirrhosis (K70,K73-K74)	2,745	1.6	25.5
10	Influenza and pneumonia (J09-J18)	2,197	1.3	20.4
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,382	0.8	12.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,376	0.8	12.8
13	Aortic aneurysm and dissection (I71)	1,094	0.6	10.2
14	Alzheimer's disease (G30)	1,074	0.6	10.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,006	0.6	9.4
...	All other causes (Residual)	22,257	12.7	207.0
All races, both sexes, 70-74 years				
...	All causes	214,247	100.0	2,491.3
1	Malignant neoplasms (C00-C97)	72,842	34.0	847.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	49,899	23.3	580.2
3	Chronic lower respiratory diseases (J40-J47)	16,689	7.8	194.1
4	Cerebrovascular diseases (I60-I69)	10,546	4.9	122.6
5	Diabetes mellitus (E10-E14)	8,233	3.8	95.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,566	2.1	53.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,504	2.1	52.4
8	Septicemia (A40-A41)	3,549	1.7	41.3
9	Influenza and pneumonia (J09-J18)	3,350	1.6	39.0
10	Alzheimer's disease (G30)	2,910	1.4	33.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	2,422	1.1	28.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,757	0.8	20.4
13	Parkinson's disease (G20-G21)	1,610	0.8	18.7
14	Aortic aneurysm and dissection (I71)	1,522	0.7	17.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,396	0.7	16.2
...	All other causes (Residual)	28,452	13.3	330.8

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 75-79 years				
...	All causes	289,029	100.0	3,945.9
1	Malignant neoplasms (C00-C97)	82,988	28.7	1,133.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	71,589	24.8	977.3
3	Chronic lower respiratory diseases (J40-J47)	22,758	7.9	310.7
4	Cerebrovascular diseases (I60-I69)	16,973	5.9	231.7
5	Diabetes mellitus (E10-E14)	10,012	3.5	136.7
6	Alzheimer's disease (G30)	7,575	2.6	103.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,385	2.2	87.2
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,922	2.0	80.8
9	Influenza and pneumonia (J09-J18)	5,740	2.0	78.4
10	Septicemia (A40-A41)	4,629	1.6	63.2
11	Parkinson's disease (G20-G21)	3,697	1.3	50.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,574	0.9	35.1
13	Chronic liver disease and cirrhosis (K70,K73-K74)	2,143	0.7	29.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,066	0.7	28.2
15	Aortic aneurysm and dissection (I71)	1,992	0.7	27.2
...	All other causes (Residual)	41,986	14.5	573.2
All races, both sexes, 80-84 years				
...	All causes	363,653	100.0	6,381.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	99,668	27.4	1,749.0
2	Malignant neoplasms (C00-C97)	80,620	22.2	1,414.7
3	Chronic lower respiratory diseases (J40-J47)	25,283	7.0	443.7
4	Cerebrovascular diseases (I60-I69)	25,006	6.9	438.8
5	Alzheimer's disease (G30)	15,434	4.2	270.8
6	Diabetes mellitus (E10-E14)	11,177	3.1	196.1
7	Influenza and pneumonia (J09-J18)	9,119	2.5	160.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,326	2.3	146.1
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,814	2.1	137.1
10	Septicemia (A40-A41)	5,774	1.6	101.3
11	Parkinson's disease (G20-G21)	5,666	1.6	99.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,868	1.1	67.9
13	Pneumonitis due to solids and liquids (J69)	3,311	0.9	58.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,651	0.7	46.5
15	Aortic aneurysm and dissection (I71)	2,354	0.6	41.3
...	All other causes (Residual)	57,582	15.8	1,010.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 85-89 years				
...	All causes	359,224	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	110,176	30.7	---
2	Malignant neoplasms (C00-C97)	55,872	15.6	---
3	Cerebrovascular diseases (I60-I69)	28,161	7.8	---
4	Alzheimer's disease (G30)	21,111	5.9	---
5	Chronic lower respiratory diseases (J40-J47)	19,724	5.5	---
6	Influenza and pneumonia (J09-J18)	11,206	3.1	---
7	Diabetes mellitus (E10-E14)	9,085	2.5	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,434	2.3	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,064	2.2	---
10	Septicemia (A40-A41)	5,335	1.5	---
11	Parkinson's disease (G20-G21)	4,955	1.4	---
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,687	1.3	---
13	Pneumonitis due to solids and liquids (J69)	4,070	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,370	0.7	---
15	Aortic aneurysm and dissection (I71)	1,964	0.5	---
...	All other causes (Residual)	64,010	17.8	---
All races, both sexes, 90-94 years				
...	All causes	240,180	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	82,401	34.3	---
2	Malignant neoplasms (C00-C97)	24,574	10.2	---
3	Cerebrovascular diseases (I60-I69)	19,235	8.0	---
4	Alzheimer's disease (G30)	17,075	7.1	---
5	Chronic lower respiratory diseases (J40-J47)	9,810	4.1	---
6	Influenza and pneumonia (J09-J18)	9,081	3.8	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,326	2.2	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,249	2.2	---
9	Diabetes mellitus (E10-E14)	4,613	1.9	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,716	1.5	---
11	Septicemia (A40-A41)	3,059	1.3	---
12	Pneumonitis due to solids and liquids (J69)	2,970	1.2	---
13	Parkinson's disease (G20-G21)	2,383	1.0	---
14	Atherosclerosis (I70)	1,608	0.7	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,374	0.6	---
...	All other causes (Residual)	47,706	19.9	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 95-99 years				
...	All causes	94,477	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	35,010	37.1	---
2	Cerebrovascular diseases (I60-I69)	7,315	7.7	---
3	Alzheimer's disease (G30)	7,268	7.7	---
4	Malignant neoplasms (C00-C97)	6,374	6.7	---
5	Influenza and pneumonia (J09-J18)	4,119	4.4	---
6	Chronic lower respiratory diseases (J40-J47)	2,936	3.1	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,115	2.2	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,861	2.0	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,695	1.8	---
10	Diabetes mellitus (E10-E14)	1,294	1.4	---
11	Pneumonitis due to solids and liquids (J69)	1,165	1.2	---
12	Septicemia (A40-A41)	1,033	1.1	---
13	Atherosclerosis (I70)	848	0.9	---
14	Parkinson's disease (G20-G21)	514	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	415	0.4	---
...	All other causes (Residual)	20,515	21.7	---
All races, both sexes, 100 years and over				
...	All causes	19,766	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,662	38.8	---
2	Alzheimer's disease (G30)	1,350	6.8	---
3	Cerebrovascular diseases (I60-I69)	1,264	6.4	---
4	Influenza and pneumonia (J09-J18)	1,129	5.7	---
5	Malignant neoplasms (C00-C97)	836	4.2	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	438	2.2	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	400	2.0	---
8	Chronic lower respiratory diseases (J40-J47)	387	2.0	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	375	1.9	---
10	Atherosclerosis (I70)	278	1.4	---
11	Pneumonitis due to solids and liquids (J69)	246	1.2	---
12	Diabetes mellitus (E10-E14)	235	1.2	---
13	Septicemia (A40-A41)	187	0.9	---
14	Anemias (D50-D64)	65	0.3	---
15	Parkinson's disease (G20-G21)	59	0.3	---
...	All other causes (Residual)	4,855	24.6	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, all ages				
...	All causes	1,203,968	100.0	809.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	309,821	25.7	208.4
2	Malignant neoplasms (C00-C97)	292,857	24.3	197.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	79,827	6.6	53.7
4	Chronic lower respiratory diseases (J40-J47)	61,235	5.1	41.2
5	Cerebrovascular diseases (I60-I69)	54,111	4.5	36.4
6	Diabetes mellitus (E10-E14)	35,478	2.9	23.9
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	27,269	2.3	18.3
8	Influenza and pneumonia (J09-J18)	24,071	2.0	16.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22,616	1.9	15.2
10	Alzheimer's disease (G30)	21,800	1.8	14.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	19,151	1.6	12.9
12	Septicemia (A40-A41)	15,839	1.3	10.7
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	14,538	1.2	9.8
14	Parkinson's disease (G20-G21)	11,554	1.0	7.8
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9,417	0.8	6.3
...	All other causes (Residual)	204,384	17.0	137.5
All races, male, 1-4 years				
...	All causes	2,634	100.0	31.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	987	37.5	11.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	286	10.9	3.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	213	8.1	2.5
4	Malignant neoplasms (C00-C97)	191	7.3	2.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	82	3.1	1.0
6	Influenza and pneumonia (J09-J18)	55	2.1	0.7
7	Septicemia (A40-A41)	44	1.7	0.5
8	Certain conditions originating in the perinatal period (P00-P96)	34	1.3	0.4
9	Chronic lower respiratory diseases (J40-J47)	30	1.1	0.4
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	28	1.1	0.3
11	Cerebrovascular diseases (I60-I69)	20	0.8	0.2
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	0.4	*
13	Meningitis (G00,G03)	10	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	9	0.3	*
15	Acute bronchitis and bronchiolitis (J20-J21)	8	0.3	*
...	All other causes (Residual)	626	23.8	7.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 5-9 years				
...	All causes	1,519	100.0	15.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	572	37.7	5.6
2	Malignant neoplasms (C00-C97)	255	16.8	2.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	97	6.4	1.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	66	4.3	0.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	63	4.1	0.6
6	Chronic lower respiratory diseases (J40-J47)	36	2.4	0.4
7	Influenza and pneumonia (J09-J18)	25	1.6	0.2
8	Septicemia (A40-A41)	24	1.6	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	21	1.4	0.2
10	Cerebrovascular diseases (I60-I69)	17	1.1	*
11	Anemias (D50-D64)	8	0.5	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.5	*
13	Diabetes mellitus (E10-E14)	6	0.4	*
13	Certain conditions originating in the perinatal period (P00-P96)	6	0.4	*
15	Meningococcal infection (A39)	5	0.3	*
...	All other causes (Residual)	310	20.4	3.1
All races, male, 10-14 years				
...	All causes	2,066	100.0	19.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	774	37.5	7.4
2	Malignant neoplasms (C00-C97)	244	11.8	2.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	144	7.0	1.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	128	6.2	1.2
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	97	4.7	0.9
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	70	3.4	0.7
7	Chronic lower respiratory diseases (J40-J47)	42	2.0	0.4
8	Influenza and pneumonia (J09-J18)	27	1.3	0.3
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	26	1.3	0.3
10	Septicemia (A40-A41)	21	1.0	0.2
11	Cerebrovascular diseases (I60-I69)	20	1.0	0.2
12	Anemias (D50-D64)	13	0.6	*
13	Pneumonitis due to solids and liquids (J69)	10	0.5	*
14	Meningitis (G00,G03)	9	0.4	*
15	Certain conditions originating in the perinatal period (P00-P96)	8	0.4	*
...	All other causes (Residual)	433	21.0	4.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 15-19 years				
...	All causes	9,558	100.0	86.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,542	47.5	41.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,932	20.2	17.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,221	12.8	11.1
4	Malignant neoplasms (C00-C97)	395	4.1	3.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	230	2.4	2.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	113	1.2	1.0
7	Chronic lower respiratory diseases (J40-J47)	43	0.4	0.4
8	Cerebrovascular diseases (I60-I69)	40	0.4	0.4
9	Diabetes mellitus (E10-E14)	31	0.3	0.3
10	Septicemia (A40-A41)	24	0.3	0.2
10	Legal intervention (Y35,Y89.0)	24	0.3	0.2
12	Influenza and pneumonia (J09-J18)	23	0.2	0.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	0.2	*
14	Anemias (D50-D64)	15	0.2	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	14	0.1	*
...	All other causes (Residual)	895	9.4	8.1
All races, male, 20-24 years				
...	All causes	15,758	100.0	145.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,363	46.7	67.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,897	18.4	26.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,260	14.3	20.8
4	Malignant neoplasms (C00-C97)	592	3.8	5.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	470	3.0	4.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	128	0.8	1.2
7	Human immunodeficiency virus (HIV) disease (B20-B24)	74	0.5	0.7
8	Cerebrovascular diseases (I60-I69)	73	0.5	0.7
9	Legal intervention (Y35,Y89.0)	70	0.4	0.6
10	Influenza and pneumonia (J09-J18)	68	0.4	0.6
11	Diabetes mellitus (E10-E14)	67	0.4	0.6
12	Septicemia (A40-A41)	61	0.4	0.6
13	Chronic lower respiratory diseases (J40-J47)	53	0.3	0.5
14	Anemias (D50-D64)	34	0.2	0.3
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	34	0.2	0.3
...	All other causes (Residual)	1,514	9.6	14.0

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 25-29 years				
...	All causes	15,107	100.0	140.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,246	41.3	58.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,306	15.3	21.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,190	14.5	20.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	847	5.6	7.9
5	Malignant neoplasms (C00-C97)	715	4.7	6.6
6	Human immunodeficiency virus (HIV) disease (B20-B24)	245	1.6	2.3
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	134	0.9	1.2
8	Diabetes mellitus (E10-E14)	123	0.8	1.1
9	Cerebrovascular diseases (I60-I69)	90	0.6	0.8
9	Influenza and pneumonia (J09-J18)	90	0.6	0.8
11	Chronic lower respiratory diseases (J40-J47)	76	0.5	0.7
12	Legal intervention (Y35,Y89.0)	71	0.5	0.7
13	Chronic liver disease and cirrhosis (K70,K73-K74)	68	0.5	0.6
14	Septicemia (A40-A41)	66	0.4	0.6
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	56	0.4	0.5
...	All other causes (Residual)	1,784	11.8	16.6
All races, male, 30-34 years				
...	All causes	14,685	100.0	148.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,138	35.0	51.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,091	14.2	21.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,713	11.7	17.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,330	9.1	13.4
5	Malignant neoplasms (C00-C97)	977	6.7	9.9
6	Human immunodeficiency virus (HIV) disease (B20-B24)	408	2.8	4.1
7	Diabetes mellitus (E10-E14)	236	1.6	2.4
8	Chronic liver disease and cirrhosis (K70,K73-K74)	193	1.3	1.9
9	Cerebrovascular diseases (I60-I69)	162	1.1	1.6
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	109	0.7	1.1
11	Influenza and pneumonia (J09-J18)	97	0.7	1.0
12	Septicemia (A40-A41)	87	0.6	0.9
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	85	0.6	0.9
14	Chronic lower respiratory diseases (J40-J47)	74	0.5	0.7
15	Legal intervention (Y35,Y89.0)	55	0.4	0.6
...	All other causes (Residual)	1,930	13.1	19.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 35-39 years				
...	All causes	19,755	100.0	185.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,582	28.3	52.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,704	13.7	25.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,360	11.9	22.1
4	Malignant neoplasms (C00-C97)	1,757	8.9	16.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,254	6.3	11.8
6	Human immunodeficiency virus (HIV) disease (B20-B24)	923	4.7	8.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	492	2.5	4.6
8	Diabetes mellitus (E10-E14)	479	2.4	4.5
9	Cerebrovascular diseases (I60-I69)	401	2.0	3.8
10	Septicemia (A40-A41)	169	0.9	1.6
11	Influenza and pneumonia (J09-J18)	154	0.8	1.4
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	151	0.8	1.4
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	119	0.6	1.1
14	Aortic aneurysm and dissection (I71)	110	0.6	1.0
15	Chronic lower respiratory diseases (J40-J47)	108	0.5	1.0
...	All other causes (Residual)	2,992	15.1	28.1
All races, male, 40-44 years				
...	All causes	30,350	100.0	276.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,440	21.2	58.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,641	18.6	51.5
3	Malignant neoplasms (C00-C97)	3,944	13.0	36.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,792	9.2	25.5
5	Human immunodeficiency virus (HIV) disease (B20-B24)	1,583	5.2	14.4
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1,230	4.1	11.2
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,035	3.4	9.4
8	Diabetes mellitus (E10-E14)	749	2.5	6.8
9	Cerebrovascular diseases (I60-I69)	742	2.4	6.8
10	Septicemia (A40-A41)	294	1.0	2.7
11	Influenza and pneumonia (J09-J18)	289	1.0	2.6
12	Chronic lower respiratory diseases (J40-J47)	284	0.9	2.6
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	277	0.9	2.5
14	Viral hepatitis (B15-B19)	239	0.8	2.2
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	170	0.6	1.6
...	All other causes (Residual)	4,641	15.3	42.3

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 45-49 years				
...	All causes	47,904	100.0	423.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,624	22.2	94.0
2	Malignant neoplasms (C00-C97)	8,901	18.6	78.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,369	15.4	65.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,043	6.4	26.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,429	5.1	21.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	1,624	3.4	14.4
7	Cerebrovascular diseases (I60-I69)	1,410	2.9	12.5
8	Diabetes mellitus (E10-E14)	1,376	2.9	12.2
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	881	1.8	7.8
10	Viral hepatitis (B15-B19)	712	1.5	6.3
11	Chronic lower respiratory diseases (J40-J47)	703	1.5	6.2
12	Septicemia (A40-A41)	540	1.1	4.8
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	485	1.0	4.3
14	Influenza and pneumonia (J09-J18)	466	1.0	4.1
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	271	0.6	2.4
...	All other causes (Residual)	7,070	14.8	62.6
All races, male, 50-54 years				
...	All causes	66,552	100.0	646.6
1	Malignant neoplasms (C00-C97)	16,469	24.7	160.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	16,287	24.5	158.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,648	10.0	64.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	3,359	5.0	32.6
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,781	4.2	27.0
6	Diabetes mellitus (E10-E14)	2,137	3.2	20.8
7	Cerebrovascular diseases (I60-I69)	2,097	3.2	20.4
8	Chronic lower respiratory diseases (J40-J47)	1,422	2.1	13.8
9	Human immunodeficiency virus (HIV) disease (B20-B24)	1,404	2.1	13.6
10	Viral hepatitis (B15-B19)	1,285	1.9	12.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	775	1.2	7.5
12	Septicemia (A40-A41)	765	1.1	7.4
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	691	1.0	6.7
14	Influenza and pneumonia (J09-J18)	660	1.0	6.4
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	489	0.7	4.8
...	All other causes (Residual)	9,283	13.9	90.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 55-59 years				
...	All causes	81,590	100.0	922.2
1	Malignant neoplasms (C00-C97)	24,740	30.3	279.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	21,227	26.0	239.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,944	6.1	55.9
4	Chronic liver disease and cirrhosis (K70,K73-K74)	3,278	4.0	37.1
5	Diabetes mellitus (E10-E14)	3,042	3.7	34.4
6	Cerebrovascular diseases (I60-I69)	2,757	3.4	31.2
7	Chronic lower respiratory diseases (J40-J47)	2,478	3.0	28.0
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,212	2.7	25.0
9	Viral hepatitis (B15-B19)	1,227	1.5	13.9
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,110	1.4	12.5
11	Septicemia (A40-A41)	1,051	1.3	11.9
12	Human immunodeficiency virus (HIV) disease (B20-B24)	840	1.0	9.5
13	Influenza and pneumonia (J09-J18)	834	1.0	9.4
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	577	0.7	6.5
15	Aortic aneurysm and dissection (I71)	453	0.6	5.1
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	453	0.6	5.1
...	All other causes (Residual)	10,367	12.7	117.2
All races, male, 60-64 years				
...	All causes	92,028	100.0	1,328.4
1	Malignant neoplasms (C00-C97)	31,819	34.6	459.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	24,331	26.4	351.2
3	Chronic lower respiratory diseases (J40-J47)	4,255	4.6	61.4
4	Diabetes mellitus (E10-E14)	3,602	3.9	52.0
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,413	3.7	49.3
6	Cerebrovascular diseases (I60-I69)	3,235	3.5	46.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,454	2.7	35.4
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,614	1.8	23.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,368	1.5	19.7
10	Septicemia (A40-A41)	1,152	1.3	16.6
11	Influenza and pneumonia (J09-J18)	1,047	1.1	15.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	696	0.8	10.0
13	Aortic aneurysm and dissection (I71)	640	0.7	9.2
14	Viral hepatitis (B15-B19)	534	0.6	7.7
15	Human immunodeficiency virus (HIV) disease (B20-B24)	504	0.5	7.3
...	All other causes (Residual)	11,364	12.3	164.0

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 65-69 years				
...	All causes	100,492	100.0	2,002.2
1	Malignant neoplasms (C00-C97)	36,166	36.0	720.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	25,585	25.5	509.8
3	Chronic lower respiratory diseases (J40-J47)	6,207	6.2	123.7
4	Cerebrovascular diseases (I60-I69)	3,978	4.0	79.3
5	Diabetes mellitus (E10-E14)	3,914	3.9	78.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,749	2.7	54.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,809	1.8	36.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,693	1.7	33.7
9	Septicemia (A40-A41)	1,442	1.4	28.7
10	Influenza and pneumonia (J09-J18)	1,232	1.2	24.5
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,125	1.1	22.4
12	Aortic aneurysm and dissection (I71)	751	0.7	15.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	735	0.7	14.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	583	0.6	11.6
15	Parkinson's disease (G20-G21)	474	0.5	9.4
...	All other causes (Residual)	12,049	12.0	240.1
All races, male, 70-74 years				
...	All causes	117,852	100.0	3,046.9
1	Malignant neoplasms (C00-C97)	39,758	33.7	1,027.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	29,947	25.4	774.2
3	Chronic lower respiratory diseases (J40-J47)	8,472	7.2	219.0
4	Cerebrovascular diseases (I60-I69)	5,373	4.6	138.9
5	Diabetes mellitus (E10-E14)	4,365	3.7	112.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,844	2.4	73.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,351	2.0	60.8
8	Septicemia (A40-A41)	1,814	1.5	46.9
9	Influenza and pneumonia (J09-J18)	1,812	1.5	46.8
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,410	1.2	36.5
11	Alzheimer's disease (G30)	1,221	1.0	31.6
12	Parkinson's disease (G20-G21)	1,033	0.9	26.7
13	Aortic aneurysm and dissection (I71)	947	0.8	24.5
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	878	0.7	22.7
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	866	0.7	22.4
...	All other causes (Residual)	14,761	12.5	381.6

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 75-79 years				
...	All causes	149,669	100.0	4,817.2
1	Malignant neoplasms (C00-C97)	44,488	29.7	1,431.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	39,346	26.3	1,266.4
3	Chronic lower respiratory diseases (J40-J47)	11,303	7.6	363.8
4	Cerebrovascular diseases (I60-I69)	7,868	5.3	253.2
5	Diabetes mellitus (E10-E14)	4,841	3.2	155.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,305	2.2	106.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,253	2.2	104.7
8	Influenza and pneumonia (J09-J18)	2,955	2.0	95.1
9	Alzheimer's disease (G30)	2,949	2.0	94.9
10	Parkinson's disease (G20-G21)	2,323	1.6	74.8
11	Septicemia (A40-A41)	2,200	1.5	70.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,194	0.8	38.4
13	Chronic liver disease and cirrhosis (K70,K73-K74)	1,186	0.8	38.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,161	0.8	37.4
15	Pneumonitis due to solids and liquids (J69)	1,125	0.8	36.2
...	All other causes (Residual)	20,172	13.5	649.3
All races, male, 80-84 years				
...	All causes	171,134	100.0	7,758.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	48,659	28.4	2,206.1
2	Malignant neoplasms (C00-C97)	41,439	24.2	1,878.7
3	Chronic lower respiratory diseases (J40-J47)	12,136	7.1	550.2
4	Cerebrovascular diseases (I60-I69)	9,832	5.7	445.8
5	Alzheimer's disease (G30)	5,301	3.1	240.3
6	Diabetes mellitus (E10-E14)	5,033	2.9	228.2
7	Influenza and pneumonia (J09-J18)	4,457	2.6	202.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,182	2.4	189.6
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,963	2.3	179.7
10	Parkinson's disease (G20-G21)	3,371	2.0	152.8
11	Septicemia (A40-A41)	2,568	1.5	116.4
12	Pneumonitis due to solids and liquids (J69)	1,801	1.1	81.7
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,473	0.9	66.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,409	0.8	63.9
15	Aortic aneurysm and dissection (I71)	1,222	0.7	55.4
...	All other causes (Residual)	24,288	14.2	1,101.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 85-89 years				
...	All causes	146,114	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	45,558	31.2	---
2	Malignant neoplasms (C00-C97)	27,005	18.5	---
3	Cerebrovascular diseases (I60-I69)	9,423	6.4	---
4	Chronic lower respiratory diseases (J40-J47)	8,729	6.0	---
5	Alzheimer's disease (G30)	6,324	4.3	---
6	Influenza and pneumonia (J09-J18)	4,876	3.3	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,853	2.6	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,655	2.5	---
9	Diabetes mellitus (E10-E14)	3,620	2.5	---
10	Parkinson's disease (G20-G21)	2,720	1.9	---
11	Pneumonitis due to solids and liquids (J69)	2,121	1.5	---
12	Septicemia (A40-A41)	2,064	1.4	---
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,486	1.0	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,055	0.7	---
15	Aortic aneurysm and dissection (I71)	937	0.6	---
...	All other causes (Residual)	22,688	15.5	---
All races, male, 90-94 years				
...	All causes	77,477	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	27,021	34.9	---
2	Malignant neoplasms (C00-C97)	10,512	13.6	---
3	Cerebrovascular diseases (I60-I69)	4,955	6.4	---
4	Alzheimer's disease (G30)	3,920	5.1	---
5	Chronic lower respiratory diseases (J40-J47)	3,737	4.8	---
6	Influenza and pneumonia (J09-J18)	3,319	4.3	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,123	2.7	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,865	2.4	---
9	Diabetes mellitus (E10-E14)	1,487	1.9	---
10	Pneumonitis due to solids and liquids (J69)	1,341	1.7	---
11	Parkinson's disease (G20-G21)	1,123	1.4	---
12	Septicemia (A40-A41)	1,021	1.3	---
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	949	1.2	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	526	0.7	---
15	Atherosclerosis (I70)	483	0.6	---
...	All other causes (Residual)	13,095	16.9	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 95-99 years				
...	All causes	22,102	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,281	37.5	---
2	Malignant neoplasms (C00-C97)	2,205	10.0	---
3	Cerebrovascular diseases (I60-I69)	1,376	6.2	---
4	Influenza and pneumonia (J09-J18)	1,191	5.4	---
5	Alzheimer's disease (G30)	1,120	5.1	---
6	Chronic lower respiratory diseases (J40-J47)	927	4.2	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	609	2.8	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	600	2.7	---
9	Pneumonitis due to solids and liquids (J69)	419	1.9	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	326	1.5	---
11	Diabetes mellitus (E10-E14)	312	1.4	---
12	Septicemia (A40-A41)	253	1.1	---
13	Parkinson's disease (G20-G21)	194	0.9	---
14	Atherosclerosis (I70)	165	0.7	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	109	0.5	---
...	All other causes (Residual)	4,015	18.2	---
All races, male, 100 years and over				
...	All causes	3,173	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,260	39.7	---
2	Influenza and pneumonia (J09-J18)	256	8.1	---
3	Malignant neoplasms (C00-C97)	242	7.6	---
4	Cerebrovascular diseases (I60-I69)	161	5.1	---
5	Alzheimer's disease (G30)	115	3.6	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	101	3.2	---
7	Chronic lower respiratory diseases (J40-J47)	87	2.7	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	81	2.6	---
9	Pneumonitis due to solids and liquids (J69)	63	2.0	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	44	1.4	---
11	Diabetes mellitus (E10-E14)	43	1.4	---
12	Atherosclerosis (I70)	38	1.2	---
13	Septicemia (A40-A41)	26	0.8	---
14	Parkinson's disease (G20-G21)	14	0.4	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	0.3	---
15	Anemias (D50-D64)	10	0.3	---
...	All other causes (Residual)	622	19.6	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, all ages				
...	All causes	1,219,744	100.0	797.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	306,246	25.1	200.2
2	Malignant neoplasms (C00-C97)	270,018	22.1	176.5
3	Cerebrovascular diseases (I60-I69)	81,841	6.7	53.5
4	Chronic lower respiratory diseases (J40-J47)	66,689	5.5	43.6
5	Alzheimer's disease (G30)	52,832	4.3	34.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	43,879	3.6	28.7
7	Diabetes mellitus (E10-E14)	35,904	2.9	23.5
8	Influenza and pneumonia (J09-J18)	28,646	2.3	18.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23,832	2.0	15.6
10	Septicemia (A40-A41)	18,989	1.6	12.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	14,548	1.2	9.5
12	Chronic liver disease and cirrhosis (K70,K73-K74)	10,014	0.8	6.5
13	Parkinson's disease (G20-G21)	8,504	0.7	5.6
14	Pneumonitis due to solids and liquids (J69)	8,054	0.7	5.3
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7,329	0.6	4.8
...	All other causes (Residual)	242,419	19.9	158.5
All races, female, 1-4 years				
...	All causes	2,069	100.0	25.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	601	29.0	7.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	260	12.6	3.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	185	8.9	2.3
4	Malignant neoplasms (C00-C97)	173	8.4	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	91	4.4	1.1
6	Influenza and pneumonia (J09-J18)	54	2.6	0.7
7	Certain conditions originating in the perinatal period (P00-P96)	36	1.7	0.4
8	Septicemia (A40-A41)	34	1.6	0.4
9	Cerebrovascular diseases (I60-I69)	32	1.5	0.4
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	31	1.5	0.4
11	Chronic lower respiratory diseases (J40-J47)	27	1.3	0.3
12	Complications of medical and surgical care (Y40-Y84,Y88)	14	0.7	*
13	Anemias (D50-D64)	11	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	0.5	*
15	Acute bronchitis and bronchiolitis (J20-J21)	8	0.4	*
...	All other causes (Residual)	501	24.2	6.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 5-9 years				
...	All causes	1,192	100.0	12.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	393	33.0	4.1
2	Malignant neoplasms (C00-C97)	225	18.9	2.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	99	8.3	1.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	67	5.6	0.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	47	3.9	0.5
6	Influenza and pneumonia (J09-J18)	23	1.9	0.2
7	Cerebrovascular diseases (I60-I69)	21	1.8	0.2
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	1.7	0.2
9	Chronic lower respiratory diseases (J40-J47)	18	1.5	*
10	Septicemia (A40-A41)	12	1.0	*
11	Meningitis (G00,G03)	6	0.5	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.5	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.5	*
14	Anemias (D50-D64)	5	0.4	*
15	Diabetes mellitus (E10-E14)	4	0.3	*
15	Diseases of appendix (K35-K38)	4	0.3	*
15	Certain conditions originating in the perinatal period (P00-P96)	4	0.3	*
...	All other causes (Residual)	232	19.5	2.4
All races, female, 10-14 years				
...	All causes	1,370	100.0	13.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	455	33.2	4.6
2	Malignant neoplasms (C00-C97)	235	17.2	2.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	81	5.9	0.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	69	5.0	0.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	61	4.5	0.6
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	52	3.8	0.5
7	Influenza and pneumonia (J09-J18)	28	2.0	0.3
8	Cerebrovascular diseases (I60-I69)	25	1.8	0.3
9	Chronic lower respiratory diseases (J40-J47)	22	1.6	0.2
10	Septicemia (A40-A41)	17	1.2	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	1.2	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	7	0.5	*
12	Meningitis (G00,G03)	7	0.5	*
14	Anemias (D50-D64)	6	0.4	*
14	Diabetes mellitus (E10-E14)	6	0.4	*
...	All other causes (Residual)	282	20.6	2.8

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 15-19 years				
...	All causes	3,741	100.0	35.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,951	52.2	18.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	292	7.8	2.8
3	Malignant neoplasms (C00-C97)	278	7.4	2.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	260	7.0	2.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	116	3.1	1.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	78	2.1	0.7
7	Pregnancy, childbirth and the puerperium (O00-O99)	44	1.2	0.4
8	Cerebrovascular diseases (I60-I69)	31	0.8	0.3
9	Septicemia (A40-A41)	29	0.8	0.3
9	Influenza and pneumonia (J09-J18)	29	0.8	0.3
11	Chronic lower respiratory diseases (J40-J47)	28	0.7	0.3
12	Diabetes mellitus (E10-E14)	24	0.6	0.2
13	Anemias (D50-D64)	20	0.5	0.2
14	Human immunodeficiency virus (HIV) disease (B20-B24)	19	0.5	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	0.4	*
...	All other causes (Residual)	526	14.1	5.0
All races, female, 20-24 years				
...	All causes	4,925	100.0	48.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,041	41.4	20.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	430	8.7	4.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	399	8.1	3.9
4	Malignant neoplasms (C00-C97)	388	7.9	3.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	268	5.4	2.6
6	Pregnancy, childbirth and the puerperium (O00-O99)	116	2.4	1.1
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	83	1.7	0.8
8	Human immunodeficiency virus (HIV) disease (B20-B24)	53	1.1	0.5
9	Cerebrovascular diseases (I60-I69)	51	1.0	0.5
10	Septicemia (A40-A41)	46	0.9	0.5
10	Diabetes mellitus (E10-E14)	46	0.9	0.5
12	Influenza and pneumonia (J09-J18)	43	0.9	0.4
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	31	0.6	0.3
14	Chronic lower respiratory diseases (J40-J47)	25	0.5	0.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	22	0.4	0.2
...	All other causes (Residual)	883	17.9	8.7

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 25-29 years				
...	All causes	5,824	100.0	56.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,888	32.4	18.4
2	Malignant neoplasms (C00-C97)	603	10.4	5.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	483	8.3	4.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	422	7.2	4.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	411	7.1	4.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	167	2.9	1.6
7	Pregnancy, childbirth and the puerperium (O00-O99)	158	2.7	1.5
8	Diabetes mellitus (E10-E14)	92	1.6	0.9
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	92	1.6	0.9
10	Cerebrovascular diseases (I60-I69)	90	1.5	0.9
11	Influenza and pneumonia (J09-J18)	65	1.1	0.6
12	Septicemia (A40-A41)	61	1.0	0.6
13	Chronic lower respiratory diseases (J40-J47)	51	0.9	0.5
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	48	0.8	0.5
15	Chronic liver disease and cirrhosis (K70,K73-K74)	43	0.7	0.4
...	All other causes (Residual)	1,150	19.7	11.2
All races, female, 30-34 years				
...	All causes	6,956	100.0	72.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,705	24.5	17.7
2	Malignant neoplasms (C00-C97)	1,168	16.8	12.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	624	9.0	6.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	514	7.4	5.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	328	4.7	3.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	271	3.9	2.8
7	Pregnancy, childbirth and the puerperium (O00-O99)	168	2.4	1.7
8	Cerebrovascular diseases (I60-I69)	163	2.3	1.7
9	Diabetes mellitus (E10-E14)	159	2.3	1.7
10	Septicemia (A40-A41)	83	1.2	0.9
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	82	1.2	0.9
12	Chronic liver disease and cirrhosis (K70,K73-K74)	80	1.2	0.8
13	Influenza and pneumonia (J09-J18)	79	1.1	0.8
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	72	1.0	0.7
15	Chronic lower respiratory diseases (J40-J47)	62	0.9	0.6
...	All other causes (Residual)	1,398	20.1	14.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 35-39 years				
...	All causes	11,126	100.0	105.7
1	Malignant neoplasms (C00-C97)	2,480	22.3	23.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,155	19.4	20.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,194	10.7	11.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	662	6.0	6.3
5	Human immunodeficiency virus (HIV) disease (B20-B24)	459	4.1	4.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	365	3.3	3.5
7	Cerebrovascular diseases (I60-I69)	332	3.0	3.2
8	Diabetes mellitus (E10-E14)	270	2.4	2.6
9	Chronic liver disease and cirrhosis (K70,K73-K74)	250	2.2	2.4
10	Septicemia (A40-A41)	159	1.4	1.5
11	Pregnancy, childbirth and the puerperium (O00-O99)	156	1.4	1.5
12	Influenza and pneumonia (J09-J18)	133	1.2	1.3
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	130	1.2	1.2
14	Chronic lower respiratory diseases (J40-J47)	120	1.1	1.1
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	114	1.0	1.1
...	All other causes (Residual)	2,147	19.3	20.4
All races, female, 40-44 years				
...	All causes	18,375	100.0	166.7
1	Malignant neoplasms (C00-C97)	5,107	27.8	46.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,754	15.0	25.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,300	12.5	20.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	908	4.9	8.2
5	Cerebrovascular diseases (I60-I69)	658	3.6	6.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	607	3.3	5.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	598	3.3	5.4
8	Diabetes mellitus (E10-E14)	486	2.6	4.4
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	398	2.2	3.6
10	Septicemia (A40-A41)	288	1.6	2.6
11	Chronic lower respiratory diseases (J40-J47)	284	1.5	2.6
12	Influenza and pneumonia (J09-J18)	208	1.1	1.9
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	196	1.1	1.8
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	142	0.8	1.3
15	Viral hepatitis (B15-B19)	134	0.7	1.2
...	All other causes (Residual)	3,307	18.0	30.0

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 45-49 years				
...	All causes	29,834	100.0	258.1
1	Malignant neoplasms (C00-C97)	9,877	33.1	85.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,241	14.2	36.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,348	11.2	29.0
4	Cerebrovascular diseases (I60-I69)	1,206	4.0	10.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,111	3.7	9.6
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,008	3.4	8.7
7	Diabetes mellitus (E10-E14)	785	2.6	6.8
8	Chronic lower respiratory diseases (J40-J47)	694	2.3	6.0
9	Human immunodeficiency virus (HIV) disease (B20-B24)	643	2.2	5.6
10	Septicemia (A40-A41)	462	1.5	4.0
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	398	1.3	3.4
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	353	1.2	3.1
13	Influenza and pneumonia (J09-J18)	334	1.1	2.9
14	Viral hepatitis (B15-B19)	307	1.0	2.7
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	164	0.5	1.4
...	All other causes (Residual)	4,903	16.4	42.4
All races, female, 50-54 years				
...	All causes	40,396	100.0	376.8
1	Malignant neoplasms (C00-C97)	14,920	36.9	139.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,282	15.6	58.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,950	7.3	27.5
4	Cerebrovascular diseases (I60-I69)	1,672	4.1	15.6
5	Diabetes mellitus (E10-E14)	1,455	3.6	13.6
6	Chronic lower respiratory diseases (J40-J47)	1,334	3.3	12.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,313	3.3	12.2
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	946	2.3	8.8
9	Septicemia (A40-A41)	664	1.6	6.2
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	575	1.4	5.4
11	Viral hepatitis (B15-B19)	511	1.3	4.8
12	Human immunodeficiency virus (HIV) disease (B20-B24)	485	1.2	4.5
13	Influenza and pneumonia (J09-J18)	449	1.1	4.2
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	311	0.8	2.9
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	215	0.5	2.0
...	All other causes (Residual)	6,314	15.6	58.9

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 55-59 years				
...	All causes	50,868	100.0	541.8
1	Malignant neoplasms (C00-C97)	20,736	40.8	220.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,566	16.8	91.2
3	Chronic lower respiratory diseases (J40-J47)	2,163	4.3	23.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,156	4.2	23.0
5	Diabetes mellitus (E10-E14)	2,043	4.0	21.8
6	Cerebrovascular diseases (I60-I69)	1,988	3.9	21.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,207	2.4	12.9
8	Septicemia (A40-A41)	907	1.8	9.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	867	1.7	9.2
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	750	1.5	8.0
11	Influenza and pneumonia (J09-J18)	562	1.1	6.0
12	Viral hepatitis (B15-B19)	414	0.8	4.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	374	0.7	4.0
14	Human immunodeficiency virus (HIV) disease (B20-B24)	260	0.5	2.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	254	0.5	2.7
...	All other causes (Residual)	7,621	15.0	81.2
All races, female, 60-64 years				
...	All causes	62,624	100.0	829.7
1	Malignant neoplasms (C00-C97)	25,876	41.3	342.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,403	18.2	151.1
3	Chronic lower respiratory diseases (J40-J47)	3,881	6.2	51.4
4	Diabetes mellitus (E10-E14)	2,617	4.2	34.7
5	Cerebrovascular diseases (I60-I69)	2,520	4.0	33.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,680	2.7	22.3
7	Septicemia (A40-A41)	1,121	1.8	14.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,095	1.7	14.5
9	Chronic liver disease and cirrhosis (K70,K73-K74)	1,065	1.7	14.1
10	Influenza and pneumonia (J09-J18)	709	1.1	9.4
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	493	0.8	6.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	477	0.8	6.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	324	0.5	4.3
14	Alzheimer's disease (G30)	275	0.4	3.6
15	Viral hepatitis (B15-B19)	238	0.4	3.2
...	All other causes (Residual)	8,850	14.1	117.3

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 65-69 years				
...	All causes	74,499	100.0	1,299.4
1	Malignant neoplasms (C00-C97)	29,458	39.5	513.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,105	18.9	246.0
3	Chronic lower respiratory diseases (J40-J47)	5,768	7.7	100.6
4	Cerebrovascular diseases (I60-I69)	3,483	4.7	60.7
5	Diabetes mellitus (E10-E14)	2,965	4.0	51.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,555	2.1	27.1
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,438	1.9	25.1
8	Septicemia (A40-A41)	1,354	1.8	23.6
9	Influenza and pneumonia (J09-J18)	965	1.3	16.8
10	Chronic liver disease and cirrhosis (K70,K73-K74)	936	1.3	16.3
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	641	0.9	11.2
12	Alzheimer's disease (G30)	601	0.8	10.5
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	423	0.6	7.4
14	Aortic aneurysm and dissection (I71)	343	0.5	6.0
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	257	0.3	4.5
...	All other causes (Residual)	10,207	13.7	178.0
All races, female, 70-74 years				
...	All causes	96,395	100.0	2,037.2
1	Malignant neoplasms (C00-C97)	33,084	34.3	699.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	19,952	20.7	421.7
3	Chronic lower respiratory diseases (J40-J47)	8,217	8.5	173.7
4	Cerebrovascular diseases (I60-I69)	5,173	5.4	109.3
5	Diabetes mellitus (E10-E14)	3,868	4.0	81.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,153	2.2	45.5
7	Septicemia (A40-A41)	1,735	1.8	36.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,722	1.8	36.4
9	Alzheimer's disease (G30)	1,689	1.8	35.7
10	Influenza and pneumonia (J09-J18)	1,538	1.6	32.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1,012	1.0	21.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	891	0.9	18.8
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	625	0.6	13.2
14	Parkinson's disease (G20-G21)	577	0.6	12.2
15	Aortic aneurysm and dissection (I71)	575	0.6	12.2
...	All other causes (Residual)	13,584	14.1	287.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 75-79 years				
...	All causes	139,360	100.0	3,304.0
1	Malignant neoplasms (C00-C97)	38,500	27.6	912.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	32,243	23.1	764.4
3	Chronic lower respiratory diseases (J40-J47)	11,455	8.2	271.6
4	Cerebrovascular diseases (I60-I69)	9,105	6.5	215.9
5	Diabetes mellitus (E10-E14)	5,171	3.7	122.6
6	Alzheimer's disease (G30)	4,626	3.3	109.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,132	2.2	74.3
8	Influenza and pneumonia (J09-J18)	2,785	2.0	66.0
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,617	1.9	62.0
10	Septicemia (A40-A41)	2,429	1.7	57.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,380	1.0	32.7
12	Parkinson's disease (G20-G21)	1,374	1.0	32.6
13	Chronic liver disease and cirrhosis (K70,K73-K74)	957	0.7	22.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	905	0.6	21.5
15	Aortic aneurysm and dissection (I71)	873	0.6	20.7
...	All other causes (Residual)	21,808	15.6	517.0
All races, female, 80-84 years				
...	All causes	192,519	100.0	5,511.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	51,009	26.5	1,460.4
2	Malignant neoplasms (C00-C97)	39,181	20.4	1,121.7
3	Cerebrovascular diseases (I60-I69)	15,174	7.9	434.4
4	Chronic lower respiratory diseases (J40-J47)	13,147	6.8	376.4
5	Alzheimer's disease (G30)	10,133	5.3	290.1
6	Diabetes mellitus (E10-E14)	6,144	3.2	175.9
7	Influenza and pneumonia (J09-J18)	4,662	2.4	133.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,144	2.2	118.6
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,851	2.0	110.3
10	Septicemia (A40-A41)	3,206	1.7	91.8
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,395	1.2	68.6
12	Parkinson's disease (G20-G21)	2,295	1.2	65.7
13	Pneumonitis due to solids and liquids (J69)	1,510	0.8	43.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,242	0.6	35.6
15	Aortic aneurysm and dissection (I71)	1,132	0.6	32.4
...	All other causes (Residual)	33,294	17.3	953.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 85-89 years				
...	All causes	213,110	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	64,618	30.3	---
2	Malignant neoplasms (C00-C97)	28,867	13.5	---
3	Cerebrovascular diseases (I60-I69)	18,738	8.8	---
4	Alzheimer's disease (G30)	14,787	6.9	---
5	Chronic lower respiratory diseases (J40-J47)	10,995	5.2	---
6	Influenza and pneumonia (J09-J18)	6,330	3.0	---
7	Diabetes mellitus (E10-E14)	5,465	2.6	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,581	2.1	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,409	2.1	---
10	Septicemia (A40-A41)	3,271	1.5	---
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,201	1.5	---
12	Parkinson's disease (G20-G21)	2,235	1.0	---
13	Pneumonitis due to solids and liquids (J69)	1,949	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,315	0.6	---
15	Atherosclerosis (I70)	1,182	0.6	---
...	All other causes (Residual)	41,167	19.3	---
All races, female, 90-94 years				
...	All causes	162,703	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	55,380	34.0	---
2	Cerebrovascular diseases (I60-I69)	14,280	8.8	---
3	Malignant neoplasms (C00-C97)	14,062	8.6	---
4	Alzheimer's disease (G30)	13,155	8.1	---
5	Chronic lower respiratory diseases (J40-J47)	6,073	3.7	---
6	Influenza and pneumonia (J09-J18)	5,762	3.5	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,384	2.1	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,203	2.0	---
9	Diabetes mellitus (E10-E14)	3,126	1.9	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,767	1.7	---
11	Septicemia (A40-A41)	2,038	1.3	---
12	Pneumonitis due to solids and liquids (J69)	1,629	1.0	---
13	Parkinson's disease (G20-G21)	1,260	0.8	---
14	Atherosclerosis (I70)	1,125	0.7	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	848	0.5	---
...	All other causes (Residual)	34,611	21.3	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 95-99 years				
...	All causes	72,375	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	26,729	36.9	---
2	Alzheimer's disease (G30)	6,148	8.5	---
3	Cerebrovascular diseases (I60-I69)	5,939	8.2	---
4	Malignant neoplasms (C00-C97)	4,169	5.8	---
5	Influenza and pneumonia (J09-J18)	2,928	4.0	---
6	Chronic lower respiratory diseases (J40-J47)	2,009	2.8	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,515	2.1	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,369	1.9	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,252	1.7	---
10	Diabetes mellitus (E10-E14)	982	1.4	---
11	Septicemia (A40-A41)	780	1.1	---
12	Pneumonitis due to solids and liquids (J69)	746	1.0	---
13	Atherosclerosis (I70)	683	0.9	---
14	Parkinson's disease (G20-G21)	320	0.4	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	306	0.4	---
...	All other causes (Residual)	16,500	22.8	---
All races, female, 100 years and over				
...	All causes	16,593	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,402	38.6	---
2	Alzheimer's disease (G30)	1,235	7.4	---
3	Cerebrovascular diseases (I60-I69)	1,103	6.6	---
4	Influenza and pneumonia (J09-J18)	873	5.3	---
5	Malignant neoplasms (C00-C97)	594	3.6	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	394	2.4	---
7	Chronic lower respiratory diseases (J40-J47)	300	1.8	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	299	1.8	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	294	1.8	---
10	Atherosclerosis (I70)	240	1.4	---
11	Diabetes mellitus (E10-E14)	192	1.2	---
12	Pneumonitis due to solids and liquids (J69)	183	1.1	---
13	Septicemia (A40-A41)	161	1.0	---
14	Anemias (D50-D64)	55	0.3	---
15	Nutritional deficiencies (E40-E64)	51	0.3	---
...	All other causes (Residual)	4,217	25.4	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, all ages				
...	All causes	2,074,151	100.0	851.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	531,636	25.6	218.3
2	Malignant neoplasms (C00-C97)	483,939	23.3	198.7
3	Chronic lower respiratory diseases (J40-J47)	118,081	5.7	48.5
4	Cerebrovascular diseases (I60-I69)	114,695	5.5	47.1
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	106,252	5.1	43.6
6	Alzheimer's disease (G30)	68,933	3.3	28.3
7	Diabetes mellitus (E10-E14)	56,390	2.7	23.2
8	Influenza and pneumonia (J09-J18)	45,947	2.2	18.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	36,871	1.8	15.1
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	31,348	1.5	12.9
11	Septicemia (A40-A41)	27,750	1.3	11.4
12	Chronic liver disease and cirrhosis (K70,K73-K74)	25,490	1.2	10.5
13	Parkinson's disease (G20-G21)	18,864	0.9	7.7
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	18,583	0.9	7.6
15	Pneumonitis due to solids and liquids (J69)	15,229	0.7	6.3
...	All other causes (Residual)	374,143	18.0	153.6
White, both sexes, 1-4 years				
...	All causes	3,287	100.0	25.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,169	35.6	9.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	396	12.0	3.1
3	Malignant neoplasms (C00-C97)	287	8.7	2.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	216	6.6	1.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	112	3.4	0.9
6	Influenza and pneumonia (J09-J18)	73	2.2	0.6
7	Septicemia (A40-A41)	60	1.8	0.5
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	49	1.5	0.4
9	Certain conditions originating in the perinatal period (P00-P96)	40	1.2	0.3
10	Cerebrovascular diseases (I60-I69)	34	1.0	0.3
11	Chronic lower respiratory diseases (J40-J47)	22	0.7	0.2
12	Complications of medical and surgical care (Y40-Y84,Y88)	17	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	0.4	*
14	Meningitis (G00,G03)	12	0.4	*
15	Acute bronchitis and bronchiolitis (J20-J21)	9	0.3	*
...	All other causes (Residual)	777	23.6	6.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 5-9 years				
...	All causes	2,001	100.0	12.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	702	35.1	4.5
2	Malignant neoplasms (C00-C97)	393	19.6	2.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	145	7.2	0.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	94	4.7	0.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	79	3.9	0.5
6	Influenza and pneumonia (J09-J18)	36	1.8	0.2
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	32	1.6	0.2
8	Septicemia (A40-A41)	28	1.4	0.2
8	Cerebrovascular diseases (I60-I69)	28	1.4	0.2
10	Chronic lower respiratory diseases (J40-J47)	26	1.3	0.2
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.4	*
12	Anemias (D50-D64)	8	0.4	*
12	Diabetes mellitus (E10-E14)	8	0.4	*
14	Meningitis (G00,G03)	6	0.3	*
14	Certain conditions originating in the perinatal period (P00-P96)	6	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.3	*
...	All other causes (Residual)	395	19.7	2.6
White, both sexes, 10-14 years				
...	All causes	2,479	100.0	15.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	923	37.2	5.9
2	Malignant neoplasms (C00-C97)	383	15.4	2.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	129	5.2	0.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	123	5.0	0.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	102	4.1	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	87	3.5	0.6
7	Influenza and pneumonia (J09-J18)	41	1.7	0.3
8	Cerebrovascular diseases (I60-I69)	35	1.4	0.2
9	Septicemia (A40-A41)	31	1.3	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	28	1.1	0.2
10	Chronic lower respiratory diseases (J40-J47)	28	1.1	0.2
12	Meningitis (G00,G03)	13	0.5	*
13	Anemias (D50-D64)	10	0.4	*
13	Certain conditions originating in the perinatal period (P00-P96)	10	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.4	*
...	All other causes (Residual)	527	21.3	3.3

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 15-19 years				
...	All causes	9,765	100.0	58.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,475	56.1	32.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,234	12.6	7.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	846	8.7	5.1
4	Malignant neoplasms (C00-C97)	526	5.4	3.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	224	2.3	1.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	133	1.4	0.8
7	Cerebrovascular diseases (I60-I69)	52	0.5	0.3
8	Influenza and pneumonia (J09-J18)	40	0.4	0.2
9	Chronic lower respiratory diseases (J40-J47)	39	0.4	0.2
10	Septicemia (A40-A41)	37	0.4	0.2
11	Diabetes mellitus (E10-E14)	34	0.3	0.2
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	24	0.2	0.1
13	Pregnancy, childbirth and the puerperium (O00-O99)	19	0.2	*
14	Pneumonitis due to solids and liquids (J69)	15	0.2	*
15	Legal intervention (Y35,Y89.0)	14	0.1	*
...	All other causes (Residual)	1,053	10.8	6.3
White, both sexes, 20-24 years				
...	All causes	15,372	100.0	93.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,006	52.1	48.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,208	14.4	13.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,347	8.8	8.1
4	Malignant neoplasms (C00-C97)	770	5.0	4.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	482	3.1	2.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	166	1.1	1.0
7	Influenza and pneumonia (J09-J18)	89	0.6	0.5
8	Cerebrovascular diseases (I60-I69)	86	0.6	0.5
9	Diabetes mellitus (E10-E14)	75	0.5	0.5
10	Septicemia (A40-A41)	69	0.4	0.4
11	Pregnancy, childbirth and the puerperium (O00-O99)	67	0.4	0.4
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	45	0.3	0.3
13	Human immunodeficiency virus (HIV) disease (B20-B24)	43	0.3	0.3
14	Chronic lower respiratory diseases (J40-J47)	40	0.3	0.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	35	0.2	0.2
...	All other causes (Residual)	1,844	12.0	11.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 25-29 years				
...	All causes	15,382	100.0	93.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,915	45.0	41.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,277	14.8	13.8
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,127	7.3	6.8
4	Malignant neoplasms (C00-C97)	1,017	6.6	6.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	803	5.2	4.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	173	1.1	1.0
7	Cerebrovascular diseases (I60-I69)	129	0.8	0.8
8	Diabetes mellitus (E10-E14)	128	0.8	0.8
9	Human immunodeficiency virus (HIV) disease (B20-B24)	122	0.8	0.7
10	Influenza and pneumonia (J09-J18)	107	0.7	0.6
11	Pregnancy, childbirth and the puerperium (O00-O99)	91	0.6	0.6
12	Chronic liver disease and cirrhosis (K70,K73-K74)	88	0.6	0.5
13	Chronic lower respiratory diseases (J40-J47)	78	0.5	0.5
14	Septicemia (A40-A41)	77	0.5	0.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	53	0.3	0.3
...	All other causes (Residual)	2,197	14.3	13.3
White, both sexes, 30-34 years				
...	All causes	15,799	100.0	103.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,731	36.3	37.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,247	14.2	14.7
3	Malignant neoplasms (C00-C97)	1,643	10.4	10.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,270	8.0	8.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	923	5.8	6.0
6	Diabetes mellitus (E10-E14)	251	1.6	1.6
7	Human immunodeficiency virus (HIV) disease (B20-B24)	245	1.6	1.6
8	Chronic liver disease and cirrhosis (K70,K73-K74)	222	1.4	1.5
9	Cerebrovascular diseases (I60-I69)	203	1.3	1.3
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	149	0.9	1.0
11	Influenza and pneumonia (J09-J18)	118	0.7	0.8
12	Septicemia (A40-A41)	117	0.7	0.8
13	Pregnancy, childbirth and the puerperium (O00-O99)	102	0.6	0.7
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	90	0.6	0.6
15	Chronic lower respiratory diseases (J40-J47)	81	0.5	0.5
...	All other causes (Residual)	2,407	15.2	15.8

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 35-39 years				
...	All causes	22,804	100.0	135.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,446	28.3	38.4
2	Malignant neoplasms (C00-C97)	3,301	14.5	19.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,670	11.7	15.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,665	11.7	15.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	784	3.4	4.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	637	2.8	3.8
7	Human immunodeficiency virus (HIV) disease (B20-B24)	546	2.4	3.3
8	Diabetes mellitus (E10-E14)	499	2.2	3.0
9	Cerebrovascular diseases (I60-I69)	470	2.1	2.8
10	Septicemia (A40-A41)	215	0.9	1.3
11	Influenza and pneumonia (J09-J18)	204	0.9	1.2
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	184	0.8	1.1
13	Chronic lower respiratory diseases (J40-J47)	155	0.7	0.9
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	140	0.6	0.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	106	0.5	0.6
...	All other causes (Residual)	3,782	16.6	22.5
White, both sexes, 40-44 years				
...	All causes	36,767	100.0	207.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,765	21.1	43.9
2	Malignant neoplasms (C00-C97)	6,988	19.0	39.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,736	15.6	32.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,368	9.2	19.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,547	4.2	8.7
6	Human immunodeficiency virus (HIV) disease (B20-B24)	976	2.7	5.5
7	Cerebrovascular diseases (I60-I69)	886	2.4	5.0
8	Diabetes mellitus (E10-E14)	845	2.3	4.8
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	793	2.2	4.5
10	Chronic lower respiratory diseases (J40-J47)	423	1.2	2.4
11	Septicemia (A40-A41)	393	1.1	2.2
12	Influenza and pneumonia (J09-J18)	376	1.0	2.1
13	Viral hepatitis (B15-B19)	292	0.8	1.7
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	266	0.7	1.5
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	237	0.6	1.3
...	All other causes (Residual)	5,876	16.0	33.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 45-49 years				
...	All causes	59,635	100.0	318.7
1	Malignant neoplasms (C00-C97)	14,686	24.6	78.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,164	18.7	59.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,024	15.1	48.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,778	6.3	20.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	3,021	5.1	16.1
6	Cerebrovascular diseases (I60-I69)	1,682	2.8	9.0
7	Diabetes mellitus (E10-E14)	1,544	2.6	8.3
8	Chronic lower respiratory diseases (J40-J47)	1,102	1.8	5.9
9	Human immunodeficiency virus (HIV) disease (B20-B24)	982	1.6	5.2
10	Viral hepatitis (B15-B19)	861	1.4	4.6
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	713	1.2	3.8
12	Septicemia (A40-A41)	663	1.1	3.5
13	Influenza and pneumonia (J09-J18)	558	0.9	3.0
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	515	0.9	2.8
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	264	0.4	1.4
...	All other causes (Residual)	9,078	15.2	48.5
White, both sexes, 50-54 years				
...	All causes	82,598	100.0	474.9
1	Malignant neoplasms (C00-C97)	24,871	30.1	143.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	16,971	20.5	97.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,994	9.7	46.0
4	Chronic liver disease and cirrhosis (K70,K73-K74)	3,986	4.8	22.9
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,481	4.2	20.0
6	Diabetes mellitus (E10-E14)	2,579	3.1	14.8
7	Cerebrovascular diseases (I60-I69)	2,457	3.0	14.1
8	Chronic lower respiratory diseases (J40-J47)	2,293	2.8	13.2
9	Viral hepatitis (B15-B19)	1,484	1.8	8.5
10	Septicemia (A40-A41)	965	1.2	5.5
11	Influenza and pneumonia (J09-J18)	845	1.0	4.9
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	831	1.0	4.8
13	Human immunodeficiency virus (HIV) disease (B20-B24)	765	0.9	4.4
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	558	0.7	3.2
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	447	0.5	2.6
...	All other causes (Residual)	12,071	14.6	69.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 55-59 years				
...	All causes	104,284	100.0	682.5
1	Malignant neoplasms (C00-C97)	36,779	35.3	240.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	22,925	22.0	150.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,879	5.6	38.5
4	Chronic lower respiratory diseases (J40-J47)	4,037	3.9	26.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	3,781	3.6	24.7
6	Diabetes mellitus (E10-E14)	3,673	3.5	24.0
7	Cerebrovascular diseases (I60-I69)	3,192	3.1	20.9
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,802	2.7	18.3
9	Septicemia (A40-A41)	1,400	1.3	9.2
10	Viral hepatitis (B15-B19)	1,292	1.2	8.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,271	1.2	8.3
12	Influenza and pneumonia (J09-J18)	1,071	1.0	7.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	561	0.5	3.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	533	0.5	3.5
15	Aortic aneurysm and dissection (I71)	462	0.4	3.0
...	All other causes (Residual)	14,626	14.0	95.7
White, both sexes, 60-64 years				
...	All causes	127,003	100.0	1,023.5
1	Malignant neoplasms (C00-C97)	48,594	38.3	391.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	28,660	22.6	231.0
3	Chronic lower respiratory diseases (J40-J47)	7,375	5.8	59.4
4	Diabetes mellitus (E10-E14)	4,705	3.7	37.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,345	3.4	35.0
6	Cerebrovascular diseases (I60-I69)	4,246	3.3	34.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	3,098	2.4	25.0
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,990	1.6	16.0
9	Septicemia (A40-A41)	1,720	1.4	13.9
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,621	1.3	13.1
11	Influenza and pneumonia (J09-J18)	1,412	1.1	11.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	752	0.6	6.1
13	Aortic aneurysm and dissection (I71)	730	0.6	5.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	657	0.5	5.3
15	Viral hepatitis (B15-B19)	595	0.5	4.8
...	All other causes (Residual)	16,503	13.0	133.0

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 65-69 years				
...	All causes	145,663	100.0	1,581.1
1	Malignant neoplasms (C00-C97)	55,933	38.4	607.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	32,151	22.1	349.0
3	Chronic lower respiratory diseases (J40-J47)	10,944	7.5	118.8
4	Cerebrovascular diseases (I60-I69)	5,660	3.9	61.4
5	Diabetes mellitus (E10-E14)	5,264	3.6	57.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,558	2.4	38.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,448	1.7	26.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,303	1.6	25.0
9	Septicemia (A40-A41)	2,113	1.5	22.9
10	Influenza and pneumonia (J09-J18)	1,787	1.2	19.4
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,312	0.9	14.2
12	Alzheimer's disease (G30)	976	0.7	10.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	927	0.6	10.1
14	Aortic aneurysm and dissection (I71)	925	0.6	10.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	867	0.6	9.4
...	All other causes (Residual)	18,495	12.7	200.7
White, both sexes, 70-74 years				
...	All causes	181,714	100.0	2,454.7
1	Malignant neoplasms (C00-C97)	62,893	34.6	849.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	41,449	22.8	559.9
3	Chronic lower respiratory diseases (J40-J47)	15,445	8.5	208.6
4	Cerebrovascular diseases (I60-I69)	8,363	4.6	113.0
5	Diabetes mellitus (E10-E14)	6,388	3.5	86.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,971	2.2	53.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,352	1.8	45.3
8	Influenza and pneumonia (J09-J18)	2,764	1.5	37.3
9	Septicemia (A40-A41)	2,739	1.5	37.0
10	Alzheimer's disease (G30)	2,587	1.4	34.9
11	Chronic liver disease and cirrhosis (K70,K73-K74)	2,176	1.2	29.4
12	Parkinson's disease (G20-G21)	1,478	0.8	20.0
13	Aortic aneurysm and dissection (I71)	1,370	0.8	18.5
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,263	0.7	17.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,218	0.7	16.5
...	All other causes (Residual)	24,258	13.3	327.7

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 75-79 years				
...	All causes	253,164	100.0	3,933.2
1	Malignant neoplasms (C00-C97)	73,267	28.9	1,138.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	62,048	24.5	964.0
3	Chronic lower respiratory diseases (J40-J47)	21,215	8.4	329.6
4	Cerebrovascular diseases (I60-I69)	14,354	5.7	223.0
5	Diabetes mellitus (E10-E14)	8,144	3.2	126.5
6	Alzheimer's disease (G30)	6,861	2.7	106.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,324	2.1	82.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,173	2.0	80.4
9	Influenza and pneumonia (J09-J18)	4,962	2.0	77.1
10	Septicemia (A40-A41)	3,790	1.5	58.9
11	Parkinson's disease (G20-G21)	3,484	1.4	54.1
12	Chronic liver disease and cirrhosis (K70,K73-K74)	1,979	0.8	30.7
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,964	0.8	30.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,871	0.7	29.1
15	Aortic aneurysm and dissection (I71)	1,807	0.7	28.1
...	All other causes (Residual)	36,921	14.6	573.6
White, both sexes, 80-84 years				
...	All causes	325,958	100.0	6,423.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	88,993	27.3	1,753.6
2	Malignant neoplasms (C00-C97)	72,318	22.2	1,425.0
3	Chronic lower respiratory diseases (J40-J47)	23,771	7.3	468.4
4	Cerebrovascular diseases (I60-I69)	21,986	6.7	433.2
5	Alzheimer's disease (G30)	14,254	4.4	280.9
6	Diabetes mellitus (E10-E14)	9,267	2.8	182.6
7	Influenza and pneumonia (J09-J18)	8,112	2.5	159.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,260	2.2	143.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,109	2.2	140.1
10	Parkinson's disease (G20-G21)	5,348	1.6	105.4
11	Septicemia (A40-A41)	4,894	1.5	96.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,149	1.0	62.1
13	Pneumonitis due to solids and liquids (J69)	2,989	0.9	58.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,430	0.7	47.9
15	Aortic aneurysm and dissection (I71)	2,152	0.7	42.4
...	All other causes (Residual)	51,926	15.9	1,023.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 85-89 years				
...	All causes	327,592	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	100,624	30.7	---
2	Malignant neoplasms (C00-C97)	50,601	15.4	---
3	Cerebrovascular diseases (I60-I69)	25,409	7.8	---
4	Alzheimer's disease (G30)	19,709	6.0	---
5	Chronic lower respiratory diseases (J40-J47)	18,643	5.7	---
6	Influenza and pneumonia (J09-J18)	10,159	3.1	---
7	Diabetes mellitus (E10-E14)	7,748	2.4	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,609	2.3	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,307	2.2	---
10	Parkinson's disease (G20-G21)	4,706	1.4	---
11	Septicemia (A40-A41)	4,615	1.4	---
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,995	1.2	---
13	Pneumonitis due to solids and liquids (J69)	3,734	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,210	0.7	---
15	Aortic aneurysm and dissection (I71)	1,823	0.6	---
...	All other causes (Residual)	58,700	17.9	---
White, both sexes, 90-94 years				
...	All causes	220,511	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	76,087	34.5	---
2	Malignant neoplasms (C00-C97)	22,208	10.1	---
3	Cerebrovascular diseases (I60-I69)	17,531	8.0	---
4	Alzheimer's disease (G30)	15,884	7.2	---
5	Chronic lower respiratory diseases (J40-J47)	9,256	4.2	---
6	Influenza and pneumonia (J09-J18)	8,318	3.8	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,980	2.3	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,704	2.1	---
9	Diabetes mellitus (E10-E14)	3,968	1.8	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,233	1.5	---
11	Pneumonitis due to solids and liquids (J69)	2,763	1.3	---
12	Septicemia (A40-A41)	2,636	1.2	---
13	Parkinson's disease (G20-G21)	2,262	1.0	---
14	Atherosclerosis (I70)	1,492	0.7	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,286	0.6	---
...	All other causes (Residual)	43,903	19.9	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 95-99 years				
...	All causes	86,150	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	32,059	37.2	---
2	Alzheimer's disease (G30)	6,703	7.8	---
3	Cerebrovascular diseases (I60-I69)	6,688	7.8	---
4	Malignant neoplasms (C00-C97)	5,712	6.6	---
5	Influenza and pneumonia (J09-J18)	3,755	4.4	---
6	Chronic lower respiratory diseases (J40-J47)	2,734	3.2	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,998	2.3	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,631	1.9	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,451	1.7	---
10	Pneumonitis due to solids and liquids (J69)	1,079	1.3	---
11	Diabetes mellitus (E10-E14)	1,072	1.2	---
12	Septicemia (A40-A41)	882	1.0	---
13	Atherosclerosis (I70)	785	0.9	---
14	Parkinson's disease (G20-G21)	480	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	388	0.5	---
...	All other causes (Residual)	18,733	21.7	---
White, both sexes, 100 years and over				
...	All causes	17,268	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,769	39.2	---
2	Alzheimer's disease (G30)	1,188	6.9	---
3	Cerebrovascular diseases (I60-I69)	1,106	6.4	---
4	Influenza and pneumonia (J09-J18)	985	5.7	---
5	Malignant neoplasms (C00-C97)	705	4.1	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	369	2.1	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	346	2.0	---
8	Chronic lower respiratory diseases (J40-J47)	343	2.0	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	324	1.9	---
10	Atherosclerosis (I70)	254	1.5	---
11	Pneumonitis due to solids and liquids (J69)	216	1.3	---
12	Diabetes mellitus (E10-E14)	180	1.0	---
13	Septicemia (A40-A41)	147	0.9	---
14	Anemias (D50-D64)	59	0.3	---
15	Parkinson's disease (G20-G21)	49	0.3	---
...	All other causes (Residual)	4,228	24.5	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, male, all ages				
...	All causes	1,023,951	100.0	848.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	266,908	26.1	221.1
2	Malignant neoplasms (C00-C97)	252,049	24.6	208.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	68,059	6.6	56.4
4	Chronic lower respiratory diseases (J40-J47)	55,934	5.5	46.3
5	Cerebrovascular diseases (I60-I69)	44,714	4.4	37.0
6	Diabetes mellitus (E10-E14)	28,744	2.8	23.8
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	24,725	2.4	20.5
8	Influenza and pneumonia (J09-J18)	20,720	2.0	17.2
9	Alzheimer's disease (G30)	20,185	2.0	16.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18,242	1.8	15.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	16,801	1.6	13.9
12	Septicemia (A40-A41)	12,600	1.2	10.4
13	Parkinson's disease (G20-G21)	10,895	1.1	9.0
14	Pneumonitis due to solids and liquids (J69)	8,010	0.8	6.6
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7,106	0.7	5.9
...	All other causes (Residual)	168,259	16.4	139.4
White, male, 1-4 years				
...	All causes	1,847	100.0	28.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	738	40.0	11.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	211	11.4	3.2
3	Malignant neoplasms (C00-C97)	146	7.9	2.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	116	6.3	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	47	2.5	0.7
6	Septicemia (A40-A41)	33	1.8	0.5
7	Influenza and pneumonia (J09-J18)	32	1.7	0.5
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	25	1.4	0.4
9	Certain conditions originating in the perinatal period (P00-P96)	19	1.0	*
10	Cerebrovascular diseases (I60-I69)	16	0.9	*
11	Chronic lower respiratory diseases (J40-J47)	13	0.7	*
12	Meningitis (G00,G03)	7	0.4	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.3	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.3	*
15	Pneumonitis due to solids and liquids (J69)	5	0.3	*
...	All other causes (Residual)	427	23.1	6.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 5-9 years				
...	All causes	1,112	100.0	14.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	414	37.2	5.2
2	Malignant neoplasms (C00-C97)	208	18.7	2.6
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	74	6.7	0.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	44	4.0	0.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	42	3.8	0.5
6	Septicemia (A40-A41)	21	1.9	0.3
7	Influenza and pneumonia (J09-J18)	18	1.6	*
8	Chronic lower respiratory diseases (J40-J47)	16	1.4	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	15	1.3	*
10	Cerebrovascular diseases (I60-I69)	14	1.3	*
11	Anemias (D50-D64)	6	0.5	*
11	Diabetes mellitus (E10-E14)	6	0.5	*
13	Certain conditions originating in the perinatal period (P00-P96)	5	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.4	*
...	All other causes (Residual)	221	19.9	2.8
White, male, 10-14 years				
...	All causes	1,478	100.0	18.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	586	39.6	7.2
2	Malignant neoplasms (C00-C97)	196	13.3	2.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	90	6.1	1.1
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	71	4.8	0.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	62	4.2	0.8
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	48	3.2	0.6
7	Cerebrovascular diseases (I60-I69)	18	1.2	*
7	Influenza and pneumonia (J09-J18)	18	1.2	*
7	Chronic lower respiratory diseases (J40-J47)	18	1.2	*
10	Septicemia (A40-A41)	17	1.2	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	1.1	*
12	Anemias (D50-D64)	7	0.5	*
12	Meningitis (G00,G03)	7	0.5	*
12	Pneumonitis due to solids and liquids (J69)	7	0.5	*
15	Certain conditions originating in the perinatal period (P00-P96)	6	0.4	*
...	All other causes (Residual)	311	21.0	3.8

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 15-19 years				
...	All causes	6,851	100.0	80.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,800	55.5	44.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,017	14.8	11.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	684	10.0	8.0
4	Malignant neoplasms (C00-C97)	315	4.6	3.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	143	2.1	1.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	77	1.1	0.9
7	Cerebrovascular diseases (I60-I69)	28	0.4	0.3
8	Chronic lower respiratory diseases (J40-J47)	23	0.3	0.3
9	Diabetes mellitus (E10-E14)	22	0.3	0.3
10	Influenza and pneumonia (J09-J18)	19	0.3	*
11	Septicemia (A40-A41)	16	0.2	*
12	Legal intervention (Y35,Y89.0)	14	0.2	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	0.2	*
14	Pneumonitis due to solids and liquids (J69)	11	0.2	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.1	*
...	All other causes (Residual)	663	9.7	7.7
White, male, 20-24 years				
...	All causes	11,662	100.0	136.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,321	54.2	73.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,879	16.1	21.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,108	9.5	12.9
4	Malignant neoplasms (C00-C97)	466	4.0	5.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	302	2.6	3.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	98	0.8	1.1
7	Influenza and pneumonia (J09-J18)	55	0.5	0.6
8	Diabetes mellitus (E10-E14)	51	0.4	0.6
9	Cerebrovascular diseases (I60-I69)	48	0.4	0.6
10	Septicemia (A40-A41)	43	0.4	0.5
11	Legal intervention (Y35,Y89.0)	32	0.3	0.4
12	Human immunodeficiency virus (HIV) disease (B20-B24)	29	0.2	0.3
13	Chronic lower respiratory diseases (J40-J47)	26	0.2	0.3
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	0.2	0.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	0.2	*
...	All other causes (Residual)	1,163	10.0	13.6

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 25-29 years				
...	All causes	11,159	100.0	130.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,316	47.6	62.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,871	16.8	21.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	899	8.1	10.5
4	Malignant neoplasms (C00-C97)	564	5.1	6.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	541	4.8	6.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	104	0.9	1.2
7	Human immunodeficiency virus (HIV) disease (B20-B24)	89	0.8	1.0
8	Diabetes mellitus (E10-E14)	79	0.7	0.9
9	Cerebrovascular diseases (I60-I69)	69	0.6	0.8
10	Influenza and pneumonia (J09-J18)	61	0.5	0.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	53	0.5	0.6
12	Chronic lower respiratory diseases (J40-J47)	44	0.4	0.5
13	Septicemia (A40-A41)	42	0.4	0.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	38	0.3	0.4
15	Legal intervention (Y35,Y89.0)	37	0.3	0.4
...	All other causes (Residual)	1,352	12.1	15.8
White, male, 30-34 years				
...	All causes	10,839	100.0	138.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,308	39.7	54.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,805	16.7	23.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	892	8.2	11.4
4	Malignant neoplasms (C00-C97)	765	7.1	9.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	726	6.7	9.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	174	1.6	2.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	166	1.5	2.1
8	Diabetes mellitus (E10-E14)	155	1.4	2.0
9	Cerebrovascular diseases (I60-I69)	98	0.9	1.2
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	81	0.7	1.0
11	Influenza and pneumonia (J09-J18)	66	0.6	0.8
12	Septicemia (A40-A41)	61	0.6	0.8
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	49	0.5	0.6
14	Legal intervention (Y35,Y89.0)	44	0.4	0.6
15	Chronic lower respiratory diseases (J40-J47)	42	0.4	0.5
...	All other causes (Residual)	1,407	13.0	17.9

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 35-39 years				
...	All causes	14,820	100.0	173.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,645	31.3	54.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,083	14.1	24.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,897	12.8	22.2
4	Malignant neoplasms (C00-C97)	1,412	9.5	16.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	573	3.9	6.7
6	Human immunodeficiency virus (HIV) disease (B20-B24)	441	3.0	5.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	425	2.9	5.0
8	Diabetes mellitus (E10-E14)	328	2.2	3.8
9	Cerebrovascular diseases (I60-I69)	251	1.7	2.9
10	Septicemia (A40-A41)	115	0.8	1.3
11	Influenza and pneumonia (J09-J18)	111	0.7	1.3
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	95	0.6	1.1
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	88	0.6	1.0
14	Aortic aneurysm and dissection (I71)	72	0.5	0.8
15	Chronic lower respiratory diseases (J40-J47)	70	0.5	0.8
...	All other causes (Residual)	2,214	14.9	25.9
White, male, 40-44 years				
...	All causes	23,301	100.0	261.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,429	23.3	60.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,161	17.9	46.6
3	Malignant neoplasms (C00-C97)	3,110	13.3	34.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,545	10.9	28.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,052	4.5	11.8
6	Human immunodeficiency virus (HIV) disease (B20-B24)	808	3.5	9.1
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	544	2.3	6.1
8	Diabetes mellitus (E10-E14)	533	2.3	6.0
9	Cerebrovascular diseases (I60-I69)	495	2.1	5.5
10	Influenza and pneumonia (J09-J18)	212	0.9	2.4
11	Chronic lower respiratory diseases (J40-J47)	208	0.9	2.3
12	Septicemia (A40-A41)	204	0.9	2.3
13	Viral hepatitis (B15-B19)	191	0.8	2.1
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	154	0.7	1.7
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	119	0.5	1.3
...	All other causes (Residual)	3,536	15.2	39.6

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 45-49 years				
...	All causes	37,447	100.0	400.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,257	22.0	88.3
2	Malignant neoplasms (C00-C97)	7,014	18.7	75.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,202	16.6	66.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,842	7.6	30.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,116	5.7	22.6
6	Diabetes mellitus (E10-E14)	1,008	2.7	10.8
7	Cerebrovascular diseases (I60-I69)	925	2.5	9.9
8	Human immunodeficiency virus (HIV) disease (B20-B24)	791	2.1	8.5
9	Viral hepatitis (B15-B19)	605	1.6	6.5
10	Chronic lower respiratory diseases (J40-J47)	568	1.5	6.1
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	493	1.3	5.3
12	Septicemia (A40-A41)	366	1.0	3.9
13	Influenza and pneumonia (J09-J18)	312	0.8	3.3
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	300	0.8	3.2
15	Aortic aneurysm and dissection (I71)	157	0.4	1.7
...	All other causes (Residual)	5,491	14.7	58.7
White, male, 50-54 years				
...	All causes	52,109	100.0	604.8
1	Malignant neoplasms (C00-C97)	13,119	25.2	152.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,619	24.2	146.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,523	10.6	64.1
4	Chronic liver disease and cirrhosis (K70,K73-K74)	2,891	5.5	33.6
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,594	5.0	30.1
6	Diabetes mellitus (E10-E14)	1,583	3.0	18.4
7	Cerebrovascular diseases (I60-I69)	1,405	2.7	16.3
8	Chronic lower respiratory diseases (J40-J47)	1,209	2.3	14.0
9	Viral hepatitis (B15-B19)	1,081	2.1	12.5
10	Human immunodeficiency virus (HIV) disease (B20-B24)	631	1.2	7.3
11	Septicemia (A40-A41)	512	1.0	5.9
12	Influenza and pneumonia (J09-J18)	498	1.0	5.8
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	488	0.9	5.7
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	410	0.8	4.8
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	284	0.5	3.3
...	All other causes (Residual)	7,262	13.9	84.3

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 55-59 years				
...	All causes	64,772	100.0	863.5
1	Malignant neoplasms (C00-C97)	19,986	30.9	266.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	16,748	25.9	223.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,072	6.3	54.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	2,786	4.3	37.1
5	Diabetes mellitus (E10-E14)	2,270	3.5	30.3
6	Chronic lower respiratory diseases (J40-J47)	2,161	3.3	28.8
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,089	3.2	27.9
8	Cerebrovascular diseases (I60-I69)	1,851	2.9	24.7
9	Viral hepatitis (B15-B19)	979	1.5	13.1
10	Septicemia (A40-A41)	748	1.2	10.0
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	740	1.1	9.9
12	Influenza and pneumonia (J09-J18)	653	1.0	8.7
13	Human immunodeficiency virus (HIV) disease (B20-B24)	375	0.6	5.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	375	0.6	5.0
15	Aortic aneurysm and dissection (I71)	352	0.5	4.7
...	All other causes (Residual)	8,587	13.3	114.5
White, male, 60-64 years				
...	All causes	76,013	100.0	1,266.7
1	Malignant neoplasms (C00-C97)	26,831	35.3	447.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	19,898	26.2	331.6
3	Chronic lower respiratory diseases (J40-J47)	3,795	5.0	63.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,912	3.8	48.5
5	Diabetes mellitus (E10-E14)	2,821	3.7	47.0
6	Cerebrovascular diseases (I60-I69)	2,391	3.1	39.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,177	2.9	36.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,529	2.0	25.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	915	1.2	15.2
10	Septicemia (A40-A41)	868	1.1	14.5
11	Influenza and pneumonia (J09-J18)	833	1.1	13.9
12	Aortic aneurysm and dissection (I71)	557	0.7	9.3
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	453	0.6	7.5
14	Viral hepatitis (B15-B19)	416	0.5	6.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	387	0.5	6.4
...	All other causes (Residual)	9,230	12.1	153.8

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 65-69 years				
...	All causes	84,199	100.0	1,937.5
1	Malignant neoplasms (C00-C97)	30,857	36.6	710.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	21,175	25.1	487.3
3	Chronic lower respiratory diseases (J40-J47)	5,624	6.7	129.4
4	Diabetes mellitus (E10-E14)	3,131	3.7	72.0
5	Cerebrovascular diseases (I60-I69)	3,009	3.6	69.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,318	2.8	53.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,606	1.9	37.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,211	1.4	27.9
9	Septicemia (A40-A41)	1,089	1.3	25.1
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,065	1.3	24.5
11	Influenza and pneumonia (J09-J18)	988	1.2	22.7
12	Aortic aneurysm and dissection (I71)	652	0.8	15.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	502	0.6	11.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	501	0.6	11.5
15	Alzheimer's disease (G30)	440	0.5	10.1
...	All other causes (Residual)	10,031	11.9	230.8
White, male, 70-74 years				
...	All causes	100,629	100.0	2,989.5
1	Malignant neoplasms (C00-C97)	34,341	34.1	1,020.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	25,325	25.2	752.4
3	Chronic lower respiratory diseases (J40-J47)	7,775	7.7	231.0
4	Cerebrovascular diseases (I60-I69)	4,263	4.2	126.6
5	Diabetes mellitus (E10-E14)	3,522	3.5	104.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,480	2.5	73.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,814	1.8	53.9
8	Influenza and pneumonia (J09-J18)	1,487	1.5	44.2
9	Septicemia (A40-A41)	1,405	1.4	41.7
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,268	1.3	37.7
11	Alzheimer's disease (G30)	1,082	1.1	32.1
12	Parkinson's disease (G20-G21)	947	0.9	28.1
13	Aortic aneurysm and dissection (I71)	862	0.9	25.6
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	836	0.8	24.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	676	0.7	20.1
...	All other causes (Residual)	12,546	12.5	372.7

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 75-79 years				
...	All causes	132,199	100.0	4,786.6
1	Malignant neoplasms (C00-C97)	39,433	29.8	1,427.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	34,615	26.2	1,253.3
3	Chronic lower respiratory diseases (J40-J47)	10,419	7.9	377.2
4	Cerebrovascular diseases (I60-I69)	6,685	5.1	242.0
5	Diabetes mellitus (E10-E14)	4,093	3.1	148.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,969	2.2	107.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,734	2.1	99.0
8	Alzheimer's disease (G30)	2,683	2.0	97.1
9	Influenza and pneumonia (J09-J18)	2,545	1.9	92.1
10	Parkinson's disease (G20-G21)	2,200	1.7	79.7
11	Septicemia (A40-A41)	1,833	1.4	66.4
12	Chronic liver disease and cirrhosis (K70,K73-K74)	1,091	0.8	39.5
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,067	0.8	38.6
14	Aortic aneurysm and dissection (I71)	1,022	0.8	37.0
15	Pneumonitis due to solids and liquids (J69)	994	0.8	36.0
...	All other causes (Residual)	17,816	13.5	645.1
White, male, 80-84 years				
...	All causes	154,625	100.0	7,810.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	44,061	28.5	2,225.7
2	Malignant neoplasms (C00-C97)	37,258	24.1	1,882.0
3	Chronic lower respiratory diseases (J40-J47)	11,316	7.3	571.6
4	Cerebrovascular diseases (I60-I69)	8,637	5.6	436.3
5	Alzheimer's disease (G30)	4,925	3.2	248.8
6	Diabetes mellitus (E10-E14)	4,332	2.8	218.8
7	Influenza and pneumonia (J09-J18)	3,972	2.6	200.6
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,694	2.4	186.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,666	2.4	185.2
10	Parkinson's disease (G20-G21)	3,199	2.1	161.6
11	Septicemia (A40-A41)	2,235	1.4	112.9
12	Pneumonitis due to solids and liquids (J69)	1,630	1.1	82.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,303	0.8	65.8
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,209	0.8	61.1
15	Aortic aneurysm and dissection (I71)	1,136	0.7	57.4
...	All other causes (Residual)	22,052	14.3	1,113.9

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 85-89 years				
...	All causes	134,013	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	42,088	31.4	---
2	Malignant neoplasms (C00-C97)	24,389	18.2	---
3	Cerebrovascular diseases (I60-I69)	8,563	6.4	---
4	Chronic lower respiratory diseases (J40-J47)	8,195	6.1	---
5	Alzheimer's disease (G30)	5,922	4.4	---
6	Influenza and pneumonia (J09-J18)	4,398	3.3	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,442	2.6	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,434	2.6	---
9	Diabetes mellitus (E10-E14)	3,182	2.4	---
10	Parkinson's disease (G20-G21)	2,591	1.9	---
11	Pneumonitis due to solids and liquids (J69)	1,966	1.5	---
12	Septicemia (A40-A41)	1,801	1.3	---
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,264	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	987	0.7	---
15	Aortic aneurysm and dissection (I71)	876	0.7	---
...	All other causes (Residual)	20,915	15.6	---
White, male, 90-94 years				
...	All causes	71,426	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	25,185	35.3	---
2	Malignant neoplasms (C00-C97)	9,457	13.2	---
3	Cerebrovascular diseases (I60-I69)	4,515	6.3	---
4	Alzheimer's disease (G30)	3,662	5.1	---
5	Chronic lower respiratory diseases (J40-J47)	3,474	4.9	---
6	Influenza and pneumonia (J09-J18)	3,046	4.3	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,909	2.7	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,784	2.5	---
9	Diabetes mellitus (E10-E14)	1,317	1.8	---
10	Pneumonitis due to solids and liquids (J69)	1,253	1.8	---
11	Parkinson's disease (G20-G21)	1,062	1.5	---
12	Septicemia (A40-A41)	880	1.2	---
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	830	1.2	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	491	0.7	---
15	Atherosclerosis (I70)	455	0.6	---
...	All other causes (Residual)	12,106	16.9	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 95-99 years				
...	All causes	20,093	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,617	37.9	---
2	Malignant neoplasms (C00-C97)	1,954	9.7	---
3	Cerebrovascular diseases (I60-I69)	1,251	6.2	---
4	Influenza and pneumonia (J09-J18)	1,090	5.4	---
5	Alzheimer's disease (G30)	1,032	5.1	---
6	Chronic lower respiratory diseases (J40-J47)	846	4.2	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	558	2.8	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	538	2.7	---
9	Pneumonitis due to solids and liquids (J69)	388	1.9	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	271	1.3	---
11	Diabetes mellitus (E10-E14)	266	1.3	---
12	Septicemia (A40-A41)	206	1.0	---
13	Parkinson's disease (G20-G21)	183	0.9	---
14	Atherosclerosis (I70)	154	0.8	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	100	0.5	---
...	All other causes (Residual)	3,639	18.1	---
White, male, 100 years and over				
...	All causes	2,695	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,089	40.4	---
2	Influenza and pneumonia (J09-J18)	218	8.1	---
3	Malignant neoplasms (C00-C97)	195	7.2	---
4	Cerebrovascular diseases (I60-I69)	128	4.7	---
5	Alzheimer's disease (G30)	94	3.5	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	78	2.9	---
7	Chronic lower respiratory diseases (J40-J47)	74	2.7	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	72	2.7	---
9	Pneumonitis due to solids and liquids (J69)	57	2.1	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	37	1.4	---
11	Atherosclerosis (I70)	35	1.3	---
12	Diabetes mellitus (E10-E14)	34	1.3	---
13	Septicemia (A40-A41)	19	0.7	---
14	Parkinson's disease (G20-G21)	12	0.4	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	0.4	---
15	Anemias (D50-D64)	10	0.4	---
...	All other causes (Residual)	533	19.8	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, female, all ages				
...	All causes	1,050,200	100.0	854.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	264,728	25.2	215.5
2	Malignant neoplasms (C00-C97)	231,890	22.1	188.8
3	Cerebrovascular diseases (I60-I69)	69,981	6.7	57.0
4	Chronic lower respiratory diseases (J40-J47)	62,147	5.9	50.6
5	Alzheimer's disease (G30)	48,748	4.6	39.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	38,193	3.6	31.1
7	Diabetes mellitus (E10-E14)	27,646	2.6	22.5
8	Influenza and pneumonia (J09-J18)	25,227	2.4	20.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18,629	1.8	15.2
10	Septicemia (A40-A41)	15,150	1.4	12.3
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	11,477	1.1	9.3
12	Chronic liver disease and cirrhosis (K70,K73-K74)	8,689	0.8	7.1
13	Parkinson's disease (G20-G21)	7,969	0.8	6.5
14	Pneumonitis due to solids and liquids (J69)	7,219	0.7	5.9
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,623	0.6	5.4
...	All other causes (Residual)	205,884	19.6	167.6
White, female, 1-4 years				
...	All causes	1,440	100.0	23.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	431	29.9	6.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	185	12.8	3.0
3	Malignant neoplasms (C00-C97)	141	9.8	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	100	6.9	1.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	65	4.5	1.0
6	Influenza and pneumonia (J09-J18)	41	2.8	0.7
7	Septicemia (A40-A41)	27	1.9	0.4
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	24	1.7	0.4
9	Certain conditions originating in the perinatal period (P00-P96)	21	1.5	0.3
10	Cerebrovascular diseases (I60-I69)	18	1.3	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	11	0.8	*
12	Chronic lower respiratory diseases (J40-J47)	9	0.6	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.6	*
14	Meningitis (G00,G03)	5	0.3	*
14	Acute bronchitis and bronchiolitis (J20-J21)	5	0.3	*
...	All other causes (Residual)	349	24.2	5.6

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 5-9 years				
...	All causes	889	100.0	11.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	288	32.4	3.8
2	Malignant neoplasms (C00-C97)	185	20.8	2.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	71	8.0	0.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	50	5.6	0.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	37	4.2	0.5
6	Influenza and pneumonia (J09-J18)	18	2.0	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	1.9	*
8	Cerebrovascular diseases (I60-I69)	14	1.6	*
9	Chronic lower respiratory diseases (J40-J47)	10	1.1	*
10	Septicemia (A40-A41)	7	0.8	*
11	Meningitis (G00,G03)	6	0.7	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.6	*
13	Anemias (D50-D64)	2	0.2	*
13	Diabetes mellitus (E10-E14)	2	0.2	*
13	Pneumonitis due to solids and liquids (J69)	2	0.2	*
13	Diseases of appendix (K35-K38)	2	0.2	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.2	*
...	All other causes (Residual)	171	19.2	2.3
White, female, 10-14 years				
...	All causes	1,001	100.0	13.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	337	33.7	4.4
2	Malignant neoplasms (C00-C97)	187	18.7	2.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	58	5.8	0.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	40	4.0	0.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	39	3.9	0.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	33	3.3	0.4
7	Influenza and pneumonia (J09-J18)	23	2.3	0.3
8	Cerebrovascular diseases (I60-I69)	17	1.7	*
9	Septicemia (A40-A41)	14	1.4	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	1.2	*
11	Chronic lower respiratory diseases (J40-J47)	10	1.0	*
12	Meningitis (G00,G03)	6	0.6	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.5	*
14	Diabetes mellitus (E10-E14)	4	0.4	*
14	Certain conditions originating in the perinatal period (P00-P96)	4	0.4	*
...	All other causes (Residual)	212	21.2	2.8

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 15-19 years				
...	All causes	2,914	100.0	36.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,675	57.5	20.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	217	7.4	2.7
3	Malignant neoplasms (C00-C97)	211	7.2	2.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	162	5.6	2.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	81	2.8	1.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	56	1.9	0.7
7	Cerebrovascular diseases (I60-I69)	24	0.8	0.3
8	Septicemia (A40-A41)	21	0.7	0.3
8	Influenza and pneumonia (J09-J18)	21	0.7	0.3
10	Pregnancy, childbirth and the puerperium (O00-O99)	19	0.7	*
11	Chronic lower respiratory diseases (J40-J47)	16	0.5	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	0.4	*
12	Diabetes mellitus (E10-E14)	12	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.3	*
15	Anemias (D50-D64)	6	0.2	*
...	All other causes (Residual)	373	12.8	4.6
White, female, 20-24 years				
...	All causes	3,710	100.0	46.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,685	45.4	21.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	329	8.9	4.1
3	Malignant neoplasms (C00-C97)	304	8.2	3.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	239	6.4	3.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	180	4.9	2.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	68	1.8	0.9
7	Pregnancy, childbirth and the puerperium (O00-O99)	67	1.8	0.8
8	Cerebrovascular diseases (I60-I69)	38	1.0	0.5
9	Influenza and pneumonia (J09-J18)	34	0.9	0.4
10	Septicemia (A40-A41)	26	0.7	0.3
11	Diabetes mellitus (E10-E14)	24	0.6	0.3
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22	0.6	0.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	0.5	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	14	0.4	*
14	Chronic lower respiratory diseases (J40-J47)	14	0.4	*
...	All other causes (Residual)	649	17.5	8.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 25-29 years				
...	All causes	4,223	100.0	52.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,599	37.9	20.0
2	Malignant neoplasms (C00-C97)	453	10.7	5.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	406	9.6	5.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	262	6.2	3.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	228	5.4	2.9
6	Pregnancy, childbirth and the puerperium (O00-O99)	91	2.2	1.1
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	69	1.6	0.9
8	Cerebrovascular diseases (I60-I69)	60	1.4	0.8
9	Diabetes mellitus (E10-E14)	49	1.2	0.6
10	Influenza and pneumonia (J09-J18)	46	1.1	0.6
11	Septicemia (A40-A41)	35	0.8	0.4
11	Chronic liver disease and cirrhosis (K70,K73-K74)	35	0.8	0.4
13	Chronic lower respiratory diseases (J40-J47)	34	0.8	0.4
14	Human immunodeficiency virus (HIV) disease (B20-B24)	33	0.8	0.4
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	24	0.6	0.3
...	All other causes (Residual)	799	18.9	10.0
White, female, 30-34 years				
...	All causes	4,960	100.0	66.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,423	28.7	19.2
2	Malignant neoplasms (C00-C97)	878	17.7	11.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	442	8.9	6.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	378	7.6	5.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	197	4.0	2.7
6	Cerebrovascular diseases (I60-I69)	105	2.1	1.4
7	Pregnancy, childbirth and the puerperium (O00-O99)	102	2.1	1.4
8	Diabetes mellitus (E10-E14)	96	1.9	1.3
9	Human immunodeficiency virus (HIV) disease (B20-B24)	71	1.4	1.0
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	68	1.4	0.9
11	Septicemia (A40-A41)	56	1.1	0.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	56	1.1	0.8
13	Influenza and pneumonia (J09-J18)	52	1.0	0.7
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	41	0.8	0.6
15	Chronic lower respiratory diseases (J40-J47)	39	0.8	0.5
...	All other causes (Residual)	956	19.3	12.9

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 35-39 years				
...	All causes	7,984	100.0	97.0
1	Malignant neoplasms (C00-C97)	1,889	23.7	22.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,801	22.6	21.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	773	9.7	9.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	582	7.3	7.1
5	Cerebrovascular diseases (I60-I69)	219	2.7	2.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	212	2.7	2.6
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	211	2.6	2.6
8	Diabetes mellitus (E10-E14)	171	2.1	2.1
9	Human immunodeficiency virus (HIV) disease (B20-B24)	105	1.3	1.3
10	Septicemia (A40-A41)	100	1.3	1.2
11	Pregnancy, childbirth and the puerperium (O00-O99)	98	1.2	1.2
12	Influenza and pneumonia (J09-J18)	93	1.2	1.1
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	89	1.1	1.1
14	Chronic lower respiratory diseases (J40-J47)	85	1.1	1.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	54	0.7	0.7
...	All other causes (Residual)	1,502	18.8	18.2
White, female, 40-44 years				
...	All causes	13,466	100.0	153.6
1	Malignant neoplasms (C00-C97)	3,878	28.8	44.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,336	17.3	26.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,575	11.7	18.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	823	6.1	9.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	495	3.7	5.6
6	Cerebrovascular diseases (I60-I69)	391	2.9	4.5
7	Diabetes mellitus (E10-E14)	312	2.3	3.6
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	249	1.8	2.8
9	Chronic lower respiratory diseases (J40-J47)	215	1.6	2.5
10	Septicemia (A40-A41)	189	1.4	2.2
11	Human immunodeficiency virus (HIV) disease (B20-B24)	168	1.2	1.9
12	Influenza and pneumonia (J09-J18)	164	1.2	1.9
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	118	0.9	1.3
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	112	0.8	1.3
15	Viral hepatitis (B15-B19)	101	0.8	1.2
...	All other causes (Residual)	2,340	17.4	26.7

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 45-49 years				
...	All causes	22,188	100.0	237.1
1	Malignant neoplasms (C00-C97)	7,672	34.6	82.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,907	13.1	31.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,822	12.7	30.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	936	4.2	10.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	905	4.1	9.7
6	Cerebrovascular diseases (I60-I69)	757	3.4	8.1
7	Diabetes mellitus (E10-E14)	536	2.4	5.7
8	Chronic lower respiratory diseases (J40-J47)	534	2.4	5.7
9	Septicemia (A40-A41)	297	1.3	3.2
10	Viral hepatitis (B15-B19)	256	1.2	2.7
11	Influenza and pneumonia (J09-J18)	246	1.1	2.6
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	220	1.0	2.4
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	215	1.0	2.3
14	Human immunodeficiency virus (HIV) disease (B20-B24)	191	0.9	2.0
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	114	0.5	1.2
...	All other causes (Residual)	3,580	16.1	38.3
White, female, 50-54 years				
...	All causes	30,489	100.0	347.5
1	Malignant neoplasms (C00-C97)	11,752	38.5	133.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,352	14.3	49.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,471	8.1	28.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	1,095	3.6	12.5
5	Chronic lower respiratory diseases (J40-J47)	1,084	3.6	12.4
6	Cerebrovascular diseases (I60-I69)	1,052	3.5	12.0
7	Diabetes mellitus (E10-E14)	996	3.3	11.4
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	887	2.9	10.1
9	Septicemia (A40-A41)	453	1.5	5.2
10	Viral hepatitis (B15-B19)	403	1.3	4.6
11	Influenza and pneumonia (J09-J18)	347	1.1	4.0
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	343	1.1	3.9
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	163	0.5	1.9
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	157	0.5	1.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	152	0.5	1.7
...	All other causes (Residual)	4,782	15.7	54.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 55-59 years				
...	All causes	39,512	100.0	508.0
1	Malignant neoplasms (C00-C97)	16,793	42.5	215.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,177	15.6	79.4
3	Chronic lower respiratory diseases (J40-J47)	1,876	4.7	24.1
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,807	4.6	23.2
5	Diabetes mellitus (E10-E14)	1,403	3.6	18.0
6	Cerebrovascular diseases (I60-I69)	1,341	3.4	17.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	995	2.5	12.8
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	713	1.8	9.2
9	Septicemia (A40-A41)	652	1.7	8.4
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	531	1.3	6.8
11	Influenza and pneumonia (J09-J18)	418	1.1	5.4
12	Viral hepatitis (B15-B19)	313	0.8	4.0
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	215	0.5	2.8
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	186	0.5	2.4
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	146	0.4	1.9
...	All other causes (Residual)	5,946	15.0	76.4
White, female, 60-64 years				
...	All causes	50,990	100.0	795.8
1	Malignant neoplasms (C00-C97)	21,763	42.7	339.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,762	17.2	136.8
3	Chronic lower respiratory diseases (J40-J47)	3,580	7.0	55.9
4	Diabetes mellitus (E10-E14)	1,884	3.7	29.4
5	Cerebrovascular diseases (I60-I69)	1,855	3.6	29.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,433	2.8	22.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	921	1.8	14.4
8	Septicemia (A40-A41)	852	1.7	13.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	706	1.4	11.0
10	Influenza and pneumonia (J09-J18)	579	1.1	9.0
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	461	0.9	7.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	299	0.6	4.7
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	270	0.5	4.2
14	Alzheimer's disease (G30)	254	0.5	4.0
15	Viral hepatitis (B15-B19)	179	0.4	2.8
...	All other causes (Residual)	7,192	14.1	112.3

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 65-69 years				
...	All causes	61,464	100.0	1,262.8
1	Malignant neoplasms (C00-C97)	25,076	40.8	515.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,976	17.9	225.5
3	Chronic lower respiratory diseases (J40-J47)	5,320	8.7	109.3
4	Cerebrovascular diseases (I60-I69)	2,651	4.3	54.5
5	Diabetes mellitus (E10-E14)	2,133	3.5	43.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,240	2.0	25.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,092	1.8	22.4
8	Septicemia (A40-A41)	1,024	1.7	21.0
9	Chronic liver disease and cirrhosis (K70,K73-K74)	842	1.4	17.3
10	Influenza and pneumonia (J09-J18)	799	1.3	16.4
11	Alzheimer's disease (G30)	536	0.9	11.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	425	0.7	8.7
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	366	0.6	7.5
14	Aortic aneurysm and dissection (I71)	273	0.4	5.6
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	247	0.4	5.1
...	All other causes (Residual)	8,464	13.8	173.9
White, female, 70-74 years				
...	All causes	81,085	100.0	2,008.7
1	Malignant neoplasms (C00-C97)	28,552	35.2	707.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	16,124	19.9	399.4
3	Chronic lower respiratory diseases (J40-J47)	7,670	9.5	190.0
4	Cerebrovascular diseases (I60-I69)	4,100	5.1	101.6
5	Diabetes mellitus (E10-E14)	2,866	3.5	71.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,538	1.9	38.1
7	Alzheimer's disease (G30)	1,505	1.9	37.3
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,491	1.8	36.9
9	Septicemia (A40-A41)	1,334	1.6	33.0
10	Influenza and pneumonia (J09-J18)	1,277	1.6	31.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	908	1.1	22.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	634	0.8	15.7
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	542	0.7	13.4
14	Parkinson's disease (G20-G21)	531	0.7	13.2
15	Aortic aneurysm and dissection (I71)	508	0.6	12.6
...	All other causes (Residual)	11,505	14.2	285.0

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 75-79 years				
...	All causes	120,965	100.0	3,291.8
1	Malignant neoplasms (C00-C97)	33,834	28.0	920.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	27,433	22.7	746.5
3	Chronic lower respiratory diseases (J40-J47)	10,796	8.9	293.8
4	Cerebrovascular diseases (I60-I69)	7,669	6.3	208.7
5	Alzheimer's disease (G30)	4,178	3.5	113.7
6	Diabetes mellitus (E10-E14)	4,051	3.3	110.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,439	2.0	66.4
8	Influenza and pneumonia (J09-J18)	2,417	2.0	65.8
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,355	1.9	64.1
10	Septicemia (A40-A41)	1,957	1.6	53.3
11	Parkinson's disease (G20-G21)	1,284	1.1	34.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,036	0.9	28.2
13	Chronic liver disease and cirrhosis (K70,K73-K74)	888	0.7	24.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	804	0.7	21.9
15	Aortic aneurysm and dissection (I71)	785	0.6	21.4
...	All other causes (Residual)	19,039	15.7	518.1
White, female, 80-84 years				
...	All causes	171,333	100.0	5,535.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	44,932	26.2	1,451.7
2	Malignant neoplasms (C00-C97)	35,060	20.5	1,132.7
3	Cerebrovascular diseases (I60-I69)	13,349	7.8	431.3
4	Chronic lower respiratory diseases (J40-J47)	12,455	7.3	402.4
5	Alzheimer's disease (G30)	9,329	5.4	301.4
6	Diabetes mellitus (E10-E14)	4,935	2.9	159.4
7	Influenza and pneumonia (J09-J18)	4,140	2.4	133.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,566	2.1	115.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,443	2.0	111.2
10	Septicemia (A40-A41)	2,659	1.6	85.9
11	Parkinson's disease (G20-G21)	2,149	1.3	69.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,940	1.1	62.7
13	Pneumonitis due to solids and liquids (J69)	1,359	0.8	43.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,127	0.7	36.4
15	Aortic aneurysm and dissection (I71)	1,016	0.6	32.8
...	All other causes (Residual)	29,874	17.4	965.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 85-89 years				
...	All causes	193,579	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	58,536	30.2	---
2	Malignant neoplasms (C00-C97)	26,212	13.5	---
3	Cerebrovascular diseases (I60-I69)	16,846	8.7	---
4	Alzheimer's disease (G30)	13,787	7.1	---
5	Chronic lower respiratory diseases (J40-J47)	10,448	5.4	---
6	Influenza and pneumonia (J09-J18)	5,761	3.0	---
7	Diabetes mellitus (E10-E14)	4,566	2.4	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,175	2.2	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,865	2.0	---
10	Septicemia (A40-A41)	2,814	1.5	---
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,731	1.4	---
12	Parkinson's disease (G20-G21)	2,115	1.1	---
13	Pneumonitis due to solids and liquids (J69)	1,768	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,223	0.6	---
15	Atherosclerosis (I70)	1,100	0.6	---
...	All other causes (Residual)	37,632	19.4	---
White, female, 90-94 years				
...	All causes	149,085	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	50,902	34.1	---
2	Cerebrovascular diseases (I60-I69)	13,016	8.7	---
3	Malignant neoplasms (C00-C97)	12,751	8.6	---
4	Alzheimer's disease (G30)	12,222	8.2	---
5	Chronic lower respiratory diseases (J40-J47)	5,782	3.9	---
6	Influenza and pneumonia (J09-J18)	5,272	3.5	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,196	2.1	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,795	1.9	---
9	Diabetes mellitus (E10-E14)	2,651	1.8	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,403	1.6	---
11	Septicemia (A40-A41)	1,756	1.2	---
12	Pneumonitis due to solids and liquids (J69)	1,510	1.0	---
13	Parkinson's disease (G20-G21)	1,200	0.8	---
14	Atherosclerosis (I70)	1,037	0.7	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	795	0.5	---
...	All other causes (Residual)	31,797	21.3	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 95-99 years				
...	All causes	66,057	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	24,442	37.0	---
2	Alzheimer's disease (G30)	5,671	8.6	---
3	Cerebrovascular diseases (I60-I69)	5,437	8.2	---
4	Malignant neoplasms (C00-C97)	3,758	5.7	---
5	Influenza and pneumonia (J09-J18)	2,665	4.0	---
6	Chronic lower respiratory diseases (J40-J47)	1,888	2.9	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,440	2.2	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,180	1.8	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,093	1.7	---
10	Diabetes mellitus (E10-E14)	806	1.2	---
11	Pneumonitis due to solids and liquids (J69)	691	1.0	---
12	Septicemia (A40-A41)	676	1.0	---
13	Atherosclerosis (I70)	631	1.0	---
14	Parkinson's disease (G20-G21)	297	0.4	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	288	0.4	---
...	All other causes (Residual)	15,094	22.8	---
White, female, 100 years and over				
...	All causes	14,573	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,680	39.0	---
2	Alzheimer's disease (G30)	1,094	7.5	---
3	Cerebrovascular diseases (I60-I69)	978	6.7	---
4	Influenza and pneumonia (J09-J18)	767	5.3	---
5	Malignant neoplasms (C00-C97)	510	3.5	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	332	2.3	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	274	1.9	---
8	Chronic lower respiratory diseases (J40-J47)	269	1.8	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	246	1.7	---
10	Atherosclerosis (I70)	219	1.5	---
11	Pneumonitis due to solids and liquids (J69)	159	1.1	---
12	Diabetes mellitus (E10-E14)	146	1.0	---
13	Septicemia (A40-A41)	128	0.9	---
14	Anemias (D50-D64)	49	0.3	---
15	Nutritional deficiencies (E40-E64)	39	0.3	---
...	All other causes (Residual)	3,683	25.3	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, all ages				
...	All causes	289,585	100.0	723.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	71,209	24.6	177.9
2	Malignant neoplasms (C00-C97)	64,049	22.1	160.0
3	Cerebrovascular diseases (I60-I69)	17,085	5.9	42.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,559	4.7	33.9
5	Diabetes mellitus (E10-E14)	12,459	4.3	31.1
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8,870	3.1	22.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,392	2.9	21.0
8	Chronic lower respiratory diseases (J40-J47)	7,901	2.7	19.7
9	Human immunodeficiency virus (HIV) disease (B20-B24)	6,470	2.2	16.2
10	Septicemia (A40-A41)	6,297	2.2	15.7
11	Influenza and pneumonia (J09-J18)	5,155	1.8	12.9
12	Certain conditions originating in the perinatal period (P00-P96)	5,052	1.7	12.6
13	Alzheimer's disease (G30)	4,760	1.6	11.9
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,640	1.6	11.6
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,558	0.9	6.4
...	All other causes (Residual)	51,129	17.7	127.7
Black, both sexes, 1-4 years				
...	All causes	1,137	100.0	42.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	325	28.6	12.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	168	14.8	6.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	119	10.5	4.4
4	Malignant neoplasms (C00-C97)	55	4.8	2.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	43	3.8	1.6
6	Chronic lower respiratory diseases (J40-J47)	30	2.6	1.1
7	Influenza and pneumonia (J09-J18)	29	2.6	1.1
8	Certain conditions originating in the perinatal period (P00-P96)	27	2.4	1.0
9	Septicemia (A40-A41)	16	1.4	*
9	Cerebrovascular diseases (I60-I69)	16	1.4	*
11	Anemias (D50-D64)	12	1.1	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.8	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.5	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.5	*
15	Acute bronchitis and bronchiolitis (J20-J21)	5	0.4	*
...	All other causes (Residual)	271	23.8	10.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 5-9 years				
...	All causes	577	100.0	18.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	220	38.1	6.9
2	Malignant neoplasms (C00-C97)	64	11.1	2.0
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	41	7.1	1.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	37	6.4	1.2
5	Chronic lower respiratory diseases (J40-J47)	25	4.3	0.8
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	23	4.0	0.7
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	1.6	*
8	Influenza and pneumonia (J09-J18)	7	1.2	*
9	Septicemia (A40-A41)	6	1.0	*
9	Cerebrovascular diseases (I60-I69)	6	1.0	*
11	Anemias (D50-D64)	5	0.9	*
12	Certain conditions originating in the perinatal period (P00-P96)	4	0.7	*
13	Meningococcal infection (A39)	3	0.5	*
13	Diseases of appendix (K35-K38)	3	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.5	*
...	All other causes (Residual)	121	21.0	3.8
Black, both sexes, 10-14 years				
...	All causes	786	100.0	23.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	239	30.4	7.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	103	13.1	3.1
3	Malignant neoplasms (C00-C97)	79	10.1	2.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	40	5.1	1.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	40	5.1	1.2
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	36	4.6	1.1
7	Chronic lower respiratory diseases (J40-J47)	35	4.5	1.1
8	Influenza and pneumonia (J09-J18)	14	1.8	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	1.5	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	8	1.0	*
10	Anemias (D50-D64)	8	1.0	*
12	Cerebrovascular diseases (I60-I69)	7	0.9	*
13	Septicemia (A40-A41)	5	0.6	*
14	Diabetes mellitus (E10-E14)	4	0.5	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.5	*
...	All other causes (Residual)	152	19.3	4.6

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 15-19 years				
...	All causes	2,956	100.0	83.4
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,325	44.8	37.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	746	25.2	21.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	144	4.9	4.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	114	3.9	3.2
5	Malignant neoplasms (C00-C97)	106	3.6	3.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	46	1.6	1.3
7	Chronic lower respiratory diseases (J40-J47)	28	0.9	0.8
8	Anemias (D50-D64)	26	0.9	0.7
9	Human immunodeficiency virus (HIV) disease (B20-B24)	25	0.8	0.7
10	Diabetes mellitus (E10-E14)	21	0.7	0.6
11	Pregnancy, childbirth and the puerperium (O00-O99)	20	0.7	0.6
12	Cerebrovascular diseases (I60-I69)	16	0.5	*
13	Septicemia (A40-A41)	12	0.4	*
14	Influenza and pneumonia (J09-J18)	8	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.3	*
14	Legal intervention (Y35,Y89.0)	8	0.3	*
...	All other causes (Residual)	303	10.3	8.5
Black, both sexes, 20-24 years				
...	All causes	4,424	100.0	138.0
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,888	42.7	58.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,003	22.7	31.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	262	5.9	8.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	234	5.3	7.3
5	Malignant neoplasms (C00-C97)	155	3.5	4.8
6	Human immunodeficiency virus (HIV) disease (B20-B24)	84	1.9	2.6
7	Pregnancy, childbirth and the puerperium (O00-O99)	49	1.1	1.5
8	Anemias (D50-D64)	41	0.9	1.3
9	Legal intervention (Y35,Y89.0)	40	0.9	1.2
10	Chronic lower respiratory diseases (J40-J47)	37	0.8	1.2
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	37	0.8	1.2
12	Septicemia (A40-A41)	35	0.8	1.1
12	Diabetes mellitus (E10-E14)	35	0.8	1.1
14	Cerebrovascular diseases (I60-I69)	30	0.7	0.9
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20	0.5	0.6
...	All other causes (Residual)	474	10.7	14.8

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 25-29 years				
...	All causes	4,741	100.0	155.4
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,515	32.0	49.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	929	19.6	30.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	401	8.5	13.1
4	Human immunodeficiency virus (HIV) disease (B20-B24)	284	6.0	9.3
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	250	5.3	8.2
6	Malignant neoplasms (C00-C97)	234	4.9	7.7
7	Diabetes mellitus (E10-E14)	83	1.8	2.7
8	Pregnancy, childbirth and the puerperium (O00-O99)	56	1.2	1.8
9	Anemias (D50-D64)	52	1.1	1.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	48	1.0	1.6
11	Influenza and pneumonia (J09-J18)	46	1.0	1.5
12	Septicemia (A40-A41)	45	0.9	1.5
12	Chronic lower respiratory diseases (J40-J47)	45	0.9	1.5
14	Cerebrovascular diseases (I60-I69)	43	0.9	1.4
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	41	0.9	1.3
...	All other causes (Residual)	669	14.1	21.9
Black, both sexes, 30-34 years				
...	All causes	4,921	100.0	184.4
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,045	21.2	39.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	850	17.3	31.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	596	12.1	22.3
4	Human immunodeficiency virus (HIV) disease (B20-B24)	421	8.6	15.8
5	Malignant neoplasms (C00-C97)	384	7.8	14.4
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	231	4.7	8.7
7	Diabetes mellitus (E10-E14)	130	2.6	4.9
8	Cerebrovascular diseases (I60-I69)	105	2.1	3.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	65	1.3	2.4
10	Pregnancy, childbirth and the puerperium (O00-O99)	52	1.1	1.9
11	Chronic lower respiratory diseases (J40-J47)	51	1.0	1.9
12	Anemias (D50-D64)	50	1.0	1.9
13	Influenza and pneumonia (J09-J18)	48	1.0	1.8
14	Septicemia (A40-A41)	45	0.9	1.7
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	42	0.9	1.6
...	All other causes (Residual)	806	16.4	30.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 35-39 years				
...	All causes	6,831	100.0	243.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,081	15.8	38.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,021	14.9	36.5
3	Human immunodeficiency virus (HIV) disease (B20-B24)	812	11.9	29.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	777	11.4	27.7
5	Malignant neoplasms (C00-C97)	718	10.5	25.6
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	225	3.3	8.0
7	Cerebrovascular diseases (I60-I69)	216	3.2	7.7
8	Diabetes mellitus (E10-E14)	214	3.1	7.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	135	2.0	4.8
10	Septicemia (A40-A41)	108	1.6	3.9
11	Influenza and pneumonia (J09-J18)	72	1.1	2.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	66	1.0	2.4
13	Chronic lower respiratory diseases (J40-J47)	65	1.0	2.3
14	Anemias (D50-D64)	64	0.9	2.3
15	Aortic aneurysm and dissection (I71)	46	0.7	1.6
...	All other causes (Residual)	1,211	17.7	43.2
Black, both sexes, 40-44 years				
...	All causes	10,159	100.0	353.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,911	18.8	66.5
2	Malignant neoplasms (C00-C97)	1,647	16.2	57.4
3	Human immunodeficiency virus (HIV) disease (B20-B24)	1,190	11.7	41.4
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,137	11.2	39.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	589	5.8	20.5
6	Cerebrovascular diseases (I60-I69)	439	4.3	15.3
7	Diabetes mellitus (E10-E14)	348	3.4	12.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	199	2.0	6.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	195	1.9	6.8
10	Septicemia (A40-A41)	166	1.6	5.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	162	1.6	5.6
12	Chronic lower respiratory diseases (J40-J47)	128	1.3	4.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	110	1.1	3.8
14	Influenza and pneumonia (J09-J18)	106	1.0	3.7
15	Anemias (D50-D64)	78	0.8	2.7
...	All other causes (Residual)	1,754	17.3	61.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 45-49 years				
...	All causes	15,609	100.0	549.5
1	Malignant neoplasms (C00-C97)	3,363	21.5	118.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,321	21.3	116.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,399	9.0	49.2
4	Human immunodeficiency virus (HIV) disease (B20-B24)	1,250	8.0	44.0
5	Cerebrovascular diseases (I60-I69)	795	5.1	28.0
6	Diabetes mellitus (E10-E14)	544	3.5	19.1
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	480	3.1	16.9
8	Chronic liver disease and cirrhosis (K70,K73-K74)	351	2.2	12.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	343	2.2	12.1
10	Septicemia (A40-A41)	302	1.9	10.6
11	Chronic lower respiratory diseases (J40-J47)	261	1.7	9.2
12	Influenza and pneumonia (J09-J18)	208	1.3	7.3
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	201	1.3	7.1
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	175	1.1	6.2
15	Viral hepatitis (B15-B19)	123	0.8	4.3
...	All other causes (Residual)	2,493	16.0	87.8
Black, both sexes, 50-54 years				
...	All causes	21,209	100.0	861.0
1	Malignant neoplasms (C00-C97)	5,455	25.7	221.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,036	23.7	204.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,367	6.4	55.5
4	Cerebrovascular diseases (I60-I69)	1,151	5.4	46.7
5	Human immunodeficiency virus (HIV) disease (B20-B24)	1,101	5.2	44.7
6	Diabetes mellitus (E10-E14)	889	4.2	36.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	502	2.4	20.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	480	2.3	19.5
9	Septicemia (A40-A41)	419	2.0	17.0
10	Chronic lower respiratory diseases (J40-J47)	411	1.9	16.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	334	1.6	13.6
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	322	1.5	13.1
13	Viral hepatitis (B15-B19)	259	1.2	10.5
14	Influenza and pneumonia (J09-J18)	231	1.1	9.4
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	146	0.7	5.9
...	All other causes (Residual)	3,106	14.6	126.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 55-59 years				
...	All causes	24,190	100.0	1,215.4
1	Malignant neoplasms (C00-C97)	7,213	29.8	362.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,098	25.2	306.4
3	Cerebrovascular diseases (I60-I69)	1,306	5.4	65.6
4	Diabetes mellitus (E10-E14)	1,203	5.0	60.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,007	4.2	50.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	640	2.6	32.2
7	Human immunodeficiency virus (HIV) disease (B20-B24)	632	2.6	31.8
8	Chronic liver disease and cirrhosis (K70,K73-K74)	545	2.3	27.4
9	Chronic lower respiratory diseases (J40-J47)	525	2.2	26.4
10	Septicemia (A40-A41)	507	2.1	25.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	364	1.5	18.3
12	Influenza and pneumonia (J09-J18)	289	1.2	14.5
13	Viral hepatitis (B15-B19)	284	1.2	14.3
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	210	0.9	10.6
15	Aortic aneurysm and dissection (I71)	126	0.5	6.3
...	All other causes (Residual)	3,241	13.4	162.8
Black, both sexes, 60-64 years				
...	All causes	23,492	100.0	1,702.1
1	Malignant neoplasms (C00-C97)	7,549	32.1	546.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,191	26.4	448.6
3	Cerebrovascular diseases (I60-I69)	1,285	5.5	93.1
4	Diabetes mellitus (E10-E14)	1,283	5.5	93.0
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	751	3.2	54.4
6	Chronic lower respiratory diseases (J40-J47)	653	2.8	47.3
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	592	2.5	42.9
8	Septicemia (A40-A41)	497	2.1	36.0
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	388	1.7	28.1
10	Human immunodeficiency virus (HIV) disease (B20-B24)	363	1.5	26.3
11	Chronic liver disease and cirrhosis (K70,K73-K74)	322	1.4	23.3
12	Influenza and pneumonia (J09-J18)	284	1.2	20.6
13	Viral hepatitis (B15-B19)	125	0.5	9.1
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	109	0.5	7.9
15	Aortic aneurysm and dissection (I71)	106	0.5	7.7
...	All other causes (Residual)	2,994	12.7	216.9

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 65-69 years				
...	All causes	24,555	100.0	2,383.5
1	Malignant neoplasms (C00-C97)	8,028	32.7	779.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,443	26.2	625.4
3	Cerebrovascular diseases (I60-I69)	1,502	6.1	145.8
4	Diabetes mellitus (E10-E14)	1,331	5.4	129.2
5	Chronic lower respiratory diseases (J40-J47)	871	3.5	84.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	837	3.4	81.2
7	Septicemia (A40-A41)	620	2.5	60.2
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	474	1.9	46.0
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	398	1.6	38.6
10	Influenza and pneumonia (J09-J18)	332	1.4	32.2
11	Chronic liver disease and cirrhosis (K70,K73-K74)	222	0.9	21.5
12	Human immunodeficiency virus (HIV) disease (B20-B24)	150	0.6	14.6
13	Aortic aneurysm and dissection (I71)	128	0.5	12.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	105	0.4	10.2
15	Pneumonitis due to solids and liquids (J69)	83	0.3	8.1
...	All other causes (Residual)	3,031	12.3	294.2
Black, both sexes, 70-74 years				
...	All causes	26,829	100.0	3,309.7
1	Malignant neoplasms (C00-C97)	8,027	29.9	990.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,199	26.8	888.1
3	Cerebrovascular diseases (I60-I69)	1,755	6.5	216.5
4	Diabetes mellitus (E10-E14)	1,530	5.7	188.7
5	Chronic lower respiratory diseases (J40-J47)	1,026	3.8	126.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,004	3.7	123.9
7	Septicemia (A40-A41)	700	2.6	86.4
8	Influenza and pneumonia (J09-J18)	468	1.7	57.7
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	437	1.6	53.9
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	432	1.6	53.3
11	Alzheimer's disease (G30)	284	1.1	35.0
12	Chronic liver disease and cirrhosis (K70,K73-K74)	182	0.7	22.5
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	146	0.5	18.0
14	Pneumonitis due to solids and liquids (J69)	122	0.5	15.1
15	Aortic aneurysm and dissection (I71)	110	0.4	13.6
...	All other causes (Residual)	3,407	12.7	420.3

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 75-79 years				
...	All causes	29,063	100.0	4,838.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,926	27.3	1,319.5
2	Malignant neoplasms (C00-C97)	7,742	26.6	1,288.9
3	Cerebrovascular diseases (I60-I69)	2,068	7.1	344.3
4	Diabetes mellitus (E10-E14)	1,539	5.3	256.2
5	Chronic lower respiratory diseases (J40-J47)	1,213	4.2	201.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,046	3.6	174.1
7	Septicemia (A40-A41)	741	2.5	123.4
8	Alzheimer's disease (G30)	620	2.1	103.2
9	Influenza and pneumonia (J09-J18)	576	2.0	95.9
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	524	1.8	87.2
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	408	1.4	67.9
12	Pneumonitis due to solids and liquids (J69)	179	0.6	29.8
13	Parkinson's disease (G20-G21)	151	0.5	25.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	146	0.5	24.3
15	Aortic aneurysm and dissection (I71)	139	0.5	23.1
...	All other causes (Residual)	4,045	13.9	673.4
Black, both sexes, 80-84 years				
...	All causes	29,923	100.0	6,966.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,633	28.9	2,009.8
2	Malignant neoplasms (C00-C97)	6,535	21.8	1,521.4
3	Cerebrovascular diseases (I60-I69)	2,317	7.7	539.4
4	Diabetes mellitus (E10-E14)	1,529	5.1	356.0
5	Chronic lower respiratory diseases (J40-J47)	1,127	3.8	262.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,013	3.4	235.8
7	Alzheimer's disease (G30)	1,007	3.4	234.4
8	Septicemia (A40-A41)	778	2.6	181.1
9	Influenza and pneumonia (J09-J18)	736	2.5	171.3
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	571	1.9	132.9
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	356	1.2	82.9
12	Pneumonitis due to solids and liquids (J69)	259	0.9	60.3
13	Parkinson's disease (G20-G21)	210	0.7	48.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	164	0.5	38.2
15	Aortic aneurysm and dissection (I71)	132	0.4	30.7
...	All other causes (Residual)	4,556	15.2	1,060.7

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 85-89 years				
...	All causes	24,885	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,597	30.5	---
2	Malignant neoplasms (C00-C97)	4,127	16.6	---
3	Cerebrovascular diseases (I60-I69)	2,063	8.3	---
4	Alzheimer's disease (G30)	1,145	4.6	---
5	Diabetes mellitus (E10-E14)	1,044	4.2	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	969	3.9	---
7	Chronic lower respiratory diseases (J40-J47)	775	3.1	---
8	Influenza and pneumonia (J09-J18)	754	3.0	---
9	Septicemia (A40-A41)	630	2.5	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	564	2.3	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	317	1.3	---
12	Pneumonitis due to solids and liquids (J69)	261	1.0	---
13	Parkinson's disease (G20-G21)	150	0.6	---
14	Atherosclerosis (I70)	112	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	107	0.4	---
...	All other causes (Residual)	4,270	17.2	---
Black, both sexes, 90-94 years				
...	All causes	15,419	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,989	32.4	---
2	Malignant neoplasms (C00-C97)	1,883	12.2	---
3	Cerebrovascular diseases (I60-I69)	1,302	8.4	---
4	Alzheimer's disease (G30)	977	6.3	---
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	508	3.3	---
6	Influenza and pneumonia (J09-J18)	500	3.2	---
7	Diabetes mellitus (E10-E14)	499	3.2	---
8	Chronic lower respiratory diseases (J40-J47)	399	2.6	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	377	2.4	---
10	Septicemia (A40-A41)	373	2.4	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	196	1.3	---
12	Pneumonitis due to solids and liquids (J69)	152	1.0	---
13	Atherosclerosis (I70)	92	0.6	---
14	Parkinson's disease (G20-G21)	81	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	66	0.4	---
...	All other causes (Residual)	3,025	19.6	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 95-99 years				
...	All causes	6,777	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,426	35.8	---
2	Malignant neoplasms (C00-C97)	560	8.3	---
3	Cerebrovascular diseases (I60-I69)	491	7.2	---
4	Alzheimer's disease (G30)	456	6.7	---
5	Influenza and pneumonia (J09-J18)	252	3.7	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	209	3.1	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	191	2.8	---
8	Diabetes mellitus (E10-E14)	183	2.7	---
9	Chronic lower respiratory diseases (J40-J47)	147	2.2	---
10	Septicemia (A40-A41)	140	2.1	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	82	1.2	---
12	Pneumonitis due to solids and liquids (J69)	66	1.0	---
13	Atherosclerosis (I70)	55	0.8	---
14	Parkinson's disease (G20-G21)	25	0.4	---
15	Nutritional deficiencies (E40-E64)	24	0.4	---
...	All other causes (Residual)	1,470	21.7	---
Black, both sexes, 100 years and over				
...	All causes	2,113	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	761	36.0	---
2	Alzheimer's disease (G30)	146	6.9	---
3	Cerebrovascular diseases (I60-I69)	136	6.4	---
4	Malignant neoplasms (C00-C97)	112	5.3	---
5	Influenza and pneumonia (J09-J18)	105	5.0	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	65	3.1	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	60	2.8	---
8	Diabetes mellitus (E10-E14)	45	2.1	---
9	Septicemia (A40-A41)	36	1.7	---
10	Chronic lower respiratory diseases (J40-J47)	34	1.6	---
11	Pneumonitis due to solids and liquids (J69)	23	1.1	---
12	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	22	1.0	---
13	Atherosclerosis (I70)	21	1.0	---
14	Nutritional deficiencies (E40-E64)	10	0.5	---
15	Parkinson's disease (G20-G21)	8	0.4	---
...	All other causes (Residual)	529	25.0	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, all ages				
...	All causes	148,309	100.0	775.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	35,669	24.1	186.5
2	Malignant neoplasms (C00-C97)	33,069	22.3	172.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,268	6.2	48.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7,584	5.1	39.7
5	Cerebrovascular diseases (I60-I69)	7,549	5.1	39.5
6	Diabetes mellitus (E10-E14)	5,493	3.7	28.7
7	Chronic lower respiratory diseases (J40-J47)	4,207	2.8	22.0
8	Human immunodeficiency virus (HIV) disease (B20-B24)	4,186	2.8	21.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,772	2.5	19.7
10	Certain conditions originating in the perinatal period (P00-P96)	2,846	1.9	14.9
11	Septicemia (A40-A41)	2,835	1.9	14.8
12	Influenza and pneumonia (J09-J18)	2,498	1.7	13.1
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,979	1.3	10.3
14	Chronic liver disease and cirrhosis (K70,K73-K74)	1,682	1.1	8.8
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,606	1.1	8.4
...	All other causes (Residual)	24,066	16.2	125.9
Black, male, 1-4 years				
...	All causes	621	100.0	45.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	189	30.4	13.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	89	14.3	6.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	58	9.3	4.2
4	Malignant neoplasms (C00-C97)	33	5.3	2.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	28	4.5	2.0
6	Influenza and pneumonia (J09-J18)	16	2.6	*
7	Chronic lower respiratory diseases (J40-J47)	13	2.1	*
8	Certain conditions originating in the perinatal period (P00-P96)	12	1.9	*
9	Septicemia (A40-A41)	11	1.8	*
10	Anemias (D50-D64)	4	0.6	*
10	Cerebrovascular diseases (I60-I69)	4	0.6	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.6	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.5	*
13	Meningitis (G00,G03)	3	0.5	*
13	Acute bronchitis and bronchiolitis (J20-J21)	3	0.5	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.5	*
...	All other causes (Residual)	148	23.8	10.8

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 5-9 years				
...	All causes	331	100.0	20.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	134	40.5	8.2
2	Malignant neoplasms (C00-C97)	31	9.4	1.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	21	6.3	1.3
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	20	6.0	1.2
5	Chronic lower respiratory diseases (J40-J47)	17	5.1	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	16	4.8	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.8	*
8	Influenza and pneumonia (J09-J18)	4	1.2	*
9	Meningococcal infection (A39)	2	0.6	*
9	Septicemia (A40-A41)	2	0.6	*
9	Anemias (D50-D64)	2	0.6	*
9	Meningitis (G00,G03)	2	0.6	*
9	Cerebrovascular diseases (I60-I69)	2	0.6	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.6	*
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2	0.6	*
...	All other causes (Residual)	68	20.5	4.2
Black, male, 10-14 years				
...	All causes	485	100.0	28.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	145	29.9	8.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	75	15.5	4.4
3	Malignant neoplasms (C00-C97)	41	8.5	2.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	28	5.8	1.7
5	Chronic lower respiratory diseases (J40-J47)	24	4.9	1.4
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	19	3.9	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	17	3.5	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	1.9	*
8	Influenza and pneumonia (J09-J18)	9	1.9	*
10	Anemias (D50-D64)	6	1.2	*
11	Septicemia (A40-A41)	3	0.6	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.4	*
12	Diabetes mellitus (E10-E14)	2	0.4	*
12	Meningitis (G00,G03)	2	0.4	*
12	Pneumonitis due to solids and liquids (J69)	2	0.4	*
12	Diseases of appendix (K35-K38)	2	0.4	*
12	Certain conditions originating in the perinatal period (P00-P96)	2	0.4	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
...	All other causes (Residual)	95	19.6	5.6

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 15-19 years				
...	All causes	2,302	100.0	128.1
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,205	52.3	67.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	550	23.9	30.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	122	5.3	6.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	81	3.5	4.5
5	Malignant neoplasms (C00-C97)	56	2.4	3.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	27	1.2	1.5
7	Chronic lower respiratory diseases (J40-J47)	17	0.7	*
8	Anemias (D50-D64)	12	0.5	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	11	0.5	*
9	Cerebrovascular diseases (I60-I69)	11	0.5	*
11	Diabetes mellitus (E10-E14)	9	0.4	*
12	Legal intervention (Y35,Y89.0)	8	0.3	*
13	Septicemia (A40-A41)	6	0.3	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.2	*
14	Aortic aneurysm and dissection (I71)	4	0.2	*
...	All other causes (Residual)	179	7.8	10.0
Black, male, 20-24 years				
...	All causes	3,449	100.0	212.3
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,713	49.7	105.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	759	22.0	46.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	230	6.7	14.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	154	4.5	9.5
5	Malignant neoplasms (C00-C97)	89	2.6	5.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	45	1.3	2.8
7	Legal intervention (Y35,Y89.0)	38	1.1	2.3
8	Chronic lower respiratory diseases (J40-J47)	27	0.8	1.7
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	26	0.8	1.6
10	Anemias (D50-D64)	25	0.7	1.5
11	Cerebrovascular diseases (I60-I69)	19	0.6	*
12	Septicemia (A40-A41)	16	0.5	*
13	Diabetes mellitus (E10-E14)	13	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	0.3	*
15	Influenza and pneumonia (J09-J18)	9	0.3	*
...	All other causes (Residual)	275	8.0	16.9

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 25-29 years				
...	All causes	3,398	100.0	227.1
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,350	39.7	90.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	724	21.3	48.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	257	7.6	17.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	211	6.2	14.1
5	Human immunodeficiency virus (HIV) disease (B20-B24)	152	4.5	10.2
6	Malignant neoplasms (C00-C97)	109	3.2	7.3
7	Diabetes mellitus (E10-E14)	42	1.2	2.8
8	Legal intervention (Y35,Y89.0)	32	0.9	2.1
9	Anemias (D50-D64)	30	0.9	2.0
10	Chronic lower respiratory diseases (J40-J47)	28	0.8	1.9
11	Influenza and pneumonia (J09-J18)	27	0.8	1.8
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	25	0.7	1.7
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	25	0.7	1.7
14	Septicemia (A40-A41)	21	0.6	1.4
15	Cerebrovascular diseases (I60-I69)	19	0.6	*
...	All other causes (Residual)	346	10.2	23.1
Black, male, 30-34 years				
...	All causes	3,250	100.0	255.8
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	933	28.7	73.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	645	19.8	50.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	375	11.5	29.5
4	Human immunodeficiency virus (HIV) disease (B20-B24)	224	6.9	17.6
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	198	6.1	15.6
6	Malignant neoplasms (C00-C97)	154	4.7	12.1
7	Diabetes mellitus (E10-E14)	74	2.3	5.8
8	Cerebrovascular diseases (I60-I69)	52	1.6	4.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	35	1.1	2.8
10	Chronic lower respiratory diseases (J40-J47)	30	0.9	2.4
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	28	0.9	2.2
12	Anemias (D50-D64)	26	0.8	2.0
13	Influenza and pneumonia (J09-J18)	24	0.7	1.9
14	Septicemia (A40-A41)	20	0.6	1.6
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	18	0.6	*
...	All other causes (Residual)	414	12.7	32.6

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 35-39 years				
...	All causes	4,147	100.0	313.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	745	18.0	56.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	698	16.8	52.8
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	641	15.5	48.5
4	Human immunodeficiency virus (HIV) disease (B20-B24)	466	11.2	35.3
5	Malignant neoplasms (C00-C97)	253	6.1	19.1
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	178	4.3	13.5
7	Cerebrovascular diseases (I60-I69)	126	3.0	9.5
8	Diabetes mellitus (E10-E14)	125	3.0	9.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	61	1.5	4.6
10	Septicemia (A40-A41)	50	1.2	3.8
11	Influenza and pneumonia (J09-J18)	37	0.9	2.8
12	Anemias (D50-D64)	36	0.9	2.7
13	Aortic aneurysm and dissection (I71)	33	0.8	2.5
13	Chronic lower respiratory diseases (J40-J47)	33	0.8	2.5
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	32	0.8	2.4
...	All other causes (Residual)	633	15.3	47.9
Black, male, 40-44 years				
...	All causes	5,959	100.0	442.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,253	21.0	93.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	818	13.7	60.8
3	Human immunodeficiency virus (HIV) disease (B20-B24)	757	12.7	56.3
4	Malignant neoplasms (C00-C97)	654	11.0	48.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	459	7.7	34.1
6	Cerebrovascular diseases (I60-I69)	209	3.5	15.5
7	Diabetes mellitus (E10-E14)	189	3.2	14.0
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	156	2.6	11.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	116	1.9	8.6
10	Chronic liver disease and cirrhosis (K70,K73-K74)	99	1.7	7.4
11	Septicemia (A40-A41)	77	1.3	5.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	69	1.2	5.1
13	Influenza and pneumonia (J09-J18)	66	1.1	4.9
13	Chronic lower respiratory diseases (J40-J47)	66	1.1	4.9
15	Aortic aneurysm and dissection (I71)	54	0.9	4.0
...	All other causes (Residual)	917	15.4	68.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 45-49 years				
...	All causes	8,970	100.0	678.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,074	23.1	157.0
2	Malignant neoplasms (C00-C97)	1,539	17.2	116.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	986	11.0	74.6
4	Human immunodeficiency virus (HIV) disease (B20-B24)	806	9.0	61.0
5	Cerebrovascular diseases (I60-I69)	404	4.5	30.6
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	361	4.0	27.3
7	Diabetes mellitus (E10-E14)	324	3.6	24.5
8	Chronic liver disease and cirrhosis (K70,K73-K74)	208	2.3	15.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	175	2.0	13.2
10	Septicemia (A40-A41)	152	1.7	11.5
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	135	1.5	10.2
12	Influenza and pneumonia (J09-J18)	132	1.5	10.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	115	1.3	8.7
13	Chronic lower respiratory diseases (J40-J47)	115	1.3	8.7
15	Viral hepatitis (B15-B19)	78	0.9	5.9
...	All other causes (Residual)	1,366	15.2	103.4
Black, male, 50-54 years				
...	All causes	12,525	100.0	1,108.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,230	25.8	285.8
2	Malignant neoplasms (C00-C97)	2,826	22.6	250.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	963	7.7	85.2
4	Human immunodeficiency virus (HIV) disease (B20-B24)	757	6.0	67.0
5	Cerebrovascular diseases (I60-I69)	602	4.8	53.3
6	Diabetes mellitus (E10-E14)	471	3.8	41.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	352	2.8	31.1
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	262	2.1	23.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	260	2.1	23.0
10	Septicemia (A40-A41)	228	1.8	20.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	192	1.5	17.0
12	Chronic lower respiratory diseases (J40-J47)	189	1.5	16.7
13	Viral hepatitis (B15-B19)	176	1.4	15.6
14	Influenza and pneumonia (J09-J18)	139	1.1	12.3
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	110	0.9	9.7
...	All other causes (Residual)	1,768	14.1	156.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 55-59 years				
...	All causes	14,468	100.0	1,612.7
1	Malignant neoplasms (C00-C97)	3,985	27.5	444.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,913	27.0	436.2
3	Cerebrovascular diseases (I60-I69)	779	5.4	86.8
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	732	5.1	81.6
5	Diabetes mellitus (E10-E14)	649	4.5	72.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	451	3.1	50.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	385	2.7	42.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	332	2.3	37.0
9	Chronic lower respiratory diseases (J40-J47)	275	1.9	30.7
10	Septicemia (A40-A41)	273	1.9	30.4
11	Viral hepatitis (B15-B19)	206	1.4	23.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	189	1.3	21.1
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	175	1.2	19.5
14	Influenza and pneumonia (J09-J18)	158	1.1	17.6
15	Aortic aneurysm and dissection (I71)	88	0.6	9.8
...	All other causes (Residual)	1,878	13.0	209.3
Black, male, 60-64 years				
...	All causes	13,659	100.0	2,252.6
1	Malignant neoplasms (C00-C97)	4,196	30.7	692.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,849	28.2	634.8
3	Cerebrovascular diseases (I60-I69)	721	5.3	118.9
4	Diabetes mellitus (E10-E14)	644	4.7	106.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	419	3.1	69.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	398	2.9	65.6
7	Chronic lower respiratory diseases (J40-J47)	397	2.9	65.5
8	Human immunodeficiency virus (HIV) disease (B20-B24)	291	2.1	48.0
9	Septicemia (A40-A41)	255	1.9	42.1
10	Chronic liver disease and cirrhosis (K70,K73-K74)	221	1.6	36.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	217	1.6	35.8
12	Influenza and pneumonia (J09-J18)	174	1.3	28.7
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	93	0.7	15.3
14	Viral hepatitis (B15-B19)	90	0.7	14.8
15	Aortic aneurysm and dissection (I71)	61	0.4	10.1
...	All other causes (Residual)	1,633	12.0	269.3

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 65-69 years				
...	All causes	13,587	100.0	3,111.4
1	Malignant neoplasms (C00-C97)	4,393	32.3	1,006.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,721	27.4	852.1
3	Cerebrovascular diseases (I60-I69)	815	6.0	186.6
4	Diabetes mellitus (E10-E14)	635	4.7	145.4
5	Chronic lower respiratory diseases (J40-J47)	490	3.6	112.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	422	3.1	96.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	336	2.5	76.9
8	Septicemia (A40-A41)	314	2.3	71.9
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	206	1.5	47.2
10	Influenza and pneumonia (J09-J18)	195	1.4	44.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	159	1.2	36.4
12	Human immunodeficiency virus (HIV) disease (B20-B24)	115	0.8	26.3
13	Aortic aneurysm and dissection (I71)	71	0.5	16.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	59	0.4	13.5
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	55	0.4	12.6
...	All other causes (Residual)	1,601	11.8	366.6
Black, male, 70-74 years				
...	All causes	14,085	100.0	4,256.1
1	Malignant neoplasms (C00-C97)	4,354	30.9	1,315.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,874	27.5	1,170.6
3	Cerebrovascular diseases (I60-I69)	899	6.4	271.7
4	Diabetes mellitus (E10-E14)	682	4.8	206.1
5	Chronic lower respiratory diseases (J40-J47)	571	4.1	172.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	455	3.2	137.5
7	Septicemia (A40-A41)	345	2.4	104.2
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	268	1.9	81.0
9	Influenza and pneumonia (J09-J18)	261	1.9	78.9
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	205	1.5	61.9
11	Alzheimer's disease (G30)	125	0.9	37.8
12	Chronic liver disease and cirrhosis (K70,K73-K74)	113	0.8	34.1
13	Pneumonitis due to solids and liquids (J69)	78	0.6	23.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	77	0.5	23.3
15	Human immunodeficiency virus (HIV) disease (B20-B24)	60	0.4	18.1
...	All other causes (Residual)	1,718	12.2	519.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 75-79 years				
...	All causes	14,001	100.0	6,197.6
1	Malignant neoplasms (C00-C97)	4,005	28.6	1,772.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,848	27.5	1,703.3
3	Cerebrovascular diseases (I60-I69)	921	6.6	407.7
4	Chronic lower respiratory diseases (J40-J47)	686	4.9	303.7
5	Diabetes mellitus (E10-E14)	609	4.3	269.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	451	3.2	199.6
7	Septicemia (A40-A41)	319	2.3	141.2
8	Influenza and pneumonia (J09-J18)	302	2.2	133.7
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	234	1.7	103.6
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	230	1.6	101.8
11	Alzheimer's disease (G30)	229	1.6	101.4
12	Pneumonitis due to solids and liquids (J69)	107	0.8	47.4
13	Parkinson's disease (G20-G21)	86	0.6	38.1
14	Aortic aneurysm and dissection (I71)	74	0.5	32.8
15	Chronic liver disease and cirrhosis (K70,K73-K74)	70	0.5	31.0
...	All other causes (Residual)	1,830	13.1	810.1
Black, male, 80-84 years				
...	All causes	12,814	100.0	8,650.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,630	28.3	2,450.5
2	Malignant neoplasms (C00-C97)	3,240	25.3	2,187.2
3	Cerebrovascular diseases (I60-I69)	919	7.2	620.4
4	Chronic lower respiratory diseases (J40-J47)	610	4.8	411.8
5	Diabetes mellitus (E10-E14)	560	4.4	378.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	412	3.2	278.1
7	Influenza and pneumonia (J09-J18)	340	2.7	229.5
8	Alzheimer's disease (G30)	310	2.4	209.3
9	Septicemia (A40-A41)	290	2.3	195.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	206	1.6	139.1
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	164	1.3	110.7
12	Pneumonitis due to solids and liquids (J69)	137	1.1	92.5
13	Parkinson's disease (G20-G21)	115	0.9	77.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	77	0.6	52.0
15	Aortic aneurysm and dissection (I71)	44	0.3	29.7
...	All other causes (Residual)	1,760	13.7	1,188.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 85-89 years				
...	All causes	9,060	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,620	28.9	---
2	Malignant neoplasms (C00-C97)	2,021	22.3	---
3	Cerebrovascular diseases (I60-I69)	604	6.7	---
4	Chronic lower respiratory diseases (J40-J47)	359	4.0	---
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	339	3.7	---
6	Alzheimer's disease (G30)	316	3.5	---
7	Influenza and pneumonia (J09-J18)	313	3.5	---
8	Diabetes mellitus (E10-E14)	312	3.4	---
9	Septicemia (A40-A41)	227	2.5	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	162	1.8	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	150	1.7	---
12	Pneumonitis due to solids and liquids (J69)	118	1.3	---
13	Parkinson's disease (G20-G21)	71	0.8	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	42	0.5	---
15	Atherosclerosis (I70)	41	0.5	---
...	All other causes (Residual)	1,365	15.1	---
Black, male, 90-94 years				
...	All causes	4,421	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,344	30.4	---
2	Malignant neoplasms (C00-C97)	846	19.1	---
3	Cerebrovascular diseases (I60-I69)	300	6.8	---
4	Alzheimer's disease (G30)	196	4.4	---
5	Chronic lower respiratory diseases (J40-J47)	180	4.1	---
6	Influenza and pneumonia (J09-J18)	166	3.8	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	162	3.7	---
8	Diabetes mellitus (E10-E14)	121	2.7	---
9	Septicemia (A40-A41)	115	2.6	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	89	2.0	---
11	Pneumonitis due to solids and liquids (J69)	52	1.2	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	52	1.2	---
13	Parkinson's disease (G20-G21)	38	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	24	0.5	---
15	Atherosclerosis (I70)	22	0.5	---
...	All other causes (Residual)	714	16.2	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 95-99 years				
...	All causes	1,420	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	471	33.2	---
2	Malignant neoplasms (C00-C97)	198	13.9	---
3	Cerebrovascular diseases (I60-I69)	91	6.4	---
4	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	63	4.4	---
5	Alzheimer's disease (G30)	60	4.2	---
5	Influenza and pneumonia (J09-J18)	60	4.2	---
7	Chronic lower respiratory diseases (J40-J47)	57	4.0	---
8	Septicemia (A40-A41)	41	2.9	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	34	2.4	---
10	Diabetes mellitus (E10-E14)	26	1.8	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	25	1.8	---
12	Pneumonitis due to solids and liquids (J69)	17	1.2	---
13	Atherosclerosis (I70)	9	0.6	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.5	---
14	Nutritional deficiencies (E40-E64)	7	0.5	---
...	All other causes (Residual)	254	17.9	---
Black, male, 100 years and over				
...	All causes	354	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	130	36.7	---
2	Malignant neoplasms (C00-C97)	38	10.7	---
3	Cerebrovascular diseases (I60-I69)	25	7.1	---
4	Influenza and pneumonia (J09-J18)	21	5.9	---
5	Alzheimer's disease (G30)	17	4.8	---
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	4.8	---
7	Chronic lower respiratory diseases (J40-J47)	10	2.8	---
8	Septicemia (A40-A41)	7	2.0	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6	1.7	---
10	Diabetes mellitus (E10-E14)	4	1.1	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4	1.1	---
12	Parkinson's disease (G20-G21)	2	0.6	---
12	Atherosclerosis (I70)	2	0.6	---
12	Pneumonitis due to solids and liquids (J69)	2	0.6	---
15	Nutritional deficiencies (E40-E64)	1	0.3	---
15	Aortic aneurysm and dissection (I71)	1	0.3	---
15	Hernia (K40-K46)	1	0.3	---
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.3	---
...	All other causes (Residual)	65	18.4	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, all ages				
...	All causes	141,276	100.0	675.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	35,540	25.2	170.0
2	Malignant neoplasms (C00-C97)	30,980	21.9	148.2
3	Cerebrovascular diseases (I60-I69)	9,536	6.7	45.6
4	Diabetes mellitus (E10-E14)	6,966	4.9	33.3
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,620	3.3	22.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,291	3.0	20.5
7	Chronic lower respiratory diseases (J40-J47)	3,694	2.6	17.7
8	Septicemia (A40-A41)	3,462	2.5	16.6
9	Alzheimer's disease (G30)	3,459	2.4	16.5
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,661	1.9	12.7
11	Influenza and pneumonia (J09-J18)	2,657	1.9	12.7
12	Human immunodeficiency virus (HIV) disease (B20-B24)	2,284	1.6	10.9
13	Certain conditions originating in the perinatal period (P00-P96)	2,206	1.6	10.6
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,286	0.9	6.2
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	883	0.6	4.2
...	All other causes (Residual)	26,751	18.9	127.9
Black, female, 1-4 years				
...	All causes	516	100.0	39.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	136	26.4	10.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	79	15.3	6.0
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	61	11.8	4.6
4	Malignant neoplasms (C00-C97)	22	4.3	1.7
5	Chronic lower respiratory diseases (J40-J47)	17	3.3	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	15	2.9	*
6	Certain conditions originating in the perinatal period (P00-P96)	15	2.9	*
8	Influenza and pneumonia (J09-J18)	13	2.5	*
9	Cerebrovascular diseases (I60-I69)	12	2.3	*
10	Anemias (D50-D64)	8	1.6	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.2	*
12	Septicemia (A40-A41)	5	1.0	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.6	*
14	Acute bronchitis and bronchiolitis (J20-J21)	2	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
...	All other causes (Residual)	120	23.3	9.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 5-9 years				
...	All causes	246	100.0	15.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	86	35.0	5.5
2	Malignant neoplasms (C00-C97)	33	13.4	2.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	21	8.5	1.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	16	6.5	*
5	Chronic lower respiratory diseases (J40-J47)	8	3.3	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	7	2.8	*
7	Septicemia (A40-A41)	4	1.6	*
7	Cerebrovascular diseases (I60-I69)	4	1.6	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	1.2	*
9	Anemias (D50-D64)	3	1.2	*
9	Influenza and pneumonia (J09-J18)	3	1.2	*
9	Certain conditions originating in the perinatal period (P00-P96)	3	1.2	*
13	Diabetes mellitus (E10-E14)	2	0.8	*
13	Diseases of appendix (K35-K38)	2	0.8	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.8	*
...	All other causes (Residual)	49	19.9	3.1
Black, female, 10-14 years				
...	All causes	301	100.0	18.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	94	31.2	5.7
2	Malignant neoplasms (C00-C97)	38	12.6	2.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	28	9.3	1.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	21	7.0	1.3
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	19	6.3	*
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	12	4.0	*
7	Chronic lower respiratory diseases (J40-J47)	11	3.7	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	6	2.0	*
8	Cerebrovascular diseases (I60-I69)	6	2.0	*
10	Influenza and pneumonia (J09-J18)	5	1.7	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	1.0	*
12	Septicemia (A40-A41)	2	0.7	*
12	Anemias (D50-D64)	2	0.7	*
12	Diabetes mellitus (E10-E14)	2	0.7	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.7	*
...	All other causes (Residual)	50	16.6	3.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 15-19 years				
...	All causes	654	100.0	37.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	196	30.0	11.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	120	18.3	6.9
3	Malignant neoplasms (C00-C97)	50	7.6	2.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	33	5.0	1.9
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	22	3.4	1.3
6	Pregnancy, childbirth and the puerperium (O00-O99)	20	3.1	1.1
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	19	2.9	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	14	2.1	*
8	Anemias (D50-D64)	14	2.1	*
10	Diabetes mellitus (E10-E14)	12	1.8	*
11	Chronic lower respiratory diseases (J40-J47)	11	1.7	*
12	Septicemia (A40-A41)	6	0.9	*
13	Cerebrovascular diseases (I60-I69)	5	0.8	*
13	Influenza and pneumonia (J09-J18)	5	0.8	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.8	*
...	All other causes (Residual)	122	18.7	7.0
Black, female, 20-24 years				
...	All causes	975	100.0	61.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	244	25.0	15.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	175	17.9	11.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	80	8.2	5.1
4	Malignant neoplasms (C00-C97)	66	6.8	4.2
5	Pregnancy, childbirth and the puerperium (O00-O99)	49	5.0	3.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	39	4.0	2.5
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	32	3.3	2.0
8	Diabetes mellitus (E10-E14)	22	2.3	1.4
9	Septicemia (A40-A41)	19	1.9	*
10	Anemias (D50-D64)	16	1.6	*
11	Cerebrovascular diseases (I60-I69)	11	1.1	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	11	1.1	*
13	Chronic lower respiratory diseases (J40-J47)	10	1.0	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.9	*
15	Influenza and pneumonia (J09-J18)	6	0.6	*
...	All other causes (Residual)	186	19.1	11.8

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 25-29 years				
...	All causes	1,343	100.0	86.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	205	15.3	13.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	165	12.3	10.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	144	10.7	9.3
4	Human immunodeficiency virus (HIV) disease (B20-B24)	132	9.8	8.5
5	Malignant neoplasms (C00-C97)	125	9.3	8.0
6	Pregnancy, childbirth and the puerperium (O00-O99)	56	4.2	3.6
7	Diabetes mellitus (E10-E14)	41	3.1	2.6
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	39	2.9	2.5
9	Septicemia (A40-A41)	24	1.8	1.5
9	Cerebrovascular diseases (I60-I69)	24	1.8	1.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	1.7	1.5
12	Anemias (D50-D64)	22	1.6	1.4
13	Influenza and pneumonia (J09-J18)	19	1.4	*
14	Chronic lower respiratory diseases (J40-J47)	17	1.3	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	16	1.2	*
...	All other causes (Residual)	291	21.7	18.7
Black, female, 30-34 years				
...	All causes	1,671	100.0	119.5
1	Malignant neoplasms (C00-C97)	230	13.8	16.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	221	13.2	15.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	205	12.3	14.7
4	Human immunodeficiency virus (HIV) disease (B20-B24)	197	11.8	14.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	112	6.7	8.0
6	Diabetes mellitus (E10-E14)	56	3.4	4.0
7	Cerebrovascular diseases (I60-I69)	53	3.2	3.8
8	Pregnancy, childbirth and the puerperium (O00-O99)	52	3.1	3.7
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	33	2.0	2.4
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	30	1.8	2.1
11	Septicemia (A40-A41)	25	1.5	1.8
12	Anemias (D50-D64)	24	1.4	1.7
12	Influenza and pneumonia (J09-J18)	24	1.4	1.7
14	Chronic lower respiratory diseases (J40-J47)	21	1.3	1.5
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	16	1.0	*
...	All other causes (Residual)	372	22.3	26.6

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 35-39 years				
...	All causes	2,684	100.0	181.5
1	Malignant neoplasms (C00-C97)	465	17.3	31.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	383	14.3	25.9
3	Human immunodeficiency virus (HIV) disease (B20-B24)	346	12.9	23.4
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	276	10.3	18.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	136	5.1	9.2
6	Cerebrovascular diseases (I60-I69)	90	3.4	6.1
7	Diabetes mellitus (E10-E14)	89	3.3	6.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	74	2.8	5.0
9	Septicemia (A40-A41)	58	2.2	3.9
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	47	1.8	3.2
11	Pregnancy, childbirth and the puerperium (O00-O99)	43	1.6	2.9
12	Influenza and pneumonia (J09-J18)	35	1.3	2.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	34	1.3	2.3
14	Chronic lower respiratory diseases (J40-J47)	32	1.2	2.2
15	Anemias (D50-D64)	28	1.0	1.9
...	All other causes (Residual)	548	20.4	37.1
Black, female, 40-44 years				
...	All causes	4,200	100.0	275.2
1	Malignant neoplasms (C00-C97)	993	23.6	65.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	658	15.7	43.1
3	Human immunodeficiency virus (HIV) disease (B20-B24)	433	10.3	28.4
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	319	7.6	20.9
5	Cerebrovascular diseases (I60-I69)	230	5.5	15.1
6	Diabetes mellitus (E10-E14)	159	3.8	10.4
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	130	3.1	8.5
8	Septicemia (A40-A41)	89	2.1	5.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	79	1.9	5.2
10	Chronic liver disease and cirrhosis (K70,K73-K74)	63	1.5	4.1
11	Chronic lower respiratory diseases (J40-J47)	62	1.5	4.1
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	43	1.0	2.8
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	41	1.0	2.7
14	Influenza and pneumonia (J09-J18)	40	1.0	2.6
15	Anemias (D50-D64)	36	0.9	2.4
...	All other causes (Residual)	825	19.6	54.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 45-49 years				
...	All causes	6,639	100.0	437.0
1	Malignant neoplasms (C00-C97)	1,824	27.5	120.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,247	18.8	82.1
3	Human immunodeficiency virus (HIV) disease (B20-B24)	444	6.7	29.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	413	6.2	27.2
5	Cerebrovascular diseases (I60-I69)	391	5.9	25.7
6	Diabetes mellitus (E10-E14)	220	3.3	14.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	168	2.5	11.1
8	Septicemia (A40-A41)	150	2.3	9.9
9	Chronic lower respiratory diseases (J40-J47)	146	2.2	9.6
10	Chronic liver disease and cirrhosis (K70,K73-K74)	143	2.2	9.4
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	119	1.8	7.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	86	1.3	5.7
13	Influenza and pneumonia (J09-J18)	76	1.1	5.0
14	Viral hepatitis (B15-B19)	45	0.7	3.0
15	Anemias (D50-D64)	44	0.7	2.9
...	All other causes (Residual)	1,123	16.9	73.9
Black, female, 50-54 years				
...	All causes	8,684	100.0	651.5
1	Malignant neoplasms (C00-C97)	2,629	30.3	197.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,806	20.8	135.5
3	Cerebrovascular diseases (I60-I69)	549	6.3	41.2
4	Diabetes mellitus (E10-E14)	418	4.8	31.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	404	4.7	30.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	344	4.0	25.8
7	Chronic lower respiratory diseases (J40-J47)	222	2.6	16.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	220	2.5	16.5
9	Septicemia (A40-A41)	191	2.2	14.3
10	Chronic liver disease and cirrhosis (K70,K73-K74)	150	1.7	11.3
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	142	1.6	10.7
12	Influenza and pneumonia (J09-J18)	92	1.1	6.9
13	Viral hepatitis (B15-B19)	83	1.0	6.2
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	60	0.7	4.5
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	36	0.4	2.7
...	All other causes (Residual)	1,338	15.4	100.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 55-59 years				
...	All causes	9,722	100.0	889.3
1	Malignant neoplasms (C00-C97)	3,228	33.2	295.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,185	22.5	199.9
3	Diabetes mellitus (E10-E14)	554	5.7	50.7
4	Cerebrovascular diseases (I60-I69)	527	5.4	48.2
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	308	3.2	28.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	275	2.8	25.2
7	Chronic lower respiratory diseases (J40-J47)	250	2.6	22.9
8	Septicemia (A40-A41)	234	2.4	21.4
9	Human immunodeficiency virus (HIV) disease (B20-B24)	181	1.9	16.6
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	175	1.8	16.0
11	Chronic liver disease and cirrhosis (K70,K73-K74)	160	1.6	14.6
12	Influenza and pneumonia (J09-J18)	131	1.3	12.0
13	Viral hepatitis (B15-B19)	78	0.8	7.1
14	Aortic aneurysm and dissection (I71)	38	0.4	3.5
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	35	0.4	3.2
...	All other causes (Residual)	1,363	14.0	124.7
Black, female, 60-64 years				
...	All causes	9,833	100.0	1,270.7
1	Malignant neoplasms (C00-C97)	3,353	34.1	433.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,342	23.8	302.6
3	Diabetes mellitus (E10-E14)	639	6.5	82.6
4	Cerebrovascular diseases (I60-I69)	564	5.7	72.9
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	353	3.6	45.6
6	Chronic lower respiratory diseases (J40-J47)	256	2.6	33.1
7	Septicemia (A40-A41)	242	2.5	31.3
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	173	1.8	22.4
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	171	1.7	22.1
10	Influenza and pneumonia (J09-J18)	110	1.1	14.2
11	Chronic liver disease and cirrhosis (K70,K73-K74)	101	1.0	13.1
12	Human immunodeficiency virus (HIV) disease (B20-B24)	72	0.7	9.3
13	Aortic aneurysm and dissection (I71)	45	0.5	5.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	38	0.4	4.9
15	Viral hepatitis (B15-B19)	35	0.4	4.5
...	All other causes (Residual)	1,339	13.6	173.0

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 65-69 years				
...	All causes	10,968	100.0	1,848.0
1	Malignant neoplasms (C00-C97)	3,635	33.1	612.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,722	24.8	458.6
3	Diabetes mellitus (E10-E14)	696	6.3	117.3
4	Cerebrovascular diseases (I60-I69)	687	6.3	115.8
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	415	3.8	69.9
6	Chronic lower respiratory diseases (J40-J47)	381	3.5	64.2
7	Septicemia (A40-A41)	306	2.8	51.6
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	192	1.8	32.3
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	138	1.3	23.3
10	Influenza and pneumonia (J09-J18)	137	1.2	23.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	63	0.6	10.6
12	Aortic aneurysm and dissection (I71)	57	0.5	9.6
13	Alzheimer's disease (G30)	49	0.4	8.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	46	0.4	7.8
15	Human immunodeficiency virus (HIV) disease (B20-B24)	35	0.3	5.9
...	All other causes (Residual)	1,409	12.8	237.4
Black, female, 70-74 years				
...	All causes	12,744	100.0	2,656.7
1	Malignant neoplasms (C00-C97)	3,673	28.8	765.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,325	26.1	693.2
3	Cerebrovascular diseases (I60-I69)	856	6.7	178.5
4	Diabetes mellitus (E10-E14)	848	6.7	176.8
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	549	4.3	114.5
6	Chronic lower respiratory diseases (J40-J47)	455	3.6	94.9
7	Septicemia (A40-A41)	355	2.8	74.0
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	227	1.8	47.3
9	Influenza and pneumonia (J09-J18)	207	1.6	43.2
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	169	1.3	35.2
11	Alzheimer's disease (G30)	159	1.2	33.1
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	69	0.5	14.4
12	Chronic liver disease and cirrhosis (K70,K73-K74)	69	0.5	14.4
14	Aortic aneurysm and dissection (I71)	52	0.4	10.8
15	Pneumonitis due to solids and liquids (J69)	44	0.3	9.2
...	All other causes (Residual)	1,687	13.2	351.7

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 75-79 years				
...	All causes	15,062	100.0	4,019.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,078	27.1	1,088.2
2	Malignant neoplasms (C00-C97)	3,737	24.8	997.2
3	Cerebrovascular diseases (I60-I69)	1,147	7.6	306.1
4	Diabetes mellitus (E10-E14)	930	6.2	248.2
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	595	4.0	158.8
6	Chronic lower respiratory diseases (J40-J47)	527	3.5	140.6
7	Septicemia (A40-A41)	422	2.8	112.6
8	Alzheimer's disease (G30)	391	2.6	104.3
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	294	2.0	78.5
10	Influenza and pneumonia (J09-J18)	274	1.8	73.1
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	174	1.2	46.4
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	81	0.5	21.6
13	Pneumonitis due to solids and liquids (J69)	72	0.5	19.2
14	Parkinson's disease (G20-G21)	65	0.4	17.3
14	Aortic aneurysm and dissection (I71)	65	0.4	17.3
...	All other causes (Residual)	2,210	14.7	589.7
Black, female, 80-84 years				
...	All causes	17,109	100.0	6,079.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,003	29.2	1,777.9
2	Malignant neoplasms (C00-C97)	3,295	19.3	1,170.9
3	Cerebrovascular diseases (I60-I69)	1,398	8.2	496.8
4	Diabetes mellitus (E10-E14)	969	5.7	344.3
5	Alzheimer's disease (G30)	697	4.1	247.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	601	3.5	213.6
7	Chronic lower respiratory diseases (J40-J47)	517	3.0	183.7
8	Septicemia (A40-A41)	488	2.9	173.4
9	Influenza and pneumonia (J09-J18)	396	2.3	140.7
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	365	2.1	129.7
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	192	1.1	68.2
12	Pneumonitis due to solids and liquids (J69)	122	0.7	43.4
13	Parkinson's disease (G20-G21)	95	0.6	33.8
14	Aortic aneurysm and dissection (I71)	88	0.5	31.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	87	0.5	30.9
...	All other causes (Residual)	2,796	16.3	993.6

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 85-89 years				
...	All causes	15,825	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,977	31.5	---
2	Malignant neoplasms (C00-C97)	2,106	13.3	---
3	Cerebrovascular diseases (I60-I69)	1,459	9.2	---
4	Alzheimer's disease (G30)	829	5.2	---
5	Diabetes mellitus (E10-E14)	732	4.6	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	630	4.0	---
7	Influenza and pneumonia (J09-J18)	441	2.8	---
8	Chronic lower respiratory diseases (J40-J47)	416	2.6	---
9	Septicemia (A40-A41)	403	2.5	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	402	2.5	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	167	1.1	---
12	Pneumonitis due to solids and liquids (J69)	143	0.9	---
13	Parkinson's disease (G20-G21)	79	0.5	---
14	Atherosclerosis (I70)	71	0.4	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	65	0.4	---
...	All other causes (Residual)	2,905	18.4	---
Black, female, 90-94 years				
...	All causes	10,998	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,645	33.1	---
2	Malignant neoplasms (C00-C97)	1,037	9.4	---
3	Cerebrovascular diseases (I60-I69)	1,002	9.1	---
4	Alzheimer's disease (G30)	781	7.1	---
5	Diabetes mellitus (E10-E14)	378	3.4	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	346	3.1	---
7	Influenza and pneumonia (J09-J18)	334	3.0	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	288	2.6	---
9	Septicemia (A40-A41)	258	2.3	---
10	Chronic lower respiratory diseases (J40-J47)	219	2.0	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	144	1.3	---
12	Pneumonitis due to solids and liquids (J69)	100	0.9	---
13	Atherosclerosis (I70)	70	0.6	---
14	Parkinson's disease (G20-G21)	43	0.4	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	42	0.4	---
...	All other causes (Residual)	2,311	21.0	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 95-99 years				
...	All causes	5,357	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,955	36.5	---
2	Cerebrovascular diseases (I60-I69)	400	7.5	---
3	Alzheimer's disease (G30)	396	7.4	---
4	Malignant neoplasms (C00-C97)	362	6.8	---
5	Influenza and pneumonia (J09-J18)	192	3.6	---
6	Diabetes mellitus (E10-E14)	157	2.9	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	157	2.9	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	146	2.7	---
9	Septicemia (A40-A41)	99	1.8	---
10	Chronic lower respiratory diseases (J40-J47)	90	1.7	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	57	1.1	---
12	Pneumonitis due to solids and liquids (J69)	49	0.9	---
13	Atherosclerosis (I70)	46	0.9	---
14	Parkinson's disease (G20-G21)	19	0.4	---
15	Nutritional deficiencies (E40-E64)	17	0.3	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	17	0.3	---
...	All other causes (Residual)	1,198	22.4	---
Black, female, 100 years and over				
...	All causes	1,759	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	631	35.9	---
2	Alzheimer's disease (G30)	129	7.3	---
3	Cerebrovascular diseases (I60-I69)	111	6.3	---
4	Influenza and pneumonia (J09-J18)	84	4.8	---
5	Malignant neoplasms (C00-C97)	74	4.2	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	56	3.2	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	48	2.7	---
8	Diabetes mellitus (E10-E14)	41	2.3	---
9	Septicemia (A40-A41)	29	1.6	---
10	Chronic lower respiratory diseases (J40-J47)	24	1.4	---
11	Pneumonitis due to solids and liquids (J69)	21	1.2	---
12	Atherosclerosis (I70)	19	1.1	---
13	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	16	0.9	---
14	Nutritional deficiencies (E40-E64)	9	0.5	---
15	Parkinson's disease (G20-G21)	6	0.3	---
...	All other causes (Residual)	461	26.2	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, all ages				
...	All causes	14,367	100.0	444.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,648	18.4	81.8
2	Malignant neoplasms (C00-C97)	2,561	17.8	79.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,701	11.8	52.6
4	Diabetes mellitus (E10-E14)	790	5.5	24.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	709	4.9	21.9
6	Chronic lower respiratory diseases (J40-J47)	611	4.3	18.9
7	Cerebrovascular diseases (I60-I69)	586	4.1	18.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	392	2.7	12.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	292	2.0	9.0
10	Influenza and pneumonia (J09-J18)	280	1.9	8.7
11	Septicemia (A40-A41)	230	1.6	7.1
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	220	1.5	6.8
13	Alzheimer's disease (G30)	191	1.3	5.9
14	Certain conditions originating in the perinatal period (P00-P96)	158	1.1	4.9
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	124	0.9	3.8
...	All other causes (Residual)	2,874	20.0	88.8
American Indian, both sexes, 1-4 years				
...	All causes	97	100.0	54.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	43	44.3	24.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	8.2	*
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	7	7.2	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	5.2	*
5	Influenza and pneumonia (J09-J18)	2	2.1	*
5	Chronic lower respiratory diseases (J40-J47)	2	2.1	*
7	Malignant neoplasms (C00-C97)	1	1.0	*
7	Cerebrovascular diseases (I60-I69)	1	1.0	*
7	Acute bronchitis and bronchiolitis (J20-J21)	1	1.0	*
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	1.0	*
7	Certain conditions originating in the perinatal period (P00-P96)	1	1.0	*
7	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	1.0	*
...	All other causes (Residual)	24	24.7	13.6

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 5-9 years				
...	All causes	40	100.0	16.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18	45.0	*
2	Malignant neoplasms (C00-C97)	5	12.5	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	10.0	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	2.5	*
4	Chronic lower respiratory diseases (J40-J47)	1	2.5	*
4	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	2.5	*
...	All other causes (Residual)	10	25.0	*
American Indian, both sexes, 10-14 years				
...	All causes	54	100.0	19.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	28	51.9	10.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	9	16.7	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	5.6	*
4	Septicemia (A40-A41)	2	3.7	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	3.7	*
6	Malignant neoplasms (C00-C97)	1	1.9	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.9	*
6	Chronic lower respiratory diseases (J40-J47)	1	1.9	*
...	All other causes (Residual)	7	13.0	*

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 15-19 years				
...	All causes	265	100.0	86.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	133	50.2	43.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	51	19.2	16.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	21	7.9	6.9
4	Malignant neoplasms (C00-C97)	11	4.2	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	1.5	*
5	Pregnancy, childbirth and the puerperium (O00-O99)	4	1.5	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	1.1	*
8	Septicemia (A40-A41)	2	0.8	*
8	Influenza and pneumonia (J09-J18)	2	0.8	*
8	Chronic lower respiratory diseases (J40-J47)	2	0.8	*
8	Pneumonitis due to solids and liquids (J69)	2	0.8	*
12	Anemias (D50-D64)	1	0.4	*
12	Cerebrovascular diseases (I60-I69)	1	0.4	*
12	Aortic aneurysm and dissection (I71)	1	0.4	*
12	Diseases of appendix (K35-K38)	1	0.4	*
12	Legal intervention (Y35,Y89.0)	1	0.4	*
...	All other causes (Residual)	25	9.4	8.2
American Indian, both sexes, 20-24 years				
...	All causes	357	100.0	120.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	186	52.1	62.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	71	19.9	24.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	36	10.1	12.2
4	Malignant neoplasms (C00-C97)	11	3.1	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	8	2.2	*
6	Chronic liver disease and cirrhosis (K70,K73-K74)	3	0.8	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.8	*
8	Influenza and pneumonia (J09-J18)	2	0.6	*
9	Cerebrovascular diseases (I60-I69)	1	0.3	*
9	Aortic aneurysm and dissection (I71)	1	0.3	*
...	All other causes (Residual)	35	9.8	11.8

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 25-29 years				
...	All causes	349	100.0	129.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	154	44.1	57.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	54	15.5	20.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	33	9.5	12.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	21	6.0	7.8
5	Malignant neoplasms (C00-C97)	14	4.0	*
6	Chronic liver disease and cirrhosis (K70,K73-K74)	13	3.7	*
7	Pregnancy, childbirth and the puerperium (O00-O99)	4	1.1	*
8	Septicemia (A40-A41)	3	0.9	*
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.9	*
10	Diabetes mellitus (E10-E14)	2	0.6	*
10	Cerebrovascular diseases (I60-I69)	2	0.6	*
12	Viral hepatitis (B15-B19)	1	0.3	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.3	*
12	Aortic aneurysm and dissection (I71)	1	0.3	*
12	Influenza and pneumonia (J09-J18)	1	0.3	*
12	Chronic lower respiratory diseases (J40-J47)	1	0.3	*
12	Pneumonitis due to solids and liquids (J69)	1	0.3	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
12	Legal intervention (Y35,Y89.0)	1	0.3	*
12	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.3	*
...	All other causes (Residual)	37	10.6	13.8
American Indian, both sexes, 30-34 years				
...	All causes	382	100.0	166.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	155	40.6	67.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	44	11.5	19.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	39	10.2	17.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	32	8.4	13.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	24	6.3	10.4
6	Malignant neoplasms (C00-C97)	11	2.9	*
7	Human immunodeficiency virus (HIV) disease (B20-B24)	4	1.0	*
7	Diabetes mellitus (E10-E14)	4	1.0	*
9	Septicemia (A40-A41)	3	0.8	*
9	Influenza and pneumonia (J09-J18)	3	0.8	*
9	Pregnancy, childbirth and the puerperium (O00-O99)	3	0.8	*
12	Salmonella infections (A01-A02)	1	0.3	*
12	Scarlet fever and erysipelas (A38,A46)	1	0.3	*
12	Cerebrovascular diseases (I60-I69)	1	0.3	*
12	Aortic aneurysm and dissection (I71)	1	0.3	*
12	Chronic lower respiratory diseases (J40-J47)	1	0.3	*
12	Pneumonitis due to solids and liquids (J69)	1	0.3	*
...	All other causes (Residual)	54	14.1	23.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 35-39 years				
...	All causes	497	100.0	219.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	155	31.2	68.4
2	Chronic liver disease and cirrhosis (K70,K73-K74)	54	10.9	23.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	40	8.0	17.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	32	6.4	14.1
5	Malignant neoplasms (C00-C97)	27	5.4	11.9
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	25	5.0	11.0
7	Diabetes mellitus (E10-E14)	21	4.2	9.3
8	Human immunodeficiency virus (HIV) disease (B20-B24)	15	3.0	*
9	Cerebrovascular diseases (I60-I69)	9	1.8	*
10	Influenza and pneumonia (J09-J18)	6	1.2	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	1.2	*
12	Viral hepatitis (B15-B19)	4	0.8	*
12	Pregnancy, childbirth and the puerperium (O00-O99)	4	0.8	*
14	Septicemia (A40-A41)	3	0.6	*
14	Chronic lower respiratory diseases (J40-J47)	3	0.6	*
...	All other causes (Residual)	93	18.7	41.1
American Indian, both sexes, 40-44 years				
...	All causes	716	100.0	309.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	165	23.0	71.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	97	13.5	41.9
3	Chronic liver disease and cirrhosis (K70,K73-K74)	91	12.7	39.3
4	Malignant neoplasms (C00-C97)	56	7.8	24.2
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	42	5.9	18.1
6	Diabetes mellitus (E10-E14)	24	3.4	10.4
7	Cerebrovascular diseases (I60-I69)	19	2.7	*
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	18	2.5	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	16	2.2	*
10	Viral hepatitis (B15-B19)	15	2.1	*
11	Influenza and pneumonia (J09-J18)	12	1.7	*
12	Septicemia (A40-A41)	9	1.3	*
13	Chronic lower respiratory diseases (J40-J47)	7	1.0	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.7	*
15	Aortic aneurysm and dissection (I71)	3	0.4	*
15	Pregnancy, childbirth and the puerperium (O00-O99)	3	0.4	*
...	All other causes (Residual)	134	18.7	57.9

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 45-49 years				
...	All causes	937	100.0	407.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	171	18.2	74.4
2	Malignant neoplasms (C00-C97)	147	15.7	63.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	134	14.3	58.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	124	13.2	53.9
5	Diabetes mellitus (E10-E14)	39	4.2	17.0
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	26	2.8	11.3
7	Cerebrovascular diseases (I60-I69)	24	2.6	10.4
8	Human immunodeficiency virus (HIV) disease (B20-B24)	21	2.2	9.1
9	Septicemia (A40-A41)	20	2.1	8.7
10	Influenza and pneumonia (J09-J18)	17	1.8	*
10	Chronic lower respiratory diseases (J40-J47)	17	1.8	*
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	16	1.7	*
13	Viral hepatitis (B15-B19)	13	1.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	1.1	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	0.4	*
...	All other causes (Residual)	154	16.4	67.0
American Indian, both sexes, 50-54 years				
...	All causes	1,021	100.0	512.8
1	Malignant neoplasms (C00-C97)	183	17.9	91.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	181	17.7	90.9
3	Chronic liver disease and cirrhosis (K70,K73-K74)	135	13.2	67.8
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	109	10.7	54.7
5	Diabetes mellitus (E10-E14)	53	5.2	26.6
6	Chronic lower respiratory diseases (J40-J47)	31	3.0	15.6
7	Cerebrovascular diseases (I60-I69)	27	2.6	13.6
8	Septicemia (A40-A41)	26	2.5	13.1
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	24	2.4	12.1
10	Viral hepatitis (B15-B19)	20	2.0	10.0
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	1.3	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	11	1.1	*
12	Influenza and pneumonia (J09-J18)	11	1.1	*
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7	0.7	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5	0.5	*
...	All other causes (Residual)	185	18.1	92.9

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 55-59 years				
...	All causes	1,186	100.0	739.0
1	Malignant neoplasms (C00-C97)	291	24.5	181.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	235	19.8	146.4
3	Chronic liver disease and cirrhosis (K70,K73-K74)	100	8.4	62.3
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	85	7.2	53.0
5	Diabetes mellitus (E10-E14)	82	6.9	51.1
6	Cerebrovascular diseases (I60-I69)	50	4.2	31.2
7	Chronic lower respiratory diseases (J40-J47)	41	3.5	25.5
8	Viral hepatitis (B15-B19)	28	2.4	17.4
9	Septicemia (A40-A41)	22	1.9	13.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21	1.8	13.1
11	Influenza and pneumonia (J09-J18)	17	1.4	*
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11	0.9	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.5	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	6	0.5	*
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	0.5	*
...	All other causes (Residual)	185	15.6	115.3
American Indian, both sexes, 60-64 years				
...	All causes	1,181	100.0	1,024.2
1	Malignant neoplasms (C00-C97)	332	28.1	287.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	233	19.7	202.1
3	Diabetes mellitus (E10-E14)	85	7.2	73.7
4	Chronic liver disease and cirrhosis (K70,K73-K74)	67	5.7	58.1
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	65	5.5	56.4
6	Chronic lower respiratory diseases (J40-J47)	59	5.0	51.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	36	3.0	31.2
8	Cerebrovascular diseases (I60-I69)	33	2.8	28.6
9	Septicemia (A40-A41)	22	1.9	19.1
10	Influenza and pneumonia (J09-J18)	18	1.5	*
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	12	1.0	*
12	Viral hepatitis (B15-B19)	11	0.9	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10	0.8	*
14	Aortic aneurysm and dissection (I71)	6	0.5	*
15	Alzheimer's disease (G30)	5	0.4	*
...	All other causes (Residual)	187	15.8	162.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 65-69 years				
...	All causes	1,222	100.0	1,534.8
1	Malignant neoplasms (C00-C97)	311	25.5	390.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	284	23.2	356.7
3	Diabetes mellitus (E10-E14)	102	8.3	128.1
4	Chronic lower respiratory diseases (J40-J47)	85	7.0	106.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	52	4.3	65.3
6	Cerebrovascular diseases (I60-I69)	44	3.6	55.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	37	3.0	46.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	33	2.7	41.4
9	Influenza and pneumonia (J09-J18)	20	1.6	25.1
10	Septicemia (A40-A41)	18	1.5	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	13	1.1	*
12	Alzheimer's disease (G30)	9	0.7	*
13	Viral hepatitis (B15-B19)	7	0.6	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.6	*
13	Aortic aneurysm and dissection (I71)	7	0.6	*
...	All other causes (Residual)	193	15.8	242.4
American Indian, both sexes, 70-74 years				
...	All causes	1,314	100.0	2,298.0
1	Malignant neoplasms (C00-C97)	374	28.5	654.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	259	19.7	452.9
3	Diabetes mellitus (E10-E14)	106	8.1	185.4
4	Chronic lower respiratory diseases (J40-J47)	88	6.7	153.9
5	Cerebrovascular diseases (I60-I69)	70	5.3	122.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	41	3.1	71.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	41	3.1	71.7
8	Septicemia (A40-A41)	31	2.4	54.2
9	Chronic liver disease and cirrhosis (K70,K73-K74)	29	2.2	50.7
10	Influenza and pneumonia (J09-J18)	19	1.4	*
11	Alzheimer's disease (G30)	15	1.1	*
12	Parkinson's disease (G20-G21)	9	0.7	*
13	Aortic aneurysm and dissection (I71)	8	0.6	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.5	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7	0.5	*
14	Pneumonitis due to solids and liquids (J69)	7	0.5	*
...	All other causes (Residual)	203	15.4	355.0

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 75-79 years				
...	All causes	1,351	100.0	3,324.7
1	Malignant neoplasms (C00-C97)	336	24.9	826.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	314	23.2	772.7
3	Diabetes mellitus (E10-E14)	89	6.6	219.0
3	Chronic lower respiratory diseases (J40-J47)	89	6.6	219.0
5	Cerebrovascular diseases (I60-I69)	77	5.7	189.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	43	3.2	105.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	38	2.8	93.5
8	Influenza and pneumonia (J09-J18)	35	2.6	86.1
9	Alzheimer's disease (G30)	25	1.9	61.5
10	Septicemia (A40-A41)	20	1.5	49.2
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	20	1.5	49.2
12	Chronic liver disease and cirrhosis (K70,K73-K74)	12	0.9	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	0.7	*
14	Pneumonitis due to solids and liquids (J69)	9	0.7	*
15	Parkinson's disease (G20-G21)	6	0.4	*
...	All other causes (Residual)	228	16.9	561.1
American Indian, both sexes, 80-84 years				
...	All causes	1,195	100.0	4,414.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	315	26.4	1,163.6
2	Malignant neoplasms (C00-C97)	225	18.8	831.1
3	Chronic lower respiratory diseases (J40-J47)	96	8.0	354.6
4	Diabetes mellitus (E10-E14)	84	7.0	310.3
5	Cerebrovascular diseases (I60-I69)	77	6.4	284.4
6	Alzheimer's disease (G30)	40	3.3	147.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	34	2.8	125.6
8	Influenza and pneumonia (J09-J18)	32	2.7	118.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	30	2.5	110.8
10	Septicemia (A40-A41)	21	1.8	77.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	17	1.4	*
12	Pneumonitis due to solids and liquids (J69)	12	1.0	*
13	Parkinson's disease (G20-G21)	11	0.9	*
14	Aortic aneurysm and dissection (I71)	8	0.7	*
14	Chronic liver disease and cirrhosis (K70,K73-K74)	8	0.7	*
...	All other causes (Residual)	185	15.5	683.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 85-89 years				
...	All causes	958	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	247	25.8	---
2	Malignant neoplasms (C00-C97)	149	15.6	---
3	Cerebrovascular diseases (I60-I69)	84	8.8	---
4	Diabetes mellitus (E10-E14)	59	6.2	---
5	Chronic lower respiratory diseases (J40-J47)	50	5.2	---
6	Alzheimer's disease (G30)	39	4.1	---
7	Influenza and pneumonia (J09-J18)	32	3.3	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	32	3.3	---
9	Parkinson's disease (G20-G21)	17	1.8	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17	1.8	---
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	16	1.7	---
12	Septicemia (A40-A41)	13	1.4	---
12	Pneumonitis due to solids and liquids (J69)	13	1.4	---
14	Chronic liver disease and cirrhosis (K70,K73-K74)	8	0.8	---
15	Aortic aneurysm and dissection (I71)	7	0.7	---
...	All other causes (Residual)	175	18.3	---
American Indian, both sexes, 90-94 years				
...	All causes	540	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	138	25.6	---
2	Malignant neoplasms (C00-C97)	59	10.9	---
3	Cerebrovascular diseases (I60-I69)	45	8.3	---
4	Alzheimer's disease (G30)	37	6.9	---
5	Diabetes mellitus (E10-E14)	28	5.2	---
6	Influenza and pneumonia (J09-J18)	27	5.0	---
7	Chronic lower respiratory diseases (J40-J47)	22	4.1	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18	3.3	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10	1.9	---
10	Septicemia (A40-A41)	8	1.5	---
11	Parkinson's disease (G20-G21)	7	1.3	---
11	Pneumonitis due to solids and liquids (J69)	7	1.3	---
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	6	1.1	---
14	Nutritional deficiencies (E40-E64)	5	0.9	---
15	Atherosclerosis (I70)	4	0.7	---
...	All other causes (Residual)	119	22.0	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 95-99 years				
...	All causes	211	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	63	29.9	---
2	Alzheimer's disease (G30)	18	8.5	---
3	Cerebrovascular diseases (I60-I69)	15	7.1	---
4	Chronic lower respiratory diseases (J40-J47)	13	6.2	---
5	Malignant neoplasms (C00-C97)	11	5.2	---
5	Diabetes mellitus (E10-E14)	11	5.2	---
5	Influenza and pneumonia (J09-J18)	11	5.2	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5	2.4	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4	1.9	---
9	Pneumonitis due to solids and liquids (J69)	4	1.9	---
11	Aortic aneurysm and dissection (I71)	3	1.4	---
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	1.4	---
13	Anemias (D50-D64)	2	0.9	---
13	Atherosclerosis (I70)	2	0.9	---
15	Septicemia (A40-A41)	1	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.5	---
15	Hernia (K40-K46)	1	0.5	---
...	All other causes (Residual)	43	20.4	---
American Indian, both sexes, 100 years and over				
...	All causes	62	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	18	29.0	---
2	Malignant neoplasms (C00-C97)	5	8.1	---
2	Influenza and pneumonia (J09-J18)	5	8.1	---
4	Cerebrovascular diseases (I60-I69)	4	6.5	---
5	Septicemia (A40-A41)	2	3.2	---
5	Nutritional deficiencies (E40-E64)	2	3.2	---
5	Alzheimer's disease (G30)	2	3.2	---
8	Anemias (D50-D64)	1	1.6	---
8	Diabetes mellitus (E10-E14)	1	1.6	---
8	Chronic lower respiratory diseases (J40-J47)	1	1.6	---
8	Pneumonitis due to solids and liquids (J69)	1	1.6	---
8	Hernia (K40-K46)	1	1.6	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	1.6	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1	1.6	---
...	All other causes (Residual)	17	27.4	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, all ages				
...	All causes	7,885	100.0	488.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,520	19.3	94.1
2	Malignant neoplasms (C00-C97)	1,345	17.1	83.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,129	14.3	69.9
4	Chronic liver disease and cirrhosis (K70,K73-K74)	415	5.3	25.7
5	Diabetes mellitus (E10-E14)	381	4.8	23.6
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	310	3.9	19.2
7	Chronic lower respiratory diseases (J40-J47)	299	3.8	18.5
8	Cerebrovascular diseases (I60-I69)	267	3.4	16.5
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	163	2.1	10.1
10	Influenza and pneumonia (J09-J18)	150	1.9	9.3
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	126	1.6	7.8
12	Septicemia (A40-A41)	102	1.3	6.3
13	Certain conditions originating in the perinatal period (P00-P96)	83	1.1	5.1
14	Alzheimer's disease (G30)	69	0.9	4.3
15	Viral hepatitis (B15-B19)	66	0.8	4.1
...	All other causes (Residual)	1,460	18.5	90.4
American Indian, male, 1-4 years				
...	All causes	57	100.0	63.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	24	42.1	26.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	8.8	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	8.8	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	5.3	*
5	Influenza and pneumonia (J09-J18)	2	3.5	*
6	Chronic lower respiratory diseases (J40-J47)	1	1.8	*
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	1.8	*
6	Certain conditions originating in the perinatal period (P00-P96)	1	1.8	*
6	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	1.8	*
...	All other causes (Residual)	14	24.6	*

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 5-9 years				
...	All causes	25	100.0	20.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10	40.0	*
2	Malignant neoplasms (C00-C97)	4	16.0	*
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	4.0	*
3	Chronic lower respiratory diseases (J40-J47)	1	4.0	*
3	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	4.0	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	4.0	*
...	All other causes (Residual)	7	28.0	*
American Indian, male, 10-14 years				
...	All causes	36	100.0	25.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	20	55.6	14.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6	16.7	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	8.3	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	5.6	*
5	Septicemia (A40-A41)	1	2.8	*
...	All other causes (Residual)	4	11.1	*
American Indian, male, 15-19 years				
...	All causes	182	100.0	117.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	91	50.0	58.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	39	21.4	25.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	16	8.8	*
4	Malignant neoplasms (C00-C97)	7	3.8	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	1.6	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	1.6	*
7	Septicemia (A40-A41)	1	0.5	*
7	Anemias (D50-D64)	1	0.5	*
7	Aortic aneurysm and dissection (I71)	1	0.5	*
7	Influenza and pneumonia (J09-J18)	1	0.5	*
7	Chronic lower respiratory diseases (J40-J47)	1	0.5	*
7	Pneumonitis due to solids and liquids (J69)	1	0.5	*
7	Legal intervention (Y35,Y89.0)	1	0.5	*
...	All other causes (Residual)	16	8.8	*

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 20-24 years				
...	All causes	259	100.0	170.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	133	51.4	87.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	60	23.2	39.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	29	11.2	19.1
4	Malignant neoplasms (C00-C97)	9	3.5	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	0.8	*
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2	0.8	*
7	Cerebrovascular diseases (I60-I69)	1	0.4	*
7	Aortic aneurysm and dissection (I71)	1	0.4	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.4	*
...	All other causes (Residual)	21	8.1	13.8
American Indian, male, 25-29 years				
...	All causes	250	100.0	178.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	105	42.0	75.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	43	17.2	30.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	28	11.2	20.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	15	6.0	*
5	Malignant neoplasms (C00-C97)	11	4.4	*
6	Chronic liver disease and cirrhosis (K70,K73-K74)	10	4.0	*
7	Diabetes mellitus (E10-E14)	2	0.8	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	0.8	*
9	Septicemia (A40-A41)	1	0.4	*
9	Viral hepatitis (B15-B19)	1	0.4	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.4	*
9	Cerebrovascular diseases (I60-I69)	1	0.4	*
9	Aortic aneurysm and dissection (I71)	1	0.4	*
9	Influenza and pneumonia (J09-J18)	1	0.4	*
9	Chronic lower respiratory diseases (J40-J47)	1	0.4	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.4	*
9	Legal intervention (Y35,Y89.0)	1	0.4	*
...	All other causes (Residual)	25	10.0	17.9

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 30-34 years				
...	All causes	264	100.0	221.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	109	41.3	91.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	38	14.4	31.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	28	10.6	23.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	24	9.1	20.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	12	4.5	*
6	Malignant neoplasms (C00-C97)	7	2.7	*
7	Septicemia (A40-A41)	3	1.1	*
7	Human immunodeficiency virus (HIV) disease (B20-B24)	3	1.1	*
9	Influenza and pneumonia (J09-J18)	2	0.8	*
10	Salmonella infections (A01-A02)	1	0.4	*
10	Diabetes mellitus (E10-E14)	1	0.4	*
10	Cerebrovascular diseases (I60-I69)	1	0.4	*
10	Aortic aneurysm and dissection (I71)	1	0.4	*
...	All other causes (Residual)	34	12.9	28.5
American Indian, male, 35-39 years				
...	All causes	315	100.0	272.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	106	33.7	91.8
2	Chronic liver disease and cirrhosis (K70,K73-K74)	33	10.5	28.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	28	8.9	24.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	25	7.9	21.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	15	4.8	*
6	Diabetes mellitus (E10-E14)	14	4.4	*
7	Malignant neoplasms (C00-C97)	10	3.2	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	8	2.5	*
9	Cerebrovascular diseases (I60-I69)	5	1.6	*
10	Influenza and pneumonia (J09-J18)	4	1.3	*
11	Septicemia (A40-A41)	2	0.6	*
11	Viral hepatitis (B15-B19)	2	0.6	*
11	Aortic aneurysm and dissection (I71)	2	0.6	*
11	Chronic lower respiratory diseases (J40-J47)	2	0.6	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.6	*
...	All other causes (Residual)	57	18.1	49.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 40-44 years				
...	All causes	453	100.0	392.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	110	24.3	95.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	65	14.3	56.3
3	Chronic liver disease and cirrhosis (K70,K73-K74)	57	12.6	49.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	31	6.8	26.8
5	Malignant neoplasms (C00-C97)	27	6.0	23.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	17	3.8	*
7	Diabetes mellitus (E10-E14)	15	3.3	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	13	2.9	*
9	Viral hepatitis (B15-B19)	10	2.2	*
10	Cerebrovascular diseases (I60-I69)	9	2.0	*
11	Influenza and pneumonia (J09-J18)	8	1.8	*
12	Septicemia (A40-A41)	5	1.1	*
12	Chronic lower respiratory diseases (J40-J47)	5	1.1	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.7	*
15	Aortic aneurysm and dissection (I71)	2	0.4	*
15	Pneumonitis due to solids and liquids (J69)	2	0.4	*
...	All other causes (Residual)	74	16.3	64.1
American Indian, male, 45-49 years				
...	All causes	562	100.0	501.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	104	18.5	92.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	99	17.6	88.2
3	Chronic liver disease and cirrhosis (K70,K73-K74)	74	13.2	66.0
4	Malignant neoplasms (C00-C97)	67	11.9	59.7
5	Diabetes mellitus (E10-E14)	25	4.4	22.3
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	17	3.0	*
7	Human immunodeficiency virus (HIV) disease (B20-B24)	16	2.8	*
8	Cerebrovascular diseases (I60-I69)	15	2.7	*
9	Influenza and pneumonia (J09-J18)	13	2.3	*
10	Septicemia (A40-A41)	11	2.0	*
11	Viral hepatitis (B15-B19)	10	1.8	*
11	Chronic lower respiratory diseases (J40-J47)	10	1.8	*
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	10	1.8	*
14	Aortic aneurysm and dissection (I71)	3	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.5	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.5	*
...	All other causes (Residual)	82	14.6	73.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 50-54 years				
...	All causes	631	100.0	657.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	135	21.4	140.6
2	Malignant neoplasms (C00-C97)	96	15.2	100.0
3	Chronic liver disease and cirrhosis (K70,K73-K74)	79	12.5	82.3
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	73	11.6	76.0
5	Diabetes mellitus (E10-E14)	35	5.5	36.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	20	3.2	20.8
7	Cerebrovascular diseases (I60-I69)	14	2.2	*
7	Chronic lower respiratory diseases (J40-J47)	14	2.2	*
9	Septicemia (A40-A41)	13	2.1	*
10	Viral hepatitis (B15-B19)	10	1.6	*
10	Influenza and pneumonia (J09-J18)	10	1.6	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	1.3	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	6	1.0	*
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	1.0	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4	0.6	*
...	All other causes (Residual)	108	17.1	112.5
American Indian, male, 55-59 years				
...	All causes	700	100.0	906.8
1	Malignant neoplasms (C00-C97)	164	23.4	212.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	163	23.3	211.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	60	8.6	77.7
4	Chronic liver disease and cirrhosis (K70,K73-K74)	59	8.4	76.4
5	Diabetes mellitus (E10-E14)	44	6.3	57.0
6	Cerebrovascular diseases (I60-I69)	27	3.9	35.0
7	Viral hepatitis (B15-B19)	17	2.4	*
7	Chronic lower respiratory diseases (J40-J47)	17	2.4	*
9	Influenza and pneumonia (J09-J18)	11	1.6	*
10	Septicemia (A40-A41)	10	1.4	*
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	9	1.3	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	1.0	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.7	*
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	0.6	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	3	0.4	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3	0.4	*
15	Atherosclerosis (I70)	3	0.4	*
...	All other causes (Residual)	94	13.4	121.8

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 60-64 years				
...	All causes	672	100.0	1,219.3
1	Malignant neoplasms (C00-C97)	186	27.7	337.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	149	22.2	270.3
3	Diabetes mellitus (E10-E14)	48	7.1	87.1
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	35	5.2	63.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	34	5.1	61.7
6	Chronic lower respiratory diseases (J40-J47)	32	4.8	58.1
7	Cerebrovascular diseases (I60-I69)	22	3.3	39.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19	2.8	*
9	Influenza and pneumonia (J09-J18)	16	2.4	*
10	Septicemia (A40-A41)	9	1.3	*
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8	1.2	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7	1.0	*
13	Viral hepatitis (B15-B19)	6	0.9	*
14	Alzheimer's disease (G30)	5	0.7	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.4	*
15	Aortic aneurysm and dissection (I71)	3	0.4	*
15	Peptic ulcer (K25-K28)	3	0.4	*
...	All other causes (Residual)	87	12.9	157.9
American Indian, male, 65-69 years				
...	All causes	644	100.0	1,712.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	178	27.6	473.4
2	Malignant neoplasms (C00-C97)	153	23.8	406.9
3	Diabetes mellitus (E10-E14)	47	7.3	125.0
4	Chronic lower respiratory diseases (J40-J47)	44	6.8	117.0
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	33	5.1	87.8
6	Chronic liver disease and cirrhosis (K70,K73-K74)	24	3.7	63.8
7	Cerebrovascular diseases (I60-I69)	19	3.0	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	2.5	*
9	Influenza and pneumonia (J09-J18)	10	1.6	*
10	Septicemia (A40-A41)	9	1.4	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	6	0.9	*
12	Viral hepatitis (B15-B19)	5	0.8	*
12	Aortic aneurysm and dissection (I71)	5	0.8	*
14	Atherosclerosis (I70)	4	0.6	*
15	Parkinson's disease (G20-G21)	3	0.5	*
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3	0.5	*
...	All other causes (Residual)	85	13.2	226.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 70-74 years				
...	All causes	713	100.0	2,736.3
1	Malignant neoplasms (C00-C97)	207	29.0	794.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	154	21.6	591.0
3	Diabetes mellitus (E10-E14)	57	8.0	218.8
4	Chronic lower respiratory diseases (J40-J47)	48	6.7	184.2
5	Cerebrovascular diseases (I60-I69)	34	4.8	130.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	29	4.1	111.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19	2.7	*
8	Chronic liver disease and cirrhosis (K70,K73-K74)	14	2.0	*
9	Septicemia (A40-A41)	13	1.8	*
10	Influenza and pneumonia (J09-J18)	12	1.7	*
11	Aortic aneurysm and dissection (I71)	7	1.0	*
12	Parkinson's disease (G20-G21)	6	0.8	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.7	*
14	Alzheimer's disease (G30)	4	0.6	*
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4	0.6	*
...	All other causes (Residual)	100	14.0	383.8
American Indian, male, 75-79 years				
...	All causes	671	100.0	3,748.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	176	26.2	983.3
2	Malignant neoplasms (C00-C97)	175	26.1	977.7
3	Cerebrovascular diseases (I60-I69)	44	6.6	245.8
4	Chronic lower respiratory diseases (J40-J47)	41	6.1	229.1
5	Diabetes mellitus (E10-E14)	29	4.3	162.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	26	3.9	145.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	2.1	*
8	Influenza and pneumonia (J09-J18)	13	1.9	*
9	Alzheimer's disease (G30)	12	1.8	*
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10	1.5	*
11	Septicemia (A40-A41)	7	1.0	*
11	Chronic liver disease and cirrhosis (K70,K73-K74)	7	1.0	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.7	*
13	Pneumonitis due to solids and liquids (J69)	5	0.7	*
15	Parkinson's disease (G20-G21)	4	0.6	*
...	All other causes (Residual)	103	15.4	575.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 80-84 years				
...	All causes	543	100.0	4,913.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	151	27.8	1,366.4
2	Malignant neoplasms (C00-C97)	110	20.3	995.4
3	Chronic lower respiratory diseases (J40-J47)	45	8.3	407.2
4	Cerebrovascular diseases (I60-I69)	33	6.1	298.6
5	Diabetes mellitus (E10-E14)	30	5.5	271.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	24	4.4	217.2
7	Influenza and pneumonia (J09-J18)	19	3.5	*
8	Alzheimer's disease (G30)	13	2.4	*
9	Septicemia (A40-A41)	9	1.7	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	1.7	*
11	Parkinson's disease (G20-G21)	7	1.3	*
12	Aortic aneurysm and dissection (I71)	6	1.1	*
13	Pneumonitis due to solids and liquids (J69)	5	0.9	*
13	Chronic liver disease and cirrhosis (K70,K73-K74)	5	0.9	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3	0.6	*
...	All other causes (Residual)	74	13.6	669.6
American Indian, male, 85-89 years				
...	All causes	397	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	106	26.7	---
2	Malignant neoplasms (C00-C97)	77	19.4	---
3	Cerebrovascular diseases (I60-I69)	28	7.1	---
4	Diabetes mellitus (E10-E14)	21	5.3	---
4	Chronic lower respiratory diseases (J40-J47)	21	5.3	---
6	Alzheimer's disease (G30)	19	4.8	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	3.3	---
8	Influenza and pneumonia (J09-J18)	11	2.8	---
9	Parkinson's disease (G20-G21)	8	2.0	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	8	2.0	---
9	Pneumonitis due to solids and liquids (J69)	8	2.0	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8	2.0	---
13	Chronic liver disease and cirrhosis (K70,K73-K74)	3	0.8	---
14	Septicemia (A40-A41)	2	0.5	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.5	---
14	Atherosclerosis (I70)	2	0.5	---
14	Aortic aneurysm and dissection (I71)	2	0.5	---
14	Hyperplasia of prostate (N40)	2	0.5	---
...	All other causes (Residual)	56	14.1	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 90-94 years				
...	All causes	192	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	40	20.8	---
2	Malignant neoplasms (C00-C97)	25	13.0	---
3	Alzheimer's disease (G30)	12	6.3	---
3	Chronic lower respiratory diseases (J40-J47)	12	6.3	---
5	Cerebrovascular diseases (I60-I69)	11	5.7	---
6	Diabetes mellitus (E10-E14)	10	5.2	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	4.7	---
8	Influenza and pneumonia (J09-J18)	6	3.1	---
9	Septicemia (A40-A41)	4	2.1	---
9	Pneumonitis due to solids and liquids (J69)	4	2.1	---
11	Parkinson's disease (G20-G21)	3	1.6	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3	1.6	---
13	Nutritional deficiencies (E40-E64)	2	1.0	---
13	Cholelithiasis and other disorders of gallbladder (K80-K82)	2	1.0	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.5	---
15	Aortic aneurysm and dissection (I71)	1	0.5	---
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.5	---
15	Hyperplasia of prostate (N40)	1	0.5	---
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.5	---
...	All other causes (Residual)	44	22.9	---
American Indian, male, 95-99 years				
...	All causes	55	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	13	23.6	---
2	Malignant neoplasms (C00-C97)	5	9.1	---
2	Influenza and pneumonia (J09-J18)	5	9.1	---
4	Chronic lower respiratory diseases (J40-J47)	4	7.3	---
5	Alzheimer's disease (G30)	3	5.5	---
6	Diabetes mellitus (E10-E14)	2	3.6	---
6	Cerebrovascular diseases (I60-I69)	2	3.6	---
6	Aortic aneurysm and dissection (I71)	2	3.6	---
6	Pneumonitis due to solids and liquids (J69)	2	3.6	---
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2	3.6	---
11	Anemias (D50-D64)	1	1.8	---
11	Atherosclerosis (I70)	1	1.8	---
11	Hernia (K40-K46)	1	1.8	---
...	All other causes (Residual)	12	21.8	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 100 years and over				
...	All causes	23	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7	30.4	---
2	Malignant neoplasms (C00-C97)	4	17.4	---
3	Diabetes mellitus (E10-E14)	1	4.3	---
3	Nutritional deficiencies (E40-E64)	1	4.3	---
3	Cerebrovascular diseases (I60-I69)	1	4.3	---
3	Influenza and pneumonia (J09-J18)	1	4.3	---
3	Pneumonitis due to solids and liquids (J69)	1	4.3	---
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1	4.3	---
...	All other causes (Residual)	6	26.1	---
American Indian, female, all ages				
...	All causes	6,482	100.0	400.0
1	Malignant neoplasms (C00-C97)	1,216	18.8	75.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,128	17.4	69.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	572	8.8	35.3
4	Diabetes mellitus (E10-E14)	409	6.3	25.2
5	Cerebrovascular diseases (I60-I69)	319	4.9	19.7
6	Chronic lower respiratory diseases (J40-J47)	312	4.8	19.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	294	4.5	18.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	166	2.6	10.2
9	Influenza and pneumonia (J09-J18)	130	2.0	8.0
10	Septicemia (A40-A41)	128	2.0	7.9
11	Alzheimer's disease (G30)	122	1.9	7.5
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	82	1.3	5.1
13	Certain conditions originating in the perinatal period (P00-P96)	75	1.2	4.6
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	62	1.0	3.8
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	62	1.0	3.8
...	All other causes (Residual)	1,405	21.7	86.7

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 1-4 years				
...	All causes	40	100.0	46.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19	47.5	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	7.5	*
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	5.0	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	5.0	*
5	Malignant neoplasms (C00-C97)	1	2.5	*
5	Cerebrovascular diseases (I60-I69)	1	2.5	*
5	Acute bronchitis and bronchiolitis (J20-J21)	1	2.5	*
5	Chronic lower respiratory diseases (J40-J47)	1	2.5	*
...	All other causes (Residual)	10	25.0	*
American Indian, female, 5-9 years				
...	All causes	15	100.0	*
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8	53.3	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	20.0	*
3	Malignant neoplasms (C00-C97)	1	6.7	*
...	All other causes (Residual)	3	20.0	*
American Indian, female, 10-14 years				
...	All causes	18	100.0	*
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8	44.4	*
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3	16.7	*
3	Septicemia (A40-A41)	1	5.6	*
3	Malignant neoplasms (C00-C97)	1	5.6	*
3	In situ neoplasms, benign neoplasms and neoplasms of uncertain or	1	5.6	*
3	Chronic lower respiratory diseases (J40-J47)	1	5.6	*
...	All other causes (Residual)	3	16.7	*

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 15-19 years				
...	All causes	83	100.0	54.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	42	50.6	27.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	12	14.5	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	6.0	*
4	Malignant neoplasms (C00-C97)	4	4.8	*
4	Pregnancy, childbirth and the puerperium (O00-O99)	4	4.8	*
6	Septicemia (A40-A41)	1	1.2	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	1.2	*
6	Cerebrovascular diseases (I60-I69)	1	1.2	*
6	Influenza and pneumonia (J09-J18)	1	1.2	*
6	Chronic lower respiratory diseases (J40-J47)	1	1.2	*
6	Pneumonitis due to solids and liquids (J69)	1	1.2	*
6	Diseases of appendix (K35-K38)	1	1.2	*
...	All other causes (Residual)	9	10.8	*
American Indian, female, 20-24 years				
...	All causes	98	100.0	68.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	53	54.1	36.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11	11.2	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7	7.1	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	6	6.1	*
5	Malignant neoplasms (C00-C97)	2	2.0	*
5	Influenza and pneumonia (J09-J18)	2	2.0	*
5	Congenital malformations, deformations and chromosomal abnormalities	2	2.0	*
8	Chronic liver disease and cirrhosis (K70,K73-K74)	1	1.0	*
...	All other causes (Residual)	14	14.3	*

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 25-29 years				
...	All causes	99	100.0	76.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	49	49.5	38.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11	11.1	*
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	6	6.1	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	5.1	*
5	Pregnancy, childbirth and the puerperium (O00-O99)	4	4.0	*
6	Malignant neoplasms (C00-C97)	3	3.0	*
6	Chronic liver disease and cirrhosis (K70,K73-K74)	3	3.0	*
8	Septicemia (A40-A41)	2	2.0	*
9	Cerebrovascular diseases (I60-I69)	1	1.0	*
9	Pneumonitis due to solids and liquids (J69)	1	1.0	*
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	1.0	*
9	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	1.0	*
...	All other causes (Residual)	12	12.1	*
American Indian, female, 30-34 years				
...	All causes	118	100.0	106.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	46	39.0	41.6
2	Chronic liver disease and cirrhosis (K70,K73-K74)	12	10.2	*
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	11	9.3	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8	6.8	*
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6	5.1	*
6	Malignant neoplasms (C00-C97)	4	3.4	*
7	Diabetes mellitus (E10-E14)	3	2.5	*
7	Pregnancy, childbirth and the puerperium (O00-O99)	3	2.5	*
9	Scarlet fever and erysipelas (A38,A46)	1	0.8	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.8	*
9	Influenza and pneumonia (J09-J18)	1	0.8	*
9	Chronic lower respiratory diseases (J40-J47)	1	0.8	*
9	Pneumonitis due to solids and liquids (J69)	1	0.8	*
...	All other causes (Residual)	20	16.9	18.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 35-39 years				
...	All causes	182	100.0	163.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	49	26.9	44.1
2	Chronic liver disease and cirrhosis (K70,K73-K74)	21	11.5	18.9
3	Malignant neoplasms (C00-C97)	17	9.3	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	15	8.2	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	10	5.5	*
6	Human immunodeficiency virus (HIV) disease (B20-B24)	7	3.8	*
6	Diabetes mellitus (E10-E14)	7	3.8	*
8	Cerebrovascular diseases (I60-I69)	4	2.2	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	2.2	*
8	Pregnancy, childbirth and the puerperium (O00-O99)	4	2.2	*
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4	2.2	*
12	Viral hepatitis (B15-B19)	2	1.1	*
12	Influenza and pneumonia (J09-J18)	2	1.1	*
14	Septicemia (A40-A41)	1	0.5	*
14	Anemias (D50-D64)	1	0.5	*
14	Chronic lower respiratory diseases (J40-J47)	1	0.5	*
14	Pneumonitis due to solids and liquids (J69)	1	0.5	*
14	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.5	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.5	*
...	All other causes (Residual)	30	16.5	27.0
American Indian, female, 40-44 years				
...	All causes	263	100.0	226.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	55	20.9	47.4
2	Chronic liver disease and cirrhosis (K70,K73-K74)	34	12.9	29.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	32	12.2	27.6
4	Malignant neoplasms (C00-C97)	29	11.0	25.0
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11	4.2	*
6	Cerebrovascular diseases (I60-I69)	10	3.8	*
7	Diabetes mellitus (E10-E14)	9	3.4	*
8	Viral hepatitis (B15-B19)	5	1.9	*
9	Septicemia (A40-A41)	4	1.5	*
9	Influenza and pneumonia (J09-J18)	4	1.5	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	3	1.1	*
11	Pregnancy, childbirth and the puerperium (O00-O99)	3	1.1	*
13	Chronic lower respiratory diseases (J40-J47)	2	0.8	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.8	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	0.8	*
...	All other causes (Residual)	58	22.1	50.0

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 45-49 years				
...	All causes	375	100.0	318.4
1	Malignant neoplasms (C00-C97)	80	21.3	67.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	67	17.9	56.9
3	Chronic liver disease and cirrhosis (K70,K73-K74)	50	13.3	42.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	35	9.3	29.7
5	Diabetes mellitus (E10-E14)	14	3.7	*
6	Septicemia (A40-A41)	9	2.4	*
6	Cerebrovascular diseases (I60-I69)	9	2.4	*
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	9	2.4	*
9	Chronic lower respiratory diseases (J40-J47)	7	1.9	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	1.9	*
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	1.6	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	5	1.3	*
13	Influenza and pneumonia (J09-J18)	4	1.1	*
14	Viral hepatitis (B15-B19)	3	0.8	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2	0.5	*
...	All other causes (Residual)	68	18.1	57.7
American Indian, female, 50-54 years				
...	All causes	390	100.0	378.4
1	Malignant neoplasms (C00-C97)	87	22.3	84.4
2	Chronic liver disease and cirrhosis (K70,K73-K74)	56	14.4	54.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	46	11.8	44.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	36	9.2	34.9
5	Diabetes mellitus (E10-E14)	18	4.6	*
6	Chronic lower respiratory diseases (J40-J47)	17	4.4	*
7	Septicemia (A40-A41)	13	3.3	*
7	Cerebrovascular diseases (I60-I69)	13	3.3	*
9	Viral hepatitis (B15-B19)	10	2.6	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	5	1.3	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	1.3	*
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4	1.0	*
13	Pneumonitis due to solids and liquids (J69)	2	0.5	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	0.5	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.3	*
15	Alzheimer's disease (G30)	1	0.3	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.3	*
15	Aortic aneurysm and dissection (I71)	1	0.3	*
15	Influenza and pneumonia (J09-J18)	1	0.3	*
15	Peptic ulcer (K25-K28)	1	0.3	*
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.3	*
15	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.3	*
15	Inflammatory diseases of female pelvic organs (N70-N76)	1	0.3	*
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1	0.3	*
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.3	*

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
...	All other causes (Residual)	65	16.7	63.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 55-59 years				
...	All causes	486	100.0	583.5
1	Malignant neoplasms (C00-C97)	127	26.1	152.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	72	14.8	86.4
3	Chronic liver disease and cirrhosis (K70,K73-K74)	41	8.4	49.2
4	Diabetes mellitus (E10-E14)	38	7.8	45.6
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	25	5.1	30.0
6	Chronic lower respiratory diseases (J40-J47)	24	4.9	28.8
7	Cerebrovascular diseases (I60-I69)	23	4.7	27.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	2.9	*
9	Septicemia (A40-A41)	12	2.5	*
10	Viral hepatitis (B15-B19)	11	2.3	*
11	Influenza and pneumonia (J09-J18)	6	1.2	*
12	Nutritional deficiencies (E40-E64)	3	0.6	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3	0.6	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.4	*
14	Anemias (D50-D64)	2	0.4	*
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2	0.4	*
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
...	All other causes (Residual)	77	15.8	92.5
American Indian, female, 60-64 years				
...	All causes	509	100.0	845.7
1	Malignant neoplasms (C00-C97)	146	28.7	242.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	84	16.5	139.6
3	Diabetes mellitus (E10-E14)	37	7.3	61.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	33	6.5	54.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	30	5.9	49.8
6	Chronic lower respiratory diseases (J40-J47)	27	5.3	44.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	3.3	*
8	Septicemia (A40-A41)	13	2.6	*
9	Cerebrovascular diseases (I60-I69)	11	2.2	*
10	Viral hepatitis (B15-B19)	5	1.0	*
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4	0.8	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3	0.6	*
12	Aortic aneurysm and dissection (I71)	3	0.6	*
14	Influenza and pneumonia (J09-J18)	2	0.4	*
14	Pneumonitis due to solids and liquids (J69)	2	0.4	*
...	All other causes (Residual)	92	18.1	152.8

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 65-69 years				
...	All causes	578	100.0	1,375.6
1	Malignant neoplasms (C00-C97)	158	27.3	376.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	106	18.3	252.3
3	Diabetes mellitus (E10-E14)	55	9.5	130.9
4	Chronic lower respiratory diseases (J40-J47)	41	7.1	97.6
5	Cerebrovascular diseases (I60-I69)	25	4.3	59.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19	3.3	*
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	2.9	*
8	Chronic liver disease and cirrhosis (K70,K73-K74)	13	2.2	*
9	Influenza and pneumonia (J09-J18)	10	1.7	*
10	Septicemia (A40-A41)	9	1.6	*
11	Alzheimer's disease (G30)	8	1.4	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7	1.2	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.9	*
13	Pneumonitis due to solids and liquids (J69)	5	0.9	*
15	Viral hepatitis (B15-B19)	2	0.3	*
15	Aortic aneurysm and dissection (I71)	2	0.3	*
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	2	0.3	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	0.3	*
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2	0.3	*
...	All other causes (Residual)	90	15.6	214.2
American Indian, female, 70-74 years				
...	All causes	601	100.0	1,931.0
1	Malignant neoplasms (C00-C97)	167	27.8	536.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	105	17.5	337.4
3	Diabetes mellitus (E10-E14)	49	8.2	157.4
4	Chronic lower respiratory diseases (J40-J47)	40	6.7	128.5
5	Cerebrovascular diseases (I60-I69)	36	6.0	115.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22	3.7	70.7
7	Septicemia (A40-A41)	18	3.0	*
8	Chronic liver disease and cirrhosis (K70,K73-K74)	15	2.5	*
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12	2.0	*
10	Alzheimer's disease (G30)	11	1.8	*
11	Influenza and pneumonia (J09-J18)	7	1.2	*
12	Pneumonitis due to solids and liquids (J69)	5	0.8	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4	0.7	*
14	Parkinson's disease (G20-G21)	3	0.5	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.3	*
15	Anemias (D50-D64)	2	0.3	*
15	Atherosclerosis (I70)	2	0.3	*
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2	0.3	*
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	2	0.3	*
...	All other causes (Residual)	97	16.1	311.7

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 75-79 years				
...	All causes	680	100.0	2,990.9
1	Malignant neoplasms (C00-C97)	161	23.7	708.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	138	20.3	607.0
3	Diabetes mellitus (E10-E14)	60	8.8	263.9
4	Chronic lower respiratory diseases (J40-J47)	48	7.1	211.1
5	Cerebrovascular diseases (I60-I69)	33	4.9	145.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	24	3.5	105.6
7	Influenza and pneumonia (J09-J18)	22	3.2	96.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17	2.5	*
9	Septicemia (A40-A41)	13	1.9	*
9	Alzheimer's disease (G30)	13	1.9	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10	1.5	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.7	*
12	Chronic liver disease and cirrhosis (K70,K73-K74)	5	0.7	*
14	Pneumonitis due to solids and liquids (J69)	4	0.6	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.6	*
...	All other causes (Residual)	123	18.1	541.0
American Indian, female, 80-84 years				
...	All causes	652	100.0	4,069.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	164	25.2	1,023.7
2	Malignant neoplasms (C00-C97)	115	17.6	717.8
3	Diabetes mellitus (E10-E14)	54	8.3	337.1
4	Chronic lower respiratory diseases (J40-J47)	51	7.8	318.3
5	Cerebrovascular diseases (I60-I69)	44	6.7	274.6
6	Alzheimer's disease (G30)	27	4.1	168.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21	3.2	131.1
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	14	2.1	*
9	Influenza and pneumonia (J09-J18)	13	2.0	*
10	Septicemia (A40-A41)	12	1.8	*
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10	1.5	*
12	Pneumonitis due to solids and liquids (J69)	7	1.1	*
13	Parkinson's disease (G20-G21)	4	0.6	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.5	*
14	Nutritional deficiencies (E40-E64)	3	0.5	*
14	Atherosclerosis (I70)	3	0.5	*
14	Chronic liver disease and cirrhosis (K70,K73-K74)	3	0.5	*
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	3	0.5	*
...	All other causes (Residual)	101	15.5	630.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 85-89 years				
...	All causes	561	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	141	25.1	---
2	Malignant neoplasms (C00-C97)	72	12.8	---
3	Cerebrovascular diseases (I60-I69)	56	10.0	---
4	Diabetes mellitus (E10-E14)	38	6.8	---
5	Chronic lower respiratory diseases (J40-J47)	29	5.2	---
6	Influenza and pneumonia (J09-J18)	21	3.7	---
7	Alzheimer's disease (G30)	20	3.6	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19	3.4	---
9	Septicemia (A40-A41)	11	2.0	---
10	Parkinson's disease (G20-G21)	9	1.6	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9	1.6	---
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	8	1.4	---
13	Aortic aneurysm and dissection (I71)	5	0.9	---
13	Pneumonitis due to solids and liquids (J69)	5	0.9	---
13	Chronic liver disease and cirrhosis (K70,K73-K74)	5	0.9	---
...	All other causes (Residual)	113	20.1	---
American Indian, female, 90-94 years				
...	All causes	348	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	98	28.2	---
2	Malignant neoplasms (C00-C97)	34	9.8	---
2	Cerebrovascular diseases (I60-I69)	34	9.8	---
4	Alzheimer's disease (G30)	25	7.2	---
5	Influenza and pneumonia (J09-J18)	21	6.0	---
6	Diabetes mellitus (E10-E14)	18	5.2	---
7	Chronic lower respiratory diseases (J40-J47)	10	2.9	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	2.6	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7	2.0	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	6	1.7	---
11	Septicemia (A40-A41)	4	1.1	---
11	Parkinson's disease (G20-G21)	4	1.1	---
11	Atherosclerosis (I70)	4	1.1	---
14	Nutritional deficiencies (E40-E64)	3	0.9	---
14	Pneumonitis due to solids and liquids (J69)	3	0.9	---
...	All other causes (Residual)	68	19.5	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 95-99 years				
...	All causes	156	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	50	32.1	---
2	Alzheimer's disease (G30)	15	9.6	---
3	Cerebrovascular diseases (I60-I69)	13	8.3	---
4	Diabetes mellitus (E10-E14)	9	5.8	---
4	Chronic lower respiratory diseases (J40-J47)	9	5.8	---
6	Malignant neoplasms (C00-C97)	6	3.8	---
6	Influenza and pneumonia (J09-J18)	6	3.8	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4	2.6	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	1.9	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3	1.9	---
11	Pneumonitis due to solids and liquids (J69)	2	1.3	---
12	Septicemia (A40-A41)	1	0.6	---
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.6	---
12	Anemias (D50-D64)	1	0.6	---
12	Atherosclerosis (I70)	1	0.6	---
12	Aortic aneurysm and dissection (I71)	1	0.6	---
...	All other causes (Residual)	31	19.9	---
American Indian, female, 100 years and over				
...	All causes	39	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	11	28.2	---
2	Influenza and pneumonia (J09-J18)	4	10.3	---
3	Cerebrovascular diseases (I60-I69)	3	7.7	---
4	Septicemia (A40-A41)	2	5.1	---
4	Alzheimer's disease (G30)	2	5.1	---
6	Malignant neoplasms (C00-C97)	1	2.6	---
6	Anemias (D50-D64)	1	2.6	---
6	Nutritional deficiencies (E40-E64)	1	2.6	---
6	Chronic lower respiratory diseases (J40-J47)	1	2.6	---
6	Hernia (K40-K46)	1	2.6	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	2.6	---
...	All other causes (Residual)	11	28.2	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, all ages				
...	All causes	45,609	100.0	308.7
1	Malignant neoplasms (C00-C97)	12,326	27.0	83.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,574	23.2	71.6
3	Cerebrovascular diseases (I60-I69)	3,586	7.9	24.3
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,194	4.8	14.9
5	Diabetes mellitus (E10-E14)	1,743	3.8	11.8
6	Influenza and pneumonia (J09-J18)	1,335	2.9	9.0
7	Chronic lower respiratory diseases (J40-J47)	1,331	2.9	9.0
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	900	2.0	6.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	893	2.0	6.0
10	Alzheimer's disease (G30)	748	1.6	5.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	634	1.4	4.3
12	Septicemia (A40-A41)	551	1.2	3.7
13	Certain conditions originating in the perinatal period (P00-P96)	499	1.1	3.4
14	Chronic liver disease and cirrhosis (K70,K73-K74)	408	0.9	2.8
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	357	0.8	2.4
...	All other causes (Residual)	7,530	16.5	51.0
Asian or Pacific Islander, both sexes, 1-4 years				
...	All causes	182	100.0	21.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	51	28.0	6.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	23	12.6	2.7
3	Malignant neoplasms (C00-C97)	21	11.5	2.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	11	6.0	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	9	4.9	*
6	Influenza and pneumonia (J09-J18)	5	2.7	*
7	Chronic lower respiratory diseases (J40-J47)	3	1.6	*
8	Septicemia (A40-A41)	2	1.1	*
8	Certain conditions originating in the perinatal period (P00-P96)	2	1.1	*
10	Salmonella infections (A01-A02)	1	0.5	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.5	*
10	Meningitis (G00,G03)	1	0.5	*
10	Cerebrovascular diseases (I60-I69)	1	0.5	*
10	Acute bronchitis and bronchiolitis (J20-J21)	1	0.5	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.5	*
...	All other causes (Residual)	49	26.9	5.8

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 5-9 years				
...	All causes	93	100.0	9.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	25	26.9	2.6
2	Malignant neoplasms (C00-C97)	18	19.4	*
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	7	7.5	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	6.5	*
5	Influenza and pneumonia (J09-J18)	5	5.4	*
6	Cerebrovascular diseases (I60-I69)	4	4.3	*
7	Septicemia (A40-A41)	2	2.2	*
7	Chronic lower respiratory diseases (J40-J47)	2	2.2	*
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	2.2	*
7	Complications of medical and surgical care (Y40-Y84,Y88)	2	2.2	*
11	Meningococcal infection (A39)	1	1.1	*
11	Meningitis (G00,G03)	1	1.1	*
11	Hernia (K40-K46)	1	1.1	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	1.1	*
...	All other causes (Residual)	16	17.2	*
Asian or Pacific Islander, both sexes, 10-14 years				
...	All causes	117	100.0	12.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	39	33.3	4.1
2	Malignant neoplasms (C00-C97)	16	13.7	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	11	9.4	*
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8	6.8	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	4.3	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	3.4	*
7	Cerebrovascular diseases (I60-I69)	3	2.6	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	1.7	*
9	Anemias (D50-D64)	1	0.9	*
9	Pneumonitis due to solids and liquids (J69)	1	0.9	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.9	*
...	All other causes (Residual)	26	22.2	2.7

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 15-19 years				
...	All causes	313	100.0	32.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	139	44.4	14.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	52	16.6	5.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	32	10.2	3.3
4	Malignant neoplasms (C00-C97)	30	9.6	3.1
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	2.9	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	1.3	*
7	Septicemia (A40-A41)	2	0.6	*
7	Meningitis (G00,G03)	2	0.6	*
7	Cerebrovascular diseases (I60-I69)	2	0.6	*
7	Influenza and pneumonia (J09-J18)	2	0.6	*
7	Chronic lower respiratory diseases (J40-J47)	2	0.6	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.3	*
12	Pneumonitis due to solids and liquids (J69)	1	0.3	*
12	Diseases of appendix (K35-K38)	1	0.3	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
12	Pregnancy, childbirth and the puerperium (O00-O99)	1	0.3	*
12	Legal intervention (Y35,Y89.0)	1	0.3	*
...	All other causes (Residual)	31	9.9	3.2
Asian or Pacific Islander, both sexes, 20-24 years				
...	All causes	530	100.0	53.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	209	39.4	21.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	118	22.3	11.8
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	56	10.6	5.6
4	Malignant neoplasms (C00-C97)	44	8.3	4.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	14	2.6	*
6	Cerebrovascular diseases (I60-I69)	7	1.3	*
7	Influenza and pneumonia (J09-J18)	5	0.9	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	0.9	*
9	Septicemia (A40-A41)	3	0.6	*
9	Diabetes mellitus (E10-E14)	3	0.6	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.4	*
11	Aortic aneurysm and dissection (I71)	2	0.4	*
11	Pneumonitis due to solids and liquids (J69)	2	0.4	*
14	Tuberculosis (A16-A19)	1	0.2	*
14	Viral hepatitis (B15-B19)	1	0.2	*
14	Meningitis (G00,G03)	1	0.2	*
14	Chronic lower respiratory diseases (J40-J47)	1	0.2	*
14	Inflammatory diseases of female pelvic organs (N70-N76)	1	0.2	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.2	*
...	All other causes (Residual)	53	10.0	5.3

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 25-29 years				
...	All causes	459	100.0	38.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	136	29.6	11.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	92	20.0	7.7
3	Malignant neoplasms (C00-C97)	53	11.5	4.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	44	9.6	3.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	42	9.2	3.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	2.0	*
7	Pregnancy, childbirth and the puerperium (O00-O99)	7	1.5	*
8	Cerebrovascular diseases (I60-I69)	6	1.3	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	5	1.1	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.9	*
11	Chronic lower respiratory diseases (J40-J47)	3	0.7	*
12	Tuberculosis (A16-A19)	2	0.4	*
12	Septicemia (A40-A41)	2	0.4	*
12	Diabetes mellitus (E10-E14)	2	0.4	*
12	Aortic aneurysm and dissection (I71)	2	0.4	*
12	Chronic liver disease and cirrhosis (K70,K73-K74)	2	0.4	*
...	All other causes (Residual)	48	10.5	4.0
Asian or Pacific Islander, both sexes, 30-34 years				
...	All causes	539	100.0	39.6
1	Malignant neoplasms (C00-C97)	107	19.9	7.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	107	19.9	7.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	83	15.4	6.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	49	9.1	3.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	41	7.6	3.0
6	Cerebrovascular diseases (I60-I69)	16	3.0	*
7	Pregnancy, childbirth and the puerperium (O00-O99)	11	2.0	*
8	Diabetes mellitus (E10-E14)	10	1.9	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	9	1.7	*
10	Influenza and pneumonia (J09-J18)	7	1.3	*
10	Chronic liver disease and cirrhosis (K70,K73-K74)	7	1.3	*
12	Septicemia (A40-A41)	5	0.9	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.9	*
14	Aortic aneurysm and dissection (I71)	4	0.7	*
15	Chronic lower respiratory diseases (J40-J47)	3	0.6	*
...	All other causes (Residual)	75	13.9	5.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 35-39 years				
...	All causes	749	100.0	54.8
1	Malignant neoplasms (C00-C97)	191	25.5	14.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	115	15.4	8.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	107	14.3	7.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	100	13.4	7.3
5	Cerebrovascular diseases (I60-I69)	38	5.1	2.8
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	33	4.4	2.4
7	Diabetes mellitus (E10-E14)	15	2.0	*
8	Pregnancy, childbirth and the puerperium (O00-O99)	11	1.5	*
9	Chronic liver disease and cirrhosis (K70,K73-K74)	10	1.3	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	9	1.2	*
11	Viral hepatitis (B15-B19)	6	0.8	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.7	*
12	Influenza and pneumonia (J09-J18)	5	0.7	*
12	Chronic lower respiratory diseases (J40-J47)	5	0.7	*
15	Aortic aneurysm and dissection (I71)	3	0.4	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.4	*
...	All other causes (Residual)	93	12.4	6.8
Asian or Pacific Islander, both sexes, 40-44 years				
...	All causes	1,083	100.0	90.6
1	Malignant neoplasms (C00-C97)	360	33.2	30.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	197	18.2	16.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	127	11.7	10.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	91	8.4	7.6
5	Cerebrovascular diseases (I60-I69)	56	5.2	4.7
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	33	3.0	2.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	28	2.6	2.3
8	Diabetes mellitus (E10-E14)	18	1.7	*
9	Septicemia (A40-A41)	14	1.3	*
10	Viral hepatitis (B15-B19)	12	1.1	*
11	Chronic lower respiratory diseases (J40-J47)	10	0.9	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	8	0.7	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.6	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.6	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	0.6	*
...	All other causes (Residual)	110	10.2	9.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 45-49 years				
...	All causes	1,557	100.0	144.0
1	Malignant neoplasms (C00-C97)	582	37.4	53.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	246	15.8	22.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	123	7.9	11.4
4	Cerebrovascular diseases (I60-I69)	115	7.4	10.6
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	72	4.6	6.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	44	2.8	4.1
7	Diabetes mellitus (E10-E14)	34	2.2	3.1
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	25	1.6	2.3
9	Viral hepatitis (B15-B19)	22	1.4	2.0
10	Septicemia (A40-A41)	17	1.1	*
10	Influenza and pneumonia (J09-J18)	17	1.1	*
10	Chronic lower respiratory diseases (J40-J47)	17	1.1	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	1.0	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	14	0.9	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	12	0.8	*
...	All other causes (Residual)	202	13.0	18.7
Asian or Pacific Islander, both sexes, 50-54 years				
...	All causes	2,120	100.0	220.9
1	Malignant neoplasms (C00-C97)	880	41.5	91.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	381	18.0	39.7
3	Cerebrovascular diseases (I60-I69)	134	6.3	14.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	128	6.0	13.3
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	76	3.6	7.9
6	Diabetes mellitus (E10-E14)	71	3.3	7.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	49	2.3	5.1
8	Viral hepatitis (B15-B19)	33	1.6	3.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	26	1.2	2.7
10	Influenza and pneumonia (J09-J18)	22	1.0	2.3
11	Chronic lower respiratory diseases (J40-J47)	21	1.0	2.2
12	Septicemia (A40-A41)	19	0.9	*
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	19	0.9	*
14	Aortic aneurysm and dissection (I71)	16	0.8	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	14	0.7	*
...	All other causes (Residual)	231	10.9	24.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 55-59 years				
...	All causes	2,798	100.0	347.0
1	Malignant neoplasms (C00-C97)	1,193	42.6	147.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	535	19.1	66.3
3	Cerebrovascular diseases (I60-I69)	197	7.0	24.4
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	129	4.6	16.0
5	Diabetes mellitus (E10-E14)	127	4.5	15.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	59	2.1	7.3
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	48	1.7	6.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	45	1.6	5.6
9	Chronic lower respiratory diseases (J40-J47)	38	1.4	4.7
10	Viral hepatitis (B15-B19)	37	1.3	4.6
11	Septicemia (A40-A41)	29	1.0	3.6
12	Aortic aneurysm and dissection (I71)	21	0.8	2.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	20	0.7	2.5
14	Influenza and pneumonia (J09-J18)	19	0.7	*
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	18	0.6	*
...	All other causes (Residual)	283	10.1	35.1
Asian or Pacific Islander, both sexes, 60-64 years				
...	All causes	2,976	100.0	520.1
1	Malignant neoplasms (C00-C97)	1,220	41.0	213.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	650	21.8	113.6
3	Cerebrovascular diseases (I60-I69)	191	6.4	33.4
4	Diabetes mellitus (E10-E14)	146	4.9	25.5
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	91	3.1	15.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	55	1.8	9.6
7	Chronic lower respiratory diseases (J40-J47)	49	1.6	8.6
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	48	1.6	8.4
9	Influenza and pneumonia (J09-J18)	42	1.4	7.3
10	Viral hepatitis (B15-B19)	41	1.4	7.2
11	Septicemia (A40-A41)	34	1.1	5.9
12	Chronic liver disease and cirrhosis (K70,K73-K74)	32	1.1	5.6
13	Aortic aneurysm and dissection (I71)	31	1.0	5.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	29	1.0	5.1
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	23	0.8	4.0
...	All other causes (Residual)	294	9.9	51.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 65-69 years				
...	All causes	3,551	100.0	826.6
1	Malignant neoplasms (C00-C97)	1,352	38.1	314.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	812	22.9	189.0
3	Cerebrovascular diseases (I60-I69)	255	7.2	59.4
4	Diabetes mellitus (E10-E14)	182	5.1	42.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	103	2.9	24.0
6	Chronic lower respiratory diseases (J40-J47)	75	2.1	17.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	75	2.1	17.5
8	Influenza and pneumonia (J09-J18)	58	1.6	13.5
9	Septicemia (A40-A41)	45	1.3	10.5
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	38	1.1	8.8
10	Chronic liver disease and cirrhosis (K70,K73-K74)	38	1.1	8.8
12	Viral hepatitis (B15-B19)	34	1.0	7.9
12	Aortic aneurysm and dissection (I71)	34	1.0	7.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	27	0.8	6.3
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	25	0.7	5.8
...	All other causes (Residual)	398	11.2	92.6
Asian or Pacific Islander, both sexes, 70-74 years				
...	All causes	4,390	100.0	1,333.8
1	Malignant neoplasms (C00-C97)	1,548	35.3	470.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	992	22.6	301.4
3	Cerebrovascular diseases (I60-I69)	358	8.2	108.8
4	Diabetes mellitus (E10-E14)	209	4.8	63.5
5	Chronic lower respiratory diseases (J40-J47)	130	3.0	39.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	117	2.7	35.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	107	2.4	32.5
8	Influenza and pneumonia (J09-J18)	99	2.3	30.1
9	Septicemia (A40-A41)	79	1.8	24.0
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	55	1.3	16.7
11	Viral hepatitis (B15-B19)	37	0.8	11.2
12	Chronic liver disease and cirrhosis (K70,K73-K74)	35	0.8	10.6
13	Parkinson's disease (G20-G21)	34	0.8	10.3
13	Aortic aneurysm and dissection (I71)	34	0.8	10.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	25	0.6	7.6
...	All other causes (Residual)	531	12.1	161.3

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 75-79 years				
...	All causes	5,451	100.0	2,207.2
1	Malignant neoplasms (C00-C97)	1,643	30.1	665.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,301	23.9	526.8
3	Cerebrovascular diseases (I60-I69)	474	8.7	191.9
4	Chronic lower respiratory diseases (J40-J47)	241	4.4	97.6
5	Diabetes mellitus (E10-E14)	240	4.4	97.2
6	Influenza and pneumonia (J09-J18)	167	3.1	67.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	147	2.7	59.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	128	2.3	51.8
9	Septicemia (A40-A41)	78	1.4	31.6
10	Alzheimer's disease (G30)	69	1.3	27.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	66	1.2	26.7
12	Parkinson's disease (G20-G21)	56	1.0	22.7
13	Chronic liver disease and cirrhosis (K70,K73-K74)	44	0.8	17.8
14	Aortic aneurysm and dissection (I71)	42	0.8	17.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	39	0.7	15.8
...	All other causes (Residual)	716	13.1	289.9
Asian or Pacific Islander, both sexes, 80-84 years				
...	All causes	6,577	100.0	3,933.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,727	26.3	1,032.9
2	Malignant neoplasms (C00-C97)	1,542	23.4	922.3
3	Cerebrovascular diseases (I60-I69)	626	9.5	374.4
4	Diabetes mellitus (E10-E14)	297	4.5	177.6
5	Chronic lower respiratory diseases (J40-J47)	289	4.4	172.8
6	Influenza and pneumonia (J09-J18)	239	3.6	142.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	174	2.6	104.1
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	164	2.5	98.1
9	Alzheimer's disease (G30)	133	2.0	79.5
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	131	2.0	78.4
11	Parkinson's disease (G20-G21)	97	1.5	58.0
12	Septicemia (A40-A41)	81	1.2	48.4
13	Aortic aneurysm and dissection (I71)	62	0.9	37.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	52	0.8	31.1
15	Pneumonitis due to solids and liquids (J69)	51	0.8	30.5
...	All other causes (Residual)	912	13.9	545.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 85-89 years				
...	All causes	5,789	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,708	29.5	---
2	Malignant neoplasms (C00-C97)	995	17.2	---
3	Cerebrovascular diseases (I60-I69)	605	10.5	---
4	Influenza and pneumonia (J09-J18)	261	4.5	---
5	Chronic lower respiratory diseases (J40-J47)	256	4.4	---
6	Diabetes mellitus (E10-E14)	234	4.0	---
7	Alzheimer's disease (G30)	218	3.8	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	126	2.2	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	121	2.1	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	112	1.9	---
11	Parkinson's disease (G20-G21)	82	1.4	---
12	Septicemia (A40-A41)	77	1.3	---
13	Pneumonitis due to solids and liquids (J69)	62	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	47	0.8	---
15	Aortic aneurysm and dissection (I71)	36	0.6	---
...	All other causes (Residual)	849	14.7	---
Asian or Pacific Islander, both sexes, 90-94 years				
...	All causes	3,710	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,187	32.0	---
2	Malignant neoplasms (C00-C97)	424	11.4	---
3	Cerebrovascular diseases (I60-I69)	357	9.6	---
4	Influenza and pneumonia (J09-J18)	236	6.4	---
5	Alzheimer's disease (G30)	177	4.8	---
6	Chronic lower respiratory diseases (J40-J47)	133	3.6	---
7	Diabetes mellitus (E10-E14)	118	3.2	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	100	2.7	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	96	2.6	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	63	1.7	---
11	Pneumonitis due to solids and liquids (J69)	48	1.3	---
12	Septicemia (A40-A41)	42	1.1	---
13	Parkinson's disease (G20-G21)	33	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	0.5	---
14	Atherosclerosis (I70)	20	0.5	---
...	All other causes (Residual)	656	17.7	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 95-99 years				
...	All causes	1,339	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	462	34.5	---
2	Cerebrovascular diseases (I60-I69)	121	9.0	---
3	Influenza and pneumonia (J09-J18)	101	7.5	---
4	Malignant neoplasms (C00-C97)	91	6.8	---
4	Alzheimer's disease (G30)	91	6.8	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	49	3.7	---
7	Chronic lower respiratory diseases (J40-J47)	42	3.1	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	30	2.2	---
9	Diabetes mellitus (E10-E14)	28	2.1	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18	1.3	---
11	Pneumonitis due to solids and liquids (J69)	16	1.2	---
12	Septicemia (A40-A41)	10	0.7	---
12	Anemias (D50-D64)	10	0.7	---
14	Parkinson's disease (G20-G21)	9	0.7	---
15	Atherosclerosis (I70)	6	0.4	---
...	All other causes (Residual)	255	19.0	---
Asian or Pacific Islander, both sexes, 100 years and over				
...	All causes	323	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	114	35.3	---
2	Influenza and pneumonia (J09-J18)	34	10.5	---
3	Cerebrovascular diseases (I60-I69)	18	5.6	---
4	Malignant neoplasms (C00-C97)	14	4.3	---
4	Alzheimer's disease (G30)	14	4.3	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	3.1	---
7	Diabetes mellitus (E10-E14)	9	2.8	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9	2.8	---
7	Chronic lower respiratory diseases (J40-J47)	9	2.8	---
10	Pneumonitis due to solids and liquids (J69)	6	1.9	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6	1.9	---
12	Atherosclerosis (I70)	3	0.9	---
13	Septicemia (A40-A41)	2	0.6	---
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.6	---
13	Nutritional deficiencies (E40-E64)	2	0.6	---
13	Parkinson's disease (G20-G21)	2	0.6	---
...	All other causes (Residual)	69	21.4	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, all ages				
...	All causes	23,823	100.0	331.4
1	Malignant neoplasms (C00-C97)	6,394	26.8	89.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,724	24.0	79.6
3	Cerebrovascular diseases (I60-I69)	1,581	6.6	22.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,371	5.8	19.1
5	Diabetes mellitus (E10-E14)	860	3.6	12.0
6	Chronic lower respiratory diseases (J40-J47)	795	3.3	11.1
7	Influenza and pneumonia (J09-J18)	703	3.0	9.8
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	628	2.6	8.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	476	2.0	6.6
10	Septicemia (A40-A41)	302	1.3	4.2
11	Certain conditions originating in the perinatal period (P00-P96)	290	1.2	4.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	286	1.2	4.0
13	Chronic liver disease and cirrhosis (K70,K73-K74)	253	1.1	3.5
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	250	1.0	3.5
15	Alzheimer's disease (G30)	245	1.0	3.4
...	All other causes (Residual)	3,665	15.4	51.0
Asian or Pacific Islander, male, 1-4 years				
...	All causes	109	100.0	25.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	36	33.0	8.3
2	Malignant neoplasms (C00-C97)	12	11.0	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	12	11.0	*
4	Influenza and pneumonia (J09-J18)	5	4.6	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	4.6	*
6	Chronic lower respiratory diseases (J40-J47)	3	2.8	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	1.8	*
7	Certain conditions originating in the perinatal period (P00-P96)	2	1.8	*
9	Salmonella infections (A01-A02)	1	0.9	*
9	Acute bronchitis and bronchiolitis (J20-J21)	1	0.9	*
...	All other causes (Residual)	30	27.5	7.0

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 5-9 years				
...	All causes	51	100.0	10.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14	27.5	*
2	Malignant neoplasms (C00-C97)	12	23.5	*
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	7.8	*
4	Influenza and pneumonia (J09-J18)	3	5.9	*
5	Chronic lower respiratory diseases (J40-J47)	2	3.9	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	3.9	*
7	Meningococcal infection (A39)	1	2.0	*
7	Septicemia (A40-A41)	1	2.0	*
7	Meningitis (G00,G03)	1	2.0	*
7	Cerebrovascular diseases (I60-I69)	1	2.0	*
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	2.0	*
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1	2.0	*
...	All other causes (Residual)	8	15.7	*
Asian or Pacific Islander, male, 10-14 years				
...	All causes	67	100.0	13.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	23	34.3	4.7
2	Malignant neoplasms (C00-C97)	7	10.4	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	10.4	*
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4	6.0	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	6.0	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	4.5	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.5	*
7	Cerebrovascular diseases (I60-I69)	1	1.5	*
7	Pneumonitis due to solids and liquids (J69)	1	1.5	*
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	1.5	*
...	All other causes (Residual)	15	22.4	*

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 15-19 years				
...	All causes	223	100.0	45.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	101	45.3	20.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	43	19.3	8.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	27	12.1	5.5
4	Malignant neoplasms (C00-C97)	17	7.6	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	2.7	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	1.3	*
7	Chronic lower respiratory diseases (J40-J47)	2	0.9	*
8	Septicemia (A40-A41)	1	0.4	*
8	Meningitis (G00,G03)	1	0.4	*
8	Cerebrovascular diseases (I60-I69)	1	0.4	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.4	*
8	Legal intervention (Y35,Y89.0)	1	0.4	*
...	All other causes (Residual)	19	8.5	*
Asian or Pacific Islander, male, 20-24 years				
...	All causes	388	100.0	76.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	150	38.7	29.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	91	23.5	17.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	47	12.1	9.2
4	Malignant neoplasms (C00-C97)	28	7.2	5.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	12	3.1	*
6	Cerebrovascular diseases (I60-I69)	5	1.3	*
7	Influenza and pneumonia (J09-J18)	4	1.0	*
8	Diabetes mellitus (E10-E14)	3	0.8	*
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.8	*
10	Septicemia (A40-A41)	2	0.5	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or	2	0.5	*
10	Pneumonitis due to solids and liquids (J69)	2	0.5	*
13	Tuberculosis (A16-A19)	1	0.3	*
13	Viral hepatitis (B15-B19)	1	0.3	*
13	Aortic aneurysm and dissection (I71)	1	0.3	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.3	*
13	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.3	*
...	All other causes (Residual)	34	8.8	6.7

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 25-29 years				
...	All causes	300	100.0	50.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	101	33.7	17.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	65	21.7	10.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	34	11.3	5.7
4	Malignant neoplasms (C00-C97)	31	10.3	5.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	29	9.7	4.9
6	Human immunodeficiency virus (HIV) disease (B20-B24)	3	1.0	*
6	Chronic lower respiratory diseases (J40-J47)	3	1.0	*
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	1.0	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	1.0	*
10	Septicemia (A40-A41)	2	0.7	*
11	Tuberculosis (A16-A19)	1	0.3	*
11	Cerebrovascular diseases (I60-I69)	1	0.3	*
11	Aortic aneurysm and dissection (I71)	1	0.3	*
11	Influenza and pneumonia (J09-J18)	1	0.3	*
11	Legal intervention (Y35,Y89.0)	1	0.3	*
...	All other causes (Residual)	21	7.0	3.5
Asian or Pacific Islander, male, 30-34 years				
...	All causes	332	100.0	49.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	76	22.9	11.4
2	Malignant neoplasms (C00-C97)	51	15.4	7.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	50	15.1	7.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	35	10.5	5.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	30	9.0	4.5
6	Cerebrovascular diseases (I60-I69)	11	3.3	*
7	Human immunodeficiency virus (HIV) disease (B20-B24)	7	2.1	*
8	Diabetes mellitus (E10-E14)	6	1.8	*
8	Chronic liver disease and cirrhosis (K70,K73-K74)	6	1.8	*
10	Influenza and pneumonia (J09-J18)	5	1.5	*
11	Septicemia (A40-A41)	3	0.9	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.9	*
11	Aortic aneurysm and dissection (I71)	3	0.9	*
14	Anemias (D50-D64)	2	0.6	*
14	Chronic lower respiratory diseases (J40-J47)	2	0.6	*
...	All other causes (Residual)	42	12.7	6.3

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 35-39 years				
...	All causes	473	100.0	70.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	86	18.2	12.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	84	17.8	12.6
3	Malignant neoplasms (C00-C97)	82	17.3	12.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	71	15.0	10.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	25	5.3	3.7
6	Cerebrovascular diseases (I60-I69)	19	4.0	*
7	Diabetes mellitus (E10-E14)	12	2.5	*
8	Chronic liver disease and cirrhosis (K70,K73-K74)	9	1.9	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	8	1.7	*
10	Viral hepatitis (B15-B19)	6	1.3	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.8	*
12	Aortic aneurysm and dissection (I71)	3	0.6	*
12	Chronic lower respiratory diseases (J40-J47)	3	0.6	*
14	Septicemia (A40-A41)	2	0.4	*
14	Influenza and pneumonia (J09-J18)	2	0.4	*
...	All other causes (Residual)	57	12.1	8.5
Asian or Pacific Islander, male, 40-44 years				
...	All causes	637	100.0	109.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	162	25.4	27.9
2	Malignant neoplasms (C00-C97)	153	24.0	26.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	83	13.0	14.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	60	9.4	10.3
5	Cerebrovascular diseases (I60-I69)	29	4.6	5.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	22	3.5	3.8
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	15	2.4	*
8	Diabetes mellitus (E10-E14)	12	1.9	*
9	Septicemia (A40-A41)	8	1.3	*
10	Viral hepatitis (B15-B19)	7	1.1	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	5	0.8	*
11	Aortic aneurysm and dissection (I71)	5	0.8	*
11	Chronic lower respiratory diseases (J40-J47)	5	0.8	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.6	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.6	*
...	All other causes (Residual)	63	9.9	10.8

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 45-49 years				
...	All causes	925	100.0	179.1
1	Malignant neoplasms (C00-C97)	281	30.4	54.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	194	21.0	37.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	77	8.3	14.9
4	Cerebrovascular diseases (I60-I69)	66	7.1	12.8
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	49	5.3	9.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	31	3.4	6.0
7	Viral hepatitis (B15-B19)	19	2.1	*
7	Diabetes mellitus (E10-E14)	19	2.1	*
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	17	1.8	*
10	Septicemia (A40-A41)	11	1.2	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	11	1.2	*
12	Chronic lower respiratory diseases (J40-J47)	10	1.1	*
13	Influenza and pneumonia (J09-J18)	9	1.0	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7	0.8	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.8	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	0.8	*
...	All other causes (Residual)	110	11.9	21.3
Asian or Pacific Islander, male, 50-54 years				
...	All causes	1,287	100.0	286.6
1	Malignant neoplasms (C00-C97)	428	33.3	95.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	303	23.5	67.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	89	6.9	19.8
4	Cerebrovascular diseases (I60-I69)	76	5.9	16.9
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	57	4.4	12.7
6	Diabetes mellitus (E10-E14)	48	3.7	10.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	37	2.9	8.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19	1.5	*
9	Viral hepatitis (B15-B19)	18	1.4	*
10	Influenza and pneumonia (J09-J18)	13	1.0	*
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	13	1.0	*
12	Septicemia (A40-A41)	12	0.9	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	10	0.8	*
13	Aortic aneurysm and dissection (I71)	10	0.8	*
13	Chronic lower respiratory diseases (J40-J47)	10	0.8	*
...	All other causes (Residual)	144	11.2	32.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 55-59 years				
...	All causes	1,650	100.0	443.4
1	Malignant neoplasms (C00-C97)	605	36.7	162.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	403	24.4	108.3
3	Cerebrovascular diseases (I60-I69)	100	6.1	26.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	80	4.8	21.5
5	Diabetes mellitus (E10-E14)	79	4.8	21.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	48	2.9	12.9
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	33	2.0	8.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	31	1.9	8.3
9	Viral hepatitis (B15-B19)	25	1.5	6.7
9	Chronic lower respiratory diseases (J40-J47)	25	1.5	6.7
11	Septicemia (A40-A41)	20	1.2	5.4
12	Aortic aneurysm and dissection (I71)	12	0.7	*
12	Influenza and pneumonia (J09-J18)	12	0.7	*
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	12	0.7	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	11	0.7	*
...	All other causes (Residual)	154	9.3	41.4
Asian or Pacific Islander, male, 60-64 years				
...	All causes	1,684	100.0	634.8
1	Malignant neoplasms (C00-C97)	606	36.0	228.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	435	25.8	164.0
3	Cerebrovascular diseases (I60-I69)	101	6.0	38.1
4	Diabetes mellitus (E10-E14)	89	5.3	33.5
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	47	2.8	17.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	36	2.1	13.6
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	32	1.9	12.1
8	Chronic lower respiratory diseases (J40-J47)	31	1.8	11.7
9	Influenza and pneumonia (J09-J18)	24	1.4	9.0
10	Viral hepatitis (B15-B19)	22	1.3	8.3
10	Chronic liver disease and cirrhosis (K70,K73-K74)	22	1.3	8.3
12	Septicemia (A40-A41)	20	1.2	7.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	19	1.1	*
13	Aortic aneurysm and dissection (I71)	19	1.1	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	0.8	*
...	All other causes (Residual)	168	10.0	63.3

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 65-69 years				
...	All causes	2,062	100.0	1,035.6
1	Malignant neoplasms (C00-C97)	763	37.0	383.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	511	24.8	256.6
3	Cerebrovascular diseases (I60-I69)	135	6.5	67.8
4	Diabetes mellitus (E10-E14)	101	4.9	50.7
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	62	3.0	31.1
6	Chronic lower respiratory diseases (J40-J47)	49	2.4	24.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	44	2.1	22.1
8	Influenza and pneumonia (J09-J18)	39	1.9	19.6
9	Septicemia (A40-A41)	30	1.5	15.1
10	Aortic aneurysm and dissection (I71)	23	1.1	11.6
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	21	1.0	10.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	21	1.0	10.5
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	21	1.0	10.5
14	Chronic liver disease and cirrhosis (K70,K73-K74)	20	1.0	10.0
15	Viral hepatitis (B15-B19)	14	0.7	*
...	All other causes (Residual)	208	10.1	104.5
Asian or Pacific Islander, male, 70-74 years				
...	All causes	2,425	100.0	1,674.5
1	Malignant neoplasms (C00-C97)	856	35.3	591.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	594	24.5	410.2
3	Cerebrovascular diseases (I60-I69)	177	7.3	122.2
4	Diabetes mellitus (E10-E14)	104	4.3	71.8
5	Chronic lower respiratory diseases (J40-J47)	78	3.2	53.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	67	2.8	46.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	63	2.6	43.5
8	Influenza and pneumonia (J09-J18)	52	2.1	35.9
9	Septicemia (A40-A41)	51	2.1	35.2
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	29	1.2	20.0
11	Parkinson's disease (G20-G21)	21	0.9	14.5
12	Aortic aneurysm and dissection (I71)	20	0.8	13.8
13	Chronic liver disease and cirrhosis (K70,K73-K74)	15	0.6	*
14	Viral hepatitis (B15-B19)	14	0.6	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	0.5	*
...	All other causes (Residual)	271	11.2	187.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 75-79 years				
...	All causes	2,798	100.0	2,762.5
1	Malignant neoplasms (C00-C97)	875	31.3	863.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	707	25.3	698.0
3	Cerebrovascular diseases (I60-I69)	218	7.8	215.2
4	Chronic lower respiratory diseases (J40-J47)	157	5.6	155.0
5	Diabetes mellitus (E10-E14)	110	3.9	108.6
6	Influenza and pneumonia (J09-J18)	95	3.4	93.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	76	2.7	75.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	54	1.9	53.3
9	Septicemia (A40-A41)	41	1.5	40.5
10	Parkinson's disease (G20-G21)	33	1.2	32.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	26	0.9	25.7
12	Alzheimer's disease (G30)	25	0.9	24.7
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	24	0.9	23.7
14	Aortic aneurysm and dissection (I71)	21	0.8	20.7
15	Pneumonitis due to solids and liquids (J69)	19	0.7	*
...	All other causes (Residual)	317	11.3	313.0
Asian or Pacific Islander, male, 80-84 years				
...	All causes	3,152	100.0	4,713.6
1	Malignant neoplasms (C00-C97)	831	26.4	1,242.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	817	25.9	1,221.8
3	Cerebrovascular diseases (I60-I69)	243	7.7	363.4
4	Chronic lower respiratory diseases (J40-J47)	165	5.2	246.7
5	Influenza and pneumonia (J09-J18)	126	4.0	188.4
6	Diabetes mellitus (E10-E14)	111	3.5	166.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	95	3.0	142.1
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	81	2.6	121.1
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	55	1.7	82.2
10	Alzheimer's disease (G30)	53	1.7	79.3
11	Parkinson's disease (G20-G21)	50	1.6	74.8
12	Aortic aneurysm and dissection (I71)	36	1.1	53.8
13	Septicemia (A40-A41)	34	1.1	50.8
14	Pneumonitis due to solids and liquids (J69)	29	0.9	43.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	27	0.9	40.4
...	All other causes (Residual)	399	12.7	596.7

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 85-89 years				
...	All causes	2,644	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	744	28.1	---
2	Malignant neoplasms (C00-C97)	518	19.6	---
3	Cerebrovascular diseases (I60-I69)	228	8.6	---
4	Influenza and pneumonia (J09-J18)	154	5.8	---
4	Chronic lower respiratory diseases (J40-J47)	154	5.8	---
6	Diabetes mellitus (E10-E14)	105	4.0	---
7	Alzheimer's disease (G30)	67	2.5	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	63	2.4	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	59	2.2	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	52	2.0	---
11	Parkinson's disease (G20-G21)	50	1.9	---
12	Septicemia (A40-A41)	34	1.3	---
13	Pneumonitis due to solids and liquids (J69)	29	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	24	0.9	---
15	Aortic aneurysm and dissection (I71)	23	0.9	---
...	All other causes (Residual)	340	12.9	---
Asian or Pacific Islander, male, 90-94 years				
...	All causes	1,438	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	452	31.4	---
2	Malignant neoplasms (C00-C97)	184	12.8	---
3	Cerebrovascular diseases (I60-I69)	129	9.0	---
4	Influenza and pneumonia (J09-J18)	101	7.0	---
5	Chronic lower respiratory diseases (J40-J47)	71	4.9	---
6	Alzheimer's disease (G30)	50	3.5	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	43	3.0	---
8	Diabetes mellitus (E10-E14)	39	2.7	---
9	Pneumonitis due to solids and liquids (J69)	32	2.2	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	30	2.1	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	26	1.8	---
12	Septicemia (A40-A41)	22	1.5	---
13	Parkinson's disease (G20-G21)	20	1.4	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	0.7	---
15	Atherosclerosis (I70)	6	0.4	---
...	All other causes (Residual)	223	15.5	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 95-99 years				
...	All causes	534	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	180	33.7	---
2	Malignant neoplasms (C00-C97)	48	9.0	---
3	Influenza and pneumonia (J09-J18)	36	6.7	---
4	Cerebrovascular diseases (I60-I69)	32	6.0	---
5	Alzheimer's disease (G30)	25	4.7	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	21	3.9	---
7	Chronic lower respiratory diseases (J40-J47)	20	3.7	---
8	Diabetes mellitus (E10-E14)	18	3.4	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15	2.8	---
10	Pneumonitis due to solids and liquids (J69)	12	2.2	---
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	1.5	---
12	Septicemia (A40-A41)	6	1.1	---
13	Parkinson's disease (G20-G21)	5	0.9	---
14	Anemias (D50-D64)	3	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.4	---
15	Nutritional deficiencies (E40-E64)	2	0.4	---
15	Aortic aneurysm and dissection (I71)	2	0.4	---
...	All other causes (Residual)	99	18.5	---
Asian or Pacific Islander, male, 100 years and over				
...	All causes	101	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	34	33.7	---
2	Influenza and pneumonia (J09-J18)	16	15.8	---
3	Cerebrovascular diseases (I60-I69)	7	6.9	---
4	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	5.9	---
5	Malignant neoplasms (C00-C97)	5	5.0	---
6	Diabetes mellitus (E10-E14)	4	4.0	---
6	Alzheimer's disease (G30)	4	4.0	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3	3.0	---
8	Chronic lower respiratory diseases (J40-J47)	3	3.0	---
8	Pneumonitis due to solids and liquids (J69)	3	3.0	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2	2.0	---
12	Atherosclerosis (I70)	1	1.0	---
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1	1.0	---
12	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	1.0	---
...	All other causes (Residual)	11	10.9	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, all ages				
...	All causes	21,786	100.0	287.2
1	Malignant neoplasms (C00-C97)	5,932	27.2	78.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,850	22.3	63.9
3	Cerebrovascular diseases (I60-I69)	2,005	9.2	26.4
4	Diabetes mellitus (E10-E14)	883	4.1	11.6
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	823	3.8	10.8
6	Influenza and pneumonia (J09-J18)	632	2.9	8.3
7	Chronic lower respiratory diseases (J40-J47)	536	2.5	7.1
8	Alzheimer's disease (G30)	503	2.3	6.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	417	1.9	5.5
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	348	1.6	4.6
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	272	1.2	3.6
12	Septicemia (A40-A41)	249	1.1	3.3
13	Certain conditions originating in the perinatal period (P00-P96)	209	1.0	2.8
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	163	0.7	2.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	155	0.7	2.0
...	All other causes (Residual)	3,809	17.5	50.2
Asian or Pacific Islander, female, 1-4 years				
...	All causes	73	100.0	17.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15	20.5	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	11	15.1	*
3	Malignant neoplasms (C00-C97)	9	12.3	*
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	9	12.3	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	5.5	*
6	Septicemia (A40-A41)	2	2.7	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.4	*
7	Meningitis (G00,G03)	1	1.4	*
7	Cerebrovascular diseases (I60-I69)	1	1.4	*
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	1.4	*
...	All other causes (Residual)	19	26.0	*

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 5-9 years				
...	All causes	42	100.0	8.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11	26.2	*
2	Malignant neoplasms (C00-C97)	6	14.3	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	9.5	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	7.1	*
4	Cerebrovascular diseases (I60-I69)	3	7.1	*
6	Influenza and pneumonia (J09-J18)	2	4.8	*
6	Complications of medical and surgical care (Y40-Y84,Y88)	2	4.8	*
8	Septicemia (A40-A41)	1	2.4	*
8	Hernia (K40-K46)	1	2.4	*
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1	2.4	*
...	All other causes (Residual)	8	19.0	*
Asian or Pacific Islander, female, 10-14 years				
...	All causes	50	100.0	10.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	16	32.0	*
2	Malignant neoplasms (C00-C97)	9	18.0	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	8.0	*
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4	8.0	*
5	Cerebrovascular diseases (I60-I69)	2	4.0	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	2.0	*
6	Anemias (D50-D64)	1	2.0	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	2.0	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1	2.0	*
...	All other causes (Residual)	11	22.0	*

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 15-19 years				
...	All causes	90	100.0	19.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	38	42.2	8.2
2	Malignant neoplasms (C00-C97)	13	14.4	*
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	9	10.0	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	5.6	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	3.3	*
6	Influenza and pneumonia (J09-J18)	2	2.2	*
7	Septicemia (A40-A41)	1	1.1	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.1	*
7	Meningitis (G00,G03)	1	1.1	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	1.1	*
7	Cerebrovascular diseases (I60-I69)	1	1.1	*
7	Pneumonitis due to solids and liquids (J69)	1	1.1	*
7	Diseases of appendix (K35-K38)	1	1.1	*
7	Pregnancy, childbirth and the puerperium (O00-O99)	1	1.1	*
...	All other causes (Residual)	12	13.3	*
Asian or Pacific Islander, female, 20-24 years				
...	All causes	142	100.0	29.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	59	41.5	12.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	27	19.0	5.5
3	Malignant neoplasms (C00-C97)	16	11.3	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	9	6.3	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	1.4	*
5	Cerebrovascular diseases (I60-I69)	2	1.4	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	1.4	*
8	Septicemia (A40-A41)	1	0.7	*
8	Meningitis (G00,G03)	1	0.7	*
8	Aortic aneurysm and dissection (I71)	1	0.7	*
8	Influenza and pneumonia (J09-J18)	1	0.7	*
8	Chronic lower respiratory diseases (J40-J47)	1	0.7	*
8	Inflammatory diseases of female pelvic organs (N70-N76)	1	0.7	*
...	All other causes (Residual)	19	13.4	*

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 25-29 years				
...	All causes	159	100.0	26.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	35	22.0	5.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	27	17.0	4.5
3	Malignant neoplasms (C00-C97)	22	13.8	3.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	13	8.2	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	10	6.3	*
6	Pregnancy, childbirth and the puerperium (O00-O99)	7	4.4	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	3.8	*
8	Cerebrovascular diseases (I60-I69)	5	3.1	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	2	1.3	*
9	Diabetes mellitus (E10-E14)	2	1.3	*
9	Chronic liver disease and cirrhosis (K70,K73-K74)	2	1.3	*
12	Tuberculosis (A16-A19)	1	0.6	*
12	Viral hepatitis (B15-B19)	1	0.6	*
12	Anemias (D50-D64)	1	0.6	*
12	Aortic aneurysm and dissection (I71)	1	0.6	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.6	*
...	All other causes (Residual)	23	14.5	3.8
Asian or Pacific Islander, female, 30-34 years				
...	All causes	207	100.0	29.8
1	Malignant neoplasms (C00-C97)	56	27.1	8.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	33	15.9	4.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	31	15.0	4.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	14	6.8	*
5	Pregnancy, childbirth and the puerperium (O00-O99)	11	5.3	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	11	5.3	*
7	Cerebrovascular diseases (I60-I69)	5	2.4	*
8	Diabetes mellitus (E10-E14)	4	1.9	*
9	Septicemia (A40-A41)	2	1.0	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	2	1.0	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	1.0	*
9	Influenza and pneumonia (J09-J18)	2	1.0	*
13	Tuberculosis (A16-A19)	1	0.5	*
13	Nutritional deficiencies (E40-E64)	1	0.5	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.5	*
13	Aortic aneurysm and dissection (I71)	1	0.5	*
13	Chronic lower respiratory diseases (J40-J47)	1	0.5	*
13	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.5	*
13	Inflammatory diseases of female pelvic organs (N70-N76)	1	0.5	*
...	All other causes (Residual)	26	12.6	3.7

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 35-39 years				
...	All causes	276	100.0	39.5
1	Malignant neoplasms (C00-C97)	109	39.5	15.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	29	10.5	4.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	29	10.5	4.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	23	8.3	3.3
5	Cerebrovascular diseases (I60-I69)	19	6.9	*
6	Pregnancy, childbirth and the puerperium (O00-O99)	11	4.0	*
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8	2.9	*
8	Diabetes mellitus (E10-E14)	3	1.1	*
8	Influenza and pneumonia (J09-J18)	3	1.1	*
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	1.1	*
11	Chronic lower respiratory diseases (J40-J47)	2	0.7	*
12	Tuberculosis (A16-A19)	1	0.4	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.4	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.4	*
12	Parkinson's disease (G20-G21)	1	0.4	*
12	Atherosclerosis (I70)	1	0.4	*
12	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.4	*
12	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.4	*
...	All other causes (Residual)	30	10.9	4.3
Asian or Pacific Islander, female, 40-44 years				
...	All causes	446	100.0	72.6
1	Malignant neoplasms (C00-C97)	207	46.4	33.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	44	9.9	7.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	35	7.8	5.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	31	7.0	5.0
5	Cerebrovascular diseases (I60-I69)	27	6.1	4.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	18	4.0	*
7	Septicemia (A40-A41)	6	1.3	*
7	Diabetes mellitus (E10-E14)	6	1.3	*
7	Chronic liver disease and cirrhosis (K70,K73-K74)	6	1.3	*
10	Viral hepatitis (B15-B19)	5	1.1	*
10	Chronic lower respiratory diseases (J40-J47)	5	1.1	*
12	Pregnancy, childbirth and the puerperium (O00-O99)	4	0.9	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	3	0.7	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3	0.7	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.7	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.7	*
...	All other causes (Residual)	40	9.0	6.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 45-49 years				
...	All causes	632	100.0	111.9
1	Malignant neoplasms (C00-C97)	301	47.6	53.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	52	8.2	9.2
3	Cerebrovascular diseases (I60-I69)	49	7.8	8.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	46	7.3	8.1
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	23	3.6	4.1
6	Diabetes mellitus (E10-E14)	15	2.4	*
7	Chronic liver disease and cirrhosis (K70,K73-K74)	13	2.1	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	1.4	*
9	Influenza and pneumonia (J09-J18)	8	1.3	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	1.3	*
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8	1.3	*
12	Chronic lower respiratory diseases (J40-J47)	7	1.1	*
13	Septicemia (A40-A41)	6	0.9	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	0.8	*
15	Viral hepatitis (B15-B19)	3	0.5	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	3	0.5	*
15	Anemias (D50-D64)	3	0.5	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.5	*
...	All other causes (Residual)	70	11.1	12.4
Asian or Pacific Islander, female, 50-54 years				
...	All causes	833	100.0	163.1
1	Malignant neoplasms (C00-C97)	452	54.3	88.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	78	9.4	15.3
3	Cerebrovascular diseases (I60-I69)	58	7.0	11.4
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	39	4.7	7.6
5	Diabetes mellitus (E10-E14)	23	2.8	4.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	19	2.3	*
7	Viral hepatitis (B15-B19)	15	1.8	*
8	Chronic liver disease and cirrhosis (K70,K73-K74)	12	1.4	*
9	Chronic lower respiratory diseases (J40-J47)	11	1.3	*
10	Influenza and pneumonia (J09-J18)	9	1.1	*
11	Septicemia (A40-A41)	7	0.8	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.8	*
13	Aortic aneurysm and dissection (I71)	6	0.7	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	0.7	*
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	0.7	*
...	All other causes (Residual)	85	10.2	16.6

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 55-59 years				
...	All causes	1,148	100.0	264.3
1	Malignant neoplasms (C00-C97)	588	51.2	135.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	132	11.5	30.4
3	Cerebrovascular diseases (I60-I69)	97	8.4	22.3
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	49	4.3	11.3
5	Diabetes mellitus (E10-E14)	48	4.2	11.1
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	15	1.3	*
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	1.2	*
8	Chronic lower respiratory diseases (J40-J47)	13	1.1	*
9	Viral hepatitis (B15-B19)	12	1.0	*
10	Chronic liver disease and cirrhosis (K70,K73-K74)	11	1.0	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10	0.9	*
12	Septicemia (A40-A41)	9	0.8	*
12	Aortic aneurysm and dissection (I71)	9	0.8	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.6	*
14	Influenza and pneumonia (J09-J18)	7	0.6	*
...	All other causes (Residual)	127	11.1	29.2
Asian or Pacific Islander, female, 60-64 years				
...	All causes	1,292	100.0	421.0
1	Malignant neoplasms (C00-C97)	614	47.5	200.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	215	16.6	70.1
3	Cerebrovascular diseases (I60-I69)	90	7.0	29.3
4	Diabetes mellitus (E10-E14)	57	4.4	18.6
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	44	3.4	14.3
6	Viral hepatitis (B15-B19)	19	1.5	*
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19	1.5	*
8	Influenza and pneumonia (J09-J18)	18	1.4	*
8	Chronic lower respiratory diseases (J40-J47)	18	1.4	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	1.2	*
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	16	1.2	*
12	Septicemia (A40-A41)	14	1.1	*
13	Aortic aneurysm and dissection (I71)	12	0.9	*
14	Chronic liver disease and cirrhosis (K70,K73-K74)	10	0.8	*
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8	0.6	*
...	All other causes (Residual)	122	9.4	39.8

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 65-69 years				
...	All causes	1,489	100.0	646.1
1	Malignant neoplasms (C00-C97)	589	39.6	255.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	301	20.2	130.6
3	Cerebrovascular diseases (I60-I69)	120	8.1	52.1
4	Diabetes mellitus (E10-E14)	81	5.4	35.1
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	41	2.8	17.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	31	2.1	13.5
7	Chronic lower respiratory diseases (J40-J47)	26	1.7	11.3
8	Viral hepatitis (B15-B19)	20	1.3	8.7
9	Influenza and pneumonia (J09-J18)	19	1.3	*
10	Chronic liver disease and cirrhosis (K70,K73-K74)	18	1.2	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	17	1.1	*
12	Septicemia (A40-A41)	15	1.0	*
13	Aortic aneurysm and dissection (I71)	11	0.7	*
14	Alzheimer's disease (G30)	8	0.5	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.4	*
15	Parkinson's disease (G20-G21)	6	0.4	*
...	All other causes (Residual)	180	12.1	78.1
Asian or Pacific Islander, female, 70-74 years				
...	All causes	1,965	100.0	1,066.0
1	Malignant neoplasms (C00-C97)	692	35.2	375.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	398	20.3	215.9
3	Cerebrovascular diseases (I60-I69)	181	9.2	98.2
4	Diabetes mellitus (E10-E14)	105	5.3	57.0
5	Chronic lower respiratory diseases (J40-J47)	52	2.6	28.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	50	2.5	27.1
7	Influenza and pneumonia (J09-J18)	47	2.4	25.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	44	2.2	23.9
9	Septicemia (A40-A41)	28	1.4	15.2
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	26	1.3	14.1
11	Viral hepatitis (B15-B19)	23	1.2	12.5
12	Chronic liver disease and cirrhosis (K70,K73-K74)	20	1.0	10.9
13	Alzheimer's disease (G30)	14	0.7	*
13	Aortic aneurysm and dissection (I71)	14	0.7	*
15	Parkinson's disease (G20-G21)	13	0.7	*
...	All other causes (Residual)	258	13.1	140.0

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 75-79 years				
...	All causes	2,653	100.0	1,821.1
1	Malignant neoplasms (C00-C97)	768	28.9	527.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	594	22.4	407.7
3	Cerebrovascular diseases (I60-I69)	256	9.6	175.7
4	Diabetes mellitus (E10-E14)	130	4.9	89.2
5	Chronic lower respiratory diseases (J40-J47)	84	3.2	57.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	74	2.8	50.8
7	Influenza and pneumonia (J09-J18)	72	2.7	49.4
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	71	2.7	48.7
9	Alzheimer's disease (G30)	44	1.7	30.2
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	40	1.5	27.5
11	Septicemia (A40-A41)	37	1.4	25.4
12	Viral hepatitis (B15-B19)	26	1.0	17.8
12	Chronic liver disease and cirrhosis (K70,K73-K74)	26	1.0	17.8
14	Parkinson's disease (G20-G21)	23	0.9	15.8
15	Aortic aneurysm and dissection (I71)	21	0.8	14.4
...	All other causes (Residual)	387	14.6	265.6
Asian or Pacific Islander, female, 80-84 years				
...	All causes	3,425	100.0	3,413.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	910	26.6	907.0
2	Malignant neoplasms (C00-C97)	711	20.8	708.7
3	Cerebrovascular diseases (I60-I69)	383	11.2	381.7
4	Diabetes mellitus (E10-E14)	186	5.4	185.4
5	Chronic lower respiratory diseases (J40-J47)	124	3.6	123.6
6	Influenza and pneumonia (J09-J18)	113	3.3	112.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	83	2.4	82.7
8	Alzheimer's disease (G30)	80	2.3	79.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	79	2.3	78.7
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	76	2.2	75.8
11	Septicemia (A40-A41)	47	1.4	46.8
11	Parkinson's disease (G20-G21)	47	1.4	46.8
13	Aortic aneurysm and dissection (I71)	26	0.8	25.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	25	0.7	24.9
15	Pneumonitis due to solids and liquids (J69)	22	0.6	21.9
...	All other causes (Residual)	513	15.0	511.3

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 85-89 years				
...	All causes	3,145	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	964	30.7	---
2	Malignant neoplasms (C00-C97)	477	15.2	---
3	Cerebrovascular diseases (I60-I69)	377	12.0	---
4	Alzheimer's disease (G30)	151	4.8	---
5	Diabetes mellitus (E10-E14)	129	4.1	---
6	Influenza and pneumonia (J09-J18)	107	3.4	---
7	Chronic lower respiratory diseases (J40-J47)	102	3.2	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	67	2.1	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	60	1.9	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	58	1.8	---
11	Septicemia (A40-A41)	43	1.4	---
12	Pneumonitis due to solids and liquids (J69)	33	1.0	---
13	Parkinson's disease (G20-G21)	32	1.0	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	0.7	---
15	Aortic aneurysm and dissection (I71)	13	0.4	---
...	All other causes (Residual)	509	16.2	---
Asian or Pacific Islander, female, 90-94 years				
...	All causes	2,272	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	735	32.4	---
2	Malignant neoplasms (C00-C97)	240	10.6	---
3	Cerebrovascular diseases (I60-I69)	228	10.0	---
4	Influenza and pneumonia (J09-J18)	135	5.9	---
5	Alzheimer's disease (G30)	127	5.6	---
6	Diabetes mellitus (E10-E14)	79	3.5	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	70	3.1	---
8	Chronic lower respiratory diseases (J40-J47)	62	2.7	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	53	2.3	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	37	1.6	---
11	Septicemia (A40-A41)	20	0.9	---
12	Pneumonitis due to solids and liquids (J69)	16	0.7	---
13	Atherosclerosis (I70)	14	0.6	---
13	Aortic aneurysm and dissection (I71)	14	0.6	---
15	Parkinson's disease (G20-G21)	13	0.6	---
...	All other causes (Residual)	429	18.9	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; no deaths due to J09 were reported in 2007]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 95-99 years				
...	All causes	805	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	282	35.0	---
2	Cerebrovascular diseases (I60-I69)	89	11.1	---
3	Alzheimer's disease (G30)	66	8.2	---
4	Influenza and pneumonia (J09-J18)	65	8.1	---
5	Malignant neoplasms (C00-C97)	43	5.3	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	28	3.5	---
7	Chronic lower respiratory diseases (J40-J47)	22	2.7	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15	1.9	---
9	Diabetes mellitus (E10-E14)	10	1.2	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	1.2	---
11	Anemias (D50-D64)	7	0.9	---
12	Atherosclerosis (I70)	5	0.6	---
13	Septicemia (A40-A41)	4	0.5	---
13	Parkinson's disease (G20-G21)	4	0.5	---
13	Pneumonitis due to solids and liquids (J69)	4	0.5	---
...	All other causes (Residual)	151	18.8	---
Asian or Pacific Islander, female, 100 years and over				
...	All causes	222	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	80	36.0	---
2	Influenza and pneumonia (J09-J18)	18	8.1	---
3	Cerebrovascular diseases (I60-I69)	11	5.0	---
4	Alzheimer's disease (G30)	10	4.5	---
5	Malignant neoplasms (C00-C97)	9	4.1	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	6	2.7	---
6	Chronic lower respiratory diseases (J40-J47)	6	2.7	---
8	Diabetes mellitus (E10-E14)	5	2.3	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	1.8	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4	1.8	---
11	Pneumonitis due to solids and liquids (J69)	3	1.4	---
12	Septicemia (A40-A41)	2	0.9	---
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.9	---
12	Nutritional deficiencies (E40-E64)	2	0.9	---
12	Parkinson's disease (G20-G21)	2	0.9	---
12	Atherosclerosis (I70)	2	0.9	---
...	All other causes (Residual)	56	25.2	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available