

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, all ages				
...	All causes	2,468,435	100.0	799.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	597,689	24.2	193.6
2	Malignant neoplasms (C00-C97)	574,743	23.3	186.2
3	Chronic lower respiratory diseases (J40-J47)	138,080	5.6	44.7
4	Cerebrovascular diseases (I60-I69)	129,476	5.2	41.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	120,859	4.9	39.1
6	Alzheimer's disease (G30)	83,494	3.4	27.0
7	Diabetes mellitus (E10-E14)	69,071	2.8	22.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	50,476	2.0	16.3
9	Influenza and pneumonia (J09-J18)	50,097	2.0	16.2
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	38,364	1.6	12.4
11	Septicemia (A40-A41)	34,812	1.4	11.3
12	Chronic liver disease and cirrhosis (K70,K73-K74)	31,903	1.3	10.3
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	26,634	1.1	8.6
14	Parkinson's disease (G20-G21)	22,032	0.9	7.1
15	Pneumonitis due to solids and liquids (J69)	17,011	0.7	5.5
...	All other causes (Residual)	483,694	19.6	156.7
All races, both sexes, 1-4 years				
...	All causes	4,316	100.0	26.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,394	32.3	8.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	507	11.7	3.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	385	8.9	2.4
4	Malignant neoplasms (C00-C97)	346	8.0	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	159	3.7	1.0
6	Influenza and pneumonia (J09-J18)	91	2.1	0.6
7	Septicemia (A40-A41)	62	1.4	0.4
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	59	1.4	0.4
9	Certain conditions originating in the perinatal period (P00-P96)	52	1.2	0.3
10	Chronic lower respiratory diseases (J40-J47)	51	1.2	0.3
11	Cerebrovascular diseases (I60-I69)	50	1.2	0.3
12	Anemias (D50-D64)	29	0.7	0.2
13	Meningitis (G00,G03)	22	0.5	0.1
14	Complications of medical and surgical care (Y40-Y84,Y88)	19	0.4	*
15	Pneumonitis due to solids and liquids (J69)	13	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	0.3	*
...	All other causes (Residual)	1,064	24.7	6.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 5-9 years				
...	All causes	2,330	100.0	11.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	758	32.5	3.7
2	Malignant neoplasms (C00-C97)	439	18.8	2.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	163	7.0	0.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	111	4.8	0.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	68	2.9	0.3
6	Chronic lower respiratory diseases (J40-J47)	60	2.6	0.3
7	Cerebrovascular diseases (I60-I69)	47	2.0	0.2
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	37	1.6	0.2
8	Influenza and pneumonia (J09-J18)	37	1.6	0.2
10	Septicemia (A40-A41)	32	1.4	0.2
11	Certain conditions originating in the perinatal period (P00-P96)	14	0.6	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	14	0.6	*
13	Anemias (D50-D64)	13	0.6	*
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7	0.3	*
15	Meningitis (G00,G03)	6	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.3	*
...	All other causes (Residual)	518	22.2	2.5
All races, both sexes, 10-14 years				
...	All causes	2,949	100.0	14.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	885	30.0	4.3
2	Malignant neoplasms (C00-C97)	477	16.2	2.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	267	9.1	1.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	150	5.1	0.7
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	135	4.6	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	117	4.0	0.6
7	Chronic lower respiratory diseases (J40-J47)	73	2.5	0.4
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	45	1.5	0.2
9	Cerebrovascular diseases (I60-I69)	43	1.5	0.2
10	Septicemia (A40-A41)	35	1.2	0.2
11	Influenza and pneumonia (J09-J18)	34	1.2	0.2
12	Diabetes mellitus (E10-E14)	23	0.8	0.1
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	0.5	*
14	Anemias (D50-D64)	13	0.4	*
15	Certain conditions originating in the perinatal period (P00-P96)	11	0.4	*
...	All other causes (Residual)	625	21.2	3.0

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 15-19 years				
...	All causes	10,887	100.0	49.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,537	41.7	20.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,832	16.8	8.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,659	15.2	7.5
4	Malignant neoplasms (C00-C97)	601	5.5	2.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	342	3.1	1.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	202	1.9	0.9
7	Cerebrovascular diseases (I60-I69)	86	0.8	0.4
8	Chronic lower respiratory diseases (J40-J47)	60	0.6	0.3
9	Influenza and pneumonia (J09-J18)	48	0.4	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	46	0.4	0.2
11	Diabetes mellitus (E10-E14)	45	0.4	0.2
12	Septicemia (A40-A41)	44	0.4	0.2
13	Pregnancy, childbirth and the puerperium (O00-O99)	42	0.4	0.2
14	Anemias (D50-D64)	41	0.4	0.2
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	0.2	0.1
...	All other causes (Residual)	1,279	11.7	5.8
All races, both sexes, 20-24 years				
...	All causes	18,664	100.0	86.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,804	41.8	36.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,941	15.8	13.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,846	15.2	13.2
4	Malignant neoplasms (C00-C97)	1,003	5.4	4.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	686	3.7	3.2
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	210	1.1	1.0
7	Influenza and pneumonia (J09-J18)	133	0.7	0.6
8	Human immunodeficiency virus (HIV) disease (B20-B24)	125	0.7	0.6
9	Pregnancy, childbirth and the puerperium (O00-O99)	121	0.6	0.6
10	Diabetes mellitus (E10-E14)	120	0.6	0.6
11	Cerebrovascular diseases (I60-I69)	104	0.6	0.5
12	Septicemia (A40-A41)	97	0.5	0.4
13	Chronic lower respiratory diseases (J40-J47)	89	0.5	0.4
14	Anemias (D50-D64)	68	0.4	0.3
15	Legal intervention (Y35,Y89.0)	57	0.3	0.3
...	All other causes (Residual)	2,260	12.1	10.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 25-29 years				
...	All causes	20,263	100.0	96.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,606	37.5	36.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,000	14.8	14.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,414	11.9	11.4
4	Malignant neoplasms (C00-C97)	1,415	7.0	6.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,166	5.8	5.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	249	1.2	1.2
7	Diabetes mellitus (E10-E14)	233	1.1	1.1
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	208	1.0	1.0
9	Pregnancy, childbirth and the puerperium (O00-O99)	190	0.9	0.9
10	Influenza and pneumonia (J09-J18)	172	0.8	0.8
11	Cerebrovascular diseases (I60-I69)	171	0.8	0.8
12	Septicemia (A40-A41)	137	0.7	0.6
13	Chronic liver disease and cirrhosis (K70,K73-K74)	119	0.6	0.6
14	Chronic lower respiratory diseases (J40-J47)	114	0.6	0.5
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	85	0.4	0.4
...	All other causes (Residual)	2,984	14.7	14.1
All races, both sexes, 30-34 years				
...	All causes	21,996	100.0	110.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,967	31.7	34.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,735	12.4	13.7
3	Malignant neoplasms (C00-C97)	2,204	10.0	11.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,056	9.3	10.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,844	8.4	9.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	492	2.2	2.5
7	Diabetes mellitus (E10-E14)	373	1.7	1.9
8	Chronic liver disease and cirrhosis (K70,K73-K74)	368	1.7	1.8
9	Cerebrovascular diseases (I60-I69)	346	1.6	1.7
10	Influenza and pneumonia (J09-J18)	213	1.0	1.1
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	189	0.9	0.9
12	Pregnancy, childbirth and the puerperium (O00-O99)	177	0.8	0.9
13	Septicemia (A40-A41)	175	0.8	0.9
14	Chronic lower respiratory diseases (J40-J47)	158	0.7	0.8
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	158	0.7	0.8
...	All other causes (Residual)	3,541	16.1	17.7

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 35-39 years				
...	All causes	28,012	100.0	138.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,945	24.8	34.4
2	Malignant neoplasms (C00-C97)	3,970	14.2	19.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,633	13.0	18.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,084	11.0	15.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,336	4.8	6.6
6	Chronic liver disease and cirrhosis (K70,K73-K74)	774	2.8	3.8
7	Human immunodeficiency virus (HIV) disease (B20-B24)	703	2.5	3.5
8	Diabetes mellitus (E10-E14)	640	2.3	3.2
9	Cerebrovascular diseases (I60-I69)	616	2.2	3.1
10	Septicemia (A40-A41)	297	1.1	1.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	293	1.0	1.5
12	Influenza and pneumonia (J09-J18)	289	1.0	1.4
13	Chronic lower respiratory diseases (J40-J47)	256	0.9	1.3
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	218	0.8	1.1
15	Pregnancy, childbirth and the puerperium (O00-O99)	144	0.5	0.7
...	All other causes (Residual)	4,814	17.2	23.9
All races, both sexes, 40-44 years				
...	All causes	42,021	100.0	201.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,847	18.7	37.6
2	Malignant neoplasms (C00-C97)	7,839	18.7	37.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,961	16.6	33.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,487	8.3	16.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,649	3.9	7.9
6	Cerebrovascular diseases (I60-I69)	1,288	3.1	6.2
7	Human immunodeficiency virus (HIV) disease (B20-B24)	1,195	2.8	5.7
8	Diabetes mellitus (E10-E14)	1,149	2.7	5.5
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,137	2.7	5.4
10	Influenza and pneumonia (J09-J18)	484	1.2	2.3
11	Septicemia (A40-A41)	470	1.1	2.2
12	Chronic lower respiratory diseases (J40-J47)	453	1.1	2.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	433	1.0	2.1
14	Viral hepatitis (B15-B19)	267	0.6	1.3
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	253	0.6	1.2
...	All other causes (Residual)	7,109	16.9	34.0

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 45-49 years				
...	All causes	73,569	100.0	324.0
1	Malignant neoplasms (C00-C97)	17,585	23.9	77.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,114	19.2	62.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,937	13.5	43.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,372	5.9	19.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	3,607	4.9	15.9
6	Cerebrovascular diseases (I60-I69)	2,295	3.1	10.1
7	Diabetes mellitus (E10-E14)	2,202	3.0	9.7
8	Human immunodeficiency virus (HIV) disease (B20-B24)	1,640	2.2	7.2
9	Chronic lower respiratory diseases (J40-J47)	1,360	1.8	6.0
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,075	1.5	4.7
11	Septicemia (A40-A41)	890	1.2	3.9
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	809	1.1	3.6
13	Viral hepatitis (B15-B19)	799	1.1	3.5
14	Influenza and pneumonia (J09-J18)	787	1.1	3.5
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	520	0.7	2.3
...	All other causes (Residual)	11,577	15.7	51.0
All races, both sexes, 50-54 years				
...	All causes	109,638	100.0	491.7
1	Malignant neoplasms (C00-C97)	32,626	29.8	146.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	22,615	20.6	101.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,730	8.9	43.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	5,044	4.6	22.6
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,427	4.0	19.9
6	Cerebrovascular diseases (I60-I69)	3,615	3.3	16.2
7	Diabetes mellitus (E10-E14)	3,408	3.1	15.3
8	Chronic lower respiratory diseases (J40-J47)	3,092	2.8	13.9
9	Viral hepatitis (B15-B19)	1,577	1.4	7.1
10	Human immunodeficiency virus (HIV) disease (B20-B24)	1,483	1.4	6.7
11	Septicemia (A40-A41)	1,443	1.3	6.5
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,413	1.3	6.3
13	Influenza and pneumonia (J09-J18)	1,139	1.0	5.1
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	922	0.8	4.1
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	864	0.8	3.9
...	All other causes (Residual)	16,240	14.8	72.8

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 55-59 years				
...	All causes	139,961	100.0	711.7
1	Malignant neoplasms (C00-C97)	46,899	33.5	238.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	30,240	21.6	153.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,946	5.7	40.4
4	Chronic liver disease and cirrhosis (K70,K73-K74)	5,345	3.8	27.2
5	Chronic lower respiratory diseases (J40-J47)	5,294	3.8	26.9
6	Diabetes mellitus (E10-E14)	5,089	3.6	25.9
7	Cerebrovascular diseases (I60-I69)	4,605	3.3	23.4
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,760	2.7	19.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,121	1.5	10.8
10	Septicemia (A40-A41)	2,062	1.5	10.5
11	Viral hepatitis (B15-B19)	2,034	1.5	10.3
12	Influenza and pneumonia (J09-J18)	1,578	1.1	8.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,173	0.8	6.0
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1,143	0.8	5.8
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	637	0.5	3.2
...	All other causes (Residual)	20,035	14.3	101.9
All races, both sexes, 60-64 years				
...	All causes	170,841	100.0	1,015.8
1	Malignant neoplasms (C00-C97)	62,602	36.6	372.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	37,837	22.1	225.0
3	Chronic lower respiratory diseases (J40-J47)	8,948	5.2	53.2
4	Diabetes mellitus (E10-E14)	6,588	3.9	39.2
5	Cerebrovascular diseases (I60-I69)	6,088	3.6	36.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,077	3.6	36.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	4,419	2.6	26.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,961	1.7	17.6
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,624	1.5	15.6
10	Septicemia (A40-A41)	2,542	1.5	15.1
11	Influenza and pneumonia (J09-J18)	2,049	1.2	12.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,499	0.9	8.9
13	Viral hepatitis (B15-B19)	1,184	0.7	7.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	879	0.5	5.2
15	Aortic aneurysm and dissection (I71)	731	0.4	4.3
...	All other causes (Residual)	23,813	13.9	141.6

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 65-69 years				
...	All causes	189,962	100.0	1,527.6
1	Malignant neoplasms (C00-C97)	70,489	37.1	566.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	41,189	21.7	331.2
3	Chronic lower respiratory diseases (J40-J47)	13,678	7.2	110.0
4	Cerebrovascular diseases (I60-I69)	7,533	4.0	60.6
5	Diabetes mellitus (E10-E14)	7,071	3.7	56.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,710	2.5	37.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,701	1.9	29.8
8	Chronic liver disease and cirrhosis (K70,K73-K74)	3,221	1.7	25.9
9	Septicemia (A40-A41)	3,087	1.6	24.8
10	Influenza and pneumonia (J09-J18)	2,573	1.4	20.7
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,698	0.9	13.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,677	0.9	13.5
13	Alzheimer's disease (G30)	1,229	0.6	9.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,082	0.6	8.7
15	Aortic aneurysm and dissection (I71)	940	0.5	7.6
...	All other causes (Residual)	26,084	13.7	209.8
All races, both sexes, 70-74 years				
...	All causes	217,189	100.0	2,340.9
1	Malignant neoplasms (C00-C97)	74,146	34.1	799.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	47,662	21.9	513.7
3	Chronic lower respiratory diseases (J40-J47)	18,099	8.3	195.1
4	Cerebrovascular diseases (I60-I69)	10,203	4.7	110.0
5	Diabetes mellitus (E10-E14)	7,616	3.5	82.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,840	2.2	52.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,697	2.2	50.6
8	Influenza and pneumonia (J09-J18)	3,493	1.6	37.6
9	Septicemia (A40-A41)	3,458	1.6	37.3
10	Alzheimer's disease (G30)	3,062	1.4	33.0
11	Chronic liver disease and cirrhosis (K70,K73-K74)	2,499	1.2	26.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,959	0.9	21.1
13	Parkinson's disease (G20-G21)	1,794	0.8	19.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,498	0.7	16.1
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,276	0.6	13.8
...	All other causes (Residual)	30,887	14.2	332.9

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 75-79 years				
...	All causes	273,348	100.0	3,735.4
1	Malignant neoplasms (C00-C97)	78,246	28.6	1,069.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	63,514	23.2	867.9
3	Chronic lower respiratory diseases (J40-J47)	22,636	8.3	309.3
4	Cerebrovascular diseases (I60-I69)	15,140	5.5	206.9
5	Diabetes mellitus (E10-E14)	8,874	3.2	121.3
6	Alzheimer's disease (G30)	7,888	2.9	107.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,345	2.3	86.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,912	2.2	80.8
9	Influenza and pneumonia (J09-J18)	5,170	1.9	70.6
10	Septicemia (A40-A41)	4,396	1.6	60.1
11	Parkinson's disease (G20-G21)	3,872	1.4	52.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,738	1.0	37.4
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,036	0.7	27.8
14	Chronic liver disease and cirrhosis (K70,K73-K74)	2,027	0.7	27.7
15	Pneumonitis due to solids and liquids (J69)	1,794	0.7	24.5
...	All other causes (Residual)	42,760	15.6	584.3
All races, both sexes, 80-84 years				
...	All causes	352,303	100.0	6,134.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	89,566	25.4	1,559.5
2	Malignant neoplasms (C00-C97)	78,779	22.4	1,371.7
3	Chronic lower respiratory diseases (J40-J47)	25,673	7.3	447.0
4	Cerebrovascular diseases (I60-I69)	22,519	6.4	392.1
5	Alzheimer's disease (G30)	16,211	4.6	282.3
6	Diabetes mellitus (E10-E14)	9,948	2.8	173.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,773	2.5	152.8
8	Influenza and pneumonia (J09-J18)	8,199	2.3	142.8
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,941	2.3	138.3
10	Parkinson's disease (G20-G21)	5,897	1.7	102.7
11	Septicemia (A40-A41)	5,535	1.6	96.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,025	1.1	70.1
13	Pneumonitis due to solids and liquids (J69)	3,196	0.9	55.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,674	0.8	46.6
15	Aortic aneurysm and dissection (I71)	1,885	0.5	32.8
...	All other causes (Residual)	61,482	17.5	1,070.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 85-89 years				
...	All causes	379,437	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	108,543	28.6	---
2	Malignant neoplasms (C00-C97)	60,095	15.8	---
3	Cerebrovascular diseases (I60-I69)	26,883	7.1	---
4	Alzheimer's disease (G30)	24,191	6.4	---
5	Chronic lower respiratory diseases (J40-J47)	22,404	5.9	---
6	Influenza and pneumonia (J09-J18)	10,330	2.7	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9,424	2.5	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,985	2.4	---
9	Diabetes mellitus (E10-E14)	8,940	2.4	---
10	Parkinson's disease (G20-G21)	5,728	1.5	---
11	Septicemia (A40-A41)	5,459	1.4	---
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,988	1.3	---
13	Pneumonitis due to solids and liquids (J69)	3,904	1.0	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,619	0.7	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1,758	0.5	---
...	All other causes (Residual)	75,186	19.8	---
All races, both sexes, 90-94 years				
...	All causes	259,524	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	82,968	32.0	---
2	Malignant neoplasms (C00-C97)	26,847	10.3	---
3	Alzheimer's disease (G30)	19,846	7.6	---
4	Cerebrovascular diseases (I60-I69)	19,031	7.3	---
5	Chronic lower respiratory diseases (J40-J47)	11,476	4.4	---
6	Influenza and pneumonia (J09-J18)	8,323	3.2	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,203	2.4	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,136	2.4	---
9	Diabetes mellitus (E10-E14)	4,908	1.9	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,059	1.6	---
11	Septicemia (A40-A41)	3,096	1.2	---
12	Pneumonitis due to solids and liquids (J69)	3,032	1.2	---
13	Parkinson's disease (G20-G21)	2,707	1.0	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,537	0.6	---
15	Atherosclerosis (I70)	1,527	0.6	---
...	All other causes (Residual)	57,828	22.3	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 95-99 years				
...	All causes	104,684	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	36,000	34.4	---
2	Alzheimer's disease (G30)	8,534	8.2	---
3	Cerebrovascular diseases (I60-I69)	7,360	7.0	---
4	Malignant neoplasms (C00-C97)	7,167	6.8	---
5	Influenza and pneumonia (J09-J18)	3,759	3.6	---
6	Chronic lower respiratory diseases (J40-J47)	3,525	3.4	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,475	2.4	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,272	2.2	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,055	2.0	---
10	Diabetes mellitus (E10-E14)	1,601	1.5	---
11	Pneumonitis due to solids and liquids (J69)	1,191	1.1	---
12	Septicemia (A40-A41)	1,087	1.0	---
13	Atherosclerosis (I70)	764	0.7	---
14	Parkinson's disease (G20-G21)	621	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	481	0.5	---
...	All other causes (Residual)	25,792	24.6	---
All races, both sexes, 100 years and over				
...	All causes	21,829	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,896	36.2	---
2	Alzheimer's disease (G30)	1,655	7.6	---
3	Cerebrovascular diseases (I60-I69)	1,321	6.1	---
4	Influenza and pneumonia (J09-J18)	999	4.6	---
5	Malignant neoplasms (C00-C97)	901	4.1	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	546	2.5	---
7	Chronic lower respiratory diseases (J40-J47)	540	2.5	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	444	2.0	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	436	2.0	---
10	Pneumonitis due to solids and liquids (J69)	240	1.1	---
11	Diabetes mellitus (E10-E14)	233	1.1	---
12	Atherosclerosis (I70)	221	1.0	---
13	Septicemia (A40-A41)	192	0.9	---
14	Anemias (D50-D64)	89	0.4	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	77	0.4	---
...	All other causes (Residual)	6,039	27.7	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, all ages				
...	All causes	1,232,432	100.0	812.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	307,384	24.9	202.5
2	Malignant neoplasms (C00-C97)	301,037	24.4	198.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	75,921	6.2	50.0
4	Chronic lower respiratory diseases (J40-J47)	65,423	5.3	43.1
5	Cerebrovascular diseases (I60-I69)	52,367	4.2	34.5
6	Diabetes mellitus (E10-E14)	35,490	2.9	23.4
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	30,277	2.5	19.9
8	Alzheimer's disease (G30)	25,364	2.1	16.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	24,865	2.0	16.4
10	Influenza and pneumonia (J09-J18)	23,615	1.9	15.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	20,798	1.7	13.7
12	Septicemia (A40-A41)	16,069	1.3	10.6
13	Parkinson's disease (G20-G21)	12,871	1.0	8.5
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	12,774	1.0	8.4
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10,846	0.9	7.1
...	All other causes (Residual)	217,331	17.6	143.2
All races, male, 1-4 years				
...	All causes	2,460	100.0	29.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	868	35.3	10.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	231	9.4	2.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	231	9.4	2.8
4	Malignant neoplasms (C00-C97)	197	8.0	2.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	90	3.7	1.1
6	Influenza and pneumonia (J09-J18)	50	2.0	0.6
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	32	1.3	0.4
8	Septicemia (A40-A41)	30	1.2	0.4
8	Certain conditions originating in the perinatal period (P00-P96)	30	1.2	0.4
10	Chronic lower respiratory diseases (J40-J47)	28	1.1	0.3
11	Cerebrovascular diseases (I60-I69)	26	1.1	0.3
12	Anemias (D50-D64)	17	0.7	*
13	Meningitis (G00,G03)	13	0.5	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	13	0.5	*
15	Pneumonitis due to solids and liquids (J69)	9	0.4	*
...	All other causes (Residual)	595	24.2	7.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 5-9 years				
...	All causes	1,325	100.0	12.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	483	36.5	4.6
2	Malignant neoplasms (C00-C97)	239	18.0	2.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	81	6.1	0.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	62	4.7	0.6
5	Chronic lower respiratory diseases (J40-J47)	44	3.3	0.4
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	34	2.6	0.3
7	Cerebrovascular diseases (I60-I69)	27	2.0	0.3
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	1.3	*
9	Septicemia (A40-A41)	16	1.2	*
10	Influenza and pneumonia (J09-J18)	15	1.1	*
11	Certain conditions originating in the perinatal period (P00-P96)	10	0.8	*
12	Anemias (D50-D64)	7	0.5	*
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7	0.5	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.4	*
...	All other causes (Residual)	272	20.5	2.6
All races, male, 10-14 years				
...	All causes	1,729	100.0	16.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	569	32.9	5.4
2	Malignant neoplasms (C00-C97)	243	14.1	2.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	180	10.4	1.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	104	6.0	1.0
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	67	3.9	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	62	3.6	0.6
7	Chronic lower respiratory diseases (J40-J47)	46	2.7	0.4
8	Cerebrovascular diseases (I60-I69)	27	1.6	0.3
9	Septicemia (A40-A41)	22	1.3	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	1.0	*
10	Influenza and pneumonia (J09-J18)	18	1.0	*
12	Diabetes mellitus (E10-E14)	14	0.8	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.3	*
13	Certain conditions originating in the perinatal period (P00-P96)	6	0.3	*
15	Anemias (D50-D64)	4	0.2	*
...	All other causes (Residual)	343	19.8	3.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 15-19 years				
...	All causes	7,866	100.0	69.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,204	40.7	28.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,581	20.1	14.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,323	16.8	11.7
4	Malignant neoplasms (C00-C97)	360	4.6	3.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	241	3.1	2.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	127	1.6	1.1
7	Cerebrovascular diseases (I60-I69)	53	0.7	0.5
8	Chronic lower respiratory diseases (J40-J47)	36	0.5	0.3
9	Influenza and pneumonia (J09-J18)	31	0.4	0.3
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	30	0.4	0.3
11	Anemias (D50-D64)	27	0.3	0.2
12	Diabetes mellitus (E10-E14)	24	0.3	0.2
13	Septicemia (A40-A41)	22	0.3	0.2
14	Legal intervention (Y35,Y89.0)	16	0.2	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	13	0.2	*
...	All other causes (Residual)	778	9.9	6.9
All races, male, 20-24 years				
...	All causes	13,924	100.0	126.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,884	42.3	53.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,486	17.9	22.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,448	17.6	22.2
4	Malignant neoplasms (C00-C97)	637	4.6	5.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	478	3.4	4.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	124	0.9	1.1
7	Human immunodeficiency virus (HIV) disease (B20-B24)	87	0.6	0.8
8	Influenza and pneumonia (J09-J18)	74	0.5	0.7
9	Diabetes mellitus (E10-E14)	66	0.5	0.6
10	Chronic lower respiratory diseases (J40-J47)	60	0.4	0.5
11	Legal intervention (Y35,Y89.0)	56	0.4	0.5
12	Septicemia (A40-A41)	54	0.4	0.5
12	Cerebrovascular diseases (I60-I69)	54	0.4	0.5
14	Anemias (D50-D64)	43	0.3	0.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	28	0.2	0.3
...	All other causes (Residual)	1,345	9.7	12.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 25-29 years				
...	All causes	14,429	100.0	135.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,721	39.6	53.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,459	17.0	23.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,020	14.0	19.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	801	5.6	7.5
5	Malignant neoplasms (C00-C97)	767	5.3	7.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	172	1.2	1.6
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	126	0.9	1.2
8	Diabetes mellitus (E10-E14)	125	0.9	1.2
9	Influenza and pneumonia (J09-J18)	99	0.7	0.9
10	Cerebrovascular diseases (I60-I69)	87	0.6	0.8
11	Chronic lower respiratory diseases (J40-J47)	73	0.5	0.7
12	Chronic liver disease and cirrhosis (K70,K73-K74)	67	0.5	0.6
13	Septicemia (A40-A41)	60	0.4	0.6
14	Legal intervention (Y35,Y89.0)	51	0.4	0.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	48	0.3	0.5
...	All other causes (Residual)	1,753	12.1	16.5
All races, male, 30-34 years				
...	All causes	14,763	100.0	147.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,082	34.4	50.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,184	14.8	21.8
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,554	10.5	15.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,411	9.6	14.1
5	Malignant neoplasms (C00-C97)	1,017	6.9	10.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	310	2.1	3.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	240	1.6	2.4
8	Diabetes mellitus (E10-E14)	219	1.5	2.2
9	Cerebrovascular diseases (I60-I69)	177	1.2	1.8
10	Influenza and pneumonia (J09-J18)	120	0.8	1.2
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	107	0.7	1.1
12	Chronic lower respiratory diseases (J40-J47)	97	0.7	1.0
13	Septicemia (A40-A41)	86	0.6	0.9
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	77	0.5	0.8
15	Aortic aneurysm and dissection (I71)	59	0.4	0.6
...	All other causes (Residual)	2,023	13.7	20.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 35-39 years				
...	All causes	17,614	100.0	175.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,817	27.3	48.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,512	14.3	25.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,372	13.5	23.6
4	Malignant neoplasms (C00-C97)	1,742	9.9	17.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,038	5.9	10.3
6	Chronic liver disease and cirrhosis (K70,K73-K74)	505	2.9	5.0
7	Human immunodeficiency virus (HIV) disease (B20-B24)	447	2.5	4.5
8	Diabetes mellitus (E10-E14)	369	2.1	3.7
9	Cerebrovascular diseases (I60-I69)	343	1.9	3.4
10	Septicemia (A40-A41)	164	0.9	1.6
10	Influenza and pneumonia (J09-J18)	164	0.9	1.6
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	159	0.9	1.6
13	Chronic lower respiratory diseases (J40-J47)	121	0.7	1.2
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	120	0.7	1.2
15	Aortic aneurysm and dissection (I71)	101	0.6	1.0
...	All other causes (Residual)	2,640	15.0	26.3
All races, male, 40-44 years				
...	All causes	25,820	100.0	248.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,379	20.8	51.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,854	18.8	46.7
3	Malignant neoplasms (C00-C97)	3,404	13.2	32.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,661	10.3	25.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,107	4.3	10.7
6	Human immunodeficiency virus (HIV) disease (B20-B24)	847	3.3	8.1
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	840	3.3	8.1
8	Diabetes mellitus (E10-E14)	692	2.7	6.7
9	Cerebrovascular diseases (I60-I69)	686	2.7	6.6
10	Influenza and pneumonia (J09-J18)	275	1.1	2.6
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	237	0.9	2.3
12	Septicemia (A40-A41)	236	0.9	2.3
13	Chronic lower respiratory diseases (J40-J47)	213	0.8	2.0
14	Viral hepatitis (B15-B19)	189	0.7	1.8
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	157	0.6	1.5
...	All other causes (Residual)	4,043	15.7	38.9

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 45-49 years				
...	All causes	44,946	100.0	401.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,937	22.1	88.7
2	Malignant neoplasms (C00-C97)	8,345	18.6	74.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,668	14.8	59.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,375	7.5	30.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,388	5.3	21.3
6	Diabetes mellitus (E10-E14)	1,409	3.1	12.6
7	Cerebrovascular diseases (I60-I69)	1,276	2.8	11.4
8	Human immunodeficiency virus (HIV) disease (B20-B24)	1,223	2.7	10.9
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	774	1.7	6.9
10	Chronic lower respiratory diseases (J40-J47)	590	1.3	5.3
11	Viral hepatitis (B15-B19)	553	1.2	4.9
12	Septicemia (A40-A41)	469	1.0	4.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	466	1.0	4.2
14	Influenza and pneumonia (J09-J18)	443	1.0	4.0
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	316	0.7	2.8
...	All other causes (Residual)	6,714	14.9	59.9
All races, male, 50-54 years				
...	All causes	67,072	100.0	613.5
1	Malignant neoplasms (C00-C97)	16,843	25.1	154.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	16,150	24.1	147.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,510	9.7	59.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	3,508	5.2	32.1
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,358	5.0	30.7
6	Diabetes mellitus (E10-E14)	2,119	3.2	19.4
7	Cerebrovascular diseases (I60-I69)	2,023	3.0	18.5
8	Chronic lower respiratory diseases (J40-J47)	1,463	2.2	13.4
9	Human immunodeficiency virus (HIV) disease (B20-B24)	1,119	1.7	10.2
10	Viral hepatitis (B15-B19)	1,106	1.6	10.1
11	Septicemia (A40-A41)	785	1.2	7.2
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	780	1.2	7.1
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	703	1.0	6.4
14	Influenza and pneumonia (J09-J18)	644	1.0	5.9
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	532	0.8	4.9
...	All other causes (Residual)	9,429	14.1	86.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 55-59 years				
...	All causes	86,775	100.0	911.2
1	Malignant neoplasms (C00-C97)	25,848	29.8	271.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	21,531	24.8	226.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,434	6.3	57.1
4	Chronic liver disease and cirrhosis (K70,K73-K74)	3,896	4.5	40.9
5	Diabetes mellitus (E10-E14)	3,183	3.7	33.4
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,859	3.3	30.0
7	Chronic lower respiratory diseases (J40-J47)	2,761	3.2	29.0
8	Cerebrovascular diseases (I60-I69)	2,690	3.1	28.2
9	Viral hepatitis (B15-B19)	1,458	1.7	15.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,206	1.4	12.7
11	Septicemia (A40-A41)	1,106	1.3	11.6
12	Influenza and pneumonia (J09-J18)	937	1.1	9.8
13	Human immunodeficiency virus (HIV) disease (B20-B24)	865	1.0	9.1
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	753	0.9	7.9
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	463	0.5	4.9
...	All other causes (Residual)	11,785	13.6	123.7
All races, male, 60-64 years				
...	All causes	102,520	100.0	1,269.2
1	Malignant neoplasms (C00-C97)	34,852	34.0	431.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	25,902	25.3	320.7
3	Chronic lower respiratory diseases (J40-J47)	4,658	4.5	57.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,175	4.1	51.7
5	Diabetes mellitus (E10-E14)	3,939	3.8	48.8
6	Cerebrovascular diseases (I60-I69)	3,419	3.3	42.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	3,051	3.0	37.8
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,010	2.0	24.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,678	1.6	20.8
10	Septicemia (A40-A41)	1,339	1.3	16.6
11	Influenza and pneumonia (J09-J18)	1,182	1.2	14.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	906	0.9	11.2
13	Viral hepatitis (B15-B19)	848	0.8	10.5
14	Human immunodeficiency virus (HIV) disease (B20-B24)	523	0.5	6.5
14	Aortic aneurysm and dissection (I71)	523	0.5	6.5
...	All other causes (Residual)	13,515	13.2	167.3

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 65-69 years				
...	All causes	109,519	100.0	1,871.3
1	Malignant neoplasms (C00-C97)	39,048	35.7	667.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	26,731	24.4	456.7
3	Chronic lower respiratory diseases (J40-J47)	7,032	6.4	120.2
4	Cerebrovascular diseases (I60-I69)	4,121	3.8	70.4
5	Diabetes mellitus (E10-E14)	4,061	3.7	69.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,037	2.8	51.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,098	1.9	35.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,024	1.8	34.6
9	Septicemia (A40-A41)	1,582	1.4	27.0
10	Influenza and pneumonia (J09-J18)	1,478	1.3	25.3
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,345	1.2	23.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	915	0.8	15.6
13	Aortic aneurysm and dissection (I71)	658	0.6	11.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	611	0.6	10.4
15	Alzheimer's disease (G30)	540	0.5	9.2
...	All other causes (Residual)	14,238	13.0	243.3
All races, male, 70-74 years				
...	All causes	120,185	100.0	2,831.9
1	Malignant neoplasms (C00-C97)	40,637	33.8	957.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	29,105	24.2	685.8
3	Chronic lower respiratory diseases (J40-J47)	9,186	7.6	216.4
4	Cerebrovascular diseases (I60-I69)	5,163	4.3	121.7
5	Diabetes mellitus (E10-E14)	4,107	3.4	96.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,901	2.4	68.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,555	2.1	60.2
8	Influenza and pneumonia (J09-J18)	1,947	1.6	45.9
9	Septicemia (A40-A41)	1,716	1.4	40.4
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,463	1.2	34.5
11	Alzheimer's disease (G30)	1,343	1.1	31.6
12	Parkinson's disease (G20-G21)	1,180	1.0	27.8
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,073	0.9	25.3
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	990	0.8	23.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	853	0.7	20.1
...	All other causes (Residual)	15,966	13.3	376.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 75-79 years				
...	All causes	143,006	100.0	4,493.7
1	Malignant neoplasms (C00-C97)	42,186	29.5	1,325.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	35,647	24.9	1,120.1
3	Chronic lower respiratory diseases (J40-J47)	11,088	7.8	348.4
4	Cerebrovascular diseases (I60-I69)	7,000	4.9	220.0
5	Diabetes mellitus (E10-E14)	4,611	3.2	144.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,346	2.3	105.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,324	2.3	104.4
8	Alzheimer's disease (G30)	3,178	2.2	99.9
9	Influenza and pneumonia (J09-J18)	2,759	1.9	86.7
10	Parkinson's disease (G20-G21)	2,493	1.7	78.3
11	Septicemia (A40-A41)	2,131	1.5	67.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,195	0.8	37.6
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,128	0.8	35.4
14	Chronic liver disease and cirrhosis (K70,K73-K74)	1,122	0.8	35.3
15	Pneumonitis due to solids and liquids (J69)	1,045	0.7	32.8
...	All other causes (Residual)	20,753	14.5	652.1
All races, male, 80-84 years				
...	All causes	168,824	100.0	7,358.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	45,176	26.8	1,969.0
2	Malignant neoplasms (C00-C97)	40,743	24.1	1,775.8
3	Chronic lower respiratory diseases (J40-J47)	12,197	7.2	531.6
4	Cerebrovascular diseases (I60-I69)	9,166	5.4	399.5
5	Alzheimer's disease (G30)	5,887	3.5	256.6
6	Diabetes mellitus (E10-E14)	4,656	2.8	202.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,434	2.6	193.3
8	Influenza and pneumonia (J09-J18)	4,061	2.4	177.0
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,013	2.4	174.9
10	Parkinson's disease (G20-G21)	3,625	2.1	158.0
11	Septicemia (A40-A41)	2,533	1.5	110.4
12	Pneumonitis due to solids and liquids (J69)	1,846	1.1	80.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,612	1.0	70.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,413	0.8	61.6
15	Aortic aneurysm and dissection (I71)	954	0.6	41.6
...	All other causes (Residual)	26,508	15.7	1,155.3

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 85-89 years				
...	All causes	157,700	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	46,714	29.6	---
2	Malignant neoplasms (C00-C97)	29,172	18.5	---
3	Chronic lower respiratory diseases (J40-J47)	9,986	6.3	---
4	Cerebrovascular diseases (I60-I69)	9,111	5.8	---
5	Alzheimer's disease (G30)	7,486	4.7	---
6	Influenza and pneumonia (J09-J18)	4,608	2.9	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,321	2.7	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,072	2.6	---
9	Diabetes mellitus (E10-E14)	3,716	2.4	---
10	Parkinson's disease (G20-G21)	3,134	2.0	---
11	Septicemia (A40-A41)	2,192	1.4	---
12	Pneumonitis due to solids and liquids (J69)	2,071	1.3	---
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,699	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,270	0.8	---
15	Aortic aneurysm and dissection (I71)	748	0.5	---
...	All other causes (Residual)	27,400	17.4	---
All races, male, 90-94 years				
...	All causes	88,586	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	29,221	33.0	---
2	Malignant neoplasms (C00-C97)	11,834	13.4	---
3	Cerebrovascular diseases (I60-I69)	5,252	5.9	---
4	Alzheimer's disease (G30)	4,958	5.6	---
5	Chronic lower respiratory diseases (J40-J47)	4,461	5.0	---
6	Influenza and pneumonia (J09-J18)	3,252	3.7	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,578	2.9	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,308	2.6	---
9	Diabetes mellitus (E10-E14)	1,710	1.9	---
10	Pneumonitis due to solids and liquids (J69)	1,475	1.7	---
11	Parkinson's disease (G20-G21)	1,285	1.5	---
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,118	1.3	---
13	Septicemia (A40-A41)	1,080	1.2	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	619	0.7	---
15	Atherosclerosis (I70)	490	0.6	---
...	All other causes (Residual)	16,945	19.1	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 95-99 years				
...	All causes	25,973	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,246	35.6	---
2	Malignant neoplasms (C00-C97)	2,639	10.2	---
3	Alzheimer's disease (G30)	1,448	5.6	---
4	Cerebrovascular diseases (I60-I69)	1,428	5.5	---
5	Influenza and pneumonia (J09-J18)	1,117	4.3	---
6	Chronic lower respiratory diseases (J40-J47)	1,106	4.3	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	773	3.0	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	714	2.7	---
9	Pneumonitis due to solids and liquids (J69)	468	1.8	---
10	Diabetes mellitus (E10-E14)	426	1.6	---
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	417	1.6	---
12	Septicemia (A40-A41)	289	1.1	---
13	Parkinson's disease (G20-G21)	234	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	168	0.6	---
15	Atherosclerosis (I70)	164	0.6	---
...	All other causes (Residual)	5,336	20.5	---
All races, male, 100 years and over				
...	All causes	3,607	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,325	36.7	---
2	Malignant neoplasms (C00-C97)	248	6.9	---
3	Influenza and pneumonia (J09-J18)	228	6.3	---
4	Cerebrovascular diseases (I60-I69)	173	4.8	---
5	Alzheimer's disease (G30)	155	4.3	---
6	Chronic lower respiratory diseases (J40-J47)	149	4.1	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	111	3.1	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	88	2.4	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	71	2.0	---
10	Pneumonitis due to solids and liquids (J69)	67	1.9	---
11	Diabetes mellitus (E10-E14)	40	1.1	---
12	Septicemia (A40-A41)	37	1.0	---
12	Atherosclerosis (I70)	37	1.0	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	0.6	---
15	Anemias (D50-D64)	14	0.4	---
...	All other causes (Residual)	844	23.4	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, all ages				
...	All causes	1,236,003	100.0	787.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	290,305	23.5	184.9
2	Malignant neoplasms (C00-C97)	273,706	22.1	174.4
3	Cerebrovascular diseases (I60-I69)	77,109	6.2	49.1
4	Chronic lower respiratory diseases (J40-J47)	72,657	5.9	46.3
5	Alzheimer's disease (G30)	58,130	4.7	37.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	44,938	3.6	28.6
7	Diabetes mellitus (E10-E14)	33,581	2.7	21.4
8	Influenza and pneumonia (J09-J18)	26,482	2.1	16.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	25,611	2.1	16.3
10	Septicemia (A40-A41)	18,743	1.5	11.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	15,788	1.3	10.1
12	Chronic liver disease and cirrhosis (K70,K73-K74)	11,105	0.9	7.1
13	Parkinson's disease (G20-G21)	9,161	0.7	5.8
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8,087	0.7	5.2
15	Pneumonitis due to solids and liquids (J69)	7,803	0.6	5.0
...	All other causes (Residual)	262,797	21.3	167.4
All races, female, 1-4 years				
...	All causes	1,856	100.0	23.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	526	28.3	6.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	276	14.9	3.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	154	8.3	1.9
4	Malignant neoplasms (C00-C97)	149	8.0	1.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	69	3.7	0.9
6	Influenza and pneumonia (J09-J18)	41	2.2	0.5
7	Septicemia (A40-A41)	32	1.7	0.4
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	27	1.5	0.3
9	Cerebrovascular diseases (I60-I69)	24	1.3	0.3
10	Chronic lower respiratory diseases (J40-J47)	23	1.2	0.3
11	Certain conditions originating in the perinatal period (P00-P96)	22	1.2	0.3
12	Anemias (D50-D64)	12	0.6	*
13	Meningitis (G00,G03)	9	0.5	*
14	Meningococcal infection (A39)	7	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.4	*
...	All other causes (Residual)	478	25.8	6.0

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 5-9 years				
...	All causes	1,005	100.0	10.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	275	27.4	2.8
2	Malignant neoplasms (C00-C97)	200	19.9	2.0
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	82	8.2	0.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	49	4.9	0.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	34	3.4	0.3
6	Influenza and pneumonia (J09-J18)	22	2.2	0.2
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	2.0	0.2
7	Cerebrovascular diseases (I60-I69)	20	2.0	0.2
9	Septicemia (A40-A41)	16	1.6	*
9	Chronic lower respiratory diseases (J40-J47)	16	1.6	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	8	0.8	*
12	Anemias (D50-D64)	6	0.6	*
13	Certain conditions originating in the perinatal period (P00-P96)	4	0.4	*
13	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	4	0.4	*
15	Diabetes mellitus (E10-E14)	3	0.3	*
15	Meningitis (G00,G03)	3	0.3	*
15	Pneumonitis due to solids and liquids (J69)	3	0.3	*
...	All other causes (Residual)	240	23.9	2.4
All races, female, 10-14 years				
...	All causes	1,220	100.0	12.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	316	25.9	3.1
2	Malignant neoplasms (C00-C97)	234	19.2	2.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	87	7.1	0.9
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	68	5.6	0.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	55	4.5	0.5
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	46	3.8	0.5
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	27	2.2	0.3
7	Chronic lower respiratory diseases (J40-J47)	27	2.2	0.3
9	Cerebrovascular diseases (I60-I69)	16	1.3	*
9	Influenza and pneumonia (J09-J18)	16	1.3	*
11	Septicemia (A40-A41)	13	1.1	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	0.8	*
13	Anemias (D50-D64)	9	0.7	*
13	Diabetes mellitus (E10-E14)	9	0.7	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.5	*
...	All other causes (Residual)	281	23.0	2.8

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 15-19 years				
...	All causes	3,021	100.0	28.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,333	44.1	12.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	336	11.1	3.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	251	8.3	2.3
4	Malignant neoplasms (C00-C97)	241	8.0	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	101	3.3	0.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	75	2.5	0.7
7	Pregnancy, childbirth and the puerperium (O00-O99)	42	1.4	0.4
8	Cerebrovascular diseases (I60-I69)	33	1.1	0.3
9	Chronic lower respiratory diseases (J40-J47)	24	0.8	0.2
10	Septicemia (A40-A41)	22	0.7	0.2
11	Diabetes mellitus (E10-E14)	21	0.7	0.2
12	Influenza and pneumonia (J09-J18)	17	0.6	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	0.5	*
14	Anemias (D50-D64)	14	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	0.4	*
...	All other causes (Residual)	482	16.0	4.5
All races, female, 20-24 years				
...	All causes	4,740	100.0	44.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,920	40.5	18.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	493	10.4	4.7
3	Malignant neoplasms (C00-C97)	366	7.7	3.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	360	7.6	3.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	208	4.4	2.0
6	Pregnancy, childbirth and the puerperium (O00-O99)	121	2.6	1.1
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	86	1.8	0.8
8	Influenza and pneumonia (J09-J18)	59	1.2	0.6
9	Diabetes mellitus (E10-E14)	54	1.1	0.5
10	Cerebrovascular diseases (I60-I69)	50	1.1	0.5
11	Septicemia (A40-A41)	43	0.9	0.4
12	Human immunodeficiency virus (HIV) disease (B20-B24)	38	0.8	0.4
13	Chronic lower respiratory diseases (J40-J47)	29	0.6	0.3
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	29	0.6	0.3
15	Anemias (D50-D64)	25	0.5	0.2
...	All other causes (Residual)	859	18.1	8.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 25-29 years				
...	All causes	5,834	100.0	55.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,885	32.3	18.0
2	Malignant neoplasms (C00-C97)	648	11.1	6.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	541	9.3	5.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	394	6.8	3.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	365	6.3	3.5
6	Pregnancy, childbirth and the puerperium (O00-O99)	190	3.3	1.8
7	Diabetes mellitus (E10-E14)	108	1.9	1.0
8	Cerebrovascular diseases (I60-I69)	84	1.4	0.8
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	82	1.4	0.8
10	Septicemia (A40-A41)	77	1.3	0.7
10	Human immunodeficiency virus (HIV) disease (B20-B24)	77	1.3	0.7
12	Influenza and pneumonia (J09-J18)	73	1.3	0.7
13	Chronic liver disease and cirrhosis (K70,K73-K74)	52	0.9	0.5
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	46	0.8	0.4
15	Chronic lower respiratory diseases (J40-J47)	41	0.7	0.4
...	All other causes (Residual)	1,171	20.1	11.2
All races, female, 30-34 years				
...	All causes	7,233	100.0	72.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,885	26.1	18.9
2	Malignant neoplasms (C00-C97)	1,187	16.4	11.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	645	8.9	6.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	551	7.6	5.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	290	4.0	2.9
6	Human immunodeficiency virus (HIV) disease (B20-B24)	182	2.5	1.8
7	Pregnancy, childbirth and the puerperium (O00-O99)	177	2.4	1.8
8	Cerebrovascular diseases (I60-I69)	169	2.3	1.7
9	Diabetes mellitus (E10-E14)	154	2.1	1.5
10	Chronic liver disease and cirrhosis (K70,K73-K74)	128	1.8	1.3
11	Influenza and pneumonia (J09-J18)	93	1.3	0.9
12	Septicemia (A40-A41)	89	1.2	0.9
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	82	1.1	0.8
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	81	1.1	0.8
15	Chronic lower respiratory diseases (J40-J47)	61	0.8	0.6
...	All other causes (Residual)	1,459	20.2	14.6

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 35-39 years				
...	All causes	10,398	100.0	102.6
1	Malignant neoplasms (C00-C97)	2,228	21.4	22.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,128	20.5	21.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,121	10.8	11.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	712	6.8	7.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	298	2.9	2.9
6	Cerebrovascular diseases (I60-I69)	273	2.6	2.7
7	Diabetes mellitus (E10-E14)	271	2.6	2.7
8	Chronic liver disease and cirrhosis (K70,K73-K74)	269	2.6	2.7
9	Human immunodeficiency virus (HIV) disease (B20-B24)	256	2.5	2.5
10	Pregnancy, childbirth and the puerperium (O00-O99)	144	1.4	1.4
11	Chronic lower respiratory diseases (J40-J47)	135	1.3	1.3
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	134	1.3	1.3
13	Septicemia (A40-A41)	133	1.3	1.3
14	Influenza and pneumonia (J09-J18)	125	1.2	1.2
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	98	0.9	1.0
...	All other causes (Residual)	2,073	19.9	20.4
All races, female, 40-44 years				
...	All causes	16,201	100.0	154.3
1	Malignant neoplasms (C00-C97)	4,435	27.4	42.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,468	15.2	23.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,107	13.0	20.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	826	5.1	7.9
5	Cerebrovascular diseases (I60-I69)	602	3.7	5.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	542	3.3	5.2
7	Diabetes mellitus (E10-E14)	457	2.8	4.4
8	Human immunodeficiency virus (HIV) disease (B20-B24)	348	2.1	3.3
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	297	1.8	2.8
10	Chronic lower respiratory diseases (J40-J47)	240	1.5	2.3
11	Septicemia (A40-A41)	234	1.4	2.2
12	Influenza and pneumonia (J09-J18)	209	1.3	2.0
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	196	1.2	1.9
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	100	0.6	1.0
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	96	0.6	0.9
...	All other causes (Residual)	3,044	18.8	29.0

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 45-49 years				
...	All causes	28,623	100.0	248.9
1	Malignant neoplasms (C00-C97)	9,240	32.3	80.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,177	14.6	36.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,269	11.4	28.4
4	Chronic liver disease and cirrhosis (K70,K73-K74)	1,219	4.3	10.6
5	Cerebrovascular diseases (I60-I69)	1,019	3.6	8.9
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	997	3.5	8.7
7	Diabetes mellitus (E10-E14)	793	2.8	6.9
8	Chronic lower respiratory diseases (J40-J47)	770	2.7	6.7
9	Septicemia (A40-A41)	421	1.5	3.7
10	Human immunodeficiency virus (HIV) disease (B20-B24)	417	1.5	3.6
11	Influenza and pneumonia (J09-J18)	344	1.2	3.0
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	343	1.2	3.0
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	301	1.1	2.6
14	Viral hepatitis (B15-B19)	246	0.9	2.1
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	204	0.7	1.8
...	All other causes (Residual)	4,863	17.0	42.3
All races, female, 50-54 years				
...	All causes	42,566	100.0	374.5
1	Malignant neoplasms (C00-C97)	15,783	37.1	138.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,465	15.2	56.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,220	7.6	28.3
4	Chronic lower respiratory diseases (J40-J47)	1,629	3.8	14.3
5	Cerebrovascular diseases (I60-I69)	1,592	3.7	14.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1,536	3.6	13.5
7	Diabetes mellitus (E10-E14)	1,289	3.0	11.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,069	2.5	9.4
9	Septicemia (A40-A41)	658	1.5	5.8
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	633	1.5	5.6
11	Influenza and pneumonia (J09-J18)	495	1.2	4.4
12	Viral hepatitis (B15-B19)	471	1.1	4.1
13	Human immunodeficiency virus (HIV) disease (B20-B24)	364	0.9	3.2
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	332	0.8	2.9
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	219	0.5	1.9
...	All other causes (Residual)	6,811	16.0	59.9

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 55-59 years				
...	All causes	53,186	100.0	524.5
1	Malignant neoplasms (C00-C97)	21,051	39.6	207.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,709	16.4	85.9
3	Chronic lower respiratory diseases (J40-J47)	2,533	4.8	25.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,512	4.7	24.8
5	Cerebrovascular diseases (I60-I69)	1,915	3.6	18.9
6	Diabetes mellitus (E10-E14)	1,906	3.6	18.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,449	2.7	14.3
8	Septicemia (A40-A41)	956	1.8	9.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	915	1.7	9.0
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	901	1.7	8.9
11	Influenza and pneumonia (J09-J18)	641	1.2	6.3
12	Viral hepatitis (B15-B19)	576	1.1	5.7
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	420	0.8	4.1
14	Human immunodeficiency virus (HIV) disease (B20-B24)	278	0.5	2.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	232	0.4	2.3
...	All other causes (Residual)	8,192	15.4	80.8
All races, female, 60-64 years				
...	All causes	68,321	100.0	781.7
1	Malignant neoplasms (C00-C97)	27,750	40.6	317.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,935	17.5	136.5
3	Chronic lower respiratory diseases (J40-J47)	4,290	6.3	49.1
4	Cerebrovascular diseases (I60-I69)	2,669	3.9	30.5
5	Diabetes mellitus (E10-E14)	2,649	3.9	30.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,902	2.8	21.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,368	2.0	15.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,283	1.9	14.7
9	Septicemia (A40-A41)	1,203	1.8	13.8
10	Influenza and pneumonia (J09-J18)	867	1.3	9.9
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	614	0.9	7.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	593	0.9	6.8
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	366	0.5	4.2
14	Viral hepatitis (B15-B19)	336	0.5	3.8
15	Alzheimer's disease (G30)	313	0.5	3.6
...	All other causes (Residual)	10,183	14.9	116.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 65-69 years				
...	All causes	80,443	100.0	1,222.0
1	Malignant neoplasms (C00-C97)	31,441	39.1	477.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,458	18.0	219.6
3	Chronic lower respiratory diseases (J40-J47)	6,646	8.3	101.0
4	Cerebrovascular diseases (I60-I69)	3,412	4.2	51.8
5	Diabetes mellitus (E10-E14)	3,010	3.7	45.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,677	2.1	25.5
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,673	2.1	25.4
8	Septicemia (A40-A41)	1,505	1.9	22.9
9	Chronic liver disease and cirrhosis (K70,K73-K74)	1,123	1.4	17.1
10	Influenza and pneumonia (J09-J18)	1,095	1.4	16.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	762	0.9	11.6
12	Alzheimer's disease (G30)	689	0.9	10.5
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	471	0.6	7.2
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	353	0.4	5.4
15	Pneumonitis due to solids and liquids (J69)	302	0.4	4.6
...	All other causes (Residual)	11,826	14.7	179.7
All races, female, 70-74 years				
...	All causes	97,004	100.0	1,926.9
1	Malignant neoplasms (C00-C97)	33,509	34.5	665.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	18,557	19.1	368.6
3	Chronic lower respiratory diseases (J40-J47)	8,913	9.2	177.0
4	Cerebrovascular diseases (I60-I69)	5,040	5.2	100.1
5	Diabetes mellitus (E10-E14)	3,509	3.6	69.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,285	2.4	45.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,796	1.9	35.7
8	Septicemia (A40-A41)	1,742	1.8	34.6
9	Alzheimer's disease (G30)	1,719	1.8	34.1
10	Influenza and pneumonia (J09-J18)	1,546	1.6	30.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1,036	1.1	20.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	969	1.0	19.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	645	0.7	12.8
14	Parkinson's disease (G20-G21)	614	0.6	12.2
15	Pneumonitis due to solids and liquids (J69)	450	0.5	8.9
...	All other causes (Residual)	14,674	15.1	291.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 75-79 years				
...	All causes	130,342	100.0	3,151.9
1	Malignant neoplasms (C00-C97)	36,060	27.7	872.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	27,867	21.4	673.9
3	Chronic lower respiratory diseases (J40-J47)	11,548	8.9	279.2
4	Cerebrovascular diseases (I60-I69)	8,140	6.2	196.8
5	Alzheimer's disease (G30)	4,710	3.6	113.9
6	Diabetes mellitus (E10-E14)	4,263	3.3	103.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,021	2.3	73.1
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,566	2.0	62.0
9	Influenza and pneumonia (J09-J18)	2,411	1.8	58.3
10	Septicemia (A40-A41)	2,265	1.7	54.8
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,543	1.2	37.3
12	Parkinson's disease (G20-G21)	1,379	1.1	33.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	908	0.7	22.0
14	Chronic liver disease and cirrhosis (K70,K73-K74)	905	0.7	21.9
15	Pneumonitis due to solids and liquids (J69)	749	0.6	18.1
...	All other causes (Residual)	22,007	16.9	532.2
All races, female, 80-84 years				
...	All causes	183,479	100.0	5,319.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	44,390	24.2	1,287.1
2	Malignant neoplasms (C00-C97)	38,036	20.7	1,102.8
3	Chronic lower respiratory diseases (J40-J47)	13,476	7.3	390.7
4	Cerebrovascular diseases (I60-I69)	13,353	7.3	387.2
5	Alzheimer's disease (G30)	10,324	5.6	299.3
6	Diabetes mellitus (E10-E14)	5,292	2.9	153.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,339	2.4	125.8
8	Influenza and pneumonia (J09-J18)	4,138	2.3	120.0
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,928	2.1	113.9
10	Septicemia (A40-A41)	3,002	1.6	87.0
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,413	1.3	70.0
12	Parkinson's disease (G20-G21)	2,272	1.2	65.9
13	Pneumonitis due to solids and liquids (J69)	1,350	0.7	39.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,261	0.7	36.6
15	Aortic aneurysm and dissection (I71)	931	0.5	27.0
...	All other causes (Residual)	34,974	19.1	1,014.0

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 85-89 years				
...	All causes	221,737	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	61,829	27.9	---
2	Malignant neoplasms (C00-C97)	30,923	13.9	---
3	Cerebrovascular diseases (I60-I69)	17,772	8.0	---
4	Alzheimer's disease (G30)	16,705	7.5	---
5	Chronic lower respiratory diseases (J40-J47)	12,418	5.6	---
6	Influenza and pneumonia (J09-J18)	5,722	2.6	---
7	Diabetes mellitus (E10-E14)	5,224	2.4	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,103	2.3	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,913	2.2	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,289	1.5	---
11	Septicemia (A40-A41)	3,267	1.5	---
12	Parkinson's disease (G20-G21)	2,594	1.2	---
13	Pneumonitis due to solids and liquids (J69)	1,833	0.8	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,349	0.6	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1,098	0.5	---
...	All other causes (Residual)	47,698	21.5	---
All races, female, 90-94 years				
...	All causes	170,938	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	53,747	31.4	---
2	Malignant neoplasms (C00-C97)	15,013	8.8	---
3	Alzheimer's disease (G30)	14,888	8.7	---
4	Cerebrovascular diseases (I60-I69)	13,779	8.1	---
5	Chronic lower respiratory diseases (J40-J47)	7,015	4.1	---
6	Influenza and pneumonia (J09-J18)	5,071	3.0	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,828	2.2	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,625	2.1	---
9	Diabetes mellitus (E10-E14)	3,198	1.9	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,941	1.7	---
11	Septicemia (A40-A41)	2,016	1.2	---
12	Pneumonitis due to solids and liquids (J69)	1,557	0.9	---
13	Parkinson's disease (G20-G21)	1,422	0.8	---
14	Atherosclerosis (I70)	1,037	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	918	0.5	---
...	All other causes (Residual)	40,883	23.9	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 95-99 years				
...	All causes	78,711	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	26,754	34.0	---
2	Alzheimer's disease (G30)	7,086	9.0	---
3	Cerebrovascular diseases (I60-I69)	5,932	7.5	---
4	Malignant neoplasms (C00-C97)	4,528	5.8	---
5	Influenza and pneumonia (J09-J18)	2,642	3.4	---
6	Chronic lower respiratory diseases (J40-J47)	2,419	3.1	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,761	2.2	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,638	2.1	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,499	1.9	---
10	Diabetes mellitus (E10-E14)	1,175	1.5	---
11	Septicemia (A40-A41)	798	1.0	---
12	Pneumonitis due to solids and liquids (J69)	723	0.9	---
13	Atherosclerosis (I70)	600	0.8	---
14	Parkinson's disease (G20-G21)	387	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	313	0.4	---
...	All other causes (Residual)	20,456	26.0	---
All races, female, 100 years and over				
...	All causes	18,222	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,571	36.1	---
2	Alzheimer's disease (G30)	1,500	8.2	---
3	Cerebrovascular diseases (I60-I69)	1,148	6.3	---
4	Influenza and pneumonia (J09-J18)	771	4.2	---
5	Malignant neoplasms (C00-C97)	653	3.6	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	475	2.6	---
7	Chronic lower respiratory diseases (J40-J47)	391	2.1	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	356	2.0	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	325	1.8	---
10	Diabetes mellitus (E10-E14)	193	1.1	---
11	Atherosclerosis (I70)	184	1.0	---
12	Pneumonitis due to solids and liquids (J69)	173	0.9	---
13	Septicemia (A40-A41)	155	0.9	---
14	Anemias (D50-D64)	75	0.4	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	57	0.3	---
...	All other causes (Residual)	5,195	28.5	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, all ages				
...	All causes	2,114,749	100.0	861.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	514,323	24.3	209.6
2	Malignant neoplasms (C00-C97)	491,686	23.3	200.3
3	Chronic lower respiratory diseases (J40-J47)	127,176	6.0	51.8
4	Cerebrovascular diseases (I60-I69)	109,119	5.2	44.5
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	104,945	5.0	42.8
6	Alzheimer's disease (G30)	76,928	3.6	31.3
7	Diabetes mellitus (E10-E14)	54,250	2.6	22.1
8	Influenza and pneumonia (J09-J18)	43,296	2.0	17.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	40,205	1.9	16.4
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	34,690	1.6	14.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	28,014	1.3	11.4
12	Septicemia (A40-A41)	27,985	1.3	11.4
13	Parkinson's disease (G20-G21)	20,792	1.0	8.5
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	20,560	1.0	8.4
15	Pneumonitis due to solids and liquids (J69)	15,235	0.7	6.2
...	All other causes (Residual)	405,545	19.2	165.2
White, both sexes, 1-4 years				
...	All causes	3,015	100.0	24.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,054	35.0	8.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	348	11.5	2.8
3	Malignant neoplasms (C00-C97)	269	8.9	2.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	214	7.1	1.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	95	3.2	0.8
6	Influenza and pneumonia (J09-J18)	61	2.0	0.5
7	Septicemia (A40-A41)	47	1.6	0.4
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	43	1.4	0.4
9	Certain conditions originating in the perinatal period (P00-P96)	34	1.1	0.3
10	Cerebrovascular diseases (I60-I69)	30	1.0	0.2
11	Chronic lower respiratory diseases (J40-J47)	23	0.8	0.2
12	Meningitis (G00,G03)	16	0.5	*
13	Anemias (D50-D64)	14	0.5	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	13	0.4	*
15	Acute bronchitis and bronchiolitis (J20-J21)	10	0.3	*
...	All other causes (Residual)	744	24.7	6.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 5-9 years				
...	All causes	1,691	100.0	10.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	568	33.6	3.7
2	Malignant neoplasms (C00-C97)	341	20.2	2.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	124	7.3	0.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	70	4.1	0.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	42	2.5	0.3
6	Influenza and pneumonia (J09-J18)	29	1.7	0.2
7	Cerebrovascular diseases (I60-I69)	27	1.6	0.2
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	26	1.5	0.2
8	Chronic lower respiratory diseases (J40-J47)	26	1.5	0.2
10	Septicemia (A40-A41)	19	1.1	*
11	Certain conditions originating in the perinatal period (P00-P96)	10	0.6	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	9	0.5	*
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7	0.4	*
14	Meningitis (G00,G03)	6	0.4	*
15	Anemias (D50-D64)	5	0.3	*
15	Pneumonitis due to solids and liquids (J69)	5	0.3	*
...	All other causes (Residual)	377	22.3	2.4
White, both sexes, 10-14 years				
...	All causes	2,150	100.0	13.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	680	31.6	4.3
2	Malignant neoplasms (C00-C97)	366	17.0	2.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	202	9.4	1.3
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	90	4.2	0.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	78	3.6	0.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	74	3.4	0.5
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	34	1.6	0.2
7	Cerebrovascular diseases (I60-I69)	34	1.6	0.2
9	Chronic lower respiratory diseases (J40-J47)	27	1.3	0.2
10	Septicemia (A40-A41)	25	1.2	0.2
10	Influenza and pneumonia (J09-J18)	25	1.2	0.2
12	Diabetes mellitus (E10-E14)	18	0.8	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	0.5	*
14	Certain conditions originating in the perinatal period (P00-P96)	7	0.3	*
15	Anemias (D50-D64)	5	0.2	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.2	*
...	All other causes (Residual)	469	21.8	3.0

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 15-19 years				
...	All causes	7,847	100.0	47.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,748	47.8	22.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,382	17.6	8.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	667	8.5	4.0
4	Malignant neoplasms (C00-C97)	465	5.9	2.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	222	2.8	1.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	142	1.8	0.9
7	Cerebrovascular diseases (I60-I69)	57	0.7	0.3
8	Diabetes mellitus (E10-E14)	36	0.5	0.2
9	Influenza and pneumonia (J09-J18)	33	0.4	0.2
10	Septicemia (A40-A41)	32	0.4	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	32	0.4	0.2
12	Chronic lower respiratory diseases (J40-J47)	31	0.4	0.2
13	Pregnancy, childbirth and the puerperium (O00-O99)	29	0.4	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	0.2	*
15	Meningitis (G00,G03)	11	0.1	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	11	0.1	*
...	All other causes (Residual)	932	11.9	5.6
White, both sexes, 20-24 years				
...	All causes	13,662	100.0	82.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,610	48.4	40.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,410	17.6	14.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,057	7.7	6.4
4	Malignant neoplasms (C00-C97)	778	5.7	4.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	417	3.1	2.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	162	1.2	1.0
7	Influenza and pneumonia (J09-J18)	101	0.7	0.6
8	Diabetes mellitus (E10-E14)	77	0.6	0.5
8	Pregnancy, childbirth and the puerperium (O00-O99)	77	0.6	0.5
10	Cerebrovascular diseases (I60-I69)	72	0.5	0.4
11	Septicemia (A40-A41)	63	0.5	0.4
12	Chronic lower respiratory diseases (J40-J47)	51	0.4	0.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	36	0.3	0.2
14	Legal intervention (Y35,Y89.0)	33	0.2	0.2
15	Human immunodeficiency virus (HIV) disease (B20-B24)	29	0.2	0.2
...	All other causes (Residual)	1,689	12.4	10.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 25-29 years				
...	All causes	15,097	100.0	92.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,473	42.9	39.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,551	16.9	15.6
3	Malignant neoplasms (C00-C97)	1,101	7.3	6.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	969	6.4	5.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	763	5.1	4.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	151	1.0	0.9
7	Diabetes mellitus (E10-E14)	150	1.0	0.9
8	Influenza and pneumonia (J09-J18)	127	0.8	0.8
9	Pregnancy, childbirth and the puerperium (O00-O99)	121	0.8	0.7
10	Cerebrovascular diseases (I60-I69)	114	0.8	0.7
11	Septicemia (A40-A41)	94	0.6	0.6
12	Chronic liver disease and cirrhosis (K70,K73-K74)	89	0.6	0.5
13	Human immunodeficiency virus (HIV) disease (B20-B24)	80	0.5	0.5
14	Chronic lower respiratory diseases (J40-J47)	69	0.5	0.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	58	0.4	0.4
...	All other causes (Residual)	2,187	14.5	13.4
White, both sexes, 30-34 years				
...	All causes	16,328	100.0	106.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,909	36.2	38.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,366	14.5	15.4
3	Malignant neoplasms (C00-C97)	1,689	10.3	11.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,410	8.6	9.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	751	4.6	4.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	302	1.8	2.0
7	Diabetes mellitus (E10-E14)	235	1.4	1.5
7	Cerebrovascular diseases (I60-I69)	235	1.4	1.5
9	Human immunodeficiency virus (HIV) disease (B20-B24)	176	1.1	1.1
10	Influenza and pneumonia (J09-J18)	155	0.9	1.0
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	149	0.9	1.0
12	Septicemia (A40-A41)	116	0.7	0.8
13	Pregnancy, childbirth and the puerperium (O00-O99)	99	0.6	0.6
14	Chronic lower respiratory diseases (J40-J47)	91	0.6	0.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	91	0.6	0.6
...	All other causes (Residual)	2,554	15.6	16.6

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 35-39 years				
...	All causes	20,935	100.0	133.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,927	28.3	37.9
2	Malignant neoplasms (C00-C97)	2,976	14.2	19.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,713	13.0	17.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,471	11.8	15.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	618	3.0	3.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	615	2.9	3.9
7	Diabetes mellitus (E10-E14)	419	2.0	2.7
8	Cerebrovascular diseases (I60-I69)	379	1.8	2.4
9	Human immunodeficiency virus (HIV) disease (B20-B24)	281	1.3	1.8
10	Influenza and pneumonia (J09-J18)	225	1.1	1.4
11	Septicemia (A40-A41)	202	1.0	1.3
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	179	0.9	1.1
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	168	0.8	1.1
14	Chronic lower respiratory diseases (J40-J47)	160	0.8	1.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	85	0.4	0.5
...	All other causes (Residual)	3,517	16.8	22.5
White, both sexes, 40-44 years				
...	All causes	32,125	100.0	194.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,774	21.1	41.1
2	Malignant neoplasms (C00-C97)	6,092	19.0	36.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,000	15.6	30.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,142	9.8	19.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,413	4.4	8.6
6	Cerebrovascular diseases (I60-I69)	838	2.6	5.1
7	Diabetes mellitus (E10-E14)	777	2.4	4.7
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	658	2.0	4.0
9	Human immunodeficiency virus (HIV) disease (B20-B24)	514	1.6	3.1
10	Influenza and pneumonia (J09-J18)	364	1.1	2.2
11	Chronic lower respiratory diseases (J40-J47)	349	1.1	2.1
12	Septicemia (A40-A41)	324	1.0	2.0
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	246	0.8	1.5
14	Viral hepatitis (B15-B19)	228	0.7	1.4
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	192	0.6	1.2
...	All other causes (Residual)	5,214	16.2	31.6

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 45-49 years				
...	All causes	57,398	100.0	314.4
1	Malignant neoplasms (C00-C97)	13,859	24.1	75.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,500	18.3	57.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,559	14.9	46.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,055	7.1	22.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	3,146	5.5	17.2
6	Diabetes mellitus (E10-E14)	1,547	2.7	8.5
7	Cerebrovascular diseases (I60-I69)	1,509	2.6	8.3
8	Chronic lower respiratory diseases (J40-J47)	1,124	2.0	6.2
9	Human immunodeficiency virus (HIV) disease (B20-B24)	767	1.3	4.2
10	Viral hepatitis (B15-B19)	678	1.2	3.7
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	640	1.1	3.5
12	Septicemia (A40-A41)	637	1.1	3.5
13	Influenza and pneumonia (J09-J18)	587	1.0	3.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	494	0.9	2.7
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	280	0.5	1.5
...	All other causes (Residual)	9,016	15.7	49.4
White, both sexes, 50-54 years				
...	All causes	85,651	100.0	471.9
1	Malignant neoplasms (C00-C97)	25,844	30.2	142.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	17,089	20.0	94.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,254	9.6	45.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	4,327	5.1	23.8
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,143	4.8	22.8
6	Chronic lower respiratory diseases (J40-J47)	2,607	3.0	14.4
7	Diabetes mellitus (E10-E14)	2,428	2.8	13.4
8	Cerebrovascular diseases (I60-I69)	2,360	2.8	13.0
9	Viral hepatitis (B15-B19)	1,299	1.5	7.2
10	Septicemia (A40-A41)	1,021	1.2	5.6
11	Influenza and pneumonia (J09-J18)	881	1.0	4.9
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	840	1.0	4.6
13	Human immunodeficiency virus (HIV) disease (B20-B24)	663	0.8	3.7
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	590	0.7	3.3
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	526	0.6	2.9
...	All other causes (Residual)	12,779	14.9	70.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 55-59 years				
...	All causes	110,343	100.0	678.9
1	Malignant neoplasms (C00-C97)	37,547	34.0	231.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	23,311	21.1	143.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,671	6.0	41.0
4	Chronic liver disease and cirrhosis (K70,K73-K74)	4,602	4.2	28.3
5	Chronic lower respiratory diseases (J40-J47)	4,588	4.2	28.2
6	Diabetes mellitus (E10-E14)	3,713	3.4	22.8
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,541	3.2	21.8
8	Cerebrovascular diseases (I60-I69)	3,070	2.8	18.9
9	Viral hepatitis (B15-B19)	1,661	1.5	10.2
10	Septicemia (A40-A41)	1,491	1.4	9.2
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,346	1.2	8.3
12	Influenza and pneumonia (J09-J18)	1,245	1.1	7.7
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	721	0.7	4.4
14	Human immunodeficiency virus (HIV) disease (B20-B24)	480	0.4	3.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	453	0.4	2.8
...	All other causes (Residual)	15,903	14.4	97.8
White, both sexes, 60-64 years				
...	All causes	139,240	100.0	982.4
1	Malignant neoplasms (C00-C97)	51,940	37.3	366.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	30,137	21.6	212.6
3	Chronic lower respiratory diseases (J40-J47)	7,985	5.7	56.3
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,163	3.7	36.4
5	Diabetes mellitus (E10-E14)	4,993	3.6	35.2
6	Cerebrovascular diseases (I60-I69)	4,450	3.2	31.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	3,838	2.8	27.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,460	1.8	17.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,029	1.5	14.3
10	Septicemia (A40-A41)	1,893	1.4	13.4
11	Influenza and pneumonia (J09-J18)	1,654	1.2	11.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	979	0.7	6.9
13	Viral hepatitis (B15-B19)	921	0.7	6.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	751	0.5	5.3
15	Aortic aneurysm and dissection (I71)	623	0.4	4.4
...	All other causes (Residual)	19,424	14.0	137.0

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 65-69 years				
...	All causes	158,882	100.0	1,495.8
1	Malignant neoplasms (C00-C97)	60,026	37.8	565.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	33,634	21.2	316.7
3	Chronic lower respiratory diseases (J40-J47)	12,498	7.9	117.7
4	Cerebrovascular diseases (I60-I69)	5,749	3.6	54.1
5	Diabetes mellitus (E10-E14)	5,411	3.4	50.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,033	2.5	38.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,890	1.8	27.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,690	1.7	25.3
9	Septicemia (A40-A41)	2,394	1.5	22.5
10	Influenza and pneumonia (J09-J18)	2,093	1.3	19.7
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,611	1.0	15.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,143	0.7	10.8
13	Alzheimer's disease (G30)	1,094	0.7	10.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	953	0.6	9.0
15	Aortic aneurysm and dissection (I71)	826	0.5	7.8
...	All other causes (Residual)	21,837	13.7	205.6
White, both sexes, 70-74 years				
...	All causes	184,095	100.0	2,315.5
1	Malignant neoplasms (C00-C97)	63,781	34.6	802.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	39,593	21.5	498.0
3	Chronic lower respiratory diseases (J40-J47)	16,666	9.1	209.6
4	Cerebrovascular diseases (I60-I69)	8,093	4.4	101.8
5	Diabetes mellitus (E10-E14)	5,895	3.2	74.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,110	2.2	51.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,608	2.0	45.4
8	Influenza and pneumonia (J09-J18)	2,872	1.6	36.1
9	Septicemia (A40-A41)	2,730	1.5	34.3
10	Alzheimer's disease (G30)	2,705	1.5	34.0
11	Chronic liver disease and cirrhosis (K70,K73-K74)	2,279	1.2	28.7
12	Parkinson's disease (G20-G21)	1,659	0.9	20.9
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,406	0.8	17.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,314	0.7	16.5
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,212	0.7	15.2
...	All other causes (Residual)	26,172	14.2	329.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 75-79 years				
...	All causes	237,879	100.0	3,734.9
1	Malignant neoplasms (C00-C97)	68,520	28.8	1,075.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	54,416	22.9	854.4
3	Chronic lower respiratory diseases (J40-J47)	21,170	8.9	332.4
4	Cerebrovascular diseases (I60-I69)	12,754	5.4	200.2
5	Alzheimer's disease (G30)	7,128	3.0	111.9
6	Diabetes mellitus (E10-E14)	7,028	3.0	110.3
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,295	2.2	83.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,067	2.1	79.6
9	Influenza and pneumonia (J09-J18)	4,416	1.9	69.3
10	Parkinson's disease (G20-G21)	3,645	1.5	57.2
11	Septicemia (A40-A41)	3,598	1.5	56.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,096	0.9	32.9
13	Chronic liver disease and cirrhosis (K70,K73-K74)	1,869	0.8	29.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,828	0.8	28.7
15	Pneumonitis due to solids and liquids (J69)	1,586	0.7	24.9
...	All other causes (Residual)	37,463	15.7	588.2
White, both sexes, 80-84 years				
...	All causes	314,629	100.0	6,171.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	79,517	25.3	1,559.7
2	Malignant neoplasms (C00-C97)	70,296	22.3	1,378.9
3	Chronic lower respiratory diseases (J40-J47)	24,086	7.7	472.5
4	Cerebrovascular diseases (I60-I69)	19,770	6.3	387.8
5	Alzheimer's disease (G30)	14,890	4.7	292.1
6	Diabetes mellitus (E10-E14)	8,250	2.6	161.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,398	2.4	145.1
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,355	2.3	144.3
9	Influenza and pneumonia (J09-J18)	7,203	2.3	141.3
10	Parkinson's disease (G20-G21)	5,613	1.8	110.1
11	Septicemia (A40-A41)	4,703	1.5	92.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,256	1.0	63.9
13	Pneumonitis due to solids and liquids (J69)	2,883	0.9	56.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,474	0.8	48.5
15	Aortic aneurysm and dissection (I71)	1,727	0.5	33.9
...	All other causes (Residual)	55,208	17.5	1,082.9

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 85-89 years				
...	All causes	345,144	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	98,846	28.6	---
2	Malignant neoplasms (C00-C97)	54,260	15.7	---
3	Cerebrovascular diseases (I60-I69)	24,307	7.0	---
4	Alzheimer's disease (G30)	22,469	6.5	---
5	Chronic lower respiratory diseases (J40-J47)	21,053	6.1	---
6	Influenza and pneumonia (J09-J18)	9,275	2.7	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,471	2.5	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,240	2.4	---
9	Diabetes mellitus (E10-E14)	7,564	2.2	---
10	Parkinson's disease (G20-G21)	5,437	1.6	---
11	Septicemia (A40-A41)	4,694	1.4	---
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,199	1.2	---
13	Pneumonitis due to solids and liquids (J69)	3,605	1.0	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,436	0.7	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1,660	0.5	---
...	All other causes (Residual)	68,628	19.9	---
White, both sexes, 90-94 years				
...	All causes	238,152	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	76,543	32.1	---
2	Malignant neoplasms (C00-C97)	24,317	10.2	---
3	Alzheimer's disease (G30)	18,492	7.8	---
4	Cerebrovascular diseases (I60-I69)	17,372	7.3	---
5	Chronic lower respiratory diseases (J40-J47)	10,785	4.5	---
6	Influenza and pneumonia (J09-J18)	7,557	3.2	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,818	2.4	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,465	2.3	---
9	Diabetes mellitus (E10-E14)	4,203	1.8	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,498	1.5	---
11	Pneumonitis due to solids and liquids (J69)	2,828	1.2	---
12	Septicemia (A40-A41)	2,702	1.1	---
13	Parkinson's disease (G20-G21)	2,588	1.1	---
14	Atherosclerosis (I70)	1,436	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,409	0.6	---
...	All other causes (Residual)	53,139	22.3	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 95-99 years				
...	All causes	95,268	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	33,016	34.7	---
2	Alzheimer's disease (G30)	7,875	8.3	---
3	Cerebrovascular diseases (I60-I69)	6,664	7.0	---
4	Malignant neoplasms (C00-C97)	6,398	6.7	---
5	Influenza and pneumonia (J09-J18)	3,410	3.6	---
6	Chronic lower respiratory diseases (J40-J47)	3,280	3.4	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,335	2.5	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,979	2.1	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,745	1.8	---
10	Diabetes mellitus (E10-E14)	1,326	1.4	---
11	Pneumonitis due to solids and liquids (J69)	1,087	1.1	---
12	Septicemia (A40-A41)	934	1.0	---
13	Atherosclerosis (I70)	707	0.7	---
14	Parkinson's disease (G20-G21)	590	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	442	0.5	---
...	All other causes (Residual)	23,480	24.6	---
White, both sexes, 100 years and over				
...	All causes	19,169	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,994	36.5	---
2	Alzheimer's disease (G30)	1,471	7.7	---
3	Cerebrovascular diseases (I60-I69)	1,159	6.0	---
4	Influenza and pneumonia (J09-J18)	880	4.6	---
5	Malignant neoplasms (C00-C97)	770	4.0	---
6	Chronic lower respiratory diseases (J40-J47)	482	2.5	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	443	2.3	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	422	2.2	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	365	1.9	---
10	Pneumonitis due to solids and liquids (J69)	214	1.1	---
11	Atherosclerosis (I70)	195	1.0	---
12	Diabetes mellitus (E10-E14)	176	0.9	---
13	Septicemia (A40-A41)	151	0.8	---
14	Anemias (D50-D64)	82	0.4	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	73	0.4	---
...	All other causes (Residual)	5,292	27.6	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
White, male, all ages				
...	All causes	1,051,514	100.0	866.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	264,425	25.1	217.8
2	Malignant neoplasms (C00-C97)	258,272	24.6	212.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	65,360	6.2	53.8
4	Chronic lower respiratory diseases (J40-J47)	59,632	5.7	49.1
5	Cerebrovascular diseases (I60-I69)	43,424	4.1	35.8
6	Diabetes mellitus (E10-E14)	28,486	2.7	23.5
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	27,422	2.6	22.6
8	Alzheimer's disease (G30)	23,442	2.2	19.3
9	Influenza and pneumonia (J09-J18)	20,238	1.9	16.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20,172	1.9	16.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	18,352	1.7	15.1
12	Septicemia (A40-A41)	12,976	1.2	10.7
13	Parkinson's disease (G20-G21)	12,173	1.2	10.0
14	Pneumonitis due to solids and liquids (J69)	8,264	0.8	6.8
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	8,229	0.8	6.8
...	All other causes (Residual)	180,647	17.2	148.8
White, male, 1-4 years				
...	All causes	1,718	100.0	27.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	653	38.0	10.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	160	9.3	2.5
3	Malignant neoplasms (C00-C97)	149	8.7	2.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	130	7.6	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	53	3.1	0.8
6	Influenza and pneumonia (J09-J18)	33	1.9	0.5
7	Septicemia (A40-A41)	23	1.3	0.4
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	1.3	0.4
9	Certain conditions originating in the perinatal period (P00-P96)	19	1.1	*
10	Cerebrovascular diseases (I60-I69)	18	1.0	*
11	Chronic lower respiratory diseases (J40-J47)	14	0.8	*
12	Meningitis (G00,G03)	11	0.6	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	9	0.5	*
14	Anemias (D50-D64)	8	0.5	*
15	Acute bronchitis and bronchiolitis (J20-J21)	7	0.4	*
...	All other causes (Residual)	408	23.7	6.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 5-9 years				
...	All causes	957	100.0	12.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	359	37.5	4.5
2	Malignant neoplasms (C00-C97)	193	20.2	2.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	58	6.1	0.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	37	3.9	0.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	17	1.8	*
6	Cerebrovascular diseases (I60-I69)	15	1.6	*
6	Chronic lower respiratory diseases (J40-J47)	15	1.6	*
8	Septicemia (A40-A41)	11	1.1	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	1.1	*
8	Influenza and pneumonia (J09-J18)	11	1.1	*
11	Certain conditions originating in the perinatal period (P00-P96)	9	0.9	*
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7	0.7	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.5	*
14	Meningitis (G00,G03)	3	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.3	*
...	All other causes (Residual)	203	21.2	2.6
White, male, 10-14 years				
...	All causes	1,265	100.0	15.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	426	33.7	5.3
2	Malignant neoplasms (C00-C97)	193	15.3	2.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	144	11.4	1.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	53	4.2	0.7
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	46	3.6	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	37	2.9	0.5
7	Cerebrovascular diseases (I60-I69)	21	1.7	0.3
8	Chronic lower respiratory diseases (J40-J47)	16	1.3	*
9	Septicemia (A40-A41)	15	1.2	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	1.1	*
11	Influenza and pneumonia (J09-J18)	12	0.9	*
12	Diabetes mellitus (E10-E14)	11	0.9	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.3	*
13	Certain conditions originating in the perinatal period (P00-P96)	4	0.3	*
15	Anemias (D50-D64)	2	0.2	*
15	Aortic aneurysm and dissection (I71)	2	0.2	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2	0.2	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.2	*
...	All other causes (Residual)	261	20.6	3.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 15-19 years				
...	All causes	5,549	100.0	64.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,618	47.2	30.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,105	19.9	12.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	547	9.9	6.4
4	Malignant neoplasms (C00-C97)	287	5.2	3.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	160	2.9	1.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	84	1.5	1.0
7	Cerebrovascular diseases (I60-I69)	37	0.7	0.4
8	Influenza and pneumonia (J09-J18)	22	0.4	0.3
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	0.3	*
9	Diabetes mellitus (E10-E14)	19	0.3	*
11	Chronic lower respiratory diseases (J40-J47)	18	0.3	*
12	Septicemia (A40-A41)	15	0.3	*
13	Aortic aneurysm and dissection (I71)	9	0.2	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.2	*
15	Legal intervention (Y35,Y89.0)	7	0.1	*
...	All other causes (Residual)	593	10.7	6.9
White, male, 20-24 years				
...	All causes	10,112	100.0	119.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,002	49.5	58.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,011	19.9	23.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	852	8.4	10.0
4	Malignant neoplasms (C00-C97)	501	5.0	5.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	311	3.1	3.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	100	1.0	1.2
7	Influenza and pneumonia (J09-J18)	54	0.5	0.6
8	Diabetes mellitus (E10-E14)	44	0.4	0.5
9	Cerebrovascular diseases (I60-I69)	43	0.4	0.5
10	Legal intervention (Y35,Y89.0)	33	0.3	0.4
11	Septicemia (A40-A41)	31	0.3	0.4
11	Chronic lower respiratory diseases (J40-J47)	31	0.3	0.4
13	Human immunodeficiency virus (HIV) disease (B20-B24)	24	0.2	0.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	22	0.2	0.3
15	Aortic aneurysm and dissection (I71)	18	0.2	*
...	All other causes (Residual)	1,035	10.2	12.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 25-29 years				
...	All causes	10,817	100.0	130.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,898	45.3	58.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,089	19.3	25.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	755	7.0	9.1
4	Malignant neoplasms (C00-C97)	613	5.7	7.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	543	5.0	6.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	88	0.8	1.1
7	Diabetes mellitus (E10-E14)	86	0.8	1.0
8	Influenza and pneumonia (J09-J18)	76	0.7	0.9
9	Human immunodeficiency virus (HIV) disease (B20-B24)	60	0.6	0.7
10	Cerebrovascular diseases (I60-I69)	56	0.5	0.7
10	Chronic liver disease and cirrhosis (K70,K73-K74)	56	0.5	0.7
12	Septicemia (A40-A41)	43	0.4	0.5
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	40	0.4	0.5
13	Chronic lower respiratory diseases (J40-J47)	40	0.4	0.5
15	Legal intervention (Y35,Y89.0)	35	0.3	0.4
...	All other causes (Residual)	1,339	12.4	16.1
White, male, 30-34 years				
...	All causes	11,066	100.0	141.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,321	39.0	55.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,887	17.1	24.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	997	9.0	12.7
4	Malignant neoplasms (C00-C97)	801	7.2	10.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	580	5.2	7.4
6	Chronic liver disease and cirrhosis (K70,K73-K74)	208	1.9	2.7
7	Diabetes mellitus (E10-E14)	139	1.3	1.8
8	Human immunodeficiency virus (HIV) disease (B20-B24)	134	1.2	1.7
9	Cerebrovascular diseases (I60-I69)	117	1.1	1.5
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	85	0.8	1.1
11	Influenza and pneumonia (J09-J18)	82	0.7	1.0
12	Septicemia (A40-A41)	63	0.6	0.8
13	Chronic lower respiratory diseases (J40-J47)	52	0.5	0.7
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	48	0.4	0.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	45	0.4	0.6
...	All other causes (Residual)	1,507	13.6	19.3

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 35-39 years				
...	All causes	13,391	100.0	169.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,104	30.6	52.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,089	15.6	26.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,752	13.1	22.2
4	Malignant neoplasms (C00-C97)	1,375	10.3	17.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	437	3.3	5.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	414	3.1	5.2
7	Diabetes mellitus (E10-E14)	248	1.9	3.1
8	Cerebrovascular diseases (I60-I69)	219	1.6	2.8
9	Human immunodeficiency virus (HIV) disease (B20-B24)	212	1.6	2.7
10	Influenza and pneumonia (J09-J18)	126	0.9	1.6
11	Septicemia (A40-A41)	110	0.8	1.4
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	99	0.7	1.3
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	89	0.7	1.1
14	Chronic lower respiratory diseases (J40-J47)	73	0.5	0.9
15	Aortic aneurysm and dissection (I71)	60	0.4	0.8
...	All other causes (Residual)	1,984	14.8	25.1
White, male, 40-44 years				
...	All causes	20,095	100.0	241.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,637	23.1	55.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,623	18.0	43.6
3	Malignant neoplasms (C00-C97)	2,702	13.4	32.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,411	12.0	29.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	960	4.8	11.6
6	Diabetes mellitus (E10-E14)	466	2.3	5.6
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	456	2.3	5.5
8	Cerebrovascular diseases (I60-I69)	446	2.2	5.4
9	Human immunodeficiency virus (HIV) disease (B20-B24)	409	2.0	4.9
10	Influenza and pneumonia (J09-J18)	200	1.0	2.4
11	Septicemia (A40-A41)	168	0.8	2.0
12	Chronic lower respiratory diseases (J40-J47)	160	0.8	1.9
13	Viral hepatitis (B15-B19)	158	0.8	1.9
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	135	0.7	1.6
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	111	0.6	1.3
...	All other causes (Residual)	3,053	15.2	36.7

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 45-49 years				
...	All causes	35,738	100.0	392.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,595	21.3	83.4
2	Malignant neoplasms (C00-C97)	6,728	18.8	73.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,737	16.1	63.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,133	8.8	34.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,118	5.9	23.3
6	Diabetes mellitus (E10-E14)	1,019	2.9	11.2
7	Cerebrovascular diseases (I60-I69)	856	2.4	9.4
8	Human immunodeficiency virus (HIV) disease (B20-B24)	650	1.8	7.1
9	Chronic lower respiratory diseases (J40-J47)	484	1.4	5.3
10	Viral hepatitis (B15-B19)	471	1.3	5.2
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	436	1.2	4.8
12	Septicemia (A40-A41)	339	0.9	3.7
13	Influenza and pneumonia (J09-J18)	321	0.9	3.5
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	303	0.8	3.3
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	177	0.5	1.9
...	All other causes (Residual)	5,371	15.0	59.0
White, male, 50-54 years				
...	All causes	53,279	100.0	592.7
1	Malignant neoplasms (C00-C97)	13,466	25.3	149.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,536	23.5	139.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,509	10.3	61.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,153	5.9	35.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	3,047	5.7	33.9
6	Diabetes mellitus (E10-E14)	1,560	2.9	17.4
7	Cerebrovascular diseases (I60-I69)	1,343	2.5	14.9
8	Chronic lower respiratory diseases (J40-J47)	1,247	2.3	13.9
9	Viral hepatitis (B15-B19)	915	1.7	10.2
10	Septicemia (A40-A41)	571	1.1	6.4
11	Human immunodeficiency virus (HIV) disease (B20-B24)	557	1.0	6.2
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	497	0.9	5.5
13	Influenza and pneumonia (J09-J18)	486	0.9	5.4
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	435	0.8	4.8
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	356	0.7	4.0
...	All other causes (Residual)	7,601	14.3	84.6

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 55-59 years				
...	All causes	69,195	100.0	869.4
1	Malignant neoplasms (C00-C97)	20,742	30.0	260.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	17,085	24.7	214.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,549	6.6	57.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	3,363	4.9	42.3
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,691	3.9	33.8
6	Chronic lower respiratory diseases (J40-J47)	2,390	3.5	30.0
7	Diabetes mellitus (E10-E14)	2,385	3.4	30.0
8	Cerebrovascular diseases (I60-I69)	1,802	2.6	22.6
9	Viral hepatitis (B15-B19)	1,206	1.7	15.2
10	Septicemia (A40-A41)	804	1.2	10.1
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	772	1.1	9.7
12	Influenza and pneumonia (J09-J18)	740	1.1	9.3
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	482	0.7	6.1
14	Human immunodeficiency virus (HIV) disease (B20-B24)	406	0.6	5.1
15	Aortic aneurysm and dissection (I71)	291	0.4	3.7
...	All other causes (Residual)	9,487	13.7	119.2
White, male, 60-64 years				
...	All causes	84,101	100.0	1,222.1
1	Malignant neoplasms (C00-C97)	28,954	34.4	420.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	21,054	25.0	306.0
3	Chronic lower respiratory diseases (J40-J47)	4,111	4.9	59.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,524	4.2	51.2
5	Diabetes mellitus (E10-E14)	3,076	3.7	44.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	2,660	3.2	38.7
7	Cerebrovascular diseases (I60-I69)	2,505	3.0	36.4
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,881	2.2	27.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,173	1.4	17.0
10	Septicemia (A40-A41)	1,003	1.2	14.6
11	Influenza and pneumonia (J09-J18)	956	1.1	13.9
12	Viral hepatitis (B15-B19)	658	0.8	9.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	626	0.7	9.1
14	Aortic aneurysm and dissection (I71)	458	0.5	6.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	447	0.5	6.5
...	All other causes (Residual)	11,015	13.1	160.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 65-69 years				
...	All causes	92,184	100.0	1,825.2
1	Malignant neoplasms (C00-C97)	33,269	36.1	658.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	22,197	24.1	439.5
3	Chronic lower respiratory diseases (J40-J47)	6,365	6.9	126.0
4	Diabetes mellitus (E10-E14)	3,210	3.5	63.6
5	Cerebrovascular diseases (I60-I69)	3,156	3.4	62.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,608	2.8	51.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,876	2.0	37.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,510	1.6	29.9
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,272	1.4	25.2
10	Septicemia (A40-A41)	1,245	1.4	24.6
11	Influenza and pneumonia (J09-J18)	1,198	1.3	23.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	649	0.7	12.8
13	Aortic aneurysm and dissection (I71)	586	0.6	11.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	556	0.6	11.0
15	Alzheimer's disease (G30)	489	0.5	9.7
...	All other causes (Residual)	11,998	13.0	237.6
White, male, 70-74 years				
...	All causes	102,609	100.0	2,792.1
1	Malignant neoplasms (C00-C97)	35,032	34.1	953.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	24,601	24.0	669.4
3	Chronic lower respiratory diseases (J40-J47)	8,376	8.2	227.9
4	Cerebrovascular diseases (I60-I69)	4,112	4.0	111.9
5	Diabetes mellitus (E10-E14)	3,299	3.2	89.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,528	2.5	68.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,952	1.9	53.1
8	Influenza and pneumonia (J09-J18)	1,594	1.6	43.4
9	Septicemia (A40-A41)	1,385	1.3	37.7
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,337	1.3	36.4
11	Alzheimer's disease (G30)	1,183	1.2	32.2
12	Parkinson's disease (G20-G21)	1,100	1.1	29.9
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,021	1.0	27.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	764	0.7	20.8
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	721	0.7	19.6
...	All other causes (Residual)	13,604	13.3	370.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 75-79 years				
...	All causes	125,427	100.0	4,472.1
1	Malignant neoplasms (C00-C97)	37,061	29.5	1,321.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	31,021	24.7	1,106.0
3	Chronic lower respiratory diseases (J40-J47)	10,279	8.2	366.5
4	Cerebrovascular diseases (I60-I69)	5,924	4.7	211.2
5	Diabetes mellitus (E10-E14)	3,781	3.0	134.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,005	2.4	107.1
7	Alzheimer's disease (G30)	2,870	2.3	102.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,757	2.2	98.3
9	Influenza and pneumonia (J09-J18)	2,375	1.9	84.7
10	Parkinson's disease (G20-G21)	2,348	1.9	83.7
11	Septicemia (A40-A41)	1,740	1.4	62.0
12	Chronic liver disease and cirrhosis (K70,K73-K74)	1,037	0.8	37.0
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,031	0.8	36.8
14	Pneumonitis due to solids and liquids (J69)	926	0.7	33.0
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	924	0.7	32.9
...	All other causes (Residual)	18,348	14.6	654.2
White, male, 80-84 years				
...	All causes	152,116	100.0	7,379.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	40,757	26.8	1,977.1
2	Malignant neoplasms (C00-C97)	36,495	24.0	1,770.4
3	Chronic lower respiratory diseases (J40-J47)	11,352	7.5	550.7
4	Cerebrovascular diseases (I60-I69)	8,106	5.3	393.2
5	Alzheimer's disease (G30)	5,447	3.6	264.2
6	Diabetes mellitus (E10-E14)	3,996	2.6	193.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,846	2.5	186.6
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,708	2.4	179.9
9	Influenza and pneumonia (J09-J18)	3,588	2.4	174.1
10	Parkinson's disease (G20-G21)	3,465	2.3	168.1
11	Septicemia (A40-A41)	2,191	1.4	106.3
12	Pneumonitis due to solids and liquids (J69)	1,668	1.1	80.9
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,320	0.9	64.0
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,310	0.9	63.5
15	Aortic aneurysm and dissection (I71)	888	0.6	43.1
...	All other causes (Residual)	23,979	15.8	1,163.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 85-89 years				
...	All causes	144,284	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	42,953	29.8	---
2	Malignant neoplasms (C00-C97)	26,367	18.3	---
3	Chronic lower respiratory diseases (J40-J47)	9,295	6.4	---
4	Cerebrovascular diseases (I60-I69)	8,333	5.8	---
5	Alzheimer's disease (G30)	6,989	4.8	---
6	Influenza and pneumonia (J09-J18)	4,162	2.9	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,895	2.7	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,822	2.6	---
9	Diabetes mellitus (E10-E14)	3,241	2.2	---
10	Parkinson's disease (G20-G21)	2,994	2.1	---
11	Pneumonitis due to solids and liquids (J69)	1,940	1.3	---
12	Septicemia (A40-A41)	1,908	1.3	---
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,450	1.0	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,190	0.8	---
15	Aortic aneurysm and dissection (I71)	699	0.5	---
...	All other causes (Residual)	25,046	17.4	---
White, male, 90-94 years				
...	All causes	81,806	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	27,257	33.3	---
2	Malignant neoplasms (C00-C97)	10,741	13.1	---
3	Cerebrovascular diseases (I60-I69)	4,835	5.9	---
4	Alzheimer's disease (G30)	4,659	5.7	---
5	Chronic lower respiratory diseases (J40-J47)	4,152	5.1	---
6	Influenza and pneumonia (J09-J18)	2,948	3.6	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,315	2.8	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,188	2.7	---
9	Diabetes mellitus (E10-E14)	1,508	1.8	---
10	Pneumonitis due to solids and liquids (J69)	1,387	1.7	---
11	Parkinson's disease (G20-G21)	1,219	1.5	---
12	Septicemia (A40-A41)	957	1.2	---
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	953	1.2	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	578	0.7	---
15	Atherosclerosis (I70)	468	0.6	---
...	All other causes (Residual)	15,641	19.1	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 95-99 years				
...	All causes	23,751	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,575	36.1	---
2	Malignant neoplasms (C00-C97)	2,363	9.9	---
3	Alzheimer's disease (G30)	1,336	5.6	---
4	Cerebrovascular diseases (I60-I69)	1,291	5.4	---
5	Chronic lower respiratory diseases (J40-J47)	1,021	4.3	---
6	Influenza and pneumonia (J09-J18)	998	4.2	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	695	2.9	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	670	2.8	---
9	Pneumonitis due to solids and liquids (J69)	431	1.8	---
10	Diabetes mellitus (E10-E14)	372	1.6	---
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	353	1.5	---
12	Septicemia (A40-A41)	255	1.1	---
13	Parkinson's disease (G20-G21)	227	1.0	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	153	0.6	---
15	Atherosclerosis (I70)	149	0.6	---
...	All other causes (Residual)	4,862	20.5	---
White, male, 100 years and over				
...	All causes	3,117	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,163	37.3	---
2	Malignant neoplasms (C00-C97)	211	6.8	---
3	Influenza and pneumonia (J09-J18)	198	6.4	---
4	Cerebrovascular diseases (I60-I69)	149	4.8	---
5	Alzheimer's disease (G30)	132	4.2	---
6	Chronic lower respiratory diseases (J40-J47)	127	4.1	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	99	3.2	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	83	2.7	---
9	Pneumonitis due to solids and liquids (J69)	63	2.0	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	54	1.7	---
11	Atherosclerosis (I70)	30	1.0	---
12	Septicemia (A40-A41)	29	0.9	---
13	Diabetes mellitus (E10-E14)	25	0.8	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	0.6	---
15	Anemias (D50-D64)	14	0.4	---
...	All other causes (Residual)	721	23.1	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
White, female, all ages				
...	All causes	1,063,235	100.0	857.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	249,898	23.5	201.5
2	Malignant neoplasms (C00-C97)	233,414	22.0	188.2
3	Chronic lower respiratory diseases (J40-J47)	67,544	6.4	54.5
4	Cerebrovascular diseases (I60-I69)	65,695	6.2	53.0
5	Alzheimer's disease (G30)	53,486	5.0	43.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	39,585	3.7	31.9
7	Diabetes mellitus (E10-E14)	25,764	2.4	20.8
8	Influenza and pneumonia (J09-J18)	23,058	2.2	18.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20,033	1.9	16.2
10	Septicemia (A40-A41)	15,009	1.4	12.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	12,331	1.2	9.9
12	Chronic liver disease and cirrhosis (K70,K73-K74)	9,662	0.9	7.8
13	Parkinson's disease (G20-G21)	8,619	0.8	6.9
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7,268	0.7	5.9
15	Pneumonitis due to solids and liquids (J69)	6,971	0.7	5.6
...	All other causes (Residual)	224,898	21.2	181.3
White, female, 1-4 years				
...	All causes	1,297	100.0	21.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	401	30.9	6.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	188	14.5	3.1
3	Malignant neoplasms (C00-C97)	120	9.3	2.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	84	6.5	1.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	42	3.2	0.7
6	Influenza and pneumonia (J09-J18)	28	2.2	0.5
7	Septicemia (A40-A41)	24	1.9	0.4
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	1.5	0.3
9	Certain conditions originating in the perinatal period (P00-P96)	15	1.2	*
10	Cerebrovascular diseases (I60-I69)	12	0.9	*
11	Chronic lower respiratory diseases (J40-J47)	9	0.7	*
12	Meningococcal infection (A39)	7	0.5	*
13	Anemias (D50-D64)	6	0.5	*
14	Meningitis (G00,G03)	5	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.3	*
...	All other causes (Residual)	328	25.3	5.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 5-9 years				
...	All causes	734	100.0	9.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	209	28.5	2.8
2	Malignant neoplasms (C00-C97)	148	20.2	2.0
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	66	9.0	0.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	33	4.5	0.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	25	3.4	0.3
6	Influenza and pneumonia (J09-J18)	18	2.5	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	15	2.0	*
8	Cerebrovascular diseases (I60-I69)	12	1.6	*
9	Chronic lower respiratory diseases (J40-J47)	11	1.5	*
10	Septicemia (A40-A41)	8	1.1	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.5	*
11	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	4	0.5	*
13	Anemias (D50-D64)	3	0.4	*
13	Meningitis (G00,G03)	3	0.4	*
13	Pneumonitis due to solids and liquids (J69)	3	0.4	*
...	All other causes (Residual)	172	23.4	2.3
White, female, 10-14 years				
...	All causes	885	100.0	11.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	254	28.7	3.3
2	Malignant neoplasms (C00-C97)	173	19.5	2.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	58	6.6	0.8
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	44	5.0	0.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	37	4.2	0.5
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	25	2.8	0.3
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	2.3	0.3
8	Cerebrovascular diseases (I60-I69)	13	1.5	*
8	Influenza and pneumonia (J09-J18)	13	1.5	*
10	Chronic lower respiratory diseases (J40-J47)	11	1.2	*
11	Septicemia (A40-A41)	10	1.1	*
12	Diabetes mellitus (E10-E14)	7	0.8	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.8	*
14	Anemias (D50-D64)	3	0.3	*
14	Meningitis (G00,G03)	3	0.3	*
14	Pneumonitis due to solids and liquids (J69)	3	0.3	*
14	Diseases of appendix (K35-K38)	3	0.3	*
14	Certain conditions originating in the perinatal period (P00-P96)	3	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.3	*
...	All other causes (Residual)	195	22.0	2.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 15-19 years				
...	All causes	2,298	100.0	28.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,130	49.2	13.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	277	12.1	3.4
3	Malignant neoplasms (C00-C97)	178	7.7	2.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	120	5.2	1.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	62	2.7	0.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	58	2.5	0.7
7	Pregnancy, childbirth and the puerperium (O00-O99)	29	1.3	0.4
8	Cerebrovascular diseases (I60-I69)	20	0.9	0.2
9	Septicemia (A40-A41)	17	0.7	*
9	Diabetes mellitus (E10-E14)	17	0.7	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	0.6	*
11	Chronic lower respiratory diseases (J40-J47)	13	0.6	*
13	Influenza and pneumonia (J09-J18)	11	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.3	*
15	Meningitis (G00,G03)	5	0.2	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.2	*
...	All other causes (Residual)	335	14.6	4.1
White, female, 20-24 years				
...	All causes	3,550	100.0	44.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,608	45.3	20.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	399	11.2	5.0
3	Malignant neoplasms (C00-C97)	277	7.8	3.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	205	5.8	2.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	106	3.0	1.3
6	Pregnancy, childbirth and the puerperium (O00-O99)	77	2.2	1.0
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	62	1.7	0.8
8	Influenza and pneumonia (J09-J18)	47	1.3	0.6
9	Diabetes mellitus (E10-E14)	33	0.9	0.4
10	Septicemia (A40-A41)	32	0.9	0.4
11	Cerebrovascular diseases (I60-I69)	29	0.8	0.4
12	Chronic lower respiratory diseases (J40-J47)	20	0.6	0.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	0.5	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	0.4	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	13	0.4	*
...	All other causes (Residual)	611	17.2	7.6

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 25-29 years				
...	All causes	4,280	100.0	53.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,575	36.8	19.7
2	Malignant neoplasms (C00-C97)	488	11.4	6.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	462	10.8	5.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	220	5.1	2.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	214	5.0	2.7
6	Pregnancy, childbirth and the puerperium (O00-O99)	121	2.8	1.5
7	Diabetes mellitus (E10-E14)	64	1.5	0.8
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	63	1.5	0.8
9	Cerebrovascular diseases (I60-I69)	58	1.4	0.7
10	Septicemia (A40-A41)	51	1.2	0.6
10	Influenza and pneumonia (J09-J18)	51	1.2	0.6
12	Chronic liver disease and cirrhosis (K70,K73-K74)	33	0.8	0.4
13	Chronic lower respiratory diseases (J40-J47)	29	0.7	0.4
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	29	0.7	0.4
15	Human immunodeficiency virus (HIV) disease (B20-B24)	20	0.5	0.3
...	All other causes (Residual)	802	18.7	10.0
White, female, 30-34 years				
...	All causes	5,262	100.0	69.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,588	30.2	21.0
2	Malignant neoplasms (C00-C97)	888	16.9	11.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	479	9.1	6.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	413	7.8	5.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	171	3.2	2.3
6	Cerebrovascular diseases (I60-I69)	118	2.2	1.6
7	Pregnancy, childbirth and the puerperium (O00-O99)	99	1.9	1.3
8	Diabetes mellitus (E10-E14)	96	1.8	1.3
9	Chronic liver disease and cirrhosis (K70,K73-K74)	94	1.8	1.2
10	Influenza and pneumonia (J09-J18)	73	1.4	1.0
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	64	1.2	0.8
12	Septicemia (A40-A41)	53	1.0	0.7
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	43	0.8	0.6
14	Human immunodeficiency virus (HIV) disease (B20-B24)	42	0.8	0.6
15	Chronic lower respiratory diseases (J40-J47)	39	0.7	0.5
...	All other causes (Residual)	1,002	19.0	13.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 35-39 years				
...	All causes	7,544	100.0	97.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,823	24.2	23.5
2	Malignant neoplasms (C00-C97)	1,601	21.2	20.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	719	9.5	9.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	624	8.3	8.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	201	2.7	2.6
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	181	2.4	2.3
7	Diabetes mellitus (E10-E14)	171	2.3	2.2
8	Cerebrovascular diseases (I60-I69)	160	2.1	2.1
9	Influenza and pneumonia (J09-J18)	99	1.3	1.3
10	Septicemia (A40-A41)	92	1.2	1.2
11	Chronic lower respiratory diseases (J40-J47)	87	1.2	1.1
12	Pregnancy, childbirth and the puerperium (O00-O99)	82	1.1	1.1
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	80	1.1	1.0
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	79	1.0	1.0
15	Human immunodeficiency virus (HIV) disease (B20-B24)	69	0.9	0.9
...	All other causes (Residual)	1,476	19.6	19.0
White, female, 40-44 years				
...	All causes	12,030	100.0	146.9
1	Malignant neoplasms (C00-C97)	3,390	28.2	41.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,137	17.8	26.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,377	11.4	16.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	731	6.1	8.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	453	3.8	5.5
6	Cerebrovascular diseases (I60-I69)	392	3.3	4.8
7	Diabetes mellitus (E10-E14)	311	2.6	3.8
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	202	1.7	2.5
9	Chronic lower respiratory diseases (J40-J47)	189	1.6	2.3
10	Influenza and pneumonia (J09-J18)	164	1.4	2.0
11	Septicemia (A40-A41)	156	1.3	1.9
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	111	0.9	1.4
13	Human immunodeficiency virus (HIV) disease (B20-B24)	105	0.9	1.3
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	81	0.7	1.0
15	Viral hepatitis (B15-B19)	70	0.6	0.9
...	All other causes (Residual)	2,161	18.0	26.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 45-49 years				
...	All causes	21,660	100.0	236.7
1	Malignant neoplasms (C00-C97)	7,131	32.9	77.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,905	13.4	31.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,822	13.0	30.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	1,028	4.7	11.2
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	922	4.3	10.1
6	Cerebrovascular diseases (I60-I69)	653	3.0	7.1
7	Chronic lower respiratory diseases (J40-J47)	640	3.0	7.0
8	Diabetes mellitus (E10-E14)	528	2.4	5.8
9	Septicemia (A40-A41)	298	1.4	3.3
10	Influenza and pneumonia (J09-J18)	266	1.2	2.9
11	Viral hepatitis (B15-B19)	207	1.0	2.3
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	204	0.9	2.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	191	0.9	2.1
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	118	0.5	1.3
15	Human immunodeficiency virus (HIV) disease (B20-B24)	117	0.5	1.3
...	All other causes (Residual)	3,630	16.8	39.7
White, female, 50-54 years				
...	All causes	32,372	100.0	353.5
1	Malignant neoplasms (C00-C97)	12,378	38.2	135.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,553	14.1	49.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,745	8.5	30.0
4	Chronic lower respiratory diseases (J40-J47)	1,360	4.2	14.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,280	4.0	14.0
6	Cerebrovascular diseases (I60-I69)	1,017	3.1	11.1
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	990	3.1	10.8
8	Diabetes mellitus (E10-E14)	868	2.7	9.5
9	Septicemia (A40-A41)	450	1.4	4.9
10	Influenza and pneumonia (J09-J18)	395	1.2	4.3
11	Viral hepatitis (B15-B19)	384	1.2	4.2
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	343	1.1	3.7
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	170	0.5	1.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	160	0.5	1.7
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	155	0.5	1.7
...	All other causes (Residual)	5,124	15.8	55.9

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 55-59 years				
...	All causes	41,148	100.0	496.1
1	Malignant neoplasms (C00-C97)	16,805	40.8	202.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,226	15.1	75.1
3	Chronic lower respiratory diseases (J40-J47)	2,198	5.3	26.5
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,122	5.2	25.6
5	Diabetes mellitus (E10-E14)	1,328	3.2	16.0
6	Cerebrovascular diseases (I60-I69)	1,268	3.1	15.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,239	3.0	14.9
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	850	2.1	10.2
9	Septicemia (A40-A41)	687	1.7	8.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	574	1.4	6.9
11	Influenza and pneumonia (J09-J18)	505	1.2	6.1
12	Viral hepatitis (B15-B19)	455	1.1	5.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	239	0.6	2.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	186	0.5	2.2
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	178	0.4	2.1
...	All other causes (Residual)	6,288	15.3	75.8
White, female, 60-64 years				
...	All causes	55,139	100.0	756.2
1	Malignant neoplasms (C00-C97)	22,986	41.7	315.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,083	16.5	124.6
3	Chronic lower respiratory diseases (J40-J47)	3,874	7.0	53.1
4	Cerebrovascular diseases (I60-I69)	1,945	3.5	26.7
5	Diabetes mellitus (E10-E14)	1,917	3.5	26.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,639	3.0	22.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,178	2.1	16.2
8	Septicemia (A40-A41)	890	1.6	12.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	856	1.6	11.7
10	Influenza and pneumonia (J09-J18)	698	1.3	9.6
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	579	1.1	7.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	353	0.6	4.8
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	304	0.6	4.2
14	Alzheimer's disease (G30)	287	0.5	3.9
15	Viral hepatitis (B15-B19)	263	0.5	3.6
...	All other causes (Residual)	8,287	15.0	113.6

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 65-69 years				
...	All causes	66,698	100.0	1,197.3
1	Malignant neoplasms (C00-C97)	26,757	40.1	480.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,437	17.1	205.3
3	Chronic lower respiratory diseases (J40-J47)	6,133	9.2	110.1
4	Cerebrovascular diseases (I60-I69)	2,593	3.9	46.5
5	Diabetes mellitus (E10-E14)	2,201	3.3	39.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,425	2.1	25.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,180	1.8	21.2
8	Septicemia (A40-A41)	1,149	1.7	20.6
9	Chronic liver disease and cirrhosis (K70,K73-K74)	1,014	1.5	18.2
10	Influenza and pneumonia (J09-J18)	895	1.3	16.1
11	Alzheimer's disease (G30)	605	0.9	10.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	494	0.7	8.9
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	397	0.6	7.1
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	339	0.5	6.1
15	Pneumonitis due to solids and liquids (J69)	263	0.4	4.7
...	All other causes (Residual)	9,816	14.7	176.2
White, female, 70-74 years				
...	All causes	81,486	100.0	1,905.9
1	Malignant neoplasms (C00-C97)	28,749	35.3	672.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,992	18.4	350.6
3	Chronic lower respiratory diseases (J40-J47)	8,290	10.2	193.9
4	Cerebrovascular diseases (I60-I69)	3,981	4.9	93.1
5	Diabetes mellitus (E10-E14)	2,596	3.2	60.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,656	2.0	38.7
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,582	1.9	37.0
8	Alzheimer's disease (G30)	1,522	1.9	35.6
9	Septicemia (A40-A41)	1,345	1.7	31.5
10	Influenza and pneumonia (J09-J18)	1,278	1.6	29.9
11	Chronic liver disease and cirrhosis (K70,K73-K74)	942	1.2	22.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	685	0.8	16.0
13	Parkinson's disease (G20-G21)	559	0.7	13.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	550	0.7	12.9
15	Pneumonitis due to solids and liquids (J69)	382	0.5	8.9
...	All other causes (Residual)	12,377	15.2	289.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 75-79 years				
...	All causes	112,452	100.0	3,154.8
1	Malignant neoplasms (C00-C97)	31,459	28.0	882.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	23,395	20.8	656.3
3	Chronic lower respiratory diseases (J40-J47)	10,891	9.7	305.5
4	Cerebrovascular diseases (I60-I69)	6,830	6.1	191.6
5	Alzheimer's disease (G30)	4,258	3.8	119.5
6	Diabetes mellitus (E10-E14)	3,247	2.9	91.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,310	2.1	64.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,290	2.0	64.2
9	Influenza and pneumonia (J09-J18)	2,041	1.8	57.3
10	Septicemia (A40-A41)	1,858	1.7	52.1
11	Parkinson's disease (G20-G21)	1,297	1.2	36.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,172	1.0	32.9
13	Chronic liver disease and cirrhosis (K70,K73-K74)	832	0.7	23.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	797	0.7	22.4
15	Pneumonitis due to solids and liquids (J69)	660	0.6	18.5
...	All other causes (Residual)	19,115	17.0	536.3
White, female, 80-84 years				
...	All causes	162,513	100.0	5,351.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	38,760	23.9	1,276.4
2	Malignant neoplasms (C00-C97)	33,801	20.8	1,113.1
3	Chronic lower respiratory diseases (J40-J47)	12,734	7.8	419.3
4	Cerebrovascular diseases (I60-I69)	11,664	7.2	384.1
5	Alzheimer's disease (G30)	9,443	5.8	311.0
6	Diabetes mellitus (E10-E14)	4,254	2.6	140.1
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,647	2.2	120.1
8	Influenza and pneumonia (J09-J18)	3,615	2.2	119.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,552	2.2	117.0
10	Septicemia (A40-A41)	2,512	1.5	82.7
11	Parkinson's disease (G20-G21)	2,148	1.3	70.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,946	1.2	64.1
13	Pneumonitis due to solids and liquids (J69)	1,215	0.7	40.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,154	0.7	38.0
15	Aortic aneurysm and dissection (I71)	839	0.5	27.6
...	All other causes (Residual)	31,229	19.2	1,028.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 85-89 years				
...	All causes	200,860	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	55,893	27.8	---
2	Malignant neoplasms (C00-C97)	27,893	13.9	---
3	Cerebrovascular diseases (I60-I69)	15,974	8.0	---
4	Alzheimer's disease (G30)	15,480	7.7	---
5	Chronic lower respiratory diseases (J40-J47)	11,758	5.9	---
6	Influenza and pneumonia (J09-J18)	5,113	2.5	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,649	2.3	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,345	2.2	---
9	Diabetes mellitus (E10-E14)	4,323	2.2	---
10	Septicemia (A40-A41)	2,786	1.4	---
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,749	1.4	---
12	Parkinson's disease (G20-G21)	2,443	1.2	---
13	Pneumonitis due to solids and liquids (J69)	1,665	0.8	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,246	0.6	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1,037	0.5	---
...	All other causes (Residual)	43,506	21.7	---
White, female, 90-94 years				
...	All causes	156,346	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	49,286	31.5	---
2	Alzheimer's disease (G30)	13,833	8.8	---
3	Malignant neoplasms (C00-C97)	13,576	8.7	---
4	Cerebrovascular diseases (I60-I69)	12,537	8.0	---
5	Chronic lower respiratory diseases (J40-J47)	6,633	4.2	---
6	Influenza and pneumonia (J09-J18)	4,609	2.9	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,630	2.3	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,150	2.0	---
9	Diabetes mellitus (E10-E14)	2,695	1.7	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,545	1.6	---
11	Septicemia (A40-A41)	1,745	1.1	---
12	Pneumonitis due to solids and liquids (J69)	1,441	0.9	---
13	Parkinson's disease (G20-G21)	1,369	0.9	---
14	Atherosclerosis (I70)	968	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	831	0.5	---
...	All other causes (Residual)	37,498	24.0	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 95-99 years				
...	All causes	71,517	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	24,441	34.2	---
2	Alzheimer's disease (G30)	6,539	9.1	---
3	Cerebrovascular diseases (I60-I69)	5,373	7.5	---
4	Malignant neoplasms (C00-C97)	4,035	5.6	---
5	Influenza and pneumonia (J09-J18)	2,412	3.4	---
6	Chronic lower respiratory diseases (J40-J47)	2,259	3.2	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,665	2.3	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,392	1.9	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,284	1.8	---
10	Diabetes mellitus (E10-E14)	954	1.3	---
11	Septicemia (A40-A41)	679	0.9	---
12	Pneumonitis due to solids and liquids (J69)	656	0.9	---
13	Atherosclerosis (I70)	558	0.8	---
14	Parkinson's disease (G20-G21)	363	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	289	0.4	---
...	All other causes (Residual)	18,618	26.0	---
White, female, 100 years and over				
...	All causes	16,052	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,831	36.3	---
2	Alzheimer's disease (G30)	1,339	8.3	---
3	Cerebrovascular diseases (I60-I69)	1,010	6.3	---
4	Influenza and pneumonia (J09-J18)	682	4.2	---
5	Malignant neoplasms (C00-C97)	559	3.5	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	389	2.4	---
7	Chronic lower respiratory diseases (J40-J47)	355	2.2	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	339	2.1	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	266	1.7	---
10	Atherosclerosis (I70)	165	1.0	---
11	Diabetes mellitus (E10-E14)	151	0.9	---
11	Pneumonitis due to solids and liquids (J69)	151	0.9	---
13	Septicemia (A40-A41)	122	0.8	---
14	Anemias (D50-D64)	68	0.4	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	54	0.3	---
...	All other causes (Residual)	4,571	28.5	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, all ages				
...	All causes	286,959	100.0	682.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	69,083	24.1	164.2
2	Malignant neoplasms (C00-C97)	65,930	23.0	156.7
3	Cerebrovascular diseases (I60-I69)	15,965	5.6	38.0
4	Diabetes mellitus (E10-E14)	12,126	4.2	28.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,069	4.2	28.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,841	3.1	21.0
7	Chronic lower respiratory diseases (J40-J47)	8,715	3.0	20.7
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7,818	2.7	18.6
9	Septicemia (A40-A41)	6,001	2.1	14.3
10	Alzheimer's disease (G30)	5,220	1.8	12.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,116	1.8	12.2
12	Influenza and pneumonia (J09-J18)	4,936	1.7	11.7
13	Human immunodeficiency virus (HIV) disease (B20-B24)	4,662	1.6	11.1
14	Certain conditions originating in the perinatal period (P00-P96)	4,031	1.4	9.6
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,635	0.9	6.3
...	All other causes (Residual)	53,811	18.8	127.9
Black, both sexes, 1-4 years				
...	All causes	1,041	100.0	38.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	267	25.6	9.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	156	15.0	5.7
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	127	12.2	4.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	51	4.9	1.9
5	Malignant neoplasms (C00-C97)	48	4.6	1.8
6	Influenza and pneumonia (J09-J18)	25	2.4	0.9
7	Chronic lower respiratory diseases (J40-J47)	21	2.0	0.8
8	Certain conditions originating in the perinatal period (P00-P96)	16	1.5	*
9	Cerebrovascular diseases (I60-I69)	15	1.4	*
10	Anemias (D50-D64)	14	1.3	*
11	Septicemia (A40-A41)	12	1.2	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	1.0	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.5	*
14	Meningitis (G00,G03)	4	0.4	*
14	Pneumonitis due to solids and liquids (J69)	4	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.4	*
...	All other causes (Residual)	262	25.2	9.6

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 5-9 years				
...	All causes	494	100.0	15.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	154	31.2	4.7
2	Malignant neoplasms (C00-C97)	65	13.2	2.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	36	7.3	1.1
4	Chronic lower respiratory diseases (J40-J47)	29	5.9	0.9
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	25	5.1	0.8
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	21	4.3	0.6
7	Cerebrovascular diseases (I60-I69)	15	3.0	*
8	Septicemia (A40-A41)	8	1.6	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	1.4	*
9	Anemias (D50-D64)	7	1.4	*
11	Influenza and pneumonia (J09-J18)	4	0.8	*
12	Certain conditions originating in the perinatal period (P00-P96)	3	0.6	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.6	*
14	Diabetes mellitus (E10-E14)	2	0.4	*
15	Tuberculosis (A16-A19)	1	0.2	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.2	*
15	Diseases of appendix (K35-K38)	1	0.2	*
15	Hernia (K40-K46)	1	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
...	All other causes (Residual)	110	22.3	3.3
Black, both sexes, 10-14 years				
...	All causes	651	100.0	19.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	165	25.3	4.8
2	Malignant neoplasms (C00-C97)	81	12.4	2.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	65	10.0	1.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	44	6.8	1.3
5	Chronic lower respiratory diseases (J40-J47)	43	6.6	1.3
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	40	6.1	1.2
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	38	5.8	1.1
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	1.5	*
9	Influenza and pneumonia (J09-J18)	8	1.2	*
10	Septicemia (A40-A41)	7	1.1	*
10	Anemias (D50-D64)	7	1.1	*
10	Cerebrovascular diseases (I60-I69)	7	1.1	*
13	Diabetes mellitus (E10-E14)	5	0.8	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.6	*
15	Certain conditions originating in the perinatal period (P00-P96)	3	0.5	*
...	All other causes (Residual)	124	19.0	3.6

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 15-19 years				
...	All causes	2,531	100.0	67.0
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,117	44.1	29.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	582	23.0	15.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	152	6.0	4.0
4	Malignant neoplasms (C00-C97)	106	4.2	2.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	104	4.1	2.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	50	2.0	1.3
7	Anemias (D50-D64)	28	1.1	0.7
9	Chronic lower respiratory diseases (J40-J47)	28	1.1	0.7
9	Cerebrovascular diseases (I60-I69)	24	0.9	0.6
10	Human immunodeficiency virus (HIV) disease (B20-B24)	17	0.7	*
11	Septicemia (A40-A41)	11	0.4	*
11	Influenza and pneumonia (J09-J18)	11	0.4	*
11	Pregnancy, childbirth and the puerperium (O00-O99)	11	0.4	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.4	*
14	Diabetes mellitus (E10-E14)	9	0.4	*
...	All other causes (Residual)	272	10.7	7.2
Black, both sexes, 20-24 years				
...	All causes	4,144	100.0	122.4
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,699	41.0	50.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	875	21.1	25.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	317	7.6	9.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	232	5.6	6.9
5	Malignant neoplasms (C00-C97)	161	3.9	4.8
6	Human immunodeficiency virus (HIV) disease (B20-B24)	94	2.3	2.8
7	Anemias (D50-D64)	51	1.2	1.5
8	Diabetes mellitus (E10-E14)	42	1.0	1.2
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	41	1.0	1.2
10	Pregnancy, childbirth and the puerperium (O00-O99)	40	1.0	1.2
11	Chronic lower respiratory diseases (J40-J47)	34	0.8	1.0
12	Septicemia (A40-A41)	30	0.7	0.9
13	Influenza and pneumonia (J09-J18)	28	0.7	0.8
14	Cerebrovascular diseases (I60-I69)	25	0.6	0.7
15	Legal intervention (Y35,Y89.0)	20	0.5	0.6
...	All other causes (Residual)	455	11.0	13.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 25-29 years				
...	All causes	4,246	100.0	140.3
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,370	32.3	45.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	815	19.2	26.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	348	8.2	11.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	286	6.7	9.4
5	Malignant neoplasms (C00-C97)	218	5.1	7.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	165	3.9	5.5
7	Diabetes mellitus (E10-E14)	78	1.8	2.6
8	Pregnancy, childbirth and the puerperium (O00-O99)	58	1.4	1.9
9	Anemias (D50-D64)	55	1.3	1.8
10	Cerebrovascular diseases (I60-I69)	47	1.1	1.6
11	Influenza and pneumonia (J09-J18)	41	1.0	1.4
11	Chronic lower respiratory diseases (J40-J47)	41	1.0	1.4
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	40	0.9	1.3
14	Septicemia (A40-A41)	34	0.8	1.1
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	31	0.7	1.0
...	All other causes (Residual)	619	14.6	20.4
Black, both sexes, 30-34 years				
...	All causes	4,674	100.0	164.6
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,031	22.1	36.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	797	17.1	28.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	566	12.1	19.9
4	Malignant neoplasms (C00-C97)	371	7.9	13.1
5	Human immunodeficiency virus (HIV) disease (B20-B24)	307	6.6	10.8
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	218	4.7	7.7
7	Diabetes mellitus (E10-E14)	131	2.8	4.6
8	Cerebrovascular diseases (I60-I69)	90	1.9	3.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	64	1.4	2.3
10	Pregnancy, childbirth and the puerperium (O00-O99)	60	1.3	2.1
11	Chronic lower respiratory diseases (J40-J47)	57	1.2	2.0
12	Anemias (D50-D64)	54	1.2	1.9
13	Septicemia (A40-A41)	52	1.1	1.8
14	Influenza and pneumonia (J09-J18)	44	0.9	1.5
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	35	0.7	1.2
...	All other causes (Residual)	797	17.1	28.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 35-39 years				
...	All causes	5,810	100.0	208.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,007	17.3	36.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	769	13.2	27.6
3	Malignant neoplasms (C00-C97)	744	12.8	26.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	665	11.4	23.9
5	Human immunodeficiency virus (HIV) disease (B20-B24)	411	7.1	14.8
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	240	4.1	8.6
7	Cerebrovascular diseases (I60-I69)	198	3.4	7.1
8	Diabetes mellitus (E10-E14)	183	3.1	6.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	111	1.9	4.0
10	Chronic lower respiratory diseases (J40-J47)	87	1.5	3.1
11	Septicemia (A40-A41)	86	1.5	3.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	73	1.3	2.6
13	Anemias (D50-D64)	71	1.2	2.6
14	Chronic liver disease and cirrhosis (K70,K73-K74)	66	1.1	2.4
15	Pregnancy, childbirth and the puerperium (O00-O99)	47	0.8	1.7
...	All other causes (Residual)	1,052	18.1	37.8
Black, both sexes, 40-44 years				
...	All causes	8,206	100.0	291.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,719	20.9	61.0
2	Malignant neoplasms (C00-C97)	1,367	16.7	48.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	833	10.2	29.6
4	Human immunodeficiency virus (HIV) disease (B20-B24)	666	8.1	23.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	423	5.2	15.0
6	Cerebrovascular diseases (I60-I69)	361	4.4	12.8
7	Diabetes mellitus (E10-E14)	331	4.0	11.8
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	212	2.6	7.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	170	2.1	6.0
10	Septicemia (A40-A41)	126	1.5	4.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	114	1.4	4.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	102	1.2	3.6
13	Influenza and pneumonia (J09-J18)	98	1.2	3.5
14	Chronic lower respiratory diseases (J40-J47)	95	1.2	3.4
15	Anemias (D50-D64)	61	0.7	2.2
...	All other causes (Residual)	1,528	18.6	54.3

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 45-49 years				
...	All causes	13,581	100.0	458.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,195	23.5	107.9
2	Malignant neoplasms (C00-C97)	2,988	22.0	101.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,090	8.0	36.8
4	Human immunodeficiency virus (HIV) disease (B20-B24)	843	6.2	28.5
5	Cerebrovascular diseases (I60-I69)	660	4.9	22.3
6	Diabetes mellitus (E10-E14)	563	4.1	19.0
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	390	2.9	13.2
8	Chronic liver disease and cirrhosis (K70,K73-K74)	283	2.1	9.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	281	2.1	9.5
10	Septicemia (A40-A41)	228	1.7	7.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	218	1.6	7.4
12	Chronic lower respiratory diseases (J40-J47)	213	1.6	7.2
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	178	1.3	6.0
14	Influenza and pneumonia (J09-J18)	167	1.2	5.6
15	Viral hepatitis (B15-B19)	84	0.6	2.8
...	All other causes (Residual)	2,200	16.2	74.3
Black, both sexes, 50-54 years				
...	All causes	20,512	100.0	731.5
1	Malignant neoplasms (C00-C97)	5,573	27.2	198.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,924	24.0	175.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,215	5.9	43.3
4	Cerebrovascular diseases (I60-I69)	1,066	5.2	38.0
5	Diabetes mellitus (E10-E14)	831	4.1	29.6
6	Human immunodeficiency virus (HIV) disease (B20-B24)	792	3.9	28.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	534	2.6	19.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	519	2.5	18.5
9	Chronic lower respiratory diseases (J40-J47)	436	2.1	15.5
10	Septicemia (A40-A41)	379	1.8	13.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	314	1.5	11.2
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	303	1.5	10.8
13	Viral hepatitis (B15-B19)	230	1.1	8.2
14	Influenza and pneumonia (J09-J18)	225	1.1	8.0
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	159	0.8	5.7
...	All other causes (Residual)	3,012	14.7	107.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 55-59 years				
...	All causes	25,246	100.0	1,104.8
1	Malignant neoplasms (C00-C97)	7,662	30.3	335.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,150	24.4	269.1
3	Cerebrovascular diseases (I60-I69)	1,317	5.2	57.6
4	Diabetes mellitus (E10-E14)	1,169	4.6	51.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,040	4.1	45.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	699	2.8	30.6
7	Human immunodeficiency virus (HIV) disease (B20-B24)	646	2.6	28.3
8	Chronic lower respiratory diseases (J40-J47)	626	2.5	27.4
9	Chronic liver disease and cirrhosis (K70,K73-K74)	581	2.3	25.4
10	Septicemia (A40-A41)	518	2.1	22.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	405	1.6	17.7
12	Viral hepatitis (B15-B19)	297	1.2	13.0
13	Influenza and pneumonia (J09-J18)	280	1.1	12.3
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	212	0.8	9.3
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	126	0.5	5.5
...	All other causes (Residual)	3,518	13.9	153.9
Black, both sexes, 60-64 years				
...	All causes	26,570	100.0	1,523.4
1	Malignant neoplasms (C00-C97)	8,713	32.8	499.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,676	25.1	382.8
3	Cerebrovascular diseases (I60-I69)	1,335	5.0	76.5
4	Diabetes mellitus (E10-E14)	1,320	5.0	75.7
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	839	3.2	48.1
6	Chronic lower respiratory diseases (J40-J47)	837	3.2	48.0
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	729	2.7	41.8
8	Septicemia (A40-A41)	587	2.2	33.7
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	479	1.8	27.5
10	Chronic liver disease and cirrhosis (K70,K73-K74)	445	1.7	25.5
11	Human immunodeficiency virus (HIV) disease (B20-B24)	383	1.4	22.0
12	Influenza and pneumonia (J09-J18)	319	1.2	18.3
13	Viral hepatitis (B15-B19)	205	0.8	11.8
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	111	0.4	6.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	107	0.4	6.1
...	All other causes (Residual)	3,485	13.1	199.8

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 65-69 years				
...	All causes	25,781	100.0	2,148.8
1	Malignant neoplasms (C00-C97)	8,439	32.7	703.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,466	25.1	538.9
3	Cerebrovascular diseases (I60-I69)	1,469	5.7	122.4
4	Diabetes mellitus (E10-E14)	1,376	5.3	114.7
5	Chronic lower respiratory diseases (J40-J47)	1,001	3.9	83.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	889	3.4	74.1
7	Septicemia (A40-A41)	629	2.4	52.4
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	495	1.9	41.3
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	480	1.9	40.0
10	Influenza and pneumonia (J09-J18)	397	1.5	33.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	238	0.9	19.8
12	Human immunodeficiency virus (HIV) disease (B20-B24)	162	0.6	13.5
13	Alzheimer's disease (G30)	108	0.4	9.0
14	Pneumonitis due to solids and liquids (J69)	103	0.4	8.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	94	0.4	7.8
...	All other causes (Residual)	3,435	13.3	286.3
Black, both sexes, 70-74 years				
...	All causes	26,710	100.0	3,041.6
1	Malignant neoplasms (C00-C97)	8,188	30.7	932.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,718	25.2	765.0
3	Cerebrovascular diseases (I60-I69)	1,652	6.2	188.1
4	Diabetes mellitus (E10-E14)	1,405	5.3	160.0
5	Chronic lower respiratory diseases (J40-J47)	1,189	4.5	135.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,072	4.0	122.1
7	Septicemia (A40-A41)	640	2.4	72.9
8	Influenza and pneumonia (J09-J18)	473	1.8	53.9
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	471	1.8	53.6
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	420	1.6	47.8
11	Alzheimer's disease (G30)	297	1.1	33.8
12	Chronic liver disease and cirrhosis (K70,K73-K74)	147	0.6	16.7
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	139	0.5	15.8
14	Pneumonitis due to solids and liquids (J69)	131	0.5	14.9
15	Human immunodeficiency virus (HIV) disease (B20-B24)	99	0.4	11.3
...	All other causes (Residual)	3,669	13.7	417.8

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 75-79 years				
...	All causes	28,238	100.0	4,450.5
1	Malignant neoplasms (C00-C97)	7,551	26.7	1,190.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,471	26.5	1,177.5
3	Cerebrovascular diseases (I60-I69)	1,850	6.6	291.6
4	Diabetes mellitus (E10-E14)	1,511	5.4	238.1
5	Chronic lower respiratory diseases (J40-J47)	1,137	4.0	179.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,081	3.8	170.4
7	Septicemia (A40-A41)	692	2.5	109.1
8	Alzheimer's disease (G30)	608	2.2	95.8
9	Influenza and pneumonia (J09-J18)	526	1.9	82.9
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	513	1.8	80.9
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	427	1.5	67.3
12	Pneumonitis due to solids and liquids (J69)	165	0.6	26.0
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	147	0.5	23.2
14	Parkinson's disease (G20-G21)	146	0.5	23.0
15	Aortic aneurysm and dissection (I71)	100	0.4	15.8
...	All other causes (Residual)	4,313	15.3	679.8
Black, both sexes, 80-84 years				
...	All causes	29,305	100.0	6,710.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,970	27.2	1,825.0
2	Malignant neoplasms (C00-C97)	6,489	22.1	1,485.9
3	Cerebrovascular diseases (I60-I69)	2,070	7.1	474.0
4	Diabetes mellitus (E10-E14)	1,316	4.5	301.3
5	Chronic lower respiratory diseases (J40-J47)	1,193	4.1	273.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,141	3.9	261.3
7	Alzheimer's disease (G30)	1,056	3.6	241.8
8	Septicemia (A40-A41)	730	2.5	167.2
9	Influenza and pneumonia (J09-J18)	704	2.4	161.2
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	619	2.1	141.7
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	395	1.3	90.4
12	Pneumonitis due to solids and liquids (J69)	238	0.8	54.5
13	Parkinson's disease (G20-G21)	181	0.6	41.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	141	0.5	32.3
15	Aortic aneurysm and dissection (I71)	103	0.4	23.6
...	All other causes (Residual)	4,959	16.9	1,135.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 85-89 years				
...	All causes	26,078	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,413	28.4	---
2	Malignant neoplasms (C00-C97)	4,454	17.1	---
3	Cerebrovascular diseases (I60-I69)	1,869	7.2	---
4	Alzheimer's disease (G30)	1,341	5.1	---
5	Diabetes mellitus (E10-E14)	1,049	4.0	---
6	Chronic lower respiratory diseases (J40-J47)	968	3.7	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	965	3.7	---
8	Influenza and pneumonia (J09-J18)	674	2.6	---
9	Septicemia (A40-A41)	644	2.5	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	615	2.4	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	324	1.2	---
12	Pneumonitis due to solids and liquids (J69)	228	0.9	---
13	Parkinson's disease (G20-G21)	191	0.7	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	122	0.5	---
15	Atherosclerosis (I70)	92	0.4	---
...	All other causes (Residual)	5,129	19.7	---
Black, both sexes, 90-94 years				
...	All causes	16,170	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,854	30.0	---
2	Malignant neoplasms (C00-C97)	1,971	12.2	---
3	Cerebrovascular diseases (I60-I69)	1,189	7.4	---
4	Alzheimer's disease (G30)	1,067	6.6	---
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	605	3.7	---
6	Diabetes mellitus (E10-E14)	539	3.3	---
7	Influenza and pneumonia (J09-J18)	479	3.0	---
8	Chronic lower respiratory diseases (J40-J47)	466	2.9	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	426	2.6	---
10	Septicemia (A40-A41)	331	2.0	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	211	1.3	---
12	Pneumonitis due to solids and liquids (J69)	134	0.8	---
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	93	0.6	---
14	Atherosclerosis (I70)	69	0.4	---
15	Parkinson's disease (G20-G21)	66	0.4	---
...	All other causes (Residual)	3,670	22.7	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 95-99 years				
...	All causes	7,367	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,324	31.5	---
2	Malignant neoplasms (C00-C97)	624	8.5	---
3	Alzheimer's disease (G30)	528	7.2	---
3	Cerebrovascular diseases (I60-I69)	528	7.2	---
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	250	3.4	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	248	3.4	---
7	Influenza and pneumonia (J09-J18)	228	3.1	---
8	Diabetes mellitus (E10-E14)	214	2.9	---
9	Chronic lower respiratory diseases (J40-J47)	164	2.2	---
10	Septicemia (A40-A41)	133	1.8	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	95	1.3	---
12	Pneumonitis due to solids and liquids (J69)	74	1.0	---
13	Atherosclerosis (I70)	48	0.7	---
14	Anemias (D50-D64)	31	0.4	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	30	0.4	---
15	Nutritional deficiencies (E40-E64)	30	0.4	---
...	All other causes (Residual)	1,818	24.7	---
Black, both sexes, 100 years and over				
...	All causes	2,180	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	733	33.6	---
2	Alzheimer's disease (G30)	154	7.1	---
3	Cerebrovascular diseases (I60-I69)	130	6.0	---
4	Malignant neoplasms (C00-C97)	108	5.0	---
5	Influenza and pneumonia (J09-J18)	86	3.9	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	85	3.9	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	58	2.7	---
8	Diabetes mellitus (E10-E14)	48	2.2	---
9	Chronic lower respiratory diseases (J40-J47)	41	1.9	---
10	Septicemia (A40-A41)	34	1.6	---
11	Atherosclerosis (I70)	21	1.0	---
12	Pneumonitis due to solids and liquids (J69)	20	0.9	---
13	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15	0.7	---
14	Nutritional deficiencies (E40-E64)	10	0.5	---
15	Anemias (D50-D64)	7	0.3	---
...	All other causes (Residual)	630	28.9	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, all ages				
...	All causes	145,802	100.0	725.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	35,089	24.1	174.6
2	Malignant neoplasms (C00-C97)	33,967	23.3	169.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,074	5.5	40.2
4	Cerebrovascular diseases (I60-I69)	6,938	4.8	34.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6,704	4.6	33.4
6	Diabetes mellitus (E10-E14)	5,640	3.9	28.1
7	Chronic lower respiratory diseases (J40-J47)	4,532	3.1	22.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,016	2.8	20.0
9	Human immunodeficiency virus (HIV) disease (B20-B24)	3,047	2.1	15.2
10	Septicemia (A40-A41)	2,691	1.8	13.4
11	Influenza and pneumonia (J09-J18)	2,380	1.6	11.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,218	1.5	11.0
13	Certain conditions originating in the perinatal period (P00-P96)	2,216	1.5	11.0
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,755	1.2	8.7
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,715	1.2	8.5
...	All other causes (Residual)	24,820	17.0	123.5
Black, male, 1-4 years				
...	All causes	595	100.0	42.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	170	28.6	12.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	93	15.6	6.7
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	57	9.6	4.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	29	4.9	2.1
5	Malignant neoplasms (C00-C97)	27	4.5	1.9
6	Influenza and pneumonia (J09-J18)	15	2.5	*
7	Chronic lower respiratory diseases (J40-J47)	11	1.8	*
8	Anemias (D50-D64)	9	1.5	*
8	Certain conditions originating in the perinatal period (P00-P96)	9	1.5	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.0	*
11	Septicemia (A40-A41)	5	0.8	*
11	Cerebrovascular diseases (I60-I69)	5	0.8	*
13	Meningococcal infection (A39)	3	0.5	*
13	Pneumonitis due to solids and liquids (J69)	3	0.5	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.5	*
...	All other causes (Residual)	150	25.2	10.8

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 5-9 years				
...	All causes	283	100.0	16.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	98	34.6	5.8
2	Malignant neoplasms (C00-C97)	25	8.8	1.5
3	Chronic lower respiratory diseases (J40-J47)	24	8.5	1.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	22	7.8	1.3
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	17	6.0	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	15	5.3	*
7	Cerebrovascular diseases (I60-I69)	9	3.2	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	1.8	*
9	Anemias (D50-D64)	4	1.4	*
10	Septicemia (A40-A41)	3	1.1	*
11	Influenza and pneumonia (J09-J18)	2	0.7	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.4	*
12	Hernia (K40-K46)	1	0.4	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.4	*
12	Certain conditions originating in the perinatal period (P00-P96)	1	0.4	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.4	*
...	All other causes (Residual)	54	19.1	3.2
Black, male, 10-14 years				
...	All causes	384	100.0	22.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	115	29.9	6.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	47	12.2	2.7
3	Malignant neoplasms (C00-C97)	40	10.4	2.3
4	Chronic lower respiratory diseases (J40-J47)	28	7.3	1.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	24	6.3	1.4
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	23	6.0	1.3
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	18	4.7	*
8	Influenza and pneumonia (J09-J18)	6	1.6	*
9	Septicemia (A40-A41)	5	1.3	*
9	Cerebrovascular diseases (I60-I69)	5	1.3	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	1.0	*
12	Diabetes mellitus (E10-E14)	3	0.8	*
13	Anemias (D50-D64)	2	0.5	*
13	Certain conditions originating in the perinatal period (P00-P96)	2	0.5	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.3	*
15	Meningitis (G00,G03)	1	0.3	*
15	Pneumonitis due to solids and liquids (J69)	1	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
...	All other causes (Residual)	58	15.1	3.3

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 15-19 years				
...	All causes	1,965	100.0	102.5
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	991	50.4	51.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	438	22.3	22.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	131	6.7	6.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	70	3.6	3.7
5	Malignant neoplasms (C00-C97)	56	2.8	2.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	35	1.8	1.8
7	Anemias (D50-D64)	21	1.1	1.1
8	Chronic lower respiratory diseases (J40-J47)	17	0.9	*
9	Cerebrovascular diseases (I60-I69)	13	0.7	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	10	0.5	*
11	Legal intervention (Y35,Y89.0)	8	0.4	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.4	*
13	Septicemia (A40-A41)	6	0.3	*
14	Diabetes mellitus (E10-E14)	5	0.3	*
14	Influenza and pneumonia (J09-J18)	5	0.3	*
...	All other causes (Residual)	152	7.7	7.9
Black, male, 20-24 years				
...	All causes	3,164	100.0	189.1
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,557	49.2	93.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	633	20.0	37.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	266	8.4	15.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	141	4.5	8.4
5	Malignant neoplasms (C00-C97)	93	2.9	5.6
6	Human immunodeficiency virus (HIV) disease (B20-B24)	61	1.9	3.6
7	Anemias (D50-D64)	34	1.1	2.0
8	Chronic lower respiratory diseases (J40-J47)	26	0.8	1.6
9	Diabetes mellitus (E10-E14)	21	0.7	1.3
10	Septicemia (A40-A41)	20	0.6	1.2
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	20	0.6	1.2
12	Influenza and pneumonia (J09-J18)	19	0.6	*
12	Legal intervention (Y35,Y89.0)	19	0.6	*
14	Cerebrovascular diseases (I60-I69)	8	0.3	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.2	*
15	Pneumonitis due to solids and liquids (J69)	5	0.2	*
...	All other causes (Residual)	236	7.5	14.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 25-29 years				
...	All causes	2,997	100.0	206.0
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,206	40.2	82.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	602	20.1	41.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	243	8.1	16.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	217	7.2	14.9
5	Human immunodeficiency virus (HIV) disease (B20-B24)	108	3.6	7.4
6	Malignant neoplasms (C00-C97)	100	3.3	6.9
7	Diabetes mellitus (E10-E14)	36	1.2	2.5
8	Chronic lower respiratory diseases (J40-J47)	30	1.0	2.1
9	Anemias (D50-D64)	28	0.9	1.9
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	27	0.9	1.9
11	Cerebrovascular diseases (I60-I69)	25	0.8	1.7
12	Influenza and pneumonia (J09-J18)	21	0.7	1.4
13	Septicemia (A40-A41)	15	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	0.5	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	14	0.5	*
15	Legal intervention (Y35,Y89.0)	14	0.5	*
...	All other causes (Residual)	296	9.9	20.3
Black, male, 30-34 years				
...	All causes	3,074	100.0	228.4
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	926	30.1	68.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	580	18.9	43.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	359	11.7	26.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	190	6.2	14.1
5	Human immunodeficiency virus (HIV) disease (B20-B24)	167	5.4	12.4
6	Malignant neoplasms (C00-C97)	157	5.1	11.7
7	Diabetes mellitus (E10-E14)	77	2.5	5.7
8	Cerebrovascular diseases (I60-I69)	50	1.6	3.7
9	Chronic lower respiratory diseases (J40-J47)	39	1.3	2.9
10	Influenza and pneumonia (J09-J18)	29	0.9	2.2
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	28	0.9	2.1
12	Anemias (D50-D64)	24	0.8	1.8
13	Septicemia (A40-A41)	20	0.7	1.5
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	15	0.5	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	15	0.5	*
...	All other causes (Residual)	398	12.9	29.6

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 35-39 years				
...	All causes	3,440	100.0	263.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	635	18.5	48.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	567	16.5	43.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	541	15.7	41.4
4	Malignant neoplasms (C00-C97)	249	7.2	19.0
5	Human immunodeficiency virus (HIV) disease (B20-B24)	227	6.6	17.4
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	191	5.6	14.6
7	Cerebrovascular diseases (I60-I69)	103	3.0	7.9
8	Diabetes mellitus (E10-E14)	98	2.8	7.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	63	1.8	4.8
10	Septicemia (A40-A41)	49	1.4	3.7
11	Chronic lower respiratory diseases (J40-J47)	43	1.3	3.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	42	1.2	3.2
13	Anemias (D50-D64)	38	1.1	2.9
13	Aortic aneurysm and dissection (I71)	38	1.1	2.9
13	Chronic liver disease and cirrhosis (K70,K73-K74)	38	1.1	2.9
...	All other causes (Residual)	518	15.1	39.6
Black, male, 40-44 years				
...	All causes	4,675	100.0	351.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,040	22.2	78.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	568	12.1	42.7
3	Malignant neoplasms (C00-C97)	545	11.7	40.9
4	Human immunodeficiency virus (HIV) disease (B20-B24)	427	9.1	32.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	344	7.4	25.8
6	Diabetes mellitus (E10-E14)	206	4.4	15.5
7	Cerebrovascular diseases (I60-I69)	186	4.0	14.0
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	164	3.5	12.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	93	2.0	7.0
10	Chronic liver disease and cirrhosis (K70,K73-K74)	68	1.5	5.1
11	Influenza and pneumonia (J09-J18)	59	1.3	4.4
12	Septicemia (A40-A41)	56	1.2	4.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	56	1.2	4.2
14	Chronic lower respiratory diseases (J40-J47)	48	1.0	3.6
15	Aortic aneurysm and dissection (I71)	44	0.9	3.3
...	All other causes (Residual)	771	16.5	57.9

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 45-49 years				
...	All causes	7,665	100.0	549.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,027	26.4	145.4
2	Malignant neoplasms (C00-C97)	1,289	16.8	92.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	736	9.6	52.8
4	Human immunodeficiency virus (HIV) disease (B20-B24)	554	7.2	39.7
5	Cerebrovascular diseases (I60-I69)	341	4.4	24.5
6	Diabetes mellitus (E10-E14)	324	4.2	23.2
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	305	4.0	21.9
8	Chronic liver disease and cirrhosis (K70,K73-K74)	174	2.3	12.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	144	1.9	10.3
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	142	1.9	10.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	127	1.7	9.1
12	Septicemia (A40-A41)	118	1.5	8.5
13	Influenza and pneumonia (J09-J18)	93	1.2	6.7
14	Chronic lower respiratory diseases (J40-J47)	92	1.2	6.6
15	Viral hepatitis (B15-B19)	56	0.7	4.0
...	All other causes (Residual)	1,143	14.9	82.0
Black, male, 50-54 years				
...	All causes	11,728	100.0	893.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,141	26.8	239.2
2	Malignant neoplasms (C00-C97)	2,796	23.8	212.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	826	7.0	62.9
4	Cerebrovascular diseases (I60-I69)	570	4.9	43.4
5	Human immunodeficiency virus (HIV) disease (B20-B24)	536	4.6	40.8
6	Diabetes mellitus (E10-E14)	468	4.0	35.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	348	3.0	26.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	255	2.2	19.4
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	247	2.1	18.8
10	Septicemia (A40-A41)	197	1.7	15.0
11	Chronic lower respiratory diseases (J40-J47)	194	1.7	14.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	161	1.4	12.3
13	Viral hepatitis (B15-B19)	156	1.3	11.9
14	Influenza and pneumonia (J09-J18)	136	1.2	10.4
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	122	1.0	9.3
...	All other causes (Residual)	1,575	13.4	119.9

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 55-59 years				
...	All causes	14,939	100.0	1,425.7
1	Malignant neoplasms (C00-C97)	4,224	28.3	403.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,892	26.1	371.4
3	Cerebrovascular diseases (I60-I69)	760	5.1	72.5
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	724	4.8	69.1
5	Diabetes mellitus (E10-E14)	655	4.4	62.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	445	3.0	42.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	416	2.8	39.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	393	2.6	37.5
9	Chronic lower respiratory diseases (J40-J47)	325	2.2	31.0
10	Septicemia (A40-A41)	273	1.8	26.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	246	1.6	23.5
12	Viral hepatitis (B15-B19)	201	1.3	19.2
13	Influenza and pneumonia (J09-J18)	166	1.1	15.8
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	162	1.1	15.5
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	98	0.7	9.4
...	All other causes (Residual)	1,959	13.1	187.0
Black, male, 60-64 years				
...	All causes	15,519	100.0	1,977.9
1	Malignant neoplasms (C00-C97)	4,877	31.4	621.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,131	26.6	526.5
3	Cerebrovascular diseases (I60-I69)	764	4.9	97.4
4	Diabetes mellitus (E10-E14)	699	4.5	89.1
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	539	3.5	68.7
6	Chronic lower respiratory diseases (J40-J47)	471	3.0	60.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	454	2.9	57.9
8	Chronic liver disease and cirrhosis (K70,K73-K74)	311	2.0	39.6
9	Septicemia (A40-A41)	305	2.0	38.9
10	Human immunodeficiency virus (HIV) disease (B20-B24)	272	1.8	34.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	255	1.6	32.5
12	Influenza and pneumonia (J09-J18)	179	1.2	22.8
13	Viral hepatitis (B15-B19)	150	1.0	19.1
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	83	0.5	10.6
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	70	0.5	8.9
...	All other causes (Residual)	1,959	12.6	249.7

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 65-69 years				
...	All causes	14,289	100.0	2,745.1
1	Malignant neoplasms (C00-C97)	4,666	32.7	896.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,814	26.7	732.7
3	Cerebrovascular diseases (I60-I69)	809	5.7	155.4
4	Diabetes mellitus (E10-E14)	691	4.8	132.7
5	Chronic lower respiratory diseases (J40-J47)	556	3.9	106.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	448	3.1	86.1
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	326	2.3	62.6
8	Septicemia (A40-A41)	303	2.1	58.2
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	246	1.7	47.3
10	Influenza and pneumonia (J09-J18)	225	1.6	43.2
11	Chronic liver disease and cirrhosis (K70,K73-K74)	164	1.1	31.5
12	Human immunodeficiency virus (HIV) disease (B20-B24)	117	0.8	22.5
13	Pneumonitis due to solids and liquids (J69)	68	0.5	13.1
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	56	0.4	10.8
15	Viral hepatitis (B15-B19)	55	0.4	10.6
...	All other causes (Residual)	1,745	12.2	335.2
Black, male, 70-74 years				
...	All causes	14,106	100.0	3,862.0
1	Malignant neoplasms (C00-C97)	4,438	31.5	1,215.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,696	26.2	1,011.9
3	Cerebrovascular diseases (I60-I69)	808	5.7	221.2
4	Chronic lower respiratory diseases (J40-J47)	665	4.7	182.1
5	Diabetes mellitus (E10-E14)	657	4.7	179.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	526	3.7	144.0
7	Septicemia (A40-A41)	288	2.0	78.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	275	1.9	75.3
9	Influenza and pneumonia (J09-J18)	267	1.9	73.1
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	239	1.7	65.4
11	Alzheimer's disease (G30)	128	0.9	35.0
12	Chronic liver disease and cirrhosis (K70,K73-K74)	85	0.6	23.3
13	Pneumonitis due to solids and liquids (J69)	76	0.5	20.8
14	Human immunodeficiency virus (HIV) disease (B20-B24)	72	0.5	19.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	68	0.5	18.6
...	All other causes (Residual)	1,818	12.9	497.7

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 75-79 years				
...	All causes	13,898	100.0	5,677.3
1	Malignant neoplasms (C00-C97)	3,989	28.7	1,629.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,751	27.0	1,532.3
3	Cerebrovascular diseases (I60-I69)	812	5.8	331.7
4	Diabetes mellitus (E10-E14)	680	4.9	277.8
5	Chronic lower respiratory diseases (J40-J47)	613	4.4	250.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	485	3.5	198.1
7	Septicemia (A40-A41)	328	2.4	134.0
8	Influenza and pneumonia (J09-J18)	264	1.9	107.8
9	Alzheimer's disease (G30)	249	1.8	101.7
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	241	1.7	98.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	221	1.6	90.3
12	Pneumonitis due to solids and liquids (J69)	96	0.7	39.2
13	Parkinson's disease (G20-G21)	95	0.7	38.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	68	0.5	27.8
15	Chronic liver disease and cirrhosis (K70,K73-K74)	59	0.4	24.1
...	All other causes (Residual)	1,947	14.0	795.3
Black, male, 80-84 years				
...	All causes	12,704	100.0	8,414.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,401	26.8	2,252.7
2	Malignant neoplasms (C00-C97)	3,241	25.5	2,146.7
3	Cerebrovascular diseases (I60-I69)	763	6.0	505.4
4	Chronic lower respiratory diseases (J40-J47)	612	4.8	405.4
5	Diabetes mellitus (E10-E14)	499	3.9	330.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	471	3.7	312.0
7	Alzheimer's disease (G30)	362	2.8	239.8
8	Influenza and pneumonia (J09-J18)	324	2.6	214.6
9	Septicemia (A40-A41)	294	2.3	194.7
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	230	1.8	152.3
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	204	1.6	135.1
12	Pneumonitis due to solids and liquids (J69)	137	1.1	90.7
13	Parkinson's disease (G20-G21)	101	0.8	66.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	54	0.4	35.8
15	Atherosclerosis (I70)	40	0.3	26.5
...	All other causes (Residual)	1,971	15.5	1,305.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 85-89 years				
...	All causes	9,682	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,706	27.9	---
2	Malignant neoplasms (C00-C97)	2,093	21.6	---
3	Cerebrovascular diseases (I60-I69)	520	5.4	---
4	Chronic lower respiratory diseases (J40-J47)	481	5.0	---
5	Alzheimer's disease (G30)	378	3.9	---
6	Diabetes mellitus (E10-E14)	343	3.5	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	339	3.5	---
8	Influenza and pneumonia (J09-J18)	257	2.7	---
9	Septicemia (A40-A41)	230	2.4	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	182	1.9	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	148	1.5	---
12	Parkinson's disease (G20-G21)	92	1.0	---
13	Pneumonitis due to solids and liquids (J69)	90	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	51	0.5	---
15	Atherosclerosis (I70)	28	0.3	---
...	All other causes (Residual)	1,744	18.0	---
Black, male, 90-94 years				
...	All causes	4,698	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,363	29.0	---
2	Malignant neoplasms (C00-C97)	819	17.4	---
3	Cerebrovascular diseases (I60-I69)	265	5.6	---
4	Alzheimer's disease (G30)	206	4.4	---
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	202	4.3	---
6	Chronic lower respiratory diseases (J40-J47)	188	4.0	---
7	Influenza and pneumonia (J09-J18)	163	3.5	---
8	Diabetes mellitus (E10-E14)	129	2.7	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	120	2.6	---
10	Septicemia (A40-A41)	95	2.0	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	78	1.7	---
12	Pneumonitis due to solids and liquids (J69)	54	1.1	---
13	Parkinson's disease (G20-G21)	33	0.7	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	26	0.6	---
15	Atherosclerosis (I70)	16	0.3	---
...	All other causes (Residual)	941	20.0	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 95-99 years				
...	All causes	1,529	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	458	30.0	---
2	Malignant neoplasms (C00-C97)	209	13.7	---
3	Cerebrovascular diseases (I60-I69)	88	5.8	---
4	Alzheimer's disease (G30)	81	5.3	---
5	Influenza and pneumonia (J09-J18)	67	4.4	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	64	4.2	---
7	Chronic lower respiratory diseases (J40-J47)	47	3.1	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	43	2.8	---
9	Diabetes mellitus (E10-E14)	38	2.5	---
10	Septicemia (A40-A41)	28	1.8	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	25	1.6	---
12	Pneumonitis due to solids and liquids (J69)	21	1.4	---
13	Atherosclerosis (I70)	11	0.7	---
14	Nutritional deficiencies (E40-E64)	10	0.7	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.5	---
...	All other causes (Residual)	332	21.7	---
Black, male, 100 years and over				
...	All causes	334	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	112	33.5	---
2	Malignant neoplasms (C00-C97)	30	9.0	---
3	Alzheimer's disease (G30)	15	4.5	---
3	Cerebrovascular diseases (I60-I69)	15	4.5	---
3	Chronic lower respiratory diseases (J40-J47)	15	4.5	---
6	Influenza and pneumonia (J09-J18)	14	4.2	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	12	3.6	---
8	Diabetes mellitus (E10-E14)	8	2.4	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	2.4	---
10	Septicemia (A40-A41)	6	1.8	---
11	Atherosclerosis (I70)	5	1.5	---
12	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3	0.9	---
13	Parkinson's disease (G20-G21)	2	0.6	---
13	Pneumonitis due to solids and liquids (J69)	2	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.3	---
15	Nutritional deficiencies (E40-E64)	1	0.3	---
15	Pneumoconioses and chemical effects (J60-J66,J68)	1	0.3	---
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.3	---
15	Hyperplasia of prostate (N40)	1	0.3	---
...	All other causes (Residual)	82	24.6	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, all ages				
...	All causes	141,157	100.0	642.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	33,994	24.1	154.8
2	Malignant neoplasms (C00-C97)	31,963	22.6	145.5
3	Cerebrovascular diseases (I60-I69)	9,027	6.4	41.1
4	Diabetes mellitus (E10-E14)	6,486	4.6	29.5
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,825	3.4	22.0
6	Chronic lower respiratory diseases (J40-J47)	4,183	3.0	19.0
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,995	2.8	18.2
8	Alzheimer's disease (G30)	3,732	2.6	17.0
9	Septicemia (A40-A41)	3,310	2.3	15.1
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,898	2.1	13.2
11	Influenza and pneumonia (J09-J18)	2,556	1.8	11.6
12	Certain conditions originating in the perinatal period (P00-P96)	1,815	1.3	8.3
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,615	1.1	7.4
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,114	0.8	5.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	920	0.7	4.2
...	All other causes (Residual)	28,724	20.3	130.8
Black, female, 1-4 years				
...	All causes	446	100.0	33.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	97	21.7	7.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	70	15.7	5.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	63	14.1	4.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	22	4.9	1.6
5	Malignant neoplasms (C00-C97)	21	4.7	1.6
6	Cerebrovascular diseases (I60-I69)	10	2.2	*
6	Influenza and pneumonia (J09-J18)	10	2.2	*
6	Chronic lower respiratory diseases (J40-J47)	10	2.2	*
9	Septicemia (A40-A41)	7	1.6	*
9	Certain conditions originating in the perinatal period (P00-P96)	7	1.6	*
11	Anemias (D50-D64)	5	1.1	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.9	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.7	*
14	Meningitis (G00,G03)	2	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
...	All other causes (Residual)	113	25.3	8.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 5-9 years				
...	All causes	211	100.0	13.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	56	26.5	3.5
2	Malignant neoplasms (C00-C97)	40	19.0	2.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	14	6.6	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	3.8	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	6	2.8	*
5	Cerebrovascular diseases (I60-I69)	6	2.8	*
7	Septicemia (A40-A41)	5	2.4	*
7	Chronic lower respiratory diseases (J40-J47)	5	2.4	*
9	Anemias (D50-D64)	3	1.4	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.9	*
10	Diabetes mellitus (E10-E14)	2	0.9	*
10	Influenza and pneumonia (J09-J18)	2	0.9	*
10	Certain conditions originating in the perinatal period (P00-P96)	2	0.9	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.9	*
15	Tuberculosis (A16-A19)	1	0.5	*
15	Diseases of appendix (K35-K38)	1	0.5	*
...	All other causes (Residual)	56	26.5	3.5
Black, female, 10-14 years				
...	All causes	267	100.0	16.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	50	18.7	3.0
2	Malignant neoplasms (C00-C97)	41	15.4	2.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	21	7.9	1.3
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	20	7.5	1.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	18	6.7	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	16	6.0	*
7	Chronic lower respiratory diseases (J40-J47)	15	5.6	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	2.2	*
9	Anemias (D50-D64)	5	1.9	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	1.1	*
11	Septicemia (A40-A41)	2	0.7	*
11	Diabetes mellitus (E10-E14)	2	0.7	*
11	Cerebrovascular diseases (I60-I69)	2	0.7	*
11	Influenza and pneumonia (J09-J18)	2	0.7	*
15	Viral hepatitis (B15-B19)	1	0.4	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.4	*
15	Diseases of appendix (K35-K38)	1	0.4	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.4	*
15	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.4	*
15	Certain conditions originating in the perinatal period (P00-P96)	1	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.4	*
...	All other causes (Residual)	57	21.3	3.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 15-19 years				
...	All causes	566	100.0	30.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	144	25.4	7.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	126	22.3	6.8
3	Malignant neoplasms (C00-C97)	50	8.8	2.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	34	6.0	1.8
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	21	3.7	1.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	15	2.7	*
7	Cerebrovascular diseases (I60-I69)	11	1.9	*
7	Chronic lower respiratory diseases (J40-J47)	11	1.9	*
7	Pregnancy, childbirth and the puerperium (O00-O99)	11	1.9	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	7	1.2	*
10	Anemias (D50-D64)	7	1.2	*
12	Influenza and pneumonia (J09-J18)	6	1.1	*
13	Septicemia (A40-A41)	5	0.9	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.9	*
15	Diabetes mellitus (E10-E14)	4	0.7	*
...	All other causes (Residual)	109	19.3	5.9
Black, female, 20-24 years				
...	All causes	980	100.0	57.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	242	24.7	14.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	142	14.5	8.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	91	9.3	5.3
4	Malignant neoplasms (C00-C97)	68	6.9	4.0
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	51	5.2	3.0
6	Pregnancy, childbirth and the puerperium (O00-O99)	40	4.1	2.3
7	Human immunodeficiency virus (HIV) disease (B20-B24)	33	3.4	1.9
8	Diabetes mellitus (E10-E14)	21	2.1	1.2
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	21	2.1	1.2
10	Anemias (D50-D64)	17	1.7	*
10	Cerebrovascular diseases (I60-I69)	17	1.7	*
12	Septicemia (A40-A41)	10	1.0	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	1.0	*
14	Influenza and pneumonia (J09-J18)	9	0.9	*
15	Chronic lower respiratory diseases (J40-J47)	8	0.8	*
...	All other causes (Residual)	200	20.4	11.7

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 25-29 years				
...	All causes	1,249	100.0	79.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	213	17.1	13.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	164	13.1	10.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	131	10.5	8.3
4	Malignant neoplasms (C00-C97)	118	9.4	7.5
5	Pregnancy, childbirth and the puerperium (O00-O99)	58	4.6	3.7
6	Human immunodeficiency virus (HIV) disease (B20-B24)	57	4.6	3.6
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	43	3.4	2.7
8	Diabetes mellitus (E10-E14)	42	3.4	2.7
9	Anemias (D50-D64)	27	2.2	1.7
10	Cerebrovascular diseases (I60-I69)	22	1.8	1.4
11	Influenza and pneumonia (J09-J18)	20	1.6	1.3
12	Septicemia (A40-A41)	19	1.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	1.3	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	13	1.0	*
15	Chronic lower respiratory diseases (J40-J47)	11	0.9	*
...	All other causes (Residual)	295	23.6	18.8
Black, female, 30-34 years				
...	All causes	1,600	100.0	107.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	217	13.6	14.5
2	Malignant neoplasms (C00-C97)	214	13.4	14.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	207	12.9	13.9
4	Human immunodeficiency virus (HIV) disease (B20-B24)	140	8.8	9.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	105	6.6	7.0
6	Pregnancy, childbirth and the puerperium (O00-O99)	60	3.8	4.0
7	Diabetes mellitus (E10-E14)	54	3.4	3.6
8	Cerebrovascular diseases (I60-I69)	40	2.5	2.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	36	2.3	2.4
10	Septicemia (A40-A41)	32	2.0	2.1
11	Anemias (D50-D64)	30	1.9	2.0
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	28	1.8	1.9
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	20	1.3	1.3
14	Chronic lower respiratory diseases (J40-J47)	18	1.1	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	16	1.0	*
...	All other causes (Residual)	383	23.9	25.6

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 35-39 years				
...	All causes	2,370	100.0	160.5
1	Malignant neoplasms (C00-C97)	495	20.9	33.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	372	15.7	25.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	228	9.6	15.4
4	Human immunodeficiency virus (HIV) disease (B20-B24)	184	7.8	12.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	98	4.1	6.6
6	Cerebrovascular diseases (I60-I69)	95	4.0	6.4
7	Diabetes mellitus (E10-E14)	85	3.6	5.8
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	49	2.1	3.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	48	2.0	3.3
10	Pregnancy, childbirth and the puerperium (O00-O99)	47	2.0	3.2
11	Chronic lower respiratory diseases (J40-J47)	44	1.9	3.0
12	Septicemia (A40-A41)	37	1.6	2.5
13	Anemias (D50-D64)	33	1.4	2.2
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	31	1.3	2.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	28	1.2	1.9
...	All other causes (Residual)	496	20.9	33.6
Black, female, 40-44 years				
...	All causes	3,531	100.0	237.7
1	Malignant neoplasms (C00-C97)	822	23.3	55.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	679	19.2	45.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	265	7.5	17.8
4	Human immunodeficiency virus (HIV) disease (B20-B24)	239	6.8	16.1
5	Cerebrovascular diseases (I60-I69)	175	5.0	11.8
6	Diabetes mellitus (E10-E14)	125	3.5	8.4
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	79	2.2	5.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	77	2.2	5.2
9	Septicemia (A40-A41)	70	2.0	4.7
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	48	1.4	3.2
11	Chronic lower respiratory diseases (J40-J47)	47	1.3	3.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	46	1.3	3.1
12	Chronic liver disease and cirrhosis (K70,K73-K74)	46	1.3	3.1
14	Influenza and pneumonia (J09-J18)	39	1.1	2.6
15	Anemias (D50-D64)	33	0.9	2.2
15	Pregnancy, childbirth and the puerperium (O00-O99)	33	0.9	2.2
...	All other causes (Residual)	708	20.1	47.7

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 45-49 years				
...	All causes	5,916	100.0	377.8
1	Malignant neoplasms (C00-C97)	1,699	28.7	108.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,168	19.7	74.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	354	6.0	22.6
4	Cerebrovascular diseases (I60-I69)	319	5.4	20.4
5	Human immunodeficiency virus (HIV) disease (B20-B24)	289	4.9	18.5
6	Diabetes mellitus (E10-E14)	239	4.0	15.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	137	2.3	8.8
8	Chronic lower respiratory diseases (J40-J47)	121	2.0	7.7
9	Septicemia (A40-A41)	110	1.9	7.0
10	Chronic liver disease and cirrhosis (K70,K73-K74)	109	1.8	7.0
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	91	1.5	5.8
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	85	1.4	5.4
13	Influenza and pneumonia (J09-J18)	74	1.3	4.7
14	Anemias (D50-D64)	40	0.7	2.6
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	36	0.6	2.3
...	All other causes (Residual)	1,045	17.7	66.7
Black, female, 50-54 years				
...	All causes	8,784	100.0	589.3
1	Malignant neoplasms (C00-C97)	2,777	31.6	186.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,783	20.3	119.6
3	Cerebrovascular diseases (I60-I69)	496	5.6	33.3
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	389	4.4	26.1
5	Diabetes mellitus (E10-E14)	363	4.1	24.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	264	3.0	17.7
7	Human immunodeficiency virus (HIV) disease (B20-B24)	256	2.9	17.2
8	Chronic lower respiratory diseases (J40-J47)	242	2.8	16.2
9	Chronic liver disease and cirrhosis (K70,K73-K74)	186	2.1	12.5
10	Septicemia (A40-A41)	182	2.1	12.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	153	1.7	10.3
12	Influenza and pneumonia (J09-J18)	89	1.0	6.0
13	Viral hepatitis (B15-B19)	74	0.8	5.0
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	56	0.6	3.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	41	0.5	2.8
...	All other causes (Residual)	1,433	16.3	96.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 55-59 years				
...	All causes	10,307	100.0	833.0
1	Malignant neoplasms (C00-C97)	3,438	33.4	277.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,258	21.9	182.5
3	Cerebrovascular diseases (I60-I69)	557	5.4	45.0
4	Diabetes mellitus (E10-E14)	514	5.0	41.5
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	316	3.1	25.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	306	3.0	24.7
7	Chronic lower respiratory diseases (J40-J47)	301	2.9	24.3
8	Septicemia (A40-A41)	245	2.4	19.8
9	Human immunodeficiency virus (HIV) disease (B20-B24)	201	2.0	16.2
10	Chronic liver disease and cirrhosis (K70,K73-K74)	165	1.6	13.3
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	159	1.5	12.9
12	Influenza and pneumonia (J09-J18)	114	1.1	9.2
13	Viral hepatitis (B15-B19)	96	0.9	7.8
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	50	0.5	4.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	39	0.4	3.2
...	All other causes (Residual)	1,548	15.0	125.1
Black, female, 60-64 years				
...	All causes	11,051	100.0	1,151.8
1	Malignant neoplasms (C00-C97)	3,836	34.7	399.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,545	23.0	265.2
3	Diabetes mellitus (E10-E14)	621	5.6	64.7
4	Cerebrovascular diseases (I60-I69)	571	5.2	59.5
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	385	3.5	40.1
6	Chronic lower respiratory diseases (J40-J47)	366	3.3	38.1
7	Septicemia (A40-A41)	282	2.6	29.4
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	224	2.0	23.3
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	190	1.7	19.8
10	Influenza and pneumonia (J09-J18)	140	1.3	14.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	134	1.2	14.0
12	Human immunodeficiency virus (HIV) disease (B20-B24)	111	1.0	11.6
13	Viral hepatitis (B15-B19)	55	0.5	5.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	52	0.5	5.4
15	Anemias (D50-D64)	38	0.3	4.0
...	All other causes (Residual)	1,501	13.6	156.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 65-69 years				
...	All causes	11,492	100.0	1,691.9
1	Malignant neoplasms (C00-C97)	3,773	32.8	555.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,652	23.1	390.4
3	Diabetes mellitus (E10-E14)	685	6.0	100.8
4	Cerebrovascular diseases (I60-I69)	660	5.7	97.2
5	Chronic lower respiratory diseases (J40-J47)	445	3.9	65.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	441	3.8	64.9
7	Septicemia (A40-A41)	326	2.8	48.0
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	234	2.0	34.5
9	Influenza and pneumonia (J09-J18)	172	1.5	25.3
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	169	1.5	24.9
11	Chronic liver disease and cirrhosis (K70,K73-K74)	74	0.6	10.9
12	Alzheimer's disease (G30)	67	0.6	9.9
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	51	0.4	7.5
14	Human immunodeficiency virus (HIV) disease (B20-B24)	45	0.4	6.6
15	Viral hepatitis (B15-B19)	36	0.3	5.3
...	All other causes (Residual)	1,662	14.5	244.7
Black, female, 70-74 years				
...	All causes	12,604	100.0	2,457.3
1	Malignant neoplasms (C00-C97)	3,750	29.8	731.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,022	24.0	589.2
3	Cerebrovascular diseases (I60-I69)	844	6.7	164.6
4	Diabetes mellitus (E10-E14)	748	5.9	145.8
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	546	4.3	106.5
6	Chronic lower respiratory diseases (J40-J47)	524	4.2	102.2
7	Septicemia (A40-A41)	352	2.8	68.6
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	232	1.8	45.2
9	Influenza and pneumonia (J09-J18)	206	1.6	40.2
10	Alzheimer's disease (G30)	169	1.3	32.9
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	145	1.2	28.3
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	71	0.6	13.8
13	Chronic liver disease and cirrhosis (K70,K73-K74)	62	0.5	12.1
14	Pneumonitis due to solids and liquids (J69)	55	0.4	10.7
15	Aortic aneurysm and dissection (I71)	41	0.3	8.0
...	All other causes (Residual)	1,837	14.6	358.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 75-79 years				
...	All causes	14,340	100.0	3,679.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,720	25.9	954.6
2	Malignant neoplasms (C00-C97)	3,562	24.8	914.1
3	Cerebrovascular diseases (I60-I69)	1,038	7.2	266.4
4	Diabetes mellitus (E10-E14)	831	5.8	213.2
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	596	4.2	152.9
6	Chronic lower respiratory diseases (J40-J47)	524	3.7	134.5
7	Septicemia (A40-A41)	364	2.5	93.4
8	Alzheimer's disease (G30)	359	2.5	92.1
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	292	2.0	74.9
10	Influenza and pneumonia (J09-J18)	262	1.8	67.2
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	186	1.3	47.7
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	79	0.6	20.3
13	Pneumonitis due to solids and liquids (J69)	69	0.5	17.7
14	Aortic aneurysm and dissection (I71)	53	0.4	13.6
15	Parkinson's disease (G20-G21)	51	0.4	13.1
...	All other causes (Residual)	2,354	16.4	604.1
Black, female, 80-84 years				
...	All causes	16,601	100.0	5,809.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,569	27.5	1,599.0
2	Malignant neoplasms (C00-C97)	3,248	19.6	1,136.7
3	Cerebrovascular diseases (I60-I69)	1,307	7.9	457.4
4	Diabetes mellitus (E10-E14)	817	4.9	285.9
5	Alzheimer's disease (G30)	694	4.2	242.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	670	4.0	234.5
7	Chronic lower respiratory diseases (J40-J47)	581	3.5	203.3
8	Septicemia (A40-A41)	436	2.6	152.6
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	389	2.3	136.1
10	Influenza and pneumonia (J09-J18)	380	2.3	133.0
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	191	1.2	66.8
12	Pneumonitis due to solids and liquids (J69)	101	0.6	35.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	87	0.5	30.4
14	Parkinson's disease (G20-G21)	80	0.5	28.0
15	Aortic aneurysm and dissection (I71)	67	0.4	23.4
...	All other causes (Residual)	2,984	18.0	1,044.3

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 85-89 years				
...	All causes	16,396	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,707	28.7	---
2	Malignant neoplasms (C00-C97)	2,361	14.4	---
3	Cerebrovascular diseases (I60-I69)	1,349	8.2	---
4	Alzheimer's disease (G30)	963	5.9	---
5	Diabetes mellitus (E10-E14)	706	4.3	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	626	3.8	---
7	Chronic lower respiratory diseases (J40-J47)	487	3.0	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	433	2.6	---
9	Influenza and pneumonia (J09-J18)	417	2.5	---
10	Septicemia (A40-A41)	414	2.5	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	176	1.1	---
12	Pneumonitis due to solids and liquids (J69)	138	0.8	---
13	Parkinson's disease (G20-G21)	99	0.6	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	71	0.4	---
15	Atherosclerosis (I70)	64	0.4	---
...	All other causes (Residual)	3,385	20.6	---
Black, female, 90-94 years				
...	All causes	11,472	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,491	30.4	---
2	Malignant neoplasms (C00-C97)	1,152	10.0	---
3	Cerebrovascular diseases (I60-I69)	924	8.1	---
4	Alzheimer's disease (G30)	861	7.5	---
5	Diabetes mellitus (E10-E14)	410	3.6	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	403	3.5	---
7	Influenza and pneumonia (J09-J18)	316	2.8	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	306	2.7	---
9	Chronic lower respiratory diseases (J40-J47)	278	2.4	---
10	Septicemia (A40-A41)	236	2.1	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	133	1.2	---
12	Pneumonitis due to solids and liquids (J69)	80	0.7	---
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	67	0.6	---
14	Atherosclerosis (I70)	53	0.5	---
15	Anemias (D50-D64)	45	0.4	---
...	All other causes (Residual)	2,717	23.7	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 95-99 years				
...	All causes	5,838	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,866	32.0	---
2	Alzheimer's disease (G30)	447	7.7	---
3	Cerebrovascular diseases (I60-I69)	440	7.5	---
4	Malignant neoplasms (C00-C97)	415	7.1	---
5	Essential hypertension and hypertensive renal disease (I10,I12,I15)	205	3.5	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	186	3.2	---
7	Diabetes mellitus (E10-E14)	176	3.0	---
8	Influenza and pneumonia (J09-J18)	161	2.8	---
9	Chronic lower respiratory diseases (J40-J47)	117	2.0	---
10	Septicemia (A40-A41)	105	1.8	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	70	1.2	---
12	Pneumonitis due to solids and liquids (J69)	53	0.9	---
13	Atherosclerosis (I70)	37	0.6	---
14	Anemias (D50-D64)	25	0.4	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	0.4	---
...	All other causes (Residual)	1,512	25.9	---
Black, female, 100 years and over				
...	All causes	1,846	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	621	33.6	---
2	Alzheimer's disease (G30)	139	7.5	---
3	Cerebrovascular diseases (I60-I69)	115	6.2	---
4	Malignant neoplasms (C00-C97)	78	4.2	---
5	Essential hypertension and hypertensive renal disease (I10,I12,I15)	73	4.0	---
6	Influenza and pneumonia (J09-J18)	72	3.9	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	50	2.7	---
8	Diabetes mellitus (E10-E14)	40	2.2	---
9	Septicemia (A40-A41)	28	1.5	---
10	Chronic lower respiratory diseases (J40-J47)	26	1.4	---
11	Pneumonitis due to solids and liquids (J69)	18	1.0	---
12	Atherosclerosis (I70)	16	0.9	---
13	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12	0.7	---
14	Nutritional deficiencies (E40-E64)	9	0.5	---
15	Anemias (D50-D64)	7	0.4	---
...	All other causes (Residual)	542	29.4	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, all ages				
...	All causes	15,565	100.0	365.1
1	Malignant neoplasms (C00-C97)	2,962	19.0	69.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,793	17.9	65.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,701	10.9	39.9
4	Diabetes mellitus (E10-E14)	857	5.5	20.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	787	5.1	18.5
6	Chronic lower respiratory diseases (J40-J47)	702	4.5	16.5
7	Cerebrovascular diseases (I60-I69)	559	3.6	13.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	469	3.0	11.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	339	2.2	8.0
10	Influenza and pneumonia (J09-J18)	326	2.1	7.6
11	Alzheimer's disease (G30)	264	1.7	6.2
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	257	1.7	6.0
13	Septicemia (A40-A41)	244	1.6	5.7
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	144	0.9	3.4
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	130	0.8	3.0
...	All other causes (Residual)	3,031	19.5	71.1
American Indian, both sexes, 1-4 years				
...	All causes	93	100.0	29.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	31	33.3	9.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	10	10.8	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	9.7	*
4	Malignant neoplasms (C00-C97)	6	6.5	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	5.4	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	2.2	*
6	Chronic lower respiratory diseases (J40-J47)	2	2.2	*
8	Septicemia (A40-A41)	1	1.1	*
8	Meningitis (G00,G03)	1	1.1	*
8	Cerebrovascular diseases (I60-I69)	1	1.1	*
8	Influenza and pneumonia (J09-J18)	1	1.1	*
8	Acute bronchitis and bronchiolitis (J20-J21)	1	1.1	*
8	Pneumonitis due to solids and liquids (J69)	1	1.1	*
8	Certain conditions originating in the perinatal period (P00-P96)	1	1.1	*
8	Legal intervention (Y35,Y89.0)	1	1.1	*
8	Complications of medical and surgical care (Y40-Y84,Y88)	1	1.1	*
...	All other causes (Residual)	19	20.4	*

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 5-9 years				
...	All causes	46	100.0	12.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	20	43.5	5.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	13.0	*
3	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	6.5	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	6.5	*
5	Malignant neoplasms (C00-C97)	2	4.3	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	4.3	*
5	Cerebrovascular diseases (I60-I69)	2	4.3	*
8	Influenza and pneumonia (J09-J18)	1	2.2	*
8	Chronic lower respiratory diseases (J40-J47)	1	2.2	*
...	All other causes (Residual)	6	13.0	*
American Indian, both sexes, 10-14 years				
...	All causes	62	100.0	16.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	27	43.5	7.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	14	22.6	*
3	Malignant neoplasms (C00-C97)	6	9.7	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	6.5	*
5	Septicemia (A40-A41)	1	1.6	*
5	Influenza and pneumonia (J09-J18)	1	1.6	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1	1.6	*
5	Complications of medical and surgical care (Y40-Y84,Y88)	1	1.6	*
...	All other causes (Residual)	7	11.3	*
American Indian, both sexes, 15-19 years				
...	All causes	242	100.0	61.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	110	45.5	28.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	70	28.9	17.8
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	25	10.3	6.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	2.1	*
5	Cerebrovascular diseases (I60-I69)	3	1.2	*
6	Malignant neoplasms (C00-C97)	2	0.8	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	0.8	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.4	*
8	Anemias (D50-D64)	1	0.4	*
8	Pneumonitis due to solids and liquids (J69)	1	0.4	*
8	Pregnancy, childbirth and the puerperium (O00-O99)	1	0.4	*
...	All other causes (Residual)	21	8.7	5.3

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 20-24 years				
...	All causes	373	100.0	102.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	160	42.9	44.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	88	23.6	24.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	50	13.4	13.8
4	Malignant neoplasms (C00-C97)	12	3.2	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	11	2.9	*
6	Chronic liver disease and cirrhosis (K70,K73-K74)	4	1.1	*
6	Pregnancy, childbirth and the puerperium (O00-O99)	4	1.1	*
6	Legal intervention (Y35,Y89.0)	4	1.1	*
9	Septicemia (A40-A41)	2	0.5	*
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2	0.5	*
9	Influenza and pneumonia (J09-J18)	2	0.5	*
9	Chronic lower respiratory diseases (J40-J47)	2	0.5	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.5	*
14	Diabetes mellitus (E10-E14)	1	0.3	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.3	*
...	All other causes (Residual)	27	7.2	7.4
American Indian, both sexes, 25-29 years				
...	All causes	376	100.0	110.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	164	43.6	48.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	51	13.6	15.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	39	10.4	11.5
4	Malignant neoplasms (C00-C97)	21	5.6	6.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	15	4.0	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	11	2.9	*
7	Cerebrovascular diseases (I60-I69)	4	1.1	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	1.1	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	3	0.8	*
9	Diabetes mellitus (E10-E14)	3	0.8	*
11	Septicemia (A40-A41)	2	0.5	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2	0.5	*
11	Legal intervention (Y35,Y89.0)	2	0.5	*
14	Meningitis (G00,G03)	1	0.3	*
14	Pneumonitis due to solids and liquids (J69)	1	0.3	*
14	Pregnancy, childbirth and the puerperium (O00-O99)	1	0.3	*
...	All other causes (Residual)	52	13.8	15.3

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 30-34 years				
...	All causes	419	100.0	134.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	146	34.8	46.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	49	11.7	15.8
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	34	8.1	10.9
4	Chronic liver disease and cirrhosis (K70,K73-K74)	33	7.9	10.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	27	6.4	8.7
6	Malignant neoplasms (C00-C97)	24	5.7	7.7
7	Influenza and pneumonia (J09-J18)	7	1.7	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	6	1.4	*
8	Diabetes mellitus (E10-E14)	6	1.4	*
10	Chronic lower respiratory diseases (J40-J47)	5	1.2	*
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	1.2	*
12	Septicemia (A40-A41)	3	0.7	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3	0.7	*
14	Cerebrovascular diseases (I60-I69)	2	0.5	*
14	Pregnancy, childbirth and the puerperium (O00-O99)	2	0.5	*
...	All other causes (Residual)	67	16.0	21.5
American Indian, both sexes, 35-39 years				
...	All causes	513	100.0	175.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	144	28.1	49.2
2	Chronic liver disease and cirrhosis (K70,K73-K74)	79	15.4	27.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	61	11.9	20.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	37	7.2	12.7
5	Malignant neoplasms (C00-C97)	31	6.0	10.6
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	22	4.3	7.5
7	Diabetes mellitus (E10-E14)	18	3.5	*
8	Influenza and pneumonia (J09-J18)	11	2.1	*
9	Cerebrovascular diseases (I60-I69)	6	1.2	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	5	1.0	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	1.0	*
12	Chronic lower respiratory diseases (J40-J47)	4	0.8	*
12	Pregnancy, childbirth and the puerperium (O00-O99)	4	0.8	*
14	Septicemia (A40-A41)	3	0.6	*
14	Viral hepatitis (B15-B19)	3	0.6	*
...	All other causes (Residual)	80	15.6	27.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 40-44 years				
...	All causes	650	100.0	232.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	158	24.3	56.4
2	Chronic liver disease and cirrhosis (K70,K73-K74)	92	14.2	32.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	84	12.9	30.0
4	Malignant neoplasms (C00-C97)	48	7.4	17.1
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	38	5.8	13.6
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	26	4.0	9.3
7	Diabetes mellitus (E10-E14)	22	3.4	7.9
8	Septicemia (A40-A41)	11	1.7	*
9	Cerebrovascular diseases (I60-I69)	9	1.4	*
10	Influenza and pneumonia (J09-J18)	8	1.2	*
11	Viral hepatitis (B15-B19)	7	1.1	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	7	1.1	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.9	*
14	Chronic lower respiratory diseases (J40-J47)	5	0.8	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.5	*
15	Anemias (D50-D64)	3	0.5	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3	0.5	*
15	Aortic aneurysm and dissection (I71)	3	0.5	*
...	All other causes (Residual)	117	18.0	41.8
American Indian, both sexes, 45-49 years				
...	All causes	990	100.0	348.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	170	17.2	59.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	157	15.9	55.3
3	Chronic liver disease and cirrhosis (K70,K73-K74)	139	14.0	49.0
4	Malignant neoplasms (C00-C97)	122	12.3	43.0
5	Diabetes mellitus (E10-E14)	54	5.5	19.0
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	45	4.5	15.9
7	Cerebrovascular diseases (I60-I69)	20	2.0	7.0
8	Viral hepatitis (B15-B19)	19	1.9	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	16	1.6	*
9	Influenza and pneumonia (J09-J18)	16	1.6	*
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	16	1.6	*
12	Septicemia (A40-A41)	13	1.3	*
12	Chronic lower respiratory diseases (J40-J47)	13	1.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	1.1	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7	0.7	*
...	All other causes (Residual)	172	17.4	60.6

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 50-54 years				
...	All causes	1,213	100.0	477.8
1	Malignant neoplasms (C00-C97)	251	20.7	98.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	205	16.9	80.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	137	11.3	54.0
4	Chronic liver disease and cirrhosis (K70,K73-K74)	125	10.3	49.2
5	Diabetes mellitus (E10-E14)	68	5.6	26.8
6	Cerebrovascular diseases (I60-I69)	40	3.3	15.8
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	32	2.6	12.6
8	Chronic lower respiratory diseases (J40-J47)	30	2.5	11.8
9	Viral hepatitis (B15-B19)	23	1.9	9.1
10	Influenza and pneumonia (J09-J18)	20	1.6	7.9
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20	1.6	7.9
12	Septicemia (A40-A41)	18	1.5	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	14	1.2	*
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	12	1.0	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	8	0.7	*
...	All other causes (Residual)	210	17.3	82.7
American Indian, both sexes, 55-59 years				
...	All causes	1,380	100.0	699.4
1	Malignant neoplasms (C00-C97)	379	27.5	192.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	245	17.8	124.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	107	7.8	54.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	103	7.5	52.2
5	Diabetes mellitus (E10-E14)	94	6.8	47.6
6	Chronic lower respiratory diseases (J40-J47)	55	4.0	27.9
7	Cerebrovascular diseases (I60-I69)	33	2.4	16.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	33	2.4	16.7
9	Septicemia (A40-A41)	28	2.0	14.2
10	Viral hepatitis (B15-B19)	20	1.4	10.1
10	Influenza and pneumonia (J09-J18)	20	1.4	10.1
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	20	1.4	10.1
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	11	0.8	*
14	Cholelithiasis and other disorders of gallbladder (K80-K82)	7	0.5	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	6	0.4	*
15	Pneumonitis due to solids and liquids (J69)	6	0.4	*
...	All other causes (Residual)	213	15.4	108.0

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 60-64 years				
...	All causes	1,324	100.0	892.0
1	Malignant neoplasms (C00-C97)	370	27.9	249.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	277	20.9	186.6
3	Diabetes mellitus (E10-E14)	108	8.2	72.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	78	5.9	52.5
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	72	5.4	48.5
6	Chronic lower respiratory diseases (J40-J47)	66	5.0	44.5
7	Cerebrovascular diseases (I60-I69)	44	3.3	29.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	29	2.2	19.5
9	Septicemia (A40-A41)	22	1.7	14.8
10	Influenza and pneumonia (J09-J18)	20	1.5	13.5
11	Viral hepatitis (B15-B19)	14	1.1	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10	0.8	*
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8	0.6	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.5	*
14	Pneumonitis due to solids and liquids (J69)	7	0.5	*
...	All other causes (Residual)	192	14.5	129.4
American Indian, both sexes, 65-69 years				
...	All causes	1,349	100.0	1,377.8
1	Malignant neoplasms (C00-C97)	423	31.4	432.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	261	19.3	266.6
3	Diabetes mellitus (E10-E14)	106	7.9	108.3
4	Chronic lower respiratory diseases (J40-J47)	96	7.1	98.1
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	55	4.1	56.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	51	3.8	52.1
7	Cerebrovascular diseases (I60-I69)	43	3.2	43.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	30	2.2	30.6
9	Septicemia (A40-A41)	26	1.9	26.6
10	Influenza and pneumonia (J09-J18)	23	1.7	23.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	19	1.4	*
12	Alzheimer's disease (G30)	12	0.9	*
13	Viral hepatitis (B15-B19)	7	0.5	*
13	Pneumonitis due to solids and liquids (J69)	7	0.5	*
15	Aortic aneurysm and dissection (I71)	6	0.4	*
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6	0.4	*
...	All other causes (Residual)	178	13.2	181.8

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 70-74 years				
...	All causes	1,454	100.0	2,202.4
1	Malignant neoplasms (C00-C97)	412	28.3	624.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	281	19.3	425.6
3	Chronic lower respiratory diseases (J40-J47)	109	7.5	165.1
4	Diabetes mellitus (E10-E14)	95	6.5	143.9
5	Cerebrovascular diseases (I60-I69)	61	4.2	92.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	44	3.0	66.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	42	2.9	63.6
8	Chronic liver disease and cirrhosis (K70,K73-K74)	32	2.2	48.5
9	Septicemia (A40-A41)	29	2.0	43.9
10	Influenza and pneumonia (J09-J18)	27	1.9	40.9
11	Alzheimer's disease (G30)	20	1.4	30.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	13	0.9	*
12	Pneumonitis due to solids and liquids (J69)	13	0.9	*
14	Parkinson's disease (G20-G21)	11	0.8	*
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	9	0.6	*
...	All other causes (Residual)	256	17.6	387.8
American Indian, both sexes, 75-79 years				
...	All causes	1,388	100.0	3,221.2
1	Malignant neoplasms (C00-C97)	345	24.9	800.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	289	20.8	670.7
3	Diabetes mellitus (E10-E14)	106	7.6	246.0
4	Chronic lower respiratory diseases (J40-J47)	97	7.0	225.1
5	Cerebrovascular diseases (I60-I69)	74	5.3	171.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	43	3.1	99.8
7	Alzheimer's disease (G30)	41	3.0	95.1
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	41	3.0	95.1
9	Influenza and pneumonia (J09-J18)	39	2.8	90.5
10	Septicemia (A40-A41)	27	1.9	62.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	20	1.4	46.4
12	Chronic liver disease and cirrhosis (K70,K73-K74)	19	1.4	*
13	Parkinson's disease (G20-G21)	17	1.2	*
14	Aortic aneurysm and dissection (I71)	11	0.8	*
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	10	0.7	*
...	All other causes (Residual)	209	15.1	485.0

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 80-84 years				
...	All causes	1,297	100.0	4,811.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	295	22.7	1,094.3
2	Malignant neoplasms (C00-C97)	256	19.7	949.6
3	Chronic lower respiratory diseases (J40-J47)	97	7.5	359.8
4	Diabetes mellitus (E10-E14)	90	6.9	333.8
5	Cerebrovascular diseases (I60-I69)	75	5.8	278.2
6	Alzheimer's disease (G30)	58	4.5	215.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	46	3.5	170.6
8	Influenza and pneumonia (J09-J18)	33	2.5	122.4
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	33	2.5	122.4
10	Septicemia (A40-A41)	17	1.3	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	16	1.2	*
12	Chronic liver disease and cirrhosis (K70,K73-K74)	14	1.1	*
13	Parkinson's disease (G20-G21)	13	1.0	*
14	Aortic aneurysm and dissection (I71)	10	0.8	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.7	*
...	All other causes (Residual)	235	18.1	871.7
American Indian, both sexes, 85-89 years				
...	All causes	1,132	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	298	26.3	---
2	Malignant neoplasms (C00-C97)	163	14.4	---
3	Cerebrovascular diseases (I60-I69)	75	6.6	---
4	Chronic lower respiratory diseases (J40-J47)	69	6.1	---
5	Diabetes mellitus (E10-E14)	59	5.2	---
6	Alzheimer's disease (G30)	57	5.0	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	45	4.0	---
8	Influenza and pneumonia (J09-J18)	41	3.6	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	35	3.1	---
10	Septicemia (A40-A41)	19	1.7	---
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	15	1.3	---
12	Parkinson's disease (G20-G21)	11	1.0	---
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.7	---
13	Pneumonitis due to solids and liquids (J69)	8	0.7	---
15	Aortic aneurysm and dissection (I71)	5	0.4	---
...	All other causes (Residual)	224	19.8	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 90-94 years				
...	All causes	604	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	173	28.6	---
2	Malignant neoplasms (C00-C97)	64	10.6	---
3	Alzheimer's disease (G30)	48	7.9	---
4	Cerebrovascular diseases (I60-I69)	43	7.1	---
5	Chronic lower respiratory diseases (J40-J47)	34	5.6	---
6	Influenza and pneumonia (J09-J18)	32	5.3	---
7	Diabetes mellitus (E10-E14)	20	3.3	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	2.8	---
9	Septicemia (A40-A41)	14	2.3	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13	2.2	---
11	Pneumonitis due to solids and liquids (J69)	12	2.0	---
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9	1.5	---
13	Parkinson's disease (G20-G21)	6	1.0	---
14	Nutritional deficiencies (E40-E64)	4	0.7	---
14	Atherosclerosis (I70)	4	0.7	---
...	All other causes (Residual)	111	18.4	---
American Indian, both sexes, 95-99 years				
...	All causes	240	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	75	31.3	---
2	Malignant neoplasms (C00-C97)	19	7.9	---
2	Alzheimer's disease (G30)	19	7.9	---
4	Cerebrovascular diseases (I60-I69)	18	7.5	---
5	Influenza and pneumonia (J09-J18)	12	5.0	---
6	Chronic lower respiratory diseases (J40-J47)	10	4.2	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6	2.5	---
8	Diabetes mellitus (E10-E14)	5	2.1	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5	2.1	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	2.1	---
11	Septicemia (A40-A41)	4	1.7	---
12	Pneumonitis due to solids and liquids (J69)	3	1.3	---
13	Peptic ulcer (K25-K28)	2	0.8	---
13	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	2	0.8	---
15	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.4	---
...	All other causes (Residual)	54	22.5	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 100 years and over				
...	All causes	66	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	25	37.9	---
2	Alzheimer's disease (G30)	6	9.1	---
3	Malignant neoplasms (C00-C97)	4	6.1	---
3	Cerebrovascular diseases (I60-I69)	4	6.1	---
5	Influenza and pneumonia (J09-J18)	3	4.5	---
5	Chronic lower respiratory diseases (J40-J47)	3	4.5	---
7	Diabetes mellitus (E10-E14)	2	3.0	---
7	Nutritional deficiencies (E40-E64)	2	3.0	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	3.0	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2	3.0	---
11	Septicemia (A40-A41)	1	1.5	---
11	Parkinson's disease (G20-G21)	1	1.5	---
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	1.5	---
...	All other causes (Residual)	10	15.2	---
American Indian, male, all ages				
...	All causes	8,516	100.0	397.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,608	18.9	75.0
2	Malignant neoplasms (C00-C97)	1,588	18.6	74.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,150	13.5	53.7
4	Diabetes mellitus (E10-E14)	432	5.1	20.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	429	5.0	20.0
6	Chronic lower respiratory diseases (J40-J47)	349	4.1	16.3
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	344	4.0	16.1
8	Cerebrovascular diseases (I60-I69)	258	3.0	12.0
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	204	2.4	9.5
10	Influenza and pneumonia (J09-J18)	172	2.0	8.0
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	153	1.8	7.1
12	Septicemia (A40-A41)	106	1.2	4.9
13	Alzheimer's disease (G30)	89	1.0	4.2
14	Certain conditions originating in the perinatal period (P00-P96)	76	0.9	3.5
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	71	0.8	3.3
...	All other causes (Residual)	1,487	17.5	69.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 1-4 years				
...	All causes	55	100.0	34.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19	34.5	*
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7	12.7	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	10.9	*
4	Malignant neoplasms (C00-C97)	4	7.3	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	5.5	*
6	Septicemia (A40-A41)	1	1.8	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.8	*
6	Chronic lower respiratory diseases (J40-J47)	1	1.8	*
6	Certain conditions originating in the perinatal period (P00-P96)	1	1.8	*
6	Complications of medical and surgical care (Y40-Y84,Y88)	1	1.8	*
...	All other causes (Residual)	11	20.0	*
American Indian, male, 5-9 years				
...	All causes	29	100.0	15.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14	48.3	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	10.3	*
3	Malignant neoplasms (C00-C97)	2	6.9	*
3	Cerebrovascular diseases (I60-I69)	2	6.9	*
5	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	3.4	*
5	Influenza and pneumonia (J09-J18)	1	3.4	*
5	Chronic lower respiratory diseases (J40-J47)	1	3.4	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1	3.4	*
...	All other causes (Residual)	4	13.8	*
American Indian, male, 10-14 years				
...	All causes	40	100.0	21.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	21	52.5	11.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8	20.0	*
3	Malignant neoplasms (C00-C97)	2	5.0	*
4	Septicemia (A40-A41)	1	2.5	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	2.5	*
...	All other causes (Residual)	7	17.5	*

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 15-19 years				
...	All causes	176	100.0	87.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	81	46.0	40.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	49	27.8	24.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	24	13.6	11.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	2.3	*
5	Cerebrovascular diseases (I60-I69)	2	1.1	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	1.1	*
7	Malignant neoplasms (C00-C97)	1	0.6	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.6	*
7	Pneumonitis due to solids and liquids (J69)	1	0.6	*
...	All other causes (Residual)	11	6.3	*
American Indian, male, 20-24 years				
...	All causes	280	100.0	147.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	120	42.9	63.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	71	25.4	37.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	45	16.1	23.7
4	Malignant neoplasms (C00-C97)	7	2.5	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	7	2.5	*
6	Legal intervention (Y35,Y89.0)	4	1.4	*
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2	0.7	*
8	Septicemia (A40-A41)	1	0.4	*
8	Diabetes mellitus (E10-E14)	1	0.4	*
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.4	*
8	Influenza and pneumonia (J09-J18)	1	0.4	*
8	Chronic lower respiratory diseases (J40-J47)	1	0.4	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.4	*
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.4	*
8	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.4	*
...	All other causes (Residual)	16	5.7	*

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 25-29 years				
...	All causes	246	100.0	139.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	103	41.9	58.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	39	15.9	22.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	32	13.0	18.2
4	Malignant neoplasms (C00-C97)	15	6.1	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	8	3.3	*
6	Chronic liver disease and cirrhosis (K70,K73-K74)	7	2.8	*
7	Human immunodeficiency virus (HIV) disease (B20-B24)	3	1.2	*
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	1.2	*
7	Diabetes mellitus (E10-E14)	2	0.8	*
9	Cerebrovascular diseases (I60-I69)	2	0.8	*
11	Septicemia (A40-A41)	1	0.4	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.4	*
11	Pneumonitis due to solids and liquids (J69)	1	0.4	*
...	All other causes (Residual)	29	11.8	16.5
American Indian, male, 30-34 years				
...	All causes	279	100.0	174.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	96	34.4	60.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	41	14.7	25.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	27	9.7	16.9
4	Chronic liver disease and cirrhosis (K70,K73-K74)	17	6.1	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	16	5.7	*
6	Malignant neoplasms (C00-C97)	10	3.6	*
7	Human immunodeficiency virus (HIV) disease (B20-B24)	6	2.2	*
8	Chronic lower respiratory diseases (J40-J47)	4	1.4	*
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	1.4	*
10	Influenza and pneumonia (J09-J18)	3	1.1	*
11	Diabetes mellitus (E10-E14)	2	0.7	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2	0.7	*
11	Cerebrovascular diseases (I60-I69)	2	0.7	*
14	Septicemia (A40-A41)	1	0.4	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.4	*
14	Aortic aneurysm and dissection (I71)	1	0.4	*
14	Pneumonitis due to solids and liquids (J69)	1	0.4	*
14	Legal intervention (Y35,Y89.0)	1	0.4	*
...	All other causes (Residual)	44	15.8	27.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 35-39 years				
...	All causes	321	100.0	216.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	95	29.6	64.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	45	14.0	30.3
3	Chronic liver disease and cirrhosis (K70,K73-K74)	42	13.1	28.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	28	8.7	18.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	14	4.4	*
6	Malignant neoplasms (C00-C97)	13	4.0	*
6	Diabetes mellitus (E10-E14)	13	4.0	*
8	Influenza and pneumonia (J09-J18)	8	2.5	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	1.2	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	3	0.9	*
11	Viral hepatitis (B15-B19)	2	0.6	*
11	Cerebrovascular diseases (I60-I69)	2	0.6	*
11	Chronic lower respiratory diseases (J40-J47)	2	0.6	*
14	Septicemia (A40-A41)	1	0.3	*
14	Meningitis (G00,G03)	1	0.3	*
14	Aortic aneurysm and dissection (I71)	1	0.3	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.3	*
...	All other causes (Residual)	46	14.3	31.0
American Indian, male, 40-44 years				
...	All causes	428	100.0	302.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	110	25.7	77.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	65	15.2	45.9
3	Chronic liver disease and cirrhosis (K70,K73-K74)	57	13.3	40.3
4	Malignant neoplasms (C00-C97)	26	6.1	18.4
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	23	5.4	16.2
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	20	4.7	14.1
7	Diabetes mellitus (E10-E14)	10	2.3	*
8	Septicemia (A40-A41)	8	1.9	*
9	Influenza and pneumonia (J09-J18)	7	1.6	*
10	Viral hepatitis (B15-B19)	6	1.4	*
10	Cerebrovascular diseases (I60-I69)	6	1.4	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	5	1.2	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	1.2	*
14	Chronic lower respiratory diseases (J40-J47)	4	0.9	*
15	Aortic aneurysm and dissection (I71)	3	0.7	*
...	All other causes (Residual)	73	17.1	51.6

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 45-49 years				
...	All causes	607	100.0	431.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	119	19.6	84.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	112	18.5	79.6
3	Chronic liver disease and cirrhosis (K70,K73-K74)	68	11.2	48.3
4	Malignant neoplasms (C00-C97)	64	10.5	45.5
5	Diabetes mellitus (E10-E14)	39	6.4	27.7
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	32	5.3	22.7
7	Influenza and pneumonia (J09-J18)	15	2.5	*
8	Cerebrovascular diseases (I60-I69)	14	2.3	*
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	13	2.1	*
10	Viral hepatitis (B15-B19)	12	2.0	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	10	1.6	*
12	Chronic lower respiratory diseases (J40-J47)	8	1.3	*
13	Septicemia (A40-A41)	7	1.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.8	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3	0.5	*
...	All other causes (Residual)	86	14.2	61.1
American Indian, male, 50-54 years				
...	All causes	704	100.0	570.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	147	20.9	119.0
2	Malignant neoplasms (C00-C97)	121	17.2	98.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	94	13.4	76.1
4	Chronic liver disease and cirrhosis (K70,K73-K74)	69	9.8	55.9
5	Diabetes mellitus (E10-E14)	38	5.4	30.8
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	22	3.1	17.8
7	Cerebrovascular diseases (I60-I69)	17	2.4	*
8	Viral hepatitis (B15-B19)	15	2.1	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	14	2.0	*
10	Influenza and pneumonia (J09-J18)	13	1.8	*
10	Chronic lower respiratory diseases (J40-J47)	13	1.8	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	1.3	*
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	9	1.3	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	8	1.1	*
15	Septicemia (A40-A41)	6	0.9	*
...	All other causes (Residual)	109	15.5	88.3

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 55-59 years				
...	All causes	837	100.0	878.5
1	Malignant neoplasms (C00-C97)	204	24.4	214.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	170	20.3	178.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	75	9.0	78.7
4	Chronic liver disease and cirrhosis (K70,K73-K74)	70	8.4	73.5
5	Diabetes mellitus (E10-E14)	59	7.0	61.9
6	Chronic lower respiratory diseases (J40-J47)	26	3.1	27.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	2.7	24.1
8	Cerebrovascular diseases (I60-I69)	18	2.2	*
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	16	1.9	*
10	Septicemia (A40-A41)	12	1.4	*
10	Viral hepatitis (B15-B19)	12	1.4	*
10	Influenza and pneumonia (J09-J18)	12	1.4	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	6	0.7	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	5	0.6	*
14	Pneumonitis due to solids and liquids (J69)	5	0.6	*
...	All other causes (Residual)	124	14.8	130.2
American Indian, male, 60-64 years				
...	All causes	753	100.0	1,047.5
1	Malignant neoplasms (C00-C97)	213	28.3	296.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	193	25.6	268.5
3	Diabetes mellitus (E10-E14)	56	7.4	77.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	53	7.0	73.7
5	Chronic lower respiratory diseases (J40-J47)	37	4.9	51.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	37	4.9	51.5
7	Cerebrovascular diseases (I60-I69)	19	2.5	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	2.1	*
9	Influenza and pneumonia (J09-J18)	11	1.5	*
10	Septicemia (A40-A41)	9	1.2	*
11	Viral hepatitis (B15-B19)	8	1.1	*
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7	0.9	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5	0.7	*
13	Pneumonitis due to solids and liquids (J69)	5	0.7	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.5	*
...	All other causes (Residual)	80	10.6	111.3

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 65-69 years				
...	All causes	738	100.0	1,591.5
1	Malignant neoplasms (C00-C97)	239	32.4	515.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	164	22.2	353.7
3	Diabetes mellitus (E10-E14)	58	7.9	125.1
4	Chronic lower respiratory diseases (J40-J47)	50	6.8	107.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	37	5.0	79.8
6	Chronic liver disease and cirrhosis (K70,K73-K74)	28	3.8	60.4
7	Cerebrovascular diseases (I60-I69)	18	2.4	*
8	Influenza and pneumonia (J09-J18)	13	1.8	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	1.5	*
10	Septicemia (A40-A41)	8	1.1	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7	0.9	*
12	Alzheimer's disease (G30)	5	0.7	*
13	Aortic aneurysm and dissection (I71)	4	0.5	*
13	Pneumonitis due to solids and liquids (J69)	4	0.5	*
15	Viral hepatitis (B15-B19)	3	0.4	*
...	All other causes (Residual)	89	12.1	191.9
American Indian, male, 70-74 years				
...	All causes	769	100.0	2,555.9
1	Malignant neoplasms (C00-C97)	218	28.3	724.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	161	20.9	535.1
3	Chronic lower respiratory diseases (J40-J47)	58	7.5	192.8
4	Diabetes mellitus (E10-E14)	44	5.7	146.2
5	Cerebrovascular diseases (I60-I69)	34	4.4	113.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	28	3.6	93.1
7	Septicemia (A40-A41)	18	2.3	*
8	Chronic liver disease and cirrhosis (K70,K73-K74)	16	2.1	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	2.1	*
10	Influenza and pneumonia (J09-J18)	14	1.8	*
11	Alzheimer's disease (G30)	12	1.6	*
12	Pneumonitis due to solids and liquids (J69)	8	1.0	*
13	Parkinson's disease (G20-G21)	5	0.7	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5	0.7	*
15	Viral hepatitis (B15-B19)	4	0.5	*
15	Peptic ulcer (K25-K28)	4	0.5	*
...	All other causes (Residual)	124	16.1	412.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 75-79 years				
...	All causes	705	100.0	3,840.9
1	Malignant neoplasms (C00-C97)	193	27.4	1,051.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	152	21.6	828.1
3	Chronic lower respiratory diseases (J40-J47)	48	6.8	261.5
4	Diabetes mellitus (E10-E14)	44	6.2	239.7
5	Cerebrovascular diseases (I60-I69)	43	6.1	234.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	25	3.5	136.2
7	Alzheimer's disease (G30)	17	2.4	*
7	Influenza and pneumonia (J09-J18)	17	2.4	*
9	Septicemia (A40-A41)	16	2.3	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	2.1	*
11	Parkinson's disease (G20-G21)	13	1.8	*
12	Chronic liver disease and cirrhosis (K70,K73-K74)	10	1.4	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7	1.0	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.9	*
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	6	0.9	*
...	All other causes (Residual)	93	13.2	506.7
American Indian, male, 80-84 years				
...	All causes	593	100.0	5,489.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	152	25.6	1,407.1
2	Malignant neoplasms (C00-C97)	129	21.8	1,194.2
3	Chronic lower respiratory diseases (J40-J47)	45	7.6	416.6
4	Cerebrovascular diseases (I60-I69)	35	5.9	324.0
5	Diabetes mellitus (E10-E14)	32	5.4	296.2
6	Alzheimer's disease (G30)	24	4.0	222.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19	3.2	*
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18	3.0	*
9	Influenza and pneumonia (J09-J18)	12	2.0	*
10	Parkinson's disease (G20-G21)	10	1.7	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	1.3	*
12	Septicemia (A40-A41)	7	1.2	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7	1.2	*
14	Aortic aneurysm and dissection (I71)	5	0.8	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	4	0.7	*
...	All other causes (Residual)	86	14.5	796.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 85-89 years				
...	All causes	471	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	135	28.7	---
2	Malignant neoplasms (C00-C97)	78	16.6	---
3	Chronic lower respiratory diseases (J40-J47)	31	6.6	---
4	Cerebrovascular diseases (I60-I69)	29	6.2	---
5	Influenza and pneumonia (J09-J18)	22	4.7	---
6	Diabetes mellitus (E10-E14)	21	4.5	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	20	4.2	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	3.6	---
9	Alzheimer's disease (G30)	16	3.4	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7	1.5	---
11	Parkinson's disease (G20-G21)	4	0.8	---
12	Septicemia (A40-A41)	3	0.6	---
12	Aortic aneurysm and dissection (I71)	3	0.6	---
12	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	3	0.6	---
15	Anemias (D50-D64)	2	0.4	---
15	Pneumonitis due to solids and liquids (J69)	2	0.4	---
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2	0.4	---
...	All other causes (Residual)	76	16.1	---
American Indian, male, 90-94 years				
...	All causes	206	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	53	25.7	---
2	Malignant neoplasms (C00-C97)	41	19.9	---
3	Influenza and pneumonia (J09-J18)	13	6.3	---
4	Alzheimer's disease (G30)	12	5.8	---
4	Chronic lower respiratory diseases (J40-J47)	12	5.8	---
6	Cerebrovascular diseases (I60-I69)	10	4.9	---
7	Diabetes mellitus (E10-E14)	9	4.4	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	3.9	---
9	Parkinson's disease (G20-G21)	5	2.4	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5	2.4	---
11	Pneumonitis due to solids and liquids (J69)	4	1.9	---
12	Septicemia (A40-A41)	3	1.5	---
13	Nutritional deficiencies (E40-E64)	2	1.0	---
14	Anemias (D50-D64)	1	0.5	---
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.5	---
14	Hernia (K40-K46)	1	0.5	---
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.5	---
...	All other causes (Residual)	25	12.1	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 95-99 years				
...	All causes	54	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	14	25.9	---
2	Malignant neoplasms (C00-C97)	7	13.0	---
3	Chronic lower respiratory diseases (J40-J47)	4	7.4	---
4	Alzheimer's disease (G30)	3	5.6	---
4	Influenza and pneumonia (J09-J18)	3	5.6	---
6	Diabetes mellitus (E10-E14)	2	3.7	---
6	Cerebrovascular diseases (I60-I69)	2	3.7	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	3.7	---
6	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	2	3.7	---
10	Septicemia (A40-A41)	1	1.9	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	1.9	---
10	Pneumonitis due to solids and liquids (J69)	1	1.9	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1	1.9	---
...	All other causes (Residual)	11	20.4	---
American Indian, male, 100 years and over				
...	All causes	12	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	25.0	---
2	Diabetes mellitus (E10-E14)	2	16.7	---
2	Influenza and pneumonia (J09-J18)	2	16.7	---
4	Nutritional deficiencies (E40-E64)	1	8.3	---
4	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	8.3	---
4	Cerebrovascular diseases (I60-I69)	1	8.3	---
...	All other causes (Residual)	2	16.7	---
American Indian, female, all ages				
...	All causes	7,049	100.0	332.4
1	Malignant neoplasms (C00-C97)	1,374	19.5	64.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,185	16.8	55.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	551	7.8	26.0
4	Diabetes mellitus (E10-E14)	425	6.0	20.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	358	5.1	16.9
6	Chronic lower respiratory diseases (J40-J47)	353	5.0	16.6
7	Cerebrovascular diseases (I60-I69)	301	4.3	14.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	186	2.6	8.8
9	Alzheimer's disease (G30)	175	2.5	8.3
10	Influenza and pneumonia (J09-J18)	154	2.2	7.3
11	Septicemia (A40-A41)	138	2.0	6.5
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	125	1.8	5.9
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	82	1.2	3.9
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	59	0.8	2.8
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	53	0.8	2.5
...	All other causes (Residual)	1,530	21.7	72.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 1-4 years				
...	All causes	38	100.0	24.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12	31.6	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	7.9	*
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	7.9	*
4	Malignant neoplasms (C00-C97)	2	5.3	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	5.3	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	2.6	*
6	Meningitis (G00,G03)	1	2.6	*
6	Cerebrovascular diseases (I60-I69)	1	2.6	*
6	Influenza and pneumonia (J09-J18)	1	2.6	*
6	Acute bronchitis and bronchiolitis (J20-J21)	1	2.6	*
6	Chronic lower respiratory diseases (J40-J47)	1	2.6	*
6	Pneumonitis due to solids and liquids (J69)	1	2.6	*
6	Legal intervention (Y35,Y89.0)	1	2.6	*
...	All other causes (Residual)	8	21.1	*
American Indian, female, 5-9 years				
...	All causes	17	100.0	*
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6	35.3	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	17.6	*
3	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	11.8	*
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	11.8	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	11.8	*
...	All other causes (Residual)	2	11.8	*
American Indian, female, 10-14 years				
...	All causes	22	100.0	12.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6	27.3	*
1	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6	27.3	*
3	Malignant neoplasms (C00-C97)	4	18.2	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	13.6	*
5	Influenza and pneumonia (J09-J18)	1	4.5	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1	4.5	*
5	Complications of medical and surgical care (Y40-Y84,Y88)	1	4.5	*
...	All other causes (Residual)	0	0.0	0.0

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 15-19 years				
...	All causes	66	100.0	34.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	29	43.9	15.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	21	31.8	11.0
3	Malignant neoplasms (C00-C97)	1	1.5	*
3	Anemias (D50-D64)	1	1.5	*
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	1.5	*
3	Cerebrovascular diseases (I60-I69)	1	1.5	*
3	Pregnancy, childbirth and the puerperium (O00-O99)	1	1.5	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1	1.5	*
...	All other causes (Residual)	10	15.2	*
American Indian, female, 20-24 years				
...	All causes	93	100.0	53.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	40	43.0	23.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	17	18.3	*
3	Malignant neoplasms (C00-C97)	5	5.4	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	5.4	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	4.3	*
5	Pregnancy, childbirth and the puerperium (O00-O99)	4	4.3	*
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2	2.2	*
8	Septicemia (A40-A41)	1	1.1	*
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	1.1	*
8	Influenza and pneumonia (J09-J18)	1	1.1	*
8	Chronic lower respiratory diseases (J40-J47)	1	1.1	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	1.1	*
...	All other causes (Residual)	11	11.8	*
American Indian, female, 25-29 years				
...	All causes	130	100.0	79.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	61	46.9	37.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	12	9.2	*
3	Chronic liver disease and cirrhosis (K70,K73-K74)	8	6.2	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7	5.4	*
5	Malignant neoplasms (C00-C97)	6	4.6	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	2.3	*
7	Cerebrovascular diseases (I60-I69)	2	1.5	*
7	Legal intervention (Y35,Y89.0)	2	1.5	*
9	Septicemia (A40-A41)	1	0.8	*
9	Diabetes mellitus (E10-E14)	1	0.8	*
9	Meningitis (G00,G03)	1	0.8	*
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.8	*
9	Pregnancy, childbirth and the puerperium (O00-O99)	1	0.8	*
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.8	*
...	All other causes (Residual)	23	17.7	14.0

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 30-34 years				
...	All causes	140	100.0	92.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	50	35.7	33.1
2	Chronic liver disease and cirrhosis (K70,K73-K74)	16	11.4	*
3	Malignant neoplasms (C00-C97)	14	10.0	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	11	7.9	*
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8	5.7	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7	5.0	*
7	Diabetes mellitus (E10-E14)	4	2.9	*
7	Influenza and pneumonia (J09-J18)	4	2.9	*
9	Septicemia (A40-A41)	2	1.4	*
9	Pregnancy, childbirth and the puerperium (O00-O99)	2	1.4	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.7	*
11	Chronic lower respiratory diseases (J40-J47)	1	0.7	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.7	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.7	*
...	All other causes (Residual)	18	12.9	*
American Indian, female, 35-39 years				
...	All causes	192	100.0	133.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	49	25.5	34.1
2	Chronic liver disease and cirrhosis (K70,K73-K74)	37	19.3	25.7
3	Malignant neoplasms (C00-C97)	18	9.4	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	16	8.3	*
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	9	4.7	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8	4.2	*
7	Diabetes mellitus (E10-E14)	5	2.6	*
8	Cerebrovascular diseases (I60-I69)	4	2.1	*
8	Pregnancy, childbirth and the puerperium (O00-O99)	4	2.1	*
10	Influenza and pneumonia (J09-J18)	3	1.6	*
11	Septicemia (A40-A41)	2	1.0	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	2	1.0	*
11	Chronic lower respiratory diseases (J40-J47)	2	1.0	*
14	Viral hepatitis (B15-B19)	1	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.5	*
...	All other causes (Residual)	31	16.1	21.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 40-44 years				
...	All causes	222	100.0	160.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	48	21.6	34.7
2	Chronic liver disease and cirrhosis (K70,K73-K74)	35	15.8	25.3
3	Malignant neoplasms (C00-C97)	22	9.9	15.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	19	8.6	*
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	15	6.8	*
6	Diabetes mellitus (E10-E14)	12	5.4	*
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	2.7	*
8	Septicemia (A40-A41)	3	1.4	*
8	Anemias (D50-D64)	3	1.4	*
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3	1.4	*
8	Cerebrovascular diseases (I60-I69)	3	1.4	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.9	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.9	*
14	Viral hepatitis (B15-B19)	1	0.5	*
14	Influenza and pneumonia (J09-J18)	1	0.5	*
14	Chronic lower respiratory diseases (J40-J47)	1	0.5	*
14	Pneumonitis due to solids and liquids (J69)	1	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.5	*
14	Pregnancy, childbirth and the puerperium (O00-O99)	1	0.5	*
...	All other causes (Residual)	43	19.4	31.1
American Indian, female, 45-49 years				
...	All causes	383	100.0	267.6
1	Chronic liver disease and cirrhosis (K70,K73-K74)	71	18.5	49.6
2	Malignant neoplasms (C00-C97)	58	15.1	40.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	51	13.3	35.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	45	11.7	31.4
5	Diabetes mellitus (E10-E14)	15	3.9	*
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	13	3.4	*
7	Viral hepatitis (B15-B19)	7	1.8	*
8	Septicemia (A40-A41)	6	1.6	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	6	1.6	*
8	Cerebrovascular diseases (I60-I69)	6	1.6	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	1.6	*
12	Chronic lower respiratory diseases (J40-J47)	5	1.3	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4	1.0	*
14	Pneumonitis due to solids and liquids (J69)	3	0.8	*
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	0.8	*
...	All other causes (Residual)	84	21.9	58.7

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 50-54 years				
...	All causes	509	100.0	390.5
1	Malignant neoplasms (C00-C97)	130	25.5	99.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	58	11.4	44.5
3	Chronic liver disease and cirrhosis (K70,K73-K74)	56	11.0	43.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	43	8.4	33.0
5	Diabetes mellitus (E10-E14)	30	5.9	23.0
6	Cerebrovascular diseases (I60-I69)	23	4.5	17.6
7	Chronic lower respiratory diseases (J40-J47)	17	3.3	*
8	Septicemia (A40-A41)	12	2.4	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	2.2	*
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	10	2.0	*
11	Viral hepatitis (B15-B19)	8	1.6	*
12	Influenza and pneumonia (J09-J18)	7	1.4	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	1.0	*
14	Pneumonitis due to solids and liquids (J69)	4	0.8	*
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	0.6	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.6	*
...	All other causes (Residual)	89	17.5	68.3
American Indian, female, 55-59 years				
...	All causes	543	100.0	532.2
1	Malignant neoplasms (C00-C97)	175	32.2	171.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	75	13.8	73.5
3	Diabetes mellitus (E10-E14)	35	6.4	34.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	33	6.1	32.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	32	5.9	31.4
6	Chronic lower respiratory diseases (J40-J47)	29	5.3	28.4
7	Septicemia (A40-A41)	16	2.9	*
8	Cerebrovascular diseases (I60-I69)	15	2.8	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	1.8	*
10	Viral hepatitis (B15-B19)	8	1.5	*
10	Influenza and pneumonia (J09-J18)	8	1.5	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5	0.9	*
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4	0.7	*
14	Cholelithiasis and other disorders of gallbladder (K80-K82)	3	0.6	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.4	*
15	Aortic aneurysm and dissection (I71)	2	0.4	*
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	0.4	*
...	All other causes (Residual)	89	16.4	87.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 60-64 years				
...	All causes	571	100.0	745.9
1	Malignant neoplasms (C00-C97)	157	27.5	205.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	84	14.7	109.7
3	Diabetes mellitus (E10-E14)	52	9.1	67.9
4	Chronic liver disease and cirrhosis (K70,K73-K74)	41	7.2	53.6
5	Chronic lower respiratory diseases (J40-J47)	29	5.1	37.9
6	Cerebrovascular diseases (I60-I69)	25	4.4	32.7
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19	3.3	*
8	Septicemia (A40-A41)	13	2.3	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	2.3	*
10	Influenza and pneumonia (J09-J18)	9	1.6	*
11	Viral hepatitis (B15-B19)	6	1.1	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5	0.9	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.5	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.5	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.4	*
15	Nutritional deficiencies (E40-E64)	2	0.4	*
15	Alzheimer's disease (G30)	2	0.4	*
15	Pneumonitis due to solids and liquids (J69)	2	0.4	*
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	0.4	*
...	All other causes (Residual)	102	17.9	133.3
American Indian, female, 65-69 years				
...	All causes	611	100.0	1,185.6
1	Malignant neoplasms (C00-C97)	184	30.1	357.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	97	15.9	188.2
3	Diabetes mellitus (E10-E14)	48	7.9	93.1
4	Chronic lower respiratory diseases (J40-J47)	46	7.5	89.3
5	Cerebrovascular diseases (I60-I69)	25	4.1	48.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	23	3.8	44.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19	3.1	*
8	Septicemia (A40-A41)	18	2.9	*
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18	2.9	*
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	12	2.0	*
11	Influenza and pneumonia (J09-J18)	10	1.6	*
12	Alzheimer's disease (G30)	7	1.1	*
13	Viral hepatitis (B15-B19)	4	0.7	*
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4	0.7	*
15	Pneumonitis due to solids and liquids (J69)	3	0.5	*
...	All other causes (Residual)	93	15.2	180.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 70-74 years				
...	All causes	685	100.0	1,906.4
1	Malignant neoplasms (C00-C97)	194	28.3	539.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	120	17.5	334.0
3	Diabetes mellitus (E10-E14)	51	7.4	141.9
3	Chronic lower respiratory diseases (J40-J47)	51	7.4	141.9
5	Cerebrovascular diseases (I60-I69)	27	3.9	75.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	26	3.8	72.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	16	2.3	*
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	16	2.3	*
9	Influenza and pneumonia (J09-J18)	13	1.9	*
10	Septicemia (A40-A41)	11	1.6	*
11	Alzheimer's disease (G30)	8	1.2	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	8	1.2	*
13	Parkinson's disease (G20-G21)	6	0.9	*
13	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	6	0.9	*
15	Pneumonitis due to solids and liquids (J69)	5	0.7	*
...	All other causes (Residual)	127	18.5	353.4
American Indian, female, 75-79 years				
...	All causes	683	100.0	2,761.3
1	Malignant neoplasms (C00-C97)	152	22.3	614.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	137	20.1	553.9
3	Diabetes mellitus (E10-E14)	62	9.1	250.7
4	Chronic lower respiratory diseases (J40-J47)	49	7.2	198.1
5	Cerebrovascular diseases (I60-I69)	31	4.5	125.3
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	28	4.1	113.2
7	Alzheimer's disease (G30)	24	3.5	97.0
8	Influenza and pneumonia (J09-J18)	22	3.2	88.9
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	16	2.3	*
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	13	1.9	*
11	Septicemia (A40-A41)	11	1.6	*
12	Chronic liver disease and cirrhosis (K70,K73-K74)	9	1.3	*
13	Aortic aneurysm and dissection (I71)	7	1.0	*
14	Parkinson's disease (G20-G21)	4	0.6	*
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	4	0.6	*
...	All other causes (Residual)	114	16.7	460.9

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 80-84 years				
...	All causes	704	100.0	4,357.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	143	20.3	885.1
2	Malignant neoplasms (C00-C97)	127	18.0	786.0
3	Diabetes mellitus (E10-E14)	58	8.2	359.0
4	Chronic lower respiratory diseases (J40-J47)	52	7.4	321.8
5	Cerebrovascular diseases (I60-I69)	40	5.7	247.6
6	Alzheimer's disease (G30)	34	4.8	210.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	27	3.8	167.1
8	Influenza and pneumonia (J09-J18)	21	3.0	130.0
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15	2.1	*
10	Septicemia (A40-A41)	10	1.4	*
10	Chronic liver disease and cirrhosis (K70,K73-K74)	10	1.4	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9	1.3	*
13	Aortic aneurysm and dissection (I71)	5	0.7	*
14	Nutritional deficiencies (E40-E64)	4	0.6	*
14	Pneumonitis due to solids and liquids (J69)	4	0.6	*
...	All other causes (Residual)	145	20.6	897.4
American Indian, female, 85-89 years				
...	All causes	661	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	163	24.7	---
2	Malignant neoplasms (C00-C97)	85	12.9	---
3	Cerebrovascular diseases (I60-I69)	46	7.0	---
4	Alzheimer's disease (G30)	41	6.2	---
5	Diabetes mellitus (E10-E14)	38	5.7	---
5	Chronic lower respiratory diseases (J40-J47)	38	5.7	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	28	4.2	---
8	Influenza and pneumonia (J09-J18)	19	2.9	---
9	Septicemia (A40-A41)	16	2.4	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15	2.3	---
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	8	1.2	---
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	1.1	---
12	Parkinson's disease (G20-G21)	7	1.1	---
14	Pneumonitis due to solids and liquids (J69)	6	0.9	---
15	Atherosclerosis (I70)	3	0.5	---
...	All other causes (Residual)	141	21.3	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 90-94 years				
...	All causes	398	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	120	30.2	---
2	Alzheimer's disease (G30)	36	9.0	---
3	Cerebrovascular diseases (I60-I69)	33	8.3	---
4	Malignant neoplasms (C00-C97)	23	5.8	---
5	Chronic lower respiratory diseases (J40-J47)	22	5.5	---
6	Influenza and pneumonia (J09-J18)	19	4.8	---
7	Septicemia (A40-A41)	11	2.8	---
7	Diabetes mellitus (E10-E14)	11	2.8	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	2.3	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	8	2.0	---
10	Pneumonitis due to solids and liquids (J69)	8	2.0	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8	2.0	---
13	Atherosclerosis (I70)	4	1.0	---
14	Tuberculosis (A16-A19)	2	0.5	---
14	Nutritional deficiencies (E40-E64)	2	0.5	---
14	Cholelithiasis and other disorders of gallbladder (K80-K82)	2	0.5	---
...	All other causes (Residual)	80	20.1	---
American Indian, female, 95-99 years				
...	All causes	186	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	61	32.8	---
2	Alzheimer's disease (G30)	16	8.6	---
2	Cerebrovascular diseases (I60-I69)	16	8.6	---
4	Malignant neoplasms (C00-C97)	12	6.5	---
5	Influenza and pneumonia (J09-J18)	9	4.8	---
6	Chronic lower respiratory diseases (J40-J47)	6	3.2	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5	2.7	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4	2.2	---
9	Septicemia (A40-A41)	3	1.6	---
9	Diabetes mellitus (E10-E14)	3	1.6	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	1.6	---
12	Pneumonitis due to solids and liquids (J69)	2	1.1	---
12	Peptic ulcer (K25-K28)	2	1.1	---
14	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.5	---
...	All other causes (Residual)	43	23.1	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 100 years and over				
...	All causes	54	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	22	40.7	---
2	Alzheimer's disease (G30)	6	11.1	---
3	Malignant neoplasms (C00-C97)	4	7.4	---
4	Cerebrovascular diseases (I60-I69)	3	5.6	---
4	Chronic lower respiratory diseases (J40-J47)	3	5.6	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	3.7	---
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2	3.7	---
8	Septicemia (A40-A41)	1	1.9	---
8	Nutritional deficiencies (E40-E64)	1	1.9	---
8	Parkinson's disease (G20-G21)	1	1.9	---
8	Influenza and pneumonia (J09-J18)	1	1.9	---
...	All other causes (Residual)	8	14.8	---
Asian or Pacific Islander, both sexes, all ages				
...	All causes	51,162	100.0	301.1
1	Malignant neoplasms (C00-C97)	14,165	27.7	83.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,490	22.5	67.6
3	Cerebrovascular diseases (I60-I69)	3,833	7.5	22.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,144	4.2	12.6
5	Diabetes mellitus (E10-E14)	1,838	3.6	10.8
6	Influenza and pneumonia (J09-J18)	1,539	3.0	9.1
7	Chronic lower respiratory diseases (J40-J47)	1,487	2.9	8.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,091	2.1	6.4
9	Alzheimer's disease (G30)	1,082	2.1	6.4
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,061	2.1	6.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	814	1.6	4.8
12	Septicemia (A40-A41)	582	1.1	3.4
13	Chronic liver disease and cirrhosis (K70,K73-K74)	467	0.9	2.7
13	Certain conditions originating in the perinatal period (P00-P96)	467	0.9	2.7
15	Parkinson's disease (G20-G21)	374	0.7	2.2
...	All other causes (Residual)	8,728	17.1	51.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 1-4 years				
...	All causes	167	100.0	17.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	42	25.1	4.5
2	Malignant neoplasms (C00-C97)	23	13.8	2.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	23	13.8	2.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	8	4.8	*
5	Chronic lower respiratory diseases (J40-J47)	5	3.0	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	3.0	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	2.4	*
7	Cerebrovascular diseases (I60-I69)	4	2.4	*
7	Influenza and pneumonia (J09-J18)	4	2.4	*
10	Septicemia (A40-A41)	2	1.2	*
11	Anemias (D50-D64)	1	0.6	*
11	Meningitis (G00,G03)	1	0.6	*
11	Acute bronchitis and bronchiolitis (J20-J21)	1	0.6	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.6	*
11	Certain conditions originating in the perinatal period (P00-P96)	1	0.6	*
...	All other causes (Residual)	42	25.1	4.5
Asian or Pacific Islander, both sexes, 5-9 years				
...	All causes	99	100.0	8.5
1	Malignant neoplasms (C00-C97)	31	31.3	2.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	16	16.2	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	8.1	*
4	Septicemia (A40-A41)	5	5.1	*
5	Chronic lower respiratory diseases (J40-J47)	4	4.0	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	3.0	*
6	Cerebrovascular diseases (I60-I69)	3	3.0	*
6	Influenza and pneumonia (J09-J18)	3	3.0	*
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	2.0	*
9	Complications of medical and surgical care (Y40-Y84,Y88)	2	2.0	*
11	Meningococcal infection (A39)	1	1.0	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.0	*
11	Anemias (D50-D64)	1	1.0	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	1.0	*
11	Certain conditions originating in the perinatal period (P00-P96)	1	1.0	*
...	All other causes (Residual)	17	17.2	*

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 10-14 years				
...	All causes	86	100.0	7.8
1	Malignant neoplasms (C00-C97)	24	27.9	2.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13	15.1	*
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7	8.1	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	7.0	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	3.5	*
5	Chronic lower respiratory diseases (J40-J47)	3	3.5	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	3.5	*
8	Septicemia (A40-A41)	2	2.3	*
8	Cerebrovascular diseases (I60-I69)	2	2.3	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.2	*
10	Anemias (D50-D64)	1	1.2	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	1.2	*
10	Certain conditions originating in the perinatal period (P00-P96)	1	1.2	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	1	1.2	*
...	All other causes (Residual)	18	20.9	*
Asian or Pacific Islander, both sexes, 15-19 years				
...	All causes	267	100.0	22.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	97	36.3	8.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	55	20.6	4.7
3	Malignant neoplasms (C00-C97)	28	10.5	2.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	23	8.6	2.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	11	4.1	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	3.0	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	1.5	*
7	Influenza and pneumonia (J09-J18)	4	1.5	*
9	Anemias (D50-D64)	2	0.7	*
9	Cerebrovascular diseases (I60-I69)	2	0.7	*
11	Tuberculosis (A16-A19)	1	0.4	*
11	Septicemia (A40-A41)	1	0.4	*
11	Aortic aneurysm and dissection (I71)	1	0.4	*
11	Chronic lower respiratory diseases (J40-J47)	1	0.4	*
11	Pregnancy, childbirth and the puerperium (O00-O99)	1	0.4	*
11	Legal intervention (Y35,Y89.0)	1	0.4	*
...	All other causes (Residual)	27	10.1	2.3

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 20-24 years				
...	All causes	485	100.0	36.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	159	32.8	12.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	126	26.0	9.6
3	Malignant neoplasms (C00-C97)	52	10.7	4.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	40	8.2	3.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	26	5.4	2.0
6	Cerebrovascular diseases (I60-I69)	7	1.4	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	1.2	*
8	Septicemia (A40-A41)	2	0.4	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.4	*
8	Influenza and pneumonia (J09-J18)	2	0.4	*
8	Chronic lower respiratory diseases (J40-J47)	2	0.4	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
8	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
14	Tuberculosis (A16-A19)	1	0.2	*
14	Scarlet fever and erysipelas (A38,A46)	1	0.2	*
14	Meningococcal infection (A39)	1	0.2	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.2	*
14	Anemias (D50-D64)	1	0.2	*
14	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.2	*
...	All other causes (Residual)	51	10.5	3.9
Asian or Pacific Islander, both sexes, 25-29 years				
...	All causes	544	100.0	37.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	154	28.3	10.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	112	20.6	7.8
3	Malignant neoplasms (C00-C97)	75	13.8	5.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	44	8.1	3.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	36	6.6	2.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	13	2.4	*
7	Pregnancy, childbirth and the puerperium (O00-O99)	10	1.8	*
8	Septicemia (A40-A41)	7	1.3	*
9	Cerebrovascular diseases (I60-I69)	6	1.1	*
10	Influenza and pneumonia (J09-J18)	4	0.7	*
10	Chronic lower respiratory diseases (J40-J47)	4	0.7	*
12	Tuberculosis (A16-A19)	3	0.6	*
12	Aortic aneurysm and dissection (I71)	3	0.6	*
12	Chronic liver disease and cirrhosis (K70,K73-K74)	3	0.6	*
15	Viral hepatitis (B15-B19)	2	0.4	*
15	Anemias (D50-D64)	2	0.4	*
15	Diabetes mellitus (E10-E14)	2	0.4	*
15	Peptic ulcer (K25-K28)	2	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
15	Legal intervention (Y35,Y89.0)	2	0.4	*
...	All other causes (Residual)	58	10.7	4.0

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 30-34 years				
...	All causes	575	100.0	40.5
1	Malignant neoplasms (C00-C97)	120	20.9	8.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	115	20.0	8.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	102	17.7	7.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	53	9.2	3.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	28	4.9	2.0
6	Cerebrovascular diseases (I60-I69)	19	3.3	*
7	Pregnancy, childbirth and the puerperium (O00-O99)	16	2.8	*
8	Chronic liver disease and cirrhosis (K70,K73-K74)	8	1.4	*
9	Influenza and pneumonia (J09-J18)	7	1.2	*
10	Chronic lower respiratory diseases (J40-J47)	5	0.9	*
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	0.9	*
12	Septicemia (A40-A41)	4	0.7	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.7	*
12	Aortic aneurysm and dissection (I71)	4	0.7	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	3	0.5	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3	0.5	*
...	All other causes (Residual)	79	13.7	5.6
Asian or Pacific Islander, both sexes, 35-39 years				
...	All causes	754	100.0	51.9
1	Malignant neoplasms (C00-C97)	219	29.0	15.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	105	13.9	7.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	94	12.5	6.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	94	12.5	6.5
5	Cerebrovascular diseases (I60-I69)	33	4.4	2.3
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	31	4.1	2.1
7	Diabetes mellitus (E10-E14)	20	2.7	1.4
8	Chronic liver disease and cirrhosis (K70,K73-K74)	14	1.9	*
9	Pregnancy, childbirth and the puerperium (O00-O99)	11	1.5	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	1.2	*
11	Influenza and pneumonia (J09-J18)	7	0.9	*
12	Septicemia (A40-A41)	6	0.8	*
12	Viral hepatitis (B15-B19)	6	0.8	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	6	0.8	*
12	Aortic aneurysm and dissection (I71)	6	0.8	*
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	0.8	*
...	All other causes (Residual)	87	11.5	6.0

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 40-44 years				
...	All causes	1,040	100.0	80.3
1	Malignant neoplasms (C00-C97)	332	31.9	25.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	158	15.2	12.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	95	9.1	7.3
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	82	7.9	6.3
5	Cerebrovascular diseases (I60-I69)	80	7.7	6.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	30	2.9	2.3
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	30	2.9	2.3
8	Diabetes mellitus (E10-E14)	19	1.8	*
9	Influenza and pneumonia (J09-J18)	14	1.3	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	1.1	*
11	Aortic aneurysm and dissection (I71)	10	1.0	*
12	Septicemia (A40-A41)	9	0.9	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	8	0.8	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	6	0.6	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	0.6	*
...	All other causes (Residual)	150	14.4	11.6
Asian or Pacific Islander, both sexes, 45-49 years				
...	All causes	1,600	100.0	132.6
1	Malignant neoplasms (C00-C97)	616	38.5	51.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	262	16.4	21.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	118	7.4	9.8
4	Cerebrovascular diseases (I60-I69)	106	6.6	8.8
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	94	5.9	7.8
6	Chronic liver disease and cirrhosis (K70,K73-K74)	39	2.4	3.2
7	Diabetes mellitus (E10-E14)	38	2.4	3.1
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	29	1.8	2.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	1.4	1.9
10	Viral hepatitis (B15-B19)	18	1.1	*
11	Influenza and pneumonia (J09-J18)	17	1.1	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	15	0.9	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	14	0.9	*
13	Aortic aneurysm and dissection (I71)	14	0.9	*
15	Septicemia (A40-A41)	12	0.8	*
...	All other causes (Residual)	185	11.6	15.3

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 50-54 years				
...	All causes	2,262	100.0	207.2
1	Malignant neoplasms (C00-C97)	958	42.4	87.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	397	17.6	36.4
3	Cerebrovascular diseases (I60-I69)	149	6.6	13.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	124	5.5	11.4
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	93	4.1	8.5
6	Diabetes mellitus (E10-E14)	81	3.6	7.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	58	2.6	5.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	34	1.5	3.1
9	Septicemia (A40-A41)	25	1.1	2.3
9	Viral hepatitis (B15-B19)	25	1.1	2.3
11	Chronic lower respiratory diseases (J40-J47)	19	0.8	*
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	17	0.8	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	16	0.7	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	14	0.6	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	0.6	*
15	Influenza and pneumonia (J09-J18)	13	0.6	*
...	All other causes (Residual)	226	10.0	20.7
Asian or Pacific Islander, both sexes, 55-59 years				
...	All causes	2,992	100.0	322.1
1	Malignant neoplasms (C00-C97)	1,311	43.8	141.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	534	17.8	57.5
3	Cerebrovascular diseases (I60-I69)	185	6.2	19.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	128	4.3	13.8
5	Diabetes mellitus (E10-E14)	113	3.8	12.2
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	73	2.4	7.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	59	2.0	6.4
8	Viral hepatitis (B15-B19)	56	1.9	6.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	43	1.4	4.6
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	36	1.2	3.9
11	Influenza and pneumonia (J09-J18)	33	1.1	3.6
12	Septicemia (A40-A41)	25	0.8	2.7
12	Chronic lower respiratory diseases (J40-J47)	25	0.8	2.7
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	23	0.8	2.5
15	Aortic aneurysm and dissection (I71)	18	0.6	*
...	All other causes (Residual)	330	11.0	35.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 60-64 years				
...	All causes	3,707	100.0	492.9
1	Malignant neoplasms (C00-C97)	1,579	42.6	209.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	747	20.2	99.3
3	Cerebrovascular diseases (I60-I69)	259	7.0	34.4
4	Diabetes mellitus (E10-E14)	167	4.5	22.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	113	3.0	15.0
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	76	2.1	10.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	64	1.7	8.5
8	Chronic lower respiratory diseases (J40-J47)	60	1.6	8.0
9	Chronic liver disease and cirrhosis (K70,K73-K74)	58	1.6	7.7
10	Influenza and pneumonia (J09-J18)	56	1.5	7.4
11	Viral hepatitis (B15-B19)	44	1.2	5.8
12	Septicemia (A40-A41)	40	1.1	5.3
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	31	0.8	4.1
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	18	0.5	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	0.4	*
...	All other causes (Residual)	381	10.3	50.7
Asian or Pacific Islander, both sexes, 65-69 years				
...	All causes	3,950	100.0	765.6
1	Malignant neoplasms (C00-C97)	1,601	40.5	310.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	828	21.0	160.5
3	Cerebrovascular diseases (I60-I69)	272	6.9	52.7
4	Diabetes mellitus (E10-E14)	178	4.5	34.5
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	127	3.2	24.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	92	2.3	17.8
7	Chronic lower respiratory diseases (J40-J47)	83	2.1	16.1
8	Influenza and pneumonia (J09-J18)	60	1.5	11.6
9	Chronic liver disease and cirrhosis (K70,K73-K74)	42	1.1	8.1
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	41	1.0	7.9
11	Septicemia (A40-A41)	38	1.0	7.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	35	0.9	6.8
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	31	0.8	6.0
14	Viral hepatitis (B15-B19)	30	0.8	5.8
15	Aortic aneurysm and dissection (I71)	28	0.7	5.4
...	All other causes (Residual)	464	11.7	89.9

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 70-74 years				
...	All causes	4,930	100.0	1,285.7
1	Malignant neoplasms (C00-C97)	1,765	35.8	460.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,070	21.7	279.1
3	Cerebrovascular diseases (I60-I69)	397	8.1	103.5
4	Diabetes mellitus (E10-E14)	221	4.5	57.6
5	Chronic lower respiratory diseases (J40-J47)	135	2.7	35.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	123	2.5	32.1
7	Influenza and pneumonia (J09-J18)	121	2.5	31.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	118	2.4	30.8
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	69	1.4	18.0
10	Septicemia (A40-A41)	59	1.2	15.4
11	Chronic liver disease and cirrhosis (K70,K73-K74)	41	0.8	10.7
12	Alzheimer's disease (G30)	40	0.8	10.4
13	Viral hepatitis (B15-B19)	39	0.8	10.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	39	0.8	10.2
15	Parkinson's disease (G20-G21)	38	0.8	9.9
...	All other causes (Residual)	655	13.3	170.8
Asian or Pacific Islander, both sexes, 75-79 years				
...	All causes	5,843	100.0	2,155.2
1	Malignant neoplasms (C00-C97)	1,830	31.3	675.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,338	22.9	493.5
3	Cerebrovascular diseases (I60-I69)	462	7.9	170.4
4	Chronic lower respiratory diseases (J40-J47)	232	4.0	85.6
5	Diabetes mellitus (E10-E14)	229	3.9	84.5
6	Influenza and pneumonia (J09-J18)	189	3.2	69.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	154	2.6	56.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	149	2.6	55.0
9	Alzheimer's disease (G30)	111	1.9	40.9
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	109	1.9	40.2
11	Septicemia (A40-A41)	79	1.4	29.1
12	Parkinson's disease (G20-G21)	64	1.1	23.6
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	54	0.9	19.9
14	Chronic liver disease and cirrhosis (K70,K73-K74)	42	0.7	15.5
15	Pneumonitis due to solids and liquids (J69)	37	0.6	13.6
...	All other causes (Residual)	764	13.1	281.8

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 80-84 years				
...	All causes	7,072	100.0	3,895.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,784	25.2	982.6
2	Malignant neoplasms (C00-C97)	1,738	24.6	957.2
3	Cerebrovascular diseases (I60-I69)	604	8.5	332.7
4	Chronic lower respiratory diseases (J40-J47)	297	4.2	163.6
5	Diabetes mellitus (E10-E14)	292	4.1	160.8
6	Influenza and pneumonia (J09-J18)	259	3.7	142.7
7	Alzheimer's disease (G30)	207	2.9	114.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	188	2.7	103.5
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	158	2.2	87.0
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	134	1.9	73.8
11	Parkinson's disease (G20-G21)	90	1.3	49.6
12	Septicemia (A40-A41)	85	1.2	46.8
13	Pneumonitis due to solids and liquids (J69)	68	1.0	37.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	50	0.7	27.5
15	Aortic aneurysm and dissection (I71)	45	0.6	24.8
...	All other causes (Residual)	1,073	15.2	591.0
Asian or Pacific Islander, both sexes, 85-89 years				
...	All causes	7,083	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,986	28.0	---
2	Malignant neoplasms (C00-C97)	1,218	17.2	---
3	Cerebrovascular diseases (I60-I69)	632	8.9	---
4	Influenza and pneumonia (J09-J18)	340	4.8	---
5	Alzheimer's disease (G30)	324	4.6	---
6	Chronic lower respiratory diseases (J40-J47)	314	4.4	---
7	Diabetes mellitus (E10-E14)	268	3.8	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	174	2.5	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	159	2.2	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	155	2.2	---
11	Septicemia (A40-A41)	102	1.4	---
12	Parkinson's disease (G20-G21)	89	1.3	---
13	Pneumonitis due to solids and liquids (J69)	63	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	53	0.7	---
15	Aortic aneurysm and dissection (I71)	43	0.6	---
...	All other causes (Residual)	1,163	16.4	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 90-94 years				
...	All causes	4,598	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,398	30.4	---
2	Malignant neoplasms (C00-C97)	495	10.8	---
3	Cerebrovascular diseases (I60-I69)	427	9.3	---
4	Influenza and pneumonia (J09-J18)	255	5.5	---
5	Alzheimer's disease (G30)	239	5.2	---
6	Chronic lower respiratory diseases (J40-J47)	191	4.2	---
7	Diabetes mellitus (E10-E14)	146	3.2	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	126	2.7	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	116	2.5	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	94	2.0	---
11	Pneumonitis due to solids and liquids (J69)	58	1.3	---
12	Septicemia (A40-A41)	49	1.1	---
13	Parkinson's disease (G20-G21)	47	1.0	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	34	0.7	---
15	Aortic aneurysm and dissection (I71)	19	0.4	---
...	All other causes (Residual)	904	19.7	---
Asian or Pacific Islander, both sexes, 95-99 years				
...	All causes	1,809	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	585	32.3	---
2	Cerebrovascular diseases (I60-I69)	150	8.3	---
3	Malignant neoplasms (C00-C97)	126	7.0	---
4	Alzheimer's disease (G30)	112	6.2	---
5	Influenza and pneumonia (J09-J18)	109	6.0	---
6	Chronic lower respiratory diseases (J40-J47)	71	3.9	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	57	3.2	---
8	Diabetes mellitus (E10-E14)	56	3.1	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	39	2.2	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	38	2.1	---
11	Pneumonitis due to solids and liquids (J69)	27	1.5	---
12	Septicemia (A40-A41)	16	0.9	---
13	Parkinson's disease (G20-G21)	13	0.7	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.5	---
14	Atherosclerosis (I70)	9	0.5	---
...	All other causes (Residual)	392	21.7	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 100 years and over				
...	All causes	414	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	144	34.8	---
2	Influenza and pneumonia (J09-J18)	30	7.2	---
3	Cerebrovascular diseases (I60-I69)	28	6.8	---
4	Alzheimer's disease (G30)	24	5.8	---
5	Malignant neoplasms (C00-C97)	19	4.6	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	17	4.1	---
7	Chronic lower respiratory diseases (J40-J47)	14	3.4	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	2.7	---
9	Diabetes mellitus (E10-E14)	7	1.7	---
10	Septicemia (A40-A41)	6	1.4	---
10	Pneumonitis due to solids and liquids (J69)	6	1.4	---
12	Atherosclerosis (I70)	5	1.2	---
12	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5	1.2	---
14	Aortic aneurysm and dissection (I71)	2	0.5	---
14	Cholelithiasis and other disorders of gallbladder (K80-K82)	2	0.5	---
...	All other causes (Residual)	94	22.7	---
Asian or Pacific Islander, male, all ages				
...	All causes	26,600	100.0	327.0
1	Malignant neoplasms (C00-C97)	7,210	27.1	88.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,262	23.5	77.0
3	Cerebrovascular diseases (I60-I69)	1,747	6.6	21.5
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,337	5.0	16.4
5	Diabetes mellitus (E10-E14)	932	3.5	11.5
6	Chronic lower respiratory diseases (J40-J47)	910	3.4	11.2
7	Influenza and pneumonia (J09-J18)	825	3.1	10.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	756	2.8	9.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	524	2.0	6.4
10	Alzheimer's disease (G30)	345	1.3	4.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	337	1.3	4.1
12	Chronic liver disease and cirrhosis (K70,K73-K74)	302	1.1	3.7
13	Septicemia (A40-A41)	296	1.1	3.6
14	Certain conditions originating in the perinatal period (P00-P96)	278	1.0	3.4
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	218	0.8	2.7
...	All other causes (Residual)	4,321	16.2	53.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 1-4 years				
...	All causes	92	100.0	19.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	26	28.3	5.5
2	Malignant neoplasms (C00-C97)	17	18.5	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	8.7	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	5.4	*
5	Cerebrovascular diseases (I60-I69)	3	3.3	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	2.2	*
6	Influenza and pneumonia (J09-J18)	2	2.2	*
6	Chronic lower respiratory diseases (J40-J47)	2	2.2	*
9	Septicemia (A40-A41)	1	1.1	*
9	Acute bronchitis and bronchiolitis (J20-J21)	1	1.1	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	1.1	*
9	Certain conditions originating in the perinatal period (P00-P96)	1	1.1	*
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1	1.1	*
...	All other causes (Residual)	22	23.9	4.6
Asian or Pacific Islander, male, 5-9 years				
...	All causes	56	100.0	9.6
1	Malignant neoplasms (C00-C97)	19	33.9	*
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12	21.4	*
3	Chronic lower respiratory diseases (J40-J47)	4	7.1	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	5.4	*
5	Septicemia (A40-A41)	2	3.6	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	3.6	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	3.6	*
8	Anemias (D50-D64)	1	1.8	*
8	Cerebrovascular diseases (I60-I69)	1	1.8	*
8	Influenza and pneumonia (J09-J18)	1	1.8	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	1.8	*
...	All other causes (Residual)	8	14.3	*
Asian or Pacific Islander, male, 10-14 years				
...	All causes	40	100.0	7.2
1	Malignant neoplasms (C00-C97)	8	20.0	*
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7	17.5	*
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5	12.5	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	10.0	*
5	Chronic lower respiratory diseases (J40-J47)	2	5.0	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	5.0	*
7	Septicemia (A40-A41)	1	2.5	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	2.5	*
7	Cerebrovascular diseases (I60-I69)	1	2.5	*
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	2.5	*
...	All other causes (Residual)	8	20.0	*

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 15-19 years				
...	All causes	176	100.0	29.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	67	38.1	11.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	38	21.6	6.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	19	10.8	*
4	Malignant neoplasms (C00-C97)	16	9.1	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	7	4.0	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	3.4	*
7	Influenza and pneumonia (J09-J18)	4	2.3	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	1.7	*
9	Septicemia (A40-A41)	1	0.6	*
9	Cerebrovascular diseases (I60-I69)	1	0.6	*
9	Chronic lower respiratory diseases (J40-J47)	1	0.6	*
9	Legal intervention (Y35,Y89.0)	1	0.6	*
...	All other causes (Residual)	12	6.8	*
Asian or Pacific Islander, male, 20-24 years				
...	All causes	368	100.0	55.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	129	35.1	19.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	100	27.2	15.0
3	Malignant neoplasms (C00-C97)	36	9.8	5.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	32	8.7	4.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	19	5.2	*
6	Cerebrovascular diseases (I60-I69)	3	0.8	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.8	*
8	Septicemia (A40-A41)	2	0.5	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.5	*
8	Chronic lower respiratory diseases (J40-J47)	2	0.5	*
11	Tuberculosis (A16-A19)	1	0.3	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.3	*
11	Anemias (D50-D64)	1	0.3	*
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.3	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.3	*
...	All other causes (Residual)	34	9.2	5.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 25-29 years				
...	All causes	369	100.0	53.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	118	32.0	17.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	88	23.8	12.8
3	Malignant neoplasms (C00-C97)	39	10.6	5.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	33	8.9	4.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	27	7.3	3.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	2.2	*
7	Cerebrovascular diseases (I60-I69)	4	1.1	*
8	Chronic lower respiratory diseases (J40-J47)	3	0.8	*
9	Viral hepatitis (B15-B19)	2	0.5	*
9	Anemias (D50-D64)	2	0.5	*
9	Aortic aneurysm and dissection (I71)	2	0.5	*
9	Influenza and pneumonia (J09-J18)	2	0.5	*
9	Legal intervention (Y35,Y89.0)	2	0.5	*
14	Tuberculosis (A16-A19)	1	0.3	*
14	Septicemia (A40-A41)	1	0.3	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.3	*
14	Diabetes mellitus (E10-E14)	1	0.3	*
14	Parkinson's disease (G20-G21)	1	0.3	*
14	Pneumonitis due to solids and liquids (J69)	1	0.3	*
14	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
...	All other causes (Residual)	31	8.4	4.5
Asian or Pacific Islander, male, 30-34 years				
...	All causes	344	100.0	51.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	85	24.7	12.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	66	19.2	9.9
3	Malignant neoplasms (C00-C97)	49	14.2	7.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	39	11.3	5.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	21	6.1	3.1
6	Cerebrovascular diseases (I60-I69)	8	2.3	*
7	Influenza and pneumonia (J09-J18)	6	1.7	*
7	Chronic liver disease and cirrhosis (K70,K73-K74)	6	1.7	*
9	Aortic aneurysm and dissection (I71)	4	1.2	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	3	0.9	*
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.9	*
12	Septicemia (A40-A41)	2	0.6	*
12	Viral hepatitis (B15-B19)	2	0.6	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.6	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2	0.6	*
12	Chronic lower respiratory diseases (J40-J47)	2	0.6	*
12	Cholelithiasis and other disorders of gallbladder (K80-K82)	2	0.6	*
...	All other causes (Residual)	42	12.2	6.3

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 35-39 years				
...	All causes	462	100.0	67.2
1	Malignant neoplasms (C00-C97)	105	22.7	15.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	80	17.3	11.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	77	16.7	11.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	64	13.9	9.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	20	4.3	2.9
6	Cerebrovascular diseases (I60-I69)	19	4.1	*
7	Chronic liver disease and cirrhosis (K70,K73-K74)	11	2.4	*
8	Diabetes mellitus (E10-E14)	10	2.2	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	5	1.1	*
10	Septicemia (A40-A41)	4	0.9	*
10	Viral hepatitis (B15-B19)	4	0.9	*
10	Influenza and pneumonia (J09-J18)	4	0.9	*
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	0.9	*
14	Chronic lower respiratory diseases (J40-J47)	3	0.6	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.6	*
...	All other causes (Residual)	49	10.6	7.1
Asian or Pacific Islander, male, 40-44 years				
...	All causes	622	100.0	101.8
1	Malignant neoplasms (C00-C97)	131	21.1	21.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	126	20.3	20.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	64	10.3	10.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	63	10.1	10.3
5	Cerebrovascular diseases (I60-I69)	48	7.7	7.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	22	3.5	3.6
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	20	3.2	3.3
8	Diabetes mellitus (E10-E14)	10	1.6	*
9	Influenza and pneumonia (J09-J18)	9	1.4	*
10	Aortic aneurysm and dissection (I71)	8	1.3	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	6	1.0	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5	0.8	*
13	Septicemia (A40-A41)	4	0.6	*
13	Viral hepatitis (B15-B19)	4	0.6	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.6	*
13	Legal intervention (Y35,Y89.0)	4	0.6	*
...	All other causes (Residual)	94	15.1	15.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 45-49 years				
...	All causes	936	100.0	164.8
1	Malignant neoplasms (C00-C97)	264	28.2	46.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	203	21.7	35.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	76	8.1	13.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	68	7.3	12.0
5	Cerebrovascular diseases (I60-I69)	65	6.9	11.4
6	Chronic liver disease and cirrhosis (K70,K73-K74)	28	3.0	4.9
7	Diabetes mellitus (E10-E14)	27	2.9	4.8
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	20	2.1	3.5
9	Viral hepatitis (B15-B19)	14	1.5	*
9	Aortic aneurysm and dissection (I71)	14	1.5	*
9	Influenza and pneumonia (J09-J18)	14	1.5	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	1.5	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	9	1.0	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9	1.0	*
15	Chronic lower respiratory diseases (J40-J47)	6	0.6	*
...	All other causes (Residual)	105	11.2	18.5
Asian or Pacific Islander, male, 50-54 years				
...	All causes	1,361	100.0	268.6
1	Malignant neoplasms (C00-C97)	460	33.8	90.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	326	24.0	64.3
3	Cerebrovascular diseases (I60-I69)	93	6.8	18.4
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	81	6.0	16.0
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	61	4.5	12.0
6	Diabetes mellitus (E10-E14)	53	3.9	10.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	44	3.2	8.7
8	Viral hepatitis (B15-B19)	20	1.5	3.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19	1.4	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	12	0.9	*
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	12	0.9	*
12	Septicemia (A40-A41)	11	0.8	*
13	Influenza and pneumonia (J09-J18)	9	0.7	*
13	Chronic lower respiratory diseases (J40-J47)	9	0.7	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7	0.5	*
...	All other causes (Residual)	144	10.6	28.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 55-59 years				
...	All causes	1,804	100.0	427.8
1	Malignant neoplasms (C00-C97)	678	37.6	160.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	384	21.3	91.1
3	Cerebrovascular diseases (I60-I69)	110	6.1	26.1
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	86	4.8	20.4
5	Diabetes mellitus (E10-E14)	84	4.7	19.9
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	54	3.0	12.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	47	2.6	11.1
8	Viral hepatitis (B15-B19)	39	2.2	9.2
9	Chronic lower respiratory diseases (J40-J47)	20	1.1	4.7
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	19	1.1	*
10	Influenza and pneumonia (J09-J18)	19	1.1	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18	1.0	*
13	Septicemia (A40-A41)	17	0.9	*
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	13	0.7	*
15	Aortic aneurysm and dissection (I71)	11	0.6	*
...	All other causes (Residual)	205	11.4	48.6
Asian or Pacific Islander, male, 60-64 years				
...	All causes	2,147	100.0	632.2
1	Malignant neoplasms (C00-C97)	808	37.6	237.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	524	24.4	154.3
3	Cerebrovascular diseases (I60-I69)	131	6.1	38.6
4	Diabetes mellitus (E10-E14)	108	5.0	31.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	59	2.7	17.4
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	52	2.4	15.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	43	2.0	12.7
8	Chronic lower respiratory diseases (J40-J47)	39	1.8	11.5
9	Influenza and pneumonia (J09-J18)	36	1.7	10.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	35	1.6	10.3
11	Viral hepatitis (B15-B19)	32	1.5	9.4
12	Septicemia (A40-A41)	22	1.0	6.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	20	0.9	5.9
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	12	0.6	*
15	Parkinson's disease (G20-G21)	10	0.5	*
...	All other causes (Residual)	216	10.1	63.6

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 65-69 years				
...	All causes	2,308	100.0	982.5
1	Malignant neoplasms (C00-C97)	874	37.9	372.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	556	24.1	236.7
3	Cerebrovascular diseases (I60-I69)	138	6.0	58.7
4	Diabetes mellitus (E10-E14)	102	4.4	43.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	66	2.9	28.1
6	Chronic lower respiratory diseases (J40-J47)	61	2.6	26.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	55	2.4	23.4
8	Influenza and pneumonia (J09-J18)	42	1.8	17.9
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	31	1.3	13.2
10	Chronic liver disease and cirrhosis (K70,K73-K74)	30	1.3	12.8
11	Septicemia (A40-A41)	26	1.1	11.1
12	Aortic aneurysm and dissection (I71)	18	0.8	*
13	Viral hepatitis (B15-B19)	13	0.6	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	13	0.6	*
15	Parkinson's disease (G20-G21)	12	0.5	*
...	All other causes (Residual)	271	11.7	115.4
Asian or Pacific Islander, male, 70-74 years				
...	All causes	2,701	100.0	1,555.4
1	Malignant neoplasms (C00-C97)	949	35.1	546.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	647	24.0	372.6
3	Cerebrovascular diseases (I60-I69)	209	7.7	120.4
4	Diabetes mellitus (E10-E14)	107	4.0	61.6
5	Chronic lower respiratory diseases (J40-J47)	87	3.2	50.1
6	Influenza and pneumonia (J09-J18)	72	2.7	41.5
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	70	2.6	40.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	61	2.3	35.1
9	Septicemia (A40-A41)	25	0.9	14.4
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	25	0.9	14.4
9	Chronic liver disease and cirrhosis (K70,K73-K74)	25	0.9	14.4
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	22	0.8	12.7
13	Alzheimer's disease (G30)	20	0.7	11.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	0.7	*
14	Parkinson's disease (G20-G21)	18	0.7	*
14	Pneumonitis due to solids and liquids (J69)	18	0.7	*
...	All other causes (Residual)	328	12.1	188.9

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 75-79 years				
...	All causes	2,976	100.0	2,598.0
1	Malignant neoplasms (C00-C97)	943	31.7	823.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	723	24.3	631.2
3	Cerebrovascular diseases (I60-I69)	221	7.4	192.9
4	Chronic lower respiratory diseases (J40-J47)	148	5.0	129.2
5	Diabetes mellitus (E10-E14)	106	3.6	92.5
6	Influenza and pneumonia (J09-J18)	103	3.5	89.9
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	75	2.5	65.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	67	2.3	58.5
9	Septicemia (A40-A41)	47	1.6	41.0
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	43	1.4	37.5
11	Alzheimer's disease (G30)	42	1.4	36.7
12	Parkinson's disease (G20-G21)	37	1.2	32.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	0.8	20.1
14	Pneumonitis due to solids and liquids (J69)	20	0.7	17.5
15	Aortic aneurysm and dissection (I71)	19	0.6	*
...	All other causes (Residual)	359	12.1	313.4
Asian or Pacific Islander, male, 80-84 years				
...	All causes	3,411	100.0	4,791.6
1	Malignant neoplasms (C00-C97)	878	25.7	1,233.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	866	25.4	1,216.5
3	Cerebrovascular diseases (I60-I69)	262	7.7	368.0
4	Chronic lower respiratory diseases (J40-J47)	188	5.5	264.1
5	Influenza and pneumonia (J09-J18)	137	4.0	192.5
6	Diabetes mellitus (E10-E14)	129	3.8	181.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	98	2.9	137.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	83	2.4	116.6
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	65	1.9	91.3
10	Alzheimer's disease (G30)	54	1.6	75.9
11	Parkinson's disease (G20-G21)	49	1.4	68.8
12	Septicemia (A40-A41)	41	1.2	57.6
13	Pneumonitis due to solids and liquids (J69)	38	1.1	53.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	31	0.9	43.5
15	Aortic aneurysm and dissection (I71)	25	0.7	35.1
...	All other causes (Residual)	467	13.7	656.0

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 85-89 years				
...	All causes	3,263	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	920	28.2	---
2	Malignant neoplasms (C00-C97)	634	19.4	---
3	Cerebrovascular diseases (I60-I69)	229	7.0	---
4	Chronic lower respiratory diseases (J40-J47)	179	5.5	---
5	Influenza and pneumonia (J09-J18)	167	5.1	---
6	Diabetes mellitus (E10-E14)	111	3.4	---
7	Alzheimer's disease (G30)	103	3.2	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	82	2.5	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	70	2.1	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	60	1.8	---
11	Septicemia (A40-A41)	51	1.6	---
12	Parkinson's disease (G20-G21)	44	1.3	---
13	Pneumonitis due to solids and liquids (J69)	39	1.2	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	28	0.9	---
15	Aortic aneurysm and dissection (I71)	21	0.6	---
...	All other causes (Residual)	525	16.1	---
Asian or Pacific Islander, male, 90-94 years				
...	All causes	1,876	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	548	29.2	---
2	Malignant neoplasms (C00-C97)	233	12.4	---
3	Cerebrovascular diseases (I60-I69)	142	7.6	---
4	Influenza and pneumonia (J09-J18)	128	6.8	---
5	Chronic lower respiratory diseases (J40-J47)	109	5.8	---
6	Alzheimer's disease (G30)	81	4.3	---
7	Diabetes mellitus (E10-E14)	64	3.4	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	53	2.8	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	44	2.3	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	37	2.0	---
11	Pneumonitis due to solids and liquids (J69)	30	1.6	---
12	Parkinson's disease (G20-G21)	28	1.5	---
13	Septicemia (A40-A41)	25	1.3	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	15	0.8	---
15	Aortic aneurysm and dissection (I71)	8	0.4	---
15	Peptic ulcer (K25-K28)	8	0.4	---
...	All other causes (Residual)	323	17.2	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 95-99 years				
...	All causes	639	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	199	31.1	---
2	Malignant neoplasms (C00-C97)	60	9.4	---
3	Influenza and pneumonia (J09-J18)	49	7.7	---
4	Cerebrovascular diseases (I60-I69)	47	7.4	---
5	Chronic lower respiratory diseases (J40-J47)	34	5.3	---
6	Alzheimer's disease (G30)	28	4.4	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	20	3.1	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18	2.8	---
9	Pneumonitis due to solids and liquids (J69)	15	2.3	---
10	Diabetes mellitus (E10-E14)	14	2.2	---
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	1.9	---
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	1.3	---
13	Septicemia (A40-A41)	5	0.8	---
13	Parkinson's disease (G20-G21)	5	0.8	---
15	Atherosclerosis (I70)	4	0.6	---
...	All other causes (Residual)	121	18.9	---
Asian or Pacific Islander, male, 100 years and over				
...	All causes	144	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	47	32.6	---
2	Influenza and pneumonia (J09-J18)	14	9.7	---
3	Alzheimer's disease (G30)	8	5.6	---
3	Cerebrovascular diseases (I60-I69)	8	5.6	---
5	Malignant neoplasms (C00-C97)	7	4.9	---
5	Chronic lower respiratory diseases (J40-J47)	7	4.9	---
7	Diabetes mellitus (E10-E14)	5	3.5	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4	2.8	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	2.8	---
10	Septicemia (A40-A41)	2	1.4	---
10	Atherosclerosis (I70)	2	1.4	---
10	Pneumonitis due to solids and liquids (J69)	2	1.4	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2	1.4	---
14	Aortic aneurysm and dissection (I71)	1	0.7	---
14	Hyperplasia of prostate (N40)	1	0.7	---
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1	0.7	---
14	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.7	---
...	All other causes (Residual)	28	19.4	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, all ages				
...	All causes	24,562	100.0	277.3
1	Malignant neoplasms (C00-C97)	6,955	28.3	78.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,228	21.3	59.0
3	Cerebrovascular diseases (I60-I69)	2,086	8.5	23.5
4	Diabetes mellitus (E10-E14)	906	3.7	10.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	807	3.3	9.1
6	Alzheimer's disease (G30)	737	3.0	8.3
7	Influenza and pneumonia (J09-J18)	714	2.9	8.1
8	Chronic lower respiratory diseases (J40-J47)	577	2.3	6.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	567	2.3	6.4
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	477	1.9	5.4
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	305	1.2	3.4
12	Septicemia (A40-A41)	286	1.2	3.2
13	Certain conditions originating in the perinatal period (P00-P96)	189	0.8	2.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	178	0.7	2.0
15	Parkinson's disease (G20-G21)	168	0.7	1.9
...	All other causes (Residual)	4,382	17.8	49.5
Asian or Pacific Islander, female, 1-4 years				
...	All causes	75	100.0	16.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	16	21.3	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	15	20.0	*
3	Malignant neoplasms (C00-C97)	6	8.0	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	5.3	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	4.0	*
5	Chronic lower respiratory diseases (J40-J47)	3	4.0	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	2.7	*
7	Influenza and pneumonia (J09-J18)	2	2.7	*
9	Septicemia (A40-A41)	1	1.3	*
9	Anemias (D50-D64)	1	1.3	*
9	Meningitis (G00,G03)	1	1.3	*
9	Cerebrovascular diseases (I60-I69)	1	1.3	*
...	All other causes (Residual)	20	26.7	4.3

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 5-9 years				
...	All causes	43	100.0	7.4
1	Malignant neoplasms (C00-C97)	12	27.9	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	11.6	*
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4	9.3	*
4	Septicemia (A40-A41)	3	7.0	*
5	Cerebrovascular diseases (I60-I69)	2	4.7	*
5	Influenza and pneumonia (J09-J18)	2	4.7	*
5	Complications of medical and surgical care (Y40-Y84,Y88)	2	4.7	*
8	Meningococcal infection (A39)	1	2.3	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	2.3	*
8	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	2.3	*
8	Certain conditions originating in the perinatal period (P00-P96)	1	2.3	*
...	All other causes (Residual)	9	20.9	*
Asian or Pacific Islander, female, 10-14 years				
...	All causes	46	100.0	8.5
1	Malignant neoplasms (C00-C97)	16	34.8	*
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6	13.0	*
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	4.3	*
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2	4.3	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	4.3	*
6	Septicemia (A40-A41)	1	2.2	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	2.2	*
6	Anemias (D50-D64)	1	2.2	*
6	Cerebrovascular diseases (I60-I69)	1	2.2	*
6	Chronic lower respiratory diseases (J40-J47)	1	2.2	*
6	Certain conditions originating in the perinatal period (P00-P96)	1	2.2	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	2.2	*
6	Complications of medical and surgical care (Y40-Y84,Y88)	1	2.2	*
...	All other causes (Residual)	10	21.7	*

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 15-19 years				
...	All causes	91	100.0	15.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	30	33.0	5.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	17	18.7	*
3	Malignant neoplasms (C00-C97)	12	13.2	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	4.4	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	4.4	*
6	Anemias (D50-D64)	2	2.2	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	2.2	*
8	Tuberculosis (A16-A19)	1	1.1	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.1	*
8	Cerebrovascular diseases (I60-I69)	1	1.1	*
8	Aortic aneurysm and dissection (I71)	1	1.1	*
8	Pregnancy, childbirth and the puerperium (O00-O99)	1	1.1	*
...	All other causes (Residual)	15	16.5	*
Asian or Pacific Islander, female, 20-24 years				
...	All causes	117	100.0	18.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	30	25.6	4.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	26	22.2	4.0
3	Malignant neoplasms (C00-C97)	16	13.7	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8	6.8	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	7	6.0	*
6	Cerebrovascular diseases (I60-I69)	4	3.4	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	2.6	*
8	Influenza and pneumonia (J09-J18)	2	1.7	*
9	Scarlet fever and erysipelas (A38,A46)	1	0.9	*
9	Meningococcal infection (A39)	1	0.9	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.9	*
9	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.9	*
...	All other causes (Residual)	17	14.5	*

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 25-29 years				
...	All causes	175	100.0	23.4
1	Malignant neoplasms (C00-C97)	36	20.6	4.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	36	20.6	4.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	24	13.7	3.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	11	6.3	*
5	Pregnancy, childbirth and the puerperium (O00-O99)	10	5.7	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	9	5.1	*
7	Septicemia (A40-A41)	6	3.4	*
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	2.9	*
9	Tuberculosis (A16-A19)	2	1.1	*
9	Cerebrovascular diseases (I60-I69)	2	1.1	*
9	Influenza and pneumonia (J09-J18)	2	1.1	*
9	Peptic ulcer (K25-K28)	2	1.1	*
9	Chronic liver disease and cirrhosis (K70,K73-K74)	2	1.1	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.6	*
14	Diabetes mellitus (E10-E14)	1	0.6	*
14	Aortic aneurysm and dissection (I71)	1	0.6	*
14	Chronic lower respiratory diseases (J40-J47)	1	0.6	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.6	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.6	*
...	All other causes (Residual)	22	12.6	2.9
Asian or Pacific Islander, female, 30-34 years				
...	All causes	231	100.0	30.8
1	Malignant neoplasms (C00-C97)	71	30.7	9.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	36	15.6	4.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	30	13.0	4.0
4	Pregnancy, childbirth and the puerperium (O00-O99)	16	6.9	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	14	6.1	*
6	Cerebrovascular diseases (I60-I69)	11	4.8	*
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7	3.0	*
8	Chronic lower respiratory diseases (J40-J47)	3	1.3	*
9	Septicemia (A40-A41)	2	0.9	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.9	*
9	Chronic liver disease and cirrhosis (K70,K73-K74)	2	0.9	*
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	0.9	*
13	Meningitis (G00,G03)	1	0.4	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.4	*
13	Influenza and pneumonia (J09-J18)	1	0.4	*
13	Pneumonitis due to solids and liquids (J69)	1	0.4	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.4	*
...	All other causes (Residual)	29	12.6	3.9

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 35-39 years				
...	All causes	292	100.0	38.1
1	Malignant neoplasms (C00-C97)	114	39.0	14.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	30	10.3	3.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	28	9.6	3.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	14	4.8	*
4	Cerebrovascular diseases (I60-I69)	14	4.8	*
6	Pregnancy, childbirth and the puerperium (O00-O99)	11	3.8	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	11	3.8	*
8	Diabetes mellitus (E10-E14)	10	3.4	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	2.1	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	1.4	*
10	Aortic aneurysm and dissection (I71)	4	1.4	*
12	Influenza and pneumonia (J09-J18)	3	1.0	*
12	Chronic liver disease and cirrhosis (K70,K73-K74)	3	1.0	*
14	Septicemia (A40-A41)	2	0.7	*
14	Viral hepatitis (B15-B19)	2	0.7	*
14	Chronic lower respiratory diseases (J40-J47)	2	0.7	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	0.7	*
...	All other causes (Residual)	32	11.0	4.2
Asian or Pacific Islander, female, 40-44 years				
...	All causes	418	100.0	61.1
1	Malignant neoplasms (C00-C97)	201	48.1	29.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	32	7.7	4.7
2	Cerebrovascular diseases (I60-I69)	32	7.7	4.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	32	7.7	4.7
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18	4.3	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	10	2.4	*
7	Diabetes mellitus (E10-E14)	9	2.2	*
8	Chronic liver disease and cirrhosis (K70,K73-K74)	8	1.9	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	1.7	*
10	Septicemia (A40-A41)	5	1.2	*
10	Influenza and pneumonia (J09-J18)	5	1.2	*
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	1.2	*
13	Chronic lower respiratory diseases (J40-J47)	3	0.7	*
13	Pregnancy, childbirth and the puerperium (O00-O99)	3	0.7	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.5	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.5	*
15	Aortic aneurysm and dissection (I71)	2	0.5	*
...	All other causes (Residual)	42	10.0	6.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 45-49 years				
...	All causes	664	100.0	104.0
1	Malignant neoplasms (C00-C97)	352	53.0	55.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	59	8.9	9.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	42	6.3	6.6
4	Cerebrovascular diseases (I60-I69)	41	6.2	6.4
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	26	3.9	4.1
6	Diabetes mellitus (E10-E14)	11	1.7	*
6	Chronic liver disease and cirrhosis (K70,K73-K74)	11	1.7	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	1.4	*
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	9	1.4	*
10	Septicemia (A40-A41)	7	1.1	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	1.1	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	6	0.9	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	5	0.8	*
14	Viral hepatitis (B15-B19)	4	0.6	*
14	Chronic lower respiratory diseases (J40-J47)	4	0.6	*
...	All other causes (Residual)	71	10.7	11.1
Asian or Pacific Islander, female, 50-54 years				
...	All causes	901	100.0	154.0
1	Malignant neoplasms (C00-C97)	498	55.3	85.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	71	7.9	12.1
3	Cerebrovascular diseases (I60-I69)	56	6.2	9.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	43	4.8	7.4
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	32	3.6	5.5
6	Diabetes mellitus (E10-E14)	28	3.1	4.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	1.7	*
8	Septicemia (A40-A41)	14	1.6	*
8	Chronic liver disease and cirrhosis (K70,K73-K74)	14	1.6	*
10	Chronic lower respiratory diseases (J40-J47)	10	1.1	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9	1.0	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.8	*
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	0.8	*
14	Viral hepatitis (B15-B19)	5	0.6	*
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	0.6	*
...	All other causes (Residual)	87	9.7	14.9

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 55-59 years				
...	All causes	1,188	100.0	234.2
1	Malignant neoplasms (C00-C97)	633	53.3	124.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	150	12.6	29.6
3	Cerebrovascular diseases (I60-I69)	75	6.3	14.8
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	42	3.5	8.3
5	Diabetes mellitus (E10-E14)	29	2.4	5.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	25	2.1	4.9
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	19	1.6	*
8	Viral hepatitis (B15-B19)	17	1.4	*
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	17	1.4	*
10	Influenza and pneumonia (J09-J18)	14	1.2	*
11	Chronic liver disease and cirrhosis (K70,K73-K74)	12	1.0	*
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	10	0.8	*
13	Septicemia (A40-A41)	8	0.7	*
14	Aortic aneurysm and dissection (I71)	7	0.6	*
15	Anemias (D50-D64)	6	0.5	*
...	All other causes (Residual)	124	10.4	24.4
Asian or Pacific Islander, female, 60-64 years				
...	All causes	1,560	100.0	378.1
1	Malignant neoplasms (C00-C97)	771	49.4	186.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	223	14.3	54.1
3	Cerebrovascular diseases (I60-I69)	128	8.2	31.0
4	Diabetes mellitus (E10-E14)	59	3.8	14.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	54	3.5	13.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	29	1.9	7.0
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	24	1.5	5.8
8	Chronic lower respiratory diseases (J40-J47)	21	1.3	5.1
9	Influenza and pneumonia (J09-J18)	20	1.3	4.8
10	Septicemia (A40-A41)	18	1.2	*
11	Chronic liver disease and cirrhosis (K70,K73-K74)	15	1.0	*
12	Viral hepatitis (B15-B19)	12	0.8	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	11	0.7	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.4	*
15	Aortic aneurysm and dissection (I71)	6	0.4	*
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	0.4	*
...	All other causes (Residual)	156	10.0	37.8

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 65-69 years				
...	All causes	1,642	100.0	584.3
1	Malignant neoplasms (C00-C97)	727	44.3	258.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	272	16.6	96.8
3	Cerebrovascular diseases (I60-I69)	134	8.2	47.7
4	Diabetes mellitus (E10-E14)	76	4.6	27.0
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	61	3.7	21.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	37	2.3	13.2
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	22	1.3	7.8
7	Chronic lower respiratory diseases (J40-J47)	22	1.3	7.8
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	21	1.3	7.5
10	Influenza and pneumonia (J09-J18)	18	1.1	*
11	Viral hepatitis (B15-B19)	17	1.0	*
12	Septicemia (A40-A41)	12	0.7	*
12	Chronic liver disease and cirrhosis (K70,K73-K74)	12	0.7	*
14	Alzheimer's disease (G30)	10	0.6	*
14	Aortic aneurysm and dissection (I71)	10	0.6	*
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	10	0.6	*
...	All other causes (Residual)	181	11.0	64.4
Asian or Pacific Islander, female, 70-74 years				
...	All causes	2,229	100.0	1,062.5
1	Malignant neoplasms (C00-C97)	816	36.6	389.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	423	19.0	201.6
3	Cerebrovascular diseases (I60-I69)	188	8.4	89.6
4	Diabetes mellitus (E10-E14)	114	5.1	54.3
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	57	2.6	27.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	53	2.4	25.3
7	Influenza and pneumonia (J09-J18)	49	2.2	23.4
8	Chronic lower respiratory diseases (J40-J47)	48	2.2	22.9
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	44	2.0	21.0
10	Septicemia (A40-A41)	34	1.5	16.2
11	Viral hepatitis (B15-B19)	24	1.1	11.4
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	21	0.9	10.0
13	Parkinson's disease (G20-G21)	20	0.9	9.5
13	Alzheimer's disease (G30)	20	0.9	9.5
15	Chronic liver disease and cirrhosis (K70,K73-K74)	16	0.7	*
...	All other causes (Residual)	302	13.5	144.0

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 75-79 years				
...	All causes	2,867	100.0	1,831.3
1	Malignant neoplasms (C00-C97)	887	30.9	566.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	615	21.5	392.8
3	Cerebrovascular diseases (I60-I69)	241	8.4	153.9
4	Diabetes mellitus (E10-E14)	123	4.3	78.6
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	87	3.0	55.6
6	Influenza and pneumonia (J09-J18)	86	3.0	54.9
7	Chronic lower respiratory diseases (J40-J47)	84	2.9	53.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	74	2.6	47.3
9	Alzheimer's disease (G30)	69	2.4	44.1
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	66	2.3	42.2
11	Septicemia (A40-A41)	32	1.1	20.4
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	31	1.1	19.8
13	Parkinson's disease (G20-G21)	27	0.9	17.2
14	Chronic liver disease and cirrhosis (K70,K73-K74)	26	0.9	16.6
15	Viral hepatitis (B15-B19)	22	0.8	14.1
...	All other causes (Residual)	397	13.8	253.6
Asian or Pacific Islander, female, 80-84 years				
...	All causes	3,661	100.0	3,316.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	918	25.1	831.7
2	Malignant neoplasms (C00-C97)	860	23.5	779.2
3	Cerebrovascular diseases (I60-I69)	342	9.3	309.8
4	Diabetes mellitus (E10-E14)	163	4.5	147.7
5	Alzheimer's disease (G30)	153	4.2	138.6
6	Influenza and pneumonia (J09-J18)	122	3.3	110.5
7	Chronic lower respiratory diseases (J40-J47)	109	3.0	98.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	90	2.5	81.5
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	75	2.0	67.9
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	69	1.9	62.5
11	Septicemia (A40-A41)	44	1.2	39.9
12	Parkinson's disease (G20-G21)	41	1.1	37.1
13	Pneumonitis due to solids and liquids (J69)	30	0.8	27.2
14	Chronic liver disease and cirrhosis (K70,K73-K74)	23	0.6	20.8
15	Aortic aneurysm and dissection (I71)	20	0.5	18.1
...	All other causes (Residual)	602	16.4	545.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 85-89 years				
...	All causes	3,820	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,066	27.9	---
2	Malignant neoplasms (C00-C97)	584	15.3	---
3	Cerebrovascular diseases (I60-I69)	403	10.5	---
4	Alzheimer's disease (G30)	221	5.8	---
5	Influenza and pneumonia (J09-J18)	173	4.5	---
6	Diabetes mellitus (E10-E14)	157	4.1	---
7	Chronic lower respiratory diseases (J40-J47)	135	3.5	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	104	2.7	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	99	2.6	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	73	1.9	---
11	Septicemia (A40-A41)	51	1.3	---
12	Parkinson's disease (G20-G21)	45	1.2	---
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	25	0.7	---
14	Pneumonitis due to solids and liquids (J69)	24	0.6	---
15	Aortic aneurysm and dissection (I71)	22	0.6	---
...	All other causes (Residual)	638	16.7	---
Asian or Pacific Islander, female, 90-94 years				
...	All causes	2,722	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	850	31.2	---
2	Cerebrovascular diseases (I60-I69)	285	10.5	---
3	Malignant neoplasms (C00-C97)	262	9.6	---
4	Alzheimer's disease (G30)	158	5.8	---
5	Influenza and pneumonia (J09-J18)	127	4.7	---
6	Diabetes mellitus (E10-E14)	82	3.0	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	82	3.0	---
6	Chronic lower respiratory diseases (J40-J47)	82	3.0	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	63	2.3	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	57	2.1	---
11	Pneumonitis due to solids and liquids (J69)	28	1.0	---
12	Septicemia (A40-A41)	24	0.9	---
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	0.7	---
13	Parkinson's disease (G20-G21)	19	0.7	---
15	Atherosclerosis (I70)	12	0.4	---
...	All other causes (Residual)	572	21.0	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 95-99 years				
...	All causes	1,170	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	386	33.0	---
2	Cerebrovascular diseases (I60-I69)	103	8.8	---
3	Alzheimer's disease (G30)	84	7.2	---
4	Malignant neoplasms (C00-C97)	66	5.6	---
5	Influenza and pneumonia (J09-J18)	60	5.1	---
6	Diabetes mellitus (E10-E14)	42	3.6	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	37	3.2	---
7	Chronic lower respiratory diseases (J40-J47)	37	3.2	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	26	2.2	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	21	1.8	---
11	Pneumonitis due to solids and liquids (J69)	12	1.0	---
12	Septicemia (A40-A41)	11	0.9	---
13	Parkinson's disease (G20-G21)	8	0.7	---
14	Atherosclerosis (I70)	5	0.4	---
14	Aortic aneurysm and dissection (I71)	5	0.4	---
...	All other causes (Residual)	267	22.8	---
Asian or Pacific Islander, female, 100 years and over				
...	All causes	270	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	97	35.9	---
2	Cerebrovascular diseases (I60-I69)	20	7.4	---
3	Alzheimer's disease (G30)	16	5.9	---
3	Influenza and pneumonia (J09-J18)	16	5.9	---
5	Essential hypertension and hypertensive renal disease (I10,I12,I15)	13	4.8	---
6	Malignant neoplasms (C00-C97)	12	4.4	---
7	Chronic lower respiratory diseases (J40-J47)	7	2.6	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	2.6	---
9	Septicemia (A40-A41)	4	1.5	---
9	Pneumonitis due to solids and liquids (J69)	4	1.5	---
11	Atherosclerosis (I70)	3	1.1	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3	1.1	---
13	Diabetes mellitus (E10-E14)	2	0.7	---
13	Cholelithiasis and other disorders of gallbladder (K80-K82)	2	0.7	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.4	---
15	Aortic aneurysm and dissection (I71)	1	0.4	---
15	Peptic ulcer (K25-K28)	1	0.4	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.4	---
...	All other causes (Residual)	60	22.2	---

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths.

\* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available