

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, all ages				
...	All causes	2,515,458	100.0	807.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	596,577	23.7	191.5
2	Malignant neoplasms (C00-C97)	576,691	22.9	185.1
3	Chronic lower respiratory diseases (J40-J47)	142,943	5.7	45.9
4	Cerebrovascular diseases (I60-I69)	128,932	5.1	41.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	126,438	5.0	40.6
6	Alzheimer's disease (G30)	84,974	3.4	27.3
7	Diabetes mellitus (E10-E14)	73,831	2.9	23.7
8	Influenza and pneumonia (J09-J18)	53,826	2.1	17.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	45,591	1.8	14.6
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	39,518	1.6	12.7
11	Septicemia (A40-A41)	35,748	1.4	11.5
12	Chronic liver disease and cirrhosis (K70,K73-K74)	33,642	1.3	10.8
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	27,853	1.1	8.9
14	Parkinson's disease (G20-G21)	23,111	0.9	7.4
15	Pneumonitis due to solids and liquids (J69)	18,195	0.7	5.8
...	All other causes (Residual)	507,588	20.2	162.9
All races, both sexes, 1-4 years				
...	All causes	4,246	100.0	26.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,377	32.4	8.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	493	11.6	3.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	412	9.7	2.5
4	Malignant neoplasms (C00-C97)	353	8.3	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	165	3.9	1.0
6	Influenza and pneumonia (J09-J18)	112	2.6	0.7
7	Septicemia (A40-A41)	61	1.4	0.4
8	Chronic lower respiratory diseases (J40-J47)	53	1.2	0.3
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	45	1.1	0.3
10	Cerebrovascular diseases (I60-I69)	42	1.0	0.3
11	Certain conditions originating in the perinatal period (P00-P96)	34	0.8	0.2
12	Complications of medical and surgical care (Y40-Y84,Y88)	22	0.5	0.1
13	Anemias (D50-D64)	14	0.3	*
13	Acute bronchitis and bronchiolitis (J20-J21)	14	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	0.3	*
...	All other causes (Residual)	1,036	24.4	6.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 5-9 years				
...	All causes	2,451	100.0	12.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	761	31.0	3.7
2	Malignant neoplasms (C00-C97)	441	18.0	2.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	182	7.4	0.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	129	5.3	0.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	92	3.8	0.5
6	Chronic lower respiratory diseases (J40-J47)	64	2.6	0.3
7	Influenza and pneumonia (J09-J18)	63	2.6	0.3
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	40	1.6	0.2
8	Cerebrovascular diseases (I60-I69)	40	1.6	0.2
10	Septicemia (A40-A41)	38	1.6	0.2
11	Anemias (D50-D64)	28	1.1	0.1
12	Certain conditions originating in the perinatal period (P00-P96)	19	0.8	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.4	*
14	Meningitis (G00,G03)	7	0.3	*
15	Pneumonitis due to solids and liquids (J69)	6	0.2	*
15	Diseases of appendix (K35-K38)	6	0.2	*
...	All other causes (Residual)	526	21.5	2.6
All races, both sexes, 10-14 years				
...	All causes	2,950	100.0	14.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	874	29.6	4.2
2	Malignant neoplasms (C00-C97)	419	14.2	2.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	282	9.6	1.4
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	176	6.0	0.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	154	5.2	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	111	3.8	0.5
7	Chronic lower respiratory diseases (J40-J47)	72	2.4	0.3
8	Influenza and pneumonia (J09-J18)	55	1.9	0.3
9	Cerebrovascular diseases (I60-I69)	47	1.6	0.2
10	Septicemia (A40-A41)	31	1.1	0.1
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	30	1.0	0.1
12	Diabetes mellitus (E10-E14)	25	0.8	0.1
13	Anemias (D50-D64)	15	0.5	*
14	Pneumonitis due to solids and liquids (J69)	12	0.4	*
15	Certain conditions originating in the perinatal period (P00-P96)	11	0.4	*
...	All other causes (Residual)	636	21.6	3.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 15-19 years				
...	All causes	10,594	100.0	48.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,298	40.6	19.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,802	17.0	8.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,680	15.9	7.8
4	Malignant neoplasms (C00-C97)	639	6.0	3.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	318	3.0	1.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	207	2.0	1.0
7	Cerebrovascular diseases (I60-I69)	69	0.7	0.3
7	Chronic lower respiratory diseases (J40-J47)	69	0.7	0.3
9	Influenza and pneumonia (J09-J18)	66	0.6	0.3
10	Diabetes mellitus (E10-E14)	49	0.5	0.2
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	44	0.4	0.2
12	Septicemia (A40-A41)	42	0.4	0.2
13	Pregnancy, childbirth and the puerperium (O00-O99)	35	0.3	0.2
14	Anemias (D50-D64)	30	0.3	0.1
15	Legal intervention (Y35,Y89.0)	29	0.3	0.1
...	All other causes (Residual)	1,217	11.5	5.6
All races, both sexes, 20-24 years				
...	All causes	19,073	100.0	86.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,032	42.1	36.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,020	15.8	13.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,874	15.1	13.0
4	Malignant neoplasms (C00-C97)	972	5.1	4.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	680	3.6	3.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	225	1.2	1.0
7	Influenza and pneumonia (J09-J18)	154	0.8	0.7
8	Pregnancy, childbirth and the puerperium (O00-O99)	137	0.7	0.6
9	Diabetes mellitus (E10-E14)	118	0.6	0.5
10	Human immunodeficiency virus (HIV) disease (B20-B24)	117	0.6	0.5
10	Cerebrovascular diseases (I60-I69)	117	0.6	0.5
12	Chronic lower respiratory diseases (J40-J47)	101	0.5	0.5
13	Septicemia (A40-A41)	92	0.5	0.4
14	Legal intervention (Y35,Y89.0)	73	0.4	0.3
15	Anemias (D50-D64)	67	0.4	0.3
...	All other causes (Residual)	2,294	12.0	10.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 25-29 years				
...	All causes	20,572	100.0	96.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,961	38.7	37.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,091	15.0	14.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,373	11.5	11.2
4	Malignant neoplasms (C00-C97)	1,319	6.4	6.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,182	5.7	5.6
6	Diabetes mellitus (E10-E14)	237	1.2	1.1
6	Influenza and pneumonia (J09-J18)	237	1.2	1.1
8	Human immunodeficiency virus (HIV) disease (B20-B24)	229	1.1	1.1
9	Pregnancy, childbirth and the puerperium (O00-O99)	196	1.0	0.9
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	193	0.9	0.9
11	Cerebrovascular diseases (I60-I69)	173	0.8	0.8
12	Chronic liver disease and cirrhosis (K70,K73-K74)	145	0.7	0.7
13	Septicemia (A40-A41)	144	0.7	0.7
14	Chronic lower respiratory diseases (J40-J47)	103	0.5	0.5
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	80	0.4	0.4
...	All other causes (Residual)	2,909	14.1	13.7
All races, both sexes, 30-34 years				
...	All causes	23,176	100.0	113.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,557	32.6	36.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,009	13.0	14.7
3	Malignant neoplasms (C00-C97)	2,180	9.4	10.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,119	9.1	10.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,812	7.8	8.8
6	Diabetes mellitus (E10-E14)	449	1.9	2.2
7	Human immunodeficiency virus (HIV) disease (B20-B24)	437	1.9	2.1
8	Chronic liver disease and cirrhosis (K70,K73-K74)	360	1.6	1.8
9	Cerebrovascular diseases (I60-I69)	357	1.5	1.7
10	Influenza and pneumonia (J09-J18)	278	1.2	1.4
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	217	0.9	1.1
12	Septicemia (A40-A41)	207	0.9	1.0
13	Pregnancy, childbirth and the puerperium (O00-O99)	190	0.8	0.9
14	Chronic lower respiratory diseases (J40-J47)	165	0.7	0.8
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	132	0.6	0.6
...	All other causes (Residual)	3,707	16.0	18.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 35-39 years				
...	All causes	27,478	100.0	140.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,031	25.6	35.9
2	Malignant neoplasms (C00-C97)	3,920	14.3	20.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,608	13.1	18.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,997	10.9	15.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,351	4.9	6.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	753	2.7	3.8
7	Diabetes mellitus (E10-E14)	631	2.3	3.2
8	Human immunodeficiency virus (HIV) disease (B20-B24)	618	2.2	3.2
9	Cerebrovascular diseases (I60-I69)	568	2.1	2.9
10	Influenza and pneumonia (J09-J18)	323	1.2	1.6
11	Septicemia (A40-A41)	287	1.0	1.5
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	241	0.9	1.2
13	Chronic lower respiratory diseases (J40-J47)	210	0.8	1.1
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	199	0.7	1.0
15	Pregnancy, childbirth and the puerperium (O00-O99)	161	0.6	0.8
...	All other causes (Residual)	4,580	16.7	23.4
All races, both sexes, 40-44 years				
...	All causes	42,415	100.0	201.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,199	19.3	39.0
2	Malignant neoplasms (C00-C97)	7,797	18.4	37.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,027	16.6	33.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,602	8.5	17.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,696	4.0	8.1
6	Diabetes mellitus (E10-E14)	1,211	2.9	5.8
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,168	2.8	5.6
8	Cerebrovascular diseases (I60-I69)	1,150	2.7	5.5
9	Human immunodeficiency virus (HIV) disease (B20-B24)	1,001	2.4	4.8
10	Influenza and pneumonia (J09-J18)	536	1.3	2.5
11	Chronic lower respiratory diseases (J40-J47)	524	1.2	2.5
12	Septicemia (A40-A41)	507	1.2	2.4
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	427	1.0	2.0
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	271	0.6	1.3
15	Viral hepatitis (B15-B19)	236	0.6	1.1
...	All other causes (Residual)	7,063	16.7	33.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 45-49 years				
...	All causes	71,674	100.0	323.5
1	Malignant neoplasms (C00-C97)	16,635	23.2	75.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,562	18.9	61.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,107	14.1	45.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,282	6.0	19.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	3,546	4.9	16.0
6	Diabetes mellitus (E10-E14)	2,288	3.2	10.3
7	Cerebrovascular diseases (I60-I69)	2,232	3.1	10.1
8	Human immunodeficiency virus (HIV) disease (B20-B24)	1,384	1.9	6.2
9	Chronic lower respiratory diseases (J40-J47)	1,364	1.9	6.2
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,132	1.6	5.1
11	Septicemia (A40-A41)	897	1.3	4.0
12	Influenza and pneumonia (J09-J18)	850	1.2	3.8
13	Viral hepatitis (B15-B19)	765	1.1	3.5
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	727	1.0	3.3
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	543	0.8	2.5
...	All other causes (Residual)	11,360	15.8	51.3
All races, both sexes, 50-54 years				
...	All causes	111,573	100.0	494.6
1	Malignant neoplasms (C00-C97)	32,262	28.9	143.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	22,538	20.2	99.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,642	9.5	47.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	5,318	4.8	23.6
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,576	4.1	20.3
6	Diabetes mellitus (E10-E14)	3,724	3.3	16.5
7	Cerebrovascular diseases (I60-I69)	3,473	3.1	15.4
8	Chronic lower respiratory diseases (J40-J47)	3,270	2.9	14.5
9	Septicemia (A40-A41)	1,564	1.4	6.9
10	Viral hepatitis (B15-B19)	1,552	1.4	6.9
11	Human immunodeficiency virus (HIV) disease (B20-B24)	1,397	1.3	6.2
12	Influenza and pneumonia (J09-J18)	1,374	1.2	6.1
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,257	1.1	5.6
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	920	0.8	4.1
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	832	0.7	3.7
...	All other causes (Residual)	16,874	15.1	74.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 55-59 years				
...	All causes	144,272	100.0	712.3
1	Malignant neoplasms (C00-C97)	47,923	33.2	236.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	30,590	21.2	151.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,556	5.9	42.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	5,861	4.1	28.9
5	Chronic lower respiratory diseases (J40-J47)	5,663	3.9	28.0
6	Diabetes mellitus (E10-E14)	5,439	3.8	26.9
7	Cerebrovascular diseases (I60-I69)	4,696	3.3	23.2
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,808	2.6	18.8
9	Septicemia (A40-A41)	2,141	1.5	10.6
10	Viral hepatitis (B15-B19)	2,088	1.4	10.3
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,897	1.3	9.4
12	Influenza and pneumonia (J09-J18)	1,820	1.3	9.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,179	0.8	5.8
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1,110	0.8	5.5
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	665	0.5	3.3
...	All other causes (Residual)	20,836	14.4	102.9
All races, both sexes, 60-64 years				
...	All causes	179,043	100.0	1,005.5
1	Malignant neoplasms (C00-C97)	64,649	36.1	363.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	39,152	21.9	219.9
3	Chronic lower respiratory diseases (J40-J47)	9,381	5.2	52.7
4	Diabetes mellitus (E10-E14)	7,249	4.0	40.7
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,602	3.7	37.1
6	Cerebrovascular diseases (I60-I69)	6,509	3.6	36.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	4,888	2.7	27.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,857	1.6	16.0
9	Septicemia (A40-A41)	2,812	1.6	15.8
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,713	1.5	15.2
11	Influenza and pneumonia (J09-J18)	2,365	1.3	13.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,500	0.8	8.4
13	Viral hepatitis (B15-B19)	1,427	0.8	8.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	886	0.5	5.0
15	Aortic aneurysm and dissection (I71)	740	0.4	4.2
...	All other causes (Residual)	25,313	14.1	142.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 65-69 years				
...	All causes	193,475	100.0	1,502.9
1	Malignant neoplasms (C00-C97)	70,579	36.5	548.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	41,719	21.6	324.1
3	Chronic lower respiratory diseases (J40-J47)	14,033	7.3	109.0
4	Diabetes mellitus (E10-E14)	7,790	4.0	60.5
5	Cerebrovascular diseases (I60-I69)	7,478	3.9	58.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,020	2.6	39.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,329	1.7	25.9
8	Chronic liver disease and cirrhosis (K70,K73-K74)	3,277	1.7	25.5
9	Septicemia (A40-A41)	3,036	1.6	23.6
10	Influenza and pneumonia (J09-J18)	2,773	1.4	21.5
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,806	0.9	14.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,692	0.9	13.1
13	Alzheimer's disease (G30)	1,317	0.7	10.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,124	0.6	8.7
15	Aortic aneurysm and dissection (I71)	930	0.5	7.2
...	All other causes (Residual)	27,572	14.3	214.2
All races, both sexes, 70-74 years				
...	All causes	221,577	100.0	2,306.2
1	Malignant neoplasms (C00-C97)	75,014	33.9	780.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	47,989	21.7	499.5
3	Chronic lower respiratory diseases (J40-J47)	18,400	8.3	191.5
4	Cerebrovascular diseases (I60-I69)	10,106	4.6	105.2
5	Diabetes mellitus (E10-E14)	8,399	3.8	87.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,994	2.3	52.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,353	2.0	45.3
8	Influenza and pneumonia (J09-J18)	3,715	1.7	38.7
9	Septicemia (A40-A41)	3,600	1.6	37.5
10	Alzheimer's disease (G30)	3,008	1.4	31.3
11	Chronic liver disease and cirrhosis (K70,K73-K74)	2,628	1.2	27.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,044	0.9	21.3
13	Parkinson's disease (G20-G21)	1,962	0.9	20.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,461	0.7	15.2
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,373	0.6	14.3
...	All other causes (Residual)	32,531	14.7	338.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 75-79 years				
...	All causes	272,276	100.0	3,685.0
1	Malignant neoplasms (C00-C97)	77,508	28.5	1,049.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	61,737	22.7	835.6
3	Chronic lower respiratory diseases (J40-J47)	22,669	8.3	306.8
4	Cerebrovascular diseases (I60-I69)	15,182	5.6	205.5
5	Diabetes mellitus (E10-E14)	9,154	3.4	123.9
6	Alzheimer's disease (G30)	7,969	2.9	107.9
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,867	2.2	79.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,646	2.1	76.4
9	Influenza and pneumonia (J09-J18)	5,296	1.9	71.7
10	Septicemia (A40-A41)	4,450	1.6	60.2
11	Parkinson's disease (G20-G21)	3,892	1.4	52.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,689	1.0	36.4
13	Chronic liver disease and cirrhosis (K70,K73-K74)	2,165	0.8	29.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,988	0.7	26.9
15	Pneumonitis due to solids and liquids (J69)	1,945	0.7	26.3
...	All other causes (Residual)	44,119	16.2	597.1
All races, both sexes, 80-84 years				
...	All causes	353,949	100.0	6,116.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	87,763	24.8	1,516.7
2	Malignant neoplasms (C00-C97)	77,840	22.0	1,345.2
3	Chronic lower respiratory diseases (J40-J47)	26,725	7.6	461.8
4	Cerebrovascular diseases (I60-I69)	22,414	6.3	387.3
5	Alzheimer's disease (G30)	16,255	4.6	280.9
6	Diabetes mellitus (E10-E14)	10,451	3.0	180.6
7	Influenza and pneumonia (J09-J18)	8,402	2.4	145.2
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,226	2.3	142.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,712	2.2	133.3
10	Parkinson's disease (G20-G21)	6,127	1.7	105.9
11	Septicemia (A40-A41)	5,348	1.5	92.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,080	1.2	70.5
13	Pneumonitis due to solids and liquids (J69)	3,194	0.9	55.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,650	0.7	45.8
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,694	0.5	29.3
...	All other causes (Residual)	65,068	18.4	1,124.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 85-89 years				
...	All causes	383,912	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	106,952	27.9	---
2	Malignant neoplasms (C00-C97)	59,810	15.6	---
3	Cerebrovascular diseases (I60-I69)	26,211	6.8	---
4	Alzheimer's disease (G30)	24,200	6.3	---
5	Chronic lower respiratory diseases (J40-J47)	23,321	6.1	---
6	Influenza and pneumonia (J09-J18)	10,793	2.8	---
7	Diabetes mellitus (E10-E14)	9,471	2.5	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,361	2.4	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,510	2.2	---
10	Parkinson's disease (G20-G21)	5,927	1.5	---
11	Septicemia (A40-A41)	5,584	1.5	---
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,244	1.4	---
13	Pneumonitis due to solids and liquids (J69)	4,154	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,768	0.7	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1,864	0.5	---
...	All other causes (Residual)	79,742	20.8	---
All races, both sexes, 90-94 years				
...	All causes	274,213	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	84,963	31.0	---
2	Malignant neoplasms (C00-C97)	28,231	10.3	---
3	Alzheimer's disease (G30)	20,597	7.5	---
4	Cerebrovascular diseases (I60-I69)	19,232	7.0	---
5	Chronic lower respiratory diseases (J40-J47)	12,299	4.5	---
6	Influenza and pneumonia (J09-J18)	9,083	3.3	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,644	2.4	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,809	2.1	---
9	Diabetes mellitus (E10-E14)	5,289	1.9	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,626	1.7	---
11	Pneumonitis due to solids and liquids (J69)	3,394	1.2	---
12	Septicemia (A40-A41)	3,375	1.2	---
13	Parkinson's disease (G20-G21)	3,008	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,583	0.6	---
15	Atherosclerosis (I70)	1,446	0.5	---
...	All other causes (Residual)	64,634	23.6	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 95-99 years				
...	All causes	109,610	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	36,080	32.9	---
2	Alzheimer's disease (G30)	8,981	8.2	---
3	Cerebrovascular diseases (I60-I69)	7,329	6.7	---
4	Malignant neoplasms (C00-C97)	7,230	6.6	---
5	Influenza and pneumonia (J09-J18)	4,251	3.9	---
6	Chronic lower respiratory diseases (J40-J47)	3,859	3.5	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,628	2.4	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,250	2.1	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,070	1.9	---
10	Diabetes mellitus (E10-E14)	1,591	1.5	---
11	Pneumonitis due to solids and liquids (J69)	1,293	1.2	---
12	Septicemia (A40-A41)	1,143	1.0	---
13	Atherosclerosis (I70)	758	0.7	---
14	Parkinson's disease (G20-G21)	650	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	518	0.5	---
...	All other causes (Residual)	28,979	26.4	---
All races, both sexes, 100 years and over				
...	All causes	22,810	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,894	34.6	---
2	Alzheimer's disease (G30)	1,705	7.5	---
3	Cerebrovascular diseases (I60-I69)	1,371	6.0	---
4	Influenza and pneumonia (J09-J18)	1,073	4.7	---
5	Malignant neoplasms (C00-C97)	894	3.9	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	655	2.9	---
7	Chronic lower respiratory diseases (J40-J47)	563	2.5	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	518	2.3	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	367	1.6	---
10	Diabetes mellitus (E10-E14)	257	1.1	---
11	Pneumonitis due to solids and liquids (J69)	255	1.1	---
12	Atherosclerosis (I70)	227	1.0	---
13	Septicemia (A40-A41)	210	0.9	---
14	Anemias (D50-D64)	100	0.4	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	74	0.3	---
...	All other causes (Residual)	6,647	29.1	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, all ages				
...	All causes	1,254,978	100.0	818.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	308,398	24.6	201.2
2	Malignant neoplasms (C00-C97)	302,231	24.1	197.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	79,257	6.3	51.7
4	Chronic lower respiratory diseases (J40-J47)	67,521	5.4	44.0
5	Cerebrovascular diseases (I60-I69)	52,335	4.2	34.1
6	Diabetes mellitus (E10-E14)	38,324	3.1	25.0
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	31,003	2.5	20.2
8	Alzheimer's disease (G30)	25,677	2.0	16.8
9	Influenza and pneumonia (J09-J18)	25,401	2.0	16.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22,649	1.8	14.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	21,621	1.7	14.1
12	Septicemia (A40-A41)	16,484	1.3	10.8
13	Parkinson's disease (G20-G21)	13,546	1.1	8.8
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	12,745	1.0	8.3
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	11,303	0.9	7.4
...	All other causes (Residual)	226,483	18.0	147.7
All races, male, 1-4 years				
...	All causes	2,405	100.0	29.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	839	34.9	10.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	266	11.1	3.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	239	9.9	2.9
4	Malignant neoplasms (C00-C97)	183	7.6	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	84	3.5	1.0
6	Influenza and pneumonia (J09-J18)	55	2.3	0.7
7	Chronic lower respiratory diseases (J40-J47)	37	1.5	0.4
8	Cerebrovascular diseases (I60-I69)	29	1.2	0.4
9	Septicemia (A40-A41)	26	1.1	0.3
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	26	1.1	0.3
11	Certain conditions originating in the perinatal period (P00-P96)	20	0.8	0.2
12	Complications of medical and surgical care (Y40-Y84,Y88)	13	0.5	*
13	Anemias (D50-D64)	9	0.4	*
14	Acute bronchitis and bronchiolitis (J20-J21)	8	0.3	*
15	Meningitis (G00,G03)	7	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.3	*
...	All other causes (Residual)	557	23.2	6.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 5-9 years				
...	All causes	1,380	100.0	13.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	461	33.4	4.4
2	Malignant neoplasms (C00-C97)	244	17.7	2.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	109	7.9	1.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	75	5.4	0.7
5	Chronic lower respiratory diseases (J40-J47)	44	3.2	0.4
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	38	2.8	0.4
7	Influenza and pneumonia (J09-J18)	32	2.3	0.3
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	1.4	*
8	Cerebrovascular diseases (I60-I69)	19	1.4	*
10	Anemias (D50-D64)	15	1.1	*
11	Septicemia (A40-A41)	14	1.0	*
12	Certain conditions originating in the perinatal period (P00-P96)	10	0.7	*
13	Pneumonitis due to solids and liquids (J69)	6	0.4	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.4	*
15	Meningitis (G00,G03)	4	0.3	*
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4	0.3	*
...	All other causes (Residual)	280	20.3	2.7
All races, male, 10-14 years				
...	All causes	1,799	100.0	17.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	572	31.8	5.4
2	Malignant neoplasms (C00-C97)	239	13.3	2.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	197	11.0	1.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	103	5.7	1.0
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	89	4.9	0.8
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	57	3.2	0.5
7	Chronic lower respiratory diseases (J40-J47)	43	2.4	0.4
8	Influenza and pneumonia (J09-J18)	31	1.7	0.3
9	Cerebrovascular diseases (I60-I69)	27	1.5	0.3
10	Septicemia (A40-A41)	20	1.1	0.2
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	1.0	*
12	Anemias (D50-D64)	12	0.7	*
13	Diabetes mellitus (E10-E14)	10	0.6	*
14	Pneumonitis due to solids and liquids (J69)	8	0.4	*
15	Certain conditions originating in the perinatal period (P00-P96)	6	0.3	*
...	All other causes (Residual)	367	20.4	3.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 15-19 years				
...	All causes	7,618	100.0	68.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,044	40.0	27.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,445	19.0	13.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,432	18.8	12.9
4	Malignant neoplasms (C00-C97)	389	5.1	3.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	224	2.9	2.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	130	1.7	1.2
7	Chronic lower respiratory diseases (J40-J47)	48	0.6	0.4
8	Influenza and pneumonia (J09-J18)	40	0.5	0.4
9	Cerebrovascular diseases (I60-I69)	39	0.5	0.4
10	Diabetes mellitus (E10-E14)	29	0.4	0.3
11	Legal intervention (Y35,Y89.0)	28	0.4	0.3
12	Septicemia (A40-A41)	26	0.3	0.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	0.2	*
14	Anemias (D50-D64)	17	0.2	*
15	Aortic aneurysm and dissection (I71)	11	0.1	*
...	All other causes (Residual)	698	9.2	6.3
All races, male, 20-24 years				
...	All causes	14,306	100.0	126.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,128	42.8	54.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,527	17.7	22.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,496	17.4	22.1
4	Malignant neoplasms (C00-C97)	577	4.0	5.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	460	3.2	4.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	145	1.0	1.3
7	Influenza and pneumonia (J09-J18)	83	0.6	0.7
8	Human immunodeficiency virus (HIV) disease (B20-B24)	76	0.5	0.7
9	Diabetes mellitus (E10-E14)	72	0.5	0.6
9	Legal intervention (Y35,Y89.0)	72	0.5	0.6
11	Chronic lower respiratory diseases (J40-J47)	69	0.5	0.6
12	Cerebrovascular diseases (I60-I69)	65	0.5	0.6
13	Septicemia (A40-A41)	46	0.3	0.4
14	Anemias (D50-D64)	43	0.3	0.4
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	32	0.2	0.3
...	All other causes (Residual)	1,415	9.9	12.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 25-29 years				
...	All causes	14,692	100.0	136.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,005	40.9	55.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,506	17.1	23.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,990	13.5	18.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	803	5.5	7.5
5	Malignant neoplasms (C00-C97)	708	4.8	6.6
6	Human immunodeficiency virus (HIV) disease (B20-B24)	151	1.0	1.4
7	Influenza and pneumonia (J09-J18)	127	0.9	1.2
8	Diabetes mellitus (E10-E14)	123	0.8	1.1
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	114	0.8	1.1
10	Cerebrovascular diseases (I60-I69)	92	0.6	0.9
11	Septicemia (A40-A41)	77	0.5	0.7
12	Chronic liver disease and cirrhosis (K70,K73-K74)	76	0.5	0.7
13	Chronic lower respiratory diseases (J40-J47)	68	0.5	0.6
14	Legal intervention (Y35,Y89.0)	66	0.4	0.6
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	45	0.3	0.4
...	All other causes (Residual)	1,741	11.8	16.2
All races, male, 30-34 years				
...	All causes	15,393	100.0	149.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,421	35.2	52.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,356	15.3	22.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,509	9.8	14.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,407	9.1	13.7
5	Malignant neoplasms (C00-C97)	1,005	6.5	9.8
6	Human immunodeficiency virus (HIV) disease (B20-B24)	277	1.8	2.7
7	Diabetes mellitus (E10-E14)	265	1.7	2.6
8	Chronic liver disease and cirrhosis (K70,K73-K74)	230	1.5	2.2
9	Cerebrovascular diseases (I60-I69)	192	1.2	1.9
10	Influenza and pneumonia (J09-J18)	163	1.1	1.6
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	115	0.7	1.1
12	Septicemia (A40-A41)	111	0.7	1.1
13	Chronic lower respiratory diseases (J40-J47)	89	0.6	0.9
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	74	0.5	0.7
15	Legal intervention (Y35,Y89.0)	66	0.4	0.6
...	All other causes (Residual)	2,113	13.7	20.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 35-39 years				
...	All causes	17,353	100.0	177.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,905	28.3	50.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,473	14.3	25.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,319	13.4	23.8
4	Malignant neoplasms (C00-C97)	1,694	9.8	17.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,056	6.1	10.8
6	Chronic liver disease and cirrhosis (K70,K73-K74)	493	2.8	5.1
7	Diabetes mellitus (E10-E14)	382	2.2	3.9
8	Human immunodeficiency virus (HIV) disease (B20-B24)	375	2.2	3.8
9	Cerebrovascular diseases (I60-I69)	285	1.6	2.9
10	Influenza and pneumonia (J09-J18)	183	1.1	1.9
11	Septicemia (A40-A41)	153	0.9	1.6
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	144	0.8	1.5
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	114	0.7	1.2
14	Aortic aneurysm and dissection (I71)	98	0.6	1.0
15	Chronic lower respiratory diseases (J40-J47)	92	0.5	0.9
...	All other causes (Residual)	2,587	14.9	26.5
All races, male, 40-44 years				
...	All causes	25,862	100.0	247.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,554	21.5	53.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,920	19.0	47.0
3	Malignant neoplasms (C00-C97)	3,340	12.9	31.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,725	10.5	26.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,079	4.2	10.3
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	884	3.4	8.4
7	Diabetes mellitus (E10-E14)	738	2.9	7.1
8	Human immunodeficiency virus (HIV) disease (B20-B24)	685	2.6	6.5
9	Cerebrovascular diseases (I60-I69)	621	2.4	5.9
10	Influenza and pneumonia (J09-J18)	273	1.1	2.6
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	252	1.0	2.4
12	Chronic lower respiratory diseases (J40-J47)	248	1.0	2.4
13	Septicemia (A40-A41)	246	1.0	2.4
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	173	0.7	1.7
15	Viral hepatitis (B15-B19)	156	0.6	1.5
...	All other causes (Residual)	3,968	15.3	37.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 45-49 years				
...	All causes	43,514	100.0	397.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,612	22.1	87.8
2	Malignant neoplasms (C00-C97)	7,802	17.9	71.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,801	15.6	62.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,205	7.4	29.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,276	5.2	20.8
6	Diabetes mellitus (E10-E14)	1,413	3.2	12.9
7	Cerebrovascular diseases (I60-I69)	1,222	2.8	11.2
8	Human immunodeficiency virus (HIV) disease (B20-B24)	1,016	2.3	9.3
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	828	1.9	7.6
10	Chronic lower respiratory diseases (J40-J47)	611	1.4	5.6
11	Viral hepatitis (B15-B19)	548	1.3	5.0
12	Influenza and pneumonia (J09-J18)	485	1.1	4.4
13	Septicemia (A40-A41)	436	1.0	4.0
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	428	1.0	3.9
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	354	0.8	3.2
...	All other causes (Residual)	6,477	14.9	59.1
All races, male, 50-54 years				
...	All causes	68,195	100.0	616.2
1	Malignant neoplasms (C00-C97)	16,570	24.3	149.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	16,079	23.6	145.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,116	10.4	64.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	3,592	5.3	32.5
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,432	5.0	31.0
6	Diabetes mellitus (E10-E14)	2,330	3.4	21.1
7	Cerebrovascular diseases (I60-I69)	1,979	2.9	17.9
8	Chronic lower respiratory diseases (J40-J47)	1,562	2.3	14.1
9	Viral hepatitis (B15-B19)	1,089	1.6	9.8
10	Human immunodeficiency virus (HIV) disease (B20-B24)	1,036	1.5	9.4
11	Septicemia (A40-A41)	860	1.3	7.8
12	Influenza and pneumonia (J09-J18)	803	1.2	7.3
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	725	1.1	6.6
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	668	1.0	6.0
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	525	0.8	4.7
...	All other causes (Residual)	9,829	14.4	88.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 55-59 years				
...	All causes	89,129	100.0	908.0
1	Malignant neoplasms (C00-C97)	26,460	29.7	269.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	21,636	24.3	220.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,901	6.6	60.1
4	Chronic liver disease and cirrhosis (K70,K73-K74)	4,176	4.7	42.5
5	Diabetes mellitus (E10-E14)	3,314	3.7	33.8
6	Chronic lower respiratory diseases (J40-J47)	2,972	3.3	30.3
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,925	3.3	29.8
8	Cerebrovascular diseases (I60-I69)	2,735	3.1	27.9
9	Viral hepatitis (B15-B19)	1,554	1.7	15.8
10	Septicemia (A40-A41)	1,158	1.3	11.8
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,104	1.2	11.2
12	Influenza and pneumonia (J09-J18)	1,072	1.2	10.9
13	Human immunodeficiency virus (HIV) disease (B20-B24)	830	0.9	8.5
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	748	0.8	7.6
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	489	0.5	5.0
...	All other causes (Residual)	12,055	13.5	122.8
All races, male, 60-64 years				
...	All causes	107,512	100.0	1,258.6
1	Malignant neoplasms (C00-C97)	36,039	33.5	421.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	26,752	24.9	313.2
3	Chronic lower respiratory diseases (J40-J47)	4,918	4.6	57.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,540	4.2	53.1
5	Diabetes mellitus (E10-E14)	4,310	4.0	50.5
6	Cerebrovascular diseases (I60-I69)	3,670	3.4	43.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	3,413	3.2	40.0
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,093	1.9	24.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,597	1.5	18.7
10	Septicemia (A40-A41)	1,490	1.4	17.4
11	Influenza and pneumonia (J09-J18)	1,358	1.3	15.9
12	Viral hepatitis (B15-B19)	1,017	0.9	11.9
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	909	0.8	10.6
14	Aortic aneurysm and dissection (I71)	535	0.5	6.3
15	Human immunodeficiency virus (HIV) disease (B20-B24)	530	0.5	6.2
...	All other causes (Residual)	14,341	13.3	167.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 65-69 years				
...	All causes	111,373	100.0	1,834.2
1	Malignant neoplasms (C00-C97)	38,926	35.0	641.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	27,241	24.5	448.6
3	Chronic lower respiratory diseases (J40-J47)	7,178	6.4	118.2
4	Diabetes mellitus (E10-E14)	4,513	4.1	74.3
5	Cerebrovascular diseases (I60-I69)	4,039	3.6	66.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,237	2.9	53.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,140	1.9	35.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,798	1.6	29.6
9	Septicemia (A40-A41)	1,551	1.4	25.5
10	Influenza and pneumonia (J09-J18)	1,541	1.4	25.4
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,422	1.3	23.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	961	0.9	15.8
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	665	0.6	11.0
14	Aortic aneurysm and dissection (I71)	644	0.6	10.6
15	Alzheimer's disease (G30)	596	0.5	9.8
...	All other causes (Residual)	14,921	13.4	245.7
All races, male, 70-74 years				
...	All causes	122,903	100.0	2,790.5
1	Malignant neoplasms (C00-C97)	41,297	33.6	937.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	29,368	23.9	666.8
3	Chronic lower respiratory diseases (J40-J47)	9,319	7.6	211.6
4	Cerebrovascular diseases (I60-I69)	5,189	4.2	117.8
5	Diabetes mellitus (E10-E14)	4,668	3.8	106.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,058	2.5	69.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,285	1.9	51.9
8	Influenza and pneumonia (J09-J18)	2,044	1.7	46.4
9	Septicemia (A40-A41)	1,828	1.5	41.5
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,536	1.2	34.9
11	Parkinson's disease (G20-G21)	1,329	1.1	30.2
12	Alzheimer's disease (G30)	1,270	1.0	28.8
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,132	0.9	25.7
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	995	0.8	22.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	837	0.7	19.0
...	All other causes (Residual)	16,748	13.6	380.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 75-79 years				
...	All causes	142,371	100.0	4,404.0
1	Malignant neoplasms (C00-C97)	41,856	29.4	1,294.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	34,871	24.5	1,078.7
3	Chronic lower respiratory diseases (J40-J47)	11,078	7.8	342.7
4	Cerebrovascular diseases (I60-I69)	7,027	4.9	217.4
5	Diabetes mellitus (E10-E14)	4,709	3.3	145.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,303	2.3	102.2
7	Alzheimer's disease (G30)	3,223	2.3	99.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,916	2.0	90.2
9	Influenza and pneumonia (J09-J18)	2,798	2.0	86.6
10	Parkinson's disease (G20-G21)	2,521	1.8	78.0
11	Septicemia (A40-A41)	2,132	1.5	66.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,249	0.9	38.6
13	Chronic liver disease and cirrhosis (K70,K73-K74)	1,175	0.8	36.3
14	Pneumonitis due to solids and liquids (J69)	1,114	0.8	34.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,102	0.8	34.1
...	All other causes (Residual)	21,297	15.0	658.8
All races, male, 80-84 years				
...	All causes	170,322	100.0	7,277.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	44,633	26.2	1,907.2
2	Malignant neoplasms (C00-C97)	40,543	23.8	1,732.4
3	Chronic lower respiratory diseases (J40-J47)	12,658	7.4	540.9
4	Cerebrovascular diseases (I60-I69)	9,120	5.4	389.7
5	Alzheimer's disease (G30)	5,905	3.5	252.3
6	Diabetes mellitus (E10-E14)	4,967	2.9	212.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,266	2.5	182.3
8	Influenza and pneumonia (J09-J18)	4,136	2.4	176.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,916	2.3	167.3
10	Parkinson's disease (G20-G21)	3,642	2.1	155.6
11	Septicemia (A40-A41)	2,479	1.5	105.9
12	Pneumonitis due to solids and liquids (J69)	1,795	1.1	76.7
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,670	1.0	71.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,423	0.8	60.8
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	922	0.5	39.4
...	All other causes (Residual)	28,247	16.6	1,207.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 85-89 years				
...	All causes	159,865	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	46,378	29.0	---
2	Malignant neoplasms (C00-C97)	29,058	18.2	---
3	Chronic lower respiratory diseases (J40-J47)	10,313	6.5	---
4	Cerebrovascular diseases (I60-I69)	9,003	5.6	---
5	Alzheimer's disease (G30)	7,390	4.6	---
6	Influenza and pneumonia (J09-J18)	4,915	3.1	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,160	2.6	---
8	Diabetes mellitus (E10-E14)	4,028	2.5	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,969	2.5	---
10	Parkinson's disease (G20-G21)	3,324	2.1	---
11	Pneumonitis due to solids and liquids (J69)	2,211	1.4	---
12	Septicemia (A40-A41)	2,187	1.4	---
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,751	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,391	0.9	---
15	Aortic aneurysm and dissection (I71)	730	0.5	---
...	All other causes (Residual)	29,057	18.2	---
All races, male, 90-94 years				
...	All causes	94,374	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	30,448	32.3	---
2	Malignant neoplasms (C00-C97)	12,433	13.2	---
3	Cerebrovascular diseases (I60-I69)	5,218	5.5	---
4	Alzheimer's disease (G30)	5,204	5.5	---
5	Chronic lower respiratory diseases (J40-J47)	4,788	5.1	---
6	Influenza and pneumonia (J09-J18)	3,531	3.7	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,505	2.7	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,436	2.6	---
9	Diabetes mellitus (E10-E14)	1,955	2.1	---
10	Pneumonitis due to solids and liquids (J69)	1,591	1.7	---
11	Parkinson's disease (G20-G21)	1,501	1.6	---
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,270	1.3	---
13	Septicemia (A40-A41)	1,191	1.3	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	635	0.7	---
15	Atherosclerosis (I70)	447	0.5	---
...	All other causes (Residual)	19,221	20.4	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 95-99 years				
...	All causes	27,447	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,407	34.3	---
2	Malignant neoplasms (C00-C97)	2,590	9.4	---
3	Cerebrovascular diseases (I60-I69)	1,510	5.5	---
4	Alzheimer's disease (G30)	1,499	5.5	---
5	Influenza and pneumonia (J09-J18)	1,370	5.0	---
6	Chronic lower respiratory diseases (J40-J47)	1,219	4.4	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	746	2.7	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	706	2.6	---
9	Pneumonitis due to solids and liquids (J69)	504	1.8	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	457	1.7	---
11	Diabetes mellitus (E10-E14)	441	1.6	---
12	Septicemia (A40-A41)	307	1.1	---
13	Parkinson's disease (G20-G21)	259	0.9	---
14	Atherosclerosis (I70)	177	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	147	0.5	---
...	All other causes (Residual)	6,108	22.3	---
All races, male, 100 years and over				
...	All causes	3,739	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,323	35.4	---
2	Influenza and pneumonia (J09-J18)	237	6.3	---
3	Malignant neoplasms (C00-C97)	236	6.3	---
4	Cerebrovascular diseases (I60-I69)	178	4.8	---
5	Alzheimer's disease (G30)	173	4.6	---
6	Chronic lower respiratory diseases (J40-J47)	144	3.9	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	98	2.6	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	92	2.5	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	88	2.4	---
10	Pneumonitis due to solids and liquids (J69)	77	2.1	---
11	Diabetes mellitus (E10-E14)	52	1.4	---
12	Septicemia (A40-A41)	48	1.3	---
13	Atherosclerosis (I70)	33	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	21	0.6	---
15	Parkinson's disease (G20-G21)	18	0.5	---
...	All other causes (Residual)	921	24.6	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, all ages				
...	All causes	1,260,480	100.0	796.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	288,179	22.9	182.0
2	Malignant neoplasms (C00-C97)	274,460	21.8	173.4
3	Cerebrovascular diseases (I60-I69)	76,597	6.1	48.4
4	Chronic lower respiratory diseases (J40-J47)	75,422	6.0	47.6
5	Alzheimer's disease (G30)	59,297	4.7	37.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	47,181	3.7	29.8
7	Diabetes mellitus (E10-E14)	35,507	2.8	22.4
8	Influenza and pneumonia (J09-J18)	28,425	2.3	18.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22,942	1.8	14.5
10	Septicemia (A40-A41)	19,264	1.5	12.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	16,550	1.3	10.5
12	Chronic liver disease and cirrhosis (K70,K73-K74)	12,021	1.0	7.6
13	Parkinson's disease (G20-G21)	9,565	0.8	6.0
14	Pneumonitis due to solids and liquids (J69)	8,519	0.7	5.4
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8,515	0.7	5.4
...	All other causes (Residual)	278,036	22.1	175.6
All races, female, 1-4 years				
...	All causes	1,841	100.0	23.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	538	29.2	6.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	227	12.3	2.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	173	9.4	2.2
4	Malignant neoplasms (C00-C97)	170	9.2	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	81	4.4	1.0
6	Influenza and pneumonia (J09-J18)	57	3.1	0.7
7	Septicemia (A40-A41)	35	1.9	0.4
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	1.0	*
9	Chronic lower respiratory diseases (J40-J47)	16	0.9	*
10	Certain conditions originating in the perinatal period (P00-P96)	14	0.8	*
11	Cerebrovascular diseases (I60-I69)	13	0.7	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	9	0.5	*
13	Acute bronchitis and bronchiolitis (J20-J21)	6	0.3	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.3	*
15	Anemias (D50-D64)	5	0.3	*
...	All other causes (Residual)	472	25.6	6.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 5-9 years				
...	All causes	1,071	100.0	10.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	300	28.0	3.0
2	Malignant neoplasms (C00-C97)	197	18.4	2.0
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	73	6.8	0.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	54	5.0	0.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	54	5.0	0.5
6	Influenza and pneumonia (J09-J18)	31	2.9	0.3
7	Septicemia (A40-A41)	24	2.2	0.2
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	21	2.0	0.2
8	Cerebrovascular diseases (I60-I69)	21	2.0	0.2
10	Chronic lower respiratory diseases (J40-J47)	20	1.9	0.2
11	Anemias (D50-D64)	13	1.2	*
12	Certain conditions originating in the perinatal period (P00-P96)	9	0.8	*
13	Diseases of appendix (K35-K38)	4	0.4	*
14	Meningitis (G00,G03)	3	0.3	*
14	Acute bronchitis and bronchiolitis (J20-J21)	3	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.3	*
...	All other causes (Residual)	238	22.2	2.4
All races, female, 10-14 years				
...	All causes	1,151	100.0	11.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	302	26.2	3.0
2	Malignant neoplasms (C00-C97)	180	15.6	1.8
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	87	7.6	0.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	85	7.4	0.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	54	4.7	0.5
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	51	4.4	0.5
7	Chronic lower respiratory diseases (J40-J47)	29	2.5	0.3
8	Influenza and pneumonia (J09-J18)	24	2.1	0.2
9	Cerebrovascular diseases (I60-I69)	20	1.7	0.2
10	Diabetes mellitus (E10-E14)	15	1.3	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	1.0	*
12	Septicemia (A40-A41)	11	1.0	*
13	Meningitis (G00,G03)	5	0.4	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.4	*
13	Certain conditions originating in the perinatal period (P00-P96)	5	0.4	*
...	All other causes (Residual)	266	23.1	2.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 15-19 years				
...	All causes	2,976	100.0	28.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,254	42.1	11.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	370	12.4	3.5
3	Malignant neoplasms (C00-C97)	250	8.4	2.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	235	7.9	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	94	3.2	0.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	77	2.6	0.7
7	Pregnancy, childbirth and the puerperium (O00-O99)	35	1.2	0.3
8	Cerebrovascular diseases (I60-I69)	30	1.0	0.3
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	26	0.9	0.2
9	Influenza and pneumonia (J09-J18)	26	0.9	0.2
11	Chronic lower respiratory diseases (J40-J47)	21	0.7	0.2
12	Diabetes mellitus (E10-E14)	20	0.7	0.2
13	Septicemia (A40-A41)	16	0.5	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	13	0.4	*
14	Anemias (D50-D64)	13	0.4	*
...	All other causes (Residual)	496	16.7	4.7
All races, female, 20-24 years				
...	All causes	4,767	100.0	44.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,904	39.9	17.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	493	10.3	4.5
3	Malignant neoplasms (C00-C97)	395	8.3	3.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	378	7.9	3.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	220	4.6	2.0
6	Pregnancy, childbirth and the puerperium (O00-O99)	137	2.9	1.3
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	80	1.7	0.7
8	Influenza and pneumonia (J09-J18)	71	1.5	0.7
9	Cerebrovascular diseases (I60-I69)	52	1.1	0.5
10	Septicemia (A40-A41)	46	1.0	0.4
10	Diabetes mellitus (E10-E14)	46	1.0	0.4
12	Human immunodeficiency virus (HIV) disease (B20-B24)	41	0.9	0.4
13	Chronic lower respiratory diseases (J40-J47)	32	0.7	0.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	28	0.6	0.3
15	Anemias (D50-D64)	24	0.5	0.2
...	All other causes (Residual)	820	17.2	7.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 25-29 years				
...	All causes	5,880	100.0	55.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,956	33.3	18.6
2	Malignant neoplasms (C00-C97)	611	10.4	5.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	585	9.9	5.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	383	6.5	3.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	379	6.4	3.6
6	Pregnancy, childbirth and the puerperium (O00-O99)	196	3.3	1.9
7	Diabetes mellitus (E10-E14)	114	1.9	1.1
8	Influenza and pneumonia (J09-J18)	110	1.9	1.0
9	Cerebrovascular diseases (I60-I69)	81	1.4	0.8
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	79	1.3	0.8
11	Human immunodeficiency virus (HIV) disease (B20-B24)	78	1.3	0.7
12	Chronic liver disease and cirrhosis (K70,K73-K74)	69	1.2	0.7
13	Septicemia (A40-A41)	67	1.1	0.6
14	Chronic lower respiratory diseases (J40-J47)	35	0.6	0.3
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	35	0.6	0.3
...	All other causes (Residual)	1,102	18.7	10.5
All races, female, 30-34 years				
...	All causes	7,783	100.0	76.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,136	27.4	20.9
2	Malignant neoplasms (C00-C97)	1,175	15.1	11.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	712	9.1	7.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	653	8.4	6.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	303	3.9	3.0
6	Pregnancy, childbirth and the puerperium (O00-O99)	190	2.4	1.9
7	Diabetes mellitus (E10-E14)	184	2.4	1.8
8	Cerebrovascular diseases (I60-I69)	165	2.1	1.6
9	Human immunodeficiency virus (HIV) disease (B20-B24)	160	2.1	1.6
10	Chronic liver disease and cirrhosis (K70,K73-K74)	130	1.7	1.3
11	Influenza and pneumonia (J09-J18)	115	1.5	1.1
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	102	1.3	1.0
13	Septicemia (A40-A41)	96	1.2	0.9
14	Chronic lower respiratory diseases (J40-J47)	76	1.0	0.7
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	58	0.7	0.6
...	All other causes (Residual)	1,528	19.6	14.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 35-39 years				
...	All causes	10,125	100.0	103.0
1	Malignant neoplasms (C00-C97)	2,226	22.0	22.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,126	21.0	21.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,135	11.2	11.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	678	6.7	6.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	295	2.9	3.0
6	Cerebrovascular diseases (I60-I69)	283	2.8	2.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	260	2.6	2.6
8	Diabetes mellitus (E10-E14)	249	2.5	2.5
9	Human immunodeficiency virus (HIV) disease (B20-B24)	243	2.4	2.5
10	Pregnancy, childbirth and the puerperium (O00-O99)	161	1.6	1.6
11	Influenza and pneumonia (J09-J18)	140	1.4	1.4
12	Septicemia (A40-A41)	134	1.3	1.4
13	Chronic lower respiratory diseases (J40-J47)	118	1.2	1.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	97	1.0	1.0
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	85	0.8	0.9
...	All other causes (Residual)	1,895	18.7	19.3
All races, female, 40-44 years				
...	All causes	16,553	100.0	156.6
1	Malignant neoplasms (C00-C97)	4,457	26.9	42.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,645	16.0	25.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,107	12.7	19.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	877	5.3	8.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	617	3.7	5.8
6	Cerebrovascular diseases (I60-I69)	529	3.2	5.0
7	Diabetes mellitus (E10-E14)	473	2.9	4.5
8	Human immunodeficiency virus (HIV) disease (B20-B24)	316	1.9	3.0
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	284	1.7	2.7
10	Chronic lower respiratory diseases (J40-J47)	276	1.7	2.6
11	Influenza and pneumonia (J09-J18)	263	1.6	2.5
12	Septicemia (A40-A41)	261	1.6	2.5
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	175	1.1	1.7
14	Pregnancy, childbirth and the puerperium (O00-O99)	99	0.6	0.9
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	98	0.6	0.9
...	All other causes (Residual)	3,076	18.6	29.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 45-49 years				
...	All causes	28,160	100.0	251.3
1	Malignant neoplasms (C00-C97)	8,833	31.4	78.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,950	14.0	35.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,306	11.7	29.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	1,270	4.5	11.3
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,077	3.8	9.6
6	Cerebrovascular diseases (I60-I69)	1,010	3.6	9.0
7	Diabetes mellitus (E10-E14)	875	3.1	7.8
8	Chronic lower respiratory diseases (J40-J47)	753	2.7	6.7
9	Septicemia (A40-A41)	461	1.6	4.1
10	Human immunodeficiency virus (HIV) disease (B20-B24)	368	1.3	3.3
11	Influenza and pneumonia (J09-J18)	365	1.3	3.3
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	304	1.1	2.7
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	299	1.1	2.7
14	Viral hepatitis (B15-B19)	217	0.8	1.9
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	189	0.7	1.7
...	All other causes (Residual)	4,883	17.3	43.6
All races, female, 50-54 years				
...	All causes	43,378	100.0	377.4
1	Malignant neoplasms (C00-C97)	15,692	36.2	136.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,459	14.9	56.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,526	8.1	30.7
4	Chronic liver disease and cirrhosis (K70,K73-K74)	1,726	4.0	15.0
5	Chronic lower respiratory diseases (J40-J47)	1,708	3.9	14.9
6	Cerebrovascular diseases (I60-I69)	1,494	3.4	13.0
7	Diabetes mellitus (E10-E14)	1,394	3.2	12.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,144	2.6	10.0
9	Septicemia (A40-A41)	704	1.6	6.1
10	Influenza and pneumonia (J09-J18)	571	1.3	5.0
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	532	1.2	4.6
12	Viral hepatitis (B15-B19)	463	1.1	4.0
13	Human immunodeficiency virus (HIV) disease (B20-B24)	361	0.8	3.1
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	307	0.7	2.7
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	252	0.6	2.2
...	All other causes (Residual)	7,045	16.2	61.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 55-59 years				
...	All causes	55,143	100.0	528.2
1	Malignant neoplasms (C00-C97)	21,463	38.9	205.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,954	16.2	85.8
3	Chronic lower respiratory diseases (J40-J47)	2,691	4.9	25.8
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,655	4.8	25.4
5	Diabetes mellitus (E10-E14)	2,125	3.9	20.4
6	Cerebrovascular diseases (I60-I69)	1,961	3.6	18.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,685	3.1	16.1
8	Septicemia (A40-A41)	983	1.8	9.4
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	883	1.6	8.5
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	793	1.4	7.6
11	Influenza and pneumonia (J09-J18)	748	1.4	7.2
12	Viral hepatitis (B15-B19)	534	1.0	5.1
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	431	0.8	4.1
14	Human immunodeficiency virus (HIV) disease (B20-B24)	280	0.5	2.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	267	0.5	2.6
...	All other causes (Residual)	8,690	15.8	83.2
All races, female, 60-64 years				
...	All causes	71,531	100.0	772.1
1	Malignant neoplasms (C00-C97)	28,610	40.0	308.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,400	17.3	133.8
3	Chronic lower respiratory diseases (J40-J47)	4,463	6.2	48.2
4	Diabetes mellitus (E10-E14)	2,939	4.1	31.7
5	Cerebrovascular diseases (I60-I69)	2,839	4.0	30.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,062	2.9	22.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,475	2.1	15.9
8	Septicemia (A40-A41)	1,322	1.8	14.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,260	1.8	13.6
10	Influenza and pneumonia (J09-J18)	1,007	1.4	10.9
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	620	0.9	6.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	591	0.8	6.4
13	Viral hepatitis (B15-B19)	410	0.6	4.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	383	0.5	4.1
15	Alzheimer's disease (G30)	325	0.5	3.5
...	All other causes (Residual)	10,825	15.1	116.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 65-69 years				
...	All causes	82,102	100.0	1,207.1
1	Malignant neoplasms (C00-C97)	31,653	38.6	465.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,478	17.6	212.9
3	Chronic lower respiratory diseases (J40-J47)	6,855	8.3	100.8
4	Cerebrovascular diseases (I60-I69)	3,439	4.2	50.6
5	Diabetes mellitus (E10-E14)	3,277	4.0	48.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,783	2.2	26.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,531	1.9	22.5
8	Septicemia (A40-A41)	1,485	1.8	21.8
9	Influenza and pneumonia (J09-J18)	1,232	1.5	18.1
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,137	1.4	16.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	731	0.9	10.7
12	Alzheimer's disease (G30)	721	0.9	10.6
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	459	0.6	6.7
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	384	0.5	5.6
15	Pneumonitis due to solids and liquids (J69)	343	0.4	5.0
...	All other causes (Residual)	12,594	15.3	185.2
All races, female, 70-74 years				
...	All causes	98,674	100.0	1,896.3
1	Malignant neoplasms (C00-C97)	33,717	34.2	648.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	18,621	18.9	357.9
3	Chronic lower respiratory diseases (J40-J47)	9,081	9.2	174.5
4	Cerebrovascular diseases (I60-I69)	4,917	5.0	94.5
5	Diabetes mellitus (E10-E14)	3,731	3.8	71.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,068	2.1	39.7
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,936	2.0	37.2
8	Septicemia (A40-A41)	1,772	1.8	34.1
9	Alzheimer's disease (G30)	1,738	1.8	33.4
10	Influenza and pneumonia (J09-J18)	1,671	1.7	32.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1,092	1.1	21.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,049	1.1	20.2
13	Parkinson's disease (G20-G21)	633	0.6	12.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	624	0.6	12.0
15	Pneumonitis due to solids and liquids (J69)	461	0.5	8.9
...	All other causes (Residual)	15,563	15.8	299.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 75-79 years				
...	All causes	129,905	100.0	3,125.8
1	Malignant neoplasms (C00-C97)	35,652	27.4	857.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	26,866	20.7	646.4
3	Chronic lower respiratory diseases (J40-J47)	11,591	8.9	278.9
4	Cerebrovascular diseases (I60-I69)	8,155	6.3	196.2
5	Alzheimer's disease (G30)	4,746	3.7	114.2
6	Diabetes mellitus (E10-E14)	4,445	3.4	107.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,730	2.1	65.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,564	2.0	61.7
9	Influenza and pneumonia (J09-J18)	2,498	1.9	60.1
10	Septicemia (A40-A41)	2,318	1.8	55.8
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,440	1.1	34.6
12	Parkinson's disease (G20-G21)	1,371	1.1	33.0
13	Chronic liver disease and cirrhosis (K70,K73-K74)	990	0.8	23.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	886	0.7	21.3
15	Pneumonitis due to solids and liquids (J69)	831	0.6	20.0
...	All other causes (Residual)	22,822	17.6	549.1
All races, female, 80-84 years				
...	All causes	183,627	100.0	5,328.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	43,130	23.5	1,251.5
2	Malignant neoplasms (C00-C97)	37,297	20.3	1,082.2
3	Chronic lower respiratory diseases (J40-J47)	14,067	7.7	408.2
4	Cerebrovascular diseases (I60-I69)	13,294	7.2	385.8
5	Alzheimer's disease (G30)	10,350	5.6	300.3
6	Diabetes mellitus (E10-E14)	5,484	3.0	159.1
7	Influenza and pneumonia (J09-J18)	4,266	2.3	123.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,960	2.2	114.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,796	2.1	110.1
10	Septicemia (A40-A41)	2,869	1.6	83.2
11	Parkinson's disease (G20-G21)	2,485	1.4	72.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,410	1.3	69.9
13	Pneumonitis due to solids and liquids (J69)	1,399	0.8	40.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,227	0.7	35.6
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	911	0.5	26.4
...	All other causes (Residual)	36,682	20.0	1,064.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 85-89 years				
...	All causes	224,047	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	60,574	27.0	---
2	Malignant neoplasms (C00-C97)	30,752	13.7	---
3	Cerebrovascular diseases (I60-I69)	17,208	7.7	---
4	Alzheimer's disease (G30)	16,810	7.5	---
5	Chronic lower respiratory diseases (J40-J47)	13,008	5.8	---
6	Influenza and pneumonia (J09-J18)	5,878	2.6	---
7	Diabetes mellitus (E10-E14)	5,443	2.4	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,201	2.3	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,541	2.0	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,493	1.6	---
11	Septicemia (A40-A41)	3,397	1.5	---
12	Parkinson's disease (G20-G21)	2,603	1.2	---
13	Pneumonitis due to solids and liquids (J69)	1,943	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,377	0.6	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1,230	0.5	---
...	All other causes (Residual)	50,589	22.6	---
All races, female, 90-94 years				
...	All causes	179,839	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	54,515	30.3	---
2	Malignant neoplasms (C00-C97)	15,798	8.8	---
3	Alzheimer's disease (G30)	15,393	8.6	---
4	Cerebrovascular diseases (I60-I69)	14,014	7.8	---
5	Chronic lower respiratory diseases (J40-J47)	7,511	4.2	---
6	Influenza and pneumonia (J09-J18)	5,552	3.1	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,208	2.3	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,356	1.9	---
9	Diabetes mellitus (E10-E14)	3,334	1.9	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,304	1.8	---
11	Septicemia (A40-A41)	2,184	1.2	---
12	Pneumonitis due to solids and liquids (J69)	1,803	1.0	---
13	Parkinson's disease (G20-G21)	1,507	0.8	---
14	Atherosclerosis (I70)	999	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	948	0.5	---
...	All other causes (Residual)	45,413	25.3	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 95-99 years				
...	All causes	82,163	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	26,673	32.5	---
2	Alzheimer's disease (G30)	7,482	9.1	---
3	Cerebrovascular diseases (I60-I69)	5,819	7.1	---
4	Malignant neoplasms (C00-C97)	4,640	5.6	---
5	Influenza and pneumonia (J09-J18)	2,881	3.5	---
6	Chronic lower respiratory diseases (J40-J47)	2,640	3.2	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,882	2.3	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,793	2.2	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,364	1.7	---
10	Diabetes mellitus (E10-E14)	1,150	1.4	---
11	Septicemia (A40-A41)	836	1.0	---
12	Pneumonitis due to solids and liquids (J69)	789	1.0	---
13	Atherosclerosis (I70)	581	0.7	---
14	Parkinson's disease (G20-G21)	391	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	371	0.5	---
...	All other causes (Residual)	22,871	27.8	---
All races, female, 100 years and over				
...	All causes	19,071	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,571	34.5	---
2	Alzheimer's disease (G30)	1,532	8.0	---
3	Cerebrovascular diseases (I60-I69)	1,193	6.3	---
4	Influenza and pneumonia (J09-J18)	836	4.4	---
5	Malignant neoplasms (C00-C97)	658	3.5	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	563	3.0	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	420	2.2	---
8	Chronic lower respiratory diseases (J40-J47)	419	2.2	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	279	1.5	---
10	Diabetes mellitus (E10-E14)	205	1.1	---
11	Atherosclerosis (I70)	194	1.0	---
12	Pneumonitis due to solids and liquids (J69)	178	0.9	---
13	Septicemia (A40-A41)	162	0.8	---
14	Anemias (D50-D64)	84	0.4	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	53	0.3	---
...	All other causes (Residual)	5,724	30.0	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, all ages				
...	All causes	2,156,077	100.0	872.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	513,701	23.8	207.9
2	Malignant neoplasms (C00-C97)	492,348	22.8	199.3
3	Chronic lower respiratory diseases (J40-J47)	131,495	6.1	53.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	109,751	5.1	44.4
5	Cerebrovascular diseases (I60-I69)	108,542	5.0	43.9
6	Alzheimer's disease (G30)	78,055	3.6	31.6
7	Diabetes mellitus (E10-E14)	57,974	2.7	23.5
8	Influenza and pneumonia (J09-J18)	46,658	2.2	18.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	36,268	1.7	14.7
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	35,775	1.7	14.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	29,507	1.4	11.9
12	Septicemia (A40-A41)	28,838	1.3	11.7
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	21,845	1.0	8.8
14	Parkinson's disease (G20-G21)	21,688	1.0	8.8
15	Pneumonitis due to solids and liquids (J69)	16,263	0.8	6.6
...	All other causes (Residual)	427,369	19.8	173.0
White, both sexes, 1-4 years				
...	All causes	2,980	100.0	24.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,027	34.5	8.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	360	12.1	3.0
3	Malignant neoplasms (C00-C97)	275	9.2	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	237	8.0	1.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	107	3.6	0.9
6	Influenza and pneumonia (J09-J18)	66	2.2	0.5
7	Septicemia (A40-A41)	45	1.5	0.4
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	33	1.1	0.3
9	Cerebrovascular diseases (I60-I69)	23	0.8	0.2
10	Chronic lower respiratory diseases (J40-J47)	22	0.7	0.2
11	Certain conditions originating in the perinatal period (P00-P96)	21	0.7	0.2
12	Complications of medical and surgical care (Y40-Y84,Y88)	13	0.4	*
13	Acute bronchitis and bronchiolitis (J20-J21)	12	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	0.3	*
15	Meningitis (G00,G03)	8	0.3	*
...	All other causes (Residual)	721	24.2	5.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 5-9 years				
...	All causes	1,775	100.0	11.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	568	32.0	3.7
2	Malignant neoplasms (C00-C97)	351	19.8	2.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	135	7.6	0.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	83	4.7	0.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	65	3.7	0.4
6	Influenza and pneumonia (J09-J18)	47	2.6	0.3
7	Septicemia (A40-A41)	30	1.7	0.2
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	28	1.6	0.2
9	Chronic lower respiratory diseases (J40-J47)	26	1.5	0.2
10	Cerebrovascular diseases (I60-I69)	23	1.3	0.1
11	Certain conditions originating in the perinatal period (P00-P96)	12	0.7	*
12	Anemias (D50-D64)	8	0.5	*
13	Meningitis (G00,G03)	5	0.3	*
14	Meningococcal infection (A39)	4	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.2	*
...	All other causes (Residual)	386	21.7	2.5
White, both sexes, 10-14 years				
...	All causes	2,152	100.0	13.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	674	31.3	4.3
2	Malignant neoplasms (C00-C97)	325	15.1	2.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	220	10.2	1.4
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	123	5.7	0.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	95	4.4	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	77	3.6	0.5
7	Cerebrovascular diseases (I60-I69)	36	1.7	0.2
7	Influenza and pneumonia (J09-J18)	36	1.7	0.2
9	Septicemia (A40-A41)	25	1.2	0.2
10	Chronic lower respiratory diseases (J40-J47)	23	1.1	0.1
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	22	1.0	0.1
12	Diabetes mellitus (E10-E14)	20	0.9	0.1
13	Pneumonitis due to solids and liquids (J69)	8	0.4	*
14	Anemias (D50-D64)	7	0.3	*
14	Certain conditions originating in the perinatal period (P00-P96)	7	0.3	*
...	All other causes (Residual)	454	21.1	2.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 15-19 years				
...	All causes	7,734	100.0	47.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,560	46.0	21.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,516	19.6	9.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	615	8.0	3.7
4	Malignant neoplasms (C00-C97)	503	6.5	3.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	207	2.7	1.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	145	1.9	0.9
7	Cerebrovascular diseases (I60-I69)	55	0.7	0.3
8	Influenza and pneumonia (J09-J18)	54	0.7	0.3
9	Diabetes mellitus (E10-E14)	38	0.5	0.2
10	Septicemia (A40-A41)	33	0.4	0.2
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	31	0.4	0.2
12	Chronic lower respiratory diseases (J40-J47)	29	0.4	0.2
13	Pregnancy, childbirth and the puerperium (O00-O99)	23	0.3	0.1
14	Complications of medical and surgical care (Y40-Y84,Y88)	13	0.2	*
15	Legal intervention (Y35,Y89.0)	12	0.2	*
...	All other causes (Residual)	900	11.6	5.5
White, both sexes, 20-24 years				
...	All causes	13,920	100.0	82.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,831	49.1	40.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,507	18.0	14.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	996	7.2	5.9
4	Malignant neoplasms (C00-C97)	764	5.5	4.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	436	3.1	2.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	161	1.2	1.0
7	Influenza and pneumonia (J09-J18)	111	0.8	0.7
8	Cerebrovascular diseases (I60-I69)	88	0.6	0.5
9	Pregnancy, childbirth and the puerperium (O00-O99)	79	0.6	0.5
10	Diabetes mellitus (E10-E14)	77	0.6	0.5
11	Septicemia (A40-A41)	59	0.4	0.4
12	Chronic lower respiratory diseases (J40-J47)	52	0.4	0.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	40	0.3	0.2
14	Legal intervention (Y35,Y89.0)	37	0.3	0.2
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	31	0.2	0.2
...	All other causes (Residual)	1,651	11.9	9.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 25-29 years				
...	All causes	15,355	100.0	93.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,775	44.1	41.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,615	17.0	15.9
3	Malignant neoplasms (C00-C97)	1,011	6.6	6.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	922	6.0	5.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	797	5.2	4.9
6	Influenza and pneumonia (J09-J18)	182	1.2	1.1
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	154	1.0	0.9
8	Diabetes mellitus (E10-E14)	151	1.0	0.9
9	Pregnancy, childbirth and the puerperium (O00-O99)	129	0.8	0.8
10	Cerebrovascular diseases (I60-I69)	124	0.8	0.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	111	0.7	0.7
12	Septicemia (A40-A41)	95	0.6	0.6
13	Human immunodeficiency virus (HIV) disease (B20-B24)	59	0.4	0.4
14	Chronic lower respiratory diseases (J40-J47)	57	0.4	0.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	51	0.3	0.3
...	All other causes (Residual)	2,122	13.8	12.9
White, both sexes, 30-34 years				
...	All causes	17,360	100.0	109.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,470	37.3	40.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,612	15.0	16.5
3	Malignant neoplasms (C00-C97)	1,636	9.4	10.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,395	8.0	8.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	782	4.5	4.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	282	1.6	1.8
7	Diabetes mellitus (E10-E14)	281	1.6	1.8
8	Cerebrovascular diseases (I60-I69)	239	1.4	1.5
9	Influenza and pneumonia (J09-J18)	208	1.2	1.3
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	183	1.1	1.2
11	Human immunodeficiency virus (HIV) disease (B20-B24)	173	1.0	1.1
12	Septicemia (A40-A41)	142	0.8	0.9
13	Chronic lower respiratory diseases (J40-J47)	106	0.6	0.7
14	Pregnancy, childbirth and the puerperium (O00-O99)	101	0.6	0.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	77	0.4	0.5
...	All other causes (Residual)	2,673	15.4	16.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 35-39 years				
...	All causes	20,540	100.0	135.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,999	29.2	39.7
2	Malignant neoplasms (C00-C97)	2,948	14.4	19.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,606	12.7	17.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,479	12.1	16.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	640	3.1	4.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	611	3.0	4.0
7	Diabetes mellitus (E10-E14)	429	2.1	2.8
8	Cerebrovascular diseases (I60-I69)	347	1.7	2.3
9	Human immunodeficiency virus (HIV) disease (B20-B24)	255	1.2	1.7
10	Influenza and pneumonia (J09-J18)	238	1.2	1.6
11	Septicemia (A40-A41)	188	0.9	1.2
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	160	0.8	1.1
13	Chronic lower respiratory diseases (J40-J47)	130	0.6	0.9
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	128	0.6	0.8
15	Pregnancy, childbirth and the puerperium (O00-O99)	95	0.5	0.6
...	All other causes (Residual)	3,287	16.0	21.7
White, both sexes, 40-44 years				
...	All causes	32,458	100.0	196.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,005	21.6	42.3
2	Malignant neoplasms (C00-C97)	6,012	18.5	36.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,032	15.5	30.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,269	10.1	19.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,445	4.5	8.7
6	Diabetes mellitus (E10-E14)	828	2.6	5.0
7	Cerebrovascular diseases (I60-I69)	730	2.2	4.4
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	631	1.9	3.8
9	Human immunodeficiency virus (HIV) disease (B20-B24)	435	1.3	2.6
10	Chronic lower respiratory diseases (J40-J47)	399	1.2	2.4
11	Influenza and pneumonia (J09-J18)	390	1.2	2.4
12	Septicemia (A40-A41)	355	1.1	2.1
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	241	0.7	1.5
14	Viral hepatitis (B15-B19)	194	0.6	1.2
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	182	0.6	1.1
...	All other causes (Residual)	5,310	16.4	32.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 45-49 years				
...	All causes	55,764	100.0	314.9
1	Malignant neoplasms (C00-C97)	12,963	23.2	73.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,148	18.2	57.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,682	15.6	49.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,957	7.1	22.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	3,072	5.5	17.3
6	Diabetes mellitus (E10-E14)	1,641	2.9	9.3
7	Cerebrovascular diseases (I60-I69)	1,423	2.6	8.0
8	Chronic lower respiratory diseases (J40-J47)	1,114	2.0	6.3
9	Human immunodeficiency virus (HIV) disease (B20-B24)	673	1.2	3.8
10	Influenza and pneumonia (J09-J18)	657	1.2	3.7
11	Viral hepatitis (B15-B19)	654	1.2	3.7
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	653	1.2	3.7
13	Septicemia (A40-A41)	624	1.1	3.5
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	434	0.8	2.5
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	318	0.6	1.8
...	All other causes (Residual)	8,751	15.7	49.4
White, both sexes, 50-54 years				
...	All causes	87,662	100.0	478.3
1	Malignant neoplasms (C00-C97)	25,540	29.1	139.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	17,050	19.4	93.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,070	10.3	49.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	4,614	5.3	25.2
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,311	4.9	23.5
6	Chronic lower respiratory diseases (J40-J47)	2,741	3.1	15.0
7	Diabetes mellitus (E10-E14)	2,666	3.0	14.5
8	Cerebrovascular diseases (I60-I69)	2,324	2.7	12.7
9	Viral hepatitis (B15-B19)	1,288	1.5	7.0
10	Septicemia (A40-A41)	1,132	1.3	6.2
11	Influenza and pneumonia (J09-J18)	1,073	1.2	5.9
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	770	0.9	4.2
13	Human immunodeficiency virus (HIV) disease (B20-B24)	619	0.7	3.4
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	578	0.7	3.2
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	514	0.6	2.8
...	All other causes (Residual)	13,372	15.3	73.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 55-59 years				
...	All causes	114,304	100.0	685.2
1	Malignant neoplasms (C00-C97)	38,618	33.8	231.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	23,619	20.7	141.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,089	6.2	42.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	5,063	4.4	30.4
5	Chronic lower respiratory diseases (J40-J47)	4,919	4.3	29.5
6	Diabetes mellitus (E10-E14)	3,964	3.5	23.8
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,586	3.1	21.5
8	Cerebrovascular diseases (I60-I69)	3,176	2.8	19.0
9	Viral hepatitis (B15-B19)	1,687	1.5	10.1
10	Septicemia (A40-A41)	1,518	1.3	9.1
11	Influenza and pneumonia (J09-J18)	1,424	1.2	8.5
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,241	1.1	7.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	747	0.7	4.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	482	0.4	2.9
15	Human immunodeficiency virus (HIV) disease (B20-B24)	479	0.4	2.9
...	All other causes (Residual)	16,692	14.6	100.1
White, both sexes, 60-64 years				
...	All causes	145,386	100.0	972.9
1	Malignant neoplasms (C00-C97)	53,247	36.6	356.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	31,075	21.4	208.0
3	Chronic lower respiratory diseases (J40-J47)	8,400	5.8	56.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,575	3.8	37.3
5	Diabetes mellitus (E10-E14)	5,456	3.8	36.5
6	Cerebrovascular diseases (I60-I69)	4,758	3.3	31.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	4,208	2.9	28.2
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,565	1.8	17.2
9	Septicemia (A40-A41)	2,149	1.5	14.4
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,944	1.3	13.0
11	Influenza and pneumonia (J09-J18)	1,914	1.3	12.8
12	Viral hepatitis (B15-B19)	1,095	0.8	7.3
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	997	0.7	6.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	743	0.5	5.0
15	Aortic aneurysm and dissection (I71)	627	0.4	4.2
...	All other causes (Residual)	20,633	14.2	138.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 65-69 years				
...	All causes	161,837	100.0	1,475.5
1	Malignant neoplasms (C00-C97)	59,888	37.0	546.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	34,095	21.1	310.9
3	Chronic lower respiratory diseases (J40-J47)	12,833	7.9	117.0
4	Diabetes mellitus (E10-E14)	5,961	3.7	54.3
5	Cerebrovascular diseases (I60-I69)	5,767	3.6	52.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,297	2.7	39.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,912	1.8	26.5
8	Septicemia (A40-A41)	2,387	1.5	21.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,381	1.5	21.7
10	Influenza and pneumonia (J09-J18)	2,306	1.4	21.0
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,690	1.0	15.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,184	0.7	10.8
13	Alzheimer's disease (G30)	1,179	0.7	10.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	983	0.6	9.0
15	Aortic aneurysm and dissection (I71)	833	0.5	7.6
...	All other causes (Residual)	23,141	14.3	211.0
White, both sexes, 70-74 years				
...	All causes	188,202	100.0	2,291.3
1	Malignant neoplasms (C00-C97)	64,448	34.2	784.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	39,832	21.2	484.9
3	Chronic lower respiratory diseases (J40-J47)	16,935	9.0	206.2
4	Cerebrovascular diseases (I60-I69)	8,045	4.3	97.9
5	Diabetes mellitus (E10-E14)	6,548	3.5	79.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,414	2.3	53.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,274	1.7	39.9
8	Influenza and pneumonia (J09-J18)	3,113	1.7	37.9
9	Septicemia (A40-A41)	2,851	1.5	34.7
10	Alzheimer's disease (G30)	2,667	1.4	32.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	2,420	1.3	29.5
12	Parkinson's disease (G20-G21)	1,784	0.9	21.7
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,498	0.8	18.2
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,305	0.7	15.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,284	0.7	15.6
...	All other causes (Residual)	27,784	14.8	338.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 75-79 years				
...	All causes	236,226	100.0	3,697.3
1	Malignant neoplasms (C00-C97)	67,676	28.6	1,059.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	52,850	22.4	827.2
3	Chronic lower respiratory diseases (J40-J47)	21,054	8.9	329.5
4	Cerebrovascular diseases (I60-I69)	12,669	5.4	198.3
5	Diabetes mellitus (E10-E14)	7,255	3.1	113.6
6	Alzheimer's disease (G30)	7,142	3.0	111.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,242	2.2	82.0
8	Influenza and pneumonia (J09-J18)	4,494	1.9	70.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,450	1.9	69.6
10	Parkinson's disease (G20-G21)	3,618	1.5	56.6
11	Septicemia (A40-A41)	3,605	1.5	56.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,069	0.9	32.4
13	Chronic liver disease and cirrhosis (K70,K73-K74)	1,989	0.8	31.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,770	0.7	27.7
15	Pneumonitis due to solids and liquids (J69)	1,707	0.7	26.7
...	All other causes (Residual)	38,636	16.4	604.7
White, both sexes, 80-84 years				
...	All causes	315,701	100.0	6,170.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	77,855	24.7	1,521.7
2	Malignant neoplasms (C00-C97)	69,316	22.0	1,354.8
3	Chronic lower respiratory diseases (J40-J47)	25,033	7.9	489.3
4	Cerebrovascular diseases (I60-I69)	19,586	6.2	382.8
5	Alzheimer's disease (G30)	14,960	4.7	292.4
6	Diabetes mellitus (E10-E14)	8,632	2.7	168.7
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,629	2.4	149.1
8	Influenza and pneumonia (J09-J18)	7,426	2.4	145.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,477	2.1	126.6
10	Parkinson's disease (G20-G21)	5,773	1.8	112.8
11	Septicemia (A40-A41)	4,540	1.4	88.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,339	1.1	65.3
13	Pneumonitis due to solids and liquids (J69)	2,896	0.9	56.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,398	0.8	46.9
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,582	0.5	30.9
...	All other causes (Residual)	58,259	18.5	1,138.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 85-89 years				
...	All causes	349,335	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	97,542	27.9	---
2	Malignant neoplasms (C00-C97)	53,955	15.4	---
3	Cerebrovascular diseases (I60-I69)	23,673	6.8	---
4	Alzheimer's disease (G30)	22,416	6.4	---
5	Chronic lower respiratory diseases (J40-J47)	21,934	6.3	---
6	Influenza and pneumonia (J09-J18)	9,700	2.8	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,825	2.5	---
8	Diabetes mellitus (E10-E14)	7,997	2.3	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,436	2.1	---
10	Parkinson's disease (G20-G21)	5,654	1.6	---
11	Septicemia (A40-A41)	4,841	1.4	---
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,435	1.3	---
13	Pneumonitis due to solids and liquids (J69)	3,810	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,569	0.7	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1,754	0.5	---
...	All other causes (Residual)	72,794	20.8	---
White, both sexes, 90-94 years				
...	All causes	251,677	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	78,565	31.2	---
2	Malignant neoplasms (C00-C97)	25,598	10.2	---
3	Alzheimer's disease (G30)	19,089	7.6	---
4	Cerebrovascular diseases (I60-I69)	17,500	7.0	---
5	Chronic lower respiratory diseases (J40-J47)	11,555	4.6	---
6	Influenza and pneumonia (J09-J18)	8,283	3.3	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,303	2.5	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,131	2.0	---
9	Diabetes mellitus (E10-E14)	4,506	1.8	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,009	1.6	---
11	Pneumonitis due to solids and liquids (J69)	3,134	1.2	---
12	Septicemia (A40-A41)	2,970	1.2	---
13	Parkinson's disease (G20-G21)	2,857	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,450	0.6	---
15	Atherosclerosis (I70)	1,353	0.5	---
...	All other causes (Residual)	59,374	23.6	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 95-99 years				
...	All causes	100,047	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	33,219	33.2	---
2	Alzheimer's disease (G30)	8,240	8.2	---
3	Cerebrovascular diseases (I60-I69)	6,675	6.7	---
4	Malignant neoplasms (C00-C97)	6,455	6.5	---
5	Influenza and pneumonia (J09-J18)	3,871	3.9	---
6	Chronic lower respiratory diseases (J40-J47)	3,612	3.6	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,496	2.5	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,934	1.9	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,840	1.8	---
10	Diabetes mellitus (E10-E14)	1,308	1.3	---
11	Pneumonitis due to solids and liquids (J69)	1,197	1.2	---
12	Septicemia (A40-A41)	983	1.0	---
13	Atherosclerosis (I70)	707	0.7	---
14	Parkinson's disease (G20-G21)	609	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	489	0.5	---
...	All other causes (Residual)	26,412	26.4	---
White, both sexes, 100 years and over				
...	All causes	20,103	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,047	35.1	---
2	Alzheimer's disease (G30)	1,512	7.5	---
3	Cerebrovascular diseases (I60-I69)	1,197	6.0	---
4	Influenza and pneumonia (J09-J18)	948	4.7	---
5	Malignant neoplasms (C00-C97)	765	3.8	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	544	2.7	---
7	Chronic lower respiratory diseases (J40-J47)	501	2.5	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	483	2.4	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	304	1.5	---
10	Pneumonitis due to solids and liquids (J69)	231	1.1	---
11	Diabetes mellitus (E10-E14)	208	1.0	---
12	Atherosclerosis (I70)	204	1.0	---
13	Septicemia (A40-A41)	167	0.8	---
14	Anemias (D50-D64)	87	0.4	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	70	0.3	---
...	All other causes (Residual)	5,835	29.0	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, all ages				
...	All causes	1,071,966	100.0	876.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	265,596	24.8	217.1
2	Malignant neoplasms (C00-C97)	259,323	24.2	212.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	68,123	6.4	55.7
4	Chronic lower respiratory diseases (J40-J47)	61,507	5.7	50.3
5	Cerebrovascular diseases (I60-I69)	43,264	4.0	35.4
6	Diabetes mellitus (E10-E14)	30,783	2.9	25.2
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	28,103	2.6	23.0
8	Alzheimer's disease (G30)	23,640	2.2	19.3
9	Influenza and pneumonia (J09-J18)	21,953	2.0	17.9
10	Chronic liver disease and cirrhosis (K70,K73-K74)	19,067	1.8	15.6
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18,275	1.7	14.9
12	Septicemia (A40-A41)	13,371	1.2	10.9
13	Parkinson's disease (G20-G21)	12,746	1.2	10.4
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	8,698	0.8	7.1
15	Pneumonitis due to solids and liquids (J69)	8,653	0.8	7.1
...	All other causes (Residual)	188,864	17.6	154.4
White, male, 1-4 years				
...	All causes	1,689	100.0	27.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	634	37.5	10.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	193	11.4	3.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	144	8.5	2.3
4	Malignant neoplasms (C00-C97)	136	8.1	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	55	3.3	0.9
6	Influenza and pneumonia (J09-J18)	32	1.9	0.5
7	Septicemia (A40-A41)	20	1.2	0.3
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	1.1	*
9	Chronic lower respiratory diseases (J40-J47)	17	1.0	*
10	Cerebrovascular diseases (I60-I69)	14	0.8	*
11	Certain conditions originating in the perinatal period (P00-P96)	11	0.7	*
12	Acute bronchitis and bronchiolitis (J20-J21)	7	0.4	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.4	*
14	Meningitis (G00,G03)	6	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.3	*
...	All other causes (Residual)	389	23.0	6.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 5-9 years				
...	All causes	1,009	100.0	12.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	350	34.7	4.4
2	Malignant neoplasms (C00-C97)	195	19.3	2.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	82	8.1	1.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	48	4.8	0.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	23	2.3	0.3
5	Influenza and pneumonia (J09-J18)	23	2.3	0.3
7	Chronic lower respiratory diseases (J40-J47)	16	1.6	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	1.3	*
8	Cerebrovascular diseases (I60-I69)	13	1.3	*
10	Septicemia (A40-A41)	12	1.2	*
11	Anemias (D50-D64)	5	0.5	*
12	Certain conditions originating in the perinatal period (P00-P96)	4	0.4	*
13	Meningococcal infection (A39)	3	0.3	*
13	Meningitis (G00,G03)	3	0.3	*
13	Pneumonitis due to solids and liquids (J69)	3	0.3	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.3	*
...	All other causes (Residual)	213	21.1	2.7
White, male, 10-14 years				
...	All causes	1,312	100.0	16.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	441	33.6	5.4
2	Malignant neoplasms (C00-C97)	184	14.0	2.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	155	11.8	1.9
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	59	4.5	0.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	59	4.5	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	42	3.2	0.5
7	Cerebrovascular diseases (I60-I69)	20	1.5	0.2
7	Influenza and pneumonia (J09-J18)	20	1.5	0.2
9	Septicemia (A40-A41)	17	1.3	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	1.1	*
11	Chronic lower respiratory diseases (J40-J47)	13	1.0	*
12	Anemias (D50-D64)	6	0.5	*
12	Diabetes mellitus (E10-E14)	6	0.5	*
14	Pneumonitis due to solids and liquids (J69)	5	0.4	*
15	Certain conditions originating in the perinatal period (P00-P96)	4	0.3	*
...	All other causes (Residual)	267	20.4	3.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 15-19 years				
...	All causes	5,484	100.0	64.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,512	45.8	29.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,213	22.1	14.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	505	9.2	6.0
4	Malignant neoplasms (C00-C97)	307	5.6	3.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	145	2.6	1.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	94	1.7	1.1
7	Influenza and pneumonia (J09-J18)	34	0.6	0.4
8	Cerebrovascular diseases (I60-I69)	33	0.6	0.4
9	Diabetes mellitus (E10-E14)	23	0.4	0.3
10	Septicemia (A40-A41)	21	0.4	0.2
10	Chronic lower respiratory diseases (J40-J47)	21	0.4	0.2
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	0.2	*
13	Legal intervention (Y35,Y89.0)	12	0.2	*
14	Anemias (D50-D64)	8	0.1	*
15	Meningitis (G00,G03)	4	0.1	*
15	Pneumonitis due to solids and liquids (J69)	4	0.1	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.1	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.1	*
...	All other causes (Residual)	527	9.6	6.2
White, male, 20-24 years				
...	All causes	10,355	100.0	119.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,212	50.3	60.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,094	20.2	24.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	803	7.8	9.3
4	Malignant neoplasms (C00-C97)	458	4.4	5.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	307	3.0	3.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	105	1.0	1.2
7	Influenza and pneumonia (J09-J18)	64	0.6	0.7
8	Diabetes mellitus (E10-E14)	53	0.5	0.6
9	Cerebrovascular diseases (I60-I69)	50	0.5	0.6
10	Chronic lower respiratory diseases (J40-J47)	39	0.4	0.5
11	Legal intervention (Y35,Y89.0)	36	0.3	0.4
12	Septicemia (A40-A41)	34	0.3	0.4
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	0.2	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18	0.2	*
15	Pneumonitis due to solids and liquids (J69)	17	0.2	*
...	All other causes (Residual)	1,047	10.1	12.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 25-29 years				
...	All causes	10,968	100.0	130.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,119	46.7	61.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,128	19.4	25.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	708	6.5	8.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	559	5.1	6.7
5	Malignant neoplasms (C00-C97)	553	5.0	6.6
6	Influenza and pneumonia (J09-J18)	98	0.9	1.2
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	89	0.8	1.1
8	Diabetes mellitus (E10-E14)	84	0.8	1.0
9	Cerebrovascular diseases (I60-I69)	63	0.6	0.8
10	Chronic liver disease and cirrhosis (K70,K73-K74)	62	0.6	0.7
11	Septicemia (A40-A41)	56	0.5	0.7
12	Legal intervention (Y35,Y89.0)	47	0.4	0.6
13	Human immunodeficiency virus (HIV) disease (B20-B24)	37	0.3	0.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	33	0.3	0.4
14	Chronic lower respiratory diseases (J40-J47)	33	0.3	0.4
...	All other causes (Residual)	1,299	11.8	15.5
White, male, 30-34 years				
...	All causes	11,707	100.0	145.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,660	39.8	57.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,053	17.5	25.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	958	8.2	11.9
4	Malignant neoplasms (C00-C97)	785	6.7	9.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	595	5.1	7.4
6	Chronic liver disease and cirrhosis (K70,K73-K74)	185	1.6	2.3
7	Diabetes mellitus (E10-E14)	174	1.5	2.2
8	Cerebrovascular diseases (I60-I69)	136	1.2	1.7
9	Human immunodeficiency virus (HIV) disease (B20-B24)	134	1.1	1.7
10	Influenza and pneumonia (J09-J18)	128	1.1	1.6
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	102	0.9	1.3
12	Septicemia (A40-A41)	83	0.7	1.0
13	Chronic lower respiratory diseases (J40-J47)	54	0.5	0.7
14	Legal intervention (Y35,Y89.0)	48	0.4	0.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	43	0.4	0.5
...	All other causes (Residual)	1,569	13.4	19.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 35-39 years				
...	All causes	13,232	100.0	173.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,170	31.5	54.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,016	15.2	26.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,764	13.3	23.1
4	Malignant neoplasms (C00-C97)	1,330	10.1	17.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	457	3.5	6.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	413	3.1	5.4
7	Diabetes mellitus (E10-E14)	270	2.0	3.5
8	Human immunodeficiency virus (HIV) disease (B20-B24)	186	1.4	2.4
9	Cerebrovascular diseases (I60-I69)	181	1.4	2.4
10	Influenza and pneumonia (J09-J18)	133	1.0	1.7
11	Septicemia (A40-A41)	94	0.7	1.2
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	92	0.7	1.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	86	0.6	1.1
14	Aortic aneurysm and dissection (I71)	72	0.5	0.9
15	Chronic lower respiratory diseases (J40-J47)	55	0.4	0.7
...	All other causes (Residual)	1,913	14.5	25.0
White, male, 40-44 years				
...	All causes	20,164	100.0	241.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,711	23.4	56.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,594	17.8	43.1
3	Malignant neoplasms (C00-C97)	2,702	13.4	32.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,475	12.3	29.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	943	4.7	11.3
6	Diabetes mellitus (E10-E14)	514	2.5	6.2
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	439	2.2	5.3
8	Cerebrovascular diseases (I60-I69)	397	2.0	4.8
9	Human immunodeficiency virus (HIV) disease (B20-B24)	346	1.7	4.1
10	Influenza and pneumonia (J09-J18)	202	1.0	2.4
11	Chronic lower respiratory diseases (J40-J47)	193	1.0	2.3
12	Septicemia (A40-A41)	174	0.9	2.1
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	143	0.7	1.7
14	Viral hepatitis (B15-B19)	127	0.6	1.5
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	108	0.5	1.3
...	All other causes (Residual)	3,096	15.4	37.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 45-49 years				
...	All causes	34,497	100.0	389.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,363	21.3	83.2
2	Malignant neoplasms (C00-C97)	6,241	18.1	70.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,820	16.9	65.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,957	8.6	33.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,006	5.8	22.7
6	Diabetes mellitus (E10-E14)	1,051	3.0	11.9
7	Cerebrovascular diseases (I60-I69)	779	2.3	8.8
8	Human immunodeficiency virus (HIV) disease (B20-B24)	549	1.6	6.2
9	Chronic lower respiratory diseases (J40-J47)	502	1.5	5.7
10	Viral hepatitis (B15-B19)	477	1.4	5.4
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	458	1.3	5.2
12	Influenza and pneumonia (J09-J18)	382	1.1	4.3
13	Septicemia (A40-A41)	312	0.9	3.5
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	270	0.8	3.1
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	218	0.6	2.5
...	All other causes (Residual)	5,112	14.8	57.8
White, male, 50-54 years				
...	All causes	54,283	100.0	597.7
1	Malignant neoplasms (C00-C97)	13,240	24.4	145.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,498	23.0	137.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,033	11.1	66.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,234	6.0	35.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	3,143	5.8	34.6
6	Diabetes mellitus (E10-E14)	1,740	3.2	19.2
7	Cerebrovascular diseases (I60-I69)	1,332	2.5	14.7
8	Chronic lower respiratory diseases (J40-J47)	1,313	2.4	14.5
9	Viral hepatitis (B15-B19)	916	1.7	10.1
10	Septicemia (A40-A41)	628	1.2	6.9
10	Influenza and pneumonia (J09-J18)	628	1.2	6.9
12	Human immunodeficiency virus (HIV) disease (B20-B24)	493	0.9	5.4
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	455	0.8	5.0
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	399	0.7	4.4
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	328	0.6	3.6
...	All other causes (Residual)	7,903	14.6	87.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 55-59 years				
...	All causes	71,487	100.0	874.7
1	Malignant neoplasms (C00-C97)	21,492	30.1	263.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	17,179	24.0	210.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,876	6.8	59.7
4	Chronic liver disease and cirrhosis (K70,K73-K74)	3,634	5.1	44.5
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,761	3.9	33.8
6	Chronic lower respiratory diseases (J40-J47)	2,582	3.6	31.6
7	Diabetes mellitus (E10-E14)	2,457	3.4	30.1
8	Cerebrovascular diseases (I60-I69)	1,857	2.6	22.7
9	Viral hepatitis (B15-B19)	1,273	1.8	15.6
10	Influenza and pneumonia (J09-J18)	840	1.2	10.3
11	Septicemia (A40-A41)	835	1.2	10.2
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	732	1.0	9.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	501	0.7	6.1
14	Human immunodeficiency virus (HIV) disease (B20-B24)	390	0.5	4.8
15	Aortic aneurysm and dissection (I71)	346	0.5	4.2
...	All other causes (Residual)	9,732	13.6	119.1
White, male, 60-64 years				
...	All causes	87,879	100.0	1,212.1
1	Malignant neoplasms (C00-C97)	29,773	33.9	410.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	21,641	24.6	298.5
3	Chronic lower respiratory diseases (J40-J47)	4,370	5.0	60.3
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,807	4.3	52.5
5	Diabetes mellitus (E10-E14)	3,324	3.8	45.8
6	Chronic liver disease and cirrhosis (K70,K73-K74)	2,945	3.4	40.6
7	Cerebrovascular diseases (I60-I69)	2,658	3.0	36.7
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,979	2.3	27.3
9	Septicemia (A40-A41)	1,120	1.3	15.4
10	Influenza and pneumonia (J09-J18)	1,111	1.3	15.3
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,101	1.3	15.2
12	Viral hepatitis (B15-B19)	792	0.9	10.9
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	620	0.7	8.6
14	Aortic aneurysm and dissection (I71)	461	0.5	6.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	434	0.5	6.0
...	All other causes (Residual)	11,743	13.4	162.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 65-69 years				
...	All causes	93,695	100.0	1,792.0
1	Malignant neoplasms (C00-C97)	33,085	35.3	632.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	22,627	24.1	432.8
3	Chronic lower respiratory diseases (J40-J47)	6,490	6.9	124.1
4	Diabetes mellitus (E10-E14)	3,593	3.8	68.7
5	Cerebrovascular diseases (I60-I69)	3,109	3.3	59.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,760	2.9	52.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,906	2.0	36.5
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,332	1.4	25.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,300	1.4	24.9
10	Influenza and pneumonia (J09-J18)	1,291	1.4	24.7
11	Septicemia (A40-A41)	1,242	1.3	23.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	673	0.7	12.9
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	590	0.6	11.3
13	Aortic aneurysm and dissection (I71)	590	0.6	11.3
15	Parkinson's disease (G20-G21)	539	0.6	10.3
...	All other causes (Residual)	12,568	13.4	240.4
White, male, 70-74 years				
...	All causes	105,008	100.0	2,759.9
1	Malignant neoplasms (C00-C97)	35,519	33.8	933.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	24,784	23.6	651.4
3	Chronic lower respiratory diseases (J40-J47)	8,496	8.1	223.3
4	Cerebrovascular diseases (I60-I69)	4,144	3.9	108.9
5	Diabetes mellitus (E10-E14)	3,731	3.6	98.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,697	2.6	70.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,747	1.7	45.9
8	Influenza and pneumonia (J09-J18)	1,718	1.6	45.2
9	Septicemia (A40-A41)	1,471	1.4	38.7
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,399	1.3	36.8
11	Parkinson's disease (G20-G21)	1,211	1.2	31.8
12	Alzheimer's disease (G30)	1,128	1.1	29.6
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,080	1.0	28.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	755	0.7	19.8
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	736	0.7	19.3
...	All other causes (Residual)	14,392	13.7	378.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 75-79 years				
...	All causes	124,662	100.0	4,404.8
1	Malignant neoplasms (C00-C97)	36,756	29.5	1,298.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	30,349	24.3	1,072.4
3	Chronic lower respiratory diseases (J40-J47)	10,186	8.2	359.9
4	Cerebrovascular diseases (I60-I69)	5,914	4.7	209.0
5	Diabetes mellitus (E10-E14)	3,884	3.1	137.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,954	2.4	104.4
7	Alzheimer's disease (G30)	2,888	2.3	102.0
8	Influenza and pneumonia (J09-J18)	2,395	1.9	84.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,379	1.9	84.1
10	Parkinson's disease (G20-G21)	2,356	1.9	83.2
11	Septicemia (A40-A41)	1,766	1.4	62.4
12	Chronic liver disease and cirrhosis (K70,K73-K74)	1,087	0.9	38.4
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,009	0.8	35.7
14	Pneumonitis due to solids and liquids (J69)	977	0.8	34.5
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	958	0.8	33.9
...	All other causes (Residual)	18,804	15.1	664.4
White, male, 80-84 years				
...	All causes	153,672	100.0	7,332.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	40,299	26.2	1,922.8
2	Malignant neoplasms (C00-C97)	36,406	23.7	1,737.1
3	Chronic lower respiratory diseases (J40-J47)	11,782	7.7	562.2
4	Cerebrovascular diseases (I60-I69)	8,048	5.2	384.0
5	Alzheimer's disease (G30)	5,471	3.6	261.0
6	Diabetes mellitus (E10-E14)	4,258	2.8	203.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,970	2.6	189.4
8	Influenza and pneumonia (J09-J18)	3,665	2.4	174.9
9	Parkinson's disease (G20-G21)	3,440	2.2	164.1
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,420	2.2	163.2
11	Septicemia (A40-A41)	2,147	1.4	102.4
12	Pneumonitis due to solids and liquids (J69)	1,631	1.1	77.8
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,395	0.9	66.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,311	0.9	62.6
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	890	0.6	42.5
...	All other causes (Residual)	25,539	16.6	1,218.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 85-89 years				
...	All causes	146,518	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	42,850	29.2	---
2	Malignant neoplasms (C00-C97)	26,305	18.0	---
3	Chronic lower respiratory diseases (J40-J47)	9,629	6.6	---
4	Cerebrovascular diseases (I60-I69)	8,159	5.6	---
5	Alzheimer's disease (G30)	6,881	4.7	---
6	Influenza and pneumonia (J09-J18)	4,430	3.0	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,895	2.7	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,547	2.4	---
9	Diabetes mellitus (E10-E14)	3,484	2.4	---
10	Parkinson's disease (G20-G21)	3,192	2.2	---
11	Pneumonitis due to solids and liquids (J69)	2,037	1.4	---
12	Septicemia (A40-A41)	1,917	1.3	---
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,504	1.0	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,305	0.9	---
15	Aortic aneurysm and dissection (I71)	674	0.5	---
...	All other causes (Residual)	26,709	18.2	---
White, male, 90-94 years				
...	All causes	87,243	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	28,514	32.7	---
2	Malignant neoplasms (C00-C97)	11,309	13.0	---
3	Alzheimer's disease (G30)	4,844	5.6	---
4	Cerebrovascular diseases (I60-I69)	4,780	5.5	---
5	Chronic lower respiratory diseases (J40-J47)	4,458	5.1	---
6	Influenza and pneumonia (J09-J18)	3,230	3.7	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,296	2.6	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,268	2.6	---
9	Diabetes mellitus (E10-E14)	1,716	2.0	---
10	Pneumonitis due to solids and liquids (J69)	1,477	1.7	---
11	Parkinson's disease (G20-G21)	1,429	1.6	---
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,117	1.3	---
13	Septicemia (A40-A41)	1,070	1.2	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	601	0.7	---
15	Atherosclerosis (I70)	421	0.5	---
...	All other causes (Residual)	17,713	20.3	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 95-99 years				
...	All causes	25,154	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,731	34.7	---
2	Malignant neoplasms (C00-C97)	2,311	9.2	---
3	Cerebrovascular diseases (I60-I69)	1,375	5.5	---
4	Alzheimer's disease (G30)	1,369	5.4	---
5	Influenza and pneumonia (J09-J18)	1,244	4.9	---
6	Chronic lower respiratory diseases (J40-J47)	1,130	4.5	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	700	2.8	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	638	2.5	---
9	Pneumonitis due to solids and liquids (J69)	480	1.9	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	404	1.6	---
11	Diabetes mellitus (E10-E14)	373	1.5	---
12	Septicemia (A40-A41)	264	1.0	---
13	Parkinson's disease (G20-G21)	238	0.9	---
14	Atherosclerosis (I70)	162	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	140	0.6	---
...	All other causes (Residual)	5,595	22.2	---
White, male, 100 years and over				
...	All causes	3,290	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,197	36.4	---
2	Influenza and pneumonia (J09-J18)	210	6.4	---
3	Malignant neoplasms (C00-C97)	205	6.2	---
4	Cerebrovascular diseases (I60-I69)	154	4.7	---
5	Alzheimer's disease (G30)	153	4.7	---
6	Chronic lower respiratory diseases (J40-J47)	117	3.6	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	88	2.7	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	76	2.3	---
9	Pneumonitis due to solids and liquids (J69)	72	2.2	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	69	2.1	---
11	Diabetes mellitus (E10-E14)	44	1.3	---
12	Septicemia (A40-A41)	39	1.2	---
13	Atherosclerosis (I70)	30	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	21	0.6	---
15	Parkinson's disease (G20-G21)	16	0.5	---
...	All other causes (Residual)	799	24.3	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, all ages				
...	All causes	1,084,111	100.0	868.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	248,105	22.9	198.8
2	Malignant neoplasms (C00-C97)	233,025	21.5	186.8
3	Chronic lower respiratory diseases (J40-J47)	69,988	6.5	56.1
4	Cerebrovascular diseases (I60-I69)	65,278	6.0	52.3
5	Alzheimer's disease (G30)	54,415	5.0	43.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	41,628	3.8	33.4
7	Diabetes mellitus (E10-E14)	27,191	2.5	21.8
8	Influenza and pneumonia (J09-J18)	24,705	2.3	19.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17,993	1.7	14.4
10	Septicemia (A40-A41)	15,467	1.4	12.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	13,147	1.2	10.5
12	Chronic liver disease and cirrhosis (K70,K73-K74)	10,440	1.0	8.4
13	Parkinson's disease (G20-G21)	8,942	0.8	7.2
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7,672	0.7	6.1
15	Pneumonitis due to solids and liquids (J69)	7,610	0.7	6.1
...	All other causes (Residual)	238,505	22.0	191.2
White, female, 1-4 years				
...	All causes	1,291	100.0	21.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	393	30.4	6.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	167	12.9	2.8
3	Malignant neoplasms (C00-C97)	139	10.8	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	93	7.2	1.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	52	4.0	0.9
6	Influenza and pneumonia (J09-J18)	34	2.6	0.6
7	Septicemia (A40-A41)	25	1.9	0.4
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	1.1	*
9	Certain conditions originating in the perinatal period (P00-P96)	10	0.8	*
10	Cerebrovascular diseases (I60-I69)	9	0.7	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.5	*
12	Acute bronchitis and bronchiolitis (J20-J21)	5	0.4	*
12	Chronic lower respiratory diseases (J40-J47)	5	0.4	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.4	*
15	Diabetes mellitus (E10-E14)	4	0.3	*
...	All other causes (Residual)	330	25.6	5.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 5-9 years				
...	All causes	766	100.0	10.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	218	28.5	2.9
2	Malignant neoplasms (C00-C97)	156	20.4	2.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	53	6.9	0.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	42	5.5	0.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	35	4.6	0.5
6	Influenza and pneumonia (J09-J18)	24	3.1	0.3
7	Septicemia (A40-A41)	18	2.3	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	15	2.0	*
9	Cerebrovascular diseases (I60-I69)	10	1.3	*
9	Chronic lower respiratory diseases (J40-J47)	10	1.3	*
11	Certain conditions originating in the perinatal period (P00-P96)	8	1.0	*
12	Anemias (D50-D64)	3	0.4	*
13	Meningitis (G00,G03)	2	0.3	*
13	Acute bronchitis and bronchiolitis (J20-J21)	2	0.3	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.3	*
...	All other causes (Residual)	168	21.9	2.2
White, female, 10-14 years				
...	All causes	840	100.0	10.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	233	27.7	3.0
2	Malignant neoplasms (C00-C97)	141	16.8	1.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	65	7.7	0.8
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	64	7.6	0.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	36	4.3	0.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	35	4.2	0.5
7	Cerebrovascular diseases (I60-I69)	16	1.9	*
7	Influenza and pneumonia (J09-J18)	16	1.9	*
9	Diabetes mellitus (E10-E14)	14	1.7	*
10	Chronic lower respiratory diseases (J40-J47)	10	1.2	*
11	Septicemia (A40-A41)	8	1.0	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	1.0	*
13	Meningitis (G00,G03)	3	0.4	*
13	Pneumonitis due to solids and liquids (J69)	3	0.4	*
13	Certain conditions originating in the perinatal period (P00-P96)	3	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.4	*
...	All other causes (Residual)	182	21.7	2.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 15-19 years				
...	All causes	2,250	100.0	28.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,048	46.6	13.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	303	13.5	3.8
3	Malignant neoplasms (C00-C97)	196	8.7	2.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	110	4.9	1.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	62	2.8	0.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	51	2.3	0.6
7	Pregnancy, childbirth and the puerperium (O00-O99)	23	1.0	0.3
8	Cerebrovascular diseases (I60-I69)	22	1.0	0.3
9	Influenza and pneumonia (J09-J18)	20	0.9	0.3
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	0.8	*
11	Diabetes mellitus (E10-E14)	15	0.7	*
12	Septicemia (A40-A41)	12	0.5	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	9	0.4	*
14	Chronic lower respiratory diseases (J40-J47)	8	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.3	*
...	All other causes (Residual)	347	15.4	4.4
White, female, 20-24 years				
...	All causes	3,565	100.0	43.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,619	45.4	19.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	413	11.6	5.0
3	Malignant neoplasms (C00-C97)	306	8.6	3.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	193	5.4	2.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	129	3.6	1.6
6	Pregnancy, childbirth and the puerperium (O00-O99)	79	2.2	1.0
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	56	1.6	0.7
8	Influenza and pneumonia (J09-J18)	47	1.3	0.6
9	Cerebrovascular diseases (I60-I69)	38	1.1	0.5
10	Septicemia (A40-A41)	25	0.7	0.3
11	Diabetes mellitus (E10-E14)	24	0.7	0.3
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	22	0.6	0.3
13	Chronic lower respiratory diseases (J40-J47)	13	0.4	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	0.4	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	12	0.3	*
...	All other causes (Residual)	576	16.2	7.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 25-29 years				
...	All causes	4,387	100.0	54.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,656	37.7	20.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	487	11.1	6.1
3	Malignant neoplasms (C00-C97)	458	10.4	5.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	238	5.4	3.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	214	4.9	2.7
6	Pregnancy, childbirth and the puerperium (O00-O99)	129	2.9	1.6
7	Influenza and pneumonia (J09-J18)	84	1.9	1.0
8	Diabetes mellitus (E10-E14)	67	1.5	0.8
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	65	1.5	0.8
10	Cerebrovascular diseases (I60-I69)	61	1.4	0.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	49	1.1	0.6
12	Septicemia (A40-A41)	39	0.9	0.5
13	Chronic lower respiratory diseases (J40-J47)	24	0.5	0.3
14	Human immunodeficiency virus (HIV) disease (B20-B24)	22	0.5	0.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18	0.4	*
...	All other causes (Residual)	758	17.3	9.5
White, female, 30-34 years				
...	All causes	5,653	100.0	72.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,810	32.0	23.3
2	Malignant neoplasms (C00-C97)	851	15.1	11.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	559	9.9	7.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	437	7.7	5.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	187	3.3	2.4
6	Diabetes mellitus (E10-E14)	107	1.9	1.4
7	Cerebrovascular diseases (I60-I69)	103	1.8	1.3
8	Pregnancy, childbirth and the puerperium (O00-O99)	101	1.8	1.3
9	Chronic liver disease and cirrhosis (K70,K73-K74)	97	1.7	1.2
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	81	1.4	1.0
11	Influenza and pneumonia (J09-J18)	80	1.4	1.0
12	Septicemia (A40-A41)	59	1.0	0.8
13	Chronic lower respiratory diseases (J40-J47)	52	0.9	0.7
14	Human immunodeficiency virus (HIV) disease (B20-B24)	39	0.7	0.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	34	0.6	0.4
...	All other causes (Residual)	1,056	18.7	13.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 35-39 years				
...	All causes	7,308	100.0	97.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,829	25.0	24.5
2	Malignant neoplasms (C00-C97)	1,618	22.1	21.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	715	9.8	9.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	590	8.1	7.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	198	2.7	2.6
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	183	2.5	2.4
7	Cerebrovascular diseases (I60-I69)	166	2.3	2.2
8	Diabetes mellitus (E10-E14)	159	2.2	2.1
9	Influenza and pneumonia (J09-J18)	105	1.4	1.4
10	Pregnancy, childbirth and the puerperium (O00-O99)	95	1.3	1.3
11	Septicemia (A40-A41)	94	1.3	1.3
12	Chronic lower respiratory diseases (J40-J47)	75	1.0	1.0
13	Human immunodeficiency virus (HIV) disease (B20-B24)	69	0.9	0.9
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	68	0.9	0.9
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	42	0.6	0.6
...	All other causes (Residual)	1,302	17.8	17.4
White, female, 40-44 years				
...	All causes	12,294	100.0	149.6
1	Malignant neoplasms (C00-C97)	3,310	26.9	40.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,294	18.7	27.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,438	11.7	17.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	794	6.5	9.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	502	4.1	6.1
6	Cerebrovascular diseases (I60-I69)	333	2.7	4.1
7	Diabetes mellitus (E10-E14)	314	2.6	3.8
8	Chronic lower respiratory diseases (J40-J47)	206	1.7	2.5
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	192	1.6	2.3
10	Influenza and pneumonia (J09-J18)	188	1.5	2.3
11	Septicemia (A40-A41)	181	1.5	2.2
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	98	0.8	1.2
13	Human immunodeficiency virus (HIV) disease (B20-B24)	89	0.7	1.1
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	74	0.6	0.9
15	Viral hepatitis (B15-B19)	67	0.5	0.8
...	All other causes (Residual)	2,214	18.0	26.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 45-49 years				
...	All causes	21,267	100.0	240.0
1	Malignant neoplasms (C00-C97)	6,722	31.6	75.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,862	13.5	32.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,785	13.1	31.4
4	Chronic liver disease and cirrhosis (K70,K73-K74)	1,066	5.0	12.0
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,000	4.7	11.3
6	Cerebrovascular diseases (I60-I69)	644	3.0	7.3
7	Chronic lower respiratory diseases (J40-J47)	612	2.9	6.9
8	Diabetes mellitus (E10-E14)	590	2.8	6.7
9	Septicemia (A40-A41)	312	1.5	3.5
10	Influenza and pneumonia (J09-J18)	275	1.3	3.1
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	195	0.9	2.2
12	Viral hepatitis (B15-B19)	177	0.8	2.0
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	164	0.8	1.9
14	Human immunodeficiency virus (HIV) disease (B20-B24)	124	0.6	1.4
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	121	0.6	1.4
...	All other causes (Residual)	3,618	17.0	40.8
White, female, 50-54 years				
...	All causes	33,379	100.0	361.0
1	Malignant neoplasms (C00-C97)	12,300	36.8	133.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,552	13.6	49.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,037	9.1	32.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	1,471	4.4	15.9
5	Chronic lower respiratory diseases (J40-J47)	1,428	4.3	15.4
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,077	3.2	11.6
7	Cerebrovascular diseases (I60-I69)	992	3.0	10.7
8	Diabetes mellitus (E10-E14)	926	2.8	10.0
9	Septicemia (A40-A41)	504	1.5	5.5
10	Influenza and pneumonia (J09-J18)	445	1.3	4.8
11	Viral hepatitis (B15-B19)	372	1.1	4.0
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	315	0.9	3.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	186	0.6	2.0
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	179	0.5	1.9
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	170	0.5	1.8
...	All other causes (Residual)	5,425	16.3	58.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 55-59 years				
...	All causes	42,817	100.0	503.2
1	Malignant neoplasms (C00-C97)	17,126	40.0	201.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,440	15.0	75.7
3	Chronic lower respiratory diseases (J40-J47)	2,337	5.5	27.5
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,213	5.2	26.0
5	Diabetes mellitus (E10-E14)	1,507	3.5	17.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1,429	3.3	16.8
7	Cerebrovascular diseases (I60-I69)	1,319	3.1	15.5
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	825	1.9	9.7
9	Septicemia (A40-A41)	683	1.6	8.0
10	Influenza and pneumonia (J09-J18)	584	1.4	6.9
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	509	1.2	6.0
12	Viral hepatitis (B15-B19)	414	1.0	4.9
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	246	0.6	2.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	210	0.5	2.5
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	206	0.5	2.4
...	All other causes (Residual)	6,769	15.8	79.6
White, female, 60-64 years				
...	All causes	57,507	100.0	747.6
1	Malignant neoplasms (C00-C97)	23,474	40.8	305.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,434	16.4	122.6
3	Chronic lower respiratory diseases (J40-J47)	4,030	7.0	52.4
4	Diabetes mellitus (E10-E14)	2,132	3.7	27.7
5	Cerebrovascular diseases (I60-I69)	2,100	3.7	27.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,768	3.1	23.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,263	2.2	16.4
8	Septicemia (A40-A41)	1,029	1.8	13.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	843	1.5	11.0
10	Influenza and pneumonia (J09-J18)	803	1.4	10.4
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	586	1.0	7.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	377	0.7	4.9
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	309	0.5	4.0
14	Viral hepatitis (B15-B19)	303	0.5	3.9
15	Alzheimer's disease (G30)	295	0.5	3.8
...	All other causes (Residual)	8,761	15.2	113.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 65-69 years				
...	All causes	68,142	100.0	1,187.2
1	Malignant neoplasms (C00-C97)	26,803	39.3	467.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,468	16.8	199.8
3	Chronic lower respiratory diseases (J40-J47)	6,343	9.3	110.5
4	Cerebrovascular diseases (I60-I69)	2,658	3.9	46.3
5	Diabetes mellitus (E10-E14)	2,368	3.5	41.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,537	2.3	26.8
7	Septicemia (A40-A41)	1,145	1.7	19.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,081	1.6	18.8
9	Influenza and pneumonia (J09-J18)	1,015	1.5	17.7
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,006	1.5	17.5
11	Alzheimer's disease (G30)	646	0.9	11.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	511	0.7	8.9
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	393	0.6	6.8
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	358	0.5	6.2
15	Parkinson's disease (G20-G21)	291	0.4	5.1
...	All other causes (Residual)	10,519	15.4	183.3
White, female, 70-74 years				
...	All causes	83,194	100.0	1,886.9
1	Malignant neoplasms (C00-C97)	28,929	34.8	656.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,048	18.1	341.3
3	Chronic lower respiratory diseases (J40-J47)	8,439	10.1	191.4
4	Cerebrovascular diseases (I60-I69)	3,901	4.7	88.5
5	Diabetes mellitus (E10-E14)	2,817	3.4	63.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,717	2.1	38.9
7	Alzheimer's disease (G30)	1,539	1.8	34.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,527	1.8	34.6
9	Influenza and pneumonia (J09-J18)	1,395	1.7	31.6
10	Septicemia (A40-A41)	1,380	1.7	31.3
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1,021	1.2	23.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	762	0.9	17.3
13	Parkinson's disease (G20-G21)	573	0.7	13.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	529	0.6	12.0
15	Pneumonitis due to solids and liquids (J69)	398	0.5	9.0
...	All other causes (Residual)	13,219	15.9	299.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 75-79 years				
...	All causes	111,564	100.0	3,134.7
1	Malignant neoplasms (C00-C97)	30,920	27.7	868.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	22,501	20.2	632.2
3	Chronic lower respiratory diseases (J40-J47)	10,868	9.7	305.4
4	Cerebrovascular diseases (I60-I69)	6,755	6.1	189.8
5	Alzheimer's disease (G30)	4,254	3.8	119.5
6	Diabetes mellitus (E10-E14)	3,371	3.0	94.7
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,288	2.1	64.3
8	Influenza and pneumonia (J09-J18)	2,099	1.9	59.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,071	1.9	58.2
10	Septicemia (A40-A41)	1,839	1.6	51.7
11	Parkinson's disease (G20-G21)	1,262	1.1	35.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,111	1.0	31.2
13	Chronic liver disease and cirrhosis (K70,K73-K74)	902	0.8	25.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	761	0.7	21.4
15	Pneumonitis due to solids and liquids (J69)	730	0.7	20.5
...	All other causes (Residual)	19,832	17.8	557.2
White, female, 80-84 years				
...	All causes	162,029	100.0	5,364.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	37,556	23.2	1,243.3
2	Malignant neoplasms (C00-C97)	32,910	20.3	1,089.5
3	Chronic lower respiratory diseases (J40-J47)	13,251	8.2	438.7
4	Cerebrovascular diseases (I60-I69)	11,538	7.1	382.0
5	Alzheimer's disease (G30)	9,489	5.9	314.1
6	Diabetes mellitus (E10-E14)	4,374	2.7	144.8
7	Influenza and pneumonia (J09-J18)	3,761	2.3	124.5
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,659	2.3	121.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,057	1.9	101.2
10	Septicemia (A40-A41)	2,393	1.5	79.2
11	Parkinson's disease (G20-G21)	2,333	1.4	77.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,944	1.2	64.4
13	Pneumonitis due to solids and liquids (J69)	1,265	0.8	41.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,087	0.7	36.0
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	836	0.5	27.7
...	All other causes (Residual)	32,576	20.1	1,078.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 85-89 years				
...	All causes	202,817	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	54,692	27.0	---
2	Malignant neoplasms (C00-C97)	27,650	13.6	---
3	Alzheimer's disease (G30)	15,535	7.7	---
4	Cerebrovascular diseases (I60-I69)	15,514	7.6	---
5	Chronic lower respiratory diseases (J40-J47)	12,305	6.1	---
6	Influenza and pneumonia (J09-J18)	5,270	2.6	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,930	2.4	---
8	Diabetes mellitus (E10-E14)	4,513	2.2	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,889	1.9	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,931	1.4	---
11	Septicemia (A40-A41)	2,924	1.4	---
12	Parkinson's disease (G20-G21)	2,462	1.2	---
13	Pneumonitis due to solids and liquids (J69)	1,773	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,264	0.6	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1,157	0.6	---
...	All other causes (Residual)	46,008	22.7	---
White, female, 90-94 years				
...	All causes	164,434	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	50,051	30.4	---
2	Malignant neoplasms (C00-C97)	14,289	8.7	---
3	Alzheimer's disease (G30)	14,245	8.7	---
4	Cerebrovascular diseases (I60-I69)	12,720	7.7	---
5	Chronic lower respiratory diseases (J40-J47)	7,097	4.3	---
6	Influenza and pneumonia (J09-J18)	5,053	3.1	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,007	2.4	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,892	1.8	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,863	1.7	---
10	Diabetes mellitus (E10-E14)	2,790	1.7	---
11	Septicemia (A40-A41)	1,900	1.2	---
12	Pneumonitis due to solids and liquids (J69)	1,657	1.0	---
13	Parkinson's disease (G20-G21)	1,428	0.9	---
14	Atherosclerosis (I70)	932	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	849	0.5	---
...	All other causes (Residual)	41,661	25.3	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 95-99 years				
...	All causes	74,893	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	24,488	32.7	---
2	Alzheimer's disease (G30)	6,871	9.2	---
3	Cerebrovascular diseases (I60-I69)	5,300	7.1	---
4	Malignant neoplasms (C00-C97)	4,144	5.5	---
5	Influenza and pneumonia (J09-J18)	2,627	3.5	---
6	Chronic lower respiratory diseases (J40-J47)	2,482	3.3	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,796	2.4	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,530	2.0	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,202	1.6	---
10	Diabetes mellitus (E10-E14)	935	1.2	---
11	Septicemia (A40-A41)	719	1.0	---
12	Pneumonitis due to solids and liquids (J69)	717	1.0	---
13	Atherosclerosis (I70)	545	0.7	---
14	Parkinson's disease (G20-G21)	371	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	349	0.5	---
...	All other causes (Residual)	20,817	27.8	---
White, female, 100 years and over				
...	All causes	16,813	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,850	34.8	---
2	Alzheimer's disease (G30)	1,359	8.1	---
3	Cerebrovascular diseases (I60-I69)	1,043	6.2	---
4	Influenza and pneumonia (J09-J18)	738	4.4	---
5	Malignant neoplasms (C00-C97)	560	3.3	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	468	2.8	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	395	2.3	---
8	Chronic lower respiratory diseases (J40-J47)	384	2.3	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	235	1.4	---
10	Atherosclerosis (I70)	174	1.0	---
11	Diabetes mellitus (E10-E14)	164	1.0	---
12	Pneumonitis due to solids and liquids (J69)	159	0.9	---
13	Septicemia (A40-A41)	128	0.8	---
14	Anemias (D50-D64)	73	0.4	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	49	0.3	---
...	All other causes (Residual)	5,034	29.9	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, all ages				
...	All causes	290,100	100.0	679.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	68,372	23.6	160.2
2	Malignant neoplasms (C00-C97)	66,820	23.0	156.6
3	Cerebrovascular diseases (I60-I69)	15,853	5.5	37.1
4	Diabetes mellitus (E10-E14)	12,895	4.4	30.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,531	4.3	29.4
6	Chronic lower respiratory diseases (J40-J47)	9,086	3.1	21.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,087	2.8	18.9
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7,858	2.7	18.4
9	Septicemia (A40-A41)	6,028	2.1	14.1
10	Alzheimer's disease (G30)	5,482	1.9	12.8
11	Influenza and pneumonia (J09-J18)	5,167	1.8	12.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,006	1.7	11.7
13	Human immunodeficiency virus (HIV) disease (B20-B24)	4,215	1.5	9.9
14	Certain conditions originating in the perinatal period (P00-P96)	4,055	1.4	9.5
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,816	1.0	6.6
...	All other causes (Residual)	55,829	19.2	130.8
Black, both sexes, 1-4 years				
...	All causes	1,052	100.0	38.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	298	28.3	10.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	159	15.1	5.8
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	108	10.3	3.9
4	Malignant neoplasms (C00-C97)	55	5.2	2.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	54	5.1	2.0
6	Influenza and pneumonia (J09-J18)	32	3.0	1.2
7	Chronic lower respiratory diseases (J40-J47)	25	2.4	0.9
8	Cerebrovascular diseases (I60-I69)	17	1.6	*
9	Certain conditions originating in the perinatal period (P00-P96)	13	1.2	*
10	Septicemia (A40-A41)	12	1.1	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	1.0	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	8	0.8	*
13	Anemias (D50-D64)	6	0.6	*
14	Meningitis (G00,G03)	3	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.3	*
...	All other causes (Residual)	248	23.6	9.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 5-9 years				
...	All causes	543	100.0	16.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	162	29.8	4.9
2	Malignant neoplasms (C00-C97)	68	12.5	2.1
3	Chronic lower respiratory diseases (J40-J47)	37	6.8	1.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	36	6.6	1.1
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	35	6.4	1.1
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	22	4.1	0.7
7	Anemias (D50-D64)	15	2.8	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	2.2	*
8	Cerebrovascular diseases (I60-I69)	12	2.2	*
10	Influenza and pneumonia (J09-J18)	11	2.0	*
11	Septicemia (A40-A41)	7	1.3	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.9	*
13	Certain conditions originating in the perinatal period (P00-P96)	4	0.7	*
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3	0.6	*
15	Meningitis (G00,G03)	2	0.4	*
15	Pneumonitis due to solids and liquids (J69)	2	0.4	*
15	Diseases of appendix (K35-K38)	2	0.4	*
...	All other causes (Residual)	108	19.9	3.3
Black, both sexes, 10-14 years				
...	All causes	638	100.0	18.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	159	24.9	4.7
2	Malignant neoplasms (C00-C97)	65	10.2	1.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	54	8.5	1.6
4	Chronic lower respiratory diseases (J40-J47)	47	7.4	1.4
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	45	7.1	1.3
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	42	6.6	1.2
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	28	4.4	0.8
8	Influenza and pneumonia (J09-J18)	15	2.4	*
9	Cerebrovascular diseases (I60-I69)	8	1.3	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	1.1	*
10	Anemias (D50-D64)	7	1.1	*
12	Septicemia (A40-A41)	5	0.8	*
13	Diabetes mellitus (E10-E14)	4	0.6	*
13	Pneumonitis due to solids and liquids (J69)	4	0.6	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.6	*
...	All other causes (Residual)	144	22.6	4.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 15-19 years				
...	All causes	2,380	100.0	64.9
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,026	43.1	28.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	551	23.2	15.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	165	6.9	4.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	101	4.2	2.8
5	Malignant neoplasms (C00-C97)	96	4.0	2.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	55	2.3	1.5
7	Chronic lower respiratory diseases (J40-J47)	36	1.5	1.0
8	Anemias (D50-D64)	18	0.8	*
9	Legal intervention (Y35,Y89.0)	16	0.7	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	13	0.5	*
11	Pregnancy, childbirth and the puerperium (O00-O99)	11	0.5	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	0.4	*
12	Diabetes mellitus (E10-E14)	10	0.4	*
14	Cerebrovascular diseases (I60-I69)	9	0.4	*
14	Influenza and pneumonia (J09-J18)	9	0.4	*
...	All other causes (Residual)	254	10.7	6.9
Black, both sexes, 20-24 years				
...	All causes	4,310	100.0	120.3
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,800	41.8	50.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	836	19.4	23.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	339	7.9	9.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	209	4.8	5.8
5	Malignant neoplasms (C00-C97)	164	3.8	4.6
6	Human immunodeficiency virus (HIV) disease (B20-B24)	90	2.1	2.5
7	Anemias (D50-D64)	55	1.3	1.5
8	Pregnancy, childbirth and the puerperium (O00-O99)	53	1.2	1.5
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	49	1.1	1.4
10	Chronic lower respiratory diseases (J40-J47)	45	1.0	1.3
11	Diabetes mellitus (E10-E14)	38	0.9	1.1
12	Influenza and pneumonia (J09-J18)	35	0.8	1.0
13	Legal intervention (Y35,Y89.0)	31	0.7	0.9
14	Septicemia (A40-A41)	29	0.7	0.8
15	Cerebrovascular diseases (I60-I69)	23	0.5	0.6
...	All other causes (Residual)	514	11.9	14.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 25-29 years				
...	All causes	4,300	100.0	140.1
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,363	31.7	44.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	846	19.7	27.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	334	7.8	10.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	305	7.1	9.9
5	Malignant neoplasms (C00-C97)	235	5.5	7.7
6	Human immunodeficiency virus (HIV) disease (B20-B24)	166	3.9	5.4
7	Diabetes mellitus (E10-E14)	80	1.9	2.6
8	Anemias (D50-D64)	63	1.5	2.1
9	Pregnancy, childbirth and the puerperium (O00-O99)	62	1.4	2.0
10	Influenza and pneumonia (J09-J18)	48	1.1	1.6
11	Septicemia (A40-A41)	44	1.0	1.4
12	Chronic lower respiratory diseases (J40-J47)	43	1.0	1.4
13	Cerebrovascular diseases (I60-I69)	40	0.9	1.3
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	39	0.9	1.3
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	30	0.7	1.0
...	All other causes (Residual)	602	14.0	19.6
Black, both sexes, 30-34 years				
...	All causes	4,768	100.0	162.7
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	967	20.3	33.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	811	17.0	27.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	623	13.1	21.3
4	Malignant neoplasms (C00-C97)	411	8.6	14.0
5	Human immunodeficiency virus (HIV) disease (B20-B24)	252	5.3	8.6
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	251	5.3	8.6
7	Diabetes mellitus (E10-E14)	148	3.1	5.1
8	Cerebrovascular diseases (I60-I69)	95	2.0	3.2
9	Anemias (D50-D64)	70	1.5	2.4
10	Pregnancy, childbirth and the puerperium (O00-O99)	64	1.3	2.2
11	Influenza and pneumonia (J09-J18)	58	1.2	2.0
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	55	1.2	1.9
13	Septicemia (A40-A41)	53	1.1	1.8
14	Chronic lower respiratory diseases (J40-J47)	48	1.0	1.6
15	Chronic liver disease and cirrhosis (K70,K73-K74)	37	0.8	1.3
...	All other causes (Residual)	825	17.3	28.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 35-39 years				
...	All causes	5,617	100.0	206.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	980	17.4	36.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	782	13.9	28.7
3	Malignant neoplasms (C00-C97)	735	13.1	27.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	656	11.7	24.1
5	Human immunodeficiency virus (HIV) disease (B20-B24)	353	6.3	13.0
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	227	4.0	8.3
7	Cerebrovascular diseases (I60-I69)	182	3.2	6.7
8	Diabetes mellitus (E10-E14)	162	2.9	6.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	106	1.9	3.9
10	Septicemia (A40-A41)	85	1.5	3.1
11	Chronic lower respiratory diseases (J40-J47)	74	1.3	2.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	71	1.3	2.6
13	Influenza and pneumonia (J09-J18)	62	1.1	2.3
14	Chronic liver disease and cirrhosis (K70,K73-K74)	56	1.0	2.1
15	Pregnancy, childbirth and the puerperium (O00-O99)	54	1.0	2.0
...	All other causes (Residual)	1,032	18.4	37.9
Black, both sexes, 40-44 years				
...	All causes	8,172	100.0	289.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,738	21.3	61.5
2	Malignant neoplasms (C00-C97)	1,381	16.9	48.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	913	11.2	32.3
4	Human immunodeficiency virus (HIV) disease (B20-B24)	551	6.7	19.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	477	5.8	16.9
6	Cerebrovascular diseases (I60-I69)	331	4.1	11.7
7	Diabetes mellitus (E10-E14)	329	4.0	11.6
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	195	2.4	6.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	164	2.0	5.8
10	Septicemia (A40-A41)	132	1.6	4.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	125	1.5	4.4
12	Influenza and pneumonia (J09-J18)	121	1.5	4.3
13	Chronic lower respiratory diseases (J40-J47)	118	1.4	4.2
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	116	1.4	4.1
15	Anemias (D50-D64)	58	0.7	2.1
...	All other causes (Residual)	1,423	17.4	50.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 45-49 years				
...	All causes	13,183	100.0	449.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,965	22.5	101.2
2	Malignant neoplasms (C00-C97)	2,887	21.9	98.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,133	8.6	38.7
4	Human immunodeficiency virus (HIV) disease (B20-B24)	694	5.3	23.7
5	Cerebrovascular diseases (I60-I69)	659	5.0	22.5
6	Diabetes mellitus (E10-E14)	545	4.1	18.6
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	427	3.2	14.6
8	Chronic liver disease and cirrhosis (K70,K73-K74)	282	2.1	9.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	269	2.0	9.2
10	Septicemia (A40-A41)	237	1.8	8.1
11	Chronic lower respiratory diseases (J40-J47)	219	1.7	7.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	208	1.6	7.1
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	201	1.5	6.9
14	Influenza and pneumonia (J09-J18)	153	1.2	5.2
15	Anemias (D50-D64)	97	0.7	3.3
...	All other causes (Residual)	2,207	16.7	75.3
Black, both sexes, 50-54 years				
...	All causes	20,348	100.0	713.3
1	Malignant neoplasms (C00-C97)	5,511	27.1	193.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,846	23.8	169.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,277	6.3	44.8
4	Cerebrovascular diseases (I60-I69)	985	4.8	34.5
5	Diabetes mellitus (E10-E14)	905	4.4	31.7
6	Human immunodeficiency virus (HIV) disease (B20-B24)	760	3.7	26.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	520	2.6	18.2
8	Chronic lower respiratory diseases (J40-J47)	480	2.4	16.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	458	2.3	16.1
10	Septicemia (A40-A41)	394	1.9	13.8
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	309	1.5	10.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	295	1.4	10.3
13	Influenza and pneumonia (J09-J18)	250	1.2	8.8
14	Viral hepatitis (B15-B19)	200	1.0	7.0
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	155	0.8	5.4
...	All other causes (Residual)	3,003	14.8	105.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 55-59 years				
...	All causes	25,447	100.0	1,066.4
1	Malignant neoplasms (C00-C97)	7,668	30.1	321.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,133	24.1	257.0
3	Cerebrovascular diseases (I60-I69)	1,273	5.0	53.3
4	Diabetes mellitus (E10-E14)	1,238	4.9	51.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,207	4.7	50.6
6	Chronic lower respiratory diseases (J40-J47)	647	2.5	27.1
7	Human immunodeficiency virus (HIV) disease (B20-B24)	620	2.4	26.0
8	Chronic liver disease and cirrhosis (K70,K73-K74)	616	2.4	25.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	591	2.3	24.8
10	Septicemia (A40-A41)	549	2.2	23.0
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	398	1.6	16.7
12	Viral hepatitis (B15-B19)	329	1.3	13.8
13	Influenza and pneumonia (J09-J18)	327	1.3	13.7
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	218	0.9	9.1
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	120	0.5	5.0
...	All other causes (Residual)	3,513	13.8	147.2
Black, both sexes, 60-64 years				
...	All causes	28,352	100.0	1,499.0
1	Malignant neoplasms (C00-C97)	9,282	32.7	490.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,020	24.8	371.2
3	Diabetes mellitus (E10-E14)	1,533	5.4	81.1
4	Cerebrovascular diseases (I60-I69)	1,435	5.1	75.9
5	Chronic lower respiratory diseases (J40-J47)	840	3.0	44.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	831	2.9	43.9
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	822	2.9	43.5
8	Septicemia (A40-A41)	586	2.1	31.0
9	Chronic liver disease and cirrhosis (K70,K73-K74)	544	1.9	28.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	454	1.6	24.0
11	Human immunodeficiency virus (HIV) disease (B20-B24)	380	1.3	20.1
11	Influenza and pneumonia (J09-J18)	380	1.3	20.1
13	Viral hepatitis (B15-B19)	272	1.0	14.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	115	0.4	6.1
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	115	0.4	6.1
...	All other causes (Residual)	3,743	13.2	197.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 65-69 years				
...	All causes	26,151	100.0	2,096.2
1	Malignant neoplasms (C00-C97)	8,607	32.9	689.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,519	24.9	522.5
3	Diabetes mellitus (E10-E14)	1,521	5.8	121.9
4	Cerebrovascular diseases (I60-I69)	1,372	5.2	110.0
5	Chronic lower respiratory diseases (J40-J47)	1,012	3.9	81.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	826	3.2	66.2
7	Septicemia (A40-A41)	582	2.2	46.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	547	2.1	43.8
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	440	1.7	35.3
10	Influenza and pneumonia (J09-J18)	383	1.5	30.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	257	1.0	20.6
12	Human immunodeficiency virus (HIV) disease (B20-B24)	177	0.7	14.2
13	Pneumonitis due to solids and liquids (J69)	126	0.5	10.1
14	Alzheimer's disease (G30)	118	0.5	9.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	99	0.4	7.9
...	All other causes (Residual)	3,565	13.6	285.8
Black, both sexes, 70-74 years				
...	All causes	27,131	100.0	2,966.4
1	Malignant neoplasms (C00-C97)	8,388	30.9	917.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,825	25.2	746.2
3	Cerebrovascular diseases (I60-I69)	1,662	6.1	181.7
4	Diabetes mellitus (E10-E14)	1,488	5.5	162.7
5	Chronic lower respiratory diseases (J40-J47)	1,210	4.5	132.3
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	933	3.4	102.0
7	Septicemia (A40-A41)	667	2.5	72.9
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	470	1.7	51.4
8	Influenza and pneumonia (J09-J18)	470	1.7	51.4
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	408	1.5	44.6
11	Alzheimer's disease (G30)	289	1.1	31.6
12	Chronic liver disease and cirrhosis (K70,K73-K74)	150	0.6	16.4
13	Pneumonitis due to solids and liquids (J69)	131	0.5	14.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	126	0.5	13.8
14	Parkinson's disease (G20-G21)	126	0.5	13.8
...	All other causes (Residual)	3,788	14.0	414.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 75-79 years				
...	All causes	28,615	100.0	4,304.8
1	Malignant neoplasms (C00-C97)	7,657	26.8	1,151.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,277	25.4	1,094.7
3	Cerebrovascular diseases (I60-I69)	1,906	6.7	286.7
4	Diabetes mellitus (E10-E14)	1,515	5.3	227.9
5	Chronic lower respiratory diseases (J40-J47)	1,284	4.5	193.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,044	3.6	157.1
7	Septicemia (A40-A41)	720	2.5	108.3
8	Alzheimer's disease (G30)	690	2.4	103.8
9	Influenza and pneumonia (J09-J18)	580	2.0	87.3
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	520	1.8	78.2
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	403	1.4	60.6
12	Pneumonitis due to solids and liquids (J69)	190	0.7	28.6
13	Parkinson's disease (G20-G21)	176	0.6	26.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	153	0.5	23.0
15	Chronic liver disease and cirrhosis (K70,K73-K74)	105	0.4	15.8
...	All other causes (Residual)	4,395	15.4	661.2
Black, both sexes, 80-84 years				
...	All causes	29,339	100.0	6,572.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,758	26.4	1,737.9
2	Malignant neoplasms (C00-C97)	6,452	22.0	1,445.4
3	Cerebrovascular diseases (I60-I69)	2,071	7.1	463.9
4	Diabetes mellitus (E10-E14)	1,424	4.9	319.0
5	Chronic lower respiratory diseases (J40-J47)	1,233	4.2	276.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,038	3.5	232.5
7	Alzheimer's disease (G30)	1,032	3.5	231.2
8	Septicemia (A40-A41)	685	2.3	153.5
9	Influenza and pneumonia (J09-J18)	655	2.2	146.7
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	588	2.0	131.7
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	367	1.3	82.2
12	Pneumonitis due to solids and liquids (J69)	220	0.7	49.3
13	Parkinson's disease (G20-G21)	211	0.7	47.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	182	0.6	40.8
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	96	0.3	21.5
...	All other causes (Residual)	5,327	18.2	1,193.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 85-89 years				
...	All causes	26,234	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,206	27.5	---
2	Malignant neoplasms (C00-C97)	4,441	16.9	---
3	Cerebrovascular diseases (I60-I69)	1,855	7.1	---
4	Alzheimer's disease (G30)	1,393	5.3	---
5	Diabetes mellitus (E10-E14)	1,106	4.2	---
6	Chronic lower respiratory diseases (J40-J47)	994	3.8	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	883	3.4	---
8	Influenza and pneumonia (J09-J18)	711	2.7	---
9	Septicemia (A40-A41)	640	2.4	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	599	2.3	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	302	1.2	---
12	Pneumonitis due to solids and liquids (J69)	246	0.9	---
13	Parkinson's disease (G20-G21)	170	0.6	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	132	0.5	---
15	Atherosclerosis (I70)	109	0.4	---
...	All other causes (Residual)	5,447	20.8	---
Black, both sexes, 90-94 years				
...	All causes	16,725	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,776	28.6	---
2	Malignant neoplasms (C00-C97)	1,997	11.9	---
3	Cerebrovascular diseases (I60-I69)	1,245	7.4	---
4	Alzheimer's disease (G30)	1,164	7.0	---
5	Diabetes mellitus (E10-E14)	590	3.5	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	541	3.2	---
7	Chronic lower respiratory diseases (J40-J47)	477	2.9	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	463	2.8	---
9	Influenza and pneumonia (J09-J18)	462	2.8	---
10	Septicemia (A40-A41)	351	2.1	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	213	1.3	---
12	Pneumonitis due to solids and liquids (J69)	175	1.0	---
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	83	0.5	---
14	Parkinson's disease (G20-G21)	72	0.4	---
14	Atherosclerosis (I70)	72	0.4	---
...	All other causes (Residual)	4,044	24.2	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 95-99 years				
...	All causes	7,288	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,163	29.7	---
2	Malignant neoplasms (C00-C97)	598	8.2	---
3	Alzheimer's disease (G30)	561	7.7	---
4	Cerebrovascular diseases (I60-I69)	487	6.7	---
5	Essential hypertension and hypertensive renal disease (I10,I12,I15)	240	3.3	---
6	Influenza and pneumonia (J09-J18)	236	3.2	---
7	Diabetes mellitus (E10-E14)	218	3.0	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	195	2.7	---
9	Chronic lower respiratory diseases (J40-J47)	162	2.2	---
10	Septicemia (A40-A41)	142	1.9	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	84	1.2	---
12	Pneumonitis due to solids and liquids (J69)	64	0.9	---
13	Atherosclerosis (I70)	39	0.5	---
14	Anemias (D50-D64)	26	0.4	---
15	Nutritional deficiencies (E40-E64)	25	0.3	---
15	Parkinson's disease (G20-G21)	25	0.3	---
...	All other causes (Residual)	2,023	27.8	---
Black, both sexes, 100 years and over				
...	All causes	2,196	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	690	31.4	---
2	Alzheimer's disease (G30)	158	7.2	---
3	Cerebrovascular diseases (I60-I69)	137	6.2	---
4	Malignant neoplasms (C00-C97)	106	4.8	---
5	Influenza and pneumonia (J09-J18)	86	3.9	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	84	3.8	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	50	2.3	---
8	Chronic lower respiratory diseases (J40-J47)	43	2.0	---
9	Diabetes mellitus (E10-E14)	41	1.9	---
10	Septicemia (A40-A41)	36	1.6	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	28	1.3	---
12	Atherosclerosis (I70)	23	1.0	---
13	Pneumonitis due to solids and liquids (J69)	22	1.0	---
14	Anemias (D50-D64)	12	0.5	---
15	Nutritional deficiencies (E40-E64)	9	0.4	---
...	All other causes (Residual)	671	30.6	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, all ages				
...	All causes	146,884	100.0	719.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	34,913	23.8	171.0
2	Malignant neoplasms (C00-C97)	33,978	23.1	166.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,387	5.7	41.1
4	Cerebrovascular diseases (I60-I69)	7,039	4.8	34.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6,739	4.6	33.0
6	Diabetes mellitus (E10-E14)	6,048	4.1	29.6
7	Chronic lower respiratory diseases (J40-J47)	4,702	3.2	23.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,756	2.6	18.4
9	Human immunodeficiency virus (HIV) disease (B20-B24)	2,744	1.9	13.4
10	Septicemia (A40-A41)	2,700	1.8	13.2
11	Influenza and pneumonia (J09-J18)	2,417	1.6	11.8
12	Certain conditions originating in the perinatal period (P00-P96)	2,252	1.5	11.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,200	1.5	10.8
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,828	1.2	9.0
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,775	1.2	8.7
...	All other causes (Residual)	25,406	17.3	124.4
Black, male, 1-4 years				
...	All causes	596	100.0	42.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	173	29.0	12.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	87	14.6	6.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	59	9.9	4.2
4	Malignant neoplasms (C00-C97)	31	5.2	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	28	4.7	2.0
6	Influenza and pneumonia (J09-J18)	18	3.0	*
7	Chronic lower respiratory diseases (J40-J47)	16	2.7	*
8	Cerebrovascular diseases (I60-I69)	14	2.3	*
9	Certain conditions originating in the perinatal period (P00-P96)	9	1.5	*
10	Septicemia (A40-A41)	6	1.0	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.0	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	6	1.0	*
13	Anemias (D50-D64)	3	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.3	*
15	Meningitis (G00,G03)	1	0.2	*
15	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
15	Pneumonitis due to solids and liquids (J69)	1	0.2	*
...	All other causes (Residual)	135	22.7	9.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 5-9 years				
...	All causes	304	100.0	18.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	97	31.9	5.8
2	Malignant neoplasms (C00-C97)	39	12.8	2.3
3	Chronic lower respiratory diseases (J40-J47)	28	9.2	1.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	21	6.9	1.3
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	19	6.3	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	13	4.3	*
7	Anemias (D50-D64)	9	3.0	*
8	Influenza and pneumonia (J09-J18)	7	2.3	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	2.0	*
10	Cerebrovascular diseases (I60-I69)	5	1.6	*
11	Certain conditions originating in the perinatal period (P00-P96)	4	1.3	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	1.0	*
13	Septicemia (A40-A41)	2	0.7	*
13	Pneumonitis due to solids and liquids (J69)	2	0.7	*
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2	0.7	*
...	All other causes (Residual)	47	15.5	2.8
Black, male, 10-14 years				
...	All causes	397	100.0	22.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	105	26.4	6.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	41	10.3	2.4
3	Malignant neoplasms (C00-C97)	38	9.6	2.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	32	8.1	1.8
5	Chronic lower respiratory diseases (J40-J47)	29	7.3	1.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	25	6.3	1.4
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	13	3.3	*
8	Influenza and pneumonia (J09-J18)	9	2.3	*
9	Cerebrovascular diseases (I60-I69)	6	1.5	*
10	Anemias (D50-D64)	5	1.3	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	1.0	*
12	Diabetes mellitus (E10-E14)	3	0.8	*
12	Pneumonitis due to solids and liquids (J69)	3	0.8	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.8	*
15	Septicemia (A40-A41)	2	0.5	*
...	All other causes (Residual)	79	19.9	4.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 15-19 years				
...	All causes	1,813	100.0	97.1
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	907	50.0	48.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	410	22.6	21.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	129	7.1	6.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	73	4.0	3.9
5	Malignant neoplasms (C00-C97)	57	3.1	3.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	32	1.8	1.7
7	Chronic lower respiratory diseases (J40-J47)	26	1.4	1.4
8	Legal intervention (Y35,Y89.0)	15	0.8	*
9	Anemias (D50-D64)	8	0.4	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.3	*
10	Diabetes mellitus (E10-E14)	5	0.3	*
10	Influenza and pneumonia (J09-J18)	5	0.3	*
13	Septicemia (A40-A41)	4	0.2	*
13	Cerebrovascular diseases (I60-I69)	4	0.2	*
13	Aortic aneurysm and dissection (I71)	4	0.2	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.2	*
...	All other causes (Residual)	125	6.9	6.7
Black, male, 20-24 years				
...	All causes	3,311	100.0	185.6
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,629	49.2	91.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	627	18.9	35.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	292	8.8	16.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	127	3.8	7.1
5	Malignant neoplasms (C00-C97)	94	2.8	5.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	61	1.8	3.4
7	Anemias (D50-D64)	37	1.1	2.1
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	32	1.0	1.8
9	Legal intervention (Y35,Y89.0)	31	0.9	1.7
10	Chronic lower respiratory diseases (J40-J47)	28	0.8	1.6
11	Diabetes mellitus (E10-E14)	18	0.5	*
12	Influenza and pneumonia (J09-J18)	15	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	0.4	*
14	Cerebrovascular diseases (I60-I69)	12	0.4	*
15	Septicemia (A40-A41)	9	0.3	*
...	All other causes (Residual)	286	8.6	16.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 25-29 years				
...	All causes	3,079	100.0	207.4
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,216	39.5	81.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	629	20.4	42.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	255	8.3	17.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	208	6.8	14.0
5	Human immunodeficiency virus (HIV) disease (B20-B24)	112	3.6	7.5
6	Malignant neoplasms (C00-C97)	111	3.6	7.5
7	Anemias (D50-D64)	38	1.2	2.6
8	Diabetes mellitus (E10-E14)	34	1.1	2.3
9	Chronic lower respiratory diseases (J40-J47)	32	1.0	2.2
10	Influenza and pneumonia (J09-J18)	26	0.8	1.8
11	Cerebrovascular diseases (I60-I69)	23	0.7	1.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	0.7	1.5
13	Septicemia (A40-A41)	18	0.6	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	17	0.6	*
15	Legal intervention (Y35,Y89.0)	16	0.5	*
...	All other causes (Residual)	321	10.4	21.6
Black, male, 30-34 years				
...	All causes	3,024	100.0	217.3
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	870	28.8	62.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	563	18.6	40.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	369	12.2	26.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	199	6.6	14.3
5	Malignant neoplasms (C00-C97)	153	5.1	11.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	133	4.4	9.6
7	Diabetes mellitus (E10-E14)	79	2.6	5.7
8	Cerebrovascular diseases (I60-I69)	44	1.5	3.2
9	Anemias (D50-D64)	37	1.2	2.7
10	Influenza and pneumonia (J09-J18)	29	1.0	2.1
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	29	1.0	2.1
12	Chronic lower respiratory diseases (J40-J47)	26	0.9	1.9
13	Chronic liver disease and cirrhosis (K70,K73-K74)	24	0.8	1.7
14	Septicemia (A40-A41)	22	0.7	1.6
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	18	0.6	*
...	All other causes (Residual)	429	14.2	30.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 35-39 years				
...	All causes	3,323	100.0	259.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	597	18.0	46.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	564	17.0	44.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	556	16.7	43.5
4	Malignant neoplasms (C00-C97)	271	8.2	21.2
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	194	5.8	15.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	184	5.5	14.4
7	Diabetes mellitus (E10-E14)	87	2.6	6.8
8	Cerebrovascular diseases (I60-I69)	86	2.6	6.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	55	1.7	4.3
10	Septicemia (A40-A41)	49	1.5	3.8
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	42	1.3	3.3
12	Influenza and pneumonia (J09-J18)	37	1.1	2.9
13	Chronic lower respiratory diseases (J40-J47)	35	1.1	2.7
14	Anemias (D50-D64)	28	0.8	2.2
15	Aortic aneurysm and dissection (I71)	26	0.8	2.0
15	Chronic liver disease and cirrhosis (K70,K73-K74)	26	0.8	2.0
...	All other causes (Residual)	486	14.6	38.0
Black, male, 40-44 years				
...	All causes	4,639	100.0	348.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,137	24.5	85.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	638	13.8	47.9
3	Malignant neoplasms (C00-C97)	483	10.4	36.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	397	8.6	29.8
5	Human immunodeficiency virus (HIV) disease (B20-B24)	326	7.0	24.5
6	Diabetes mellitus (E10-E14)	189	4.1	14.2
7	Cerebrovascular diseases (I60-I69)	180	3.9	13.5
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	144	3.1	10.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	94	2.0	7.1
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	70	1.5	5.3
11	Septicemia (A40-A41)	67	1.4	5.0
11	Chronic liver disease and cirrhosis (K70,K73-K74)	67	1.4	5.0
13	Influenza and pneumonia (J09-J18)	58	1.3	4.4
14	Chronic lower respiratory diseases (J40-J47)	52	1.1	3.9
15	Aortic aneurysm and dissection (I71)	41	0.9	3.1
...	All other causes (Residual)	696	15.0	52.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 45-49 years				
...	All causes	7,404	100.0	536.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,900	25.7	137.6
2	Malignant neoplasms (C00-C97)	1,217	16.4	88.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	774	10.5	56.0
4	Human immunodeficiency virus (HIV) disease (B20-B24)	453	6.1	32.8
5	Cerebrovascular diseases (I60-I69)	356	4.8	25.8
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	332	4.5	24.0
7	Diabetes mellitus (E10-E14)	305	4.1	22.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	165	2.2	11.9
9	Chronic liver disease and cirrhosis (K70,K73-K74)	158	2.1	11.4
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	140	1.9	10.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	124	1.7	9.0
12	Septicemia (A40-A41)	108	1.5	7.8
13	Chronic lower respiratory diseases (J40-J47)	93	1.3	6.7
14	Influenza and pneumonia (J09-J18)	79	1.1	5.7
15	Viral hepatitis (B15-B19)	50	0.7	3.6
...	All other causes (Residual)	1,150	15.5	83.3
Black, male, 50-54 years				
...	All causes	11,789	100.0	882.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,102	26.3	232.2
2	Malignant neoplasms (C00-C97)	2,739	23.2	205.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	877	7.4	65.6
4	Cerebrovascular diseases (I60-I69)	564	4.8	42.2
5	Human immunodeficiency virus (HIV) disease (B20-B24)	526	4.5	39.4
6	Diabetes mellitus (E10-E14)	496	4.2	37.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	330	2.8	24.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	260	2.2	19.5
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	245	2.1	18.3
10	Chronic lower respiratory diseases (J40-J47)	224	1.9	16.8
11	Septicemia (A40-A41)	213	1.8	15.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	179	1.5	13.4
13	Influenza and pneumonia (J09-J18)	141	1.2	10.6
14	Viral hepatitis (B15-B19)	131	1.1	9.8
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	125	1.1	9.4
...	All other causes (Residual)	1,637	13.9	122.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 55-59 years				
...	All causes	14,971	100.0	1,365.2
1	Malignant neoplasms (C00-C97)	4,121	27.5	375.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,853	25.7	351.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	852	5.7	77.7
4	Cerebrovascular diseases (I60-I69)	753	5.0	68.7
5	Diabetes mellitus (E10-E14)	715	4.8	65.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	432	2.9	39.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	413	2.8	37.7
8	Chronic lower respiratory diseases (J40-J47)	339	2.3	30.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	336	2.2	30.6
10	Septicemia (A40-A41)	286	1.9	26.1
11	Viral hepatitis (B15-B19)	227	1.5	20.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	221	1.5	20.2
13	Influenza and pneumonia (J09-J18)	189	1.3	17.2
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	179	1.2	16.3
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	92	0.6	8.4
...	All other causes (Residual)	1,963	13.1	179.0
Black, male, 60-64 years				
...	All causes	16,628	100.0	1,956.6
1	Malignant neoplasms (C00-C97)	5,191	31.2	610.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,423	26.6	520.4
3	Diabetes mellitus (E10-E14)	833	5.0	98.0
4	Cerebrovascular diseases (I60-I69)	831	5.0	97.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	590	3.5	69.4
6	Chronic lower respiratory diseases (J40-J47)	476	2.9	56.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	451	2.7	53.1
8	Chronic liver disease and cirrhosis (K70,K73-K74)	382	2.3	44.9
9	Septicemia (A40-A41)	331	2.0	38.9
10	Human immunodeficiency virus (HIV) disease (B20-B24)	278	1.7	32.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	261	1.6	30.7
12	Influenza and pneumonia (J09-J18)	210	1.3	24.7
13	Viral hepatitis (B15-B19)	184	1.1	21.7
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	100	0.6	11.8
15	Pneumonitis due to solids and liquids (J69)	67	0.4	7.9
...	All other causes (Residual)	2,020	12.1	237.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 65-69 years				
...	All causes	14,525	100.0	2,675.3
1	Malignant neoplasms (C00-C97)	4,710	32.4	867.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,858	26.6	710.6
3	Cerebrovascular diseases (I60-I69)	759	5.2	139.8
4	Diabetes mellitus (E10-E14)	750	5.2	138.1
5	Chronic lower respiratory diseases (J40-J47)	584	4.0	107.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	435	3.0	80.1
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	364	2.5	67.0
8	Septicemia (A40-A41)	281	1.9	51.8
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	250	1.7	46.0
10	Influenza and pneumonia (J09-J18)	201	1.4	37.0
11	Chronic liver disease and cirrhosis (K70,K73-K74)	166	1.1	30.6
12	Human immunodeficiency virus (HIV) disease (B20-B24)	124	0.9	22.8
13	Pneumonitis due to solids and liquids (J69)	72	0.5	13.3
14	Viral hepatitis (B15-B19)	58	0.4	10.7
14	Alzheimer's disease (G30)	58	0.4	10.7
...	All other causes (Residual)	1,855	12.8	341.7
Black, male, 70-74 years				
...	All causes	14,383	100.0	3,770.3
1	Malignant neoplasms (C00-C97)	4,572	31.8	1,198.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,773	26.2	989.1
3	Cerebrovascular diseases (I60-I69)	831	5.8	217.8
4	Diabetes mellitus (E10-E14)	744	5.2	195.0
5	Chronic lower respiratory diseases (J40-J47)	676	4.7	177.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	459	3.2	120.3
7	Septicemia (A40-A41)	305	2.1	80.0
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	251	1.7	65.8
9	Influenza and pneumonia (J09-J18)	241	1.7	63.2
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	225	1.6	59.0
11	Alzheimer's disease (G30)	127	0.9	33.3
12	Chronic liver disease and cirrhosis (K70,K73-K74)	97	0.7	25.4
13	Parkinson's disease (G20-G21)	84	0.6	22.0
14	Pneumonitis due to solids and liquids (J69)	78	0.5	20.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	57	0.4	14.9
...	All other causes (Residual)	1,863	13.0	488.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 75-79 years				
...	All causes	13,863	100.0	5,353.6
1	Malignant neoplasms (C00-C97)	3,950	28.5	1,525.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,622	26.1	1,398.7
3	Cerebrovascular diseases (I60-I69)	850	6.1	328.3
4	Chronic lower respiratory diseases (J40-J47)	698	5.0	269.6
5	Diabetes mellitus (E10-E14)	644	4.6	248.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	470	3.4	181.5
7	Septicemia (A40-A41)	310	2.2	119.7
8	Alzheimer's disease (G30)	279	2.0	107.7
8	Influenza and pneumonia (J09-J18)	279	2.0	107.7
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	244	1.8	94.2
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	233	1.7	90.0
12	Pneumonitis due to solids and liquids (J69)	104	0.8	40.2
13	Parkinson's disease (G20-G21)	101	0.7	39.0
14	Chronic liver disease and cirrhosis (K70,K73-K74)	57	0.4	22.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	55	0.4	21.2
...	All other causes (Residual)	1,967	14.2	759.6
Black, male, 80-84 years				
...	All causes	12,504	100.0	8,030.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,345	26.8	2,148.4
2	Malignant neoplasms (C00-C97)	3,122	25.0	2,005.2
3	Cerebrovascular diseases (I60-I69)	745	6.0	478.5
4	Chronic lower respiratory diseases (J40-J47)	613	4.9	393.7
5	Diabetes mellitus (E10-E14)	538	4.3	345.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	407	3.3	261.4
7	Alzheimer's disease (G30)	342	2.7	219.7
8	Influenza and pneumonia (J09-J18)	307	2.5	197.2
9	Septicemia (A40-A41)	282	2.3	181.1
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	221	1.8	141.9
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	172	1.4	110.5
12	Parkinson's disease (G20-G21)	121	1.0	77.7
13	Pneumonitis due to solids and liquids (J69)	118	0.9	75.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	78	0.6	50.1
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	35	0.3	22.5
...	All other causes (Residual)	2,058	16.5	1,321.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 85-89 years				
...	All causes	9,625	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,558	26.6	---
2	Malignant neoplasms (C00-C97)	2,048	21.3	---
3	Cerebrovascular diseases (I60-I69)	578	6.0	---
4	Chronic lower respiratory diseases (J40-J47)	458	4.8	---
5	Diabetes mellitus (E10-E14)	401	4.2	---
6	Alzheimer's disease (G30)	378	3.9	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	326	3.4	---
8	Influenza and pneumonia (J09-J18)	294	3.1	---
9	Septicemia (A40-A41)	221	2.3	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	178	1.8	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	143	1.5	---
12	Pneumonitis due to solids and liquids (J69)	117	1.2	---
13	Parkinson's disease (G20-G21)	81	0.8	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	51	0.5	---
15	Atherosclerosis (I70)	42	0.4	---
...	All other causes (Residual)	1,751	18.2	---
Black, male, 90-94 years				
...	All causes	4,818	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,323	27.5	---
2	Malignant neoplasms (C00-C97)	799	16.6	---
3	Cerebrovascular diseases (I60-I69)	282	5.9	---
4	Alzheimer's disease (G30)	250	5.2	---
5	Chronic lower respiratory diseases (J40-J47)	192	4.0	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	167	3.5	---
7	Diabetes mellitus (E10-E14)	159	3.3	---
8	Influenza and pneumonia (J09-J18)	148	3.1	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	104	2.2	---
10	Septicemia (A40-A41)	101	2.1	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	82	1.7	---
12	Pneumonitis due to solids and liquids (J69)	70	1.5	---
13	Parkinson's disease (G20-G21)	34	0.7	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	0.4	---
15	Atherosclerosis (I70)	16	0.3	---
...	All other causes (Residual)	1,071	22.2	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 95-99 years				
...	All causes	1,502	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	441	29.4	---
2	Malignant neoplasms (C00-C97)	205	13.6	---
3	Alzheimer's disease (G30)	79	5.3	---
4	Cerebrovascular diseases (I60-I69)	78	5.2	---
5	Influenza and pneumonia (J09-J18)	61	4.1	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	52	3.5	---
7	Chronic lower respiratory diseases (J40-J47)	49	3.3	---
8	Diabetes mellitus (E10-E14)	43	2.9	---
9	Septicemia (A40-A41)	37	2.5	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	37	2.5	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	24	1.6	---
12	Parkinson's disease (G20-G21)	12	0.8	---
12	Pneumonitis due to solids and liquids (J69)	12	0.8	---
14	Atherosclerosis (I70)	11	0.7	---
15	Hyperplasia of prostate (N40)	7	0.5	---
...	All other causes (Residual)	354	23.6	---
Black, male, 100 years and over				
...	All causes	309	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	94	30.4	---
2	Malignant neoplasms (C00-C97)	19	6.1	---
2	Chronic lower respiratory diseases (J40-J47)	19	6.1	---
4	Influenza and pneumonia (J09-J18)	18	5.8	---
5	Alzheimer's disease (G30)	16	5.2	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	4.9	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	12	3.9	---
8	Cerebrovascular diseases (I60-I69)	11	3.6	---
9	Septicemia (A40-A41)	6	1.9	---
10	Diabetes mellitus (E10-E14)	5	1.6	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5	1.6	---
12	Pneumonitis due to solids and liquids (J69)	4	1.3	---
13	Atherosclerosis (I70)	3	1.0	---
14	Anemias (D50-D64)	2	0.6	---
14	Nutritional deficiencies (E40-E64)	2	0.6	---
14	Parkinson's disease (G20-G21)	2	0.6	---
...	All other causes (Residual)	76	24.6	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, all ages				
...	All causes	143,216	100.0	643.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	33,459	23.4	150.3
2	Malignant neoplasms (C00-C97)	32,842	22.9	147.5
3	Cerebrovascular diseases (I60-I69)	8,814	6.2	39.6
4	Diabetes mellitus (E10-E14)	6,847	4.8	30.8
5	Chronic lower respiratory diseases (J40-J47)	4,384	3.1	19.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,331	3.0	19.5
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,144	2.9	18.6
8	Alzheimer's disease (G30)	3,914	2.7	17.6
9	Septicemia (A40-A41)	3,328	2.3	15.0
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,806	2.0	12.6
11	Influenza and pneumonia (J09-J18)	2,750	1.9	12.4
12	Certain conditions originating in the perinatal period (P00-P96)	1,803	1.3	8.1
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,471	1.0	6.6
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,119	0.8	5.0
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,041	0.7	4.7
...	All other causes (Residual)	30,163	21.1	135.5
Black, female, 1-4 years				
...	All causes	456	100.0	33.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	125	27.4	9.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	72	15.8	5.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	49	10.7	3.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	26	5.7	1.9
5	Malignant neoplasms (C00-C97)	24	5.3	1.8
6	Influenza and pneumonia (J09-J18)	14	3.1	*
7	Chronic lower respiratory diseases (J40-J47)	9	2.0	*
8	Septicemia (A40-A41)	6	1.3	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	1.1	*
10	Certain conditions originating in the perinatal period (P00-P96)	4	0.9	*
11	Anemias (D50-D64)	3	0.7	*
11	Cerebrovascular diseases (I60-I69)	3	0.7	*
13	Meningitis (G00,G03)	2	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
15	Atherosclerosis (I70)	1	0.2	*
15	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
15	Pneumonitis due to solids and liquids (J69)	1	0.2	*
15	Peptic ulcer (K25-K28)	1	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.2	*
...	All other causes (Residual)	106	23.2	7.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 5-9 years				
...	All causes	239	100.0	14.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	65	27.2	4.0
2	Malignant neoplasms (C00-C97)	29	12.1	1.8
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	16	6.7	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	15	6.3	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	9	3.8	*
5	Chronic lower respiratory diseases (J40-J47)	9	3.8	*
7	Cerebrovascular diseases (I60-I69)	7	2.9	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	2.5	*
8	Anemias (D50-D64)	6	2.5	*
10	Septicemia (A40-A41)	5	2.1	*
11	Influenza and pneumonia (J09-J18)	4	1.7	*
12	Diseases of appendix (K35-K38)	2	0.8	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.8	*
14	Meningitis (G00,G03)	1	0.4	*
14	Acute bronchitis and bronchiolitis (J20-J21)	1	0.4	*
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.4	*
...	All other causes (Residual)	60	25.1	3.7
Black, female, 10-14 years				
...	All causes	241	100.0	14.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	54	22.4	3.2
2	Malignant neoplasms (C00-C97)	27	11.2	1.6
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	20	8.3	1.2
4	Chronic lower respiratory diseases (J40-J47)	18	7.5	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	15	6.2	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	13	5.4	*
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	10	4.1	*
8	Influenza and pneumonia (J09-J18)	6	2.5	*
9	Septicemia (A40-A41)	3	1.2	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	1.2	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	1.2	*
12	Anemias (D50-D64)	2	0.8	*
12	Meningitis (G00,G03)	2	0.8	*
12	Cerebrovascular diseases (I60-I69)	2	0.8	*
15	Diabetes mellitus (E10-E14)	1	0.4	*
15	Pneumonitis due to solids and liquids (J69)	1	0.4	*
15	Certain conditions originating in the perinatal period (P00-P96)	1	0.4	*
...	All other causes (Residual)	60	24.9	3.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 15-19 years				
...	All causes	567	100.0	31.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	141	24.9	7.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	119	21.0	6.6
3	Malignant neoplasms (C00-C97)	39	6.9	2.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	36	6.3	2.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	28	4.9	1.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	23	4.1	1.3
7	Pregnancy, childbirth and the puerperium (O00-O99)	11	1.9	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	10	1.8	*
8	Anemias (D50-D64)	10	1.8	*
8	Chronic lower respiratory diseases (J40-J47)	10	1.8	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.9	*
11	Diabetes mellitus (E10-E14)	5	0.9	*
11	Cerebrovascular diseases (I60-I69)	5	0.9	*
14	Influenza and pneumonia (J09-J18)	4	0.7	*
14	Pneumonitis due to solids and liquids (J69)	4	0.7	*
...	All other causes (Residual)	117	20.6	6.5
Black, female, 20-24 years				
...	All causes	999	100.0	55.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	209	20.9	11.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	171	17.1	9.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	82	8.2	4.6
4	Malignant neoplasms (C00-C97)	70	7.0	3.9
5	Pregnancy, childbirth and the puerperium (O00-O99)	53	5.3	2.9
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	47	4.7	2.6
7	Human immunodeficiency virus (HIV) disease (B20-B24)	29	2.9	1.6
8	Septicemia (A40-A41)	20	2.0	1.1
8	Diabetes mellitus (E10-E14)	20	2.0	1.1
8	Influenza and pneumonia (J09-J18)	20	2.0	1.1
11	Anemias (D50-D64)	18	1.8	*
12	Chronic lower respiratory diseases (J40-J47)	17	1.7	*
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	17	1.7	*
14	Cerebrovascular diseases (I60-I69)	11	1.1	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.8	*
...	All other causes (Residual)	207	20.7	11.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 25-29 years				
...	All causes	1,221	100.0	77.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	217	17.8	13.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	147	12.0	9.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	126	10.3	8.0
4	Malignant neoplasms (C00-C97)	124	10.2	7.8
5	Pregnancy, childbirth and the puerperium (O00-O99)	62	5.1	3.9
6	Human immunodeficiency virus (HIV) disease (B20-B24)	54	4.4	3.4
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	50	4.1	3.2
8	Diabetes mellitus (E10-E14)	46	3.8	2.9
9	Septicemia (A40-A41)	26	2.1	1.6
10	Anemias (D50-D64)	25	2.0	1.6
11	Influenza and pneumonia (J09-J18)	22	1.8	1.4
12	Cerebrovascular diseases (I60-I69)	17	1.4	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	1.3	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	13	1.1	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	13	1.1	*
...	All other causes (Residual)	263	21.5	16.6
Black, female, 30-34 years				
...	All causes	1,744	100.0	113.4
1	Malignant neoplasms (C00-C97)	258	14.8	16.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	254	14.6	16.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	248	14.2	16.1
4	Human immunodeficiency virus (HIV) disease (B20-B24)	119	6.8	7.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	97	5.6	6.3
6	Diabetes mellitus (E10-E14)	69	4.0	4.5
7	Pregnancy, childbirth and the puerperium (O00-O99)	64	3.7	4.2
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	52	3.0	3.4
9	Cerebrovascular diseases (I60-I69)	51	2.9	3.3
10	Anemias (D50-D64)	33	1.9	2.1
11	Septicemia (A40-A41)	31	1.8	2.0
12	Influenza and pneumonia (J09-J18)	29	1.7	1.9
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	26	1.5	1.7
14	Chronic lower respiratory diseases (J40-J47)	22	1.3	1.4
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	19	1.1	*
...	All other causes (Residual)	372	21.3	24.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 35-39 years				
...	All causes	2,294	100.0	159.3
1	Malignant neoplasms (C00-C97)	464	20.2	32.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	383	16.7	26.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	218	9.5	15.1
4	Human immunodeficiency virus (HIV) disease (B20-B24)	169	7.4	11.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	100	4.4	6.9
6	Cerebrovascular diseases (I60-I69)	96	4.2	6.7
7	Diabetes mellitus (E10-E14)	75	3.3	5.2
8	Pregnancy, childbirth and the puerperium (O00-O99)	54	2.4	3.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	51	2.2	3.5
10	Chronic lower respiratory diseases (J40-J47)	39	1.7	2.7
11	Septicemia (A40-A41)	36	1.6	2.5
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	33	1.4	2.3
13	Chronic liver disease and cirrhosis (K70,K73-K74)	30	1.3	2.1
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	29	1.3	2.0
15	Influenza and pneumonia (J09-J18)	25	1.1	1.7
...	All other causes (Residual)	492	21.4	34.2
Black, female, 40-44 years				
...	All causes	3,533	100.0	236.4
1	Malignant neoplasms (C00-C97)	898	25.4	60.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	601	17.0	40.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	275	7.8	18.4
4	Human immunodeficiency virus (HIV) disease (B20-B24)	225	6.4	15.1
5	Cerebrovascular diseases (I60-I69)	151	4.3	10.1
6	Diabetes mellitus (E10-E14)	140	4.0	9.4
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	80	2.3	5.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	70	2.0	4.7
9	Chronic lower respiratory diseases (J40-J47)	66	1.9	4.4
10	Septicemia (A40-A41)	65	1.8	4.3
11	Influenza and pneumonia (J09-J18)	63	1.8	4.2
12	Chronic liver disease and cirrhosis (K70,K73-K74)	58	1.6	3.9
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	51	1.4	3.4
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	46	1.3	3.1
15	Anemias (D50-D64)	32	0.9	2.1
...	All other causes (Residual)	712	20.2	47.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 45-49 years				
...	All causes	5,779	100.0	372.8
1	Malignant neoplasms (C00-C97)	1,670	28.9	107.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,065	18.4	68.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	359	6.2	23.2
4	Cerebrovascular diseases (I60-I69)	303	5.2	19.5
5	Human immunodeficiency virus (HIV) disease (B20-B24)	241	4.2	15.5
6	Diabetes mellitus (E10-E14)	240	4.2	15.5
7	Septicemia (A40-A41)	129	2.2	8.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	129	2.2	8.3
7	Chronic lower respiratory diseases (J40-J47)	126	2.2	8.1
10	Chronic liver disease and cirrhosis (K70,K73-K74)	124	2.1	8.0
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	95	1.6	6.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	84	1.5	5.4
13	Influenza and pneumonia (J09-J18)	74	1.3	4.8
14	Anemias (D50-D64)	52	0.9	3.4
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	36	0.6	2.3
...	All other causes (Residual)	1,052	18.2	67.9
Black, female, 50-54 years				
...	All causes	8,559	100.0	564.3
1	Malignant neoplasms (C00-C97)	2,772	32.4	182.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,744	20.4	115.0
3	Cerebrovascular diseases (I60-I69)	421	4.9	27.8
4	Diabetes mellitus (E10-E14)	409	4.8	27.0
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	400	4.7	26.4
6	Chronic lower respiratory diseases (J40-J47)	256	3.0	16.9
7	Human immunodeficiency virus (HIV) disease (B20-B24)	234	2.7	15.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	198	2.3	13.1
9	Chronic liver disease and cirrhosis (K70,K73-K74)	190	2.2	12.5
10	Septicemia (A40-A41)	181	2.1	11.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	116	1.4	7.6
12	Influenza and pneumonia (J09-J18)	109	1.3	7.2
13	Viral hepatitis (B15-B19)	69	0.8	4.5
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	64	0.7	4.2
15	Anemias (D50-D64)	33	0.4	2.2
15	Aortic aneurysm and dissection (I71)	33	0.4	2.2
...	All other causes (Residual)	1,330	15.5	87.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 55-59 years				
...	All causes	10,476	100.0	812.3
1	Malignant neoplasms (C00-C97)	3,547	33.9	275.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,280	21.8	176.8
3	Diabetes mellitus (E10-E14)	523	5.0	40.6
4	Cerebrovascular diseases (I60-I69)	520	5.0	40.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	355	3.4	27.5
6	Chronic lower respiratory diseases (J40-J47)	308	2.9	23.9
7	Septicemia (A40-A41)	263	2.5	20.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	255	2.4	19.8
9	Chronic liver disease and cirrhosis (K70,K73-K74)	203	1.9	15.7
10	Human immunodeficiency virus (HIV) disease (B20-B24)	188	1.8	14.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	177	1.7	13.7
12	Influenza and pneumonia (J09-J18)	138	1.3	10.7
13	Viral hepatitis (B15-B19)	102	1.0	7.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	46	0.4	3.6
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	39	0.4	3.0
...	All other causes (Residual)	1,532	14.6	118.8
Black, female, 60-64 years				
...	All causes	11,724	100.0	1,125.6
1	Malignant neoplasms (C00-C97)	4,091	34.9	392.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,597	22.2	249.3
3	Diabetes mellitus (E10-E14)	700	6.0	67.2
4	Cerebrovascular diseases (I60-I69)	604	5.2	58.0
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	380	3.2	36.5
6	Chronic lower respiratory diseases (J40-J47)	364	3.1	34.9
7	Septicemia (A40-A41)	255	2.2	24.5
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	232	2.0	22.3
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	193	1.6	18.5
10	Influenza and pneumonia (J09-J18)	170	1.5	16.3
11	Chronic liver disease and cirrhosis (K70,K73-K74)	162	1.4	15.6
12	Human immunodeficiency virus (HIV) disease (B20-B24)	102	0.9	9.8
13	Viral hepatitis (B15-B19)	88	0.8	8.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	61	0.5	5.9
15	Pneumonitis due to solids and liquids (J69)	42	0.4	4.0
...	All other causes (Residual)	1,683	14.4	161.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 65-69 years				
...	All causes	11,626	100.0	1,649.9
1	Malignant neoplasms (C00-C97)	3,897	33.5	553.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,661	22.9	377.6
3	Diabetes mellitus (E10-E14)	771	6.6	109.4
4	Cerebrovascular diseases (I60-I69)	613	5.3	87.0
5	Chronic lower respiratory diseases (J40-J47)	428	3.7	60.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	391	3.4	55.5
7	Septicemia (A40-A41)	301	2.6	42.7
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	190	1.6	27.0
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	183	1.6	26.0
10	Influenza and pneumonia (J09-J18)	182	1.6	25.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	91	0.8	12.9
12	Alzheimer's disease (G30)	60	0.5	8.5
13	Pneumonitis due to solids and liquids (J69)	54	0.5	7.7
14	Human immunodeficiency virus (HIV) disease (B20-B24)	53	0.5	7.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	51	0.4	7.2
...	All other causes (Residual)	1,700	14.6	241.3
Black, female, 70-74 years				
...	All causes	12,748	100.0	2,391.1
1	Malignant neoplasms (C00-C97)	3,816	29.9	715.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,052	23.9	572.5
3	Cerebrovascular diseases (I60-I69)	831	6.5	155.9
4	Diabetes mellitus (E10-E14)	744	5.8	139.6
5	Chronic lower respiratory diseases (J40-J47)	534	4.2	100.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	474	3.7	88.9
7	Septicemia (A40-A41)	362	2.8	67.9
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	245	1.9	46.0
9	Influenza and pneumonia (J09-J18)	229	1.8	43.0
10	Alzheimer's disease (G30)	162	1.3	30.4
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	157	1.2	29.4
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	69	0.5	12.9
13	Pneumonitis due to solids and liquids (J69)	53	0.4	9.9
13	Chronic liver disease and cirrhosis (K70,K73-K74)	53	0.4	9.9
15	Aortic aneurysm and dissection (I71)	50	0.4	9.4
...	All other causes (Residual)	1,917	15.0	359.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 75-79 years				
...	All causes	14,752	100.0	3,635.5
1	Malignant neoplasms (C00-C97)	3,707	25.1	913.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,655	24.8	900.8
3	Cerebrovascular diseases (I60-I69)	1,056	7.2	260.2
4	Diabetes mellitus (E10-E14)	871	5.9	214.7
5	Chronic lower respiratory diseases (J40-J47)	586	4.0	144.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	574	3.9	141.5
7	Alzheimer's disease (G30)	411	2.8	101.3
8	Septicemia (A40-A41)	410	2.8	101.0
9	Influenza and pneumonia (J09-J18)	301	2.0	74.2
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	276	1.9	68.0
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	170	1.2	41.9
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	98	0.7	24.2
13	Pneumonitis due to solids and liquids (J69)	86	0.6	21.2
14	Parkinson's disease (G20-G21)	75	0.5	18.5
15	Aortic aneurysm and dissection (I71)	59	0.4	14.5
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	59	0.4	14.5
...	All other causes (Residual)	2,358	16.0	581.1
Black, female, 80-84 years				
...	All causes	16,835	100.0	5,791.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,413	26.2	1,518.1
2	Malignant neoplasms (C00-C97)	3,330	19.8	1,145.5
3	Cerebrovascular diseases (I60-I69)	1,326	7.9	456.1
4	Diabetes mellitus (E10-E14)	886	5.3	304.8
5	Alzheimer's disease (G30)	690	4.1	237.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	631	3.7	217.1
7	Chronic lower respiratory diseases (J40-J47)	620	3.7	213.3
8	Septicemia (A40-A41)	403	2.4	138.6
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	367	2.2	126.2
10	Influenza and pneumonia (J09-J18)	348	2.1	119.7
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	195	1.2	67.1
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	104	0.6	35.8
13	Pneumonitis due to solids and liquids (J69)	102	0.6	35.1
14	Parkinson's disease (G20-G21)	90	0.5	31.0
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	61	0.4	21.0
...	All other causes (Residual)	3,269	19.4	1,124.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 85-89 years				
...	All causes	16,609	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,648	28.0	---
2	Malignant neoplasms (C00-C97)	2,393	14.4	---
3	Cerebrovascular diseases (I60-I69)	1,277	7.7	---
4	Alzheimer's disease (G30)	1,015	6.1	---
5	Diabetes mellitus (E10-E14)	705	4.2	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	557	3.4	---
7	Chronic lower respiratory diseases (J40-J47)	536	3.2	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	421	2.5	---
9	Septicemia (A40-A41)	419	2.5	---
10	Influenza and pneumonia (J09-J18)	417	2.5	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	159	1.0	---
12	Pneumonitis due to solids and liquids (J69)	129	0.8	---
13	Parkinson's disease (G20-G21)	89	0.5	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	81	0.5	---
15	Atherosclerosis (I70)	67	0.4	---
...	All other causes (Residual)	3,696	22.3	---
Black, female, 90-94 years				
...	All causes	11,907	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,453	29.0	---
2	Malignant neoplasms (C00-C97)	1,198	10.1	---
3	Cerebrovascular diseases (I60-I69)	963	8.1	---
4	Alzheimer's disease (G30)	914	7.7	---
5	Diabetes mellitus (E10-E14)	431	3.6	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	374	3.1	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	359	3.0	---
8	Influenza and pneumonia (J09-J18)	314	2.6	---
9	Chronic lower respiratory diseases (J40-J47)	285	2.4	---
10	Septicemia (A40-A41)	250	2.1	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	131	1.1	---
12	Pneumonitis due to solids and liquids (J69)	105	0.9	---
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	63	0.5	---
14	Atherosclerosis (I70)	56	0.5	---
15	Anemias (D50-D64)	48	0.4	---
...	All other causes (Residual)	2,963	24.9	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 95-99 years				
...	All causes	5,786	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,722	29.8	---
2	Alzheimer's disease (G30)	482	8.3	---
3	Cerebrovascular diseases (I60-I69)	409	7.1	---
4	Malignant neoplasms (C00-C97)	393	6.8	---
5	Essential hypertension and hypertensive renal disease (I10,I12,I15)	203	3.5	---
6	Diabetes mellitus (E10-E14)	175	3.0	---
6	Influenza and pneumonia (J09-J18)	175	3.0	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	143	2.5	---
9	Chronic lower respiratory diseases (J40-J47)	113	2.0	---
10	Septicemia (A40-A41)	105	1.8	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	60	1.0	---
12	Pneumonitis due to solids and liquids (J69)	52	0.9	---
13	Atherosclerosis (I70)	28	0.5	---
14	Anemias (D50-D64)	20	0.3	---
15	Nutritional deficiencies (E40-E64)	19	0.3	---
...	All other causes (Residual)	1,687	29.2	---
Black, female, 100 years and over				
...	All causes	1,887	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	596	31.6	---
2	Alzheimer's disease (G30)	142	7.5	---
3	Cerebrovascular diseases (I60-I69)	126	6.7	---
4	Malignant neoplasms (C00-C97)	87	4.6	---
5	Essential hypertension and hypertensive renal disease (I10,I12,I15)	72	3.8	---
6	Influenza and pneumonia (J09-J18)	68	3.6	---
7	Diabetes mellitus (E10-E14)	36	1.9	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	35	1.9	---
9	Septicemia (A40-A41)	30	1.6	---
10	Chronic lower respiratory diseases (J40-J47)	24	1.3	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	23	1.2	---
12	Atherosclerosis (I70)	20	1.1	---
13	Pneumonitis due to solids and liquids (J69)	18	1.0	---
14	Anemias (D50-D64)	10	0.5	---
15	Nutritional deficiencies (E40-E64)	7	0.4	---
...	All other causes (Residual)	593	31.4	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, all ages				
...	All causes	15,945	100.0	366.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,918	18.3	67.1
2	Malignant neoplasms (C00-C97)	2,810	17.6	64.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,829	11.5	42.1
4	Diabetes mellitus (E10-E14)	927	5.8	21.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	821	5.1	18.9
6	Chronic lower respiratory diseases (J40-J47)	657	4.1	15.1
7	Cerebrovascular diseases (I60-I69)	600	3.8	13.8
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	459	2.9	10.6
9	Influenza and pneumonia (J09-J18)	349	2.2	8.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	287	1.8	6.6
11	Septicemia (A40-A41)	269	1.7	6.2
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	254	1.6	5.8
13	Alzheimer's disease (G30)	226	1.4	5.2
14	Certain conditions originating in the perinatal period (P00-P96)	126	0.8	2.9
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	124	0.8	2.9
...	All other causes (Residual)	3,289	20.6	75.7
American Indian, both sexes, 1-4 years				
...	All causes	85	100.0	27.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	31	36.5	9.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	10.6	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7	8.2	*
4	Malignant neoplasms (C00-C97)	5	5.9	*
5	Influenza and pneumonia (J09-J18)	3	3.5	*
6	Septicemia (A40-A41)	2	2.4	*
6	Cerebrovascular diseases (I60-I69)	2	2.4	*
6	Chronic lower respiratory diseases (J40-J47)	2	2.4	*
9	Meningococcal infection (A39)	1	1.2	*
9	Infections of kidney (N10-N12,N13.6,N15.1)	1	1.2	*
9	Complications of medical and surgical care (Y40-Y84,Y88)	1	1.2	*
...	All other causes (Residual)	21	24.7	6.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 5-9 years				
...	All causes	38	100.0	10.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11	28.9	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	13.2	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	10.5	*
4	Malignant neoplasms (C00-C97)	3	7.9	*
4	Cerebrovascular diseases (I60-I69)	3	7.9	*
6	Septicemia (A40-A41)	1	2.6	*
6	Influenza and pneumonia (J09-J18)	1	2.6	*
6	Pneumonitis due to solids and liquids (J69)	1	2.6	*
6	Complications of medical and surgical care (Y40-Y84,Y88)	1	2.6	*
...	All other causes (Residual)	8	21.1	*
American Indian, both sexes, 10-14 years				
...	All causes	58	100.0	15.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	21	36.2	5.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11	19.0	*
3	Malignant neoplasms (C00-C97)	6	10.3	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	6.9	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	3.4	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.7	*
6	Cerebrovascular diseases (I60-I69)	1	1.7	*
6	Chronic lower respiratory diseases (J40-J47)	1	1.7	*
6	Certain conditions originating in the perinatal period (P00-P96)	1	1.7	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1	1.7	*
...	All other causes (Residual)	9	15.5	*

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 15-19 years				
...	All causes	233	100.0	60.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	109	46.8	28.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	60	25.8	15.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	19	8.2	*
4	Malignant neoplasms (C00-C97)	6	2.6	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	2.1	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	1.3	*
7	Influenza and pneumonia (J09-J18)	2	0.9	*
8	Diabetes mellitus (E10-E14)	1	0.4	*
8	Cerebrovascular diseases (I60-I69)	1	0.4	*
8	Aortic aneurysm and dissection (I71)	1	0.4	*
8	Chronic lower respiratory diseases (J40-J47)	1	0.4	*
8	Peptic ulcer (K25-K28)	1	0.4	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.4	*
8	Pregnancy, childbirth and the puerperium (O00-O99)	1	0.4	*
8	Legal intervention (Y35,Y89.0)	1	0.4	*
8	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.4	*
...	All other causes (Residual)	20	8.6	5.2
American Indian, both sexes, 20-24 years				
...	All causes	356	100.0	94.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	170	47.8	45.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	75	21.1	19.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	34	9.6	9.0
4	Malignant neoplasms (C00-C97)	8	2.2	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	8	2.2	*
6	Chronic liver disease and cirrhosis (K70,K73-K74)	6	1.7	*
7	Influenza and pneumonia (J09-J18)	3	0.8	*
7	Pregnancy, childbirth and the puerperium (O00-O99)	3	0.8	*
7	Legal intervention (Y35,Y89.0)	3	0.8	*
10	Cerebrovascular diseases (I60-I69)	2	0.6	*
10	Chronic lower respiratory diseases (J40-J47)	2	0.6	*
10	Pneumonitis due to solids and liquids (J69)	2	0.6	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.3	*
13	Anemias (D50-D64)	1	0.3	*
13	Diabetes mellitus (E10-E14)	1	0.3	*
...	All other causes (Residual)	37	10.4	9.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 25-29 years				
...	All causes	408	100.0	118.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	193	47.3	56.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	54	13.2	15.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	28	6.9	8.1
4	Chronic liver disease and cirrhosis (K70,K73-K74)	22	5.4	6.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	15	3.7	*
6	Malignant neoplasms (C00-C97)	13	3.2	*
7	Septicemia (A40-A41)	5	1.2	*
7	Diabetes mellitus (E10-E14)	5	1.2	*
9	Cerebrovascular diseases (I60-I69)	3	0.7	*
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.7	*
11	Influenza and pneumonia (J09-J18)	2	0.5	*
11	Pregnancy, childbirth and the puerperium (O00-O99)	2	0.5	*
11	Legal intervention (Y35,Y89.0)	2	0.5	*
14	Meningococcal infection (A39)	1	0.2	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.2	*
14	Peptic ulcer (K25-K28)	1	0.2	*
14	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.2	*
14	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.2	*
14	Certain conditions originating in the perinatal period (P00-P96)	1	0.2	*
...	All other causes (Residual)	55	13.5	15.9
American Indian, both sexes, 30-34 years				
...	All causes	433	100.0	134.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	157	36.3	48.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	40	9.2	12.5
3	Chronic liver disease and cirrhosis (K70,K73-K74)	38	8.8	11.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	32	7.4	10.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	27	6.2	8.4
6	Malignant neoplasms (C00-C97)	21	4.8	6.5
7	Diabetes mellitus (E10-E14)	12	2.8	*
8	Pregnancy, childbirth and the puerperium (O00-O99)	8	1.8	*
9	Cerebrovascular diseases (I60-I69)	7	1.6	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	6	1.4	*
10	Influenza and pneumonia (J09-J18)	6	1.4	*
12	Chronic lower respiratory diseases (J40-J47)	4	0.9	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.7	*
14	Septicemia (A40-A41)	2	0.5	*
14	Cholelithiasis and other disorders of gallbladder (K80-K82)	2	0.5	*
...	All other causes (Residual)	68	15.7	21.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 35-39 years				
...	All causes	528	100.0	181.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	145	27.5	49.8
2	Chronic liver disease and cirrhosis (K70,K73-K74)	69	13.1	23.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	54	10.2	18.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	49	9.3	16.8
5	Diabetes mellitus (E10-E14)	27	5.1	9.3
6	Malignant neoplasms (C00-C97)	25	4.7	8.6
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	23	4.4	7.9
8	Influenza and pneumonia (J09-J18)	10	1.9	*
9	Septicemia (A40-A41)	6	1.1	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	5	0.9	*
10	Cerebrovascular diseases (I60-I69)	5	0.9	*
12	Viral hepatitis (B15-B19)	4	0.8	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4	0.8	*
14	Pregnancy, childbirth and the puerperium (O00-O99)	3	0.6	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.6	*
...	All other causes (Residual)	96	18.2	33.0
American Indian, both sexes, 40-44 years				
...	All causes	673	100.0	236.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	148	22.0	52.0
2	Chronic liver disease and cirrhosis (K70,K73-K74)	100	14.9	35.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	84	12.5	29.5
4	Malignant neoplasms (C00-C97)	54	8.0	19.0
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	53	7.9	18.6
6	Diabetes mellitus (E10-E14)	35	5.2	12.3
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	24	3.6	8.4
8	Cerebrovascular diseases (I60-I69)	16	2.4	*
9	Influenza and pneumonia (J09-J18)	10	1.5	*
10	Septicemia (A40-A41)	8	1.2	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	1.0	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	5	0.7	*
13	Viral hepatitis (B15-B19)	4	0.6	*
14	Chronic lower respiratory diseases (J40-J47)	3	0.4	*
14	Cholelithiasis and other disorders of gallbladder (K80-K82)	3	0.4	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.4	*
...	All other causes (Residual)	116	17.2	40.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 45-49 years				
...	All causes	1,023	100.0	361.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	172	16.8	60.8
2	Chronic liver disease and cirrhosis (K70,K73-K74)	144	14.1	50.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	139	13.6	49.1
4	Malignant neoplasms (C00-C97)	138	13.5	48.8
5	Diabetes mellitus (E10-E14)	48	4.7	17.0
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	36	3.5	12.7
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	33	3.2	11.7
8	Cerebrovascular diseases (I60-I69)	30	2.9	10.6
9	Septicemia (A40-A41)	18	1.8	*
10	Influenza and pneumonia (J09-J18)	17	1.7	*
11	Chronic lower respiratory diseases (J40-J47)	14	1.4	*
12	Viral hepatitis (B15-B19)	13	1.3	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	1.1	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	7	0.7	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	6	0.6	*
...	All other causes (Residual)	197	19.3	69.6
American Indian, both sexes, 50-54 years				
...	All causes	1,254	100.0	477.7
1	Malignant neoplasms (C00-C97)	241	19.2	91.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	221	17.6	84.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	178	14.2	67.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	124	9.9	47.2
5	Diabetes mellitus (E10-E14)	76	6.1	29.0
6	Cerebrovascular diseases (I60-I69)	29	2.3	11.0
7	Chronic lower respiratory diseases (J40-J47)	28	2.2	10.7
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	28	2.2	10.7
9	Septicemia (A40-A41)	24	1.9	9.1
10	Viral hepatitis (B15-B19)	22	1.8	8.4
11	Influenza and pneumonia (J09-J18)	21	1.7	8.0
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	1.0	*
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	10	0.8	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	9	0.7	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	0.4	*
...	All other causes (Residual)	225	17.9	85.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 55-59 years				
...	All causes	1,347	100.0	641.1
1	Malignant neoplasms (C00-C97)	271	20.1	129.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	261	19.4	124.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	130	9.7	61.9
4	Diabetes mellitus (E10-E14)	110	8.2	52.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	109	8.1	51.9
6	Chronic lower respiratory diseases (J40-J47)	53	3.9	25.2
7	Cerebrovascular diseases (I60-I69)	34	2.5	16.2
8	Septicemia (A40-A41)	32	2.4	15.2
8	Influenza and pneumonia (J09-J18)	32	2.4	15.2
10	Viral hepatitis (B15-B19)	30	2.2	14.3
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	1.7	10.9
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	22	1.6	10.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	13	1.0	*
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	13	1.0	*
15	Pneumonitis due to solids and liquids (J69)	6	0.4	*
...	All other causes (Residual)	208	15.4	99.0
American Indian, both sexes, 60-64 years				
...	All causes	1,435	100.0	887.6
1	Malignant neoplasms (C00-C97)	392	27.3	242.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	308	21.5	190.5
3	Diabetes mellitus (E10-E14)	99	6.9	61.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	87	6.1	53.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	82	5.7	50.7
6	Chronic lower respiratory diseases (J40-J47)	72	5.0	44.5
7	Cerebrovascular diseases (I60-I69)	55	3.8	34.0
8	Septicemia (A40-A41)	33	2.3	20.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	27	1.9	16.7
10	Influenza and pneumonia (J09-J18)	22	1.5	13.6
11	Viral hepatitis (B15-B19)	18	1.3	*
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8	0.6	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7	0.5	*
14	Pneumonitis due to solids and liquids (J69)	6	0.4	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.3	*
...	All other causes (Residual)	214	14.9	132.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 65-69 years				
...	All causes	1,404	100.0	1,343.8
1	Malignant neoplasms (C00-C97)	379	27.0	362.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	300	21.4	287.1
3	Diabetes mellitus (E10-E14)	100	7.1	95.7
4	Chronic lower respiratory diseases (J40-J47)	97	6.9	92.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	58	4.1	55.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	57	4.1	54.6
7	Cerebrovascular diseases (I60-I69)	52	3.7	49.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	38	2.7	36.4
9	Septicemia (A40-A41)	27	1.9	25.8
10	Influenza and pneumonia (J09-J18)	25	1.8	23.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	15	1.1	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.6	*
12	Alzheimer's disease (G30)	9	0.6	*
14	Viral hepatitis (B15-B19)	8	0.6	*
14	Aortic aneurysm and dissection (I71)	8	0.6	*
...	All other causes (Residual)	222	15.8	212.5
American Indian, both sexes, 70-74 years				
...	All causes	1,452	100.0	2,037.9
1	Malignant neoplasms (C00-C97)	403	27.8	565.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	298	20.5	418.3
3	Diabetes mellitus (E10-E14)	118	8.1	165.6
4	Chronic lower respiratory diseases (J40-J47)	104	7.2	146.0
5	Cerebrovascular diseases (I60-I69)	63	4.3	88.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	49	3.4	68.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	43	3.0	60.4
8	Influenza and pneumonia (J09-J18)	33	2.3	46.3
9	Septicemia (A40-A41)	25	1.7	35.1
10	Chronic liver disease and cirrhosis (K70,K73-K74)	21	1.4	29.5
11	Alzheimer's disease (G30)	19	1.3	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.8	*
12	Parkinson's disease (G20-G21)	11	0.8	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7	0.5	*
14	Aortic aneurysm and dissection (I71)	7	0.5	*
14	Pneumonitis due to solids and liquids (J69)	7	0.5	*
...	All other causes (Residual)	233	16.0	327.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 75-79 years				
...	All causes	1,443	100.0	3,096.4
1	Malignant neoplasms (C00-C97)	345	23.9	740.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	304	21.1	652.3
3	Cerebrovascular diseases (I60-I69)	111	7.7	238.2
4	Diabetes mellitus (E10-E14)	110	7.6	236.0
5	Chronic lower respiratory diseases (J40-J47)	84	5.8	180.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	41	2.8	88.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	32	2.2	68.7
8	Alzheimer's disease (G30)	31	2.1	66.5
9	Septicemia (A40-A41)	30	2.1	64.4
10	Influenza and pneumonia (J09-J18)	26	1.8	55.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	22	1.5	47.2
12	Parkinson's disease (G20-G21)	15	1.0	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	15	1.0	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	0.8	*
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	9	0.6	*
...	All other causes (Residual)	256	17.7	549.3
American Indian, both sexes, 80-84 years				
...	All causes	1,350	100.0	4,662.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	349	25.9	1,205.2
2	Malignant neoplasms (C00-C97)	251	18.6	866.8
3	Chronic lower respiratory diseases (J40-J47)	94	7.0	324.6
4	Diabetes mellitus (E10-E14)	80	5.9	276.3
5	Cerebrovascular diseases (I60-I69)	65	4.8	224.5
6	Alzheimer's disease (G30)	50	3.7	172.7
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	39	2.9	134.7
8	Influenza and pneumonia (J09-J18)	38	2.8	131.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	35	2.6	120.9
10	Parkinson's disease (G20-G21)	22	1.6	76.0
11	Septicemia (A40-A41)	18	1.3	*
12	Pneumonitis due to solids and liquids (J69)	17	1.3	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	16	1.2	*
14	Chronic liver disease and cirrhosis (K70,K73-K74)	12	0.9	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.5	*
...	All other causes (Residual)	257	19.0	887.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 85-89 years				
...	All causes	1,082	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	283	26.2	---
2	Malignant neoplasms (C00-C97)	166	15.3	---
3	Cerebrovascular diseases (I60-I69)	61	5.6	---
4	Diabetes mellitus (E10-E14)	60	5.5	---
4	Chronic lower respiratory diseases (J40-J47)	60	5.5	---
6	Alzheimer's disease (G30)	58	5.4	---
7	Influenza and pneumonia (J09-J18)	41	3.8	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	26	2.4	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	25	2.3	---
10	Septicemia (A40-A41)	19	1.8	---
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	18	1.7	---
12	Pneumonitis due to solids and liquids (J69)	12	1.1	---
13	Parkinson's disease (G20-G21)	9	0.8	---
14	Chronic liver disease and cirrhosis (K70,K73-K74)	7	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.6	---
...	All other causes (Residual)	231	21.3	---
American Indian, both sexes, 90-94 years				
...	All causes	654	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	169	25.8	---
2	Malignant neoplasms (C00-C97)	60	9.2	---
3	Cerebrovascular diseases (I60-I69)	41	6.3	---
4	Influenza and pneumonia (J09-J18)	36	5.5	---
5	Alzheimer's disease (G30)	35	5.4	---
6	Diabetes mellitus (E10-E14)	31	4.7	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	24	3.7	---
8	Chronic lower respiratory diseases (J40-J47)	22	3.4	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19	2.9	---
10	Pneumonitis due to solids and liquids (J69)	17	2.6	---
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10	1.5	---
12	Septicemia (A40-A41)	9	1.4	---
12	Parkinson's disease (G20-G21)	9	1.4	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.9	---
15	Atherosclerosis (I70)	3	0.5	---
15	Aortic aneurysm and dissection (I71)	3	0.5	---
15	Chronic liver disease and cirrhosis (K70,K73-K74)	3	0.5	---
...	All other causes (Residual)	157	24.0	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 95-99 years				
...	All causes	265	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	80	30.2	---
2	Malignant neoplasms (C00-C97)	18	6.8	---
3	Alzheimer's disease (G30)	17	6.4	---
4	Influenza and pneumonia (J09-J18)	16	6.0	---
5	Cerebrovascular diseases (I60-I69)	15	5.7	---
6	Diabetes mellitus (E10-E14)	13	4.9	---
6	Chronic lower respiratory diseases (J40-J47)	13	4.9	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6	2.3	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5	1.9	---
10	Atherosclerosis (I70)	4	1.5	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	1.5	---
12	Septicemia (A40-A41)	3	1.1	---
12	Pneumonitis due to solids and liquids (J69)	3	1.1	---
14	Hernia (K40-K46)	2	0.8	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.4	---
15	Anemias (D50-D64)	1	0.4	---
15	Nutritional deficiencies (E40-E64)	1	0.4	---
15	Parkinson's disease (G20-G21)	1	0.4	---
15	Hyperplasia of prostate (N40)	1	0.4	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.4	---
...	All other causes (Residual)	60	22.6	---
American Indian, both sexes, 100 years and over				
...	All causes	58	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	14	24.1	---
2	Malignant neoplasms (C00-C97)	4	6.9	---
2	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4	6.9	---
2	Influenza and pneumonia (J09-J18)	4	6.9	---
5	Alzheimer's disease (G30)	3	5.2	---
5	Cerebrovascular diseases (I60-I69)	3	5.2	---
7	Septicemia (A40-A41)	2	3.4	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	3.4	---
9	Diabetes mellitus (E10-E14)	1	1.7	---
9	Aortic aneurysm and dissection (I71)	1	1.7	---
9	Chronic lower respiratory diseases (J40-J47)	1	1.7	---
9	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	1.7	---
9	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	1.7	---
...	All other causes (Residual)	17	29.3	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, all ages				
...	All causes	8,638	100.0	395.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,680	19.4	76.9
2	Malignant neoplasms (C00-C97)	1,486	17.2	68.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,227	14.2	56.1
4	Diabetes mellitus (E10-E14)	453	5.2	20.7
4	Chronic liver disease and cirrhosis (K70,K73-K74)	453	5.2	20.7
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	352	4.1	16.1
7	Chronic lower respiratory diseases (J40-J47)	304	3.5	13.9
8	Cerebrovascular diseases (I60-I69)	257	3.0	11.8
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	196	2.3	9.0
10	Influenza and pneumonia (J09-J18)	177	2.0	8.1
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	129	1.5	5.9
12	Septicemia (A40-A41)	109	1.3	5.0
13	Certain conditions originating in the perinatal period (P00-P96)	73	0.8	3.3
14	Viral hepatitis (B15-B19)	71	0.8	3.2
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	66	0.8	3.0
...	All other causes (Residual)	1,605	18.6	73.4
American Indian, male, 1-4 years				
...	All causes	46	100.0	28.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	21	45.7	13.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	10.9	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	8.7	*
4	Malignant neoplasms (C00-C97)	3	6.5	*
5	Chronic lower respiratory diseases (J40-J47)	2	4.3	*
6	Cerebrovascular diseases (I60-I69)	1	2.2	*
6	Influenza and pneumonia (J09-J18)	1	2.2	*
...	All other causes (Residual)	9	19.6	*
American Indian, male, 5-9 years				
...	All causes	16	100.0	*
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	25.0	*
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2	12.5	*
3	Malignant neoplasms (C00-C97)	1	6.3	*
3	Cerebrovascular diseases (I60-I69)	1	6.3	*
3	Influenza and pneumonia (J09-J18)	1	6.3	*
3	Pneumonitis due to solids and liquids (J69)	1	6.3	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1	6.3	*
3	Complications of medical and surgical care (Y40-Y84,Y88)	1	6.3	*
...	All other causes (Residual)	4	25.0	*

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 10-14 years				
...	All causes	34	100.0	17.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13	38.2	*
2	Malignant neoplasms (C00-C97)	5	14.7	*
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5	14.7	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	8.8	*
5	Cerebrovascular diseases (I60-I69)	1	2.9	*
5	Chronic lower respiratory diseases (J40-J47)	1	2.9	*
5	Certain conditions originating in the perinatal period (P00-P96)	1	2.9	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1	2.9	*
...	All other causes (Residual)	4	11.8	*
American Indian, male, 15-19 years				
...	All causes	156	100.0	78.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	70	44.9	35.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	45	28.8	22.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	15	9.6	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	2.6	*
5	Malignant neoplasms (C00-C97)	3	1.9	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	1.3	*
7	Diabetes mellitus (E10-E14)	1	0.6	*
7	Aortic aneurysm and dissection (I71)	1	0.6	*
7	Influenza and pneumonia (J09-J18)	1	0.6	*
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.6	*
7	Legal intervention (Y35,Y89.0)	1	0.6	*
7	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.6	*
...	All other causes (Residual)	11	7.1	*

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 20-24 years				
...	All causes	266	100.0	135.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	122	45.9	62.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	64	24.1	32.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	29	10.9	14.7
4	Malignant neoplasms (C00-C97)	5	1.9	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	1.9	*
6	Chronic liver disease and cirrhosis (K70,K73-K74)	4	1.5	*
7	Legal intervention (Y35,Y89.0)	3	1.1	*
8	Influenza and pneumonia (J09-J18)	2	0.8	*
8	Chronic lower respiratory diseases (J40-J47)	2	0.8	*
8	Pneumonitis due to solids and liquids (J69)	2	0.8	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.4	*
11	Anemias (D50-D64)	1	0.4	*
...	All other causes (Residual)	26	9.8	13.2
American Indian, male, 25-29 years				
...	All causes	287	100.0	159.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	136	47.4	75.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	45	15.7	25.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	25	8.7	13.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	10	3.5	*
5	Malignant neoplasms (C00-C97)	8	2.8	*
5	Chronic liver disease and cirrhosis (K70,K73-K74)	8	2.8	*
7	Diabetes mellitus (E10-E14)	4	1.4	*
8	Septicemia (A40-A41)	3	1.0	*
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	1.0	*
10	Cerebrovascular diseases (I60-I69)	2	0.7	*
10	Legal intervention (Y35,Y89.0)	2	0.7	*
12	Meningococcal infection (A39)	1	0.3	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.3	*
12	Peptic ulcer (K25-K28)	1	0.3	*
12	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.3	*
12	Certain conditions originating in the perinatal period (P00-P96)	1	0.3	*
...	All other causes (Residual)	36	12.5	20.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 30-34 years				
...	All causes	271	100.0	164.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	105	38.7	63.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	29	10.7	17.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	25	9.2	15.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	23	8.5	13.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	20	7.4	12.1
6	Malignant neoplasms (C00-C97)	9	3.3	*
7	Diabetes mellitus (E10-E14)	8	3.0	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	4	1.5	*
9	Cerebrovascular diseases (I60-I69)	3	1.1	*
9	Influenza and pneumonia (J09-J18)	3	1.1	*
11	Chronic lower respiratory diseases (J40-J47)	2	0.7	*
12	Septicemia (A40-A41)	1	0.4	*
12	Viral hepatitis (B15-B19)	1	0.4	*
12	Pneumonitis due to solids and liquids (J69)	1	0.4	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.4	*
12	Legal intervention (Y35,Y89.0)	1	0.4	*
...	All other causes (Residual)	35	12.9	21.2
American Indian, male, 35-39 years				
...	All causes	329	100.0	221.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	98	29.8	65.9
2	Chronic liver disease and cirrhosis (K70,K73-K74)	40	12.2	26.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	38	11.6	25.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	31	9.4	20.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	18	5.5	*
6	Diabetes mellitus (E10-E14)	17	5.2	*
7	Malignant neoplasms (C00-C97)	10	3.0	*
8	Influenza and pneumonia (J09-J18)	6	1.8	*
9	Septicemia (A40-A41)	3	0.9	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	3	0.9	*
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3	0.9	*
12	Viral hepatitis (B15-B19)	2	0.6	*
13	Meningitis (G00,G03)	1	0.3	*
13	Cerebrovascular diseases (I60-I69)	1	0.3	*
13	Pneumonitis due to solids and liquids (J69)	1	0.3	*
13	Diseases of appendix (K35-K38)	1	0.3	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.3	*
13	Legal intervention (Y35,Y89.0)	1	0.3	*
...	All other causes (Residual)	53	16.1	35.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 40-44 years				
...	All causes	405	100.0	281.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	103	25.4	71.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	57	14.1	39.6
3	Chronic liver disease and cirrhosis (K70,K73-K74)	47	11.6	32.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	42	10.4	29.2
5	Malignant neoplasms (C00-C97)	23	5.7	16.0
6	Diabetes mellitus (E10-E14)	20	4.9	13.9
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	20	4.9	13.9
8	Cerebrovascular diseases (I60-I69)	6	1.5	*
8	Influenza and pneumonia (J09-J18)	6	1.5	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	5	1.2	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	1.0	*
12	Viral hepatitis (B15-B19)	3	0.7	*
13	Pneumonitis due to solids and liquids (J69)	2	0.5	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	0.5	*
15	Septicemia (A40-A41)	1	0.2	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.2	*
15	Chronic lower respiratory diseases (J40-J47)	1	0.2	*
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.2	*
15	Legal intervention (Y35,Y89.0)	1	0.2	*
...	All other causes (Residual)	60	14.8	41.7
American Indian, male, 45-49 years				
...	All causes	605	100.0	429.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	120	19.8	85.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	100	16.5	70.9
3	Chronic liver disease and cirrhosis (K70,K73-K74)	74	12.2	52.5
4	Malignant neoplasms (C00-C97)	66	10.9	46.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	27	4.5	19.2
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	24	4.0	17.0
7	Diabetes mellitus (E10-E14)	22	3.6	15.6
8	Cerebrovascular diseases (I60-I69)	12	2.0	*
8	Influenza and pneumonia (J09-J18)	12	2.0	*
10	Septicemia (A40-A41)	9	1.5	*
11	Viral hepatitis (B15-B19)	8	1.3	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	1.3	*
13	Chronic lower respiratory diseases (J40-J47)	7	1.2	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	6	1.0	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4	0.7	*
...	All other causes (Residual)	106	17.5	75.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 50-54 years				
...	All causes	742	100.0	579.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	147	19.8	114.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	126	17.0	98.4
3	Malignant neoplasms (C00-C97)	124	16.7	96.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	72	9.7	56.2
5	Diabetes mellitus (E10-E14)	40	5.4	31.2
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	21	2.8	16.4
7	Cerebrovascular diseases (I60-I69)	16	2.2	*
8	Septicemia (A40-A41)	13	1.8	*
8	Viral hepatitis (B15-B19)	13	1.8	*
10	Chronic lower respiratory diseases (J40-J47)	12	1.6	*
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	10	1.3	*
12	Influenza and pneumonia (J09-J18)	9	1.2	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	8	1.1	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.7	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	0.5	*
...	All other causes (Residual)	122	16.4	95.2
American Indian, male, 55-59 years				
...	All causes	811	100.0	799.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	191	23.6	188.4
2	Malignant neoplasms (C00-C97)	152	18.7	149.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	85	10.5	83.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	67	8.3	66.1
5	Diabetes mellitus (E10-E14)	59	7.3	58.2
6	Viral hepatitis (B15-B19)	24	3.0	23.7
6	Chronic lower respiratory diseases (J40-J47)	24	3.0	23.7
8	Influenza and pneumonia (J09-J18)	22	2.7	21.7
9	Cerebrovascular diseases (I60-I69)	18	2.2	*
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	16	2.0	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	1.6	*
12	Septicemia (A40-A41)	12	1.5	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10	1.2	*
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	10	1.2	*
15	Peptic ulcer (K25-K28)	3	0.4	*
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	3	0.4	*
...	All other causes (Residual)	102	12.6	100.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 60-64 years				
...	All causes	795	100.0	1,017.9
1	Malignant neoplasms (C00-C97)	202	25.4	258.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	187	23.5	239.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	61	7.7	78.1
4	Diabetes mellitus (E10-E14)	56	7.0	71.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	51	6.4	65.3
6	Cerebrovascular diseases (I60-I69)	34	4.3	43.5
7	Chronic lower respiratory diseases (J40-J47)	32	4.0	41.0
8	Septicemia (A40-A41)	15	1.9	*
9	Viral hepatitis (B15-B19)	12	1.5	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	1.4	*
11	Influenza and pneumonia (J09-J18)	10	1.3	*
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6	0.8	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4	0.5	*
13	Pneumonitis due to solids and liquids (J69)	4	0.5	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	3	0.4	*
...	All other causes (Residual)	107	13.5	137.0
American Indian, male, 65-69 years				
...	All causes	797	100.0	1,611.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	210	26.3	424.6
2	Malignant neoplasms (C00-C97)	208	26.1	420.6
3	Chronic lower respiratory diseases (J40-J47)	52	6.5	105.1
4	Diabetes mellitus (E10-E14)	49	6.1	99.1
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	44	5.5	89.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	37	4.6	74.8
7	Cerebrovascular diseases (I60-I69)	21	2.6	42.5
8	Influenza and pneumonia (J09-J18)	16	2.0	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	1.6	*
10	Septicemia (A40-A41)	8	1.0	*
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	8	1.0	*
12	Aortic aneurysm and dissection (I71)	6	0.8	*
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6	0.8	*
14	Viral hepatitis (B15-B19)	4	0.5	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.5	*
14	Parkinson's disease (G20-G21)	4	0.5	*
...	All other causes (Residual)	107	13.4	216.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 70-74 years				
...	All causes	773	100.0	2,355.1
1	Malignant neoplasms (C00-C97)	223	28.8	679.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	177	22.9	539.3
3	Diabetes mellitus (E10-E14)	51	6.6	155.4
4	Chronic lower respiratory diseases (J40-J47)	44	5.7	134.1
5	Cerebrovascular diseases (I60-I69)	34	4.4	103.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	28	3.6	85.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	3.0	70.1
8	Influenza and pneumonia (J09-J18)	22	2.8	67.0
9	Septicemia (A40-A41)	16	2.1	*
10	Chronic liver disease and cirrhosis (K70,K73-K74)	14	1.8	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.9	*
12	Parkinson's disease (G20-G21)	6	0.8	*
13	Aortic aneurysm and dissection (I71)	4	0.5	*
14	Alzheimer's disease (G30)	3	0.4	*
14	Atherosclerosis (I70)	3	0.4	*
14	Pneumonitis due to solids and liquids (J69)	3	0.4	*
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3	0.4	*
...	All other causes (Residual)	112	14.5	341.2
American Indian, male, 75-79 years				
...	All causes	728	100.0	3,621.7
1	Malignant neoplasms (C00-C97)	195	26.8	970.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	183	25.1	910.4
3	Diabetes mellitus (E10-E14)	54	7.4	268.6
4	Chronic lower respiratory diseases (J40-J47)	44	6.0	218.9
5	Cerebrovascular diseases (I60-I69)	41	5.6	204.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	23	3.2	114.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	1.9	*
8	Influenza and pneumonia (J09-J18)	12	1.6	*
9	Septicemia (A40-A41)	10	1.4	*
9	Alzheimer's disease (G30)	10	1.4	*
11	Chronic liver disease and cirrhosis (K70,K73-K74)	9	1.2	*
12	Cholelithiasis and other disorders of gallbladder (K80-K82)	7	1.0	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.8	*
13	Parkinson's disease (G20-G21)	6	0.8	*
15	Aortic aneurysm and dissection (I71)	5	0.7	*
...	All other causes (Residual)	109	15.0	542.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 80-84 years				
...	All causes	621	100.0	5,302.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	157	25.3	1,340.6
2	Malignant neoplasms (C00-C97)	129	20.8	1,101.5
3	Chronic lower respiratory diseases (J40-J47)	42	6.8	358.6
4	Diabetes mellitus (E10-E14)	34	5.5	290.3
5	Cerebrovascular diseases (I60-I69)	31	5.0	264.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	26	4.2	222.0
7	Alzheimer's disease (G30)	17	2.7	*
8	Parkinson's disease (G20-G21)	14	2.3	*
8	Influenza and pneumonia (J09-J18)	14	2.3	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	2.3	*
11	Pneumonitis due to solids and liquids (J69)	8	1.3	*
12	Septicemia (A40-A41)	6	1.0	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.8	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5	0.8	*
13	Chronic liver disease and cirrhosis (K70,K73-K74)	5	0.8	*
...	All other causes (Residual)	114	18.4	973.4
American Indian, male, 85-89 years				
...	All causes	445	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	120	27.0	---
2	Malignant neoplasms (C00-C97)	79	17.8	---
3	Diabetes mellitus (E10-E14)	23	5.2	---
3	Influenza and pneumonia (J09-J18)	23	5.2	---
5	Chronic lower respiratory diseases (J40-J47)	22	4.9	---
6	Cerebrovascular diseases (I60-I69)	21	4.7	---
7	Alzheimer's disease (G30)	19	4.3	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	2.7	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11	2.5	---
10	Septicemia (A40-A41)	7	1.6	---
10	Pneumonitis due to solids and liquids (J69)	7	1.6	---
12	Parkinson's disease (G20-G21)	4	0.9	---
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.7	---
13	Chronic liver disease and cirrhosis (K70,K73-K74)	3	0.7	---
15	Nutritional deficiencies (E40-E64)	2	0.4	---
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2	0.4	---
15	Atherosclerosis (I70)	2	0.4	---
15	Pneumoconioses and chemical effects (J60-J66,J68)	2	0.4	---
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	2	0.4	---
...	All other causes (Residual)	81	18.2	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 90-94 years				
...	All causes	227	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	55	24.2	---
2	Malignant neoplasms (C00-C97)	33	14.5	---
3	Influenza and pneumonia (J09-J18)	12	5.3	---
4	Diabetes mellitus (E10-E14)	11	4.8	---
4	Alzheimer's disease (G30)	11	4.8	---
4	Cerebrovascular diseases (I60-I69)	11	4.8	---
7	Chronic lower respiratory diseases (J40-J47)	9	4.0	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9	4.0	---
9	Pneumonitis due to solids and liquids (J69)	7	3.1	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	3.1	---
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5	2.2	---
12	Parkinson's disease (G20-G21)	4	1.8	---
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.9	---
13	Atherosclerosis (I70)	2	0.9	---
13	Chronic liver disease and cirrhosis (K70,K73-K74)	2	0.9	---
...	All other causes (Residual)	47	20.7	---
American Indian, male, 95-99 years				
...	All causes	74	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	21	28.4	---
2	Chronic lower respiratory diseases (J40-J47)	7	9.5	---
3	Malignant neoplasms (C00-C97)	5	6.8	---
3	Influenza and pneumonia (J09-J18)	5	6.8	---
5	Diabetes mellitus (E10-E14)	4	5.4	---
6	Cerebrovascular diseases (I60-I69)	3	4.1	---
7	Alzheimer's disease (G30)	2	2.7	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2	2.7	---
7	Atherosclerosis (I70)	2	2.7	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	2.7	---
11	Nutritional deficiencies (E40-E64)	1	1.4	---
11	Parkinson's disease (G20-G21)	1	1.4	---
11	Pneumonitis due to solids and liquids (J69)	1	1.4	---
11	Hernia (K40-K46)	1	1.4	---
11	Hyperplasia of prostate (N40)	1	1.4	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1	1.4	---
...	All other causes (Residual)	15	20.3	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 100 years and over				
...	All causes	8	100.0	---
1	Malignant neoplasms (C00-C97)	3	37.5	---
2	Septicemia (A40-A41)	1	12.5	---
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	12.5	---
2	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	12.5	---
...	All other causes (Residual)	2	25.0	---
American Indian, female, all ages				
...	All causes	7,307	100.0	338.1
1	Malignant neoplasms (C00-C97)	1,324	18.1	61.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,238	16.9	57.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	602	8.2	27.9
4	Diabetes mellitus (E10-E14)	474	6.5	21.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	368	5.0	17.0
6	Chronic lower respiratory diseases (J40-J47)	353	4.8	16.3
7	Cerebrovascular diseases (I60-I69)	343	4.7	15.9
8	Influenza and pneumonia (J09-J18)	172	2.4	8.0
9	Alzheimer's disease (G30)	161	2.2	7.4
10	Septicemia (A40-A41)	160	2.2	7.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	158	2.2	7.3
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	107	1.5	5.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	72	1.0	3.3
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	58	0.8	2.7
15	Certain conditions originating in the perinatal period (P00-P96)	53	0.7	2.5
...	All other causes (Residual)	1,664	22.8	77.0
American Indian, female, 1-4 years				
...	All causes	39	100.0	25.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10	25.6	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	10.3	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	7.7	*
4	Septicemia (A40-A41)	2	5.1	*
4	Malignant neoplasms (C00-C97)	2	5.1	*
4	Influenza and pneumonia (J09-J18)	2	5.1	*
7	Meningococcal infection (A39)	1	2.6	*
7	Cerebrovascular diseases (I60-I69)	1	2.6	*
7	Infections of kidney (N10-N12,N13.6,N15.1)	1	2.6	*
7	Complications of medical and surgical care (Y40-Y84,Y88)	1	2.6	*
...	All other causes (Residual)	12	30.8	*

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 5-9 years				
...	All causes	22	100.0	11.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9	40.9	*
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	13.6	*
3	Malignant neoplasms (C00-C97)	2	9.1	*
3	Cerebrovascular diseases (I60-I69)	2	9.1	*
5	Septicemia (A40-A41)	1	4.5	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	4.5	*
...	All other causes (Residual)	4	18.2	*
American Indian, female, 10-14 years				
...	All causes	24	100.0	13.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8	33.3	*
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6	25.0	*
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	8.3	*
4	Malignant neoplasms (C00-C97)	1	4.2	*
4	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	4.2	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	4.2	*
...	All other causes (Residual)	5	20.8	*
American Indian, female, 15-19 years				
...	All causes	77	100.0	40.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	39	50.6	20.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	15	19.5	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	5.2	*
4	Malignant neoplasms (C00-C97)	3	3.9	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	1.3	*
5	Cerebrovascular diseases (I60-I69)	1	1.3	*
5	Influenza and pneumonia (J09-J18)	1	1.3	*
5	Chronic lower respiratory diseases (J40-J47)	1	1.3	*
5	Peptic ulcer (K25-K28)	1	1.3	*
5	Pregnancy, childbirth and the puerperium (O00-O99)	1	1.3	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	1.3	*
...	All other causes (Residual)	9	11.7	*

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 20-24 years				
...	All causes	90	100.0	50.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	48	53.3	26.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11	12.2	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	5.6	*
4	Malignant neoplasms (C00-C97)	3	3.3	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	3.3	*
4	Pregnancy, childbirth and the puerperium (O00-O99)	3	3.3	*
7	Cerebrovascular diseases (I60-I69)	2	2.2	*
9	Chronic liver disease and cirrhosis (K70,K73-K74)	2	2.2	*
9	Diabetes mellitus (E10-E14)	1	1.1	*
9	Influenza and pneumonia (J09-J18)	1	1.1	*
...	All other causes (Residual)	11	12.2	*
American Indian, female, 25-29 years				
...	All causes	121	100.0	73.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	57	47.1	34.4
2	Chronic liver disease and cirrhosis (K70,K73-K74)	14	11.6	*
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	9	7.4	*
4	Malignant neoplasms (C00-C97)	5	4.1	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	4.1	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	2.5	*
7	Septicemia (A40-A41)	2	1.7	*
7	Influenza and pneumonia (J09-J18)	2	1.7	*
7	Pregnancy, childbirth and the puerperium (O00-O99)	2	1.7	*
10	Diabetes mellitus (E10-E14)	1	0.8	*
10	Cerebrovascular diseases (I60-I69)	1	0.8	*
10	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.8	*
...	All other causes (Residual)	19	15.7	*

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 30-34 years				
...	All causes	162	100.0	103.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	52	32.1	33.3
2	Chronic liver disease and cirrhosis (K70,K73-K74)	18	11.1	*
3	Malignant neoplasms (C00-C97)	12	7.4	*
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11	6.8	*
5	Pregnancy, childbirth and the puerperium (O00-O99)	8	4.9	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7	4.3	*
7	Diabetes mellitus (E10-E14)	4	2.5	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	2.5	*
7	Cerebrovascular diseases (I60-I69)	4	2.5	*
10	Influenza and pneumonia (J09-J18)	3	1.9	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	2	1.2	*
11	Chronic lower respiratory diseases (J40-J47)	2	1.2	*
11	Cholelithiasis and other disorders of gallbladder (K80-K82)	2	1.2	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	1.2	*
15	Septicemia (A40-A41)	1	0.6	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.6	*
...	All other causes (Residual)	29	17.9	18.6
American Indian, female, 35-39 years				
...	All causes	199	100.0	139.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	47	23.6	32.9
2	Chronic liver disease and cirrhosis (K70,K73-K74)	29	14.6	20.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	18	9.0	*
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	16	8.0	*
5	Malignant neoplasms (C00-C97)	15	7.5	*
6	Diabetes mellitus (E10-E14)	10	5.0	*
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	2.5	*
8	Cerebrovascular diseases (I60-I69)	4	2.0	*
8	Influenza and pneumonia (J09-J18)	4	2.0	*
10	Septicemia (A40-A41)	3	1.5	*
10	Pregnancy, childbirth and the puerperium (O00-O99)	3	1.5	*
12	Viral hepatitis (B15-B19)	2	1.0	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	2	1.0	*
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	1.0	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.5	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.5	*
15	Chronic lower respiratory diseases (J40-J47)	1	0.5	*
15	Hernia (K40-K46)	1	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.5	*
15	Inflammatory diseases of female pelvic organs (N70-N76)	1	0.5	*
15	Legal intervention (Y35,Y89.0)	1	0.5	*
...	All other causes (Residual)	32	16.1	22.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 40-44 years				
...	All causes	268	100.0	190.7
1	Chronic liver disease and cirrhosis (K70,K73-K74)	53	19.8	37.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	45	16.8	32.0
3	Malignant neoplasms (C00-C97)	31	11.6	22.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	27	10.1	19.2
5	Diabetes mellitus (E10-E14)	15	5.6	*
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11	4.1	*
7	Cerebrovascular diseases (I60-I69)	10	3.7	*
8	Septicemia (A40-A41)	7	2.6	*
9	Influenza and pneumonia (J09-J18)	4	1.5	*
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	1.5	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	1.1	*
12	Tuberculosis (A16-A19)	2	0.7	*
12	Chronic lower respiratory diseases (J40-J47)	2	0.7	*
12	Cholelithiasis and other disorders of gallbladder (K80-K82)	2	0.7	*
15	Viral hepatitis (B15-B19)	1	0.4	*
15	Meningitis (G00,G03)	1	0.4	*
15	Aortic aneurysm and dissection (I71)	1	0.4	*
15	Peptic ulcer (K25-K28)	1	0.4	*
15	Pregnancy, childbirth and the puerperium (O00-O99)	1	0.4	*
15	Certain conditions originating in the perinatal period (P00-P96)	1	0.4	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.4	*
...	All other causes (Residual)	45	16.8	32.0
American Indian, female, 45-49 years				
...	All causes	418	100.0	294.6
1	Malignant neoplasms (C00-C97)	72	17.2	50.7
2	Chronic liver disease and cirrhosis (K70,K73-K74)	70	16.7	49.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	52	12.4	36.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	39	9.3	27.5
5	Diabetes mellitus (E10-E14)	26	6.2	18.3
6	Cerebrovascular diseases (I60-I69)	18	4.3	*
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	12	2.9	*
8	Septicemia (A40-A41)	9	2.2	*
9	Chronic lower respiratory diseases (J40-J47)	7	1.7	*
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	1.4	*
11	Viral hepatitis (B15-B19)	5	1.2	*
11	Influenza and pneumonia (J09-J18)	5	1.2	*
13	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	4	1.0	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.7	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2	0.5	*
...	All other causes (Residual)	88	21.1	62.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 50-54 years				
...	All causes	512	100.0	380.9
1	Malignant neoplasms (C00-C97)	117	22.9	87.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	74	14.5	55.1
3	Chronic liver disease and cirrhosis (K70,K73-K74)	52	10.2	38.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	52	10.2	38.7
5	Diabetes mellitus (E10-E14)	36	7.0	26.8
6	Chronic lower respiratory diseases (J40-J47)	16	3.1	*
7	Cerebrovascular diseases (I60-I69)	13	2.5	*
8	Influenza and pneumonia (J09-J18)	12	2.3	*
9	Septicemia (A40-A41)	11	2.1	*
10	Viral hepatitis (B15-B19)	9	1.8	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	1.6	*
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7	1.4	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.2	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.2	*
13	Anemias (D50-D64)	1	0.2	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.2	*
13	Aortic aneurysm and dissection (I71)	1	0.2	*
13	Pneumonitis due to solids and liquids (J69)	1	0.2	*
13	Peptic ulcer (K25-K28)	1	0.2	*
13	Diseases of appendix (K35-K38)	1	0.2	*
13	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.2	*
13	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.2	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.2	*
13	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.2	*
...	All other causes (Residual)	93	18.2	69.2
American Indian, female, 55-59 years				
...	All causes	536	100.0	493.0
1	Malignant neoplasms (C00-C97)	119	22.2	109.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	70	13.1	64.4
3	Diabetes mellitus (E10-E14)	51	9.5	46.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	45	8.4	41.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	42	7.8	38.6
6	Chronic lower respiratory diseases (J40-J47)	29	5.4	26.7
7	Septicemia (A40-A41)	20	3.7	18.4
8	Cerebrovascular diseases (I60-I69)	16	3.0	*
9	Influenza and pneumonia (J09-J18)	10	1.9	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	1.9	*
11	Viral hepatitis (B15-B19)	6	1.1	*
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6	1.1	*
13	Pneumonitis due to solids and liquids (J69)	5	0.9	*
14	Nutritional deficiencies (E40-E64)	3	0.6	*
14	Alzheimer's disease (G30)	3	0.6	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3	0.6	*
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	0.6	*
...	All other causes (Residual)	95	17.7	87.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 60-64 years				
...	All causes	640	100.0	765.7
1	Malignant neoplasms (C00-C97)	190	29.7	227.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	121	18.9	144.8
3	Diabetes mellitus (E10-E14)	43	6.7	51.4
4	Chronic lower respiratory diseases (J40-J47)	40	6.3	47.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	36	5.6	43.1
6	Cerebrovascular diseases (I60-I69)	21	3.3	25.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	21	3.3	25.1
8	Septicemia (A40-A41)	18	2.8	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	2.5	*
10	Influenza and pneumonia (J09-J18)	12	1.9	*
11	Viral hepatitis (B15-B19)	6	0.9	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.5	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3	0.5	*
14	Pneumonitis due to solids and liquids (J69)	2	0.3	*
14	Cholelithiasis and other disorders of gallbladder (K80-K82)	2	0.3	*
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2	0.3	*
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	0.3	*
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	2	0.3	*
...	All other causes (Residual)	100	15.6	119.6
American Indian, female, 65-69 years				
...	All causes	607	100.0	1,103.1
1	Malignant neoplasms (C00-C97)	171	28.2	310.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	90	14.8	163.6
3	Diabetes mellitus (E10-E14)	51	8.4	92.7
4	Chronic lower respiratory diseases (J40-J47)	45	7.4	81.8
5	Cerebrovascular diseases (I60-I69)	31	5.1	56.3
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	25	4.1	45.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	20	3.3	36.3
8	Septicemia (A40-A41)	19	3.1	*
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14	2.3	*
10	Influenza and pneumonia (J09-J18)	9	1.5	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7	1.2	*
12	Alzheimer's disease (G30)	6	1.0	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.8	*
13	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	5	0.8	*
15	Viral hepatitis (B15-B19)	4	0.7	*
...	All other causes (Residual)	105	17.3	190.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 70-74 years				
...	All causes	679	100.0	1,767.0
1	Malignant neoplasms (C00-C97)	180	26.5	468.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	121	17.8	314.9
3	Diabetes mellitus (E10-E14)	67	9.9	174.4
4	Chronic lower respiratory diseases (J40-J47)	60	8.8	156.1
5	Cerebrovascular diseases (I60-I69)	29	4.3	75.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	21	3.1	54.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20	2.9	52.0
8	Alzheimer's disease (G30)	16	2.4	*
9	Influenza and pneumonia (J09-J18)	11	1.6	*
10	Septicemia (A40-A41)	9	1.3	*
11	Chronic liver disease and cirrhosis (K70,K73-K74)	7	1.0	*
12	Nutritional deficiencies (E40-E64)	5	0.7	*
12	Parkinson's disease (G20-G21)	5	0.7	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5	0.7	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.6	*
15	Pneumonitis due to solids and liquids (J69)	4	0.6	*
...	All other causes (Residual)	115	16.9	299.3
American Indian, female, 75-79 years				
...	All causes	715	100.0	2,698.0
1	Malignant neoplasms (C00-C97)	150	21.0	566.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	121	16.9	456.6
3	Cerebrovascular diseases (I60-I69)	70	9.8	264.1
4	Diabetes mellitus (E10-E14)	56	7.8	211.3
5	Chronic lower respiratory diseases (J40-J47)	40	5.6	150.9
6	Alzheimer's disease (G30)	21	2.9	79.2
7	Septicemia (A40-A41)	20	2.8	75.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18	2.5	*
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18	2.5	*
10	Influenza and pneumonia (J09-J18)	14	2.0	*
11	Chronic liver disease and cirrhosis (K70,K73-K74)	13	1.8	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	11	1.5	*
13	Parkinson's disease (G20-G21)	9	1.3	*
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	7	1.0	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.8	*
...	All other causes (Residual)	141	19.7	532.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 80-84 years				
...	All causes	729	100.0	4,227.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	192	26.3	1,113.3
2	Malignant neoplasms (C00-C97)	122	16.7	707.4
3	Chronic lower respiratory diseases (J40-J47)	52	7.1	301.5
4	Diabetes mellitus (E10-E14)	46	6.3	266.7
5	Cerebrovascular diseases (I60-I69)	34	4.7	197.1
6	Alzheimer's disease (G30)	33	4.5	191.3
7	Influenza and pneumonia (J09-J18)	24	3.3	139.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21	2.9	121.8
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13	1.8	*
10	Septicemia (A40-A41)	12	1.6	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	11	1.5	*
12	Pneumonitis due to solids and liquids (J69)	9	1.2	*
13	Parkinson's disease (G20-G21)	8	1.1	*
14	Chronic liver disease and cirrhosis (K70,K73-K74)	7	1.0	*
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	5	0.7	*
...	All other causes (Residual)	140	19.2	811.8
American Indian, female, 85-89 years				
...	All causes	637	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	163	25.6	---
2	Malignant neoplasms (C00-C97)	87	13.7	---
3	Cerebrovascular diseases (I60-I69)	40	6.3	---
4	Alzheimer's disease (G30)	39	6.1	---
5	Chronic lower respiratory diseases (J40-J47)	38	6.0	---
6	Diabetes mellitus (E10-E14)	37	5.8	---
7	Influenza and pneumonia (J09-J18)	18	2.8	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	16	2.5	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	2.2	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14	2.2	---
11	Septicemia (A40-A41)	12	1.9	---
12	Parkinson's disease (G20-G21)	5	0.8	---
12	Pneumonitis due to solids and liquids (J69)	5	0.8	---
14	Chronic liver disease and cirrhosis (K70,K73-K74)	4	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.5	---
15	Nutritional deficiencies (E40-E64)	3	0.5	---
...	All other causes (Residual)	139	21.8	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 90-94 years				
...	All causes	427	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	114	26.7	---
2	Cerebrovascular diseases (I60-I69)	30	7.0	---
3	Malignant neoplasms (C00-C97)	27	6.3	---
4	Alzheimer's disease (G30)	24	5.6	---
4	Influenza and pneumonia (J09-J18)	24	5.6	---
6	Diabetes mellitus (E10-E14)	20	4.7	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15	3.5	---
8	Chronic lower respiratory diseases (J40-J47)	13	3.0	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	2.8	---
10	Pneumonitis due to solids and liquids (J69)	10	2.3	---
11	Septicemia (A40-A41)	8	1.9	---
12	Parkinson's disease (G20-G21)	5	1.2	---
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5	1.2	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.9	---
15	Aortic aneurysm and dissection (I71)	3	0.7	---
...	All other causes (Residual)	113	26.5	---
American Indian, female, 95-99 years				
...	All causes	191	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	59	30.9	---
2	Alzheimer's disease (G30)	15	7.9	---
3	Malignant neoplasms (C00-C97)	13	6.8	---
4	Cerebrovascular diseases (I60-I69)	12	6.3	---
5	Influenza and pneumonia (J09-J18)	11	5.8	---
6	Diabetes mellitus (E10-E14)	9	4.7	---
7	Chronic lower respiratory diseases (J40-J47)	6	3.1	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5	2.6	---
9	Septicemia (A40-A41)	3	1.6	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3	1.6	---
11	Atherosclerosis (I70)	2	1.0	---
11	Pneumonitis due to solids and liquids (J69)	2	1.0	---
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	1.0	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.5	---
14	Anemias (D50-D64)	1	0.5	---
14	Hernia (K40-K46)	1	0.5	---
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.5	---
...	All other causes (Residual)	45	23.6	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 100 years and over				
...	All causes	50	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	13	26.0	---
2	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4	8.0	---
2	Influenza and pneumonia (J09-J18)	4	8.0	---
4	Alzheimer's disease (G30)	3	6.0	---
4	Cerebrovascular diseases (I60-I69)	3	6.0	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	4.0	---
7	Septicemia (A40-A41)	1	2.0	---
7	Malignant neoplasms (C00-C97)	1	2.0	---
7	Diabetes mellitus (E10-E14)	1	2.0	---
7	Aortic aneurysm and dissection (I71)	1	2.0	---
7	Chronic lower respiratory diseases (J40-J47)	1	2.0	---
7	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	2.0	---
...	All other causes (Residual)	15	30.0	---
Asian or Pacific Islander, both sexes, all ages				
...	All causes	53,336	100.0	305.3
1	Malignant neoplasms (C00-C97)	14,713	27.6	84.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,586	21.7	66.3
3	Cerebrovascular diseases (I60-I69)	3,937	7.4	22.5
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,327	4.4	13.3
5	Diabetes mellitus (E10-E14)	2,035	3.8	11.6
6	Chronic lower respiratory diseases (J40-J47)	1,705	3.2	9.8
7	Influenza and pneumonia (J09-J18)	1,652	3.1	9.5
8	Alzheimer's disease (G30)	1,211	2.3	6.9
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,043	2.0	6.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	949	1.8	5.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	878	1.6	5.0
12	Septicemia (A40-A41)	613	1.1	3.5
13	Chronic liver disease and cirrhosis (K70,K73-K74)	498	0.9	2.9
14	Certain conditions originating in the perinatal period (P00-P96)	482	0.9	2.8
15	Parkinson's disease (G20-G21)	460	0.9	2.6
...	All other causes (Residual)	9,247	17.3	52.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 1-4 years				
...	All causes	129	100.0	13.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	21	16.3	2.2
2	Malignant neoplasms (C00-C97)	18	14.0	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	16	12.4	*
4	Influenza and pneumonia (J09-J18)	11	8.5	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	9	7.0	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	3.1	*
6	Chronic lower respiratory diseases (J40-J47)	4	3.1	*
8	Septicemia (A40-A41)	2	1.6	*
8	Anemias (D50-D64)	2	1.6	*
10	Tuberculosis (A16-A19)	1	0.8	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.8	*
...	All other causes (Residual)	40	31.0	4.2
Asian or Pacific Islander, both sexes, 5-9 years				
...	All causes	95	100.0	8.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	20	21.1	1.7
2	Malignant neoplasms (C00-C97)	19	20.0	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	7.4	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	6.3	*
5	Anemias (D50-D64)	5	5.3	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	5.3	*
7	Influenza and pneumonia (J09-J18)	4	4.2	*
8	Certain conditions originating in the perinatal period (P00-P96)	3	3.2	*
9	Cerebrovascular diseases (I60-I69)	2	2.1	*
10	Salmonella infections (A01-A02)	1	1.1	*
10	Chronic lower respiratory diseases (J40-J47)	1	1.1	*
10	Diseases of appendix (K35-K38)	1	1.1	*
...	All other causes (Residual)	21	22.1	1.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 10-14 years				
...	All causes	102	100.0	9.1
1	Malignant neoplasms (C00-C97)	23	22.5	2.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	20	19.6	1.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	9	8.8	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	3.9	*
4	Influenza and pneumonia (J09-J18)	4	3.9	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	3.9	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	3.9	*
8	Cerebrovascular diseases (I60-I69)	2	2.0	*
8	Pregnancy, childbirth and the puerperium (O00-O99)	2	2.0	*
8	Certain conditions originating in the perinatal period (P00-P96)	2	2.0	*
11	Septicemia (A40-A41)	1	1.0	*
11	Anemias (D50-D64)	1	1.0	*
11	Diabetes mellitus (E10-E14)	1	1.0	*
11	Chronic lower respiratory diseases (J40-J47)	1	1.0	*
...	All other causes (Residual)	24	23.5	2.1
Asian or Pacific Islander, both sexes, 15-19 years				
...	All causes	247	100.0	21.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	78	31.6	6.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	61	24.7	5.3
3	Malignant neoplasms (C00-C97)	34	13.8	3.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	20	8.1	1.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	2.0	*
6	Cerebrovascular diseases (I60-I69)	4	1.6	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	1.6	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	1.2	*
8	Aortic aneurysm and dissection (I71)	3	1.2	*
8	Chronic lower respiratory diseases (J40-J47)	3	1.2	*
11	Septicemia (A40-A41)	2	0.8	*
12	Viral hepatitis (B15-B19)	1	0.4	*
12	Anemias (D50-D64)	1	0.4	*
12	Meningitis (G00,G03)	1	0.4	*
12	Influenza and pneumonia (J09-J18)	1	0.4	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.4	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.4	*
...	All other causes (Residual)	24	9.7	2.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 20-24 years				
...	All causes	487	100.0	36.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	195	40.0	14.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	99	20.3	7.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	44	9.0	3.3
4	Malignant neoplasms (C00-C97)	36	7.4	2.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	27	5.5	2.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	15	3.1	*
7	Influenza and pneumonia (J09-J18)	5	1.0	*
8	Septicemia (A40-A41)	4	0.8	*
8	Cerebrovascular diseases (I60-I69)	4	0.8	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.6	*
11	Diabetes mellitus (E10-E14)	2	0.4	*
11	Aortic aneurysm and dissection (I71)	2	0.4	*
11	Chronic lower respiratory diseases (J40-J47)	2	0.4	*
11	Pneumonitis due to solids and liquids (J69)	2	0.4	*
11	Pregnancy, childbirth and the puerperium (O00-O99)	2	0.4	*
11	Legal intervention (Y35,Y89.0)	2	0.4	*
...	All other causes (Residual)	43	8.8	3.2
Asian or Pacific Islander, both sexes, 25-29 years				
...	All causes	509	100.0	34.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	147	28.9	10.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	117	23.0	8.0
3	Malignant neoplasms (C00-C97)	60	11.8	4.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	60	11.8	4.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	36	7.1	2.5
6	Cerebrovascular diseases (I60-I69)	6	1.2	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	1.2	*
8	Influenza and pneumonia (J09-J18)	5	1.0	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.8	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	3	0.6	*
10	Chronic lower respiratory diseases (J40-J47)	3	0.6	*
10	Chronic liver disease and cirrhosis (K70,K73-K74)	3	0.6	*
10	Pregnancy, childbirth and the puerperium (O00-O99)	3	0.6	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
15	Tuberculosis (A16-A19)	1	0.2	*
15	Diabetes mellitus (E10-E14)	1	0.2	*
15	Legal intervention (Y35,Y89.0)	1	0.2	*
...	All other causes (Residual)	51	10.0	3.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 30-34 years				
...	All causes	615	100.0	42.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	119	19.3	8.2
2	Malignant neoplasms (C00-C97)	112	18.2	7.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	106	17.2	7.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	74	12.0	5.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	31	5.0	2.1
6	Pregnancy, childbirth and the puerperium (O00-O99)	17	2.8	*
7	Cerebrovascular diseases (I60-I69)	16	2.6	*
8	Septicemia (A40-A41)	10	1.6	*
9	Diabetes mellitus (E10-E14)	8	1.3	*
10	Chronic lower respiratory diseases (J40-J47)	7	1.1	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	6	1.0	*
11	Influenza and pneumonia (J09-J18)	6	1.0	*
13	Viral hepatitis (B15-B19)	4	0.7	*
13	Anemias (D50-D64)	4	0.7	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.7	*
...	All other causes (Residual)	91	14.8	6.3
Asian or Pacific Islander, both sexes, 35-39 years				
...	All causes	793	100.0	54.1
1	Malignant neoplasms (C00-C97)	212	26.7	14.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	110	13.9	7.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	105	13.2	7.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	100	12.6	6.8
5	Cerebrovascular diseases (I60-I69)	34	4.3	2.3
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	32	4.0	2.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	17	2.1	*
8	Diabetes mellitus (E10-E14)	13	1.6	*
8	Influenza and pneumonia (J09-J18)	13	1.6	*
10	Pregnancy, childbirth and the puerperium (O00-O99)	9	1.1	*
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	1.1	*
12	Septicemia (A40-A41)	8	1.0	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.8	*
14	Viral hepatitis (B15-B19)	5	0.6	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	5	0.6	*
14	Chronic lower respiratory diseases (J40-J47)	5	0.6	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.6	*
...	All other causes (Residual)	105	13.2	7.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 40-44 years				
...	All causes	1,112	100.0	81.8
1	Malignant neoplasms (C00-C97)	350	31.5	25.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	173	15.6	12.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	133	12.0	9.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	85	7.6	6.3
5	Cerebrovascular diseases (I60-I69)	73	6.6	5.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	36	3.2	2.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	26	2.3	1.9
8	Diabetes mellitus (E10-E14)	19	1.7	*
9	Influenza and pneumonia (J09-J18)	15	1.3	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	1.3	*
11	Septicemia (A40-A41)	12	1.1	*
11	Viral hepatitis (B15-B19)	12	1.1	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	10	0.9	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	0.9	*
13	Aortic aneurysm and dissection (I71)	10	0.9	*
...	All other causes (Residual)	133	12.0	9.8
Asian or Pacific Islander, both sexes, 45-49 years				
...	All causes	1,704	100.0	138.2
1	Malignant neoplasms (C00-C97)	647	38.0	52.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	310	18.2	25.1
3	Cerebrovascular diseases (I60-I69)	120	7.0	9.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	120	7.0	9.7
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	88	5.2	7.1
6	Diabetes mellitus (E10-E14)	54	3.2	4.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	48	2.8	3.9
8	Influenza and pneumonia (J09-J18)	23	1.3	1.9
9	Viral hepatitis (B15-B19)	20	1.2	1.6
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	19	1.1	*
11	Septicemia (A40-A41)	18	1.1	*
12	Chronic lower respiratory diseases (J40-J47)	17	1.0	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	0.8	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	11	0.6	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	10	0.6	*
...	All other causes (Residual)	186	10.9	15.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 50-54 years				
...	All causes	2,309	100.0	206.8
1	Malignant neoplasms (C00-C97)	970	42.0	86.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	421	18.2	37.7
3	Cerebrovascular diseases (I60-I69)	135	5.8	12.1
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	117	5.1	10.5
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	82	3.6	7.3
6	Diabetes mellitus (E10-E14)	77	3.3	6.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	60	2.6	5.4
8	Viral hepatitis (B15-B19)	42	1.8	3.8
9	Influenza and pneumonia (J09-J18)	30	1.3	2.7
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	23	1.0	2.1
11	Chronic lower respiratory diseases (J40-J47)	21	0.9	1.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	20	0.9	1.8
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	0.7	*
14	Septicemia (A40-A41)	14	0.6	*
14	Aortic aneurysm and dissection (I71)	14	0.6	*
...	All other causes (Residual)	267	11.6	23.9
Asian or Pacific Islander, both sexes, 55-59 years				
...	All causes	3,174	100.0	324.5
1	Malignant neoplasms (C00-C97)	1,366	43.0	139.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	577	18.2	59.0
3	Cerebrovascular diseases (I60-I69)	213	6.7	21.8
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	130	4.1	13.3
5	Diabetes mellitus (E10-E14)	127	4.0	13.0
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	80	2.5	8.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	73	2.3	7.5
8	Chronic lower respiratory diseases (J40-J47)	44	1.4	4.5
9	Septicemia (A40-A41)	42	1.3	4.3
9	Viral hepatitis (B15-B19)	42	1.3	4.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	42	1.3	4.3
12	Influenza and pneumonia (J09-J18)	37	1.2	3.8
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	25	0.8	2.6
13	Aortic aneurysm and dissection (I71)	25	0.8	2.6
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	21	0.7	2.1
...	All other causes (Residual)	330	10.4	33.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 60-64 years				
...	All causes	3,870	100.0	477.4
1	Malignant neoplasms (C00-C97)	1,728	44.7	213.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	749	19.4	92.4
3	Cerebrovascular diseases (I60-I69)	261	6.7	32.2
4	Diabetes mellitus (E10-E14)	161	4.2	19.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	123	3.2	15.2
6	Chronic lower respiratory diseases (J40-J47)	69	1.8	8.5
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	65	1.7	8.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	55	1.4	6.8
9	Influenza and pneumonia (J09-J18)	49	1.3	6.0
9	Chronic liver disease and cirrhosis (K70,K73-K74)	49	1.3	6.0
11	Septicemia (A40-A41)	44	1.1	5.4
12	Viral hepatitis (B15-B19)	42	1.1	5.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	42	1.1	5.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	0.6	2.8
15	Aortic aneurysm and dissection (I71)	19	0.5	*
...	All other causes (Residual)	391	10.1	48.2
Asian or Pacific Islander, both sexes, 65-69 years				
...	All causes	4,083	100.0	737.5
1	Malignant neoplasms (C00-C97)	1,705	41.8	308.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	805	19.7	145.4
3	Cerebrovascular diseases (I60-I69)	287	7.0	51.8
4	Diabetes mellitus (E10-E14)	208	5.1	37.6
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	118	2.9	21.3
6	Chronic lower respiratory diseases (J40-J47)	91	2.2	16.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	84	2.1	15.2
8	Influenza and pneumonia (J09-J18)	59	1.4	10.7
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	53	1.3	9.6
10	Chronic liver disease and cirrhosis (K70,K73-K74)	51	1.2	9.2
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	48	1.2	8.7
12	Septicemia (A40-A41)	40	1.0	7.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	33	0.8	6.0
14	Viral hepatitis (B15-B19)	19	0.5	*
14	Parkinson's disease (G20-G21)	19	0.5	*
14	Pneumonitis due to solids and liquids (J69)	19	0.5	*
...	All other causes (Residual)	444	10.9	80.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 70-74 years				
...	All causes	4,792	100.0	1,173.4
1	Malignant neoplasms (C00-C97)	1,775	37.0	434.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,034	21.6	253.2
3	Cerebrovascular diseases (I60-I69)	336	7.0	82.3
4	Diabetes mellitus (E10-E14)	245	5.1	60.0
5	Chronic lower respiratory diseases (J40-J47)	151	3.2	37.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	123	2.6	30.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	103	2.1	25.2
8	Influenza and pneumonia (J09-J18)	99	2.1	24.2
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	69	1.4	16.9
10	Septicemia (A40-A41)	57	1.2	14.0
11	Parkinson's disease (G20-G21)	41	0.9	10.0
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	40	0.8	9.8
13	Chronic liver disease and cirrhosis (K70,K73-K74)	37	0.8	9.1
14	Viral hepatitis (B15-B19)	33	0.7	8.1
14	Alzheimer's disease (G30)	33	0.7	8.1
...	All other causes (Residual)	616	12.9	150.8
Asian or Pacific Islander, both sexes, 75-79 years				
...	All causes	5,992	100.0	2,078.8
1	Malignant neoplasms (C00-C97)	1,830	30.5	634.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,306	21.8	453.1
3	Cerebrovascular diseases (I60-I69)	496	8.3	172.1
4	Diabetes mellitus (E10-E14)	274	4.6	95.1
5	Chronic lower respiratory diseases (J40-J47)	247	4.1	85.7
6	Influenza and pneumonia (J09-J18)	196	3.3	68.0
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	181	3.0	62.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	120	2.0	41.6
9	Alzheimer's disease (G30)	106	1.8	36.8
10	Septicemia (A40-A41)	95	1.6	33.0
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	85	1.4	29.5
12	Parkinson's disease (G20-G21)	83	1.4	28.8
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	53	0.9	18.4
14	Chronic liver disease and cirrhosis (K70,K73-K74)	49	0.8	17.0
15	Pneumonitis due to solids and liquids (J69)	41	0.7	14.2
...	All other causes (Residual)	830	13.9	287.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 80-84 years				
...	All causes	7,559	100.0	3,880.5
1	Malignant neoplasms (C00-C97)	1,821	24.1	934.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,801	23.8	924.6
3	Cerebrovascular diseases (I60-I69)	692	9.2	355.2
4	Chronic lower respiratory diseases (J40-J47)	365	4.8	187.4
5	Diabetes mellitus (E10-E14)	315	4.2	161.7
6	Influenza and pneumonia (J09-J18)	283	3.7	145.3
7	Alzheimer's disease (G30)	213	2.8	109.3
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	191	2.5	98.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	162	2.1	83.2
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	137	1.8	70.3
11	Parkinson's disease (G20-G21)	121	1.6	62.1
12	Septicemia (A40-A41)	105	1.4	53.9
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	63	0.8	32.3
14	Pneumonitis due to solids and liquids (J69)	61	0.8	31.3
15	Chronic liver disease and cirrhosis (K70,K73-K74)	40	0.5	20.5
...	All other causes (Residual)	1,189	15.7	610.4
Asian or Pacific Islander, both sexes, 85-89 years				
...	All causes	7,261	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,921	26.5	---
2	Malignant neoplasms (C00-C97)	1,248	17.2	---
3	Cerebrovascular diseases (I60-I69)	622	8.6	---
4	Influenza and pneumonia (J09-J18)	341	4.7	---
5	Alzheimer's disease (G30)	333	4.6	---
5	Chronic lower respiratory diseases (J40-J47)	333	4.6	---
7	Diabetes mellitus (E10-E14)	308	4.2	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	209	2.9	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	192	2.6	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	165	2.3	---
11	Parkinson's disease (G20-G21)	94	1.3	---
12	Pneumonitis due to solids and liquids (J69)	86	1.2	---
13	Septicemia (A40-A41)	84	1.2	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	61	0.8	---
15	Aortic aneurysm and dissection (I71)	45	0.6	---
...	All other causes (Residual)	1,219	16.8	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 90-94 years				
...	All causes	5,157	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,453	28.2	---
2	Malignant neoplasms (C00-C97)	576	11.2	---
3	Cerebrovascular diseases (I60-I69)	446	8.6	---
4	Alzheimer's disease (G30)	309	6.0	---
5	Influenza and pneumonia (J09-J18)	302	5.9	---
6	Chronic lower respiratory diseases (J40-J47)	245	4.8	---
7	Diabetes mellitus (E10-E14)	162	3.1	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	144	2.8	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	118	2.3	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	104	2.0	---
11	Parkinson's disease (G20-G21)	70	1.4	---
12	Pneumonitis due to solids and liquids (J69)	68	1.3	---
13	Septicemia (A40-A41)	45	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	44	0.9	---
15	Aortic aneurysm and dissection (I71)	21	0.4	---
...	All other causes (Residual)	1,050	20.4	---
Asian or Pacific Islander, both sexes, 95-99 years				
...	All causes	2,010	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	618	30.7	---
2	Alzheimer's disease (G30)	163	8.1	---
3	Malignant neoplasms (C00-C97)	159	7.9	---
4	Cerebrovascular diseases (I60-I69)	152	7.6	---
5	Influenza and pneumonia (J09-J18)	128	6.4	---
6	Chronic lower respiratory diseases (J40-J47)	72	3.6	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	71	3.5	---
8	Diabetes mellitus (E10-E14)	52	2.6	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	42	2.1	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	31	1.5	---
11	Pneumonitis due to solids and liquids (J69)	29	1.4	---
12	Septicemia (A40-A41)	15	0.7	---
12	Parkinson's disease (G20-G21)	15	0.7	---
14	Anemias (D50-D64)	14	0.7	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	10	0.5	---
...	All other causes (Residual)	439	21.8	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 100 years and over				
...	All causes	453	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	143	31.6	---
2	Influenza and pneumonia (J09-J18)	35	7.7	---
3	Cerebrovascular diseases (I60-I69)	34	7.5	---
4	Alzheimer's disease (G30)	32	7.1	---
5	Essential hypertension and hypertensive renal disease (I10,I12,I15)	23	5.1	---
6	Malignant neoplasms (C00-C97)	19	4.2	---
7	Chronic lower respiratory diseases (J40-J47)	18	4.0	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	2.4	---
9	Diabetes mellitus (E10-E14)	7	1.5	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7	1.5	---
11	Septicemia (A40-A41)	5	1.1	---
12	Pneumonitis due to solids and liquids (J69)	2	0.4	---
13	Anemias (D50-D64)	1	0.2	---
13	Nutritional deficiencies (E40-E64)	1	0.2	---
13	Parkinson's disease (G20-G21)	1	0.2	---
13	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.2	---
13	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.2	---
13	Hyperplasia of prostate (N40)	1	0.2	---
...	All other causes (Residual)	111	24.5	---
Asian or Pacific Islander, male, all ages				
...	All causes	27,490	100.0	328.6
1	Malignant neoplasms (C00-C97)	7,444	27.1	89.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,209	22.6	74.2
3	Cerebrovascular diseases (I60-I69)	1,775	6.5	21.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,520	5.5	18.2
5	Diabetes mellitus (E10-E14)	1,040	3.8	12.4
6	Chronic lower respiratory diseases (J40-J47)	1,008	3.7	12.0
7	Influenza and pneumonia (J09-J18)	854	3.1	10.2
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	720	2.6	8.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	489	1.8	5.8
10	Alzheimer's disease (G30)	404	1.5	4.8
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	353	1.3	4.2
12	Chronic liver disease and cirrhosis (K70,K73-K74)	326	1.2	3.9
13	Septicemia (A40-A41)	304	1.1	3.6
14	Certain conditions originating in the perinatal period (P00-P96)	278	1.0	3.3
15	Parkinson's disease (G20-G21)	263	1.0	3.1
...	All other causes (Residual)	4,503	16.4	53.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 1-4 years				
...	All causes	74	100.0	15.4
1	Malignant neoplasms (C00-C97)	13	17.6	*
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11	14.9	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	12.2	*
4	Influenza and pneumonia (J09-J18)	4	5.4	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	5.4	*
6	Anemias (D50-D64)	2	2.7	*
6	Chronic lower respiratory diseases (J40-J47)	2	2.7	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.4	*
8	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	1.4	*
...	All other causes (Residual)	27	36.5	5.6
Asian or Pacific Islander, male, 5-9 years				
...	All causes	51	100.0	8.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12	23.5	*
2	Malignant neoplasms (C00-C97)	9	17.6	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	9.8	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	7.8	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	3.9	*
5	Certain conditions originating in the perinatal period (P00-P96)	2	3.9	*
7	Salmonella infections (A01-A02)	1	2.0	*
7	Anemias (D50-D64)	1	2.0	*
7	Influenza and pneumonia (J09-J18)	1	2.0	*
...	All other causes (Residual)	14	27.5	*
Asian or Pacific Islander, male, 10-14 years				
...	All causes	56	100.0	9.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13	23.2	*
2	Malignant neoplasms (C00-C97)	12	21.4	*
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5	8.9	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	3.6	*
4	Influenza and pneumonia (J09-J18)	2	3.6	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	3.6	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	3.6	*
8	Septicemia (A40-A41)	1	1.8	*
8	Anemias (D50-D64)	1	1.8	*
8	Diabetes mellitus (E10-E14)	1	1.8	*
8	Certain conditions originating in the perinatal period (P00-P96)	1	1.8	*
...	All other causes (Residual)	14	25.0	*

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 15-19 years				
...	All causes	165	100.0	28.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	52	31.5	8.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	45	27.3	7.7
3	Malignant neoplasms (C00-C97)	22	13.3	3.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	18	10.9	*
5	Aortic aneurysm and dissection (I71)	3	1.8	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	1.2	*
6	Cerebrovascular diseases (I60-I69)	2	1.2	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	1.2	*
9	Septicemia (A40-A41)	1	0.6	*
9	Viral hepatitis (B15-B19)	1	0.6	*
9	Anemias (D50-D64)	1	0.6	*
9	Chronic lower respiratory diseases (J40-J47)	1	0.6	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.6	*
9	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.6	*
...	All other causes (Residual)	13	7.9	*
Asian or Pacific Islander, male, 20-24 years				
...	All causes	374	100.0	54.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	167	44.7	24.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	77	20.6	11.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	35	9.4	5.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	21	5.6	3.1
5	Malignant neoplasms (C00-C97)	20	5.3	2.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	2.1	*
7	Septicemia (A40-A41)	3	0.8	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.8	*
7	Cerebrovascular diseases (I60-I69)	3	0.8	*
10	Aortic aneurysm and dissection (I71)	2	0.5	*
10	Influenza and pneumonia (J09-J18)	2	0.5	*
10	Legal intervention (Y35,Y89.0)	2	0.5	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.3	*
13	Diabetes mellitus (E10-E14)	1	0.3	*
13	Pneumonitis due to solids and liquids (J69)	1	0.3	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.3	*
...	All other causes (Residual)	26	7.0	3.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 25-29 years				
...	All causes	358	100.0	50.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	121	33.8	17.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	78	21.8	11.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	41	11.5	5.8
4	Malignant neoplasms (C00-C97)	36	10.1	5.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	26	7.3	3.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	1.4	*
7	Cerebrovascular diseases (I60-I69)	4	1.1	*
8	Influenza and pneumonia (J09-J18)	3	0.8	*
8	Chronic lower respiratory diseases (J40-J47)	3	0.8	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.6	*
10	Chronic liver disease and cirrhosis (K70,K73-K74)	2	0.6	*
12	Tuberculosis (A16-A19)	1	0.3	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.3	*
12	Diabetes mellitus (E10-E14)	1	0.3	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
12	Legal intervention (Y35,Y89.0)	1	0.3	*
...	All other causes (Residual)	32	8.9	4.5
Asian or Pacific Islander, male, 30-34 years				
...	All causes	391	100.0	57.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	93	23.8	13.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	75	19.2	11.0
3	Malignant neoplasms (C00-C97)	58	14.8	8.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	57	14.6	8.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	19	4.9	*
6	Cerebrovascular diseases (I60-I69)	9	2.3	*
7	Chronic lower respiratory diseases (J40-J47)	7	1.8	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	6	1.5	*
9	Septicemia (A40-A41)	5	1.3	*
10	Viral hepatitis (B15-B19)	4	1.0	*
10	Diabetes mellitus (E10-E14)	4	1.0	*
12	Influenza and pneumonia (J09-J18)	3	0.8	*
12	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	3	0.8	*
14	Tuberculosis (A16-A19)	2	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.5	*
...	All other causes (Residual)	44	11.3	6.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 35-39 years				
...	All causes	469	100.0	67.7
1	Malignant neoplasms (C00-C97)	83	17.7	12.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	81	17.3	11.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	73	15.6	10.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	71	15.1	10.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	25	5.3	3.6
6	Cerebrovascular diseases (I60-I69)	17	3.6	*
7	Chronic liver disease and cirrhosis (K70,K73-K74)	14	3.0	*
8	Diabetes mellitus (E10-E14)	8	1.7	*
9	Septicemia (A40-A41)	7	1.5	*
9	Influenza and pneumonia (J09-J18)	7	1.5	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	1.3	*
12	Viral hepatitis (B15-B19)	4	0.9	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.9	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.4	*
14	Chronic lower respiratory diseases (J40-J47)	2	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
...	All other causes (Residual)	63	13.4	9.1
Asian or Pacific Islander, male, 40-44 years				
...	All causes	654	100.0	102.0
1	Malignant neoplasms (C00-C97)	132	20.2	20.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	132	20.2	20.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	102	15.6	15.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	64	9.8	10.0
5	Cerebrovascular diseases (I60-I69)	38	5.8	5.9
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	28	4.3	4.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	22	3.4	3.4
8	Diabetes mellitus (E10-E14)	15	2.3	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	1.7	*
10	Viral hepatitis (B15-B19)	10	1.5	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	8	1.2	*
12	Influenza and pneumonia (J09-J18)	7	1.1	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.9	*
13	Aortic aneurysm and dissection (I71)	6	0.9	*
15	Septicemia (A40-A41)	4	0.6	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4	0.6	*
...	All other causes (Residual)	65	9.9	10.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 45-49 years				
...	All causes	1,008	100.0	173.2
1	Malignant neoplasms (C00-C97)	278	27.6	47.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	249	24.7	42.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	87	8.6	15.0
4	Cerebrovascular diseases (I60-I69)	75	7.4	12.9
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	59	5.9	10.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	38	3.8	6.5
7	Diabetes mellitus (E10-E14)	35	3.5	6.0
8	Viral hepatitis (B15-B19)	13	1.3	*
9	Influenza and pneumonia (J09-J18)	12	1.2	*
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	11	1.1	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	1.0	*
12	Chronic lower respiratory diseases (J40-J47)	9	0.9	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	8	0.8	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	8	0.8	*
15	Septicemia (A40-A41)	7	0.7	*
...	All other causes (Residual)	109	10.8	18.7
Asian or Pacific Islander, male, 50-54 years				
...	All causes	1,381	100.0	265.7
1	Malignant neoplasms (C00-C97)	467	33.8	89.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	332	24.0	63.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	80	5.8	15.4
4	Cerebrovascular diseases (I60-I69)	67	4.9	12.9
5	Diabetes mellitus (E10-E14)	54	3.9	10.4
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	52	3.8	10.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	47	3.4	9.0
8	Viral hepatitis (B15-B19)	29	2.1	5.6
9	Influenza and pneumonia (J09-J18)	25	1.8	4.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	16	1.2	*
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	14	1.0	*
12	Aortic aneurysm and dissection (I71)	13	0.9	*
12	Chronic lower respiratory diseases (J40-J47)	13	0.9	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	9	0.7	*
15	Septicemia (A40-A41)	6	0.4	*
...	All other causes (Residual)	157	11.4	30.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 55-59 years				
...	All causes	1,860	100.0	417.3
1	Malignant neoplasms (C00-C97)	695	37.4	155.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	413	22.2	92.7
3	Cerebrovascular diseases (I60-I69)	107	5.8	24.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	88	4.7	19.7
5	Diabetes mellitus (E10-E14)	83	4.5	18.6
6	Chronic liver disease and cirrhosis (K70,K73-K74)	62	3.3	13.9
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	56	3.0	12.6
8	Viral hepatitis (B15-B19)	30	1.6	6.7
9	Chronic lower respiratory diseases (J40-J47)	27	1.5	6.1
10	Septicemia (A40-A41)	25	1.3	5.6
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	1.2	5.2
12	Influenza and pneumonia (J09-J18)	21	1.1	4.7
13	Aortic aneurysm and dissection (I71)	19	1.0	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	0.9	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	16	0.9	*
...	All other causes (Residual)	179	9.6	40.2
Asian or Pacific Islander, male, 60-64 years				
...	All causes	2,210	100.0	607.4
1	Malignant neoplasms (C00-C97)	873	39.5	240.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	501	22.7	137.7
3	Cerebrovascular diseases (I60-I69)	147	6.7	40.4
4	Diabetes mellitus (E10-E14)	97	4.4	26.7
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	82	3.7	22.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	45	2.0	12.4
7	Chronic lower respiratory diseases (J40-J47)	40	1.8	11.0
8	Chronic liver disease and cirrhosis (K70,K73-K74)	35	1.6	9.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	34	1.5	9.3
10	Viral hepatitis (B15-B19)	29	1.3	8.0
11	Influenza and pneumonia (J09-J18)	27	1.2	7.4
12	Septicemia (A40-A41)	24	1.1	6.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	24	1.1	6.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	0.6	*
15	Aortic aneurysm and dissection (I71)	9	0.4	*
...	All other causes (Residual)	230	10.4	63.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 65-69 years				
...	All causes	2,356	100.0	938.1
1	Malignant neoplasms (C00-C97)	923	39.2	367.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	546	23.2	217.4
3	Cerebrovascular diseases (I60-I69)	150	6.4	59.7
4	Diabetes mellitus (E10-E14)	121	5.1	48.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	69	2.9	27.5
6	Chronic lower respiratory diseases (J40-J47)	52	2.2	20.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	50	2.1	19.9
8	Influenza and pneumonia (J09-J18)	33	1.4	13.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	33	1.4	13.1
10	Chronic liver disease and cirrhosis (K70,K73-K74)	31	1.3	12.3
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	30	1.3	11.9
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	1.0	9.2
13	Septicemia (A40-A41)	20	0.8	8.0
14	Parkinson's disease (G20-G21)	12	0.5	*
15	Pneumonitis due to solids and liquids (J69)	11	0.5	*
...	All other causes (Residual)	252	10.7	100.3
Asian or Pacific Islander, male, 70-74 years				
...	All causes	2,739	100.0	1,477.5
1	Malignant neoplasms (C00-C97)	983	35.9	530.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	634	23.1	342.0
3	Cerebrovascular diseases (I60-I69)	180	6.6	97.1
4	Diabetes mellitus (E10-E14)	142	5.2	76.6
5	Chronic lower respiratory diseases (J40-J47)	103	3.8	55.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	82	3.0	44.2
7	Influenza and pneumonia (J09-J18)	63	2.3	34.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	56	2.0	30.2
9	Septicemia (A40-A41)	36	1.3	19.4
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	32	1.2	17.3
11	Parkinson's disease (G20-G21)	28	1.0	15.1
12	Chronic liver disease and cirrhosis (K70,K73-K74)	26	0.9	14.0
13	Aortic aneurysm and dissection (I71)	19	0.7	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	0.7	*
15	Viral hepatitis (B15-B19)	16	0.6	*
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	16	0.6	*
...	All other causes (Residual)	305	11.1	164.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 75-79 years				
...	All causes	3,118	100.0	2,523.3
1	Malignant neoplasms (C00-C97)	955	30.6	772.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	717	23.0	580.2
3	Cerebrovascular diseases (I60-I69)	222	7.1	179.7
4	Chronic lower respiratory diseases (J40-J47)	150	4.8	121.4
5	Diabetes mellitus (E10-E14)	127	4.1	102.8
6	Influenza and pneumonia (J09-J18)	112	3.6	90.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	93	3.0	75.3
8	Parkinson's disease (G20-G21)	58	1.9	46.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	53	1.7	42.9
10	Septicemia (A40-A41)	46	1.5	37.2
10	Alzheimer's disease (G30)	46	1.5	37.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	43	1.4	34.8
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	32	1.0	25.9
14	Pneumonitis due to solids and liquids (J69)	29	0.9	23.5
15	Chronic liver disease and cirrhosis (K70,K73-K74)	22	0.7	17.8
...	All other causes (Residual)	413	13.2	334.2
Asian or Pacific Islander, male, 80-84 years				
...	All causes	3,525	100.0	4,574.7
1	Malignant neoplasms (C00-C97)	886	25.1	1,149.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	832	23.6	1,079.8
3	Cerebrovascular diseases (I60-I69)	296	8.4	384.1
4	Chronic lower respiratory diseases (J40-J47)	221	6.3	286.8
5	Influenza and pneumonia (J09-J18)	150	4.3	194.7
6	Diabetes mellitus (E10-E14)	137	3.9	177.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	98	2.8	127.2
8	Alzheimer's disease (G30)	75	2.1	97.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	75	2.1	97.3
10	Parkinson's disease (G20-G21)	67	1.9	87.0
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	49	1.4	63.6
12	Septicemia (A40-A41)	44	1.2	57.1
13	Pneumonitis due to solids and liquids (J69)	38	1.1	49.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	29	0.8	37.6
15	Aortic aneurysm and dissection (I71)	19	0.5	*
...	All other causes (Residual)	509	14.4	660.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 85-89 years				
...	All causes	3,277	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	850	25.9	---
2	Malignant neoplasms (C00-C97)	626	19.1	---
3	Cerebrovascular diseases (I60-I69)	245	7.5	---
4	Chronic lower respiratory diseases (J40-J47)	204	6.2	---
5	Influenza and pneumonia (J09-J18)	168	5.1	---
6	Diabetes mellitus (E10-E14)	120	3.7	---
7	Alzheimer's disease (G30)	112	3.4	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	111	3.4	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	84	2.6	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	67	2.0	---
11	Pneumonitis due to solids and liquids (J69)	50	1.5	---
12	Parkinson's disease (G20-G21)	47	1.4	---
13	Septicemia (A40-A41)	42	1.3	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	32	1.0	---
15	Aortic aneurysm and dissection (I71)	28	0.9	---
...	All other causes (Residual)	491	15.0	---
Asian or Pacific Islander, male, 90-94 years				
...	All causes	2,086	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	556	26.7	---
2	Malignant neoplasms (C00-C97)	292	14.0	---
3	Cerebrovascular diseases (I60-I69)	145	7.0	---
4	Influenza and pneumonia (J09-J18)	141	6.8	---
5	Chronic lower respiratory diseases (J40-J47)	129	6.2	---
6	Alzheimer's disease (G30)	99	4.7	---
7	Diabetes mellitus (E10-E14)	69	3.3	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	63	3.0	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	49	2.3	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	44	2.1	---
11	Pneumonitis due to solids and liquids (J69)	37	1.8	---
12	Parkinson's disease (G20-G21)	34	1.6	---
13	Septicemia (A40-A41)	19	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	0.6	---
15	Atherosclerosis (I70)	8	0.4	---
15	Aortic aneurysm and dissection (I71)	8	0.4	---
...	All other causes (Residual)	381	18.3	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 95-99 years				
...	All causes	717	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	214	29.8	---
2	Malignant neoplasms (C00-C97)	69	9.6	---
3	Influenza and pneumonia (J09-J18)	60	8.4	---
4	Cerebrovascular diseases (I60-I69)	54	7.5	---
5	Alzheimer's disease (G30)	49	6.8	---
6	Chronic lower respiratory diseases (J40-J47)	33	4.6	---
7	Diabetes mellitus (E10-E14)	21	2.9	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	21	2.9	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	14	2.0	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	2.0	---
11	Pneumonitis due to solids and liquids (J69)	11	1.5	---
12	Parkinson's disease (G20-G21)	8	1.1	---
13	Septicemia (A40-A41)	6	0.8	---
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.8	---
13	Anemias (D50-D64)	6	0.8	---
...	All other causes (Residual)	131	18.3	---
Asian or Pacific Islander, male, 100 years and over				
...	All causes	132	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	31	23.5	---
2	Cerebrovascular diseases (I60-I69)	13	9.8	---
3	Malignant neoplasms (C00-C97)	9	6.8	---
3	Influenza and pneumonia (J09-J18)	9	6.8	---
5	Chronic lower respiratory diseases (J40-J47)	8	6.1	---
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5	3.8	---
7	Alzheimer's disease (G30)	4	3.0	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4	3.0	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	3.0	---
10	Diabetes mellitus (E10-E14)	3	2.3	---
11	Septicemia (A40-A41)	2	1.5	---
12	Pneumonitis due to solids and liquids (J69)	1	0.8	---
12	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.8	---
12	Hyperplasia of prostate (N40)	1	0.8	---
...	All other causes (Residual)	37	28.0	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, all ages				
...	All causes	25,846	100.0	283.8
1	Malignant neoplasms (C00-C97)	7,269	28.1	79.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,377	20.8	59.1
3	Cerebrovascular diseases (I60-I69)	2,162	8.4	23.7
4	Diabetes mellitus (E10-E14)	995	3.8	10.9
5	Alzheimer's disease (G30)	807	3.1	8.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	807	3.1	8.9
7	Influenza and pneumonia (J09-J18)	798	3.1	8.8
8	Chronic lower respiratory diseases (J40-J47)	697	2.7	7.7
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	525	2.0	5.8
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	460	1.8	5.1
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	323	1.2	3.5
12	Septicemia (A40-A41)	309	1.2	3.4
13	Certain conditions originating in the perinatal period (P00-P96)	204	0.8	2.2
14	Parkinson's disease (G20-G21)	197	0.8	2.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	191	0.7	2.1
...	All other causes (Residual)	4,725	18.3	51.9
Asian or Pacific Islander, female, 1-4 years				
...	All causes	55	100.0	11.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10	18.2	*
2	Influenza and pneumonia (J09-J18)	7	12.7	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	12.7	*
4	Malignant neoplasms (C00-C97)	5	9.1	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	9.1	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	5.5	*
7	Septicemia (A40-A41)	2	3.6	*
7	Chronic lower respiratory diseases (J40-J47)	2	3.6	*
9	Tuberculosis (A16-A19)	1	1.8	*
...	All other causes (Residual)	13	23.6	*

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 5-9 years				
...	All causes	44	100.0	7.5
1	Malignant neoplasms (C00-C97)	10	22.7	*
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8	18.2	*
3	Anemias (D50-D64)	4	9.1	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	6.8	*
4	Influenza and pneumonia (J09-J18)	3	6.8	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	6.8	*
7	Cerebrovascular diseases (I60-I69)	2	4.5	*
8	Chronic lower respiratory diseases (J40-J47)	1	2.3	*
8	Diseases of appendix (K35-K38)	1	2.3	*
8	Certain conditions originating in the perinatal period (P00-P96)	1	2.3	*
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1	2.3	*
...	All other causes (Residual)	7	15.9	*
Asian or Pacific Islander, female, 10-14 years				
...	All causes	46	100.0	8.3
1	Malignant neoplasms (C00-C97)	11	23.9	*
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7	15.2	*
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4	8.7	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	4.3	*
4	Cerebrovascular diseases (I60-I69)	2	4.3	*
4	Influenza and pneumonia (J09-J18)	2	4.3	*
4	Pregnancy, childbirth and the puerperium (O00-O99)	2	4.3	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	4.3	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	4.3	*
10	Chronic lower respiratory diseases (J40-J47)	1	2.2	*
10	Certain conditions originating in the perinatal period (P00-P96)	1	2.2	*
...	All other causes (Residual)	10	21.7	*

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 15-19 years				
...	All causes	82	100.0	14.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	26	31.7	4.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	16	19.5	*
3	Malignant neoplasms (C00-C97)	12	14.6	*
4	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	3.7	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	3.7	*
6	Cerebrovascular diseases (I60-I69)	2	2.4	*
6	Chronic lower respiratory diseases (J40-J47)	2	2.4	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	2.4	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	2.4	*
10	Septicemia (A40-A41)	1	1.2	*
10	Meningitis (G00,G03)	1	1.2	*
10	Influenza and pneumonia (J09-J18)	1	1.2	*
...	All other causes (Residual)	11	13.4	*
Asian or Pacific Islander, female, 20-24 years				
...	All causes	113	100.0	17.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	28	24.8	4.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	22	19.5	3.3
3	Malignant neoplasms (C00-C97)	16	14.2	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	9	8.0	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	6.2	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	6	5.3	*
7	Influenza and pneumonia (J09-J18)	3	2.7	*
8	Chronic lower respiratory diseases (J40-J47)	2	1.8	*
8	Pregnancy, childbirth and the puerperium (O00-O99)	2	1.8	*
10	Septicemia (A40-A41)	1	0.9	*
10	Diabetes mellitus (E10-E14)	1	0.9	*
10	Cerebrovascular diseases (I60-I69)	1	0.9	*
10	Pneumonitis due to solids and liquids (J69)	1	0.9	*
10	Certain conditions originating in the perinatal period (P00-P96)	1	0.9	*
...	All other causes (Residual)	13	11.5	*

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 25-29 years				
...	All causes	151	100.0	20.1
1	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	39	25.8	5.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	26	17.2	3.5
3	Malignant neoplasms (C00-C97)	24	15.9	3.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	19	12.6	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	10	6.6	*
6	Pregnancy, childbirth and the puerperium (O00-O99)	3	2.0	*
7	Human immunodeficiency virus (HIV) disease (B20-B24)	2	1.3	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	1.3	*
7	Cerebrovascular diseases (I60-I69)	2	1.3	*
7	Influenza and pneumonia (J09-J18)	2	1.3	*
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.7	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.7	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.7	*
...	All other causes (Residual)	19	12.6	*
Asian or Pacific Islander, female, 30-34 years				
...	All causes	224	100.0	29.0
1	Malignant neoplasms (C00-C97)	54	24.1	7.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	31	13.8	4.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	26	11.6	3.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	17	7.6	*
4	Pregnancy, childbirth and the puerperium (O00-O99)	17	7.6	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	12	5.4	*
7	Cerebrovascular diseases (I60-I69)	7	3.1	*
8	Septicemia (A40-A41)	5	2.2	*
9	Anemias (D50-D64)	4	1.8	*
9	Diabetes mellitus (E10-E14)	4	1.8	*
11	Influenza and pneumonia (J09-J18)	3	1.3	*
12	Chronic liver disease and cirrhosis (K70,K73-K74)	2	0.9	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.9	*
14	Tuberculosis (A16-A19)	1	0.4	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.4	*
14	Aortic aneurysm and dissection (I71)	1	0.4	*
14	Pneumonitis due to solids and liquids (J69)	1	0.4	*
14	Inflammatory diseases of female pelvic organs (N70-N76)	1	0.4	*
14	Certain conditions originating in the perinatal period (P00-P96)	1	0.4	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.4	*
...	All other causes (Residual)	32	14.3	4.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 35-39 years				
...	All causes	324	100.0	42.0
1	Malignant neoplasms (C00-C97)	129	39.8	16.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	39	12.0	5.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	32	9.9	4.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	19	5.9	*
5	Cerebrovascular diseases (I60-I69)	17	5.2	*
6	Pregnancy, childbirth and the puerperium (O00-O99)	9	2.8	*
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7	2.2	*
8	Influenza and pneumonia (J09-J18)	6	1.9	*
9	Diabetes mellitus (E10-E14)	5	1.5	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	3	0.9	*
10	Chronic lower respiratory diseases (J40-J47)	3	0.9	*
10	Chronic liver disease and cirrhosis (K70,K73-K74)	3	0.9	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.9	*
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.9	*
15	Tuberculosis (A16-A19)	2	0.6	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.6	*
15	Nutritional deficiencies (E40-E64)	2	0.6	*
15	Infections of kidney (N10-N12,N13.6,N15.1)	2	0.6	*
...	All other causes (Residual)	38	11.7	4.9
Asian or Pacific Islander, female, 40-44 years				
...	All causes	458	100.0	63.7
1	Malignant neoplasms (C00-C97)	218	47.6	30.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	41	9.0	5.7
3	Cerebrovascular diseases (I60-I69)	35	7.6	4.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	31	6.8	4.3
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	21	4.6	2.9
6	Septicemia (A40-A41)	8	1.7	*
6	Influenza and pneumonia (J09-J18)	8	1.7	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8	1.7	*
9	Pregnancy, childbirth and the puerperium (O00-O99)	7	1.5	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.9	*
10	Diabetes mellitus (E10-E14)	4	0.9	*
10	Aortic aneurysm and dissection (I71)	4	0.9	*
10	Chronic liver disease and cirrhosis (K70,K73-K74)	4	0.9	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.9	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3	0.7	*
15	Pneumonitis due to solids and liquids (J69)	3	0.7	*
...	All other causes (Residual)	55	12.0	7.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 45-49 years				
...	All causes	696	100.0	107.0
1	Malignant neoplasms (C00-C97)	369	53.0	56.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	61	8.8	9.4
3	Cerebrovascular diseases (I60-I69)	45	6.5	6.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	33	4.7	5.1
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	29	4.2	4.5
6	Diabetes mellitus (E10-E14)	19	2.7	*
7	Septicemia (A40-A41)	11	1.6	*
7	Influenza and pneumonia (J09-J18)	11	1.6	*
9	Chronic liver disease and cirrhosis (K70,K73-K74)	10	1.4	*
10	Chronic lower respiratory diseases (J40-J47)	8	1.1	*
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8	1.1	*
12	Viral hepatitis (B15-B19)	7	1.0	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3	0.4	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.4	*
15	Tuberculosis (A16-A19)	2	0.3	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.3	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	0.3	*
...	All other causes (Residual)	73	10.5	11.2
Asian or Pacific Islander, female, 50-54 years				
...	All causes	928	100.0	155.6
1	Malignant neoplasms (C00-C97)	503	54.2	84.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	89	9.6	14.9
3	Cerebrovascular diseases (I60-I69)	68	7.3	11.4
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	37	4.0	6.2
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	30	3.2	5.0
6	Diabetes mellitus (E10-E14)	23	2.5	3.9
7	Viral hepatitis (B15-B19)	13	1.4	*
7	Chronic liver disease and cirrhosis (K70,K73-K74)	13	1.4	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	1.2	*
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	9	1.0	*
11	Septicemia (A40-A41)	8	0.9	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.9	*
11	Chronic lower respiratory diseases (J40-J47)	8	0.9	*
14	Influenza and pneumonia (J09-J18)	5	0.5	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4	0.4	*
...	All other causes (Residual)	99	10.7	16.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 55-59 years				
...	All causes	1,314	100.0	246.8
1	Malignant neoplasms (C00-C97)	671	51.1	126.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	164	12.5	30.8
3	Cerebrovascular diseases (I60-I69)	106	8.1	19.9
4	Diabetes mellitus (E10-E14)	44	3.3	8.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	42	3.2	7.9
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	24	1.8	4.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19	1.4	*
8	Septicemia (A40-A41)	17	1.3	*
8	Chronic lower respiratory diseases (J40-J47)	17	1.3	*
10	Influenza and pneumonia (J09-J18)	16	1.2	*
11	Viral hepatitis (B15-B19)	12	0.9	*
12	Chronic liver disease and cirrhosis (K70,K73-K74)	11	0.8	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.7	*
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7	0.5	*
15	Aortic aneurysm and dissection (I71)	6	0.5	*
...	All other causes (Residual)	149	11.3	28.0
Asian or Pacific Islander, female, 60-64 years				
...	All causes	1,660	100.0	371.6
1	Malignant neoplasms (C00-C97)	855	51.5	191.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	248	14.9	55.5
3	Cerebrovascular diseases (I60-I69)	114	6.9	25.5
4	Diabetes mellitus (E10-E14)	64	3.9	14.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	41	2.5	9.2
6	Chronic lower respiratory diseases (J40-J47)	29	1.7	6.5
7	Influenza and pneumonia (J09-J18)	22	1.3	4.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21	1.3	4.7
9	Septicemia (A40-A41)	20	1.2	4.5
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	20	1.2	4.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	18	1.1	*
12	Chronic liver disease and cirrhosis (K70,K73-K74)	14	0.8	*
13	Viral hepatitis (B15-B19)	13	0.8	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	0.6	*
14	Aortic aneurysm and dissection (I71)	10	0.6	*
...	All other causes (Residual)	161	9.7	36.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 65-69 years				
...	All causes	1,727	100.0	571.0
1	Malignant neoplasms (C00-C97)	782	45.3	258.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	259	15.0	85.6
3	Cerebrovascular diseases (I60-I69)	137	7.9	45.3
4	Diabetes mellitus (E10-E14)	87	5.0	28.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	49	2.8	16.2
6	Chronic lower respiratory diseases (J40-J47)	39	2.3	12.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	34	2.0	11.2
8	Influenza and pneumonia (J09-J18)	26	1.5	8.6
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	23	1.3	7.6
10	Septicemia (A40-A41)	20	1.2	6.6
10	Chronic liver disease and cirrhosis (K70,K73-K74)	20	1.2	6.6
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	15	0.9	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	0.6	*
14	Viral hepatitis (B15-B19)	9	0.5	*
14	Alzheimer's disease (G30)	9	0.5	*
...	All other causes (Residual)	208	12.0	68.8
Asian or Pacific Islander, female, 70-74 years				
...	All causes	2,053	100.0	920.6
1	Malignant neoplasms (C00-C97)	792	38.6	355.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	400	19.5	179.4
3	Cerebrovascular diseases (I60-I69)	156	7.6	70.0
4	Diabetes mellitus (E10-E14)	103	5.0	46.2
5	Chronic lower respiratory diseases (J40-J47)	48	2.3	21.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	47	2.3	21.1
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	41	2.0	18.4
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	37	1.8	16.6
9	Influenza and pneumonia (J09-J18)	36	1.8	16.1
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	22	1.1	9.9
11	Septicemia (A40-A41)	21	1.0	9.4
11	Alzheimer's disease (G30)	21	1.0	9.4
13	Viral hepatitis (B15-B19)	17	0.8	*
14	Parkinson's disease (G20-G21)	13	0.6	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	11	0.5	*
...	All other causes (Residual)	288	14.0	129.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 75-79 years				
...	All causes	2,874	100.0	1,745.2
1	Malignant neoplasms (C00-C97)	875	30.4	531.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	589	20.5	357.7
3	Cerebrovascular diseases (I60-I69)	274	9.5	166.4
4	Diabetes mellitus (E10-E14)	147	5.1	89.3
5	Chronic lower respiratory diseases (J40-J47)	97	3.4	58.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	88	3.1	53.4
7	Influenza and pneumonia (J09-J18)	84	2.9	51.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	67	2.3	40.7
9	Alzheimer's disease (G30)	60	2.1	36.4
10	Septicemia (A40-A41)	49	1.7	29.8
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	42	1.5	25.5
12	Chronic liver disease and cirrhosis (K70,K73-K74)	27	0.9	16.4
13	Parkinson's disease (G20-G21)	25	0.9	15.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	21	0.7	12.8
15	Viral hepatitis (B15-B19)	19	0.7	*
...	All other causes (Residual)	410	14.3	249.0
Asian or Pacific Islander, female, 80-84 years				
...	All causes	4,034	100.0	3,426.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	969	24.0	823.0
2	Malignant neoplasms (C00-C97)	935	23.2	794.1
3	Cerebrovascular diseases (I60-I69)	396	9.8	336.3
4	Diabetes mellitus (E10-E14)	178	4.4	151.2
5	Chronic lower respiratory diseases (J40-J47)	144	3.6	122.3
6	Alzheimer's disease (G30)	138	3.4	117.2
7	Influenza and pneumonia (J09-J18)	133	3.3	113.0
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	93	2.3	79.0
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	88	2.2	74.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	87	2.2	73.9
11	Septicemia (A40-A41)	61	1.5	51.8
12	Parkinson's disease (G20-G21)	54	1.3	45.9
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	34	0.8	28.9
14	Chronic liver disease and cirrhosis (K70,K73-K74)	26	0.6	22.1
15	Pneumonitis due to solids and liquids (J69)	23	0.6	19.5
...	All other causes (Residual)	675	16.7	573.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 85-89 years				
...	All causes	3,984	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,071	26.9	---
2	Malignant neoplasms (C00-C97)	622	15.6	---
3	Cerebrovascular diseases (I60-I69)	377	9.5	---
4	Alzheimer's disease (G30)	221	5.5	---
5	Diabetes mellitus (E10-E14)	188	4.7	---
6	Influenza and pneumonia (J09-J18)	173	4.3	---
7	Chronic lower respiratory diseases (J40-J47)	129	3.2	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	125	3.1	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	98	2.5	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	81	2.0	---
11	Parkinson's disease (G20-G21)	47	1.2	---
12	Septicemia (A40-A41)	42	1.1	---
13	Pneumonitis due to solids and liquids (J69)	36	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	29	0.7	---
15	Aortic aneurysm and dissection (I71)	17	0.4	---
15	Chronic liver disease and cirrhosis (K70,K73-K74)	17	0.4	---
...	All other causes (Residual)	711	17.8	---
Asian or Pacific Islander, female, 90-94 years				
...	All causes	3,071	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	897	29.2	---
2	Cerebrovascular diseases (I60-I69)	301	9.8	---
3	Malignant neoplasms (C00-C97)	284	9.2	---
4	Alzheimer's disease (G30)	210	6.8	---
5	Influenza and pneumonia (J09-J18)	161	5.2	---
6	Chronic lower respiratory diseases (J40-J47)	116	3.8	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	100	3.3	---
8	Diabetes mellitus (E10-E14)	93	3.0	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	55	1.8	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	55	1.8	---
11	Parkinson's disease (G20-G21)	36	1.2	---
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	32	1.0	---
13	Pneumonitis due to solids and liquids (J69)	31	1.0	---
14	Septicemia (A40-A41)	26	0.8	---
15	Aortic aneurysm and dissection (I71)	13	0.4	---
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	13	0.4	---
...	All other causes (Residual)	648	21.1	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 95-99 years				
...	All causes	1,293	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	404	31.2	---
2	Alzheimer's disease (G30)	114	8.8	---
3	Cerebrovascular diseases (I60-I69)	98	7.6	---
4	Malignant neoplasms (C00-C97)	90	7.0	---
5	Influenza and pneumonia (J09-J18)	68	5.3	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	57	4.4	---
7	Chronic lower respiratory diseases (J40-J47)	39	3.0	---
8	Diabetes mellitus (E10-E14)	31	2.4	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	21	1.6	---
10	Pneumonitis due to solids and liquids (J69)	18	1.4	---
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	1.3	---
12	Septicemia (A40-A41)	9	0.7	---
13	Anemias (D50-D64)	8	0.6	---
14	Parkinson's disease (G20-G21)	7	0.5	---
15	Atherosclerosis (I70)	6	0.5	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	6	0.5	---
...	All other causes (Residual)	300	23.2	---
Asian or Pacific Islander, female, 100 years and over				
...	All causes	321	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	112	34.9	---
2	Alzheimer's disease (G30)	28	8.7	---
3	Influenza and pneumonia (J09-J18)	26	8.1	---
4	Cerebrovascular diseases (I60-I69)	21	6.5	---
5	Essential hypertension and hypertensive renal disease (I10,I12,I15)	19	5.9	---
6	Malignant neoplasms (C00-C97)	10	3.1	---
6	Chronic lower respiratory diseases (J40-J47)	10	3.1	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	2.2	---
9	Diabetes mellitus (E10-E14)	4	1.2	---
10	Septicemia (A40-A41)	3	0.9	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2	0.6	---
12	Anemias (D50-D64)	1	0.3	---
12	Nutritional deficiencies (E40-E64)	1	0.3	---
12	Parkinson's disease (G20-G21)	1	0.3	---
12	Pneumonitis due to solids and liquids (J69)	1	0.3	---
12	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.3	---
...	All other causes (Residual)	74	23.1	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011