

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, all ages				
...	All causes	2,543,279	100.0	810.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	599,711	23.6	191.0
2	Malignant neoplasms (C00-C97)	582,623	22.9	185.6
3	Chronic lower respiratory diseases (J40-J47)	143,489	5.6	45.7
4	Cerebrovascular diseases (I60-I69)	128,546	5.1	40.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	127,792	5.0	40.7
6	Alzheimer's disease (G30)	83,637	3.3	26.6
7	Diabetes mellitus (E10-E14)	73,932	2.9	23.6
8	Influenza and pneumonia (J09-J18)	50,636	2.0	16.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	45,622	1.8	14.5
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	40,600	1.6	12.9
11	Septicemia (A40-A41)	35,842	1.4	11.4
12	Chronic liver disease and cirrhosis (K70,K73-K74)	34,979	1.4	11.1
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	29,115	1.1	9.3
14	Parkinson's disease (G20-G21)	23,818	0.9	7.6
15	Pneumonitis due to solids and liquids (J69)	17,897	0.7	5.7
...	All other causes (Residual)	525,040	20.6	167.3
All races, both sexes, 1-4 years				
...	All causes	4,218	100.0	26.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,353	32.1	8.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	501	11.9	3.1
3	Malignant neoplasms (C00-C97)	392	9.3	2.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	339	8.0	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	154	3.7	1.0
6	Influenza and pneumonia (J09-J18)	93	2.2	0.6
7	Septicemia (A40-A41)	62	1.5	0.4
8	Cerebrovascular diseases (I60-I69)	56	1.3	0.3
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	55	1.3	0.3
10	Chronic lower respiratory diseases (J40-J47)	51	1.2	0.3
11	Certain conditions originating in the perinatal period (P00-P96)	42	1.0	0.3
12	Acute bronchitis and bronchiolitis (J20-J21)	17	0.4	*
13	Anemias (D50-D64)	15	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	12	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	0.3	*
...	All other causes (Residual)	1,065	25.2	6.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 5-9 years				
...	All causes	2,334	100.0	11.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	743	31.8	3.6
2	Malignant neoplasms (C00-C97)	440	18.9	2.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	167	7.2	0.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	138	5.9	0.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	67	2.9	0.3
6	Chronic lower respiratory diseases (J40-J47)	63	2.7	0.3
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	47	2.0	0.2
8	Influenza and pneumonia (J09-J18)	44	1.9	0.2
9	Cerebrovascular diseases (I60-I69)	34	1.5	0.2
10	Septicemia (A40-A41)	26	1.1	0.1
11	Certain conditions originating in the perinatal period (P00-P96)	18	0.8	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	14	0.6	*
13	Diabetes mellitus (E10-E14)	12	0.5	*
14	Anemias (D50-D64)	10	0.4	*
15	Meningitis (G00,G03)	9	0.4	*
...	All other causes (Residual)	502	21.5	2.5
All races, both sexes, 10-14 years				
...	All causes	2,866	100.0	13.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	807	28.2	3.9
2	Malignant neoplasms (C00-C97)	472	16.5	2.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	306	10.7	1.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	173	6.0	0.8
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	160	5.6	0.8
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	108	3.8	0.5
7	Chronic lower respiratory diseases (J40-J47)	56	2.0	0.3
8	Cerebrovascular diseases (I60-I69)	51	1.8	0.2
9	Influenza and pneumonia (J09-J18)	41	1.4	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	40	1.4	0.2
11	Septicemia (A40-A41)	26	0.9	0.1
12	Diabetes mellitus (E10-E14)	22	0.8	0.1
13	Anemias (D50-D64)	18	0.6	*
14	Certain conditions originating in the perinatal period (P00-P96)	16	0.6	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	8	0.3	*
...	All other causes (Residual)	562	19.6	2.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 15-19 years				
...	All causes	10,074	100.0	47.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,995	39.7	18.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,782	17.7	8.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,614	16.0	7.6
4	Malignant neoplasms (C00-C97)	628	6.2	2.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	302	3.0	1.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	201	2.0	0.9
7	Cerebrovascular diseases (I60-I69)	63	0.6	0.3
8	Chronic lower respiratory diseases (J40-J47)	54	0.5	0.3
9	Influenza and pneumonia (J09-J18)	47	0.5	0.2
10	Diabetes mellitus (E10-E14)	44	0.4	0.2
11	Septicemia (A40-A41)	40	0.4	0.2
12	Pregnancy, childbirth and the puerperium (O00-O99)	37	0.4	0.2
13	Legal intervention (Y35,Y89.0)	35	0.3	0.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	30	0.3	0.1
14	Anemias (D50-D64)	30	0.3	0.1
...	All other causes (Residual)	1,172	11.6	5.5
All races, both sexes, 20-24 years				
...	All causes	19,108	100.0	84.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,913	41.4	35.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,090	16.2	13.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,000	15.7	13.3
4	Malignant neoplasms (C00-C97)	946	5.0	4.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	654	3.4	2.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	222	1.2	1.0
7	Diabetes mellitus (E10-E14)	152	0.8	0.7
8	Pregnancy, childbirth and the puerperium (O00-O99)	132	0.7	0.6
9	Cerebrovascular diseases (I60-I69)	120	0.6	0.5
10	Human immunodeficiency virus (HIV) disease (B20-B24)	105	0.5	0.5
11	Influenza and pneumonia (J09-J18)	100	0.5	0.4
12	Chronic lower respiratory diseases (J40-J47)	89	0.5	0.4
13	Legal intervention (Y35,Y89.0)	83	0.4	0.4
14	Septicemia (A40-A41)	80	0.4	0.4
15	Anemias (D50-D64)	59	0.3	0.3
...	All other causes (Residual)	2,363	12.4	10.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 25-29 years				
...	All causes	20,992	100.0	98.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,975	38.0	37.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,184	15.2	14.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,388	11.4	11.2
4	Malignant neoplasms (C00-C97)	1,384	6.6	6.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,177	5.6	5.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	247	1.2	1.2
7	Diabetes mellitus (E10-E14)	246	1.2	1.1
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	201	1.0	0.9
9	Pregnancy, childbirth and the puerperium (O00-O99)	191	0.9	0.9
10	Cerebrovascular diseases (I60-I69)	185	0.9	0.9
11	Chronic liver disease and cirrhosis (K70,K73-K74)	171	0.8	0.8
12	Septicemia (A40-A41)	148	0.7	0.7
13	Influenza and pneumonia (J09-J18)	141	0.7	0.7
14	Chronic lower respiratory diseases (J40-J47)	132	0.6	0.6
15	Legal intervention (Y35,Y89.0)	85	0.4	0.4
...	All other causes (Residual)	3,137	14.9	14.7
All races, both sexes, 30-34 years				
...	All causes	23,599	100.0	112.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,876	33.4	37.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,032	12.8	14.5
3	Malignant neoplasms (C00-C97)	2,290	9.7	11.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,054	8.7	9.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,954	8.3	9.3
6	Chronic liver disease and cirrhosis (K70,K73-K74)	426	1.8	2.0
7	Human immunodeficiency virus (HIV) disease (B20-B24)	405	1.7	1.9
8	Diabetes mellitus (E10-E14)	400	1.7	1.9
9	Cerebrovascular diseases (I60-I69)	350	1.5	1.7
10	Influenza and pneumonia (J09-J18)	215	0.9	1.0
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	200	0.8	1.0
12	Pregnancy, childbirth and the puerperium (O00-O99)	199	0.8	1.0
13	Septicemia (A40-A41)	177	0.8	0.8
14	Chronic lower respiratory diseases (J40-J47)	157	0.7	0.8
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	150	0.6	0.7
...	All other causes (Residual)	3,714	15.7	17.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 35-39 years				
...	All causes	27,372	100.0	140.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,081	25.9	36.3
2	Malignant neoplasms (C00-C97)	3,717	13.6	19.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,639	13.3	18.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,110	11.4	16.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,442	5.3	7.4
6	Chronic liver disease and cirrhosis (K70,K73-K74)	757	2.8	3.9
7	Diabetes mellitus (E10-E14)	661	2.4	3.4
8	Cerebrovascular diseases (I60-I69)	585	2.1	3.0
9	Human immunodeficiency virus (HIV) disease (B20-B24)	520	1.9	2.7
10	Septicemia (A40-A41)	261	1.0	1.3
11	Influenza and pneumonia (J09-J18)	247	0.9	1.3
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	241	0.9	1.2
13	Chronic lower respiratory diseases (J40-J47)	221	0.8	1.1
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	184	0.7	0.9
15	Pregnancy, childbirth and the puerperium (O00-O99)	170	0.6	0.9
...	All other causes (Residual)	4,536	16.6	23.3
All races, both sexes, 40-44 years				
...	All causes	41,790	100.0	198.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,953	19.0	37.8
2	Malignant neoplasms (C00-C97)	7,620	18.2	36.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,850	16.4	32.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,648	8.7	17.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,712	4.1	8.1
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,263	3.0	6.0
7	Diabetes mellitus (E10-E14)	1,206	2.9	5.7
8	Cerebrovascular diseases (I60-I69)	1,145	2.7	5.4
9	Human immunodeficiency virus (HIV) disease (B20-B24)	825	2.0	3.9
10	Chronic lower respiratory diseases (J40-J47)	522	1.2	2.5
11	Septicemia (A40-A41)	496	1.2	2.4
12	Influenza and pneumonia (J09-J18)	438	1.0	2.1
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	408	1.0	1.9
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	226	0.5	1.1
15	Viral hepatitis (B15-B19)	217	0.5	1.0
...	All other causes (Residual)	7,261	17.4	34.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 45-49 years				
...	All causes	68,479	100.0	315.7
1	Malignant neoplasms (C00-C97)	16,014	23.4	73.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,862	18.8	59.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,706	14.2	44.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,180	6.1	19.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	3,449	5.0	15.9
6	Diabetes mellitus (E10-E14)	2,192	3.2	10.1
7	Cerebrovascular diseases (I60-I69)	2,190	3.2	10.1
8	Chronic lower respiratory diseases (J40-J47)	1,264	1.8	5.8
9	Human immunodeficiency virus (HIV) disease (B20-B24)	1,244	1.8	5.7
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,061	1.5	4.9
11	Septicemia (A40-A41)	890	1.3	4.1
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	736	1.1	3.4
13	Influenza and pneumonia (J09-J18)	695	1.0	3.2
14	Viral hepatitis (B15-B19)	639	0.9	2.9
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	477	0.7	2.2
...	All other causes (Residual)	10,880	15.9	50.2
All races, both sexes, 50-54 years				
...	All causes	110,984	100.0	491.5
1	Malignant neoplasms (C00-C97)	32,014	28.8	141.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	22,403	20.2	99.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,688	9.6	47.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	5,428	4.9	24.0
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,682	4.2	20.7
6	Diabetes mellitus (E10-E14)	3,555	3.2	15.7
7	Cerebrovascular diseases (I60-I69)	3,464	3.1	15.3
8	Chronic lower respiratory diseases (J40-J47)	3,269	2.9	14.5
9	Septicemia (A40-A41)	1,450	1.3	6.4
10	Viral hepatitis (B15-B19)	1,428	1.3	6.3
11	Human immunodeficiency virus (HIV) disease (B20-B24)	1,338	1.2	5.9
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,328	1.2	5.9
13	Influenza and pneumonia (J09-J18)	1,130	1.0	5.0
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	994	0.9	4.4
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	858	0.8	3.8
...	All other causes (Residual)	16,955	15.3	75.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 55-59 years				
...	All causes	148,978	100.0	717.2
1	Malignant neoplasms (C00-C97)	48,954	32.9	235.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	31,457	21.1	151.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,036	6.1	43.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	6,041	4.1	29.1
5	Chronic lower respiratory diseases (J40-J47)	5,893	4.0	28.4
6	Diabetes mellitus (E10-E14)	5,377	3.6	25.9
7	Cerebrovascular diseases (I60-I69)	4,796	3.2	23.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,085	2.7	19.7
9	Viral hepatitis (B15-B19)	2,240	1.5	10.8
10	Septicemia (A40-A41)	2,120	1.4	10.2
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,979	1.3	9.5
12	Influenza and pneumonia (J09-J18)	1,741	1.2	8.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,291	0.9	6.2
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1,094	0.7	5.3
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	699	0.5	3.4
...	All other causes (Residual)	22,175	14.9	106.8
All races, both sexes, 60-64 years				
...	All causes	180,628	100.0	1,014.0
1	Malignant neoplasms (C00-C97)	64,176	35.5	360.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	39,771	22.0	223.3
3	Chronic lower respiratory diseases (J40-J47)	9,319	5.2	52.3
4	Diabetes mellitus (E10-E14)	7,176	4.0	40.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,786	3.8	38.1
6	Cerebrovascular diseases (I60-I69)	6,274	3.5	35.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	5,189	2.9	29.1
8	Septicemia (A40-A41)	2,862	1.6	16.1
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,844	1.6	16.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,786	1.5	15.6
11	Influenza and pneumonia (J09-J18)	2,188	1.2	12.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,715	0.9	9.6
13	Viral hepatitis (B15-B19)	1,637	0.9	9.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	828	0.5	4.6
15	Human immunodeficiency virus (HIV) disease (B20-B24)	699	0.4	3.9
...	All other causes (Residual)	26,378	14.6	148.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 65-69 years				
...	All causes	205,490	100.0	1,470.2
1	Malignant neoplasms (C00-C97)	74,619	36.3	533.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	44,335	21.6	317.2
3	Chronic lower respiratory diseases (J40-J47)	14,632	7.1	104.7
4	Diabetes mellitus (E10-E14)	8,304	4.0	59.4
5	Cerebrovascular diseases (I60-I69)	7,836	3.8	56.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,507	2.7	39.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	3,854	1.9	27.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,560	1.7	25.5
9	Septicemia (A40-A41)	3,305	1.6	23.6
10	Influenza and pneumonia (J09-J18)	2,736	1.3	19.6
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,976	1.0	14.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,848	0.9	13.2
13	Alzheimer's disease (G30)	1,304	0.6	9.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,122	0.5	8.0
15	Parkinson's disease (G20-G21)	901	0.4	6.4
...	All other causes (Residual)	29,651	14.4	212.1
All races, both sexes, 70-74 years				
...	All causes	226,856	100.0	2,266.7
1	Malignant neoplasms (C00-C97)	77,020	34.0	769.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	48,808	21.5	487.7
3	Chronic lower respiratory diseases (J40-J47)	18,943	8.4	189.3
4	Cerebrovascular diseases (I60-I69)	10,332	4.6	103.2
5	Diabetes mellitus (E10-E14)	8,409	3.7	84.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,051	2.2	50.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,439	2.0	44.4
8	Septicemia (A40-A41)	3,708	1.6	37.1
9	Influenza and pneumonia (J09-J18)	3,533	1.6	35.3
10	Alzheimer's disease (G30)	2,996	1.3	29.9
11	Chronic liver disease and cirrhosis (K70,K73-K74)	2,762	1.2	27.6
12	Parkinson's disease (G20-G21)	2,042	0.9	20.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,016	0.9	20.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,536	0.7	15.3
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,391	0.6	13.9
...	All other causes (Residual)	33,870	14.9	338.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 75-79 years				
...	All causes	271,936	100.0	3,630.9
1	Malignant neoplasms (C00-C97)	77,543	28.5	1,035.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	61,317	22.5	818.7
3	Chronic lower respiratory diseases (J40-J47)	22,475	8.3	300.1
4	Cerebrovascular diseases (I60-I69)	14,585	5.4	194.7
5	Diabetes mellitus (E10-E14)	9,074	3.3	121.2
6	Alzheimer's disease (G30)	7,566	2.8	101.0
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,956	2.2	79.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,609	2.1	74.9
9	Influenza and pneumonia (J09-J18)	5,044	1.9	67.3
10	Septicemia (A40-A41)	4,457	1.6	59.5
11	Parkinson's disease (G20-G21)	3,963	1.5	52.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,766	1.0	36.9
13	Chronic liver disease and cirrhosis (K70,K73-K74)	2,229	0.8	29.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,988	0.7	26.5
15	Pneumonitis due to solids and liquids (J69)	1,835	0.7	24.5
...	All other causes (Residual)	45,529	16.7	607.9
All races, both sexes, 80-84 years				
...	All causes	348,492	100.0	6,026.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	85,174	24.4	1,472.8
2	Malignant neoplasms (C00-C97)	76,649	22.0	1,325.4
3	Chronic lower respiratory diseases (J40-J47)	25,832	7.4	446.7
4	Cerebrovascular diseases (I60-I69)	21,549	6.2	372.6
5	Alzheimer's disease (G30)	15,715	4.5	271.7
6	Diabetes mellitus (E10-E14)	10,272	2.9	177.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,347	2.4	144.3
8	Influenza and pneumonia (J09-J18)	7,996	2.3	138.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,648	2.2	132.2
10	Parkinson's disease (G20-G21)	6,154	1.8	106.4
11	Septicemia (A40-A41)	5,346	1.5	92.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,102	1.2	70.9
13	Pneumonitis due to solids and liquids (J69)	3,021	0.9	52.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,631	0.8	45.5
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,665	0.5	28.8
...	All other causes (Residual)	66,391	19.1	1,148.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 85-89 years				
...	All causes	384,262	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	105,723	27.5	---
2	Malignant neoplasms (C00-C97)	60,075	15.6	---
3	Cerebrovascular diseases (I60-I69)	25,926	6.7	---
4	Alzheimer's disease (G30)	23,592	6.1	---
5	Chronic lower respiratory diseases (J40-J47)	23,112	6.0	---
6	Influenza and pneumonia (J09-J18)	10,238	2.7	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,393	2.4	---
8	Diabetes mellitus (E10-E14)	9,387	2.4	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,211	2.1	---
10	Parkinson's disease (G20-G21)	6,162	1.6	---
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,555	1.4	---
12	Septicemia (A40-A41)	5,493	1.4	---
13	Pneumonitis due to solids and liquids (J69)	3,987	1.0	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,784	0.7	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1,776	0.5	---
...	All other causes (Residual)	82,848	21.6	---
All races, both sexes, 90-94 years				
...	All causes	284,273	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	87,292	30.7	---
2	Malignant neoplasms (C00-C97)	29,096	10.2	---
3	Alzheimer's disease (G30)	20,850	7.3	---
4	Cerebrovascular diseases (I60-I69)	19,939	7.0	---
5	Chronic lower respiratory diseases (J40-J47)	12,826	4.5	---
6	Influenza and pneumonia (J09-J18)	8,780	3.1	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,101	2.5	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,801	2.0	---
9	Diabetes mellitus (E10-E14)	5,493	1.9	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,920	1.7	---
11	Pneumonitis due to solids and liquids (J69)	3,396	1.2	---
12	Septicemia (A40-A41)	3,371	1.2	---
13	Parkinson's disease (G20-G21)	3,196	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,599	0.6	---
15	Atherosclerosis (I70)	1,476	0.5	---
...	All other causes (Residual)	69,137	24.3	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 95-99 years				
...	All causes	112,809	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	36,950	32.8	---
2	Alzheimer's disease (G30)	8,873	7.9	---
3	Cerebrovascular diseases (I60-I69)	7,576	6.7	---
4	Malignant neoplasms (C00-C97)	7,523	6.7	---
5	Influenza and pneumonia (J09-J18)	3,996	3.5	---
6	Chronic lower respiratory diseases (J40-J47)	3,955	3.5	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,809	2.5	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,463	2.2	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,084	1.8	---
10	Diabetes mellitus (E10-E14)	1,669	1.5	---
11	Pneumonitis due to solids and liquids (J69)	1,344	1.2	---
12	Septicemia (A40-A41)	1,140	1.0	---
13	Atherosclerosis (I70)	769	0.7	---
14	Parkinson's disease (G20-G21)	712	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	510	0.5	---
...	All other causes (Residual)	30,436	27.0	---
All races, both sexes, 100 years and over				
...	All causes	23,963	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,241	34.4	---
2	Alzheimer's disease (G30)	1,794	7.5	---
3	Cerebrovascular diseases (I60-I69)	1,384	5.8	---
4	Influenza and pneumonia (J09-J18)	1,032	4.3	---
5	Malignant neoplasms (C00-C97)	972	4.1	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	642	2.7	---
7	Chronic lower respiratory diseases (J40-J47)	600	2.5	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	534	2.2	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	388	1.6	---
10	Diabetes mellitus (E10-E14)	273	1.1	---
10	Pneumonitis due to solids and liquids (J69)	273	1.1	---
12	Atherosclerosis (I70)	238	1.0	---
13	Septicemia (A40-A41)	202	0.8	---
14	Anemias (D50-D64)	88	0.4	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	76	0.3	---
...	All other causes (Residual)	7,226	30.2	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, all ages				
...	All causes	1,273,722	100.0	824.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	312,491	24.5	202.3
2	Malignant neoplasms (C00-C97)	305,670	24.0	197.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	80,010	6.3	51.8
4	Chronic lower respiratory diseases (J40-J47)	67,673	5.3	43.8
5	Cerebrovascular diseases (I60-I69)	52,638	4.1	34.1
6	Diabetes mellitus (E10-E14)	38,584	3.0	25.0
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	31,780	2.5	20.6
8	Alzheimer's disease (G30)	25,653	2.0	16.6
9	Influenza and pneumonia (J09-J18)	24,013	1.9	15.5
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22,731	1.8	14.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	22,623	1.8	14.6
12	Septicemia (A40-A41)	16,789	1.3	10.9
13	Parkinson's disease (G20-G21)	14,121	1.1	9.1
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	13,208	1.0	8.5
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	11,859	0.9	7.7
...	All other causes (Residual)	233,879	18.4	151.4
All races, male, 1-4 years				
...	All causes	2,395	100.0	29.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	806	33.7	9.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	275	11.5	3.4
3	Malignant neoplasms (C00-C97)	218	9.1	2.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	197	8.2	2.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	80	3.3	1.0
6	Influenza and pneumonia (J09-J18)	55	2.3	0.7
7	Cerebrovascular diseases (I60-I69)	36	1.5	0.4
8	Chronic lower respiratory diseases (J40-J47)	35	1.5	0.4
9	Septicemia (A40-A41)	29	1.2	0.4
10	Certain conditions originating in the perinatal period (P00-P96)	27	1.1	0.3
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	0.8	0.2
12	Acute bronchitis and bronchiolitis (J20-J21)	13	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.3	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.3	*
15	Anemias (D50-D64)	5	0.2	*
15	Pneumonitis due to solids and liquids (J69)	5	0.2	*
...	All other causes (Residual)	580	24.2	7.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 5-9 years				
...	All causes	1,320	100.0	12.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	457	34.6	4.4
2	Malignant neoplasms (C00-C97)	254	19.2	2.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	83	6.3	0.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	75	5.7	0.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	33	2.5	0.3
6	Chronic lower respiratory diseases (J40-J47)	31	2.3	0.3
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	1.7	0.2
8	Influenza and pneumonia (J09-J18)	21	1.6	0.2
9	Septicemia (A40-A41)	17	1.3	*
10	Cerebrovascular diseases (I60-I69)	15	1.1	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	8	0.6	*
12	Anemias (D50-D64)	7	0.5	*
13	Diabetes mellitus (E10-E14)	6	0.5	*
13	Certain conditions originating in the perinatal period (P00-P96)	6	0.5	*
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5	0.4	*
...	All other causes (Residual)	279	21.1	2.7
All races, male, 10-14 years				
...	All causes	1,716	100.0	16.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	525	30.6	5.0
2	Malignant neoplasms (C00-C97)	259	15.1	2.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	221	12.9	2.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	106	6.2	1.0
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	87	5.1	0.8
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	63	3.7	0.6
7	Chronic lower respiratory diseases (J40-J47)	31	1.8	0.3
8	Cerebrovascular diseases (I60-I69)	26	1.5	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	25	1.5	0.2
10	Influenza and pneumonia (J09-J18)	19	1.1	*
11	Septicemia (A40-A41)	15	0.9	*
12	Diabetes mellitus (E10-E14)	12	0.7	*
13	Anemias (D50-D64)	9	0.5	*
14	Certain conditions originating in the perinatal period (P00-P96)	5	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.2	*
...	All other causes (Residual)	309	18.0	2.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 15-19 years				
...	All causes	7,224	100.0	65.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,814	39.0	25.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,403	19.4	12.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,372	19.0	12.5
4	Malignant neoplasms (C00-C97)	379	5.2	3.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	204	2.8	1.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	117	1.6	1.1
7	Cerebrovascular diseases (I60-I69)	35	0.5	0.3
7	Chronic lower respiratory diseases (J40-J47)	35	0.5	0.3
7	Legal intervention (Y35,Y89.0)	35	0.5	0.3
10	Influenza and pneumonia (J09-J18)	26	0.4	0.2
11	Diabetes mellitus (E10-E14)	20	0.3	0.2
12	Septicemia (A40-A41)	17	0.2	*
12	Anemias (D50-D64)	17	0.2	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	0.2	*
15	Pneumonitis due to solids and liquids (J69)	10	0.1	*
...	All other causes (Residual)	724	10.0	6.6
All races, male, 20-24 years				
...	All causes	14,232	100.0	123.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,035	42.4	52.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,592	18.2	22.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,546	17.9	22.0
4	Malignant neoplasms (C00-C97)	545	3.8	4.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	440	3.1	3.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	133	0.9	1.2
7	Diabetes mellitus (E10-E14)	86	0.6	0.7
8	Legal intervention (Y35,Y89.0)	77	0.5	0.7
9	Human immunodeficiency virus (HIV) disease (B20-B24)	73	0.5	0.6
9	Cerebrovascular diseases (I60-I69)	73	0.5	0.6
11	Influenza and pneumonia (J09-J18)	64	0.4	0.6
12	Chronic lower respiratory diseases (J40-J47)	63	0.4	0.5
13	Septicemia (A40-A41)	37	0.3	0.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	33	0.2	0.3
14	Anemias (D50-D64)	33	0.2	0.3
...	All other causes (Residual)	1,402	9.9	12.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 25-29 years				
...	All causes	14,951	100.0	137.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,959	39.9	54.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,597	17.4	23.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,040	13.6	18.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	797	5.3	7.3
5	Malignant neoplasms (C00-C97)	753	5.0	6.9
6	Human immunodeficiency virus (HIV) disease (B20-B24)	184	1.2	1.7
7	Diabetes mellitus (E10-E14)	136	0.9	1.3
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	129	0.9	1.2
9	Chronic liver disease and cirrhosis (K70,K73-K74)	108	0.7	1.0
10	Cerebrovascular diseases (I60-I69)	100	0.7	0.9
11	Septicemia (A40-A41)	82	0.5	0.8
12	Legal intervention (Y35,Y89.0)	81	0.5	0.7
13	Influenza and pneumonia (J09-J18)	80	0.5	0.7
14	Chronic lower respiratory diseases (J40-J47)	79	0.5	0.7
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	46	0.3	0.4
...	All other causes (Residual)	1,780	11.9	16.4
All races, male, 30-34 years				
...	All causes	15,832	100.0	150.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,716	36.1	54.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,388	15.1	22.8
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,611	10.2	15.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,417	9.0	13.5
5	Malignant neoplasms (C00-C97)	1,049	6.6	10.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	272	1.7	2.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	271	1.7	2.6
8	Diabetes mellitus (E10-E14)	223	1.4	2.1
9	Cerebrovascular diseases (I60-I69)	197	1.2	1.9
10	Influenza and pneumonia (J09-J18)	125	0.8	1.2
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	115	0.7	1.1
12	Septicemia (A40-A41)	99	0.6	0.9
13	Chronic lower respiratory diseases (J40-J47)	91	0.6	0.9
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	76	0.5	0.7
15	Legal intervention (Y35,Y89.0)	60	0.4	0.6
...	All other causes (Residual)	2,122	13.4	20.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 35-39 years				
...	All causes	17,368	100.0	178.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,030	29.0	51.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,473	14.2	25.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,404	13.8	24.7
4	Malignant neoplasms (C00-C97)	1,583	9.1	16.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,165	6.7	12.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	491	2.8	5.1
7	Diabetes mellitus (E10-E14)	398	2.3	4.1
8	Cerebrovascular diseases (I60-I69)	334	1.9	3.4
9	Human immunodeficiency virus (HIV) disease (B20-B24)	304	1.8	3.1
10	Septicemia (A40-A41)	144	0.8	1.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	132	0.8	1.4
12	Influenza and pneumonia (J09-J18)	125	0.7	1.3
13	Chronic lower respiratory diseases (J40-J47)	101	0.6	1.0
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	100	0.6	1.0
15	Aortic aneurysm and dissection (I71)	84	0.5	0.9
...	All other causes (Residual)	2,500	14.4	25.7
All races, male, 40-44 years				
...	All causes	25,561	100.0	244.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,443	21.3	52.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,758	18.6	45.5
3	Malignant neoplasms (C00-C97)	3,266	12.8	31.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,781	10.9	26.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,110	4.3	10.6
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	973	3.8	9.3
7	Diabetes mellitus (E10-E14)	758	3.0	7.2
8	Cerebrovascular diseases (I60-I69)	628	2.5	6.0
9	Human immunodeficiency virus (HIV) disease (B20-B24)	549	2.1	5.2
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	244	1.0	2.3
11	Septicemia (A40-A41)	242	0.9	2.3
12	Chronic lower respiratory diseases (J40-J47)	240	0.9	2.3
13	Influenza and pneumonia (J09-J18)	233	0.9	2.2
14	Aortic aneurysm and dissection (I71)	141	0.6	1.3
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	137	0.5	1.3
...	All other causes (Residual)	4,058	15.9	38.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 45-49 years				
...	All causes	41,387	100.0	385.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,970	21.7	83.6
2	Malignant neoplasms (C00-C97)	7,527	18.2	70.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,493	15.7	60.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,086	7.5	28.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,264	5.5	21.1
6	Diabetes mellitus (E10-E14)	1,382	3.3	12.9
7	Cerebrovascular diseases (I60-I69)	1,186	2.9	11.1
8	Human immunodeficiency virus (HIV) disease (B20-B24)	891	2.2	8.3
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	777	1.9	7.2
10	Chronic lower respiratory diseases (J40-J47)	573	1.4	5.3
11	Septicemia (A40-A41)	476	1.2	4.4
12	Viral hepatitis (B15-B19)	455	1.1	4.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	432	1.0	4.0
14	Influenza and pneumonia (J09-J18)	377	0.9	3.5
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	291	0.7	2.7
...	All other causes (Residual)	6,207	15.0	57.9
All races, male, 50-54 years				
...	All causes	67,827	100.0	612.1
1	Malignant neoplasms (C00-C97)	16,476	24.3	148.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,957	23.5	144.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,098	10.5	64.1
4	Chronic liver disease and cirrhosis (K70,K73-K74)	3,715	5.5	33.5
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,492	5.1	31.5
6	Diabetes mellitus (E10-E14)	2,222	3.3	20.1
7	Cerebrovascular diseases (I60-I69)	1,915	2.8	17.3
8	Chronic lower respiratory diseases (J40-J47)	1,560	2.3	14.1
9	Viral hepatitis (B15-B19)	1,017	1.5	9.2
10	Human immunodeficiency virus (HIV) disease (B20-B24)	1,003	1.5	9.1
11	Septicemia (A40-A41)	774	1.1	7.0
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	768	1.1	6.9
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	753	1.1	6.8
14	Influenza and pneumonia (J09-J18)	643	0.9	5.8
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	577	0.9	5.2
...	All other causes (Residual)	9,857	14.5	89.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 55-59 years				
...	All causes	91,743	100.0	911.2
1	Malignant neoplasms (C00-C97)	26,871	29.3	266.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	22,221	24.2	220.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,123	6.7	60.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	4,301	4.7	42.7
5	Diabetes mellitus (E10-E14)	3,311	3.6	32.9
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,116	3.4	30.9
7	Chronic lower respiratory diseases (J40-J47)	2,984	3.3	29.6
8	Cerebrovascular diseases (I60-I69)	2,810	3.1	27.9
9	Viral hepatitis (B15-B19)	1,575	1.7	15.6
10	Septicemia (A40-A41)	1,164	1.3	11.6
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,147	1.3	11.4
12	Influenza and pneumonia (J09-J18)	1,007	1.1	10.0
13	Human immunodeficiency virus (HIV) disease (B20-B24)	840	0.9	8.3
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	810	0.9	8.0
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	511	0.6	5.1
...	All other causes (Residual)	12,952	14.1	128.6
All races, male, 60-64 years				
...	All causes	109,208	100.0	1,279.6
1	Malignant neoplasms (C00-C97)	35,799	32.8	419.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	27,336	25.0	320.3
3	Chronic lower respiratory diseases (J40-J47)	4,897	4.5	57.4
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,681	4.3	54.8
5	Diabetes mellitus (E10-E14)	4,323	4.0	50.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	3,638	3.3	42.6
7	Cerebrovascular diseases (I60-I69)	3,600	3.3	42.2
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,216	2.0	26.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,553	1.4	18.2
10	Septicemia (A40-A41)	1,539	1.4	18.0
11	Influenza and pneumonia (J09-J18)	1,294	1.2	15.2
12	Viral hepatitis (B15-B19)	1,179	1.1	13.8
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,060	1.0	12.4
14	Human immunodeficiency virus (HIV) disease (B20-B24)	531	0.5	6.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	487	0.4	5.7
...	All other causes (Residual)	15,075	13.8	176.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 65-69 years				
...	All causes	118,915	100.0	1,799.9
1	Malignant neoplasms (C00-C97)	41,329	34.8	625.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	29,105	24.5	440.5
3	Chronic lower respiratory diseases (J40-J47)	7,451	6.3	112.8
4	Diabetes mellitus (E10-E14)	4,824	4.1	73.0
5	Cerebrovascular diseases (I60-I69)	4,288	3.6	64.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,587	3.0	54.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,475	2.1	37.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,940	1.6	29.4
9	Septicemia (A40-A41)	1,746	1.5	26.4
10	Influenza and pneumonia (J09-J18)	1,559	1.3	23.6
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,538	1.3	23.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,005	0.8	15.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	641	0.5	9.7
14	Parkinson's disease (G20-G21)	608	0.5	9.2
15	Aortic aneurysm and dissection (I71)	590	0.5	8.9
...	All other causes (Residual)	16,229	13.6	245.6
All races, male, 70-74 years				
...	All causes	126,020	100.0	2,741.9
1	Malignant neoplasms (C00-C97)	42,324	33.6	920.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	30,008	23.8	652.9
3	Chronic lower respiratory diseases (J40-J47)	9,450	7.5	205.6
4	Cerebrovascular diseases (I60-I69)	5,312	4.2	115.6
5	Diabetes mellitus (E10-E14)	4,665	3.7	101.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,118	2.5	67.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,373	1.9	51.6
8	Influenza and pneumonia (J09-J18)	1,948	1.5	42.4
9	Septicemia (A40-A41)	1,816	1.4	39.5
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,633	1.3	35.5
11	Parkinson's disease (G20-G21)	1,346	1.1	29.3
12	Alzheimer's disease (G30)	1,291	1.0	28.1
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,162	0.9	25.3
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,064	0.8	23.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	935	0.7	20.3
...	All other causes (Residual)	17,575	13.9	382.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 75-79 years				
...	All causes	142,424	100.0	4,327.1
1	Malignant neoplasms (C00-C97)	41,802	29.4	1,270.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	34,752	24.4	1,055.8
3	Chronic lower respiratory diseases (J40-J47)	11,111	7.8	337.6
4	Cerebrovascular diseases (I60-I69)	6,787	4.8	206.2
5	Diabetes mellitus (E10-E14)	4,718	3.3	143.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,303	2.3	100.4
7	Alzheimer's disease (G30)	2,973	2.1	90.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,937	2.1	89.2
9	Influenza and pneumonia (J09-J18)	2,658	1.9	80.8
10	Parkinson's disease (G20-G21)	2,543	1.8	77.3
11	Septicemia (A40-A41)	2,161	1.5	65.7
12	Chronic liver disease and cirrhosis (K70,K73-K74)	1,218	0.9	37.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,213	0.9	36.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,121	0.8	34.1
15	Pneumonitis due to solids and liquids (J69)	1,078	0.8	32.8
...	All other causes (Residual)	22,049	15.5	669.9
All races, male, 80-84 years				
...	All causes	168,257	100.0	7,139.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	43,649	25.9	1,852.1
2	Malignant neoplasms (C00-C97)	39,959	23.7	1,695.6
3	Chronic lower respiratory diseases (J40-J47)	12,254	7.3	520.0
4	Cerebrovascular diseases (I60-I69)	8,899	5.3	377.6
5	Alzheimer's disease (G30)	5,732	3.4	243.2
6	Diabetes mellitus (E10-E14)	4,988	3.0	211.7
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,270	2.5	181.2
8	Influenza and pneumonia (J09-J18)	4,023	2.4	170.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,894	2.3	165.2
10	Parkinson's disease (G20-G21)	3,887	2.3	164.9
11	Septicemia (A40-A41)	2,404	1.4	102.0
12	Pneumonitis due to solids and liquids (J69)	1,713	1.0	72.7
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,666	1.0	70.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,425	0.8	60.5
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	900	0.5	38.2
...	All other causes (Residual)	28,594	17.0	1,213.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 85-89 years				
...	All causes	161,326	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	46,584	28.9	---
2	Malignant neoplasms (C00-C97)	29,169	18.1	---
3	Chronic lower respiratory diseases (J40-J47)	10,226	6.3	---
4	Cerebrovascular diseases (I60-I69)	8,928	5.5	---
5	Alzheimer's disease (G30)	7,406	4.6	---
6	Influenza and pneumonia (J09-J18)	4,631	2.9	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,245	2.6	---
8	Diabetes mellitus (E10-E14)	3,989	2.5	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,848	2.4	---
10	Parkinson's disease (G20-G21)	3,446	2.1	---
11	Septicemia (A40-A41)	2,290	1.4	---
12	Pneumonitis due to solids and liquids (J69)	2,120	1.3	---
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,878	1.2	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,352	0.8	---
15	Aortic aneurysm and dissection (I71)	715	0.4	---
...	All other causes (Residual)	30,499	18.9	---
All races, male, 90-94 years				
...	All causes	99,249	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	31,852	32.1	---
2	Malignant neoplasms (C00-C97)	12,991	13.1	---
3	Cerebrovascular diseases (I60-I69)	5,544	5.6	---
4	Alzheimer's disease (G30)	5,425	5.5	---
5	Chronic lower respiratory diseases (J40-J47)	5,003	5.0	---
6	Influenza and pneumonia (J09-J18)	3,529	3.6	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,721	2.7	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,436	2.5	---
9	Diabetes mellitus (E10-E14)	2,017	2.0	---
10	Pneumonitis due to solids and liquids (J69)	1,689	1.7	---
11	Parkinson's disease (G20-G21)	1,613	1.6	---
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,382	1.4	---
13	Septicemia (A40-A41)	1,271	1.3	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	651	0.7	---
15	Atherosclerosis (I70)	504	0.5	---
...	All other causes (Residual)	20,621	20.8	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 95-99 years				
...	All causes	29,435	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,054	34.2	---
2	Malignant neoplasms (C00-C97)	2,815	9.6	---
3	Cerebrovascular diseases (I60-I69)	1,672	5.7	---
4	Alzheimer's disease (G30)	1,645	5.6	---
5	Chronic lower respiratory diseases (J40-J47)	1,307	4.4	---
6	Influenza and pneumonia (J09-J18)	1,277	4.3	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	801	2.7	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	720	2.4	---
9	Pneumonitis due to solids and liquids (J69)	545	1.9	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	533	1.8	---
11	Diabetes mellitus (E10-E14)	459	1.6	---
12	Septicemia (A40-A41)	317	1.1	---
13	Parkinson's disease (G20-G21)	274	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	191	0.6	---
15	Atherosclerosis (I70)	182	0.6	---
...	All other causes (Residual)	6,643	22.6	---
All races, male, 100 years and over				
...	All causes	4,093	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,515	37.0	---
2	Malignant neoplasms (C00-C97)	259	6.3	---
3	Influenza and pneumonia (J09-J18)	222	5.4	---
4	Alzheimer's disease (G30)	200	4.9	---
5	Cerebrovascular diseases (I60-I69)	198	4.8	---
6	Chronic lower respiratory diseases (J40-J47)	136	3.3	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	102	2.5	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	99	2.4	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	94	2.3	---
10	Pneumonitis due to solids and liquids (J69)	87	2.1	---
11	Septicemia (A40-A41)	43	1.1	---
12	Diabetes mellitus (E10-E14)	42	1.0	---
13	Atherosclerosis (I70)	39	1.0	---
14	Parkinson's disease (G20-G21)	22	0.5	---
15	Anemias (D50-D64)	18	0.4	---
...	All other causes (Residual)	1,017	24.8	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, all ages				
...	All causes	1,269,557	100.0	796.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	287,220	22.6	180.2
2	Malignant neoplasms (C00-C97)	276,953	21.8	173.7
3	Cerebrovascular diseases (I60-I69)	75,908	6.0	47.6
4	Chronic lower respiratory diseases (J40-J47)	75,816	6.0	47.6
5	Alzheimer's disease (G30)	57,984	4.6	36.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	47,782	3.8	30.0
7	Diabetes mellitus (E10-E14)	35,348	2.8	22.2
8	Influenza and pneumonia (J09-J18)	26,623	2.1	16.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22,891	1.8	14.4
10	Septicemia (A40-A41)	19,053	1.5	12.0
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	17,256	1.4	10.8
12	Chronic liver disease and cirrhosis (K70,K73-K74)	12,356	1.0	7.8
13	Parkinson's disease (G20-G21)	9,697	0.8	6.1
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8,820	0.7	5.5
15	Pneumonitis due to solids and liquids (J69)	8,184	0.6	5.1
...	All other causes (Residual)	287,666	22.7	180.4
All races, female, 1-4 years				
...	All causes	1,823	100.0	23.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	547	30.0	7.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	226	12.4	2.9
3	Malignant neoplasms (C00-C97)	174	9.5	2.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	142	7.8	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	74	4.1	0.9
6	Influenza and pneumonia (J09-J18)	38	2.1	0.5
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	35	1.9	0.4
8	Septicemia (A40-A41)	33	1.8	0.4
9	Cerebrovascular diseases (I60-I69)	20	1.1	0.3
10	Chronic lower respiratory diseases (J40-J47)	16	0.9	*
11	Certain conditions originating in the perinatal period (P00-P96)	15	0.8	*
12	Anemias (D50-D64)	10	0.5	*
13	Meningitis (G00,G03)	6	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.3	*
15	Acute bronchitis and bronchiolitis (J20-J21)	4	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.2	*
...	All other causes (Residual)	474	26.0	6.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 5-9 years				
...	All causes	1,014	100.0	10.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	286	28.2	2.9
2	Malignant neoplasms (C00-C97)	186	18.3	1.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	84	8.3	0.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	63	6.2	0.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	34	3.4	0.3
6	Chronic lower respiratory diseases (J40-J47)	32	3.2	0.3
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	24	2.4	0.2
8	Influenza and pneumonia (J09-J18)	23	2.3	0.2
9	Cerebrovascular diseases (I60-I69)	19	1.9	*
10	Certain conditions originating in the perinatal period (P00-P96)	12	1.2	*
11	Septicemia (A40-A41)	9	0.9	*
12	Diabetes mellitus (E10-E14)	6	0.6	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.6	*
14	Meningitis (G00,G03)	5	0.5	*
15	Anemias (D50-D64)	3	0.3	*
...	All other causes (Residual)	222	21.9	2.2
All races, female, 10-14 years				
...	All causes	1,150	100.0	11.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	282	24.5	2.8
2	Malignant neoplasms (C00-C97)	213	18.5	2.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	85	7.4	0.8
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	73	6.3	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	67	5.8	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	45	3.9	0.4
7	Cerebrovascular diseases (I60-I69)	25	2.2	0.2
7	Chronic lower respiratory diseases (J40-J47)	25	2.2	0.2
9	Influenza and pneumonia (J09-J18)	22	1.9	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	15	1.3	*
11	Septicemia (A40-A41)	11	1.0	*
11	Certain conditions originating in the perinatal period (P00-P96)	11	1.0	*
13	Diabetes mellitus (E10-E14)	10	0.9	*
14	Anemias (D50-D64)	9	0.8	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.3	*
...	All other causes (Residual)	253	22.0	2.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 15-19 years				
...	All causes	2,850	100.0	27.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,181	41.4	11.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	410	14.4	3.9
3	Malignant neoplasms (C00-C97)	249	8.7	2.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	211	7.4	2.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	98	3.4	0.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	84	2.9	0.8
7	Pregnancy, childbirth and the puerperium (O00-O99)	37	1.3	0.4
8	Cerebrovascular diseases (I60-I69)	28	1.0	0.3
9	Diabetes mellitus (E10-E14)	24	0.8	0.2
10	Septicemia (A40-A41)	23	0.8	0.2
11	Influenza and pneumonia (J09-J18)	21	0.7	0.2
12	Chronic lower respiratory diseases (J40-J47)	19	0.7	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	0.5	*
14	Anemias (D50-D64)	13	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	0.4	*
...	All other causes (Residual)	428	15.0	4.1
All races, female, 20-24 years				
...	All causes	4,876	100.0	44.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,878	38.5	17.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	544	11.2	4.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	408	8.4	3.7
4	Malignant neoplasms (C00-C97)	401	8.2	3.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	214	4.4	1.9
6	Pregnancy, childbirth and the puerperium (O00-O99)	132	2.7	1.2
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	89	1.8	0.8
8	Diabetes mellitus (E10-E14)	66	1.4	0.6
9	Cerebrovascular diseases (I60-I69)	47	1.0	0.4
10	Septicemia (A40-A41)	43	0.9	0.4
11	Influenza and pneumonia (J09-J18)	36	0.7	0.3
12	Human immunodeficiency virus (HIV) disease (B20-B24)	32	0.7	0.3
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	28	0.6	0.3
14	Anemias (D50-D64)	26	0.5	0.2
14	Chronic lower respiratory diseases (J40-J47)	26	0.5	0.2
...	All other causes (Residual)	906	18.6	8.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 25-29 years				
...	All causes	6,041	100.0	57.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,016	33.4	19.1
2	Malignant neoplasms (C00-C97)	631	10.4	6.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	587	9.7	5.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	380	6.3	3.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	348	5.8	3.3
6	Pregnancy, childbirth and the puerperium (O00-O99)	191	3.2	1.8
7	Diabetes mellitus (E10-E14)	110	1.8	1.0
8	Cerebrovascular diseases (I60-I69)	85	1.4	0.8
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	72	1.2	0.7
10	Septicemia (A40-A41)	66	1.1	0.6
11	Human immunodeficiency virus (HIV) disease (B20-B24)	63	1.0	0.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	63	1.0	0.6
13	Influenza and pneumonia (J09-J18)	61	1.0	0.6
14	Chronic lower respiratory diseases (J40-J47)	53	0.9	0.5
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	36	0.6	0.3
...	All other causes (Residual)	1,279	21.2	12.1
All races, female, 30-34 years				
...	All causes	7,767	100.0	74.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,160	27.8	20.7
2	Malignant neoplasms (C00-C97)	1,241	16.0	11.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	644	8.3	6.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	637	8.2	6.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	343	4.4	3.3
6	Pregnancy, childbirth and the puerperium (O00-O99)	199	2.6	1.9
7	Diabetes mellitus (E10-E14)	177	2.3	1.7
8	Chronic liver disease and cirrhosis (K70,K73-K74)	155	2.0	1.5
9	Cerebrovascular diseases (I60-I69)	153	2.0	1.5
10	Human immunodeficiency virus (HIV) disease (B20-B24)	133	1.7	1.3
11	Influenza and pneumonia (J09-J18)	90	1.2	0.9
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	85	1.1	0.8
13	Septicemia (A40-A41)	78	1.0	0.7
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	74	1.0	0.7
15	Chronic lower respiratory diseases (J40-J47)	66	0.8	0.6
...	All other causes (Residual)	1,532	19.7	14.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 35-39 years				
...	All causes	10,004	100.0	102.4
1	Malignant neoplasms (C00-C97)	2,134	21.3	21.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,051	20.5	21.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,166	11.7	11.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	706	7.1	7.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	277	2.8	2.8
6	Chronic liver disease and cirrhosis (K70,K73-K74)	266	2.7	2.7
7	Diabetes mellitus (E10-E14)	263	2.6	2.7
8	Cerebrovascular diseases (I60-I69)	251	2.5	2.6
9	Human immunodeficiency virus (HIV) disease (B20-B24)	216	2.2	2.2
10	Pregnancy, childbirth and the puerperium (O00-O99)	170	1.7	1.7
11	Influenza and pneumonia (J09-J18)	122	1.2	1.2
12	Chronic lower respiratory diseases (J40-J47)	120	1.2	1.2
13	Septicemia (A40-A41)	117	1.2	1.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	109	1.1	1.1
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	84	0.8	0.9
...	All other causes (Residual)	1,952	19.5	20.0
All races, female, 40-44 years				
...	All causes	16,229	100.0	153.5
1	Malignant neoplasms (C00-C97)	4,354	26.8	41.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,510	15.5	23.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,092	12.9	19.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	867	5.3	8.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	602	3.7	5.7
6	Cerebrovascular diseases (I60-I69)	517	3.2	4.9
7	Diabetes mellitus (E10-E14)	448	2.8	4.2
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	290	1.8	2.7
9	Chronic lower respiratory diseases (J40-J47)	282	1.7	2.7
10	Human immunodeficiency virus (HIV) disease (B20-B24)	276	1.7	2.6
11	Septicemia (A40-A41)	254	1.6	2.4
12	Influenza and pneumonia (J09-J18)	205	1.3	1.9
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	164	1.0	1.6
14	Pregnancy, childbirth and the puerperium (O00-O99)	103	0.6	1.0
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	90	0.6	0.9
...	All other causes (Residual)	3,175	19.6	30.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 45-49 years				
...	All causes	27,092	100.0	247.1
1	Malignant neoplasms (C00-C97)	8,487	31.3	77.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,892	14.4	35.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,213	11.9	29.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	1,185	4.4	10.8
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,094	4.0	10.0
6	Cerebrovascular diseases (I60-I69)	1,004	3.7	9.2
7	Diabetes mellitus (E10-E14)	810	3.0	7.4
8	Chronic lower respiratory diseases (J40-J47)	691	2.6	6.3
9	Septicemia (A40-A41)	414	1.5	3.8
10	Human immunodeficiency virus (HIV) disease (B20-B24)	353	1.3	3.2
11	Influenza and pneumonia (J09-J18)	318	1.2	2.9
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	304	1.1	2.8
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	284	1.0	2.6
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	186	0.7	1.7
15	Viral hepatitis (B15-B19)	184	0.7	1.7
...	All other causes (Residual)	4,673	17.2	42.6
All races, female, 50-54 years				
...	All causes	43,157	100.0	375.3
1	Malignant neoplasms (C00-C97)	15,538	36.0	135.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,446	14.9	56.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,590	8.3	31.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	1,713	4.0	14.9
5	Chronic lower respiratory diseases (J40-J47)	1,709	4.0	14.9
6	Cerebrovascular diseases (I60-I69)	1,549	3.6	13.5
7	Diabetes mellitus (E10-E14)	1,333	3.1	11.6
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,190	2.8	10.3
9	Septicemia (A40-A41)	676	1.6	5.9
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	560	1.3	4.9
11	Influenza and pneumonia (J09-J18)	487	1.1	4.2
12	Viral hepatitis (B15-B19)	411	1.0	3.6
13	Human immunodeficiency virus (HIV) disease (B20-B24)	335	0.8	2.9
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	281	0.7	2.4
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	241	0.6	2.1
...	All other causes (Residual)	7,098	16.4	61.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 55-59 years				
...	All causes	57,235	100.0	534.7
1	Malignant neoplasms (C00-C97)	22,083	38.6	206.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,236	16.1	86.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,913	5.1	27.2
4	Chronic lower respiratory diseases (J40-J47)	2,909	5.1	27.2
5	Diabetes mellitus (E10-E14)	2,066	3.6	19.3
6	Cerebrovascular diseases (I60-I69)	1,986	3.5	18.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,740	3.0	16.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	969	1.7	9.1
9	Septicemia (A40-A41)	956	1.7	8.9
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	832	1.5	7.8
11	Influenza and pneumonia (J09-J18)	734	1.3	6.9
12	Viral hepatitis (B15-B19)	665	1.2	6.2
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	481	0.8	4.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	313	0.5	2.9
15	Human immunodeficiency virus (HIV) disease (B20-B24)	254	0.4	2.4
...	All other causes (Residual)	9,098	15.9	85.0
All races, female, 60-64 years				
...	All causes	71,420	100.0	769.7
1	Malignant neoplasms (C00-C97)	28,377	39.7	305.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,435	17.4	134.0
3	Chronic lower respiratory diseases (J40-J47)	4,422	6.2	47.7
4	Diabetes mellitus (E10-E14)	2,853	4.0	30.7
5	Cerebrovascular diseases (I60-I69)	2,674	3.7	28.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,105	2.9	22.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,551	2.2	16.7
8	Septicemia (A40-A41)	1,323	1.9	14.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,233	1.7	13.3
10	Influenza and pneumonia (J09-J18)	894	1.3	9.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	655	0.9	7.1
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	628	0.9	6.8
13	Viral hepatitis (B15-B19)	458	0.6	4.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	341	0.5	3.7
15	Alzheimer's disease (G30)	336	0.5	3.6
...	All other causes (Residual)	11,135	15.6	120.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 65-69 years				
...	All causes	86,575	100.0	1,174.6
1	Malignant neoplasms (C00-C97)	33,290	38.5	451.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,230	17.6	206.6
3	Chronic lower respiratory diseases (J40-J47)	7,181	8.3	97.4
4	Cerebrovascular diseases (I60-I69)	3,548	4.1	48.1
5	Diabetes mellitus (E10-E14)	3,480	4.0	47.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,920	2.2	26.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,620	1.9	22.0
8	Septicemia (A40-A41)	1,559	1.8	21.2
9	Chronic liver disease and cirrhosis (K70,K73-K74)	1,379	1.6	18.7
10	Influenza and pneumonia (J09-J18)	1,177	1.4	16.0
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	843	1.0	11.4
12	Alzheimer's disease (G30)	741	0.9	10.1
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	481	0.6	6.5
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	438	0.5	5.9
15	Pneumonitis due to solids and liquids (J69)	328	0.4	4.5
...	All other causes (Residual)	13,360	15.4	181.3
All races, female, 70-74 years				
...	All causes	100,836	100.0	1,863.2
1	Malignant neoplasms (C00-C97)	34,696	34.4	641.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	18,800	18.6	347.4
3	Chronic lower respiratory diseases (J40-J47)	9,493	9.4	175.4
4	Cerebrovascular diseases (I60-I69)	5,020	5.0	92.8
5	Diabetes mellitus (E10-E14)	3,744	3.7	69.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,066	2.0	38.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,933	1.9	35.7
8	Septicemia (A40-A41)	1,892	1.9	35.0
9	Alzheimer's disease (G30)	1,705	1.7	31.5
10	Influenza and pneumonia (J09-J18)	1,585	1.6	29.3
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1,129	1.1	20.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	952	0.9	17.6
13	Parkinson's disease (G20-G21)	696	0.7	12.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	601	0.6	11.1
15	Pneumonitis due to solids and liquids (J69)	450	0.4	8.3
...	All other causes (Residual)	16,074	15.9	297.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 75-79 years				
...	All causes	129,512	100.0	3,085.0
1	Malignant neoplasms (C00-C97)	35,741	27.6	851.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	26,565	20.5	632.8
3	Chronic lower respiratory diseases (J40-J47)	11,364	8.8	270.7
4	Cerebrovascular diseases (I60-I69)	7,798	6.0	185.7
5	Alzheimer's disease (G30)	4,593	3.5	109.4
6	Diabetes mellitus (E10-E14)	4,356	3.4	103.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,672	2.1	63.6
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,653	2.0	63.2
9	Influenza and pneumonia (J09-J18)	2,386	1.8	56.8
10	Septicemia (A40-A41)	2,296	1.8	54.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,553	1.2	37.0
12	Parkinson's disease (G20-G21)	1,420	1.1	33.8
13	Chronic liver disease and cirrhosis (K70,K73-K74)	1,011	0.8	24.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	867	0.7	20.7
15	Pneumonitis due to solids and liquids (J69)	757	0.6	18.0
...	All other causes (Residual)	23,480	18.1	559.3
All races, female, 80-84 years				
...	All causes	180,235	100.0	5,260.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	41,525	23.0	1,211.9
2	Malignant neoplasms (C00-C97)	36,690	20.4	1,070.8
3	Chronic lower respiratory diseases (J40-J47)	13,578	7.5	396.3
4	Cerebrovascular diseases (I60-I69)	12,650	7.0	369.2
5	Alzheimer's disease (G30)	9,983	5.5	291.4
6	Diabetes mellitus (E10-E14)	5,284	2.9	154.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,077	2.3	119.0
8	Influenza and pneumonia (J09-J18)	3,973	2.2	116.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,754	2.1	109.6
10	Septicemia (A40-A41)	2,942	1.6	85.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,436	1.4	71.1
12	Parkinson's disease (G20-G21)	2,267	1.3	66.2
13	Pneumonitis due to solids and liquids (J69)	1,308	0.7	38.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,206	0.7	35.2
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	891	0.5	26.0
...	All other causes (Residual)	37,671	20.9	1,099.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 85-89 years				
...	All causes	222,936	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	59,139	26.5	---
2	Malignant neoplasms (C00-C97)	30,906	13.9	---
3	Cerebrovascular diseases (I60-I69)	16,998	7.6	---
4	Alzheimer's disease (G30)	16,186	7.3	---
5	Chronic lower respiratory diseases (J40-J47)	12,886	5.8	---
6	Influenza and pneumonia (J09-J18)	5,607	2.5	---
7	Diabetes mellitus (E10-E14)	5,398	2.4	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,148	2.3	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,363	2.0	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,677	1.6	---
11	Septicemia (A40-A41)	3,203	1.4	---
12	Parkinson's disease (G20-G21)	2,716	1.2	---
13	Pneumonitis due to solids and liquids (J69)	1,867	0.8	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,432	0.6	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1,141	0.5	---
...	All other causes (Residual)	52,269	23.4	---
All races, female, 90-94 years				
...	All causes	185,024	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	55,440	30.0	---
2	Malignant neoplasms (C00-C97)	16,105	8.7	---
3	Alzheimer's disease (G30)	15,425	8.3	---
4	Cerebrovascular diseases (I60-I69)	14,395	7.8	---
5	Chronic lower respiratory diseases (J40-J47)	7,823	4.2	---
6	Influenza and pneumonia (J09-J18)	5,251	2.8	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,380	2.4	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,538	1.9	---
9	Diabetes mellitus (E10-E14)	3,476	1.9	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,365	1.8	---
11	Septicemia (A40-A41)	2,100	1.1	---
12	Pneumonitis due to solids and liquids (J69)	1,707	0.9	---
13	Parkinson's disease (G20-G21)	1,583	0.9	---
14	Atherosclerosis (I70)	972	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	948	0.5	---
...	All other causes (Residual)	48,516	26.2	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 95-99 years				
...	All causes	83,374	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	26,896	32.3	---
2	Alzheimer's disease (G30)	7,228	8.7	---
3	Cerebrovascular diseases (I60-I69)	5,904	7.1	---
4	Malignant neoplasms (C00-C97)	4,708	5.6	---
5	Influenza and pneumonia (J09-J18)	2,719	3.3	---
6	Chronic lower respiratory diseases (J40-J47)	2,648	3.2	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,008	2.4	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,930	2.3	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,364	1.6	---
10	Diabetes mellitus (E10-E14)	1,210	1.5	---
11	Septicemia (A40-A41)	823	1.0	---
12	Pneumonitis due to solids and liquids (J69)	799	1.0	---
13	Atherosclerosis (I70)	587	0.7	---
14	Parkinson's disease (G20-G21)	438	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	319	0.4	---
...	All other causes (Residual)	23,793	28.5	---
All races, female, 100 years and over				
...	All causes	19,870	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,726	33.9	---
2	Alzheimer's disease (G30)	1,594	8.0	---
3	Cerebrovascular diseases (I60-I69)	1,186	6.0	---
4	Influenza and pneumonia (J09-J18)	810	4.1	---
5	Malignant neoplasms (C00-C97)	713	3.6	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	540	2.7	---
7	Chronic lower respiratory diseases (J40-J47)	464	2.3	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	435	2.2	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	294	1.5	---
10	Diabetes mellitus (E10-E14)	231	1.2	---
11	Atherosclerosis (I70)	199	1.0	---
12	Pneumonitis due to solids and liquids (J69)	186	0.9	---
13	Septicemia (A40-A41)	159	0.8	---
14	Anemias (D50-D64)	70	0.4	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	60	0.3	---
...	All other causes (Residual)	6,203	31.2	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, all ages				
...	All causes	2,175,178	100.0	876.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	514,372	23.6	207.2
2	Malignant neoplasms (C00-C97)	496,885	22.8	200.2
3	Chronic lower respiratory diseases (J40-J47)	131,782	6.1	53.1
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	110,789	5.1	44.6
5	Cerebrovascular diseases (I60-I69)	107,964	5.0	43.5
6	Alzheimer's disease (G30)	76,590	3.5	30.9
7	Diabetes mellitus (E10-E14)	57,806	2.7	23.3
8	Influenza and pneumonia (J09-J18)	43,585	2.0	17.6
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	36,606	1.7	14.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	36,066	1.7	14.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	30,747	1.4	12.4
12	Septicemia (A40-A41)	28,945	1.3	11.7
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	22,908	1.1	9.2
14	Parkinson's disease (G20-G21)	22,352	1.0	9.0
15	Pneumonitis due to solids and liquids (J69)	15,938	0.7	6.4
...	All other causes (Residual)	441,843	20.3	178.0
White, both sexes, 1-4 years				
...	All causes	2,958	100.0	24.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,004	33.9	8.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	366	12.4	3.0
3	Malignant neoplasms (C00-C97)	301	10.2	2.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	189	6.4	1.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	101	3.4	0.8
6	Influenza and pneumonia (J09-J18)	67	2.3	0.6
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	44	1.5	0.4
8	Septicemia (A40-A41)	37	1.3	0.3
9	Cerebrovascular diseases (I60-I69)	35	1.2	0.3
10	Certain conditions originating in the perinatal period (P00-P96)	24	0.8	0.2
11	Chronic lower respiratory diseases (J40-J47)	20	0.7	0.2
12	Acute bronchitis and bronchiolitis (J20-J21)	11	0.4	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.3	*
14	Pneumonitis due to solids and liquids (J69)	7	0.2	*
15	Anemias (D50-D64)	6	0.2	*
15	Diabetes mellitus (E10-E14)	6	0.2	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.2	*
...	All other causes (Residual)	726	24.5	6.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 5-9 years				
...	All causes	1,642	100.0	10.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	491	29.9	3.2
2	Malignant neoplasms (C00-C97)	352	21.4	2.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	111	6.8	0.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	94	5.7	0.6
5	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	44	2.7	0.3
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	43	2.6	0.3
7	Influenza and pneumonia (J09-J18)	36	2.2	0.2
8	Cerebrovascular diseases (I60-I69)	27	1.6	0.2
9	Septicemia (A40-A41)	21	1.3	0.1
10	Chronic lower respiratory diseases (J40-J47)	19	1.2	*
11	Certain conditions originating in the perinatal period (P00-P96)	14	0.9	*
12	Meningitis (G00,G03)	9	0.5	*
13	Diabetes mellitus (E10-E14)	7	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.4	*
15	Anemias (D50-D64)	5	0.3	*
...	All other causes (Residual)	363	22.1	2.3
White, both sexes, 10-14 years				
...	All causes	2,069	100.0	13.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	596	28.8	3.8
2	Malignant neoplasms (C00-C97)	360	17.4	2.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	245	11.8	1.6
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	118	5.7	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	102	4.9	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	73	3.5	0.5
7	Cerebrovascular diseases (I60-I69)	40	1.9	0.3
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	30	1.4	0.2
9	Influenza and pneumonia (J09-J18)	27	1.3	0.2
10	Septicemia (A40-A41)	22	1.1	0.1
10	Chronic lower respiratory diseases (J40-J47)	22	1.1	0.1
12	Diabetes mellitus (E10-E14)	10	0.5	*
12	Certain conditions originating in the perinatal period (P00-P96)	10	0.5	*
14	Anemias (D50-D64)	6	0.3	*
15	Meningitis (G00,G03)	5	0.2	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.2	*
...	All other causes (Residual)	398	19.2	2.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 15-19 years				
...	All causes	7,296	100.0	45.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,330	45.6	20.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,500	20.6	9.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	543	7.4	3.3
4	Malignant neoplasms (C00-C97)	489	6.7	3.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	192	2.6	1.2
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	147	2.0	0.9
7	Cerebrovascular diseases (I60-I69)	46	0.6	0.3
8	Influenza and pneumonia (J09-J18)	40	0.5	0.2
9	Septicemia (A40-A41)	31	0.4	0.2
10	Chronic lower respiratory diseases (J40-J47)	28	0.4	0.2
11	Diabetes mellitus (E10-E14)	27	0.4	0.2
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	26	0.4	0.2
13	Legal intervention (Y35,Y89.0)	20	0.3	0.1
14	Pregnancy, childbirth and the puerperium (O00-O99)	18	0.2	*
15	Pneumonitis due to solids and liquids (J69)	13	0.2	*
...	All other causes (Residual)	846	11.6	5.2
White, both sexes, 20-24 years				
...	All causes	13,805	100.0	80.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,671	48.3	39.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,501	18.1	14.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,036	7.5	6.1
4	Malignant neoplasms (C00-C97)	729	5.3	4.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	421	3.0	2.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	174	1.3	1.0
7	Diabetes mellitus (E10-E14)	104	0.8	0.6
8	Cerebrovascular diseases (I60-I69)	87	0.6	0.5
9	Pregnancy, childbirth and the puerperium (O00-O99)	80	0.6	0.5
10	Influenza and pneumonia (J09-J18)	70	0.5	0.4
11	Septicemia (A40-A41)	50	0.4	0.3
12	Chronic lower respiratory diseases (J40-J47)	47	0.3	0.3
13	Legal intervention (Y35,Y89.0)	45	0.3	0.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	36	0.3	0.2
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	29	0.2	0.2
...	All other causes (Residual)	1,725	12.5	10.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 25-29 years				
...	All causes	15,674	100.0	95.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,760	43.1	41.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,689	17.2	16.4
3	Malignant neoplasms (C00-C97)	1,087	6.9	6.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	942	6.0	5.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	790	5.0	4.8
6	Diabetes mellitus (E10-E14)	162	1.0	1.0
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	158	1.0	1.0
8	Chronic liver disease and cirrhosis (K70,K73-K74)	131	0.8	0.8
9	Cerebrovascular diseases (I60-I69)	128	0.8	0.8
10	Pregnancy, childbirth and the puerperium (O00-O99)	112	0.7	0.7
11	Influenza and pneumonia (J09-J18)	101	0.6	0.6
12	Septicemia (A40-A41)	99	0.6	0.6
13	Human immunodeficiency virus (HIV) disease (B20-B24)	82	0.5	0.5
13	Chronic lower respiratory diseases (J40-J47)	82	0.5	0.5
15	Legal intervention (Y35,Y89.0)	51	0.3	0.3
...	All other causes (Residual)	2,300	14.7	14.0
White, both sexes, 30-34 years				
...	All causes	17,696	100.0	110.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,726	38.0	41.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,613	14.8	16.3
3	Malignant neoplasms (C00-C97)	1,704	9.6	10.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,379	7.8	8.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	802	4.5	5.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	353	2.0	2.2
7	Diabetes mellitus (E10-E14)	254	1.4	1.6
8	Cerebrovascular diseases (I60-I69)	246	1.4	1.5
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	169	1.0	1.1
10	Influenza and pneumonia (J09-J18)	158	0.9	1.0
11	Human immunodeficiency virus (HIV) disease (B20-B24)	136	0.8	0.8
12	Septicemia (A40-A41)	117	0.7	0.7
12	Pregnancy, childbirth and the puerperium (O00-O99)	117	0.7	0.7
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	90	0.5	0.6
15	Chronic lower respiratory diseases (J40-J47)	86	0.5	0.5
...	All other causes (Residual)	2,746	15.5	17.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 35-39 years				
...	All causes	20,401	100.0	136.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,004	29.4	40.0
2	Malignant neoplasms (C00-C97)	2,802	13.7	18.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,730	13.4	18.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,402	11.8	16.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	660	3.2	4.4
6	Chronic liver disease and cirrhosis (K70,K73-K74)	627	3.1	4.2
7	Diabetes mellitus (E10-E14)	435	2.1	2.9
8	Cerebrovascular diseases (I60-I69)	381	1.9	2.5
9	Human immunodeficiency virus (HIV) disease (B20-B24)	188	0.9	1.3
10	Septicemia (A40-A41)	187	0.9	1.2
11	Influenza and pneumonia (J09-J18)	173	0.8	1.2
12	Chronic lower respiratory diseases (J40-J47)	151	0.7	1.0
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	146	0.7	1.0
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	146	0.7	1.0
15	Pregnancy, childbirth and the puerperium (O00-O99)	98	0.5	0.7
...	All other causes (Residual)	3,271	16.0	21.8
White, both sexes, 40-44 years				
...	All causes	31,827	100.0	193.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,766	21.3	41.1
2	Malignant neoplasms (C00-C97)	5,874	18.5	35.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,846	15.2	29.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,249	10.2	19.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,438	4.5	8.7
6	Diabetes mellitus (E10-E14)	841	2.6	5.1
7	Cerebrovascular diseases (I60-I69)	759	2.4	4.6
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	656	2.1	4.0
9	Chronic lower respiratory diseases (J40-J47)	375	1.2	2.3
10	Human immunodeficiency virus (HIV) disease (B20-B24)	362	1.1	2.2
11	Influenza and pneumonia (J09-J18)	331	1.0	2.0
12	Septicemia (A40-A41)	326	1.0	2.0
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	218	0.7	1.3
14	Viral hepatitis (B15-B19)	179	0.6	1.1
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	174	0.5	1.1
...	All other causes (Residual)	5,433	17.1	33.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 45-49 years				
...	All causes	53,097	100.0	307.7
1	Malignant neoplasms (C00-C97)	12,509	23.6	72.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,449	17.8	54.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,280	15.6	48.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,851	7.3	22.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	3,016	5.7	17.5
6	Diabetes mellitus (E10-E14)	1,512	2.8	8.8
7	Cerebrovascular diseases (I60-I69)	1,407	2.6	8.2
8	Chronic lower respiratory diseases (J40-J47)	1,006	1.9	5.8
9	Septicemia (A40-A41)	634	1.2	3.7
10	Human immunodeficiency virus (HIV) disease (B20-B24)	611	1.2	3.5
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	602	1.1	3.5
12	Viral hepatitis (B15-B19)	538	1.0	3.1
13	Influenza and pneumonia (J09-J18)	525	1.0	3.0
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	417	0.8	2.4
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	259	0.5	1.5
...	All other causes (Residual)	8,481	16.0	49.1
White, both sexes, 50-54 years				
...	All causes	87,042	100.0	475.7
1	Malignant neoplasms (C00-C97)	25,306	29.1	138.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	16,986	19.5	92.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,082	10.4	49.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	4,749	5.5	26.0
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,381	5.0	23.9
6	Chronic lower respiratory diseases (J40-J47)	2,766	3.2	15.1
7	Diabetes mellitus (E10-E14)	2,552	2.9	13.9
8	Cerebrovascular diseases (I60-I69)	2,265	2.6	12.4
9	Viral hepatitis (B15-B19)	1,211	1.4	6.6
10	Septicemia (A40-A41)	1,045	1.2	5.7
11	Influenza and pneumonia (J09-J18)	844	1.0	4.6
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	811	0.9	4.4
13	Human immunodeficiency virus (HIV) disease (B20-B24)	632	0.7	3.5
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	616	0.7	3.4
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	497	0.6	2.7
...	All other causes (Residual)	13,299	15.3	72.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 55-59 years				
...	All causes	117,751	100.0	690.9
1	Malignant neoplasms (C00-C97)	39,218	33.3	230.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	24,140	20.5	141.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,525	6.4	44.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	5,275	4.5	31.0
5	Chronic lower respiratory diseases (J40-J47)	5,138	4.4	30.1
6	Diabetes mellitus (E10-E14)	3,922	3.3	23.0
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,828	3.3	22.5
8	Cerebrovascular diseases (I60-I69)	3,316	2.8	19.5
9	Viral hepatitis (B15-B19)	1,831	1.6	10.7
10	Septicemia (A40-A41)	1,541	1.3	9.0
11	Influenza and pneumonia (J09-J18)	1,337	1.1	7.8
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,261	1.1	7.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	828	0.7	4.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	553	0.5	3.2
15	Human immunodeficiency virus (HIV) disease (B20-B24)	500	0.4	2.9
...	All other causes (Residual)	17,538	14.9	102.9
White, both sexes, 60-64 years				
...	All causes	145,906	100.0	982.8
1	Malignant neoplasms (C00-C97)	52,800	36.2	355.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	31,209	21.4	210.2
3	Chronic lower respiratory diseases (J40-J47)	8,252	5.7	55.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,720	3.9	38.5
5	Diabetes mellitus (E10-E14)	5,386	3.7	36.3
6	Chronic liver disease and cirrhosis (K70,K73-K74)	4,508	3.1	30.4
7	Cerebrovascular diseases (I60-I69)	4,487	3.1	30.2
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,700	1.9	18.2
9	Septicemia (A40-A41)	2,173	1.5	14.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,884	1.3	12.7
11	Influenza and pneumonia (J09-J18)	1,735	1.2	11.7
12	Viral hepatitis (B15-B19)	1,272	0.9	8.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,161	0.8	7.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	711	0.5	4.8
15	Pneumonitis due to solids and liquids (J69)	581	0.4	3.9
...	All other causes (Residual)	21,327	14.6	143.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 65-69 years				
...	All causes	172,271	100.0	1,445.2
1	Malignant neoplasms (C00-C97)	63,528	36.9	532.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	36,291	21.1	304.4
3	Chronic lower respiratory diseases (J40-J47)	13,373	7.8	112.2
4	Diabetes mellitus (E10-E14)	6,373	3.7	53.5
5	Cerebrovascular diseases (I60-I69)	6,084	3.5	51.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,807	2.8	40.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	3,418	2.0	28.7
8	Septicemia (A40-A41)	2,606	1.5	21.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,583	1.5	21.7
10	Influenza and pneumonia (J09-J18)	2,258	1.3	18.9
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,859	1.1	15.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,308	0.8	11.0
13	Alzheimer's disease (G30)	1,174	0.7	9.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	972	0.6	8.2
15	Parkinson's disease (G20-G21)	816	0.5	6.8
...	All other causes (Residual)	24,821	14.4	208.2
White, both sexes, 70-74 years				
...	All causes	192,435	100.0	2,249.3
1	Malignant neoplasms (C00-C97)	66,155	34.4	773.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	40,544	21.1	473.9
3	Chronic lower respiratory diseases (J40-J47)	17,357	9.0	202.9
4	Cerebrovascular diseases (I60-I69)	8,199	4.3	95.8
5	Diabetes mellitus (E10-E14)	6,486	3.4	75.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,404	2.3	51.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,349	1.7	39.1
8	Septicemia (A40-A41)	2,941	1.5	34.4
9	Influenza and pneumonia (J09-J18)	2,897	1.5	33.9
10	Alzheimer's disease (G30)	2,635	1.4	30.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	2,477	1.3	29.0
12	Parkinson's disease (G20-G21)	1,881	1.0	22.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,483	0.8	17.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,350	0.7	15.8
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,314	0.7	15.4
...	All other causes (Residual)	28,963	15.1	338.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 75-79 years				
...	All causes	235,269	100.0	3,643.6
1	Malignant neoplasms (C00-C97)	67,553	28.7	1,046.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	52,216	22.2	808.7
3	Chronic lower respiratory diseases (J40-J47)	20,855	8.9	323.0
4	Cerebrovascular diseases (I60-I69)	12,098	5.1	187.4
5	Diabetes mellitus (E10-E14)	7,145	3.0	110.7
6	Alzheimer's disease (G30)	6,793	2.9	105.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,359	2.3	83.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,423	1.9	68.5
9	Influenza and pneumonia (J09-J18)	4,276	1.8	66.2
10	Parkinson's disease (G20-G21)	3,709	1.6	57.4
11	Septicemia (A40-A41)	3,647	1.6	56.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,105	0.9	32.6
13	Chronic liver disease and cirrhosis (K70,K73-K74)	2,031	0.9	31.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,790	0.8	27.7
15	Pneumonitis due to solids and liquids (J69)	1,602	0.7	24.8
...	All other causes (Residual)	39,667	16.9	614.3
White, both sexes, 80-84 years				
...	All causes	310,210	100.0	6,088.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	75,487	24.3	1,481.6
2	Malignant neoplasms (C00-C97)	68,025	21.9	1,335.1
3	Chronic lower respiratory diseases (J40-J47)	24,203	7.8	475.0
4	Cerebrovascular diseases (I60-I69)	18,834	6.1	369.7
5	Alzheimer's disease (G30)	14,367	4.6	282.0
6	Diabetes mellitus (E10-E14)	8,367	2.7	164.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,709	2.5	151.3
8	Influenza and pneumonia (J09-J18)	7,039	2.3	138.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,398	2.1	125.6
10	Parkinson's disease (G20-G21)	5,788	1.9	113.6
11	Septicemia (A40-A41)	4,528	1.5	88.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,343	1.1	65.6
13	Pneumonitis due to solids and liquids (J69)	2,704	0.9	53.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,401	0.8	47.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,533	0.5	30.1
...	All other causes (Residual)	59,484	19.2	1,167.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 85-89 years				
...	All causes	348,333	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	95,913	27.5	---
2	Malignant neoplasms (C00-C97)	54,111	15.5	---
3	Cerebrovascular diseases (I60-I69)	23,198	6.7	---
4	Alzheimer's disease (G30)	21,780	6.3	---
5	Chronic lower respiratory diseases (J40-J47)	21,699	6.2	---
6	Influenza and pneumonia (J09-J18)	9,143	2.6	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,855	2.5	---
8	Diabetes mellitus (E10-E14)	7,921	2.3	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,114	2.0	---
10	Parkinson's disease (G20-G21)	5,839	1.7	---
11	Septicemia (A40-A41)	4,759	1.4	---
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,740	1.4	---
13	Pneumonitis due to solids and liquids (J69)	3,640	1.0	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,583	0.7	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1,665	0.5	---
...	All other causes (Residual)	75,373	21.6	---
White, both sexes, 90-94 years				
...	All causes	260,450	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	80,411	30.9	---
2	Malignant neoplasms (C00-C97)	26,315	10.1	---
3	Alzheimer's disease (G30)	19,337	7.4	---
4	Cerebrovascular diseases (I60-I69)	18,195	7.0	---
5	Chronic lower respiratory diseases (J40-J47)	12,073	4.6	---
6	Influenza and pneumonia (J09-J18)	7,957	3.1	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,788	2.6	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,101	2.0	---
9	Diabetes mellitus (E10-E14)	4,662	1.8	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,301	1.7	---
11	Pneumonitis due to solids and liquids (J69)	3,142	1.2	---
12	Parkinson's disease (G20-G21)	3,038	1.2	---
13	Septicemia (A40-A41)	2,947	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,492	0.6	---
15	Atherosclerosis (I70)	1,389	0.5	---
...	All other causes (Residual)	63,302	24.3	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 95-99 years				
...	All causes	102,679	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	33,972	33.1	---
2	Alzheimer's disease (G30)	8,096	7.9	---
3	Cerebrovascular diseases (I60-I69)	6,846	6.7	---
4	Malignant neoplasms (C00-C97)	6,776	6.6	---
5	Chronic lower respiratory diseases (J40-J47)	3,670	3.6	---
6	Influenza and pneumonia (J09-J18)	3,583	3.5	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,689	2.6	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,117	2.1	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,801	1.8	---
10	Diabetes mellitus (E10-E14)	1,418	1.4	---
11	Pneumonitis due to solids and liquids (J69)	1,235	1.2	---
12	Septicemia (A40-A41)	968	0.9	---
13	Atherosclerosis (I70)	724	0.7	---
14	Parkinson's disease (G20-G21)	662	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	476	0.5	---
...	All other causes (Residual)	27,646	26.9	---
White, both sexes, 100 years and over				
...	All causes	21,012	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,270	34.6	---
2	Alzheimer's disease (G30)	1,551	7.4	---
3	Cerebrovascular diseases (I60-I69)	1,223	5.8	---
4	Influenza and pneumonia (J09-J18)	895	4.3	---
5	Malignant neoplasms (C00-C97)	831	4.0	---
6	Chronic lower respiratory diseases (J40-J47)	544	2.6	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	536	2.6	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	505	2.4	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	322	1.5	---
10	Pneumonitis due to solids and liquids (J69)	239	1.1	---
11	Atherosclerosis (I70)	226	1.1	---
12	Diabetes mellitus (E10-E14)	215	1.0	---
13	Septicemia (A40-A41)	169	0.8	---
14	Anemias (D50-D64)	82	0.4	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	70	0.3	---
15	Parkinson's disease (G20-G21)	70	0.3	---
...	All other causes (Residual)	6,264	29.8	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, all ages				
...	All causes	1,085,250	100.0	882.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	267,970	24.7	218.0
2	Malignant neoplasms (C00-C97)	261,975	24.1	213.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	68,677	6.3	55.9
4	Chronic lower respiratory diseases (J40-J47)	61,452	5.7	50.0
5	Cerebrovascular diseases (I60-I69)	43,435	4.0	35.3
6	Diabetes mellitus (E10-E14)	30,832	2.8	25.1
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	28,681	2.6	23.3
8	Alzheimer's disease (G30)	23,607	2.2	19.2
9	Influenza and pneumonia (J09-J18)	20,575	1.9	16.7
10	Chronic liver disease and cirrhosis (K70,K73-K74)	19,973	1.8	16.2
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18,269	1.7	14.9
12	Septicemia (A40-A41)	13,654	1.3	11.1
13	Parkinson's disease (G20-G21)	13,281	1.2	10.8
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9,098	0.8	7.4
15	Pneumonitis due to solids and liquids (J69)	8,662	0.8	7.0
...	All other causes (Residual)	195,109	18.0	158.7
White, male, 1-4 years				
...	All causes	1,683	100.0	27.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	591	35.1	9.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	208	12.4	3.4
3	Malignant neoplasms (C00-C97)	165	9.8	2.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	109	6.5	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	47	2.8	0.8
6	Influenza and pneumonia (J09-J18)	42	2.5	0.7
7	Cerebrovascular diseases (I60-I69)	25	1.5	0.4
8	Septicemia (A40-A41)	19	1.1	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	1.0	*
9	Certain conditions originating in the perinatal period (P00-P96)	16	1.0	*
11	Chronic lower respiratory diseases (J40-J47)	11	0.7	*
12	Acute bronchitis and bronchiolitis (J20-J21)	8	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.4	*
14	Diabetes mellitus (E10-E14)	4	0.2	*
14	Pneumonitis due to solids and liquids (J69)	4	0.2	*
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	4	0.2	*
...	All other causes (Residual)	408	24.2	6.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 5-9 years				
...	All causes	916	100.0	11.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	292	31.9	3.7
2	Malignant neoplasms (C00-C97)	204	22.3	2.6
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	51	5.6	0.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	48	5.2	0.6
5	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	22	2.4	0.3
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	21	2.3	0.3
7	Influenza and pneumonia (J09-J18)	18	2.0	*
8	Septicemia (A40-A41)	15	1.6	*
9	Cerebrovascular diseases (I60-I69)	12	1.3	*
9	Chronic lower respiratory diseases (J40-J47)	12	1.3	*
11	Certain conditions originating in the perinatal period (P00-P96)	6	0.7	*
12	Meningitis (G00,G03)	4	0.4	*
13	Anemias (D50-D64)	3	0.3	*
13	Diabetes mellitus (E10-E14)	3	0.3	*
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3	0.3	*
...	All other causes (Residual)	202	22.1	2.5
White, male, 10-14 years				
...	All causes	1,216	100.0	15.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	384	31.6	4.8
2	Malignant neoplasms (C00-C97)	195	16.0	2.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	175	14.4	2.2
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	63	5.2	0.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	58	4.8	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	40	3.3	0.5
7	Cerebrovascular diseases (I60-I69)	21	1.7	0.3
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	1.6	*
9	Septicemia (A40-A41)	13	1.1	*
10	Influenza and pneumonia (J09-J18)	11	0.9	*
11	Chronic lower respiratory diseases (J40-J47)	8	0.7	*
12	Diabetes mellitus (E10-E14)	7	0.6	*
13	Meningitis (G00,G03)	3	0.2	*
13	Diseases of appendix (K35-K38)	3	0.2	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.2	*
...	All other causes (Residual)	213	17.5	2.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 15-19 years				
...	All causes	5,119	100.0	61.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,320	45.3	27.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,165	22.8	14.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	445	8.7	5.3
4	Malignant neoplasms (C00-C97)	289	5.6	3.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	127	2.5	1.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	87	1.7	1.0
7	Cerebrovascular diseases (I60-I69)	25	0.5	0.3
8	Influenza and pneumonia (J09-J18)	22	0.4	0.3
9	Chronic lower respiratory diseases (J40-J47)	20	0.4	0.2
9	Legal intervention (Y35,Y89.0)	20	0.4	0.2
11	Septicemia (A40-A41)	14	0.3	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	0.3	*
13	Diabetes mellitus (E10-E14)	12	0.2	*
14	Pneumonitis due to solids and liquids (J69)	9	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.1	*
...	All other causes (Residual)	544	10.6	6.5
White, male, 20-24 years				
...	All causes	10,243	100.0	116.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,132	50.1	58.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,088	20.4	23.8
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	834	8.1	9.5
4	Malignant neoplasms (C00-C97)	416	4.1	4.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	283	2.8	3.2
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	99	1.0	1.1
7	Diabetes mellitus (E10-E14)	64	0.6	0.7
8	Cerebrovascular diseases (I60-I69)	52	0.5	0.6
9	Influenza and pneumonia (J09-J18)	46	0.4	0.5
10	Legal intervention (Y35,Y89.0)	42	0.4	0.5
11	Chronic lower respiratory diseases (J40-J47)	29	0.3	0.3
12	Septicemia (A40-A41)	24	0.2	0.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	22	0.2	0.3
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	0.2	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	15	0.1	*
...	All other causes (Residual)	1,081	10.6	12.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 25-29 years				
...	All causes	11,184	100.0	133.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,069	45.3	60.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,189	19.6	26.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	743	6.6	8.8
4	Malignant neoplasms (C00-C97)	606	5.4	7.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	561	5.0	6.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	96	0.9	1.1
7	Diabetes mellitus (E10-E14)	92	0.8	1.1
8	Chronic liver disease and cirrhosis (K70,K73-K74)	88	0.8	1.0
9	Human immunodeficiency virus (HIV) disease (B20-B24)	71	0.6	0.8
10	Cerebrovascular diseases (I60-I69)	70	0.6	0.8
11	Septicemia (A40-A41)	60	0.5	0.7
12	Influenza and pneumonia (J09-J18)	57	0.5	0.7
13	Legal intervention (Y35,Y89.0)	49	0.4	0.6
14	Chronic lower respiratory diseases (J40-J47)	46	0.4	0.5
15	Aortic aneurysm and dissection (I71)	27	0.2	0.3
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	27	0.2	0.3
...	All other causes (Residual)	1,333	11.9	15.8
White, male, 30-34 years				
...	All causes	11,923	100.0	145.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,871	40.9	59.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,050	17.2	25.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	964	8.1	11.8
4	Malignant neoplasms (C00-C97)	803	6.7	9.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	597	5.0	7.3
6	Chronic liver disease and cirrhosis (K70,K73-K74)	234	2.0	2.9
7	Diabetes mellitus (E10-E14)	157	1.3	1.9
8	Cerebrovascular diseases (I60-I69)	138	1.2	1.7
9	Human immunodeficiency virus (HIV) disease (B20-B24)	108	0.9	1.3
10	Influenza and pneumonia (J09-J18)	95	0.8	1.2
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	94	0.8	1.1
12	Septicemia (A40-A41)	64	0.5	0.8
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	49	0.4	0.6
14	Chronic lower respiratory diseases (J40-J47)	46	0.4	0.6
15	Legal intervention (Y35,Y89.0)	45	0.4	0.5
...	All other causes (Residual)	1,608	13.5	19.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 35-39 years				
...	All causes	13,156	100.0	173.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,255	32.3	56.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,111	16.0	27.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,680	12.8	22.2
4	Malignant neoplasms (C00-C97)	1,258	9.6	16.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	476	3.6	6.3
6	Chronic liver disease and cirrhosis (K70,K73-K74)	410	3.1	5.4
7	Diabetes mellitus (E10-E14)	266	2.0	3.5
8	Cerebrovascular diseases (I60-I69)	218	1.7	2.9
9	Human immunodeficiency virus (HIV) disease (B20-B24)	147	1.1	1.9
10	Septicemia (A40-A41)	109	0.8	1.4
11	Influenza and pneumonia (J09-J18)	88	0.7	1.2
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	81	0.6	1.1
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	80	0.6	1.1
14	Chronic lower respiratory diseases (J40-J47)	68	0.5	0.9
15	Aortic aneurysm and dissection (I71)	61	0.5	0.8
...	All other causes (Residual)	1,848	14.0	24.4
White, male, 40-44 years				
...	All causes	19,753	100.0	238.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,632	23.4	55.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,415	17.3	41.2
3	Malignant neoplasms (C00-C97)	2,585	13.1	31.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,480	12.6	29.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	938	4.7	11.3
6	Diabetes mellitus (E10-E14)	524	2.7	6.3
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	477	2.4	5.7
8	Cerebrovascular diseases (I60-I69)	424	2.1	5.1
9	Human immunodeficiency virus (HIV) disease (B20-B24)	286	1.4	3.4
10	Influenza and pneumonia (J09-J18)	178	0.9	2.1
11	Chronic lower respiratory diseases (J40-J47)	168	0.9	2.0
12	Septicemia (A40-A41)	164	0.8	2.0
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	134	0.7	1.6
14	Viral hepatitis (B15-B19)	112	0.6	1.3
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	102	0.5	1.2
...	All other causes (Residual)	3,134	15.9	37.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 45-49 years				
...	All causes	32,605	100.0	377.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,771	20.8	78.5
2	Malignant neoplasms (C00-C97)	5,975	18.3	69.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,545	17.0	64.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,837	8.7	32.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,023	6.2	23.4
6	Diabetes mellitus (E10-E14)	982	3.0	11.4
7	Cerebrovascular diseases (I60-I69)	753	2.3	8.7
8	Human immunodeficiency virus (HIV) disease (B20-B24)	493	1.5	5.7
9	Chronic lower respiratory diseases (J40-J47)	449	1.4	5.2
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	415	1.3	4.8
11	Viral hepatitis (B15-B19)	383	1.2	4.4
12	Septicemia (A40-A41)	348	1.1	4.0
13	Influenza and pneumonia (J09-J18)	282	0.9	3.3
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	257	0.8	3.0
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	168	0.5	1.9
...	All other causes (Residual)	4,924	15.1	57.1
White, male, 50-54 years				
...	All causes	53,962	100.0	594.9
1	Malignant neoplasms (C00-C97)	13,180	24.4	145.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,398	23.0	136.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,009	11.1	66.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	3,294	6.1	36.3
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,271	6.1	36.1
6	Diabetes mellitus (E10-E14)	1,632	3.0	18.0
7	Chronic lower respiratory diseases (J40-J47)	1,304	2.4	14.4
8	Cerebrovascular diseases (I60-I69)	1,267	2.3	14.0
9	Viral hepatitis (B15-B19)	870	1.6	9.6
10	Septicemia (A40-A41)	561	1.0	6.2
11	Human immunodeficiency virus (HIV) disease (B20-B24)	527	1.0	5.8
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	493	0.9	5.4
13	Influenza and pneumonia (J09-J18)	486	0.9	5.4
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	453	0.8	5.0
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	360	0.7	4.0
...	All other causes (Residual)	7,857	14.6	86.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 55-59 years				
...	All causes	73,372	100.0	878.7
1	Malignant neoplasms (C00-C97)	21,714	29.6	260.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	17,470	23.8	209.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,051	6.9	60.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	3,774	5.1	45.2
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,917	4.0	34.9
6	Chronic lower respiratory diseases (J40-J47)	2,593	3.5	31.1
7	Diabetes mellitus (E10-E14)	2,503	3.4	30.0
8	Cerebrovascular diseases (I60-I69)	1,959	2.7	23.5
9	Viral hepatitis (B15-B19)	1,314	1.8	15.7
10	Septicemia (A40-A41)	867	1.2	10.4
11	Influenza and pneumonia (J09-J18)	767	1.0	9.2
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	750	1.0	9.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	526	0.7	6.3
14	Human immunodeficiency virus (HIV) disease (B20-B24)	418	0.6	5.0
15	Aortic aneurysm and dissection (I71)	343	0.5	4.1
...	All other causes (Residual)	10,406	14.2	124.6
White, male, 60-64 years				
...	All causes	88,887	100.0	1,234.8
1	Malignant neoplasms (C00-C97)	29,496	33.2	409.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	21,903	24.6	304.3
3	Chronic lower respiratory diseases (J40-J47)	4,318	4.9	60.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,934	4.4	54.7
5	Diabetes mellitus (E10-E14)	3,309	3.7	46.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	3,155	3.5	43.8
7	Cerebrovascular diseases (I60-I69)	2,610	2.9	36.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,103	2.4	29.2
9	Septicemia (A40-A41)	1,178	1.3	16.4
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,057	1.2	14.7
11	Influenza and pneumonia (J09-J18)	1,031	1.2	14.3
12	Viral hepatitis (B15-B19)	933	1.0	13.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	741	0.8	10.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	425	0.5	5.9
15	Aortic aneurysm and dissection (I71)	389	0.4	5.4
...	All other causes (Residual)	12,305	13.8	170.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 65-69 years				
...	All causes	100,195	100.0	1,759.1
1	Malignant neoplasms (C00-C97)	35,246	35.2	618.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	24,256	24.2	425.8
3	Chronic lower respiratory diseases (J40-J47)	6,746	6.7	118.4
4	Diabetes mellitus (E10-E14)	3,799	3.8	66.7
5	Cerebrovascular diseases (I60-I69)	3,349	3.3	58.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,128	3.1	54.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,194	2.2	38.5
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,445	1.4	25.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,426	1.4	25.0
10	Septicemia (A40-A41)	1,374	1.4	24.1
11	Influenza and pneumonia (J09-J18)	1,289	1.3	22.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	721	0.7	12.7
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	569	0.6	10.0
14	Parkinson's disease (G20-G21)	554	0.6	9.7
15	Alzheimer's disease (G30)	512	0.5	9.0
...	All other causes (Residual)	13,587	13.6	238.5
White, male, 70-74 years				
...	All causes	107,691	100.0	2,712.1
1	Malignant neoplasms (C00-C97)	36,558	33.9	920.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	25,378	23.6	639.1
3	Chronic lower respiratory diseases (J40-J47)	8,547	7.9	215.2
4	Cerebrovascular diseases (I60-I69)	4,196	3.9	105.7
5	Diabetes mellitus (E10-E14)	3,704	3.4	93.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,711	2.5	68.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,836	1.7	46.2
8	Influenza and pneumonia (J09-J18)	1,604	1.5	40.4
9	Septicemia (A40-A41)	1,482	1.4	37.3
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,458	1.4	36.7
11	Parkinson's disease (G20-G21)	1,247	1.2	31.4
12	Alzheimer's disease (G30)	1,137	1.1	28.6
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,109	1.0	27.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	842	0.8	21.2
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	790	0.7	19.9
...	All other causes (Residual)	15,092	14.0	380.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 75-79 years				
...	All causes	124,028	100.0	4,317.5
1	Malignant neoplasms (C00-C97)	36,625	29.5	1,275.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	29,976	24.2	1,043.5
3	Chronic lower respiratory diseases (J40-J47)	10,197	8.2	355.0
4	Cerebrovascular diseases (I60-I69)	5,648	4.6	196.6
5	Diabetes mellitus (E10-E14)	3,840	3.1	133.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,963	2.4	103.1
7	Alzheimer's disease (G30)	2,676	2.2	93.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,386	1.9	83.1
9	Parkinson's disease (G20-G21)	2,378	1.9	82.8
10	Influenza and pneumonia (J09-J18)	2,246	1.8	78.2
11	Septicemia (A40-A41)	1,796	1.4	62.5
12	Chronic liver disease and cirrhosis (K70,K73-K74)	1,110	0.9	38.6
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,024	0.8	35.6
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	984	0.8	34.3
15	Pneumonitis due to solids and liquids (J69)	933	0.8	32.5
...	All other causes (Residual)	19,246	15.5	670.0
White, male, 80-84 years				
...	All causes	151,151	100.0	7,184.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	39,327	26.0	1,869.3
2	Malignant neoplasms (C00-C97)	35,686	23.6	1,696.2
3	Chronic lower respiratory diseases (J40-J47)	11,400	7.5	541.9
4	Cerebrovascular diseases (I60-I69)	7,835	5.2	372.4
5	Alzheimer's disease (G30)	5,247	3.5	249.4
6	Diabetes mellitus (E10-E14)	4,196	2.8	199.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,942	2.6	187.4
8	Parkinson's disease (G20-G21)	3,667	2.4	174.3
9	Influenza and pneumonia (J09-J18)	3,567	2.4	169.5
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,356	2.2	159.5
11	Septicemia (A40-A41)	2,070	1.4	98.4
12	Pneumonitis due to solids and liquids (J69)	1,547	1.0	73.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,367	0.9	65.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,320	0.9	62.7
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	858	0.6	40.8
...	All other causes (Residual)	25,766	17.0	1,224.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 85-89 years				
...	All causes	147,587	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	42,895	29.1	---
2	Malignant neoplasms (C00-C97)	26,388	17.9	---
3	Chronic lower respiratory diseases (J40-J47)	9,519	6.4	---
4	Cerebrovascular diseases (I60-I69)	8,042	5.4	---
5	Alzheimer's disease (G30)	6,903	4.7	---
6	Influenza and pneumonia (J09-J18)	4,143	2.8	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,984	2.7	---
8	Diabetes mellitus (E10-E14)	3,519	2.4	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,442	2.3	---
10	Parkinson's disease (G20-G21)	3,266	2.2	---
11	Septicemia (A40-A41)	2,008	1.4	---
12	Pneumonitis due to solids and liquids (J69)	1,937	1.3	---
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,607	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,268	0.9	---
15	Aortic aneurysm and dissection (I71)	670	0.5	---
...	All other causes (Residual)	27,996	19.0	---
White, male, 90-94 years				
...	All causes	91,564	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	29,647	32.4	---
2	Malignant neoplasms (C00-C97)	11,784	12.9	---
3	Alzheimer's disease (G30)	5,077	5.5	---
4	Cerebrovascular diseases (I60-I69)	5,071	5.5	---
5	Chronic lower respiratory diseases (J40-J47)	4,647	5.1	---
6	Influenza and pneumonia (J09-J18)	3,217	3.5	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,594	2.8	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,203	2.4	---
9	Diabetes mellitus (E10-E14)	1,781	1.9	---
10	Pneumonitis due to solids and liquids (J69)	1,593	1.7	---
11	Parkinson's disease (G20-G21)	1,550	1.7	---
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,196	1.3	---
13	Septicemia (A40-A41)	1,129	1.2	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	613	0.7	---
15	Atherosclerosis (I70)	477	0.5	---
...	All other causes (Residual)	18,985	20.7	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 95-99 years				
...	All causes	26,937	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,315	34.6	---
2	Malignant neoplasms (C00-C97)	2,547	9.5	---
3	Alzheimer's disease (G30)	1,514	5.6	---
4	Cerebrovascular diseases (I60-I69)	1,513	5.6	---
5	Chronic lower respiratory diseases (J40-J47)	1,200	4.5	---
6	Influenza and pneumonia (J09-J18)	1,140	4.2	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	767	2.8	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	636	2.4	---
9	Pneumonitis due to solids and liquids (J69)	511	1.9	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	471	1.7	---
11	Diabetes mellitus (E10-E14)	403	1.5	---
12	Septicemia (A40-A41)	270	1.0	---
13	Parkinson's disease (G20-G21)	253	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	177	0.7	---
15	Atherosclerosis (I70)	167	0.6	---
...	All other causes (Residual)	6,053	22.5	---
White, male, 100 years and over				
...	All causes	3,566	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,351	37.9	---
2	Malignant neoplasms (C00-C97)	223	6.3	---
3	Influenza and pneumonia (J09-J18)	190	5.3	---
4	Cerebrovascular diseases (I60-I69)	175	4.9	---
5	Alzheimer's disease (G30)	164	4.6	---
6	Chronic lower respiratory diseases (J40-J47)	114	3.2	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	92	2.6	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	79	2.2	---
9	Pneumonitis due to solids and liquids (J69)	76	2.1	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	74	2.1	---
11	Atherosclerosis (I70)	39	1.1	---
12	Septicemia (A40-A41)	36	1.0	---
13	Diabetes mellitus (E10-E14)	35	1.0	---
14	Parkinson's disease (G20-G21)	21	0.6	---
15	Anemias (D50-D64)	17	0.5	---
...	All other causes (Residual)	880	24.7	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, all ages				
...	All causes	1,089,928	100.0	869.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	246,402	22.6	196.7
2	Malignant neoplasms (C00-C97)	234,910	21.6	187.5
3	Chronic lower respiratory diseases (J40-J47)	70,330	6.5	56.1
4	Cerebrovascular diseases (I60-I69)	64,529	5.9	51.5
5	Alzheimer's disease (G30)	52,983	4.9	42.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	42,112	3.9	33.6
7	Diabetes mellitus (E10-E14)	26,974	2.5	21.5
8	Influenza and pneumonia (J09-J18)	23,010	2.1	18.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17,797	1.6	14.2
10	Septicemia (A40-A41)	15,291	1.4	12.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	13,810	1.3	11.0
12	Chronic liver disease and cirrhosis (K70,K73-K74)	10,774	1.0	8.6
13	Parkinson's disease (G20-G21)	9,071	0.8	7.2
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7,925	0.7	6.3
15	Pneumonitis due to solids and liquids (J69)	7,276	0.7	5.8
...	All other causes (Residual)	246,734	22.6	196.9
White, female, 1-4 years				
...	All causes	1,275	100.0	21.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	413	32.4	7.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	158	12.4	2.7
3	Malignant neoplasms (C00-C97)	136	10.7	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	80	6.3	1.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	54	4.2	0.9
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	28	2.2	0.5
7	Influenza and pneumonia (J09-J18)	25	2.0	0.4
8	Septicemia (A40-A41)	18	1.4	*
9	Cerebrovascular diseases (I60-I69)	10	0.8	*
10	Chronic lower respiratory diseases (J40-J47)	9	0.7	*
11	Certain conditions originating in the perinatal period (P00-P96)	8	0.6	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.4	*
13	Anemias (D50-D64)	3	0.2	*
13	Meningitis (G00,G03)	3	0.2	*
13	Acute bronchitis and bronchiolitis (J20-J21)	3	0.2	*
13	Pneumonitis due to solids and liquids (J69)	3	0.2	*
...	All other causes (Residual)	319	25.0	5.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 5-9 years				
...	All causes	726	100.0	9.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	199	27.4	2.6
2	Malignant neoplasms (C00-C97)	148	20.4	2.0
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	60	8.3	0.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	46	6.3	0.6
5	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	22	3.0	0.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	22	3.0	0.3
7	Influenza and pneumonia (J09-J18)	18	2.5	*
8	Cerebrovascular diseases (I60-I69)	15	2.1	*
9	Certain conditions originating in the perinatal period (P00-P96)	8	1.1	*
10	Chronic lower respiratory diseases (J40-J47)	7	1.0	*
11	Septicemia (A40-A41)	6	0.8	*
12	Meningitis (G00,G03)	5	0.7	*
13	Diabetes mellitus (E10-E14)	4	0.6	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.6	*
15	Anemias (D50-D64)	2	0.3	*
15	Aortic aneurysm and dissection (I71)	2	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.3	*
...	All other causes (Residual)	156	21.5	2.1
White, female, 10-14 years				
...	All causes	853	100.0	11.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	212	24.9	2.8
2	Malignant neoplasms (C00-C97)	165	19.3	2.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	70	8.2	0.9
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	55	6.4	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	44	5.2	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	33	3.9	0.4
7	Cerebrovascular diseases (I60-I69)	19	2.2	*
8	Influenza and pneumonia (J09-J18)	16	1.9	*
9	Chronic lower respiratory diseases (J40-J47)	14	1.6	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	1.3	*
11	Septicemia (A40-A41)	9	1.1	*
12	Certain conditions originating in the perinatal period (P00-P96)	8	0.9	*
13	Anemias (D50-D64)	4	0.5	*
14	Diabetes mellitus (E10-E14)	3	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.4	*
...	All other causes (Residual)	187	21.9	2.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 15-19 years				
...	All causes	2,177	100.0	27.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,010	46.4	12.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	335	15.4	4.3
3	Malignant neoplasms (C00-C97)	200	9.2	2.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	98	4.5	1.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	65	3.0	0.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	60	2.8	0.8
7	Cerebrovascular diseases (I60-I69)	21	1.0	0.3
8	Influenza and pneumonia (J09-J18)	18	0.8	*
8	Pregnancy, childbirth and the puerperium (O00-O99)	18	0.8	*
10	Septicemia (A40-A41)	17	0.8	*
11	Diabetes mellitus (E10-E14)	15	0.7	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	0.6	*
13	Chronic lower respiratory diseases (J40-J47)	8	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.3	*
15	Anemias (D50-D64)	5	0.2	*
...	All other causes (Residual)	289	13.3	3.7
White, female, 20-24 years				
...	All causes	3,562	100.0	42.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,539	43.2	18.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	413	11.6	5.0
3	Malignant neoplasms (C00-C97)	313	8.8	3.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	202	5.7	2.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	138	3.9	1.7
6	Pregnancy, childbirth and the puerperium (O00-O99)	80	2.2	1.0
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	75	2.1	0.9
8	Diabetes mellitus (E10-E14)	40	1.1	0.5
9	Cerebrovascular diseases (I60-I69)	35	1.0	0.4
10	Septicemia (A40-A41)	26	0.7	0.3
11	Influenza and pneumonia (J09-J18)	24	0.7	0.3
12	Chronic lower respiratory diseases (J40-J47)	18	0.5	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	0.4	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	11	0.3	*
...	All other causes (Residual)	621	17.4	7.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 25-29 years				
...	All causes	4,490	100.0	56.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,691	37.7	21.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	500	11.1	6.2
3	Malignant neoplasms (C00-C97)	481	10.7	6.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	229	5.1	2.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	199	4.4	2.5
6	Pregnancy, childbirth and the puerperium (O00-O99)	112	2.5	1.4
7	Diabetes mellitus (E10-E14)	70	1.6	0.9
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	62	1.4	0.8
9	Cerebrovascular diseases (I60-I69)	58	1.3	0.7
10	Influenza and pneumonia (J09-J18)	44	1.0	0.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	43	1.0	0.5
12	Septicemia (A40-A41)	39	0.9	0.5
13	Chronic lower respiratory diseases (J40-J47)	36	0.8	0.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	21	0.5	0.3
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	0.3	*
...	All other causes (Residual)	891	19.8	11.1
White, female, 30-34 years				
...	All causes	5,773	100.0	73.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,855	32.1	23.5
2	Malignant neoplasms (C00-C97)	901	15.6	11.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	563	9.8	7.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	415	7.2	5.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	205	3.6	2.6
6	Chronic liver disease and cirrhosis (K70,K73-K74)	119	2.1	1.5
7	Pregnancy, childbirth and the puerperium (O00-O99)	117	2.0	1.5
8	Cerebrovascular diseases (I60-I69)	108	1.9	1.4
9	Diabetes mellitus (E10-E14)	97	1.7	1.2
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	75	1.3	1.0
11	Influenza and pneumonia (J09-J18)	63	1.1	0.8
12	Septicemia (A40-A41)	53	0.9	0.7
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	41	0.7	0.5
14	Chronic lower respiratory diseases (J40-J47)	40	0.7	0.5
15	Human immunodeficiency virus (HIV) disease (B20-B24)	28	0.5	0.4
...	All other causes (Residual)	1,093	18.9	13.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 35-39 years				
...	All causes	7,245	100.0	97.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,749	24.1	23.6
2	Malignant neoplasms (C00-C97)	1,544	21.3	20.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	722	10.0	9.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	619	8.5	8.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	217	3.0	2.9
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	184	2.5	2.5
7	Diabetes mellitus (E10-E14)	169	2.3	2.3
8	Cerebrovascular diseases (I60-I69)	163	2.2	2.2
9	Pregnancy, childbirth and the puerperium (O00-O99)	98	1.4	1.3
10	Influenza and pneumonia (J09-J18)	85	1.2	1.1
11	Chronic lower respiratory diseases (J40-J47)	83	1.1	1.1
12	Septicemia (A40-A41)	78	1.1	1.1
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	66	0.9	0.9
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	65	0.9	0.9
15	Human immunodeficiency virus (HIV) disease (B20-B24)	41	0.6	0.6
...	All other causes (Residual)	1,362	18.8	18.4
White, female, 40-44 years				
...	All causes	12,074	100.0	147.9
1	Malignant neoplasms (C00-C97)	3,289	27.2	40.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,134	17.7	26.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,431	11.9	17.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	769	6.4	9.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	500	4.1	6.1
6	Cerebrovascular diseases (I60-I69)	335	2.8	4.1
7	Diabetes mellitus (E10-E14)	317	2.6	3.9
8	Chronic lower respiratory diseases (J40-J47)	207	1.7	2.5
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	179	1.5	2.2
10	Septicemia (A40-A41)	162	1.3	2.0
11	Influenza and pneumonia (J09-J18)	153	1.3	1.9
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	84	0.7	1.0
13	Human immunodeficiency virus (HIV) disease (B20-B24)	76	0.6	0.9
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	72	0.6	0.9
15	Viral hepatitis (B15-B19)	67	0.6	0.8
...	All other causes (Residual)	2,299	19.0	28.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 45-49 years				
...	All causes	20,492	100.0	237.5
1	Malignant neoplasms (C00-C97)	6,534	31.9	75.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,735	13.3	31.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,678	13.1	31.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,014	4.9	11.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	993	4.8	11.5
6	Cerebrovascular diseases (I60-I69)	654	3.2	7.6
7	Chronic lower respiratory diseases (J40-J47)	557	2.7	6.5
8	Diabetes mellitus (E10-E14)	530	2.6	6.1
9	Septicemia (A40-A41)	286	1.4	3.3
10	Influenza and pneumonia (J09-J18)	243	1.2	2.8
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	187	0.9	2.2
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	160	0.8	1.9
13	Viral hepatitis (B15-B19)	155	0.8	1.8
14	Human immunodeficiency virus (HIV) disease (B20-B24)	118	0.6	1.4
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	104	0.5	1.2
...	All other causes (Residual)	3,544	17.3	41.1
White, female, 50-54 years				
...	All causes	33,080	100.0	358.5
1	Malignant neoplasms (C00-C97)	12,126	36.7	131.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,588	13.9	49.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,073	9.3	33.3
4	Chronic lower respiratory diseases (J40-J47)	1,462	4.4	15.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,455	4.4	15.8
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,110	3.4	12.0
7	Cerebrovascular diseases (I60-I69)	998	3.0	10.8
8	Diabetes mellitus (E10-E14)	920	2.8	10.0
9	Septicemia (A40-A41)	484	1.5	5.2
10	Influenza and pneumonia (J09-J18)	358	1.1	3.9
11	Viral hepatitis (B15-B19)	341	1.0	3.7
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	318	1.0	3.4
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	163	0.5	1.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	150	0.5	1.6
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	150	0.5	1.6
...	All other causes (Residual)	5,384	16.3	58.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 55-59 years				
...	All causes	44,379	100.0	510.5
1	Malignant neoplasms (C00-C97)	17,504	39.4	201.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,670	15.0	76.7
3	Chronic lower respiratory diseases (J40-J47)	2,545	5.7	29.3
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,474	5.6	28.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,501	3.4	17.3
6	Diabetes mellitus (E10-E14)	1,419	3.2	16.3
7	Cerebrovascular diseases (I60-I69)	1,357	3.1	15.6
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	911	2.1	10.5
9	Septicemia (A40-A41)	674	1.5	7.8
10	Influenza and pneumonia (J09-J18)	570	1.3	6.6
11	Viral hepatitis (B15-B19)	517	1.2	5.9
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	511	1.2	5.9
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	302	0.7	3.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	260	0.6	3.0
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	176	0.4	2.0
...	All other causes (Residual)	6,988	15.7	80.4
White, female, 60-64 years				
...	All causes	57,019	100.0	745.6
1	Malignant neoplasms (C00-C97)	23,304	40.9	304.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,306	16.3	121.7
3	Chronic lower respiratory diseases (J40-J47)	3,934	6.9	51.4
4	Diabetes mellitus (E10-E14)	2,077	3.6	27.2
5	Cerebrovascular diseases (I60-I69)	1,877	3.3	24.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,786	3.1	23.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,353	2.4	17.7
8	Septicemia (A40-A41)	995	1.7	13.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	827	1.5	10.8
10	Influenza and pneumonia (J09-J18)	704	1.2	9.2
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	597	1.0	7.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	420	0.7	5.5
13	Viral hepatitis (B15-B19)	339	0.6	4.4
14	Alzheimer's disease (G30)	299	0.5	3.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	286	0.5	3.7
...	All other causes (Residual)	8,915	15.6	116.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 65-69 years				
...	All causes	72,076	100.0	1,158.0
1	Malignant neoplasms (C00-C97)	28,282	39.2	454.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,035	16.7	193.4
3	Chronic lower respiratory diseases (J40-J47)	6,627	9.2	106.5
4	Cerebrovascular diseases (I60-I69)	2,735	3.8	43.9
5	Diabetes mellitus (E10-E14)	2,574	3.6	41.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,679	2.3	27.0
7	Septicemia (A40-A41)	1,232	1.7	19.8
8	Chronic liver disease and cirrhosis (K70,K73-K74)	1,224	1.7	19.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,157	1.6	18.6
10	Influenza and pneumonia (J09-J18)	969	1.3	15.6
11	Alzheimer's disease (G30)	662	0.9	10.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	587	0.8	9.4
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	414	0.6	6.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	403	0.6	6.5
15	Pneumonitis due to solids and liquids (J69)	277	0.4	4.5
...	All other causes (Residual)	11,219	15.6	180.2
White, female, 70-74 years				
...	All causes	84,744	100.0	1,848.5
1	Malignant neoplasms (C00-C97)	29,597	34.9	645.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,166	17.9	330.8
3	Chronic lower respiratory diseases (J40-J47)	8,810	10.4	192.2
4	Cerebrovascular diseases (I60-I69)	4,003	4.7	87.3
5	Diabetes mellitus (E10-E14)	2,782	3.3	60.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,693	2.0	36.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,513	1.8	33.0
8	Alzheimer's disease (G30)	1,498	1.8	32.7
9	Septicemia (A40-A41)	1,459	1.7	31.8
10	Influenza and pneumonia (J09-J18)	1,293	1.5	28.2
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1,019	1.2	22.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	693	0.8	15.1
13	Parkinson's disease (G20-G21)	634	0.7	13.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	508	0.6	11.1
15	Pneumonitis due to solids and liquids (J69)	404	0.5	8.8
...	All other causes (Residual)	13,672	16.1	298.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 75-79 years				
...	All causes	111,241	100.0	3,103.4
1	Malignant neoplasms (C00-C97)	30,928	27.8	862.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	22,240	20.0	620.5
3	Chronic lower respiratory diseases (J40-J47)	10,658	9.6	297.3
4	Cerebrovascular diseases (I60-I69)	6,450	5.8	179.9
5	Alzheimer's disease (G30)	4,117	3.7	114.9
6	Diabetes mellitus (E10-E14)	3,305	3.0	92.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,396	2.2	66.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,037	1.8	56.8
9	Influenza and pneumonia (J09-J18)	2,030	1.8	56.6
10	Septicemia (A40-A41)	1,851	1.7	51.6
11	Parkinson's disease (G20-G21)	1,331	1.2	37.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,181	1.1	32.9
13	Chronic liver disease and cirrhosis (K70,K73-K74)	921	0.8	25.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	766	0.7	21.4
15	Pneumonitis due to solids and liquids (J69)	669	0.6	18.7
...	All other causes (Residual)	20,361	18.3	568.0
White, female, 80-84 years				
...	All causes	159,059	100.0	5,317.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	36,160	22.7	1,208.9
2	Malignant neoplasms (C00-C97)	32,339	20.3	1,081.2
3	Chronic lower respiratory diseases (J40-J47)	12,803	8.0	428.0
4	Cerebrovascular diseases (I60-I69)	10,999	6.9	367.7
5	Alzheimer's disease (G30)	9,120	5.7	304.9
6	Diabetes mellitus (E10-E14)	4,171	2.6	139.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,767	2.4	125.9
8	Influenza and pneumonia (J09-J18)	3,472	2.2	116.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,042	1.9	101.7
10	Septicemia (A40-A41)	2,458	1.5	82.2
11	Parkinson's disease (G20-G21)	2,121	1.3	70.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,976	1.2	66.1
13	Pneumonitis due to solids and liquids (J69)	1,157	0.7	38.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,081	0.7	36.1
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	819	0.5	27.4
...	All other causes (Residual)	33,574	21.1	1,122.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 85-89 years				
...	All causes	200,746	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	53,018	26.4	---
2	Malignant neoplasms (C00-C97)	27,723	13.8	---
3	Cerebrovascular diseases (I60-I69)	15,156	7.5	---
4	Alzheimer's disease (G30)	14,877	7.4	---
5	Chronic lower respiratory diseases (J40-J47)	12,180	6.1	---
6	Influenza and pneumonia (J09-J18)	5,000	2.5	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,871	2.4	---
8	Diabetes mellitus (E10-E14)	4,402	2.2	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,672	1.8	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,133	1.6	---
11	Septicemia (A40-A41)	2,751	1.4	---
12	Parkinson's disease (G20-G21)	2,573	1.3	---
13	Pneumonitis due to solids and liquids (J69)	1,703	0.8	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,315	0.7	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1,073	0.5	---
...	All other causes (Residual)	47,299	23.6	---
White, female, 90-94 years				
...	All causes	168,886	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	50,764	30.1	---
2	Malignant neoplasms (C00-C97)	14,531	8.6	---
3	Alzheimer's disease (G30)	14,260	8.4	---
4	Cerebrovascular diseases (I60-I69)	13,124	7.8	---
5	Chronic lower respiratory diseases (J40-J47)	7,426	4.4	---
6	Influenza and pneumonia (J09-J18)	4,740	2.8	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,194	2.5	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,105	1.8	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,898	1.7	---
10	Diabetes mellitus (E10-E14)	2,881	1.7	---
11	Septicemia (A40-A41)	1,818	1.1	---
12	Pneumonitis due to solids and liquids (J69)	1,549	0.9	---
13	Parkinson's disease (G20-G21)	1,488	0.9	---
14	Atherosclerosis (I70)	912	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	879	0.5	---
...	All other causes (Residual)	44,317	26.2	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 95-99 years				
...	All causes	75,742	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	24,657	32.6	---
2	Alzheimer's disease (G30)	6,582	8.7	---
3	Cerebrovascular diseases (I60-I69)	5,333	7.0	---
4	Malignant neoplasms (C00-C97)	4,229	5.6	---
5	Chronic lower respiratory diseases (J40-J47)	2,470	3.3	---
6	Influenza and pneumonia (J09-J18)	2,443	3.2	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,922	2.5	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,646	2.2	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,165	1.5	---
10	Diabetes mellitus (E10-E14)	1,015	1.3	---
11	Pneumonitis due to solids and liquids (J69)	724	1.0	---
12	Septicemia (A40-A41)	698	0.9	---
13	Atherosclerosis (I70)	557	0.7	---
14	Parkinson's disease (G20-G21)	409	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	299	0.4	---
...	All other causes (Residual)	21,593	28.5	---
White, female, 100 years and over				
...	All causes	17,446	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,919	33.9	---
2	Alzheimer's disease (G30)	1,387	8.0	---
3	Cerebrovascular diseases (I60-I69)	1,048	6.0	---
4	Influenza and pneumonia (J09-J18)	705	4.0	---
5	Malignant neoplasms (C00-C97)	608	3.5	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	457	2.6	---
7	Chronic lower respiratory diseases (J40-J47)	430	2.5	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	413	2.4	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	248	1.4	---
10	Atherosclerosis (I70)	187	1.1	---
11	Diabetes mellitus (E10-E14)	180	1.0	---
12	Pneumonitis due to solids and liquids (J69)	163	0.9	---
13	Septicemia (A40-A41)	133	0.8	---
14	Anemias (D50-D64)	65	0.4	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	57	0.3	---
...	All other causes (Residual)	5,446	31.2	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, all ages				
...	All causes	295,222	100.0	683.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	70,123	23.8	162.3
2	Malignant neoplasms (C00-C97)	67,379	22.8	156.0
3	Cerebrovascular diseases (I60-I69)	15,894	5.4	36.8
4	Diabetes mellitus (E10-E14)	12,983	4.4	30.1
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,709	4.3	29.4
6	Chronic lower respiratory diseases (J40-J47)	9,375	3.2	21.7
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8,241	2.8	19.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,207	2.8	19.0
9	Septicemia (A40-A41)	5,983	2.0	13.8
10	Alzheimer's disease (G30)	5,451	1.8	12.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,200	1.8	12.0
12	Influenza and pneumonia (J09-J18)	5,000	1.7	11.6
13	Certain conditions originating in the perinatal period (P00-P96)	3,958	1.3	9.2
14	Human immunodeficiency virus (HIV) disease (B20-B24)	3,921	1.3	9.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,791	0.9	6.5
...	All other causes (Residual)	58,007	19.6	134.3
Black, both sexes, 1-4 years				
...	All causes	1,021	100.0	37.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	280	27.4	10.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	133	13.0	4.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	108	10.6	3.9
4	Malignant neoplasms (C00-C97)	65	6.4	2.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	44	4.3	1.6
6	Chronic lower respiratory diseases (J40-J47)	30	2.9	1.1
7	Influenza and pneumonia (J09-J18)	21	2.1	0.8
8	Septicemia (A40-A41)	18	1.8	*
8	Cerebrovascular diseases (I60-I69)	18	1.8	*
10	Certain conditions originating in the perinatal period (P00-P96)	17	1.7	*
11	Anemias (D50-D64)	7	0.7	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.6	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.6	*
14	Acute bronchitis and bronchiolitis (J20-J21)	4	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.3	*
...	All other causes (Residual)	261	25.6	9.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 5-9 years				
...	All causes	551	100.0	16.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	189	34.3	5.7
2	Malignant neoplasms (C00-C97)	64	11.6	1.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	45	8.2	1.3
4	Chronic lower respiratory diseases (J40-J47)	41	7.4	1.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	40	7.3	1.2
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	19	3.4	*
7	Complications of medical and surgical care (Y40-Y84,Y88)	8	1.5	*
8	Cerebrovascular diseases (I60-I69)	6	1.1	*
8	Influenza and pneumonia (J09-J18)	6	1.1	*
10	Diabetes mellitus (E10-E14)	5	0.9	*
11	Septicemia (A40-A41)	4	0.7	*
12	Anemias (D50-D64)	3	0.5	*
12	Certain conditions originating in the perinatal period (P00-P96)	3	0.5	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.4	*
14	Hernia (K40-K46)	2	0.4	*
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2	0.4	*
...	All other causes (Residual)	112	20.3	3.3
Black, both sexes, 10-14 years				
...	All causes	645	100.0	19.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	170	26.4	5.0
2	Malignant neoplasms (C00-C97)	84	13.0	2.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	70	10.9	2.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	41	6.4	1.2
5	Chronic lower respiratory diseases (J40-J47)	34	5.3	1.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	32	5.0	0.9
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	25	3.9	0.7
8	Diabetes mellitus (E10-E14)	12	1.9	*
9	Anemias (D50-D64)	10	1.6	*
9	Influenza and pneumonia (J09-J18)	10	1.6	*
11	Cerebrovascular diseases (I60-I69)	9	1.4	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	1.1	*
13	Certain conditions originating in the perinatal period (P00-P96)	5	0.8	*
14	Septicemia (A40-A41)	3	0.5	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.5	*
...	All other causes (Residual)	130	20.2	3.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 15-19 years				
...	All causes	2,316	100.0	64.6
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,035	44.7	28.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	486	21.0	13.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	167	7.2	4.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	95	4.1	2.6
5	Malignant neoplasms (C00-C97)	94	4.1	2.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	45	1.9	1.3
7	Chronic lower respiratory diseases (J40-J47)	23	1.0	0.6
8	Anemias (D50-D64)	18	0.8	*
9	Pregnancy, childbirth and the puerperium (O00-O99)	17	0.7	*
10	Diabetes mellitus (E10-E14)	16	0.7	*
11	Legal intervention (Y35,Y89.0)	14	0.6	*
12	Cerebrovascular diseases (I60-I69)	13	0.6	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	9	0.4	*
14	Septicemia (A40-A41)	8	0.3	*
15	Influenza and pneumonia (J09-J18)	5	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.2	*
...	All other causes (Residual)	266	11.5	7.4
Black, both sexes, 20-24 years				
...	All causes	4,389	100.0	117.5
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,871	42.6	50.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	910	20.7	24.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	357	8.1	9.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	194	4.4	5.2
5	Malignant neoplasms (C00-C97)	156	3.6	4.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	78	1.8	2.1
7	Pregnancy, childbirth and the puerperium (O00-O99)	51	1.2	1.4
8	Anemias (D50-D64)	47	1.1	1.3
9	Diabetes mellitus (E10-E14)	42	1.0	1.1
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	40	0.9	1.1
11	Chronic lower respiratory diseases (J40-J47)	39	0.9	1.0
12	Legal intervention (Y35,Y89.0)	31	0.7	0.8
13	Cerebrovascular diseases (I60-I69)	30	0.7	0.8
14	Influenza and pneumonia (J09-J18)	27	0.6	0.7
15	Septicemia (A40-A41)	25	0.6	0.7
...	All other causes (Residual)	491	11.2	13.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 25-29 years				
...	All causes	4,267	100.0	136.7
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,360	31.9	43.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	866	20.3	27.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	322	7.5	10.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	296	6.9	9.5
5	Malignant neoplasms (C00-C97)	214	5.0	6.9
6	Human immunodeficiency virus (HIV) disease (B20-B24)	160	3.7	5.1
7	Diabetes mellitus (E10-E14)	75	1.8	2.4
8	Pregnancy, childbirth and the puerperium (O00-O99)	69	1.6	2.2
9	Anemias (D50-D64)	58	1.4	1.9
10	Cerebrovascular diseases (I60-I69)	44	1.0	1.4
10	Chronic lower respiratory diseases (J40-J47)	44	1.0	1.4
12	Septicemia (A40-A41)	41	1.0	1.3
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	37	0.9	1.2
14	Influenza and pneumonia (J09-J18)	35	0.8	1.1
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	32	0.7	1.0
...	All other causes (Residual)	614	14.4	19.7
Black, both sexes, 30-34 years				
...	All causes	4,838	100.0	161.7
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,073	22.2	35.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	868	17.9	29.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	575	11.9	19.2
4	Malignant neoplasms (C00-C97)	439	9.1	14.7
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	266	5.5	8.9
6	Human immunodeficiency virus (HIV) disease (B20-B24)	263	5.4	8.8
7	Diabetes mellitus (E10-E14)	129	2.7	4.3
8	Cerebrovascular diseases (I60-I69)	80	1.7	2.7
9	Chronic lower respiratory diseases (J40-J47)	69	1.4	2.3
10	Pregnancy, childbirth and the puerperium (O00-O99)	66	1.4	2.2
11	Anemias (D50-D64)	55	1.1	1.8
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	55	1.1	1.8
13	Septicemia (A40-A41)	49	1.0	1.6
14	Influenza and pneumonia (J09-J18)	46	1.0	1.5
15	Chronic liver disease and cirrhosis (K70,K73-K74)	30	0.6	1.0
...	All other causes (Residual)	775	16.0	25.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 35-39 years				
...	All causes	5,636	100.0	207.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,073	19.0	39.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	799	14.2	29.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	739	13.1	27.2
4	Malignant neoplasms (C00-C97)	666	11.8	24.5
5	Human immunodeficiency virus (HIV) disease (B20-B24)	318	5.6	11.7
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	218	3.9	8.0
7	Diabetes mellitus (E10-E14)	196	3.5	7.2
8	Cerebrovascular diseases (I60-I69)	162	2.9	6.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	83	1.5	3.1
10	Anemias (D50-D64)	71	1.3	2.6
11	Influenza and pneumonia (J09-J18)	66	1.2	2.4
12	Chronic lower respiratory diseases (J40-J47)	65	1.2	2.4
13	Septicemia (A40-A41)	61	1.1	2.2
14	Pregnancy, childbirth and the puerperium (O00-O99)	56	1.0	2.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	55	1.0	2.0
...	All other causes (Residual)	1,008	17.9	37.1
Black, both sexes, 40-44 years				
...	All causes	8,057	100.0	283.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,726	21.4	60.7
2	Malignant neoplasms (C00-C97)	1,282	15.9	45.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	886	11.0	31.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	551	6.8	19.4
5	Human immunodeficiency virus (HIV) disease (B20-B24)	442	5.5	15.5
6	Diabetes mellitus (E10-E14)	312	3.9	11.0
7	Cerebrovascular diseases (I60-I69)	308	3.8	10.8
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	260	3.2	9.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	173	2.1	6.1
10	Septicemia (A40-A41)	155	1.9	5.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	141	1.8	5.0
12	Chronic lower respiratory diseases (J40-J47)	131	1.6	4.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	108	1.3	3.8
14	Influenza and pneumonia (J09-J18)	86	1.1	3.0
15	Anemias (D50-D64)	75	0.9	2.6
...	All other causes (Residual)	1,421	17.6	50.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 45-49 years				
...	All causes	12,757	100.0	441.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,994	23.5	103.5
2	Malignant neoplasms (C00-C97)	2,741	21.5	94.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,111	8.7	38.4
4	Cerebrovascular diseases (I60-I69)	659	5.2	22.8
5	Human immunodeficiency virus (HIV) disease (B20-B24)	618	4.8	21.4
6	Diabetes mellitus (E10-E14)	568	4.5	19.6
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	409	3.2	14.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	291	2.3	10.1
9	Chronic liver disease and cirrhosis (K70,K73-K74)	259	2.0	9.0
10	Chronic lower respiratory diseases (J40-J47)	244	1.9	8.4
11	Septicemia (A40-A41)	223	1.7	7.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	200	1.6	6.9
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	193	1.5	6.7
14	Influenza and pneumonia (J09-J18)	147	1.2	5.1
15	Viral hepatitis (B15-B19)	69	0.5	2.4
...	All other causes (Residual)	2,031	15.9	70.2
Black, both sexes, 50-54 years				
...	All causes	20,198	100.0	703.7
1	Malignant neoplasms (C00-C97)	5,468	27.1	190.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,773	23.6	166.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,268	6.3	44.2
4	Cerebrovascular diseases (I60-I69)	1,008	5.0	35.1
5	Diabetes mellitus (E10-E14)	842	4.2	29.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	689	3.4	24.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	469	2.3	16.3
8	Chronic liver disease and cirrhosis (K70,K73-K74)	461	2.3	16.1
9	Chronic lower respiratory diseases (J40-J47)	447	2.2	15.6
10	Septicemia (A40-A41)	365	1.8	12.7
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	349	1.7	12.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	335	1.7	11.7
13	Influenza and pneumonia (J09-J18)	246	1.2	8.6
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	180	0.9	6.3
15	Viral hepatitis (B15-B19)	171	0.8	6.0
...	All other causes (Residual)	3,127	15.5	108.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 55-59 years				
...	All causes	26,463	100.0	1,063.9
1	Malignant neoplasms (C00-C97)	7,981	30.2	320.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,465	24.4	259.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,271	4.8	51.1
4	Cerebrovascular diseases (I60-I69)	1,248	4.7	50.2
5	Diabetes mellitus (E10-E14)	1,219	4.6	49.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	662	2.5	26.6
7	Chronic lower respiratory diseases (J40-J47)	654	2.5	26.3
8	Human immunodeficiency virus (HIV) disease (B20-B24)	582	2.2	23.4
9	Chronic liver disease and cirrhosis (K70,K73-K74)	565	2.1	22.7
10	Septicemia (A40-A41)	522	2.0	21.0
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	423	1.6	17.0
12	Influenza and pneumonia (J09-J18)	335	1.3	13.5
13	Viral hepatitis (B15-B19)	334	1.3	13.4
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	219	0.8	8.8
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	138	0.5	5.5
...	All other causes (Residual)	3,845	14.5	154.6
Black, both sexes, 60-64 years				
...	All causes	29,195	100.0	1,494.4
1	Malignant neoplasms (C00-C97)	9,261	31.7	474.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,455	25.5	381.6
3	Diabetes mellitus (E10-E14)	1,473	5.0	75.4
4	Cerebrovascular diseases (I60-I69)	1,470	5.0	75.2
5	Chronic lower respiratory diseases (J40-J47)	913	3.1	46.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	826	2.8	42.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	822	2.8	42.1
8	Septicemia (A40-A41)	622	2.1	31.8
9	Chronic liver disease and cirrhosis (K70,K73-K74)	545	1.9	27.9
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	489	1.7	25.0
11	Human immunodeficiency virus (HIV) disease (B20-B24)	384	1.3	19.7
12	Influenza and pneumonia (J09-J18)	373	1.3	19.1
13	Viral hepatitis (B15-B19)	302	1.0	15.5
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	125	0.4	6.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	98	0.3	5.0
15	Pneumonitis due to solids and liquids (J69)	98	0.3	5.0
...	All other causes (Residual)	3,939	13.5	201.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 65-69 years				
...	All causes	27,328	100.0	2,039.7
1	Malignant neoplasms (C00-C97)	8,931	32.7	666.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,863	25.1	512.2
3	Diabetes mellitus (E10-E14)	1,554	5.7	116.0
4	Cerebrovascular diseases (I60-I69)	1,406	5.1	104.9
5	Chronic lower respiratory diseases (J40-J47)	1,062	3.9	79.3
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	856	3.1	63.9
7	Septicemia (A40-A41)	620	2.3	46.3
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	538	2.0	40.2
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	479	1.8	35.8
10	Influenza and pneumonia (J09-J18)	385	1.4	28.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	308	1.1	23.0
12	Human immunodeficiency virus (HIV) disease (B20-B24)	199	0.7	14.9
13	Viral hepatitis (B15-B19)	122	0.4	9.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	111	0.4	8.3
15	Alzheimer's disease (G30)	107	0.4	8.0
...	All other causes (Residual)	3,787	13.9	282.7
Black, both sexes, 70-74 years				
...	All causes	27,823	100.0	2,949.2
1	Malignant neoplasms (C00-C97)	8,623	31.0	914.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,881	24.7	729.4
3	Cerebrovascular diseases (I60-I69)	1,659	6.0	175.9
4	Diabetes mellitus (E10-E14)	1,562	5.6	165.6
5	Chronic lower respiratory diseases (J40-J47)	1,322	4.8	140.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	932	3.3	98.8
7	Septicemia (A40-A41)	676	2.4	71.7
8	Influenza and pneumonia (J09-J18)	483	1.7	51.2
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	462	1.7	49.0
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	453	1.6	48.0
11	Alzheimer's disease (G30)	302	1.1	32.0
12	Chronic liver disease and cirrhosis (K70,K73-K74)	178	0.6	18.9
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	136	0.5	14.4
14	Pneumonitis due to solids and liquids (J69)	125	0.4	13.3
15	Parkinson's disease (G20-G21)	118	0.4	12.5
...	All other causes (Residual)	3,911	14.1	414.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 75-79 years				
...	All causes	28,931	100.0	4,268.2
1	Malignant neoplasms (C00-C97)	7,658	26.5	1,129.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,402	25.6	1,092.0
3	Cerebrovascular diseases (I60-I69)	1,902	6.6	280.6
4	Diabetes mellitus (E10-E14)	1,525	5.3	225.0
5	Chronic lower respiratory diseases (J40-J47)	1,294	4.5	190.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,016	3.5	149.9
7	Septicemia (A40-A41)	693	2.4	102.2
8	Alzheimer's disease (G30)	647	2.2	95.5
9	Influenza and pneumonia (J09-J18)	563	1.9	83.1
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	560	1.9	82.6
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	409	1.4	60.3
12	Pneumonitis due to solids and liquids (J69)	182	0.6	26.9
13	Parkinson's disease (G20-G21)	168	0.6	24.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	150	0.5	22.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	120	0.4	17.7
...	All other causes (Residual)	4,642	16.0	684.8
Black, both sexes, 80-84 years				
...	All causes	29,325	100.0	6,482.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,585	25.9	1,676.6
2	Malignant neoplasms (C00-C97)	6,451	22.0	1,426.0
3	Cerebrovascular diseases (I60-I69)	2,002	6.8	442.5
4	Diabetes mellitus (E10-E14)	1,452	5.0	321.0
5	Chronic lower respiratory diseases (J40-J47)	1,208	4.1	267.0
6	Alzheimer's disease (G30)	1,058	3.6	233.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,024	3.5	226.4
8	Septicemia (A40-A41)	698	2.4	154.3
9	Influenza and pneumonia (J09-J18)	627	2.1	138.6
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	615	2.1	135.9
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	399	1.4	88.2
12	Pneumonitis due to solids and liquids (J69)	237	0.8	52.4
13	Parkinson's disease (G20-G21)	228	0.8	50.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	170	0.6	37.6
15	Atherosclerosis (I70)	92	0.3	20.3
...	All other causes (Residual)	5,479	18.7	1,211.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 85-89 years				
...	All causes	26,885	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,413	27.6	---
2	Malignant neoplasms (C00-C97)	4,468	16.6	---
3	Cerebrovascular diseases (I60-I69)	1,927	7.2	---
4	Alzheimer's disease (G30)	1,398	5.2	---
5	Diabetes mellitus (E10-E14)	1,120	4.2	---
6	Chronic lower respiratory diseases (J40-J47)	989	3.7	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	920	3.4	---
8	Influenza and pneumonia (J09-J18)	677	2.5	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	614	2.3	---
10	Septicemia (A40-A41)	607	2.3	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	306	1.1	---
12	Pneumonitis due to solids and liquids (J69)	254	0.9	---
13	Parkinson's disease (G20-G21)	175	0.7	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	147	0.5	---
15	Atherosclerosis (I70)	97	0.4	---
...	All other causes (Residual)	5,773	21.5	---
Black, both sexes, 90-94 years				
...	All causes	17,561	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,081	28.9	---
2	Malignant neoplasms (C00-C97)	2,041	11.6	---
3	Cerebrovascular diseases (I60-I69)	1,241	7.1	---
4	Alzheimer's disease (G30)	1,099	6.3	---
5	Diabetes mellitus (E10-E14)	627	3.6	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	545	3.1	---
7	Chronic lower respiratory diseases (J40-J47)	528	3.0	---
8	Influenza and pneumonia (J09-J18)	479	2.7	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	472	2.7	---
10	Septicemia (A40-A41)	348	2.0	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	187	1.1	---
12	Pneumonitis due to solids and liquids (J69)	166	0.9	---
13	Parkinson's disease (G20-G21)	95	0.5	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	71	0.4	---
15	Atherosclerosis (I70)	65	0.4	---
...	All other causes (Residual)	4,516	25.7	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 95-99 years				
...	All causes	7,589	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,235	29.5	---
2	Alzheimer's disease (G30)	585	7.7	---
3	Malignant neoplasms (C00-C97)	564	7.4	---
4	Cerebrovascular diseases (I60-I69)	545	7.2	---
5	Essential hypertension and hypertensive renal disease (I10,I12,I15)	255	3.4	---
6	Influenza and pneumonia (J09-J18)	246	3.2	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	219	2.9	---
8	Diabetes mellitus (E10-E14)	204	2.7	---
9	Chronic lower respiratory diseases (J40-J47)	187	2.5	---
10	Septicemia (A40-A41)	141	1.9	---
11	Pneumonitis due to solids and liquids (J69)	69	0.9	---
12	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	67	0.9	---
13	Anemias (D50-D64)	31	0.4	---
14	Parkinson's disease (G20-G21)	29	0.4	---
14	Atherosclerosis (I70)	29	0.4	---
...	All other causes (Residual)	2,183	28.8	---
Black, both sexes, 100 years and over				
...	All causes	2,309	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	787	34.1	---
2	Alzheimer's disease (G30)	184	8.0	---
3	Cerebrovascular diseases (I60-I69)	120	5.2	---
4	Malignant neoplasms (C00-C97)	115	5.0	---
5	Influenza and pneumonia (J09-J18)	81	3.5	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	80	3.5	---
7	Diabetes mellitus (E10-E14)	49	2.1	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	46	2.0	---
9	Chronic lower respiratory diseases (J40-J47)	44	1.9	---
10	Septicemia (A40-A41)	30	1.3	---
11	Pneumonitis due to solids and liquids (J69)	26	1.1	---
12	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19	0.8	---
13	Atherosclerosis (I70)	9	0.4	---
14	Cholelithiasis and other disorders of gallbladder (K80-K82)	8	0.3	---
15	Anemias (D50-D64)	6	0.3	---
...	All other causes (Residual)	705	30.5	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, all ages				
...	All causes	150,586	100.0	728.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	36,140	24.0	174.7
2	Malignant neoplasms (C00-C97)	34,427	22.9	166.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,583	5.7	41.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7,129	4.7	34.5
5	Cerebrovascular diseases (I60-I69)	7,061	4.7	34.1
6	Diabetes mellitus (E10-E14)	6,145	4.1	29.7
7	Chronic lower respiratory diseases (J40-J47)	4,885	3.2	23.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,818	2.5	18.5
9	Septicemia (A40-A41)	2,709	1.8	13.1
10	Human immunodeficiency virus (HIV) disease (B20-B24)	2,546	1.7	12.3
11	Influenza and pneumonia (J09-J18)	2,404	1.6	11.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,310	1.5	11.2
13	Certain conditions originating in the perinatal period (P00-P96)	2,235	1.5	10.8
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,908	1.3	9.2
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,807	1.2	8.7
...	All other causes (Residual)	26,479	17.6	128.0
Black, male, 1-4 years				
...	All causes	578	100.0	41.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	169	29.2	12.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	79	13.7	5.7
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	51	8.8	3.7
4	Malignant neoplasms (C00-C97)	39	6.7	2.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	28	4.8	2.0
6	Chronic lower respiratory diseases (J40-J47)	24	4.2	1.7
7	Influenza and pneumonia (J09-J18)	12	2.1	*
8	Certain conditions originating in the perinatal period (P00-P96)	11	1.9	*
9	Septicemia (A40-A41)	9	1.6	*
9	Cerebrovascular diseases (I60-I69)	9	1.6	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	6	1.0	*
12	Acute bronchitis and bronchiolitis (J20-J21)	4	0.7	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.3	*
14	Anemias (D50-D64)	1	0.2	*
14	Pneumonitis due to solids and liquids (J69)	1	0.2	*
14	Diseases of appendix (K35-K38)	1	0.2	*
14	Hernia (K40-K46)	1	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
...	All other causes (Residual)	130	22.5	9.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 5-9 years				
...	All causes	322	100.0	18.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	118	36.6	6.9
2	Malignant neoplasms (C00-C97)	36	11.2	2.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	30	9.3	1.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	25	7.8	1.5
5	Chronic lower respiratory diseases (J40-J47)	17	5.3	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	8	2.5	*
7	Complications of medical and surgical care (Y40-Y84,Y88)	6	1.9	*
8	Anemias (D50-D64)	3	0.9	*
8	Diabetes mellitus (E10-E14)	3	0.9	*
10	Septicemia (A40-A41)	2	0.6	*
10	Cerebrovascular diseases (I60-I69)	2	0.6	*
10	Influenza and pneumonia (J09-J18)	2	0.6	*
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2	0.6	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.3	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.3	*
14	Hernia (K40-K46)	1	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
...	All other causes (Residual)	64	19.9	3.8
Black, male, 10-14 years				
...	All causes	409	100.0	23.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	120	29.3	6.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	48	11.7	2.8
3	Malignant neoplasms (C00-C97)	47	11.5	2.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	31	7.6	1.8
5	Chronic lower respiratory diseases (J40-J47)	23	5.6	1.3
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	16	3.9	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	16	3.9	*
8	Influenza and pneumonia (J09-J18)	6	1.5	*
9	Anemias (D50-D64)	5	1.2	*
9	Diabetes mellitus (E10-E14)	5	1.2	*
9	Cerebrovascular diseases (I60-I69)	5	1.2	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	1.0	*
13	Certain conditions originating in the perinatal period (P00-P96)	3	0.7	*
14	Septicemia (A40-A41)	2	0.5	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.5	*
...	All other causes (Residual)	76	18.6	4.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 15-19 years				
...	All causes	1,776	100.0	97.3
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	930	52.4	50.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	365	20.6	20.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	126	7.1	6.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	64	3.6	3.5
5	Malignant neoplasms (C00-C97)	57	3.2	3.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	25	1.4	1.4
7	Chronic lower respiratory diseases (J40-J47)	14	0.8	*
7	Legal intervention (Y35,Y89.0)	14	0.8	*
9	Anemias (D50-D64)	11	0.6	*
10	Diabetes mellitus (E10-E14)	7	0.4	*
11	Cerebrovascular diseases (I60-I69)	6	0.3	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.2	*
12	Aortic aneurysm and dissection (I71)	4	0.2	*
14	Septicemia (A40-A41)	3	0.2	*
14	Influenza and pneumonia (J09-J18)	3	0.2	*
...	All other causes (Residual)	143	8.1	7.8
Black, male, 20-24 years				
...	All causes	3,351	100.0	179.0
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,687	50.3	90.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	655	19.5	35.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	297	8.9	15.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	129	3.8	6.9
5	Malignant neoplasms (C00-C97)	91	2.7	4.9
6	Human immunodeficiency virus (HIV) disease (B20-B24)	57	1.7	3.0
7	Chronic lower respiratory diseases (J40-J47)	32	1.0	1.7
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	30	0.9	1.6
9	Legal intervention (Y35,Y89.0)	29	0.9	1.5
10	Anemias (D50-D64)	23	0.7	1.2
11	Diabetes mellitus (E10-E14)	20	0.6	1.1
12	Cerebrovascular diseases (I60-I69)	18	0.5	*
13	Influenza and pneumonia (J09-J18)	17	0.5	*
14	Septicemia (A40-A41)	10	0.3	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.1	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.1	*
...	All other causes (Residual)	246	7.3	13.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 25-29 years				
...	All causes	3,048	100.0	200.7
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,226	40.2	80.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	635	20.8	41.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	255	8.4	16.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	194	6.4	12.8
5	Human immunodeficiency virus (HIV) disease (B20-B24)	110	3.6	7.2
6	Malignant neoplasms (C00-C97)	97	3.2	6.4
7	Diabetes mellitus (E10-E14)	41	1.3	2.7
8	Anemias (D50-D64)	34	1.1	2.2
9	Chronic lower respiratory diseases (J40-J47)	28	0.9	1.8
10	Legal intervention (Y35,Y89.0)	26	0.9	1.7
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	24	0.8	1.6
12	Cerebrovascular diseases (I60-I69)	22	0.7	1.4
13	Influenza and pneumonia (J09-J18)	20	0.7	1.3
14	Septicemia (A40-A41)	19	0.6	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	16	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	0.5	*
...	All other causes (Residual)	285	9.4	18.8
Black, male, 30-34 years				
...	All causes	3,219	100.0	225.8
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	951	29.5	66.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	642	19.9	45.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	373	11.6	26.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	227	7.1	15.9
5	Malignant neoplasms (C00-C97)	171	5.3	12.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	158	4.9	11.1
7	Diabetes mellitus (E10-E14)	60	1.9	4.2
8	Cerebrovascular diseases (I60-I69)	49	1.5	3.4
9	Chronic lower respiratory diseases (J40-J47)	43	1.3	3.0
10	Septicemia (A40-A41)	29	0.9	2.0
11	Anemias (D50-D64)	25	0.8	1.8
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	25	0.8	1.8
13	Influenza and pneumonia (J09-J18)	21	0.7	1.5
14	Aortic aneurysm and dissection (I71)	18	0.6	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	18	0.6	*
...	All other causes (Residual)	409	12.7	28.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 35-39 years				
...	All causes	3,386	100.0	264.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	663	19.6	51.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	660	19.5	51.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	572	16.9	44.7
4	Malignant neoplasms (C00-C97)	227	6.7	17.7
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	172	5.1	13.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	148	4.4	11.6
7	Diabetes mellitus (E10-E14)	113	3.3	8.8
8	Cerebrovascular diseases (I60-I69)	94	2.8	7.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	43	1.3	3.4
10	Influenza and pneumonia (J09-J18)	33	1.0	2.6
11	Chronic lower respiratory diseases (J40-J47)	31	0.9	2.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	30	0.9	2.3
12	Chronic liver disease and cirrhosis (K70,K73-K74)	30	0.9	2.3
14	Septicemia (A40-A41)	29	0.9	2.3
14	Anemias (D50-D64)	29	0.9	2.3
...	All other causes (Residual)	512	15.1	40.0
Black, male, 40-44 years				
...	All causes	4,633	100.0	345.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,133	24.5	84.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	612	13.2	45.7
3	Malignant neoplasms (C00-C97)	470	10.1	35.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	455	9.8	34.0
5	Human immunodeficiency virus (HIV) disease (B20-B24)	246	5.3	18.4
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	196	4.2	14.6
7	Diabetes mellitus (E10-E14)	189	4.1	14.1
8	Cerebrovascular diseases (I60-I69)	165	3.6	12.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	98	2.1	7.3
10	Chronic liver disease and cirrhosis (K70,K73-K74)	89	1.9	6.6
11	Septicemia (A40-A41)	70	1.5	5.2
12	Chronic lower respiratory diseases (J40-J47)	68	1.5	5.1
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	58	1.3	4.3
14	Aortic aneurysm and dissection (I71)	43	0.9	3.2
15	Influenza and pneumonia (J09-J18)	41	0.9	3.1
...	All other causes (Residual)	700	15.1	52.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 45-49 years				
...	All causes	7,209	100.0	529.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,879	26.1	137.9
2	Malignant neoplasms (C00-C97)	1,205	16.7	88.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	735	10.2	53.9
4	Human immunodeficiency virus (HIV) disease (B20-B24)	385	5.3	28.3
5	Cerebrovascular diseases (I60-I69)	366	5.1	26.9
6	Diabetes mellitus (E10-E14)	326	4.5	23.9
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	324	4.5	23.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	161	2.2	11.8
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	147	2.0	10.8
10	Chronic liver disease and cirrhosis (K70,K73-K74)	145	2.0	10.6
11	Septicemia (A40-A41)	118	1.6	8.7
12	Chronic lower respiratory diseases (J40-J47)	115	1.6	8.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	110	1.5	8.1
14	Influenza and pneumonia (J09-J18)	82	1.1	6.0
15	Viral hepatitis (B15-B19)	50	0.7	3.7
...	All other causes (Residual)	1,061	14.7	77.9
Black, male, 50-54 years				
...	All causes	11,643	100.0	865.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,059	26.3	227.5
2	Malignant neoplasms (C00-C97)	2,721	23.4	202.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	872	7.5	64.8
4	Cerebrovascular diseases (I60-I69)	537	4.6	39.9
5	Diabetes mellitus (E10-E14)	488	4.2	36.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	463	4.0	34.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	294	2.5	21.9
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	277	2.4	20.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	249	2.1	18.5
10	Chronic lower respiratory diseases (J40-J47)	227	1.9	16.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	198	1.7	14.7
12	Septicemia (A40-A41)	187	1.6	13.9
13	Influenza and pneumonia (J09-J18)	139	1.2	10.3
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	139	1.2	10.3
15	Viral hepatitis (B15-B19)	117	1.0	8.7
...	All other causes (Residual)	1,676	14.4	124.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 55-59 years				
...	All causes	15,536	100.0	1,356.6
1	Malignant neoplasms (C00-C97)	4,275	27.5	373.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,141	26.7	361.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	911	5.9	79.5
4	Cerebrovascular diseases (I60-I69)	723	4.7	63.1
5	Diabetes mellitus (E10-E14)	670	4.3	58.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	414	2.7	36.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	387	2.5	33.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	360	2.3	31.4
9	Chronic lower respiratory diseases (J40-J47)	332	2.1	29.0
10	Septicemia (A40-A41)	263	1.7	23.0
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	258	1.7	22.5
12	Viral hepatitis (B15-B19)	215	1.4	18.8
13	Influenza and pneumonia (J09-J18)	201	1.3	17.6
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	187	1.2	16.3
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	106	0.7	9.3
...	All other causes (Residual)	2,093	13.5	182.8
Black, male, 60-64 years				
...	All causes	17,126	100.0	1,954.6
1	Malignant neoplasms (C00-C97)	5,202	30.4	593.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,671	27.3	533.1
3	Diabetes mellitus (E10-E14)	822	4.8	93.8
4	Cerebrovascular diseases (I60-I69)	814	4.8	92.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	593	3.5	67.7
6	Chronic lower respiratory diseases (J40-J47)	494	2.9	56.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	455	2.7	51.9
8	Chronic liver disease and cirrhosis (K70,K73-K74)	396	2.3	45.2
9	Septicemia (A40-A41)	325	1.9	37.1
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	278	1.6	31.7
11	Human immunodeficiency virus (HIV) disease (B20-B24)	269	1.6	30.7
12	Influenza and pneumonia (J09-J18)	216	1.3	24.7
13	Viral hepatitis (B15-B19)	210	1.2	24.0
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	103	0.6	11.8
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	61	0.4	7.0
...	All other causes (Residual)	2,217	12.9	253.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 65-69 years				
...	All causes	15,428	100.0	2,639.5
1	Malignant neoplasms (C00-C97)	4,929	31.9	843.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,107	26.6	702.7
3	Diabetes mellitus (E10-E14)	827	5.4	141.5
4	Cerebrovascular diseases (I60-I69)	752	4.9	128.7
5	Chronic lower respiratory diseases (J40-J47)	605	3.9	103.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	453	2.9	77.5
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	362	2.3	61.9
8	Septicemia (A40-A41)	327	2.1	55.9
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	252	1.6	43.1
10	Influenza and pneumonia (J09-J18)	216	1.4	37.0
11	Chronic liver disease and cirrhosis (K70,K73-K74)	210	1.4	35.9
12	Human immunodeficiency virus (HIV) disease (B20-B24)	156	1.0	26.7
13	Viral hepatitis (B15-B19)	74	0.5	12.7
14	Pneumonitis due to solids and liquids (J69)	67	0.4	11.5
15	Aortic aneurysm and dissection (I71)	64	0.4	10.9
...	All other causes (Residual)	2,027	13.1	346.8
Black, male, 70-74 years				
...	All causes	14,690	100.0	3,728.2
1	Malignant neoplasms (C00-C97)	4,565	31.1	1,158.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,775	25.7	958.1
3	Cerebrovascular diseases (I60-I69)	862	5.9	218.8
4	Diabetes mellitus (E10-E14)	780	5.3	198.0
5	Chronic lower respiratory diseases (J40-J47)	748	5.1	189.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	455	3.1	115.5
7	Septicemia (A40-A41)	291	2.0	73.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	283	1.9	71.8
9	Influenza and pneumonia (J09-J18)	253	1.7	64.2
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	235	1.6	59.6
11	Alzheimer's disease (G30)	126	0.9	32.0
12	Chronic liver disease and cirrhosis (K70,K73-K74)	118	0.8	29.9
13	Pneumonitis due to solids and liquids (J69)	87	0.6	22.1
14	Parkinson's disease (G20-G21)	81	0.6	20.6
15	Human immunodeficiency virus (HIV) disease (B20-B24)	76	0.5	19.3
...	All other causes (Residual)	1,955	13.3	496.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 75-79 years				
...	All causes	14,332	100.0	5,402.9
1	Malignant neoplasms (C00-C97)	3,937	27.5	1,484.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,816	26.6	1,438.6
3	Cerebrovascular diseases (I60-I69)	873	6.1	329.1
4	Chronic lower respiratory diseases (J40-J47)	710	5.0	267.7
5	Diabetes mellitus (E10-E14)	665	4.6	250.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	472	3.3	177.9
7	Septicemia (A40-A41)	311	2.2	117.2
8	Influenza and pneumonia (J09-J18)	292	2.0	110.1
9	Alzheimer's disease (G30)	259	1.8	97.6
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	246	1.7	92.7
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	246	1.7	92.7
12	Parkinson's disease (G20-G21)	116	0.8	43.7
13	Pneumonitis due to solids and liquids (J69)	113	0.8	42.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	75	0.5	28.3
15	Chronic liver disease and cirrhosis (K70,K73-K74)	72	0.5	27.1
...	All other causes (Residual)	2,129	14.9	802.6
Black, male, 80-84 years				
...	All causes	12,885	100.0	8,109.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,300	25.6	2,076.9
2	Malignant neoplasms (C00-C97)	3,213	24.9	2,022.2
3	Cerebrovascular diseases (I60-I69)	760	5.9	478.3
4	Chronic lower respiratory diseases (J40-J47)	597	4.6	375.7
5	Diabetes mellitus (E10-E14)	584	4.5	367.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	434	3.4	273.2
7	Alzheimer's disease (G30)	386	3.0	242.9
8	Influenza and pneumonia (J09-J18)	283	2.2	178.1
9	Septicemia (A40-A41)	275	2.1	173.1
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	244	1.9	153.6
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	217	1.7	136.6
12	Parkinson's disease (G20-G21)	136	1.1	85.6
13	Pneumonitis due to solids and liquids (J69)	124	1.0	78.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	75	0.6	47.2
15	Atherosclerosis (I70)	46	0.4	29.0
...	All other causes (Residual)	2,211	17.2	1,391.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 85-89 years				
...	All causes	9,834	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,630	26.7	---
2	Malignant neoplasms (C00-C97)	2,085	21.2	---
3	Cerebrovascular diseases (I60-I69)	582	5.9	---
4	Chronic lower respiratory diseases (J40-J47)	459	4.7	---
5	Alzheimer's disease (G30)	380	3.9	---
6	Diabetes mellitus (E10-E14)	341	3.5	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	332	3.4	---
8	Influenza and pneumonia (J09-J18)	292	3.0	---
9	Septicemia (A40-A41)	228	2.3	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	197	2.0	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	147	1.5	---
12	Pneumonitis due to solids and liquids (J69)	131	1.3	---
13	Parkinson's disease (G20-G21)	98	1.0	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	57	0.6	---
15	Atherosclerosis (I70)	37	0.4	---
...	All other causes (Residual)	1,838	18.7	---
Black, male, 90-94 years				
...	All causes	5,182	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,480	28.6	---
2	Malignant neoplasms (C00-C97)	831	16.0	---
3	Cerebrovascular diseases (I60-I69)	295	5.7	---
4	Chronic lower respiratory diseases (J40-J47)	236	4.6	---
5	Alzheimer's disease (G30)	218	4.2	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	170	3.3	---
7	Influenza and pneumonia (J09-J18)	159	3.1	---
8	Diabetes mellitus (E10-E14)	156	3.0	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	125	2.4	---
10	Septicemia (A40-A41)	119	2.3	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	72	1.4	---
12	Pneumonitis due to solids and liquids (J69)	56	1.1	---
13	Parkinson's disease (G20-G21)	36	0.7	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	26	0.5	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	18	0.3	---
...	All other causes (Residual)	1,185	22.9	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 95-99 years				
...	All causes	1,647	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	499	30.3	---
2	Malignant neoplasms (C00-C97)	195	11.8	---
3	Cerebrovascular diseases (I60-I69)	94	5.7	---
4	Alzheimer's disease (G30)	89	5.4	---
5	Influenza and pneumonia (J09-J18)	71	4.3	---
6	Chronic lower respiratory diseases (J40-J47)	63	3.8	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	58	3.5	---
8	Diabetes mellitus (E10-E14)	40	2.4	---
9	Septicemia (A40-A41)	36	2.2	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	34	2.1	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17	1.0	---
12	Pneumonitis due to solids and liquids (J69)	12	0.7	---
13	Parkinson's disease (G20-G21)	11	0.7	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.5	---
14	Atherosclerosis (I70)	9	0.5	---
...	All other causes (Residual)	410	24.9	---
Black, male, 100 years and over				
...	All causes	349	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	111	31.8	---
2	Malignant neoplasms (C00-C97)	26	7.4	---
3	Alzheimer's disease (G30)	25	7.2	---
4	Essential hypertension and hypertensive renal disease (I10,I12,I15)	15	4.3	---
4	Chronic lower respiratory diseases (J40-J47)	15	4.3	---
6	Influenza and pneumonia (J09-J18)	14	4.0	---
7	Cerebrovascular diseases (I60-I69)	13	3.7	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	3.7	---
9	Pneumonitis due to solids and liquids (J69)	8	2.3	---
10	Diabetes mellitus (E10-E14)	7	2.0	---
11	Septicemia (A40-A41)	6	1.7	---
12	Cholelithiasis and other disorders of gallbladder (K80-K82)	4	1.1	---
13	Nutritional deficiencies (E40-E64)	2	0.6	---
13	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.3	---
15	Anemias (D50-D64)	1	0.3	---
15	Parkinson's disease (G20-G21)	1	0.3	---
15	Aortic aneurysm and dissection (I71)	1	0.3	---
15	Diseases of appendix (K35-K38)	1	0.3	---
15	Hernia (K40-K46)	1	0.3	---
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.3	---
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1	0.3	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.3	---
...	All other causes (Residual)	79	22.6	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, all ages				
...	All causes	144,636	100.0	642.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	33,983	23.5	150.9
2	Malignant neoplasms (C00-C97)	32,952	22.8	146.3
3	Cerebrovascular diseases (I60-I69)	8,833	6.1	39.2
4	Diabetes mellitus (E10-E14)	6,838	4.7	30.4
5	Chronic lower respiratory diseases (J40-J47)	4,490	3.1	19.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,389	3.0	19.5
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,126	2.9	18.3
8	Alzheimer's disease (G30)	3,899	2.7	17.3
9	Septicemia (A40-A41)	3,274	2.3	14.5
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,890	2.0	12.8
11	Influenza and pneumonia (J09-J18)	2,596	1.8	11.5
12	Certain conditions originating in the perinatal period (P00-P96)	1,723	1.2	7.7
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,375	1.0	6.1
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,112	0.8	4.9
15	Chronic liver disease and cirrhosis (K70,K73-K74)	984	0.7	4.4
...	All other causes (Residual)	31,172	21.6	138.4
Black, female, 1-4 years				
...	All causes	443	100.0	33.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	111	25.1	8.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	57	12.9	4.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	54	12.2	4.0
4	Malignant neoplasms (C00-C97)	26	5.9	1.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	16	3.6	*
6	Septicemia (A40-A41)	9	2.0	*
6	Cerebrovascular diseases (I60-I69)	9	2.0	*
6	Influenza and pneumonia (J09-J18)	9	2.0	*
9	Anemias (D50-D64)	6	1.4	*
9	Chronic lower respiratory diseases (J40-J47)	6	1.4	*
9	Certain conditions originating in the perinatal period (P00-P96)	6	1.4	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.9	*
13	Meningitis (G00,G03)	2	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.5	*
15	Tuberculosis (A16-A19)	1	0.2	*
15	Nutritional deficiencies (E40-E64)	1	0.2	*
15	Aortic aneurysm and dissection (I71)	1	0.2	*
15	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.2	*
...	All other causes (Residual)	122	27.5	9.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 5-9 years				
...	All causes	229	100.0	13.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	71	31.0	4.3
2	Malignant neoplasms (C00-C97)	28	12.2	1.7
3	Chronic lower respiratory diseases (J40-J47)	24	10.5	1.5
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	15	6.6	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	15	6.6	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	11	4.8	*
7	Cerebrovascular diseases (I60-I69)	4	1.7	*
7	Influenza and pneumonia (J09-J18)	4	1.7	*
9	Certain conditions originating in the perinatal period (P00-P96)	3	1.3	*
10	Septicemia (A40-A41)	2	0.9	*
10	Diabetes mellitus (E10-E14)	2	0.9	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.9	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.4	*
13	Diseases of appendix (K35-K38)	1	0.4	*
13	Hernia (K40-K46)	1	0.4	*
...	All other causes (Residual)	45	19.7	2.7
Black, female, 10-14 years				
...	All causes	236	100.0	14.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	50	21.2	3.0
2	Malignant neoplasms (C00-C97)	37	15.7	2.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	22	9.3	1.3
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	16	6.8	*
5	Chronic lower respiratory diseases (J40-J47)	11	4.7	*
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	10	4.2	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	9	3.8	*
8	Diabetes mellitus (E10-E14)	7	3.0	*
9	Anemias (D50-D64)	5	2.1	*
10	Cerebrovascular diseases (I60-I69)	4	1.7	*
10	Influenza and pneumonia (J09-J18)	4	1.7	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	1.3	*
13	Certain conditions originating in the perinatal period (P00-P96)	2	0.8	*
14	Septicemia (A40-A41)	1	0.4	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.4	*
...	All other causes (Residual)	53	22.5	3.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 15-19 years				
...	All causes	540	100.0	30.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	121	22.4	6.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	105	19.4	6.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	41	7.6	2.3
4	Malignant neoplasms (C00-C97)	37	6.9	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	31	5.7	1.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	20	3.7	1.1
7	Pregnancy, childbirth and the puerperium (O00-O99)	17	3.1	*
8	Diabetes mellitus (E10-E14)	9	1.7	*
8	Chronic lower respiratory diseases (J40-J47)	9	1.7	*
10	Anemias (D50-D64)	7	1.3	*
10	Cerebrovascular diseases (I60-I69)	7	1.3	*
12	Septicemia (A40-A41)	5	0.9	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	5	0.9	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.7	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.4	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2	0.4	*
15	Influenza and pneumonia (J09-J18)	2	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
...	All other causes (Residual)	114	21.1	6.5
Black, female, 20-24 years				
...	All causes	1,038	100.0	55.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	255	24.6	13.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	184	17.7	9.9
3	Malignant neoplasms (C00-C97)	65	6.3	3.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	65	6.3	3.5
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	60	5.8	3.2
6	Pregnancy, childbirth and the puerperium (O00-O99)	51	4.9	2.7
7	Anemias (D50-D64)	24	2.3	1.3
8	Diabetes mellitus (E10-E14)	22	2.1	1.2
9	Human immunodeficiency virus (HIV) disease (B20-B24)	21	2.0	1.1
10	Septicemia (A40-A41)	15	1.4	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	1.4	*
12	Cerebrovascular diseases (I60-I69)	12	1.2	*
13	Influenza and pneumonia (J09-J18)	10	1.0	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	10	1.0	*
15	Chronic lower respiratory diseases (J40-J47)	7	0.7	*
...	All other causes (Residual)	222	21.4	11.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 25-29 years				
...	All causes	1,219	100.0	76.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	231	18.9	14.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	134	11.0	8.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	128	10.5	8.0
4	Malignant neoplasms (C00-C97)	117	9.6	7.3
5	Pregnancy, childbirth and the puerperium (O00-O99)	69	5.7	4.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	50	4.1	3.1
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	41	3.4	2.6
8	Diabetes mellitus (E10-E14)	34	2.8	2.1
9	Anemias (D50-D64)	24	2.0	1.5
10	Septicemia (A40-A41)	22	1.8	1.4
10	Cerebrovascular diseases (I60-I69)	22	1.8	1.4
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21	1.7	1.3
13	Chronic lower respiratory diseases (J40-J47)	16	1.3	*
14	Influenza and pneumonia (J09-J18)	15	1.2	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	0.7	*
...	All other causes (Residual)	287	23.5	17.9
Black, female, 30-34 years				
...	All causes	1,619	100.0	103.3
1	Malignant neoplasms (C00-C97)	268	16.6	17.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	226	14.0	14.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	202	12.5	12.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	122	7.5	7.8
5	Human immunodeficiency virus (HIV) disease (B20-B24)	105	6.5	6.7
6	Diabetes mellitus (E10-E14)	69	4.3	4.4
7	Pregnancy, childbirth and the puerperium (O00-O99)	66	4.1	4.2
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	39	2.4	2.5
9	Cerebrovascular diseases (I60-I69)	31	1.9	2.0
10	Anemias (D50-D64)	30	1.9	1.9
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	30	1.9	1.9
12	Chronic lower respiratory diseases (J40-J47)	26	1.6	1.7
13	Influenza and pneumonia (J09-J18)	25	1.5	1.6
14	Septicemia (A40-A41)	20	1.2	1.3
15	Chronic liver disease and cirrhosis (K70,K73-K74)	18	1.1	*
...	All other causes (Residual)	342	21.1	21.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 35-39 years				
...	All causes	2,250	100.0	156.4
1	Malignant neoplasms (C00-C97)	439	19.5	30.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	410	18.2	28.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	227	10.1	15.8
4	Human immunodeficiency virus (HIV) disease (B20-B24)	170	7.6	11.8
5	Diabetes mellitus (E10-E14)	83	3.7	5.8
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	79	3.5	5.5
7	Cerebrovascular diseases (I60-I69)	68	3.0	4.7
8	Pregnancy, childbirth and the puerperium (O00-O99)	56	2.5	3.9
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	46	2.0	3.2
10	Anemias (D50-D64)	42	1.9	2.9
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	40	1.8	2.8
12	Chronic lower respiratory diseases (J40-J47)	34	1.5	2.4
13	Influenza and pneumonia (J09-J18)	33	1.5	2.3
14	Septicemia (A40-A41)	32	1.4	2.2
15	Chronic liver disease and cirrhosis (K70,K73-K74)	25	1.1	1.7
...	All other causes (Residual)	466	20.7	32.4
Black, female, 40-44 years				
...	All causes	3,424	100.0	227.8
1	Malignant neoplasms (C00-C97)	812	23.7	54.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	593	17.3	39.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	274	8.0	18.2
4	Human immunodeficiency virus (HIV) disease (B20-B24)	196	5.7	13.0
5	Cerebrovascular diseases (I60-I69)	143	4.2	9.5
6	Diabetes mellitus (E10-E14)	123	3.6	8.2
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	96	2.8	6.4
8	Septicemia (A40-A41)	85	2.5	5.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	75	2.2	5.0
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	64	1.9	4.3
11	Chronic lower respiratory diseases (J40-J47)	63	1.8	4.2
12	Chronic liver disease and cirrhosis (K70,K73-K74)	52	1.5	3.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	50	1.5	3.3
14	Influenza and pneumonia (J09-J18)	45	1.3	3.0
15	Anemias (D50-D64)	39	1.1	2.6
15	Pregnancy, childbirth and the puerperium (O00-O99)	39	1.1	2.6
...	All other causes (Residual)	675	19.7	44.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 45-49 years				
...	All causes	5,548	100.0	362.5
1	Malignant neoplasms (C00-C97)	1,536	27.7	100.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,115	20.1	72.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	376	6.8	24.6
4	Cerebrovascular diseases (I60-I69)	293	5.3	19.1
5	Diabetes mellitus (E10-E14)	242	4.4	15.8
6	Human immunodeficiency virus (HIV) disease (B20-B24)	233	4.2	15.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	130	2.3	8.5
8	Chronic lower respiratory diseases (J40-J47)	129	2.3	8.4
9	Chronic liver disease and cirrhosis (K70,K73-K74)	114	2.1	7.4
10	Septicemia (A40-A41)	105	1.9	6.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	90	1.6	5.9
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	85	1.5	5.6
13	Influenza and pneumonia (J09-J18)	65	1.2	4.2
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	46	0.8	3.0
15	Anemias (D50-D64)	32	0.6	2.1
...	All other causes (Residual)	957	17.2	62.5
Black, female, 50-54 years				
...	All causes	8,555	100.0	560.8
1	Malignant neoplasms (C00-C97)	2,747	32.1	180.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,714	20.0	112.4
3	Cerebrovascular diseases (I60-I69)	471	5.5	30.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	396	4.6	26.0
5	Diabetes mellitus (E10-E14)	354	4.1	23.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	226	2.6	14.8
7	Chronic lower respiratory diseases (J40-J47)	220	2.6	14.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	220	2.6	14.4
9	Septicemia (A40-A41)	178	2.1	11.7
10	Chronic liver disease and cirrhosis (K70,K73-K74)	167	2.0	10.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	137	1.6	9.0
12	Influenza and pneumonia (J09-J18)	107	1.3	7.0
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	72	0.8	4.7
14	Viral hepatitis (B15-B19)	54	0.6	3.5
15	Anemias (D50-D64)	41	0.5	2.7
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	41	0.5	2.7
...	All other causes (Residual)	1,410	16.5	92.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 55-59 years				
...	All causes	10,927	100.0	814.1
1	Malignant neoplasms (C00-C97)	3,706	33.9	276.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,324	21.3	173.1
3	Diabetes mellitus (E10-E14)	549	5.0	40.9
4	Cerebrovascular diseases (I60-I69)	525	4.8	39.1
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	360	3.3	26.8
6	Chronic lower respiratory diseases (J40-J47)	322	2.9	24.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	302	2.8	22.5
8	Septicemia (A40-A41)	259	2.4	19.3
9	Chronic liver disease and cirrhosis (K70,K73-K74)	178	1.6	13.3
10	Human immunodeficiency virus (HIV) disease (B20-B24)	168	1.5	12.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	165	1.5	12.3
12	Influenza and pneumonia (J09-J18)	134	1.2	10.0
13	Viral hepatitis (B15-B19)	119	1.1	8.9
14	Anemias (D50-D64)	50	0.5	3.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	45	0.4	3.4
...	All other causes (Residual)	1,721	15.7	128.2
Black, female, 60-64 years				
...	All causes	12,069	100.0	1,120.2
1	Malignant neoplasms (C00-C97)	4,059	33.6	376.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,784	23.1	258.4
3	Cerebrovascular diseases (I60-I69)	656	5.4	60.9
4	Diabetes mellitus (E10-E14)	651	5.4	60.4
5	Chronic lower respiratory diseases (J40-J47)	419	3.5	38.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	367	3.0	34.1
7	Septicemia (A40-A41)	297	2.5	27.6
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	233	1.9	21.6
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	211	1.7	19.6
10	Influenza and pneumonia (J09-J18)	157	1.3	14.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	149	1.2	13.8
12	Human immunodeficiency virus (HIV) disease (B20-B24)	115	1.0	10.7
13	Viral hepatitis (B15-B19)	92	0.8	8.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	47	0.4	4.4
15	Pneumonitis due to solids and liquids (J69)	39	0.3	3.6
...	All other causes (Residual)	1,793	14.9	166.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 65-69 years				
...	All causes	11,900	100.0	1,575.5
1	Malignant neoplasms (C00-C97)	4,002	33.6	529.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,756	23.2	364.9
3	Diabetes mellitus (E10-E14)	727	6.1	96.3
4	Cerebrovascular diseases (I60-I69)	654	5.5	86.6
5	Chronic lower respiratory diseases (J40-J47)	457	3.8	60.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	403	3.4	53.4
7	Septicemia (A40-A41)	293	2.5	38.8
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	227	1.9	30.1
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	176	1.5	23.3
10	Influenza and pneumonia (J09-J18)	169	1.4	22.4
11	Chronic liver disease and cirrhosis (K70,K73-K74)	98	0.8	13.0
12	Alzheimer's disease (G30)	70	0.6	9.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	58	0.5	7.7
14	Viral hepatitis (B15-B19)	48	0.4	6.4
15	Anemias (D50-D64)	46	0.4	6.1
...	All other causes (Residual)	1,716	14.4	227.2
Black, female, 70-74 years				
...	All causes	13,133	100.0	2,390.6
1	Malignant neoplasms (C00-C97)	4,058	30.9	738.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,106	23.7	565.4
3	Cerebrovascular diseases (I60-I69)	797	6.1	145.1
4	Diabetes mellitus (E10-E14)	782	6.0	142.3
5	Chronic lower respiratory diseases (J40-J47)	574	4.4	104.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	477	3.6	86.8
7	Septicemia (A40-A41)	385	2.9	70.1
8	Influenza and pneumonia (J09-J18)	230	1.8	41.9
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	227	1.7	41.3
10	Alzheimer's disease (G30)	176	1.3	32.0
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	170	1.3	30.9
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	68	0.5	12.4
13	Chronic liver disease and cirrhosis (K70,K73-K74)	60	0.5	10.9
14	Aortic aneurysm and dissection (I71)	51	0.4	9.3
15	Pneumonitis due to solids and liquids (J69)	38	0.3	6.9
...	All other causes (Residual)	1,934	14.7	352.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 75-79 years				
...	All causes	14,599	100.0	3,538.6
1	Malignant neoplasms (C00-C97)	3,721	25.5	901.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,586	24.6	869.2
3	Cerebrovascular diseases (I60-I69)	1,029	7.0	249.4
4	Diabetes mellitus (E10-E14)	860	5.9	208.5
5	Chronic lower respiratory diseases (J40-J47)	584	4.0	141.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	544	3.7	131.9
7	Alzheimer's disease (G30)	388	2.7	94.0
8	Septicemia (A40-A41)	382	2.6	92.6
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	314	2.2	76.1
10	Influenza and pneumonia (J09-J18)	271	1.9	65.7
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	163	1.1	39.5
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	75	0.5	18.2
13	Pneumonitis due to solids and liquids (J69)	69	0.5	16.7
14	Parkinson's disease (G20-G21)	52	0.4	12.6
15	Chronic liver disease and cirrhosis (K70,K73-K74)	48	0.3	11.6
...	All other causes (Residual)	2,513	17.2	609.1
Black, female, 80-84 years				
...	All causes	16,440	100.0	5,601.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,285	26.1	1,459.9
2	Malignant neoplasms (C00-C97)	3,238	19.7	1,103.2
3	Cerebrovascular diseases (I60-I69)	1,242	7.6	423.2
4	Diabetes mellitus (E10-E14)	868	5.3	295.7
5	Alzheimer's disease (G30)	672	4.1	229.0
6	Chronic lower respiratory diseases (J40-J47)	611	3.7	208.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	590	3.6	201.0
8	Septicemia (A40-A41)	423	2.6	144.1
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	371	2.3	126.4
10	Influenza and pneumonia (J09-J18)	344	2.1	117.2
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	182	1.1	62.0
12	Pneumonitis due to solids and liquids (J69)	113	0.7	38.5
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	95	0.6	32.4
14	Parkinson's disease (G20-G21)	92	0.6	31.3
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	54	0.3	18.4
...	All other causes (Residual)	3,260	19.8	1,110.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 85-89 years				
...	All causes	17,051	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,783	28.1	---
2	Malignant neoplasms (C00-C97)	2,383	14.0	---
3	Cerebrovascular diseases (I60-I69)	1,345	7.9	---
4	Alzheimer's disease (G30)	1,018	6.0	---
5	Diabetes mellitus (E10-E14)	779	4.6	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	588	3.4	---
7	Chronic lower respiratory diseases (J40-J47)	530	3.1	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	417	2.4	---
9	Influenza and pneumonia (J09-J18)	385	2.3	---
10	Septicemia (A40-A41)	379	2.2	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	159	0.9	---
12	Pneumonitis due to solids and liquids (J69)	123	0.7	---
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	90	0.5	---
14	Parkinson's disease (G20-G21)	77	0.5	---
15	Atherosclerosis (I70)	60	0.4	---
...	All other causes (Residual)	3,935	23.1	---
Black, female, 90-94 years				
...	All causes	12,379	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,601	29.1	---
2	Malignant neoplasms (C00-C97)	1,210	9.8	---
3	Cerebrovascular diseases (I60-I69)	946	7.6	---
4	Alzheimer's disease (G30)	881	7.1	---
5	Diabetes mellitus (E10-E14)	471	3.8	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	375	3.0	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	347	2.8	---
8	Influenza and pneumonia (J09-J18)	320	2.6	---
9	Chronic lower respiratory diseases (J40-J47)	292	2.4	---
10	Septicemia (A40-A41)	229	1.8	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	115	0.9	---
12	Pneumonitis due to solids and liquids (J69)	110	0.9	---
13	Parkinson's disease (G20-G21)	59	0.5	---
14	Atherosclerosis (I70)	48	0.4	---
15	Anemias (D50-D64)	46	0.4	---
...	All other causes (Residual)	3,329	26.9	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 95-99 years				
...	All causes	5,942	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,736	29.2	---
2	Alzheimer's disease (G30)	496	8.3	---
3	Cerebrovascular diseases (I60-I69)	451	7.6	---
4	Malignant neoplasms (C00-C97)	369	6.2	---
5	Essential hypertension and hypertensive renal disease (I10,I12,I15)	221	3.7	---
6	Influenza and pneumonia (J09-J18)	175	2.9	---
7	Diabetes mellitus (E10-E14)	164	2.8	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	161	2.7	---
9	Chronic lower respiratory diseases (J40-J47)	124	2.1	---
10	Septicemia (A40-A41)	105	1.8	---
11	Pneumonitis due to solids and liquids (J69)	57	1.0	---
12	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	50	0.8	---
13	Anemias (D50-D64)	27	0.5	---
14	Atherosclerosis (I70)	20	0.3	---
15	Parkinson's disease (G20-G21)	18	0.3	---
...	All other causes (Residual)	1,768	29.8	---
Black, female, 100 years and over				
...	All causes	1,960	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	676	34.5	---
2	Alzheimer's disease (G30)	159	8.1	---
3	Cerebrovascular diseases (I60-I69)	107	5.5	---
4	Malignant neoplasms (C00-C97)	89	4.5	---
5	Influenza and pneumonia (J09-J18)	67	3.4	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	65	3.3	---
7	Diabetes mellitus (E10-E14)	42	2.1	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	33	1.7	---
9	Chronic lower respiratory diseases (J40-J47)	29	1.5	---
10	Septicemia (A40-A41)	24	1.2	---
11	Pneumonitis due to solids and liquids (J69)	18	0.9	---
12	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17	0.9	---
13	Atherosclerosis (I70)	9	0.5	---
14	Anemias (D50-D64)	5	0.3	---
15	Aortic aneurysm and dissection (I71)	4	0.2	---
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	4	0.2	---
...	All other causes (Residual)	612	31.2	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, all ages				
...	All causes	16,527	100.0	375.7
1	Malignant neoplasms (C00-C97)	3,019	18.3	68.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,950	17.8	67.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,922	11.6	43.7
4	Diabetes mellitus (E10-E14)	985	6.0	22.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	918	5.6	20.9
6	Chronic lower respiratory diseases (J40-J47)	708	4.3	16.1
7	Cerebrovascular diseases (I60-I69)	580	3.5	13.2
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	485	2.9	11.0
9	Influenza and pneumonia (J09-J18)	306	1.9	7.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	295	1.8	6.7
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	263	1.6	6.0
12	Septicemia (A40-A41)	255	1.5	5.8
13	Alzheimer's disease (G30)	217	1.3	4.9
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	150	0.9	3.4
15	Certain conditions originating in the perinatal period (P00-P96)	129	0.8	2.9
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	129	0.8	2.9
...	All other causes (Residual)	3,216	19.5	73.1
American Indian, both sexes, 1-4 years				
...	All causes	89	100.0	28.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	39	43.8	12.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	11	12.4	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	6.7	*
4	Malignant neoplasms (C00-C97)	5	5.6	*
5	Septicemia (A40-A41)	3	3.4	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	1.1	*
6	Cerebrovascular diseases (I60-I69)	1	1.1	*
6	Influenza and pneumonia (J09-J18)	1	1.1	*
6	Hernia (K40-K46)	1	1.1	*
...	All other causes (Residual)	21	23.6	6.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 5-9 years				
...	All causes	50	100.0	12.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	33	66.0	8.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	4.0	*
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	4.0	*
4	Malignant neoplasms (C00-C97)	1	2.0	*
4	Anemias (D50-D64)	1	2.0	*
4	Influenza and pneumonia (J09-J18)	1	2.0	*
4	Certain conditions originating in the perinatal period (P00-P96)	1	2.0	*
...	All other causes (Residual)	9	18.0	*
American Indian, both sexes, 10-14 years				
...	All causes	51	100.0	13.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18	35.3	*
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	10	19.6	*
3	Malignant neoplasms (C00-C97)	5	9.8	*
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	9.8	*
5	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	3.9	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	3.9	*
7	Cerebrovascular diseases (I60-I69)	1	2.0	*
7	Influenza and pneumonia (J09-J18)	1	2.0	*
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1	2.0	*
...	All other causes (Residual)	6	11.8	*
American Indian, both sexes, 15-19 years				
...	All causes	203	100.0	53.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	95	46.8	25.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	56	27.6	14.8
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	19	9.4	*
4	Malignant neoplasms (C00-C97)	5	2.5	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	2.5	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	1.0	*
7	Septicemia (A40-A41)	1	0.5	*
7	Diabetes mellitus (E10-E14)	1	0.5	*
7	Cerebrovascular diseases (I60-I69)	1	0.5	*
7	Influenza and pneumonia (J09-J18)	1	0.5	*
7	Pregnancy, childbirth and the puerperium (O00-O99)	1	0.5	*
...	All other causes (Residual)	16	7.9	*

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 20-24 years				
...	All causes	401	100.0	103.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	174	43.4	44.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	96	23.9	24.8
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	45	11.2	11.6
4	Malignant neoplasms (C00-C97)	10	2.5	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	9	2.2	*
6	Chronic liver disease and cirrhosis (K70,K73-K74)	6	1.5	*
7	Legal intervention (Y35,Y89.0)	4	1.0	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.7	*
8	Diabetes mellitus (E10-E14)	3	0.7	*
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.7	*
11	Septicemia (A40-A41)	2	0.5	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2	0.5	*
11	Pneumonitis due to solids and liquids (J69)	2	0.5	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.5	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.2	*
15	Chronic lower respiratory diseases (J40-J47)	1	0.2	*
15	Pregnancy, childbirth and the puerperium (O00-O99)	1	0.2	*
...	All other causes (Residual)	37	9.2	9.5
American Indian, both sexes, 25-29 years				
...	All causes	469	100.0	134.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	189	40.3	54.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	71	15.1	20.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	38	8.1	10.9
4	Chronic liver disease and cirrhosis (K70,K73-K74)	28	6.0	8.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	25	5.3	7.2
6	Malignant neoplasms (C00-C97)	15	3.2	*
7	Pregnancy, childbirth and the puerperium (O00-O99)	6	1.3	*
8	Septicemia (A40-A41)	4	0.9	*
8	Cerebrovascular diseases (I60-I69)	4	0.9	*
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	0.9	*
8	Legal intervention (Y35,Y89.0)	4	0.9	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	3	0.6	*
12	Influenza and pneumonia (J09-J18)	3	0.6	*
14	Diabetes mellitus (E10-E14)	2	0.4	*
14	Pneumonitis due to solids and liquids (J69)	2	0.4	*
...	All other causes (Residual)	71	15.1	20.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 30-34 years				
...	All causes	465	100.0	142.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	165	35.5	50.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	47	10.1	14.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	44	9.5	13.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	37	8.0	11.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	37	8.0	11.3
6	Malignant neoplasms (C00-C97)	19	4.1	*
7	Diabetes mellitus (E10-E14)	11	2.4	*
8	Influenza and pneumonia (J09-J18)	8	1.7	*
9	Septicemia (A40-A41)	7	1.5	*
10	Cerebrovascular diseases (I60-I69)	4	0.9	*
10	Pregnancy, childbirth and the puerperium (O00-O99)	4	0.9	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.6	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.4	*
13	Chronic lower respiratory diseases (J40-J47)	2	0.4	*
13	Pneumonitis due to solids and liquids (J69)	2	0.4	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
13	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	2	0.4	*
...	All other causes (Residual)	69	14.8	21.1
American Indian, both sexes, 35-39 years				
...	All causes	510	100.0	173.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	158	31.0	53.8
2	Chronic liver disease and cirrhosis (K70,K73-K74)	61	12.0	20.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	47	9.2	16.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	46	9.0	15.7
5	Malignant neoplasms (C00-C97)	30	5.9	10.2
6	Diabetes mellitus (E10-E14)	18	3.5	*
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	15	2.9	*
8	Septicemia (A40-A41)	9	1.8	*
9	Viral hepatitis (B15-B19)	6	1.2	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	1.2	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	5	1.0	*
11	Pregnancy, childbirth and the puerperium (O00-O99)	5	1.0	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	1.0	*
14	Influenza and pneumonia (J09-J18)	3	0.6	*
15	Cerebrovascular diseases (I60-I69)	2	0.4	*
15	Chronic lower respiratory diseases (J40-J47)	2	0.4	*
15	Infections of kidney (N10-N12,N13.6,N15.1)	2	0.4	*
...	All other causes (Residual)	90	17.6	30.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 40-44 years				
...	All causes	688	100.0	239.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	172	25.0	59.9
2	Chronic liver disease and cirrhosis (K70,K73-K74)	109	15.8	37.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	90	13.1	31.3
4	Malignant neoplasms (C00-C97)	58	8.4	20.2
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	32	4.7	11.1
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	24	3.5	8.4
7	Diabetes mellitus (E10-E14)	22	3.2	7.7
8	Cerebrovascular diseases (I60-I69)	12	1.7	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	8	1.2	*
10	Septicemia (A40-A41)	7	1.0	*
10	Viral hepatitis (B15-B19)	7	1.0	*
10	Influenza and pneumonia (J09-J18)	7	1.0	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	1.0	*
14	Chronic lower respiratory diseases (J40-J47)	5	0.7	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2	0.3	*
15	Pregnancy, childbirth and the puerperium (O00-O99)	2	0.3	*
15	Legal intervention (Y35,Y89.0)	2	0.3	*
...	All other causes (Residual)	122	17.7	42.5
American Indian, both sexes, 45-49 years				
...	All causes	965	100.0	344.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	188	19.5	67.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	140	14.5	50.0
3	Chronic liver disease and cirrhosis (K70,K73-K74)	131	13.6	46.8
4	Malignant neoplasms (C00-C97)	115	11.9	41.1
5	Diabetes mellitus (E10-E14)	58	6.0	20.7
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	45	4.7	16.1
7	Cerebrovascular diseases (I60-I69)	23	2.4	8.2
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	23	2.4	8.2
9	Septicemia (A40-A41)	17	1.8	*
10	Viral hepatitis (B15-B19)	13	1.3	*
10	Influenza and pneumonia (J09-J18)	13	1.3	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	1.1	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	7	0.7	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7	0.7	*
15	Chronic lower respiratory diseases (J40-J47)	6	0.6	*
...	All other causes (Residual)	168	17.4	60.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 50-54 years				
...	All causes	1,299	100.0	486.7
1	Malignant neoplasms (C00-C97)	236	18.2	88.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	205	15.8	76.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	193	14.9	72.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	171	13.2	64.1
5	Diabetes mellitus (E10-E14)	72	5.5	27.0
6	Chronic lower respiratory diseases (J40-J47)	34	2.6	12.7
7	Cerebrovascular diseases (I60-I69)	26	2.0	9.7
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	25	1.9	9.4
9	Viral hepatitis (B15-B19)	21	1.6	7.9
10	Septicemia (A40-A41)	20	1.5	7.5
11	Influenza and pneumonia (J09-J18)	18	1.4	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	1.2	*
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	13	1.0	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	8	0.6	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	6	0.5	*
...	All other causes (Residual)	236	18.2	88.4
American Indian, both sexes, 55-59 years				
...	All causes	1,482	100.0	672.2
1	Malignant neoplasms (C00-C97)	366	24.7	166.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	238	16.1	107.9
3	Chronic liver disease and cirrhosis (K70,K73-K74)	132	8.9	59.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	132	8.9	59.9
5	Diabetes mellitus (E10-E14)	107	7.2	48.5
6	Chronic lower respiratory diseases (J40-J47)	61	4.1	27.7
7	Cerebrovascular diseases (I60-I69)	39	2.6	17.7
8	Viral hepatitis (B15-B19)	32	2.2	14.5
9	Influenza and pneumonia (J09-J18)	29	2.0	13.2
10	Septicemia (A40-A41)	25	1.7	11.3
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	23	1.6	10.4
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22	1.5	10.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	13	0.9	*
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	12	0.8	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	7	0.5	*
...	All other causes (Residual)	244	16.5	110.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 60-64 years				
...	All causes	1,440	100.0	868.9
1	Malignant neoplasms (C00-C97)	389	27.0	234.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	299	20.8	180.4
3	Diabetes mellitus (E10-E14)	126	8.8	76.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	98	6.8	59.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	82	5.7	49.5
6	Chronic lower respiratory diseases (J40-J47)	80	5.6	48.3
7	Cerebrovascular diseases (I60-I69)	51	3.5	30.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	1.6	13.9
9	Influenza and pneumonia (J09-J18)	21	1.5	12.7
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	20	1.4	12.1
11	Septicemia (A40-A41)	18	1.3	*
12	Viral hepatitis (B15-B19)	16	1.1	*
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	13	0.9	*
14	Pneumonitis due to solids and liquids (J69)	5	0.3	*
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	0.3	*
...	All other causes (Residual)	194	13.5	117.1
American Indian, both sexes, 65-69 years				
...	All causes	1,572	100.0	1,373.4
1	Malignant neoplasms (C00-C97)	465	29.6	406.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	324	20.6	283.1
3	Diabetes mellitus (E10-E14)	139	8.8	121.4
4	Chronic lower respiratory diseases (J40-J47)	102	6.5	89.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	69	4.4	60.3
6	Cerebrovascular diseases (I60-I69)	53	3.4	46.3
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	50	3.2	43.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	29	1.8	25.3
9	Septicemia (A40-A41)	24	1.5	21.0
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	20	1.3	17.5
11	Influenza and pneumonia (J09-J18)	18	1.1	*
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	12	0.8	*
13	Pneumonitis due to solids and liquids (J69)	11	0.7	*
14	Viral hepatitis (B15-B19)	9	0.6	*
15	Parkinson's disease (G20-G21)	6	0.4	*
15	Alzheimer's disease (G30)	6	0.4	*
...	All other causes (Residual)	235	14.9	205.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 70-74 years				
...	All causes	1,555	100.0	2,079.7
1	Malignant neoplasms (C00-C97)	427	27.5	571.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	323	20.8	432.0
3	Diabetes mellitus (E10-E14)	122	7.8	163.2
4	Chronic lower respiratory diseases (J40-J47)	106	6.8	141.8
5	Cerebrovascular diseases (I60-I69)	81	5.2	108.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	63	4.1	84.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	42	2.7	56.2
8	Chronic liver disease and cirrhosis (K70,K73-K74)	39	2.5	52.2
9	Influenza and pneumonia (J09-J18)	28	1.8	37.4
10	Septicemia (A40-A41)	27	1.7	36.1
11	Alzheimer's disease (G30)	15	1.0	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10	0.6	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.5	*
13	Parkinson's disease (G20-G21)	7	0.5	*
13	Pneumonitis due to solids and liquids (J69)	7	0.5	*
...	All other causes (Residual)	251	16.1	335.7
American Indian, both sexes, 75-79 years				
...	All causes	1,474	100.0	3,040.4
1	Malignant neoplasms (C00-C97)	361	24.5	744.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	344	23.3	709.6
3	Chronic lower respiratory diseases (J40-J47)	104	7.1	214.5
4	Diabetes mellitus (E10-E14)	102	6.9	210.4
5	Cerebrovascular diseases (I60-I69)	74	5.0	152.6
6	Influenza and pneumonia (J09-J18)	41	2.8	84.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	33	2.2	68.1
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	32	2.2	66.0
9	Septicemia (A40-A41)	29	2.0	59.8
10	Chronic liver disease and cirrhosis (K70,K73-K74)	25	1.7	51.6
11	Alzheimer's disease (G30)	24	1.6	49.5
12	Pneumonitis due to solids and liquids (J69)	13	0.9	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	12	0.8	*
14	Parkinson's disease (G20-G21)	11	0.7	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.5	*
...	All other causes (Residual)	261	17.7	538.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 80-84 years				
...	All causes	1,387	100.0	4,642.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	349	25.2	1,168.1
2	Malignant neoplasms (C00-C97)	273	19.7	913.7
3	Diabetes mellitus (E10-E14)	100	7.2	334.7
4	Chronic lower respiratory diseases (J40-J47)	94	6.8	314.6
5	Cerebrovascular diseases (I60-I69)	76	5.5	254.4
6	Alzheimer's disease (G30)	47	3.4	157.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	43	3.1	143.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	36	2.6	120.5
9	Influenza and pneumonia (J09-J18)	31	2.2	103.8
10	Septicemia (A40-A41)	28	2.0	93.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	17	1.2	*
12	Parkinson's disease (G20-G21)	16	1.2	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	14	1.0	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.6	*
14	Pneumonitis due to solids and liquids (J69)	8	0.6	*
...	All other causes (Residual)	247	17.8	826.7
American Indian, both sexes, 85-89 years				
...	All causes	1,126	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	268	23.8	---
2	Malignant neoplasms (C00-C97)	160	14.2	---
3	Cerebrovascular diseases (I60-I69)	71	6.3	---
4	Diabetes mellitus (E10-E14)	64	5.7	---
5	Chronic lower respiratory diseases (J40-J47)	63	5.6	---
6	Alzheimer's disease (G30)	57	5.1	---
7	Influenza and pneumonia (J09-J18)	42	3.7	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	36	3.2	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	28	2.5	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	18	1.6	---
11	Septicemia (A40-A41)	17	1.5	---
12	Parkinson's disease (G20-G21)	16	1.4	---
13	Pneumonitis due to solids and liquids (J69)	12	1.1	---
14	Chronic liver disease and cirrhosis (K70,K73-K74)	8	0.7	---
15	Anemias (D50-D64)	6	0.5	---
...	All other causes (Residual)	260	23.1	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 90-94 years				
...	All causes	647	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	157	24.3	---
2	Malignant neoplasms (C00-C97)	61	9.4	---
3	Alzheimer's disease (G30)	44	6.8	---
4	Cerebrovascular diseases (I60-I69)	42	6.5	---
5	Chronic lower respiratory diseases (J40-J47)	37	5.7	---
6	Diabetes mellitus (E10-E14)	32	4.9	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	25	3.9	---
8	Influenza and pneumonia (J09-J18)	20	3.1	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14	2.2	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	13	2.0	---
11	Septicemia (A40-A41)	12	1.9	---
12	Pneumonitis due to solids and liquids (J69)	11	1.7	---
13	Anemias (D50-D64)	4	0.6	---
13	Parkinson's disease (G20-G21)	4	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.5	---
15	Peptic ulcer (K25-K28)	3	0.5	---
15	Chronic liver disease and cirrhosis (K70,K73-K74)	3	0.5	---
...	All other causes (Residual)	162	25.0	---
American Indian, both sexes, 95-99 years				
...	All causes	220	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	62	28.2	---
2	Cerebrovascular diseases (I60-I69)	16	7.3	---
3	Alzheimer's disease (G30)	13	5.9	---
4	Malignant neoplasms (C00-C97)	12	5.5	---
5	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9	4.1	---
5	Influenza and pneumonia (J09-J18)	9	4.1	---
5	Chronic lower respiratory diseases (J40-J47)	9	4.1	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7	3.2	---
9	Diabetes mellitus (E10-E14)	6	2.7	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	2.3	---
11	Septicemia (A40-A41)	2	0.9	---
11	Parkinson's disease (G20-G21)	2	0.9	---
13	Tuberculosis (A16-A19)	1	0.5	---
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.5	---
13	Anemias (D50-D64)	1	0.5	---
13	Nutritional deficiencies (E40-E64)	1	0.5	---
13	Pneumonitis due to solids and liquids (J69)	1	0.5	---
13	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.5	---
13	Hyperplasia of prostate (N40)	1	0.5	---
...	All other causes (Residual)	61	27.7	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 100 years and over				
...	All causes	64	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	18	28.1	---
2	Influenza and pneumonia (J09-J18)	7	10.9	---
3	Alzheimer's disease (G30)	6	9.4	---
4	Malignant neoplasms (C00-C97)	4	6.3	---
5	Cerebrovascular diseases (I60-I69)	2	3.1	---
5	Pneumonitis due to solids and liquids (J69)	2	3.1	---
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2	3.1	---
8	Parkinson's disease (G20-G21)	1	1.6	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	1.6	---
8	Chronic lower respiratory diseases (J40-J47)	1	1.6	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	1.6	---
...	All other causes (Residual)	19	29.7	---
American Indian, male, all ages				
...	All causes	9,067	100.0	410.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,731	19.1	78.3
2	Malignant neoplasms (C00-C97)	1,559	17.2	70.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,278	14.1	57.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	517	5.7	23.4
5	Diabetes mellitus (E10-E14)	514	5.7	23.3
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	386	4.3	17.5
7	Chronic lower respiratory diseases (J40-J47)	370	4.1	16.7
8	Cerebrovascular diseases (I60-I69)	266	2.9	12.0
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	207	2.3	9.4
10	Influenza and pneumonia (J09-J18)	144	1.6	6.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	143	1.6	6.5
12	Septicemia (A40-A41)	114	1.3	5.2
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	74	0.8	3.3
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	73	0.8	3.3
15	Certain conditions originating in the perinatal period (P00-P96)	72	0.8	3.3
...	All other causes (Residual)	1,619	17.9	73.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 1-4 years				
...	All causes	54	100.0	34.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	28	51.9	17.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	11.1	*
3	Malignant neoplasms (C00-C97)	4	7.4	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	5.6	*
5	Cerebrovascular diseases (I60-I69)	1	1.9	*
...	All other causes (Residual)	12	22.2	*
American Indian, male, 5-9 years				
...	All causes	34	100.0	17.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	28	82.4	14.3
2	Malignant neoplasms (C00-C97)	1	2.9	*
2	Influenza and pneumonia (J09-J18)	1	2.9	*
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1	2.9	*
...	All other causes (Residual)	3	8.8	*
American Indian, male, 10-14 years				
...	All causes	31	100.0	16.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9	29.0	*
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8	25.8	*
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	12.9	*
4	Malignant neoplasms (C00-C97)	3	9.7	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	6.5	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	3.2	*
...	All other causes (Residual)	4	12.9	*
American Indian, male, 15-19 years				
...	All causes	143	100.0	73.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	61	42.7	31.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	43	30.1	22.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	14	9.8	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	3.5	*
5	Malignant neoplasms (C00-C97)	3	2.1	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	1.4	*
7	Diabetes mellitus (E10-E14)	1	0.7	*
7	Cerebrovascular diseases (I60-I69)	1	0.7	*
7	Influenza and pneumonia (J09-J18)	1	0.7	*
...	All other causes (Residual)	12	8.4	*

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 20-24 years				
...	All causes	284	100.0	140.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	126	44.4	62.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	73	25.7	36.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	35	12.3	17.3
4	Malignant neoplasms (C00-C97)	6	2.1	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	1.8	*
5	Chronic liver disease and cirrhosis (K70,K73-K74)	5	1.8	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	1.1	*
7	Legal intervention (Y35,Y89.0)	3	1.1	*
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2	0.7	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.7	*
11	Septicemia (A40-A41)	1	0.4	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.4	*
11	Diabetes mellitus (E10-E14)	1	0.4	*
11	Pneumonitis due to solids and liquids (J69)	1	0.4	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.4	*
...	All other causes (Residual)	19	6.7	*
American Indian, male, 25-29 years				
...	All causes	307	100.0	169.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	130	42.3	71.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	54	17.6	29.8
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	29	9.4	16.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	17	5.5	*
5	Chronic liver disease and cirrhosis (K70,K73-K74)	16	5.2	*
6	Malignant neoplasms (C00-C97)	9	2.9	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	1.3	*
7	Legal intervention (Y35,Y89.0)	4	1.3	*
9	Cerebrovascular diseases (I60-I69)	2	0.7	*
9	Influenza and pneumonia (J09-J18)	2	0.7	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.3	*
11	Meningitis (G00,G03)	1	0.3	*
11	Chronic lower respiratory diseases (J40-J47)	1	0.3	*
11	Pneumonitis due to solids and liquids (J69)	1	0.3	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.3	*
...	All other causes (Residual)	34	11.1	18.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 30-34 years				
...	All causes	306	100.0	181.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	113	36.9	67.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	39	12.7	23.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	36	11.8	21.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	28	9.2	16.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	21	6.9	12.5
6	Malignant neoplasms (C00-C97)	9	2.9	*
7	Diabetes mellitus (E10-E14)	6	2.0	*
7	Influenza and pneumonia (J09-J18)	6	2.0	*
9	Septicemia (A40-A41)	4	1.3	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.7	*
10	Chronic lower respiratory diseases (J40-J47)	2	0.7	*
10	Pneumonitis due to solids and liquids (J69)	2	0.7	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.3	*
13	Meningitis (G00,G03)	1	0.3	*
13	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.3	*
...	All other causes (Residual)	35	11.4	20.8
American Indian, male, 35-39 years				
...	All causes	312	100.0	207.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	109	34.9	72.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	39	12.5	26.0
3	Chronic liver disease and cirrhosis (K70,K73-K74)	38	12.2	25.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	29	9.3	19.3
5	Diabetes mellitus (E10-E14)	12	3.8	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	11	3.5	*
7	Malignant neoplasms (C00-C97)	10	3.2	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	1.6	*
9	Septicemia (A40-A41)	3	1.0	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	3	1.0	*
11	Viral hepatitis (B15-B19)	2	0.6	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.3	*
12	Cerebrovascular diseases (I60-I69)	1	0.3	*
12	Influenza and pneumonia (J09-J18)	1	0.3	*
12	Chronic lower respiratory diseases (J40-J47)	1	0.3	*
12	Peptic ulcer (K25-K28)	1	0.3	*
12	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.3	*
12	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.3	*
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.3	*
12	Legal intervention (Y35,Y89.0)	1	0.3	*
...	All other causes (Residual)	42	13.5	28.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 40-44 years				
...	All causes	423	100.0	290.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	108	25.5	74.2
2	Chronic liver disease and cirrhosis (K70,K73-K74)	63	14.9	43.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	59	13.9	40.5
4	Malignant neoplasms (C00-C97)	32	7.6	22.0
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	28	6.6	19.2
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	20	4.7	13.7
7	Diabetes mellitus (E10-E14)	16	3.8	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	6	1.4	*
9	Cerebrovascular diseases (I60-I69)	5	1.2	*
10	Septicemia (A40-A41)	4	0.9	*
10	Influenza and pneumonia (J09-J18)	4	0.9	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.7	*
13	Viral hepatitis (B15-B19)	2	0.5	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2	0.5	*
13	Legal intervention (Y35,Y89.0)	2	0.5	*
...	All other causes (Residual)	69	16.3	47.4
American Indian, male, 45-49 years				
...	All causes	565	100.0	404.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	122	21.6	87.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	86	15.2	61.6
3	Chronic liver disease and cirrhosis (K70,K73-K74)	65	11.5	46.5
4	Malignant neoplasms (C00-C97)	56	9.9	40.1
5	Diabetes mellitus (E10-E14)	38	6.7	27.2
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	37	6.5	26.5
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	19	3.4	*
8	Cerebrovascular diseases (I60-I69)	10	1.8	*
9	Viral hepatitis (B15-B19)	8	1.4	*
9	Influenza and pneumonia (J09-J18)	8	1.4	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	1.2	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	6	1.1	*
13	Septicemia (A40-A41)	4	0.7	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4	0.7	*
15	Pneumonitis due to solids and liquids (J69)	3	0.5	*
15	Peptic ulcer (K25-K28)	3	0.5	*
...	All other causes (Residual)	89	15.8	63.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 50-54 years				
...	All causes	787	100.0	603.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	150	19.1	115.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	126	16.0	96.7
3	Malignant neoplasms (C00-C97)	122	15.5	93.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	94	11.9	72.1
5	Diabetes mellitus (E10-E14)	39	5.0	29.9
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	20	2.5	15.3
7	Cerebrovascular diseases (I60-I69)	19	2.4	*
8	Chronic lower respiratory diseases (J40-J47)	16	2.0	*
9	Viral hepatitis (B15-B19)	14	1.8	*
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	13	1.7	*
11	Septicemia (A40-A41)	12	1.5	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	1.3	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	6	0.8	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	6	0.8	*
13	Influenza and pneumonia (J09-J18)	6	0.8	*
...	All other causes (Residual)	134	17.0	102.8
American Indian, male, 55-59 years				
...	All causes	904	100.0	848.7
1	Malignant neoplasms (C00-C97)	194	21.5	182.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	166	18.4	155.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	95	10.5	89.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	87	9.6	81.7
5	Diabetes mellitus (E10-E14)	60	6.6	56.3
6	Chronic lower respiratory diseases (J40-J47)	37	4.1	34.7
7	Viral hepatitis (B15-B19)	20	2.2	18.8
7	Cerebrovascular diseases (I60-I69)	20	2.2	18.8
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	19	2.1	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	1.9	*
11	Influenza and pneumonia (J09-J18)	16	1.8	*
12	Septicemia (A40-A41)	15	1.7	*
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7	0.8	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	6	0.7	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.4	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.4	*
15	Aortic aneurysm and dissection (I71)	4	0.4	*
15	Pneumonitis due to solids and liquids (J69)	4	0.4	*
...	All other causes (Residual)	129	14.3	121.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 60-64 years				
...	All causes	842	100.0	1,060.5
1	Malignant neoplasms (C00-C97)	208	24.7	262.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	203	24.1	255.7
3	Diabetes mellitus (E10-E14)	73	8.7	91.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	71	8.4	89.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	50	5.9	63.0
6	Chronic lower respiratory diseases (J40-J47)	40	4.8	50.4
7	Cerebrovascular diseases (I60-I69)	30	3.6	37.8
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	14	1.7	*
9	Influenza and pneumonia (J09-J18)	13	1.5	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	1.3	*
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11	1.3	*
12	Septicemia (A40-A41)	9	1.1	*
13	Viral hepatitis (B15-B19)	7	0.8	*
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	0.6	*
15	Pneumonitis due to solids and liquids (J69)	4	0.5	*
...	All other causes (Residual)	93	11.0	117.1
American Indian, male, 65-69 years				
...	All causes	878	100.0	1,617.6
1	Malignant neoplasms (C00-C97)	260	29.6	479.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	196	22.3	361.1
3	Diabetes mellitus (E10-E14)	75	8.5	138.2
4	Chronic lower respiratory diseases (J40-J47)	46	5.2	84.7
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	35	4.0	64.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	31	3.5	57.1
7	Cerebrovascular diseases (I60-I69)	29	3.3	53.4
8	Septicemia (A40-A41)	17	1.9	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	1.6	*
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11	1.3	*
11	Influenza and pneumonia (J09-J18)	10	1.1	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9	1.0	*
13	Pneumonitis due to solids and liquids (J69)	7	0.8	*
14	Viral hepatitis (B15-B19)	5	0.6	*
15	Atherosclerosis (I70)	4	0.5	*
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	0.5	*
...	All other causes (Residual)	125	14.2	230.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 70-74 years				
...	All causes	853	100.0	2,469.7
1	Malignant neoplasms (C00-C97)	236	27.7	683.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	205	24.0	593.5
3	Diabetes mellitus (E10-E14)	62	7.3	179.5
4	Chronic lower respiratory diseases (J40-J47)	53	6.2	153.4
5	Cerebrovascular diseases (I60-I69)	46	5.4	133.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	45	5.3	130.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	20	2.3	57.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19	2.2	*
9	Septicemia (A40-A41)	12	1.4	*
9	Influenza and pneumonia (J09-J18)	12	1.4	*
11	Alzheimer's disease (G30)	7	0.8	*
12	Parkinson's disease (G20-G21)	4	0.5	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4	0.5	*
12	Pneumonitis due to solids and liquids (J69)	4	0.5	*
15	Viral hepatitis (B15-B19)	3	0.4	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.4	*
15	Aortic aneurysm and dissection (I71)	3	0.4	*
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3	0.4	*
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	3	0.4	*
...	All other causes (Residual)	109	12.8	315.6
American Indian, male, 75-79 years				
...	All causes	754	100.0	3,582.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	197	26.1	935.9
2	Malignant neoplasms (C00-C97)	172	22.8	817.1
3	Chronic lower respiratory diseases (J40-J47)	63	8.4	299.3
4	Diabetes mellitus (E10-E14)	53	7.0	251.8
5	Cerebrovascular diseases (I60-I69)	30	4.0	142.5
6	Influenza and pneumonia (J09-J18)	21	2.8	99.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18	2.4	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	2.1	*
9	Septicemia (A40-A41)	13	1.7	*
10	Chronic liver disease and cirrhosis (K70,K73-K74)	12	1.6	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7	0.9	*
11	Pneumonitis due to solids and liquids (J69)	7	0.9	*
13	Alzheimer's disease (G30)	6	0.8	*
14	Parkinson's disease (G20-G21)	5	0.7	*
15	Atherosclerosis (I70)	4	0.5	*
...	All other causes (Residual)	130	17.2	617.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 80-84 years				
...	All causes	633	100.0	5,252.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	180	28.4	1,493.5
2	Malignant neoplasms (C00-C97)	121	19.1	1,004.0
3	Chronic lower respiratory diseases (J40-J47)	53	8.4	439.8
4	Diabetes mellitus (E10-E14)	44	7.0	365.1
5	Cerebrovascular diseases (I60-I69)	27	4.3	224.0
6	Alzheimer's disease (G30)	19	3.0	*
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18	2.8	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	2.7	*
9	Influenza and pneumonia (J09-J18)	14	2.2	*
10	Septicemia (A40-A41)	11	1.7	*
10	Chronic liver disease and cirrhosis (K70,K73-K74)	11	1.7	*
12	Parkinson's disease (G20-G21)	10	1.6	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4	0.6	*
14	Nutritional deficiencies (E40-E64)	3	0.5	*
14	Cholelithiasis and other disorders of gallbladder (K80-K82)	3	0.5	*
...	All other causes (Residual)	98	15.5	813.1
American Indian, male, 85-89 years				
...	All causes	443	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	115	26.0	---
2	Malignant neoplasms (C00-C97)	77	17.4	---
3	Chronic lower respiratory diseases (J40-J47)	36	8.1	---
4	Cerebrovascular diseases (I60-I69)	23	5.2	---
5	Diabetes mellitus (E10-E14)	21	4.7	---
6	Influenza and pneumonia (J09-J18)	14	3.2	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13	2.9	---
8	Parkinson's disease (G20-G21)	10	2.3	---
9	Alzheimer's disease (G30)	9	2.0	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7	1.6	---
10	Pneumonitis due to solids and liquids (J69)	7	1.6	---
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	1.4	---
13	Septicemia (A40-A41)	5	1.1	---
14	Chronic liver disease and cirrhosis (K70,K73-K74)	3	0.7	---
14	Cholelithiasis and other disorders of gallbladder (K80-K82)	3	0.7	---
...	All other causes (Residual)	94	21.2	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 90-94 years				
...	All causes	234	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	59	25.2	---
2	Malignant neoplasms (C00-C97)	31	13.2	---
3	Cerebrovascular diseases (I60-I69)	19	8.1	---
4	Chronic lower respiratory diseases (J40-J47)	17	7.3	---
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	5.1	---
6	Diabetes mellitus (E10-E14)	10	4.3	---
7	Alzheimer's disease (G30)	8	3.4	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7	3.0	---
8	Influenza and pneumonia (J09-J18)	7	3.0	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7	3.0	---
11	Pneumonitis due to solids and liquids (J69)	4	1.7	---
12	Septicemia (A40-A41)	3	1.3	---
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.9	---
13	Parkinson's disease (G20-G21)	2	0.9	---
15	Anemias (D50-D64)	1	0.4	---
15	Peptic ulcer (K25-K28)	1	0.4	---
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.4	---
15	Hyperplasia of prostate (N40)	1	0.4	---
...	All other causes (Residual)	42	17.9	---
American Indian, male, 95-99 years				
...	All causes	59	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	20	33.9	---
2	Malignant neoplasms (C00-C97)	3	5.1	---
2	Diabetes mellitus (E10-E14)	3	5.1	---
2	Influenza and pneumonia (J09-J18)	3	5.1	---
5	Alzheimer's disease (G30)	2	3.4	---
5	Cerebrovascular diseases (I60-I69)	2	3.4	---
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	3.4	---
8	Tuberculosis (A16-A19)	1	1.7	---
8	Septicemia (A40-A41)	1	1.7	---
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.7	---
8	Nutritional deficiencies (E40-E64)	1	1.7	---
8	Parkinson's disease (G20-G21)	1	1.7	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	1.7	---
8	Chronic lower respiratory diseases (J40-J47)	1	1.7	---
8	Hyperplasia of prostate (N40)	1	1.7	---
...	All other causes (Residual)	16	27.1	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 100 years and over				
...	All causes	13	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	30.8	---
2	Influenza and pneumonia (J09-J18)	2	15.4	---
3	Malignant neoplasms (C00-C97)	1	7.7	---
3	Chronic lower respiratory diseases (J40-J47)	1	7.7	---
3	Pneumonitis due to solids and liquids (J69)	1	7.7	---
3	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	7.7	---
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1	7.7	---
...	All other causes (Residual)	2	15.4	---
American Indian, female, all ages				
...	All causes	7,460	100.0	340.9
1	Malignant neoplasms (C00-C97)	1,460	19.6	66.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,219	16.3	55.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	644	8.6	29.4
4	Diabetes mellitus (E10-E14)	471	6.3	21.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	401	5.4	18.3
6	Chronic lower respiratory diseases (J40-J47)	338	4.5	15.4
7	Cerebrovascular diseases (I60-I69)	314	4.2	14.4
8	Alzheimer's disease (G30)	162	2.2	7.4
8	Influenza and pneumonia (J09-J18)	162	2.2	7.4
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	152	2.0	6.9
11	Septicemia (A40-A41)	141	1.9	6.4
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	99	1.3	4.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	76	1.0	3.5
14	Certain conditions originating in the perinatal period (P00-P96)	57	0.8	2.6
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	56	0.8	2.6
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	56	0.8	2.6
...	All other causes (Residual)	1,652	22.1	75.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 1-4 years				
...	All causes	35	100.0	22.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11	31.4	*
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	14.3	*
3	Septicemia (A40-A41)	3	8.6	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	8.6	*
5	Malignant neoplasms (C00-C97)	1	2.9	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	2.9	*
5	Influenza and pneumonia (J09-J18)	1	2.9	*
5	Hernia (K40-K46)	1	2.9	*
...	All other causes (Residual)	9	25.7	*
American Indian, female, 5-9 years				
...	All causes	16	100.0	*
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5	31.3	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	12.5	*
3	Anemias (D50-D64)	1	6.3	*
3	Certain conditions originating in the perinatal period (P00-P96)	1	6.3	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1	6.3	*
...	All other causes (Residual)	6	37.5	*
American Indian, female, 10-14 years				
...	All causes	20	100.0	10.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9	45.0	*
2	Malignant neoplasms (C00-C97)	2	10.0	*
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2	10.0	*
4	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	5.0	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	5.0	*
4	Cerebrovascular diseases (I60-I69)	1	5.0	*
4	Influenza and pneumonia (J09-J18)	1	5.0	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1	5.0	*
...	All other causes (Residual)	2	10.0	*
American Indian, female, 15-19 years				
...	All causes	60	100.0	32.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	34	56.7	18.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	13	21.7	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	8.3	*
4	Malignant neoplasms (C00-C97)	2	3.3	*
5	Septicemia (A40-A41)	1	1.7	*
5	Pregnancy, childbirth and the puerperium (O00-O99)	1	1.7	*
...	All other causes (Residual)	4	6.7	*

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 20-24 years				
...	All causes	117	100.0	63.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	48	41.0	25.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	23	19.7	12.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	10	8.5	*
4	Malignant neoplasms (C00-C97)	4	3.4	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	3.4	*
6	Diabetes mellitus (E10-E14)	2	1.7	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	1.7	*
8	Septicemia (A40-A41)	1	0.9	*
8	Chronic lower respiratory diseases (J40-J47)	1	0.9	*
8	Pneumonitis due to solids and liquids (J69)	1	0.9	*
8	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.9	*
8	Pregnancy, childbirth and the puerperium (O00-O99)	1	0.9	*
8	Legal intervention (Y35,Y89.0)	1	0.9	*
...	All other causes (Residual)	18	15.4	*
American Indian, female, 25-29 years				
...	All causes	162	100.0	97.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	59	36.4	35.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	17	10.5	*
3	Chronic liver disease and cirrhosis (K70,K73-K74)	12	7.4	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	9	5.6	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	8	4.9	*
6	Malignant neoplasms (C00-C97)	6	3.7	*
6	Pregnancy, childbirth and the puerperium (O00-O99)	6	3.7	*
8	Septicemia (A40-A41)	4	2.5	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	2	1.2	*
9	Diabetes mellitus (E10-E14)	2	1.2	*
9	Cerebrovascular diseases (I60-I69)	2	1.2	*
12	Viral hepatitis (B15-B19)	1	0.6	*
12	Anemias (D50-D64)	1	0.6	*
12	Influenza and pneumonia (J09-J18)	1	0.6	*
12	Pneumonitis due to solids and liquids (J69)	1	0.6	*
...	All other causes (Residual)	31	19.1	18.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 30-34 years				
...	All causes	159	100.0	100.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	52	32.7	32.8
2	Chronic liver disease and cirrhosis (K70,K73-K74)	16	10.1	*
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11	6.9	*
4	Malignant neoplasms (C00-C97)	10	6.3	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	9	5.7	*
6	Diabetes mellitus (E10-E14)	5	3.1	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	3.1	*
8	Cerebrovascular diseases (I60-I69)	4	2.5	*
8	Pregnancy, childbirth and the puerperium (O00-O99)	4	2.5	*
10	Septicemia (A40-A41)	3	1.9	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	1.3	*
11	Influenza and pneumonia (J09-J18)	2	1.3	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	1.3	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.6	*
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.6	*
...	All other causes (Residual)	32	20.1	20.2
American Indian, female, 35-39 years				
...	All causes	198	100.0	137.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	49	24.7	34.1
2	Chronic liver disease and cirrhosis (K70,K73-K74)	23	11.6	16.0
3	Malignant neoplasms (C00-C97)	20	10.1	13.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	17	8.6	*
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8	4.0	*
6	Septicemia (A40-A41)	6	3.0	*
6	Diabetes mellitus (E10-E14)	6	3.0	*
8	Pregnancy, childbirth and the puerperium (O00-O99)	5	2.5	*
9	Viral hepatitis (B15-B19)	4	2.0	*
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	2.0	*
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	2.0	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	2	1.0	*
12	Influenza and pneumonia (J09-J18)	2	1.0	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.5	*
14	Cerebrovascular diseases (I60-I69)	1	0.5	*
14	Chronic lower respiratory diseases (J40-J47)	1	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.5	*
14	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.5	*
...	All other causes (Residual)	43	21.7	29.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 40-44 years				
...	All causes	265	100.0	187.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	64	24.2	45.2
2	Chronic liver disease and cirrhosis (K70,K73-K74)	46	17.4	32.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	31	11.7	21.9
4	Malignant neoplasms (C00-C97)	26	9.8	18.4
5	Cerebrovascular diseases (I60-I69)	7	2.6	*
6	Diabetes mellitus (E10-E14)	6	2.3	*
7	Viral hepatitis (B15-B19)	5	1.9	*
8	Chronic lower respiratory diseases (J40-J47)	4	1.5	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	1.5	*
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4	1.5	*
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	1.5	*
12	Septicemia (A40-A41)	3	1.1	*
12	Influenza and pneumonia (J09-J18)	3	1.1	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.8	*
14	Pregnancy, childbirth and the puerperium (O00-O99)	2	0.8	*
...	All other causes (Residual)	54	20.4	38.1
American Indian, female, 45-49 years				
...	All causes	400	100.0	285.5
1	Chronic liver disease and cirrhosis (K70,K73-K74)	66	16.5	47.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	66	16.5	47.1
3	Malignant neoplasms (C00-C97)	59	14.8	42.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	54	13.5	38.5
5	Diabetes mellitus (E10-E14)	20	5.0	14.3
6	Septicemia (A40-A41)	13	3.3	*
6	Cerebrovascular diseases (I60-I69)	13	3.3	*
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8	2.0	*
9	Viral hepatitis (B15-B19)	5	1.3	*
9	Influenza and pneumonia (J09-J18)	5	1.3	*
11	Chronic lower respiratory diseases (J40-J47)	4	1.0	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	1.0	*
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	1.0	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3	0.8	*
15	Pneumonitis due to solids and liquids (J69)	2	0.5	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	0.5	*
...	All other causes (Residual)	72	18.0	51.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 50-54 years				
...	All causes	512	100.0	374.8
1	Malignant neoplasms (C00-C97)	114	22.3	83.5
2	Chronic liver disease and cirrhosis (K70,K73-K74)	77	15.0	56.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	67	13.1	49.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	55	10.7	40.3
5	Diabetes mellitus (E10-E14)	33	6.4	24.2
6	Chronic lower respiratory diseases (J40-J47)	18	3.5	*
7	Influenza and pneumonia (J09-J18)	12	2.3	*
8	Septicemia (A40-A41)	8	1.6	*
9	Viral hepatitis (B15-B19)	7	1.4	*
9	Cerebrovascular diseases (I60-I69)	7	1.4	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	1.0	*
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5	1.0	*
13	Pneumonitis due to solids and liquids (J69)	3	0.6	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
...	All other causes (Residual)	97	18.9	71.0
American Indian, female, 55-59 years				
...	All causes	578	100.0	507.2
1	Malignant neoplasms (C00-C97)	172	29.8	150.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	72	12.5	63.2
3	Diabetes mellitus (E10-E14)	47	8.1	41.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	45	7.8	39.5
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	37	6.4	32.5
6	Chronic lower respiratory diseases (J40-J47)	24	4.2	21.1
7	Cerebrovascular diseases (I60-I69)	19	3.3	*
8	Influenza and pneumonia (J09-J18)	13	2.2	*
9	Viral hepatitis (B15-B19)	12	2.1	*
10	Septicemia (A40-A41)	10	1.7	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7	1.2	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.9	*
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	0.9	*
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4	0.7	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	3	0.5	*
15	Atherosclerosis (I70)	3	0.5	*
...	All other causes (Residual)	100	17.3	87.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 60-64 years				
...	All causes	598	100.0	692.8
1	Malignant neoplasms (C00-C97)	181	30.3	209.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	96	16.1	111.2
3	Diabetes mellitus (E10-E14)	53	8.9	61.4
4	Chronic lower respiratory diseases (J40-J47)	40	6.7	46.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	32	5.4	37.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	27	4.5	31.3
7	Cerebrovascular diseases (I60-I69)	21	3.5	24.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	2.0	*
9	Septicemia (A40-A41)	9	1.5	*
9	Viral hepatitis (B15-B19)	9	1.5	*
11	Influenza and pneumonia (J09-J18)	8	1.3	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	6	1.0	*
13	Alzheimer's disease (G30)	2	0.3	*
13	Atherosclerosis (I70)	2	0.3	*
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2	0.3	*
...	All other causes (Residual)	98	16.4	113.5
American Indian, female, 65-69 years				
...	All causes	694	100.0	1,153.2
1	Malignant neoplasms (C00-C97)	205	29.5	340.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	128	18.4	212.7
3	Diabetes mellitus (E10-E14)	64	9.2	106.3
4	Chronic lower respiratory diseases (J40-J47)	56	8.1	93.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	38	5.5	63.1
6	Cerebrovascular diseases (I60-I69)	24	3.5	39.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	2.2	*
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15	2.2	*
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	11	1.6	*
10	Influenza and pneumonia (J09-J18)	8	1.2	*
11	Septicemia (A40-A41)	7	1.0	*
12	Viral hepatitis (B15-B19)	4	0.6	*
12	Parkinson's disease (G20-G21)	4	0.6	*
12	Alzheimer's disease (G30)	4	0.6	*
12	Pneumonitis due to solids and liquids (J69)	4	0.6	*
...	All other causes (Residual)	107	15.4	177.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 70-74 years				
...	All causes	702	100.0	1,745.0
1	Malignant neoplasms (C00-C97)	191	27.2	474.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	118	16.8	293.3
3	Diabetes mellitus (E10-E14)	60	8.5	149.1
4	Chronic lower respiratory diseases (J40-J47)	53	7.5	131.7
5	Cerebrovascular diseases (I60-I69)	35	5.0	87.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	3.3	57.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	19	2.7	*
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18	2.6	*
9	Influenza and pneumonia (J09-J18)	16	2.3	*
10	Septicemia (A40-A41)	15	2.1	*
11	Alzheimer's disease (G30)	8	1.1	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	6	0.9	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.6	*
13	Nutritional deficiencies (E40-E64)	4	0.6	*
15	Parkinson's disease (G20-G21)	3	0.4	*
15	Aortic aneurysm and dissection (I71)	3	0.4	*
15	Pneumonitis due to solids and liquids (J69)	3	0.4	*
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	3	0.4	*
...	All other causes (Residual)	120	17.1	298.3
American Indian, female, 75-79 years				
...	All causes	720	100.0	2,624.7
1	Malignant neoplasms (C00-C97)	189	26.3	689.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	147	20.4	535.9
3	Diabetes mellitus (E10-E14)	49	6.8	178.6
4	Cerebrovascular diseases (I60-I69)	44	6.1	160.4
5	Chronic lower respiratory diseases (J40-J47)	41	5.7	149.5
6	Influenza and pneumonia (J09-J18)	20	2.8	72.9
7	Alzheimer's disease (G30)	18	2.5	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	2.4	*
9	Septicemia (A40-A41)	16	2.2	*
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14	1.9	*
11	Chronic liver disease and cirrhosis (K70,K73-K74)	13	1.8	*
12	Parkinson's disease (G20-G21)	6	0.8	*
12	Pneumonitis due to solids and liquids (J69)	6	0.8	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.7	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5	0.7	*
...	All other causes (Residual)	130	18.1	473.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 80-84 years				
...	All causes	754	100.0	4,229.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	169	22.4	948.1
2	Malignant neoplasms (C00-C97)	152	20.2	852.7
3	Diabetes mellitus (E10-E14)	56	7.4	314.1
4	Cerebrovascular diseases (I60-I69)	49	6.5	274.9
5	Chronic lower respiratory diseases (J40-J47)	41	5.4	230.0
6	Alzheimer's disease (G30)	28	3.7	157.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	26	3.4	145.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18	2.4	*
9	Septicemia (A40-A41)	17	2.3	*
9	Influenza and pneumonia (J09-J18)	17	2.3	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10	1.3	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.8	*
12	Parkinson's disease (G20-G21)	6	0.8	*
12	Pneumonitis due to solids and liquids (J69)	6	0.8	*
12	Chronic liver disease and cirrhosis (K70,K73-K74)	6	0.8	*
12	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	6	0.8	*
...	All other causes (Residual)	141	18.7	791.0
American Indian, female, 85-89 years				
...	All causes	683	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	153	22.4	---
2	Malignant neoplasms (C00-C97)	83	12.2	---
3	Alzheimer's disease (G30)	48	7.0	---
3	Cerebrovascular diseases (I60-I69)	48	7.0	---
5	Diabetes mellitus (E10-E14)	43	6.3	---
6	Influenza and pneumonia (J09-J18)	28	4.1	---
7	Chronic lower respiratory diseases (J40-J47)	27	4.0	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	23	3.4	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22	3.2	---
10	Septicemia (A40-A41)	12	1.8	---
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	11	1.6	---
12	Parkinson's disease (G20-G21)	6	0.9	---
13	Anemias (D50-D64)	5	0.7	---
13	Pneumonitis due to solids and liquids (J69)	5	0.7	---
13	Chronic liver disease and cirrhosis (K70,K73-K74)	5	0.7	---
...	All other causes (Residual)	164	24.0	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 90-94 years				
...	All causes	413	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	98	23.7	---
2	Alzheimer's disease (G30)	36	8.7	---
3	Malignant neoplasms (C00-C97)	30	7.3	---
4	Cerebrovascular diseases (I60-I69)	23	5.6	---
5	Diabetes mellitus (E10-E14)	22	5.3	---
6	Chronic lower respiratory diseases (J40-J47)	20	4.8	---
7	Influenza and pneumonia (J09-J18)	13	3.1	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	3.1	---
7	Septicemia (A40-A41)	9	2.2	---
10	Pneumonitis due to solids and liquids (J69)	7	1.7	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7	1.7	---
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	6	1.5	---
13	Anemias (D50-D64)	3	0.7	---
14	Parkinson's disease (G20-G21)	2	0.5	---
14	Aortic aneurysm and dissection (I71)	2	0.5	---
14	Peptic ulcer (K25-K28)	2	0.5	---
14	Chronic liver disease and cirrhosis (K70,K73-K74)	2	0.5	---
...	All other causes (Residual)	118	28.6	---
American Indian, female, 95-99 years				
...	All causes	161	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	42	26.1	---
2	Cerebrovascular diseases (I60-I69)	14	8.7	---
3	Alzheimer's disease (G30)	11	6.8	---
4	Malignant neoplasms (C00-C97)	9	5.6	---
5	Essential hypertension and hypertensive renal disease (I10,I12,I15)	8	5.0	---
5	Chronic lower respiratory diseases (J40-J47)	8	5.0	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7	4.3	---
8	Influenza and pneumonia (J09-J18)	6	3.7	---
9	Diabetes mellitus (E10-E14)	3	1.9	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	1.9	---
11	Septicemia (A40-A41)	1	0.6	---
11	Anemias (D50-D64)	1	0.6	---
11	Parkinson's disease (G20-G21)	1	0.6	---
11	Pneumonitis due to solids and liquids (J69)	1	0.6	---
11	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.6	---
...	All other causes (Residual)	45	28.0	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 100 years and over				
...	All causes	51	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	14	27.5	---
2	Alzheimer's disease (G30)	6	11.8	---
3	Influenza and pneumonia (J09-J18)	5	9.8	---
4	Malignant neoplasms (C00-C97)	3	5.9	---
5	Cerebrovascular diseases (I60-I69)	2	3.9	---
6	Parkinson's disease (G20-G21)	1	2.0	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	2.0	---
6	Pneumonitis due to solids and liquids (J69)	1	2.0	---
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1	2.0	---
...	All other causes (Residual)	17	33.3	---
Asian or Pacific Islander, both sexes, all ages				
...	All causes	56,352	100.0	311.5
1	Malignant neoplasms (C00-C97)	15,340	27.2	84.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,266	21.8	67.8
3	Cerebrovascular diseases (I60-I69)	4,108	7.3	22.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,372	4.2	13.1
5	Diabetes mellitus (E10-E14)	2,158	3.8	11.9
6	Influenza and pneumonia (J09-J18)	1,745	3.1	9.6
7	Chronic lower respiratory diseases (J40-J47)	1,624	2.9	9.0
8	Alzheimer's disease (G30)	1,379	2.4	7.6
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,152	2.0	6.4
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,054	1.9	5.8
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	857	1.5	4.7
12	Septicemia (A40-A41)	659	1.2	3.6
13	Chronic liver disease and cirrhosis (K70,K73-K74)	523	0.9	2.9
14	Certain conditions originating in the perinatal period (P00-P96)	476	0.8	2.6
15	Parkinson's disease (G20-G21)	472	0.8	2.6
...	All other causes (Residual)	10,167	18.0	56.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 1-4 years				
...	All causes	150	100.0	15.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	30	20.0	3.1
2	Malignant neoplasms (C00-C97)	21	14.0	2.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	21	14.0	2.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	8	5.3	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	4.0	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	3.3	*
7	Septicemia (A40-A41)	4	2.7	*
7	Influenza and pneumonia (J09-J18)	4	2.7	*
9	Anemias (D50-D64)	2	1.3	*
9	Cerebrovascular diseases (I60-I69)	2	1.3	*
9	Acute bronchitis and bronchiolitis (J20-J21)	2	1.3	*
12	Tuberculosis (A16-A19)	1	0.7	*
12	Meningitis (G00,G03)	1	0.7	*
12	Chronic lower respiratory diseases (J40-J47)	1	0.7	*
12	Certain conditions originating in the perinatal period (P00-P96)	1	0.7	*
12	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.7	*
...	All other causes (Residual)	40	26.7	4.1
Asian or Pacific Islander, both sexes, 5-9 years				
...	All causes	91	100.0	7.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	30	33.0	2.5
2	Malignant neoplasms (C00-C97)	23	25.3	1.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	9.9	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	5.5	*
5	Chronic lower respiratory diseases (J40-J47)	3	3.3	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	2.2	*
7	Septicemia (A40-A41)	1	1.1	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.1	*
7	Anemias (D50-D64)	1	1.1	*
7	Cerebrovascular diseases (I60-I69)	1	1.1	*
7	Influenza and pneumonia (J09-J18)	1	1.1	*
...	All other causes (Residual)	14	15.4	*

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 10-14 years				
...	All causes	101	100.0	8.8
1	Malignant neoplasms (C00-C97)	23	22.8	2.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	23	22.8	2.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	10	9.9	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	7.9	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	5.0	*
6	Influenza and pneumonia (J09-J18)	3	3.0	*
7	Anemias (D50-D64)	2	2.0	*
8	Septicemia (A40-A41)	1	1.0	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.0	*
8	Cerebrovascular diseases (I60-I69)	1	1.0	*
8	Certain conditions originating in the perinatal period (P00-P96)	1	1.0	*
...	All other causes (Residual)	23	22.8	2.0
Asian or Pacific Islander, both sexes, 15-19 years				
...	All causes	259	100.0	22.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	84	32.4	7.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	59	22.8	5.1
3	Malignant neoplasms (C00-C97)	40	15.4	3.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	17	6.6	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	10	3.9	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	2.7	*
7	Cerebrovascular diseases (I60-I69)	3	1.2	*
7	Chronic lower respiratory diseases (J40-J47)	3	1.2	*
9	Anemias (D50-D64)	2	0.8	*
10	Influenza and pneumonia (J09-J18)	1	0.4	*
10	Pregnancy, childbirth and the puerperium (O00-O99)	1	0.4	*
10	Legal intervention (Y35,Y89.0)	1	0.4	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.4	*
...	All other causes (Residual)	30	11.6	2.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 20-24 years				
...	All causes	513	100.0	36.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	158	30.8	11.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	136	26.5	9.7
3	Malignant neoplasms (C00-C97)	51	9.9	3.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	48	9.4	3.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	30	5.8	2.1
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.2	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	1.0	*
8	Septicemia (A40-A41)	3	0.6	*
8	Diabetes mellitus (E10-E14)	3	0.6	*
8	Cerebrovascular diseases (I60-I69)	3	0.6	*
8	Influenza and pneumonia (J09-J18)	3	0.6	*
8	Legal intervention (Y35,Y89.0)	3	0.6	*
13	Tuberculosis (A16-A19)	2	0.4	*
13	Aortic aneurysm and dissection (I71)	2	0.4	*
13	Chronic lower respiratory diseases (J40-J47)	2	0.4	*
...	All other causes (Residual)	58	11.3	4.2
Asian or Pacific Islander, both sexes, 25-29 years				
...	All causes	582	100.0	38.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	160	27.5	10.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	128	22.0	8.5
3	Malignant neoplasms (C00-C97)	68	11.7	4.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	48	8.2	3.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	40	6.9	2.7
6	Cerebrovascular diseases (I60-I69)	9	1.5	*
7	Diabetes mellitus (E10-E14)	7	1.2	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	1.2	*
9	Chronic lower respiratory diseases (J40-J47)	5	0.9	*
10	Septicemia (A40-A41)	4	0.7	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.7	*
10	Anemias (D50-D64)	4	0.7	*
10	Aortic aneurysm and dissection (I71)	4	0.7	*
10	Pregnancy, childbirth and the puerperium (O00-O99)	4	0.7	*
15	Meningococcal infection (A39)	3	0.5	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	3	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.5	*
...	All other causes (Residual)	81	13.9	5.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 30-34 years				
...	All causes	600	100.0	39.3
1	Malignant neoplasms (C00-C97)	128	21.3	8.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	117	19.5	7.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	106	17.7	7.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	63	10.5	4.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	35	5.8	2.3
6	Cerebrovascular diseases (I60-I69)	20	3.3	1.3
7	Pregnancy, childbirth and the puerperium (O00-O99)	12	2.0	*
8	Diabetes mellitus (E10-E14)	6	1.0	*
8	Chronic liver disease and cirrhosis (K70,K73-K74)	6	1.0	*
10	Septicemia (A40-A41)	4	0.7	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.7	*
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	0.7	*
13	Aortic aneurysm and dissection (I71)	3	0.5	*
13	Influenza and pneumonia (J09-J18)	3	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.5	*
...	All other causes (Residual)	86	14.3	5.6
Asian or Pacific Islander, both sexes, 35-39 years				
...	All causes	825	100.0	55.6
1	Malignant neoplasms (C00-C97)	219	26.5	14.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	120	14.5	8.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	118	14.3	7.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	115	13.9	7.7
5	Cerebrovascular diseases (I60-I69)	40	4.8	2.7
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	28	3.4	1.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	14	1.7	*
8	Diabetes mellitus (E10-E14)	12	1.5	*
9	Pregnancy, childbirth and the puerperium (O00-O99)	11	1.3	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	9	1.1	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	0.8	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.7	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.7	*
14	Viral hepatitis (B15-B19)	5	0.6	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5	0.6	*
14	Influenza and pneumonia (J09-J18)	5	0.6	*
...	All other causes (Residual)	105	12.7	7.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 40-44 years				
...	All causes	1,218	100.0	84.9
1	Malignant neoplasms (C00-C97)	406	33.3	28.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	188	15.4	13.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	129	10.6	9.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	107	8.8	7.5
5	Cerebrovascular diseases (I60-I69)	66	5.4	4.6
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	32	2.6	2.2
7	Diabetes mellitus (E10-E14)	31	2.5	2.2
8	Chronic liver disease and cirrhosis (K70,K73-K74)	24	2.0	1.7
9	Influenza and pneumonia (J09-J18)	14	1.1	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	13	1.1	*
11	Viral hepatitis (B15-B19)	11	0.9	*
11	Chronic lower respiratory diseases (J40-J47)	11	0.9	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	11	0.9	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	0.8	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	0.8	*
...	All other causes (Residual)	155	12.7	10.8
Asian or Pacific Islander, both sexes, 45-49 years				
...	All causes	1,660	100.0	131.8
1	Malignant neoplasms (C00-C97)	649	39.1	51.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	279	16.8	22.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	127	7.7	10.1
4	Cerebrovascular diseases (I60-I69)	101	6.1	8.0
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	91	5.5	7.2
6	Diabetes mellitus (E10-E14)	54	3.3	4.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	43	2.6	3.4
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	27	1.6	2.1
9	Viral hepatitis (B15-B19)	19	1.1	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	1.0	*
11	Septicemia (A40-A41)	16	1.0	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	0.8	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	11	0.7	*
14	Aortic aneurysm and dissection (I71)	10	0.6	*
14	Influenza and pneumonia (J09-J18)	10	0.6	*
...	All other causes (Residual)	193	11.6	15.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 50-54 years				
...	All causes	2,445	100.0	213.5
1	Malignant neoplasms (C00-C97)	1,004	41.1	87.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	439	18.0	38.3
3	Cerebrovascular diseases (I60-I69)	165	6.7	14.4
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	145	5.9	12.7
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	96	3.9	8.4
6	Diabetes mellitus (E10-E14)	89	3.6	7.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	47	1.9	4.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	33	1.3	2.9
9	Viral hepatitis (B15-B19)	25	1.0	2.2
10	Influenza and pneumonia (J09-J18)	22	0.9	1.9
10	Chronic lower respiratory diseases (J40-J47)	22	0.9	1.9
12	Septicemia (A40-A41)	20	0.8	1.7
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	18	0.7	*
14	Aortic aneurysm and dissection (I71)	17	0.7	*
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	16	0.7	*
...	All other causes (Residual)	287	11.7	25.1
Asian or Pacific Islander, both sexes, 55-59 years				
...	All causes	3,282	100.0	321.3
1	Malignant neoplasms (C00-C97)	1,389	42.3	136.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	614	18.7	60.1
3	Cerebrovascular diseases (I60-I69)	193	5.9	18.9
4	Diabetes mellitus (E10-E14)	129	3.9	12.6
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	108	3.3	10.6
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	96	2.9	9.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	69	2.1	6.8
8	Viral hepatitis (B15-B19)	43	1.3	4.2
9	Influenza and pneumonia (J09-J18)	40	1.2	3.9
9	Chronic lower respiratory diseases (J40-J47)	40	1.2	3.9
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	34	1.0	3.3
12	Septicemia (A40-A41)	32	1.0	3.1
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	28	0.9	2.7
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	27	0.8	2.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	25	0.8	2.4
15	Aortic aneurysm and dissection (I71)	25	0.8	2.4
...	All other causes (Residual)	390	11.9	38.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 60-64 years				
...	All causes	4,087	100.0	481.7
1	Malignant neoplasms (C00-C97)	1,726	42.2	203.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	808	19.8	95.2
3	Cerebrovascular diseases (I60-I69)	266	6.5	31.3
4	Diabetes mellitus (E10-E14)	191	4.7	22.5
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	142	3.5	16.7
6	Chronic lower respiratory diseases (J40-J47)	74	1.8	8.7
7	Influenza and pneumonia (J09-J18)	59	1.4	7.0
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	59	1.4	7.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	57	1.4	6.7
10	Chronic liver disease and cirrhosis (K70,K73-K74)	54	1.3	6.4
11	Septicemia (A40-A41)	49	1.2	5.8
12	Viral hepatitis (B15-B19)	47	1.1	5.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	45	1.1	5.3
14	Aortic aneurysm and dissection (I71)	27	0.7	3.2
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	23	0.6	2.7
...	All other causes (Residual)	460	11.3	54.2
Asian or Pacific Islander, both sexes, 65-69 years				
...	All causes	4,319	100.0	716.4
1	Malignant neoplasms (C00-C97)	1,695	39.2	281.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	857	19.8	142.2
3	Cerebrovascular diseases (I60-I69)	293	6.8	48.6
4	Diabetes mellitus (E10-E14)	238	5.5	39.5
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	112	2.6	18.6
6	Chronic lower respiratory diseases (J40-J47)	95	2.2	15.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	92	2.1	15.3
8	Influenza and pneumonia (J09-J18)	75	1.7	12.4
9	Chronic liver disease and cirrhosis (K70,K73-K74)	59	1.4	9.8
10	Septicemia (A40-A41)	55	1.3	9.1
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	48	1.1	8.0
12	Viral hepatitis (B15-B19)	42	1.0	7.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	41	0.9	6.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	34	0.8	5.6
15	Aortic aneurysm and dissection (I71)	25	0.6	4.1
...	All other causes (Residual)	558	12.9	92.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 70-74 years				
...	All causes	5,043	100.0	1,160.3
1	Malignant neoplasms (C00-C97)	1,815	36.0	417.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,060	21.0	243.9
3	Cerebrovascular diseases (I60-I69)	393	7.8	90.4
4	Diabetes mellitus (E10-E14)	239	4.7	55.0
5	Chronic lower respiratory diseases (J40-J47)	158	3.1	36.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	131	2.6	30.1
7	Influenza and pneumonia (J09-J18)	125	2.5	28.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	116	2.3	26.7
9	Chronic liver disease and cirrhosis (K70,K73-K74)	68	1.3	15.6
10	Septicemia (A40-A41)	64	1.3	14.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	61	1.2	14.0
12	Alzheimer's disease (G30)	44	0.9	10.1
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	43	0.9	9.9
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	37	0.7	8.5
15	Parkinson's disease (G20-G21)	36	0.7	8.3
...	All other causes (Residual)	653	12.9	150.2
Asian or Pacific Islander, both sexes, 75-79 years				
...	All causes	6,262	100.0	2,045.4
1	Malignant neoplasms (C00-C97)	1,971	31.5	643.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,355	21.6	442.6
3	Cerebrovascular diseases (I60-I69)	511	8.2	166.9
4	Diabetes mellitus (E10-E14)	302	4.8	98.6
5	Chronic lower respiratory diseases (J40-J47)	222	3.5	72.5
6	Influenza and pneumonia (J09-J18)	164	2.6	53.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	156	2.5	51.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	137	2.2	44.8
9	Alzheimer's disease (G30)	102	1.6	33.3
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	89	1.4	29.1
11	Septicemia (A40-A41)	88	1.4	28.7
12	Parkinson's disease (G20-G21)	75	1.2	24.5
13	Chronic liver disease and cirrhosis (K70,K73-K74)	53	0.8	17.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	40	0.6	13.1
15	Pneumonitis due to solids and liquids (J69)	38	0.6	12.4
...	All other causes (Residual)	959	15.3	313.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 80-84 years				
...	All causes	7,570	100.0	3,679.6
1	Malignant neoplasms (C00-C97)	1,900	25.1	923.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,753	23.2	852.1
3	Cerebrovascular diseases (I60-I69)	637	8.4	309.6
4	Diabetes mellitus (E10-E14)	353	4.7	171.6
5	Chronic lower respiratory diseases (J40-J47)	327	4.3	158.9
6	Influenza and pneumonia (J09-J18)	299	3.9	145.3
7	Alzheimer's disease (G30)	243	3.2	118.1
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	203	2.7	98.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	183	2.4	89.0
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	130	1.7	63.2
11	Parkinson's disease (G20-G21)	122	1.6	59.3
12	Septicemia (A40-A41)	92	1.2	44.7
13	Pneumonitis due to solids and liquids (J69)	72	1.0	35.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	52	0.7	25.3
15	Chronic liver disease and cirrhosis (K70,K73-K74)	49	0.6	23.8
...	All other causes (Residual)	1,155	15.3	561.4
Asian or Pacific Islander, both sexes, 85-89 years				
...	All causes	7,918	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,129	26.9	---
2	Malignant neoplasms (C00-C97)	1,336	16.9	---
3	Cerebrovascular diseases (I60-I69)	730	9.2	---
4	Influenza and pneumonia (J09-J18)	376	4.7	---
5	Chronic lower respiratory diseases (J40-J47)	361	4.6	---
6	Alzheimer's disease (G30)	357	4.5	---
7	Diabetes mellitus (E10-E14)	282	3.6	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	196	2.5	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	183	2.3	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	149	1.9	---
11	Parkinson's disease (G20-G21)	132	1.7	---
12	Septicemia (A40-A41)	110	1.4	---
13	Pneumonitis due to solids and liquids (J69)	81	1.0	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	49	0.6	---
15	Aortic aneurysm and dissection (I71)	33	0.4	---
...	All other causes (Residual)	1,414	17.9	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 90-94 years				
...	All causes	5,615	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,643	29.3	---
2	Malignant neoplasms (C00-C97)	679	12.1	---
3	Cerebrovascular diseases (I60-I69)	461	8.2	---
4	Alzheimer's disease (G30)	370	6.6	---
5	Influenza and pneumonia (J09-J18)	324	5.8	---
6	Chronic lower respiratory diseases (J40-J47)	188	3.3	---
7	Diabetes mellitus (E10-E14)	172	3.1	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	134	2.4	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	130	2.3	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	112	2.0	---
11	Pneumonitis due to solids and liquids (J69)	77	1.4	---
12	Septicemia (A40-A41)	64	1.1	---
13	Parkinson's disease (G20-G21)	59	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	33	0.6	---
15	Aortic aneurysm and dissection (I71)	32	0.6	---
...	All other causes (Residual)	1,137	20.2	---
Asian or Pacific Islander, both sexes, 95-99 years				
...	All causes	2,321	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	681	29.3	---
2	Alzheimer's disease (G30)	179	7.7	---
3	Malignant neoplasms (C00-C97)	171	7.4	---
4	Cerebrovascular diseases (I60-I69)	169	7.3	---
5	Influenza and pneumonia (J09-J18)	158	6.8	---
6	Chronic lower respiratory diseases (J40-J47)	89	3.8	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	82	3.5	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	59	2.5	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	46	2.0	---
10	Diabetes mellitus (E10-E14)	41	1.8	---
11	Pneumonitis due to solids and liquids (J69)	39	1.7	---
12	Septicemia (A40-A41)	29	1.2	---
13	Parkinson's disease (G20-G21)	19	0.8	---
14	Atherosclerosis (I70)	16	0.7	---
15	Anemias (D50-D64)	10	0.4	---
...	All other causes (Residual)	533	23.0	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 100 years and over				
...	All causes	578	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	166	28.7	---
2	Alzheimer's disease (G30)	53	9.2	---
3	Influenza and pneumonia (J09-J18)	49	8.5	---
4	Cerebrovascular diseases (I60-I69)	39	6.7	---
5	Essential hypertension and hypertensive renal disease (I10,I12,I15)	25	4.3	---
6	Malignant neoplasms (C00-C97)	22	3.8	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19	3.3	---
8	Chronic lower respiratory diseases (J40-J47)	11	1.9	---
9	Diabetes mellitus (E10-E14)	9	1.6	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8	1.4	---
11	Pneumonitis due to solids and liquids (J69)	6	1.0	---
12	Septicemia (A40-A41)	3	0.5	---
12	Atherosclerosis (I70)	3	0.5	---
12	Aortic aneurysm and dissection (I71)	3	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.3	---
...	All other causes (Residual)	160	27.7	---
Asian or Pacific Islander, male, all ages				
...	All causes	28,819	100.0	332.8
1	Malignant neoplasms (C00-C97)	7,709	26.7	89.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,650	23.1	76.8
3	Cerebrovascular diseases (I60-I69)	1,876	6.5	21.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,472	5.1	17.0
5	Diabetes mellitus (E10-E14)	1,093	3.8	12.6
6	Chronic lower respiratory diseases (J40-J47)	966	3.4	11.2
7	Influenza and pneumonia (J09-J18)	890	3.1	10.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	805	2.8	9.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	501	1.7	5.8
10	Alzheimer's disease (G30)	439	1.5	5.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	377	1.3	4.4
12	Chronic liver disease and cirrhosis (K70,K73-K74)	326	1.1	3.8
13	Septicemia (A40-A41)	312	1.1	3.6
14	Certain conditions originating in the perinatal period (P00-P96)	262	0.9	3.0
15	Parkinson's disease (G20-G21)	252	0.9	2.9
...	All other causes (Residual)	4,889	17.0	56.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 1-4 years				
...	All causes	80	100.0	16.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18	22.5	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	13	16.3	*
3	Malignant neoplasms (C00-C97)	10	12.5	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	6.3	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	3.8	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	2.5	*
7	Tuberculosis (A16-A19)	1	1.3	*
7	Septicemia (A40-A41)	1	1.3	*
7	Anemias (D50-D64)	1	1.3	*
7	Cerebrovascular diseases (I60-I69)	1	1.3	*
7	Influenza and pneumonia (J09-J18)	1	1.3	*
7	Acute bronchitis and bronchiolitis (J20-J21)	1	1.3	*
...	All other causes (Residual)	23	28.8	4.7
Asian or Pacific Islander, male, 5-9 years				
...	All causes	48	100.0	7.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19	39.6	*
2	Malignant neoplasms (C00-C97)	13	27.1	*
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	8.3	*
4	Chronic lower respiratory diseases (J40-J47)	2	4.2	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	4.2	*
6	Anemias (D50-D64)	1	2.1	*
6	Cerebrovascular diseases (I60-I69)	1	2.1	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1	2.1	*
...	All other causes (Residual)	5	10.4	*
Asian or Pacific Islander, male, 10-14 years				
...	All causes	60	100.0	10.3
1	Malignant neoplasms (C00-C97)	14	23.3	*
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12	20.0	*
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7	11.7	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	10.0	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	5.0	*
6	Anemias (D50-D64)	2	3.3	*
6	Influenza and pneumonia (J09-J18)	2	3.3	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.7	*
...	All other causes (Residual)	13	21.7	*

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 15-19 years				
...	All causes	186	100.0	31.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	68	36.6	11.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	38	20.4	6.4
3	Malignant neoplasms (C00-C97)	30	16.1	5.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	14	7.5	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	8	4.3	*
6	Cerebrovascular diseases (I60-I69)	3	1.6	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	1.6	*
8	Anemias (D50-D64)	1	0.5	*
8	Chronic lower respiratory diseases (J40-J47)	1	0.5	*
8	Legal intervention (Y35,Y89.0)	1	0.5	*
8	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.5	*
...	All other causes (Residual)	18	9.7	*
Asian or Pacific Islander, male, 20-24 years				
...	All causes	354	100.0	50.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	122	34.5	17.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	88	24.9	12.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	36	10.2	5.1
4	Malignant neoplasms (C00-C97)	32	9.0	4.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	23	6.5	3.3
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.8	*
6	Cerebrovascular diseases (I60-I69)	3	0.8	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.8	*
6	Legal intervention (Y35,Y89.0)	3	0.8	*
10	Septicemia (A40-A41)	2	0.6	*
10	Chronic lower respiratory diseases (J40-J47)	2	0.6	*
12	Diabetes mellitus (E10-E14)	1	0.3	*
12	Aortic aneurysm and dissection (I71)	1	0.3	*
12	Influenza and pneumonia (J09-J18)	1	0.3	*
...	All other causes (Residual)	34	9.6	4.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 25-29 years				
...	All causes	412	100.0	56.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	125	30.3	17.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	99	24.0	13.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	42	10.2	5.7
4	Malignant neoplasms (C00-C97)	41	10.0	5.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	25	6.1	3.4
6	Cerebrovascular diseases (I60-I69)	6	1.5	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	1.2	*
8	Chronic lower respiratory diseases (J40-J47)	4	1.0	*
9	Septicemia (A40-A41)	3	0.7	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.7	*
9	Diabetes mellitus (E10-E14)	3	0.7	*
9	Aortic aneurysm and dissection (I71)	3	0.7	*
13	Meningococcal infection (A39)	2	0.5	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.5	*
13	Anemias (D50-D64)	2	0.5	*
13	Chronic liver disease and cirrhosis (K70,K73-K74)	2	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.5	*
13	Legal intervention (Y35,Y89.0)	2	0.5	*
...	All other causes (Residual)	41	10.0	5.6
Asian or Pacific Islander, male, 30-34 years				
...	All causes	384	100.0	53.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	90	23.4	12.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	75	19.5	10.4
3	Malignant neoplasms (C00-C97)	66	17.2	9.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	52	13.5	7.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	24	6.3	3.3
6	Cerebrovascular diseases (I60-I69)	10	2.6	*
7	Human immunodeficiency virus (HIV) disease (B20-B24)	4	1.0	*
7	Chronic liver disease and cirrhosis (K70,K73-K74)	4	1.0	*
9	Aortic aneurysm and dissection (I71)	3	0.8	*
9	Influenza and pneumonia (J09-J18)	3	0.8	*
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.8	*
12	Septicemia (A40-A41)	2	0.5	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.5	*
14	Viral hepatitis (B15-B19)	1	0.3	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.3	*
14	Pneumonitis due to solids and liquids (J69)	1	0.3	*
14	Diseases of appendix (K35-K38)	1	0.3	*
14	Legal intervention (Y35,Y89.0)	1	0.3	*
...	All other causes (Residual)	41	10.7	5.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 35-39 years				
...	All causes	514	100.0	73.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	101	19.6	14.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	94	18.3	13.4
3	Malignant neoplasms (C00-C97)	88	17.1	12.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	82	16.0	11.7
5	Cerebrovascular diseases (I60-I69)	21	4.1	3.0
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	18	3.5	*
7	Chronic liver disease and cirrhosis (K70,K73-K74)	13	2.5	*
8	Diabetes mellitus (E10-E14)	7	1.4	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	6	1.2	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	1.0	*
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	1.0	*
12	Viral hepatitis (B15-B19)	4	0.8	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4	0.8	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.8	*
15	Septicemia (A40-A41)	3	0.6	*
15	Influenza and pneumonia (J09-J18)	3	0.6	*
15	Legal intervention (Y35,Y89.0)	3	0.6	*
...	All other causes (Residual)	53	10.3	7.6
Asian or Pacific Islander, male, 40-44 years				
...	All causes	752	100.0	111.4
1	Malignant neoplasms (C00-C97)	179	23.8	26.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	151	20.1	22.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	91	12.1	13.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	77	10.2	11.4
5	Cerebrovascular diseases (I60-I69)	34	4.5	5.0
6	Diabetes mellitus (E10-E14)	29	3.9	4.3
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	21	2.8	3.1
8	Chronic liver disease and cirrhosis (K70,K73-K74)	20	2.7	3.0
9	Human immunodeficiency virus (HIV) disease (B20-B24)	11	1.5	*
10	Viral hepatitis (B15-B19)	10	1.3	*
10	Influenza and pneumonia (J09-J18)	10	1.3	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	1.2	*
13	Aortic aneurysm and dissection (I71)	6	0.8	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	0.8	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.7	*
...	All other causes (Residual)	93	12.4	13.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 45-49 years				
...	All causes	1,008	100.0	169.8
1	Malignant neoplasms (C00-C97)	291	28.9	49.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	234	23.2	39.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	91	9.0	15.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	65	6.4	10.9
5	Cerebrovascular diseases (I60-I69)	57	5.7	9.6
6	Diabetes mellitus (E10-E14)	36	3.6	6.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	31	3.1	5.2
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	19	1.9	*
9	Viral hepatitis (B15-B19)	14	1.4	*
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9	0.9	*
11	Aortic aneurysm and dissection (I71)	8	0.8	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	7	0.7	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.7	*
12	Chronic lower respiratory diseases (J40-J47)	7	0.7	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.7	*
...	All other causes (Residual)	125	12.4	21.1
Asian or Pacific Islander, male, 50-54 years				
...	All causes	1,435	100.0	268.5
1	Malignant neoplasms (C00-C97)	453	31.6	84.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	350	24.4	65.5
3	Cerebrovascular diseases (I60-I69)	92	6.4	17.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	91	6.3	17.0
5	Diabetes mellitus (E10-E14)	63	4.4	11.8
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	62	4.3	11.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	33	2.3	6.2
8	Viral hepatitis (B15-B19)	16	1.1	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	1.1	*
10	Septicemia (A40-A41)	14	1.0	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	13	0.9	*
11	Aortic aneurysm and dissection (I71)	13	0.9	*
11	Chronic lower respiratory diseases (J40-J47)	13	0.9	*
14	Influenza and pneumonia (J09-J18)	12	0.8	*
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	10	0.7	*
...	All other causes (Residual)	184	12.8	34.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 55-59 years				
...	All causes	1,931	100.0	414.0
1	Malignant neoplasms (C00-C97)	688	35.6	147.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	444	23.0	95.2
3	Cerebrovascular diseases (I60-I69)	108	5.6	23.2
4	Diabetes mellitus (E10-E14)	78	4.0	16.7
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	74	3.8	15.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	66	3.4	14.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	53	2.7	11.4
8	Viral hepatitis (B15-B19)	26	1.3	5.6
9	Influenza and pneumonia (J09-J18)	23	1.2	4.9
10	Chronic lower respiratory diseases (J40-J47)	22	1.1	4.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	20	1.0	4.3
11	Aortic aneurysm and dissection (I71)	20	1.0	4.3
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20	1.0	4.3
14	Septicemia (A40-A41)	19	1.0	*
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	19	1.0	*
...	All other causes (Residual)	251	13.0	53.8
Asian or Pacific Islander, male, 60-64 years				
...	All causes	2,353	100.0	618.6
1	Malignant neoplasms (C00-C97)	893	38.0	234.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	559	23.8	146.9
3	Cerebrovascular diseases (I60-I69)	146	6.2	38.4
4	Diabetes mellitus (E10-E14)	119	5.1	31.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	83	3.5	21.8
6	Chronic lower respiratory diseases (J40-J47)	45	1.9	11.8
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	41	1.7	10.8
8	Chronic liver disease and cirrhosis (K70,K73-K74)	37	1.6	9.7
9	Influenza and pneumonia (J09-J18)	34	1.4	8.9
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	30	1.3	7.9
11	Viral hepatitis (B15-B19)	29	1.2	7.6
12	Septicemia (A40-A41)	27	1.1	7.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	27	1.1	7.1
14	Aortic aneurysm and dissection (I71)	16	0.7	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.4	*
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	9	0.4	*
...	All other causes (Residual)	249	10.6	65.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 65-69 years				
...	All causes	2,414	100.0	886.9
1	Malignant neoplasms (C00-C97)	894	37.0	328.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	546	22.6	200.6
3	Cerebrovascular diseases (I60-I69)	158	6.5	58.1
4	Diabetes mellitus (E10-E14)	123	5.1	45.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	62	2.6	22.8
6	Chronic lower respiratory diseases (J40-J47)	54	2.2	19.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	47	1.9	17.3
8	Influenza and pneumonia (J09-J18)	44	1.8	16.2
9	Chronic liver disease and cirrhosis (K70,K73-K74)	40	1.7	14.7
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	33	1.4	12.1
11	Septicemia (A40-A41)	28	1.2	10.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	23	1.0	8.5
13	Viral hepatitis (B15-B19)	21	0.9	7.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	0.7	*
14	Aortic aneurysm and dissection (I71)	16	0.7	*
...	All other causes (Residual)	309	12.8	113.5
Asian or Pacific Islander, male, 70-74 years				
...	All causes	2,786	100.0	1,416.8
1	Malignant neoplasms (C00-C97)	965	34.6	490.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	650	23.3	330.6
3	Cerebrovascular diseases (I60-I69)	208	7.5	105.8
4	Diabetes mellitus (E10-E14)	119	4.3	60.5
5	Chronic lower respiratory diseases (J40-J47)	102	3.7	51.9
6	Influenza and pneumonia (J09-J18)	79	2.8	40.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	79	2.8	40.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	63	2.3	32.0
9	Chronic liver disease and cirrhosis (K70,K73-K74)	37	1.3	18.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	35	1.3	17.8
11	Septicemia (A40-A41)	31	1.1	15.8
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	22	0.8	11.2
13	Alzheimer's disease (G30)	21	0.8	10.7
14	Aortic aneurysm and dissection (I71)	20	0.7	10.2
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	17	0.6	*
...	All other causes (Residual)	338	12.1	171.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 75-79 years				
...	All causes	3,310	100.0	2,498.4
1	Malignant neoplasms (C00-C97)	1,068	32.3	806.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	763	23.1	575.9
3	Cerebrovascular diseases (I60-I69)	236	7.1	178.1
4	Diabetes mellitus (E10-E14)	160	4.8	120.8
5	Chronic lower respiratory diseases (J40-J47)	141	4.3	106.4
6	Influenza and pneumonia (J09-J18)	99	3.0	74.7
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	76	2.3	57.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	63	1.9	47.6
9	Parkinson's disease (G20-G21)	44	1.3	33.2
10	Septicemia (A40-A41)	41	1.2	30.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	36	1.1	27.2
12	Alzheimer's disease (G30)	32	1.0	24.2
13	Pneumonitis due to solids and liquids (J69)	25	0.8	18.9
14	Chronic liver disease and cirrhosis (K70,K73-K74)	24	0.7	18.1
15	Aortic aneurysm and dissection (I71)	21	0.6	15.9
...	All other causes (Residual)	481	14.5	363.1
Asian or Pacific Islander, male, 80-84 years				
...	All causes	3,588	100.0	4,382.4
1	Malignant neoplasms (C00-C97)	939	26.2	1,146.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	842	23.5	1,028.4
3	Cerebrovascular diseases (I60-I69)	277	7.7	338.3
4	Chronic lower respiratory diseases (J40-J47)	204	5.7	249.2
5	Diabetes mellitus (E10-E14)	164	4.6	200.3
6	Influenza and pneumonia (J09-J18)	159	4.4	194.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	93	2.6	113.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	87	2.4	106.3
9	Alzheimer's disease (G30)	80	2.2	97.7
10	Parkinson's disease (G20-G21)	74	2.1	90.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	51	1.4	62.3
12	Septicemia (A40-A41)	48	1.3	58.6
13	Pneumonitis due to solids and liquids (J69)	40	1.1	48.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	28	0.8	34.2
15	Aortic aneurysm and dissection (I71)	26	0.7	31.8
...	All other causes (Residual)	476	13.3	581.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 85-89 years				
...	All causes	3,462	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	944	27.3	---
2	Malignant neoplasms (C00-C97)	619	17.9	---
3	Cerebrovascular diseases (I60-I69)	281	8.1	---
4	Chronic lower respiratory diseases (J40-J47)	212	6.1	---
5	Influenza and pneumonia (J09-J18)	182	5.3	---
6	Alzheimer's disease (G30)	114	3.3	---
7	Diabetes mellitus (E10-E14)	108	3.1	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	101	2.9	---
9	Parkinson's disease (G20-G21)	72	2.1	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	68	2.0	---
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	67	1.9	---
12	Septicemia (A40-A41)	49	1.4	---
13	Pneumonitis due to solids and liquids (J69)	45	1.3	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	25	0.7	---
15	Aortic aneurysm and dissection (I71)	14	0.4	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	14	0.4	---
...	All other causes (Residual)	547	15.8	---
Asian or Pacific Islander, male, 90-94 years				
...	All causes	2,269	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	666	29.4	---
2	Malignant neoplasms (C00-C97)	345	15.2	---
3	Cerebrovascular diseases (I60-I69)	159	7.0	---
4	Influenza and pneumonia (J09-J18)	146	6.4	---
5	Alzheimer's disease (G30)	122	5.4	---
6	Chronic lower respiratory diseases (J40-J47)	103	4.5	---
7	Diabetes mellitus (E10-E14)	70	3.1	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	54	2.4	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	51	2.2	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	48	2.1	---
11	Pneumonitis due to solids and liquids (J69)	36	1.6	---
12	Parkinson's disease (G20-G21)	25	1.1	---
13	Septicemia (A40-A41)	20	0.9	---
14	Aortic aneurysm and dissection (I71)	12	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	0.4	---
15	Atherosclerosis (I70)	10	0.4	---
...	All other causes (Residual)	392	17.3	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 95-99 years				
...	All causes	792	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	220	27.8	---
2	Malignant neoplasms (C00-C97)	70	8.8	---
3	Cerebrovascular diseases (I60-I69)	63	8.0	---
3	Influenza and pneumonia (J09-J18)	63	8.0	---
5	Chronic lower respiratory diseases (J40-J47)	43	5.4	---
6	Alzheimer's disease (G30)	40	5.1	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	27	3.4	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	24	3.0	---
9	Pneumonitis due to solids and liquids (J69)	22	2.8	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17	2.1	---
11	Diabetes mellitus (E10-E14)	13	1.6	---
12	Septicemia (A40-A41)	10	1.3	---
13	Parkinson's disease (G20-G21)	9	1.1	---
14	Atherosclerosis (I70)	6	0.8	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.5	---
...	All other causes (Residual)	161	20.3	---
Asian or Pacific Islander, male, 100 years and over				
...	All causes	165	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	49	29.7	---
2	Influenza and pneumonia (J09-J18)	16	9.7	---
3	Alzheimer's disease (G30)	11	6.7	---
4	Cerebrovascular diseases (I60-I69)	10	6.1	---
5	Malignant neoplasms (C00-C97)	9	5.5	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	8	4.8	---
7	Chronic lower respiratory diseases (J40-J47)	6	3.6	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	3.6	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4	2.4	---
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	1.2	---
10	Pneumonitis due to solids and liquids (J69)	2	1.2	---
12	Tuberculosis (A16-A19)	1	0.6	---
12	Septicemia (A40-A41)	1	0.6	---
12	Aortic aneurysm and dissection (I71)	1	0.6	---
12	Hyperplasia of prostate (N40)	1	0.6	---
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1	0.6	---
...	All other causes (Residual)	37	22.4	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, all ages				
...	All causes	27,533	100.0	292.0
1	Malignant neoplasms (C00-C97)	7,631	27.7	80.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,616	20.4	59.6
3	Cerebrovascular diseases (I60-I69)	2,232	8.1	23.7
4	Diabetes mellitus (E10-E14)	1,065	3.9	11.3
5	Alzheimer's disease (G30)	940	3.4	10.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	900	3.3	9.5
7	Influenza and pneumonia (J09-J18)	855	3.1	9.1
8	Chronic lower respiratory diseases (J40-J47)	658	2.4	7.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	553	2.0	5.9
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	480	1.7	5.1
11	Septicemia (A40-A41)	347	1.3	3.7
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	347	1.3	3.7
13	Parkinson's disease (G20-G21)	220	0.8	2.3
14	Certain conditions originating in the perinatal period (P00-P96)	214	0.8	2.3
15	Chronic liver disease and cirrhosis (K70,K73-K74)	197	0.7	2.1
...	All other causes (Residual)	5,278	19.2	56.0
Asian or Pacific Islander, female, 1-4 years				
...	All causes	70	100.0	14.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12	17.1	*
2	Malignant neoplasms (C00-C97)	11	15.7	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	11.4	*
4	Septicemia (A40-A41)	3	4.3	*
4	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	4.3	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	4.3	*
4	Influenza and pneumonia (J09-J18)	3	4.3	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	4.3	*
9	Anemias (D50-D64)	1	1.4	*
9	Meningitis (G00,G03)	1	1.4	*
9	Cerebrovascular diseases (I60-I69)	1	1.4	*
9	Acute bronchitis and bronchiolitis (J20-J21)	1	1.4	*
9	Chronic lower respiratory diseases (J40-J47)	1	1.4	*
9	Certain conditions originating in the perinatal period (P00-P96)	1	1.4	*
9	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	1.4	*
...	All other causes (Residual)	17	24.3	*

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 5-9 years				
...	All causes	43	100.0	7.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11	25.6	*
2	Malignant neoplasms (C00-C97)	10	23.3	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	16.3	*
4	Septicemia (A40-A41)	1	2.3	*
4	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	2.3	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	2.3	*
4	Influenza and pneumonia (J09-J18)	1	2.3	*
4	Chronic lower respiratory diseases (J40-J47)	1	2.3	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1	2.3	*
...	All other causes (Residual)	9	20.9	*
Asian or Pacific Islander, female, 10-14 years				
...	All causes	41	100.0	7.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11	26.8	*
2	Malignant neoplasms (C00-C97)	9	22.0	*
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3	7.3	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	4.9	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	4.9	*
6	Septicemia (A40-A41)	1	2.4	*
6	Cerebrovascular diseases (I60-I69)	1	2.4	*
6	Influenza and pneumonia (J09-J18)	1	2.4	*
6	Certain conditions originating in the perinatal period (P00-P96)	1	2.4	*
...	All other causes (Residual)	10	24.4	*
Asian or Pacific Islander, female, 15-19 years				
...	All causes	73	100.0	12.7
1	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	21	28.8	3.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	16	21.9	*
3	Malignant neoplasms (C00-C97)	10	13.7	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	5.5	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	4.1	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	2.7	*
6	Chronic lower respiratory diseases (J40-J47)	2	2.7	*
8	Anemias (D50-D64)	1	1.4	*
8	Influenza and pneumonia (J09-J18)	1	1.4	*
8	Pregnancy, childbirth and the puerperium (O00-O99)	1	1.4	*
...	All other causes (Residual)	12	16.4	*

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 20-24 years				
...	All causes	159	100.0	23.1
1	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	48	30.2	7.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	36	22.6	5.2
3	Malignant neoplasms (C00-C97)	19	11.9	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	12	7.5	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	7	4.4	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	1.9	*
7	Tuberculosis (A16-A19)	2	1.3	*
7	Diabetes mellitus (E10-E14)	2	1.3	*
7	Influenza and pneumonia (J09-J18)	2	1.3	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	1.3	*
11	Septicemia (A40-A41)	1	0.6	*
11	Aortic aneurysm and dissection (I71)	1	0.6	*
11	Pneumonitis due to solids and liquids (J69)	1	0.6	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.6	*
...	All other causes (Residual)	22	13.8	3.2
Asian or Pacific Islander, female, 25-29 years				
...	All causes	170	100.0	21.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	35	20.6	4.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	29	17.1	3.7
3	Malignant neoplasms (C00-C97)	27	15.9	3.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	15	8.8	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	3.5	*
6	Diabetes mellitus (E10-E14)	4	2.4	*
6	Pregnancy, childbirth and the puerperium (O00-O99)	4	2.4	*
8	Cerebrovascular diseases (I60-I69)	3	1.8	*
9	Anemias (D50-D64)	2	1.2	*
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	1.2	*
11	Meningococcal infection (A39)	1	0.6	*
11	Septicemia (A40-A41)	1	0.6	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.6	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.6	*
11	Aortic aneurysm and dissection (I71)	1	0.6	*
11	Influenza and pneumonia (J09-J18)	1	0.6	*
11	Chronic lower respiratory diseases (J40-J47)	1	0.6	*
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.6	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.6	*
11	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.6	*
...	All other causes (Residual)	33	19.4	4.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 30-34 years				
...	All causes	216	100.0	26.8
1	Malignant neoplasms (C00-C97)	62	28.7	7.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	31	14.4	3.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	27	12.5	3.3
4	Pregnancy, childbirth and the puerperium (O00-O99)	12	5.6	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	11	5.1	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	11	5.1	*
7	Cerebrovascular diseases (I60-I69)	10	4.6	*
8	Diabetes mellitus (E10-E14)	6	2.8	*
9	Tuberculosis (A16-A19)	2	0.9	*
9	Septicemia (A40-A41)	2	0.9	*
9	Chronic liver disease and cirrhosis (K70,K73-K74)	2	0.9	*
12	Meningococcal infection (A39)	1	0.5	*
12	Viral hepatitis (B15-B19)	1	0.5	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.5	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.5	*
12	Atherosclerosis (I70)	1	0.5	*
12	Pneumonitis due to solids and liquids (J69)	1	0.5	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.5	*
12	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.5	*
12	Inflammatory diseases of female pelvic organs (N70-N76)	1	0.5	*
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.5	*
...	All other causes (Residual)	30	13.9	3.7
Asian or Pacific Islander, female, 35-39 years				
...	All causes	311	100.0	39.7
1	Malignant neoplasms (C00-C97)	131	42.1	16.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	33	10.6	4.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	26	8.4	3.3
4	Cerebrovascular diseases (I60-I69)	19	6.1	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	17	5.5	*
6	Pregnancy, childbirth and the puerperium (O00-O99)	11	3.5	*
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	10	3.2	*
8	Diabetes mellitus (E10-E14)	5	1.6	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	3	1.0	*
10	Aortic aneurysm and dissection (I71)	2	0.6	*
10	Influenza and pneumonia (J09-J18)	2	0.6	*
10	Chronic lower respiratory diseases (J40-J47)	2	0.6	*
10	Pneumonitis due to solids and liquids (J69)	2	0.6	*
10	Cholelithiasis and other disorders of gallbladder (K80-K82)	2	0.6	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.6	*
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	0.6	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.6	*
...	All other causes (Residual)	40	12.9	5.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 40-44 years				
...	All causes	466	100.0	61.4
1	Malignant neoplasms (C00-C97)	227	48.7	29.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	38	8.2	5.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	37	7.9	4.9
4	Cerebrovascular diseases (I60-I69)	32	6.9	4.2
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	30	6.4	4.0
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	11	2.4	*
7	Chronic lower respiratory diseases (J40-J47)	8	1.7	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	1.1	*
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	1.1	*
10	Septicemia (A40-A41)	4	0.9	*
10	Influenza and pneumonia (J09-J18)	4	0.9	*
10	Chronic liver disease and cirrhosis (K70,K73-K74)	4	0.9	*
10	Pregnancy, childbirth and the puerperium (O00-O99)	4	0.9	*
14	Aortic aneurysm and dissection (I71)	3	0.6	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.4	*
15	Diabetes mellitus (E10-E14)	2	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
...	All other causes (Residual)	48	10.3	6.3
Asian or Pacific Islander, female, 45-49 years				
...	All causes	652	100.0	98.0
1	Malignant neoplasms (C00-C97)	358	54.9	53.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	45	6.9	6.8
3	Cerebrovascular diseases (I60-I69)	44	6.7	6.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	36	5.5	5.4
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	26	4.0	3.9
6	Diabetes mellitus (E10-E14)	18	2.8	*
7	Chronic liver disease and cirrhosis (K70,K73-K74)	12	1.8	*
8	Septicemia (A40-A41)	10	1.5	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	1.5	*
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8	1.2	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.9	*
12	Viral hepatitis (B15-B19)	5	0.8	*
12	Influenza and pneumonia (J09-J18)	5	0.8	*
14	Anemias (D50-D64)	2	0.3	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2	0.3	*
14	Aortic aneurysm and dissection (I71)	2	0.3	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	0.3	*
...	All other causes (Residual)	61	9.4	9.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 50-54 years				
...	All causes	1,010	100.0	165.5
1	Malignant neoplasms (C00-C97)	551	54.6	90.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	89	8.8	14.6
3	Cerebrovascular diseases (I60-I69)	73	7.2	12.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	54	5.3	8.8
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	34	3.4	5.6
6	Diabetes mellitus (E10-E14)	26	2.6	4.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	1.7	*
8	Chronic liver disease and cirrhosis (K70,K73-K74)	14	1.4	*
9	Influenza and pneumonia (J09-J18)	10	1.0	*
10	Viral hepatitis (B15-B19)	9	0.9	*
10	Chronic lower respiratory diseases (J40-J47)	9	0.9	*
12	Septicemia (A40-A41)	6	0.6	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.6	*
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	0.6	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5	0.5	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	0.5	*
...	All other causes (Residual)	96	9.5	15.7
Asian or Pacific Islander, female, 55-59 years				
...	All causes	1,351	100.0	243.3
1	Malignant neoplasms (C00-C97)	701	51.9	126.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	170	12.6	30.6
3	Cerebrovascular diseases (I60-I69)	85	6.3	15.3
4	Diabetes mellitus (E10-E14)	51	3.8	9.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	42	3.1	7.6
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	22	1.6	4.0
7	Chronic lower respiratory diseases (J40-J47)	18	1.3	*
8	Viral hepatitis (B15-B19)	17	1.3	*
8	Influenza and pneumonia (J09-J18)	17	1.3	*
10	Chronic liver disease and cirrhosis (K70,K73-K74)	16	1.2	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	1.0	*
12	Septicemia (A40-A41)	13	1.0	*
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	9	0.7	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	0.6	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.5	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7	0.5	*
...	All other causes (Residual)	154	11.4	27.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 60-64 years				
...	All causes	1,734	100.0	370.4
1	Malignant neoplasms (C00-C97)	833	48.0	178.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	249	14.4	53.2
3	Cerebrovascular diseases (I60-I69)	120	6.9	25.6
4	Diabetes mellitus (E10-E14)	72	4.2	15.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	59	3.4	12.6
6	Chronic lower respiratory diseases (J40-J47)	29	1.7	6.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	27	1.6	5.8
8	Influenza and pneumonia (J09-J18)	25	1.4	5.3
9	Septicemia (A40-A41)	22	1.3	4.7
10	Viral hepatitis (B15-B19)	18	1.0	*
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	18	1.0	*
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	18	1.0	*
13	Chronic liver disease and cirrhosis (K70,K73-K74)	17	1.0	*
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	14	0.8	*
15	Aortic aneurysm and dissection (I71)	11	0.6	*
...	All other causes (Residual)	202	11.6	43.2
Asian or Pacific Islander, female, 65-69 years				
...	All causes	1,905	100.0	576.0
1	Malignant neoplasms (C00-C97)	801	42.0	242.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	311	16.3	94.0
3	Cerebrovascular diseases (I60-I69)	135	7.1	40.8
4	Diabetes mellitus (E10-E14)	115	6.0	34.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	50	2.6	15.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	45	2.4	13.6
7	Chronic lower respiratory diseases (J40-J47)	41	2.2	12.4
8	Influenza and pneumonia (J09-J18)	31	1.6	9.4
9	Septicemia (A40-A41)	27	1.4	8.2
10	Viral hepatitis (B15-B19)	21	1.1	6.4
11	Chronic liver disease and cirrhosis (K70,K73-K74)	19	1.0	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	0.9	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	18	0.9	*
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	15	0.8	*
15	Parkinson's disease (G20-G21)	10	0.5	*
...	All other causes (Residual)	248	13.0	75.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 70-74 years				
...	All causes	2,257	100.0	948.4
1	Malignant neoplasms (C00-C97)	850	37.7	357.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	410	18.2	172.3
3	Cerebrovascular diseases (I60-I69)	185	8.2	77.7
4	Diabetes mellitus (E10-E14)	120	5.3	50.4
5	Chronic lower respiratory diseases (J40-J47)	56	2.5	23.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	53	2.3	22.3
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	52	2.3	21.9
8	Influenza and pneumonia (J09-J18)	46	2.0	19.3
9	Septicemia (A40-A41)	33	1.5	13.9
10	Chronic liver disease and cirrhosis (K70,K73-K74)	31	1.4	13.0
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	26	1.2	10.9
12	Alzheimer's disease (G30)	23	1.0	9.7
13	Parkinson's disease (G20-G21)	22	1.0	9.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	21	0.9	8.8
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	20	0.9	8.4
...	All other causes (Residual)	309	13.7	129.8
Asian or Pacific Islander, female, 75-79 years				
...	All causes	2,952	100.0	1,699.9
1	Malignant neoplasms (C00-C97)	903	30.6	520.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	592	20.1	340.9
3	Cerebrovascular diseases (I60-I69)	275	9.3	158.4
4	Diabetes mellitus (E10-E14)	142	4.8	81.8
5	Chronic lower respiratory diseases (J40-J47)	81	2.7	46.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	80	2.7	46.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	74	2.5	42.6
8	Alzheimer's disease (G30)	70	2.4	40.3
9	Influenza and pneumonia (J09-J18)	65	2.2	37.4
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	53	1.8	30.5
11	Septicemia (A40-A41)	47	1.6	27.1
12	Parkinson's disease (G20-G21)	31	1.1	17.9
13	Chronic liver disease and cirrhosis (K70,K73-K74)	29	1.0	16.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	21	0.7	12.1
15	Viral hepatitis (B15-B19)	17	0.6	*
...	All other causes (Residual)	472	16.0	271.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 80-84 years				
...	All causes	3,982	100.0	3,215.0
1	Malignant neoplasms (C00-C97)	961	24.1	775.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	911	22.9	735.5
3	Cerebrovascular diseases (I60-I69)	360	9.0	290.7
4	Diabetes mellitus (E10-E14)	189	4.7	152.6
5	Alzheimer's disease (G30)	163	4.1	131.6
6	Influenza and pneumonia (J09-J18)	140	3.5	113.0
7	Chronic lower respiratory diseases (J40-J47)	123	3.1	99.3
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	110	2.8	88.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	96	2.4	77.5
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	79	2.0	63.8
11	Parkinson's disease (G20-G21)	48	1.2	38.8
12	Septicemia (A40-A41)	44	1.1	35.5
13	Pneumonitis due to solids and liquids (J69)	32	0.8	25.8
14	Chronic liver disease and cirrhosis (K70,K73-K74)	31	0.8	25.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	24	0.6	19.4
...	All other causes (Residual)	671	16.9	541.8
Asian or Pacific Islander, female, 85-89 years				
...	All causes	4,456	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,185	26.6	---
2	Malignant neoplasms (C00-C97)	717	16.1	---
3	Cerebrovascular diseases (I60-I69)	449	10.1	---
4	Alzheimer's disease (G30)	243	5.5	---
5	Influenza and pneumonia (J09-J18)	194	4.4	---
6	Diabetes mellitus (E10-E14)	174	3.9	---
7	Chronic lower respiratory diseases (J40-J47)	149	3.3	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	116	2.6	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	95	2.1	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	81	1.8	---
11	Septicemia (A40-A41)	61	1.4	---
12	Parkinson's disease (G20-G21)	60	1.3	---
13	Pneumonitis due to solids and liquids (J69)	36	0.8	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	24	0.5	---
15	Aortic aneurysm and dissection (I71)	19	0.4	---
...	All other causes (Residual)	853	19.1	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 90-94 years				
...	All causes	3,346	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	977	29.2	---
2	Malignant neoplasms (C00-C97)	334	10.0	---
3	Cerebrovascular diseases (I60-I69)	302	9.0	---
4	Alzheimer's disease (G30)	248	7.4	---
5	Influenza and pneumonia (J09-J18)	178	5.3	---
6	Diabetes mellitus (E10-E14)	102	3.0	---
7	Chronic lower respiratory diseases (J40-J47)	85	2.5	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	80	2.4	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	79	2.4	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	64	1.9	---
11	Septicemia (A40-A41)	44	1.3	---
12	Pneumonitis due to solids and liquids (J69)	41	1.2	---
13	Parkinson's disease (G20-G21)	34	1.0	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	0.7	---
15	Aortic aneurysm and dissection (I71)	20	0.6	---
...	All other causes (Residual)	735	22.0	---
Asian or Pacific Islander, female, 95-99 years				
...	All causes	1,529	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	461	30.2	---
2	Alzheimer's disease (G30)	139	9.1	---
3	Cerebrovascular diseases (I60-I69)	106	6.9	---
4	Malignant neoplasms (C00-C97)	101	6.6	---
5	Influenza and pneumonia (J09-J18)	95	6.2	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	55	3.6	---
7	Chronic lower respiratory diseases (J40-J47)	46	3.0	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	35	2.3	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	29	1.9	---
10	Diabetes mellitus (E10-E14)	28	1.8	---
11	Septicemia (A40-A41)	19	1.2	---
12	Pneumonitis due to solids and liquids (J69)	17	1.1	---
13	Parkinson's disease (G20-G21)	10	0.7	---
13	Atherosclerosis (I70)	10	0.7	---
15	Anemias (D50-D64)	8	0.5	---
...	All other causes (Residual)	370	24.2	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 100 years and over				
...	All causes	413	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	117	28.3	---
2	Alzheimer's disease (G30)	42	10.2	---
3	Influenza and pneumonia (J09-J18)	33	8.0	---
4	Cerebrovascular diseases (I60-I69)	29	7.0	---
5	Essential hypertension and hypertensive renal disease (I10,I12,I15)	17	4.1	---
6	Malignant neoplasms (C00-C97)	13	3.1	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	3.1	---
8	Diabetes mellitus (E10-E14)	9	2.2	---
9	Chronic lower respiratory diseases (J40-J47)	5	1.2	---
10	Pneumonitis due to solids and liquids (J69)	4	1.0	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4	1.0	---
12	Atherosclerosis (I70)	3	0.7	---
13	Septicemia (A40-A41)	2	0.5	---
13	Aortic aneurysm and dissection (I71)	2	0.5	---
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.2	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.2	---
...	All other causes (Residual)	118	28.6	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012