

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, all ages				
...	All causes	2,403,351	100.0	873.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	710,760	29.6	258.2
2	Malignant neoplasms (C00-C97)	553,091	23.0	200.9
3	Cerebrovascular diseases (I60-I69)	167,661	7.0	60.9
4	Chronic lower respiratory diseases (J40-J47)	122,009	5.1	44.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	97,900	4.1	35.6
6	Diabetes mellitus (E10-E14)	69,301	2.9	25.2
7	Influenza and pneumonia (J10-J18)	65,313	2.7	23.7
8	Alzheimer's disease (G30)	49,558	2.1	18.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	37,251	1.5	13.5
10	Septicemia (A40-A41)	31,224	1.3	11.3
11	Intentional self-harm (suicide) (X60-X84,Y87.0)	29,350	1.2	10.7
12	Chronic liver disease and cirrhosis (K70,K73-K74)	26,552	1.1	9.6
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	18,073	0.8	6.6
14	Assault (homicide) (X85-Y09,Y87.1)	16,765	0.7	6.1
15	Pneumonitis due to solids and liquids (J69)	16,636	0.7	6.0
...	All other causes (Residual)	391,907	16.3	142.4
All races, both sexes, 1-4 years				
...	All causes	4,979	100.0	32.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,826	36.7	12.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	495	9.9	3.3
3	Malignant neoplasms (C00-C97)	420	8.4	2.8
4	Assault (homicide) (X85-Y09,Y87.1)	356	7.2	2.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	181	3.6	1.2
6	Influenza and pneumonia (J10-J18)	103	2.1	0.7
7	Septicemia (A40-A41)	99	2.0	0.7
8	Certain conditions originating in the perinatal period (P00-P96)	79	1.6	0.5
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	53	1.1	0.3
10	Chronic lower respiratory diseases (J40-J47)	51	1.0	0.3
11	Cerebrovascular diseases (I60-I69)	47	0.9	0.3
12	Meningitis (G00,G03)	36	0.7	0.2
13	Anemias (D50-D64)	35	0.7	0.2
14	Complications of medical and surgical care (Y40-Y84,Y88)	22	0.4	0.1
15	Meningococcal infection (A39)	19	0.4	*
...	All other causes (Residual)	1,157	23.2	7.6

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 5-14 years				
...	All causes	7,413	100.0	18.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,979	40.2	7.5
2	Malignant neoplasms (C00-C97)	1,014	13.7	2.6
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	399	5.4	1.0
4	Assault (homicide) (X85-Y09,Y87.1)	371	5.0	0.9
5	Intentional self-harm (suicide) (X60-X84,Y87.0)	307	4.1	0.8
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	271	3.7	0.7
7	Chronic lower respiratory diseases (J40-J47)	139	1.9	0.4
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	99	1.3	0.2
9	Influenza and pneumonia (J10-J18)	87	1.2	0.2
10	Cerebrovascular diseases (I60-I69)	76	1.0	0.2
11	Septicemia (A40-A41)	63	0.8	0.2
12	Anemias (D50-D64)	52	0.7	0.1
13	Human immunodeficiency virus (HIV) disease (B20-B24)	46	0.6	0.1
14	Certain conditions originating in the perinatal period (P00-P96)	41	0.6	0.1
15	Complications of medical and surgical care (Y40-Y84,Y88)	31	0.4	0.1
...	All other causes (Residual)	1,438	19.4	3.6
All races, both sexes, 15-24 years				
...	All causes	31,307	100.0	81.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14,113	45.1	36.8
2	Assault (homicide) (X85-Y09,Y87.1)	4,939	15.8	12.9
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	3,994	12.8	10.4
4	Malignant neoplasms (C00-C97)	1,713	5.5	4.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,031	3.3	2.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	441	1.4	1.1
7	Cerebrovascular diseases (I60-I69)	199	0.6	0.5
8	Chronic lower respiratory diseases (J40-J47)	190	0.6	0.5
9	Influenza and pneumonia (J10-J18)	189	0.6	0.5
10	Human immunodeficiency virus (HIV) disease (B20-B24)	179	0.6	0.5
11	Diabetes mellitus (E10-E14)	162	0.5	0.4
12	Septicemia (A40-A41)	100	0.3	0.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	95	0.3	0.2
13	Pregnancy, childbirth and the puerperium (O00-099)	95	0.3	0.2
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	88	0.3	0.2
...	All other causes (Residual)	3,779	12.1	9.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 25-34 years				
...	All causes	40,451	100.0	108.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,769	29.1	31.4
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	4,792	11.8	12.8
3	Assault (homicide) (X85-Y09,Y87.1)	4,164	10.3	11.1
4	Malignant neoplasms (C00-C97)	3,916	9.7	10.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,958	7.3	7.9
6	Human immunodeficiency virus (HIV) disease (B20-B24)	2,437	6.0	6.5
7	Diabetes mellitus (E10-E14)	623	1.5	1.7
8	Cerebrovascular diseases (I60-I69)	602	1.5	1.6
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	477	1.2	1.3
10	Chronic liver disease and cirrhosis (K70,K73-K74)	415	1.0	1.1
11	Influenza and pneumonia (J10-J18)	364	0.9	1.0
12	Chronic lower respiratory diseases (J40-J47)	296	0.7	0.8
13	Septicemia (A40-A41)	280	0.7	0.7
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	244	0.6	0.7
15	Pregnancy, childbirth and the puerperium (O00-099)	182	0.4	0.5
...	All other causes (Residual)	6,932	17.1	18.5
All races, both sexes, 35-44 years				
...	All causes	89,798	100.0	200.0
1	Malignant neoplasms (C00-C97)	16,520	18.4	36.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15,413	17.2	34.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,181	14.7	29.4
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	6,562	7.3	14.6
5	Human immunodeficiency virus (HIV) disease (B20-B24)	5,919	6.6	13.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	3,371	3.8	7.5
7	Assault (homicide) (X85-Y09,Y87.1)	3,219	3.6	7.2
8	Cerebrovascular diseases (I60-I69)	2,599	2.9	5.8
9	Diabetes mellitus (E10-E14)	1,926	2.1	4.3
10	Influenza and pneumonia (J10-J18)	1,068	1.2	2.4
11	Chronic lower respiratory diseases (J40-J47)	930	1.0	2.1
12	Septicemia (A40-A41)	877	1.0	2.0
13	Viral hepatitis (B15-B19)	779	0.9	1.7
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	740	0.8	1.6
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	569	0.6	1.3
...	All other causes (Residual)	16,125	18.0	35.9

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 45-54 years				
...	All causes	160,341	100.0	431.6
1	Malignant neoplasms (C00-C97)	48,034	30.0	129.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	35,480	22.1	95.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,278	7.7	33.0
4	Chronic liver disease and cirrhosis (K70,K73-K74)	6,654	4.1	17.9
5	Cerebrovascular diseases (I60-I69)	6,011	3.7	16.2
6	Intentional self-harm (suicide) (X60-X84,Y87.0)	5,437	3.4	14.6
7	Diabetes mellitus (E10-E14)	4,954	3.1	13.3
8	Human immunodeficiency virus (HIV) disease (B20-B24)	4,142	2.6	11.1
9	Chronic lower respiratory diseases (J40-J47)	3,251	2.0	8.8
10	Viral hepatitis (B15-B19)	1,894	1.2	5.1
11	Septicemia (A40-A41)	1,845	1.2	5.0
12	Influenza and pneumonia (J10-J18)	1,774	1.1	4.8
13	Assault (homicide) (X85-Y09,Y87.1)	1,755	1.1	4.7
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,651	1.0	4.4
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	878	0.5	2.4
...	All other causes (Residual)	24,303	15.2	65.4
All races, both sexes, 55-64 years				
...	All causes	240,846	100.0	1,004.6
1	Malignant neoplasms (C00-C97)	89,005	37.0	371.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	63,399	26.3	264.5
3	Chronic lower respiratory diseases (J40-J47)	10,739	4.5	44.8
4	Cerebrovascular diseases (I60-I69)	9,956	4.1	41.5
5	Diabetes mellitus (E10-E14)	9,186	3.8	38.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,505	3.1	31.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	5,774	2.4	24.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,100	1.3	12.9
9	Intentional self-harm (suicide) (X60-X84,Y87.0)	2,945	1.2	12.3
10	Septicemia (A40-A41)	2,899	1.2	12.1
11	Influenza and pneumonia (J10-J18)	2,879	1.2	12.0
12	Aortic aneurysm and dissection (I71)	1,493	0.6	6.2
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,440	0.6	6.0
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1,239	0.5	5.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,224	0.5	5.1
...	All other causes (Residual)	28,063	11.7	117.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 65-74 years				
...	All causes	441,209	100.0	2,428.6
1	Malignant neoplasms (C00-C97)	150,131	34.0	826.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	122,405	27.7	673.8
3	Chronic lower respiratory diseases (J40-J47)	31,157	7.1	171.5
4	Cerebrovascular diseases (I60-I69)	23,649	5.4	130.2
5	Diabetes mellitus (E10-E14)	16,674	3.8	91.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,698	1.7	42.4
7	Influenza and pneumonia (J10-J18)	7,189	1.6	39.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,990	1.6	38.5
9	Septicemia (A40-A41)	5,704	1.3	31.4
10	Chronic liver disease and cirrhosis (K70,K73-K74)	5,482	1.2	30.2
11	Aortic aneurysm and dissection (I71)	4,012	0.9	22.1
12	Alzheimer's disease (G30)	3,431	0.8	18.9
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,784	0.6	15.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,546	0.6	14.0
15	Intentional self-harm (suicide) (X60-X84,Y87.0)	2,292	0.5	12.6
...	All other causes (Residual)	49,065	11.1	270.1
All races, both sexes, 75-84 years				
...	All causes	700,445	100.0	5,688.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	220,060	31.4	1,787.1
2	Malignant neoplasms (C00-C97)	165,099	23.6	1,340.8
3	Cerebrovascular diseases (I60-I69)	57,020	8.1	463.1
4	Chronic lower respiratory diseases (J40-J47)	47,722	6.8	387.6
5	Diabetes mellitus (E10-E14)	22,184	3.2	180.2
6	Influenza and pneumonia (J10-J18)	19,821	2.8	161.0
7	Alzheimer's disease (G30)	17,253	2.5	140.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12,458	1.8	101.2
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,758	1.7	95.5
10	Septicemia (A40-A41)	9,938	1.4	80.7
11	Parkinson's disease (G20-G21)	7,652	1.1	62.1
12	Aortic aneurysm and dissection (I71)	5,835	0.8	47.4
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	5,623	0.8	45.7
14	Pneumonitis due to solids and liquids (J69)	5,505	0.8	44.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,645	0.7	37.7
...	All other causes (Residual)	87,872	12.5	713.6

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 85 years and over				
...	All causes	658,171	100.0	15,321.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	251,242	38.2	5,848.7
2	Malignant neoplasms (C00-C97)	77,136	11.7	1,795.6
3	Cerebrovascular diseases (I60-I69)	67,376	10.2	1,568.4
4	Influenza and pneumonia (J10-J18)	31,547	4.8	734.4
5	Alzheimer's disease (G30)	28,309	4.3	659.0
6	Chronic lower respiratory diseases (J40-J47)	27,496	4.2	640.1
7	Diabetes mellitus (E10-E14)	13,556	2.1	315.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11,777	1.8	274.2
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,595	1.8	269.9
10	Septicemia (A40-A41)	9,144	1.4	212.9
11	Atherosclerosis (I70)	8,005	1.2	186.3
12	Pneumonitis due to solids and liquids (J69)	7,954	1.2	185.2
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	6,907	1.0	160.8
14	Parkinson's disease (G20-G21)	5,590	0.8	130.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3,600	0.5	83.8
...	All other causes (Residual)	96,937	14.7	2,256.6
All races, male, all ages				
...	All causes	1,177,578	100.0	874.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	344,807	29.3	256.1
2	Malignant neoplasms (C00-C97)	286,082	24.3	212.5
3	Cerebrovascular diseases (I60-I69)	64,769	5.5	48.1
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	63,817	5.4	47.4
5	Chronic lower respiratory diseases (J40-J47)	60,004	5.1	44.6
6	Diabetes mellitus (E10-E14)	31,602	2.7	23.5
7	Influenza and pneumonia (J10-J18)	28,658	2.4	21.3
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	23,618	2.0	17.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17,811	1.5	13.2
10	Chronic liver disease and cirrhosis (K70,K73-K74)	17,214	1.5	12.8
11	Alzheimer's disease (G30)	14,438	1.2	10.7
12	Septicemia (A40-A41)	13,537	1.1	10.1
13	Assault (homicide) (X85-Y09,Y87.1)	12,820	1.1	9.5
14	Human immunodeficiency virus (HIV) disease (B20-B24)	10,868	0.9	8.1
15	Aortic aneurysm and dissection (I71)	9,661	0.8	7.2
...	All other causes (Residual)	177,872	15.1	132.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 1-4 years				
...	All causes	2,824	100.0	36.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,098	38.9	14.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	259	9.2	3.3
3	Malignant neoplasms (C00-C97)	233	8.3	3.0
4	Assault (homicide) (X85-Y09,Y87.1)	198	7.0	2.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	107	3.8	1.4
6	Influenza and pneumonia (J10-J18)	54	1.9	0.7
7	Septicemia (A40-A41)	53	1.9	0.7
8	Certain conditions originating in the perinatal period (P00-P96)	52	1.8	0.7
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	33	1.2	0.4
10	Chronic lower respiratory diseases (J40-J47)	28	1.0	0.4
11	Meningitis (G00,G03)	20	0.7	0.3
12	Cerebrovascular diseases (I60-I69)	19	0.7	*
13	Anemias (D50-D64)	16	0.6	*
14	Meningococcal infection (A39)	12	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	11	0.4	*
...	All other causes (Residual)	631	22.3	8.1
All races, male, 5-14 years				
...	All causes	4,401	100.0	21.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,861	42.3	9.2
2	Malignant neoplasms (C00-C97)	569	12.9	2.8
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	244	5.5	1.2
4	Assault (homicide) (X85-Y09,Y87.1)	227	5.2	1.1
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	207	4.7	1.0
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	163	3.7	0.8
7	Chronic lower respiratory diseases (J40-J47)	80	1.8	0.4
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	49	1.1	0.2
9	Influenza and pneumonia (J10-J18)	48	1.1	0.2
10	Septicemia (A40-A41)	38	0.9	0.2
11	Cerebrovascular diseases (I60-I69)	33	0.7	0.2
12	Anemias (D50-D64)	29	0.7	0.1
13	Certain conditions originating in the perinatal period (P00-P96)	23	0.5	0.1
14	Human immunodeficiency virus (HIV) disease (B20-B24)	22	0.5	0.1
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	0.4	*
...	All other causes (Residual)	791	18.0	3.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 15-24 years				
...	All causes	23,071	100.0	117.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,460	45.3	53.3
2	Assault (homicide) (X85-Y09,Y87.1)	4,203	18.2	21.4
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	3,424	14.8	17.4
4	Malignant neoplasms (C00-C97)	1,022	4.4	5.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	638	2.8	3.2
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	268	1.2	1.4
7	Chronic lower respiratory diseases (J40-J47)	120	0.5	0.6
8	Cerebrovascular diseases (I60-I69)	106	0.5	0.5
9	Influenza and pneumonia (J10-J18)	102	0.4	0.5
10	Human immunodeficiency virus (HIV) disease (B20-B24)	99	0.4	0.5
11	Diabetes mellitus (E10-E14)	94	0.4	0.5
12	Legal intervention (Y35,Y89.0)	71	0.3	0.4
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	62	0.3	0.3
14	Septicemia (A40-A41)	60	0.3	0.3
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	43	0.2	0.2
...	All other causes (Residual)	2,299	10.0	11.7
All races, male, 25-34 years				
...	All causes	27,890	100.0	149.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,059	32.5	48.7
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	3,938	14.1	21.2
3	Assault (homicide) (X85-Y09,Y87.1)	3,361	12.1	18.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,922	6.9	10.3
5	Malignant neoplasms (C00-C97)	1,854	6.6	10.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	1,602	5.7	8.6
7	Diabetes mellitus (E10-E14)	376	1.3	2.0
8	Cerebrovascular diseases (I60-I69)	308	1.1	1.7
9	Chronic liver disease and cirrhosis (K70,K73-K74)	280	1.0	1.5
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	279	1.0	1.5
11	Influenza and pneumonia (J10-J18)	204	0.7	1.1
12	Chronic lower respiratory diseases (J40-J47)	165	0.6	0.9
13	Septicemia (A40-A41)	158	0.6	0.8
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	129	0.5	0.7
15	Legal intervention (Y35,Y89.0)	108	0.4	0.6
...	All other causes (Residual)	4,147	14.9	22.3

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 35-44 years				
...	All causes	57,297	100.0	256.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,444	20.0	51.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,286	16.2	41.6
3	Malignant neoplasms (C00-C97)	7,344	12.8	32.9
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	5,114	8.9	22.9
5	Human immunodeficiency virus (HIV) disease (B20-B24)	4,441	7.8	19.9
6	Assault (homicide) (X85-Y09,Y87.1)	2,319	4.0	10.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,314	4.0	10.4
8	Cerebrovascular diseases (I60-I69)	1,297	2.3	5.8
9	Diabetes mellitus (E10-E14)	1,151	2.0	5.2
10	Influenza and pneumonia (J10-J18)	630	1.1	2.8
11	Viral hepatitis (B15-B19)	540	0.9	2.4
12	Septicemia (A40-A41)	451	0.8	2.0
13	Chronic lower respiratory diseases (J40-J47)	437	0.8	2.0
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	429	0.7	1.9
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	316	0.6	1.4
...	All other causes (Residual)	9,784	17.1	43.9
All races, male, 45-54 years				
...	All causes	100,398	100.0	552.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	25,927	25.8	142.7
2	Malignant neoplasms (C00-C97)	24,215	24.1	133.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,063	9.0	49.9
4	Chronic liver disease and cirrhosis (K70,K73-K74)	4,975	5.0	27.4
5	Intentional self-harm (suicide) (X60-X84,Y87.0)	4,149	4.1	22.8
6	Human immunodeficiency virus (HIV) disease (B20-B24)	3,296	3.3	18.1
7	Cerebrovascular diseases (I60-I69)	3,235	3.2	17.8
8	Diabetes mellitus (E10-E14)	2,821	2.8	15.5
9	Chronic lower respiratory diseases (J40-J47)	1,661	1.7	9.1
10	Viral hepatitis (B15-B19)	1,425	1.4	7.8
11	Assault (homicide) (X85-Y09,Y87.1)	1,284	1.3	7.1
12	Influenza and pneumonia (J10-J18)	1,066	1.1	5.9
13	Septicemia (A40-A41)	1,039	1.0	5.7
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	982	1.0	5.4
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	555	0.6	3.1
...	All other causes (Residual)	14,705	14.6	80.9

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 55-64 years				
...	All causes	143,321	100.0	1,253.5
1	Malignant neoplasms (C00-C97)	48,427	33.8	423.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	43,285	30.2	378.6
3	Chronic lower respiratory diseases (J40-J47)	5,564	3.9	48.7
4	Cerebrovascular diseases (I60-I69)	5,496	3.8	48.1
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,133	3.6	44.9
6	Diabetes mellitus (E10-E14)	4,943	3.4	43.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	4,005	2.8	35.0
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	2,265	1.6	19.8
9	Influenza and pneumonia (J10-J18)	1,680	1.2	14.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,651	1.2	14.4
11	Septicemia (A40-A41)	1,513	1.1	13.2
12	Aortic aneurysm and dissection (I71)	1,111	0.8	9.7
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,013	0.7	8.9
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	818	0.6	7.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	657	0.5	5.7
...	All other causes (Residual)	15,760	11.0	137.8
All races, male, 65-74 years				
...	All causes	247,408	100.0	3,015.7
1	Malignant neoplasms (C00-C97)	83,191	33.6	1,014.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	74,589	30.1	909.2
3	Chronic lower respiratory diseases (J40-J47)	16,210	6.6	197.6
4	Cerebrovascular diseases (I60-I69)	12,042	4.9	146.8
5	Diabetes mellitus (E10-E14)	8,226	3.3	100.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,639	1.9	56.5
7	Influenza and pneumonia (J10-J18)	4,013	1.6	48.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,564	1.4	43.4
9	Chronic liver disease and cirrhosis (K70,K73-K74)	3,317	1.3	40.4
10	Septicemia (A40-A41)	2,867	1.2	34.9
11	Aortic aneurysm and dissection (I71)	2,730	1.1	33.3
12	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,888	0.8	23.0
13	Alzheimer's disease (G30)	1,554	0.6	18.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,415	0.6	17.2
15	Parkinson's disease (G20-G21)	1,338	0.5	16.3
...	All other causes (Residual)	25,825	10.4	314.8

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 75-84 years				
...	All causes	340,219	100.0	6,854.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	109,693	32.2	2,210.1
2	Malignant neoplasms (C00-C97)	85,907	25.3	1,730.9
3	Cerebrovascular diseases (I60-I69)	23,946	7.0	482.5
4	Chronic lower respiratory diseases (J40-J47)	23,838	7.0	480.3
5	Influenza and pneumonia (J10-J18)	9,751	2.9	196.5
6	Diabetes mellitus (E10-E14)	9,738	2.9	196.2
7	Alzheimer's disease (G30)	6,228	1.8	125.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,213	1.8	125.2
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,063	1.8	122.2
10	Parkinson's disease (G20-G21)	4,530	1.3	91.3
11	Septicemia (A40-A41)	4,482	1.3	90.3
12	Aortic aneurysm and dissection (I71)	3,443	1.0	69.4
13	Pneumonitis due to solids and liquids (J69)	3,038	0.9	61.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,319	0.7	46.7
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,204	0.6	44.4
...	All other causes (Residual)	38,826	11.4	782.3
All races, male, 85 years and over				
...	All causes	214,742	100.0	16,605.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	78,896	36.7	6,100.8
2	Malignant neoplasms (C00-C97)	33,260	15.5	2,571.9
3	Cerebrovascular diseases (I60-I69)	18,212	8.5	1,408.3
4	Chronic lower respiratory diseases (J40-J47)	11,876	5.5	918.3
5	Influenza and pneumonia (J10-J18)	10,945	5.1	846.3
6	Alzheimer's disease (G30)	6,406	3.0	495.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,671	2.2	361.2
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,413	2.1	341.2
9	Diabetes mellitus (E10-E14)	4,234	2.0	327.4
10	Pneumonitis due to solids and liquids (J69)	3,387	1.6	261.9
11	Septicemia (A40-A41)	2,736	1.3	211.6
12	Parkinson's disease (G20-G21)	2,728	1.3	210.9
13	Atherosclerosis (I70)	2,166	1.0	167.5
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,747	0.8	135.1
15	Aortic aneurysm and dissection (I71)	1,562	0.7	120.8
...	All other causes (Residual)	27,503	12.8	2,126.7

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, all ages				
...	All causes	1,225,773	100.0	871.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	365,953	29.9	260.2
2	Malignant neoplasms (C00-C97)	267,009	21.8	189.9
3	Cerebrovascular diseases (I60-I69)	102,892	8.4	73.2
4	Chronic lower respiratory diseases (J40-J47)	62,005	5.1	44.1
5	Diabetes mellitus (E10-E14)	37,699	3.1	26.8
6	Influenza and pneumonia (J10-J18)	36,655	3.0	26.1
7	Alzheimer's disease (G30)	35,120	2.9	25.0
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	34,083	2.8	24.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19,440	1.6	13.8
10	Septicemia (A40-A41)	17,687	1.4	12.6
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	11,178	0.9	7.9
12	Chronic liver disease and cirrhosis (K70,K73-K74)	9,338	0.8	6.6
13	Atherosclerosis (I70)	9,036	0.7	6.4
14	Pneumonitis due to solids and liquids (J69)	8,246	0.7	5.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6,987	0.6	5.0
...	All other causes (Residual)	202,445	16.5	143.9
All races, female, 1-4 years				
...	All causes	2,155	100.0	29.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	728	33.8	9.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	236	11.0	3.2
3	Malignant neoplasms (C00-C97)	187	8.7	2.5
4	Assault (homicide) (X85-Y09,Y87.1)	158	7.3	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	74	3.4	1.0
6	Influenza and pneumonia (J10-J18)	49	2.3	0.7
7	Septicemia (A40-A41)	46	2.1	0.6
8	Cerebrovascular diseases (I60-I69)	28	1.3	0.4
9	Certain conditions originating in the perinatal period (P00-P96)	27	1.3	0.4
10	Chronic lower respiratory diseases (J40-J47)	23	1.1	0.3
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	0.9	0.3
12	Anemias (D50-D64)	19	0.9	*
13	Meningitis (G00,G03)	16	0.7	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	11	0.5	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	8	0.4	*
...	All other causes (Residual)	525	24.4	7.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 5-14 years				
...	All causes	3,012	100.0	15.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,118	37.1	5.8
2	Malignant neoplasms (C00-C97)	445	14.8	2.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	192	6.4	1.0
4	Assault (homicide) (X85-Y09,Y87.1)	144	4.8	0.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	108	3.6	0.6
6	Intentional self-harm (suicide) (X60-X84,Y87.0)	63	2.1	0.3
7	Chronic lower respiratory diseases (J40-J47)	59	2.0	0.3
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	50	1.7	0.3
9	Cerebrovascular diseases (I60-I69)	43	1.4	0.2
10	Influenza and pneumonia (J10-J18)	39	1.3	0.2
11	Septicemia (A40-A41)	25	0.8	0.1
12	Human immunodeficiency virus (HIV) disease (B20-B24)	24	0.8	0.1
13	Anemias (D50-D64)	23	0.8	0.1
14	Certain conditions originating in the perinatal period (P00-P96)	18	0.6	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	16	0.5	*
...	All other causes (Residual)	645	21.4	3.3
All races, female, 15-24 years				
...	All causes	8,236	100.0	44.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,653	44.4	19.5
2	Assault (homicide) (X85-Y09,Y87.1)	736	8.9	3.9
3	Malignant neoplasms (C00-C97)	691	8.4	3.7
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	570	6.9	3.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	393	4.8	2.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	173	2.1	0.9
7	Pregnancy, childbirth and the puerperium (O00-099)	95	1.2	0.5
8	Cerebrovascular diseases (I60-I69)	93	1.1	0.5
9	Influenza and pneumonia (J10-J18)	87	1.1	0.5
10	Human immunodeficiency virus (HIV) disease (B20-B24)	80	1.0	0.4
11	Chronic lower respiratory diseases (J40-J47)	70	0.8	0.4
12	Diabetes mellitus (E10-E14)	68	0.8	0.4
13	Anemias (D50-D64)	46	0.6	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	45	0.5	0.2
15	Septicemia (A40-A41)	40	0.5	0.2
...	All other causes (Residual)	1,396	16.9	7.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 25-34 years				
...	All causes	12,561	100.0	66.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,710	21.6	14.4
2	Malignant neoplasms (C00-C97)	2,062	16.4	11.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,036	8.2	5.5
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	854	6.8	4.5
5	Human immunodeficiency virus (HIV) disease (B20-B24)	835	6.6	4.4
6	Assault (homicide) (X85-Y09,Y87.1)	803	6.4	4.3
7	Cerebrovascular diseases (I60-I69)	294	2.3	1.6
8	Diabetes mellitus (E10-E14)	247	2.0	1.3
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	198	1.6	1.1
10	Pregnancy, childbirth and the puerperium (O00-099)	182	1.4	1.0
11	Influenza and pneumonia (J10-J18)	160	1.3	0.8
12	Chronic liver disease and cirrhosis (K70,K73-K74)	135	1.1	0.7
13	Chronic lower respiratory diseases (J40-J47)	131	1.0	0.7
14	Septicemia (A40-A41)	122	1.0	0.6
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	115	0.9	0.6
...	All other causes (Residual)	2,677	21.3	14.2
All races, female, 35-44 years				
...	All causes	32,501	100.0	143.9
1	Malignant neoplasms (C00-C97)	9,176	28.2	40.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,969	12.2	17.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,895	12.0	17.2
4	Human immunodeficiency virus (HIV) disease (B20-B24)	1,478	4.5	6.5
5	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,448	4.5	6.4
6	Cerebrovascular diseases (I60-I69)	1,302	4.0	5.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,057	3.3	4.7
8	Assault (homicide) (X85-Y09,Y87.1)	900	2.8	4.0
9	Diabetes mellitus (E10-E14)	775	2.4	3.4
10	Chronic lower respiratory diseases (J40-J47)	493	1.5	2.2
11	Influenza and pneumonia (J10-J18)	438	1.3	1.9
12	Septicemia (A40-A41)	426	1.3	1.9
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	311	1.0	1.4
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	253	0.8	1.1
15	Viral hepatitis (B15-B19)	239	0.7	1.1
...	All other causes (Residual)	6,341	19.5	28.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 45-54 years				
...	All causes	59,943	100.0	315.8
1	Malignant neoplasms (C00-C97)	23,819	39.7	125.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,553	15.9	50.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,215	5.4	16.9
4	Cerebrovascular diseases (I60-I69)	2,776	4.6	14.6
5	Diabetes mellitus (E10-E14)	2,133	3.6	11.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1,679	2.8	8.8
7	Chronic lower respiratory diseases (J40-J47)	1,590	2.7	8.4
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,288	2.1	6.8
9	Human immunodeficiency virus (HIV) disease (B20-B24)	846	1.4	4.5
10	Septicemia (A40-A41)	806	1.3	4.2
11	Influenza and pneumonia (J10-J18)	708	1.2	3.7
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	669	1.1	3.5
13	Assault (homicide) (X85-Y09,Y87.1)	471	0.8	2.5
14	Viral hepatitis (B15-B19)	469	0.8	2.5
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	323	0.5	1.7
...	All other causes (Residual)	9,598	16.0	50.6
All races, female, 55-64 years				
...	All causes	97,525	100.0	777.7
1	Malignant neoplasms (C00-C97)	40,578	41.6	323.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,114	20.6	160.4
3	Chronic lower respiratory diseases (J40-J47)	5,175	5.3	41.3
4	Cerebrovascular diseases (I60-I69)	4,460	4.6	35.6
5	Diabetes mellitus (E10-E14)	4,243	4.4	33.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,372	2.4	18.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,769	1.8	14.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,449	1.5	11.6
9	Septicemia (A40-A41)	1,386	1.4	11.1
10	Influenza and pneumonia (J10-J18)	1,199	1.2	9.6
11	Intentional self-harm (suicide) (X60-X84,Y87.0)	680	0.7	5.4
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	622	0.6	5.0
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	567	0.6	4.5
14	Aortic aneurysm and dissection (I71)	382	0.4	3.0
15	Viral hepatitis (B15-B19)	327	0.3	2.6
...	All other causes (Residual)	12,202	12.5	97.3

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 65-74 years				
...	All causes	193,801	100.0	1,945.1
1	Malignant neoplasms (C00-C97)	66,940	34.5	671.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	47,816	24.7	479.9
3	Chronic lower respiratory diseases (J40-J47)	14,947	7.7	150.0
4	Cerebrovascular diseases (I60-I69)	11,607	6.0	116.5
5	Diabetes mellitus (E10-E14)	8,448	4.4	84.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,426	1.8	34.4
7	Influenza and pneumonia (J10-J18)	3,176	1.6	31.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,059	1.6	30.7
9	Septicemia (A40-A41)	2,837	1.5	28.5
10	Chronic liver disease and cirrhosis (K70,K73-K74)	2,165	1.1	21.7
11	Alzheimer's disease (G30)	1,877	1.0	18.8
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,476	0.8	14.8
13	Aortic aneurysm and dissection (I71)	1,282	0.7	12.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,131	0.6	11.4
15	Parkinson's disease (G20-G21)	770	0.4	7.7
...	All other causes (Residual)	22,844	11.8	229.3
All races, female, 75-84 years				
...	All causes	360,226	100.0	4,900.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	110,367	30.6	1,501.5
2	Malignant neoplasms (C00-C97)	79,192	22.0	1,077.4
3	Cerebrovascular diseases (I60-I69)	33,074	9.2	450.0
4	Chronic lower respiratory diseases (J40-J47)	23,884	6.6	324.9
5	Diabetes mellitus (E10-E14)	12,446	3.5	169.3
6	Alzheimer's disease (G30)	11,025	3.1	150.0
7	Influenza and pneumonia (J10-J18)	10,070	2.8	137.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,245	1.7	85.0
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,695	1.6	77.5
10	Septicemia (A40-A41)	5,456	1.5	74.2
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	3,419	0.9	46.5
12	Parkinson's disease (G20-G21)	3,122	0.9	42.5
13	Pneumonitis due to solids and liquids (J69)	2,467	0.7	33.6
14	Aortic aneurysm and dissection (I71)	2,392	0.7	32.5
15	Atherosclerosis (I70)	2,342	0.7	31.9
...	All other causes (Residual)	49,030	13.6	667.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 85 years and over				
...	All causes	443,429	100.0	14,768.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	172,346	38.9	5,740.1
2	Cerebrovascular diseases (I60-I69)	49,164	11.1	1,637.4
3	Malignant neoplasms (C00-C97)	43,876	9.9	1,461.3
4	Alzheimer's disease (G30)	21,903	4.9	729.5
5	Influenza and pneumonia (J10-J18)	20,602	4.6	686.2
6	Chronic lower respiratory diseases (J40-J47)	15,620	3.5	520.2
7	Diabetes mellitus (E10-E14)	9,322	2.1	310.5
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,182	1.6	239.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,106	1.6	236.7
10	Septicemia (A40-A41)	6,408	1.4	213.4
11	Atherosclerosis (I70)	5,839	1.3	194.5
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	5,160	1.2	171.9
13	Pneumonitis due to solids and liquids (J69)	4,567	1.0	152.1
14	Parkinson's disease (G20-G21)	2,862	0.6	95.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,264	0.5	75.4
...	All other causes (Residual)	69,208	15.6	2,305.0
White, both sexes, all ages				
...	All causes	2,071,287	100.0	915.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	621,719	30.0	274.8
2	Malignant neoplasms (C00-C97)	480,011	23.2	212.2
3	Cerebrovascular diseases (I60-I69)	144,580	7.0	63.9
4	Chronic lower respiratory diseases (J40-J47)	112,840	5.4	49.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	82,592	4.0	36.5
6	Influenza and pneumonia (J10-J18)	57,914	2.8	25.6
7	Diabetes mellitus (E10-E14)	55,561	2.7	24.6
8	Alzheimer's disease (G30)	46,460	2.2	20.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	29,598	1.4	13.1
10	Intentional self-harm (suicide) (X60-X84,Y87.0)	26,475	1.3	11.7
11	Septicemia (A40-A41)	24,804	1.2	11.0
12	Chronic liver disease and cirrhosis (K70,K73-K74)	22,982	1.1	10.2
13	Parkinson's disease (G20-G21)	14,942	0.7	6.6
14	Pneumonitis due to solids and liquids (J69)	14,869	0.7	6.6
15	Aortic aneurysm and dissection (I71)	14,210	0.7	6.3
...	All other causes (Residual)	321,730	15.5	142.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 1-4 years				
...	All causes	3,494	100.0	29.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,335	38.2	11.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	347	9.9	2.9
3	Malignant neoplasms (C00-C97)	330	9.4	2.7
4	Assault (homicide) (X85-Y09,Y87.1)	199	5.7	1.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	126	3.6	1.0
6	Influenza and pneumonia (J10-J18)	69	2.0	0.6
7	Septicemia (A40-A41)	66	1.9	0.5
8	Certain conditions originating in the perinatal period (P00-P96)	46	1.3	0.4
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	44	1.3	0.4
10	Cerebrovascular diseases (I60-I69)	30	0.9	0.2
11	Chronic lower respiratory diseases (J40-J47)	28	0.8	0.2
12	Meningitis (G00,G03)	23	0.7	0.2
13	Anemias (D50-D64)	15	0.4	*
14	Meningococcal infection (A39)	14	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	12	0.3	*
...	All other causes (Residual)	810	23.2	6.7
White, both sexes, 5-14 years				
...	All causes	5,450	100.0	17.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,219	40.7	7.1
2	Malignant neoplasms (C00-C97)	818	15.0	2.6
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	292	5.4	0.9
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	254	4.7	0.8
5	Assault (homicide) (X85-Y09,Y87.1)	240	4.4	0.8
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	177	3.2	0.6
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	76	1.4	0.2
8	Influenza and pneumonia (J10-J18)	70	1.3	0.2
9	Chronic lower respiratory diseases (J40-J47)	63	1.2	0.2
10	Cerebrovascular diseases (I60-I69)	55	1.0	0.2
11	Septicemia (A40-A41)	49	0.9	0.2
12	Certain conditions originating in the perinatal period (P00-P96)	30	0.6	0.1
13	Complications of medical and surgical care (Y40-Y84,Y88)	23	0.4	0.1
14	Human immunodeficiency virus (HIV) disease (B20-B24)	21	0.4	0.1
14	Anemias (D50-D64)	21	0.4	0.1
14	Meningitis (G00,G03)	21	0.4	0.1
...	All other causes (Residual)	1,021	18.7	3.3

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 15-24 years				
...	All causes	23,018	100.0	75.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,823	51.4	38.8
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	3,315	14.4	10.9
3	Assault (homicide) (X85-Y09,Y87.1)	1,981	8.6	6.5
4	Malignant neoplasms (C00-C97)	1,333	5.8	4.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	656	2.8	2.2
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	332	1.4	1.1
7	Cerebrovascular diseases (I60-I69)	148	0.6	0.5
8	Influenza and pneumonia (J10-J18)	138	0.6	0.5
9	Diabetes mellitus (E10-E14)	105	0.5	0.3
10	Chronic lower respiratory diseases (J40-J47)	103	0.4	0.3
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	73	0.3	0.2
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	63	0.3	0.2
13	Septicemia (A40-A41)	60	0.3	0.2
14	Pregnancy, childbirth and the puerperium (O00-099)	55	0.2	0.2
15	Human immunodeficiency virus (HIV) disease (B20-B24)	53	0.2	0.2
...	All other causes (Residual)	2,780	12.1	9.1
White, both sexes, 25-34 years				
...	All causes	28,719	100.0	96.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,519	33.1	31.9
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	4,084	14.2	13.7
3	Malignant neoplasms (C00-C97)	3,072	10.7	10.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,937	6.7	6.5
5	Assault (homicide) (X85-Y09,Y87.1)	1,814	6.3	6.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	983	3.4	3.3
7	Diabetes mellitus (E10-E14)	420	1.5	1.4
7	Cerebrovascular diseases (I60-I69)	420	1.5	1.4
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	391	1.4	1.3
10	Chronic liver disease and cirrhosis (K70,K73-K74)	312	1.1	1.0
11	Influenza and pneumonia (J10-J18)	261	0.9	0.9
12	Chronic lower respiratory diseases (J40-J47)	166	0.6	0.6
13	Septicemia (A40-A41)	165	0.6	0.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	136	0.5	0.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	129	0.4	0.4
...	All other causes (Residual)	4,910	17.1	16.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 35-44 years				
...	All causes	66,496	100.0	180.3
1	Malignant neoplasms (C00-C97)	12,764	19.2	34.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,663	19.0	34.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,532	14.3	25.8
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	5,942	8.9	16.1
5	Human immunodeficiency virus (HIV) disease (B20-B24)	2,785	4.2	7.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	2,745	4.1	7.4
7	Assault (homicide) (X85-Y09,Y87.1)	1,725	2.6	4.7
8	Cerebrovascular diseases (I60-I69)	1,645	2.5	4.5
9	Diabetes mellitus (E10-E14)	1,419	2.1	3.8
10	Influenza and pneumonia (J10-J18)	767	1.2	2.1
11	Viral hepatitis (B15-B19)	625	0.9	1.7
12	Chronic lower respiratory diseases (J40-J47)	624	0.9	1.7
13	Septicemia (A40-A41)	578	0.9	1.6
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	459	0.7	1.2
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	412	0.6	1.1
...	All other causes (Residual)	11,811	17.8	32.0
White, both sexes, 45-54 years				
...	All causes	122,674	100.0	391.8
1	Malignant neoplasms (C00-C97)	38,478	31.4	122.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	26,991	22.0	86.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,901	8.1	31.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	5,515	4.5	17.6
5	Intentional self-harm (suicide) (X60-X84,Y87.0)	5,035	4.1	16.1
6	Cerebrovascular diseases (I60-I69)	3,914	3.2	12.5
7	Diabetes mellitus (E10-E14)	3,514	2.9	11.2
8	Chronic lower respiratory diseases (J40-J47)	2,584	2.1	8.3
9	Human immunodeficiency virus (HIV) disease (B20-B24)	1,856	1.5	5.9
10	Viral hepatitis (B15-B19)	1,462	1.2	4.7
11	Influenza and pneumonia (J10-J18)	1,297	1.1	4.1
12	Septicemia (A40-A41)	1,219	1.0	3.9
13	Assault (homicide) (X85-Y09,Y87.1)	1,048	0.9	3.3
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	936	0.8	3.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	558	0.5	1.8
...	All other causes (Residual)	18,366	15.0	58.7

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 55-64 years				
...	All causes	195,541	100.0	948.9
1	Malignant neoplasms (C00-C97)	74,759	38.2	362.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	50,581	25.9	245.4
3	Chronic lower respiratory diseases (J40-J47)	9,532	4.9	46.3
4	Cerebrovascular diseases (I60-I69)	7,239	3.7	35.1
5	Diabetes mellitus (E10-E14)	6,879	3.5	33.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,275	3.2	30.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	4,948	2.5	24.0
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	2,775	1.4	13.5
9	Influenza and pneumonia (J10-J18)	2,258	1.2	11.0
10	Septicemia (A40-A41)	2,066	1.1	10.0
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,009	1.0	9.7
12	Aortic aneurysm and dissection (I71)	1,257	0.6	6.1
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,058	0.5	5.1
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	851	0.4	4.1
15	Viral hepatitis (B15-B19)	665	0.3	3.2
...	All other causes (Residual)	22,389	11.4	108.6
White, both sexes, 65-74 years				
...	All causes	376,986	100.0	2,375.1
1	Malignant neoplasms (C00-C97)	130,570	34.6	822.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	103,449	27.4	651.8
3	Chronic lower respiratory diseases (J40-J47)	28,765	7.6	181.2
4	Cerebrovascular diseases (I60-I69)	19,224	5.1	121.1
5	Diabetes mellitus (E10-E14)	13,075	3.5	82.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,617	1.8	41.7
7	Influenza and pneumonia (J10-J18)	5,969	1.6	37.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,144	1.4	32.4
9	Chronic liver disease and cirrhosis (K70,K73-K74)	4,925	1.3	31.0
10	Septicemia (A40-A41)	4,418	1.2	27.8
11	Aortic aneurysm and dissection (I71)	3,651	1.0	23.0
12	Alzheimer's disease (G30)	3,139	0.8	19.8
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,239	0.6	14.1
14	Intentional self-harm (suicide) (X60-X84,Y87.0)	2,168	0.6	13.7
15	Parkinson's disease (G20-G21)	1,982	0.5	12.5
...	All other causes (Residual)	41,651	11.0	262.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 75-84 years				
...	All causes	627,729	100.0	5,652.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	196,833	31.4	1,772.4
2	Malignant neoplasms (C00-C97)	148,126	23.6	1,333.8
3	Cerebrovascular diseases (I60-I69)	50,367	8.0	453.5
4	Chronic lower respiratory diseases (J40-J47)	45,025	7.2	405.4
5	Diabetes mellitus (E10-E14)	18,478	2.9	166.4
6	Influenza and pneumonia (J10-J18)	17,811	2.8	160.4
7	Alzheimer's disease (G30)	16,124	2.6	145.2
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,740	1.7	96.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10,470	1.7	94.3
10	Septicemia (A40-A41)	8,223	1.3	74.0
11	Parkinson's disease (G20-G21)	7,306	1.2	65.8
12	Aortic aneurysm and dissection (I71)	5,398	0.9	48.6
13	Pneumonitis due to solids and liquids (J69)	4,948	0.8	44.6
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	4,498	0.7	40.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,302	0.7	38.7
...	All other causes (Residual)	79,080	12.6	712.1
White, both sexes, 85 years and over				
...	All causes	602,761	100.0	15,532.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	231,050	38.3	5,953.9
2	Malignant neoplasms (C00-C97)	69,679	11.6	1,795.6
3	Cerebrovascular diseases (I60-I69)	61,462	10.2	1,583.8
4	Influenza and pneumonia (J10-J18)	29,101	4.8	749.9
5	Alzheimer's disease (G30)	26,683	4.4	687.6
6	Chronic lower respiratory diseases (J40-J47)	25,926	4.3	668.1
7	Diabetes mellitus (E10-E14)	11,645	1.9	300.1
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,852	1.8	279.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10,311	1.7	265.7
10	Septicemia (A40-A41)	7,790	1.3	200.7
11	Atherosclerosis (I70)	7,491	1.2	193.0
12	Pneumonitis due to solids and liquids (J69)	7,331	1.2	188.9
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	5,880	1.0	151.5
14	Parkinson's disease (G20-G21)	5,354	0.9	138.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3,352	0.6	86.4
...	All other causes (Residual)	88,854	14.7	2,289.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, male, all ages				
...	All causes	1,007,191	100.0	905.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	301,551	29.9	271.2
2	Malignant neoplasms (C00-C97)	247,403	24.6	222.5
3	Cerebrovascular diseases (I60-I69)	54,938	5.5	49.4
4	Chronic lower respiratory diseases (J40-J47)	54,816	5.4	49.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	53,329	5.3	48.0
6	Diabetes mellitus (E10-E14)	26,009	2.6	23.4
7	Influenza and pneumonia (J10-J18)	25,002	2.5	22.5
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	21,293	2.1	19.1
9	Chronic liver disease and cirrhosis (K70,K73-K74)	15,002	1.5	13.5
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14,385	1.4	12.9
11	Alzheimer's disease (G30)	13,524	1.3	12.2
12	Septicemia (A40-A41)	10,716	1.1	9.6
13	Aortic aneurysm and dissection (I71)	8,797	0.9	7.9
14	Parkinson's disease (G20-G21)	8,367	0.8	7.5
15	Pneumonitis due to solids and liquids (J69)	7,488	0.7	6.7
...	All other causes (Residual)	144,571	14.4	130.0
White, male, 1-4 years				
...	All causes	2,004	100.0	32.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	808	40.3	13.1
2	Malignant neoplasms (C00-C97)	190	9.5	3.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	180	9.0	2.9
4	Assault (homicide) (X85-Y09,Y87.1)	114	5.7	1.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	80	4.0	1.3
6	Septicemia (A40-A41)	35	1.7	0.6
7	Influenza and pneumonia (J10-J18)	30	1.5	0.5
7	Certain conditions originating in the perinatal period (P00-P96)	30	1.5	0.5
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	25	1.2	0.4
10	Meningitis (G00,G03)	14	0.7	*
10	Cerebrovascular diseases (I60-I69)	14	0.7	*
12	Chronic lower respiratory diseases (J40-J47)	13	0.6	*
13	Meningococcal infection (A39)	8	0.4	*
13	Anemias (D50-D64)	8	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	8	0.4	*
...	All other causes (Residual)	447	22.3	7.3

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 5-14 years				
...	All causes	3,255	100.0	20.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,378	42.3	8.6
2	Malignant neoplasms (C00-C97)	456	14.0	2.9
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	200	6.1	1.3
4	Assault (homicide) (X85-Y09,Y87.1)	148	4.5	0.9
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	147	4.5	0.9
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	111	3.4	0.7
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	39	1.2	0.2
7	Influenza and pneumonia (J10-J18)	39	1.2	0.2
9	Chronic lower respiratory diseases (J40-J47)	33	1.0	0.2
10	Septicemia (A40-A41)	30	0.9	0.2
11	Cerebrovascular diseases (I60-I69)	26	0.8	0.2
12	Certain conditions originating in the perinatal period (P00-P96)	20	0.6	0.1
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	0.4	*
14	Anemias (D50-D64)	12	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	11	0.3	*
...	All other causes (Residual)	591	18.2	3.7
White, male, 15-24 years				
...	All causes	16,868	100.0	107.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,770	52.0	56.0
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	2,853	16.9	18.2
3	Assault (homicide) (X85-Y09,Y87.1)	1,583	9.4	10.1
4	Malignant neoplasms (C00-C97)	798	4.7	5.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	413	2.4	2.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	201	1.2	1.3
7	Cerebrovascular diseases (I60-I69)	84	0.5	0.5
8	Influenza and pneumonia (J10-J18)	79	0.5	0.5
9	Chronic lower respiratory diseases (J40-J47)	65	0.4	0.4
10	Diabetes mellitus (E10-E14)	64	0.4	0.4
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	48	0.3	0.3
12	Legal intervention (Y35,Y89.0)	42	0.2	0.3
13	Septicemia (A40-A41)	39	0.2	0.2
14	Human immunodeficiency virus (HIV) disease (B20-B24)	38	0.2	0.2
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	33	0.2	0.2
...	All other causes (Residual)	1,758	10.4	11.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 25-34 years				
...	All causes	20,140	100.0	134.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,369	36.6	49.2
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	3,363	16.7	22.4
3	Malignant neoplasms (C00-C97)	1,500	7.4	10.0
4	Assault (homicide) (X85-Y09,Y87.1)	1,356	6.7	9.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,316	6.5	8.8
6	Human immunodeficiency virus (HIV) disease (B20-B24)	749	3.7	5.0
7	Diabetes mellitus (E10-E14)	267	1.3	1.8
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	229	1.1	1.5
9	Cerebrovascular diseases (I60-I69)	225	1.1	1.5
10	Chronic liver disease and cirrhosis (K70,K73-K74)	224	1.1	1.5
11	Influenza and pneumonia (J10-J18)	138	0.7	0.9
12	Septicemia (A40-A41)	95	0.5	0.6
13	Chronic lower respiratory diseases (J40-J47)	84	0.4	0.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	73	0.4	0.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	70	0.3	0.5
15	Legal intervention (Y35,Y89.0)	70	0.3	0.5
...	All other causes (Residual)	3,012	15.0	20.1
White, male, 35-44 years				
...	All causes	43,378	100.0	234.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,462	21.8	51.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,020	16.2	37.9
3	Malignant neoplasms (C00-C97)	5,739	13.2	31.0
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	4,620	10.7	25.0
5	Human immunodeficiency virus (HIV) disease (B20-B24)	2,300	5.3	12.4
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1,921	4.4	10.4
7	Assault (homicide) (X85-Y09,Y87.1)	1,202	2.8	6.5
8	Diabetes mellitus (E10-E14)	866	2.0	4.7
9	Cerebrovascular diseases (I60-I69)	842	1.9	4.5
10	Influenza and pneumonia (J10-J18)	455	1.0	2.5
11	Viral hepatitis (B15-B19)	439	1.0	2.4
12	Septicemia (A40-A41)	311	0.7	1.7
13	Chronic lower respiratory diseases (J40-J47)	292	0.7	1.6
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	259	0.6	1.4
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	243	0.6	1.3
...	All other causes (Residual)	7,407	17.1	40.0

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 45-54 years				
...	All causes	77,866	100.0	502.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,480	26.3	132.2
2	Malignant neoplasms (C00-C97)	19,353	24.9	125.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,322	9.4	47.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	4,205	5.4	27.1
5	Intentional self-harm (suicide) (X60-X84,Y87.0)	3,834	4.9	24.8
6	Cerebrovascular diseases (I60-I69)	2,138	2.7	13.8
7	Diabetes mellitus (E10-E14)	2,077	2.7	13.4
8	Human immunodeficiency virus (HIV) disease (B20-B24)	1,590	2.0	10.3
9	Chronic lower respiratory diseases (J40-J47)	1,321	1.7	8.5
10	Viral hepatitis (B15-B19)	1,123	1.4	7.3
11	Influenza and pneumonia (J10-J18)	759	1.0	4.9
12	Assault (homicide) (X85-Y09,Y87.1)	718	0.9	4.6
13	Septicemia (A40-A41)	693	0.9	4.5
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	558	0.7	3.6
15	Aortic aneurysm and dissection (I71)	384	0.5	2.5
...	All other causes (Residual)	11,311	14.5	73.0
White, male, 55-64 years				
...	All causes	117,114	100.0	1,177.7
1	Malignant neoplasms (C00-C97)	40,457	34.5	406.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	35,422	30.2	356.2
3	Chronic lower respiratory diseases (J40-J47)	4,893	4.2	49.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,242	3.6	42.7
5	Cerebrovascular diseases (I60-I69)	3,998	3.4	40.2
6	Diabetes mellitus (E10-E14)	3,823	3.3	38.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	3,475	3.0	34.9
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	2,133	1.8	21.5
9	Influenza and pneumonia (J10-J18)	1,302	1.1	13.1
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,106	0.9	11.1
11	Septicemia (A40-A41)	1,097	0.9	11.0
12	Aortic aneurysm and dissection (I71)	956	0.8	9.6
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	564	0.5	5.7
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	503	0.4	5.1
15	Human immunodeficiency virus (HIV) disease (B20-B24)	499	0.4	5.0
...	All other causes (Residual)	12,644	10.8	127.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 65-74 years				
...	All causes	213,379	100.0	2,949.8
1	Malignant neoplasms (C00-C97)	72,284	33.9	999.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	64,464	30.2	891.2
3	Chronic lower respiratory diseases (J40-J47)	14,786	6.9	204.4
4	Cerebrovascular diseases (I60-I69)	9,828	4.6	135.9
5	Diabetes mellitus (E10-E14)	6,739	3.2	93.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,977	1.9	55.0
7	Influenza and pneumonia (J10-J18)	3,314	1.6	45.8
8	Chronic liver disease and cirrhosis (K70,K73-K74)	3,006	1.4	41.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,769	1.3	38.3
10	Aortic aneurysm and dissection (I71)	2,533	1.2	35.0
11	Septicemia (A40-A41)	2,264	1.1	31.3
12	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,784	0.8	24.7
13	Alzheimer's disease (G30)	1,432	0.7	19.8
14	Parkinson's disease (G20-G21)	1,262	0.6	17.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,247	0.6	17.2
...	All other causes (Residual)	21,690	10.2	299.8
White, male, 75-84 years				
...	All causes	306,370	100.0	6,817.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	99,295	32.4	2,209.6
2	Malignant neoplasms (C00-C97)	76,714	25.0	1,707.1
3	Chronic lower respiratory diseases (J40-J47)	22,254	7.3	495.2
4	Cerebrovascular diseases (I60-I69)	21,210	6.9	472.0
5	Influenza and pneumonia (J10-J18)	8,742	2.9	194.5
6	Diabetes mellitus (E10-E14)	8,419	2.7	187.4
7	Alzheimer's disease (G30)	5,854	1.9	130.3
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,493	1.8	122.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,353	1.7	119.1
10	Parkinson's disease (G20-G21)	4,307	1.4	95.8
11	Septicemia (A40-A41)	3,737	1.2	83.2
12	Aortic aneurysm and dissection (I71)	3,227	1.1	71.8
13	Pneumonitis due to solids and liquids (J69)	2,746	0.9	61.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,168	0.7	48.2
15	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,817	0.6	40.4
...	All other causes (Residual)	35,034	11.4	779.6

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 85 years and over				
...	All causes	196,409	100.0	16,897.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	72,735	37.0	6,257.6
2	Malignant neoplasms (C00-C97)	29,863	15.2	2,569.2
3	Cerebrovascular diseases (I60-I69)	16,526	8.4	1,421.8
4	Chronic lower respiratory diseases (J40-J47)	11,058	5.6	951.4
5	Influenza and pneumonia (J10-J18)	10,045	5.1	864.2
6	Alzheimer's disease (G30)	6,010	3.1	517.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,171	2.1	358.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,100	2.1	352.7
9	Diabetes mellitus (E10-E14)	3,740	1.9	321.8
10	Pneumonitis due to solids and liquids (J69)	3,149	1.6	270.9
11	Parkinson's disease (G20-G21)	2,609	1.3	224.5
12	Septicemia (A40-A41)	2,326	1.2	200.1
13	Atherosclerosis (I70)	2,011	1.0	173.0
14	Aortic aneurysm and dissection (I71)	1,472	0.7	126.6
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,469	0.7	126.4
...	All other causes (Residual)	25,125	12.8	2,161.6
White, female, all ages				
...	All causes	1,064,096	100.0	924.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	320,168	30.1	278.3
2	Malignant neoplasms (C00-C97)	232,608	21.9	202.2
3	Cerebrovascular diseases (I60-I69)	89,642	8.4	77.9
4	Chronic lower respiratory diseases (J40-J47)	58,024	5.5	50.4
5	Alzheimer's disease (G30)	32,936	3.1	28.6
6	Influenza and pneumonia (J10-J18)	32,912	3.1	28.6
7	Diabetes mellitus (E10-E14)	29,552	2.8	25.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	29,263	2.8	25.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15,213	1.4	13.2
10	Septicemia (A40-A41)	14,088	1.3	12.2
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	8,655	0.8	7.5
12	Atherosclerosis (I70)	8,307	0.8	7.2
13	Chronic liver disease and cirrhosis (K70,K73-K74)	7,980	0.7	6.9
14	Pneumonitis due to solids and liquids (J69)	7,381	0.7	6.4
15	Parkinson's disease (G20-G21)	6,575	0.6	5.7
...	All other causes (Residual)	170,792	16.1	148.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 1-4 years				
...	All causes	1,490	100.0	25.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	527	35.4	9.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	167	11.2	2.8
3	Malignant neoplasms (C00-C97)	140	9.4	2.4
4	Assault (homicide) (X85-Y09,Y87.1)	85	5.7	1.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	46	3.1	0.8
6	Influenza and pneumonia (J10-J18)	39	2.6	0.7
7	Septicemia (A40-A41)	31	2.1	0.5
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	1.3	*
9	Cerebrovascular diseases (I60-I69)	16	1.1	*
9	Certain conditions originating in the perinatal period (P00-P96)	16	1.1	*
11	Chronic lower respiratory diseases (J40-J47)	15	1.0	*
12	Meningitis (G00,G03)	9	0.6	*
13	Anemias (D50-D64)	7	0.5	*
14	Meningococcal infection (A39)	6	0.4	*
15	Pneumonitis due to solids and liquids (J69)	5	0.3	*
...	All other causes (Residual)	362	24.3	6.2
White, female, 5-14 years				
...	All causes	2,195	100.0	14.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	841	38.3	5.5
2	Malignant neoplasms (C00-C97)	362	16.5	2.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	145	6.6	1.0
4	Assault (homicide) (X85-Y09,Y87.1)	92	4.2	0.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	66	3.0	0.4
6	Intentional self-harm (suicide) (X60-X84,Y87.0)	54	2.5	0.4
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	37	1.7	0.2
8	Influenza and pneumonia (J10-J18)	31	1.4	0.2
9	Chronic lower respiratory diseases (J40-J47)	30	1.4	0.2
10	Cerebrovascular diseases (I60-I69)	29	1.3	0.2
11	Septicemia (A40-A41)	19	0.9	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	12	0.5	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	12	0.5	*
14	Meningitis (G00,G03)	11	0.5	*
15	Certain conditions originating in the perinatal period (P00-P96)	10	0.5	*
...	All other causes (Residual)	444	20.2	2.9

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 15-24 years				
...	All causes	6,150	100.0	41.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,053	49.6	20.6
2	Malignant neoplasms (C00-C97)	535	8.7	3.6
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	462	7.5	3.1
4	Assault (homicide) (X85-Y09,Y87.1)	398	6.5	2.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	243	4.0	1.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	131	2.1	0.9
7	Cerebrovascular diseases (I60-I69)	64	1.0	0.4
8	Influenza and pneumonia (J10-J18)	59	1.0	0.4
9	Pregnancy, childbirth and the puerperium (O00-099)	55	0.9	0.4
10	Diabetes mellitus (E10-E14)	41	0.7	0.3
11	Chronic lower respiratory diseases (J40-J47)	38	0.6	0.3
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	30	0.5	0.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	25	0.4	0.2
14	Septicemia (A40-A41)	21	0.3	0.1
15	Meningococcal infection (A39)	17	0.3	*
...	All other causes (Residual)	978	15.9	6.6
White, female, 25-34 years				
...	All causes	8,579	100.0	57.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,150	25.1	14.4
2	Malignant neoplasms (C00-C97)	1,572	18.3	10.6
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	721	8.4	4.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	621	7.2	4.2
5	Assault (homicide) (X85-Y09,Y87.1)	458	5.3	3.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	234	2.7	1.6
7	Cerebrovascular diseases (I60-I69)	195	2.3	1.3
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	162	1.9	1.1
9	Diabetes mellitus (E10-E14)	153	1.8	1.0
10	Influenza and pneumonia (J10-J18)	123	1.4	0.8
11	Pregnancy, childbirth and the puerperium (O00-099)	108	1.3	0.7
12	Chronic liver disease and cirrhosis (K70,K73-K74)	88	1.0	0.6
13	Chronic lower respiratory diseases (J40-J47)	82	1.0	0.6
14	Septicemia (A40-A41)	70	0.8	0.5
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	63	0.7	0.4
...	All other causes (Residual)	1,779	20.7	12.0

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 35-44 years				
...	All causes	23,118	100.0	125.8
1	Malignant neoplasms (C00-C97)	7,025	30.4	38.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,201	13.8	17.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,512	10.9	13.7
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,322	5.7	7.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	824	3.6	4.5
6	Cerebrovascular diseases (I60-I69)	803	3.5	4.4
7	Diabetes mellitus (E10-E14)	553	2.4	3.0
8	Assault (homicide) (X85-Y09,Y87.1)	523	2.3	2.8
9	Human immunodeficiency virus (HIV) disease (B20-B24)	485	2.1	2.6
10	Chronic lower respiratory diseases (J40-J47)	332	1.4	1.8
11	Influenza and pneumonia (J10-J18)	312	1.3	1.7
12	Septicemia (A40-A41)	267	1.2	1.5
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	200	0.9	1.1
14	Viral hepatitis (B15-B19)	186	0.8	1.0
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	169	0.7	0.9
...	All other causes (Residual)	4,404	19.1	24.0
White, female, 45-54 years				
...	All causes	44,808	100.0	283.2
1	Malignant neoplasms (C00-C97)	19,125	42.7	120.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,511	14.5	41.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,579	5.8	16.3
4	Cerebrovascular diseases (I60-I69)	1,776	4.0	11.2
5	Diabetes mellitus (E10-E14)	1,437	3.2	9.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1,310	2.9	8.3
7	Chronic lower respiratory diseases (J40-J47)	1,263	2.8	8.0
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,201	2.7	7.6
9	Influenza and pneumonia (J10-J18)	538	1.2	3.4
10	Septicemia (A40-A41)	526	1.2	3.3
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	378	0.8	2.4
12	Viral hepatitis (B15-B19)	339	0.8	2.1
13	Assault (homicide) (X85-Y09,Y87.1)	330	0.7	2.1
14	Human immunodeficiency virus (HIV) disease (B20-B24)	266	0.6	1.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	239	0.5	1.5
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	239	0.5	1.5
...	All other causes (Residual)	6,751	15.1	42.7

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 55-64 years				
...	All causes	78,427	100.0	735.5
1	Malignant neoplasms (C00-C97)	34,302	43.7	321.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,159	19.3	142.2
3	Chronic lower respiratory diseases (J40-J47)	4,639	5.9	43.5
4	Cerebrovascular diseases (I60-I69)	3,241	4.1	30.4
5	Diabetes mellitus (E10-E14)	3,056	3.9	28.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,033	2.6	19.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,473	1.9	13.8
8	Septicemia (A40-A41)	969	1.2	9.1
9	Influenza and pneumonia (J10-J18)	956	1.2	9.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	903	1.2	8.5
11	Intentional self-harm (suicide) (X60-X84,Y87.0)	642	0.8	6.0
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	494	0.6	4.6
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	348	0.4	3.3
14	Aortic aneurysm and dissection (I71)	301	0.4	2.8
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	259	0.3	2.4
...	All other causes (Residual)	9,652	12.3	90.5
White, female, 65-74 years				
...	All causes	163,607	100.0	1,893.9
1	Malignant neoplasms (C00-C97)	58,286	35.6	674.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	38,985	23.8	451.3
3	Chronic lower respiratory diseases (J40-J47)	13,979	8.5	161.8
4	Cerebrovascular diseases (I60-I69)	9,396	5.7	108.8
5	Diabetes mellitus (E10-E14)	6,336	3.9	73.3
6	Influenza and pneumonia (J10-J18)	2,655	1.6	30.7
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,640	1.6	30.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,375	1.5	27.5
9	Septicemia (A40-A41)	2,154	1.3	24.9
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,919	1.2	22.2
11	Alzheimer's disease (G30)	1,707	1.0	19.8
12	Aortic aneurysm and dissection (I71)	1,118	0.7	12.9
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	992	0.6	11.5
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	980	0.6	11.3
15	Parkinson's disease (G20-G21)	720	0.4	8.3
...	All other causes (Residual)	19,365	11.8	224.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 75-84 years				
...	All causes	321,359	100.0	4,860.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	97,538	30.4	1,475.2
2	Malignant neoplasms (C00-C97)	71,412	22.2	1,080.1
3	Cerebrovascular diseases (I60-I69)	29,157	9.1	441.0
4	Chronic lower respiratory diseases (J40-J47)	22,771	7.1	344.4
5	Alzheimer's disease (G30)	10,270	3.2	155.3
6	Diabetes mellitus (E10-E14)	10,059	3.1	152.1
7	Influenza and pneumonia (J10-J18)	9,069	2.8	137.2
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,247	1.6	79.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,117	1.6	77.4
10	Septicemia (A40-A41)	4,486	1.4	67.8
11	Parkinson's disease (G20-G21)	2,999	0.9	45.4
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,715	0.8	41.1
13	Pneumonitis due to solids and liquids (J69)	2,202	0.7	33.3
14	Aortic aneurysm and dissection (I71)	2,171	0.7	32.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,134	0.7	32.3
...	All other causes (Residual)	44,012	13.7	665.7
White, female, 85 years and over				
...	All causes	406,352	100.0	14,948.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	158,315	39.0	5,824.0
2	Cerebrovascular diseases (I60-I69)	44,936	11.1	1,653.1
3	Malignant neoplasms (C00-C97)	39,816	9.8	1,464.7
4	Alzheimer's disease (G30)	20,673	5.1	760.5
5	Influenza and pneumonia (J10-J18)	19,056	4.7	701.0
6	Chronic lower respiratory diseases (J40-J47)	14,868	3.7	547.0
7	Diabetes mellitus (E10-E14)	7,905	1.9	290.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,752	1.7	248.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,140	1.5	225.9
10	Atherosclerosis (I70)	5,480	1.3	201.6
11	Septicemia (A40-A41)	5,464	1.3	201.0
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	4,411	1.1	162.3
13	Pneumonitis due to solids and liquids (J69)	4,182	1.0	153.8
14	Parkinson's disease (G20-G21)	2,745	0.7	101.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,104	0.5	77.4
...	All other causes (Residual)	63,505	15.6	2,336.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, all ages				
...	All causes	285,826	100.0	809.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	77,523	27.1	219.6
2	Malignant neoplasms (C00-C97)	61,945	21.7	175.5
3	Cerebrovascular diseases (I60-I69)	19,221	6.7	54.4
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,277	4.3	34.8
5	Diabetes mellitus (E10-E14)	12,021	4.2	34.1
6	Assault (homicide) (X85-Y09,Y87.1)	7,867	2.8	22.3
7	Human immunodeficiency virus (HIV) disease (B20-B24)	7,848	2.7	22.2
8	Chronic lower respiratory diseases (J40-J47)	7,607	2.7	21.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,911	2.4	19.6
10	Influenza and pneumonia (J10-J18)	5,990	2.1	17.0
11	Septicemia (A40-A41)	5,928	2.1	16.8
12	Certain conditions originating in the perinatal period (P00-P96)	5,021	1.8	14.2
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	3,887	1.4	11.0
14	Chronic liver disease and cirrhosis (K70,K73-K74)	2,737	1.0	7.8
15	Alzheimer's disease (G30)	2,728	1.0	7.7
...	All other causes (Residual)	46,315	16.2	131.2
Black, both sexes, 1-4 years				
...	All causes	1,248	100.0	56.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	405	32.5	18.2
2	Assault (homicide) (X85-Y09,Y87.1)	145	11.6	6.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	126	10.1	5.7
4	Malignant neoplasms (C00-C97)	70	5.6	3.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	49	3.9	2.2
6	Influenza and pneumonia (J10-J18)	30	2.4	1.3
7	Certain conditions originating in the perinatal period (P00-P96)	27	2.2	1.2
8	Septicemia (A40-A41)	26	2.1	1.2
9	Chronic lower respiratory diseases (J40-J47)	21	1.7	0.9
10	Anemias (D50-D64)	19	1.5	*
11	Cerebrovascular diseases (I60-I69)	14	1.1	*
12	Meningitis (G00,G03)	13	1.0	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	12	1.0	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.7	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	8	0.6	*
...	All other causes (Residual)	274	22.0	12.3

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 5-14 years				
...	All causes	1,643	100.0	26.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	631	38.4	10.1
2	Malignant neoplasms (C00-C97)	150	9.1	2.4
3	Assault (homicide) (X85-Y09,Y87.1)	124	7.5	2.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	91	5.5	1.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	76	4.6	1.2
6	Chronic lower respiratory diseases (J40-J47)	70	4.3	1.1
7	Intentional self-harm (suicide) (X60-X84,Y87.0)	40	2.4	0.6
8	Anemias (D50-D64)	31	1.9	0.5
9	Human immunodeficiency virus (HIV) disease (B20-B24)	23	1.4	0.4
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	1.2	*
11	Cerebrovascular diseases (I60-I69)	15	0.9	*
12	Septicemia (A40-A41)	11	0.7	*
12	Influenza and pneumonia (J10-J18)	11	0.7	*
14	Certain conditions originating in the perinatal period (P00-P96)	10	0.6	*
15	Diabetes mellitus (E10-E14)	9	0.5	*
15	Meningitis (G00,G03)	9	0.5	*
...	All other causes (Residual)	323	19.7	5.2
Black, both sexes, 15-24 years				
...	All causes	7,049	100.0	120.8
1	Assault (homicide) (X85-Y09,Y87.1)	2,819	40.0	48.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,695	24.0	29.0
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	480	6.8	8.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	333	4.7	5.7
5	Malignant neoplasms (C00-C97)	301	4.3	5.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	125	1.8	2.1
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	91	1.3	1.6
8	Chronic lower respiratory diseases (J40-J47)	81	1.1	1.4
9	Anemias (D50-D64)	62	0.9	1.1
10	Diabetes mellitus (E10-E14)	55	0.8	0.9
11	Influenza and pneumonia (J10-J18)	44	0.6	0.8
12	Cerebrovascular diseases (I60-I69)	43	0.6	0.7
13	Pregnancy, childbirth and the puerperium (O00-099)	36	0.5	0.6
14	Septicemia (A40-A41)	33	0.5	0.6
15	Legal intervention (Y35,Y89.0)	28	0.4	0.5
...	All other causes (Residual)	823	11.7	14.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 25-34 years				
...	All causes	10,239	100.0	195.5
1	Assault (homicide) (X85-Y09,Y87.1)	2,224	21.7	42.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,770	17.3	33.8
3	Human immunodeficiency virus (HIV) disease (B20-B24)	1,432	14.0	27.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	937	9.2	17.9
5	Malignant neoplasms (C00-C97)	649	6.3	12.4
6	Intentional self-harm (suicide) (X60-X84,Y87.0)	476	4.6	9.1
7	Diabetes mellitus (E10-E14)	189	1.8	3.6
8	Cerebrovascular diseases (I60-I69)	160	1.6	3.1
9	Anemias (D50-D64)	124	1.2	2.4
10	Chronic lower respiratory diseases (J40-J47)	116	1.1	2.2
11	Septicemia (A40-A41)	108	1.1	2.1
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	100	1.0	1.9
13	Influenza and pneumonia (J10-J18)	93	0.9	1.8
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	77	0.8	1.5
15	Pregnancy, childbirth and the puerperium (O00-099)	65	0.6	1.2
...	All other causes (Residual)	1,719	16.8	32.8
Black, both sexes, 35-44 years				
...	All causes	20,557	100.0	360.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,305	16.1	57.9
2	Malignant neoplasms (C00-C97)	3,108	15.1	54.5
3	Human immunodeficiency virus (HIV) disease (B20-B24)	3,080	15.0	54.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,287	11.1	40.1
5	Assault (homicide) (X85-Y09,Y87.1)	1,376	6.7	24.1
6	Cerebrovascular diseases (I60-I69)	836	4.1	14.7
7	Diabetes mellitus (E10-E14)	462	2.2	8.1
8	Chronic liver disease and cirrhosis (K70,K73-K74)	451	2.2	7.9
9	Intentional self-harm (suicide) (X60-X84,Y87.0)	438	2.1	7.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	306	1.5	5.4
11	Chronic lower respiratory diseases (J40-J47)	275	1.3	4.8
12	Septicemia (A40-A41)	272	1.3	4.8
13	Influenza and pneumonia (J10-J18)	257	1.3	4.5
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	179	0.9	3.1
15	Anemias (D50-D64)	141	0.7	2.5
...	All other causes (Residual)	3,784	18.4	66.3

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 45-54 years				
...	All causes	33,289	100.0	806.1
1	Malignant neoplasms (C00-C97)	8,147	24.5	197.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,680	23.1	186.0
3	Human immunodeficiency virus (HIV) disease (B20-B24)	2,257	6.8	54.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,988	6.0	48.1
5	Cerebrovascular diseases (I60-I69)	1,836	5.5	44.5
6	Diabetes mellitus (E10-E14)	1,279	3.8	31.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	937	2.8	22.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	669	2.0	16.2
9	Assault (homicide) (X85-Y09,Y87.1)	628	1.9	15.2
10	Chronic lower respiratory diseases (J40-J47)	599	1.8	14.5
11	Septicemia (A40-A41)	575	1.7	13.9
12	Influenza and pneumonia (J10-J18)	425	1.3	10.3
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	409	1.2	9.9
14	Viral hepatitis (B15-B19)	336	1.0	8.1
15	Intentional self-harm (suicide) (X60-X84,Y87.0)	275	0.8	6.7
...	All other causes (Residual)	5,249	15.8	127.1
Black, both sexes, 55-64 years				
...	All causes	39,281	100.0	1,628.6
1	Malignant neoplasms (C00-C97)	12,119	30.9	502.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,385	29.0	472.0
3	Cerebrovascular diseases (I60-I69)	2,291	5.8	95.0
4	Diabetes mellitus (E10-E14)	2,015	5.1	83.5
5	Chronic lower respiratory diseases (J40-J47)	1,073	2.7	44.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	997	2.5	41.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	991	2.5	41.1
8	Septicemia (A40-A41)	768	2.0	31.8
9	Human immunodeficiency virus (HIV) disease (B20-B24)	636	1.6	26.4
10	Chronic liver disease and cirrhosis (K70,K73-K74)	633	1.6	26.2
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	559	1.4	23.2
12	Influenza and pneumonia (J10-J18)	538	1.4	22.3
13	Assault (homicide) (X85-Y09,Y87.1)	204	0.5	8.5
14	Aortic aneurysm and dissection (I71)	200	0.5	8.3
15	Viral hepatitis (B15-B19)	143	0.4	5.9
...	All other causes (Residual)	4,729	12.0	196.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 65-74 years				
...	All causes	55,195	100.0	3,251.5
1	Malignant neoplasms (C00-C97)	16,636	30.1	980.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	16,590	30.1	977.3
3	Cerebrovascular diseases (I60-I69)	3,689	6.7	217.3
4	Diabetes mellitus (E10-E14)	3,118	5.6	183.7
5	Chronic lower respiratory diseases (J40-J47)	2,014	3.6	118.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,653	3.0	97.4
7	Septicemia (A40-A41)	1,189	2.2	70.0
8	Influenza and pneumonia (J10-J18)	997	1.8	58.7
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	868	1.6	51.1
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	806	1.5	47.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	434	0.8	25.6
12	Pneumonitis due to solids and liquids (J69)	290	0.5	17.1
13	Aortic aneurysm and dissection (I71)	281	0.5	16.6
14	Alzheimer's disease (G30)	261	0.5	15.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	248	0.4	14.6
...	All other causes (Residual)	6,121	11.1	360.6
Black, both sexes, 75-84 years				
...	All causes	61,444	100.0	6,826.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	19,802	32.2	2,200.0
2	Malignant neoplasms (C00-C97)	14,361	23.4	1,595.5
3	Cerebrovascular diseases (I60-I69)	5,399	8.8	599.8
4	Diabetes mellitus (E10-E14)	3,215	5.2	357.2
5	Chronic lower respiratory diseases (J40-J47)	2,152	3.5	239.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,789	2.9	198.8
7	Septicemia (A40-A41)	1,590	2.6	176.6
8	Influenza and pneumonia (J10-J18)	1,569	2.6	174.3
9	Alzheimer's disease (G30)	1,008	1.6	112.0
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	982	1.6	109.1
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	778	1.3	86.4
12	Pneumonitis due to solids and liquids (J69)	472	0.8	52.4
13	Aortic aneurysm and dissection (I71)	333	0.5	37.0
14	Atherosclerosis (I70)	323	0.5	35.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	284	0.5	31.6
...	All other causes (Residual)	7,387	12.0	820.7

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 85 years and over				
...	All causes	47,038	100.0	14,752.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	17,229	36.6	5,403.4
2	Malignant neoplasms (C00-C97)	6,390	13.6	2,004.0
3	Cerebrovascular diseases (I60-I69)	4,891	10.4	1,533.9
4	Influenza and pneumonia (J10-J18)	1,921	4.1	602.5
5	Diabetes mellitus (E10-E14)	1,678	3.6	526.3
6	Alzheimer's disease (G30)	1,423	3.0	446.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,299	2.8	407.4
8	Septicemia (A40-A41)	1,255	2.7	393.6
9	Chronic lower respiratory diseases (J40-J47)	1,195	2.5	374.8
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	906	1.9	284.1
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	579	1.2	181.6
12	Pneumonitis due to solids and liquids (J69)	511	1.1	160.3
13	Atherosclerosis (I70)	442	0.9	138.6
14	Nutritional deficiencies (E40-E64)	214	0.5	67.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	205	0.4	64.3
...	All other causes (Residual)	6,900	14.7	2,164.0
Black, male, all ages				
...	All causes	145,184	100.0	865.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	36,740	25.3	219.0
2	Malignant neoplasms (C00-C97)	32,817	22.6	195.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,531	5.9	50.9
4	Cerebrovascular diseases (I60-I69)	8,026	5.5	47.8
5	Assault (homicide) (X85-Y09,Y87.1)	6,482	4.5	38.6
6	Human immunodeficiency virus (HIV) disease (B20-B24)	5,400	3.7	32.2
7	Diabetes mellitus (E10-E14)	4,771	3.3	28.4
8	Chronic lower respiratory diseases (J40-J47)	4,238	2.9	25.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,074	2.1	18.3
10	Influenza and pneumonia (J10-J18)	2,915	2.0	17.4
11	Certain conditions originating in the perinatal period (P00-P96)	2,844	2.0	17.0
12	Septicemia (A40-A41)	2,587	1.8	15.4
13	Chronic liver disease and cirrhosis (K70,K73-K74)	1,730	1.2	10.3
14	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,636	1.1	9.8
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,592	1.1	9.5
...	All other causes (Residual)	21,801	15.0	130.0

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 1-4 years				
...	All causes	692	100.0	61.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	239	34.5	21.1
2	Assault (homicide) (X85-Y09,Y87.1)	75	10.8	6.6
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	70	10.1	6.2
4	Malignant neoplasms (C00-C97)	34	4.9	3.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	24	3.5	2.1
6	Influenza and pneumonia (J10-J18)	21	3.0	1.9
7	Certain conditions originating in the perinatal period (P00-P96)	18	2.6	*
8	Chronic lower respiratory diseases (J40-J47)	14	2.0	*
9	Septicemia (A40-A41)	13	1.9	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	1.2	*
10	Anemias (D50-D64)	8	1.2	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	6	0.9	*
12	Meningitis (G00,G03)	6	0.9	*
14	Meningococcal infection (A39)	4	0.6	*
15	Cerebrovascular diseases (I60-I69)	3	0.4	*
...	All other causes (Residual)	149	21.5	13.2
Black, male, 5-14 years				
...	All causes	974	100.0	30.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	414	42.5	13.0
2	Malignant neoplasms (C00-C97)	87	8.9	2.7
3	Assault (homicide) (X85-Y09,Y87.1)	72	7.4	2.3
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	52	5.3	1.6
5	Chronic lower respiratory diseases (J40-J47)	43	4.4	1.4
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	42	4.3	1.3
7	Intentional self-harm (suicide) (X60-X84,Y87.0)	35	3.6	1.1
8	Anemias (D50-D64)	17	1.7	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	12	1.2	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.8	*
11	Influenza and pneumonia (J10-J18)	7	0.7	*
12	Septicemia (A40-A41)	6	0.6	*
12	Meningitis (G00,G03)	6	0.6	*
14	Cerebrovascular diseases (I60-I69)	5	0.5	*
15	Diabetes mellitus (E10-E14)	4	0.4	*
...	All other causes (Residual)	164	16.8	5.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 15-24 years				
...	All causes	5,318	100.0	181.4
1	Assault (homicide) (X85-Y09,Y87.1)	2,501	47.0	85.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,261	23.7	43.0
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	415	7.8	14.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	193	3.6	6.6
5	Malignant neoplasms (C00-C97)	179	3.4	6.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	60	1.1	2.0
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	58	1.1	2.0
8	Chronic lower respiratory diseases (J40-J47)	51	1.0	1.7
9	Anemias (D50-D64)	30	0.6	1.0
10	Diabetes mellitus (E10-E14)	28	0.5	1.0
10	Legal intervention (Y35,Y89.0)	28	0.5	1.0
12	Influenza and pneumonia (J10-J18)	21	0.4	0.7
13	Cerebrovascular diseases (I60-I69)	19	0.4	*
14	Septicemia (A40-A41)	18	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	0.2	*
...	All other causes (Residual)	446	8.4	15.2
Black, male, 25-34 years				
...	All causes	6,750	100.0	271.8
1	Assault (homicide) (X85-Y09,Y87.1)	1,911	28.3	76.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,340	19.9	54.0
3	Human immunodeficiency virus (HIV) disease (B20-B24)	835	12.4	33.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	547	8.1	22.0
5	Intentional self-harm (suicide) (X60-X84,Y87.0)	402	6.0	16.2
6	Malignant neoplasms (C00-C97)	262	3.9	10.5
7	Diabetes mellitus (E10-E14)	99	1.5	4.0
8	Chronic lower respiratory diseases (J40-J47)	72	1.1	2.9
9	Cerebrovascular diseases (I60-I69)	71	1.1	2.9
10	Septicemia (A40-A41)	59	0.9	2.4
10	Influenza and pneumonia (J10-J18)	59	0.9	2.4
12	Anemias (D50-D64)	55	0.8	2.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	50	0.7	2.0
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	46	0.7	1.9
15	Legal intervention (Y35,Y89.0)	33	0.5	1.3
...	All other causes (Residual)	909	13.5	36.6

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 35-44 years				
...	All causes	12,251	100.0	456.7
1	Human immunodeficiency virus (HIV) disease (B20-B24)	2,096	17.1	78.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,009	16.4	74.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,678	13.7	62.5
4	Malignant neoplasms (C00-C97)	1,309	10.7	48.8
5	Assault (homicide) (X85-Y09,Y87.1)	1,040	8.5	38.8
6	Cerebrovascular diseases (I60-I69)	397	3.2	14.8
7	Intentional self-harm (suicide) (X60-X84,Y87.0)	356	2.9	13.3
8	Chronic liver disease and cirrhosis (K70,K73-K74)	284	2.3	10.6
9	Diabetes mellitus (E10-E14)	254	2.1	9.5
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	175	1.4	6.5
11	Influenza and pneumonia (J10-J18)	152	1.2	5.7
12	Chronic lower respiratory diseases (J40-J47)	130	1.1	4.8
13	Septicemia (A40-A41)	125	1.0	4.7
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	100	0.8	3.7
15	Viral hepatitis (B15-B19)	78	0.6	2.9
...	All other causes (Residual)	2,068	16.9	77.1
Black, male, 45-54 years				
...	All causes	19,913	100.0	1,060.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,837	24.3	257.5
2	Malignant neoplasms (C00-C97)	4,192	21.1	223.2
3	Human immunodeficiency virus (HIV) disease (B20-B24)	1,682	8.4	89.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,475	7.4	78.5
5	Cerebrovascular diseases (I60-I69)	969	4.9	51.6
6	Diabetes mellitus (E10-E14)	654	3.3	34.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	639	3.2	34.0
8	Assault (homicide) (X85-Y09,Y87.1)	508	2.6	27.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	398	2.0	21.2
10	Septicemia (A40-A41)	319	1.6	17.0
11	Chronic lower respiratory diseases (J40-J47)	294	1.5	15.7
12	Influenza and pneumonia (J10-J18)	272	1.4	14.5
13	Viral hepatitis (B15-B19)	241	1.2	12.8
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	233	1.2	12.4
15	Intentional self-harm (suicide) (X60-X84,Y87.0)	219	1.1	11.7
...	All other causes (Residual)	2,981	15.0	158.7

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 55-64 years				
...	All causes	22,678	100.0	2,172.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,882	30.3	659.4
2	Malignant neoplasms (C00-C97)	6,829	30.1	654.3
3	Cerebrovascular diseases (I60-I69)	1,258	5.5	120.5
4	Diabetes mellitus (E10-E14)	962	4.2	92.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	750	3.3	71.9
6	Chronic lower respiratory diseases (J40-J47)	595	2.6	57.0
7	Human immunodeficiency virus (HIV) disease (B20-B24)	501	2.2	48.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	489	2.2	46.9
9	Chronic liver disease and cirrhosis (K70,K73-K74)	414	1.8	39.7
10	Septicemia (A40-A41)	387	1.7	37.1
11	Influenza and pneumonia (J10-J18)	327	1.4	31.3
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	303	1.3	29.0
13	Assault (homicide) (X85-Y09,Y87.1)	160	0.7	15.3
14	Aortic aneurysm and dissection (I71)	125	0.6	12.0
15	Viral hepatitis (B15-B19)	88	0.4	8.4
...	All other causes (Residual)	2,608	11.5	249.9
Black, male, 65-74 years				
...	All causes	29,071	100.0	4,065.5
1	Malignant neoplasms (C00-C97)	9,321	32.1	1,303.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,672	29.8	1,212.8
3	Cerebrovascular diseases (I60-I69)	1,835	6.3	256.6
4	Diabetes mellitus (E10-E14)	1,264	4.3	176.8
5	Chronic lower respiratory diseases (J40-J47)	1,209	4.2	169.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	716	2.5	100.1
7	Influenza and pneumonia (J10-J18)	571	2.0	79.9
8	Septicemia (A40-A41)	555	1.9	77.6
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	543	1.9	75.9
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	345	1.2	48.2
11	Chronic liver disease and cirrhosis (K70,K73-K74)	258	0.9	36.1
12	Pneumonitis due to solids and liquids (J69)	190	0.7	26.6
13	Human immunodeficiency virus (HIV) disease (B20-B24)	168	0.6	23.5
14	Aortic aneurysm and dissection (I71)	151	0.5	21.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	131	0.5	18.3
...	All other causes (Residual)	3,142	10.8	439.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 75-84 years				
...	All causes	28,026	100.0	8,240.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,579	30.6	2,522.4
2	Malignant neoplasms (C00-C97)	7,767	27.7	2,283.6
3	Cerebrovascular diseases (I60-I69)	2,177	7.8	640.1
4	Chronic lower respiratory diseases (J40-J47)	1,223	4.4	359.6
5	Diabetes mellitus (E10-E14)	1,113	4.0	327.2
6	Influenza and pneumonia (J10-J18)	773	2.8	227.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	765	2.7	224.9
8	Septicemia (A40-A41)	684	2.4	201.1
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	438	1.6	128.8
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	360	1.3	105.8
11	Alzheimer's disease (G30)	322	1.1	94.7
12	Pneumonitis due to solids and liquids (J69)	246	0.9	72.3
13	Parkinson's disease (G20-G21)	158	0.6	46.5
14	Aortic aneurysm and dissection (I71)	155	0.6	45.6
15	Atherosclerosis (I70)	133	0.5	39.1
...	All other causes (Residual)	3,133	11.2	921.2
Black, male, 85 years and over				
...	All causes	14,560	100.0	15,494.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,885	33.6	5,198.6
2	Malignant neoplasms (C00-C97)	2,831	19.4	3,012.7
3	Cerebrovascular diseases (I60-I69)	1,265	8.7	1,346.2
4	Influenza and pneumonia (J10-J18)	655	4.5	697.0
5	Chronic lower respiratory diseases (J40-J47)	601	4.1	639.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	425	2.9	452.3
7	Diabetes mellitus (E10-E14)	392	2.7	417.2
8	Septicemia (A40-A41)	371	2.5	394.8
9	Alzheimer's disease (G30)	322	2.2	342.7
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	239	1.6	254.3
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	234	1.6	249.0
12	Pneumonitis due to solids and liquids (J69)	182	1.3	193.7
13	Atherosclerosis (I70)	128	0.9	136.2
14	Parkinson's disease (G20-G21)	75	0.5	79.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	66	0.5	70.2
...	All other causes (Residual)	1,889	13.0	2,010.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, all ages				
...	All causes	140,642	100.0	759.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	40,783	29.0	220.1
2	Malignant neoplasms (C00-C97)	29,128	20.7	157.2
3	Cerebrovascular diseases (I60-I69)	11,195	8.0	60.4
4	Diabetes mellitus (E10-E14)	7,250	5.2	39.1
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,837	2.7	20.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,746	2.7	20.2
7	Chronic lower respiratory diseases (J40-J47)	3,369	2.4	18.2
8	Septicemia (A40-A41)	3,341	2.4	18.0
9	Influenza and pneumonia (J10-J18)	3,075	2.2	16.6
10	Human immunodeficiency virus (HIV) disease (B20-B24)	2,448	1.7	13.2
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,295	1.6	12.4
12	Certain conditions originating in the perinatal period (P00-P96)	2,177	1.5	11.8
13	Alzheimer's disease (G30)	1,952	1.4	10.5
14	Assault (homicide) (X85-Y09,Y87.1)	1,385	1.0	7.5
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,007	0.7	5.4
...	All other causes (Residual)	23,654	16.8	127.7
Black, female, 1-4 years				
...	All causes	556	100.0	50.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	166	29.9	15.2
2	Assault (homicide) (X85-Y09,Y87.1)	70	12.6	6.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	56	10.1	5.1
4	Malignant neoplasms (C00-C97)	36	6.5	3.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	25	4.5	2.3
6	Septicemia (A40-A41)	13	2.3	*
7	Anemias (D50-D64)	11	2.0	*
7	Cerebrovascular diseases (I60-I69)	11	2.0	*
9	Influenza and pneumonia (J10-J18)	9	1.6	*
9	Certain conditions originating in the perinatal period (P00-P96)	9	1.6	*
11	Meningitis (G00,G03)	7	1.3	*
11	Chronic lower respiratory diseases (J40-J47)	7	1.3	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	6	1.1	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	6	1.1	*
15	Acute bronchitis and bronchiolitis (J20-J21)	3	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.5	*
...	All other causes (Residual)	118	21.2	10.8

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 5-14 years				
...	All causes	669	100.0	21.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	217	32.4	7.0
2	Malignant neoplasms (C00-C97)	63	9.4	2.0
3	Assault (homicide) (X85-Y09,Y87.1)	52	7.8	1.7
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	39	5.8	1.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	34	5.1	1.1
6	Chronic lower respiratory diseases (J40-J47)	27	4.0	0.9
7	Anemias (D50-D64)	14	2.1	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	11	1.6	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	1.6	*
10	Cerebrovascular diseases (I60-I69)	10	1.5	*
11	Certain conditions originating in the perinatal period (P00-P96)	7	1.0	*
12	Meningococcal infection (A39)	5	0.7	*
12	Septicemia (A40-A41)	5	0.7	*
12	Diabetes mellitus (E10-E14)	5	0.7	*
12	Intentional self-harm (suicide) (X60-X84,Y87.0)	5	0.7	*
...	All other causes (Residual)	164	24.5	5.3
Black, female, 15-24 years				
...	All causes	1,731	100.0	59.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	434	25.1	14.9
2	Assault (homicide) (X85-Y09,Y87.1)	318	18.4	11.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	140	8.1	4.8
4	Malignant neoplasms (C00-C97)	122	7.0	4.2
5	Human immunodeficiency virus (HIV) disease (B20-B24)	65	3.8	2.2
5	Intentional self-harm (suicide) (X60-X84,Y87.0)	65	3.8	2.2
7	Pregnancy, childbirth and the puerperium (O00-099)	36	2.1	1.2
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	33	1.9	1.1
9	Anemias (D50-D64)	32	1.8	1.1
10	Chronic lower respiratory diseases (J40-J47)	30	1.7	1.0
11	Diabetes mellitus (E10-E14)	27	1.6	0.9
12	Cerebrovascular diseases (I60-I69)	24	1.4	0.8
13	Influenza and pneumonia (J10-J18)	23	1.3	0.8
14	Septicemia (A40-A41)	15	0.9	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	0.8	*
...	All other causes (Residual)	353	20.4	12.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 25-34 years				
...	All causes	3,489	100.0	126.7
1	Human immunodeficiency virus (HIV) disease (B20-B24)	597	17.1	21.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	430	12.3	15.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	390	11.2	14.2
4	Malignant neoplasms (C00-C97)	387	11.1	14.1
5	Assault (homicide) (X85-Y09,Y87.1)	313	9.0	11.4
6	Diabetes mellitus (E10-E14)	90	2.6	3.3
7	Cerebrovascular diseases (I60-I69)	89	2.6	3.2
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	74	2.1	2.7
9	Anemias (D50-D64)	69	2.0	2.5
10	Pregnancy, childbirth and the puerperium (O00-099)	65	1.9	2.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	50	1.4	1.8
12	Septicemia (A40-A41)	49	1.4	1.8
13	Chronic lower respiratory diseases (J40-J47)	44	1.3	1.6
14	Influenza and pneumonia (J10-J18)	34	1.0	1.2
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	31	0.9	1.1
...	All other causes (Residual)	777	22.3	28.2
Black, female, 35-44 years				
...	All causes	8,306	100.0	274.7
1	Malignant neoplasms (C00-C97)	1,799	21.7	59.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,296	15.6	42.9
3	Human immunodeficiency virus (HIV) disease (B20-B24)	984	11.8	32.5
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	609	7.3	20.1
5	Cerebrovascular diseases (I60-I69)	439	5.3	14.5
6	Assault (homicide) (X85-Y09,Y87.1)	336	4.0	11.1
7	Diabetes mellitus (E10-E14)	208	2.5	6.9
8	Chronic liver disease and cirrhosis (K70,K73-K74)	167	2.0	5.5
9	Septicemia (A40-A41)	147	1.8	4.9
10	Chronic lower respiratory diseases (J40-J47)	145	1.7	4.8
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	131	1.6	4.3
12	Influenza and pneumonia (J10-J18)	105	1.3	3.5
13	Anemias (D50-D64)	86	1.0	2.8
14	Intentional self-harm (suicide) (X60-X84,Y87.0)	82	1.0	2.7
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	79	1.0	2.6
...	All other causes (Residual)	1,693	20.4	56.0

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 45-54 years				
...	All causes	13,376	100.0	594.1
1	Malignant neoplasms (C00-C97)	3,955	29.6	175.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,843	21.3	126.3
3	Cerebrovascular diseases (I60-I69)	867	6.5	38.5
4	Diabetes mellitus (E10-E14)	625	4.7	27.8
5	Human immunodeficiency virus (HIV) disease (B20-B24)	575	4.3	25.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	513	3.8	22.8
7	Chronic lower respiratory diseases (J40-J47)	305	2.3	13.5
8	Chronic liver disease and cirrhosis (K70,K73-K74)	298	2.2	13.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	271	2.0	12.0
10	Septicemia (A40-A41)	256	1.9	11.4
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	176	1.3	7.8
12	Influenza and pneumonia (J10-J18)	153	1.1	6.8
13	Assault (homicide) (X85-Y09,Y87.1)	120	0.9	5.3
14	Viral hepatitis (B15-B19)	95	0.7	4.2
15	Anemias (D50-D64)	64	0.5	2.8
...	All other causes (Residual)	2,260	16.9	100.4
Black, female, 55-64 years				
...	All causes	16,603	100.0	1,213.4
1	Malignant neoplasms (C00-C97)	5,290	31.9	386.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,503	27.1	329.1
3	Diabetes mellitus (E10-E14)	1,053	6.3	77.0
4	Cerebrovascular diseases (I60-I69)	1,033	6.2	75.5
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	502	3.0	36.7
6	Chronic lower respiratory diseases (J40-J47)	478	2.9	34.9
7	Septicemia (A40-A41)	381	2.3	27.8
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	256	1.5	18.7
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	247	1.5	18.1
10	Chronic liver disease and cirrhosis (K70,K73-K74)	219	1.3	16.0
11	Influenza and pneumonia (J10-J18)	211	1.3	15.4
12	Human immunodeficiency virus (HIV) disease (B20-B24)	135	0.8	9.9
13	Aortic aneurysm and dissection (I71)	75	0.5	5.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	59	0.4	4.3
15	Viral hepatitis (B15-B19)	55	0.3	4.0
...	All other causes (Residual)	2,106	12.7	153.9

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 65-74 years				
...	All causes	26,124	100.0	2,659.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,918	30.3	805.9
2	Malignant neoplasms (C00-C97)	7,315	28.0	744.5
3	Diabetes mellitus (E10-E14)	1,854	7.1	188.7
3	Cerebrovascular diseases (I60-I69)	1,854	7.1	188.7
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	937	3.6	95.4
6	Chronic lower respiratory diseases (J40-J47)	805	3.1	81.9
7	Septicemia (A40-A41)	634	2.4	64.5
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	461	1.8	46.9
9	Influenza and pneumonia (J10-J18)	426	1.6	43.4
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	325	1.2	33.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	176	0.7	17.9
12	Alzheimer's disease (G30)	148	0.6	15.1
13	Aortic aneurysm and dissection (I71)	130	0.5	13.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	117	0.4	11.9
15	Pneumonitis due to solids and liquids (J69)	100	0.4	10.2
...	All other causes (Residual)	2,924	11.2	297.6
Black, female, 75-84 years				
...	All causes	33,418	100.0	5,967.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,223	33.6	2,004.2
2	Malignant neoplasms (C00-C97)	6,594	19.7	1,177.6
3	Cerebrovascular diseases (I60-I69)	3,222	9.6	575.4
4	Diabetes mellitus (E10-E14)	2,102	6.3	375.4
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,024	3.1	182.9
6	Chronic lower respiratory diseases (J40-J47)	929	2.8	165.9
7	Septicemia (A40-A41)	906	2.7	161.8
8	Influenza and pneumonia (J10-J18)	796	2.4	142.1
9	Alzheimer's disease (G30)	686	2.1	122.5
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	622	1.9	111.1
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	340	1.0	60.7
12	Pneumonitis due to solids and liquids (J69)	226	0.7	40.4
13	Atherosclerosis (I70)	190	0.6	33.9
14	Aortic aneurysm and dissection (I71)	178	0.5	31.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	165	0.5	29.5
...	All other causes (Residual)	4,215	12.6	752.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 85 years and over				
...	All causes	32,478	100.0	14,441.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,344	38.0	5,489.0
2	Cerebrovascular diseases (I60-I69)	3,626	11.2	1,612.4
3	Malignant neoplasms (C00-C97)	3,559	11.0	1,582.6
4	Diabetes mellitus (E10-E14)	1,286	4.0	571.8
5	Influenza and pneumonia (J10-J18)	1,266	3.9	562.9
6	Alzheimer's disease (G30)	1,101	3.4	489.6
7	Septicemia (A40-A41)	884	2.7	393.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	874	2.7	388.6
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	672	2.1	298.8
10	Chronic lower respiratory diseases (J40-J47)	594	1.8	264.1
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	340	1.0	151.2
12	Pneumonitis due to solids and liquids (J69)	329	1.0	146.3
13	Atherosclerosis (I70)	314	1.0	139.6
14	Nutritional deficiencies (E40-E64)	159	0.5	70.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	139	0.4	61.8
15	Aortic aneurysm and dissection (I71)	139	0.4	61.8
...	All other causes (Residual)	4,852	14.9	2,157.5
American Indian, both sexes, all ages				
...	All causes	11,363	100.0	466.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,417	21.3	99.2
2	Malignant neoplasms (C00-C97)	1,914	16.8	78.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,353	11.9	55.5
4	Diabetes mellitus (E10-E14)	616	5.4	25.3
5	Cerebrovascular diseases (I60-I69)	572	5.0	23.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	534	4.7	21.9
7	Chronic lower respiratory diseases (J40-J47)	429	3.8	17.6
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	297	2.6	12.2
9	Influenza and pneumonia (J10-J18)	289	2.5	11.9
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	215	1.9	8.8
11	Assault (homicide) (X85-Y09,Y87.1)	203	1.8	8.3
12	Septicemia (A40-A41)	162	1.4	6.6
13	Certain conditions originating in the perinatal period (P00-P96)	124	1.1	5.1
14	Alzheimer's disease (G30)	95	0.8	3.9
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	95	0.8	3.9
...	All other causes (Residual)	2,048	18.0	84.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 1-4 years				
...	All causes	91	100.0	55.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	39	42.9	23.9
2	Assault (homicide) (X85-Y09,Y87.1)	8	8.8	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	4.4	*
4	Septicemia (A40-A41)	3	3.3	*
4	Malignant neoplasms (C00-C97)	3	3.3	*
6	Chronic lower respiratory diseases (J40-J47)	2	2.2	*
6	Certain conditions originating in the perinatal period (P00-P96)	2	2.2	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	1	1.1	*
8	Parkinson's disease (G20-G21)	1	1.1	*
8	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	1.1	*
8	Cerebrovascular diseases (I60-I69)	1	1.1	*
8	Diseases of appendix (K35-K38)	1	1.1	*
...	All other causes (Residual)	25	27.5	15.3
American Indian, both sexes, 5-14 years				
...	All causes	113	100.0	24.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	70	61.9	15.0
2	Malignant neoplasms (C00-C97)	8	7.1	*
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	7	6.2	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	4.4	*
5	Influenza and pneumonia (J10-J18)	3	2.7	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	2.7	*
7	Cerebrovascular diseases (I60-I69)	2	1.8	*
8	Septicemia (A40-A41)	1	0.9	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.9	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.9	*
8	Assault (homicide) (X85-Y09,Y87.1)	1	0.9	*
8	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.9	*
...	All other causes (Residual)	10	8.8	*

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 15-24 years				
...	All causes	519	100.0	117.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	299	57.6	67.9
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	89	17.1	20.2
3	Assault (homicide) (X85-Y09,Y87.1)	51	9.8	11.6
4	Malignant neoplasms (C00-C97)	13	2.5	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	11	2.1	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.8	*
7	Chronic lower respiratory diseases (J40-J47)	3	0.6	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.6	*
9	Septicemia (A40-A41)	2	0.4	*
9	Diabetes mellitus (E10-E14)	2	0.4	*
9	Influenza and pneumonia (J10-J18)	2	0.4	*
9	Chronic liver disease and cirrhosis (K70,K73-K74)	2	0.4	*
9	Pregnancy, childbirth and the puerperium (O00-099)	2	0.4	*
14	Meningococcal infection (A39)	1	0.2	*
14	Cerebrovascular diseases (I60-I69)	1	0.2	*
14	Certain conditions originating in the perinatal period (P00-P96)	1	0.2	*
14	Legal intervention (Y35,Y89.0)	1	0.2	*
...	All other causes (Residual)	32	6.2	7.3
American Indian, both sexes, 25-34 years				
...	All causes	596	100.0	158.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	239	40.1	63.5
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	78	13.1	20.7
3	Assault (homicide) (X85-Y09,Y87.1)	59	9.9	15.7
4	Chronic liver disease and cirrhosis (K70,K73-K74)	39	6.5	10.4
5	Malignant neoplasms (C00-C97)	26	4.4	6.9
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	24	4.0	6.4
7	Human immunodeficiency virus (HIV) disease (B20-B24)	8	1.3	*
8	Diabetes mellitus (E10-E14)	7	1.2	*
9	Cerebrovascular diseases (I60-I69)	5	0.8	*
9	Influenza and pneumonia (J10-J18)	5	0.8	*
11	Viral hepatitis (B15-B19)	4	0.7	*
11	Chronic lower respiratory diseases (J40-J47)	4	0.7	*
11	Pregnancy, childbirth and the puerperium (O00-099)	4	0.7	*
14	Septicemia (A40-A41)	3	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.5	*
...	All other causes (Residual)	88	14.8	23.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 35-44 years				
...	All causes	1,081	100.0	299.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	256	23.7	70.9
2	Chronic liver disease and cirrhosis (K70,K73-K74)	137	12.7	37.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	122	11.3	33.8
4	Malignant neoplasms (C00-C97)	105	9.7	29.1
5	Intentional self-harm (suicide) (X60-X84,Y87.0)	67	6.2	18.5
6	Assault (homicide) (X85-Y09,Y87.1)	46	4.3	12.7
7	Influenza and pneumonia (J10-J18)	28	2.6	7.8
8	Human immunodeficiency virus (HIV) disease (B20-B24)	24	2.2	6.6
9	Diabetes mellitus (E10-E14)	23	2.1	6.4
10	Cerebrovascular diseases (I60-I69)	22	2.0	6.1
11	Viral hepatitis (B15-B19)	15	1.4	*
12	Septicemia (A40-A41)	13	1.2	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	1.1	*
14	Chronic lower respiratory diseases (J40-J47)	10	0.9	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.4	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	0.4	*
...	All other causes (Residual)	193	17.9	53.4
American Indian, both sexes, 45-54 years				
...	All causes	1,354	100.0	508.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	249	18.4	93.6
2	Malignant neoplasms (C00-C97)	220	16.2	82.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	197	14.5	74.0
4	Chronic liver disease and cirrhosis (K70,K73-K74)	147	10.9	55.3
5	Diabetes mellitus (E10-E14)	78	5.8	29.3
6	Cerebrovascular diseases (I60-I69)	47	3.5	17.7
7	Intentional self-harm (suicide) (X60-X84,Y87.0)	31	2.3	11.7
8	Viral hepatitis (B15-B19)	25	1.8	9.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22	1.6	8.3
10	Septicemia (A40-A41)	21	1.6	7.9
10	Influenza and pneumonia (J10-J18)	21	1.6	7.9
12	Human immunodeficiency virus (HIV) disease (B20-B24)	18	1.3	*
12	Assault (homicide) (X85-Y09,Y87.1)	18	1.3	*
14	Chronic lower respiratory diseases (J40-J47)	17	1.3	*
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	5	0.4	*
15	Peptic ulcer (K25-K28)	5	0.4	*
...	All other causes (Residual)	233	17.2	87.6

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 55-64 years				
...	All causes	1,698	100.0	1,113.5
1	Malignant neoplasms (C00-C97)	417	24.6	273.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	394	23.2	258.4
3	Diabetes mellitus (E10-E14)	134	7.9	87.9
4	Chronic liver disease and cirrhosis (K70,K73-K74)	123	7.2	80.7
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	94	5.5	61.6
6	Cerebrovascular diseases (I60-I69)	79	4.7	51.8
7	Chronic lower respiratory diseases (J40-J47)	71	4.2	46.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	39	2.3	25.6
9	Septicemia (A40-A41)	29	1.7	19.0
10	Influenza and pneumonia (J10-J18)	24	1.4	15.7
11	Intentional self-harm (suicide) (X60-X84,Y87.0)	16	0.9	*
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	14	0.8	*
13	Viral hepatitis (B15-B19)	10	0.6	*
14	Pneumonitis due to solids and liquids (J69)	7	0.4	*
15	Tuberculosis (A16-A19)	6	0.4	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.4	*
15	Assault (homicide) (X85-Y09,Y87.1)	6	0.4	*
...	All other causes (Residual)	229	13.5	150.2
American Indian, both sexes, 65-74 years				
...	All causes	2,141	100.0	2,314.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	548	25.6	592.5
2	Malignant neoplasms (C00-C97)	539	25.2	582.7
3	Diabetes mellitus (E10-E14)	186	8.7	201.1
4	Chronic lower respiratory diseases (J40-J47)	124	5.8	134.1
5	Cerebrovascular diseases (I60-I69)	108	5.0	116.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	65	3.0	70.3
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	63	2.9	68.1
8	Chronic liver disease and cirrhosis (K70,K73-K74)	59	2.8	63.8
9	Influenza and pneumonia (J10-J18)	45	2.1	48.7
10	Septicemia (A40-A41)	33	1.5	35.7
11	Aortic aneurysm and dissection (I71)	21	1.0	22.7
12	Atherosclerosis (I70)	16	0.7	*
13	Alzheimer's disease (G30)	14	0.7	*
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	13	0.6	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	0.5	*
...	All other causes (Residual)	297	13.9	321.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 75-84 years				
...	All causes	2,062	100.0	3,925.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	597	29.0	1,136.6
2	Malignant neoplasms (C00-C97)	412	20.0	784.4
3	Cerebrovascular diseases (I60-I69)	178	8.6	338.9
4	Diabetes mellitus (E10-E14)	133	6.5	253.2
5	Chronic lower respiratory diseases (J40-J47)	126	6.1	239.9
6	Influenza and pneumonia (J10-J18)	71	3.4	135.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	51	2.5	97.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	46	2.2	87.6
9	Alzheimer's disease (G30)	39	1.9	74.2
10	Septicemia (A40-A41)	34	1.6	64.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	25	1.2	47.6
12	Parkinson's disease (G20-G21)	20	1.0	38.1
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	16	0.8	*
14	Pneumonitis due to solids and liquids (J69)	14	0.7	*
15	Atherosclerosis (I70)	13	0.6	*
...	All other causes (Residual)	287	13.9	546.4
American Indian, both sexes, 85 years and over				
...	All causes	1,382	100.0	6,376.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	459	33.2	2,117.9
2	Malignant neoplasms (C00-C97)	169	12.2	779.8
3	Cerebrovascular diseases (I60-I69)	129	9.3	595.2
4	Influenza and pneumonia (J10-J18)	84	6.1	387.6
5	Chronic lower respiratory diseases (J40-J47)	71	5.1	327.6
6	Diabetes mellitus (E10-E14)	53	3.8	244.6
7	Alzheimer's disease (G30)	39	2.8	180.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	27	2.0	124.6
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	24	1.7	110.7
10	Septicemia (A40-A41)	22	1.6	101.5
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	21	1.5	96.9
12	Atherosclerosis (I70)	17	1.2	*
13	Nutritional deficiencies (E40-E64)	15	1.1	*
13	Pneumonitis due to solids and liquids (J69)	15	1.1	*
15	Parkinson's disease (G20-G21)	13	0.9	*
...	All other causes (Residual)	224	16.2	1,033.6

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, all ages				
...	All causes	6,185	100.0	512.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,341	21.7	111.2
2	Malignant neoplasms (C00-C97)	997	16.1	82.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	900	14.6	74.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	298	4.8	24.7
5	Diabetes mellitus (E10-E14)	275	4.4	22.8
6	Cerebrovascular diseases (I60-I69)	250	4.0	20.7
7	Intentional self-harm (suicide) (X60-X84,Y87.0)	237	3.8	19.6
8	Chronic lower respiratory diseases (J40-J47)	228	3.7	18.9
9	Assault (homicide) (X85-Y09,Y87.1)	159	2.6	13.2
10	Influenza and pneumonia (J10-J18)	149	2.4	12.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	98	1.6	8.1
12	Septicemia (A40-A41)	74	1.2	6.1
13	Certain conditions originating in the perinatal period (P00-P96)	72	1.2	6.0
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	56	0.9	4.6
15	Human immunodeficiency virus (HIV) disease (B20-B24)	43	0.7	3.6
...	All other causes (Residual)	1,008	16.3	83.6
American Indian, male, 1-4 years				
...	All causes	49	100.0	59.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	25	51.0	30.3
2	Assault (homicide) (X85-Y09,Y87.1)	6	12.2	*
3	Septicemia (A40-A41)	2	4.1	*
3	Malignant neoplasms (C00-C97)	2	4.1	*
5	Cerebrovascular diseases (I60-I69)	1	2.0	*
5	Chronic lower respiratory diseases (J40-J47)	1	2.0	*
5	Certain conditions originating in the perinatal period (P00-P96)	1	2.0	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	2.0	*
...	All other causes (Residual)	10	20.4	*

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 5-14 years				
...	All causes	61	100.0	25.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	37	60.7	15.6
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	6	9.8	*
3	Malignant neoplasms (C00-C97)	4	6.6	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	4.9	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	3.3	*
6	Septicemia (A40-A41)	1	1.6	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.6	*
6	Assault (homicide) (X85-Y09,Y87.1)	1	1.6	*
...	All other causes (Residual)	6	9.8	*
American Indian, male, 15-24 years				
...	All causes	369	100.0	167.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	202	54.7	91.4
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	71	19.2	32.1
3	Assault (homicide) (X85-Y09,Y87.1)	46	12.5	20.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	8	2.2	*
5	Malignant neoplasms (C00-C97)	7	1.9	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.8	*
6	Chronic lower respiratory diseases (J40-J47)	3	0.8	*
8	Diabetes mellitus (E10-E14)	2	0.5	*
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	0.5	*
10	Septicemia (A40-A41)	1	0.3	*
10	Influenza and pneumonia (J10-J18)	1	0.3	*
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.3	*
10	Legal intervention (Y35,Y89.0)	1	0.3	*
...	All other causes (Residual)	21	5.7	9.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 25-34 years				
...	All causes	410	100.0	212.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	172	42.0	89.1
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	59	14.4	30.6
3	Assault (homicide) (X85-Y09,Y87.1)	46	11.2	23.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	23	5.6	11.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	16	3.9	*
6	Malignant neoplasms (C00-C97)	13	3.2	*
7	Diabetes mellitus (E10-E14)	6	1.5	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	5	1.2	*
9	Influenza and pneumonia (J10-J18)	3	0.7	*
9	Chronic lower respiratory diseases (J40-J47)	3	0.7	*
11	Septicemia (A40-A41)	2	0.5	*
11	Viral hepatitis (B15-B19)	2	0.5	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.5	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.2	*
14	Pneumonitis due to solids and liquids (J69)	1	0.2	*
14	Legal intervention (Y35,Y89.0)	1	0.2	*
...	All other causes (Residual)	55	13.4	28.5
American Indian, male, 35-44 years				
...	All causes	675	100.0	375.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	166	24.6	92.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	83	12.3	46.1
3	Chronic liver disease and cirrhosis (K70,K73-K74)	78	11.6	43.3
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	53	7.9	29.5
5	Malignant neoplasms (C00-C97)	49	7.3	27.2
6	Assault (homicide) (X85-Y09,Y87.1)	32	4.7	17.8
7	Human immunodeficiency virus (HIV) disease (B20-B24)	20	3.0	11.1
8	Influenza and pneumonia (J10-J18)	18	2.7	*
9	Diabetes mellitus (E10-E14)	16	2.4	*
10	Cerebrovascular diseases (I60-I69)	11	1.6	*
11	Viral hepatitis (B15-B19)	8	1.2	*
12	Septicemia (A40-A41)	7	1.0	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	1.0	*
14	Chronic lower respiratory diseases (J40-J47)	4	0.6	*
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2	0.3	*
...	All other causes (Residual)	121	17.9	67.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 45-54 years				
...	All causes	858	100.0	667.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	179	20.9	139.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	150	17.5	116.6
3	Malignant neoplasms (C00-C97)	116	13.5	90.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	91	10.6	70.7
5	Diabetes mellitus (E10-E14)	41	4.8	31.9
6	Intentional self-harm (suicide) (X60-X84,Y87.0)	24	2.8	18.7
7	Cerebrovascular diseases (I60-I69)	22	2.6	17.1
8	Viral hepatitis (B15-B19)	16	1.9	*
8	Influenza and pneumonia (J10-J18)	16	1.9	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	14	1.6	*
10	Assault (homicide) (X85-Y09,Y87.1)	14	1.6	*
12	Septicemia (A40-A41)	12	1.4	*
12	Chronic lower respiratory diseases (J40-J47)	12	1.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	1.3	*
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	3	0.3	*
15	Peptic ulcer (K25-K28)	3	0.3	*
...	All other causes (Residual)	134	15.6	104.2
American Indian, male, 55-64 years				
...	All causes	964	100.0	1,347.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	252	26.1	352.2
2	Malignant neoplasms (C00-C97)	226	23.4	315.9
3	Chronic liver disease and cirrhosis (K70,K73-K74)	71	7.4	99.2
4	Diabetes mellitus (E10-E14)	68	7.1	95.0
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	56	5.8	78.3
6	Cerebrovascular diseases (I60-I69)	43	4.5	60.1
7	Chronic lower respiratory diseases (J40-J47)	41	4.3	57.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21	2.2	29.3
9	Influenza and pneumonia (J10-J18)	16	1.7	*
10	Intentional self-harm (suicide) (X60-X84,Y87.0)	15	1.6	*
11	Septicemia (A40-A41)	14	1.5	*
12	Viral hepatitis (B15-B19)	5	0.5	*
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	5	0.5	*
14	Tuberculosis (A16-A19)	4	0.4	*
14	Pneumonitis due to solids and liquids (J69)	4	0.4	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	0.4	*
14	Assault (homicide) (X85-Y09,Y87.1)	4	0.4	*
...	All other causes (Residual)	115	11.9	160.7

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 65-74 years				
...	All causes	1,113	100.0	2,676.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	336	30.2	808.0
2	Malignant neoplasms (C00-C97)	291	26.1	699.8
3	Diabetes mellitus (E10-E14)	70	6.3	168.3
4	Cerebrovascular diseases (I60-I69)	65	5.8	156.3
5	Chronic lower respiratory diseases (J40-J47)	50	4.5	120.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	36	3.2	86.6
7	Influenza and pneumonia (J10-J18)	29	2.6	69.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	28	2.5	67.3
9	Chronic liver disease and cirrhosis (K70,K73-K74)	21	1.9	50.5
10	Septicemia (A40-A41)	15	1.3	*
11	Aortic aneurysm and dissection (I71)	11	1.0	*
12	Intentional self-harm (suicide) (X60-X84,Y87.0)	8	0.7	*
13	Pneumonitis due to solids and liquids (J69)	7	0.6	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.5	*
14	Atherosclerosis (I70)	6	0.5	*
...	All other causes (Residual)	134	12.0	322.2
American Indian, male, 75-84 years				
...	All causes	989	100.0	4,442.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	306	30.9	1,374.4
2	Malignant neoplasms (C00-C97)	213	21.5	956.7
3	Chronic lower respiratory diseases (J40-J47)	77	7.8	345.8
4	Cerebrovascular diseases (I60-I69)	69	7.0	309.9
5	Diabetes mellitus (E10-E14)	50	5.1	224.6
6	Influenza and pneumonia (J10-J18)	33	3.3	148.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	32	3.2	143.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	1.6	*
9	Septicemia (A40-A41)	14	1.4	*
10	Alzheimer's disease (G30)	13	1.3	*
11	Parkinson's disease (G20-G21)	12	1.2	*
11	Chronic liver disease and cirrhosis (K70,K73-K74)	12	1.2	*
13	Pneumonitis due to solids and liquids (J69)	8	0.8	*
14	Aortic aneurysm and dissection (I71)	7	0.7	*
15	Anemias (D50-D64)	6	0.6	*
15	Atherosclerosis (I70)	6	0.6	*
...	All other causes (Residual)	115	11.6	516.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 85 years and over				
...	All causes	501	100.0	7,299.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	155	30.9	2,258.2
2	Malignant neoplasms (C00-C97)	74	14.8	1,078.1
3	Cerebrovascular diseases (I60-I69)	39	7.8	568.2
4	Chronic lower respiratory diseases (J40-J47)	36	7.2	524.5
5	Influenza and pneumonia (J10-J18)	31	6.2	451.6
6	Diabetes mellitus (E10-E14)	22	4.4	320.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	2.6	*
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12	2.4	*
9	Alzheimer's disease (G30)	9	1.8	*
10	Parkinson's disease (G20-G21)	8	1.6	*
11	Pneumonitis due to solids and liquids (J69)	7	1.4	*
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	6	1.2	*
13	Septicemia (A40-A41)	5	1.0	*
14	Atherosclerosis (I70)	4	0.8	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.4	*
15	Nutritional deficiencies (E40-E64)	2	0.4	*
15	Assault (homicide) (X85-Y09,Y87.1)	2	0.4	*
...	All other causes (Residual)	74	14.8	1,078.1
American Indian, female, all ages				
...	All causes	5,178	100.0	421.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,076	20.8	87.5
2	Malignant neoplasms (C00-C97)	917	17.7	74.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	453	8.7	36.8
4	Diabetes mellitus (E10-E14)	341	6.6	27.7
5	Cerebrovascular diseases (I60-I69)	322	6.2	26.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	236	4.6	19.2
7	Chronic lower respiratory diseases (J40-J47)	201	3.9	16.3
8	Influenza and pneumonia (J10-J18)	140	2.7	11.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	117	2.3	9.5
10	Septicemia (A40-A41)	88	1.7	7.2
11	Alzheimer's disease (G30)	69	1.3	5.6
12	Intentional self-harm (suicide) (X60-X84,Y87.0)	60	1.2	4.9
13	Certain conditions originating in the perinatal period (P00-P96)	52	1.0	4.2
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	49	0.9	4.0
15	Assault (homicide) (X85-Y09,Y87.1)	44	0.8	3.6
...	All other causes (Residual)	1,013	19.6	82.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 1-4 years				
...	All causes	42	100.0	52.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14	33.3	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	7.1	*
3	Assault (homicide) (X85-Y09,Y87.1)	2	4.8	*
4	Septicemia (A40-A41)	1	2.4	*
4	Human immunodeficiency virus (HIV) disease (B20-B24)	1	2.4	*
4	Malignant neoplasms (C00-C97)	1	2.4	*
4	Parkinson's disease (G20-G21)	1	2.4	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	2.4	*
4	Chronic lower respiratory diseases (J40-J47)	1	2.4	*
4	Diseases of appendix (K35-K38)	1	2.4	*
4	Certain conditions originating in the perinatal period (P00-P96)	1	2.4	*
...	All other causes (Residual)	15	35.7	*
American Indian, female, 5-14 years				
...	All causes	52	100.0	22.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	33	63.5	14.4
2	Malignant neoplasms (C00-C97)	4	7.7	*
3	Influenza and pneumonia (J10-J18)	3	5.8	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	3.8	*
4	Cerebrovascular diseases (I60-I69)	2	3.8	*
6	Human immunodeficiency virus (HIV) disease (B20-B24)	1	1.9	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	1.9	*
6	Intentional self-harm (suicide) (X60-X84,Y87.0)	1	1.9	*
6	Complications of medical and surgical care (Y40-Y84,Y88)	1	1.9	*
...	All other causes (Residual)	4	7.7	*

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 15-24 years				
...	All causes	150	100.0	68.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	97	64.7	44.2
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	18	12.0	*
3	Malignant neoplasms (C00-C97)	6	4.0	*
4	Assault (homicide) (X85-Y09,Y87.1)	5	3.3	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	2.0	*
6	Pregnancy, childbirth and the puerperium (O00-099)	2	1.3	*
7	Meningococcal infection (A39)	1	0.7	*
7	Septicemia (A40-A41)	1	0.7	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.7	*
7	Cerebrovascular diseases (I60-I69)	1	0.7	*
7	Influenza and pneumonia (J10-J18)	1	0.7	*
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.7	*
7	Certain conditions originating in the perinatal period (P00-P96)	1	0.7	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.7	*
...	All other causes (Residual)	11	7.3	*
American Indian, female, 25-34 years				
...	All causes	186	100.0	101.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	67	36.0	36.6
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	19	10.2	*
3	Chronic liver disease and cirrhosis (K70,K73-K74)	16	8.6	*
4	Malignant neoplasms (C00-C97)	13	7.0	*
4	Assault (homicide) (X85-Y09,Y87.1)	13	7.0	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	8	4.3	*
7	Cerebrovascular diseases (I60-I69)	5	2.7	*
8	Pregnancy, childbirth and the puerperium (O00-099)	4	2.2	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	3	1.6	*
10	Viral hepatitis (B15-B19)	2	1.1	*
10	Influenza and pneumonia (J10-J18)	2	1.1	*
12	Septicemia (A40-A41)	1	0.5	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.5	*
12	Diabetes mellitus (E10-E14)	1	0.5	*
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1	0.5	*
12	Chronic lower respiratory diseases (J40-J47)	1	0.5	*
12	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.5	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.5	*
12	Inflammatory diseases of female pelvic organs (N70-N76)	1	0.5	*
...	All other causes (Residual)	26	14.0	14.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
	American Indian, female, 35-44 years			
...	All causes	406	100.0	224.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	90	22.2	49.6
2	Chronic liver disease and cirrhosis (K70,K73-K74)	59	14.5	32.5
3	Malignant neoplasms (C00-C97)	56	13.8	30.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	39	9.6	21.5
5	Intentional self-harm (suicide) (X60-X84,Y87.0)	14	3.4	*
5	Assault (homicide) (X85-Y09,Y87.1)	14	3.4	*
7	Cerebrovascular diseases (I60-I69)	11	2.7	*
8	Influenza and pneumonia (J10-J18)	10	2.5	*
9	Viral hepatitis (B15-B19)	7	1.7	*
9	Diabetes mellitus (E10-E14)	7	1.7	*
11	Septicemia (A40-A41)	6	1.5	*
11	Chronic lower respiratory diseases (J40-J47)	6	1.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	1.2	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	4	1.0	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.7	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.7	*
...	All other causes (Residual)	72	17.7	39.7
	American Indian, female, 45-54 years			
...	All causes	496	100.0	360.9
1	Malignant neoplasms (C00-C97)	104	21.0	75.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	70	14.1	50.9
3	Chronic liver disease and cirrhosis (K70,K73-K74)	56	11.3	40.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	47	9.5	34.2
5	Diabetes mellitus (E10-E14)	37	7.5	26.9
6	Cerebrovascular diseases (I60-I69)	25	5.0	18.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	2.2	*
8	Septicemia (A40-A41)	9	1.8	*
8	Viral hepatitis (B15-B19)	9	1.8	*
10	Intentional self-harm (suicide) (X60-X84,Y87.0)	7	1.4	*
11	Influenza and pneumonia (J10-J18)	5	1.0	*
11	Chronic lower respiratory diseases (J40-J47)	5	1.0	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.8	*
13	Assault (homicide) (X85-Y09,Y87.1)	4	0.8	*
15	Pneumonitis due to solids and liquids (J69)	3	0.6	*
...	All other causes (Residual)	100	20.2	72.8

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 55-64 years				
...	All causes	734	100.0	906.8
1	Malignant neoplasms (C00-C97)	191	26.0	236.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	142	19.3	175.4
3	Diabetes mellitus (E10-E14)	66	9.0	81.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	52	7.1	64.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	38	5.2	46.9
6	Cerebrovascular diseases (I60-I69)	36	4.9	44.5
7	Chronic lower respiratory diseases (J40-J47)	30	4.1	37.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18	2.5	*
9	Septicemia (A40-A41)	15	2.0	*
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	9	1.2	*
11	Influenza and pneumonia (J10-J18)	8	1.1	*
12	Viral hepatitis (B15-B19)	5	0.7	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.4	*
13	Alzheimer's disease (G30)	3	0.4	*
13	Pneumonitis due to solids and liquids (J69)	3	0.4	*
...	All other causes (Residual)	115	15.7	142.1
American Indian, female, 65-74 years				
...	All causes	1,028	100.0	2,019.2
1	Malignant neoplasms (C00-C97)	248	24.1	487.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	212	20.6	416.4
3	Diabetes mellitus (E10-E14)	116	11.3	227.8
4	Chronic lower respiratory diseases (J40-J47)	74	7.2	145.3
5	Cerebrovascular diseases (I60-I69)	43	4.2	84.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	38	3.7	74.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	37	3.6	72.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	27	2.6	53.0
9	Septicemia (A40-A41)	18	1.8	*
10	Influenza and pneumonia (J10-J18)	16	1.6	*
11	Alzheimer's disease (G30)	10	1.0	*
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	10	1.0	*
11	Atherosclerosis (I70)	10	1.0	*
11	Aortic aneurysm and dissection (I71)	10	1.0	*
15	Parkinson's disease (G20-G21)	5	0.5	*
...	All other causes (Residual)	154	15.0	302.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 75-84 years				
...	All causes	1,073	100.0	3,545.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	291	27.1	961.6
2	Malignant neoplasms (C00-C97)	199	18.5	657.6
3	Cerebrovascular diseases (I60-I69)	109	10.2	360.2
4	Diabetes mellitus (E10-E14)	83	7.7	274.3
5	Chronic lower respiratory diseases (J40-J47)	49	4.6	161.9
6	Influenza and pneumonia (J10-J18)	38	3.5	125.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	30	2.8	99.1
8	Alzheimer's disease (G30)	26	2.4	85.9
9	Septicemia (A40-A41)	20	1.9	66.1
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19	1.8	*
11	Chronic liver disease and cirrhosis (K70,K73-K74)	13	1.2	*
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	11	1.0	*
13	Parkinson's disease (G20-G21)	8	0.7	*
14	Atherosclerosis (I70)	7	0.7	*
15	Pneumonitis due to solids and liquids (J69)	6	0.6	*
...	All other causes (Residual)	164	15.3	541.9
American Indian, female, 85 years and over				
...	All causes	881	100.0	5,949.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	304	34.5	2,052.9
2	Malignant neoplasms (C00-C97)	95	10.8	641.5
3	Cerebrovascular diseases (I60-I69)	90	10.2	607.8
4	Influenza and pneumonia (J10-J18)	53	6.0	357.9
5	Chronic lower respiratory diseases (J40-J47)	35	4.0	236.4
6	Diabetes mellitus (E10-E14)	31	3.5	209.3
7	Alzheimer's disease (G30)	30	3.4	202.6
8	Septicemia (A40-A41)	17	1.9	*
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	15	1.7	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	1.6	*
11	Nutritional deficiencies (E40-E64)	13	1.5	*
11	Atherosclerosis (I70)	13	1.5	*
13	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12	1.4	*
14	Pneumonitis due to solids and liquids (J69)	8	0.9	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.8	*
...	All other causes (Residual)	144	16.3	972.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, all ages				
...	All causes	34,875	100.0	309.4
1	Malignant neoplasms (C00-C97)	9,221	26.4	81.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,101	26.1	80.7
3	Cerebrovascular diseases (I60-I69)	3,288	9.4	29.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,678	4.8	14.9
5	Chronic lower respiratory diseases (J40-J47)	1,133	3.2	10.1
6	Influenza and pneumonia (J10-J18)	1,120	3.2	9.9
7	Diabetes mellitus (E10-E14)	1,103	3.2	9.8
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	616	1.8	5.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	527	1.5	4.7
10	Certain conditions originating in the perinatal period (P00-P96)	392	1.1	3.5
11	Assault (homicide) (X85-Y09,Y87.1)	356	1.0	3.2
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	338	1.0	3.0
13	Septicemia (A40-A41)	330	0.9	2.9
14	Chronic liver disease and cirrhosis (K70,K73-K74)	299	0.9	2.7
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	297	0.9	2.6
...	All other causes (Residual)	5,076	14.6	45.0
Asian or Pacific Islander, both sexes, 1-4 years				
...	All causes	146	100.0	19.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	47	32.2	6.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	18	12.3	*
3	Malignant neoplasms (C00-C97)	17	11.6	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	3.4	*
5	Septicemia (A40-A41)	4	2.7	*
5	Influenza and pneumonia (J10-J18)	4	2.7	*
5	Certain conditions originating in the perinatal period (P00-P96)	4	2.7	*
5	Assault (homicide) (X85-Y09,Y87.1)	4	2.7	*
9	Cerebrovascular diseases (I60-I69)	2	1.4	*
9	Complications of medical and surgical care (Y40-Y84,Y88)	2	1.4	*
11	Viral hepatitis (B15-B19)	1	0.7	*
11	Anemias (D50-D64)	1	0.7	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.7	*
...	All other causes (Residual)	36	24.7	4.9

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 5-14 years				
...	All causes	207	100.0	11.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	59	28.5	3.4
2	Malignant neoplasms (C00-C97)	38	18.4	2.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	13	6.3	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	13	6.3	*
5	Chronic lower respiratory diseases (J40-J47)	6	2.9	*
5	Intentional self-harm (suicide) (X60-X84,Y87.0)	6	2.9	*
5	Assault (homicide) (X85-Y09,Y87.1)	6	2.9	*
8	Cerebrovascular diseases (I60-I69)	4	1.9	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	1.4	*
9	Influenza and pneumonia (J10-J18)	3	1.4	*
11	Septicemia (A40-A41)	2	1.0	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	2	1.0	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.5	*
13	Acute bronchitis and bronchiolitis (J20-J21)	1	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.5	*
13	Certain conditions originating in the perinatal period (P00-P96)	1	0.5	*
...	All other causes (Residual)	48	23.2	2.7
Asian or Pacific Islander, both sexes, 15-24 years				
...	All causes	721	100.0	44.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	296	41.1	18.2
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	110	15.3	6.8
3	Assault (homicide) (X85-Y09,Y87.1)	88	12.2	5.4
4	Malignant neoplasms (C00-C97)	66	9.2	4.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	31	4.3	1.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	15	2.1	*
7	Cerebrovascular diseases (I60-I69)	7	1.0	*
8	Septicemia (A40-A41)	5	0.7	*
8	Influenza and pneumonia (J10-J18)	5	0.7	*
10	Tuberculosis (A16-A19)	3	0.4	*
10	Meningococcal infection (A39)	3	0.4	*
10	Chronic lower respiratory diseases (J40-J47)	3	0.4	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.3	*
13	Anemias (D50-D64)	2	0.3	*
13	Pregnancy, childbirth and the puerperium (O00-099)	2	0.3	*
...	All other causes (Residual)	83	11.5	5.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 25-34 years				
...	All causes	897	100.0	46.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	241	26.9	12.3
2	Malignant neoplasms (C00-C97)	169	18.8	8.7
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	154	17.2	7.9
4	Assault (homicide) (X85-Y09,Y87.1)	67	7.5	3.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	60	6.7	3.1
6	Cerebrovascular diseases (I60-I69)	17	1.9	*
7	Human immunodeficiency virus (HIV) disease (B20-B24)	14	1.6	*
8	Chronic lower respiratory diseases (J40-J47)	10	1.1	*
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	1.0	*
10	Diabetes mellitus (E10-E14)	7	0.8	*
10	Chronic liver disease and cirrhosis (K70,K73-K74)	7	0.8	*
12	Aortic aneurysm and dissection (I71)	5	0.6	*
12	Influenza and pneumonia (J10-J18)	5	0.6	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.6	*
12	Pregnancy, childbirth and the puerperium (O00-099)	5	0.6	*
...	All other causes (Residual)	122	13.6	6.2
Asian or Pacific Islander, both sexes, 35-44 years				
...	All causes	1,664	100.0	86.0
1	Malignant neoplasms (C00-C97)	543	32.6	28.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	222	13.3	11.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	207	12.4	10.7
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	115	6.9	5.9
5	Cerebrovascular diseases (I60-I69)	96	5.8	5.0
6	Assault (homicide) (X85-Y09,Y87.1)	72	4.3	3.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	38	2.3	2.0
8	Human immunodeficiency virus (HIV) disease (B20-B24)	30	1.8	1.6
9	Diabetes mellitus (E10-E14)	22	1.3	1.1
10	Viral hepatitis (B15-B19)	21	1.3	1.1
10	Chronic lower respiratory diseases (J40-J47)	21	1.3	1.1
12	Influenza and pneumonia (J10-J18)	16	1.0	*
13	Septicemia (A40-A41)	14	0.8	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	13	0.8	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	0.6	*
...	All other causes (Residual)	224	13.5	11.6

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 45-54 years				
...	All causes	3,024	100.0	208.9
1	Malignant neoplasms (C00-C97)	1,189	39.3	82.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	560	18.5	38.7
3	Cerebrovascular diseases (I60-I69)	214	7.1	14.8
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	192	6.3	13.3
5	Intentional self-harm (suicide) (X60-X84,Y87.0)	96	3.2	6.6
6	Diabetes mellitus (E10-E14)	83	2.7	5.7
7	Viral hepatitis (B15-B19)	71	2.3	4.9
8	Assault (homicide) (X85-Y09,Y87.1)	61	2.0	4.2
9	Chronic liver disease and cirrhosis (K70,K73-K74)	55	1.8	3.8
10	Chronic lower respiratory diseases (J40-J47)	51	1.7	3.5
11	Influenza and pneumonia (J10-J18)	31	1.0	2.1
12	Septicemia (A40-A41)	30	1.0	2.1
13	Aortic aneurysm and dissection (I71)	28	0.9	1.9
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	26	0.9	1.8
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	24	0.8	1.7
...	All other causes (Residual)	313	10.4	21.6
Asian or Pacific Islander, both sexes, 55-64 years				
...	All causes	4,326	100.0	539.7
1	Malignant neoplasms (C00-C97)	1,710	39.5	213.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,039	24.0	129.6
3	Cerebrovascular diseases (I60-I69)	347	8.0	43.3
4	Diabetes mellitus (E10-E14)	158	3.7	19.7
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	139	3.2	17.3
6	Viral hepatitis (B15-B19)	86	2.0	10.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	70	1.6	8.7
8	Chronic lower respiratory diseases (J40-J47)	63	1.5	7.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	61	1.4	7.6
10	Influenza and pneumonia (J10-J18)	59	1.4	7.4
11	Intentional self-harm (suicide) (X60-X84,Y87.0)	51	1.2	6.4
12	Septicemia (A40-A41)	36	0.8	4.5
13	Aortic aneurysm and dissection (I71)	33	0.8	4.1
14	Assault (homicide) (X85-Y09,Y87.1)	31	0.7	3.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	25	0.6	3.1
...	All other causes (Residual)	418	9.7	52.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 65-74 years				
...	All causes	6,887	100.0	1,362.9
1	Malignant neoplasms (C00-C97)	2,386	34.6	472.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,818	26.4	359.8
3	Cerebrovascular diseases (I60-I69)	628	9.1	124.3
4	Diabetes mellitus (E10-E14)	295	4.3	58.4
5	Chronic lower respiratory diseases (J40-J47)	254	3.7	50.3
6	Influenza and pneumonia (J10-J18)	178	2.6	35.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	150	2.2	29.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	128	1.9	25.3
9	Viral hepatitis (B15-B19)	68	1.0	13.5
10	Septicemia (A40-A41)	64	0.9	12.7
10	Chronic liver disease and cirrhosis (K70,K73-K74)	64	0.9	12.7
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	59	0.9	11.7
12	Aortic aneurysm and dissection (I71)	59	0.9	11.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	49	0.7	9.7
15	Intentional self-harm (suicide) (X60-X84,Y87.0)	31	0.5	6.1
...	All other causes (Residual)	656	9.5	129.8
Asian or Pacific Islander, both sexes, 75-84 years				
...	All causes	9,210	100.0	3,607.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,828	30.7	1,107.6
2	Malignant neoplasms (C00-C97)	2,200	23.9	861.6
3	Cerebrovascular diseases (I60-I69)	1,076	11.7	421.4
4	Chronic lower respiratory diseases (J40-J47)	419	4.5	164.1
5	Influenza and pneumonia (J10-J18)	370	4.0	144.9
6	Diabetes mellitus (E10-E14)	358	3.9	140.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	189	2.1	74.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	153	1.7	59.9
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	127	1.4	49.7
10	Aortic aneurysm and dissection (I71)	92	1.0	36.0
11	Septicemia (A40-A41)	91	1.0	35.6
12	Alzheimer's disease (G30)	82	0.9	32.1
13	Parkinson's disease (G20-G21)	75	0.8	29.4
14	Pneumonitis due to solids and liquids (J69)	71	0.8	27.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	52	0.6	20.4
...	All other causes (Residual)	1,027	11.2	402.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 85 years and over				
...	All causes	6,990	100.0	9,376.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,504	35.8	3,359.0
2	Malignant neoplasms (C00-C97)	898	12.8	1,204.6
3	Cerebrovascular diseases (I60-I69)	894	12.8	1,199.3
4	Influenza and pneumonia (J10-J18)	441	6.3	591.6
5	Chronic lower respiratory diseases (J40-J47)	304	4.3	407.8
6	Diabetes mellitus (E10-E14)	180	2.6	241.5
7	Alzheimer's disease (G30)	164	2.3	220.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	140	2.0	187.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	140	2.0	187.8
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	100	1.4	134.1
11	Pneumonitis due to solids and liquids (J69)	97	1.4	130.1
12	Septicemia (A40-A41)	77	1.1	103.3
13	Parkinson's disease (G20-G21)	58	0.8	77.8
14	Atherosclerosis (I70)	55	0.8	73.8
14	Aortic aneurysm and dissection (I71)	55	0.8	73.8
...	All other causes (Residual)	883	12.6	1,184.5
Asian or Pacific Islander, male, all ages				
...	All causes	19,018	100.0	349.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,175	27.2	95.0
2	Malignant neoplasms (C00-C97)	4,865	25.6	89.3
3	Cerebrovascular diseases (I60-I69)	1,555	8.2	28.5
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,057	5.6	19.4
5	Chronic lower respiratory diseases (J40-J47)	722	3.8	13.3
6	Influenza and pneumonia (J10-J18)	592	3.1	10.9
7	Diabetes mellitus (E10-E14)	547	2.9	10.0
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	452	2.4	8.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	254	1.3	4.7
9	Assault (homicide) (X85-Y09,Y87.1)	254	1.3	4.7
11	Certain conditions originating in the perinatal period (P00-P96)	224	1.2	4.1
12	Chronic liver disease and cirrhosis (K70,K73-K74)	184	1.0	3.4
13	Aortic aneurysm and dissection (I71)	172	0.9	3.2
14	Viral hepatitis (B15-B19)	161	0.8	3.0
15	Septicemia (A40-A41)	160	0.8	2.9
...	All other causes (Residual)	2,644	13.9	48.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 1-4 years				
...	All causes	79	100.0	21.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	26	32.9	7.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	10.1	*
3	Malignant neoplasms (C00-C97)	7	8.9	*
4	Septicemia (A40-A41)	3	3.8	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	3.8	*
4	Influenza and pneumonia (J10-J18)	3	3.8	*
4	Certain conditions originating in the perinatal period (P00-P96)	3	3.8	*
4	Assault (homicide) (X85-Y09,Y87.1)	3	3.8	*
9	Viral hepatitis (B15-B19)	1	1.3	*
9	Cerebrovascular diseases (I60-I69)	1	1.3	*
9	Complications of medical and surgical care (Y40-Y84,Y88)	1	1.3	*
...	All other causes (Residual)	20	25.3	5.4
Asian or Pacific Islander, male, 5-14 years				
...	All causes	111	100.0	12.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	32	28.8	3.6
2	Malignant neoplasms (C00-C97)	22	19.8	2.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	7	6.3	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	5.4	*
4	Assault (homicide) (X85-Y09,Y87.1)	6	5.4	*
6	Chronic lower respiratory diseases (J40-J47)	4	3.6	*
7	Intentional self-harm (suicide) (X60-X84,Y87.0)	3	2.7	*
8	Cerebrovascular diseases (I60-I69)	2	1.8	*
8	Influenza and pneumonia (J10-J18)	2	1.8	*
8	Complications of medical and surgical care (Y40-Y84,Y88)	2	1.8	*
11	Septicemia (A40-A41)	1	0.9	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.9	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.9	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.9	*
...	All other causes (Residual)	21	18.9	2.3

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 15-24 years				
...	All causes	516	100.0	63.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	227	44.0	28.0
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	85	16.5	10.5
3	Assault (homicide) (X85-Y09,Y87.1)	73	14.1	9.0
4	Malignant neoplasms (C00-C97)	38	7.4	4.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	24	4.7	3.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	1.4	*
7	Cerebrovascular diseases (I60-I69)	3	0.6	*
8	Tuberculosis (A16-A19)	2	0.4	*
8	Meningococcal infection (A39)	2	0.4	*
8	Septicemia (A40-A41)	2	0.4	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.4	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.2	*
12	Anemias (D50-D64)	1	0.2	*
12	Influenza and pneumonia (J10-J18)	1	0.2	*
12	Chronic lower respiratory diseases (J40-J47)	1	0.2	*
12	Pneumonitis due to solids and liquids (J69)	1	0.2	*
12	Diseases of appendix (K35-K38)	1	0.2	*
12	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.2	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
...	All other causes (Residual)	43	8.3	5.3
Asian or Pacific Islander, male, 25-34 years				
...	All causes	590	100.0	62.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	178	30.2	19.0
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	114	19.3	12.2
3	Malignant neoplasms (C00-C97)	79	13.4	8.4
4	Assault (homicide) (X85-Y09,Y87.1)	48	8.1	5.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	43	7.3	4.6
6	Human immunodeficiency virus (HIV) disease (B20-B24)	13	2.2	*
7	Cerebrovascular diseases (I60-I69)	12	2.0	*
8	Chronic lower respiratory diseases (J40-J47)	6	1.0	*
8	Chronic liver disease and cirrhosis (K70,K73-K74)	6	1.0	*
10	Diabetes mellitus (E10-E14)	4	0.7	*
10	Influenza and pneumonia (J10-J18)	4	0.7	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.7	*
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	0.7	*
10	Legal intervention (Y35,Y89.0)	4	0.7	*
15	Viral hepatitis (B15-B19)	3	0.5	*
15	Aortic aneurysm and dissection (I71)	3	0.5	*
...	All other causes (Residual)	65	11.0	6.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 35-44 years				
...	All causes	993	100.0	106.8
1	Malignant neoplasms (C00-C97)	247	24.9	26.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	174	17.5	18.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	138	13.9	14.8
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	85	8.6	9.1
5	Cerebrovascular diseases (I60-I69)	47	4.7	5.1
6	Assault (homicide) (X85-Y09,Y87.1)	45	4.5	4.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	31	3.1	3.3
8	Human immunodeficiency virus (HIV) disease (B20-B24)	25	2.5	2.7
9	Viral hepatitis (B15-B19)	15	1.5	*
9	Diabetes mellitus (E10-E14)	15	1.5	*
11	Chronic lower respiratory diseases (J40-J47)	11	1.1	*
12	Septicemia (A40-A41)	8	0.8	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.8	*
14	Aortic aneurysm and dissection (I71)	6	0.6	*
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	5	0.5	*
15	Influenza and pneumonia (J10-J18)	5	0.5	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	0.5	*
...	All other causes (Residual)	123	12.4	13.2
Asian or Pacific Islander, male, 45-54 years				
...	All causes	1,761	100.0	260.6
1	Malignant neoplasms (C00-C97)	554	31.5	82.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	431	24.5	63.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	116	6.6	17.2
4	Cerebrovascular diseases (I60-I69)	106	6.0	15.7
5	Intentional self-harm (suicide) (X60-X84,Y87.0)	72	4.1	10.7
6	Diabetes mellitus (E10-E14)	49	2.8	7.3
7	Viral hepatitis (B15-B19)	45	2.6	6.7
8	Assault (homicide) (X85-Y09,Y87.1)	44	2.5	6.5
9	Chronic liver disease and cirrhosis (K70,K73-K74)	40	2.3	5.9
10	Chronic lower respiratory diseases (J40-J47)	34	1.9	5.0
11	Influenza and pneumonia (J10-J18)	19	1.1	*
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	18	1.0	*
13	Aortic aneurysm and dissection (I71)	16	0.9	*
14	Septicemia (A40-A41)	15	0.9	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	0.9	*
...	All other causes (Residual)	187	10.6	27.7

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 55-64 years				
...	All causes	2,565	100.0	684.4
1	Malignant neoplasms (C00-C97)	915	35.7	244.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	729	28.4	194.5
3	Cerebrovascular diseases (I60-I69)	197	7.7	52.6
4	Diabetes mellitus (E10-E14)	90	3.5	24.0
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	85	3.3	22.7
6	Viral hepatitis (B15-B19)	52	2.0	13.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	45	1.8	12.0
8	Influenza and pneumonia (J10-J18)	35	1.4	9.3
8	Chronic lower respiratory diseases (J40-J47)	35	1.4	9.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	35	1.4	9.3
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	35	1.4	9.3
12	Aortic aneurysm and dissection (I71)	29	1.1	7.7
13	Assault (homicide) (X85-Y09,Y87.1)	23	0.9	6.1
14	Septicemia (A40-A41)	15	0.6	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	0.5	*
...	All other causes (Residual)	231	9.0	61.6
Asian or Pacific Islander, male, 65-74 years				
...	All causes	3,845	100.0	1,800.3
1	Malignant neoplasms (C00-C97)	1,295	33.7	606.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,117	29.1	523.0
3	Cerebrovascular diseases (I60-I69)	314	8.2	147.0
4	Chronic lower respiratory diseases (J40-J47)	165	4.3	77.3
5	Diabetes mellitus (E10-E14)	153	4.0	71.6
6	Influenza and pneumonia (J10-J18)	99	2.6	46.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	83	2.2	38.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	51	1.3	23.9
9	Aortic aneurysm and dissection (I71)	35	0.9	16.4
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	34	0.9	15.9
11	Septicemia (A40-A41)	33	0.9	15.5
12	Chronic liver disease and cirrhosis (K70,K73-K74)	32	0.8	15.0
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	31	0.8	14.5
14	Viral hepatitis (B15-B19)	30	0.8	14.0
15	Intentional self-harm (suicide) (X60-X84,Y87.0)	20	0.5	9.4
...	All other causes (Residual)	353	9.2	165.3

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 75-84 years				
...	All causes	4,834	100.0	4,510.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,513	31.3	1,411.7
2	Malignant neoplasms (C00-C97)	1,213	25.1	1,131.8
3	Cerebrovascular diseases (I60-I69)	490	10.1	457.2
4	Chronic lower respiratory diseases (J40-J47)	284	5.9	265.0
5	Influenza and pneumonia (J10-J18)	203	4.2	189.4
6	Diabetes mellitus (E10-E14)	156	3.2	145.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	100	2.1	93.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	79	1.6	73.7
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	56	1.2	52.3
10	Aortic aneurysm and dissection (I71)	54	1.1	50.4
11	Parkinson's disease (G20-G21)	53	1.1	49.5
12	Septicemia (A40-A41)	47	1.0	43.9
13	Alzheimer's disease (G30)	39	0.8	36.4
14	Pneumonitis due to solids and liquids (J69)	38	0.8	35.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	29	0.6	27.1
...	All other causes (Residual)	480	9.9	447.9
Asian or Pacific Islander, male, 85 years and over				
...	All causes	3,272	100.0	10,894.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,121	34.3	3,732.3
2	Malignant neoplasms (C00-C97)	492	15.0	1,638.1
3	Cerebrovascular diseases (I60-I69)	382	11.7	1,271.8
4	Influenza and pneumonia (J10-J18)	214	6.5	712.5
5	Chronic lower respiratory diseases (J40-J47)	181	5.5	602.6
6	Diabetes mellitus (E10-E14)	80	2.4	266.4
7	Alzheimer's disease (G30)	65	2.0	216.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	62	1.9	206.4
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	62	1.9	206.4
10	Pneumonitis due to solids and liquids (J69)	49	1.5	163.1
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	38	1.2	126.5
12	Parkinson's disease (G20-G21)	36	1.1	119.9
13	Septicemia (A40-A41)	34	1.0	113.2
14	Aortic aneurysm and dissection (I71)	29	0.9	96.6
15	Atherosclerosis (I70)	23	0.7	76.6
...	All other causes (Residual)	404	12.3	1,345.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, all ages				
...	All causes	15,857	100.0	272.2
1	Malignant neoplasms (C00-C97)	4,356	27.5	74.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,926	24.8	67.4
3	Cerebrovascular diseases (I60-I69)	1,733	10.9	29.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	621	3.9	10.7
5	Diabetes mellitus (E10-E14)	556	3.5	9.5
6	Influenza and pneumonia (J10-J18)	528	3.3	9.1
7	Chronic lower respiratory diseases (J40-J47)	411	2.6	7.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	273	1.7	4.7
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	179	1.1	3.1
10	Septicemia (A40-A41)	170	1.1	2.9
11	Certain conditions originating in the perinatal period (P00-P96)	168	1.1	2.9
12	Intentional self-harm (suicide) (X60-X84,Y87.0)	164	1.0	2.8
13	Alzheimer's disease (G30)	163	1.0	2.8
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	148	0.9	2.5
15	Viral hepatitis (B15-B19)	124	0.8	2.1
...	All other causes (Residual)	2,337	14.7	40.1
Asian or Pacific Islander, female, 1-4 years				
...	All causes	67	100.0	18.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	21	31.3	5.8
2	Malignant neoplasms (C00-C97)	10	14.9	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	10	14.9	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	3.0	*
5	Septicemia (A40-A41)	1	1.5	*
5	Anemias (D50-D64)	1	1.5	*
5	Cerebrovascular diseases (I60-I69)	1	1.5	*
5	Influenza and pneumonia (J10-J18)	1	1.5	*
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	1.5	*
5	Certain conditions originating in the perinatal period (P00-P96)	1	1.5	*
5	Assault (homicide) (X85-Y09,Y87.1)	1	1.5	*
5	Complications of medical and surgical care (Y40-Y84,Y88)	1	1.5	*
...	All other causes (Residual)	16	23.9	*

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 5-14 years				
...	All causes	96	100.0	11.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	27	28.1	3.2
2	Malignant neoplasms (C00-C97)	16	16.7	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	7.3	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	6	6.3	*
5	Intentional self-harm (suicide) (X60-X84,Y87.0)	3	3.1	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	2.1	*
6	Cerebrovascular diseases (I60-I69)	2	2.1	*
6	Chronic lower respiratory diseases (J40-J47)	2	2.1	*
9	Septicemia (A40-A41)	1	1.0	*
9	Influenza and pneumonia (J10-J18)	1	1.0	*
9	Acute bronchitis and bronchiolitis (J20-J21)	1	1.0	*
9	Certain conditions originating in the perinatal period (P00-P96)	1	1.0	*
...	All other causes (Residual)	27	28.1	3.2
Asian or Pacific Islander, female, 15-24 years				
...	All causes	205	100.0	25.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	69	33.7	8.5
2	Malignant neoplasms (C00-C97)	28	13.7	3.4
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	25	12.2	3.1
4	Assault (homicide) (X85-Y09,Y87.1)	15	7.3	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	3.9	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	7	3.4	*
7	Cerebrovascular diseases (I60-I69)	4	2.0	*
7	Influenza and pneumonia (J10-J18)	4	2.0	*
9	Septicemia (A40-A41)	3	1.5	*
10	Chronic lower respiratory diseases (J40-J47)	2	1.0	*
10	Pregnancy, childbirth and the puerperium (O00-099)	2	1.0	*
12	Tuberculosis (A16-A19)	1	0.5	*
12	Meningococcal infection (A39)	1	0.5	*
12	Anemias (D50-D64)	1	0.5	*
12	Aortic aneurysm and dissection (I71)	1	0.5	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.5	*
...	All other causes (Residual)	33	16.1	4.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 25-34 years				
...	All causes	307	100.0	30.3
1	Malignant neoplasms (C00-C97)	90	29.3	8.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	63	20.5	6.2
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	40	13.0	3.9
4	Assault (homicide) (X85-Y09,Y87.1)	19	6.2	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	17	5.5	*
6	Cerebrovascular diseases (I60-I69)	5	1.6	*
6	Pregnancy, childbirth and the puerperium (O00-099)	5	1.6	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	1.6	*
9	Chronic lower respiratory diseases (J40-J47)	4	1.3	*
10	Diabetes mellitus (E10-E14)	3	1.0	*
11	Septicemia (A40-A41)	2	0.7	*
11	Aortic aneurysm and dissection (I71)	2	0.7	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.3	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.3	*
13	Nutritional deficiencies (E40-E64)	1	0.3	*
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1	0.3	*
13	Influenza and pneumonia (J10-J18)	1	0.3	*
13	Acute bronchitis and bronchiolitis (J20-J21)	1	0.3	*
13	Peptic ulcer (K25-K28)	1	0.3	*
13	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.3	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.3	*
...	All other causes (Residual)	42	13.7	4.1
Asian or Pacific Islander, female, 35-44 years				
...	All causes	671	100.0	66.7
1	Malignant neoplasms (C00-C97)	296	44.1	29.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	69	10.3	6.9
3	Cerebrovascular diseases (I60-I69)	49	7.3	4.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	48	7.2	4.8
5	Intentional self-harm (suicide) (X60-X84,Y87.0)	30	4.5	3.0
6	Assault (homicide) (X85-Y09,Y87.1)	27	4.0	2.7
7	Influenza and pneumonia (J10-J18)	11	1.6	*
8	Chronic lower respiratory diseases (J40-J47)	10	1.5	*
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	1.2	*
10	Diabetes mellitus (E10-E14)	7	1.0	*
10	Chronic liver disease and cirrhosis (K70,K73-K74)	7	1.0	*
12	Septicemia (A40-A41)	6	0.9	*
12	Viral hepatitis (B15-B19)	6	0.9	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.9	*
12	Pregnancy, childbirth and the puerperium (O00-099)	6	0.9	*
...	All other causes (Residual)	85	12.7	8.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 45-54 years				
...	All causes	1,263	100.0	163.6
1	Malignant neoplasms (C00-C97)	635	50.3	82.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	129	10.2	16.7
3	Cerebrovascular diseases (I60-I69)	108	8.6	14.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	76	6.0	9.8
5	Diabetes mellitus (E10-E14)	34	2.7	4.4
6	Viral hepatitis (B15-B19)	26	2.1	3.4
7	Intentional self-harm (suicide) (X60-X84,Y87.0)	24	1.9	3.1
8	Chronic lower respiratory diseases (J40-J47)	17	1.3	*
8	Assault (homicide) (X85-Y09,Y87.1)	17	1.3	*
10	Septicemia (A40-A41)	15	1.2	*
10	Chronic liver disease and cirrhosis (K70,K73-K74)	15	1.2	*
12	Aortic aneurysm and dissection (I71)	12	1.0	*
12	Influenza and pneumonia (J10-J18)	12	1.0	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.7	*
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	8	0.6	*
...	All other causes (Residual)	126	10.0	16.3
Asian or Pacific Islander, female, 55-64 years				
...	All causes	1,761	100.0	412.7
1	Malignant neoplasms (C00-C97)	795	45.1	186.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	310	17.6	72.6
3	Cerebrovascular diseases (I60-I69)	150	8.5	35.2
4	Diabetes mellitus (E10-E14)	68	3.9	15.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	54	3.1	12.7
6	Viral hepatitis (B15-B19)	34	1.9	8.0
7	Chronic lower respiratory diseases (J40-J47)	28	1.6	6.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	26	1.5	6.1
9	Chronic liver disease and cirrhosis (K70,K73-K74)	25	1.4	5.9
10	Influenza and pneumonia (J10-J18)	24	1.4	5.6
11	Septicemia (A40-A41)	21	1.2	4.9
12	Intentional self-harm (suicide) (X60-X84,Y87.0)	16	0.9	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.6	*
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	9	0.5	*
15	Assault (homicide) (X85-Y09,Y87.1)	8	0.5	*
...	All other causes (Residual)	182	10.3	42.7

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 65-74 years				
...	All causes	3,042	100.0	1,042.8
1	Malignant neoplasms (C00-C97)	1,091	35.9	374.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	701	23.0	240.3
3	Cerebrovascular diseases (I60-I69)	314	10.3	107.6
4	Diabetes mellitus (E10-E14)	142	4.7	48.7
5	Chronic lower respiratory diseases (J40-J47)	89	2.9	30.5
6	Influenza and pneumonia (J10-J18)	79	2.6	27.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	77	2.5	26.4
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	67	2.2	23.0
9	Viral hepatitis (B15-B19)	38	1.2	13.0
10	Chronic liver disease and cirrhosis (K70,K73-K74)	32	1.1	11.0
11	Septicemia (A40-A41)	31	1.0	10.6
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	25	0.8	8.6
13	Aortic aneurysm and dissection (I71)	24	0.8	8.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	0.6	*
15	Alzheimer's disease (G30)	12	0.4	*
...	All other causes (Residual)	302	9.9	103.5
Asian or Pacific Islander, female, 75-84 years				
...	All causes	4,376	100.0	2,953.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,315	30.1	887.5
2	Malignant neoplasms (C00-C97)	987	22.6	666.2
3	Cerebrovascular diseases (I60-I69)	586	13.4	395.5
4	Diabetes mellitus (E10-E14)	202	4.6	136.3
5	Influenza and pneumonia (J10-J18)	167	3.8	112.7
6	Chronic lower respiratory diseases (J40-J47)	135	3.1	91.1
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	89	2.0	60.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	74	1.7	49.9
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	71	1.6	47.9
10	Septicemia (A40-A41)	44	1.0	29.7
11	Alzheimer's disease (G30)	43	1.0	29.0
12	Aortic aneurysm and dissection (I71)	38	0.9	25.6
13	Pneumonitis due to solids and liquids (J69)	33	0.8	22.3
14	Chronic liver disease and cirrhosis (K70,K73-K74)	29	0.7	19.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	0.5	15.5
...	All other causes (Residual)	540	12.3	364.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
	Asian or Pacific Islander, female, 85 years and over			
...	All causes	3,718	100.0	8,353.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,383	37.2	3,107.1
2	Cerebrovascular diseases (I60-I69)	512	13.8	1,150.3
3	Malignant neoplasms (C00-C97)	406	10.9	912.1
4	Influenza and pneumonia (J10-J18)	227	6.1	510.0
5	Chronic lower respiratory diseases (J40-J47)	123	3.3	276.3
6	Diabetes mellitus (E10-E14)	100	2.7	224.7
7	Alzheimer's disease (G30)	99	2.7	222.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	78	2.1	175.2
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	78	2.1	175.2
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	62	1.7	139.3
11	Pneumonitis due to solids and liquids (J69)	48	1.3	107.8
12	Septicemia (A40-A41)	43	1.2	96.6
13	Atherosclerosis (I70)	32	0.9	71.9
14	Aortic aneurysm and dissection (I71)	26	0.7	58.4
15	Parkinson's disease (G20-G21)	22	0.6	49.4
...	All other causes (Residual)	479	12.9	1,076.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable