

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Percent of		
		Number	total deaths	Rate
All races, both sexes, all ages				
...	All causes	2,416,425	100.0	848.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	700,142	29.0	245.8
2	Malignant neoplasms (C00-C97)	553,768	22.9	194.4
3	Cerebrovascular diseases (I60-I69)	163,538	6.8	57.4
4	Chronic lower respiratory diseases (J40-J47)	123,013	5.1	43.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	101,537	4.2	35.7
6	Diabetes mellitus (E10-E14)	71,372	3.0	25.1
7	Influenza and pneumonia (J10-J18)	62,034	2.6	21.8
8	Alzheimer's disease (G30)	53,852	2.2	18.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	39,480	1.6	13.9
10	Septicemia (A40-A41)	32,238	1.3	11.3
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	30,622	1.3	10.8
12	Chronic liver disease and cirrhosis (K70,K73-K74)	27,035	1.1	9.5
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	20,308	0.8	7.1
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	19,250	0.8	6.8
15	Pneumonitis due to solids and liquids (J69)	17,301	0.7	6.1
...	All other causes (Residual)	400,935	16.6	140.8
All races, both sexes, 1-4 years				
...	All causes	5,107	100.0	33.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,714	33.6	11.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	557	10.9	3.6
3	Malignant neoplasms (C00-C97)	420	8.2	2.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	415	8.1	2.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	225	4.4	1.5
6	Influenza and pneumonia (J10-J18)	112	2.2	0.7
7	Septicemia (A40-A41)	108	2.1	0.7
8	Certain conditions originating in the perinatal period (P00-P96)	72	1.4	0.5
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	58	1.1	0.4
10	Cerebrovascular diseases (I60-I69)	54	1.1	0.4
11	Chronic lower respiratory diseases (J40-J47)	43	0.8	0.3
12	Meningitis (G00,G03)	26	0.5	0.2
13	Complications of medical and surgical care (Y40-Y84,Y88)	24	0.5	0.2
14	Anemias (D50-D64)	22	0.4	0.1
15	Meningococcal infection (A39)	19	0.4	*
...	All other causes (Residual)	1,238	24.2	8.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Percent of		
		Number	total deaths	Rate
All races, both sexes, 5-14 years				
...	All causes	7,095	100.0	17.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,836	40.0	6.9
2	Malignant neoplasms (C00-C97)	1,008	14.2	2.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	376	5.3	0.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	326	4.6	0.8
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	279	3.9	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	272	3.8	0.7
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	105	1.5	0.3
8	Chronic lower respiratory diseases (J40-J47)	104	1.5	0.3
9	Influenza and pneumonia (J10-J18)	92	1.3	0.2
10	Cerebrovascular diseases (I60-I69)	80	1.1	0.2
11	Septicemia (A40-A41)	69	1.0	0.2
12	Anemias (D50-D64)	55	0.8	0.1
13	Diabetes mellitus (E10-E14)	39	0.5	0.1
14	Human immunodeficiency virus (HIV) disease (B20-B24)	38	0.5	0.1
15	Certain conditions originating in the perinatal period (P00-P96)	36	0.5	0.1
...	All other causes (Residual)	1,380	19.5	3.4
All races, both sexes, 15-24 years				
...	All causes	32,252	100.0	80.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14,411	44.7	36.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5,297	16.4	13.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,971	12.3	9.9
4	Malignant neoplasms (C00-C97)	1,704	5.3	4.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	999	3.1	2.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	505	1.6	1.3
7	Human immunodeficiency virus (HIV) disease (B20-B24)	225	0.7	0.6
8	Cerebrovascular diseases (I60-I69)	196	0.6	0.5
9	Influenza and pneumonia (J10-J18)	181	0.6	0.5
10	Chronic lower respiratory diseases (J40-J47)	171	0.5	0.4
11	Diabetes mellitus (E10-E14)	151	0.5	0.4
12	Septicemia (A40-A41)	133	0.4	0.3
13	Pregnancy, childbirth and the puerperium (O00-099)	110	0.3	0.3
14	Anemias (D50-D64)	109	0.3	0.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	102	0.3	0.3
...	All other causes (Residual)	3,987	12.4	10.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Percent of		
		Number	total deaths	Rate
All races, both sexes, 25-34 years				
...	All causes	41,683	100.0	105.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,839	28.4	29.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5,204	12.5	13.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,070	12.2	12.8
4	Malignant neoplasms (C00-C97)	3,994	9.6	10.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,160	7.6	8.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	2,101	5.0	5.3
7	Cerebrovascular diseases (I60-I69)	601	1.4	1.5
8	Diabetes mellitus (E10-E14)	595	1.4	1.5
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	458	1.1	1.2
10	Chronic liver disease and cirrhosis (K70,K73-K74)	387	0.9	1.0
11	Influenza and pneumonia (J10-J18)	339	0.8	0.9
12	Chronic lower respiratory diseases (J40-J47)	290	0.7	0.7
13	Septicemia (A40-A41)	281	0.7	0.7
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	225	0.5	0.6
15	Pregnancy, childbirth and the puerperium (O00-099)	190	0.5	0.5
...	All other causes (Residual)	6,949	16.7	17.5
All races, both sexes, 35-44 years				
...	All causes	91,674	100.0	203.6
1	Malignant neoplasms (C00-C97)	16,569	18.1	36.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15,945	17.4	35.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,326	14.5	29.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,635	7.2	14.7
5	Human immunodeficiency virus (HIV) disease (B20-B24)	5,867	6.4	13.0
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4,268	4.7	9.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	3,336	3.6	7.4
8	Cerebrovascular diseases (I60-I69)	2,491	2.7	5.5
9	Diabetes mellitus (E10-E14)	1,958	2.1	4.3
10	Influenza and pneumonia (J10-J18)	983	1.1	2.2
11	Chronic lower respiratory diseases (J40-J47)	969	1.1	2.2
12	Septicemia (A40-A41)	829	0.9	1.8
13	Viral hepatitis (B15-B19)	797	0.9	1.8
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	768	0.8	1.7
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	586	0.6	1.3
...	All other causes (Residual)	16,347	17.8	36.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Percent of		
		Number	total deaths	Rate
All races, both sexes, 45-54 years				
...	All causes	168,065	100.0	428.9
1	Malignant neoplasms (C00-C97)	49,562	29.5	126.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	36,399	21.7	92.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,344	7.9	34.1
4	Chronic liver disease and cirrhosis (K70,K73-K74)	7,259	4.3	18.5
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,942	3.5	15.2
6	Cerebrovascular diseases (I60-I69)	5,910	3.5	15.1
7	Diabetes mellitus (E10-E14)	5,343	3.2	13.6
8	Human immunodeficiency virus (HIV) disease (B20-B24)	4,120	2.5	10.5
9	Chronic lower respiratory diseases (J40-J47)	3,324	2.0	8.5
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,467	1.5	6.3
11	Viral hepatitis (B15-B19)	2,060	1.2	5.3
12	Septicemia (A40-A41)	1,977	1.2	5.0
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,822	1.1	4.6
14	Influenza and pneumonia (J10-J18)	1,801	1.1	4.6
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	931	0.6	2.4
...	All other causes (Residual)	25,804	15.4	65.8
All races, both sexes, 55-64 years				
...	All causes	244,139	100.0	964.6
1	Malignant neoplasms (C00-C97)	90,223	37.0	356.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	62,486	25.6	246.9
3	Chronic lower respiratory diseases (J40-J47)	11,166	4.6	44.1
4	Cerebrovascular diseases (I60-I69)	9,608	3.9	38.0
5	Diabetes mellitus (E10-E14)	9,570	3.9	37.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,658	3.1	30.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	5,750	2.4	22.7
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,317	1.4	13.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,284	1.3	13.0
10	Septicemia (A40-A41)	3,111	1.3	12.3
11	Influenza and pneumonia (J10-J18)	2,704	1.1	10.7
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,462	0.6	5.8
13	Aortic aneurysm and dissection (I71)	1,394	0.6	5.5
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1,317	0.5	5.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,253	0.5	5.0
...	All other causes (Residual)	29,836	12.2	117.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Percent of total deaths		
		Number	Rate	
All races, both sexes, 65-74 years				
...	All causes	430,960	100.0	2,353.3
1	Malignant neoplasms (C00-C97)	147,018	34.1	802.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	116,299	27.0	635.1
3	Chronic lower respiratory diseases (J40-J47)	30,751	7.1	167.9
4	Cerebrovascular diseases (I60-I69)	22,598	5.2	123.4
5	Diabetes mellitus (E10-E14)	16,731	3.9	91.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,835	1.8	42.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,356	1.7	40.2
8	Influenza and pneumonia (J10-J18)	6,650	1.5	36.3
9	Septicemia (A40-A41)	5,998	1.4	32.8
10	Chronic liver disease and cirrhosis (K70,K73-K74)	5,486	1.3	30.0
11	Aortic aneurysm and dissection (I71)	3,517	0.8	19.2
12	Alzheimer's disease (G30)	3,422	0.8	18.7
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,832	0.7	15.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,574	0.6	14.1
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,432	0.6	13.3
...	All other causes (Residual)	49,461	11.5	270.1
All races, both sexes, 75-84 years				
...	All causes	701,929	100.0	5,582.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	216,992	30.9	1,725.7
2	Malignant neoplasms (C00-C97)	165,445	23.6	1,315.8
3	Cerebrovascular diseases (I60-I69)	55,822	8.0	443.9
4	Chronic lower respiratory diseases (J40-J47)	47,762	6.8	379.8
5	Diabetes mellitus (E10-E14)	22,805	3.2	181.4
6	Influenza and pneumonia (J10-J18)	18,677	2.7	148.5
7	Alzheimer's disease (G30)	18,542	2.6	147.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13,097	1.9	104.2
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,688	1.8	100.9
10	Septicemia (A40-A41)	10,351	1.5	82.3
11	Parkinson's disease (G20-G21)	8,125	1.2	64.6
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	5,993	0.9	47.7
13	Pneumonitis due to solids and liquids (J69)	5,759	0.8	45.8
14	Aortic aneurysm and dissection (I71)	5,751	0.8	45.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,654	0.7	37.0
...	All other causes (Residual)	89,466	12.7	711.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Percent of		
		Number	total deaths	Rate
All races, both sexes, 85 years and over				
...	All causes	665,531	100.0	15,112.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	249,439	37.5	5,664.2
2	Malignant neoplasms (C00-C97)	77,751	11.7	1,765.6
3	Cerebrovascular diseases (I60-I69)	66,066	9.9	1,500.2
4	Alzheimer's disease (G30)	31,281	4.7	710.3
5	Influenza and pneumonia (J10-J18)	30,191	4.5	685.6
6	Chronic lower respiratory diseases (J40-J47)	28,391	4.3	644.7
7	Diabetes mellitus (E10-E14)	14,171	2.1	321.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12,668	1.9	287.7
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,171	1.8	276.4
10	Septicemia (A40-A41)	9,069	1.4	205.9
11	Pneumonitis due to solids and liquids (J69)	8,342	1.3	189.4
12	Atherosclerosis (I70)	7,888	1.2	179.1
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	7,572	1.1	171.9
14	Parkinson's disease (G20-G21)	5,909	0.9	134.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3,643	0.5	82.7
...	All other causes (Residual)	100,979	15.2	2,293.0
All races, male, all ages				
...	All causes	1,183,421	100.0	846.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	339,095	28.7	242.5
2	Malignant neoplasms (C00-C97)	287,075	24.3	205.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	66,060	5.6	47.2
4	Cerebrovascular diseases (I60-I69)	63,177	5.3	45.2
5	Chronic lower respiratory diseases (J40-J47)	59,697	5.0	42.7
6	Diabetes mellitus (E10-E14)	32,841	2.8	23.5
7	Influenza and pneumonia (J10-J18)	27,342	2.3	19.6
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	24,672	2.1	17.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18,852	1.6	13.5
10	Chronic liver disease and cirrhosis (K70,K73-K74)	17,393	1.5	12.4
11	Alzheimer's disease (G30)	15,762	1.3	11.3
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	15,555	1.3	11.1
13	Septicemia (A40-A41)	14,307	1.2	10.2
14	Human immunodeficiency virus (HIV) disease (B20-B24)	10,590	0.9	7.6
15	Parkinson's disease (G20-G21)	9,405	0.8	6.7
...	All other causes (Residual)	181,598	15.3	129.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Percent of		
		Number	total deaths	Rate
All races, male, 1-4 years				
...	All causes	2,899	100.0	37.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,045	36.0	13.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	282	9.7	3.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	238	8.2	3.0
4	Malignant neoplasms (C00-C97)	229	7.9	2.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	121	4.2	1.5
6	Septicemia (A40-A41)	60	2.1	0.8
7	Influenza and pneumonia (J10-J18)	57	2.0	0.7
8	Certain conditions originating in the perinatal period (P00-P96)	44	1.5	0.6
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	30	1.0	0.4
10	Chronic lower respiratory diseases (J40-J47)	29	1.0	0.4
11	Cerebrovascular diseases (I60-I69)	25	0.9	0.3
12	Meningitis (G00,G03)	16	0.6	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	14	0.5	*
14	Anemias (D50-D64)	12	0.4	*
15	Meningococcal infection (A39)	9	0.3	*
...	All other causes (Residual)	688	23.7	8.8
All races, male, 5-14 years				
...	All causes	4,168	100.0	19.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,803	43.3	8.6
2	Malignant neoplasms (C00-C97)	533	12.8	2.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	214	5.1	1.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	210	5.0	1.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	193	4.6	0.9
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	138	3.3	0.7
7	Chronic lower respiratory diseases (J40-J47)	61	1.5	0.3
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	52	1.2	0.2
9	Influenza and pneumonia (J10-J18)	47	1.1	0.2
10	Cerebrovascular diseases (I60-I69)	42	1.0	0.2
11	Septicemia (A40-A41)	39	0.9	0.2
12	Anemias (D50-D64)	27	0.6	0.1
13	Certain conditions originating in the perinatal period (P00-P96)	24	0.6	0.1
14	Human immunodeficiency virus (HIV) disease (B20-B24)	23	0.6	0.1
14	Diabetes mellitus (E10-E14)	23	0.6	0.1
...	All other causes (Residual)	739	17.7	3.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Percent of		
		Number	total deaths	Rate
All races, male, 15-24 years				
...	All causes	23,963	100.0	117.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,832	45.2	52.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4,541	19.0	22.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,409	14.2	16.6
4	Malignant neoplasms (C00-C97)	1,024	4.3	5.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	647	2.7	3.2
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	306	1.3	1.5
7	Influenza and pneumonia (J10-J18)	106	0.4	0.5
8	Human immunodeficiency virus (HIV) disease (B20-B24)	105	0.4	0.5
9	Cerebrovascular diseases (I60-I69)	103	0.4	0.5
10	Chronic lower respiratory diseases (J40-J47)	99	0.4	0.5
11	Diabetes mellitus (E10-E14)	85	0.4	0.4
12	Legal intervention (Y35,Y89.0)	84	0.4	0.4
13	Septicemia (A40-A41)	75	0.3	0.4
14	Anemias (D50-D64)	59	0.2	0.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	53	0.2	0.3
...	All other causes (Residual)	2,435	10.2	11.9
All races, male, 25-34 years				
...	All causes	28,757	100.0	143.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,092	31.6	45.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,199	14.6	21.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4,157	14.5	20.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,068	7.2	10.3
5	Malignant neoplasms (C00-C97)	1,860	6.5	9.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	1,421	4.9	7.1
7	Diabetes mellitus (E10-E14)	341	1.2	1.7
8	Cerebrovascular diseases (I60-I69)	311	1.1	1.6
9	Chronic liver disease and cirrhosis (K70,K73-K74)	259	0.9	1.3
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	249	0.9	1.2
11	Influenza and pneumonia (J10-J18)	191	0.7	1.0
12	Chronic lower respiratory diseases (J40-J47)	146	0.5	0.7
13	Septicemia (A40-A41)	139	0.5	0.7
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	112	0.4	0.6
15	Legal intervention (Y35,Y89.0)	104	0.4	0.5
...	All other causes (Residual)	4,108	14.3	20.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Percent of		
		Number	total deaths	Rate
All races, male, 35-44 years				
...	All causes	58,164	100.0	259.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,612	20.0	51.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,346	16.1	41.7
3	Malignant neoplasms (C00-C97)	7,297	12.5	32.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,180	8.9	23.1
5	Human immunodeficiency virus (HIV) disease (B20-B24)	4,360	7.5	19.5
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,122	5.4	13.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,206	3.8	9.8
8	Cerebrovascular diseases (I60-I69)	1,267	2.2	5.7
9	Diabetes mellitus (E10-E14)	1,181	2.0	5.3
10	Influenza and pneumonia (J10-J18)	556	1.0	2.5
11	Viral hepatitis (B15-B19)	553	1.0	2.5
12	Chronic lower respiratory diseases (J40-J47)	455	0.8	2.0
13	Septicemia (A40-A41)	451	0.8	2.0
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	414	0.7	1.8
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	317	0.5	1.4
...	All other causes (Residual)	9,847	16.9	44.0
All races, male, 45-54 years				
...	All causes	104,848	100.0	545.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	26,280	25.1	136.6
2	Malignant neoplasms (C00-C97)	25,071	23.9	130.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,743	9.3	50.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	5,295	5.1	27.5
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,504	4.3	23.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	3,241	3.1	16.8
7	Cerebrovascular diseases (I60-I69)	3,205	3.1	16.7
8	Diabetes mellitus (E10-E14)	3,154	3.0	16.4
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,832	1.7	9.5
10	Chronic lower respiratory diseases (J40-J47)	1,702	1.6	8.8
11	Viral hepatitis (B15-B19)	1,570	1.5	8.2
12	Influenza and pneumonia (J10-J18)	1,099	1.0	5.7
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,053	1.0	5.5
14	Septicemia (A40-A41)	1,043	1.0	5.4
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	569	0.5	3.0
...	All other causes (Residual)	15,487	14.8	80.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Percent of		
		Number	total deaths	Rate
All races, male, 55-64 years				
...	All causes	144,958	100.0	1,192.7
1	Malignant neoplasms (C00-C97)	49,244	34.0	405.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	42,519	29.3	349.8
3	Chronic lower respiratory diseases (J40-J47)	5,702	3.9	46.9
4	Cerebrovascular diseases (I60-I69)	5,277	3.6	43.4
5	Diabetes mellitus (E10-E14)	5,241	3.6	43.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,228	3.6	43.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	4,028	2.8	33.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,563	1.8	21.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,714	1.2	14.1
10	Septicemia (A40-A41)	1,606	1.1	13.2
11	Influenza and pneumonia (J10-J18)	1,587	1.1	13.1
12	Human immunodeficiency virus (HIV) disease (B20-B24)	1,051	0.7	8.6
13	Aortic aneurysm and dissection (I71)	1,031	0.7	8.5
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	784	0.5	6.5
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	723	0.5	5.9
...	All other causes (Residual)	16,660	11.5	137.1
All races, male, 65-74 years				
...	All causes	241,581	100.0	2,911.5
1	Malignant neoplasms (C00-C97)	81,699	33.8	984.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	70,637	29.2	851.3
3	Chronic lower respiratory diseases (J40-J47)	15,877	6.6	191.3
4	Cerebrovascular diseases (I60-I69)	11,649	4.8	140.4
5	Diabetes mellitus (E10-E14)	8,437	3.5	101.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,751	2.0	57.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,826	1.6	46.1
8	Influenza and pneumonia (J10-J18)	3,732	1.5	45.0
9	Chronic liver disease and cirrhosis (K70,K73-K74)	3,335	1.4	40.2
10	Septicemia (A40-A41)	3,044	1.3	36.7
11	Aortic aneurysm and dissection (I71)	2,382	1.0	28.7
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,040	0.8	24.6
13	Alzheimer's disease (G30)	1,472	0.6	17.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,439	0.6	17.3
15	Parkinson's disease (G20-G21)	1,394	0.6	16.8
...	All other causes (Residual)	25,867	10.7	311.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Percent of		
		Number	total deaths	Rate
All races, male, 75-84 years				
...	All causes	340,742	100.0	6,833.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	108,576	31.9	2,177.3
2	Malignant neoplasms (C00-C97)	86,126	25.3	1,727.1
3	Chronic lower respiratory diseases (J40-J47)	23,691	7.0	475.1
4	Cerebrovascular diseases (I60-I69)	23,301	6.8	467.3
5	Diabetes mellitus (E10-E14)	9,918	2.9	198.9
6	Influenza and pneumonia (J10-J18)	9,294	2.7	186.4
7	Alzheimer's disease (G30)	6,795	2.0	136.3
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,652	2.0	133.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,485	1.9	130.0
10	Parkinson's disease (G20-G21)	4,837	1.4	97.0
11	Septicemia (A40-A41)	4,762	1.4	95.5
12	Aortic aneurysm and dissection (I71)	3,313	1.0	66.4
13	Pneumonitis due to solids and liquids (J69)	3,145	0.9	63.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,403	0.7	48.2
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,282	0.7	45.8
...	All other causes (Residual)	39,162	11.5	785.3
All races, male, 85 years and over				
...	All causes	217,533	100.0	16,744.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	78,473	36.1	6,040.5
2	Malignant neoplasms (C00-C97)	33,954	15.6	2,613.6
3	Cerebrovascular diseases (I60-I69)	17,930	8.2	1,380.2
4	Chronic lower respiratory diseases (J40-J47)	11,912	5.5	916.9
5	Influenza and pneumonia (J10-J18)	10,502	4.8	808.4
6	Alzheimer's disease (G30)	7,224	3.3	556.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,114	2.4	393.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,657	2.1	358.5
9	Diabetes mellitus (E10-E14)	4,457	2.0	343.1
10	Pneumonitis due to solids and liquids (J69)	3,511	1.6	270.3
11	Parkinson's disease (G20-G21)	2,915	1.3	224.4
12	Septicemia (A40-A41)	2,914	1.3	224.3
13	Atherosclerosis (I70)	2,172	1.0	167.2
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,959	0.9	150.8
15	Aortic aneurysm and dissection (I71)	1,491	0.7	114.8
...	All other causes (Residual)	28,348	13.0	2,182.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Percent of		
		Number	total deaths	Rate
All races, female, all ages				
...	All causes	1,233,004	100.0	850.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	361,047	29.3	249.0
2	Malignant neoplasms (C00-C97)	266,693	21.6	183.9
3	Cerebrovascular diseases (I60-I69)	100,361	8.1	69.2
4	Chronic lower respiratory diseases (J40-J47)	63,316	5.1	43.7
5	Diabetes mellitus (E10-E14)	38,531	3.1	26.6
6	Alzheimer's disease (G30)	38,090	3.1	26.3
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	35,477	2.9	24.5
8	Influenza and pneumonia (J10-J18)	34,692	2.8	23.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20,628	1.7	14.2
10	Septicemia (A40-A41)	17,931	1.5	12.4
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	11,996	1.0	8.3
12	Chronic liver disease and cirrhosis (K70,K73-K74)	9,642	0.8	6.7
13	Atherosclerosis (I70)	8,943	0.7	6.2
14	Pneumonitis due to solids and liquids (J69)	8,725	0.7	6.0
15	Parkinson's disease (G20-G21)	7,139	0.6	4.9
...	All other causes (Residual)	209,793	17.0	144.7
All races, female, 1-4 years				
...	All causes	2,208	100.0	29.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	669	30.3	8.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	275	12.5	3.7
3	Malignant neoplasms (C00-C97)	191	8.7	2.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	177	8.0	2.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	104	4.7	1.4
6	Influenza and pneumonia (J10-J18)	55	2.5	0.7
7	Septicemia (A40-A41)	48	2.2	0.6
8	Cerebrovascular diseases (I60-I69)	29	1.3	0.4
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	28	1.3	0.4
9	Certain conditions originating in the perinatal period (P00-P96)	28	1.3	0.4
11	Chronic lower respiratory diseases (J40-J47)	14	0.6	*
12	Meningococcal infection (A39)	10	0.5	*
12	Anemias (D50-D64)	10	0.5	*
12	Meningitis (G00,G03)	10	0.5	*
12	Acute bronchitis and bronchiolitis (J20-J21)	10	0.5	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	10	0.5	*
...	All other causes (Residual)	540	24.5	7.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Percent of		
		Number	total deaths	Rate
All races, female, 5-14 years				
...	All causes	2,927	100.0	14.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,033	35.3	5.2
2	Malignant neoplasms (C00-C97)	475	16.2	2.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	166	5.7	0.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	134	4.6	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	133	4.5	0.7
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	65	2.2	0.3
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	53	1.8	0.3
8	Influenza and pneumonia (J10-J18)	45	1.5	0.2
9	Chronic lower respiratory diseases (J40-J47)	43	1.5	0.2
10	Cerebrovascular diseases (I60-I69)	38	1.3	0.2
11	Septicemia (A40-A41)	30	1.0	0.1
12	Anemias (D50-D64)	28	1.0	0.1
13	Diabetes mellitus (E10-E14)	16	0.5	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	15	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	0.4	*
15	Certain conditions originating in the perinatal period (P00-P96)	12	0.4	*
...	All other causes (Residual)	629	21.5	3.1
All races, female, 15-24 years				
...	All causes	8,289	100.0	42.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,579	43.2	18.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	756	9.1	3.9
3	Malignant neoplasms (C00-C97)	680	8.2	3.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	562	6.8	2.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	352	4.2	1.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	199	2.4	1.0
7	Human immunodeficiency virus (HIV) disease (B20-B24)	120	1.4	0.6
8	Pregnancy, childbirth and the puerperium (O00-099)	110	1.3	0.6
9	Cerebrovascular diseases (I60-I69)	93	1.1	0.5
10	Influenza and pneumonia (J10-J18)	75	0.9	0.4
11	Chronic lower respiratory diseases (J40-J47)	72	0.9	0.4
12	Diabetes mellitus (E10-E14)	66	0.8	0.3
13	Septicemia (A40-A41)	58	0.7	0.3
14	Anemias (D50-D64)	50	0.6	0.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	49	0.6	0.3
...	All other causes (Residual)	1,468	17.7	7.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Percent of		
		Number	total deaths	Rate
All races, female, 25-34 years				
...	All causes	12,926	100.0	66.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,747	21.3	14.0
2	Malignant neoplasms (C00-C97)	2,134	16.5	10.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,092	8.4	5.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,047	8.1	5.3
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	871	6.7	4.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	680	5.3	3.5
7	Cerebrovascular diseases (I60-I69)	290	2.2	1.5
8	Diabetes mellitus (E10-E14)	254	2.0	1.3
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	209	1.6	1.1
10	Pregnancy, childbirth and the puerperium (O00-099)	190	1.5	1.0
11	Influenza and pneumonia (J10-J18)	148	1.1	0.8
12	Chronic lower respiratory diseases (J40-J47)	144	1.1	0.7
13	Septicemia (A40-A41)	142	1.1	0.7
14	Chronic liver disease and cirrhosis (K70,K73-K74)	128	1.0	0.7
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	113	0.9	0.6
...	All other causes (Residual)	2,737	21.2	14.0
All races, female, 35-44 years				
...	All causes	33,510	100.0	148.2
1	Malignant neoplasms (C00-C97)	9,272	27.7	41.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,333	12.9	19.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,980	11.9	17.6
4	Human immunodeficiency virus (HIV) disease (B20-B24)	1,507	4.5	6.7
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,455	4.3	6.4
6	Cerebrovascular diseases (I60-I69)	1,224	3.7	5.4
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,146	3.4	5.1
8	Chronic liver disease and cirrhosis (K70,K73-K74)	1,130	3.4	5.0
9	Diabetes mellitus (E10-E14)	777	2.3	3.4
10	Chronic lower respiratory diseases (J40-J47)	514	1.5	2.3
11	Influenza and pneumonia (J10-J18)	427	1.3	1.9
12	Septicemia (A40-A41)	378	1.1	1.7
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	354	1.1	1.6
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	269	0.8	1.2
15	Viral hepatitis (B15-B19)	244	0.7	1.1
...	All other causes (Residual)	6,500	19.4	28.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Percent of		
		Number	total deaths	Rate
All races, female, 45-54 years				
...	All causes	63,217	100.0	316.8
1	Malignant neoplasms (C00-C97)	24,491	38.7	122.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,119	16.0	50.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,601	5.7	18.0
4	Cerebrovascular diseases (I60-I69)	2,705	4.3	13.6
5	Diabetes mellitus (E10-E14)	2,189	3.5	11.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1,964	3.1	9.8
7	Chronic lower respiratory diseases (J40-J47)	1,622	2.6	8.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,438	2.3	7.2
9	Septicemia (A40-A41)	934	1.5	4.7
10	Human immunodeficiency virus (HIV) disease (B20-B24)	879	1.4	4.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	769	1.2	3.9
12	Influenza and pneumonia (J10-J18)	702	1.1	3.5
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	635	1.0	3.2
14	Viral hepatitis (B15-B19)	490	0.8	2.5
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	362	0.6	1.8
...	All other causes (Residual)	10,317	16.3	51.7
All races, female, 55-64 years				
...	All causes	99,181	100.0	754.0
1	Malignant neoplasms (C00-C97)	40,979	41.3	311.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	19,967	20.1	151.8
3	Chronic lower respiratory diseases (J40-J47)	5,464	5.5	41.5
4	Cerebrovascular diseases (I60-I69)	4,331	4.4	32.9
5	Diabetes mellitus (E10-E14)	4,329	4.4	32.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,430	2.5	18.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,722	1.7	13.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,570	1.6	11.9
9	Septicemia (A40-A41)	1,505	1.5	11.4
10	Influenza and pneumonia (J10-J18)	1,117	1.1	8.5
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	754	0.8	5.7
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	678	0.7	5.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	580	0.6	4.4
14	Viral hepatitis (B15-B19)	371	0.4	2.8
15	Aortic aneurysm and dissection (I71)	363	0.4	2.8
...	All other causes (Residual)	13,021	13.1	99.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Percent of		
		Number	total deaths	Rate
All races, female, 65-74 years				
...	All causes	189,379	100.0	1,890.8
1	Malignant neoplasms (C00-C97)	65,319	34.5	652.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	45,662	24.1	455.9
3	Chronic lower respiratory diseases (J40-J47)	14,874	7.9	148.5
4	Cerebrovascular diseases (I60-I69)	10,949	5.8	109.3
5	Diabetes mellitus (E10-E14)	8,294	4.4	82.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,530	1.9	35.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,084	1.6	30.8
8	Septicemia (A40-A41)	2,954	1.6	29.5
9	Influenza and pneumonia (J10-J18)	2,918	1.5	29.1
10	Chronic liver disease and cirrhosis (K70,K73-K74)	2,151	1.1	21.5
11	Alzheimer's disease (G30)	1,950	1.0	19.5
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,453	0.8	14.5
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,135	0.6	11.3
13	Aortic aneurysm and dissection (I71)	1,135	0.6	11.3
15	Parkinson's disease (G20-G21)	753	0.4	7.5
...	All other causes (Residual)	23,218	12.3	231.8
All races, female, 75-84 years				
...	All causes	361,187	100.0	4,760.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	108,416	30.0	1,428.9
2	Malignant neoplasms (C00-C97)	79,319	22.0	1,045.4
3	Cerebrovascular diseases (I60-I69)	32,521	9.0	428.6
4	Chronic lower respiratory diseases (J40-J47)	24,071	6.7	317.3
5	Diabetes mellitus (E10-E14)	12,887	3.6	169.9
6	Alzheimer's disease (G30)	11,747	3.3	154.8
7	Influenza and pneumonia (J10-J18)	9,383	2.6	123.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,612	1.8	87.1
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,036	1.7	79.6
10	Septicemia (A40-A41)	5,589	1.5	73.7
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	3,711	1.0	48.9
12	Parkinson's disease (G20-G21)	3,288	0.9	43.3
13	Pneumonitis due to solids and liquids (J69)	2,614	0.7	34.5
14	Aortic aneurysm and dissection (I71)	2,438	0.7	32.1
15	Atherosclerosis (I70)	2,365	0.7	31.2
...	All other causes (Residual)	50,190	13.9	661.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Percent of		
		Number	total deaths	Rate
All races, female, 85 years and over				
...	All causes	447,998	100.0	14,429.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	170,966	38.2	5,506.8
2	Cerebrovascular diseases (I60-I69)	48,136	10.7	1,550.4
3	Malignant neoplasms (C00-C97)	43,797	9.8	1,410.7
4	Alzheimer's disease (G30)	24,057	5.4	774.9
5	Influenza and pneumonia (J10-J18)	19,689	4.4	634.2
6	Chronic lower respiratory diseases (J40-J47)	16,479	3.7	530.8
7	Diabetes mellitus (E10-E14)	9,714	2.2	312.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,554	1.7	243.3
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,514	1.7	242.0
10	Septicemia (A40-A41)	6,155	1.4	198.3
11	Atherosclerosis (I70)	5,716	1.3	184.1
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	5,613	1.3	180.8
13	Pneumonitis due to solids and liquids (J69)	4,831	1.1	155.6
14	Parkinson's disease (G20-G21)	2,994	0.7	96.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,245	0.5	72.3
...	All other causes (Residual)	72,538	16.2	2,336.4
White, both sexes, all ages				
...	All causes	2,079,691	100.0	895.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	610,638	29.4	262.8
2	Malignant neoplasms (C00-C97)	479,651	23.1	206.4
3	Cerebrovascular diseases (I60-I69)	140,465	6.8	60.5
4	Chronic lower respiratory diseases (J40-J47)	113,819	5.5	49.0
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	85,964	4.1	37.0
6	Diabetes mellitus (E10-E14)	57,180	2.7	24.6
7	Influenza and pneumonia (J10-J18)	54,774	2.6	23.6
8	Alzheimer's disease (G30)	50,348	2.4	21.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	31,345	1.5	13.5
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	27,710	1.3	11.9
11	Septicemia (A40-A41)	25,806	1.2	11.1
12	Chronic liver disease and cirrhosis (K70,K73-K74)	23,408	1.1	10.1
13	Parkinson's disease (G20-G21)	15,757	0.8	6.8
14	Pneumonitis due to solids and liquids (J69)	15,425	0.7	6.6
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	14,752	0.7	6.3
...	All other causes (Residual)	332,649	16.0	143.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Percent of		
		Number	total deaths	Rate
White, both sexes, 1-4 years				
...	All causes	3,670	100.0	30.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,309	35.7	11.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	387	10.5	3.2
3	Malignant neoplasms (C00-C97)	353	9.6	3.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	228	6.2	1.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	141	3.8	1.2
6	Septicemia (A40-A41)	80	2.2	0.7
7	Influenza and pneumonia (J10-J18)	78	2.1	0.7
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	48	1.3	0.4
9	Certain conditions originating in the perinatal period (P00-P96)	47	1.3	0.4
10	Cerebrovascular diseases (I60-I69)	39	1.1	0.3
11	Chronic lower respiratory diseases (J40-J47)	20	0.5	0.2
12	Meningitis (G00,G03)	19	0.5	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	16	0.4	*
14	Meningococcal infection (A39)	14	0.4	*
15	Acute bronchitis and bronchiolitis (J20-J21)	13	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	0.4	*
...	All other causes (Residual)	865	23.6	7.2
White, both sexes, 5-14 years				
...	All causes	5,170	100.0	16.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,060	39.8	6.4
2	Malignant neoplasms (C00-C97)	780	15.1	2.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	282	5.5	0.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	218	4.2	0.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	218	4.2	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	187	3.6	0.6
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	77	1.5	0.2
8	Influenza and pneumonia (J10-J18)	67	1.3	0.2
9	Cerebrovascular diseases (I60-I69)	59	1.1	0.2
10	Septicemia (A40-A41)	54	1.0	0.2
11	Chronic lower respiratory diseases (J40-J47)	48	0.9	0.2
12	Anemias (D50-D64)	27	0.5	0.1
13	Diabetes mellitus (E10-E14)	21	0.4	0.1
13	Certain conditions originating in the perinatal period (P00-P96)	21	0.4	0.1
15	Meningococcal infection (A39)	20	0.4	0.1
...	All other causes (Residual)	1,031	19.9	3.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Percent of		
		Number	total deaths	Rate
White, both sexes, 15-24 years				
...	All causes	23,748	100.0	75.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,038	50.7	38.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,334	14.0	10.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,264	9.5	7.2
4	Malignant neoplasms (C00-C97)	1,359	5.7	4.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	639	2.7	2.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	374	1.6	1.2
7	Cerebrovascular diseases (I60-I69)	138	0.6	0.4
8	Influenza and pneumonia (J10-J18)	126	0.5	0.4
9	Chronic lower respiratory diseases (J40-J47)	85	0.4	0.3
10	Septicemia (A40-A41)	82	0.3	0.3
10	Diabetes mellitus (E10-E14)	82	0.3	0.3
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	74	0.3	0.2
13	Pregnancy, childbirth and the puerperium (O00-099)	64	0.3	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	61	0.3	0.2
15	Human immunodeficiency virus (HIV) disease (B20-B24)	60	0.3	0.2
15	Legal intervention (Y35,Y89.0)	60	0.3	0.2
...	All other causes (Residual)	2,908	12.2	9.2
White, both sexes, 25-34 years				
...	All causes	29,994	100.0	95.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,727	32.4	30.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,356	14.5	13.8
3	Malignant neoplasms (C00-C97)	3,094	10.3	9.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,623	8.7	8.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,086	7.0	6.6
6	Human immunodeficiency virus (HIV) disease (B20-B24)	814	2.7	2.6
7	Cerebrovascular diseases (I60-I69)	416	1.4	1.3
8	Diabetes mellitus (E10-E14)	383	1.3	1.2
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	375	1.3	1.2
10	Chronic liver disease and cirrhosis (K70,K73-K74)	307	1.0	1.0
11	Influenza and pneumonia (J10-J18)	232	0.8	0.7
12	Septicemia (A40-A41)	169	0.6	0.5
13	Chronic lower respiratory diseases (J40-J47)	156	0.5	0.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	126	0.4	0.4
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	125	0.4	0.4
...	All other causes (Residual)	5,005	16.7	15.9

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Percent of		
		Number	total deaths	Rate
White, both sexes, 35-44 years				
...	All causes	68,137	100.0	185.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,143	19.3	35.8
2	Malignant neoplasms (C00-C97)	12,959	19.0	35.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,510	14.0	25.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,017	8.8	16.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,745	4.0	7.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	2,681	3.9	7.3
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,672	3.9	7.3
8	Cerebrovascular diseases (I60-I69)	1,605	2.4	4.4
9	Diabetes mellitus (E10-E14)	1,399	2.1	3.8
10	Influenza and pneumonia (J10-J18)	678	1.0	1.8
11	Chronic lower respiratory diseases (J40-J47)	665	1.0	1.8
12	Viral hepatitis (B15-B19)	646	0.9	1.8
13	Septicemia (A40-A41)	531	0.8	1.4
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	479	0.7	1.3
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	432	0.6	1.2
...	All other causes (Residual)	11,975	17.6	32.6
White, both sexes, 45-54 years				
...	All causes	128,699	100.0	393.2
1	Malignant neoplasms (C00-C97)	39,686	30.8	121.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	27,472	21.3	83.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,825	8.4	33.1
4	Chronic liver disease and cirrhosis (K70,K73-K74)	5,989	4.7	18.3
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,540	4.3	16.9
6	Diabetes mellitus (E10-E14)	3,842	3.0	11.7
7	Cerebrovascular diseases (I60-I69)	3,736	2.9	11.4
8	Chronic lower respiratory diseases (J40-J47)	2,703	2.1	8.3
9	Human immunodeficiency virus (HIV) disease (B20-B24)	1,792	1.4	5.5
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,639	1.3	5.0
11	Viral hepatitis (B15-B19)	1,632	1.3	5.0
12	Influenza and pneumonia (J10-J18)	1,316	1.0	4.0
13	Septicemia (A40-A41)	1,312	1.0	4.0
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,043	0.8	3.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	610	0.5	1.9
...	All other causes (Residual)	19,562	15.2	59.8

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Percent of		
		Number	total deaths	Rate
White, both sexes, 55-64 years				
...	All causes	197,853	100.0	913.5
1	Malignant neoplasms (C00-C97)	75,603	38.2	349.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	49,719	25.1	229.6
3	Chronic lower respiratory diseases (J40-J47)	9,967	5.0	46.0
4	Diabetes mellitus (E10-E14)	7,093	3.6	32.7
5	Cerebrovascular diseases (I60-I69)	6,874	3.5	31.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,367	3.2	29.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	4,924	2.5	22.7
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,106	1.6	14.3
9	Septicemia (A40-A41)	2,240	1.1	10.3
10	Influenza and pneumonia (J10-J18)	2,141	1.1	9.9
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,101	1.1	9.7
12	Aortic aneurysm and dissection (I71)	1,167	0.6	5.4
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,075	0.5	5.0
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	863	0.4	4.0
15	Viral hepatitis (B15-B19)	778	0.4	3.6
...	All other causes (Residual)	23,835	12.0	110.0
White, both sexes, 65-74 years				
...	All causes	367,128	100.0	2,298.9
1	Malignant neoplasms (C00-C97)	127,404	34.7	797.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	97,764	26.6	612.2
3	Chronic lower respiratory diseases (J40-J47)	28,385	7.7	177.7
4	Cerebrovascular diseases (I60-I69)	18,047	4.9	113.0
5	Diabetes mellitus (E10-E14)	13,085	3.6	81.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,778	1.8	42.4
7	Influenza and pneumonia (J10-J18)	5,644	1.5	35.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,458	1.5	34.2
9	Chronic liver disease and cirrhosis (K70,K73-K74)	4,939	1.3	30.9
10	Septicemia (A40-A41)	4,677	1.3	29.3
11	Aortic aneurysm and dissection (I71)	3,181	0.9	19.9
12	Alzheimer's disease (G30)	3,136	0.9	19.6
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,288	0.6	14.3
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,282	0.6	14.3
15	Parkinson's disease (G20-G21)	2,011	0.5	12.6
...	All other causes (Residual)	42,049	11.5	263.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Percent of		
		Number	total deaths	Rate
White, both sexes, 75-84 years				
...	All causes	627,752	100.0	5,560.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	193,525	30.8	1,714.3
2	Malignant neoplasms (C00-C97)	148,005	23.6	1,311.1
3	Cerebrovascular diseases (I60-I69)	49,265	7.8	436.4
4	Chronic lower respiratory diseases (J40-J47)	44,996	7.2	398.6
5	Diabetes mellitus (E10-E14)	19,070	3.0	168.9
6	Alzheimer's disease (G30)	17,226	2.7	152.6
7	Influenza and pneumonia (J10-J18)	16,621	2.6	147.2
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,588	1.8	102.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10,936	1.7	96.9
10	Septicemia (A40-A41)	8,708	1.4	77.1
11	Parkinson's disease (G20-G21)	7,772	1.2	68.8
12	Aortic aneurysm and dissection (I71)	5,293	0.8	46.9
13	Pneumonitis due to solids and liquids (J69)	5,172	0.8	45.8
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	4,776	0.8	42.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,292	0.7	38.0
...	All other causes (Residual)	80,507	12.8	713.2
White, both sexes, 85 years and over				
...	All causes	609,265	100.0	15,287.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	229,241	37.6	5,752.1
2	Malignant neoplasms (C00-C97)	70,350	11.5	1,765.2
3	Cerebrovascular diseases (I60-I69)	60,225	9.9	1,511.2
4	Alzheimer's disease (G30)	29,415	4.8	738.1
5	Influenza and pneumonia (J10-J18)	27,699	4.5	695.0
6	Chronic lower respiratory diseases (J40-J47)	26,772	4.4	671.8
7	Diabetes mellitus (E10-E14)	12,197	2.0	306.0
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,379	1.9	285.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11,088	1.8	278.2
10	Septicemia (A40-A41)	7,782	1.3	195.3
11	Pneumonitis due to solids and liquids (J69)	7,603	1.2	190.8
12	Atherosclerosis (I70)	7,409	1.2	185.9
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	6,510	1.1	163.3
14	Parkinson's disease (G20-G21)	5,652	0.9	141.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3,344	0.5	83.9
...	All other causes (Residual)	92,599	15.2	2,323.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Percent of		
		Number	total deaths	Rate
White, male, all ages				
...	All causes	1,011,218	100.0	881.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	295,556	29.2	257.8
2	Malignant neoplasms (C00-C97)	248,146	24.5	216.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	55,493	5.5	48.4
4	Chronic lower respiratory diseases (J40-J47)	54,561	5.4	47.6
5	Cerebrovascular diseases (I60-I69)	53,428	5.3	46.6
6	Diabetes mellitus (E10-E14)	26,917	2.7	23.5
7	Influenza and pneumonia (J10-J18)	23,744	2.3	20.7
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	22,328	2.2	19.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15,241	1.5	13.3
10	Chronic liver disease and cirrhosis (K70,K73-K74)	15,048	1.5	13.1
11	Alzheimer's disease (G30)	14,714	1.5	12.8
12	Septicemia (A40-A41)	11,371	1.1	9.9
13	Parkinson's disease (G20-G21)	8,982	0.9	7.8
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8,254	0.8	7.2
15	Aortic aneurysm and dissection (I71)	8,224	0.8	7.2
...	All other causes (Residual)	149,211	14.8	130.1
White, male, 1-4 years				
...	All causes	2,094	100.0	34.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	815	38.9	13.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	195	9.3	3.2
3	Malignant neoplasms (C00-C97)	189	9.0	3.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	132	6.3	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	74	3.5	1.2
6	Septicemia (A40-A41)	46	2.2	0.8
7	Influenza and pneumonia (J10-J18)	41	2.0	0.7
8	Certain conditions originating in the perinatal period (P00-P96)	28	1.3	0.5
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	24	1.1	0.4
10	Cerebrovascular diseases (I60-I69)	18	0.9	*
11	Meningitis (G00,G03)	13	0.6	*
12	Chronic lower respiratory diseases (J40-J47)	12	0.6	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	10	0.5	*
14	Meningococcal infection (A39)	7	0.3	*
15	Anemias (D50-D64)	5	0.2	*
15	Acute bronchitis and bronchiolitis (J20-J21)	5	0.2	*
...	All other causes (Residual)	480	22.9	7.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Percent of		
		Number	total deaths	Rate
White, male, 5-14 years				
...	All causes	3,014	100.0	18.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,285	42.6	7.8
2	Malignant neoplasms (C00-C97)	398	13.2	2.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	169	5.6	1.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	168	5.6	1.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	119	3.9	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	98	3.3	0.6
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	37	1.2	0.2
8	Cerebrovascular diseases (I60-I69)	34	1.1	0.2
8	Influenza and pneumonia (J10-J18)	34	1.1	0.2
10	Septicemia (A40-A41)	32	1.1	0.2
11	Chronic lower respiratory diseases (J40-J47)	28	0.9	0.2
12	Anemias (D50-D64)	16	0.5	*
12	Certain conditions originating in the perinatal period (P00-P96)	16	0.5	*
14	Diabetes mellitus (E10-E14)	15	0.5	*
15	Meningococcal infection (A39)	12	0.4	*
...	All other causes (Residual)	553	18.3	3.4
White, male, 15-24 years				
...	All causes	17,529	100.0	108.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,067	51.7	55.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,860	16.3	17.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,811	10.3	11.2
4	Malignant neoplasms (C00-C97)	824	4.7	5.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	414	2.4	2.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	222	1.3	1.4
7	Cerebrovascular diseases (I60-I69)	81	0.5	0.5
8	Influenza and pneumonia (J10-J18)	79	0.5	0.5
9	Legal intervention (Y35,Y89.0)	58	0.3	0.4
10	Diabetes mellitus (E10-E14)	50	0.3	0.3
11	Septicemia (A40-A41)	46	0.3	0.3
12	Chronic lower respiratory diseases (J40-J47)	45	0.3	0.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	37	0.2	0.2
14	Human immunodeficiency virus (HIV) disease (B20-B24)	35	0.2	0.2
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	30	0.2	0.2
...	All other causes (Residual)	1,870	10.7	11.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Percent of		
		Number	total deaths	Rate
White, male, 25-34 years				
...	All causes	20,978	100.0	130.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,518	35.8	46.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,604	17.2	22.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,988	9.5	12.3
4	Malignant neoplasms (C00-C97)	1,493	7.1	9.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,408	6.7	8.7
6	Human immunodeficiency virus (HIV) disease (B20-B24)	628	3.0	3.9
7	Cerebrovascular diseases (I60-I69)	220	1.0	1.4
8	Diabetes mellitus (E10-E14)	212	1.0	1.3
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	212	1.0	1.3
10	Chronic liver disease and cirrhosis (K70,K73-K74)	211	1.0	1.3
11	Influenza and pneumonia (J10-J18)	127	0.6	0.8
12	Septicemia (A40-A41)	85	0.4	0.5
13	Legal intervention (Y35,Y89.0)	77	0.4	0.5
14	Chronic lower respiratory diseases (J40-J47)	74	0.4	0.5
15	Aortic aneurysm and dissection (I71)	73	0.3	0.5
...	All other causes (Residual)	3,048	14.5	18.9
White, male, 35-44 years				
...	All causes	44,243	100.0	239.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,647	21.8	52.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,981	15.8	37.8
3	Malignant neoplasms (C00-C97)	5,783	13.1	31.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,691	10.6	25.4
5	Human immunodeficiency virus (HIV) disease (B20-B24)	2,238	5.1	12.1
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,972	4.5	10.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,826	4.1	9.9
8	Diabetes mellitus (E10-E14)	876	2.0	4.7
9	Cerebrovascular diseases (I60-I69)	834	1.9	4.5
10	Viral hepatitis (B15-B19)	456	1.0	2.5
11	Influenza and pneumonia (J10-J18)	382	0.9	2.1
12	Chronic lower respiratory diseases (J40-J47)	316	0.7	1.7
13	Septicemia (A40-A41)	309	0.7	1.7
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	261	0.6	1.4
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	256	0.6	1.4
...	All other causes (Residual)	7,415	16.8	40.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Percent of		
		Number	total deaths	Rate
White, male, 45-54 years				
...	All causes	81,403	100.0	501.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,620	25.3	127.0
2	Malignant neoplasms (C00-C97)	20,069	24.7	123.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,897	9.7	48.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	4,431	5.4	27.3
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,218	5.2	26.0
6	Diabetes mellitus (E10-E14)	2,326	2.9	14.3
7	Cerebrovascular diseases (I60-I69)	2,055	2.5	12.7
8	Human immunodeficiency virus (HIV) disease (B20-B24)	1,540	1.9	9.5
9	Chronic lower respiratory diseases (J40-J47)	1,391	1.7	8.6
10	Viral hepatitis (B15-B19)	1,271	1.6	7.8
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,196	1.5	7.4
12	Influenza and pneumonia (J10-J18)	791	1.0	4.9
13	Septicemia (A40-A41)	695	0.9	4.3
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	613	0.8	3.8
15	Aortic aneurysm and dissection (I71)	375	0.5	2.3
...	All other causes (Residual)	11,915	14.6	73.4
White, male, 55-64 years				
...	All causes	118,353	100.0	1,127.5
1	Malignant neoplasms (C00-C97)	41,155	34.8	392.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	34,724	29.3	330.8
3	Chronic lower respiratory diseases (J40-J47)	5,043	4.3	48.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,319	3.6	41.1
5	Diabetes mellitus (E10-E14)	4,031	3.4	38.4
6	Cerebrovascular diseases (I60-I69)	3,790	3.2	36.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	3,442	2.9	32.8
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,396	2.0	22.8
9	Influenza and pneumonia (J10-J18)	1,239	1.0	11.8
10	Septicemia (A40-A41)	1,160	1.0	11.1
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,144	1.0	10.9
12	Aortic aneurysm and dissection (I71)	877	0.7	8.4
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	583	0.5	5.6
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	518	0.4	4.9
15	Viral hepatitis (B15-B19)	508	0.4	4.8
...	All other causes (Residual)	13,424	11.3	127.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Percent of		
		Number	total deaths	Rate
White, male, 65-74 years				
...	All causes	207,793	100.0	2,842.3
1	Malignant neoplasms (C00-C97)	70,875	34.1	969.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	60,615	29.2	829.1
3	Chronic lower respiratory diseases (J40-J47)	14,499	7.0	198.3
4	Cerebrovascular diseases (I60-I69)	9,393	4.5	128.5
5	Diabetes mellitus (E10-E14)	6,917	3.3	94.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,119	2.0	56.3
7	Influenza and pneumonia (J10-J18)	3,144	1.5	43.0
8	Chronic liver disease and cirrhosis (K70,K73-K74)	3,003	1.4	41.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,950	1.4	40.4
10	Septicemia (A40-A41)	2,378	1.1	32.5
11	Aortic aneurysm and dissection (I71)	2,185	1.1	29.9
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,921	0.9	26.3
13	Alzheimer's disease (G30)	1,349	0.6	18.5
14	Parkinson's disease (G20-G21)	1,311	0.6	17.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,304	0.6	17.8
...	All other causes (Residual)	21,830	10.5	298.6
White, male, 75-84 years				
...	All causes	306,278	100.0	6,799.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	98,006	32.0	2,175.8
2	Malignant neoplasms (C00-C97)	76,782	25.1	1,704.6
3	Chronic lower respiratory diseases (J40-J47)	22,042	7.2	489.4
4	Cerebrovascular diseases (I60-I69)	20,664	6.7	458.8
5	Diabetes mellitus (E10-E14)	8,520	2.8	189.2
6	Influenza and pneumonia (J10-J18)	8,225	2.7	182.6
7	Alzheimer's disease (G30)	6,327	2.1	140.5
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,050	2.0	134.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,565	1.8	123.5
10	Parkinson's disease (G20-G21)	4,641	1.5	103.0
11	Septicemia (A40-A41)	4,029	1.3	89.4
12	Aortic aneurysm and dissection (I71)	3,087	1.0	68.5
13	Pneumonitis due to solids and liquids (J69)	2,849	0.9	63.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,227	0.7	49.4
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,820	0.6	40.4
...	All other causes (Residual)	35,444	11.6	786.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Percent of		
		Number	total deaths	Rate
White, male, 85 years and over				
...	All causes	199,179	100.0	16,935.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	72,415	36.4	6,157.2
2	Malignant neoplasms (C00-C97)	30,550	15.3	2,597.6
3	Cerebrovascular diseases (I60-I69)	16,303	8.2	1,386.2
4	Chronic lower respiratory diseases (J40-J47)	11,098	5.6	943.6
5	Influenza and pneumonia (J10-J18)	9,578	4.8	814.4
6	Alzheimer's disease (G30)	6,789	3.4	577.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,574	2.3	388.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,329	2.2	368.1
9	Diabetes mellitus (E10-E14)	3,967	2.0	337.3
10	Pneumonitis due to solids and liquids (J69)	3,239	1.6	275.4
11	Parkinson's disease (G20-G21)	2,795	1.4	237.6
12	Septicemia (A40-A41)	2,491	1.3	211.8
13	Atherosclerosis (I70)	2,045	1.0	173.9
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,672	0.8	142.2
15	Aortic aneurysm and dissection (I71)	1,415	0.7	120.3
...	All other causes (Residual)	25,919	13.0	2,203.8
White, female, all ages				
...	All causes	1,068,473	100.0	907.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	315,082	29.5	267.7
2	Malignant neoplasms (C00-C97)	231,505	21.7	196.7
3	Cerebrovascular diseases (I60-I69)	87,037	8.1	74.0
4	Chronic lower respiratory diseases (J40-J47)	59,258	5.5	50.3
5	Alzheimer's disease (G30)	35,634	3.3	30.3
6	Influenza and pneumonia (J10-J18)	31,030	2.9	26.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	30,471	2.9	25.9
8	Diabetes mellitus (E10-E14)	30,263	2.8	25.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16,104	1.5	13.7
10	Septicemia (A40-A41)	14,435	1.4	12.3
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	9,401	0.9	8.0
12	Chronic liver disease and cirrhosis (K70,K73-K74)	8,360	0.8	7.1
13	Atherosclerosis (I70)	8,265	0.8	7.0
14	Pneumonitis due to solids and liquids (J69)	7,720	0.7	6.6
15	Parkinson's disease (G20-G21)	6,775	0.6	5.8
...	All other causes (Residual)	177,133	16.6	150.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Percent of		
		Number	total deaths	Rate
White, female, 1-4 years				
...	All causes	1,576	100.0	27.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	494	31.3	8.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	192	12.2	3.3
3	Malignant neoplasms (C00-C97)	164	10.4	2.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	96	6.1	1.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	67	4.3	1.1
6	Influenza and pneumonia (J10-J18)	37	2.3	0.6
7	Septicemia (A40-A41)	34	2.2	0.6
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	24	1.5	0.4
9	Cerebrovascular diseases (I60-I69)	21	1.3	0.4
10	Certain conditions originating in the perinatal period (P00-P96)	19	1.2	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.6	*
12	Acute bronchitis and bronchiolitis (J20-J21)	8	0.5	*
12	Chronic lower respiratory diseases (J40-J47)	8	0.5	*
12	Pneumonitis due to solids and liquids (J69)	8	0.5	*
15	Meningococcal infection (A39)	7	0.4	*
...	All other causes (Residual)	388	24.6	6.7
White, female, 5-14 years				
...	All causes	2,156	100.0	13.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	775	35.9	5.0
2	Malignant neoplasms (C00-C97)	382	17.7	2.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	113	5.2	0.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	99	4.6	0.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	89	4.1	0.6
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	50	2.3	0.3
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	40	1.9	0.3
8	Influenza and pneumonia (J10-J18)	33	1.5	0.2
9	Cerebrovascular diseases (I60-I69)	25	1.2	0.2
10	Septicemia (A40-A41)	22	1.0	0.1
11	Chronic lower respiratory diseases (J40-J47)	20	0.9	0.1
12	Anemias (D50-D64)	11	0.5	*
13	Meningococcal infection (A39)	8	0.4	*
13	Meningitis (G00,G03)	8	0.4	*
15	Diabetes mellitus (E10-E14)	6	0.3	*
15	Pneumonitis due to solids and liquids (J69)	6	0.3	*
...	All other causes (Residual)	469	21.8	3.0

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Percent of		
		Number	total deaths	Rate
White, female, 15-24 years				
...	All causes	6,219	100.0	40.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,971	47.8	19.5
2	Malignant neoplasms (C00-C97)	535	8.6	3.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	474	7.6	3.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	453	7.3	3.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	225	3.6	1.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	152	2.4	1.0
7	Pregnancy, childbirth and the puerperium (O00-099)	64	1.0	0.4
8	Cerebrovascular diseases (I60-I69)	57	0.9	0.4
9	Influenza and pneumonia (J10-J18)	47	0.8	0.3
10	Chronic lower respiratory diseases (J40-J47)	40	0.6	0.3
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	37	0.6	0.2
12	Septicemia (A40-A41)	36	0.6	0.2
13	Diabetes mellitus (E10-E14)	32	0.5	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	31	0.5	0.2
15	Human immunodeficiency virus (HIV) disease (B20-B24)	25	0.4	0.2
...	All other causes (Residual)	1,040	16.7	6.8
White, female, 25-34 years				
...	All causes	9,016	100.0	58.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,209	24.5	14.4
2	Malignant neoplasms (C00-C97)	1,601	17.8	10.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	752	8.3	4.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	678	7.5	4.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	635	7.0	4.1
6	Cerebrovascular diseases (I60-I69)	196	2.2	1.3
7	Human immunodeficiency virus (HIV) disease (B20-B24)	186	2.1	1.2
8	Diabetes mellitus (E10-E14)	171	1.9	1.1
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	163	1.8	1.1
10	Pregnancy, childbirth and the puerperium (O00-099)	114	1.3	0.7
11	Influenza and pneumonia (J10-J18)	105	1.2	0.7
12	Chronic liver disease and cirrhosis (K70,K73-K74)	96	1.1	0.6
13	Septicemia (A40-A41)	84	0.9	0.5
14	Chronic lower respiratory diseases (J40-J47)	82	0.9	0.5
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	70	0.8	0.5
...	All other causes (Residual)	1,874	20.8	12.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Percent of		
		Number	total deaths	Rate
White, female, 35-44 years				
...	All causes	23,894	100.0	131.0
1	Malignant neoplasms (C00-C97)	7,176	30.0	39.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,496	14.6	19.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,529	10.6	13.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,326	5.5	7.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	919	3.8	5.0
6	Cerebrovascular diseases (I60-I69)	771	3.2	4.2
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	700	2.9	3.8
8	Diabetes mellitus (E10-E14)	523	2.2	2.9
9	Human immunodeficiency virus (HIV) disease (B20-B24)	443	1.9	2.4
10	Chronic lower respiratory diseases (J40-J47)	349	1.5	1.9
11	Influenza and pneumonia (J10-J18)	296	1.2	1.6
12	Septicemia (A40-A41)	222	0.9	1.2
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	218	0.9	1.2
14	Viral hepatitis (B15-B19)	190	0.8	1.0
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	176	0.7	1.0
...	All other causes (Residual)	4,560	19.1	25.0
White, female, 45-54 years				
...	All causes	47,296	100.0	286.8
1	Malignant neoplasms (C00-C97)	19,617	41.5	118.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,852	14.5	41.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,928	6.2	17.8
4	Cerebrovascular diseases (I60-I69)	1,681	3.6	10.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,558	3.3	9.4
6	Diabetes mellitus (E10-E14)	1,516	3.2	9.2
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,322	2.8	8.0
8	Chronic lower respiratory diseases (J40-J47)	1,312	2.8	8.0
9	Septicemia (A40-A41)	617	1.3	3.7
10	Influenza and pneumonia (J10-J18)	525	1.1	3.2
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	443	0.9	2.7
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	430	0.9	2.6
13	Viral hepatitis (B15-B19)	361	0.8	2.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	268	0.6	1.6
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	263	0.6	1.6
...	All other causes (Residual)	7,603	16.1	46.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Percent of		
		Number	total deaths	Rate
White, female, 55-64 years				
...	All causes	79,500	100.0	712.2
1	Malignant neoplasms (C00-C97)	34,448	43.3	308.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,995	18.9	134.3
3	Chronic lower respiratory diseases (J40-J47)	4,924	6.2	44.1
4	Cerebrovascular diseases (I60-I69)	3,084	3.9	27.6
5	Diabetes mellitus (E10-E14)	3,062	3.9	27.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,048	2.6	18.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,482	1.9	13.3
8	Septicemia (A40-A41)	1,080	1.4	9.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	957	1.2	8.6
10	Influenza and pneumonia (J10-J18)	902	1.1	8.1
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	710	0.9	6.4
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	492	0.6	4.4
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	387	0.5	3.5
14	Aortic aneurysm and dissection (I71)	290	0.4	2.6
15	Alzheimer's disease (G30)	289	0.4	2.6
...	All other causes (Residual)	10,350	13.0	92.7
White, female, 65-74 years				
...	All causes	159,335	100.0	1,840.2
1	Malignant neoplasms (C00-C97)	56,529	35.5	652.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	37,149	23.3	429.0
3	Chronic lower respiratory diseases (J40-J47)	13,886	8.7	160.4
4	Cerebrovascular diseases (I60-I69)	8,654	5.4	99.9
5	Diabetes mellitus (E10-E14)	6,168	3.9	71.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,659	1.7	30.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,508	1.6	29.0
8	Influenza and pneumonia (J10-J18)	2,500	1.6	28.9
9	Septicemia (A40-A41)	2,299	1.4	26.6
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,936	1.2	22.4
11	Alzheimer's disease (G30)	1,787	1.1	20.6
12	Aortic aneurysm and dissection (I71)	996	0.6	11.5
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	984	0.6	11.4
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	977	0.6	11.3
15	Parkinson's disease (G20-G21)	700	0.4	8.1
...	All other causes (Residual)	19,603	12.3	226.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Percent of		
		Number	total deaths	Rate
White, female, 75-84 years				
...	All causes	321,474	100.0	4,738.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	95,519	29.7	1,407.9
2	Malignant neoplasms (C00-C97)	71,223	22.2	1,049.8
3	Cerebrovascular diseases (I60-I69)	28,601	8.9	421.6
4	Chronic lower respiratory diseases (J40-J47)	22,954	7.1	338.3
5	Alzheimer's disease (G30)	10,899	3.4	160.6
6	Diabetes mellitus (E10-E14)	10,550	3.3	155.5
7	Influenza and pneumonia (J10-J18)	8,396	2.6	123.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,538	1.7	81.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,371	1.7	79.2
10	Septicemia (A40-A41)	4,679	1.5	69.0
11	Parkinson's disease (G20-G21)	3,131	1.0	46.2
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,956	0.9	43.6
13	Pneumonitis due to solids and liquids (J69)	2,323	0.7	34.2
14	Aortic aneurysm and dissection (I71)	2,206	0.7	32.5
15	Atherosclerosis (I70)	2,176	0.7	32.1
...	All other causes (Residual)	44,952	14.0	662.6
White, female, 85 years and over				
...	All causes	410,086	100.0	14,597.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	156,826	38.2	5,582.5
2	Cerebrovascular diseases (I60-I69)	43,922	10.7	1,563.5
3	Malignant neoplasms (C00-C97)	39,800	9.7	1,416.7
4	Alzheimer's disease (G30)	22,626	5.5	805.4
5	Influenza and pneumonia (J10-J18)	18,121	4.4	645.0
6	Chronic lower respiratory diseases (J40-J47)	15,674	3.8	557.9
7	Diabetes mellitus (E10-E14)	8,230	2.0	293.0
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,050	1.7	251.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,514	1.6	231.9
10	Atherosclerosis (I70)	5,364	1.3	190.9
11	Septicemia (A40-A41)	5,291	1.3	188.3
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	4,838	1.2	172.2
13	Pneumonitis due to solids and liquids (J69)	4,364	1.1	155.3
14	Parkinson's disease (G20-G21)	2,857	0.7	101.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,052	0.5	73.0
...	All other causes (Residual)	66,557	16.2	2,369.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Percent of		
		Number	total deaths	Rate
Black, both sexes, all ages				
...	All causes	287,709	100.0	773.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	77,674	27.0	208.8
2	Malignant neoplasms (C00-C97)	62,170	21.6	167.1
3	Cerebrovascular diseases (I60-I69)	19,002	6.6	51.1
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,462	4.3	33.5
5	Diabetes mellitus (E10-E14)	12,305	4.3	33.1
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8,226	2.9	22.1
7	Human immunodeficiency virus (HIV) disease (B20-B24)	7,844	2.7	21.1
8	Chronic lower respiratory diseases (J40-J47)	7,589	2.6	20.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,274	2.5	19.6
10	Septicemia (A40-A41)	5,880	2.0	15.8
11	Influenza and pneumonia (J10-J18)	5,771	2.0	15.5
12	Certain conditions originating in the perinatal period (P00-P96)	4,847	1.7	13.0
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	4,038	1.4	10.9
14	Alzheimer's disease (G30)	3,114	1.1	8.4
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,775	1.0	7.5
...	All other causes (Residual)	46,738	16.2	125.7
Black, both sexes, 1-4 years				
...	All causes	1,181	100.0	47.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	339	28.7	13.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	169	14.3	6.8
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	131	11.1	5.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	74	6.3	3.0
5	Malignant neoplasms (C00-C97)	48	4.1	1.9
6	Influenza and pneumonia (J10-J18)	26	2.2	1.0
7	Chronic lower respiratory diseases (J40-J47)	23	1.9	0.9
7	Certain conditions originating in the perinatal period (P00-P96)	23	1.9	0.9
9	Septicemia (A40-A41)	22	1.9	0.9
10	Cerebrovascular diseases (I60-I69)	12	1.0	*
11	Anemias (D50-D64)	11	0.9	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	10	0.8	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.6	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.4	*
14	Meningitis (G00,G03)	5	0.4	*
...	All other causes (Residual)	276	23.4	11.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 5-14 years				
...	All causes	1,587	100.0	23.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	644	40.6	9.4
2	Malignant neoplasms (C00-C97)	176	11.1	2.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	94	5.9	1.4
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	75	4.7	1.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	73	4.6	1.1
6	Chronic lower respiratory diseases (J40-J47)	54	3.4	0.8
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	41	2.6	0.6
8	Human immunodeficiency virus (HIV) disease (B20-B24)	26	1.6	0.4
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	26	1.6	0.4
8	Anemias (D50-D64)	26	1.6	0.4
11	Influenza and pneumonia (J10-J18)	17	1.1	*
12	Diabetes mellitus (E10-E14)	15	0.9	*
13	Cerebrovascular diseases (I60-I69)	14	0.9	*
14	Certain conditions originating in the perinatal period (P00-P96)	11	0.7	*
15	Septicemia (A40-A41)	10	0.6	*
...	All other causes (Residual)	285	18.0	4.2
Black, both sexes, 15-24 years				
...	All causes	7,147	100.0	117.7
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,870	40.2	47.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,742	24.4	28.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	433	6.1	7.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	326	4.6	5.4
5	Malignant neoplasms (C00-C97)	262	3.7	4.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	163	2.3	2.7
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	114	1.6	1.9
8	Chronic lower respiratory diseases (J40-J47)	77	1.1	1.3
9	Anemias (D50-D64)	73	1.0	1.2
10	Diabetes mellitus (E10-E14)	60	0.8	1.0
11	Cerebrovascular diseases (I60-I69)	51	0.7	0.8
12	Influenza and pneumonia (J10-J18)	45	0.6	0.7
12	Pregnancy, childbirth and the puerperium (O00-099)	45	0.6	0.7
14	Septicemia (A40-A41)	41	0.6	0.7
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	27	0.4	0.4
...	All other causes (Residual)	818	11.4	13.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Percent of		
		Number	total deaths	Rate
Black, both sexes, 25-34 years				
...	All causes	10,028	100.0	185.0
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,371	23.6	43.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,638	16.3	30.2
3	Human immunodeficiency virus (HIV) disease (B20-B24)	1,258	12.5	23.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	970	9.7	17.9
5	Malignant neoplasms (C00-C97)	697	7.0	12.9
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	492	4.9	9.1
7	Diabetes mellitus (E10-E14)	202	2.0	3.7
8	Cerebrovascular diseases (I60-I69)	158	1.6	2.9
9	Chronic lower respiratory diseases (J40-J47)	129	1.3	2.4
10	Anemias (D50-D64)	119	1.2	2.2
11	Septicemia (A40-A41)	98	1.0	1.8
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	95	0.9	1.8
13	Influenza and pneumonia (J10-J18)	89	0.9	1.6
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	68	0.7	1.3
15	Pregnancy, childbirth and the puerperium (O00-099)	66	0.7	1.2
...	All other causes (Residual)	1,578	15.7	29.1
Black, both sexes, 35-44 years				
...	All causes	20,687	100.0	356.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,451	16.7	59.5
2	Human immunodeficiency virus (HIV) disease (B20-B24)	3,119	15.1	53.7
3	Malignant neoplasms (C00-C97)	2,981	14.4	51.4
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,321	11.2	40.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,450	7.0	25.0
6	Cerebrovascular diseases (I60-I69)	765	3.7	13.2
7	Diabetes mellitus (E10-E14)	511	2.5	8.8
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	425	2.1	7.3
9	Chronic liver disease and cirrhosis (K70,K73-K74)	422	2.0	7.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	309	1.5	5.3
11	Chronic lower respiratory diseases (J40-J47)	279	1.3	4.8
12	Influenza and pneumonia (J10-J18)	276	1.3	4.8
13	Septicemia (A40-A41)	269	1.3	4.6
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	179	0.9	3.1
15	Anemias (D50-D64)	154	0.7	2.7
...	All other causes (Residual)	3,776	18.3	65.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Percent of		
		Number	total deaths	Rate
Black, both sexes, 45-54 years				
...	All causes	34,518	100.0	771.9
1	Malignant neoplasms (C00-C97)	8,245	23.9	184.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,044	23.3	179.9
3	Human immunodeficiency virus (HIV) disease (B20-B24)	2,290	6.6	51.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,133	6.2	47.7
5	Cerebrovascular diseases (I60-I69)	1,906	5.5	42.6
6	Diabetes mellitus (E10-E14)	1,324	3.8	29.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,011	2.9	22.6
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	719	2.1	16.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	706	2.0	15.8
10	Septicemia (A40-A41)	600	1.7	13.4
11	Chronic lower respiratory diseases (J40-J47)	558	1.6	12.5
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	443	1.3	9.9
13	Influenza and pneumonia (J10-J18)	436	1.3	9.7
14	Viral hepatitis (B15-B19)	355	1.0	7.9
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	277	0.8	6.2
...	All other causes (Residual)	5,471	15.8	122.3
Black, both sexes, 55-64 years				
...	All causes	39,976	100.0	1,576.8
1	Malignant neoplasms (C00-C97)	12,323	30.8	486.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,348	28.4	447.6
3	Cerebrovascular diseases (I60-I69)	2,310	5.8	91.1
4	Diabetes mellitus (E10-E14)	2,154	5.4	85.0
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,064	2.7	42.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,054	2.6	41.6
7	Chronic lower respiratory diseases (J40-J47)	1,031	2.6	40.7
8	Septicemia (A40-A41)	814	2.0	32.1
9	Human immunodeficiency virus (HIV) disease (B20-B24)	703	1.8	27.7
10	Chronic liver disease and cirrhosis (K70,K73-K74)	657	1.6	25.9
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	558	1.4	22.0
12	Influenza and pneumonia (J10-J18)	480	1.2	18.9
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	225	0.6	8.9
14	Aortic aneurysm and dissection (I71)	192	0.5	7.6
15	Viral hepatitis (B15-B19)	170	0.4	6.7
...	All other causes (Residual)	4,893	12.2	193.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 65-74 years				
...	All causes	54,379	100.0	3,257.6
1	Malignant neoplasms (C00-C97)	16,547	30.4	991.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	16,120	29.6	965.7
3	Cerebrovascular diseases (I60-I69)	3,714	6.8	222.5
4	Diabetes mellitus (E10-E14)	3,145	5.8	188.4
5	Chronic lower respiratory diseases (J40-J47)	1,985	3.7	118.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,711	3.1	102.5
7	Septicemia (A40-A41)	1,198	2.2	71.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	852	1.6	51.0
9	Influenza and pneumonia (J10-J18)	806	1.5	48.3
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	792	1.5	47.4
11	Chronic liver disease and cirrhosis (K70,K73-K74)	417	0.8	25.0
12	Alzheimer's disease (G30)	254	0.5	15.2
13	Pneumonitis due to solids and liquids (J69)	252	0.5	15.1
14	Aortic aneurysm and dissection (I71)	244	0.4	14.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	239	0.4	14.3
...	All other causes (Residual)	6,103	11.2	365.6
Black, both sexes, 75-84 years				
...	All causes	62,233	100.0	6,585.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	19,924	32.0	2,108.4
2	Malignant neoplasms (C00-C97)	14,590	23.4	1,543.9
3	Cerebrovascular diseases (I60-I69)	5,295	8.5	560.3
4	Diabetes mellitus (E10-E14)	3,168	5.1	335.2
5	Chronic lower respiratory diseases (J40-J47)	2,212	3.6	234.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,910	3.1	202.1
7	Influenza and pneumonia (J10-J18)	1,597	2.6	169.0
8	Septicemia (A40-A41)	1,514	2.4	160.2
9	Alzheimer's disease (G30)	1,178	1.9	124.7
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,064	1.7	112.6
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	836	1.3	88.5
12	Pneumonitis due to solids and liquids (J69)	503	0.8	53.2
13	Aortic aneurysm and dissection (I71)	355	0.6	37.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	298	0.5	31.5
15	Atherosclerosis (I70)	290	0.5	30.7
...	All other causes (Residual)	7,499	12.0	793.6

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Percent of		
		Number	total deaths	Rate
Black, both sexes, 85 years and over				
...	All causes	47,387	100.0	14,550.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	17,181	36.3	5,275.7
2	Malignant neoplasms (C00-C97)	6,291	13.3	1,931.7
3	Cerebrovascular diseases (I60-I69)	4,731	10.0	1,452.7
4	Influenza and pneumonia (J10-J18)	1,893	4.0	581.3
5	Diabetes mellitus (E10-E14)	1,725	3.6	529.7
6	Alzheimer's disease (G30)	1,650	3.5	506.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,399	3.0	429.6
8	Chronic lower respiratory diseases (J40-J47)	1,224	2.6	375.8
9	Septicemia (A40-A41)	1,188	2.5	364.8
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	938	2.0	288.0
11	Pneumonitis due to solids and liquids (J69)	604	1.3	185.5
12	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	598	1.3	183.6
13	Atherosclerosis (I70)	409	0.9	125.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	239	0.5	73.4
15	Parkinson's disease (G20-G21)	192	0.4	59.0
...	All other causes (Residual)	7,125	15.0	2,187.8
Black, male, all ages				
...	All causes	145,908	100.0	823.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	37,016	25.4	209.0
2	Malignant neoplasms (C00-C97)	32,679	22.4	184.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,537	5.9	48.2
4	Cerebrovascular diseases (I60-I69)	7,907	5.4	44.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6,780	4.6	38.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	5,328	3.7	30.1
7	Diabetes mellitus (E10-E14)	5,049	3.5	28.5
8	Chronic lower respiratory diseases (J40-J47)	4,187	2.9	23.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,186	2.2	18.0
10	Influenza and pneumonia (J10-J18)	2,813	1.9	15.9
11	Certain conditions originating in the perinatal period (P00-P96)	2,746	1.9	15.5
12	Septicemia (A40-A41)	2,635	1.8	14.9
13	Chronic liver disease and cirrhosis (K70,K73-K74)	1,843	1.3	10.4
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,715	1.2	9.7
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,627	1.1	9.2
...	All other causes (Residual)	21,860	15.0	123.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Percent of		
		Number	total deaths	Rate
Black, male, 1-4 years				
...	All causes	672	100.0	53.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	194	28.9	15.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	101	15.0	8.0
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	69	10.3	5.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	41	6.1	3.2
5	Malignant neoplasms (C00-C97)	30	4.5	2.4
6	Chronic lower respiratory diseases (J40-J47)	17	2.5	*
7	Influenza and pneumonia (J10-J18)	14	2.1	*
7	Certain conditions originating in the perinatal period (P00-P96)	14	2.1	*
9	Septicemia (A40-A41)	12	1.8	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	8	1.2	*
11	Anemias (D50-D64)	7	1.0	*
12	Cerebrovascular diseases (I60-I69)	5	0.7	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.6	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.4	*
...	All other causes (Residual)	150	22.3	11.9
Black, male, 5-14 years				
...	All causes	960	100.0	27.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	442	46.0	12.8
2	Malignant neoplasms (C00-C97)	102	10.6	2.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	68	7.1	2.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	35	3.6	1.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	35	3.6	1.0
6	Chronic lower respiratory diseases (J40-J47)	32	3.3	0.9
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	32	3.3	0.9
8	Human immunodeficiency virus (HIV) disease (B20-B24)	15	1.6	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	1.4	*
10	Anemias (D50-D64)	10	1.0	*
11	Influenza and pneumonia (J10-J18)	9	0.9	*
12	Diabetes mellitus (E10-E14)	6	0.6	*
13	Cerebrovascular diseases (I60-I69)	5	0.5	*
13	Certain conditions originating in the perinatal period (P00-P96)	5	0.5	*
15	Meningococcal infection (A39)	3	0.3	*
15	Septicemia (A40-A41)	3	0.3	*
15	Meningitis (G00,G03)	3	0.3	*
...	All other causes (Residual)	142	14.8	4.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Percent of		
		Number	total deaths	Rate
Black, male, 15-24 years				
...	All causes	5,479	100.0	180.7
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,600	47.5	85.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,309	23.9	43.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	395	7.2	13.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	209	3.8	6.9
5	Malignant neoplasms (C00-C97)	153	2.8	5.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	72	1.3	2.4
7	Human immunodeficiency virus (HIV) disease (B20-B24)	69	1.3	2.3
8	Chronic lower respiratory diseases (J40-J47)	48	0.9	1.6
9	Anemias (D50-D64)	41	0.7	1.4
10	Diabetes mellitus (E10-E14)	30	0.5	1.0
11	Septicemia (A40-A41)	25	0.5	0.8
12	Legal intervention (Y35,Y89.0)	24	0.4	0.8
13	Influenza and pneumonia (J10-J18)	23	0.4	0.8
14	Cerebrovascular diseases (I60-I69)	20	0.4	0.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	0.2	*
...	All other causes (Residual)	448	8.2	14.8
Black, male, 25-34 years				
...	All causes	6,673	100.0	259.2
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,028	30.4	78.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,220	18.3	47.4
3	Human immunodeficiency virus (HIV) disease (B20-B24)	767	11.5	29.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	586	8.8	22.8
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	425	6.4	16.5
6	Malignant neoplasms (C00-C97)	270	4.0	10.5
7	Diabetes mellitus (E10-E14)	123	1.8	4.8
8	Cerebrovascular diseases (I60-I69)	79	1.2	3.1
9	Anemias (D50-D64)	74	1.1	2.9
10	Chronic lower respiratory diseases (J40-J47)	70	1.0	2.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	56	0.8	2.2
12	Influenza and pneumonia (J10-J18)	53	0.8	2.1
13	Septicemia (A40-A41)	46	0.7	1.8
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	31	0.5	1.2
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	29	0.4	1.1
...	All other causes (Residual)	816	12.2	31.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Percent of		
		Number	total deaths	Rate
Black, male, 35-44 years				
...	All causes	12,135	100.0	445.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,092	17.2	76.7
2	Human immunodeficiency virus (HIV) disease (B20-B24)	2,071	17.1	75.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,633	13.5	59.9
4	Malignant neoplasms (C00-C97)	1,217	10.0	44.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,051	8.7	38.5
6	Cerebrovascular diseases (I60-I69)	361	3.0	13.2
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	336	2.8	12.3
8	Diabetes mellitus (E10-E14)	278	2.3	10.2
9	Chronic liver disease and cirrhosis (K70,K73-K74)	274	2.3	10.0
10	Influenza and pneumonia (J10-J18)	156	1.3	5.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	141	1.2	5.2
12	Septicemia (A40-A41)	128	1.1	4.7
12	Chronic lower respiratory diseases (J40-J47)	128	1.1	4.7
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	104	0.9	3.8
15	Viral hepatitis (B15-B19)	79	0.7	2.9
...	All other causes (Residual)	2,086	17.2	76.5
Black, male, 45-54 years				
...	All causes	20,594	100.0	996.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,016	24.4	242.6
2	Malignant neoplasms (C00-C97)	4,235	20.6	204.8
3	Human immunodeficiency virus (HIV) disease (B20-B24)	1,668	8.1	80.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,597	7.8	77.2
5	Cerebrovascular diseases (I60-I69)	1,008	4.9	48.8
6	Diabetes mellitus (E10-E14)	713	3.5	34.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	713	3.5	34.5
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	556	2.7	26.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	400	1.9	19.3
10	Septicemia (A40-A41)	309	1.5	14.9
11	Chronic lower respiratory diseases (J40-J47)	276	1.3	13.3
12	Influenza and pneumonia (J10-J18)	275	1.3	13.3
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	266	1.3	12.9
14	Viral hepatitis (B15-B19)	243	1.2	11.8
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	204	1.0	9.9
...	All other causes (Residual)	3,115	15.1	150.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Percent of		
		Number	total deaths	Rate
Black, male, 55-64 years				
...	All causes	22,905	100.0	2,025.3
1	Malignant neoplasms (C00-C97)	6,833	29.8	604.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,810	29.7	602.2
3	Cerebrovascular diseases (I60-I69)	1,265	5.5	111.9
4	Diabetes mellitus (E10-E14)	1,043	4.6	92.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	764	3.3	67.6
6	Chronic lower respiratory diseases (J40-J47)	563	2.5	49.8
7	Human immunodeficiency virus (HIV) disease (B20-B24)	533	2.3	47.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	514	2.2	45.4
9	Chronic liver disease and cirrhosis (K70,K73-K74)	469	2.0	41.5
10	Septicemia (A40-A41)	417	1.8	36.9
11	Influenza and pneumonia (J10-J18)	298	1.3	26.3
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	288	1.3	25.5
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	170	0.7	15.0
14	Aortic aneurysm and dissection (I71)	127	0.6	11.2
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	106	0.5	9.4
...	All other causes (Residual)	2,705	11.8	239.2
Black, male, 65-74 years				
...	All causes	28,775	100.0	4,166.6
1	Malignant neoplasms (C00-C97)	9,222	32.0	1,335.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,604	29.9	1,245.8
3	Cerebrovascular diseases (I60-I69)	1,859	6.5	269.2
4	Diabetes mellitus (E10-E14)	1,296	4.5	187.7
5	Chronic lower respiratory diseases (J40-J47)	1,160	4.0	168.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	787	2.7	114.0
7	Septicemia (A40-A41)	591	2.1	85.6
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	517	1.8	74.9
9	Influenza and pneumonia (J10-J18)	479	1.7	69.4
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	376	1.3	54.4
11	Chronic liver disease and cirrhosis (K70,K73-K74)	270	0.9	39.1
12	Human immunodeficiency virus (HIV) disease (B20-B24)	150	0.5	21.7
13	Pneumonitis due to solids and liquids (J69)	141	0.5	20.4
14	Aortic aneurysm and dissection (I71)	138	0.5	20.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	119	0.4	17.2
...	All other causes (Residual)	3,066	10.7	444.0

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Percent of		
		Number	total deaths	Rate
Black, male, 75-84 years				
...	All causes	28,395	100.0	8,355.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,731	30.7	2,569.3
2	Malignant neoplasms (C00-C97)	7,782	27.4	2,290.0
3	Cerebrovascular diseases (I60-I69)	2,086	7.3	613.9
4	Chronic lower respiratory diseases (J40-J47)	1,294	4.6	380.8
5	Diabetes mellitus (E10-E14)	1,162	4.1	341.9
6	Influenza and pneumonia (J10-J18)	823	2.9	242.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	793	2.8	233.4
8	Septicemia (A40-A41)	663	2.3	195.1
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	462	1.6	136.0
10	Alzheimer's disease (G30)	413	1.5	121.5
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	402	1.4	118.3
12	Pneumonitis due to solids and liquids (J69)	256	0.9	75.3
13	Aortic aneurysm and dissection (I71)	161	0.6	47.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	141	0.5	41.5
15	Parkinson's disease (G20-G21)	127	0.4	37.4
...	All other causes (Residual)	3,099	10.9	912.0
Black, male, 85 years and over				
...	All causes	14,489	100.0	16,439.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,812	33.2	5,459.9
2	Malignant neoplasms (C00-C97)	2,829	19.5	3,209.9
3	Cerebrovascular diseases (I60-I69)	1,189	8.2	1,349.1
4	Influenza and pneumonia (J10-J18)	630	4.3	714.8
5	Chronic lower respiratory diseases (J40-J47)	592	4.1	671.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	451	3.1	511.7
7	Diabetes mellitus (E10-E14)	397	2.7	450.5
8	Septicemia (A40-A41)	374	2.6	424.4
9	Alzheimer's disease (G30)	365	2.5	414.1
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	243	1.7	275.7
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	225	1.6	255.3
12	Pneumonitis due to solids and liquids (J69)	215	1.5	243.9
13	Atherosclerosis (I70)	100	0.7	113.5
14	Parkinson's disease (G20-G21)	88	0.6	99.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	78	0.5	88.5
...	All other causes (Residual)	1,901	13.1	2,157.0

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, all ages				
...	All causes	141,801	100.0	727.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	40,658	28.7	208.6
2	Malignant neoplasms (C00-C97)	29,491	20.8	151.3
3	Cerebrovascular diseases (I60-I69)	11,095	7.8	56.9
4	Diabetes mellitus (E10-E14)	7,256	5.1	37.2
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,088	2.9	21.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,925	2.8	20.1
7	Chronic lower respiratory diseases (J40-J47)	3,402	2.4	17.5
8	Septicemia (A40-A41)	3,245	2.3	16.7
9	Influenza and pneumonia (J10-J18)	2,958	2.1	15.2
10	Human immunodeficiency virus (HIV) disease (B20-B24)	2,516	1.8	12.9
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,323	1.6	11.9
12	Alzheimer's disease (G30)	2,204	1.6	11.3
13	Certain conditions originating in the perinatal period (P00-P96)	2,101	1.5	10.8
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,446	1.0	7.4
15	Chronic liver disease and cirrhosis (K70,K73-K74)	932	0.7	4.8
...	All other causes (Residual)	24,161	17.0	124.0
Black, female, 1-4 years				
...	All causes	509	100.0	41.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	145	28.5	11.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	68	13.4	5.6
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	62	12.2	5.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	33	6.5	2.7
5	Malignant neoplasms (C00-C97)	18	3.5	*
6	Influenza and pneumonia (J10-J18)	12	2.4	*
7	Septicemia (A40-A41)	10	2.0	*
8	Certain conditions originating in the perinatal period (P00-P96)	9	1.8	*
9	Cerebrovascular diseases (I60-I69)	7	1.4	*
10	Chronic lower respiratory diseases (J40-J47)	6	1.2	*
11	Anemias (D50-D64)	4	0.8	*
12	Meningococcal infection (A39)	3	0.6	*
12	Meningitis (G00,G03)	3	0.6	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.6	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.4	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.4	*
...	All other causes (Residual)	122	24.0	10.0

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 5-14 years				
...	All causes	627	100.0	18.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	202	32.2	6.0
2	Malignant neoplasms (C00-C97)	74	11.8	2.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	40	6.4	1.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	38	6.1	1.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	26	4.1	0.8
6	Chronic lower respiratory diseases (J40-J47)	22	3.5	0.7
7	Anemias (D50-D64)	16	2.6	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	2.1	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	11	1.8	*
10	Diabetes mellitus (E10-E14)	9	1.4	*
10	Cerebrovascular diseases (I60-I69)	9	1.4	*
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	9	1.4	*
13	Influenza and pneumonia (J10-J18)	8	1.3	*
14	Septicemia (A40-A41)	7	1.1	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	1.1	*
...	All other causes (Residual)	136	21.7	4.1
Black, female, 15-24 years				
...	All causes	1,668	100.0	54.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	433	26.0	14.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	270	16.2	8.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	117	7.0	3.8
4	Malignant neoplasms (C00-C97)	109	6.5	3.6
5	Human immunodeficiency virus (HIV) disease (B20-B24)	94	5.6	3.1
6	Pregnancy, childbirth and the puerperium (O00-099)	45	2.7	1.5
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	42	2.5	1.4
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	38	2.3	1.3
9	Anemias (D50-D64)	32	1.9	1.1
10	Cerebrovascular diseases (I60-I69)	31	1.9	1.0
11	Diabetes mellitus (E10-E14)	30	1.8	1.0
12	Chronic lower respiratory diseases (J40-J47)	29	1.7	1.0
13	Influenza and pneumonia (J10-J18)	22	1.3	0.7
14	Septicemia (A40-A41)	16	1.0	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	0.9	*
...	All other causes (Residual)	345	20.7	11.3

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Percent of		
		Number	total deaths	Rate
Black, female, 25-34 years				
...	All causes	3,355	100.0	117.9
1	Human immunodeficiency virus (HIV) disease (B20-B24)	491	14.6	17.3
2	Malignant neoplasms (C00-C97)	427	12.7	15.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	418	12.5	14.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	384	11.4	13.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	343	10.2	12.1
6	Diabetes mellitus (E10-E14)	79	2.4	2.8
6	Cerebrovascular diseases (I60-I69)	79	2.4	2.8
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	67	2.0	2.4
9	Pregnancy, childbirth and the puerperium (O00-099)	66	2.0	2.3
10	Chronic lower respiratory diseases (J40-J47)	59	1.8	2.1
11	Septicemia (A40-A41)	52	1.5	1.8
12	Anemias (D50-D64)	45	1.3	1.6
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	39	1.2	1.4
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	37	1.1	1.3
15	Influenza and pneumonia (J10-J18)	36	1.1	1.3
...	All other causes (Residual)	733	21.8	25.8
Black, female, 35-44 years				
...	All causes	8,552	100.0	278.0
1	Malignant neoplasms (C00-C97)	1,764	20.6	57.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,359	15.9	44.2
3	Human immunodeficiency virus (HIV) disease (B20-B24)	1,048	12.3	34.1
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	688	8.0	22.4
5	Cerebrovascular diseases (I60-I69)	404	4.7	13.1
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	399	4.7	13.0
7	Diabetes mellitus (E10-E14)	233	2.7	7.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	168	2.0	5.5
9	Chronic lower respiratory diseases (J40-J47)	151	1.8	4.9
10	Chronic liver disease and cirrhosis (K70,K73-K74)	148	1.7	4.8
11	Septicemia (A40-A41)	141	1.6	4.6
12	Influenza and pneumonia (J10-J18)	120	1.4	3.9
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	89	1.0	2.9
14	Anemias (D50-D64)	88	1.0	2.9
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	75	0.9	2.4
...	All other causes (Residual)	1,677	19.6	54.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Percent of		
		Number	total deaths	Rate
Black, female, 45-54 years				
...	All causes	13,924	100.0	579.1
1	Malignant neoplasms (C00-C97)	4,010	28.8	166.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,028	21.7	125.9
3	Cerebrovascular diseases (I60-I69)	898	6.4	37.3
4	Human immunodeficiency virus (HIV) disease (B20-B24)	622	4.5	25.9
5	Diabetes mellitus (E10-E14)	611	4.4	25.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	536	3.8	22.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	306	2.2	12.7
8	Chronic liver disease and cirrhosis (K70,K73-K74)	298	2.1	12.4
9	Septicemia (A40-A41)	291	2.1	12.1
10	Chronic lower respiratory diseases (J40-J47)	282	2.0	11.7
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	177	1.3	7.4
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	163	1.2	6.8
13	Influenza and pneumonia (J10-J18)	161	1.2	6.7
14	Viral hepatitis (B15-B19)	112	0.8	4.7
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	73	0.5	3.0
...	All other causes (Residual)	2,356	16.9	98.0
Black, female, 55-64 years				
...	All causes	17,071	100.0	1,215.5
1	Malignant neoplasms (C00-C97)	5,490	32.2	390.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,538	26.6	323.1
3	Diabetes mellitus (E10-E14)	1,111	6.5	79.1
4	Cerebrovascular diseases (I60-I69)	1,045	6.1	74.4
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	540	3.2	38.5
6	Chronic lower respiratory diseases (J40-J47)	468	2.7	33.3
7	Septicemia (A40-A41)	397	2.3	28.3
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	300	1.8	21.4
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	270	1.6	19.2
10	Chronic liver disease and cirrhosis (K70,K73-K74)	188	1.1	13.4
11	Influenza and pneumonia (J10-J18)	182	1.1	13.0
12	Human immunodeficiency virus (HIV) disease (B20-B24)	170	1.0	12.1
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	75	0.4	5.3
14	Viral hepatitis (B15-B19)	69	0.4	4.9
15	Aortic aneurysm and dissection (I71)	65	0.4	4.6
...	All other causes (Residual)	2,163	12.7	154.0

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Percent of		
		Number	total deaths	Rate
Black, female, 65-74 years				
...	All causes	25,604	100.0	2,616.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,516	29.4	768.0
2	Malignant neoplasms (C00-C97)	7,325	28.6	748.4
3	Cerebrovascular diseases (I60-I69)	1,855	7.2	189.5
4	Diabetes mellitus (E10-E14)	1,849	7.2	188.9
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	924	3.6	94.4
6	Chronic lower respiratory diseases (J40-J47)	825	3.2	84.3
7	Septicemia (A40-A41)	607	2.4	62.0
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	416	1.6	42.5
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	335	1.3	34.2
10	Influenza and pneumonia (J10-J18)	327	1.3	33.4
11	Chronic liver disease and cirrhosis (K70,K73-K74)	147	0.6	15.0
12	Alzheimer's disease (G30)	143	0.6	14.6
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	120	0.5	12.3
14	Pneumonitis due to solids and liquids (J69)	111	0.4	11.3
15	Aortic aneurysm and dissection (I71)	106	0.4	10.8
...	All other causes (Residual)	2,998	11.7	306.3
Black, female, 75-84 years				
...	All causes	33,838	100.0	5,591.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,193	33.1	1,849.6
2	Malignant neoplasms (C00-C97)	6,808	20.1	1,125.0
3	Cerebrovascular diseases (I60-I69)	3,209	9.5	530.3
4	Diabetes mellitus (E10-E14)	2,006	5.9	331.5
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,117	3.3	184.6
6	Chronic lower respiratory diseases (J40-J47)	918	2.7	151.7
7	Septicemia (A40-A41)	851	2.5	140.6
8	Influenza and pneumonia (J10-J18)	774	2.3	127.9
9	Alzheimer's disease (G30)	765	2.3	126.4
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	662	2.0	109.4
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	374	1.1	61.8
12	Pneumonitis due to solids and liquids (J69)	247	0.7	40.8
13	Aortic aneurysm and dissection (I71)	194	0.6	32.1
14	Atherosclerosis (I70)	164	0.5	27.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	157	0.5	25.9
...	All other causes (Residual)	4,399	13.0	726.9

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Percent of		
		Number	total deaths	Rate
Black, female, 85 years and over				
...	All causes	32,898	100.0	13,849.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,369	37.6	5,207.3
2	Cerebrovascular diseases (I60-I69)	3,542	10.8	1,491.2
3	Malignant neoplasms (C00-C97)	3,462	10.5	1,457.5
4	Diabetes mellitus (E10-E14)	1,328	4.0	559.1
5	Alzheimer's disease (G30)	1,285	3.9	541.0
6	Influenza and pneumonia (J10-J18)	1,263	3.8	531.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	948	2.9	399.1
8	Septicemia (A40-A41)	814	2.5	342.7
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	695	2.1	292.6
10	Chronic lower respiratory diseases (J40-J47)	632	1.9	266.1
11	Pneumonitis due to solids and liquids (J69)	389	1.2	163.8
12	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	373	1.1	157.0
13	Atherosclerosis (I70)	309	0.9	130.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	161	0.5	67.8
15	Aortic aneurysm and dissection (I71)	137	0.4	57.7
...	All other causes (Residual)	5,191	15.8	2,185.4
American Indian, both sexes, all ages				
...	All causes	11,977	100.0	392.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,402	20.1	78.6
2	Malignant neoplasms (C00-C97)	2,155	18.0	70.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,361	11.4	44.6
4	Diabetes mellitus (E10-E14)	644	5.4	21.1
5	Cerebrovascular diseases (I60-I69)	574	4.8	18.8
6	Chronic liver disease and cirrhosis (K70,K73-K74)	533	4.5	17.5
7	Chronic lower respiratory diseases (J40-J47)	427	3.6	14.0
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	321	2.7	10.5
9	Influenza and pneumonia (J10-J18)	318	2.7	10.4
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	236	2.0	7.7
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	211	1.8	6.9
12	Septicemia (A40-A41)	155	1.3	5.1
13	Certain conditions originating in the perinatal period (P00-P96)	126	1.1	4.1
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	98	0.8	3.2
15	Alzheimer's disease (G30)	93	0.8	3.0
...	All other causes (Residual)	2,323	19.4	76.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 1-4 years				
...	All causes	103	100.0	48.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	30	29.1	14.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	15	14.6	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	12	11.7	*
4	Malignant neoplasms (C00-C97)	4	3.9	*
5	Septicemia (A40-A41)	3	2.9	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	2.9	*
5	Influenza and pneumonia (J10-J18)	3	2.9	*
5	Acute bronchitis and bronchiolitis (J20-J21)	3	2.9	*
9	Meningitis (G00,G03)	2	1.9	*
10	Meningococcal infection (A39)	1	1.0	*
10	Diseases of appendix (K35-K38)	1	1.0	*
10	Infections of kidney (N10-N12,N13.6,N15.1)	1	1.0	*
10	Certain conditions originating in the perinatal period (P00-P96)	1	1.0	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	1	1.0	*
...	All other causes (Residual)	23	22.3	10.7
American Indian, both sexes, 5-14 years				
...	All causes	129	100.0	22.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	73	56.6	12.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	12	9.3	*
3	Malignant neoplasms (C00-C97)	5	3.9	*
4	Influenza and pneumonia (J10-J18)	4	3.1	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	3.1	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	2.3	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	2.3	*
8	Septicemia (A40-A41)	2	1.6	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.8	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.8	*
9	Anemias (D50-D64)	1	0.8	*
9	Diabetes mellitus (E10-E14)	1	0.8	*
9	Cerebrovascular diseases (I60-I69)	1	0.8	*
9	Chronic lower respiratory diseases (J40-J47)	1	0.8	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.8	*
9	Certain conditions originating in the perinatal period (P00-P96)	1	0.8	*
9	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.8	*
...	All other causes (Residual)	14	10.9	*

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Percent of		
		Number	total deaths	Rate
American Indian, both sexes, 15-24 years				
...	All causes	557	100.0	102.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	293	52.6	53.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	86	15.4	15.8
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	59	10.6	10.9
4	Malignant neoplasms (C00-C97)	22	3.9	4.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	10	1.8	*
6	Diabetes mellitus (E10-E14)	6	1.1	*
6	Chronic liver disease and cirrhosis (K70,K73-K74)	6	1.1	*
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	0.9	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.7	*
10	Septicemia (A40-A41)	3	0.5	*
10	Influenza and pneumonia (J10-J18)	3	0.5	*
10	Chronic lower respiratory diseases (J40-J47)	3	0.5	*
13	Anemias (D50-D64)	2	0.4	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
13	Legal intervention (Y35,Y89.0)	2	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
...	All other causes (Residual)	49	8.8	9.0
American Indian, both sexes, 25-34 years				
...	All causes	596	100.0	132.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	216	36.2	47.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	76	12.8	16.8
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	54	9.1	12.0
4	Malignant neoplasms (C00-C97)	34	5.7	7.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	29	4.9	6.4
6	Chronic liver disease and cirrhosis (K70,K73-K74)	25	4.2	5.5
7	Human immunodeficiency virus (HIV) disease (B20-B24)	11	1.8	*
8	Influenza and pneumonia (J10-J18)	9	1.5	*
9	Cerebrovascular diseases (I60-I69)	5	0.8	*
9	Pregnancy, childbirth and the puerperium (O00-099)	5	0.8	*
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	0.8	*
12	Septicemia (A40-A41)	4	0.7	*
12	Viral hepatitis (B15-B19)	4	0.7	*
12	Diabetes mellitus (E10-E14)	4	0.7	*
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2	0.3	*
15	Chronic lower respiratory diseases (J40-J47)	2	0.3	*
15	Legal intervention (Y35,Y89.0)	2	0.3	*
...	All other causes (Residual)	109	18.3	24.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Percent of		
		Number	total deaths	Rate
American Indian, both sexes, 35-44 years				
...	All causes	1,132	100.0	241.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	260	23.0	55.4
2	Chronic liver disease and cirrhosis (K70,K73-K74)	135	11.9	28.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	125	11.0	26.6
4	Malignant neoplasms (C00-C97)	114	10.1	24.3
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	80	7.1	17.0
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	36	3.2	7.7
7	Human immunodeficiency virus (HIV) disease (B20-B24)	35	3.1	7.5
8	Cerebrovascular diseases (I60-I69)	30	2.7	6.4
9	Influenza and pneumonia (J10-J18)	19	1.7	*
10	Diabetes mellitus (E10-E14)	18	1.6	*
11	Septicemia (A40-A41)	15	1.3	*
12	Viral hepatitis (B15-B19)	13	1.1	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	1.1	*
14	Chronic lower respiratory diseases (J40-J47)	10	0.9	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.5	*
...	All other causes (Residual)	223	19.7	47.5
American Indian, both sexes, 45-54 years				
...	All causes	1,559	100.0	432.8
1	Malignant neoplasms (C00-C97)	282	18.1	78.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	270	17.3	75.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	193	12.4	53.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	187	12.0	51.9
5	Diabetes mellitus (E10-E14)	90	5.8	25.0
6	Cerebrovascular diseases (I60-I69)	50	3.2	13.9
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	39	2.5	10.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	32	2.1	8.9
9	Septicemia (A40-A41)	29	1.9	8.1
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	26	1.7	7.2
11	Viral hepatitis (B15-B19)	22	1.4	6.1
11	Human immunodeficiency virus (HIV) disease (B20-B24)	22	1.4	6.1
13	Influenza and pneumonia (J10-J18)	20	1.3	5.6
14	Chronic lower respiratory diseases (J40-J47)	18	1.2	*
15	Pneumonitis due to solids and liquids (J69)	5	0.3	*
15	Peptic ulcer (K25-K28)	5	0.3	*
...	All other causes (Residual)	269	17.3	74.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 55-64 years				
...	All causes	1,768	100.0	896.2
1	Malignant neoplasms (C00-C97)	470	26.6	238.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	416	23.5	210.9
3	Diabetes mellitus (E10-E14)	142	8.0	72.0
4	Chronic liver disease and cirrhosis (K70,K73-K74)	100	5.7	50.7
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	98	5.5	49.7
6	Chronic lower respiratory diseases (J40-J47)	72	4.1	36.5
7	Cerebrovascular diseases (I60-I69)	54	3.1	27.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	51	2.9	25.9
9	Influenza and pneumonia (J10-J18)	25	1.4	12.7
10	Septicemia (A40-A41)	21	1.2	10.6
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	17	1.0	*
12	Aortic aneurysm and dissection (I71)	8	0.5	*
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	6	0.3	*
13	Peptic ulcer (K25-K28)	6	0.3	*
15	Viral hepatitis (B15-B19)	5	0.3	*
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	0.3	*
...	All other causes (Residual)	272	15.4	137.9
American Indian, both sexes, 65-74 years				
...	All causes	2,206	100.0	2,072.3
1	Malignant neoplasms (C00-C97)	596	27.0	559.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	554	25.1	520.4
3	Diabetes mellitus (E10-E14)	171	7.8	160.6
4	Cerebrovascular diseases (I60-I69)	141	6.4	132.5
5	Chronic lower respiratory diseases (J40-J47)	122	5.5	114.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	61	2.8	57.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	58	2.6	54.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	54	2.4	50.7
9	Influenza and pneumonia (J10-J18)	42	1.9	39.5
10	Septicemia (A40-A41)	39	1.8	36.6
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	0.7	*
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	16	0.7	*
13	Aortic aneurysm and dissection (I71)	12	0.5	*
14	Parkinson's disease (G20-G21)	10	0.5	*
14	Alzheimer's disease (G30)	10	0.5	*
14	Pneumonitis due to solids and liquids (J69)	10	0.5	*
...	All other causes (Residual)	294	13.3	276.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Percent of		
		Number	total deaths	Rate
American Indian, both sexes, 75-84 years				
...	All causes	2,146	100.0	4,218.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	568	26.5	1,116.5
2	Malignant neoplasms (C00-C97)	453	21.1	890.4
3	Cerebrovascular diseases (I60-I69)	160	7.5	314.5
4	Diabetes mellitus (E10-E14)	156	7.3	306.6
5	Chronic lower respiratory diseases (J40-J47)	129	6.0	253.6
6	Influenza and pneumonia (J10-J18)	85	4.0	167.1
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	63	2.9	123.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	52	2.4	102.2
9	Alzheimer's disease (G30)	40	1.9	78.6
10	Septicemia (A40-A41)	21	1.0	41.3
10	Pneumonitis due to solids and liquids (J69)	21	1.0	41.3
12	Chronic liver disease and cirrhosis (K70,K73-K74)	18	0.8	*
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	16	0.7	*
14	Aortic aneurysm and dissection (I71)	15	0.7	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	0.7	*
15	Parkinson's disease (G20-G21)	14	0.7	*
15	Atherosclerosis (I70)	14	0.7	*
...	All other causes (Residual)	307	14.3	603.5
American Indian, both sexes, 85 years and over				
...	All causes	1,402	100.0	8,897.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	415	29.6	2,633.6
2	Malignant neoplasms (C00-C97)	172	12.3	1,091.5
3	Cerebrovascular diseases (I60-I69)	131	9.3	831.3
4	Influenza and pneumonia (J10-J18)	91	6.5	577.5
5	Chronic lower respiratory diseases (J40-J47)	68	4.9	431.5
6	Diabetes mellitus (E10-E14)	56	4.0	355.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	42	3.0	266.5
8	Alzheimer's disease (G30)	41	2.9	260.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	28	2.0	177.7
10	Pneumonitis due to solids and liquids (J69)	18	1.3	*
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	16	1.1	*
11	Atherosclerosis (I70)	16	1.1	*
13	Septicemia (A40-A41)	14	1.0	*
14	Anemias (D50-D64)	10	0.7	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.6	*
15	Parkinson's disease (G20-G21)	9	0.6	*
...	All other causes (Residual)	266	19.0	1,688.0

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Percent of		
		Number	total deaths	Rate
American Indian, male, all ages				
...	All causes	6,466	100.0	424.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,358	21.0	89.1
2	Malignant neoplasms (C00-C97)	1,103	17.1	72.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	908	14.0	59.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	309	4.8	20.3
5	Diabetes mellitus (E10-E14)	276	4.3	18.1
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	259	4.0	17.0
7	Cerebrovascular diseases (I60-I69)	217	3.4	14.2
8	Chronic lower respiratory diseases (J40-J47)	200	3.1	13.1
9	Influenza and pneumonia (J10-J18)	160	2.5	10.5
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	146	2.3	9.6
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	105	1.6	6.9
12	Septicemia (A40-A41)	83	1.3	5.4
13	Certain conditions originating in the perinatal period (P00-P96)	72	1.1	4.7
14	Human immunodeficiency virus (HIV) disease (B20-B24)	57	0.9	3.7
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	46	0.7	3.0
...	All other causes (Residual)	1,167	18.0	76.6
American Indian, male, 1-4 years				
...	All causes	52	100.0	47.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15	28.8	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	9.6	*
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	9.6	*
4	Malignant neoplasms (C00-C97)	3	5.8	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	3.8	*
5	Acute bronchitis and bronchiolitis (J20-J21)	2	3.8	*
7	Meningococcal infection (A39)	1	1.9	*
7	Septicemia (A40-A41)	1	1.9	*
7	Meningitis (G00,G03)	1	1.9	*
7	Diseases of appendix (K35-K38)	1	1.9	*
7	Certain conditions originating in the perinatal period (P00-P96)	1	1.9	*
...	All other causes (Residual)	15	28.8	*

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Percent of		
		Number	total deaths	Rate
American Indian, male, 5-14 years				
...	All causes	75	100.0	25.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	46	61.3	15.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7	9.3	*
3	Malignant neoplasms (C00-C97)	3	4.0	*
4	Septicemia (A40-A41)	2	2.7	*
5	Human immunodeficiency virus (HIV) disease (B20-B24)	1	1.3	*
5	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.3	*
5	Diabetes mellitus (E10-E14)	1	1.3	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	1.3	*
5	Cerebrovascular diseases (I60-I69)	1	1.3	*
5	Influenza and pneumonia (J10-J18)	1	1.3	*
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	1.3	*
5	Certain conditions originating in the perinatal period (P00-P96)	1	1.3	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	1.3	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1	1.3	*
5	Complications of medical and surgical care (Y40-Y84,Y88)	1	1.3	*
...	All other causes (Residual)	6	8.0	*
American Indian, male, 15-24 years				
...	All causes	400	100.0	142.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	203	50.8	72.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	69	17.3	24.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	45	11.3	16.1
4	Malignant neoplasms (C00-C97)	15	3.8	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	8	2.0	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	1.3	*
7	Chronic liver disease and cirrhosis (K70,K73-K74)	4	1.0	*
8	Diabetes mellitus (E10-E14)	3	0.8	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.5	*
9	Chronic lower respiratory diseases (J40-J47)	2	0.5	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.5	*
9	Legal intervention (Y35,Y89.0)	2	0.5	*
13	Septicemia (A40-A41)	1	0.3	*
13	Viral hepatitis (B15-B19)	1	0.3	*
13	Meningitis (G00,G03)	1	0.3	*
13	Influenza and pneumonia (J10-J18)	1	0.3	*
13	Pneumonitis due to solids and liquids (J69)	1	0.3	*
...	All other causes (Residual)	35	8.8	12.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Percent of		
		Number	total deaths	Rate
American Indian, male, 25-34 years				
...	All causes	417	100.0	180.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	163	39.1	70.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	63	15.1	27.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	39	9.4	16.8
4	Malignant neoplasms (C00-C97)	19	4.6	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	18	4.3	*
6	Chronic liver disease and cirrhosis (K70,K73-K74)	17	4.1	*
7	Human immunodeficiency virus (HIV) disease (B20-B24)	10	2.4	*
8	Influenza and pneumonia (J10-J18)	6	1.4	*
9	Septicemia (A40-A41)	2	0.5	*
9	Viral hepatitis (B15-B19)	2	0.5	*
9	Diabetes mellitus (E10-E14)	2	0.5	*
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1	0.2	*
12	Cerebrovascular diseases (I60-I69)	1	0.2	*
12	Chronic lower respiratory diseases (J40-J47)	1	0.2	*
12	Pneumonitis due to solids and liquids (J69)	1	0.2	*
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.2	*
12	Legal intervention (Y35,Y89.0)	1	0.2	*
...	All other causes (Residual)	70	16.8	30.2
American Indian, male, 35-44 years				
...	All causes	722	100.0	311.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	188	26.0	81.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	85	11.8	36.7
3	Chronic liver disease and cirrhosis (K70,K73-K74)	82	11.4	35.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	65	9.0	28.0
5	Malignant neoplasms (C00-C97)	53	7.3	22.9
6	Human immunodeficiency virus (HIV) disease (B20-B24)	24	3.3	10.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	24	3.3	10.4
8	Cerebrovascular diseases (I60-I69)	13	1.8	*
8	Influenza and pneumonia (J10-J18)	13	1.8	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	1.1	*
11	Septicemia (A40-A41)	7	1.0	*
11	Diabetes mellitus (E10-E14)	7	1.0	*
13	Viral hepatitis (B15-B19)	6	0.8	*
14	Legal intervention (Y35,Y89.0)	5	0.7	*
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	3	0.4	*
15	Chronic lower respiratory diseases (J40-J47)	3	0.4	*
...	All other causes (Residual)	136	18.8	58.7

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 45-54 years				
...	All causes	940	100.0	536.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	191	20.3	109.1
2	Malignant neoplasms (C00-C97)	135	14.4	77.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	130	13.8	74.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	99	10.5	56.5
5	Diabetes mellitus (E10-E14)	51	5.4	29.1
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	33	3.5	18.8
7	Cerebrovascular diseases (I60-I69)	22	2.3	12.6
8	Human immunodeficiency virus (HIV) disease (B20-B24)	19	2.0	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19	2.0	*
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	19	2.0	*
11	Septicemia (A40-A41)	17	1.8	*
12	Viral hepatitis (B15-B19)	16	1.7	*
13	Influenza and pneumonia (J10-J18)	12	1.3	*
14	Chronic lower respiratory diseases (J40-J47)	8	0.9	*
15	Peptic ulcer (K25-K28)	4	0.4	*
...	All other causes (Residual)	165	17.6	94.2
American Indian, male, 55-64 years				
...	All causes	1,004	100.0	1,053.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	287	28.6	301.1
2	Malignant neoplasms (C00-C97)	244	24.3	256.0
3	Chronic liver disease and cirrhosis (K70,K73-K74)	68	6.8	71.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	68	6.8	71.3
5	Diabetes mellitus (E10-E14)	65	6.5	68.2
6	Chronic lower respiratory diseases (J40-J47)	34	3.4	35.7
7	Cerebrovascular diseases (I60-I69)	23	2.3	24.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	1.6	*
9	Influenza and pneumonia (J10-J18)	13	1.3	*
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	13	1.3	*
11	Septicemia (A40-A41)	11	1.1	*
12	Aortic aneurysm and dissection (I71)	6	0.6	*
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	0.5	*
14	Viral hepatitis (B15-B19)	3	0.3	*
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	3	0.3	*
14	Atherosclerosis (I70)	3	0.3	*
14	Peptic ulcer (K25-K28)	3	0.3	*
...	All other causes (Residual)	139	13.8	145.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Percent of		
		Number	total deaths	Rate
American Indian, male, 65-74 years				
...	All causes	1,165	100.0	2,393.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	332	28.5	682.1
2	Malignant neoplasms (C00-C97)	328	28.2	673.9
3	Diabetes mellitus (E10-E14)	68	5.8	139.7
4	Cerebrovascular diseases (I60-I69)	64	5.5	131.5
5	Chronic lower respiratory diseases (J40-J47)	56	4.8	115.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	33	2.8	67.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	29	2.5	59.6
8	Influenza and pneumonia (J10-J18)	23	2.0	47.3
9	Septicemia (A40-A41)	20	1.7	41.1
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19	1.6	*
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	8	0.7	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.6	*
12	Aortic aneurysm and dissection (I71)	7	0.6	*
14	Parkinson's disease (G20-G21)	6	0.5	*
15	Atherosclerosis (I70)	5	0.4	*
15	Peptic ulcer (K25-K28)	5	0.4	*
...	All other causes (Residual)	155	13.3	318.4
American Indian, male, 75-84 years				
...	All causes	983	100.0	4,775.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	285	29.0	1,384.5
2	Malignant neoplasms (C00-C97)	225	22.9	1,093.0
3	Diabetes mellitus (E10-E14)	63	6.4	306.0
3	Chronic lower respiratory diseases (J40-J47)	63	6.4	306.0
5	Cerebrovascular diseases (I60-I69)	51	5.2	247.8
6	Influenza and pneumonia (J10-J18)	42	4.3	204.0
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	34	3.5	165.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	24	2.4	116.6
9	Alzheimer's disease (G30)	17	1.7	*
10	Septicemia (A40-A41)	13	1.3	*
11	Parkinson's disease (G20-G21)	10	1.0	*
12	Aortic aneurysm and dissection (I71)	8	0.8	*
12	Pneumonitis due to solids and liquids (J69)	8	0.8	*
12	Chronic liver disease and cirrhosis (K70,K73-K74)	8	0.8	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.6	*
15	Atherosclerosis (I70)	6	0.6	*
...	All other causes (Residual)	120	12.2	582.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 85 years and over				
...	All causes	492	100.0	9,758.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	146	29.7	2,895.7
2	Malignant neoplasms (C00-C97)	75	15.2	1,487.5
3	Cerebrovascular diseases (I60-I69)	42	8.5	833.0
4	Influenza and pneumonia (J10-J18)	38	7.7	753.7
5	Chronic lower respiratory diseases (J40-J47)	31	6.3	614.8
6	Diabetes mellitus (E10-E14)	16	3.3	*
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	2.8	*
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14	2.8	*
9	Septicemia (A40-A41)	7	1.4	*
9	Alzheimer's disease (G30)	7	1.4	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	1.0	*
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	5	1.0	*
11	Atherosclerosis (I70)	5	1.0	*
14	Aortic aneurysm and dissection (I71)	4	0.8	*
14	Pneumonitis due to solids and liquids (J69)	4	0.8	*
...	All other causes (Residual)	79	16.1	1,566.8
American Indian, female, all ages				
...	All causes	5,511	100.0	360.2
1	Malignant neoplasms (C00-C97)	1,052	19.1	68.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,044	18.9	68.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	453	8.2	29.6
4	Diabetes mellitus (E10-E14)	368	6.7	24.1
5	Cerebrovascular diseases (I60-I69)	357	6.5	23.3
6	Chronic lower respiratory diseases (J40-J47)	227	4.1	14.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	224	4.1	14.6
8	Influenza and pneumonia (J10-J18)	158	2.9	10.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	131	2.4	8.6
10	Septicemia (A40-A41)	72	1.3	4.7
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	65	1.2	4.2
12	Alzheimer's disease (G30)	64	1.2	4.2
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	62	1.1	4.1
14	Certain conditions originating in the perinatal period (P00-P96)	54	1.0	3.5
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	52	0.9	3.4
...	All other causes (Residual)	1,128	20.5	73.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 1-4 years				
...	All causes	51	100.0	48.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15	29.4	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	10	19.6	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7	13.7	*
4	Influenza and pneumonia (J10-J18)	3	5.9	*
5	Septicemia (A40-A41)	2	3.9	*
6	Malignant neoplasms (C00-C97)	1	2.0	*
6	Meningitis (G00,G03)	1	2.0	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	2.0	*
6	Acute bronchitis and bronchiolitis (J20-J21)	1	2.0	*
6	Infections of kidney (N10-N12,N13.6,N15.1)	1	2.0	*
6	Complications of medical and surgical care (Y40-Y84,Y88)	1	2.0	*
...	All other causes (Residual)	8	15.7	*
American Indian, female, 5-14 years				
...	All causes	54	100.0	18.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	27	50.0	9.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5	9.3	*
3	Influenza and pneumonia (J10-J18)	3	5.6	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	5.6	*
5	Malignant neoplasms (C00-C97)	2	3.7	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	3.7	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	3.7	*
8	Anemias (D50-D64)	1	1.9	*
8	Chronic lower respiratory diseases (J40-J47)	1	1.9	*
...	All other causes (Residual)	8	14.8	*

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 15-24 years				
...	All causes	157	100.0	59.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	90	57.3	34.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	17	10.8	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	14	8.9	*
4	Malignant neoplasms (C00-C97)	7	4.5	*
5	Diabetes mellitus (E10-E14)	3	1.9	*
6	Septicemia (A40-A41)	2	1.3	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	1.3	*
6	Anemias (D50-D64)	2	1.3	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	1.3	*
6	Influenza and pneumonia (J10-J18)	2	1.3	*
6	Chronic liver disease and cirrhosis (K70,K73-K74)	2	1.3	*
6	Complications of medical and surgical care (Y40-Y84,Y88)	2	1.3	*
13	Chronic lower respiratory diseases (J40-J47)	1	0.6	*
13	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.6	*
...	All other causes (Residual)	10	6.4	*
American Indian, female, 25-34 years				
...	All causes	179	100.0	81.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	53	29.6	24.1
2	Malignant neoplasms (C00-C97)	15	8.4	*
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	15	8.4	*
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	13	7.3	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	11	6.1	*
6	Chronic liver disease and cirrhosis (K70,K73-K74)	8	4.5	*
7	Pregnancy, childbirth and the puerperium (O00-099)	5	2.8	*
8	Cerebrovascular diseases (I60-I69)	4	2.2	*
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	2.2	*
10	Influenza and pneumonia (J10-J18)	3	1.7	*
11	Septicemia (A40-A41)	2	1.1	*
11	Viral hepatitis (B15-B19)	2	1.1	*
11	Diabetes mellitus (E10-E14)	2	1.1	*
14	Tuberculosis (A16-A19)	1	0.6	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.6	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.6	*
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1	0.6	*
14	Chronic lower respiratory diseases (J40-J47)	1	0.6	*
14	Peptic ulcer (K25-K28)	1	0.6	*
14	Diseases of appendix (K35-K38)	1	0.6	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.6	*
14	Legal intervention (Y35,Y89.0)	1	0.6	*
...	All other causes (Residual)	33	18.4	15.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Percent of		
		Number	total deaths	Rate
American Indian, female, 35-44 years				
...	All causes	410	100.0	172.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	72	17.6	30.3
2	Malignant neoplasms (C00-C97)	61	14.9	25.7
3	Chronic liver disease and cirrhosis (K70,K73-K74)	53	12.9	22.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	40	9.8	16.8
5	Cerebrovascular diseases (I60-I69)	17	4.1	*
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	15	3.7	*
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	12	2.9	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	11	2.7	*
8	Diabetes mellitus (E10-E14)	11	2.7	*
10	Septicemia (A40-A41)	8	2.0	*
11	Viral hepatitis (B15-B19)	7	1.7	*
11	Chronic lower respiratory diseases (J40-J47)	7	1.7	*
13	Influenza and pneumonia (J10-J18)	6	1.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	1.2	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	1.0	*
...	All other causes (Residual)	81	19.8	34.1
American Indian, female, 45-54 years				
...	All causes	619	100.0	334.3
1	Malignant neoplasms (C00-C97)	147	23.7	79.4
2	Chronic liver disease and cirrhosis (K70,K73-K74)	88	14.2	47.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	79	12.8	42.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	63	10.2	34.0
5	Diabetes mellitus (E10-E14)	39	6.3	21.1
6	Cerebrovascular diseases (I60-I69)	28	4.5	15.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	2.1	*
8	Septicemia (A40-A41)	12	1.9	*
9	Chronic lower respiratory diseases (J40-J47)	10	1.6	*
10	Influenza and pneumonia (J10-J18)	8	1.3	*
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7	1.1	*
12	Viral hepatitis (B15-B19)	6	1.0	*
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6	1.0	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	3	0.5	*
14	Pneumonitis due to solids and liquids (J69)	3	0.5	*
...	All other causes (Residual)	107	17.3	57.8

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Percent of		
		Number	total deaths	Rate
American Indian, female, 55-64 years				
...	All causes	764	100.0	749.4
1	Malignant neoplasms (C00-C97)	226	29.6	221.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	129	16.9	126.5
3	Diabetes mellitus (E10-E14)	77	10.1	75.5
4	Chronic lower respiratory diseases (J40-J47)	38	5.0	37.3
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	35	4.6	34.3
6	Chronic liver disease and cirrhosis (K70,K73-K74)	32	4.2	31.4
7	Cerebrovascular diseases (I60-I69)	31	4.1	30.4
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	30	3.9	29.4
9	Influenza and pneumonia (J10-J18)	12	1.6	*
10	Septicemia (A40-A41)	10	1.3	*
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4	0.5	*
12	Parkinson's disease (G20-G21)	3	0.4	*
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	3	0.4	*
12	Peptic ulcer (K25-K28)	3	0.4	*
15	Viral hepatitis (B15-B19)	2	0.3	*
15	Aortic aneurysm and dissection (I71)	2	0.3	*
...	All other causes (Residual)	127	16.6	124.6
American Indian, female, 65-74 years				
...	All causes	1,041	100.0	1,801.7
1	Malignant neoplasms (C00-C97)	268	25.7	463.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	222	21.3	384.2
3	Diabetes mellitus (E10-E14)	103	9.9	178.3
4	Cerebrovascular diseases (I60-I69)	77	7.4	133.3
5	Chronic lower respiratory diseases (J40-J47)	66	6.3	114.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	35	3.4	60.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	29	2.8	50.2
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	28	2.7	48.5
9	Septicemia (A40-A41)	19	1.8	*
9	Influenza and pneumonia (J10-J18)	19	1.8	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.9	*
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	8	0.8	*
13	Viral hepatitis (B15-B19)	6	0.6	*
13	Alzheimer's disease (G30)	6	0.6	*
13	Pneumonitis due to solids and liquids (J69)	6	0.6	*
...	All other causes (Residual)	140	13.4	242.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Percent of		
		Number	total deaths	Rate
American Indian, female, 75-84 years				
...	All causes	1,163	100.0	3,839.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	283	24.3	934.3
2	Malignant neoplasms (C00-C97)	228	19.6	752.7
3	Cerebrovascular diseases (I60-I69)	109	9.4	359.9
4	Diabetes mellitus (E10-E14)	93	8.0	307.0
5	Chronic lower respiratory diseases (J40-J47)	66	5.7	217.9
6	Influenza and pneumonia (J10-J18)	43	3.7	142.0
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	29	2.5	95.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	28	2.4	92.4
9	Alzheimer's disease (G30)	23	2.0	75.9
10	Pneumonitis due to solids and liquids (J69)	13	1.1	*
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	12	1.0	*
12	Chronic liver disease and cirrhosis (K70,K73-K74)	10	0.9	*
13	Septicemia (A40-A41)	8	0.7	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.7	*
13	Atherosclerosis (I70)	8	0.7	*
...	All other causes (Residual)	202	17.4	666.9
American Indian, female, 85 years and over				
...	All causes	910	100.0	8,492.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	269	29.6	2,510.3
2	Malignant neoplasms (C00-C97)	97	10.7	905.2
3	Cerebrovascular diseases (I60-I69)	89	9.8	830.5
4	Influenza and pneumonia (J10-J18)	53	5.8	494.6
5	Diabetes mellitus (E10-E14)	40	4.4	373.3
6	Chronic lower respiratory diseases (J40-J47)	37	4.1	345.3
7	Alzheimer's disease (G30)	34	3.7	317.3
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	28	3.1	261.3
9	Pneumonitis due to solids and liquids (J69)	14	1.5	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	1.5	*
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	11	1.2	*
11	Atherosclerosis (I70)	11	1.2	*
13	Anemias (D50-D64)	8	0.9	*
13	Nutritional deficiencies (E40-E64)	8	0.9	*
15	Septicemia (A40-A41)	7	0.8	*
15	Parkinson's disease (G20-G21)	7	0.8	*
...	All other causes (Residual)	183	20.1	1,707.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, all ages				
...	All causes	37,048	100.0	303.8
1	Malignant neoplasms (C00-C97)	9,792	26.4	80.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,428	25.4	77.3
3	Cerebrovascular diseases (I60-I69)	3,497	9.4	28.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,750	4.7	14.4
5	Diabetes mellitus (E10-E14)	1,243	3.4	10.2
6	Chronic lower respiratory diseases (J40-J47)	1,178	3.2	9.7
7	Influenza and pneumonia (J10-J18)	1,171	3.2	9.6
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	634	1.7	5.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	625	1.7	5.1
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	543	1.5	4.5
11	Septicemia (A40-A41)	397	1.1	3.3
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	397	1.1	3.3
13	Certain conditions originating in the perinatal period (P00-P96)	363	1.0	3.0
14	Chronic liver disease and cirrhosis (K70,K73-K74)	319	0.9	2.6
15	Alzheimer's disease (G30)	297	0.8	2.4
...	All other causes (Residual)	5,414	14.6	44.4
Asian or Pacific Islander, both sexes, 1-4 years				
...	All causes	153	100.0	22.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	36	23.5	5.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	24	15.7	3.5
3	Malignant neoplasms (C00-C97)	15	9.8	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	7	4.6	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	3.9	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	3.3	*
6	Influenza and pneumonia (J10-J18)	5	3.3	*
8	Septicemia (A40-A41)	3	2.0	*
8	Cerebrovascular diseases (I60-I69)	3	2.0	*
10	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.7	*
10	Certain conditions originating in the perinatal period (P00-P96)	1	0.7	*
...	All other causes (Residual)	47	30.7	6.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 5-14 years				
...	All causes	209	100.0	12.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	59	28.2	3.5
2	Malignant neoplasms (C00-C97)	47	22.5	2.8
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	16	7.7	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	10	4.8	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	9	4.3	*
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8	3.8	*
7	Cerebrovascular diseases (I60-I69)	6	2.9	*
8	Influenza and pneumonia (J10-J18)	4	1.9	*
9	Septicemia (A40-A41)	3	1.4	*
9	Certain conditions originating in the perinatal period (P00-P96)	3	1.4	*
11	Diabetes mellitus (E10-E14)	2	1.0	*
11	Meningitis (G00,G03)	2	1.0	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.5	*
13	Anemias (D50-D64)	1	0.5	*
13	Chronic lower respiratory diseases (J40-J47)	1	0.5	*
13	Hernia (K40-K46)	1	0.5	*
...	All other causes (Residual)	36	17.2	2.1
Asian or Pacific Islander, both sexes, 15-24 years				
...	All causes	800	100.0	43.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	338	42.3	18.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	118	14.8	6.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	104	13.0	5.6
4	Malignant neoplasms (C00-C97)	61	7.6	3.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	24	3.0	1.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	12	1.5	*
7	Septicemia (A40-A41)	7	0.9	*
7	Cerebrovascular diseases (I60-I69)	7	0.9	*
7	Influenza and pneumonia (J10-J18)	7	0.9	*
10	Chronic lower respiratory diseases (J40-J47)	6	0.8	*
11	Tuberculosis (A16-A19)	3	0.4	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.4	*
11	Diabetes mellitus (E10-E14)	3	0.4	*
14	Meningococcal infection (A39)	2	0.3	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.3	*
14	Meningitis (G00,G03)	2	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.3	*
...	All other causes (Residual)	99	12.4	5.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 25-34 years				
...	All causes	1,065	100.0	47.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	258	24.2	11.5
2	Malignant neoplasms (C00-C97)	169	15.9	7.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	156	14.6	6.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	146	13.7	6.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	75	7.0	3.3
6	Cerebrovascular diseases (I60-I69)	22	2.1	1.0
7	Human immunodeficiency virus (HIV) disease (B20-B24)	18	1.7	*
8	Septicemia (A40-A41)	10	0.9	*
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	10	0.9	*
10	Influenza and pneumonia (J10-J18)	9	0.8	*
11	Chronic liver disease and cirrhosis (K70,K73-K74)	8	0.8	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.6	*
12	Diabetes mellitus (E10-E14)	6	0.6	*
12	Legal intervention (Y35,Y89.0)	6	0.6	*
15	Pregnancy, childbirth and the puerperium (O00-099)	5	0.5	*
...	All other causes (Residual)	161	15.1	7.2
Asian or Pacific Islander, both sexes, 35-44 years				
...	All causes	1,718	100.0	84.2
1	Malignant neoplasms (C00-C97)	515	30.0	25.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	240	14.0	11.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	221	12.9	10.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	113	6.6	5.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	110	6.4	5.4
6	Cerebrovascular diseases (I60-I69)	91	5.3	4.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	34	2.0	1.7
8	Human immunodeficiency virus (HIV) disease (B20-B24)	32	1.9	1.6
9	Diabetes mellitus (E10-E14)	30	1.7	1.5
10	Viral hepatitis (B15-B19)	17	1.0	*
11	Chronic lower respiratory diseases (J40-J47)	15	0.9	*
12	Septicemia (A40-A41)	14	0.8	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	0.8	*
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	14	0.8	*
15	Pregnancy, childbirth and the puerperium (O00-099)	12	0.7	*
...	All other causes (Residual)	246	14.3	12.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Percent of		
		Number	total deaths	Rate
Asian or Pacific Islander, both sexes, 45-54 years				
...	All causes	3,289	100.0	202.7
1	Malignant neoplasms (C00-C97)	1,349	41.0	83.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	613	18.6	37.8
3	Cerebrovascular diseases (I60-I69)	218	6.6	13.4
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	193	5.9	11.9
5	Diabetes mellitus (E10-E14)	87	2.6	5.4
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	86	2.6	5.3
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	83	2.5	5.1
8	Chronic liver disease and cirrhosis (K70,K73-K74)	72	2.2	4.4
9	Viral hepatitis (B15-B19)	51	1.6	3.1
10	Chronic lower respiratory diseases (J40-J47)	45	1.4	2.8
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	41	1.2	2.5
12	Septicemia (A40-A41)	36	1.1	2.2
13	Influenza and pneumonia (J10-J18)	29	0.9	1.8
14	Aortic aneurysm and dissection (I71)	25	0.8	1.5
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	23	0.7	1.4
...	All other causes (Residual)	338	10.3	20.8
Asian or Pacific Islander, both sexes, 55-64 years				
...	All causes	4,542	100.0	495.3
1	Malignant neoplasms (C00-C97)	1,827	40.2	199.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,003	22.1	109.4
3	Cerebrovascular diseases (I60-I69)	370	8.1	40.4
4	Diabetes mellitus (E10-E14)	181	4.0	19.7
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	129	2.8	14.1
6	Chronic lower respiratory diseases (J40-J47)	96	2.1	10.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	78	1.7	8.5
8	Chronic liver disease and cirrhosis (K70,K73-K74)	69	1.5	7.5
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	63	1.4	6.9
10	Viral hepatitis (B15-B19)	61	1.3	6.7
11	Influenza and pneumonia (J10-J18)	58	1.3	6.3
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	44	1.0	4.8
13	Septicemia (A40-A41)	36	0.8	3.9
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	35	0.8	3.8
15	Aortic aneurysm and dissection (I71)	27	0.6	2.9
...	All other causes (Residual)	465	10.2	50.7

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Percent of		
		Number	total deaths	Rate
Asian or Pacific Islander, both sexes, 65-74 years				
...	All causes	7,247	100.0	1,275.9
1	Malignant neoplasms (C00-C97)	2,471	34.1	435.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,861	25.7	327.6
3	Cerebrovascular diseases (I60-I69)	696	9.6	122.5
4	Diabetes mellitus (E10-E14)	330	4.6	58.1
5	Chronic lower respiratory diseases (J40-J47)	259	3.6	45.6
6	Influenza and pneumonia (J10-J18)	158	2.2	27.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	144	2.0	25.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	133	1.8	23.4
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	85	1.2	15.0
10	Septicemia (A40-A41)	84	1.2	14.8
11	Viral hepatitis (B15-B19)	80	1.1	14.1
11	Aortic aneurysm and dissection (I71)	80	1.1	14.1
13	Chronic liver disease and cirrhosis (K70,K73-K74)	72	1.0	12.7
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	53	0.7	9.3
15	Parkinson's disease (G20-G21)	35	0.5	6.2
...	All other causes (Residual)	706	9.7	124.3
Asian or Pacific Islander, both sexes, 75-84 years				
...	All causes	9,798	100.0	3,385.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,975	30.4	1,027.9
2	Malignant neoplasms (C00-C97)	2,397	24.5	828.2
3	Cerebrovascular diseases (I60-I69)	1,102	11.2	380.8
4	Chronic lower respiratory diseases (J40-J47)	425	4.3	146.8
5	Diabetes mellitus (E10-E14)	411	4.2	142.0
6	Influenza and pneumonia (J10-J18)	374	3.8	129.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	201	2.1	69.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	199	2.0	68.8
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	137	1.4	47.3
10	Septicemia (A40-A41)	108	1.1	37.3
11	Parkinson's disease (G20-G21)	101	1.0	34.9
12	Alzheimer's disease (G30)	98	1.0	33.9
13	Aortic aneurysm and dissection (I71)	88	0.9	30.4
14	Pneumonitis due to solids and liquids (J69)	63	0.6	21.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	50	0.5	17.3
...	All other causes (Residual)	1,069	10.9	369.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Percent of		
		Number	total deaths	Rate
Asian or Pacific Islander, both sexes, 85 years and over				
...	All causes	7,477	100.0	9,714.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,602	34.8	3,380.8
2	Cerebrovascular diseases (I60-I69)	979	13.1	1,272.0
3	Malignant neoplasms (C00-C97)	938	12.5	1,218.7
4	Influenza and pneumonia (J10-J18)	508	6.8	660.0
5	Chronic lower respiratory diseases (J40-J47)	327	4.4	424.9
6	Diabetes mellitus (E10-E14)	193	2.6	250.8
7	Alzheimer's disease (G30)	175	2.3	227.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	153	2.0	198.8
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	152	2.0	197.5
10	Pneumonitis due to solids and liquids (J69)	117	1.6	152.0
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	108	1.4	140.3
12	Septicemia (A40-A41)	85	1.1	110.4
13	Parkinson's disease (G20-G21)	56	0.7	72.8
14	Atherosclerosis (I70)	54	0.7	70.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	51	0.7	66.3
...	All other causes (Residual)	979	13.1	1,272.0
Asian or Pacific Islander, male, all ages				
...	All causes	19,829	100.0	335.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,165	26.0	87.3
2	Malignant neoplasms (C00-C97)	5,147	26.0	87.0
3	Cerebrovascular diseases (I60-I69)	1,625	8.2	27.5
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,122	5.7	19.0
5	Chronic lower respiratory diseases (J40-J47)	749	3.8	12.7
6	Influenza and pneumonia (J10-J18)	625	3.2	10.6
7	Diabetes mellitus (E10-E14)	599	3.0	10.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	458	2.3	7.7
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	375	1.9	6.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	320	1.6	5.4
11	Septicemia (A40-A41)	218	1.1	3.7
12	Certain conditions originating in the perinatal period (P00-P96)	204	1.0	3.4
13	Chronic liver disease and cirrhosis (K70,K73-K74)	193	1.0	3.3
14	Aortic aneurysm and dissection (I71)	189	1.0	3.2
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	163	0.8	2.8
...	All other causes (Residual)	2,677	13.5	45.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 1-4 years				
...	All causes	81	100.0	23.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	21	25.9	6.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	13	16.0	*
3	Malignant neoplasms (C00-C97)	7	8.6	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	4.9	*
5	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	3.7	*
6	Cerebrovascular diseases (I60-I69)	2	2.5	*
6	Influenza and pneumonia (J10-J18)	2	2.5	*
8	Septicemia (A40-A41)	1	1.2	*
8	Infections of kidney (N10-N12,N13.6,N15.1)	1	1.2	*
8	Certain conditions originating in the perinatal period (P00-P96)	1	1.2	*
...	All other causes (Residual)	26	32.1	7.6
Asian or Pacific Islander, male, 5-14 years				
...	All causes	119	100.0	13.6
1	Malignant neoplasms (C00-C97)	30	25.2	3.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	30	25.2	3.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7	5.9	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	4.2	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	4.2	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	3.4	*
7	Influenza and pneumonia (J10-J18)	3	2.5	*
8	Septicemia (A40-A41)	2	1.7	*
8	Meningitis (G00,G03)	2	1.7	*
8	Cerebrovascular diseases (I60-I69)	2	1.7	*
8	Certain conditions originating in the perinatal period (P00-P96)	2	1.7	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.8	*
12	Anemias (D50-D64)	1	0.8	*
12	Diabetes mellitus (E10-E14)	1	0.8	*
12	Chronic lower respiratory diseases (J40-J47)	1	0.8	*
...	All other causes (Residual)	23	19.3	2.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Percent of		
		Number	total deaths	Rate
Asian or Pacific Islander, male, 15-24 years				
...	All causes	555	100.0	59.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	253	45.6	27.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	85	15.3	9.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	85	15.3	9.1
4	Malignant neoplasms (C00-C97)	32	5.8	3.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	16	2.9	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	1.3	*
7	Chronic lower respiratory diseases (J40-J47)	4	0.7	*
8	Septicemia (A40-A41)	3	0.5	*
8	Influenza and pneumonia (J10-J18)	3	0.5	*
10	Tuberculosis (A16-A19)	2	0.4	*
10	Diabetes mellitus (E10-E14)	2	0.4	*
10	Cerebrovascular diseases (I60-I69)	2	0.4	*
13	Meningococcal infection (A39)	1	0.2	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.2	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.2	*
13	Meningitis (G00,G03)	1	0.2	*
13	Aortic aneurysm and dissection (I71)	1	0.2	*
13	Pneumonitis due to solids and liquids (J69)	1	0.2	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
...	All other causes (Residual)	53	9.5	5.7
Asian or Pacific Islander, male, 25-34 years				
...	All causes	689	100.0	62.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	191	27.7	17.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	107	15.5	9.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	102	14.8	9.2
4	Malignant neoplasms (C00-C97)	78	11.3	7.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	56	8.1	5.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	16	2.3	*
7	Cerebrovascular diseases (I60-I69)	11	1.6	*
8	Septicemia (A40-A41)	6	0.9	*
8	Legal intervention (Y35,Y89.0)	6	0.9	*
10	Influenza and pneumonia (J10-J18)	5	0.7	*
10	Chronic liver disease and cirrhosis (K70,K73-K74)	5	0.7	*
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	0.7	*
13	Diabetes mellitus (E10-E14)	4	0.6	*
14	Tuberculosis (A16-A19)	2	0.3	*
14	Viral hepatitis (B15-B19)	2	0.3	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.3	*
...	All other causes (Residual)	91	13.2	8.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Percent of		
		Number	total deaths	Rate
Asian or Pacific Islander, male, 35-44 years				
...	All causes	1,064	100.0	108.2
1	Malignant neoplasms (C00-C97)	244	22.9	24.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	188	17.7	19.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	144	13.5	14.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	88	8.3	9.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	75	7.0	7.6
6	Cerebrovascular diseases (I60-I69)	59	5.5	6.0
7	Human immunodeficiency virus (HIV) disease (B20-B24)	27	2.5	2.7
8	Chronic liver disease and cirrhosis (K70,K73-K74)	24	2.3	2.4
9	Diabetes mellitus (E10-E14)	20	1.9	2.0
10	Viral hepatitis (B15-B19)	12	1.1	*
11	Aortic aneurysm and dissection (I71)	9	0.8	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.8	*
13	Chronic lower respiratory diseases (J40-J47)	8	0.8	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	0.8	*
15	Septicemia (A40-A41)	7	0.7	*
...	All other causes (Residual)	142	13.3	14.4
Asian or Pacific Islander, male, 45-54 years				
...	All causes	1,911	100.0	253.6
1	Malignant neoplasms (C00-C97)	632	33.1	83.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	453	23.7	60.1
3	Cerebrovascular diseases (I60-I69)	120	6.3	15.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	119	6.2	15.8
5	Diabetes mellitus (E10-E14)	64	3.3	8.5
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	61	3.2	8.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	52	2.7	6.9
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	49	2.6	6.5
9	Viral hepatitis (B15-B19)	40	2.1	5.3
10	Chronic lower respiratory diseases (J40-J47)	27	1.4	3.6
11	Septicemia (A40-A41)	22	1.2	2.9
12	Influenza and pneumonia (J10-J18)	21	1.1	2.8
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21	1.1	2.8
14	Aortic aneurysm and dissection (I71)	19	1.0	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	14	0.7	*
...	All other causes (Residual)	197	10.3	26.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Percent of		
		Number	total deaths	Rate
Asian or Pacific Islander, male, 55-64 years				
...	All causes	2,696	100.0	625.5
1	Malignant neoplasms (C00-C97)	1,012	37.5	234.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	698	25.9	162.0
3	Cerebrovascular diseases (I60-I69)	199	7.4	46.2
4	Diabetes mellitus (E10-E14)	102	3.8	23.7
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	77	2.9	17.9
6	Chronic lower respiratory diseases (J40-J47)	62	2.3	14.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	49	1.8	11.4
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	48	1.8	11.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	40	1.5	9.3
10	Influenza and pneumonia (J10-J18)	37	1.4	8.6
11	Viral hepatitis (B15-B19)	31	1.1	7.2
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	30	1.1	7.0
13	Aortic aneurysm and dissection (I71)	21	0.8	4.9
14	Septicemia (A40-A41)	18	0.7	*
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	17	0.6	*
...	All other causes (Residual)	255	9.5	59.2
Asian or Pacific Islander, male, 65-74 years				
...	All causes	3,848	100.0	1,556.0
1	Malignant neoplasms (C00-C97)	1,274	33.1	515.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,086	28.2	439.1
3	Cerebrovascular diseases (I60-I69)	333	8.7	134.7
4	Chronic lower respiratory diseases (J40-J47)	162	4.2	65.5
5	Diabetes mellitus (E10-E14)	156	4.1	63.1
6	Influenza and pneumonia (J10-J18)	86	2.2	34.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	82	2.1	33.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	70	1.8	28.3
9	Septicemia (A40-A41)	55	1.4	22.2
10	Aortic aneurysm and dissection (I71)	52	1.4	21.0
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	41	1.1	16.6
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	33	0.9	13.3
12	Chronic liver disease and cirrhosis (K70,K73-K74)	33	0.9	13.3
14	Viral hepatitis (B15-B19)	29	0.8	11.7
15	Parkinson's disease (G20-G21)	18	0.5	*
...	All other causes (Residual)	338	8.8	136.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Percent of total deaths		
		Number	Rate	
Asian or Pacific Islander, male, 75-84 years				
...	All causes	5,086	100.0	4,168.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,554	30.6	1,273.8
2	Malignant neoplasms (C00-C97)	1,337	26.3	1,095.9
3	Cerebrovascular diseases (I60-I69)	500	9.8	409.8
4	Chronic lower respiratory diseases (J40-J47)	292	5.7	239.3
5	Influenza and pneumonia (J10-J18)	204	4.0	167.2
6	Diabetes mellitus (E10-E14)	173	3.4	141.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	106	2.1	86.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	103	2.0	84.4
9	Parkinson's disease (G20-G21)	59	1.2	48.4
10	Septicemia (A40-A41)	57	1.1	46.7
10	Aortic aneurysm and dissection (I71)	57	1.1	46.7
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	56	1.1	45.9
13	Alzheimer's disease (G30)	38	0.7	31.1
14	Pneumonitis due to solids and liquids (J69)	32	0.6	26.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	29	0.6	23.8
...	All other causes (Residual)	489	9.6	400.8
Asian or Pacific Islander, male, 85 years and over				
...	All causes	3,373	100.0	11,308.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,100	32.6	3,688.1
2	Malignant neoplasms (C00-C97)	500	14.8	1,676.4
3	Cerebrovascular diseases (I60-I69)	396	11.7	1,327.7
4	Influenza and pneumonia (J10-J18)	256	7.6	858.3
5	Chronic lower respiratory diseases (J40-J47)	191	5.7	640.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	89	2.6	298.4
7	Diabetes mellitus (E10-E14)	77	2.3	258.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	75	2.2	251.5
9	Alzheimer's disease (G30)	63	1.9	211.2
10	Pneumonitis due to solids and liquids (J69)	53	1.6	177.7
11	Septicemia (A40-A41)	42	1.2	140.8
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	39	1.2	130.8
13	Parkinson's disease (G20-G21)	30	0.9	100.6
14	Aortic aneurysm and dissection (I71)	29	0.9	97.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	0.7	77.1
...	All other causes (Residual)	410	12.2	1,374.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, all ages				
...	All causes	17,219	100.0	274.4
1	Malignant neoplasms (C00-C97)	4,645	27.0	74.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,263	24.8	67.9
3	Cerebrovascular diseases (I60-I69)	1,872	10.9	29.8
4	Diabetes mellitus (E10-E14)	644	3.7	10.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	628	3.6	10.0
6	Influenza and pneumonia (J10-J18)	546	3.2	8.7
7	Chronic lower respiratory diseases (J40-J47)	429	2.5	6.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	305	1.8	4.9
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	234	1.4	3.7
10	Alzheimer's disease (G30)	188	1.1	3.0
11	Septicemia (A40-A41)	179	1.0	2.9
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	176	1.0	2.8
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	168	1.0	2.7
14	Certain conditions originating in the perinatal period (P00-P96)	159	0.9	2.5
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	142	0.8	2.3
...	All other causes (Residual)	2,641	15.3	42.1
Asian or Pacific Islander, female, 1-4 years				
...	All causes	72	100.0	21.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15	20.8	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	11	15.3	*
3	Malignant neoplasms (C00-C97)	8	11.1	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	8.3	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	4.2	*
5	Influenza and pneumonia (J10-J18)	3	4.2	*
7	Septicemia (A40-A41)	2	2.8	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	2.8	*
9	Cerebrovascular diseases (I60-I69)	1	1.4	*
...	All other causes (Residual)	21	29.2	6.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 5-14 years				
...	All causes	90	100.0	10.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	29	32.2	3.5
2	Malignant neoplasms (C00-C97)	17	18.9	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	11	12.2	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	5.6	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	5.6	*
6	Cerebrovascular diseases (I60-I69)	4	4.4	*
7	Septicemia (A40-A41)	1	1.1	*
7	Diabetes mellitus (E10-E14)	1	1.1	*
7	Influenza and pneumonia (J10-J18)	1	1.1	*
7	Hernia (K40-K46)	1	1.1	*
7	Certain conditions originating in the perinatal period (P00-P96)	1	1.1	*
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1	1.1	*
...	All other causes (Residual)	13	14.4	*
Asian or Pacific Islander, female, 15-24 years				
...	All causes	245	100.0	26.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	85	34.7	9.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	33	13.5	3.6
3	Malignant neoplasms (C00-C97)	29	11.8	3.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	19	7.8	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	8	3.3	*
6	Cerebrovascular diseases (I60-I69)	5	2.0	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	2.0	*
8	Septicemia (A40-A41)	4	1.6	*
8	Influenza and pneumonia (J10-J18)	4	1.6	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.8	*
10	Chronic lower respiratory diseases (J40-J47)	2	0.8	*
12	Tuberculosis (A16-A19)	1	0.4	*
12	Meningococcal infection (A39)	1	0.4	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.4	*
12	Anemias (D50-D64)	1	0.4	*
12	Diabetes mellitus (E10-E14)	1	0.4	*
12	Meningitis (G00,G03)	1	0.4	*
12	Peptic ulcer (K25-K28)	1	0.4	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.4	*
12	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.4	*
12	Pregnancy, childbirth and the puerperium (O00-099)	1	0.4	*
...	All other causes (Residual)	39	15.9	4.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 25-34 years				
...	All causes	376	100.0	32.9
1	Malignant neoplasms (C00-C97)	91	24.2	8.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	67	17.8	5.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	54	14.4	4.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	39	10.4	3.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	19	5.1	*
6	Cerebrovascular diseases (I60-I69)	11	2.9	*
7	Pregnancy, childbirth and the puerperium (O00-099)	5	1.3	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	1.3	*
9	Septicemia (A40-A41)	4	1.1	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	1.1	*
9	Influenza and pneumonia (J10-J18)	4	1.1	*
12	Chronic liver disease and cirrhosis (K70,K73-K74)	3	0.8	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.8	*
14	Tuberculosis (A16-A19)	2	0.5	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.5	*
14	Diabetes mellitus (E10-E14)	2	0.5	*
14	Chronic lower respiratory diseases (J40-J47)	2	0.5	*
...	All other causes (Residual)	59	15.7	5.2
Asian or Pacific Islander, female, 35-44 years				
...	All causes	654	100.0	61.9
1	Malignant neoplasms (C00-C97)	271	41.4	25.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	77	11.8	7.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	52	8.0	4.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	35	5.4	3.3
5	Cerebrovascular diseases (I60-I69)	32	4.9	3.0
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	25	3.8	2.4
7	Pregnancy, childbirth and the puerperium (O00-099)	12	1.8	*
8	Diabetes mellitus (E10-E14)	10	1.5	*
8	Chronic liver disease and cirrhosis (K70,K73-K74)	10	1.5	*
10	Septicemia (A40-A41)	7	1.1	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	1.1	*
10	Chronic lower respiratory diseases (J40-J47)	7	1.1	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	0.9	*
14	Viral hepatitis (B15-B19)	5	0.8	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	5	0.8	*
14	Influenza and pneumonia (J10-J18)	5	0.8	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.8	*
...	All other causes (Residual)	83	12.7	7.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Percent of		
		Number	total deaths	Rate
Asian or Pacific Islander, female, 45-54 years				
...	All causes	1,378	100.0	158.5
1	Malignant neoplasms (C00-C97)	717	52.0	82.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	160	11.6	18.4
3	Cerebrovascular diseases (I60-I69)	98	7.1	11.3
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	74	5.4	8.5
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	37	2.7	4.3
6	Diabetes mellitus (E10-E14)	23	1.7	2.6
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	22	1.6	2.5
8	Chronic liver disease and cirrhosis (K70,K73-K74)	20	1.5	2.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20	1.5	2.3
10	Chronic lower respiratory diseases (J40-J47)	18	1.3	*
11	Septicemia (A40-A41)	14	1.0	*
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	12	0.9	*
13	Viral hepatitis (B15-B19)	11	0.8	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.6	*
14	Influenza and pneumonia (J10-J18)	8	0.6	*
...	All other causes (Residual)	136	9.9	15.6
Asian or Pacific Islander, female, 55-64 years				
...	All causes	1,846	100.0	379.9
1	Malignant neoplasms (C00-C97)	815	44.1	167.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	305	16.5	62.8
3	Cerebrovascular diseases (I60-I69)	171	9.3	35.2
4	Diabetes mellitus (E10-E14)	79	4.3	16.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	52	2.8	10.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	38	2.1	7.8
7	Chronic lower respiratory diseases (J40-J47)	34	1.8	7.0
8	Viral hepatitis (B15-B19)	30	1.6	6.2
9	Influenza and pneumonia (J10-J18)	21	1.1	4.3
10	Chronic liver disease and cirrhosis (K70,K73-K74)	20	1.1	4.1
11	Septicemia (A40-A41)	18	1.0	*
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	18	1.0	*
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	15	0.8	*
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	14	0.8	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	0.7	*
...	All other causes (Residual)	204	11.1	42.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Percent of		
		Number	total deaths	Rate
Asian or Pacific Islander, female, 65-74 years				
...	All causes	3,399	100.0	1,059.9
1	Malignant neoplasms (C00-C97)	1,197	35.2	373.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	775	22.8	241.7
3	Cerebrovascular diseases (I60-I69)	363	10.7	113.2
4	Diabetes mellitus (E10-E14)	174	5.1	54.3
5	Chronic lower respiratory diseases (J40-J47)	97	2.9	30.2
6	Influenza and pneumonia (J10-J18)	72	2.1	22.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	63	1.9	19.6
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	62	1.8	19.3
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	52	1.5	16.2
10	Viral hepatitis (B15-B19)	51	1.5	15.9
11	Chronic liver disease and cirrhosis (K70,K73-K74)	39	1.1	12.2
12	Septicemia (A40-A41)	29	0.9	9.0
13	Aortic aneurysm and dissection (I71)	28	0.8	8.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	22	0.6	6.9
15	Parkinson's disease (G20-G21)	17	0.5	*
...	All other causes (Residual)	358	10.5	111.6
Asian or Pacific Islander, female, 75-84 years				
...	All causes	4,712	100.0	2,814.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,421	30.2	848.7
2	Malignant neoplasms (C00-C97)	1,060	22.5	633.1
3	Cerebrovascular diseases (I60-I69)	602	12.8	359.6
4	Diabetes mellitus (E10-E14)	238	5.1	142.2
5	Influenza and pneumonia (J10-J18)	170	3.6	101.5
6	Chronic lower respiratory diseases (J40-J47)	133	2.8	79.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	96	2.0	57.3
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	95	2.0	56.7
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	81	1.7	48.4
10	Alzheimer's disease (G30)	60	1.3	35.8
11	Septicemia (A40-A41)	51	1.1	30.5
12	Parkinson's disease (G20-G21)	42	0.9	25.1
13	Aortic aneurysm and dissection (I71)	31	0.7	18.5
13	Pneumonitis due to solids and liquids (J69)	31	0.7	18.5
15	Chronic liver disease and cirrhosis (K70,K73-K74)	26	0.6	15.5
...	All other causes (Residual)	575	12.2	343.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2001

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Percent of		
		Number	total deaths	Rate
	Asian or Pacific Islander, female, 85 years and over			
...	All causes	4,104	100.0	8,706.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,502	36.6	3,186.3
2	Cerebrovascular diseases (I60-I69)	583	14.2	1,236.8
3	Malignant neoplasms (C00-C97)	438	10.7	929.2
4	Influenza and pneumonia (J10-J18)	252	6.1	534.6
5	Chronic lower respiratory diseases (J40-J47)	136	3.3	288.5
6	Diabetes mellitus (E10-E14)	116	2.8	246.1
7	Alzheimer's disease (G30)	112	2.7	237.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	78	1.9	165.5
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	69	1.7	146.4
10	Pneumonitis due to solids and liquids (J69)	64	1.6	135.8
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	63	1.5	133.6
12	Septicemia (A40-A41)	43	1.0	91.2
13	Atherosclerosis (I70)	32	0.8	67.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	28	0.7	59.4
15	Parkinson's disease (G20-G21)	26	0.6	55.2
...	All other causes (Residual)	562	13.7	1,192.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable