

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
	All races, both sexes, all ages			
...	All causes	2,443,387	100.0	847.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	696,947	28.5	241.7
2	Malignant neoplasms (C00-C97)	557,271	22.8	193.2
3	Cerebrovascular diseases (I60-I69)	162,672	6.7	56.4
4	Chronic lower respiratory diseases (J40-J47)	124,816	5.1	43.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	106,742	4.4	37.0
6	Diabetes mellitus (E10-E14)	73,249	3.0	25.4
7	Influenza and pneumonia (J10-J18)	65,681	2.7	22.8
8	Alzheimer's disease (G30)	58,866	2.4	20.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	40,974	1.7	14.2
10	Septicemia (A40-A41)	33,865	1.4	11.7
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	31,655	1.3	11.0
12	Chronic liver disease and cirrhosis (K70,K73-K74)	27,257	1.1	9.5
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	20,261	0.8	7.0
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	17,638	0.7	6.1
15	Pneumonitis due to solids and liquids (J69)	17,593	0.7	6.1
...	All other causes (Residual)	407,900	16.7	141.5
	All races, both sexes, 1-4 years			
...	All causes	4,858	100.0	31.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,641	33.8	10.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	530	10.9	3.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	423	8.7	2.7
4	Malignant neoplasms (C00-C97)	402	8.3	2.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	165	3.4	1.1
6	Influenza and pneumonia (J10-J18)	110	2.3	0.7
7	Septicemia (A40-A41)	79	1.6	0.5
8	Chronic lower respiratory diseases (J40-J47)	65	1.3	0.4
8	Certain conditions originating in the perinatal period (P00-P96)	65	1.3	0.4
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior	60	1.2	0.4
11	Cerebrovascular diseases (I60-I69)	53	1.1	0.3
12	Acute bronchitis and bronchiolitis (J20-J21)	23	0.5	0.1
13	Anemias (D50-D64)	21	0.4	0.1
14	Meningitis (G00,G03)	19	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	19	0.4	*
...	All other causes (Residual)	1,183	24.4	7.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 5-14 years				
...	All causes	7,150	100.0	17.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,718	38.0	6.6
2	Malignant neoplasms (C00-C97)	1,072	15.0	2.6
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	417	5.8	1.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	356	5.0	0.9
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	264	3.7	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	255	3.6	0.6
7	Chronic lower respiratory diseases (J40-J47)	136	1.9	0.3
8	Septicemia (A40-A41)	95	1.3	0.2
9	Cerebrovascular diseases (I60-I69)	91	1.3	0.2
9	Influenza and pneumonia (J10-J18)	91	1.3	0.2
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	89	1.2	0.2
12	Anemias (D50-D64)	53	0.7	0.1
13	Complications of medical and surgical care (Y40-Y84,Y88)	35	0.5	0.1
14	Diabetes mellitus (E10-E14)	34	0.5	0.1
15	Meningitis (G00,G03)	31	0.4	0.1
15	Certain conditions originating in the perinatal period (P00-P96)	31	0.4	0.1
...	All other causes (Residual)	1,382	19.3	3.4
All races, both sexes, 15-24 years				
...	All causes	33,046	100.0	81.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15,412	46.6	38.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5,219	15.8	12.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,010	12.1	9.9
4	Malignant neoplasms (C00-C97)	1,730	5.2	4.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,022	3.1	2.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	492	1.5	1.2
7	Chronic lower respiratory diseases (J40-J47)	192	0.6	0.5
8	Human immunodeficiency virus (HIV) disease (B20-B24)	178	0.5	0.4
9	Diabetes mellitus (E10-E14)	171	0.5	0.4
9	Cerebrovascular diseases (I60-I69)	171	0.5	0.4
11	Influenza and pneumonia (J10-J18)	167	0.5	0.4
12	Septicemia (A40-A41)	118	0.4	0.3
13	Anemias (D50-D64)	111	0.3	0.3
14	Legal intervention (Y35,Y89.0)	91	0.3	0.2
15	Pregnancy, childbirth and the puerperium (O00-099)	89	0.3	0.2
...	All other causes (Residual)	3,873	11.7	9.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 25-34 years				
...	All causes	41,355	100.0	103.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,569	30.4	31.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,046	12.2	12.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4,489	10.9	11.2
4	Malignant neoplasms (C00-C97)	3,872	9.4	9.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,165	7.7	7.9
6	Human immunodeficiency virus (HIV) disease (B20-B24)	1,839	4.4	4.6
7	Diabetes mellitus (E10-E14)	642	1.6	1.6
8	Cerebrovascular diseases (I60-I69)	567	1.4	1.4
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	475	1.1	1.2
10	Chronic liver disease and cirrhosis (K70,K73-K74)	374	0.9	0.9
11	Influenza and pneumonia (J10-J18)	345	0.8	0.9
12	Septicemia (A40-A41)	314	0.8	0.8
13	Chronic lower respiratory diseases (J40-J47)	301	0.7	0.8
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	269	0.7	0.7
15	Anemias (D50-D64)	180	0.4	0.5
...	All other causes (Residual)	6,908	16.7	17.3
All races, both sexes, 35-44 years				
...	All causes	91,140	100.0	202.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	16,710	18.3	37.2
2	Malignant neoplasms (C00-C97)	16,085	17.6	35.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,688	15.0	30.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,851	7.5	15.3
5	Human immunodeficiency virus (HIV) disease (B20-B24)	5,707	6.3	12.7
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,239	3.6	7.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	3,154	3.5	7.0
8	Cerebrovascular diseases (I60-I69)	2,425	2.7	5.4
9	Diabetes mellitus (E10-E14)	2,164	2.4	4.8
10	Chronic lower respiratory diseases (J40-J47)	1,008	1.1	2.2
11	Influenza and pneumonia (J10-J18)	971	1.1	2.2
12	Septicemia (A40-A41)	856	0.9	1.9
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	749	0.8	1.7
14	Viral hepatitis (B15-B19)	740	0.8	1.6
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	572	0.6	1.3
...	All other causes (Residual)	16,221	17.8	36.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 45-54 years				
...	All causes	172,385	100.0	430.1
1	Malignant neoplasms (C00-C97)	49,637	28.8	123.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	37,570	21.8	93.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14,675	8.5	36.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	7,216	4.2	18.0
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,308	3.7	15.7
6	Cerebrovascular diseases (I60-I69)	6,055	3.5	15.1
7	Diabetes mellitus (E10-E14)	5,496	3.2	13.7
8	Human immunodeficiency virus (HIV) disease (B20-B24)	4,474	2.6	11.2
9	Chronic lower respiratory diseases (J40-J47)	3,475	2.0	8.7
10	Viral hepatitis (B15-B19)	2,331	1.4	5.8
11	Septicemia (A40-A41)	2,074	1.2	5.2
12	Influenza and pneumonia (J10-J18)	1,918	1.1	4.8
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,915	1.1	4.8
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,893	1.1	4.7
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	923	0.5	2.3
...	All other causes (Residual)	26,425	15.3	65.9
All races, both sexes, 55-64 years				
...	All causes	253,342	100.0	952.4
1	Malignant neoplasms (C00-C97)	93,391	36.9	351.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	64,234	25.4	241.5
3	Chronic lower respiratory diseases (J40-J47)	11,280	4.5	42.4
4	Diabetes mellitus (E10-E14)	10,022	4.0	37.7
5	Cerebrovascular diseases (I60-I69)	9,897	3.9	37.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,345	3.3	31.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	6,097	2.4	22.9
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,618	1.4	13.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,455	1.4	13.0
10	Septicemia (A40-A41)	3,360	1.3	12.6
11	Influenza and pneumonia (J10-J18)	2,987	1.2	11.2
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,526	0.6	5.7
13	Aortic aneurysm and dissection (I71)	1,460	0.6	5.5
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1,347	0.5	5.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior	1,193	0.5	4.5
...	All other causes (Residual)	31,130	12.3	117.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 65-74 years				
...	All causes	422,990	100.0	2,314.7
1	Malignant neoplasms (C00-C97)	144,757	34.2	792.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	112,547	26.6	615.9
3	Chronic lower respiratory diseases (J40-J47)	29,788	7.0	163.0
4	Cerebrovascular diseases (I60-I69)	21,992	5.2	120.3
5	Diabetes mellitus (E10-E14)	16,709	4.0	91.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,086	1.9	44.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,164	1.7	39.2
8	Influenza and pneumonia (J10-J18)	6,847	1.6	37.5
9	Septicemia (A40-A41)	6,336	1.5	34.7
10	Chronic liver disease and cirrhosis (K70,K73-K74)	5,381	1.3	29.4
11	Alzheimer's disease (G30)	3,602	0.9	19.7
12	Aortic aneurysm and dissection (I71)	3,482	0.8	19.1
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,922	0.7	16.0
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,463	0.6	13.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha'	2,348	0.6	12.8
...	All other causes (Residual)	48,566	11.5	265.8
All races, both sexes, 75-84 years				
...	All causes	707,654	100.0	5,556.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	213,581	30.2	1,677.2
2	Malignant neoplasms (C00-C97)	167,062	23.6	1,311.9
3	Cerebrovascular diseases (I60-I69)	54,889	7.8	431.0
4	Chronic lower respiratory diseases (J40-J47)	49,241	7.0	386.7
5	Diabetes mellitus (E10-E14)	23,282	3.3	182.8
6	Alzheimer's disease (G30)	20,135	2.8	158.1
7	Influenza and pneumonia (J10-J18)	19,984	2.8	156.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13,896	2.0	109.1
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,904	1.8	101.3
10	Septicemia (A40-A41)	11,010	1.6	86.5
11	Parkinson's disease (G20-G21)	8,139	1.2	63.9
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	6,138	0.9	48.2
13	Pneumonitis due to solids and liquids (J69)	5,900	0.8	46.3
14	Aortic aneurysm and dissection (I71)	5,518	0.8	43.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha'	4,597	0.6	36.1
...	All other causes (Residual)	91,378	12.9	717.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 85 years and over				
...	All causes	681,076	100.0	14,828.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	250,173	36.7	5,446.8
2	Malignant neoplasms (C00-C97)	79,182	11.6	1,723.9
3	Cerebrovascular diseases (I60-I69)	66,412	9.8	1,445.9
4	Alzheimer's disease (G30)	34,552	5.1	752.3
5	Influenza and pneumonia (J10-J18)	31,995	4.7	696.6
6	Chronic lower respiratory diseases (J40-J47)	29,284	4.3	637.6
7	Diabetes mellitus (E10-E14)	14,724	2.2	320.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13,256	1.9	288.6
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,651	1.9	275.4
10	Septicemia (A40-A41)	9,324	1.4	203.0
11	Pneumonitis due to solids and liquids (J69)	8,543	1.3	186.0
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	8,285	1.2	180.4
13	Atherosclerosis (I70)	7,723	1.1	168.1
14	Parkinson's disease (G20-G21)	6,212	0.9	135.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior	3,613	0.5	78.7
...	All other causes (Residual)	105,147	15.4	2,289.3
All races, male, all ages				
...	All causes	1,199,264	100.0	846.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	340,933	28.4	240.7
2	Malignant neoplasms (C00-C97)	288,768	24.1	203.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	69,257	5.8	48.9
4	Cerebrovascular diseases (I60-I69)	62,622	5.2	44.2
5	Chronic lower respiratory diseases (J40-J47)	60,713	5.1	42.9
6	Diabetes mellitus (E10-E14)	34,301	2.9	24.2
7	Influenza and pneumonia (J10-J18)	28,918	2.4	20.4
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	25,409	2.1	17.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19,695	1.6	13.9
10	Chronic liver disease and cirrhosis (K70,K73-K74)	17,401	1.5	12.3
11	Alzheimer's disease (G30)	16,989	1.4	12.0
12	Septicemia (A40-A41)	14,947	1.2	10.6
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	13,640	1.1	9.6
14	Human immunodeficiency virus (HIV) disease (B20-B24)	10,468	0.9	7.4
15	Parkinson's disease (G20-G21)	9,593	0.8	6.8
...	All other causes (Residual)	185,610	15.5	131.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 1-4 years				
...	All causes	2,806	100.0	35.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,024	36.5	12.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	282	10.0	3.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	234	8.3	2.9
4	Malignant neoplasms (C00-C97)	216	7.7	2.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	87	3.1	1.1
6	Influenza and pneumonia (J10-J18)	63	2.2	0.8
7	Chronic lower respiratory diseases (J40-J47)	44	1.6	0.6
8	Septicemia (A40-A41)	41	1.5	0.5
9	Certain conditions originating in the perinatal period (P00-P96)	40	1.4	0.5
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	30	1.1	0.4
10	Cerebrovascular diseases (I60-I69)	30	1.1	0.4
12	Meningitis (G00,G03)	13	0.5	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	11	0.4	*
14	Meningococcal infection (A39)	9	0.3	*
14	Anemias (D50-D64)	9	0.3	*
14	Acute bronchitis and bronchiolitis (J20-J21)	9	0.3	*
...	All other causes (Residual)	664	23.7	8.3
All races, male, 5-14 years				
...	All causes	4,198	100.0	20.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,713	40.8	8.2
2	Malignant neoplasms (C00-C97)	600	14.3	2.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	215	5.1	1.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	199	4.7	0.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	197	4.7	0.9
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	141	3.4	0.7
7	Chronic lower respiratory diseases (J40-J47)	74	1.8	0.4
8	Septicemia (A40-A41)	52	1.2	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	49	1.2	0.2
10	Cerebrovascular diseases (I60-I69)	47	1.1	0.2
11	Influenza and pneumonia (J10-J18)	38	0.9	0.2
12	Anemias (D50-D64)	26	0.6	0.1
12	Complications of medical and surgical care (Y40-Y84,Y88)	26	0.6	0.1
14	Diabetes mellitus (E10-E14)	22	0.5	0.1
15	Certain conditions originating in the perinatal period (P00-P96)	19	0.5	*
...	All other causes (Residual)	780	18.6	3.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 15-24 years				
...	All causes	24,416	100.0	117.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,438	46.8	54.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4,474	18.3	21.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,432	14.1	16.5
4	Malignant neoplasms (C00-C97)	1,016	4.2	4.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	686	2.8	3.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	282	1.2	1.4
7	Chronic lower respiratory diseases (J40-J47)	120	0.5	0.6
8	Cerebrovascular diseases (I60-I69)	103	0.4	0.5
9	Diabetes mellitus (E10-E14)	98	0.4	0.5
10	Influenza and pneumonia (J10-J18)	97	0.4	0.5
11	Human immunodeficiency virus (HIV) disease (B20-B24)	93	0.4	0.4
12	Legal intervention (Y35,Y89.0)	87	0.4	0.4
13	Anemias (D50-D64)	67	0.3	0.3
14	Septicemia (A40-A41)	59	0.2	0.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	50	0.2	0.2
...	All other causes (Residual)	2,314	9.5	11.1
All races, male, 25-34 years				
...	All causes	28,736	100.0	142.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,635	33.5	47.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,135	14.4	20.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,669	12.8	18.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,131	7.4	10.5
5	Malignant neoplasms (C00-C97)	1,868	6.5	9.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	1,187	4.1	5.9
7	Diabetes mellitus (E10-E14)	385	1.3	1.9
8	Cerebrovascular diseases (I60-I69)	284	1.0	1.4
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	284	1.0	1.4
10	Chronic liver disease and cirrhosis (K70,K73-K74)	236	0.8	1.2
11	Influenza and pneumonia (J10-J18)	190	0.7	0.9
12	Septicemia (A40-A41)	166	0.6	0.8
13	Chronic lower respiratory diseases (J40-J47)	165	0.6	0.8
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	129	0.4	0.6
15	Anemias (D50-D64)	101	0.4	0.5
...	All other causes (Residual)	4,171	14.5	20.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 35-44 years				
...	All causes	57,593	100.0	257.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,012	20.9	53.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,638	16.7	43.1
3	Malignant neoplasms (C00-C97)	7,056	12.3	31.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,305	9.2	23.7
5	Human immunodeficiency virus (HIV) disease (B20-B24)	4,198	7.3	18.8
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,386	4.1	10.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,079	3.6	9.3
8	Diabetes mellitus (E10-E14)	1,330	2.3	5.9
9	Cerebrovascular diseases (I60-I69)	1,178	2.0	5.3
10	Influenza and pneumonia (J10-J18)	546	0.9	2.4
11	Viral hepatitis (B15-B19)	519	0.9	2.3
12	Chronic lower respiratory diseases (J40-J47)	497	0.9	2.2
13	Septicemia (A40-A41)	441	0.8	2.0
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	407	0.7	1.8
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	296	0.5	1.3
...	All other causes (Residual)	9,705	16.9	43.4
All races, male, 45-54 years				
...	All causes	107,722	100.0	547.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	27,239	25.3	138.4
2	Malignant neoplasms (C00-C97)	25,192	23.4	128.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,492	9.7	53.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	5,285	4.9	26.9
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,796	4.5	24.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	3,485	3.2	17.7
7	Cerebrovascular diseases (I60-I69)	3,278	3.0	16.7
8	Diabetes mellitus (E10-E14)	3,225	3.0	16.4
9	Chronic lower respiratory diseases (J40-J47)	1,795	1.7	9.1
10	Viral hepatitis (B15-B19)	1,755	1.6	8.9
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,394	1.3	7.1
12	Influenza and pneumonia (J10-J18)	1,138	1.1	5.8
13	Septicemia (A40-A41)	1,124	1.0	5.7
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,086	1.0	5.5
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	560	0.5	2.8
...	All other causes (Residual)	15,878	14.7	80.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 55-64 years				
...	All causes	151,363	100.0	1,184.0
1	Malignant neoplasms (C00-C97)	51,115	33.8	399.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	43,900	29.0	343.4
3	Chronic lower respiratory diseases (J40-J47)	5,783	3.8	45.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,781	3.8	45.2
5	Diabetes mellitus (E10-E14)	5,683	3.8	44.5
6	Cerebrovascular diseases (I60-I69)	5,456	3.6	42.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	4,228	2.8	33.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,837	1.9	22.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,824	1.2	14.3
10	Septicemia (A40-A41)	1,748	1.2	13.7
11	Influenza and pneumonia (J10-J18)	1,738	1.1	13.6
12	Aortic aneurysm and dissection (I71)	1,092	0.7	8.5
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,083	0.7	8.5
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	850	0.6	6.6
15	Viral hepatitis (B15-B19)	702	0.5	5.5
...	All other causes (Residual)	17,543	11.6	137.2
All races, male, 65-74 years				
...	All causes	237,021	100.0	2,855.3
1	Malignant neoplasms (C00-C97)	80,085	33.8	964.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	68,658	29.0	827.1
3	Chronic lower respiratory diseases (J40-J47)	15,339	6.5	184.8
4	Cerebrovascular diseases (I60-I69)	11,207	4.7	135.0
5	Diabetes mellitus (E10-E14)	8,417	3.6	101.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,031	2.1	60.6
7	Influenza and pneumonia (J10-J18)	3,850	1.6	46.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,802	1.6	45.8
9	Chronic liver disease and cirrhosis (K70,K73-K74)	3,197	1.3	38.5
10	Septicemia (A40-A41)	3,154	1.3	38.0
11	Aortic aneurysm and dissection (I71)	2,369	1.0	28.5
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,053	0.9	24.7
13	Alzheimer's disease (G30)	1,547	0.7	18.6
14	Parkinson's disease (G20-G21)	1,487	0.6	17.9
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,419	0.6	17.1
...	All other causes (Residual)	25,406	10.7	306.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 75-84 years				
...	All causes	343,504	100.0	6,760.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	107,217	31.2	2,110.1
2	Malignant neoplasms (C00-C97)	86,952	25.3	1,711.3
3	Chronic lower respiratory diseases (J40-J47)	24,432	7.1	480.8
4	Cerebrovascular diseases (I60-I69)	22,655	6.6	445.9
5	Diabetes mellitus (E10-E14)	10,295	3.0	202.6
6	Influenza and pneumonia (J10-J18)	9,895	2.9	194.7
7	Alzheimer's disease (G30)	7,227	2.1	142.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,920	2.0	136.2
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,687	1.9	131.6
10	Septicemia (A40-A41)	5,028	1.5	99.0
11	Parkinson's disease (G20-G21)	4,811	1.4	94.7
12	Pneumonitis due to solids and liquids (J69)	3,243	0.9	63.8
13	Aortic aneurysm and dissection (I71)	3,185	0.9	62.7
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,405	0.7	47.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior	2,332	0.7	45.9
...	All other causes (Residual)	40,220	11.7	791.6
All races, male, 85 years and over				
...	All causes	225,906	100.0	16,254.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	80,935	35.8	5,823.5
2	Malignant neoplasms (C00-C97)	34,621	15.3	2,491.1
3	Cerebrovascular diseases (I60-I69)	18,316	8.1	1,317.9
4	Chronic lower respiratory diseases (J40-J47)	12,436	5.5	894.8
5	Influenza and pneumonia (J10-J18)	11,199	5.0	805.8
6	Alzheimer's disease (G30)	7,950	3.5	572.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,358	2.4	385.5
8	Diabetes mellitus (E10-E14)	4,844	2.1	348.5
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,817	2.1	346.6
10	Pneumonitis due to solids and liquids (J69)	3,731	1.7	268.5
11	Parkinson's disease (G20-G21)	3,059	1.4	220.1
12	Septicemia (A40-A41)	2,973	1.3	213.9
13	Atherosclerosis (I70)	2,177	1.0	156.6
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,132	0.9	153.4
15	Aortic aneurysm and dissection (I71)	1,481	0.7	106.6
...	All other causes (Residual)	29,877	13.2	2,149.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, all ages				
...	All causes	1,244,123	100.0	848.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	356,014	28.6	242.7
2	Malignant neoplasms (C00-C97)	268,503	21.6	183.0
3	Cerebrovascular diseases (I60-I69)	100,050	8.0	68.2
4	Chronic lower respiratory diseases (J40-J47)	64,103	5.2	43.7
5	Alzheimer's disease (G30)	41,877	3.4	28.5
6	Diabetes mellitus (E10-E14)	38,948	3.1	26.5
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	37,485	3.0	25.6
8	Influenza and pneumonia (J10-J18)	36,763	3.0	25.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21,279	1.7	14.5
10	Septicemia (A40-A41)	18,918	1.5	12.9
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	12,614	1.0	8.6
12	Chronic liver disease and cirrhosis (K70,K73-K74)	9,856	0.8	6.7
13	Pneumonitis due to solids and liquids (J69)	8,717	0.7	5.9
14	Atherosclerosis (I70)	8,650	0.7	5.9
15	Parkinson's disease (G20-G21)	7,366	0.6	5.0
...	All other causes (Residual)	212,980	17.1	145.2
All races, female, 1-4 years				
...	All causes	2,052	100.0	27.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	617	30.1	8.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	248	12.1	3.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	189	9.2	2.5
4	Malignant neoplasms (C00-C97)	186	9.1	2.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	78	3.8	1.0
6	Influenza and pneumonia (J10-J18)	47	2.3	0.6
7	Septicemia (A40-A41)	38	1.9	0.5
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	30	1.5	0.4
9	Certain conditions originating in the perinatal period (P00-P96)	25	1.2	0.3
10	Cerebrovascular diseases (I60-I69)	23	1.1	0.3
11	Chronic lower respiratory diseases (J40-J47)	21	1.0	0.3
12	Acute bronchitis and bronchiolitis (J20-J21)	14	0.7	*
13	Anemias (D50-D64)	12	0.6	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	8	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.3	*
...	All other causes (Residual)	509	24.8	6.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 5-14 years				
...	All causes	2,952	100.0	14.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,005	34.0	5.0
2	Malignant neoplasms (C00-C97)	472	16.0	2.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	202	6.8	1.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	159	5.4	0.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	114	3.9	0.6
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	65	2.2	0.3
7	Chronic lower respiratory diseases (J40-J47)	62	2.1	0.3
8	Influenza and pneumonia (J10-J18)	53	1.8	0.3
9	Cerebrovascular diseases (I60-I69)	44	1.5	0.2
10	Septicemia (A40-A41)	43	1.5	0.2
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha'	40	1.4	0.2
12	Anemias (D50-D64)	27	0.9	0.1
13	Meningitis (G00,G03)	15	0.5	*
14	Diabetes mellitus (E10-E14)	12	0.4	*
14	Certain conditions originating in the perinatal period (P00-P96)	12	0.4	*
...	All other causes (Residual)	627	21.2	3.1
All races, female, 15-24 years				
...	All causes	8,630	100.0	43.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,974	46.0	20.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	745	8.6	3.8
3	Malignant neoplasms (C00-C97)	714	8.3	3.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	578	6.7	2.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	336	3.9	1.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	210	2.4	1.1
7	Pregnancy, childbirth and the puerperium (O00-099)	89	1.0	0.5
8	Human immunodeficiency virus (HIV) disease (B20-B24)	85	1.0	0.4
9	Diabetes mellitus (E10-E14)	73	0.8	0.4
10	Chronic lower respiratory diseases (J40-J47)	72	0.8	0.4
11	Influenza and pneumonia (J10-J18)	70	0.8	0.4
12	Cerebrovascular diseases (I60-I69)	68	0.8	0.3
13	Septicemia (A40-A41)	59	0.7	0.3
14	Anemias (D50-D64)	44	0.5	0.2
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	43	0.5	0.2
...	All other causes (Residual)	1,470	17.0	7.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 25-34 years				
...	All causes	12,619	100.0	64.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,934	23.3	14.9
2	Malignant neoplasms (C00-C97)	2,004	15.9	10.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,034	8.2	5.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	911	7.2	4.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	820	6.5	4.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	652	5.2	3.3
7	Cerebrovascular diseases (I60-I69)	283	2.2	1.4
8	Diabetes mellitus (E10-E14)	257	2.0	1.3
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	191	1.5	1.0
10	Pregnancy, childbirth and the puerperium (O00-099)	175	1.4	0.9
11	Influenza and pneumonia (J10-J18)	155	1.2	0.8
12	Septicemia (A40-A41)	148	1.2	0.8
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	140	1.1	0.7
14	Chronic liver disease and cirrhosis (K70,K73-K74)	138	1.1	0.7
15	Chronic lower respiratory diseases (J40-J47)	136	1.1	0.7
...	All other causes (Residual)	2,641	20.9	13.4
All races, female, 35-44 years				
...	All causes	33,547	100.0	148.8
1	Malignant neoplasms (C00-C97)	9,029	26.9	40.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,698	14.0	20.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,050	12.1	18.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,546	4.6	6.9
5	Human immunodeficiency virus (HIV) disease (B20-B24)	1,509	4.5	6.7
6	Cerebrovascular diseases (I60-I69)	1,247	3.7	5.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,075	3.2	4.8
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	853	2.5	3.8
9	Diabetes mellitus (E10-E14)	834	2.5	3.7
10	Chronic lower respiratory diseases (J40-J47)	511	1.5	2.3
11	Influenza and pneumonia (J10-J18)	425	1.3	1.9
12	Septicemia (A40-A41)	415	1.2	1.8
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	342	1.0	1.5
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	276	0.8	1.2
15	Viral hepatitis (B15-B19)	221	0.7	1.0
...	All other causes (Residual)	6,516	19.4	28.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 45-54 years				
...	All causes	64,663	100.0	316.9
1	Malignant neoplasms (C00-C97)	24,445	37.8	119.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,331	16.0	50.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,183	6.5	20.5
4	Cerebrovascular diseases (I60-I69)	2,777	4.3	13.6
5	Diabetes mellitus (E10-E14)	2,271	3.5	11.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1,931	3.0	9.5
7	Chronic lower respiratory diseases (J40-J47)	1,680	2.6	8.2
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,512	2.3	7.4
9	Human immunodeficiency virus (HIV) disease (B20-B24)	989	1.5	4.8
10	Septicemia (A40-A41)	950	1.5	4.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	807	1.2	4.0
12	Influenza and pneumonia (J10-J18)	780	1.2	3.8
13	Viral hepatitis (B15-B19)	576	0.9	2.8
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	521	0.8	2.6
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	363	0.6	1.8
...	All other causes (Residual)	10,547	16.3	51.7
All races, female, 55-64 years				
...	All causes	101,979	100.0	738.0
1	Malignant neoplasms (C00-C97)	42,276	41.5	306.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,334	19.9	147.2
3	Chronic lower respiratory diseases (J40-J47)	5,497	5.4	39.8
4	Cerebrovascular diseases (I60-I69)	4,441	4.4	32.1
5	Diabetes mellitus (E10-E14)	4,339	4.3	31.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,564	2.5	18.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,869	1.8	13.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,631	1.6	11.8
9	Septicemia (A40-A41)	1,612	1.6	11.7
10	Influenza and pneumonia (J10-J18)	1,249	1.2	9.0
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	781	0.8	5.7
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	676	0.7	4.9
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior	513	0.5	3.7
14	Viral hepatitis (B15-B19)	370	0.4	2.7
15	Aortic aneurysm and dissection (I71)	368	0.4	2.7
...	All other causes (Residual)	13,459	13.2	97.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 65-74 years				
...	All causes	185,969	100.0	1,864.7
1	Malignant neoplasms (C00-C97)	64,672	34.8	648.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	43,889	23.6	440.1
3	Chronic lower respiratory diseases (J40-J47)	14,449	7.8	144.9
4	Cerebrovascular diseases (I60-I69)	10,785	5.8	108.1
5	Diabetes mellitus (E10-E14)	8,292	4.5	83.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,362	1.8	33.7
7	Septicemia (A40-A41)	3,182	1.7	31.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,055	1.6	30.6
9	Influenza and pneumonia (J10-J18)	2,997	1.6	30.1
10	Chronic liver disease and cirrhosis (K70,K73-K74)	2,184	1.2	21.9
11	Alzheimer's disease (G30)	2,055	1.1	20.6
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,503	0.8	15.1
13	Aortic aneurysm and dissection (I71)	1,113	0.6	11.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha'	1,033	0.6	10.4
15	Parkinson's disease (G20-G21)	739	0.4	7.4
...	All other causes (Residual)	22,659	12.2	227.2
All races, female, 75-84 years				
...	All causes	364,150	100.0	4,757.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	106,364	29.2	1,389.7
2	Malignant neoplasms (C00-C97)	80,110	22.0	1,046.7
3	Cerebrovascular diseases (I60-I69)	32,234	8.9	421.2
4	Chronic lower respiratory diseases (J40-J47)	24,809	6.8	324.1
5	Diabetes mellitus (E10-E14)	12,987	3.6	169.7
6	Alzheimer's disease (G30)	12,908	3.5	168.7
7	Influenza and pneumonia (J10-J18)	10,089	2.8	131.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,976	1.9	91.1
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,217	1.7	81.2
10	Septicemia (A40-A41)	5,982	1.6	78.2
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	3,733	1.0	48.8
12	Parkinson's disease (G20-G21)	3,328	0.9	43.5
13	Pneumonitis due to solids and liquids (J69)	2,657	0.7	34.7
14	Aortic aneurysm and dissection (I71)	2,333	0.6	30.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha'	2,265	0.6	29.6
...	All other causes (Residual)	51,158	14.0	668.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 85 years and over				
...	All causes	455,170	100.0	14,209.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	169,238	37.2	5,283.3
2	Cerebrovascular diseases (I60-I69)	48,096	10.6	1,501.5
3	Malignant neoplasms (C00-C97)	44,561	9.8	1,391.1
4	Alzheimer's disease (G30)	26,602	5.8	830.5
5	Influenza and pneumonia (J10-J18)	20,796	4.6	649.2
6	Chronic lower respiratory diseases (J40-J47)	16,848	3.7	526.0
7	Diabetes mellitus (E10-E14)	9,880	2.2	308.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,898	1.7	246.6
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,834	1.7	244.6
10	Septicemia (A40-A41)	6,351	1.4	198.3
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	6,153	1.4	192.1
12	Atherosclerosis (I70)	5,546	1.2	173.1
13	Pneumonitis due to solids and liquids (J69)	4,812	1.1	150.2
14	Parkinson's disease (G20-G21)	3,153	0.7	98.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha'	2,208	0.5	68.9
...	All other causes (Residual)	75,194	16.5	2,347.4
White, both sexes, all ages				
...	All causes	2,102,589	100.0	895.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	606,876	28.9	258.5
2	Malignant neoplasms (C00-C97)	482,481	22.9	205.5
3	Cerebrovascular diseases (I60-I69)	139,719	6.6	59.5
4	Chronic lower respiratory diseases (J40-J47)	115,395	5.5	49.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	90,866	4.3	38.7
6	Diabetes mellitus (E10-E14)	58,459	2.8	24.9
7	Influenza and pneumonia (J10-J18)	58,346	2.8	24.9
8	Alzheimer's disease (G30)	55,058	2.6	23.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	32,615	1.6	13.9
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	28,731	1.4	12.2
11	Septicemia (A40-A41)	27,110	1.3	11.5
12	Chronic liver disease and cirrhosis (K70,K73-K74)	23,809	1.1	10.1
13	Parkinson's disease (G20-G21)	16,135	0.8	6.9
14	Pneumonitis due to solids and liquids (J69)	15,773	0.8	6.7
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	15,587	0.7	6.6
...	All other causes (Residual)	335,629	16.0	143.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 1-4 years				
...	All causes	3,406	100.0	28.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,199	35.2	9.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	395	11.6	3.3
3	Malignant neoplasms (C00-C97)	306	9.0	2.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	239	7.0	2.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	115	3.4	0.9
6	Influenza and pneumonia (J10-J18)	63	1.8	0.5
7	Septicemia (A40-A41)	58	1.7	0.5
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior	43	1.3	0.4
9	Cerebrovascular diseases (I60-I69)	40	1.2	0.3
10	Certain conditions originating in the perinatal period (P00-P96)	36	1.1	0.3
11	Chronic lower respiratory diseases (J40-J47)	30	0.9	0.2
12	Acute bronchitis and bronchiolitis (J20-J21)	20	0.6	0.2
13	Complications of medical and surgical care (Y40-Y84,Y88)	14	0.4	*
14	Meningitis (G00,G03)	13	0.4	*
15	Anemias (D50-D64)	12	0.4	*
...	All other causes (Residual)	823	24.2	6.8
White, both sexes, 5-14 years				
...	All causes	5,138	100.0	16.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,965	38.2	6.2
2	Malignant neoplasms (C00-C97)	846	16.5	2.7
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	321	6.2	1.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	214	4.2	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	195	3.8	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	176	3.4	0.6
7	Septicemia (A40-A41)	69	1.3	0.2
7	Influenza and pneumonia (J10-J18)	69	1.3	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior	67	1.3	0.2
10	Cerebrovascular diseases (I60-I69)	63	1.2	0.2
11	Chronic lower respiratory diseases (J40-J47)	55	1.1	0.2
12	Anemias (D50-D64)	23	0.4	0.1
13	Certain conditions originating in the perinatal period (P00-P96)	21	0.4	0.1
13	Complications of medical and surgical care (Y40-Y84,Y88)	21	0.4	0.1
15	Diabetes mellitus (E10-E14)	20	0.4	0.1
15	Meningitis (G00,G03)	20	0.4	0.1
...	All other causes (Residual)	993	19.3	3.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 15-24 years				
...	All causes	24,641	100.0	77.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,048	53.0	40.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,406	13.8	10.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,137	8.7	6.7
4	Malignant neoplasms (C00-C97)	1,399	5.7	4.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	667	2.7	2.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	383	1.6	1.2
7	Cerebrovascular diseases (I60-I69)	130	0.5	0.4
8	Influenza and pneumonia (J10-J18)	123	0.5	0.4
9	Diabetes mellitus (E10-E14)	109	0.4	0.3
9	Chronic lower respiratory diseases (J40-J47)	109	0.4	0.3
11	Septicemia (A40-A41)	86	0.3	0.3
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha'	68	0.3	0.2
13	Human immunodeficiency virus (HIV) disease (B20-B24)	52	0.2	0.2
14	Legal intervention (Y35,Y89.0)	49	0.2	0.2
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	42	0.2	0.1
...	All other causes (Residual)	2,833	11.5	8.9
White, both sexes, 25-34 years				
...	All causes	29,571	100.0	93.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,378	35.1	32.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,298	14.5	13.6
3	Malignant neoplasms (C00-C97)	2,997	10.1	9.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,040	6.9	6.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,865	6.3	5.9
6	Human immunodeficiency virus (HIV) disease (B20-B24)	688	2.3	2.2
7	Diabetes mellitus (E10-E14)	439	1.5	1.4
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	384	1.3	1.2
9	Cerebrovascular diseases (I60-I69)	376	1.3	1.2
10	Chronic liver disease and cirrhosis (K70,K73-K74)	285	1.0	0.9
11	Influenza and pneumonia (J10-J18)	248	0.8	0.8
12	Septicemia (A40-A41)	202	0.7	0.6
13	Chronic lower respiratory diseases (J40-J47)	189	0.6	0.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	152	0.5	0.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha'	121	0.4	0.4
...	All other causes (Residual)	4,909	16.6	15.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 35-44 years				
...	All causes	68,093	100.0	186.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,901	20.4	38.1
2	Malignant neoplasms (C00-C97)	12,582	18.5	34.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,967	14.6	27.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,219	9.1	17.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,656	3.9	7.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	2,570	3.8	7.0
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,739	2.6	4.8
8	Diabetes mellitus (E10-E14)	1,554	2.3	4.3
9	Cerebrovascular diseases (I60-I69)	1,547	2.3	4.2
10	Influenza and pneumonia (J10-J18)	697	1.0	1.9
11	Chronic lower respiratory diseases (J40-J47)	696	1.0	1.9
12	Viral hepatitis (B15-B19)	600	0.9	1.6
13	Septicemia (A40-A41)	574	0.8	1.6
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	463	0.7	1.3
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	417	0.6	1.1
...	All other causes (Residual)	11,911	17.5	32.6
White, both sexes, 45-54 years				
...	All causes	131,816	100.0	395.3
1	Malignant neoplasms (C00-C97)	39,516	30.0	118.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	28,279	21.5	84.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,069	9.2	36.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	6,020	4.6	18.1
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,885	4.5	17.6
6	Diabetes mellitus (E10-E14)	3,908	3.0	11.7
7	Cerebrovascular diseases (I60-I69)	3,874	2.9	11.6
8	Chronic lower respiratory diseases (J40-J47)	2,812	2.1	8.4
9	Human immunodeficiency virus (HIV) disease (B20-B24)	1,921	1.5	5.8
10	Viral hepatitis (B15-B19)	1,881	1.4	5.6
11	Influenza and pneumonia (J10-J18)	1,456	1.1	4.4
12	Septicemia (A40-A41)	1,410	1.1	4.2
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,117	0.8	3.3
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,082	0.8	3.2
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	605	0.5	1.8
...	All other causes (Residual)	19,981	15.2	59.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 55-64 years				
...	All causes	205,414	100.0	902.5
1	Malignant neoplasms (C00-C97)	78,140	38.0	343.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	51,089	24.9	224.5
3	Chronic lower respiratory diseases (J40-J47)	10,053	4.9	44.2
4	Diabetes mellitus (E10-E14)	7,440	3.6	32.7
5	Cerebrovascular diseases (I60-I69)	7,146	3.5	31.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,985	3.4	30.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	5,259	2.6	23.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,414	1.7	15.0
9	Septicemia (A40-A41)	2,410	1.2	10.6
10	Influenza and pneumonia (J10-J18)	2,357	1.1	10.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,248	1.1	9.9
12	Aortic aneurysm and dissection (I71)	1,231	0.6	5.4
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	1,023	0.5	4.5
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	925	0.5	4.1
15	Viral hepatitis (B15-B19)	804	0.4	3.5
...	All other causes (Residual)	24,890	12.1	109.4
White, both sexes, 65-74 years				
...	All causes	360,047	100.0	2,267.6
1	Malignant neoplasms (C00-C97)	125,458	34.8	790.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	94,492	26.2	595.1
3	Chronic lower respiratory diseases (J40-J47)	27,501	7.6	173.2
4	Cerebrovascular diseases (I60-I69)	17,572	4.9	110.7
5	Diabetes mellitus (E10-E14)	12,942	3.6	81.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,967	1.9	43.9
7	Influenza and pneumonia (J10-J18)	5,830	1.6	36.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,303	1.5	33.4
9	Septicemia (A40-A41)	4,938	1.4	31.1
10	Chronic liver disease and cirrhosis (K70,K73-K74)	4,877	1.4	30.7
11	Alzheimer's disease (G30)	3,282	0.9	20.7
12	Aortic aneurysm and dissection (I71)	3,152	0.9	19.9
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,329	0.6	14.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	2,091	0.6	13.2
15	Parkinson's disease (G20-G21)	2,061	0.6	13.0
...	All other causes (Residual)	41,252	11.5	259.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 75-84 years				
...	All causes	632,353	100.0	5,544.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	190,122	30.1	1,666.9
2	Malignant neoplasms (C00-C97)	149,525	23.6	1,311.0
3	Cerebrovascular diseases (I60-I69)	48,298	7.6	423.5
4	Chronic lower respiratory diseases (J40-J47)	46,258	7.3	405.6
5	Diabetes mellitus (E10-E14)	19,433	3.1	170.4
6	Alzheimer's disease (G30)	18,791	3.0	164.8
7	Influenza and pneumonia (J10-J18)	17,898	2.8	156.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,816	1.9	103.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11,659	1.8	102.2
10	Septicemia (A40-A41)	9,202	1.5	80.7
11	Parkinson's disease (G20-G21)	7,780	1.2	68.2
12	Pneumonitis due to solids and liquids (J69)	5,308	0.8	46.5
13	Aortic aneurysm and dissection (I71)	5,128	0.8	45.0
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	4,862	0.8	42.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior	4,222	0.7	37.0
...	All other causes (Residual)	82,051	13.0	719.4
White, both sexes, 85 years and over				
...	All causes	623,455	100.0	15,015.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	229,565	36.8	5,529.1
2	Malignant neoplasms (C00-C97)	71,652	11.5	1,725.7
3	Cerebrovascular diseases (I60-I69)	60,600	9.7	1,459.5
4	Alzheimer's disease (G30)	32,459	5.2	781.8
5	Influenza and pneumonia (J10-J18)	29,440	4.7	709.1
6	Chronic lower respiratory diseases (J40-J47)	27,662	4.4	666.2
7	Diabetes mellitus (E10-E14)	12,610	2.0	303.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,877	1.9	286.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11,575	1.9	278.8
10	Septicemia (A40-A41)	7,968	1.3	191.9
11	Pneumonitis due to solids and liquids (J69)	7,868	1.3	189.5
12	Atherosclerosis (I70)	7,254	1.2	174.7
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	7,075	1.1	170.4
14	Parkinson's disease (G20-G21)	5,948	1.0	143.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior	3,350	0.5	80.7
...	All other causes (Residual)	96,552	15.5	2,325.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, male, all ages				
...	All causes	1,025,196	100.0	884.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	296,904	29.0	256.0
2	Malignant neoplasms (C00-C97)	249,867	24.4	215.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	58,467	5.7	50.4
4	Chronic lower respiratory diseases (J40-J47)	55,409	5.4	47.8
5	Cerebrovascular diseases (I60-I69)	52,959	5.2	45.7
6	Diabetes mellitus (E10-E14)	28,110	2.7	24.2
7	Influenza and pneumonia (J10-J18)	25,381	2.5	21.9
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	23,049	2.2	19.9
9	Alzheimer's disease (G30)	15,874	1.5	13.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15,850	1.5	13.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	15,278	1.5	13.2
12	Septicemia (A40-A41)	11,919	1.2	10.3
13	Parkinson's disease (G20-G21)	9,130	0.9	7.9
14	Aortic aneurysm and dissection (I71)	8,132	0.8	7.0
15	Pneumonitis due to solids and liquids (J69)	7,954	0.8	6.9
...	All other causes (Residual)	150,913	14.7	130.1
White, male, 1-4 years				
...	All causes	1,958	100.0	31.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	736	37.6	11.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	211	10.8	3.4
3	Malignant neoplasms (C00-C97)	168	8.6	2.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	133	6.8	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	65	3.3	1.0
6	Influenza and pneumonia (J10-J18)	35	1.8	0.6
7	Septicemia (A40-A41)	34	1.7	0.5
8	Cerebrovascular diseases (I60-I69)	22	1.1	0.4
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	21	1.1	0.3
9	Chronic lower respiratory diseases (J40-J47)	21	1.1	0.3
11	Certain conditions originating in the perinatal period (P00-P96)	19	1.0	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	9	0.5	*
13	Meningitis (G00,G03)	8	0.4	*
14	Acute bronchitis and bronchiolitis (J20-J21)	7	0.4	*
15	Meningococcal infection (A39)	6	0.3	*
...	All other causes (Residual)	463	23.6	7.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 5-14 years				
...	All causes	3,005	100.0	18.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,240	41.3	7.6
2	Malignant neoplasms (C00-C97)	459	15.3	2.8
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	159	5.3	1.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	159	5.3	1.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	99	3.3	0.6
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	98	3.3	0.6
7	Septicemia (A40-A41)	42	1.4	0.3
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	37	1.2	0.2
9	Chronic lower respiratory diseases (J40-J47)	31	1.0	0.2
10	Cerebrovascular diseases (I60-I69)	29	1.0	0.2
11	Influenza and pneumonia (J10-J18)	26	0.9	0.2
12	Diabetes mellitus (E10-E14)	17	0.6	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	16	0.5	*
14	Certain conditions originating in the perinatal period (P00-P96)	12	0.4	*
15	Anemias (D50-D64)	11	0.4	*
15	Meningitis (G00,G03)	11	0.4	*
...	All other causes (Residual)	559	18.6	3.4
White, male, 15-24 years				
...	All causes	18,082	100.0	109.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,679	53.5	58.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,919	16.1	17.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,746	9.7	10.6
4	Malignant neoplasms (C00-C97)	830	4.6	5.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	457	2.5	2.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	218	1.2	1.3
7	Cerebrovascular diseases (I60-I69)	79	0.4	0.5
8	Influenza and pneumonia (J10-J18)	74	0.4	0.4
9	Diabetes mellitus (E10-E14)	60	0.3	0.4
9	Chronic lower respiratory diseases (J40-J47)	60	0.3	0.4
11	Legal intervention (Y35,Y89.0)	46	0.3	0.3
12	Septicemia (A40-A41)	45	0.2	0.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	38	0.2	0.2
14	Human immunodeficiency virus (HIV) disease (B20-B24)	31	0.2	0.2
15	Aortic aneurysm and dissection (I71)	24	0.1	0.1
...	All other causes (Residual)	1,776	9.8	10.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 25-34 years				
...	All causes	20,798	100.0	128.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,986	38.4	49.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,495	16.8	21.6
3	Malignant neoplasms (C00-C97)	1,472	7.1	9.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,435	6.9	8.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,420	6.8	8.8
6	Human immunodeficiency virus (HIV) disease (B20-B24)	532	2.6	3.3
7	Diabetes mellitus (E10-E14)	266	1.3	1.6
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	231	1.1	1.4
9	Cerebrovascular diseases (I60-I69)	195	0.9	1.2
10	Chronic liver disease and cirrhosis (K70,K73-K74)	193	0.9	1.2
11	Influenza and pneumonia (J10-J18)	137	0.7	0.8
12	Septicemia (A40-A41)	109	0.5	0.7
13	Chronic lower respiratory diseases (J40-J47)	101	0.5	0.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	76	0.4	0.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha'	70	0.3	0.4
...	All other causes (Residual)	3,080	14.8	19.0
White, male, 35-44 years				
...	All causes	43,955	100.0	239.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,040	22.8	54.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,288	16.6	39.7
3	Malignant neoplasms (C00-C97)	5,604	12.7	30.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,813	10.9	26.2
5	Human immunodeficiency virus (HIV) disease (B20-B24)	2,123	4.8	11.6
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1,783	4.1	9.7
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,240	2.8	6.8
8	Diabetes mellitus (E10-E14)	978	2.2	5.3
9	Cerebrovascular diseases (I60-I69)	773	1.8	4.2
10	Viral hepatitis (B15-B19)	429	1.0	2.3
11	Influenza and pneumonia (J10-J18)	393	0.9	2.1
12	Chronic lower respiratory diseases (J40-J47)	336	0.8	1.8
13	Septicemia (A40-A41)	294	0.7	1.6
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	242	0.6	1.3
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	229	0.5	1.2
...	All other causes (Residual)	7,390	16.8	40.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 45-54 years				
...	All causes	83,651	100.0	505.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	21,283	25.4	128.6
2	Malignant neoplasms (C00-C97)	20,157	24.1	121.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,664	10.4	52.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	4,489	5.4	27.1
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,477	5.4	27.0
6	Diabetes mellitus (E10-E14)	2,368	2.8	14.3
7	Cerebrovascular diseases (I60-I69)	2,132	2.5	12.9
8	Human immunodeficiency virus (HIV) disease (B20-B24)	1,619	1.9	9.8
9	Chronic lower respiratory diseases (J40-J47)	1,453	1.7	8.8
10	Viral hepatitis (B15-B19)	1,432	1.7	8.7
11	Influenza and pneumonia (J10-J18)	857	1.0	5.2
12	Septicemia (A40-A41)	776	0.9	4.7
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	757	0.9	4.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	622	0.7	3.8
15	Aortic aneurysm and dissection (I71)	382	0.5	2.3
...	All other causes (Residual)	12,183	14.6	73.6
White, male, 55-64 years				
...	All causes	123,551	100.0	1,118.6
1	Malignant neoplasms (C00-C97)	42,632	34.5	386.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	35,789	29.0	324.0
3	Chronic lower respiratory diseases (J40-J47)	5,080	4.1	46.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,807	3.9	43.5
5	Diabetes mellitus (E10-E14)	4,371	3.5	39.6
6	Cerebrovascular diseases (I60-I69)	3,937	3.2	35.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	3,672	3.0	33.2
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,684	2.2	24.3
9	Influenza and pneumonia (J10-J18)	1,358	1.1	12.3
10	Septicemia (A40-A41)	1,290	1.0	11.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,186	1.0	10.7
12	Aortic aneurysm and dissection (I71)	926	0.7	8.4
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	582	0.5	5.3
14	Human immunodeficiency virus (HIV) disease (B20-B24)	546	0.4	4.9
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	542	0.4	4.9
...	All other causes (Residual)	14,149	11.5	128.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 65-74 years				
...	All causes	203,735	100.0	2,795.4
1	Malignant neoplasms (C00-C97)	69,588	34.2	954.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	58,873	28.9	807.8
3	Chronic lower respiratory diseases (J40-J47)	14,015	6.9	192.3
4	Cerebrovascular diseases (I60-I69)	9,021	4.4	123.8
5	Diabetes mellitus (E10-E14)	6,800	3.3	93.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,316	2.1	59.2
7	Influenza and pneumonia (J10-J18)	3,291	1.6	45.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,914	1.4	40.0
9	Chronic liver disease and cirrhosis (K70,K73-K74)	2,901	1.4	39.8
10	Septicemia (A40-A41)	2,482	1.2	34.1
11	Aortic aneurysm and dissection (I71)	2,174	1.1	29.8
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,953	1.0	26.8
13	Alzheimer's disease (G30)	1,409	0.7	19.3
14	Parkinson's disease (G20-G21)	1,384	0.7	19.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha'	1,189	0.6	16.3
...	All other causes (Residual)	21,425	10.5	294.0
White, male, 75-84 years				
...	All causes	308,654	100.0	6,738.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	96,734	31.3	2,112.0
2	Malignant neoplasms (C00-C97)	77,649	25.2	1,695.3
3	Chronic lower respiratory diseases (J40-J47)	22,681	7.3	495.2
4	Cerebrovascular diseases (I60-I69)	20,037	6.5	437.5
5	Diabetes mellitus (E10-E14)	8,920	2.9	194.7
6	Influenza and pneumonia (J10-J18)	8,843	2.9	193.1
7	Alzheimer's disease (G30)	6,746	2.2	147.3
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,090	2.0	133.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,972	1.9	130.4
10	Parkinson's disease (G20-G21)	4,604	1.5	100.5
11	Septicemia (A40-A41)	4,193	1.4	91.5
12	Aortic aneurysm and dissection (I71)	3,000	1.0	65.5
13	Pneumonitis due to solids and liquids (J69)	2,931	0.9	64.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha'	2,155	0.7	47.0
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,901	0.6	41.5
...	All other causes (Residual)	36,198	11.7	790.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 85 years and over				
...	All causes	207,144	100.0	16,473.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	74,691	36.1	5,939.8
2	Malignant neoplasms (C00-C97)	31,270	15.1	2,486.8
3	Cerebrovascular diseases (I60-I69)	16,692	8.1	1,327.4
4	Chronic lower respiratory diseases (J40-J47)	11,611	5.6	923.4
5	Influenza and pneumonia (J10-J18)	10,260	5.0	815.9
6	Alzheimer's disease (G30)	7,479	3.6	594.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,746	2.3	377.4
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,484	2.2	356.6
9	Diabetes mellitus (E10-E14)	4,329	2.1	344.3
10	Pneumonitis due to solids and liquids (J69)	3,453	1.7	274.6
11	Parkinson's disease (G20-G21)	2,926	1.4	232.7
12	Septicemia (A40-A41)	2,552	1.2	202.9
13	Atherosclerosis (I70)	2,052	1.0	163.2
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,826	0.9	145.2
15	Aortic aneurysm and dissection (I71)	1,406	0.7	111.8
...	All other causes (Residual)	27,367	13.2	2,176.4
White, female, all ages				
...	All causes	1,077,393	100.0	907.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	309,972	28.8	261.0
2	Malignant neoplasms (C00-C97)	232,614	21.6	195.8
3	Cerebrovascular diseases (I60-I69)	86,760	8.1	73.0
4	Chronic lower respiratory diseases (J40-J47)	59,986	5.6	50.5
5	Alzheimer's disease (G30)	39,184	3.6	33.0
6	Influenza and pneumonia (J10-J18)	32,965	3.1	27.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	32,399	3.0	27.3
8	Diabetes mellitus (E10-E14)	30,349	2.8	25.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16,765	1.6	14.1
10	Septicemia (A40-A41)	15,191	1.4	12.8
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	9,868	0.9	8.3
12	Chronic liver disease and cirrhosis (K70,K73-K74)	8,531	0.8	7.2
13	Atherosclerosis (I70)	7,943	0.7	6.7
14	Pneumonitis due to solids and liquids (J69)	7,819	0.7	6.6
15	Parkinson's disease (G20-G21)	7,005	0.7	5.9
...	All other causes (Residual)	180,042	16.7	151.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 1-4 years				
...	All causes	1,448	100.0	24.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	463	32.0	7.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	184	12.7	3.1
3	Malignant neoplasms (C00-C97)	138	9.5	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	106	7.3	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	50	3.5	0.8
6	Influenza and pneumonia (J10-J18)	28	1.9	0.5
7	Septicemia (A40-A41)	24	1.7	0.4
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	22	1.5	0.4
9	Cerebrovascular diseases (I60-I69)	18	1.2	*
10	Certain conditions originating in the perinatal period (P00-P96)	17	1.2	*
11	Acute bronchitis and bronchiolitis (J20-J21)	13	0.9	*
12	Chronic lower respiratory diseases (J40-J47)	9	0.6	*
13	Anemias (D50-D64)	7	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.4	*
15	Meningitis (G00,G03)	5	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.3	*
...	All other causes (Residual)	353	24.4	6.0
White, female, 5-14 years				
...	All causes	2,133	100.0	13.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	725	34.0	4.7
2	Malignant neoplasms (C00-C97)	387	18.1	2.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	162	7.6	1.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	97	4.5	0.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	77	3.6	0.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	55	2.6	0.4
7	Influenza and pneumonia (J10-J18)	43	2.0	0.3
8	Cerebrovascular diseases (I60-I69)	34	1.6	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	30	1.4	0.2
10	Septicemia (A40-A41)	27	1.3	0.2
11	Chronic lower respiratory diseases (J40-J47)	24	1.1	0.2
12	Anemias (D50-D64)	12	0.6	*
13	Meningitis (G00,G03)	9	0.4	*
13	Certain conditions originating in the perinatal period (P00-P96)	9	0.4	*
15	Pneumonitis due to solids and liquids (J69)	6	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.3	*
...	All other causes (Residual)	430	20.2	2.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 15-24 years				
...	All causes	6,559	100.0	42.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,369	51.4	21.8
2	Malignant neoplasms (C00-C97)	569	8.7	3.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	487	7.4	3.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	391	6.0	2.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	210	3.2	1.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	165	2.5	1.1
7	Cerebrovascular diseases (I60-I69)	51	0.8	0.3
8	Diabetes mellitus (E10-E14)	49	0.7	0.3
8	Influenza and pneumonia (J10-J18)	49	0.7	0.3
8	Chronic lower respiratory diseases (J40-J47)	49	0.7	0.3
11	Septicemia (A40-A41)	41	0.6	0.3
12	Pregnancy, childbirth and the puerperium (O00-099)	40	0.6	0.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	30	0.5	0.2
14	Human immunodeficiency virus (HIV) disease (B20-B24)	21	0.3	0.1
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20	0.3	0.1
...	All other causes (Residual)	1,018	15.5	6.6
White, female, 25-34 years				
...	All causes	8,773	100.0	56.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,392	27.3	15.5
2	Malignant neoplasms (C00-C97)	1,525	17.4	9.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	803	9.2	5.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	620	7.1	4.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	430	4.9	2.8
6	Cerebrovascular diseases (I60-I69)	181	2.1	1.2
7	Diabetes mellitus (E10-E14)	173	2.0	1.1
8	Human immunodeficiency virus (HIV) disease (B20-B24)	156	1.8	1.0
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	153	1.7	1.0
10	Influenza and pneumonia (J10-J18)	111	1.3	0.7
11	Pregnancy, childbirth and the puerperium (O00-099)	97	1.1	0.6
12	Septicemia (A40-A41)	93	1.1	0.6
13	Chronic liver disease and cirrhosis (K70,K73-K74)	92	1.0	0.6
14	Chronic lower respiratory diseases (J40-J47)	88	1.0	0.6
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	76	0.9	0.5
...	All other causes (Residual)	1,783	20.3	11.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 35-44 years				
...	All causes	24,138	100.0	133.2
1	Malignant neoplasms (C00-C97)	6,978	28.9	38.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,861	16.0	21.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,679	11.1	14.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,406	5.8	7.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	873	3.6	4.8
6	Cerebrovascular diseases (I60-I69)	774	3.2	4.3
7	Diabetes mellitus (E10-E14)	576	2.4	3.2
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	499	2.1	2.8
9	Human immunodeficiency virus (HIV) disease (B20-B24)	447	1.9	2.5
10	Chronic lower respiratory diseases (J40-J47)	360	1.5	2.0
11	Influenza and pneumonia (J10-J18)	304	1.3	1.7
12	Septicemia (A40-A41)	280	1.2	1.5
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	221	0.9	1.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	188	0.8	1.0
15	Viral hepatitis (B15-B19)	171	0.7	0.9
...	All other causes (Residual)	4,521	18.7	25.0
White, female, 45-54 years				
...	All causes	48,165	100.0	286.8
1	Malignant neoplasms (C00-C97)	19,359	40.2	115.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,996	14.5	41.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,405	7.1	20.3
4	Cerebrovascular diseases (I60-I69)	1,742	3.6	10.4
5	Diabetes mellitus (E10-E14)	1,540	3.2	9.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1,531	3.2	9.1
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,408	2.9	8.4
8	Chronic lower respiratory diseases (J40-J47)	1,359	2.8	8.1
9	Septicemia (A40-A41)	634	1.3	3.8
10	Influenza and pneumonia (J10-J18)	599	1.2	3.6
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	460	1.0	2.7
12	Viral hepatitis (B15-B19)	449	0.9	2.7
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	360	0.7	2.1
14	Human immunodeficiency virus (HIV) disease (B20-B24)	302	0.6	1.8
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	286	0.6	1.7
...	All other causes (Residual)	7,735	16.1	46.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 55-64 years				
...	All causes	81,863	100.0	698.7
1	Malignant neoplasms (C00-C97)	35,508	43.4	303.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,300	18.7	130.6
3	Chronic lower respiratory diseases (J40-J47)	4,973	6.1	42.4
4	Cerebrovascular diseases (I60-I69)	3,209	3.9	27.4
5	Diabetes mellitus (E10-E14)	3,069	3.7	26.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,178	2.7	18.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,587	1.9	13.5
8	Septicemia (A40-A41)	1,120	1.4	9.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,062	1.3	9.1
10	Influenza and pneumonia (J10-J18)	999	1.2	8.5
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	730	0.9	6.2
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	441	0.5	3.8
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	383	0.5	3.3
14	Aortic aneurysm and dissection (I71)	305	0.4	2.6
15	Viral hepatitis (B15-B19)	277	0.3	2.4
...	All other causes (Residual)	10,722	13.1	91.5
White, female, 65-74 years				
...	All causes	156,312	100.0	1,819.7
1	Malignant neoplasms (C00-C97)	55,870	35.7	650.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	35,619	22.8	414.7
3	Chronic lower respiratory diseases (J40-J47)	13,486	8.6	157.0
4	Cerebrovascular diseases (I60-I69)	8,551	5.5	99.5
5	Diabetes mellitus (E10-E14)	6,142	3.9	71.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,651	1.7	30.9
7	Influenza and pneumonia (J10-J18)	2,539	1.6	29.6
8	Septicemia (A40-A41)	2,456	1.6	28.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,389	1.5	27.8
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,976	1.3	23.0
11	Alzheimer's disease (G30)	1,873	1.2	21.8
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,021	0.7	11.9
13	Aortic aneurysm and dissection (I71)	978	0.6	11.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	902	0.6	10.5
15	Parkinson's disease (G20-G21)	677	0.4	7.9
...	All other causes (Residual)	19,182	12.3	223.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 75-84 years				
...	All causes	323,699	100.0	4,742.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	93,388	28.9	1,368.2
2	Malignant neoplasms (C00-C97)	71,876	22.2	1,053.1
3	Cerebrovascular diseases (I60-I69)	28,261	8.7	414.1
4	Chronic lower respiratory diseases (J40-J47)	23,577	7.3	345.4
5	Alzheimer's disease (G30)	12,045	3.7	176.5
6	Diabetes mellitus (E10-E14)	10,513	3.2	154.0
7	Influenza and pneumonia (J10-J18)	9,055	2.8	132.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,726	1.8	83.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,687	1.8	83.3
10	Septicemia (A40-A41)	5,009	1.5	73.4
11	Parkinson's disease (G20-G21)	3,176	1.0	46.5
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,961	0.9	43.4
13	Pneumonitis due to solids and liquids (J69)	2,377	0.7	34.8
14	Aortic aneurysm and dissection (I71)	2,128	0.7	31.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior	2,067	0.6	30.3
...	All other causes (Residual)	45,853	14.2	671.8
White, female, 85 years and over				
...	All causes	416,311	100.0	14,382.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	154,874	37.2	5,350.6
2	Cerebrovascular diseases (I60-I69)	43,908	10.5	1,516.9
3	Malignant neoplasms (C00-C97)	40,382	9.7	1,395.1
4	Alzheimer's disease (G30)	24,980	6.0	863.0
5	Influenza and pneumonia (J10-J18)	19,180	4.6	662.6
6	Chronic lower respiratory diseases (J40-J47)	16,051	3.9	554.5
7	Diabetes mellitus (E10-E14)	8,281	2.0	286.1
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,393	1.8	255.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,829	1.6	235.9
10	Septicemia (A40-A41)	5,416	1.3	187.1
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	5,249	1.3	181.3
12	Atherosclerosis (I70)	5,202	1.2	179.7
13	Pneumonitis due to solids and liquids (J69)	4,415	1.1	152.5
14	Parkinson's disease (G20-G21)	3,022	0.7	104.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior	2,040	0.5	70.5
...	All other causes (Residual)	69,089	16.6	2,386.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, all ages				
...	All causes	290,051	100.0	768.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	77,621	26.8	205.6
2	Malignant neoplasms (C00-C97)	62,617	21.6	165.9
3	Cerebrovascular diseases (I60-I69)	18,856	6.5	50.0
4	Diabetes mellitus (E10-E14)	12,687	4.4	33.6
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,513	4.3	33.1
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8,287	2.9	22.0
7	Human immunodeficiency virus (HIV) disease (B20-B24)	7,835	2.7	20.8
8	Chronic lower respiratory diseases (J40-J47)	7,831	2.7	20.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,488	2.6	19.8
10	Septicemia (A40-A41)	6,137	2.1	16.3
11	Influenza and pneumonia (J10-J18)	5,871	2.0	15.6
12	Certain conditions originating in the perinatal period (P00-P96)	4,931	1.7	13.1
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	4,196	1.4	11.1
14	Alzheimer's disease (G30)	3,341	1.2	8.9
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,588	0.9	6.9
...	All other causes (Residual)	47,252	16.3	125.2
Black, both sexes, 1-4 years				
...	All causes	1,196	100.0	47.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	359	30.0	14.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	161	13.5	6.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	114	9.5	4.5
4	Malignant neoplasms (C00-C97)	74	6.2	2.9
5	Influenza and pneumonia (J10-J18)	41	3.4	1.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	40	3.3	1.6
7	Chronic lower respiratory diseases (J40-J47)	30	2.5	1.2
8	Certain conditions originating in the perinatal period (P00-P96)	26	2.2	1.0
9	Septicemia (A40-A41)	17	1.4	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	14	1.2	*
11	Cerebrovascular diseases (I60-I69)	11	0.9	*
12	Anemias (D50-D64)	9	0.8	*
13	Meningitis (G00,G03)	6	0.5	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	5	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.4	*
...	All other causes (Residual)	284	23.7	11.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 5-14 years				
...	All causes	1,666	100.0	24.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	621	37.3	9.1
2	Malignant neoplasms (C00-C97)	172	10.3	2.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	135	8.1	2.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	83	5.0	1.2
5	Chronic lower respiratory diseases (J40-J47)	76	4.6	1.1
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	65	3.9	1.0
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	35	2.1	0.5
8	Anemias (D50-D64)	26	1.6	0.4
9	Cerebrovascular diseases (I60-I69)	24	1.4	0.4
10	Septicemia (A40-A41)	20	1.2	0.3
11	Human immunodeficiency virus (HIV) disease (B20-B24)	19	1.1	*
11	Influenza and pneumonia (J10-J18)	19	1.1	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	18	1.1	*
14	Diabetes mellitus (E10-E14)	14	0.8	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	14	0.8	*
...	All other causes (Residual)	325	19.5	4.8
Black, both sexes, 15-24 years				
...	All causes	7,045	100.0	113.7
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,900	41.2	46.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,726	24.5	27.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	403	5.7	6.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	306	4.3	4.9
5	Malignant neoplasms (C00-C97)	250	3.5	4.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	124	1.8	2.0
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	100	1.4	1.6
8	Anemias (D50-D64)	79	1.1	1.3
8	Chronic lower respiratory diseases (J40-J47)	79	1.1	1.3
10	Diabetes mellitus (E10-E14)	59	0.8	1.0
11	Pregnancy, childbirth and the puerperium (O00-099)	46	0.7	0.7
12	Cerebrovascular diseases (I60-I69)	38	0.5	0.6
12	Influenza and pneumonia (J10-J18)	38	0.5	0.6
14	Legal intervention (Y35,Y89.0)	37	0.5	0.6
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	36	0.5	0.6
...	All other causes (Residual)	824	11.7	13.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 25-34 years				
...	All causes	10,171	100.0	186.8
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,469	24.3	45.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,711	16.8	31.4
3	Human immunodeficiency virus (HIV) disease (B20-B24)	1,123	11.0	20.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,004	9.9	18.4
5	Malignant neoplasms (C00-C97)	669	6.6	12.3
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	504	5.0	9.3
7	Diabetes mellitus (E10-E14)	195	1.9	3.6
8	Cerebrovascular diseases (I60-I69)	168	1.7	3.1
9	Anemias (D50-D64)	130	1.3	2.4
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	109	1.1	2.0
11	Chronic lower respiratory diseases (J40-J47)	104	1.0	1.9
12	Septicemia (A40-A41)	103	1.0	1.9
13	Influenza and pneumonia (J10-J18)	83	0.8	1.5
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	75	0.7	1.4
15	Pregnancy, childbirth and the puerperium (O00-099)	66	0.6	1.2
...	All other causes (Residual)	1,658	16.3	30.5
Black, both sexes, 35-44 years				
...	All causes	20,235	100.0	348.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,356	16.6	57.8
2	Human immunodeficiency virus (HIV) disease (B20-B24)	3,063	15.1	52.8
3	Malignant neoplasms (C00-C97)	2,903	14.3	50.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,268	11.2	39.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,377	6.8	23.7
6	Cerebrovascular diseases (I60-I69)	744	3.7	12.8
7	Diabetes mellitus (E10-E14)	560	2.8	9.6
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	442	2.2	7.6
9	Chronic liver disease and cirrhosis (K70,K73-K74)	318	1.6	5.5
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	309	1.5	5.3
11	Chronic lower respiratory diseases (J40-J47)	298	1.5	5.1
12	Septicemia (A40-A41)	258	1.3	4.4
13	Influenza and pneumonia (J10-J18)	252	1.2	4.3
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	191	0.9	3.3
15	Anemias (D50-D64)	145	0.7	2.5
...	All other causes (Residual)	3,751	18.5	64.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 45-54 years				
...	All causes	35,623	100.0	765.8
1	Malignant neoplasms (C00-C97)	8,450	23.7	181.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,413	23.6	180.9
3	Human immunodeficiency virus (HIV) disease (B20-B24)	2,509	7.0	53.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,179	6.1	46.8
5	Cerebrovascular diseases (I60-I69)	1,893	5.3	40.7
6	Diabetes mellitus (E10-E14)	1,399	3.9	30.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	968	2.7	20.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	752	2.1	16.2
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	722	2.0	15.5
10	Septicemia (A40-A41)	600	1.7	12.9
11	Chronic lower respiratory diseases (J40-J47)	599	1.7	12.9
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	417	1.2	9.0
13	Influenza and pneumonia (J10-J18)	403	1.1	8.7
14	Viral hepatitis (B15-B19)	374	1.0	8.0
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	273	0.8	5.9
...	All other causes (Residual)	5,672	15.9	121.9
Black, both sexes, 55-64 years				
...	All causes	41,336	100.0	1,565.2
1	Malignant neoplasms (C00-C97)	12,825	31.0	485.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,696	28.3	442.9
3	Cerebrovascular diseases (I60-I69)	2,324	5.6	88.0
4	Diabetes mellitus (E10-E14)	2,224	5.4	84.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,093	2.6	41.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,082	2.6	41.0
7	Chronic lower respiratory diseases (J40-J47)	1,076	2.6	40.7
8	Septicemia (A40-A41)	876	2.1	33.2
9	Human immunodeficiency virus (HIV) disease (B20-B24)	688	1.7	26.1
10	Chronic liver disease and cirrhosis (K70,K73-K74)	660	1.6	25.0
11	Influenza and pneumonia (J10-J18)	560	1.4	21.2
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	554	1.3	21.0
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	218	0.5	8.3
14	Aortic aneurysm and dissection (I71)	197	0.5	7.5
15	Viral hepatitis (B15-B19)	187	0.5	7.1
...	All other causes (Residual)	5,076	12.3	192.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 65-74 years				
...	All causes	53,314	100.0	3,159.3
1	Malignant neoplasms (C00-C97)	16,245	30.5	962.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,601	29.3	924.5
3	Cerebrovascular diseases (I60-I69)	3,630	6.8	215.1
4	Diabetes mellitus (E10-E14)	3,175	6.0	188.1
5	Chronic lower respiratory diseases (J40-J47)	1,916	3.6	113.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,690	3.2	100.1
7	Septicemia (A40-A41)	1,260	2.4	74.7
8	Influenza and pneumonia (J10-J18)	848	1.6	50.3
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	832	1.6	49.3
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	803	1.5	47.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	365	0.7	21.6
12	Alzheimer's disease (G30)	283	0.5	16.8
13	Pneumonitis due to solids and liquids (J69)	250	0.5	14.8
14	Aortic aneurysm and dissection (I71)	246	0.5	14.6
15	Human immunodeficiency virus (HIV) disease (B20-B24)	243	0.5	14.4
...	All other causes (Residual)	5,927	11.1	351.2
Black, both sexes, 75-84 years				
...	All causes	62,862	100.0	6,518.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	19,757	31.4	2,048.8
2	Malignant neoplasms (C00-C97)	14,664	23.3	1,520.7
3	Cerebrovascular diseases (I60-I69)	5,326	8.5	552.3
4	Diabetes mellitus (E10-E14)	3,249	5.2	336.9
5	Chronic lower respiratory diseases (J40-J47)	2,398	3.8	248.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,959	3.1	203.2
7	Septicemia (A40-A41)	1,651	2.6	171.2
8	Influenza and pneumonia (J10-J18)	1,603	2.6	166.2
9	Alzheimer's disease (G30)	1,187	1.9	123.1
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,138	1.8	118.0
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	806	1.3	83.6
12	Pneumonitis due to solids and liquids (J69)	503	0.8	52.2
13	Atherosclerosis (I70)	352	0.6	36.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior	302	0.5	31.3
15	Aortic aneurysm and dissection (I71)	288	0.5	29.9
...	All other causes (Residual)	7,679	12.2	796.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 85 years and over				
...	All causes	48,022	100.0	14,260.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	17,225	35.9	5,115.2
2	Malignant neoplasms (C00-C97)	6,350	13.2	1,885.7
3	Cerebrovascular diseases (I60-I69)	4,660	9.7	1,383.9
4	Influenza and pneumonia (J10-J18)	1,937	4.0	575.2
5	Alzheimer's disease (G30)	1,831	3.8	543.7
6	Diabetes mellitus (E10-E14)	1,811	3.8	537.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,487	3.1	441.6
8	Chronic lower respiratory diseases (J40-J47)	1,240	2.6	368.2
9	Septicemia (A40-A41)	1,236	2.6	367.0
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,040	2.2	308.8
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	584	1.2	173.4
12	Pneumonitis due to solids and liquids (J69)	556	1.2	165.1
13	Atherosclerosis (I70)	394	0.8	117.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior	214	0.4	63.6
15	Nutritional deficiencies (E40-E64)	204	0.4	60.6
...	All other causes (Residual)	7,253	15.1	2,153.9
Black, male, all ages				
...	All causes	146,835	100.0	816.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	37,094	25.3	206.3
2	Malignant neoplasms (C00-C97)	32,627	22.2	181.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,612	5.9	47.9
4	Cerebrovascular diseases (I60-I69)	7,828	5.3	43.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6,896	4.7	38.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	5,301	3.6	29.5
7	Diabetes mellitus (E10-E14)	5,207	3.5	29.0
8	Chronic lower respiratory diseases (J40-J47)	4,341	3.0	24.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,427	2.3	19.1
10	Influenza and pneumonia (J10-J18)	2,768	1.9	15.4
11	Certain conditions originating in the perinatal period (P00-P96)	2,725	1.9	15.2
12	Septicemia (A40-A41)	2,703	1.8	15.0
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,717	1.2	9.6
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,633	1.1	9.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,622	1.1	9.0
...	All other causes (Residual)	22,334	15.2	124.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 1-4 years				
...	All causes	702	100.0	54.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	239	34.0	18.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	92	13.1	7.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	59	8.4	4.6
4	Malignant neoplasms (C00-C97)	34	4.8	2.6
5	Influenza and pneumonia (J10-J18)	22	3.1	1.7
6	Chronic lower respiratory diseases (J40-J47)	20	2.8	1.6
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	18	2.6	*
7	Certain conditions originating in the perinatal period (P00-P96)	18	2.6	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	8	1.1	*
10	Cerebrovascular diseases (I60-I69)	7	1.0	*
11	Septicemia (A40-A41)	5	0.7	*
11	Meningitis (G00,G03)	5	0.7	*
13	Anemias (D50-D64)	4	0.6	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.4	*
15	Meningococcal infection (A39)	2	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.3	*
...	All other causes (Residual)	164	23.4	12.7
Black, male, 5-14 years				
...	All causes	998	100.0	28.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	400	40.1	11.6
2	Malignant neoplasms (C00-C97)	106	10.6	3.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	86	8.6	2.5
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	48	4.8	1.4
5	Chronic lower respiratory diseases (J40-J47)	41	4.1	1.2
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	33	3.3	1.0
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	29	2.9	0.8
8	Cerebrovascular diseases (I60-I69)	16	1.6	*
9	Anemias (D50-D64)	14	1.4	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	10	1.0	*
10	Influenza and pneumonia (J10-J18)	10	1.0	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	10	1.0	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	9	0.9	*
14	Septicemia (A40-A41)	7	0.7	*
15	Certain conditions originating in the perinatal period (P00-P96)	6	0.6	*
...	All other causes (Residual)	173	17.3	5.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 15-24 years				
...	All causes	5,364	100.0	172.6
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,582	48.1	83.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,298	24.2	41.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	351	6.5	11.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	192	3.6	6.2
5	Malignant neoplasms (C00-C97)	140	2.6	4.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	60	1.1	1.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	60	1.1	1.9
8	Chronic lower respiratory diseases (J40-J47)	59	1.1	1.9
9	Anemias (D50-D64)	50	0.9	1.6
10	Diabetes mellitus (E10-E14)	36	0.7	1.2
10	Legal intervention (Y35,Y89.0)	36	0.7	1.2
12	Cerebrovascular diseases (I60-I69)	22	0.4	0.7
13	Influenza and pneumonia (J10-J18)	20	0.4	0.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	0.3	*
15	Septicemia (A40-A41)	10	0.2	*
...	All other causes (Residual)	433	8.1	13.9
Black, male, 25-34 years				
...	All causes	6,848	100.0	264.5
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,129	31.1	82.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,309	19.1	50.6
3	Human immunodeficiency virus (HIV) disease (B20-B24)	634	9.3	24.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	619	9.0	23.9
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	447	6.5	17.3
6	Malignant neoplasms (C00-C97)	290	4.2	11.2
7	Diabetes mellitus (E10-E14)	113	1.7	4.4
8	Cerebrovascular diseases (I60-I69)	74	1.1	2.9
9	Anemias (D50-D64)	73	1.1	2.8
10	Chronic lower respiratory diseases (J40-J47)	58	0.8	2.2
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	49	0.7	1.9
12	Septicemia (A40-A41)	48	0.7	1.9
13	Influenza and pneumonia (J10-J18)	45	0.7	1.7
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	43	0.6	1.7
15	Legal intervention (Y35,Y89.0)	28	0.4	1.1
...	All other causes (Residual)	889	13.0	34.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 35-44 years				
...	All causes	11,849	100.0	434.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,060	17.4	75.6
2	Human immunodeficiency virus (HIV) disease (B20-B24)	2,017	17.0	74.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,592	13.4	58.4
4	Malignant neoplasms (C00-C97)	1,172	9.9	43.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,058	8.9	38.8
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	354	3.0	13.0
7	Cerebrovascular diseases (I60-I69)	332	2.8	12.2
8	Diabetes mellitus (E10-E14)	319	2.7	11.7
9	Chronic liver disease and cirrhosis (K70,K73-K74)	183	1.5	6.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	163	1.4	6.0
11	Chronic lower respiratory diseases (J40-J47)	155	1.3	5.7
12	Influenza and pneumonia (J10-J18)	138	1.2	5.1
13	Septicemia (A40-A41)	133	1.1	4.9
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	113	1.0	4.1
15	Viral hepatitis (B15-B19)	68	0.6	2.5
15	Anemias (D50-D64)	68	0.6	2.5
...	All other causes (Residual)	1,924	16.2	70.6
Black, male, 45-54 years				
...	All causes	21,122	100.0	983.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,286	25.0	246.0
2	Malignant neoplasms (C00-C97)	4,240	20.1	197.3
3	Human immunodeficiency virus (HIV) disease (B20-B24)	1,827	8.6	85.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,552	7.3	72.2
5	Cerebrovascular diseases (I60-I69)	1,000	4.7	46.5
6	Diabetes mellitus (E10-E14)	748	3.5	34.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	645	3.1	30.0
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	582	2.8	27.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	432	2.0	20.1
10	Septicemia (A40-A41)	315	1.5	14.7
11	Chronic lower respiratory diseases (J40-J47)	309	1.5	14.4
12	Viral hepatitis (B15-B19)	266	1.3	12.4
13	Influenza and pneumonia (J10-J18)	241	1.1	11.2
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	237	1.1	11.0
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	219	1.0	10.2
...	All other causes (Residual)	3,223	15.3	150.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 55-64 years				
...	All causes	24,000	100.0	2,039.2
1	Malignant neoplasms (C00-C97)	7,183	29.9	610.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,124	29.7	605.3
3	Cerebrovascular diseases (I60-I69)	1,298	5.4	110.3
4	Diabetes mellitus (E10-E14)	1,114	4.6	94.7
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	808	3.4	68.7
6	Chronic lower respiratory diseases (J40-J47)	615	2.6	52.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	578	2.4	49.1
8	Human immunodeficiency virus (HIV) disease (B20-B24)	523	2.2	44.4
9	Chronic liver disease and cirrhosis (K70,K73-K74)	445	1.9	37.8
10	Septicemia (A40-A41)	423	1.8	35.9
11	Influenza and pneumonia (J10-J18)	341	1.4	29.0
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	276	1.2	23.5
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	181	0.8	15.4
14	Aortic aneurysm and dissection (I71)	142	0.6	12.1
15	Viral hepatitis (B15-B19)	124	0.5	10.5
...	All other causes (Residual)	2,825	11.8	240.0
Black, male, 65-74 years				
...	All causes	28,198	100.0	4,024.5
1	Malignant neoplasms (C00-C97)	8,931	31.7	1,274.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,357	29.6	1,192.7
3	Cerebrovascular diseases (I60-I69)	1,842	6.5	262.9
4	Diabetes mellitus (E10-E14)	1,332	4.7	190.1
5	Chronic lower respiratory diseases (J40-J47)	1,107	3.9	158.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	806	2.9	115.0
7	Septicemia (A40-A41)	603	2.1	86.1
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	539	1.9	76.9
9	Influenza and pneumonia (J10-J18)	462	1.6	65.9
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	364	1.3	52.0
11	Chronic liver disease and cirrhosis (K70,K73-K74)	227	0.8	32.4
12	Human immunodeficiency virus (HIV) disease (B20-B24)	184	0.7	26.3
13	Pneumonitis due to solids and liquids (J69)	152	0.5	21.7
14	Aortic aneurysm and dissection (I71)	133	0.5	19.0
15	Alzheimer's disease (G30)	126	0.4	18.0
...	All other causes (Residual)	3,033	10.8	432.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 75-84 years				
...	All causes	28,478	100.0	8,169.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,539	30.0	2,449.6
2	Malignant neoplasms (C00-C97)	7,749	27.2	2,223.0
3	Cerebrovascular diseases (I60-I69)	2,049	7.2	587.8
4	Chronic lower respiratory diseases (J40-J47)	1,367	4.8	392.2
5	Diabetes mellitus (E10-E14)	1,150	4.0	329.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	820	2.9	235.2
7	Influenza and pneumonia (J10-J18)	798	2.8	228.9
8	Septicemia (A40-A41)	740	2.6	212.3
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	452	1.6	129.7
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	444	1.6	127.4
11	Alzheimer's disease (G30)	427	1.5	122.5
12	Pneumonitis due to solids and liquids (J69)	256	0.9	73.4
13	Atherosclerosis (I70)	154	0.5	44.2
14	Parkinson's disease (G20-G21)	138	0.5	39.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha'	135	0.5	38.7
...	All other causes (Residual)	3,260	11.4	935.2
Black, male, 85 years and over				
...	All causes	14,584	100.0	15,635.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,781	32.8	5,125.7
2	Malignant neoplasms (C00-C97)	2,776	19.0	2,976.1
3	Cerebrovascular diseases (I60-I69)	1,168	8.0	1,252.2
4	Influenza and pneumonia (J10-J18)	643	4.4	689.4
5	Chronic lower respiratory diseases (J40-J47)	602	4.1	645.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	524	3.6	561.8
7	Diabetes mellitus (E10-E14)	389	2.7	417.0
8	Alzheimer's disease (G30)	385	2.6	412.8
9	Septicemia (A40-A41)	367	2.5	393.5
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	256	1.8	274.5
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	243	1.7	260.5
12	Pneumonitis due to solids and liquids (J69)	218	1.5	233.7
13	Atherosclerosis (I70)	103	0.7	110.4
14	Parkinson's disease (G20-G21)	83	0.6	89.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha'	74	0.5	79.3
...	All other causes (Residual)	1,972	13.5	2,114.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, all ages				
...	All causes	143,216	100.0	724.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	40,527	28.3	205.0
2	Malignant neoplasms (C00-C97)	29,990	20.9	151.7
3	Cerebrovascular diseases (I60-I69)	11,028	7.7	55.8
4	Diabetes mellitus (E10-E14)	7,480	5.2	37.8
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,061	2.8	20.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,901	2.7	19.7
7	Chronic lower respiratory diseases (J40-J47)	3,490	2.4	17.7
8	Septicemia (A40-A41)	3,434	2.4	17.4
9	Influenza and pneumonia (J10-J18)	3,103	2.2	15.7
10	Human immunodeficiency virus (HIV) disease (B20-B24)	2,534	1.8	12.8
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,479	1.7	12.5
12	Alzheimer's disease (G30)	2,382	1.7	12.0
13	Certain conditions originating in the perinatal period (P00-P96)	2,206	1.5	11.2
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,391	1.0	7.0
15	Chronic liver disease and cirrhosis (K70,K73-K74)	966	0.7	4.9
...	All other causes (Residual)	24,244	16.9	122.6
Black, female, 1-4 years				
...	All causes	494	100.0	39.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	120	24.3	9.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	69	14.0	5.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	55	11.1	4.4
4	Malignant neoplasms (C00-C97)	40	8.1	3.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	22	4.5	1.8
6	Influenza and pneumonia (J10-J18)	19	3.8	*
7	Septicemia (A40-A41)	12	2.4	*
8	Chronic lower respiratory diseases (J40-J47)	10	2.0	*
9	Certain conditions originating in the perinatal period (P00-P96)	8	1.6	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	6	1.2	*
11	Anemias (D50-D64)	5	1.0	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.8	*
12	Cerebrovascular diseases (I60-I69)	4	0.8	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.6	*
15	Pneumonitis due to solids and liquids (J69)	2	0.4	*
...	All other causes (Residual)	115	23.3	9.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 5-14 years				
...	All causes	668	100.0	19.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	221	33.1	6.6
2	Malignant neoplasms (C00-C97)	66	9.9	2.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	49	7.3	1.5
4	Chronic lower respiratory diseases (J40-J47)	35	5.2	1.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	35	5.2	1.0
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	32	4.8	1.0
7	Septicemia (A40-A41)	13	1.9	*
8	Anemias (D50-D64)	12	1.8	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	9	1.3	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	9	1.3	*
9	Diabetes mellitus (E10-E14)	9	1.3	*
9	Influenza and pneumonia (J10-J18)	9	1.3	*
13	Cerebrovascular diseases (I60-I69)	8	1.2	*
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6	0.9	*
15	Meningitis (G00,G03)	5	0.7	*
...	All other causes (Residual)	150	22.5	4.5
Black, female, 15-24 years				
...	All causes	1,681	100.0	54.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	428	25.5	13.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	318	18.9	10.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	114	6.8	3.7
4	Malignant neoplasms (C00-C97)	110	6.5	3.6
5	Human immunodeficiency virus (HIV) disease (B20-B24)	64	3.8	2.1
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	52	3.1	1.7
7	Pregnancy, childbirth and the puerperium (O00-099)	46	2.7	1.5
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	40	2.4	1.3
9	Anemias (D50-D64)	29	1.7	0.9
10	Diabetes mellitus (E10-E14)	23	1.4	0.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21	1.2	0.7
12	Chronic lower respiratory diseases (J40-J47)	20	1.2	0.6
13	Influenza and pneumonia (J10-J18)	18	1.1	*
14	Cerebrovascular diseases (I60-I69)	16	1.0	*
15	Septicemia (A40-A41)	14	0.8	*
...	All other causes (Residual)	368	21.9	11.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 25-34 years				
...	All causes	3,323	100.0	116.4
1	Human immunodeficiency virus (HIV) disease (B20-B24)	489	14.7	17.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	402	12.1	14.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	385	11.6	13.5
4	Malignant neoplasms (C00-C97)	379	11.4	13.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	340	10.2	11.9
6	Cerebrovascular diseases (I60-I69)	94	2.8	3.3
7	Diabetes mellitus (E10-E14)	82	2.5	2.9
8	Pregnancy, childbirth and the puerperium (O00-099)	66	2.0	2.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	60	1.8	2.1
10	Anemias (D50-D64)	57	1.7	2.0
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	57	1.7	2.0
12	Septicemia (A40-A41)	55	1.7	1.9
13	Chronic lower respiratory diseases (J40-J47)	46	1.4	1.6
14	Influenza and pneumonia (J10-J18)	38	1.1	1.3
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	32	1.0	1.1
...	All other causes (Residual)	741	22.3	26.0
Black, female, 35-44 years				
...	All causes	8,386	100.0	272.3
1	Malignant neoplasms (C00-C97)	1,731	20.6	56.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,296	15.5	42.1
3	Human immunodeficiency virus (HIV) disease (B20-B24)	1,046	12.5	34.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	676	8.1	22.0
5	Cerebrovascular diseases (I60-I69)	412	4.9	13.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	319	3.8	10.4
7	Diabetes mellitus (E10-E14)	241	2.9	7.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	146	1.7	4.7
9	Chronic lower respiratory diseases (J40-J47)	143	1.7	4.6
10	Chronic liver disease and cirrhosis (K70,K73-K74)	135	1.6	4.4
11	Septicemia (A40-A41)	125	1.5	4.1
12	Influenza and pneumonia (J10-J18)	114	1.4	3.7
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	88	1.0	2.9
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	78	0.9	2.5
15	Anemias (D50-D64)	77	0.9	2.5
...	All other causes (Residual)	1,759	21.0	57.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 45-54 years				
...	All causes	14,501	100.0	579.4
1	Malignant neoplasms (C00-C97)	4,210	29.0	168.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,127	21.6	124.9
3	Cerebrovascular diseases (I60-I69)	893	6.2	35.7
4	Human immunodeficiency virus (HIV) disease (B20-B24)	682	4.7	27.2
5	Diabetes mellitus (E10-E14)	651	4.5	26.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	627	4.3	25.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	323	2.2	12.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	320	2.2	12.8
9	Chronic lower respiratory diseases (J40-J47)	290	2.0	11.6
10	Septicemia (A40-A41)	285	2.0	11.4
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	180	1.2	7.2
12	Influenza and pneumonia (J10-J18)	162	1.1	6.5
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	140	1.0	5.6
14	Viral hepatitis (B15-B19)	108	0.7	4.3
15	Anemias (D50-D64)	89	0.6	3.6
...	All other causes (Residual)	2,414	16.6	96.4
Black, female, 55-64 years				
...	All causes	17,336	100.0	1,184.2
1	Malignant neoplasms (C00-C97)	5,642	32.5	385.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,572	26.4	312.3
3	Diabetes mellitus (E10-E14)	1,110	6.4	75.8
4	Cerebrovascular diseases (I60-I69)	1,026	5.9	70.1
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	504	2.9	34.4
6	Chronic lower respiratory diseases (J40-J47)	461	2.7	31.5
7	Septicemia (A40-A41)	453	2.6	30.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	285	1.6	19.5
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	278	1.6	19.0
10	Influenza and pneumonia (J10-J18)	219	1.3	15.0
11	Chronic liver disease and cirrhosis (K70,K73-K74)	215	1.2	14.7
12	Human immunodeficiency virus (HIV) disease (B20-B24)	165	1.0	11.3
13	Viral hepatitis (B15-B19)	63	0.4	4.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior	61	0.4	4.2
15	Aortic aneurysm and dissection (I71)	55	0.3	3.8
...	All other causes (Residual)	2,227	12.8	152.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 65-74 years				
...	All causes	25,116	100.0	2,545.0
1	Malignant neoplasms (C00-C97)	7,314	29.1	741.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,244	28.8	734.0
3	Diabetes mellitus (E10-E14)	1,843	7.3	186.7
4	Cerebrovascular diseases (I60-I69)	1,788	7.1	181.2
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	884	3.5	89.6
6	Chronic lower respiratory diseases (J40-J47)	809	3.2	82.0
7	Septicemia (A40-A41)	657	2.6	66.6
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	439	1.7	44.5
9	Influenza and pneumonia (J10-J18)	386	1.5	39.1
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	293	1.2	29.7
11	Alzheimer's disease (G30)	157	0.6	15.9
12	Chronic liver disease and cirrhosis (K70,K73-K74)	138	0.5	14.0
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	113	0.4	11.5
13	Aortic aneurysm and dissection (I71)	113	0.4	11.5
15	Pneumonitis due to solids and liquids (J69)	98	0.4	9.9
...	All other causes (Residual)	2,840	11.3	287.8
Black, female, 75-84 years				
...	All causes	34,384	100.0	5,584.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,218	32.6	1,821.9
2	Malignant neoplasms (C00-C97)	6,915	20.1	1,123.1
3	Cerebrovascular diseases (I60-I69)	3,277	9.5	532.2
4	Diabetes mellitus (E10-E14)	2,099	6.1	340.9
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,139	3.3	185.0
6	Chronic lower respiratory diseases (J40-J47)	1,031	3.0	167.4
7	Septicemia (A40-A41)	911	2.6	148.0
8	Influenza and pneumonia (J10-J18)	805	2.3	130.7
9	Alzheimer's disease (G30)	760	2.2	123.4
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	694	2.0	112.7
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	354	1.0	57.5
12	Pneumonitis due to solids and liquids (J69)	247	0.7	40.1
13	Atherosclerosis (I70)	198	0.6	32.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	167	0.5	27.1
15	Aortic aneurysm and dissection (I71)	166	0.5	27.0
...	All other causes (Residual)	4,403	12.8	715.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 85 years and over				
...	All causes	33,438	100.0	13,734.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,444	37.2	5,111.2
2	Malignant neoplasms (C00-C97)	3,574	10.7	1,468.0
3	Cerebrovascular diseases (I60-I69)	3,492	10.4	1,434.3
4	Alzheimer's disease (G30)	1,446	4.3	593.9
5	Diabetes mellitus (E10-E14)	1,422	4.3	584.1
6	Influenza and pneumonia (J10-J18)	1,294	3.9	531.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	963	2.9	395.5
8	Septicemia (A40-A41)	869	2.6	356.9
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	784	2.3	322.0
10	Chronic lower respiratory diseases (J40-J47)	638	1.9	262.0
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	341	1.0	140.1
12	Pneumonitis due to solids and liquids (J69)	338	1.0	138.8
13	Atherosclerosis (I70)	291	0.9	119.5
14	Nutritional deficiencies (E40-E64)	160	0.5	65.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	140	0.4	57.5
...	All other causes (Residual)	5,242	15.7	2,153.1
American Indian, both sexes, all ages				
...	All causes	12,415	100.0	403.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,467	19.9	80.2
2	Malignant neoplasms (C00-C97)	2,175	17.5	70.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,488	12.0	48.4
4	Diabetes mellitus (E10-E14)	744	6.0	24.2
5	Cerebrovascular diseases (I60-I69)	567	4.6	18.4
6	Chronic liver disease and cirrhosis (K70,K73-K74)	547	4.4	17.8
7	Chronic lower respiratory diseases (J40-J47)	452	3.6	14.7
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	324	2.6	10.5
9	Influenza and pneumonia (J10-J18)	293	2.4	9.5
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	267	2.2	8.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	222	1.8	7.2
12	Septicemia (A40-A41)	195	1.6	6.3
13	Certain conditions originating in the perinatal period (P00-P96)	141	1.1	4.6
14	Alzheimer's disease (G30)	115	0.9	3.7
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	93	0.7	3.0
...	All other causes (Residual)	2,325	18.7	75.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 1-4 years				
...	All causes	90	100.0	45.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	33	36.7	16.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	12	13.3	*
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	6	6.7	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	5.6	*
5	Septicemia (A40-A41)	3	3.3	*
5	Malignant neoplasms (C00-C97)	3	3.3	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha'	2	2.2	*
8	Meningococcal infection (A39)	1	1.1	*
8	Influenza and pneumonia (J10-J18)	1	1.1	*
8	Acute bronchitis and bronchiolitis (J20-J21)	1	1.1	*
8	Chronic lower respiratory diseases (J40-J47)	1	1.1	*
...	All other causes (Residual)	22	24.4	11.0
American Indian, both sexes, 5-14 years				
...	All causes	126	100.0	21.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	60	47.6	10.3
2	Malignant neoplasms (C00-C97)	12	9.5	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	11	8.7	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	4.8	*
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6	4.8	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	4.0	*
7	Septicemia (A40-A41)	2	1.6	*
7	Cerebrovascular diseases (I60-I69)	2	1.6	*
9	Meningococcal infection (A39)	1	0.8	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha'	1	0.8	*
9	Anemias (D50-D64)	1	0.8	*
9	Meningitis (G00,G03)	1	0.8	*
9	Chronic lower respiratory diseases (J40-J47)	1	0.8	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.8	*
9	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.8	*
...	All other causes (Residual)	15	11.9	*

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 15-24 years				
...	All causes	583	100.0	104.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	300	51.5	53.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	100	17.2	17.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	70	12.0	12.6
4	Malignant neoplasms (C00-C97)	20	3.4	3.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	17	2.9	*
6	Influenza and pneumonia (J10-J18)	3	0.5	*
6	Legal intervention (Y35,Y89.0)	3	0.5	*
8	Septicemia (A40-A41)	2	0.3	*
8	Diabetes mellitus (E10-E14)	2	0.3	*
8	Cerebrovascular diseases (I60-I69)	2	0.3	*
8	Chronic lower respiratory diseases (J40-J47)	2	0.3	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.3	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.2	*
13	Anemias (D50-D64)	1	0.2	*
13	Meningitis (G00,G03)	1	0.2	*
13	Pneumonitis due to solids and liquids (J69)	1	0.2	*
13	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.2	*
13	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.2	*
13	Certain conditions originating in the perinatal period (P00-P96)	1	0.2	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.2	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
...	All other causes (Residual)	51	8.7	9.2
American Indian, both sexes, 25-34 years				
...	All causes	652	100.0	141.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	254	39.0	55.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	84	12.9	18.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	63	9.7	13.7
4	Malignant neoplasms (C00-C97)	35	5.4	7.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	35	5.4	7.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	34	5.2	7.4
7	Human immunodeficiency virus (HIV) disease (B20-B24)	13	2.0	*
8	Influenza and pneumonia (J10-J18)	7	1.1	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.9	*
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	0.9	*
11	Cerebrovascular diseases (I60-I69)	5	0.8	*
12	Septicemia (A40-A41)	4	0.6	*
13	Diabetes mellitus (E10-E14)	3	0.5	*
14	Viral hepatitis (B15-B19)	2	0.3	*
14	Anemias (D50-D64)	2	0.3	*
14	Hernia (K40-K46)	2	0.3	*
14	Legal intervention (Y35,Y89.0)	2	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.3	*
...	All other causes (Residual)	93	14.3	20.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 35-44 years				
...	All causes	1,169	100.0	248.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	333	28.5	70.8
2	Chronic liver disease and cirrhosis (K70,K73-K74)	142	12.1	30.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	107	9.2	22.7
4	Malignant neoplasms (C00-C97)	100	8.6	21.3
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	68	5.8	14.5
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	52	4.4	11.1
7	Human immunodeficiency virus (HIV) disease (B20-B24)	36	3.1	7.7
8	Diabetes mellitus (E10-E14)	33	2.8	7.0
8	Cerebrovascular diseases (I60-I69)	33	2.8	7.0
10	Septicemia (A40-A41)	13	1.1	*
10	Influenza and pneumonia (J10-J18)	13	1.1	*
12	Viral hepatitis (B15-B19)	12	1.0	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	0.9	*
14	Chronic lower respiratory diseases (J40-J47)	6	0.5	*
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	4	0.3	*
15	Pneumonitis due to solids and liquids (J69)	4	0.3	*
...	All other causes (Residual)	203	17.4	43.1
American Indian, both sexes, 45-54 years				
...	All causes	1,602	100.0	428.9
1	Malignant neoplasms (C00-C97)	293	18.3	78.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	246	15.4	65.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	215	13.4	57.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	162	10.1	43.4
5	Diabetes mellitus (E10-E14)	104	6.5	27.8
6	Cerebrovascular diseases (I60-I69)	54	3.4	14.5
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	45	2.8	12.0
8	Septicemia (A40-A41)	29	1.8	7.8
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	29	1.8	7.8
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	26	1.6	7.0
11	Viral hepatitis (B15-B19)	25	1.6	6.7
11	Chronic lower respiratory diseases (J40-J47)	25	1.6	6.7
13	Influenza and pneumonia (J10-J18)	23	1.4	6.2
14	Human immunodeficiency virus (HIV) disease (B20-B24)	11	0.7	*
15	Pneumonitis due to solids and liquids (J69)	6	0.4	*
...	All other causes (Residual)	309	19.3	82.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
	American Indian, both sexes, 55-64 years			
...	All causes	1,886	100.0	898.0
1	Malignant neoplasms (C00-C97)	485	25.7	230.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	412	21.8	196.2
3	Diabetes mellitus (E10-E14)	167	8.9	79.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	112	5.9	53.3
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	112	5.9	53.3
6	Chronic lower respiratory diseases (J40-J47)	72	3.8	34.3
7	Cerebrovascular diseases (I60-I69)	71	3.8	33.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	43	2.3	20.5
9	Septicemia (A40-A41)	27	1.4	12.9
10	Influenza and pneumonia (J10-J18)	25	1.3	11.9
11	Viral hepatitis (B15-B19)	13	0.7	*
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	13	0.7	*
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	11	0.6	*
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	9	0.5	*
15	Pneumonitis due to solids and liquids (J69)	8	0.4	*
...	All other causes (Residual)	306	16.2	145.7
	American Indian, both sexes, 65-74 years			
...	All causes	2,291	100.0	2,076.1
1	Malignant neoplasms (C00-C97)	589	25.7	533.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	542	23.7	491.2
3	Diabetes mellitus (E10-E14)	223	9.7	202.1
4	Chronic lower respiratory diseases (J40-J47)	135	5.9	122.3
5	Cerebrovascular diseases (I60-I69)	116	5.1	105.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	79	3.4	71.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	67	2.9	60.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	56	2.4	50.7
9	Septicemia (A40-A41)	46	2.0	41.7
10	Influenza and pneumonia (J10-J18)	35	1.5	31.7
11	Aortic aneurysm and dissection (I71)	19	0.8	*
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	18	0.8	*
13	Atherosclerosis (I70)	17	0.7	*
14	Parkinson's disease (G20-G21)	14	0.6	*
14	Alzheimer's disease (G30)	14	0.6	*
...	All other causes (Residual)	321	14.0	290.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 75-84 years				
...	All causes	2,215	100.0	4,110.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	633	28.6	1,174.6
2	Malignant neoplasms (C00-C97)	460	20.8	853.6
3	Diabetes mellitus (E10-E14)	162	7.3	300.6
4	Cerebrovascular diseases (I60-I69)	158	7.1	293.2
5	Chronic lower respiratory diseases (J40-J47)	139	6.3	257.9
6	Influenza and pneumonia (J10-J18)	81	3.7	150.3
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	57	2.6	105.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	46	2.1	85.4
9	Alzheimer's disease (G30)	42	1.9	77.9
10	Septicemia (A40-A41)	40	1.8	74.2
11	Chronic liver disease and cirrhosis (K70,K73-K74)	25	1.1	46.4
12	Pneumonitis due to solids and liquids (J69)	23	1.0	42.7
13	Parkinson's disease (G20-G21)	22	1.0	40.8
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	17	0.8	*
15	Atherosclerosis (I70)	16	0.7	*
...	All other causes (Residual)	294	13.3	545.5
American Indian, both sexes, 85 years and over				
...	All causes	1,451	100.0	8,299.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	459	31.6	2,625.4
2	Malignant neoplasms (C00-C97)	177	12.2	1,012.4
3	Cerebrovascular diseases (I60-I69)	125	8.6	715.0
4	Influenza and pneumonia (J10-J18)	99	6.8	566.3
5	Chronic lower respiratory diseases (J40-J47)	70	4.8	400.4
6	Alzheimer's disease (G30)	58	4.0	331.8
7	Diabetes mellitus (E10-E14)	50	3.4	286.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	30	2.1	171.6
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	29	2.0	165.9
10	Septicemia (A40-A41)	25	1.7	143.0
11	Pneumonitis due to solids and liquids (J69)	22	1.5	125.8
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	20	1.4	114.4
13	Parkinson's disease (G20-G21)	14	1.0	*
14	Nutritional deficiencies (E40-E64)	13	0.9	*
15	Atherosclerosis (I70)	8	0.6	*
...	All other causes (Residual)	252	17.4	1,441.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
	American Indian, male, all ages			
...	All causes	6,750	100.0	439.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,412	20.9	92.0
2	Malignant neoplasms (C00-C97)	1,081	16.0	70.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,003	14.9	65.3
4	Diabetes mellitus (E10-E14)	336	5.0	21.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	319	4.7	20.8
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	258	3.8	16.8
7	Cerebrovascular diseases (I60-I69)	236	3.5	15.4
8	Chronic lower respiratory diseases (J40-J47)	220	3.3	14.3
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	185	2.7	12.0
10	Influenza and pneumonia (J10-J18)	133	2.0	8.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	98	1.5	6.4
12	Septicemia (A40-A41)	95	1.4	6.2
13	Certain conditions originating in the perinatal period (P00-P96)	86	1.3	5.6
14	Human immunodeficiency virus (HIV) disease (B20-B24)	50	0.7	3.3
15	Pneumonitis due to solids and liquids (J69)	44	0.7	2.9
...	All other causes (Residual)	1,194	17.7	77.8
	American Indian, male, 1-4 years			
...	All causes	49	100.0	48.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19	38.8	*
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	10.2	*
3	Septicemia (A40-A41)	2	4.1	*
3	Malignant neoplasms (C00-C97)	2	4.1	*
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	4.1	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	4.1	*
7	Meningococcal infection (A39)	1	2.0	*
7	Influenza and pneumonia (J10-J18)	1	2.0	*
7	Acute bronchitis and bronchiolitis (J20-J21)	1	2.0	*
7	Chronic lower respiratory diseases (J40-J47)	1	2.0	*
...	All other causes (Residual)	13	26.5	*

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
	American Indian, male, 5-14 years			
...	All causes	65	100.0	22.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	31	47.7	10.5
2	Malignant neoplasms (C00-C97)	7	10.8	*
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	6.2	*
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4	6.2	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	4.6	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	4.6	*
7	Septicemia (A40-A41)	2	3.1	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	1	1.5	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	1.5	*
8	Infections of kidney (N10-N12,N13.6,N15.1)	1	1.5	*
...	All other causes (Residual)	8	12.3	*
	American Indian, male, 15-24 years			
...	All causes	416	100.0	145.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	212	51.0	74.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	80	19.2	27.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	54	13.0	18.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	14	3.4	*
5	Malignant neoplasms (C00-C97)	10	2.4	*
6	Legal intervention (Y35,Y89.0)	3	0.7	*
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.5	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.2	*
8	Diabetes mellitus (E10-E14)	1	0.2	*
8	Cerebrovascular diseases (I60-I69)	1	0.2	*
8	Influenza and pneumonia (J10-J18)	1	0.2	*
8	Pneumonitis due to solids and liquids (J69)	1	0.2	*
8	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.2	*
8	Certain conditions originating in the perinatal period (P00-P96)	1	0.2	*
8	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
...	All other causes (Residual)	33	7.9	11.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
	American Indian, male, 25-34 years			
...	All causes	457	100.0	193.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	186	40.7	78.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	74	16.2	31.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	46	10.1	19.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	25	5.5	10.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	15	3.3	*
6	Malignant neoplasms (C00-C97)	14	3.1	*
7	Human immunodeficiency virus (HIV) disease (B20-B24)	10	2.2	*
8	Septicemia (A40-A41)	4	0.9	*
8	Cerebrovascular diseases (I60-I69)	4	0.9	*
8	Influenza and pneumonia (J10-J18)	4	0.9	*
11	Diabetes mellitus (E10-E14)	3	0.7	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.7	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.7	*
14	Viral hepatitis (B15-B19)	2	0.4	*
14	Hernia (K40-K46)	2	0.4	*
14	Legal intervention (Y35,Y89.0)	2	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
...	All other causes (Residual)	58	12.7	24.5
	American Indian, male, 35-44 years			
...	All causes	749	100.0	321.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	228	30.4	97.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	82	10.9	35.2
2	Chronic liver disease and cirrhosis (K70,K73-K74)	82	10.9	35.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	52	6.9	22.3
5	Malignant neoplasms (C00-C97)	44	5.9	18.9
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	40	5.3	17.2
7	Human immunodeficiency virus (HIV) disease (B20-B24)	29	3.9	12.4
8	Diabetes mellitus (E10-E14)	21	2.8	9.0
9	Cerebrovascular diseases (I60-I69)	14	1.9	*
10	Influenza and pneumonia (J10-J18)	9	1.2	*
11	Viral hepatitis (B15-B19)	8	1.1	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	1.1	*
13	Septicemia (A40-A41)	6	0.8	*
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	4	0.5	*
15	Pneumonitis due to solids and liquids (J69)	3	0.4	*
...	All other causes (Residual)	119	15.9	51.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
	American Indian, male, 45-54 years			
...	All causes	978	100.0	539.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	189	19.3	104.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	149	15.2	82.2
3	Malignant neoplasms (C00-C97)	138	14.1	76.1
4	Chronic liver disease and cirrhosis (K70,K73-K74)	105	10.7	57.9
5	Diabetes mellitus (E10-E14)	57	5.8	31.4
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	30	3.1	16.5
7	Cerebrovascular diseases (I60-I69)	28	2.9	15.4
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	23	2.4	12.7
9	Viral hepatitis (B15-B19)	17	1.7	*
10	Septicemia (A40-A41)	16	1.6	*
11	Influenza and pneumonia (J10-J18)	14	1.4	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	1.4	*
13	Chronic lower respiratory diseases (J40-J47)	12	1.2	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	8	0.8	*
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	4	0.4	*
15	Pneumonitis due to solids and liquids (J69)	4	0.4	*
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	4	0.4	*
...	All other causes (Residual)	166	17.0	91.5
	American Indian, male, 55-64 years			
...	All causes	1,074	100.0	1,059.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	277	25.8	273.2
2	Malignant neoplasms (C00-C97)	265	24.7	261.4
3	Diabetes mellitus (E10-E14)	85	7.9	83.8
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	72	6.7	71.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	65	6.1	64.1
6	Cerebrovascular diseases (I60-I69)	35	3.3	34.5
6	Chronic lower respiratory diseases (J40-J47)	35	3.3	34.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20	1.9	19.7
9	Influenza and pneumonia (J10-J18)	13	1.2	*
10	Septicemia (A40-A41)	11	1.0	*
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	10	0.9	*
12	Viral hepatitis (B15-B19)	7	0.7	*
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	7	0.7	*
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	0.5	*
15	Atherosclerosis (I70)	4	0.4	*
15	Pneumonitis due to solids and liquids (J69)	4	0.4	*
...	All other causes (Residual)	159	14.8	156.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 65-74 years				
...	All causes	1,201	100.0	2,366.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	324	27.0	638.4
2	Malignant neoplasms (C00-C97)	307	25.6	604.9
3	Diabetes mellitus (E10-E14)	96	8.0	189.2
4	Chronic lower respiratory diseases (J40-J47)	64	5.3	126.1
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	56	4.7	110.3
6	Cerebrovascular diseases (I60-I69)	49	4.1	96.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	38	3.2	74.9
8	Septicemia (A40-A41)	23	1.9	45.3
9	Influenza and pneumonia (J10-J18)	19	1.6	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19	1.6	*
11	Aortic aneurysm and dissection (I71)	15	1.2	*
12	Parkinson's disease (G20-G21)	10	0.8	*
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	7	0.6	*
13	Atherosclerosis (I70)	7	0.6	*
15	Alzheimer's disease (G30)	5	0.4	*
15	Pneumonitis due to solids and liquids (J69)	5	0.4	*
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	5	0.4	*
...	All other causes (Residual)	152	12.7	299.5
American Indian, male, 75-84 years				
...	All causes	1,048	100.0	4,748.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	314	30.0	1,422.7
2	Malignant neoplasms (C00-C97)	236	22.5	1,069.3
3	Chronic lower respiratory diseases (J40-J47)	77	7.3	348.9
4	Cerebrovascular diseases (I60-I69)	61	5.8	276.4
5	Diabetes mellitus (E10-E14)	57	5.4	258.3
6	Influenza and pneumonia (J10-J18)	33	3.1	149.5
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	30	2.9	135.9
8	Septicemia (A40-A41)	23	2.2	104.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20	1.9	90.6
10	Parkinson's disease (G20-G21)	16	1.5	*
11	Alzheimer's disease (G30)	15	1.4	*
12	Pneumonitis due to solids and liquids (J69)	13	1.2	*
13	Chronic liver disease and cirrhosis (K70,K73-K74)	12	1.1	*
14	Aortic aneurysm and dissection (I71)	7	0.7	*
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	6	0.6	*
...	All other causes (Residual)	128	12.2	579.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 85 years and over				
...	All causes	516	100.0	9,219.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	177	34.3	3,162.4
2	Malignant neoplasms (C00-C97)	58	11.2	1,036.3
3	Cerebrovascular diseases (I60-I69)	43	8.3	768.3
4	Influenza and pneumonia (J10-J18)	36	7.0	643.2
5	Chronic lower respiratory diseases (J40-J47)	28	5.4	500.3
6	Diabetes mellitus (E10-E14)	16	3.1	*
7	Alzheimer's disease (G30)	14	2.7	*
8	Pneumonitis due to solids and liquids (J69)	13	2.5	*
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11	2.1	*
10	Parkinson's disease (G20-G21)	9	1.7	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	1.7	*
12	Septicemia (A40-A41)	6	1.2	*
13	Nutritional deficiencies (E40-E64)	4	0.8	*
14	Tuberculosis (A16-A19)	2	0.4	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	2	0.4	*
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2	0.4	*
14	Atherosclerosis (I70)	2	0.4	*
14	Aortic aneurysm and dissection (I71)	2	0.4	*
...	All other causes (Residual)	82	15.9	1,465.1
American Indian, female, all ages				
...	All causes	5,665	100.0	367.7
1	Malignant neoplasms (C00-C97)	1,094	19.3	71.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,055	18.6	68.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	485	8.6	31.5
4	Diabetes mellitus (E10-E14)	408	7.2	26.5
5	Cerebrovascular diseases (I60-I69)	331	5.8	21.5
6	Chronic lower respiratory diseases (J40-J47)	232	4.1	15.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	228	4.0	14.8
8	Influenza and pneumonia (J10-J18)	160	2.8	10.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	124	2.2	8.0
10	Septicemia (A40-A41)	100	1.8	6.5
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	82	1.4	5.3
12	Alzheimer's disease (G30)	80	1.4	5.2
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	66	1.2	4.3
14	Certain conditions originating in the perinatal period (P00-P96)	55	1.0	3.6
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	51	0.9	3.3
...	All other causes (Residual)	1,114	19.7	72.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
	American Indian, female, 1-4 years			
...	All causes	41	100.0	42.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14	34.1	*
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7	17.1	*
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	9.8	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	7.3	*
5	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	2	4.9	*
6	Septicemia (A40-A41)	1	2.4	*
6	Malignant neoplasms (C00-C97)	1	2.4	*
...	All other causes (Residual)	9	22.0	*
	American Indian, female, 5-14 years			
...	All causes	61	100.0	21.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	29	47.5	10.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8	13.1	*
3	Malignant neoplasms (C00-C97)	5	8.2	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	4.9	*
5	Cerebrovascular diseases (I60-I69)	2	3.3	*
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2	3.3	*
7	Meningococcal infection (A39)	1	1.6	*
7	Anemias (D50-D64)	1	1.6	*
7	Meningitis (G00,G03)	1	1.6	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	1.6	*
7	Chronic lower respiratory diseases (J40-J47)	1	1.6	*
...	All other causes (Residual)	7	11.5	*

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
	American Indian, female, 15-24 years			
...	All causes	167	100.0	61.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	88	52.7	32.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	20	12.0	7.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	16	9.6	*
4	Malignant neoplasms (C00-C97)	10	6.0	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	1.8	*
6	Septicemia (A40-A41)	2	1.2	*
6	Influenza and pneumonia (J10-J18)	2	1.2	*
6	Chronic lower respiratory diseases (J40-J47)	2	1.2	*
9	Anemias (D50-D64)	1	0.6	*
9	Diabetes mellitus (E10-E14)	1	0.6	*
9	Meningitis (G00,G03)	1	0.6	*
9	Cerebrovascular diseases (I60-I69)	1	0.6	*
9	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.6	*
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.6	*
...	All other causes (Residual)	18	10.8	*
	American Indian, female, 25-34 years			
...	All causes	195	100.0	87.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	68	34.9	30.5
2	Malignant neoplasms (C00-C97)	21	10.8	9.4
3	Chronic liver disease and cirrhosis (K70,K73-K74)	20	10.3	9.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	17	8.7	*
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	10	5.1	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	9	4.6	*
7	Human immunodeficiency virus (HIV) disease (B20-B24)	3	1.5	*
7	Influenza and pneumonia (J10-J18)	3	1.5	*
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	1.5	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	1.5	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	1	0.5	*
11	Anemias (D50-D64)	1	0.5	*
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1	0.5	*
11	Cerebrovascular diseases (I60-I69)	1	0.5	*
11	Aortic aneurysm and dissection (I71)	1	0.5	*
...	All other causes (Residual)	33	16.9	14.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 35-44 years				
...	All causes	420	100.0	176.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	105	25.0	44.2
2	Chronic liver disease and cirrhosis (K70,K73-K74)	60	14.3	25.3
3	Malignant neoplasms (C00-C97)	56	13.3	23.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	25	6.0	10.5
5	Cerebrovascular diseases (I60-I69)	19	4.5	*
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	16	3.8	*
7	Diabetes mellitus (E10-E14)	12	2.9	*
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	12	2.9	*
9	Septicemia (A40-A41)	7	1.7	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	7	1.7	*
11	Viral hepatitis (B15-B19)	4	1.0	*
11	Influenza and pneumonia (J10-J18)	4	1.0	*
11	Chronic lower respiratory diseases (J40-J47)	4	1.0	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha'	3	0.7	*
15	Tuberculosis (A16-A19)	2	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.5	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	0.5	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.5	*
...	All other causes (Residual)	78	18.6	32.8
American Indian, female, 45-54 years				
...	All causes	624	100.0	324.7
1	Malignant neoplasms (C00-C97)	155	24.8	80.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	66	10.6	34.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	57	9.1	29.7
3	Chronic liver disease and cirrhosis (K70,K73-K74)	57	9.1	29.7
5	Diabetes mellitus (E10-E14)	47	7.5	24.5
6	Cerebrovascular diseases (I60-I69)	26	4.2	13.5
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	15	2.4	*
8	Septicemia (A40-A41)	13	2.1	*
8	Chronic lower respiratory diseases (J40-J47)	13	2.1	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	1.9	*
11	Influenza and pneumonia (J10-J18)	9	1.4	*
12	Viral hepatitis (B15-B19)	8	1.3	*
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	1.0	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	3	0.5	*
14	Infections of kidney (N10-N12,N13.6,N15.1)	3	0.5	*
...	All other causes (Residual)	134	21.5	69.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
	American Indian, female, 55-64 years			
...	All causes	812	100.0	747.5
1	Malignant neoplasms (C00-C97)	220	27.1	202.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	135	16.6	124.3
3	Diabetes mellitus (E10-E14)	82	10.1	75.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	47	5.8	43.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	40	4.9	36.8
6	Chronic lower respiratory diseases (J40-J47)	37	4.6	34.1
7	Cerebrovascular diseases (I60-I69)	36	4.4	33.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	2.8	21.2
9	Septicemia (A40-A41)	16	2.0	*
10	Influenza and pneumonia (J10-J18)	12	1.5	*
11	Viral hepatitis (B15-B19)	6	0.7	*
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	4	0.5	*
12	Aortic aneurysm and dissection (I71)	4	0.5	*
12	Pneumonitis due to solids and liquids (J69)	4	0.5	*
12	Peptic ulcer (K25-K28)	4	0.5	*
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	0.5	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.5	*
...	All other causes (Residual)	134	16.5	123.4
	American Indian, female, 65-74 years			
...	All causes	1,090	100.0	1,828.9
1	Malignant neoplasms (C00-C97)	282	25.9	473.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	218	20.0	365.8
3	Diabetes mellitus (E10-E14)	127	11.7	213.1
4	Chronic lower respiratory diseases (J40-J47)	71	6.5	119.1
5	Cerebrovascular diseases (I60-I69)	67	6.1	112.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	37	3.4	62.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	29	2.7	48.7
8	Septicemia (A40-A41)	23	2.1	38.6
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	23	2.1	38.6
10	Influenza and pneumonia (J10-J18)	16	1.5	*
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	11	1.0	*
12	Atherosclerosis (I70)	10	0.9	*
13	Alzheimer's disease (G30)	9	0.8	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	5	0.5	*
15	Tuberculosis (A16-A19)	4	0.4	*
15	Parkinson's disease (G20-G21)	4	0.4	*
15	Aortic aneurysm and dissection (I71)	4	0.4	*
...	All other causes (Residual)	150	13.8	251.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
	American Indian, female, 75-84 years			
...	All causes	1,167	100.0	3,667.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	319	27.3	1,002.5
2	Malignant neoplasms (C00-C97)	224	19.2	703.9
3	Diabetes mellitus (E10-E14)	105	9.0	330.0
4	Cerebrovascular diseases (I60-I69)	97	8.3	304.8
5	Chronic lower respiratory diseases (J40-J47)	62	5.3	194.8
6	Influenza and pneumonia (J10-J18)	48	4.1	150.8
7	Alzheimer's disease (G30)	27	2.3	84.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	27	2.3	84.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	26	2.2	81.7
10	Septicemia (A40-A41)	17	1.5	*
11	Chronic liver disease and cirrhosis (K70,K73-K74)	13	1.1	*
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	11	0.9	*
12	Atherosclerosis (I70)	11	0.9	*
14	Pneumonitis due to solids and liquids (J69)	10	0.9	*
15	Parkinson's disease (G20-G21)	6	0.5	*
15	Aortic aneurysm and dissection (I71)	6	0.5	*
...	All other causes (Residual)	158	13.5	496.5
	American Indian, female, 85 years and over			
...	All causes	935	100.0	7,866.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	282	30.2	2,372.5
2	Malignant neoplasms (C00-C97)	119	12.7	1,001.2
3	Cerebrovascular diseases (I60-I69)	82	8.8	689.9
4	Influenza and pneumonia (J10-J18)	63	6.7	530.0
5	Alzheimer's disease (G30)	44	4.7	370.2
6	Chronic lower respiratory diseases (J40-J47)	42	4.5	353.4
7	Diabetes mellitus (E10-E14)	34	3.6	286.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21	2.2	176.7
9	Septicemia (A40-A41)	19	2.0	*
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	18	1.9	*
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18	1.9	*
12	Nutritional deficiencies (E40-E64)	9	1.0	*
12	Pneumonitis due to solids and liquids (J69)	9	1.0	*
14	Atherosclerosis (I70)	6	0.6	*
15	Parkinson's disease (G20-G21)	5	0.5	*
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	5	0.5	*
...	All other causes (Residual)	159	17.0	1,337.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, all ages				
...	All causes	38,332	100.0	299.5
1	Malignant neoplasms (C00-C97)	9,998	26.1	78.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,983	26.0	78.0
3	Cerebrovascular diseases (I60-I69)	3,530	9.2	27.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,875	4.9	14.7
5	Diabetes mellitus (E10-E14)	1,359	3.5	10.6
6	Influenza and pneumonia (J10-J18)	1,171	3.1	9.1
7	Chronic lower respiratory diseases (J40-J47)	1,138	3.0	8.9
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	661	1.7	5.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	649	1.7	5.1
10	Septicemia (A40-A41)	423	1.1	3.3
11	Certain conditions originating in the perinatal period (P00-P96)	405	1.1	3.2
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	402	1.0	3.1
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	399	1.0	3.1
14	Alzheimer's disease (G30)	352	0.9	2.8
15	Chronic liver disease and cirrhosis (K70,K73-K74)	313	0.8	2.4
...	All other causes (Residual)	5,674	14.8	44.3
Asian or Pacific Islander, both sexes, 1-4 years				
...	All causes	166	100.0	23.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	50	30.1	7.0
2	Malignant neoplasms (C00-C97)	19	11.4	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	16	9.6	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	11	6.6	*
5	Influenza and pneumonia (J10-J18)	5	3.0	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	2.4	*
6	Chronic lower respiratory diseases (J40-J47)	4	2.4	*
8	Certain conditions originating in the perinatal period (P00-P96)	3	1.8	*
9	Viral hepatitis (B15-B19)	2	1.2	*
9	Cerebrovascular diseases (I60-I69)	2	1.2	*
11	Shigellosis and amebiasis (A03,A06)	1	0.6	*
11	Septicemia (A40-A41)	1	0.6	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior	1	0.6	*
...	All other causes (Residual)	47	28.3	6.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 5-14 years				
...	All causes	220	100.0	12.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	72	32.7	4.1
2	Malignant neoplasms (C00-C97)	42	19.1	2.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	15	6.8	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	9	4.1	*
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	9	4.1	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	3.2	*
7	Septicemia (A40-A41)	4	1.8	*
7	Chronic lower respiratory diseases (J40-J47)	4	1.8	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha'	3	1.4	*
9	Anemias (D50-D64)	3	1.4	*
9	Influenza and pneumonia (J10-J18)	3	1.4	*
12	Cerebrovascular diseases (I60-I69)	2	0.9	*
12	Certain conditions originating in the perinatal period (P00-P96)	2	0.9	*
14	Meningococcal infection (A39)	1	0.5	*
14	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.5	*
...	All other causes (Residual)	43	19.5	2.4
Asian or Pacific Islander, both sexes, 15-24 years				
...	All causes	777	100.0	41.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	338	43.5	18.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	112	14.4	6.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	101	13.0	5.4
4	Malignant neoplasms (C00-C97)	61	7.9	3.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	32	4.1	1.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	1.0	*
7	Septicemia (A40-A41)	6	0.8	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha'	4	0.5	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.5	*
10	Aortic aneurysm and dissection (I71)	3	0.4	*
10	Influenza and pneumonia (J10-J18)	3	0.4	*
10	Pregnancy, childbirth and the puerperium (O00-099)	3	0.4	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.4	*
14	Tuberculosis (A16-A19)	2	0.3	*
14	Chronic lower respiratory diseases (J40-J47)	2	0.3	*
14	Legal intervention (Y35,Y89.0)	2	0.3	*
...	All other causes (Residual)	93	12.0	4.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 25-34 years				
...	All causes	961	100.0	40.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	226	23.5	9.4
2	Malignant neoplasms (C00-C97)	171	17.8	7.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	160	16.6	6.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	92	9.6	3.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	87	9.1	3.6
6	Cerebrovascular diseases (I60-I69)	18	1.9	*
7	Human immunodeficiency virus (HIV) disease (B20-B24)	15	1.6	*
8	Pregnancy, childbirth and the puerperium (O00-099)	12	1.2	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior	10	1.0	*
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	10	1.0	*
11	Influenza and pneumonia (J10-J18)	7	0.7	*
11	Chronic lower respiratory diseases (J40-J47)	7	0.7	*
13	Septicemia (A40-A41)	5	0.5	*
13	Viral hepatitis (B15-B19)	5	0.5	*
13	Anemias (D50-D64)	5	0.5	*
13	Diabetes mellitus (E10-E14)	5	0.5	*
...	All other causes (Residual)	126	13.1	5.3
Asian or Pacific Islander, both sexes, 35-44 years				
...	All causes	1,643	100.0	76.1
1	Malignant neoplasms (C00-C97)	500	30.4	23.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	258	15.7	12.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	208	12.7	9.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	122	7.4	5.7
5	Cerebrovascular diseases (I60-I69)	101	6.1	4.7
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	71	4.3	3.3
7	Human immunodeficiency virus (HIV) disease (B20-B24)	38	2.3	1.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	38	2.3	1.8
9	Diabetes mellitus (E10-E14)	17	1.0	*
10	Viral hepatitis (B15-B19)	16	1.0	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	0.8	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	13	0.8	*
13	Septicemia (A40-A41)	11	0.7	*
14	Influenza and pneumonia (J10-J18)	9	0.5	*
15	Chronic lower respiratory diseases (J40-J47)	8	0.5	*
...	All other causes (Residual)	220	13.4	10.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 45-54 years				
...	All causes	3,344	100.0	195.3
1	Malignant neoplasms (C00-C97)	1,378	41.2	80.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	632	18.9	36.9
3	Cerebrovascular diseases (I60-I69)	234	7.0	13.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	212	6.3	12.4
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	105	3.1	6.1
6	Diabetes mellitus (E10-E14)	85	2.5	5.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	66	2.0	3.9
8	Viral hepatitis (B15-B19)	51	1.5	3.0
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	47	1.4	2.7
10	Chronic lower respiratory diseases (J40-J47)	39	1.2	2.3
11	Influenza and pneumonia (J10-J18)	36	1.1	2.1
12	Septicemia (A40-A41)	35	1.0	2.0
13	Human immunodeficiency virus (HIV) disease (B20-B24)	33	1.0	1.9
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	33	1.0	1.9
15	Aortic aneurysm and dissection (I71)	19	0.6	*
...	All other causes (Residual)	339	10.1	19.8
Asian or Pacific Islander, both sexes, 55-64 years				
...	All causes	4,706	100.0	475.5
1	Malignant neoplasms (C00-C97)	1,941	41.2	196.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,037	22.0	104.8
3	Cerebrovascular diseases (I60-I69)	356	7.6	36.0
4	Diabetes mellitus (E10-E14)	191	4.1	19.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	155	3.3	15.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	82	1.7	8.3
7	Chronic lower respiratory diseases (J40-J47)	79	1.7	8.0
8	Viral hepatitis (B15-B19)	68	1.4	6.9
9	Chronic liver disease and cirrhosis (K70,K73-K74)	66	1.4	6.7
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	64	1.4	6.5
11	Septicemia (A40-A41)	47	1.0	4.7
12	Influenza and pneumonia (J10-J18)	45	1.0	4.5
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	36	0.8	3.6
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	27	0.6	2.7
15	Aortic aneurysm and dissection (I71)	26	0.6	2.6
...	All other causes (Residual)	486	10.3	49.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 65-74 years				
...	All causes	7,338	100.0	1,226.7
1	Malignant neoplasms (C00-C97)	2,465	33.6	412.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,912	26.1	319.6
3	Cerebrovascular diseases (I60-I69)	674	9.2	112.7
4	Diabetes mellitus (E10-E14)	369	5.0	61.7
5	Chronic lower respiratory diseases (J40-J47)	236	3.2	39.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	208	2.8	34.8
7	Influenza and pneumonia (J10-J18)	134	1.8	22.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	115	1.6	19.2
9	Septicemia (A40-A41)	92	1.3	15.4
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	74	1.0	12.4
11	Chronic liver disease and cirrhosis (K70,K73-K74)	72	1.0	12.0
12	Viral hepatitis (B15-B19)	66	0.9	11.0
13	Aortic aneurysm and dissection (I71)	65	0.9	10.9
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	49	0.7	8.2
15	Parkinson's disease (G20-G21)	37	0.5	6.2
...	All other causes (Residual)	770	10.5	128.7
Asian or Pacific Islander, both sexes, 75-84 years				
...	All causes	10,224	100.0	3,290.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,069	30.0	987.7
2	Malignant neoplasms (C00-C97)	2,413	23.6	776.6
3	Cerebrovascular diseases (I60-I69)	1,107	10.8	356.3
4	Chronic lower respiratory diseases (J40-J47)	446	4.4	143.5
5	Diabetes mellitus (E10-E14)	438	4.3	141.0
6	Influenza and pneumonia (J10-J18)	402	3.9	129.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	232	2.3	74.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	225	2.2	72.4
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	121	1.2	38.9
10	Septicemia (A40-A41)	117	1.1	37.7
11	Alzheimer's disease (G30)	115	1.1	37.0
12	Parkinson's disease (G20-G21)	92	0.9	29.6
13	Aortic aneurysm and dissection (I71)	89	0.9	28.6
14	Pneumonitis due to solids and liquids (J69)	66	0.6	21.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha'	65	0.6	20.9
...	All other causes (Residual)	1,227	12.0	394.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 85 years and over				
...	All causes	8,148	100.0	9,378.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,924	35.9	3,365.6
2	Cerebrovascular diseases (I60-I69)	1,027	12.6	1,182.1
3	Malignant neoplasms (C00-C97)	1,003	12.3	1,154.5
4	Influenza and pneumonia (J10-J18)	519	6.4	597.4
5	Chronic lower respiratory diseases (J40-J47)	312	3.8	359.1
6	Diabetes mellitus (E10-E14)	253	3.1	291.2
7	Alzheimer's disease (G30)	204	2.5	234.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	164	2.0	188.8
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	161	2.0	185.3
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	150	1.8	172.7
11	Pneumonitis due to solids and liquids (J69)	97	1.2	111.6
12	Septicemia (A40-A41)	95	1.2	109.3
13	Parkinson's disease (G20-G21)	76	0.9	87.5
14	Atherosclerosis (I70)	67	0.8	77.1
15	Aortic aneurysm and dissection (I71)	57	0.7	65.6
...	All other causes (Residual)	1,039	12.8	1,195.9
Asian or Pacific Islander, male, all ages				
...	All causes	20,483	100.0	331.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,523	27.0	89.4
2	Malignant neoplasms (C00-C97)	5,193	25.4	84.0
3	Cerebrovascular diseases (I60-I69)	1,599	7.8	25.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,175	5.7	19.0
5	Chronic lower respiratory diseases (J40-J47)	743	3.6	12.0
6	Diabetes mellitus (E10-E14)	648	3.2	10.5
7	Influenza and pneumonia (J10-J18)	636	3.1	10.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	469	2.3	7.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	320	1.6	5.2
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	277	1.4	4.5
11	Septicemia (A40-A41)	230	1.1	3.7
12	Certain conditions originating in the perinatal period (P00-P96)	225	1.1	3.6
13	Chronic liver disease and cirrhosis (K70,K73-K74)	182	0.9	2.9
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	181	0.9	2.9
15	Aortic aneurysm and dissection (I71)	174	0.8	2.8
...	All other causes (Residual)	2,908	14.2	47.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 1-4 years				
...	All causes	97	100.0	27.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	30	30.9	8.4
2	Malignant neoplasms (C00-C97)	12	12.4	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	10	10.3	*
4	Influenza and pneumonia (J10-J18)	5	5.2	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	4.1	*
6	Certain conditions originating in the perinatal period (P00-P96)	3	3.1	*
7	Viral hepatitis (B15-B19)	2	2.1	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	2.1	*
7	Chronic lower respiratory diseases (J40-J47)	2	2.1	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha'	1	1.0	*
10	Cerebrovascular diseases (I60-I69)	1	1.0	*
...	All other causes (Residual)	25	25.8	7.0
Asian or Pacific Islander, male, 5-14 years				
...	All causes	130	100.0	14.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	42	32.3	4.7
2	Malignant neoplasms (C00-C97)	28	21.5	3.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	10	7.7	*
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7	5.4	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	3.8	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	3.8	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha'	2	1.5	*
7	Cerebrovascular diseases (I60-I69)	2	1.5	*
7	Influenza and pneumonia (J10-J18)	2	1.5	*
7	Chronic lower respiratory diseases (J40-J47)	2	1.5	*
11	Septicemia (A40-A41)	1	0.8	*
11	Anemias (D50-D64)	1	0.8	*
11	Certain conditions originating in the perinatal period (P00-P96)	1	0.8	*
...	All other causes (Residual)	22	16.9	2.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 15-24 years				
...	All causes	554	100.0	58.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	249	44.9	26.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	92	16.6	9.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	82	14.8	8.7
4	Malignant neoplasms (C00-C97)	36	6.5	3.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	23	4.2	2.4
6	Septicemia (A40-A41)	4	0.7	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	0.7	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha'	3	0.5	*
8	Aortic aneurysm and dissection (I71)	3	0.5	*
10	Influenza and pneumonia (J10-J18)	2	0.4	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
10	Legal intervention (Y35,Y89.0)	2	0.4	*
13	Tuberculosis (A16-A19)	1	0.2	*
13	Meningococcal infection (A39)	1	0.2	*
13	Viral hepatitis (B15-B19)	1	0.2	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.2	*
13	Diabetes mellitus (E10-E14)	1	0.2	*
13	Cerebrovascular diseases (I60-I69)	1	0.2	*
13	Chronic lower respiratory diseases (J40-J47)	1	0.2	*
13	Pneumonitis due to solids and liquids (J69)	1	0.2	*
...	All other causes (Residual)	44	7.9	4.7
Asian or Pacific Islander, male, 25-34 years				
...	All causes	633	100.0	54.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	154	24.3	13.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	119	18.8	10.2
3	Malignant neoplasms (C00-C97)	92	14.5	7.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	67	10.6	5.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	59	9.3	5.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	11	1.7	*
6	Cerebrovascular diseases (I60-I69)	11	1.7	*
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	1.1	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha'	6	0.9	*
10	Septicemia (A40-A41)	5	0.8	*
10	Chronic lower respiratory diseases (J40-J47)	5	0.8	*
12	Viral hepatitis (B15-B19)	4	0.6	*
12	Influenza and pneumonia (J10-J18)	4	0.6	*
14	Anemias (D50-D64)	3	0.5	*
14	Diabetes mellitus (E10-E14)	3	0.5	*
...	All other causes (Residual)	83	13.1	7.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 35-44 years				
...	All causes	1,040	100.0	100.0
1	Malignant neoplasms (C00-C97)	236	22.7	22.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	208	20.0	20.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	152	14.6	14.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	86	8.3	8.3
5	Cerebrovascular diseases (I60-I69)	59	5.7	5.7
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	48	4.6	4.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	31	3.0	3.0
8	Human immunodeficiency virus (HIV) disease (B20-B24)	29	2.8	2.8
9	Viral hepatitis (B15-B19)	14	1.3	*
10	Diabetes mellitus (E10-E14)	12	1.2	*
11	Septicemia (A40-A41)	8	0.8	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.7	*
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	0.7	*
14	Influenza and pneumonia (J10-J18)	6	0.6	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha'	5	0.5	*
...	All other causes (Residual)	132	12.7	12.7
Asian or Pacific Islander, male, 45-54 years				
...	All causes	1,971	100.0	248.4
1	Malignant neoplasms (C00-C97)	657	33.3	82.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	481	24.4	60.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	127	6.4	16.0
4	Cerebrovascular diseases (I60-I69)	118	6.0	14.9
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	70	3.6	8.8
6	Diabetes mellitus (E10-E14)	52	2.6	6.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	46	2.3	5.8
8	Viral hepatitis (B15-B19)	40	2.0	5.0
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	32	1.6	4.0
10	Human immunodeficiency virus (HIV) disease (B20-B24)	31	1.6	3.9
11	Influenza and pneumonia (J10-J18)	26	1.3	3.3
12	Chronic lower respiratory diseases (J40-J47)	21	1.1	2.6
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18	0.9	*
14	Septicemia (A40-A41)	17	0.9	*
15	Aortic aneurysm and dissection (I71)	14	0.7	*
...	All other causes (Residual)	221	11.2	27.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 55-64 years				
...	All causes	2,738	100.0	594.5
1	Malignant neoplasms (C00-C97)	1,035	37.8	224.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	710	25.9	154.2
3	Cerebrovascular diseases (I60-I69)	186	6.8	40.4
4	Diabetes mellitus (E10-E14)	113	4.1	24.5
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	94	3.4	20.4
6	Chronic lower respiratory diseases (J40-J47)	53	1.9	11.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	46	1.7	10.0
8	Viral hepatitis (B15-B19)	44	1.6	9.6
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	44	1.6	9.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	40	1.5	8.7
11	Influenza and pneumonia (J10-J18)	26	0.9	5.6
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	25	0.9	5.4
13	Septicemia (A40-A41)	24	0.9	5.2
14	Aortic aneurysm and dissection (I71)	22	0.8	4.8
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	19	0.7	*
...	All other causes (Residual)	257	9.4	55.8
Asian or Pacific Islander, male, 65-74 years				
...	All causes	3,887	100.0	1,487.1
1	Malignant neoplasms (C00-C97)	1,259	32.4	481.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,104	28.4	422.4
3	Cerebrovascular diseases (I60-I69)	295	7.6	112.9
4	Diabetes mellitus (E10-E14)	189	4.9	72.3
5	Chronic lower respiratory diseases (J40-J47)	153	3.9	58.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	120	3.1	45.9
7	Influenza and pneumonia (J10-J18)	78	2.0	29.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	63	1.6	24.1
9	Aortic aneurysm and dissection (I71)	47	1.2	18.0
10	Septicemia (A40-A41)	46	1.2	17.6
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	42	1.1	16.1
12	Chronic liver disease and cirrhosis (K70,K73-K74)	31	0.8	11.9
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	29	0.7	11.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	23	0.6	8.8
15	Viral hepatitis (B15-B19)	19	0.5	*
...	All other causes (Residual)	389	10.0	148.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 75-84 years				
...	All causes	5,324	100.0	4,090.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,630	30.6	1,252.4
2	Malignant neoplasms (C00-C97)	1,318	24.8	1,012.7
3	Cerebrovascular diseases (I60-I69)	508	9.5	390.3
4	Chronic lower respiratory diseases (J40-J47)	307	5.8	235.9
5	Influenza and pneumonia (J10-J18)	221	4.2	169.8
6	Diabetes mellitus (E10-E14)	168	3.2	129.1
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	115	2.2	88.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	108	2.0	83.0
9	Septicemia (A40-A41)	72	1.4	55.3
10	Aortic aneurysm and dissection (I71)	56	1.1	43.0
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	54	1.0	41.5
12	Parkinson's disease (G20-G21)	53	1.0	40.7
13	Pneumonitis due to solids and liquids (J69)	43	0.8	33.0
14	Alzheimer's disease (G30)	39	0.7	30.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha'	37	0.7	28.4
...	All other causes (Residual)	595	11.2	457.2
Asian or Pacific Islander, male, 85 years and over				
...	All causes	3,662	100.0	10,938.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,286	35.1	3,841.3
2	Malignant neoplasms (C00-C97)	517	14.1	1,544.3
3	Cerebrovascular diseases (I60-I69)	413	11.3	1,233.6
4	Influenza and pneumonia (J10-J18)	260	7.1	776.6
5	Chronic lower respiratory diseases (J40-J47)	195	5.3	582.5
6	Diabetes mellitus (E10-E14)	110	3.0	328.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	79	2.2	236.0
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	79	2.2	236.0
9	Alzheimer's disease (G30)	72	2.0	215.1
10	Septicemia (A40-A41)	48	1.3	143.4
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	48	1.3	143.4
12	Pneumonitis due to solids and liquids (J69)	47	1.3	140.4
13	Parkinson's disease (G20-G21)	41	1.1	122.5
14	Aortic aneurysm and dissection (I71)	27	0.7	80.6
15	Atherosclerosis (I70)	20	0.5	59.7
...	All other causes (Residual)	420	11.5	1,254.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, all ages				
...	All causes	17,849	100.0	269.7
1	Malignant neoplasms (C00-C97)	4,805	26.9	72.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,460	25.0	67.4
3	Cerebrovascular diseases (I60-I69)	1,931	10.8	29.2
4	Diabetes mellitus (E10-E14)	711	4.0	10.7
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	700	3.9	10.6
6	Influenza and pneumonia (J10-J18)	535	3.0	8.1
7	Chronic lower respiratory diseases (J40-J47)	395	2.2	6.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	329	1.8	5.0
9	Alzheimer's disease (G30)	231	1.3	3.5
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	221	1.2	3.3
11	Septicemia (A40-A41)	193	1.1	2.9
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	192	1.1	2.9
13	Certain conditions originating in the perinatal period (P00-P96)	180	1.0	2.7
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	132	0.7	2.0
15	Chronic liver disease and cirrhosis (K70,K73-K74)	131	0.7	2.0
...	All other causes (Residual)	2,703	15.1	40.8
Asian or Pacific Islander, female, 1-4 years				
...	All causes	69	100.0	19.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	20	29.0	5.7
2	Malignant neoplasms (C00-C97)	7	10.1	*
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7	10.1	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	8.7	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	2.9	*
5	Chronic lower respiratory diseases (J40-J47)	2	2.9	*
7	Shigellosis and amebiasis (A03,A06)	1	1.4	*
7	Septicemia (A40-A41)	1	1.4	*
7	Cerebrovascular diseases (I60-I69)	1	1.4	*
...	All other causes (Residual)	22	31.9	6.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
	Asian or Pacific Islander, female, 5-14 years			
...	All causes	90	100.0	10.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	30	33.3	3.5
2	Malignant neoplasms (C00-C97)	14	15.6	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	5.6	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	4.4	*
5	Septicemia (A40-A41)	3	3.3	*
6	Anemias (D50-D64)	2	2.2	*
6	Chronic lower respiratory diseases (J40-J47)	2	2.2	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	2.2	*
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2	2.2	*
10	Meningococcal infection (A39)	1	1.1	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha'	1	1.1	*
10	Influenza and pneumonia (J10-J18)	1	1.1	*
10	Infections of kidney (N10-N12,N13.6,N15.1)	1	1.1	*
10	Certain conditions originating in the perinatal period (P00-P96)	1	1.1	*
...	All other causes (Residual)	21	23.3	2.4
	Asian or Pacific Islander, female, 15-24 years			
...	All causes	223	100.0	23.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	89	39.9	9.5
2	Malignant neoplasms (C00-C97)	25	11.2	2.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	20	9.0	2.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	19	8.5	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	9	4.0	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	1.8	*
7	Pregnancy, childbirth and the puerperium (O00-099)	3	1.3	*
7	Complications of medical and surgical care (Y40-Y84,Y88)	3	1.3	*
9	Septicemia (A40-A41)	2	0.9	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.9	*
11	Tuberculosis (A16-A19)	1	0.4	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha'	1	0.4	*
11	Anemias (D50-D64)	1	0.4	*
11	Meningitis (G00,G03)	1	0.4	*
11	Influenza and pneumonia (J10-J18)	1	0.4	*
11	Chronic lower respiratory diseases (J40-J47)	1	0.4	*
11	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.4	*
...	All other causes (Residual)	40	17.9	4.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 25-34 years				
...	All causes	328	100.0	26.6
1	Malignant neoplasms (C00-C97)	79	24.1	6.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	72	22.0	5.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	41	12.5	3.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	33	10.1	2.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	20	6.1	1.6
6	Pregnancy, childbirth and the puerperium (O00-099)	12	3.7	*
7	Cerebrovascular diseases (I60-I69)	7	2.1	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	4	1.2	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	4	1.2	*
10	Influenza and pneumonia (J10-J18)	3	0.9	*
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.9	*
12	Anemias (D50-D64)	2	0.6	*
12	Diabetes mellitus (E10-E14)	2	0.6	*
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2	0.6	*
12	Chronic lower respiratory diseases (J40-J47)	2	0.6	*
12	Chronic liver disease and cirrhosis (K70,K73-K74)	2	0.6	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.6	*
...	All other causes (Residual)	38	11.6	3.1
Asian or Pacific Islander, female, 35-44 years				
...	All causes	603	100.0	53.9
1	Malignant neoplasms (C00-C97)	264	43.8	23.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	56	9.3	5.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	50	8.3	4.5
4	Cerebrovascular diseases (I60-I69)	42	7.0	3.8
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	36	6.0	3.2
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	23	3.8	2.1
7	Human immunodeficiency virus (HIV) disease (B20-B24)	9	1.5	*
8	Chronic liver disease and cirrhosis (K70,K73-K74)	7	1.2	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	1.0	*
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	1.0	*
11	Diabetes mellitus (E10-E14)	5	0.8	*
12	Chronic lower respiratory diseases (J40-J47)	4	0.7	*
13	Septicemia (A40-A41)	3	0.5	*
13	Anemias (D50-D64)	3	0.5	*
13	Influenza and pneumonia (J10-J18)	3	0.5	*
13	Pregnancy, childbirth and the puerperium (O00-099)	3	0.5	*
...	All other causes (Residual)	83	13.8	7.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
	Asian or Pacific Islander, female, 45-54 years			
...	All causes	1,373	100.0	149.5
1	Malignant neoplasms (C00-C97)	721	52.5	78.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	151	11.0	16.4
3	Cerebrovascular diseases (I60-I69)	116	8.4	12.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	85	6.2	9.3
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	35	2.5	3.8
6	Diabetes mellitus (E10-E14)	33	2.4	3.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	20	1.5	2.2
8	Septicemia (A40-A41)	18	1.3	*
8	Chronic lower respiratory diseases (J40-J47)	18	1.3	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	1.1	*
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	15	1.1	*
12	Viral hepatitis (B15-B19)	11	0.8	*
13	Influenza and pneumonia (J10-J18)	10	0.7	*
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	6	0.4	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha'	5	0.4	*
15	Aortic aneurysm and dissection (I71)	5	0.4	*
...	All other causes (Residual)	109	7.9	11.9
	Asian or Pacific Islander, female, 55-64 years			
...	All causes	1,968	100.0	372.0
1	Malignant neoplasms (C00-C97)	906	46.0	171.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	327	16.6	61.8
3	Cerebrovascular diseases (I60-I69)	170	8.6	32.1
4	Diabetes mellitus (E10-E14)	78	4.0	14.7
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	61	3.1	11.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	42	2.1	7.9
7	Chronic lower respiratory diseases (J40-J47)	26	1.3	4.9
8	Viral hepatitis (B15-B19)	24	1.2	4.5
9	Septicemia (A40-A41)	23	1.2	4.3
10	Chronic liver disease and cirrhosis (K70,K73-K74)	20	1.0	3.8
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	20	1.0	3.8
12	Influenza and pneumonia (J10-J18)	19	1.0	*
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	11	0.6	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha'	10	0.5	*
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8	0.4	*
...	All other causes (Residual)	223	11.3	42.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
	Asian or Pacific Islander, female, 65-74 years			
...	All causes	3,451	100.0	1,024.7
1	Malignant neoplasms (C00-C97)	1,206	34.9	358.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	808	23.4	239.9
3	Cerebrovascular diseases (I60-I69)	379	11.0	112.5
4	Diabetes mellitus (E10-E14)	180	5.2	53.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	88	2.5	26.1
6	Chronic lower respiratory diseases (J40-J47)	83	2.4	24.6
7	Influenza and pneumonia (J10-J18)	56	1.6	16.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	52	1.5	15.4
9	Viral hepatitis (B15-B19)	47	1.4	14.0
10	Septicemia (A40-A41)	46	1.3	13.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	41	1.2	12.2
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	32	0.9	9.5
13	Parkinson's disease (G20-G21)	22	0.6	6.5
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	20	0.6	5.9
15	Aortic aneurysm and dissection (I71)	18	0.5	*
...	All other causes (Residual)	373	10.8	110.8
	Asian or Pacific Islander, female, 75-84 years			
...	All causes	4,900	100.0	2,713.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,439	29.4	796.9
2	Malignant neoplasms (C00-C97)	1,095	22.3	606.4
3	Cerebrovascular diseases (I60-I69)	599	12.2	331.7
4	Diabetes mellitus (E10-E14)	270	5.5	149.5
5	Influenza and pneumonia (J10-J18)	181	3.7	100.2
6	Chronic lower respiratory diseases (J40-J47)	139	2.8	77.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	124	2.5	68.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	110	2.2	60.9
9	Alzheimer's disease (G30)	76	1.6	42.1
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	67	1.4	37.1
11	Septicemia (A40-A41)	45	0.9	24.9
12	Parkinson's disease (G20-G21)	39	0.8	21.6
13	Aortic aneurysm and dissection (I71)	33	0.7	18.3
13	Chronic liver disease and cirrhosis (K70,K73-K74)	33	0.7	18.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha'	28	0.6	15.5
...	All other causes (Residual)	622	12.7	344.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2002

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
	Asian or Pacific Islander, female, 85 years and over			
...	All causes	4,486	100.0	8,400.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,638	36.5	3,067.4
2	Cerebrovascular diseases (I60-I69)	614	13.7	1,149.8
3	Malignant neoplasms (C00-C97)	486	10.8	910.1
4	Influenza and pneumonia (J10-J18)	259	5.8	485.0
5	Diabetes mellitus (E10-E14)	143	3.2	267.8
6	Alzheimer's disease (G30)	132	2.9	247.2
7	Chronic lower respiratory diseases (J40-J47)	117	2.6	219.1
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	102	2.3	191.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	85	1.9	159.2
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	82	1.8	153.6
11	Pneumonitis due to solids and liquids (J69)	50	1.1	93.6
12	Septicemia (A40-A41)	47	1.0	88.0
12	Atherosclerosis (I70)	47	1.0	88.0
14	Parkinson's disease (G20-G21)	35	0.8	65.5
15	Aortic aneurysm and dissection (I71)	30	0.7	56.2
...	All other causes (Residual)	619	13.8	1,159.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable