

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, all ages				
...	All causes	2,448,017	100.0	825.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	652,091	26.6	220.0
2	Malignant neoplasms (C00-C97)	559,312	22.8	188.7
3	Cerebrovascular diseases (I60-I69)	143,579	5.9	48.4
4	Chronic lower respiratory diseases (J40-J47)	130,933	5.3	44.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	117,809	4.8	39.7
6	Diabetes mellitus (E10-E14)	75,119	3.1	25.3
7	Alzheimer's disease (G30)	71,599	2.9	24.2
8	Influenza and pneumonia (J10-J18)	63,001	2.6	21.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	43,901	1.8	14.8
10	Septicemia (A40-A41)	34,136	1.4	11.5
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	32,637	1.3	11.0
12	Chronic liver disease and cirrhosis (K70,K73-K74)	27,530	1.1	9.3
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	24,902	1.0	8.4
14	Parkinson's disease (G20-G21)	19,544	0.8	6.6
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	18,124	0.7	6.1
...	All other causes (Residual)	433,800	17.7	146.4
All races, both sexes, 1-4 years				
...	All causes	4,756	100.0	29.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,664	35.0	10.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	522	11.0	3.2
3	Malignant neoplasms (C00-C97)	377	7.9	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	375	7.9	2.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	151	3.2	0.9
6	Influenza and pneumonia (J10-J18)	110	2.3	0.7
7	Septicemia (A40-A41)	85	1.8	0.5
8	Cerebrovascular diseases (I60-I69)	62	1.3	0.4
9	Certain conditions originating in the perinatal period (P00-P96)	58	1.2	0.4
10	Chronic lower respiratory diseases (J40-J47)	56	1.2	0.3
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	52	1.1	0.3
12	Meningitis (G00,G03)	32	0.7	0.2
13	Anemias (D50-D64)	23	0.5	0.1
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	16	0.3	*
...	All other causes (Residual)	1,157	24.3	7.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 5-14 years				
...	All causes	6,602	100.0	16.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,415	36.6	6.0
2	Malignant neoplasms (C00-C97)	1,000	15.1	2.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	396	6.0	1.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	341	5.2	0.8
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	272	4.1	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	252	3.8	0.6
7	Influenza and pneumonia (J10-J18)	106	1.6	0.3
8	Chronic lower respiratory diseases (J40-J47)	104	1.6	0.3
9	Cerebrovascular diseases (I60-I69)	95	1.4	0.2
10	Septicemia (A40-A41)	81	1.2	0.2
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	76	1.2	0.2
12	Diabetes mellitus (E10-E14)	38	0.6	0.1
13	Anemias (D50-D64)	26	0.4	0.1
13	Meningitis (G00,G03)	26	0.4	0.1
13	Certain conditions originating in the perinatal period (P00-P96)	26	0.4	0.1
...	All other causes (Residual)	1,348	20.4	3.3
All races, both sexes, 15-24 years				
...	All causes	34,234	100.0	81.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15,753	46.0	37.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5,466	16.0	13.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,212	12.3	10.0
4	Malignant neoplasms (C00-C97)	1,717	5.0	4.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,119	3.3	2.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	504	1.5	1.2
7	Diabetes mellitus (E10-E14)	202	0.6	0.5
8	Cerebrovascular diseases (I60-I69)	196	0.6	0.5
9	Pregnancy, childbirth and the puerperium (O00-099)	183	0.5	0.4
10	Influenza and pneumonia (J10-J18)	172	0.5	0.4
11	Human immunodeficiency virus (HIV) disease (B20-B24)	159	0.5	0.4
12	Septicemia (A40-A41)	148	0.4	0.4
12	Chronic lower respiratory diseases (J40-J47)	148	0.4	0.4
14	Anemias (D50-D64)	103	0.3	0.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	93	0.3	0.2
...	All other causes (Residual)	4,059	11.9	9.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 25-34 years				
...	All causes	41,925	100.0	104.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,997	33.4	34.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,990	11.9	12.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4,752	11.3	11.8
4	Malignant neoplasms (C00-C97)	3,601	8.6	9.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,249	7.7	8.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	1,318	3.1	3.3
7	Diabetes mellitus (E10-E14)	617	1.5	1.5
8	Cerebrovascular diseases (I60-I69)	546	1.3	1.4
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	436	1.0	1.1
10	Influenza and pneumonia (J10-J18)	354	0.8	0.9
11	Pregnancy, childbirth and the puerperium (O00-099)	312	0.7	0.8
12	Septicemia (A40-A41)	311	0.7	0.8
12	Chronic liver disease and cirrhosis (K70,K73-K74)	311	0.7	0.8
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	285	0.7	0.7
15	Chronic lower respiratory diseases (J40-J47)	258	0.6	0.6
...	All other causes (Residual)	6,588	15.7	16.4
All races, both sexes, 35-44 years				
...	All causes	84,785	100.0	193.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	16,919	20.0	38.6
2	Malignant neoplasms (C00-C97)	14,566	17.2	33.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,688	15.0	28.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,550	7.7	14.9
5	Human immunodeficiency virus (HIV) disease (B20-B24)	4,363	5.1	9.9
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,109	3.7	7.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,688	3.2	6.1
8	Cerebrovascular diseases (I60-I69)	2,260	2.7	5.2
9	Diabetes mellitus (E10-E14)	2,045	2.4	4.7
10	Influenza and pneumonia (J10-J18)	934	1.1	2.1
11	Chronic lower respiratory diseases (J40-J47)	890	1.0	2.0
12	Septicemia (A40-A41)	840	1.0	1.9
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	742	0.9	1.7
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	517	0.6	1.2
15	Viral hepatitis (B15-B19)	509	0.6	1.2
...	All other causes (Residual)	15,165	17.9	34.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 45-54 years				
...	All causes	183,530	100.0	432.0
1	Malignant neoplasms (C00-C97)	50,405	27.5	118.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	38,103	20.8	89.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18,339	10.0	43.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	7,517	4.1	17.7
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,991	3.8	16.5
6	Cerebrovascular diseases (I60-I69)	6,381	3.5	15.0
7	Diabetes mellitus (E10-E14)	5,691	3.1	13.4
8	Human immunodeficiency virus (HIV) disease (B20-B24)	4,516	2.5	10.6
9	Chronic lower respiratory diseases (J40-J47)	3,977	2.2	9.4
10	Viral hepatitis (B15-B19)	2,314	1.3	5.4
11	Septicemia (A40-A41)	2,211	1.2	5.2
12	Influenza and pneumonia (J10-J18)	2,183	1.2	5.1
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,060	1.1	4.8
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,028	1.1	4.8
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,166	0.6	2.7
...	All other causes (Residual)	29,648	16.2	69.8
All races, both sexes, 55-64 years				
...	All causes	275,301	100.0	906.9
1	Malignant neoplasms (C00-C97)	99,240	36.0	326.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	65,208	23.7	214.8
3	Chronic lower respiratory diseases (J40-J47)	12,747	4.6	42.0
4	Diabetes mellitus (E10-E14)	11,301	4.1	37.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,853	3.9	35.8
6	Cerebrovascular diseases (I60-I69)	10,028	3.6	33.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	7,126	2.6	23.5
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,210	1.5	13.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,141	1.5	13.6
10	Septicemia (A40-A41)	3,912	1.4	12.9
11	Influenza and pneumonia (J10-J18)	3,422	1.2	11.3
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,954	0.7	6.4
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,612	0.6	5.3
14	Aortic aneurysm and dissection (I71)	1,456	0.5	4.8
15	Viral hepatitis (B15-B19)	1,377	0.5	4.5
...	All other causes (Residual)	36,714	13.3	120.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 65-74 years				
...	All causes	398,355	100.0	2,137.1
1	Malignant neoplasms (C00-C97)	138,446	34.8	742.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	96,729	24.3	518.9
3	Chronic lower respiratory diseases (J40-J47)	29,910	7.5	160.5
4	Cerebrovascular diseases (I60-I69)	18,839	4.7	101.1
5	Diabetes mellitus (E10-E14)	16,183	4.1	86.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,632	2.2	46.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,320	1.8	39.3
8	Influenza and pneumonia (J10-J18)	6,623	1.7	35.5
9	Septicemia (A40-A41)	6,073	1.5	32.6
10	Chronic liver disease and cirrhosis (K70,K73-K74)	5,066	1.3	27.2
11	Alzheimer's disease (G30)	3,813	1.0	20.5
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	3,306	0.8	17.7
13	Aortic aneurysm and dissection (I71)	2,870	0.7	15.4
14	Parkinson's disease (G20-G21)	2,414	0.6	13.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,350	0.6	12.6
...	All other causes (Residual)	49,781	12.5	267.1
All races, both sexes, 75-84 years				
...	All causes	686,665	100.0	5,260.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	190,693	27.8	1,460.8
2	Malignant neoplasms (C00-C97)	166,421	24.2	1,274.8
3	Chronic lower respiratory diseases (J40-J47)	50,333	7.3	385.6
4	Cerebrovascular diseases (I60-I69)	46,859	6.8	359.0
5	Alzheimer's disease (G30)	23,139	3.4	177.3
6	Diabetes mellitus (E10-E14)	23,136	3.4	177.2
7	Influenza and pneumonia (J10-J18)	18,563	2.7	142.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14,403	2.1	110.3
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,854	2.0	106.1
10	Septicemia (A40-A41)	10,626	1.5	81.4
11	Parkinson's disease (G20-G21)	9,294	1.4	71.2
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	7,256	1.1	55.6
13	Pneumonitis due to solids and liquids (J69)	5,558	0.8	42.6
14	Aortic aneurysm and dissection (I71)	4,971	0.7	38.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,594	0.7	35.2
...	All other causes (Residual)	96,965	14.1	742.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 85 years and over				
...	All causes	703,169	100.0	13,798.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	243,504	34.6	4,778.4
2	Malignant neoplasms (C00-C97)	83,455	11.9	1,637.7
3	Cerebrovascular diseases (I60-I69)	58,183	8.3	1,141.8
4	Alzheimer's disease (G30)	43,906	6.2	861.6
5	Chronic lower respiratory diseases (J40-J47)	32,473	4.6	637.2
6	Influenza and pneumonia (J10-J18)	30,267	4.3	593.9
7	Diabetes mellitus (E10-E14)	15,903	2.3	312.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14,693	2.1	288.3
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14,243	2.0	279.5
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	10,703	1.5	210.0
11	Septicemia (A40-A41)	9,544	1.4	187.3
12	Pneumonitis due to solids and liquids (J69)	8,352	1.2	163.9
13	Parkinson's disease (G20-G21)	7,322	1.0	143.7
14	Atherosclerosis (I70)	6,773	1.0	132.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3,909	0.6	76.7
...	All other causes (Residual)	119,939	17.1	2,353.6
All races, male, all ages				
...	All causes	1,207,675	100.0	827.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	322,841	26.7	221.1
2	Malignant neoplasms (C00-C97)	290,422	24.0	198.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	76,375	6.3	52.3
4	Chronic lower respiratory diseases (J40-J47)	62,435	5.2	42.8
5	Cerebrovascular diseases (I60-I69)	56,586	4.7	38.8
6	Diabetes mellitus (E10-E14)	36,538	3.0	25.0
7	Influenza and pneumonia (J10-J18)	28,052	2.3	19.2
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	25,907	2.1	17.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21,268	1.8	14.6
10	Alzheimer's disease (G30)	20,559	1.7	14.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	17,937	1.5	12.3
12	Septicemia (A40-A41)	15,322	1.3	10.5
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	14,376	1.2	9.8
14	Parkinson's disease (G20-G21)	11,247	0.9	7.7
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	9,458	0.8	6.5
...	All other causes (Residual)	198,352	16.4	135.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 1-4 years				
...	All causes	2,765	100.0	33.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,027	37.1	12.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	264	9.5	3.2
3	Malignant neoplasms (C00-C97)	217	7.8	2.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	214	7.7	2.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	81	2.9	1.0
6	Influenza and pneumonia (J10-J18)	56	2.0	0.7
7	Septicemia (A40-A41)	47	1.7	0.6
8	Cerebrovascular diseases (I60-I69)	41	1.5	0.5
9	Certain conditions originating in the perinatal period (P00-P96)	36	1.3	0.4
10	Chronic lower respiratory diseases (J40-J47)	33	1.2	0.4
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	31	1.1	0.4
12	Meningitis (G00,G03)	21	0.8	0.3
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	10	0.4	*
15	Anemias (D50-D64)	9	0.3	*
...	All other causes (Residual)	666	24.1	8.0
All races, male, 5-14 years				
...	All causes	3,853	100.0	18.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,493	38.7	7.2
2	Malignant neoplasms (C00-C97)	557	14.5	2.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	204	5.3	1.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	204	5.3	1.0
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	195	5.1	0.9
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	131	3.4	0.6
7	Chronic lower respiratory diseases (J40-J47)	63	1.6	0.3
8	Cerebrovascular diseases (I60-I69)	55	1.4	0.3
9	Influenza and pneumonia (J10-J18)	53	1.4	0.3
10	Septicemia (A40-A41)	49	1.3	0.2
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	36	0.9	0.2
12	Diabetes mellitus (E10-E14)	15	0.4	*
12	Meningitis (G00,G03)	15	0.4	*
14	Anemias (D50-D64)	14	0.4	*
15	Certain conditions originating in the perinatal period (P00-P96)	11	0.3	*
...	All other causes (Residual)	758	19.7	3.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 15-24 years				
...	All causes	25,509	100.0	117.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,827	46.4	54.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4,765	18.7	22.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,498	13.7	16.2
4	Malignant neoplasms (C00-C97)	1,035	4.1	4.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	770	3.0	3.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	326	1.3	1.5
7	Influenza and pneumonia (J10-J18)	117	0.5	0.5
8	Diabetes mellitus (E10-E14)	115	0.5	0.5
9	Chronic lower respiratory diseases (J40-J47)	95	0.4	0.4
10	Cerebrovascular diseases (I60-I69)	94	0.4	0.4
11	Human immunodeficiency virus (HIV) disease (B20-B24)	89	0.3	0.4
12	Septicemia (A40-A41)	82	0.3	0.4
12	Legal intervention (Y35,Y89.0)	82	0.3	0.4
14	Anemias (D50-D64)	57	0.2	0.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	54	0.2	0.2
...	All other causes (Residual)	2,503	9.8	11.6
All races, male, 25-34 years				
...	All causes	29,283	100.0	143.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,665	36.4	52.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,066	13.9	19.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4,008	13.7	19.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,206	7.5	10.8
5	Malignant neoplasms (C00-C97)	1,805	6.2	8.8
6	Human immunodeficiency virus (HIV) disease (B20-B24)	809	2.8	4.0
7	Diabetes mellitus (E10-E14)	372	1.3	1.8
8	Cerebrovascular diseases (I60-I69)	301	1.0	1.5
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	244	0.8	1.2
10	Influenza and pneumonia (J10-J18)	196	0.7	1.0
10	Chronic liver disease and cirrhosis (K70,K73-K74)	196	0.7	1.0
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	153	0.5	0.7
13	Septicemia (A40-A41)	152	0.5	0.7
14	Chronic lower respiratory diseases (J40-J47)	130	0.4	0.6
15	Legal intervention (Y35,Y89.0)	109	0.4	0.5
...	All other causes (Residual)	3,871	13.2	19.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 35-44 years				
...	All causes	53,309	100.0	243.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,026	22.6	54.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,931	16.8	40.7
3	Malignant neoplasms (C00-C97)	6,338	11.9	28.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,063	9.5	23.1
5	Human immunodeficiency virus (HIV) disease (B20-B24)	3,146	5.9	14.3
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,317	4.3	10.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,786	3.4	8.1
8	Diabetes mellitus (E10-E14)	1,270	2.4	5.8
9	Cerebrovascular diseases (I60-I69)	1,148	2.2	5.2
10	Influenza and pneumonia (J10-J18)	540	1.0	2.5
11	Chronic lower respiratory diseases (J40-J47)	426	0.8	1.9
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	423	0.8	1.9
13	Septicemia (A40-A41)	414	0.8	1.9
14	Viral hepatitis (B15-B19)	349	0.7	1.6
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	272	0.5	1.2
...	All other causes (Residual)	8,860	16.6	40.4
All races, male, 45-54 years				
...	All causes	114,472	100.0	547.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	27,486	24.0	131.5
2	Malignant neoplasms (C00-C97)	25,403	22.2	121.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,002	11.4	62.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	5,423	4.7	26.0
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,269	4.6	25.2
6	Diabetes mellitus (E10-E14)	3,467	3.0	16.6
7	Cerebrovascular diseases (I60-I69)	3,446	3.0	16.5
8	Human immunodeficiency virus (HIV) disease (B20-B24)	3,417	3.0	16.4
9	Chronic lower respiratory diseases (J40-J47)	2,034	1.8	9.7
10	Viral hepatitis (B15-B19)	1,697	1.5	8.1
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,579	1.4	7.6
12	Influenza and pneumonia (J10-J18)	1,299	1.1	6.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,237	1.1	5.9
14	Septicemia (A40-A41)	1,175	1.0	5.6
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	733	0.6	3.5
...	All other causes (Residual)	17,805	15.6	85.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 55-64 years				
...	All causes	165,429	100.0	1,131.0
1	Malignant neoplasms (C00-C97)	54,039	32.7	369.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	44,895	27.1	306.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,507	4.5	51.3
4	Chronic lower respiratory diseases (J40-J47)	6,589	4.0	45.0
5	Diabetes mellitus (E10-E14)	6,570	4.0	44.9
6	Cerebrovascular diseases (I60-I69)	5,636	3.4	38.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	5,094	3.1	34.8
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,250	2.0	22.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,352	1.4	16.1
10	Septicemia (A40-A41)	2,074	1.3	14.2
11	Influenza and pneumonia (J10-J18)	1,988	1.2	13.6
12	Human immunodeficiency virus (HIV) disease (B20-B24)	1,292	0.8	8.8
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,151	0.7	7.9
14	Aortic aneurysm and dissection (I71)	1,063	0.6	7.3
15	Viral hepatitis (B15-B19)	977	0.6	6.7
...	All other causes (Residual)	20,952	12.7	143.2
All races, male, 65-74 years				
...	All causes	222,807	100.0	2,612.2
1	Malignant neoplasms (C00-C97)	76,685	34.4	899.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	59,050	26.5	692.3
3	Chronic lower respiratory diseases (J40-J47)	15,414	6.9	180.7
4	Cerebrovascular diseases (I60-I69)	9,688	4.3	113.6
5	Diabetes mellitus (E10-E14)	8,505	3.8	99.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,343	2.4	62.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,733	1.7	43.8
8	Influenza and pneumonia (J10-J18)	3,648	1.6	42.8
9	Septicemia (A40-A41)	3,095	1.4	36.3
10	Chronic liver disease and cirrhosis (K70,K73-K74)	3,052	1.4	35.8
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,940	0.9	22.7
12	Aortic aneurysm and dissection (I71)	1,911	0.9	22.4
13	Alzheimer's disease (G30)	1,648	0.7	19.3
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,639	0.7	19.2
15	Parkinson's disease (G20-G21)	1,596	0.7	18.7
...	All other causes (Residual)	25,860	11.6	303.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 75-84 years				
...	All causes	335,235	100.0	6,349.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	96,580	28.8	1,829.4
2	Malignant neoplasms (C00-C97)	87,096	26.0	1,649.7
3	Chronic lower respiratory diseases (J40-J47)	24,479	7.3	463.7
4	Cerebrovascular diseases (I60-I69)	19,688	5.9	372.9
5	Diabetes mellitus (E10-E14)	10,701	3.2	202.7
6	Influenza and pneumonia (J10-J18)	9,203	2.7	174.3
7	Alzheimer's disease (G30)	8,326	2.5	157.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,337	2.2	139.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,177	2.1	135.9
10	Parkinson's disease (G20-G21)	5,658	1.7	107.2
11	Septicemia (A40-A41)	4,823	1.4	91.4
12	Pneumonitis due to solids and liquids (J69)	3,057	0.9	57.9
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,917	0.9	55.3
14	Aortic aneurysm and dissection (I71)	2,778	0.8	52.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,419	0.7	45.8
...	All other causes (Residual)	42,996	12.8	814.4
All races, male, 85 years and over				
...	All causes	238,796	100.0	14,889.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	82,489	34.5	5,143.4
2	Malignant neoplasms (C00-C97)	37,197	15.6	2,319.3
3	Cerebrovascular diseases (I60-I69)	16,412	6.9	1,023.3
4	Chronic lower respiratory diseases (J40-J47)	13,150	5.5	819.9
5	Influenza and pneumonia (J10-J18)	10,802	4.5	673.5
6	Alzheimer's disease (G30)	10,262	4.3	639.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,038	2.5	376.5
8	Diabetes mellitus (E10-E14)	5,523	2.3	344.4
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,521	2.3	344.2
10	Pneumonitis due to solids and liquids (J69)	3,677	1.5	229.3
11	Parkinson's disease (G20-G21)	3,666	1.5	228.6
12	Septicemia (A40-A41)	3,238	1.4	201.9
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,716	1.1	169.3
14	Atherosclerosis (I70)	1,952	0.8	121.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,486	0.6	92.7
...	All other causes (Residual)	34,667	14.5	2,161.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, all ages				
...	All causes	1,240,342	100.0	824.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	329,250	26.5	218.9
2	Malignant neoplasms (C00-C97)	268,890	21.7	178.8
3	Cerebrovascular diseases (I60-I69)	86,993	7.0	57.8
4	Chronic lower respiratory diseases (J40-J47)	68,498	5.5	45.5
5	Alzheimer's disease (G30)	51,040	4.1	33.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	41,434	3.3	27.5
7	Diabetes mellitus (E10-E14)	38,581	3.1	25.7
8	Influenza and pneumonia (J10-J18)	34,949	2.8	23.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22,633	1.8	15.0
10	Septicemia (A40-A41)	18,814	1.5	12.5
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	15,444	1.2	10.3
12	Chronic liver disease and cirrhosis (K70,K73-K74)	9,593	0.8	6.4
13	Pneumonitis due to solids and liquids (J69)	8,519	0.7	5.7
14	Parkinson's disease (G20-G21)	8,297	0.7	5.5
15	Atherosclerosis (I70)	7,369	0.6	4.9
...	All other causes (Residual)	230,038	18.5	152.9
All races, female, 1-4 years				
...	All causes	1,991	100.0	25.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	637	32.0	8.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	258	13.0	3.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	161	8.1	2.0
4	Malignant neoplasms (C00-C97)	160	8.0	2.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	70	3.5	0.9
6	Influenza and pneumonia (J10-J18)	54	2.7	0.7
7	Septicemia (A40-A41)	38	1.9	0.5
8	Chronic lower respiratory diseases (J40-J47)	23	1.2	0.3
9	Certain conditions originating in the perinatal period (P00-P96)	22	1.1	0.3
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	21	1.1	0.3
10	Cerebrovascular diseases (I60-I69)	21	1.1	0.3
12	Anemias (D50-D64)	14	0.7	*
13	Meningitis (G00,G03)	11	0.6	*
14	Acute bronchitis and bronchiolitis (J20-J21)	6	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.3	*
...	All other causes (Residual)	489	24.6	6.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 5-14 years				
...	All causes	2,749	100.0	13.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	922	33.5	4.7
2	Malignant neoplasms (C00-C97)	443	16.1	2.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	201	7.3	1.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	137	5.0	0.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	121	4.4	0.6
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	68	2.5	0.3
7	Influenza and pneumonia (J10-J18)	53	1.9	0.3
8	Chronic lower respiratory diseases (J40-J47)	41	1.5	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	40	1.5	0.2
9	Cerebrovascular diseases (I60-I69)	40	1.5	0.2
11	Septicemia (A40-A41)	32	1.2	0.2
12	Diabetes mellitus (E10-E14)	23	0.8	0.1
13	Certain conditions originating in the perinatal period (P00-P96)	15	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	0.5	*
15	Anemias (D50-D64)	12	0.4	*
...	All other causes (Residual)	588	21.4	3.0
All races, female, 15-24 years				
...	All causes	8,725	100.0	42.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,926	45.0	19.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	714	8.2	3.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	701	8.0	3.4
4	Malignant neoplasms (C00-C97)	682	7.8	3.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	349	4.0	1.7
6	Pregnancy, childbirth and the puerperium (O00-099)	183	2.1	0.9
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	178	2.0	0.9
8	Cerebrovascular diseases (I60-I69)	102	1.2	0.5
9	Diabetes mellitus (E10-E14)	87	1.0	0.4
10	Human immunodeficiency virus (HIV) disease (B20-B24)	70	0.8	0.3
11	Septicemia (A40-A41)	66	0.8	0.3
12	Influenza and pneumonia (J10-J18)	55	0.6	0.3
13	Chronic lower respiratory diseases (J40-J47)	53	0.6	0.3
14	Anemias (D50-D64)	46	0.5	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	46	0.5	0.2
...	All other causes (Residual)	1,467	16.8	7.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 25-34 years				
...	All causes	12,642	100.0	64.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,332	26.4	16.9
2	Malignant neoplasms (C00-C97)	1,796	14.2	9.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,043	8.3	5.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	924	7.3	4.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	744	5.9	3.8
6	Human immunodeficiency virus (HIV) disease (B20-B24)	509	4.0	2.6
7	Pregnancy, childbirth and the puerperium (O00-099)	312	2.5	1.6
8	Diabetes mellitus (E10-E14)	245	1.9	1.2
8	Cerebrovascular diseases (I60-I69)	245	1.9	1.2
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	192	1.5	1.0
11	Septicemia (A40-A41)	159	1.3	0.8
12	Influenza and pneumonia (J10-J18)	158	1.2	0.8
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	132	1.0	0.7
14	Chronic lower respiratory diseases (J40-J47)	128	1.0	0.6
15	Chronic liver disease and cirrhosis (K70,K73-K74)	115	0.9	0.6
...	All other causes (Residual)	2,608	20.6	13.2
All races, female, 35-44 years				
...	All causes	31,476	100.0	143.6
1	Malignant neoplasms (C00-C97)	8,228	26.1	37.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,893	15.5	22.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,757	11.9	17.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,487	4.7	6.8
5	Human immunodeficiency virus (HIV) disease (B20-B24)	1,217	3.9	5.6
6	Cerebrovascular diseases (I60-I69)	1,112	3.5	5.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	902	2.9	4.1
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	792	2.5	3.6
9	Diabetes mellitus (E10-E14)	775	2.5	3.5
10	Chronic lower respiratory diseases (J40-J47)	464	1.5	2.1
11	Septicemia (A40-A41)	426	1.4	1.9
12	Influenza and pneumonia (J10-J18)	394	1.3	1.8
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	319	1.0	1.5
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	245	0.8	1.1
15	Pregnancy, childbirth and the puerperium (O00-099)	207	0.7	0.9
...	All other causes (Residual)	6,258	19.9	28.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 45-54 years				
...	All causes	69,058	100.0	319.9
1	Malignant neoplasms (C00-C97)	25,002	36.2	115.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,617	15.4	49.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,337	7.7	24.7
4	Cerebrovascular diseases (I60-I69)	2,935	4.3	13.6
5	Diabetes mellitus (E10-E14)	2,224	3.2	10.3
6	Chronic liver disease and cirrhosis (K70,K73-K74)	2,094	3.0	9.7
7	Chronic lower respiratory diseases (J40-J47)	1,943	2.8	9.0
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,722	2.5	8.0
9	Human immunodeficiency virus (HIV) disease (B20-B24)	1,099	1.6	5.1
10	Septicemia (A40-A41)	1,036	1.5	4.8
11	Influenza and pneumonia (J10-J18)	884	1.3	4.1
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	791	1.1	3.7
13	Viral hepatitis (B15-B19)	617	0.9	2.9
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	481	0.7	2.2
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	433	0.6	2.0
...	All other causes (Residual)	11,843	17.1	54.9
All races, female, 55-64 years				
...	All causes	109,872	100.0	698.5
1	Malignant neoplasms (C00-C97)	45,201	41.1	287.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,313	18.5	129.1
3	Chronic lower respiratory diseases (J40-J47)	6,158	5.6	39.2
4	Diabetes mellitus (E10-E14)	4,731	4.3	30.1
5	Cerebrovascular diseases (I60-I69)	4,392	4.0	27.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,346	3.0	21.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,032	1.8	12.9
8	Septicemia (A40-A41)	1,838	1.7	11.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,789	1.6	11.4
10	Influenza and pneumonia (J10-J18)	1,434	1.3	9.1
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	960	0.9	6.1
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	803	0.7	5.1
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	564	0.5	3.6
14	Viral hepatitis (B15-B19)	400	0.4	2.5
15	Aortic aneurysm and dissection (I71)	393	0.4	2.5
...	All other causes (Residual)	15,518	14.1	98.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 65-74 years				
...	All causes	175,548	100.0	1,736.3
1	Malignant neoplasms (C00-C97)	61,761	35.2	610.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	37,679	21.5	372.7
3	Chronic lower respiratory diseases (J40-J47)	14,496	8.3	143.4
4	Cerebrovascular diseases (I60-I69)	9,151	5.2	90.5
5	Diabetes mellitus (E10-E14)	7,678	4.4	75.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,587	2.0	35.5
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,289	1.9	32.5
8	Septicemia (A40-A41)	2,978	1.7	29.5
9	Influenza and pneumonia (J10-J18)	2,975	1.7	29.4
10	Alzheimer's disease (G30)	2,165	1.2	21.4
11	Chronic liver disease and cirrhosis (K70,K73-K74)	2,014	1.1	19.9
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,667	0.9	16.5
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,008	0.6	10.0
14	Aortic aneurysm and dissection (I71)	959	0.5	9.5
15	Parkinson's disease (G20-G21)	818	0.5	8.1
...	All other causes (Residual)	23,323	13.3	230.7
All races, female, 75-84 years				
...	All causes	351,430	100.0	4,520.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	94,113	26.8	1,210.5
2	Malignant neoplasms (C00-C97)	79,325	22.6	1,020.3
3	Cerebrovascular diseases (I60-I69)	27,171	7.7	349.5
4	Chronic lower respiratory diseases (J40-J47)	25,854	7.4	332.5
5	Alzheimer's disease (G30)	14,813	4.2	190.5
6	Diabetes mellitus (E10-E14)	12,435	3.5	159.9
7	Influenza and pneumonia (J10-J18)	9,360	2.7	120.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,226	2.1	92.9
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,517	1.9	83.8
10	Septicemia (A40-A41)	5,803	1.7	74.6
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	4,339	1.2	55.8
12	Parkinson's disease (G20-G21)	3,636	1.0	46.8
13	Pneumonitis due to solids and liquids (J69)	2,501	0.7	32.2
14	Aortic aneurysm and dissection (I71)	2,193	0.6	28.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,175	0.6	28.0
...	All other causes (Residual)	53,969	15.4	694.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 85 years and over				
...	All causes	464,373	100.0	13,297.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	161,015	34.7	4,610.8
2	Malignant neoplasms (C00-C97)	46,258	10.0	1,324.6
3	Cerebrovascular diseases (I60-I69)	41,771	9.0	1,196.1
4	Alzheimer's disease (G30)	33,644	7.2	963.4
5	Influenza and pneumonia (J10-J18)	19,465	4.2	557.4
6	Chronic lower respiratory diseases (J40-J47)	19,323	4.2	553.3
7	Diabetes mellitus (E10-E14)	10,380	2.2	297.2
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,722	1.9	249.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,655	1.9	247.8
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	7,987	1.7	228.7
11	Septicemia (A40-A41)	6,306	1.4	180.6
12	Atherosclerosis (I70)	4,821	1.0	138.1
13	Pneumonitis due to solids and liquids (J69)	4,675	1.0	133.9
14	Parkinson's disease (G20-G21)	3,656	0.8	104.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,423	0.5	69.4
...	All other causes (Residual)	85,272	18.4	2,441.8
White, both sexes, all ages				
...	All causes	2,098,097	100.0	873.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	564,796	26.9	235.2
2	Malignant neoplasms (C00-C97)	482,132	23.0	200.8
3	Cerebrovascular diseases (I60-I69)	121,868	5.8	50.7
4	Chronic lower respiratory diseases (J40-J47)	120,884	5.8	50.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	100,406	4.8	41.8
6	Alzheimer's disease (G30)	66,191	3.2	27.6
7	Diabetes mellitus (E10-E14)	59,755	2.8	24.9
8	Influenza and pneumonia (J10-J18)	55,540	2.6	23.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	34,806	1.7	14.5
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	29,527	1.4	12.3
11	Septicemia (A40-A41)	27,194	1.3	11.3
12	Chronic liver disease and cirrhosis (K70,K73-K74)	24,003	1.1	10.0
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	19,254	0.9	8.0
14	Parkinson's disease (G20-G21)	18,496	0.9	7.7
15	Pneumonitis due to solids and liquids (J69)	15,338	0.7	6.4
...	All other causes (Residual)	357,907	17.1	149.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 1-4 years				
...	All causes	3,408	100.0	27.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,262	37.0	10.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	385	11.3	3.1
3	Malignant neoplasms (C00-C97)	293	8.6	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	217	6.4	1.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	93	2.7	0.7
6	Influenza and pneumonia (J10-J18)	76	2.2	0.6
7	Septicemia (A40-A41)	59	1.7	0.5
8	Cerebrovascular diseases (I60-I69)	45	1.3	0.4
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	42	1.2	0.3
10	Certain conditions originating in the perinatal period (P00-P96)	35	1.0	0.3
11	Chronic lower respiratory diseases (J40-J47)	24	0.7	0.2
12	Meningitis (G00,G03)	20	0.6	0.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	13	0.4	*
15	Acute bronchitis and bronchiolitis (J20-J21)	12	0.4	*
...	All other causes (Residual)	819	24.0	6.5
White, both sexes, 5-14 years				
...	All causes	4,710	100.0	15.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,785	37.9	5.7
2	Malignant neoplasms (C00-C97)	787	16.7	2.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	285	6.1	0.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	209	4.4	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	168	3.6	0.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	159	3.4	0.5
7	Influenza and pneumonia (J10-J18)	77	1.6	0.2
8	Cerebrovascular diseases (I60-I69)	64	1.4	0.2
9	Septicemia (A40-A41)	57	1.2	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	57	1.2	0.2
11	Chronic lower respiratory diseases (J40-J47)	45	1.0	0.1
12	Diabetes mellitus (E10-E14)	25	0.5	0.1
13	Certain conditions originating in the perinatal period (P00-P96)	22	0.5	0.1
14	Meningitis (G00,G03)	18	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18	0.4	*
...	All other causes (Residual)	934	19.8	3.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 15-24 years				
...	All causes	25,410	100.0	77.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,325	52.4	40.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,540	13.9	10.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,172	8.5	6.6
4	Malignant neoplasms (C00-C97)	1,365	5.4	4.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	736	2.9	2.2
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	386	1.5	1.2
7	Cerebrovascular diseases (I60-I69)	136	0.5	0.4
8	Influenza and pneumonia (J10-J18)	134	0.5	0.4
9	Diabetes mellitus (E10-E14)	124	0.5	0.4
10	Pregnancy, childbirth and the puerperium (O00-099)	110	0.4	0.3
11	Septicemia (A40-A41)	95	0.4	0.3
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	76	0.3	0.2
13	Chronic lower respiratory diseases (J40-J47)	71	0.3	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	60	0.2	0.2
15	Legal intervention (Y35,Y89.0)	59	0.2	0.2
...	All other causes (Residual)	3,021	11.9	9.2
White, both sexes, 25-34 years				
...	All causes	30,103	100.0	95.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,628	38.6	36.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,229	14.0	13.4
3	Malignant neoplasms (C00-C97)	2,694	8.9	8.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,084	6.9	6.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,022	6.7	6.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	464	1.5	1.5
7	Diabetes mellitus (E10-E14)	372	1.2	1.2
8	Cerebrovascular diseases (I60-I69)	356	1.2	1.1
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	339	1.1	1.1
10	Influenza and pneumonia (J10-J18)	262	0.9	0.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	231	0.8	0.7
12	Septicemia (A40-A41)	191	0.6	0.6
13	Pregnancy, childbirth and the puerperium (O00-099)	179	0.6	0.6
14	Chronic lower respiratory diseases (J40-J47)	152	0.5	0.5
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	152	0.5	0.5
...	All other causes (Residual)	4,748	15.8	15.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 35-44 years				
...	All causes	63,584	100.0	180.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14,172	22.3	40.1
2	Malignant neoplasms (C00-C97)	11,352	17.9	32.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,101	14.3	25.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,917	9.3	16.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,284	3.6	6.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	1,958	3.1	5.5
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,615	2.5	4.6
8	Cerebrovascular diseases (I60-I69)	1,450	2.3	4.1
9	Diabetes mellitus (E10-E14)	1,428	2.2	4.0
10	Influenza and pneumonia (J10-J18)	683	1.1	1.9
11	Chronic lower respiratory diseases (J40-J47)	647	1.0	1.8
12	Septicemia (A40-A41)	552	0.9	1.6
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	427	0.7	1.2
14	Viral hepatitis (B15-B19)	421	0.7	1.2
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	401	0.6	1.1
...	All other causes (Residual)	11,176	17.6	31.6
White, both sexes, 45-54 years				
...	All causes	140,224	100.0	399.6
1	Malignant neoplasms (C00-C97)	39,717	28.3	113.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	28,374	20.2	80.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15,135	10.8	43.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,505	4.6	18.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	6,342	4.5	18.1
6	Cerebrovascular diseases (I60-I69)	4,083	2.9	11.6
7	Diabetes mellitus (E10-E14)	4,079	2.9	11.6
8	Chronic lower respiratory diseases (J40-J47)	3,208	2.3	9.1
9	Human immunodeficiency virus (HIV) disease (B20-B24)	1,947	1.4	5.5
10	Viral hepatitis (B15-B19)	1,895	1.4	5.4
11	Influenza and pneumonia (J10-J18)	1,623	1.2	4.6
12	Septicemia (A40-A41)	1,499	1.1	4.3
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,221	0.9	3.5
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,182	0.8	3.4
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	606	0.4	1.7
...	All other causes (Residual)	22,808	16.3	65.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 55-64 years				
...	All causes	222,218	100.0	860.7
1	Malignant neoplasms (C00-C97)	82,439	37.1	319.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	51,407	23.1	199.1
3	Chronic lower respiratory diseases (J40-J47)	11,372	5.1	44.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,032	4.1	35.0
5	Diabetes mellitus (E10-E14)	8,458	3.8	32.8
6	Cerebrovascular diseases (I60-I69)	7,140	3.2	27.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	6,138	2.8	23.8
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,979	1.8	15.4
9	Septicemia (A40-A41)	2,877	1.3	11.1
10	Influenza and pneumonia (J10-J18)	2,700	1.2	10.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,663	1.2	10.3
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,211	0.5	4.7
13	Aortic aneurysm and dissection (I71)	1,209	0.5	4.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,109	0.5	4.3
15	Viral hepatitis (B15-B19)	1,059	0.5	4.1
...	All other causes (Residual)	29,425	13.2	114.0
White, both sexes, 65-74 years				
...	All causes	336,362	100.0	2,094.5
1	Malignant neoplasms (C00-C97)	119,280	35.5	742.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	80,273	23.9	499.9
3	Chronic lower respiratory diseases (J40-J47)	27,517	8.2	171.3
4	Cerebrovascular diseases (I60-I69)	14,829	4.4	92.3
5	Diabetes mellitus (E10-E14)	12,476	3.7	77.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,424	2.2	46.2
7	Influenza and pneumonia (J10-J18)	5,490	1.6	34.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,343	1.6	33.3
9	Septicemia (A40-A41)	4,751	1.4	29.6
10	Chronic liver disease and cirrhosis (K70,K73-K74)	4,581	1.4	28.5
11	Alzheimer's disease (G30)	3,383	1.0	21.1
12	Aortic aneurysm and dissection (I71)	2,539	0.8	15.8
13	Parkinson's disease (G20-G21)	2,224	0.7	13.8
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,222	0.7	13.8
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,215	0.7	13.8
...	All other causes (Residual)	41,815	12.4	260.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 75-84 years				
...	All causes	610,820	100.0	5,263.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	168,711	27.6	1,453.7
2	Malignant neoplasms (C00-C97)	148,732	24.3	1,281.6
3	Chronic lower respiratory diseases (J40-J47)	47,232	7.7	407.0
4	Cerebrovascular diseases (I60-I69)	40,832	6.7	351.8
5	Alzheimer's disease (G30)	21,272	3.5	183.3
6	Diabetes mellitus (E10-E14)	19,212	3.1	165.5
7	Influenza and pneumonia (J10-J18)	16,547	2.7	142.6
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,524	2.1	107.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11,995	2.0	103.4
10	Septicemia (A40-A41)	8,829	1.4	76.1
11	Parkinson's disease (G20-G21)	8,797	1.4	75.8
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	5,805	1.0	50.0
13	Pneumonitis due to solids and liquids (J69)	4,932	0.8	42.5
14	Aortic aneurysm and dissection (I71)	4,532	0.7	39.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,220	0.7	36.4
...	All other causes (Residual)	86,648	14.2	746.6
White, both sexes, 85 years and over				
...	All causes	642,539	100.0	14,021.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	223,603	34.8	4,879.3
2	Malignant neoplasms (C00-C97)	75,404	11.7	1,645.4
3	Cerebrovascular diseases (I60-I69)	52,854	8.2	1,153.3
4	Alzheimer's disease (G30)	40,860	6.4	891.6
5	Chronic lower respiratory diseases (J40-J47)	30,593	4.8	667.6
6	Influenza and pneumonia (J10-J18)	27,788	4.3	606.4
7	Diabetes mellitus (E10-E14)	13,579	2.1	296.3
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,333	2.1	290.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12,844	2.0	280.3
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	9,178	1.4	200.3
11	Septicemia (A40-A41)	8,123	1.3	177.3
12	Pneumonitis due to solids and liquids (J69)	7,630	1.2	166.5
13	Parkinson's disease (G20-G21)	7,006	1.1	152.9
14	Atherosclerosis (I70)	6,356	1.0	138.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3,648	0.6	79.6
...	All other causes (Residual)	109,740	17.1	2,394.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, male, all ages				
...	All causes	1,028,152	100.0	864.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	279,324	27.2	234.9
2	Malignant neoplasms (C00-C97)	250,478	24.4	210.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	64,600	6.3	54.3
4	Chronic lower respiratory diseases (J40-J47)	56,911	5.5	47.9
5	Cerebrovascular diseases (I60-I69)	47,194	4.6	39.7
6	Diabetes mellitus (E10-E14)	29,628	2.9	24.9
7	Influenza and pneumonia (J10-J18)	24,425	2.4	20.5
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	23,478	2.3	19.7
9	Alzheimer's disease (G30)	18,990	1.8	16.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17,137	1.7	14.4
11	Chronic liver disease and cirrhosis (K70,K73-K74)	15,679	1.5	13.2
12	Septicemia (A40-A41)	12,172	1.2	10.2
13	Parkinson's disease (G20-G21)	10,649	1.0	9.0
14	Pneumonitis due to solids and liquids (J69)	7,759	0.8	6.5
15	Aortic aneurysm and dissection (I71)	7,203	0.7	6.1
...	All other causes (Residual)	162,525	15.8	136.7
White, male, 1-4 years				
...	All causes	1,996	100.0	30.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	793	39.7	12.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	201	10.1	3.1
3	Malignant neoplasms (C00-C97)	168	8.4	2.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	119	6.0	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	55	2.8	0.9
6	Influenza and pneumonia (J10-J18)	36	1.8	0.6
7	Septicemia (A40-A41)	33	1.7	0.5
8	Cerebrovascular diseases (I60-I69)	28	1.4	0.4
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	26	1.3	0.4
10	Certain conditions originating in the perinatal period (P00-P96)	20	1.0	0.3
11	Meningitis (G00,G03)	13	0.7	*
12	Chronic lower respiratory diseases (J40-J47)	11	0.6	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.5	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	9	0.5	*
15	Acute bronchitis and bronchiolitis (J20-J21)	8	0.4	*
...	All other causes (Residual)	467	23.4	7.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 5-14 years				
...	All causes	2,749	100.0	17.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,101	40.1	6.8
2	Malignant neoplasms (C00-C97)	429	15.6	2.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	160	5.8	1.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	145	5.3	0.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	96	3.5	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	90	3.3	0.6
7	Influenza and pneumonia (J10-J18)	36	1.3	0.2
8	Cerebrovascular diseases (I60-I69)	35	1.3	0.2
9	Septicemia (A40-A41)	33	1.2	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	29	1.1	0.2
11	Chronic lower respiratory diseases (J40-J47)	26	0.9	0.2
12	Diabetes mellitus (E10-E14)	11	0.4	*
12	Certain conditions originating in the perinatal period (P00-P96)	11	0.4	*
14	Meningitis (G00,G03)	9	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	8	0.3	*
...	All other causes (Residual)	530	19.3	3.3
White, male, 15-24 years				
...	All causes	18,792	100.0	110.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,035	53.4	58.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,945	15.7	17.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,800	9.6	10.6
4	Malignant neoplasms (C00-C97)	826	4.4	4.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	509	2.7	3.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	252	1.3	1.5
7	Influenza and pneumonia (J10-J18)	90	0.5	0.5
8	Diabetes mellitus (E10-E14)	73	0.4	0.4
9	Cerebrovascular diseases (I60-I69)	65	0.3	0.4
10	Legal intervention (Y35,Y89.0)	58	0.3	0.3
11	Septicemia (A40-A41)	56	0.3	0.3
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	47	0.3	0.3
13	Chronic lower respiratory diseases (J40-J47)	46	0.2	0.3
14	Human immunodeficiency virus (HIV) disease (B20-B24)	32	0.2	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	32	0.2	0.2
...	All other causes (Residual)	1,926	10.2	11.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 25-34 years				
...	All causes	21,232	100.0	130.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,876	41.8	54.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,447	16.2	21.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,587	7.5	9.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,469	6.9	9.1
5	Malignant neoplasms (C00-C97)	1,378	6.5	8.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	337	1.6	2.1
7	Diabetes mellitus (E10-E14)	227	1.1	1.4
8	Cerebrovascular diseases (I60-I69)	197	0.9	1.2
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	193	0.9	1.2
10	Chronic liver disease and cirrhosis (K70,K73-K74)	152	0.7	0.9
11	Influenza and pneumonia (J10-J18)	142	0.7	0.9
12	Septicemia (A40-A41)	97	0.5	0.6
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	82	0.4	0.5
14	Chronic lower respiratory diseases (J40-J47)	72	0.3	0.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	69	0.3	0.4
...	All other causes (Residual)	2,907	13.7	17.9
White, male, 35-44 years				
...	All causes	40,824	100.0	228.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,096	24.7	56.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,644	16.3	37.2
3	Malignant neoplasms (C00-C97)	5,072	12.4	28.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,572	11.2	25.6
5	Human immunodeficiency virus (HIV) disease (B20-B24)	1,608	3.9	9.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1,539	3.8	8.6
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,134	2.8	6.3
8	Diabetes mellitus (E10-E14)	912	2.2	5.1
9	Cerebrovascular diseases (I60-I69)	754	1.8	4.2
10	Influenza and pneumonia (J10-J18)	391	1.0	2.2
11	Chronic lower respiratory diseases (J40-J47)	311	0.8	1.7
12	Viral hepatitis (B15-B19)	295	0.7	1.7
13	Septicemia (A40-A41)	276	0.7	1.5
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	238	0.6	1.3
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	220	0.5	1.2
...	All other causes (Residual)	6,762	16.6	37.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 45-54 years				
...	All causes	88,882	100.0	509.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	21,173	23.8	121.3
2	Malignant neoplasms (C00-C97)	20,194	22.7	115.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,714	12.1	61.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,906	5.5	28.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	4,633	5.2	26.5
6	Diabetes mellitus (E10-E14)	2,555	2.9	14.6
7	Cerebrovascular diseases (I60-I69)	2,239	2.5	12.8
8	Chronic lower respiratory diseases (J40-J47)	1,641	1.8	9.4
9	Human immunodeficiency virus (HIV) disease (B20-B24)	1,613	1.8	9.2
10	Viral hepatitis (B15-B19)	1,409	1.6	8.1
11	Influenza and pneumonia (J10-J18)	959	1.1	5.5
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	881	1.0	5.0
13	Septicemia (A40-A41)	800	0.9	4.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	768	0.9	4.4
15	Aortic aneurysm and dissection (I71)	405	0.5	2.3
...	All other causes (Residual)	13,992	15.7	80.2
White, male, 55-64 years				
...	All causes	134,272	100.0	1,068.1
1	Malignant neoplasms (C00-C97)	44,816	33.4	356.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	36,235	27.0	288.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,185	4.6	49.2
4	Chronic lower respiratory diseases (J40-J47)	5,821	4.3	46.3
5	Diabetes mellitus (E10-E14)	5,045	3.8	40.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	4,407	3.3	35.1
7	Cerebrovascular diseases (I60-I69)	3,986	3.0	31.7
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,080	2.3	24.5
9	Influenza and pneumonia (J10-J18)	1,545	1.2	12.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,545	1.2	12.3
11	Septicemia (A40-A41)	1,544	1.1	12.3
12	Aortic aneurysm and dissection (I71)	903	0.7	7.2
13	Viral hepatitis (B15-B19)	760	0.6	6.0
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	727	0.5	5.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	644	0.5	5.1
...	All other causes (Residual)	17,029	12.7	135.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 65-74 years				
...	All causes	189,550	100.0	2,552.7
1	Malignant neoplasms (C00-C97)	66,080	34.9	889.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	49,895	26.3	671.9
3	Chronic lower respiratory diseases (J40-J47)	14,050	7.4	189.2
4	Cerebrovascular diseases (I60-I69)	7,645	4.0	103.0
5	Diabetes mellitus (E10-E14)	6,778	3.6	91.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,586	2.4	61.8
7	Influenza and pneumonia (J10-J18)	3,014	1.6	40.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,867	1.5	38.6
9	Chronic liver disease and cirrhosis (K70,K73-K74)	2,761	1.5	37.2
10	Septicemia (A40-A41)	2,423	1.3	32.6
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,848	1.0	24.9
12	Aortic aneurysm and dissection (I71)	1,723	0.9	23.2
13	Parkinson's disease (G20-G21)	1,472	0.8	19.8
14	Alzheimer's disease (G30)	1,465	0.8	19.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,206	0.6	16.2
...	All other causes (Residual)	21,737	11.5	292.7
White, male, 75-84 years				
...	All causes	300,228	100.0	6,343.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	86,700	28.9	1,831.8
2	Malignant neoplasms (C00-C97)	77,915	26.0	1,646.2
3	Chronic lower respiratory diseases (J40-J47)	22,693	7.6	479.5
4	Cerebrovascular diseases (I60-I69)	17,264	5.8	364.8
5	Diabetes mellitus (E10-E14)	9,170	3.1	193.7
6	Influenza and pneumonia (J10-J18)	8,225	2.7	173.8
7	Alzheimer's disease (G30)	7,669	2.6	162.0
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,627	2.2	140.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,139	2.0	129.7
10	Parkinson's disease (G20-G21)	5,374	1.8	113.5
11	Septicemia (A40-A41)	4,058	1.4	85.7
12	Pneumonitis due to solids and liquids (J69)	2,711	0.9	57.3
13	Aortic aneurysm and dissection (I71)	2,563	0.9	54.2
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,326	0.8	49.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,251	0.7	47.6
...	All other causes (Residual)	38,543	12.8	814.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 85 years and over				
...	All causes	218,990	100.0	15,156.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	76,410	34.9	5,288.4
2	Malignant neoplasms (C00-C97)	33,559	15.3	2,322.7
3	Cerebrovascular diseases (I60-I69)	14,935	6.8	1,033.7
4	Chronic lower respiratory diseases (J40-J47)	12,225	5.6	846.1
5	Influenza and pneumonia (J10-J18)	9,889	4.5	684.4
6	Alzheimer's disease (G30)	9,559	4.4	661.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,399	2.5	373.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,153	2.4	356.6
9	Diabetes mellitus (E10-E14)	4,857	2.2	336.2
10	Parkinson's disease (G20-G21)	3,506	1.6	242.7
11	Pneumonitis due to solids and liquids (J69)	3,384	1.5	234.2
12	Septicemia (A40-A41)	2,756	1.3	190.7
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,314	1.1	160.2
14	Atherosclerosis (I70)	1,854	0.8	128.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,401	0.6	97.0
...	All other causes (Residual)	31,789	14.5	2,200.1
White, female, all ages				
...	All causes	1,069,945	100.0	882.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	285,472	26.7	235.5
2	Malignant neoplasms (C00-C97)	231,654	21.7	191.1
3	Cerebrovascular diseases (I60-I69)	74,674	7.0	61.6
4	Chronic lower respiratory diseases (J40-J47)	63,973	6.0	52.8
5	Alzheimer's disease (G30)	47,201	4.4	38.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	35,806	3.3	29.5
7	Influenza and pneumonia (J10-J18)	31,115	2.9	25.7
8	Diabetes mellitus (E10-E14)	30,127	2.8	24.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17,669	1.7	14.6
10	Septicemia (A40-A41)	15,022	1.4	12.4
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	12,236	1.1	10.1
12	Chronic liver disease and cirrhosis (K70,K73-K74)	8,324	0.8	6.9
13	Parkinson's disease (G20-G21)	7,847	0.7	6.5
14	Pneumonitis due to solids and liquids (J69)	7,579	0.7	6.3
15	Atherosclerosis (I70)	6,752	0.6	5.6
...	All other causes (Residual)	194,494	18.2	160.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 1-4 years				
...	All causes	1,412	100.0	22.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	469	33.2	7.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	184	13.0	3.0
3	Malignant neoplasms (C00-C97)	125	8.9	2.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	98	6.9	1.6
5	Influenza and pneumonia (J10-J18)	40	2.8	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	38	2.7	0.6
7	Septicemia (A40-A41)	26	1.8	0.4
8	Cerebrovascular diseases (I60-I69)	17	1.2	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	1.1	*
10	Certain conditions originating in the perinatal period (P00-P96)	15	1.1	*
11	Chronic lower respiratory diseases (J40-J47)	13	0.9	*
12	Anemias (D50-D64)	7	0.5	*
12	Meningitis (G00,G03)	7	0.5	*
14	Meningococcal infection (A39)	5	0.4	*
15	Acute bronchitis and bronchiolitis (J20-J21)	4	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.3	*
...	All other causes (Residual)	340	24.1	5.5
White, female, 5-14 years				
...	All causes	1,961	100.0	12.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	684	34.9	4.5
2	Malignant neoplasms (C00-C97)	358	18.3	2.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	140	7.1	0.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	72	3.7	0.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	69	3.5	0.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	49	2.5	0.3
7	Influenza and pneumonia (J10-J18)	41	2.1	0.3
8	Cerebrovascular diseases (I60-I69)	29	1.5	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	28	1.4	0.2
10	Septicemia (A40-A41)	24	1.2	0.2
11	Chronic lower respiratory diseases (J40-J47)	19	1.0	*
12	Diabetes mellitus (E10-E14)	14	0.7	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	0.6	*
13	Certain conditions originating in the perinatal period (P00-P96)	11	0.6	*
15	Meningitis (G00,G03)	9	0.5	*
...	All other causes (Residual)	403	20.6	2.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 15-24 years				
...	All causes	6,618	100.0	41.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,290	49.7	20.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	595	9.0	3.7
3	Malignant neoplasms (C00-C97)	539	8.1	3.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	372	5.6	2.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	227	3.4	1.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	134	2.0	0.8
7	Pregnancy, childbirth and the puerperium (O00-099)	110	1.7	0.7
8	Cerebrovascular diseases (I60-I69)	71	1.1	0.4
9	Diabetes mellitus (E10-E14)	51	0.8	0.3
10	Influenza and pneumonia (J10-J18)	44	0.7	0.3
11	Septicemia (A40-A41)	39	0.6	0.2
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	29	0.4	0.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	28	0.4	0.2
14	Chronic lower respiratory diseases (J40-J47)	25	0.4	0.2
15	Human immunodeficiency virus (HIV) disease (B20-B24)	18	0.3	*
...	All other causes (Residual)	1,046	15.8	6.6
White, female, 25-34 years				
...	All causes	8,871	100.0	58.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,752	31.0	18.0
2	Malignant neoplasms (C00-C97)	1,316	14.8	8.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	782	8.8	5.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	615	6.9	4.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	435	4.9	2.8
6	Pregnancy, childbirth and the puerperium (O00-099)	179	2.0	1.2
7	Cerebrovascular diseases (I60-I69)	159	1.8	1.0
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	146	1.6	1.0
9	Diabetes mellitus (E10-E14)	145	1.6	0.9
10	Human immunodeficiency virus (HIV) disease (B20-B24)	127	1.4	0.8
11	Influenza and pneumonia (J10-J18)	120	1.4	0.8
12	Septicemia (A40-A41)	94	1.1	0.6
13	Chronic lower respiratory diseases (J40-J47)	80	0.9	0.5
14	Chronic liver disease and cirrhosis (K70,K73-K74)	79	0.9	0.5
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	70	0.8	0.5
...	All other causes (Residual)	1,772	20.0	11.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 35-44 years				
...	All causes	22,760	100.0	130.4
1	Malignant neoplasms (C00-C97)	6,280	27.6	36.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,076	17.9	23.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,457	10.8	14.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,345	5.9	7.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	745	3.3	4.3
6	Cerebrovascular diseases (I60-I69)	696	3.1	4.0
7	Diabetes mellitus (E10-E14)	516	2.3	3.0
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	481	2.1	2.8
9	Human immunodeficiency virus (HIV) disease (B20-B24)	350	1.5	2.0
10	Chronic lower respiratory diseases (J40-J47)	336	1.5	1.9
11	Influenza and pneumonia (J10-J18)	292	1.3	1.7
12	Septicemia (A40-A41)	276	1.2	1.6
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	207	0.9	1.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	163	0.7	0.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	136	0.6	0.8
...	All other causes (Residual)	4,404	19.3	25.2
White, female, 45-54 years				
...	All causes	51,342	100.0	291.1
1	Malignant neoplasms (C00-C97)	19,523	38.0	110.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,201	14.0	40.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,421	8.6	25.1
4	Cerebrovascular diseases (I60-I69)	1,844	3.6	10.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,709	3.3	9.7
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,599	3.1	9.1
7	Chronic lower respiratory diseases (J40-J47)	1,567	3.1	8.9
8	Diabetes mellitus (E10-E14)	1,524	3.0	8.6
9	Septicemia (A40-A41)	699	1.4	4.0
10	Influenza and pneumonia (J10-J18)	664	1.3	3.8
11	Viral hepatitis (B15-B19)	486	0.9	2.8
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	453	0.9	2.6
13	Human immunodeficiency virus (HIV) disease (B20-B24)	334	0.7	1.9
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	301	0.6	1.7
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	294	0.6	1.7
...	All other causes (Residual)	8,723	17.0	49.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 55-64 years				
...	All causes	87,946	100.0	663.9
1	Malignant neoplasms (C00-C97)	37,623	42.8	284.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,172	17.3	114.5
3	Chronic lower respiratory diseases (J40-J47)	5,551	6.3	41.9
4	Diabetes mellitus (E10-E14)	3,413	3.9	25.8
5	Cerebrovascular diseases (I60-I69)	3,154	3.6	23.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,847	3.2	21.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,731	2.0	13.1
8	Septicemia (A40-A41)	1,333	1.5	10.1
9	Influenza and pneumonia (J10-J18)	1,155	1.3	8.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,118	1.3	8.4
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	899	1.0	6.8
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	484	0.6	3.7
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	465	0.5	3.5
14	Alzheimer's disease (G30)	327	0.4	2.5
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	312	0.4	2.4
...	All other causes (Residual)	12,362	14.1	93.3
White, female, 65-74 years				
...	All causes	146,812	100.0	1,700.4
1	Malignant neoplasms (C00-C97)	53,200	36.2	616.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	30,378	20.7	351.8
3	Chronic lower respiratory diseases (J40-J47)	13,467	9.2	156.0
4	Cerebrovascular diseases (I60-I69)	7,184	4.9	83.2
5	Diabetes mellitus (E10-E14)	5,698	3.9	66.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,838	1.9	32.9
7	Influenza and pneumonia (J10-J18)	2,476	1.7	28.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,476	1.7	28.7
9	Septicemia (A40-A41)	2,328	1.6	27.0
10	Alzheimer's disease (G30)	1,918	1.3	22.2
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1,820	1.2	21.1
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,121	0.8	13.0
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	861	0.6	10.0
14	Aortic aneurysm and dissection (I71)	816	0.6	9.5
15	Parkinson's disease (G20-G21)	752	0.5	8.7
...	All other causes (Residual)	19,479	13.3	225.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 75-84 years				
...	All causes	310,592	100.0	4,519.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	82,011	26.4	1,193.3
2	Malignant neoplasms (C00-C97)	70,817	22.8	1,030.5
3	Chronic lower respiratory diseases (J40-J47)	24,539	7.9	357.1
4	Cerebrovascular diseases (I60-I69)	23,568	7.6	342.9
5	Alzheimer's disease (G30)	13,603	4.4	197.9
6	Diabetes mellitus (E10-E14)	10,042	3.2	146.1
7	Influenza and pneumonia (J10-J18)	8,322	2.7	121.1
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,897	1.9	85.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,856	1.9	85.2
10	Septicemia (A40-A41)	4,771	1.5	69.4
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	3,479	1.1	50.6
12	Parkinson's disease (G20-G21)	3,423	1.1	49.8
13	Pneumonitis due to solids and liquids (J69)	2,221	0.7	32.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,969	0.6	28.7
14	Aortic aneurysm and dissection (I71)	1,969	0.6	28.7
...	All other causes (Residual)	48,105	15.5	700.0
White, female, 85 years and over				
...	All causes	423,549	100.0	13,498.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	147,193	34.8	4,691.0
2	Malignant neoplasms (C00-C97)	41,845	9.9	1,333.6
3	Cerebrovascular diseases (I60-I69)	37,919	9.0	1,208.5
4	Alzheimer's disease (G30)	31,301	7.4	997.5
5	Chronic lower respiratory diseases (J40-J47)	18,368	4.3	585.4
6	Influenza and pneumonia (J10-J18)	17,899	4.2	570.4
7	Diabetes mellitus (E10-E14)	8,722	2.1	278.0
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,180	1.9	260.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,445	1.8	237.3
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	6,864	1.6	218.8
11	Septicemia (A40-A41)	5,367	1.3	171.0
12	Atherosclerosis (I70)	4,502	1.1	143.5
13	Pneumonitis due to solids and liquids (J69)	4,246	1.0	135.3
14	Parkinson's disease (G20-G21)	3,500	0.8	111.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,247	0.5	71.6
...	All other causes (Residual)	77,951	18.4	2,484.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, all ages				
...	All causes	292,808	100.0	749.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	74,159	25.3	189.8
2	Malignant neoplasms (C00-C97)	63,165	21.6	161.7
3	Cerebrovascular diseases (I60-I69)	17,541	6.0	44.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,652	4.7	34.9
5	Diabetes mellitus (E10-E14)	12,970	4.4	33.2
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8,669	3.0	22.2
7	Chronic lower respiratory diseases (J40-J47)	8,229	2.8	21.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,075	2.8	20.7
9	Human immunodeficiency virus (HIV) disease (B20-B24)	7,022	2.4	18.0
10	Septicemia (A40-A41)	6,221	2.1	15.9
11	Influenza and pneumonia (J10-J18)	5,780	2.0	14.8
12	Certain conditions originating in the perinatal period (P00-P96)	5,050	1.7	12.9
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	4,953	1.7	12.7
14	Alzheimer's disease (G30)	4,620	1.6	11.8
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,530	0.9	6.5
...	All other causes (Residual)	50,172	17.1	128.4
Black, both sexes, 1-4 years				
...	All causes	1,097	100.0	41.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	327	29.8	12.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	140	12.8	5.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	106	9.7	4.0
4	Malignant neoplasms (C00-C97)	64	5.8	2.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	45	4.1	1.7
6	Chronic lower respiratory diseases (J40-J47)	31	2.8	1.2
7	Influenza and pneumonia (J10-J18)	29	2.6	1.1
8	Certain conditions originating in the perinatal period (P00-P96)	23	2.1	0.9
9	Septicemia (A40-A41)	20	1.8	0.8
10	Cerebrovascular diseases (I60-I69)	14	1.3	*
11	Anemias (D50-D64)	11	1.0	*
12	Meningitis (G00,G03)	6	0.5	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.5	*
13	Pneumonitis due to solids and liquids (J69)	5	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.3	*
...	All other causes (Residual)	268	24.4	10.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 5-14 years				
...	All causes	1,543	100.0	23.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	501	32.5	7.6
2	Malignant neoplasms (C00-C97)	172	11.1	2.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	150	9.7	2.3
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	88	5.7	1.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	77	5.0	1.2
6	Chronic lower respiratory diseases (J40-J47)	53	3.4	0.8
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	41	2.7	0.6
8	Cerebrovascular diseases (I60-I69)	24	1.6	0.4
9	Influenza and pneumonia (J10-J18)	22	1.4	0.3
10	Septicemia (A40-A41)	21	1.4	0.3
11	Anemias (D50-D64)	15	1.0	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	0.9	*
13	Diabetes mellitus (E10-E14)	12	0.8	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	9	0.6	*
15	Meningitis (G00,G03)	7	0.5	*
...	All other causes (Residual)	337	21.8	5.1
Black, both sexes, 15-24 years				
...	All causes	7,371	100.0	112.3
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,077	41.7	46.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,780	24.1	27.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	437	5.9	6.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	334	4.5	5.1
5	Malignant neoplasms (C00-C97)	264	3.6	4.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	109	1.5	1.7
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	100	1.4	1.5
8	Anemias (D50-D64)	77	1.0	1.2
9	Chronic lower respiratory diseases (J40-J47)	75	1.0	1.1
10	Diabetes mellitus (E10-E14)	73	1.0	1.1
11	Pregnancy, childbirth and the puerperium (O00-099)	69	0.9	1.1
12	Cerebrovascular diseases (I60-I69)	51	0.7	0.8
13	Septicemia (A40-A41)	44	0.6	0.7
14	Influenza and pneumonia (J10-J18)	31	0.4	0.5
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	29	0.4	0.4
...	All other causes (Residual)	821	11.1	12.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 25-34 years				
...	All causes	10,036	100.0	179.2
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,591	25.8	46.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,796	17.9	32.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,022	10.2	18.3
4	Human immunodeficiency virus (HIV) disease (B20-B24)	836	8.3	14.9
5	Malignant neoplasms (C00-C97)	688	6.9	12.3
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	485	4.8	8.7
7	Diabetes mellitus (E10-E14)	219	2.2	3.9
8	Cerebrovascular diseases (I60-I69)	163	1.6	2.9
9	Anemias (D50-D64)	131	1.3	2.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	121	1.2	2.2
11	Pregnancy, childbirth and the puerperium (O00-099)	117	1.2	2.1
12	Septicemia (A40-A41)	108	1.1	1.9
13	Chronic lower respiratory diseases (J40-J47)	97	1.0	1.7
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	81	0.8	1.4
15	Influenza and pneumonia (J10-J18)	78	0.8	1.4
...	All other causes (Residual)	1,503	15.0	26.8
Black, both sexes, 35-44 years				
...	All causes	18,186	100.0	318.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,189	17.5	55.8
2	Malignant neoplasms (C00-C97)	2,560	14.1	44.8
3	Human immunodeficiency virus (HIV) disease (B20-B24)	2,337	12.9	40.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,208	12.1	38.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,369	7.5	24.0
6	Cerebrovascular diseases (I60-I69)	689	3.8	12.1
7	Diabetes mellitus (E10-E14)	541	3.0	9.5
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	419	2.3	7.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	313	1.7	5.5
10	Septicemia (A40-A41)	262	1.4	4.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	239	1.3	4.2
12	Chronic lower respiratory diseases (J40-J47)	219	1.2	3.8
13	Influenza and pneumonia (J10-J18)	217	1.2	3.8
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	197	1.1	3.4
15	Anemias (D50-D64)	147	0.8	2.6
...	All other causes (Residual)	3,280	18.0	57.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 45-54 years				
...	All causes	37,749	100.0	744.2
1	Malignant neoplasms (C00-C97)	8,901	23.6	175.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,716	23.1	171.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,689	7.1	53.0
4	Human immunodeficiency virus (HIV) disease (B20-B24)	2,520	6.7	49.7
5	Cerebrovascular diseases (I60-I69)	2,004	5.3	39.5
6	Diabetes mellitus (E10-E14)	1,410	3.7	27.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	892	2.4	17.6
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	797	2.1	15.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	754	2.0	14.9
10	Chronic lower respiratory diseases (J40-J47)	703	1.9	13.9
11	Septicemia (A40-A41)	658	1.7	13.0
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	532	1.4	10.5
13	Influenza and pneumonia (J10-J18)	499	1.3	9.8
14	Viral hepatitis (B15-B19)	355	0.9	7.0
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	314	0.8	6.2
...	All other causes (Residual)	6,005	15.9	118.4
Black, both sexes, 55-64 years				
...	All causes	45,416	100.0	1,483.3
1	Malignant neoplasms (C00-C97)	13,960	30.7	455.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,115	26.7	395.7
3	Diabetes mellitus (E10-E14)	2,440	5.4	79.7
4	Cerebrovascular diseases (I60-I69)	2,430	5.4	79.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,471	3.2	48.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,346	3.0	44.0
7	Chronic lower respiratory diseases (J40-J47)	1,207	2.7	39.4
8	Septicemia (A40-A41)	942	2.1	30.8
9	Human immunodeficiency virus (HIV) disease (B20-B24)	911	2.0	29.8
10	Chronic liver disease and cirrhosis (K70,K73-K74)	763	1.7	24.9
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	691	1.5	22.6
12	Influenza and pneumonia (J10-J18)	611	1.3	20.0
13	Viral hepatitis (B15-B19)	257	0.6	8.4
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	248	0.5	8.1
15	Aortic aneurysm and dissection (I71)	213	0.5	7.0
...	All other causes (Residual)	5,811	12.8	189.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 65-74 years				
...	All causes	51,718	100.0	2,928.1
1	Malignant neoplasms (C00-C97)	15,719	30.4	889.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,005	27.1	792.9
3	Cerebrovascular diseases (I60-I69)	3,235	6.3	183.2
4	Diabetes mellitus (E10-E14)	3,128	6.0	177.1
5	Chronic lower respiratory diseases (J40-J47)	2,020	3.9	114.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,756	3.4	99.4
7	Septicemia (A40-A41)	1,183	2.3	67.0
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	958	1.9	54.2
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	927	1.8	52.5
10	Influenza and pneumonia (J10-J18)	918	1.8	52.0
11	Alzheimer's disease (G30)	390	0.8	22.1
12	Chronic liver disease and cirrhosis (K70,K73-K74)	357	0.7	20.2
13	Aortic aneurysm and dissection (I71)	255	0.5	14.4
14	Human immunodeficiency virus (HIV) disease (B20-B24)	246	0.5	13.9
15	Pneumonitis due to solids and liquids (J69)	235	0.5	13.3
...	All other causes (Residual)	6,386	12.3	361.5
Black, both sexes, 75-84 years				
...	All causes	61,998	100.0	6,134.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	18,351	29.6	1,815.7
2	Malignant neoplasms (C00-C97)	14,295	23.1	1,414.4
3	Cerebrovascular diseases (I60-I69)	4,746	7.7	469.6
4	Diabetes mellitus (E10-E14)	3,251	5.2	321.7
5	Chronic lower respiratory diseases (J40-J47)	2,439	3.9	241.3
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,103	3.4	208.1
7	Alzheimer's disease (G30)	1,596	2.6	157.9
8	Septicemia (A40-A41)	1,590	2.6	157.3
9	Influenza and pneumonia (J10-J18)	1,514	2.4	149.8
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,234	2.0	122.1
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	980	1.6	97.0
12	Pneumonitis due to solids and liquids (J69)	530	0.9	52.4
13	Parkinson's disease (G20-G21)	342	0.6	33.8
14	Aortic aneurysm and dissection (I71)	312	0.5	30.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	282	0.5	27.9
...	All other causes (Residual)	8,433	13.6	834.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 85 years and over				
...	All causes	48,952	100.0	13,083.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	16,189	33.1	4,326.7
2	Malignant neoplasms (C00-C97)	6,528	13.3	1,744.7
3	Cerebrovascular diseases (I60-I69)	4,141	8.5	1,106.7
4	Alzheimer's disease (G30)	2,580	5.3	689.5
5	Diabetes mellitus (E10-E14)	1,895	3.9	506.5
6	Influenza and pneumonia (J10-J18)	1,763	3.6	471.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,588	3.2	424.4
8	Chronic lower respiratory diseases (J40-J47)	1,372	2.8	366.7
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,276	2.6	341.0
10	Septicemia (A40-A41)	1,260	2.6	336.8
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	654	1.3	174.8
12	Pneumonitis due to solids and liquids (J69)	573	1.2	153.1
13	Atherosclerosis (I70)	366	0.7	97.8
14	Parkinson's disease (G20-G21)	212	0.4	56.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	204	0.4	54.5
...	All other causes (Residual)	8,351	17.1	2,231.9
Black, male, all ages				
...	All causes	149,108	100.0	799.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	36,343	24.4	194.8
2	Malignant neoplasms (C00-C97)	32,726	21.9	175.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,329	6.3	50.0
4	Cerebrovascular diseases (I60-I69)	7,519	5.0	40.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7,412	5.0	39.7
6	Diabetes mellitus (E10-E14)	5,730	3.8	30.7
7	Human immunodeficiency virus (HIV) disease (B20-B24)	4,684	3.1	25.1
8	Chronic lower respiratory diseases (J40-J47)	4,464	3.0	23.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,645	2.4	19.5
10	Certain conditions originating in the perinatal period (P00-P96)	2,828	1.9	15.2
11	Septicemia (A40-A41)	2,786	1.9	14.9
12	Influenza and pneumonia (J10-J18)	2,729	1.8	14.6
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,128	1.4	11.4
14	Chronic liver disease and cirrhosis (K70,K73-K74)	1,656	1.1	8.9
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,621	1.1	8.7
...	All other causes (Residual)	23,508	15.8	126.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 1-4 years				
...	All causes	623	100.0	46.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	184	29.5	13.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	88	14.1	6.6
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	48	7.7	3.6
4	Malignant neoplasms (C00-C97)	37	5.9	2.8
5	Chronic lower respiratory diseases (J40-J47)	22	3.5	1.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	19	3.0	*
7	Influenza and pneumonia (J10-J18)	17	2.7	*
8	Certain conditions originating in the perinatal period (P00-P96)	16	2.6	*
9	Septicemia (A40-A41)	13	2.1	*
10	Cerebrovascular diseases (I60-I69)	10	1.6	*
11	Anemias (D50-D64)	5	0.8	*
12	Meningitis (G00,G03)	3	0.5	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.5	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.3	*
14	Pneumonitis due to solids and liquids (J69)	2	0.3	*
...	All other causes (Residual)	154	24.7	11.5
Black, male, 5-14 years				
...	All causes	909	100.0	27.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	314	34.5	9.3
2	Malignant neoplasms (C00-C97)	103	11.3	3.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	97	10.7	2.9
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	41	4.5	1.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	36	4.0	1.1
6	Chronic lower respiratory diseases (J40-J47)	34	3.7	1.0
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	27	3.0	0.8
8	Cerebrovascular diseases (I60-I69)	15	1.7	*
9	Septicemia (A40-A41)	14	1.5	*
10	Influenza and pneumonia (J10-J18)	13	1.4	*
11	Anemias (D50-D64)	9	1.0	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	6	0.7	*
12	Meningitis (G00,G03)	6	0.7	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.4	*
15	Diabetes mellitus (E10-E14)	3	0.3	*
15	Diseases of appendix (K35-K38)	3	0.3	*
...	All other causes (Residual)	184	20.2	5.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 15-24 years				
...	All causes	5,709	100.0	172.1
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,790	48.9	84.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,334	23.4	40.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	382	6.7	11.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	229	4.0	6.9
5	Malignant neoplasms (C00-C97)	155	2.7	4.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	64	1.1	1.9
7	Human immunodeficiency virus (HIV) disease (B20-B24)	57	1.0	1.7
8	Chronic lower respiratory diseases (J40-J47)	48	0.8	1.4
9	Anemias (D50-D64)	45	0.8	1.4
10	Diabetes mellitus (E10-E14)	39	0.7	1.2
11	Septicemia (A40-A41)	24	0.4	0.7
11	Cerebrovascular diseases (I60-I69)	24	0.4	0.7
11	Legal intervention (Y35,Y89.0)	24	0.4	0.7
14	Influenza and pneumonia (J10-J18)	23	0.4	0.7
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	0.2	*
...	All other causes (Residual)	460	8.1	13.9
Black, male, 25-34 years				
...	All causes	6,843	100.0	254.3
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,319	33.9	86.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,369	20.0	50.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	633	9.3	23.5
4	Human immunodeficiency virus (HIV) disease (B20-B24)	458	6.7	17.0
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	402	5.9	14.9
6	Malignant neoplasms (C00-C97)	321	4.7	11.9
7	Diabetes mellitus (E10-E14)	128	1.9	4.8
8	Cerebrovascular diseases (I60-I69)	89	1.3	3.3
9	Anemias (D50-D64)	69	1.0	2.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	61	0.9	2.3
11	Chronic lower respiratory diseases (J40-J47)	53	0.8	2.0
12	Septicemia (A40-A41)	50	0.7	1.9
13	Influenza and pneumonia (J10-J18)	45	0.7	1.7
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	43	0.6	1.6
15	Legal intervention (Y35,Y89.0)	39	0.6	1.4
...	All other causes (Residual)	764	11.2	28.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 35-44 years				
...	All causes	10,626	100.0	395.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,989	18.7	74.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,557	14.7	57.9
3	Human immunodeficiency virus (HIV) disease (B20-B24)	1,489	14.0	55.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,091	10.3	40.6
5	Malignant neoplasms (C00-C97)	973	9.2	36.2
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	337	3.2	12.5
7	Cerebrovascular diseases (I60-I69)	332	3.1	12.4
8	Diabetes mellitus (E10-E14)	310	2.9	11.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	165	1.6	6.1
10	Chronic liver disease and cirrhosis (K70,K73-K74)	144	1.4	5.4
11	Influenza and pneumonia (J10-J18)	129	1.2	4.8
12	Septicemia (A40-A41)	122	1.1	4.5
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	104	1.0	3.9
14	Chronic lower respiratory diseases (J40-J47)	103	1.0	3.8
15	Aortic aneurysm and dissection (I71)	81	0.8	3.0
...	All other causes (Residual)	1,700	16.0	63.3
Black, male, 45-54 years				
...	All causes	22,248	100.0	948.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,569	25.0	237.4
2	Malignant neoplasms (C00-C97)	4,365	19.6	186.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,936	8.7	82.5
4	Human immunodeficiency virus (HIV) disease (B20-B24)	1,760	7.9	75.0
5	Cerebrovascular diseases (I60-I69)	1,050	4.7	44.8
6	Diabetes mellitus (E10-E14)	780	3.5	33.3
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	633	2.8	27.0
8	Chronic liver disease and cirrhosis (K70,K73-K74)	603	2.7	25.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	443	2.0	18.9
10	Chronic lower respiratory diseases (J40-J47)	354	1.6	15.1
11	Septicemia (A40-A41)	346	1.6	14.8
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	310	1.4	13.2
13	Influenza and pneumonia (J10-J18)	302	1.4	12.9
14	Viral hepatitis (B15-B19)	240	1.1	10.2
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	240	1.1	10.2
...	All other causes (Residual)	3,317	14.9	141.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 55-64 years				
...	All causes	26,711	100.0	1,954.3
1	Malignant neoplasms (C00-C97)	7,767	29.1	568.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,505	28.1	549.1
3	Cerebrovascular diseases (I60-I69)	1,417	5.3	103.7
4	Diabetes mellitus (E10-E14)	1,287	4.8	94.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,094	4.1	80.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	750	2.8	54.9
7	Human immunodeficiency virus (HIV) disease (B20-B24)	696	2.6	50.9
8	Chronic lower respiratory diseases (J40-J47)	669	2.5	48.9
9	Chronic liver disease and cirrhosis (K70,K73-K74)	536	2.0	39.2
10	Septicemia (A40-A41)	488	1.8	35.7
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	400	1.5	29.3
12	Influenza and pneumonia (J10-J18)	377	1.4	27.6
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	196	0.7	14.3
14	Viral hepatitis (B15-B19)	175	0.7	12.8
15	Aortic aneurysm and dissection (I71)	138	0.5	10.1
...	All other causes (Residual)	3,216	12.0	235.3
Black, male, 65-74 years				
...	All causes	27,616	100.0	3,747.3
1	Malignant neoplasms (C00-C97)	8,724	31.6	1,183.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,676	27.8	1,041.6
3	Cerebrovascular diseases (I60-I69)	1,653	6.0	224.3
4	Diabetes mellitus (E10-E14)	1,430	5.2	194.0
5	Chronic lower respiratory diseases (J40-J47)	1,159	4.2	157.3
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	756	2.7	102.6
7	Septicemia (A40-A41)	594	2.2	80.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	594	2.2	80.6
9	Influenza and pneumonia (J10-J18)	512	1.9	69.5
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	472	1.7	64.0
11	Chronic liver disease and cirrhosis (K70,K73-K74)	230	0.8	31.2
12	Human immunodeficiency virus (HIV) disease (B20-B24)	184	0.7	25.0
13	Alzheimer's disease (G30)	163	0.6	22.1
14	Pneumonitis due to solids and liquids (J69)	146	0.5	19.8
15	Aortic aneurysm and dissection (I71)	133	0.5	18.0
...	All other causes (Residual)	3,190	11.6	432.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 75-84 years				
...	All causes	28,069	100.0	7,667.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,069	28.7	2,204.1
2	Malignant neoplasms (C00-C97)	7,386	26.3	2,017.5
3	Cerebrovascular diseases (I60-I69)	1,844	6.6	503.7
4	Chronic lower respiratory diseases (J40-J47)	1,375	4.9	375.6
5	Diabetes mellitus (E10-E14)	1,253	4.5	342.3
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	899	3.2	245.6
7	Influenza and pneumonia (J10-J18)	692	2.5	189.0
8	Septicemia (A40-A41)	661	2.4	180.6
9	Alzheimer's disease (G30)	549	2.0	150.0
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	530	1.9	144.8
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	497	1.8	135.8
12	Pneumonitis due to solids and liquids (J69)	290	1.0	79.2
13	Parkinson's disease (G20-G21)	198	0.7	54.1
14	Aortic aneurysm and dissection (I71)	143	0.5	39.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	124	0.4	33.9
...	All other causes (Residual)	3,559	12.7	972.1
Black, male, 85 years and over				
...	All causes	14,856	100.0	13,809.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,551	30.6	4,230.5
2	Malignant neoplasms (C00-C97)	2,887	19.4	2,683.7
3	Cerebrovascular diseases (I60-I69)	1,058	7.1	983.5
4	Chronic lower respiratory diseases (J40-J47)	640	4.3	594.9
5	Influenza and pneumonia (J10-J18)	572	3.9	531.7
6	Alzheimer's disease (G30)	558	3.8	518.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	519	3.5	482.4
8	Diabetes mellitus (E10-E14)	500	3.4	464.8
9	Septicemia (A40-A41)	403	2.7	374.6
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	311	2.1	289.1
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	242	1.6	225.0
12	Pneumonitis due to solids and liquids (J69)	217	1.5	201.7
13	Parkinson's disease (G20-G21)	105	0.7	97.6
14	Atherosclerosis (I70)	81	0.5	75.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	59	0.4	54.8
...	All other causes (Residual)	2,153	14.5	2,001.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, all ages				
...	All causes	143,700	100.0	703.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	37,816	26.3	185.2
2	Malignant neoplasms (C00-C97)	30,439	21.2	149.1
3	Cerebrovascular diseases (I60-I69)	10,022	7.0	49.1
4	Diabetes mellitus (E10-E14)	7,240	5.0	35.5
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,430	3.1	21.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,323	3.0	21.2
7	Chronic lower respiratory diseases (J40-J47)	3,765	2.6	18.4
8	Septicemia (A40-A41)	3,435	2.4	16.8
9	Alzheimer's disease (G30)	3,328	2.3	16.3
10	Influenza and pneumonia (J10-J18)	3,051	2.1	14.9
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,825	2.0	13.8
12	Human immunodeficiency virus (HIV) disease (B20-B24)	2,338	1.6	11.5
13	Certain conditions originating in the perinatal period (P00-P96)	2,222	1.5	10.9
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,257	0.9	6.2
15	Chronic liver disease and cirrhosis (K70,K73-K74)	874	0.6	4.3
...	All other causes (Residual)	26,335	18.3	129.0
Black, female, 1-4 years				
...	All causes	474	100.0	36.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	143	30.2	11.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	58	12.2	4.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	52	11.0	4.0
4	Malignant neoplasms (C00-C97)	27	5.7	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	26	5.5	2.0
6	Influenza and pneumonia (J10-J18)	12	2.5	*
7	Chronic lower respiratory diseases (J40-J47)	9	1.9	*
8	Septicemia (A40-A41)	7	1.5	*
8	Certain conditions originating in the perinatal period (P00-P96)	7	1.5	*
10	Anemias (D50-D64)	6	1.3	*
11	Cerebrovascular diseases (I60-I69)	4	0.8	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.6	*
12	Meningitis (G00,G03)	3	0.6	*
12	Pneumonitis due to solids and liquids (J69)	3	0.6	*
15	Acute bronchitis and bronchiolitis (J20-J21)	2	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
...	All other causes (Residual)	110	23.2	8.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 5-14 years				
...	All causes	634	100.0	19.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	187	29.5	5.7
2	Malignant neoplasms (C00-C97)	69	10.9	2.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	53	8.4	1.6
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	47	7.4	1.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	41	6.5	1.3
6	Chronic lower respiratory diseases (J40-J47)	19	3.0	*
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	14	2.2	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	1.6	*
9	Diabetes mellitus (E10-E14)	9	1.4	*
9	Cerebrovascular diseases (I60-I69)	9	1.4	*
9	Influenza and pneumonia (J10-J18)	9	1.4	*
12	Septicemia (A40-A41)	7	1.1	*
13	Anemias (D50-D64)	6	0.9	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	3	0.5	*
14	Certain conditions originating in the perinatal period (P00-P96)	3	0.5	*
...	All other causes (Residual)	148	23.3	4.5
Black, female, 15-24 years				
...	All causes	1,662	100.0	51.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	446	26.8	13.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	287	17.3	8.8
3	Malignant neoplasms (C00-C97)	109	6.6	3.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	105	6.3	3.2
5	Pregnancy, childbirth and the puerperium (O00-099)	69	4.2	2.1
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	55	3.3	1.7
7	Human immunodeficiency virus (HIV) disease (B20-B24)	52	3.1	1.6
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	36	2.2	1.1
9	Diabetes mellitus (E10-E14)	34	2.0	1.0
10	Anemias (D50-D64)	32	1.9	1.0
11	Cerebrovascular diseases (I60-I69)	27	1.6	0.8
11	Chronic lower respiratory diseases (J40-J47)	27	1.6	0.8
13	Septicemia (A40-A41)	20	1.2	0.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18	1.1	*
15	Influenza and pneumonia (J10-J18)	8	0.5	*
...	All other causes (Residual)	337	20.3	10.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 25-34 years				
...	All causes	3,193	100.0	109.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	427	13.4	14.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	389	12.2	13.4
3	Human immunodeficiency virus (HIV) disease (B20-B24)	378	11.8	13.0
4	Malignant neoplasms (C00-C97)	367	11.5	12.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	272	8.5	9.3
6	Pregnancy, childbirth and the puerperium (000-099)	117	3.7	4.0
7	Diabetes mellitus (E10-E14)	91	2.8	3.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	83	2.6	2.9
9	Cerebrovascular diseases (I60-I69)	74	2.3	2.5
10	Anemias (D50-D64)	62	1.9	2.1
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	60	1.9	2.1
12	Septicemia (A40-A41)	58	1.8	2.0
13	Chronic lower respiratory diseases (J40-J47)	44	1.4	1.5
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	38	1.2	1.3
15	Influenza and pneumonia (J10-J18)	33	1.0	1.1
...	All other causes (Residual)	700	21.9	24.1
Black, female, 35-44 years				
...	All causes	7,560	100.0	250.0
1	Malignant neoplasms (C00-C97)	1,587	21.0	52.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,200	15.9	39.7
3	Human immunodeficiency virus (HIV) disease (B20-B24)	848	11.2	28.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	651	8.6	21.5
5	Cerebrovascular diseases (I60-I69)	357	4.7	11.8
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	278	3.7	9.2
7	Diabetes mellitus (E10-E14)	231	3.1	7.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	148	2.0	4.9
9	Septicemia (A40-A41)	140	1.9	4.6
10	Chronic lower respiratory diseases (J40-J47)	116	1.5	3.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	95	1.3	3.1
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	93	1.2	3.1
13	Influenza and pneumonia (J10-J18)	88	1.2	2.9
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	82	1.1	2.7
15	Anemias (D50-D64)	68	0.9	2.2
...	All other causes (Residual)	1,578	20.9	52.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 45-54 years				
...	All causes	15,501	100.0	568.4
1	Malignant neoplasms (C00-C97)	4,536	29.3	166.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,147	20.3	115.4
3	Cerebrovascular diseases (I60-I69)	954	6.2	35.0
4	Human immunodeficiency virus (HIV) disease (B20-B24)	760	4.9	27.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	753	4.9	27.6
6	Diabetes mellitus (E10-E14)	630	4.1	23.1
7	Chronic lower respiratory diseases (J40-J47)	349	2.3	12.8
8	Septicemia (A40-A41)	312	2.0	11.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	311	2.0	11.4
10	Chronic liver disease and cirrhosis (K70,K73-K74)	289	1.9	10.6
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	222	1.4	8.1
12	Influenza and pneumonia (J10-J18)	197	1.3	7.2
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	164	1.1	6.0
14	Viral hepatitis (B15-B19)	115	0.7	4.2
15	Anemias (D50-D64)	78	0.5	2.9
...	All other causes (Residual)	2,684	17.3	98.4
Black, female, 55-64 years				
...	All causes	18,705	100.0	1,103.6
1	Malignant neoplasms (C00-C97)	6,193	33.1	365.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,610	24.6	272.0
3	Diabetes mellitus (E10-E14)	1,153	6.2	68.0
4	Cerebrovascular diseases (I60-I69)	1,013	5.4	59.8
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	596	3.2	35.2
6	Chronic lower respiratory diseases (J40-J47)	538	2.9	31.7
7	Septicemia (A40-A41)	454	2.4	26.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	377	2.0	22.2
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	291	1.6	17.2
10	Influenza and pneumonia (J10-J18)	234	1.3	13.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	227	1.2	13.4
12	Human immunodeficiency virus (HIV) disease (B20-B24)	215	1.1	12.7
13	Viral hepatitis (B15-B19)	82	0.4	4.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	80	0.4	4.7
15	Aortic aneurysm and dissection (I71)	75	0.4	4.4
...	All other causes (Residual)	2,567	13.7	151.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 65-74 years				
...	All causes	24,102	100.0	2,341.5
1	Malignant neoplasms (C00-C97)	6,995	29.0	679.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,329	26.3	614.9
3	Diabetes mellitus (E10-E14)	1,698	7.0	165.0
4	Cerebrovascular diseases (I60-I69)	1,582	6.6	153.7
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,000	4.1	97.1
6	Chronic lower respiratory diseases (J40-J47)	861	3.6	83.6
7	Septicemia (A40-A41)	589	2.4	57.2
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	486	2.0	47.2
9	Influenza and pneumonia (J10-J18)	406	1.7	39.4
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	333	1.4	32.4
11	Alzheimer's disease (G30)	227	0.9	22.1
12	Chronic liver disease and cirrhosis (K70,K73-K74)	127	0.5	12.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	123	0.5	11.9
14	Aortic aneurysm and dissection (I71)	122	0.5	11.9
15	Pneumonitis due to solids and liquids (J69)	89	0.4	8.6
...	All other causes (Residual)	3,135	13.0	304.6
Black, female, 75-84 years				
...	All causes	33,929	100.0	5,263.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,282	30.3	1,595.1
2	Malignant neoplasms (C00-C97)	6,909	20.4	1,071.9
3	Cerebrovascular diseases (I60-I69)	2,902	8.6	450.2
4	Diabetes mellitus (E10-E14)	1,998	5.9	310.0
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,204	3.5	186.8
6	Chronic lower respiratory diseases (J40-J47)	1,064	3.1	165.1
7	Alzheimer's disease (G30)	1,047	3.1	162.4
8	Septicemia (A40-A41)	929	2.7	144.1
9	Influenza and pneumonia (J10-J18)	822	2.4	127.5
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	737	2.2	114.3
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	450	1.3	69.8
12	Pneumonitis due to solids and liquids (J69)	240	0.7	37.2
13	Aortic aneurysm and dissection (I71)	169	0.5	26.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	158	0.5	24.5
15	Parkinson's disease (G20-G21)	144	0.4	22.3
...	All other causes (Residual)	4,874	14.4	756.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 85 years and over				
...	All causes	34,096	100.0	12,789.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,638	34.1	4,365.6
2	Malignant neoplasms (C00-C97)	3,641	10.7	1,365.8
3	Cerebrovascular diseases (I60-I69)	3,083	9.0	1,156.5
4	Alzheimer's disease (G30)	2,022	5.9	758.5
5	Diabetes mellitus (E10-E14)	1,395	4.1	523.3
6	Influenza and pneumonia (J10-J18)	1,191	3.5	446.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,069	3.1	401.0
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	965	2.8	362.0
9	Septicemia (A40-A41)	857	2.5	321.5
10	Chronic lower respiratory diseases (J40-J47)	732	2.1	274.6
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	412	1.2	154.5
12	Pneumonitis due to solids and liquids (J69)	356	1.0	133.5
13	Atherosclerosis (I70)	285	0.8	106.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	145	0.4	54.4
15	Aortic aneurysm and dissection (I71)	133	0.4	49.9
...	All other causes (Residual)	6,172	18.1	2,315.2
American Indian, both sexes, all ages				
...	All causes	13,918	100.0	440.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,659	19.1	84.1
2	Malignant neoplasms (C00-C97)	2,465	17.7	78.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,626	11.7	51.4
4	Diabetes mellitus (E10-E14)	818	5.9	25.9
5	Cerebrovascular diseases (I60-I69)	627	4.5	19.8
6	Chronic liver disease and cirrhosis (K70,K73-K74)	596	4.3	18.9
7	Chronic lower respiratory diseases (J40-J47)	518	3.7	16.4
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	392	2.8	12.4
9	Influenza and pneumonia (J10-J18)	354	2.5	11.2
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	265	1.9	8.4
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	257	1.8	8.1
12	Septicemia (A40-A41)	217	1.6	6.9
13	Alzheimer's disease (G30)	173	1.2	5.5
14	Certain conditions originating in the perinatal period (P00-P96)	130	0.9	4.1
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	117	0.8	3.7
...	All other causes (Residual)	2,704	19.4	85.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 1-4 years				
...	All causes	100	100.0	59.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	44	44.0	26.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	11	11.0	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	10	10.0	*
4	Malignant neoplasms (C00-C97)	4	4.0	*
5	Meningitis (G00,G03)	3	3.0	*
6	Septicemia (A40-A41)	1	1.0	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	1.0	*
6	Influenza and pneumonia (J10-J18)	1	1.0	*
6	Chronic lower respiratory diseases (J40-J47)	1	1.0	*
...	All other causes (Residual)	24	24.0	14.2
American Indian, both sexes, 5-14 years				
...	All causes	110	100.0	19.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	55	50.0	9.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	14	12.7	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7	6.4	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	5.5	*
5	Malignant neoplasms (C00-C97)	4	3.6	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	1.8	*
6	Chronic lower respiratory diseases (J40-J47)	2	1.8	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.9	*
8	Meningitis (G00,G03)	1	0.9	*
8	Influenza and pneumonia (J10-J18)	1	0.9	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.9	*
...	All other causes (Residual)	16	14.5	*
American Indian, both sexes, 15-24 years				
...	All causes	635	100.0	107.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	321	50.6	54.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	128	20.2	21.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	82	12.9	13.9
4	Malignant neoplasms (C00-C97)	15	2.4	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	13	2.0	*
6	Diabetes mellitus (E10-E14)	4	0.6	*
7	Septicemia (A40-A41)	3	0.5	*
7	Chronic liver disease and cirrhosis (K70,K73-K74)	3	0.5	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.3	*
9	Cerebrovascular diseases (I60-I69)	2	0.3	*
9	Pregnancy, childbirth and the puerperium (O00-099)	2	0.3	*
12	Salmonella infections (A01-A02)	1	0.2	*
12	Anemias (D50-D64)	1	0.2	*
12	Influenza and pneumonia (J10-J18)	1	0.2	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.2	*
...	All other causes (Residual)	55	8.7	9.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 25-34 years				
...	All causes	720	100.0	150.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	324	45.0	67.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	91	12.6	19.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	60	8.3	12.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	46	6.4	9.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	36	5.0	7.5
6	Malignant neoplasms (C00-C97)	29	4.0	6.1
7	Diabetes mellitus (E10-E14)	17	2.4	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	10	1.4	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.8	*
10	Influenza and pneumonia (J10-J18)	5	0.7	*
11	Septicemia (A40-A41)	4	0.6	*
11	Cerebrovascular diseases (I60-I69)	4	0.6	*
13	Chronic lower respiratory diseases (J40-J47)	3	0.4	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.4	*
15	Legal intervention (Y35,Y89.0)	2	0.3	*
...	All other causes (Residual)	80	11.1	16.8
American Indian, both sexes, 35-44 years				
...	All causes	1,223	100.0	265.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	305	24.9	66.2
2	Chronic liver disease and cirrhosis (K70,K73-K74)	125	10.2	27.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	120	9.8	26.1
4	Malignant neoplasms (C00-C97)	116	9.5	25.2
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	84	6.9	18.2
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	45	3.7	9.8
7	Diabetes mellitus (E10-E14)	41	3.4	8.9
8	Human immunodeficiency virus (HIV) disease (B20-B24)	35	2.9	7.6
9	Cerebrovascular diseases (I60-I69)	32	2.6	6.9
10	Influenza and pneumonia (J10-J18)	18	1.5	*
11	Septicemia (A40-A41)	16	1.3	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	1.2	*
13	Chronic lower respiratory diseases (J40-J47)	12	1.0	*
14	Viral hepatitis (B15-B19)	10	0.8	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	0.7	*
...	All other causes (Residual)	241	19.7	52.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 45-54 years				
...	All causes	1,927	100.0	474.2
1	Malignant neoplasms (C00-C97)	340	17.6	83.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	330	17.1	81.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	258	13.4	63.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	196	10.2	48.2
5	Diabetes mellitus (E10-E14)	98	5.1	24.1
6	Cerebrovascular diseases (I60-I69)	64	3.3	15.7
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	53	2.8	13.0
8	Chronic lower respiratory diseases (J40-J47)	33	1.7	8.1
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	29	1.5	7.1
10	Influenza and pneumonia (J10-J18)	27	1.4	6.6
11	Human immunodeficiency virus (HIV) disease (B20-B24)	26	1.3	6.4
12	Septicemia (A40-A41)	24	1.2	5.9
13	Viral hepatitis (B15-B19)	23	1.2	5.7
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20	1.0	4.9
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	10	0.5	*
...	All other causes (Residual)	396	20.6	97.4
American Indian, both sexes, 55-64 years				
...	All causes	2,240	100.0	903.8
1	Malignant neoplasms (C00-C97)	580	25.9	234.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	485	21.7	195.7
3	Diabetes mellitus (E10-E14)	177	7.9	71.4
4	Chronic liver disease and cirrhosis (K70,K73-K74)	143	6.4	57.7
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	135	6.0	54.5
6	Chronic lower respiratory diseases (J40-J47)	91	4.1	36.7
7	Cerebrovascular diseases (I60-I69)	89	4.0	35.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	47	2.1	19.0
9	Septicemia (A40-A41)	45	2.0	18.2
10	Influenza and pneumonia (J10-J18)	43	1.9	17.4
11	Viral hepatitis (B15-B19)	14	0.6	*
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	12	0.5	*
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	11	0.5	*
14	Atherosclerosis (I70)	10	0.4	*
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	7	0.3	*
...	All other causes (Residual)	351	15.7	141.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 65-74 years				
...	All causes	2,487	100.0	1,999.6
1	Malignant neoplasms (C00-C97)	676	27.2	543.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	552	22.2	443.8
3	Diabetes mellitus (E10-E14)	234	9.4	188.1
4	Cerebrovascular diseases (I60-I69)	142	5.7	114.2
4	Chronic lower respiratory diseases (J40-J47)	142	5.7	114.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	75	3.0	60.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	65	2.6	52.3
8	Chronic liver disease and cirrhosis (K70,K73-K74)	61	2.5	49.0
9	Influenza and pneumonia (J10-J18)	50	2.0	40.2
10	Septicemia (A40-A41)	47	1.9	37.8
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	28	1.1	22.5
12	Pneumonitis due to solids and liquids (J69)	15	0.6	*
13	Aortic aneurysm and dissection (I71)	13	0.5	*
14	Atherosclerosis (I70)	12	0.5	*
15	Parkinson's disease (G20-G21)	10	0.4	*
15	Alzheimer's disease (G30)	10	0.4	*
...	All other causes (Residual)	355	14.3	285.4
American Indian, both sexes, 75-84 years				
...	All causes	2,433	100.0	3,928.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	597	24.5	963.9
2	Malignant neoplasms (C00-C97)	518	21.3	836.3
3	Chronic lower respiratory diseases (J40-J47)	164	6.7	264.8
4	Cerebrovascular diseases (I60-I69)	163	6.7	263.2
5	Diabetes mellitus (E10-E14)	159	6.5	256.7
6	Influenza and pneumonia (J10-J18)	98	4.0	158.2
7	Alzheimer's disease (G30)	69	2.8	111.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	62	2.5	100.1
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	60	2.5	96.9
10	Septicemia (A40-A41)	45	1.8	72.7
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	38	1.6	61.4
12	Parkinson's disease (G20-G21)	31	1.3	50.0
13	Chronic liver disease and cirrhosis (K70,K73-K74)	27	1.1	43.6
14	Pneumonitis due to solids and liquids (J69)	17	0.7	*
15	Atherosclerosis (I70)	16	0.7	*
...	All other causes (Residual)	369	15.2	595.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 85 years and over				
...	All causes	1,684	100.0	7,505.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	508	30.2	2,264.0
2	Malignant neoplasms (C00-C97)	183	10.9	815.6
3	Cerebrovascular diseases (I60-I69)	129	7.7	574.9
4	Influenza and pneumonia (J10-J18)	105	6.2	468.0
5	Alzheimer's disease (G30)	94	5.6	418.9
6	Diabetes mellitus (E10-E14)	88	5.2	392.2
7	Chronic lower respiratory diseases (J40-J47)	69	4.1	307.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	46	2.7	205.0
9	Septicemia (A40-A41)	30	1.8	133.7
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	30	1.8	133.7
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	27	1.6	120.3
12	Pneumonitis due to solids and liquids (J69)	21	1.2	93.6
13	Parkinson's disease (G20-G21)	18	1.1	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	0.8	*
15	Atherosclerosis (I70)	10	0.6	*
...	All other causes (Residual)	313	18.6	1,395.0
American Indian, male, all ages				
...	All causes	7,607	100.0	481.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,471	19.3	93.2
2	Malignant neoplasms (C00-C97)	1,297	17.1	82.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,118	14.7	70.8
4	Diabetes mellitus (E10-E14)	412	5.4	26.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	348	4.6	22.0
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	313	4.1	19.8
7	Chronic lower respiratory diseases (J40-J47)	257	3.4	16.3
8	Cerebrovascular diseases (I60-I69)	249	3.3	15.8
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	193	2.5	12.2
10	Influenza and pneumonia (J10-J18)	180	2.4	11.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	105	1.4	6.7
12	Septicemia (A40-A41)	104	1.4	6.6
13	Certain conditions originating in the perinatal period (P00-P96)	75	1.0	4.8
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	61	0.8	3.9
15	Human immunodeficiency virus (HIV) disease (B20-B24)	59	0.8	3.7
...	All other causes (Residual)	1,365	17.9	86.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 1-4 years				
...	All causes	62	100.0	72.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	31	50.0	36.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	6.5	*
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	6.5	*
4	Malignant neoplasms (C00-C97)	3	4.8	*
5	Meningitis (G00,G03)	2	3.2	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	1.6	*
...	All other causes (Residual)	17	27.4	*
American Indian, male, 5-14 years				
...	All causes	64	100.0	22.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	36	56.3	12.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11	17.2	*
3	Malignant neoplasms (C00-C97)	4	6.3	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	4.7	*
5	Human immunodeficiency virus (HIV) disease (B20-B24)	1	1.6	*
5	Influenza and pneumonia (J10-J18)	1	1.6	*
5	Chronic lower respiratory diseases (J40-J47)	1	1.6	*
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	1.6	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	1.6	*
...	All other causes (Residual)	5	7.8	*
American Indian, male, 15-24 years				
...	All causes	439	100.0	145.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	223	50.8	73.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	99	22.6	32.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	67	15.3	22.1
4	Malignant neoplasms (C00-C97)	10	2.3	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	7	1.6	*
6	Diabetes mellitus (E10-E14)	2	0.5	*
6	Chronic liver disease and cirrhosis (K70,K73-K74)	2	0.5	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.2	*
8	Cerebrovascular diseases (I60-I69)	1	0.2	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
...	All other causes (Residual)	26	5.9	8.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 25-34 years				
...	All causes	512	100.0	206.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	240	46.9	96.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	73	14.3	29.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	45	8.8	18.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	33	6.4	13.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	22	4.3	8.9
6	Malignant neoplasms (C00-C97)	16	3.1	*
7	Diabetes mellitus (E10-E14)	11	2.1	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	7	1.4	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	1.0	*
10	Influenza and pneumonia (J10-J18)	4	0.8	*
11	Cerebrovascular diseases (I60-I69)	2	0.4	*
11	Legal intervention (Y35,Y89.0)	2	0.4	*
13	Tuberculosis (A16-A19)	1	0.2	*
13	Septicemia (A40-A41)	1	0.2	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.2	*
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1	0.2	*
13	Aortic aneurysm and dissection (I71)	1	0.2	*
13	Chronic lower respiratory diseases (J40-J47)	1	0.2	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.2	*
...	All other causes (Residual)	45	8.8	18.1
American Indian, male, 35-44 years				
...	All causes	777	100.0	336.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	213	27.4	92.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	84	10.8	36.4
3	Chronic liver disease and cirrhosis (K70,K73-K74)	70	9.0	30.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	68	8.8	29.5
5	Malignant neoplasms (C00-C97)	62	8.0	26.9
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	32	4.1	13.9
7	Diabetes mellitus (E10-E14)	27	3.5	11.7
8	Human immunodeficiency virus (HIV) disease (B20-B24)	22	2.8	9.5
9	Cerebrovascular diseases (I60-I69)	16	2.1	*
10	Influenza and pneumonia (J10-J18)	12	1.5	*
11	Septicemia (A40-A41)	10	1.3	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	1.2	*
13	Viral hepatitis (B15-B19)	7	0.9	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	0.5	*
14	Legal intervention (Y35,Y89.0)	4	0.5	*
...	All other causes (Residual)	137	17.6	59.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 45-54 years				
...	All causes	1,160	100.0	588.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	221	19.1	112.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	181	15.6	91.9
3	Malignant neoplasms (C00-C97)	161	13.9	81.7
4	Chronic liver disease and cirrhosis (K70,K73-K74)	116	10.0	58.9
5	Diabetes mellitus (E10-E14)	63	5.4	32.0
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	42	3.6	21.3
7	Cerebrovascular diseases (I60-I69)	27	2.3	13.7
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	25	2.2	12.7
9	Human immunodeficiency virus (HIV) disease (B20-B24)	24	2.1	12.2
10	Viral hepatitis (B15-B19)	18	1.6	*
10	Chronic lower respiratory diseases (J40-J47)	18	1.6	*
12	Influenza and pneumonia (J10-J18)	16	1.4	*
13	Septicemia (A40-A41)	12	1.0	*
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	9	0.8	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.5	*
...	All other causes (Residual)	221	19.1	112.2
American Indian, male, 55-64 years				
...	All causes	1,341	100.0	1,124.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	328	24.5	275.0
2	Malignant neoplasms (C00-C97)	321	23.9	269.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	99	7.4	83.0
4	Chronic liver disease and cirrhosis (K70,K73-K74)	95	7.1	79.6
5	Diabetes mellitus (E10-E14)	94	7.0	78.8
6	Chronic lower respiratory diseases (J40-J47)	45	3.4	37.7
7	Cerebrovascular diseases (I60-I69)	43	3.2	36.0
8	Influenza and pneumonia (J10-J18)	24	1.8	20.1
9	Septicemia (A40-A41)	22	1.6	18.4
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	1.3	*
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11	0.8	*
12	Viral hepatitis (B15-B19)	10	0.7	*
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	9	0.7	*
14	Atherosclerosis (I70)	5	0.4	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.3	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.3	*
15	Anemias (D50-D64)	4	0.3	*
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	4	0.3	*
15	Pneumonitis due to solids and liquids (J69)	4	0.3	*
...	All other causes (Residual)	198	14.8	166.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 65-74 years				
...	All causes	1,297	100.0	2,254.1
1	Malignant neoplasms (C00-C97)	358	27.6	622.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	319	24.6	554.4
3	Diabetes mellitus (E10-E14)	115	8.9	199.9
4	Cerebrovascular diseases (I60-I69)	65	5.0	113.0
4	Chronic lower respiratory diseases (J40-J47)	65	5.0	113.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	45	3.5	78.2
7	Septicemia (A40-A41)	29	2.2	50.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	29	2.2	50.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	28	2.2	48.7
10	Influenza and pneumonia (J10-J18)	24	1.9	41.7
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	16	1.2	*
12	Aortic aneurysm and dissection (I71)	10	0.8	*
13	Pneumonitis due to solids and liquids (J69)	8	0.6	*
14	Alzheimer's disease (G30)	6	0.5	*
14	Atherosclerosis (I70)	6	0.5	*
...	All other causes (Residual)	174	13.4	302.4
American Indian, male, 75-84 years				
...	All causes	1,144	100.0	4,373.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	294	25.7	1,123.9
2	Malignant neoplasms (C00-C97)	267	23.3	1,020.7
3	Chronic lower respiratory diseases (J40-J47)	92	8.0	351.7
4	Diabetes mellitus (E10-E14)	69	6.0	263.8
5	Cerebrovascular diseases (I60-I69)	60	5.2	229.4
6	Influenza and pneumonia (J10-J18)	52	4.5	198.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	28	2.4	107.0
8	Alzheimer's disease (G30)	19	1.7	*
9	Parkinson's disease (G20-G21)	18	1.6	*
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	18	1.6	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18	1.6	*
12	Septicemia (A40-A41)	16	1.4	*
13	Pneumonitis due to solids and liquids (J69)	12	1.0	*
14	Chronic liver disease and cirrhosis (K70,K73-K74)	11	1.0	*
15	Aortic aneurysm and dissection (I71)	7	0.6	*
...	All other causes (Residual)	163	14.2	623.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 85 years and over				
...	All causes	614	100.0	8,419.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	183	29.8	2,509.3
2	Malignant neoplasms (C00-C97)	95	15.5	1,302.6
3	Influenza and pneumonia (J10-J18)	45	7.3	617.0
4	Cerebrovascular diseases (I60-I69)	34	5.5	466.2
5	Chronic lower respiratory diseases (J40-J47)	32	5.2	438.8
6	Diabetes mellitus (E10-E14)	31	5.0	425.1
7	Alzheimer's disease (G30)	23	3.7	315.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19	3.1	*
9	Septicemia (A40-A41)	14	2.3	*
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12	2.0	*
11	Parkinson's disease (G20-G21)	9	1.5	*
11	Pneumonitis due to solids and liquids (J69)	9	1.5	*
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	8	1.3	*
14	Anemias (D50-D64)	4	0.7	*
14	Atherosclerosis (I70)	4	0.7	*
...	All other causes (Residual)	92	15.0	1,261.5
American Indian, female, all ages				
...	All causes	6,311	100.0	398.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,188	18.8	75.1
2	Malignant neoplasms (C00-C97)	1,168	18.5	73.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	508	8.0	32.1
4	Diabetes mellitus (E10-E14)	406	6.4	25.7
5	Cerebrovascular diseases (I60-I69)	378	6.0	23.9
6	Chronic lower respiratory diseases (J40-J47)	261	4.1	16.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	248	3.9	15.7
8	Influenza and pneumonia (J10-J18)	174	2.8	11.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	160	2.5	10.1
10	Alzheimer's disease (G30)	125	2.0	7.9
11	Septicemia (A40-A41)	113	1.8	7.1
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	79	1.3	5.0
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	64	1.0	4.0
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	57	0.9	3.6
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	56	0.9	3.5
...	All other causes (Residual)	1,326	21.0	83.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 1-4 years				
...	All causes	38	100.0	45.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13	34.2	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	18.4	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	15.8	*
4	Septicemia (A40-A41)	1	2.6	*
4	Malignant neoplasms (C00-C97)	1	2.6	*
4	Meningitis (G00,G03)	1	2.6	*
4	Influenza and pneumonia (J10-J18)	1	2.6	*
4	Chronic lower respiratory diseases (J40-J47)	1	2.6	*
...	All other causes (Residual)	7	18.4	*
American Indian, female, 5-14 years				
...	All causes	46	100.0	16.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19	41.3	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	10.9	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	8.7	*
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3	6.5	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	4.3	*
6	Meningitis (G00,G03)	1	2.2	*
6	Chronic lower respiratory diseases (J40-J47)	1	2.2	*
...	All other causes (Residual)	11	23.9	*
American Indian, female, 15-24 years				
...	All causes	196	100.0	67.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	98	50.0	34.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	29	14.8	10.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	15	7.7	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	6	3.1	*
5	Malignant neoplasms (C00-C97)	5	2.6	*
6	Septicemia (A40-A41)	3	1.5	*
7	Diabetes mellitus (E10-E14)	2	1.0	*
7	Pregnancy, childbirth and the puerperium (O00-099)	2	1.0	*
9	Salmonella infections (A01-A02)	1	0.5	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.5	*
9	Anemias (D50-D64)	1	0.5	*
9	Cerebrovascular diseases (I60-I69)	1	0.5	*
9	Influenza and pneumonia (J10-J18)	1	0.5	*
9	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.5	*
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.5	*
...	All other causes (Residual)	29	14.8	10.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 25-34 years				
...	All causes	208	100.0	90.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	84	40.4	36.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	18	8.7	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	15	7.2	*
4	Chronic liver disease and cirrhosis (K70,K73-K74)	14	6.7	*
5	Malignant neoplasms (C00-C97)	13	6.3	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	13	6.3	*
7	Diabetes mellitus (E10-E14)	6	2.9	*
8	Septicemia (A40-A41)	3	1.4	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	3	1.4	*
10	Cerebrovascular diseases (I60-I69)	2	1.0	*
10	Chronic lower respiratory diseases (J40-J47)	2	1.0	*
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	1.0	*
13	Influenza and pneumonia (J10-J18)	1	0.5	*
13	Pneumonitis due to solids and liquids (J69)	1	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.5	*
13	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.5	*
13	Pregnancy, childbirth and the puerperium (O00-099)	1	0.5	*
...	All other causes (Residual)	28	13.5	12.2
American Indian, female, 35-44 years				
...	All causes	446	100.0	194.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	92	20.6	40.0
2	Chronic liver disease and cirrhosis (K70,K73-K74)	55	12.3	23.9
3	Malignant neoplasms (C00-C97)	54	12.1	23.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	36	8.1	15.7
5	Cerebrovascular diseases (I60-I69)	16	3.6	*
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	16	3.6	*
7	Diabetes mellitus (E10-E14)	14	3.1	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	13	2.9	*
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	13	2.9	*
10	Chronic lower respiratory diseases (J40-J47)	9	2.0	*
11	Septicemia (A40-A41)	6	1.3	*
11	Influenza and pneumonia (J10-J18)	6	1.3	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	1.3	*
14	Pregnancy, childbirth and the puerperium (O00-099)	4	0.9	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	0.9	*
...	All other causes (Residual)	102	22.9	44.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 45-54 years				
...	All causes	767	100.0	366.2
1	Malignant neoplasms (C00-C97)	179	23.3	85.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	109	14.2	52.0
3	Chronic liver disease and cirrhosis (K70,K73-K74)	80	10.4	38.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	77	10.0	36.8
5	Cerebrovascular diseases (I60-I69)	37	4.8	17.7
6	Diabetes mellitus (E10-E14)	35	4.6	16.7
7	Chronic lower respiratory diseases (J40-J47)	15	2.0	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	1.8	*
9	Septicemia (A40-A41)	12	1.6	*
10	Influenza and pneumonia (J10-J18)	11	1.4	*
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11	1.4	*
12	Viral hepatitis (B15-B19)	5	0.7	*
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	0.5	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.5	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.4	*
15	Pneumonitis due to solids and liquids (J69)	3	0.4	*
...	All other causes (Residual)	168	21.9	80.2
American Indian, female, 55-64 years				
...	All causes	899	100.0	699.4
1	Malignant neoplasms (C00-C97)	259	28.8	201.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	157	17.5	122.1
3	Diabetes mellitus (E10-E14)	83	9.2	64.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	48	5.3	37.3
5	Cerebrovascular diseases (I60-I69)	46	5.1	35.8
5	Chronic lower respiratory diseases (J40-J47)	46	5.1	35.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	36	4.0	28.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	30	3.3	23.3
9	Septicemia (A40-A41)	23	2.6	17.9
10	Influenza and pneumonia (J10-J18)	19	2.1	*
11	Atherosclerosis (I70)	5	0.6	*
12	Viral hepatitis (B15-B19)	4	0.4	*
13	Parkinson's disease (G20-G21)	3	0.3	*
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	3	0.3	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.2	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.2	*
15	Aortic aneurysm and dissection (I71)	2	0.2	*
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	2	0.2	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	0.2	*
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	0.2	*
...	All other causes (Residual)	125	13.9	97.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 65-74 years				
...	All causes	1,190	100.0	1,780.5
1	Malignant neoplasms (C00-C97)	318	26.7	475.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	233	19.6	348.6
3	Diabetes mellitus (E10-E14)	119	10.0	178.1
4	Cerebrovascular diseases (I60-I69)	77	6.5	115.2
4	Chronic lower respiratory diseases (J40-J47)	77	6.5	115.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	37	3.1	55.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	32	2.7	47.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	30	2.5	44.9
9	Influenza and pneumonia (J10-J18)	26	2.2	38.9
10	Septicemia (A40-A41)	18	1.5	*
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	12	1.0	*
12	Parkinson's disease (G20-G21)	7	0.6	*
12	Pneumonitis due to solids and liquids (J69)	7	0.6	*
14	Atherosclerosis (I70)	6	0.5	*
15	Viral hepatitis (B15-B19)	4	0.3	*
15	Alzheimer's disease (G30)	4	0.3	*
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	0.3	*
...	All other causes (Residual)	179	15.0	267.8
American Indian, female, 75-84 years				
...	All causes	1,289	100.0	3,602.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	303	23.5	846.8
2	Malignant neoplasms (C00-C97)	251	19.5	701.5
3	Cerebrovascular diseases (I60-I69)	103	8.0	287.9
4	Diabetes mellitus (E10-E14)	90	7.0	251.5
5	Chronic lower respiratory diseases (J40-J47)	72	5.6	201.2
6	Alzheimer's disease (G30)	50	3.9	139.7
7	Influenza and pneumonia (J10-J18)	46	3.6	128.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	44	3.4	123.0
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	32	2.5	89.4
10	Septicemia (A40-A41)	29	2.2	81.1
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	20	1.6	55.9
12	Chronic liver disease and cirrhosis (K70,K73-K74)	16	1.2	*
13	Parkinson's disease (G20-G21)	13	1.0	*
14	Atherosclerosis (I70)	10	0.8	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.6	*
15	Aortic aneurysm and dissection (I71)	8	0.6	*
...	All other causes (Residual)	194	15.1	542.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 85 years and over				
...	All causes	1,070	100.0	7,065.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	325	30.4	2,145.9
2	Cerebrovascular diseases (I60-I69)	95	8.9	627.3
3	Malignant neoplasms (C00-C97)	88	8.2	581.0
4	Alzheimer's disease (G30)	71	6.6	468.8
5	Influenza and pneumonia (J10-J18)	60	5.6	396.2
6	Diabetes mellitus (E10-E14)	57	5.3	376.4
7	Chronic lower respiratory diseases (J40-J47)	37	3.5	244.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	27	2.5	178.3
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	19	1.8	*
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18	1.7	*
11	Septicemia (A40-A41)	16	1.5	*
12	Pneumonitis due to solids and liquids (J69)	12	1.1	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	0.9	*
14	Parkinson's disease (G20-G21)	9	0.8	*
15	Aortic aneurysm and dissection (I71)	7	0.7	*
...	All other causes (Residual)	219	20.5	1,446.0
Asian or Pacific Islander, both sexes, all ages				
...	All causes	43,194	100.0	307.7
1	Malignant neoplasms (C00-C97)	11,550	26.7	82.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,477	24.3	74.6
3	Cerebrovascular diseases (I60-I69)	3,543	8.2	25.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,125	4.9	15.1
5	Diabetes mellitus (E10-E14)	1,576	3.6	11.2
6	Influenza and pneumonia (J10-J18)	1,327	3.1	9.5
7	Chronic lower respiratory diseases (J40-J47)	1,302	3.0	9.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	755	1.7	5.4
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	726	1.7	5.2
10	Alzheimer's disease (G30)	615	1.4	4.4
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	582	1.3	4.1
12	Septicemia (A40-A41)	504	1.2	3.6
13	Certain conditions originating in the perinatal period (P00-P96)	463	1.1	3.3
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	428	1.0	3.0
15	Chronic liver disease and cirrhosis (K70,K73-K74)	401	0.9	2.9
...	All other causes (Residual)	6,820	15.8	48.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 1-4 years				
...	All causes	151	100.0	19.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	31	20.5	3.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	20	13.2	2.5
3	Malignant neoplasms (C00-C97)	16	10.6	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	12	7.9	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8	5.3	*
6	Septicemia (A40-A41)	5	3.3	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	3.3	*
8	Influenza and pneumonia (J10-J18)	4	2.6	*
9	Meningitis (G00,G03)	3	2.0	*
9	Cerebrovascular diseases (I60-I69)	3	2.0	*
11	Anemias (D50-D64)	1	0.7	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.7	*
...	All other causes (Residual)	42	27.8	5.3
Asian or Pacific Islander, both sexes, 5-14 years				
...	All causes	239	100.0	12.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	74	31.0	4.0
2	Malignant neoplasms (C00-C97)	37	15.5	2.0
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	17	7.1	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	16	6.7	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	14	5.9	*
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8	3.3	*
7	Cerebrovascular diseases (I60-I69)	7	2.9	*
8	Influenza and pneumonia (J10-J18)	6	2.5	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	2.1	*
10	Chronic lower respiratory diseases (J40-J47)	4	1.7	*
11	Septicemia (A40-A41)	3	1.3	*
12	Meningococcal infection (A39)	1	0.4	*
12	Diabetes mellitus (E10-E14)	1	0.4	*
12	Pneumonitis due to solids and liquids (J69)	1	0.4	*
12	Diseases of appendix (K35-K38)	1	0.4	*
12	Certain conditions originating in the perinatal period (P00-P96)	1	0.4	*
...	All other causes (Residual)	43	18.0	2.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 15-24 years				
...	All causes	818	100.0	41.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	327	40.0	16.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	135	16.5	6.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	107	13.1	5.5
4	Malignant neoplasms (C00-C97)	73	8.9	3.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	36	4.4	1.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	17	2.1	*
7	Cerebrovascular diseases (I60-I69)	7	0.9	*
8	Septicemia (A40-A41)	6	0.7	*
8	Influenza and pneumonia (J10-J18)	6	0.7	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.5	*
11	Tuberculosis (A16-A19)	3	0.4	*
11	Anemias (D50-D64)	3	0.4	*
13	Viral hepatitis (B15-B19)	2	0.2	*
13	Meningitis (G00,G03)	2	0.2	*
13	Chronic lower respiratory diseases (J40-J47)	2	0.2	*
13	Pregnancy, childbirth and the puerperium (O00-099)	2	0.2	*
...	All other causes (Residual)	86	10.5	4.4
Asian or Pacific Islander, both sexes, 25-34 years				
...	All causes	1,066	100.0	41.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	249	23.4	9.8
2	Malignant neoplasms (C00-C97)	190	17.8	7.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	185	17.4	7.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	97	9.1	3.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	79	7.4	3.1
6	Cerebrovascular diseases (I60-I69)	23	2.2	0.9
7	Pregnancy, childbirth and the puerperium (O00-099)	15	1.4	*
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	13	1.2	*
9	Chronic liver disease and cirrhosis (K70,K73-K74)	11	1.0	*
10	Diabetes mellitus (E10-E14)	9	0.8	*
10	Influenza and pneumonia (J10-J18)	9	0.8	*
12	Septicemia (A40-A41)	8	0.8	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	8	0.8	*
12	Aortic aneurysm and dissection (I71)	8	0.8	*
15	Chronic lower respiratory diseases (J40-J47)	6	0.6	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.6	*
...	All other causes (Residual)	150	14.1	5.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 35-44 years				
...	All causes	1,792	100.0	75.4
1	Malignant neoplasms (C00-C97)	538	30.0	22.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	278	15.5	11.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	234	13.1	9.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	130	7.3	5.5
5	Cerebrovascular diseases (I60-I69)	89	5.0	3.7
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	80	4.5	3.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	40	2.2	1.7
8	Diabetes mellitus (E10-E14)	35	2.0	1.5
9	Human immunodeficiency virus (HIV) disease (B20-B24)	33	1.8	1.4
10	Aortic aneurysm and dissection (I71)	17	0.9	*
11	Influenza and pneumonia (J10-J18)	16	0.9	*
12	Viral hepatitis (B15-B19)	13	0.7	*
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	13	0.7	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	0.7	*
15	Chronic lower respiratory diseases (J40-J47)	12	0.7	*
15	Pregnancy, childbirth and the puerperium (O00-099)	12	0.7	*
...	All other causes (Residual)	239	13.3	10.1
Asian or Pacific Islander, both sexes, 45-54 years				
...	All causes	3,630	100.0	189.7
1	Malignant neoplasms (C00-C97)	1,447	39.9	75.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	683	18.8	35.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	257	7.1	13.4
4	Cerebrovascular diseases (I60-I69)	230	6.3	12.0
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	119	3.3	6.2
6	Diabetes mellitus (E10-E14)	104	2.9	5.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	87	2.4	4.5
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	52	1.4	2.7
9	Viral hepatitis (B15-B19)	41	1.1	2.1
10	Aortic aneurysm and dissection (I71)	34	0.9	1.8
10	Influenza and pneumonia (J10-J18)	34	0.9	1.8
12	Chronic lower respiratory diseases (J40-J47)	33	0.9	1.7
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	33	0.9	1.7
14	Septicemia (A40-A41)	30	0.8	1.6
15	Human immunodeficiency virus (HIV) disease (B20-B24)	23	0.6	1.2
...	All other causes (Residual)	423	11.7	22.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 55-64 years				
...	All causes	5,427	100.0	442.4
1	Malignant neoplasms (C00-C97)	2,261	41.7	184.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,201	22.1	97.9
3	Cerebrovascular diseases (I60-I69)	369	6.8	30.1
4	Diabetes mellitus (E10-E14)	226	4.2	18.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	215	4.0	17.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	85	1.6	6.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	82	1.5	6.7
8	Chronic lower respiratory diseases (J40-J47)	77	1.4	6.3
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	75	1.4	6.1
10	Influenza and pneumonia (J10-J18)	68	1.3	5.5
11	Septicemia (A40-A41)	48	0.9	3.9
12	Viral hepatitis (B15-B19)	47	0.9	3.8
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	45	0.8	3.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	38	0.7	3.1
15	Aortic aneurysm and dissection (I71)	30	0.6	2.4
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	30	0.6	2.4
...	All other causes (Residual)	530	9.8	43.2
Asian or Pacific Islander, both sexes, 65-74 years				
...	All causes	7,788	100.0	1,129.1
1	Malignant neoplasms (C00-C97)	2,771	35.6	401.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,899	24.4	275.3
3	Cerebrovascular diseases (I60-I69)	633	8.1	91.8
4	Diabetes mellitus (E10-E14)	345	4.4	50.0
5	Chronic lower respiratory diseases (J40-J47)	231	3.0	33.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	206	2.6	29.9
7	Influenza and pneumonia (J10-J18)	165	2.1	23.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	156	2.0	22.6
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	98	1.3	14.2
10	Septicemia (A40-A41)	92	1.2	13.3
11	Chronic liver disease and cirrhosis (K70,K73-K74)	67	0.9	9.7
12	Viral hepatitis (B15-B19)	64	0.8	9.3
13	Aortic aneurysm and dissection (I71)	63	0.8	9.1
14	Parkinson's disease (G20-G21)	53	0.7	7.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	52	0.7	7.5
...	All other causes (Residual)	893	11.5	129.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 75-84 years				
...	All causes	11,414	100.0	3,033.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,034	26.6	806.3
2	Malignant neoplasms (C00-C97)	2,876	25.2	764.3
3	Cerebrovascular diseases (I60-I69)	1,118	9.8	297.1
4	Diabetes mellitus (E10-E14)	514	4.5	136.6
5	Chronic lower respiratory diseases (J40-J47)	498	4.4	132.3
6	Influenza and pneumonia (J10-J18)	404	3.5	107.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	290	2.5	77.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	243	2.1	64.6
9	Alzheimer's disease (G30)	202	1.8	53.7
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	179	1.6	47.6
11	Septicemia (A40-A41)	162	1.4	43.1
12	Parkinson's disease (G20-G21)	124	1.1	33.0
13	Aortic aneurysm and dissection (I71)	112	1.0	29.8
14	Chronic liver disease and cirrhosis (K70,K73-K74)	89	0.8	23.7
15	Pneumonitis due to solids and liquids (J69)	79	0.7	21.0
...	All other causes (Residual)	1,490	13.1	396.0
Asian or Pacific Islander, both sexes, 85 years and over				
...	All causes	9,994	100.0	8,565.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,204	32.1	2,746.0
2	Malignant neoplasms (C00-C97)	1,340	13.4	1,148.5
3	Cerebrovascular diseases (I60-I69)	1,059	10.6	907.6
4	Influenza and pneumonia (J10-J18)	611	6.1	523.7
5	Chronic lower respiratory diseases (J40-J47)	439	4.4	376.3
6	Alzheimer's disease (G30)	372	3.7	318.8
7	Diabetes mellitus (E10-E14)	341	3.4	292.3
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	226	2.3	193.7
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	222	2.2	190.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	215	2.2	184.3
11	Septicemia (A40-A41)	131	1.3	112.3
12	Pneumonitis due to solids and liquids (J69)	128	1.3	109.7
13	Parkinson's disease (G20-G21)	86	0.9	73.7
14	Aortic aneurysm and dissection (I71)	76	0.8	65.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	44	0.4	37.7
...	All other causes (Residual)	1,500	15.0	1,285.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, all ages				
...	All causes	22,808	100.0	333.9
1	Malignant neoplasms (C00-C97)	5,921	26.0	86.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,703	25.0	83.5
3	Cerebrovascular diseases (I60-I69)	1,624	7.1	23.8
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,328	5.8	19.4
5	Chronic lower respiratory diseases (J40-J47)	803	3.5	11.8
6	Diabetes mellitus (E10-E14)	768	3.4	11.2
7	Influenza and pneumonia (J10-J18)	718	3.1	10.5
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	495	2.2	7.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	381	1.7	5.6
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	314	1.4	4.6
11	Septicemia (A40-A41)	260	1.1	3.8
12	Certain conditions originating in the perinatal period (P00-P96)	258	1.1	3.8
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	256	1.1	3.7
14	Chronic liver disease and cirrhosis (K70,K73-K74)	254	1.1	3.7
15	Alzheimer's disease (G30)	229	1.0	3.4
...	All other causes (Residual)	3,496	15.3	51.2
Asian or Pacific Islander, male, 1-4 years				
...	All causes	84	100.0	20.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19	22.6	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	11	13.1	*
3	Malignant neoplasms (C00-C97)	9	10.7	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	6	7.1	*
5	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	3.6	*
5	Meningitis (G00,G03)	3	3.6	*
5	Cerebrovascular diseases (I60-I69)	3	3.6	*
5	Influenza and pneumonia (J10-J18)	3	3.6	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	3.6	*
10	Septicemia (A40-A41)	1	1.2	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	1	1.2	*
...	All other causes (Residual)	22	26.2	5.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 5-14 years				
...	All causes	131	100.0	14.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	42	32.1	4.5
2	Malignant neoplasms (C00-C97)	21	16.0	2.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	6.1	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8	6.1	*
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6	4.6	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	3.8	*
6	Cerebrovascular diseases (I60-I69)	5	3.8	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	2.3	*
8	Influenza and pneumonia (J10-J18)	3	2.3	*
10	Septicemia (A40-A41)	2	1.5	*
10	Chronic lower respiratory diseases (J40-J47)	2	1.5	*
12	Meningococcal infection (A39)	1	0.8	*
12	Diabetes mellitus (E10-E14)	1	0.8	*
...	All other causes (Residual)	24	18.3	2.6
Asian or Pacific Islander, male, 15-24 years				
...	All causes	569	100.0	56.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	235	41.3	23.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	108	19.0	10.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	72	12.7	7.2
4	Malignant neoplasms (C00-C97)	44	7.7	4.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	25	4.4	2.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	10	1.8	*
7	Cerebrovascular diseases (I60-I69)	4	0.7	*
7	Influenza and pneumonia (J10-J18)	4	0.7	*
9	Tuberculosis (A16-A19)	2	0.4	*
9	Septicemia (A40-A41)	2	0.4	*
9	Anemias (D50-D64)	2	0.4	*
12	Viral hepatitis (B15-B19)	1	0.2	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.2	*
12	Diabetes mellitus (E10-E14)	1	0.2	*
12	Meningitis (G00,G03)	1	0.2	*
12	Chronic lower respiratory diseases (J40-J47)	1	0.2	*
12	Pneumonitis due to solids and liquids (J69)	1	0.2	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
...	All other causes (Residual)	53	9.3	5.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 25-34 years				
...	All causes	696	100.0	55.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	180	25.9	14.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	144	20.7	11.5
3	Malignant neoplasms (C00-C97)	90	12.9	7.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	71	10.2	5.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	57	8.2	4.6
6	Cerebrovascular diseases (I60-I69)	13	1.9	*
7	Aortic aneurysm and dissection (I71)	8	1.1	*
7	Chronic liver disease and cirrhosis (K70,K73-K74)	8	1.1	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	7	1.0	*
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	1.0	*
11	Diabetes mellitus (E10-E14)	6	0.9	*
12	Influenza and pneumonia (J10-J18)	5	0.7	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.7	*
14	Septicemia (A40-A41)	4	0.6	*
14	Chronic lower respiratory diseases (J40-J47)	4	0.6	*
...	All other causes (Residual)	87	12.5	6.9
Asian or Pacific Islander, male, 35-44 years				
...	All causes	1,082	100.0	93.6
1	Malignant neoplasms (C00-C97)	231	21.3	20.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	214	19.8	18.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	160	14.8	13.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	86	7.9	7.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	60	5.5	5.2
6	Cerebrovascular diseases (I60-I69)	46	4.3	4.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	33	3.0	2.9
8	Human immunodeficiency virus (HIV) disease (B20-B24)	27	2.5	2.3
9	Diabetes mellitus (E10-E14)	21	1.9	1.8
10	Aortic aneurysm and dissection (I71)	15	1.4	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	1.0	*
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	9	0.8	*
12	Chronic lower respiratory diseases (J40-J47)	9	0.8	*
14	Viral hepatitis (B15-B19)	8	0.7	*
14	Influenza and pneumonia (J10-J18)	8	0.7	*
...	All other causes (Residual)	144	13.3	12.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 45-54 years				
...	All causes	2,182	100.0	242.4
1	Malignant neoplasms (C00-C97)	683	31.3	75.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	523	24.0	58.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	171	7.8	19.0
4	Cerebrovascular diseases (I60-I69)	130	6.0	14.4
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	81	3.7	9.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	71	3.3	7.9
7	Diabetes mellitus (E10-E14)	69	3.2	7.7
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	40	1.8	4.4
9	Viral hepatitis (B15-B19)	30	1.4	3.3
10	Aortic aneurysm and dissection (I71)	28	1.3	3.1
11	Influenza and pneumonia (J10-J18)	22	1.0	2.4
12	Chronic lower respiratory diseases (J40-J47)	21	1.0	2.3
13	Human immunodeficiency virus (HIV) disease (B20-B24)	20	0.9	2.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20	0.9	2.2
15	Septicemia (A40-A41)	17	0.8	*
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	17	0.8	*
...	All other causes (Residual)	239	11.0	26.6
Asian or Pacific Islander, male, 55-64 years				
...	All causes	3,105	100.0	545.4
1	Malignant neoplasms (C00-C97)	1,135	36.6	199.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	827	26.6	145.3
3	Cerebrovascular diseases (I60-I69)	190	6.1	33.4
4	Diabetes mellitus (E10-E14)	144	4.6	25.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	129	4.2	22.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	56	1.8	9.8
7	Chronic lower respiratory diseases (J40-J47)	54	1.7	9.5
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	50	1.6	8.8
9	Influenza and pneumonia (J10-J18)	42	1.4	7.4
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	40	1.3	7.0
11	Viral hepatitis (B15-B19)	32	1.0	5.6
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	21	0.7	3.7
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	21	0.7	3.7
14	Septicemia (A40-A41)	20	0.6	3.5
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	20	0.6	3.5
14	Aortic aneurysm and dissection (I71)	20	0.6	3.5
...	All other causes (Residual)	304	9.8	53.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 65-74 years				
...	All causes	4,344	100.0	1,403.8
1	Malignant neoplasms (C00-C97)	1,523	35.1	492.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,160	26.7	374.9
3	Cerebrovascular diseases (I60-I69)	325	7.5	105.0
4	Diabetes mellitus (E10-E14)	182	4.2	58.8
5	Chronic lower respiratory diseases (J40-J47)	140	3.2	45.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	118	2.7	38.1
7	Influenza and pneumonia (J10-J18)	98	2.3	31.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	82	1.9	26.5
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	50	1.2	16.2
10	Septicemia (A40-A41)	49	1.1	15.8
11	Aortic aneurysm and dissection (I71)	45	1.0	14.5
12	Chronic liver disease and cirrhosis (K70,K73-K74)	32	0.7	10.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	31	0.7	10.0
13	Parkinson's disease (G20-G21)	31	0.7	10.0
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	26	0.6	8.4
...	All other causes (Residual)	452	10.4	146.1
Asian or Pacific Islander, male, 75-84 years				
...	All causes	5,794	100.0	3,759.2
1	Malignant neoplasms (C00-C97)	1,528	26.4	991.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,517	26.2	984.3
3	Cerebrovascular diseases (I60-I69)	520	9.0	337.4
4	Chronic lower respiratory diseases (J40-J47)	319	5.5	207.0
5	Influenza and pneumonia (J10-J18)	234	4.0	151.8
6	Diabetes mellitus (E10-E14)	209	3.6	135.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	152	2.6	98.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	121	2.1	78.5
9	Alzheimer's disease (G30)	89	1.5	57.7
10	Septicemia (A40-A41)	88	1.5	57.1
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	76	1.3	49.3
12	Parkinson's disease (G20-G21)	68	1.2	44.1
13	Aortic aneurysm and dissection (I71)	65	1.1	42.2
14	Pneumonitis due to solids and liquids (J69)	44	0.8	28.5
14	Chronic liver disease and cirrhosis (K70,K73-K74)	44	0.8	28.5
...	All other causes (Residual)	720	12.4	467.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 85 years and over				
...	All causes	4,336	100.0	9,839.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,345	31.0	3,052.0
2	Malignant neoplasms (C00-C97)	656	15.1	1,488.6
3	Cerebrovascular diseases (I60-I69)	385	8.9	873.6
4	Influenza and pneumonia (J10-J18)	296	6.8	671.7
5	Chronic lower respiratory diseases (J40-J47)	253	5.8	574.1
6	Diabetes mellitus (E10-E14)	135	3.1	306.3
7	Alzheimer's disease (G30)	122	2.8	276.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	114	2.6	258.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	101	2.3	229.2
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	83	1.9	188.3
11	Pneumonitis due to solids and liquids (J69)	67	1.5	152.0
12	Septicemia (A40-A41)	65	1.5	147.5
13	Parkinson's disease (G20-G21)	46	1.1	104.4
14	Aortic aneurysm and dissection (I71)	44	1.0	99.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	0.5	52.2
...	All other causes (Residual)	601	13.9	1,363.8
Asian or Pacific Islander, female, all ages				
...	All causes	20,386	100.0	282.8
1	Malignant neoplasms (C00-C97)	5,629	27.6	78.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,774	23.4	66.2
3	Cerebrovascular diseases (I60-I69)	1,919	9.4	26.6
4	Diabetes mellitus (E10-E14)	808	4.0	11.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	797	3.9	11.1
6	Influenza and pneumonia (J10-J18)	609	3.0	8.4
7	Chronic lower respiratory diseases (J40-J47)	499	2.4	6.9
8	Alzheimer's disease (G30)	386	1.9	5.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	374	1.8	5.2
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	326	1.6	4.5
11	Septicemia (A40-A41)	244	1.2	3.4
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	231	1.1	3.2
13	Certain conditions originating in the perinatal period (P00-P96)	205	1.0	2.8
14	Chronic liver disease and cirrhosis (K70,K73-K74)	147	0.7	2.0
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	146	0.7	2.0
...	All other causes (Residual)	3,292	16.1	45.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 1-4 years				
...	All causes	67	100.0	17.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12	17.9	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	13.4	*
3	Malignant neoplasms (C00-C97)	7	10.4	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	6	9.0	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	7.5	*
6	Septicemia (A40-A41)	4	6.0	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	3.0	*
8	Anemias (D50-D64)	1	1.5	*
8	Influenza and pneumonia (J10-J18)	1	1.5	*
...	All other causes (Residual)	20	29.9	5.2
Asian or Pacific Islander, female, 5-14 years				
...	All causes	108	100.0	11.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	32	29.6	3.5
2	Malignant neoplasms (C00-C97)	16	14.8	*
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	9	8.3	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	8.3	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8	7.4	*
6	Influenza and pneumonia (J10-J18)	3	2.8	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	1.9	*
7	Cerebrovascular diseases (I60-I69)	2	1.9	*
7	Chronic lower respiratory diseases (J40-J47)	2	1.9	*
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2	1.9	*
11	Septicemia (A40-A41)	1	0.9	*
11	Pneumonitis due to solids and liquids (J69)	1	0.9	*
11	Diseases of appendix (K35-K38)	1	0.9	*
11	Certain conditions originating in the perinatal period (P00-P96)	1	0.9	*
...	All other causes (Residual)	19	17.6	*

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 15-24 years				
...	All causes	249	100.0	26.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	92	36.9	9.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	35	14.1	3.7
3	Malignant neoplasms (C00-C97)	29	11.6	3.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	27	10.8	2.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	11	4.4	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	2.8	*
7	Septicemia (A40-A41)	4	1.6	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	1.2	*
8	Cerebrovascular diseases (I60-I69)	3	1.2	*
10	Influenza and pneumonia (J10-J18)	2	0.8	*
10	Pregnancy, childbirth and the puerperium (O00-099)	2	0.8	*
12	Tuberculosis (A16-A19)	1	0.4	*
12	Viral hepatitis (B15-B19)	1	0.4	*
12	Anemias (D50-D64)	1	0.4	*
12	Nutritional deficiencies (E40-E64)	1	0.4	*
12	Meningitis (G00,G03)	1	0.4	*
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1	0.4	*
12	Chronic lower respiratory diseases (J40-J47)	1	0.4	*
12	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.4	*
...	All other causes (Residual)	26	10.4	2.7
Asian or Pacific Islander, female, 25-34 years				
...	All causes	370	100.0	28.6
1	Malignant neoplasms (C00-C97)	100	27.0	7.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	69	18.6	5.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	41	11.1	3.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	26	7.0	2.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	22	5.9	1.7
6	Pregnancy, childbirth and the puerperium (O00-099)	15	4.1	*
7	Cerebrovascular diseases (I60-I69)	10	2.7	*
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	1.6	*
9	Septicemia (A40-A41)	4	1.1	*
9	Influenza and pneumonia (J10-J18)	4	1.1	*
11	Diabetes mellitus (E10-E14)	3	0.8	*
11	Chronic liver disease and cirrhosis (K70,K73-K74)	3	0.8	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.5	*
13	Chronic lower respiratory diseases (J40-J47)	2	0.5	*
15	Tuberculosis (A16-A19)	1	0.3	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.3	*
15	Anemias (D50-D64)	1	0.3	*
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
15	Inflammatory diseases of female pelvic organs (N70-N76)	1	0.3	*
15	Certain conditions originating in the perinatal period (P00-P96)	1	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.3	*
...	All other causes (Residual)	55	14.9	4.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 35-44 years				
...	All causes	710	100.0	58.1
1	Malignant neoplasms (C00-C97)	307	43.2	25.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	74	10.4	6.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	64	9.0	5.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	44	6.2	3.6
5	Cerebrovascular diseases (I60-I69)	43	6.1	3.5
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	20	2.8	1.6
7	Diabetes mellitus (E10-E14)	14	2.0	*
8	Pregnancy, childbirth and the puerperium (O00-099)	12	1.7	*
9	Influenza and pneumonia (J10-J18)	8	1.1	*
10	Chronic liver disease and cirrhosis (K70,K73-K74)	7	1.0	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	6	0.8	*
12	Viral hepatitis (B15-B19)	5	0.7	*
13	Septicemia (A40-A41)	4	0.6	*
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	4	0.6	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.4	*
15	Chronic lower respiratory diseases (J40-J47)	3	0.4	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.4	*
...	All other causes (Residual)	89	12.5	7.3
Asian or Pacific Islander, female, 45-54 years				
...	All causes	1,448	100.0	142.8
1	Malignant neoplasms (C00-C97)	764	52.8	75.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	160	11.0	15.8
3	Cerebrovascular diseases (I60-I69)	100	6.9	9.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	86	5.9	8.5
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	38	2.6	3.7
6	Diabetes mellitus (E10-E14)	35	2.4	3.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	16	1.1	*
8	Septicemia (A40-A41)	13	0.9	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	0.9	*
10	Influenza and pneumonia (J10-J18)	12	0.8	*
10	Chronic lower respiratory diseases (J40-J47)	12	0.8	*
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	12	0.8	*
13	Viral hepatitis (B15-B19)	11	0.8	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.6	*
15	Tuberculosis (A16-A19)	8	0.6	*
...	All other causes (Residual)	159	11.0	15.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 55-64 years				
...	All causes	2,322	100.0	353.2
1	Malignant neoplasms (C00-C97)	1,126	48.5	171.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	374	16.1	56.9
3	Cerebrovascular diseases (I60-I69)	179	7.7	27.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	86	3.7	13.1
5	Diabetes mellitus (E10-E14)	82	3.5	12.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	45	1.9	6.8
7	Septicemia (A40-A41)	28	1.2	4.3
8	Influenza and pneumonia (J10-J18)	26	1.1	4.0
8	Chronic liver disease and cirrhosis (K70,K73-K74)	26	1.1	4.0
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	25	1.1	3.8
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	25	1.1	3.8
12	Chronic lower respiratory diseases (J40-J47)	23	1.0	3.5
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	0.7	*
14	Viral hepatitis (B15-B19)	15	0.6	*
15	Aortic aneurysm and dissection (I71)	10	0.4	*
...	All other causes (Residual)	235	10.1	35.7
Asian or Pacific Islander, female, 65-74 years				
...	All causes	3,444	100.0	905.5
1	Malignant neoplasms (C00-C97)	1,248	36.2	328.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	739	21.5	194.3
3	Cerebrovascular diseases (I60-I69)	308	8.9	81.0
4	Diabetes mellitus (E10-E14)	163	4.7	42.9
5	Chronic lower respiratory diseases (J40-J47)	91	2.6	23.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	88	2.6	23.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	74	2.1	19.5
8	Influenza and pneumonia (J10-J18)	67	1.9	17.6
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	48	1.4	12.6
10	Septicemia (A40-A41)	43	1.2	11.3
11	Viral hepatitis (B15-B19)	42	1.2	11.0
12	Chronic liver disease and cirrhosis (K70,K73-K74)	35	1.0	9.2
13	Parkinson's disease (G20-G21)	22	0.6	5.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	21	0.6	5.5
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	19	0.6	*
...	All other causes (Residual)	436	12.7	114.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 75-84 years				
...	All causes	5,620	100.0	2,529.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,517	27.0	682.9
2	Malignant neoplasms (C00-C97)	1,348	24.0	606.8
3	Cerebrovascular diseases (I60-I69)	598	10.6	269.2
4	Diabetes mellitus (E10-E14)	305	5.4	137.3
5	Chronic lower respiratory diseases (J40-J47)	179	3.2	80.6
6	Influenza and pneumonia (J10-J18)	170	3.0	76.5
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	138	2.5	62.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	122	2.2	54.9
9	Alzheimer's disease (G30)	113	2.0	50.9
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	103	1.8	46.4
11	Septicemia (A40-A41)	74	1.3	33.3
12	Parkinson's disease (G20-G21)	56	1.0	25.2
13	Aortic aneurysm and dissection (I71)	47	0.8	21.2
14	Chronic liver disease and cirrhosis (K70,K73-K74)	45	0.8	20.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	40	0.7	18.0
...	All other causes (Residual)	765	13.6	344.4
Asian or Pacific Islander, female, 85 years and over				
...	All causes	5,658	100.0	7,792.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,859	32.9	2,560.3
2	Malignant neoplasms (C00-C97)	684	12.1	942.0
3	Cerebrovascular diseases (I60-I69)	674	11.9	928.3
4	Influenza and pneumonia (J10-J18)	315	5.6	433.8
5	Alzheimer's disease (G30)	250	4.4	344.3
6	Diabetes mellitus (E10-E14)	206	3.6	283.7
7	Chronic lower respiratory diseases (J40-J47)	186	3.3	256.2
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	139	2.5	191.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	114	2.0	157.0
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	112	2.0	154.3
11	Septicemia (A40-A41)	66	1.2	90.9
12	Pneumonitis due to solids and liquids (J69)	61	1.1	84.0
13	Parkinson's disease (G20-G21)	40	0.7	55.1
14	Aortic aneurysm and dissection (I71)	32	0.6	44.1
15	Atherosclerosis (I70)	28	0.5	38.6
...	All other causes (Residual)	892	15.8	1,228.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable