

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, all ages				
...	All causes	2,437,163	100.0	793.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	599,413	24.6	195.2
2	Malignant neoplasms (C00-C97)	567,628	23.3	184.9
3	Chronic lower respiratory diseases (J40-J47)	137,353	5.6	44.7
4	Cerebrovascular diseases (I60-I69)	128,842	5.3	42.0
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	118,021	4.8	38.4
6	Alzheimer's disease (G30)	79,003	3.2	25.7
7	Diabetes mellitus (E10-E14)	68,705	2.8	22.4
8	Influenza and pneumonia (J09-J18)	53,692	2.2	17.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	48,935	2.0	15.9
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	36,909	1.5	12.0
11	Septicemia (A40-A41)	35,639	1.5	11.6
12	Chronic liver disease and cirrhosis (K70,K73-K74)	30,558	1.3	10.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	25,734	1.1	8.4
14	Parkinson's disease (G20-G21)	20,565	0.8	6.7
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	16,799	0.7	5.5
...	All other causes (Residual)	469,367	19.3	152.9
All races, both sexes, 1-4 years				
...	All causes	4,450	100.0	26.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,466	32.9	8.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	464	10.4	2.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	376	8.4	2.2
4	Malignant neoplasms (C00-C97)	350	7.9	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	154	3.5	0.9
6	Influenza and pneumonia (J09-J18)	146	3.3	0.9
7	Septicemia (A40-A41)	71	1.6	0.4
8	Chronic lower respiratory diseases (J40-J47)	66	1.5	0.4
9	Certain conditions originating in the perinatal period (P00-P96)	58	1.3	0.3
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	53	1.2	0.3
11	Cerebrovascular diseases (I60-I69)	47	1.1	0.3
12	Anemias (D50-D64)	20	0.4	0.1
13	Acute bronchitis and bronchiolitis (J20-J21)	19	0.4	*
14	Meningitis (G00,G03)	18	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	18	0.4	*
...	All other causes (Residual)	1,124	25.3	6.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 5-14 years				
...	All causes	5,651	100.0	13.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,689	29.9	4.2
2	Malignant neoplasms (C00-C97)	896	15.9	2.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	364	6.4	0.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	305	5.4	0.8
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	265	4.7	0.7
6	Influenza and pneumonia (J09-J18)	228	4.0	0.6
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	217	3.8	0.5
8	Chronic lower respiratory diseases (J40-J47)	123	2.2	0.3
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	85	1.5	0.2
10	Cerebrovascular diseases (I60-I69)	74	1.3	0.2
11	Septicemia (A40-A41)	70	1.2	0.2
12	Anemias (D50-D64)	35	0.6	0.1
13	Diabetes mellitus (E10-E14)	29	0.5	0.1
14	Certain conditions originating in the perinatal period (P00-P96)	26	0.5	0.1
15	Meningitis (G00,G03)	22	0.4	0.1
...	All other causes (Residual)	1,223	21.6	3.0
All races, both sexes, 15-24 years				
...	All causes	30,416	100.0	70.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,458	41.0	28.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4,862	16.0	11.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,371	14.4	10.1
4	Malignant neoplasms (C00-C97)	1,636	5.4	3.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,035	3.4	2.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	457	1.5	1.1
7	Influenza and pneumonia (J09-J18)	418	1.4	1.0
8	Pregnancy, childbirth and the puerperium (O00-O99)	227	0.7	0.5
9	Cerebrovascular diseases (I60-I69)	193	0.6	0.4
10	Chronic lower respiratory diseases (J40-J47)	187	0.6	0.4
11	Diabetes mellitus (E10-E14)	167	0.5	0.4
12	Human immunodeficiency virus (HIV) disease (B20-B24)	149	0.5	0.3
13	Septicemia (A40-A41)	144	0.5	0.3
14	Anemias (D50-D64)	93	0.3	0.2
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	86	0.3	0.2
...	All other causes (Residual)	3,933	12.9	9.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 25-34 years				
...	All causes	42,502	100.0	102.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14,062	33.1	33.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,320	12.5	12.8
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4,222	9.9	10.2
4	Malignant neoplasms (C00-C97)	3,659	8.6	8.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,174	7.5	7.6
6	Human immunodeficiency virus (HIV) disease (B20-B24)	881	2.1	2.1
7	Influenza and pneumonia (J09-J18)	807	1.9	1.9
8	Diabetes mellitus (E10-E14)	604	1.4	1.5
9	Cerebrovascular diseases (I60-I69)	537	1.3	1.3
10	Chronic liver disease and cirrhosis (K70,K73-K74)	459	1.1	1.1
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	409	1.0	1.0
12	Pregnancy, childbirth and the puerperium (O00-O99)	403	0.9	1.0
13	Septicemia (A40-A41)	383	0.9	0.9
14	Chronic lower respiratory diseases (J40-J47)	283	0.7	0.7
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	267	0.6	0.6
...	All other causes (Residual)	7,032	16.5	16.9
All races, both sexes, 35-44 years				
...	All causes	74,665	100.0	179.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15,102	20.2	36.4
2	Malignant neoplasms (C00-C97)	12,519	16.8	30.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,081	14.8	26.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,677	8.9	16.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,762	3.7	6.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	2,481	3.3	6.0
7	Human immunodeficiency virus (HIV) disease (B20-B24)	2,425	3.2	5.8
8	Cerebrovascular diseases (I60-I69)	1,916	2.6	4.6
9	Diabetes mellitus (E10-E14)	1,872	2.5	4.5
10	Influenza and pneumonia (J09-J18)	1,314	1.8	3.2
11	Septicemia (A40-A41)	925	1.2	2.2
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	829	1.1	2.0
13	Chronic lower respiratory diseases (J40-J47)	743	1.0	1.8
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	462	0.6	1.1
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	435	0.6	1.0
...	All other causes (Residual)	13,122	17.6	31.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 45-54 years				
...	All causes	187,568	100.0	420.6
1	Malignant neoplasms (C00-C97)	50,616	27.0	113.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	36,927	19.7	82.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19,974	10.6	44.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8,598	4.6	19.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	8,377	4.5	18.8
6	Cerebrovascular diseases (I60-I69)	6,163	3.3	13.8
7	Diabetes mellitus (E10-E14)	5,725	3.1	12.8
8	Chronic lower respiratory diseases (J40-J47)	4,664	2.5	10.5
9	Human immunodeficiency virus (HIV) disease (B20-B24)	3,388	1.8	7.6
10	Influenza and pneumonia (J09-J18)	2,918	1.6	6.5
11	Viral hepatitis (B15-B19)	2,602	1.4	5.8
12	Septicemia (A40-A41)	2,433	1.3	5.5
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,352	1.3	5.3
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,066	1.1	4.6
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,390	0.7	3.1
...	All other causes (Residual)	29,375	15.7	65.9
All races, both sexes, 55-64 years				
...	All causes	303,307	100.0	871.9
1	Malignant neoplasms (C00-C97)	106,829	35.2	307.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	67,261	22.2	193.4
3	Chronic lower respiratory diseases (J40-J47)	14,160	4.7	40.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,933	4.3	37.2
5	Diabetes mellitus (E10-E14)	11,361	3.7	32.7
6	Cerebrovascular diseases (I60-I69)	10,523	3.5	30.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	9,154	3.0	26.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,808	1.9	16.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,792	1.6	13.8
10	Septicemia (A40-A41)	4,628	1.5	13.3
11	Influenza and pneumonia (J09-J18)	4,139	1.4	11.9
12	Viral hepatitis (B15-B19)	3,063	1.0	8.8
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,503	0.8	7.2
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1,854	0.6	5.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,476	0.5	4.2
...	All other causes (Residual)	42,823	14.1	123.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 65-74 years				
...	All causes	401,032	100.0	1,928.8
1	Malignant neoplasms (C00-C97)	141,871	35.4	682.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	89,765	22.4	431.7
3	Chronic lower respiratory diseases (J40-J47)	31,314	7.8	150.6
4	Cerebrovascular diseases (I60-I69)	17,578	4.4	84.5
5	Diabetes mellitus (E10-E14)	14,785	3.7	71.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,940	2.2	43.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,223	2.1	39.5
8	Septicemia (A40-A41)	6,662	1.7	32.0
9	Influenza and pneumonia (J09-J18)	6,259	1.6	30.1
10	Chronic liver disease and cirrhosis (K70,K73-K74)	5,383	1.3	25.9
11	Alzheimer's disease (G30)	4,109	1.0	19.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,455	0.9	16.6
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,917	0.7	14.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,452	0.6	11.8
15	Parkinson's disease (G20-G21)	2,380	0.6	11.4
...	All other causes (Residual)	54,939	13.7	264.2
All races, both sexes, 75-84 years				
...	All causes	627,727	100.0	4,774.4
1	Malignant neoplasms (C00-C97)	157,960	25.2	1,201.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	157,686	25.1	1,199.3
3	Chronic lower respiratory diseases (J40-J47)	49,021	7.8	372.8
4	Cerebrovascular diseases (I60-I69)	38,407	6.1	292.1
5	Alzheimer's disease (G30)	23,324	3.7	177.4
6	Diabetes mellitus (E10-E14)	18,991	3.0	144.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14,993	2.4	114.0
8	Influenza and pneumonia (J09-J18)	13,929	2.2	105.9
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,482	2.1	102.5
10	Septicemia (A40-A41)	10,313	1.6	78.4
11	Parkinson's disease (G20-G21)	9,219	1.5	70.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	6,641	1.1	50.5
13	Pneumonitis due to solids and liquids (J69)	4,651	0.7	35.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,561	0.7	34.7
15	Chronic liver disease and cirrhosis (K70,K73-K74)	3,538	0.6	26.9
...	All other causes (Residual)	101,011	16.1	768.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 85 years and over				
...	All causes	733,178	100.0	13,021.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	231,699	31.6	4,115.0
2	Malignant neoplasms (C00-C97)	91,204	12.4	1,619.8
3	Cerebrovascular diseases (I60-I69)	53,253	7.3	945.8
4	Alzheimer's disease (G30)	50,735	6.9	901.0
5	Chronic lower respiratory diseases (J40-J47)	36,763	5.0	652.9
6	Influenza and pneumonia (J09-J18)	23,281	3.2	413.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17,249	2.4	306.3
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	16,689	2.3	296.4
9	Diabetes mellitus (E10-E14)	15,168	2.1	269.4
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	11,165	1.5	198.3
11	Septicemia (A40-A41)	9,788	1.3	173.8
12	Parkinson's disease (G20-G21)	8,429	1.1	149.7
13	Pneumonitis due to solids and liquids (J69)	7,873	1.1	139.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,656	0.6	82.7
15	Atherosclerosis (I70)	4,161	0.6	73.9
...	All other causes (Residual)	151,065	20.6	2,682.9
All races, male, all ages				
...	All causes	1,217,379	100.0	803.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	307,225	25.2	202.9
2	Malignant neoplasms (C00-C97)	296,763	24.4	195.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	75,022	6.2	49.5
4	Chronic lower respiratory diseases (J40-J47)	65,119	5.3	43.0
5	Cerebrovascular diseases (I60-I69)	52,073	4.3	34.4
6	Diabetes mellitus (E10-E14)	35,054	2.9	23.1
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	29,089	2.4	19.2
8	Influenza and pneumonia (J09-J18)	25,128	2.1	16.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23,930	2.0	15.8
10	Alzheimer's disease (G30)	23,900	2.0	15.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	19,804	1.6	13.1
12	Septicemia (A40-A41)	16,371	1.3	10.8
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	13,126	1.1	8.7
14	Parkinson's disease (G20-G21)	12,088	1.0	8.0
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10,461	0.9	6.9
...	All other causes (Residual)	212,226	17.4	140.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 1-4 years				
...	All causes	2,495	100.0	28.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	925	37.1	10.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	230	9.2	2.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	188	7.5	2.2
4	Malignant neoplasms (C00-C97)	180	7.2	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	77	3.1	0.9
6	Influenza and pneumonia (J09-J18)	64	2.6	0.7
7	Septicemia (A40-A41)	40	1.6	0.5
8	Chronic lower respiratory diseases (J40-J47)	36	1.4	0.4
9	Certain conditions originating in the perinatal period (P00-P96)	33	1.3	0.4
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	30	1.2	0.3
11	Cerebrovascular diseases (I60-I69)	24	1.0	0.3
12	Anemias (D50-D64)	15	0.6	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	12	0.5	*
14	Meningitis (G00,G03)	11	0.4	*
15	Meningococcal infection (A39)	9	0.4	*
...	All other causes (Residual)	621	24.9	7.1
All races, male, 5-14 years				
...	All causes	3,248	100.0	15.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,052	32.4	5.1
2	Malignant neoplasms (C00-C97)	488	15.0	2.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	206	6.3	1.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	196	6.0	0.9
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	175	5.4	0.8
6	Influenza and pneumonia (J09-J18)	114	3.5	0.5
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	112	3.4	0.5
8	Chronic lower respiratory diseases (J40-J47)	77	2.4	0.4
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	51	1.6	0.2
10	Cerebrovascular diseases (I60-I69)	45	1.4	0.2
11	Septicemia (A40-A41)	27	0.8	0.1
12	Anemias (D50-D64)	14	0.4	*
13	Diabetes mellitus (E10-E14)	12	0.4	*
13	Meningitis (G00,G03)	12	0.4	*
15	Certain conditions originating in the perinatal period (P00-P96)	11	0.3	*
...	All other causes (Residual)	656	20.2	3.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 15-24 years				
...	All causes	22,312	100.0	100.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,239	41.4	41.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4,205	18.8	19.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,595	16.1	16.2
4	Malignant neoplasms (C00-C97)	993	4.5	4.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	687	3.1	3.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	263	1.2	1.2
7	Influenza and pneumonia (J09-J18)	224	1.0	1.0
8	Cerebrovascular diseases (I60-I69)	108	0.5	0.5
8	Chronic lower respiratory diseases (J40-J47)	108	0.5	0.5
10	Human immunodeficiency virus (HIV) disease (B20-B24)	90	0.4	0.4
10	Diabetes mellitus (E10-E14)	90	0.4	0.4
12	Septicemia (A40-A41)	81	0.4	0.4
13	Legal intervention (Y35,Y89.0)	71	0.3	0.3
14	Anemias (D50-D64)	52	0.2	0.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	47	0.2	0.2
...	All other causes (Residual)	2,459	11.0	11.1
All races, male, 25-34 years				
...	All causes	29,199	100.0	137.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,532	36.1	49.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,293	14.7	20.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,481	11.9	16.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,164	7.4	10.2
5	Malignant neoplasms (C00-C97)	1,807	6.2	8.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	561	1.9	2.6
7	Influenza and pneumonia (J09-J18)	422	1.4	2.0
8	Diabetes mellitus (E10-E14)	353	1.2	1.7
9	Chronic liver disease and cirrhosis (K70,K73-K74)	305	1.0	1.4
10	Cerebrovascular diseases (I60-I69)	298	1.0	1.4
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	238	0.8	1.1
12	Septicemia (A40-A41)	197	0.7	0.9
13	Chronic lower respiratory diseases (J40-J47)	143	0.5	0.7
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	141	0.5	0.7
15	Legal intervention (Y35,Y89.0)	112	0.4	0.5
...	All other causes (Residual)	4,152	14.2	19.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 35-44 years				
...	All causes	46,555	100.0	223.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,649	22.9	51.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,738	16.6	37.1
3	Malignant neoplasms (C00-C97)	5,398	11.6	25.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,142	11.0	24.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,122	4.6	10.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	1,689	3.6	8.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,648	3.5	7.9
8	Diabetes mellitus (E10-E14)	1,208	2.6	5.8
9	Cerebrovascular diseases (I60-I69)	1,044	2.2	5.0
10	Influenza and pneumonia (J09-J18)	732	1.6	3.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	467	1.0	2.2
12	Septicemia (A40-A41)	458	1.0	2.2
13	Chronic lower respiratory diseases (J40-J47)	355	0.8	1.7
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	278	0.6	1.3
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	277	0.6	1.3
...	All other causes (Residual)	7,350	15.8	35.2
All races, male, 45-54 years				
...	All causes	114,820	100.0	522.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	26,432	23.0	120.3
2	Malignant neoplasms (C00-C97)	25,447	22.2	115.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,454	11.7	61.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,469	5.6	29.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	5,757	5.0	26.2
6	Diabetes mellitus (E10-E14)	3,602	3.1	16.4
7	Cerebrovascular diseases (I60-I69)	3,385	2.9	15.4
8	Human immunodeficiency virus (HIV) disease (B20-B24)	2,480	2.2	11.3
9	Chronic lower respiratory diseases (J40-J47)	2,270	2.0	10.3
10	Viral hepatitis (B15-B19)	1,805	1.6	8.2
11	Influenza and pneumonia (J09-J18)	1,558	1.4	7.1
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,508	1.3	6.9
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,347	1.2	6.1
14	Septicemia (A40-A41)	1,314	1.1	6.0
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	892	0.8	4.1
...	All other causes (Residual)	17,100	14.9	77.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 55-64 years				
...	All causes	184,142	100.0	1,097.3
1	Malignant neoplasms (C00-C97)	59,058	32.1	351.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	46,802	25.4	278.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,960	4.9	53.4
4	Chronic lower respiratory diseases (J40-J47)	7,354	4.0	43.8
5	Diabetes mellitus (E10-E14)	6,837	3.7	40.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	6,474	3.5	38.6
7	Cerebrovascular diseases (I60-I69)	5,977	3.2	35.6
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,457	2.4	26.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,707	1.5	16.1
10	Septicemia (A40-A41)	2,445	1.3	14.6
11	Influenza and pneumonia (J09-J18)	2,330	1.3	13.9
12	Viral hepatitis (B15-B19)	2,228	1.2	13.3
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,527	0.8	9.1
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1,444	0.8	8.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	849	0.5	5.1
...	All other causes (Residual)	24,693	13.4	147.1
All races, male, 65-74 years				
...	All causes	225,794	100.0	2,353.6
1	Malignant neoplasms (C00-C97)	77,920	34.5	812.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	56,302	24.9	586.9
3	Chronic lower respiratory diseases (J40-J47)	15,717	7.0	163.8
4	Cerebrovascular diseases (I60-I69)	9,283	4.1	96.8
5	Diabetes mellitus (E10-E14)	8,295	3.7	86.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,610	2.5	58.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,360	1.9	45.4
8	Influenza and pneumonia (J09-J18)	3,465	1.5	36.1
9	Septicemia (A40-A41)	3,351	1.5	34.9
10	Chronic liver disease and cirrhosis (K70,K73-K74)	3,330	1.5	34.7
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,394	1.1	25.0
12	Alzheimer's disease (G30)	1,819	0.8	19.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,794	0.8	18.7
14	Parkinson's disease (G20-G21)	1,602	0.7	16.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,411	0.6	14.7
...	All other causes (Residual)	29,141	12.9	303.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 75-84 years				
...	All causes	311,032	100.0	5,709.6
1	Malignant neoplasms (C00-C97)	83,563	26.9	1,534.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	82,287	26.5	1,510.5
3	Chronic lower respiratory diseases (J40-J47)	23,745	7.6	435.9
4	Cerebrovascular diseases (I60-I69)	16,347	5.3	300.1
5	Diabetes mellitus (E10-E14)	9,057	2.9	166.3
6	Alzheimer's disease (G30)	8,659	2.8	159.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,506	2.4	137.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,273	2.3	133.5
9	Influenza and pneumonia (J09-J18)	7,038	2.3	129.2
10	Parkinson's disease (G20-G21)	5,653	1.8	103.8
11	Septicemia (A40-A41)	4,784	1.5	87.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,766	0.9	50.8
13	Pneumonitis due to solids and liquids (J69)	2,592	0.8	47.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,481	0.8	45.5
15	Aortic aneurysm and dissection (I71)	1,893	0.6	34.7
...	All other causes (Residual)	45,388	14.6	833.2
All races, male, 85 years and over				
...	All causes	262,782	100.0	14,740.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	84,387	32.1	4,733.7
2	Malignant neoplasms (C00-C97)	41,860	15.9	2,348.2
3	Cerebrovascular diseases (I60-I69)	15,472	5.9	867.9
4	Chronic lower respiratory diseases (J40-J47)	15,296	5.8	858.0
5	Alzheimer's disease (G30)	13,038	5.0	731.4
6	Influenza and pneumonia (J09-J18)	9,029	3.4	506.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,277	2.8	408.2
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,594	2.5	369.9
9	Diabetes mellitus (E10-E14)	5,597	2.1	314.0
10	Parkinson's disease (G20-G21)	4,485	1.7	251.6
11	Pneumonitis due to solids and liquids (J69)	3,695	1.4	207.3
12	Septicemia (A40-A41)	3,554	1.4	199.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,127	1.2	175.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,012	0.8	112.9
15	Atherosclerosis (I70)	1,301	0.5	73.0
...	All other causes (Residual)	46,058	17.5	2,583.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, all ages				
...	All causes	1,219,784	100.0	784.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	292,188	24.0	187.8
2	Malignant neoplasms (C00-C97)	270,865	22.2	174.1
3	Cerebrovascular diseases (I60-I69)	76,769	6.3	49.4
4	Chronic lower respiratory diseases (J40-J47)	72,234	5.9	46.4
5	Alzheimer's disease (G30)	55,103	4.5	35.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	42,999	3.5	27.6
7	Diabetes mellitus (E10-E14)	33,651	2.8	21.6
8	Influenza and pneumonia (J09-J18)	28,564	2.3	18.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	25,005	2.0	16.1
10	Septicemia (A40-A41)	19,268	1.6	12.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	15,273	1.3	9.8
12	Chronic liver disease and cirrhosis (K70,K73-K74)	10,754	0.9	6.9
13	Parkinson's disease (G20-G21)	8,477	0.7	5.4
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7,820	0.6	5.0
15	Pneumonitis due to solids and liquids (J69)	7,649	0.6	4.9
...	All other causes (Residual)	253,165	20.8	162.7
All races, female, 1-4 years				
...	All causes	1,955	100.0	23.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	541	27.7	6.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	234	12.0	2.8
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	188	9.6	2.3
4	Malignant neoplasms (C00-C97)	170	8.7	2.0
5	Influenza and pneumonia (J09-J18)	82	4.2	1.0
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	77	3.9	0.9
7	Septicemia (A40-A41)	31	1.6	0.4
8	Chronic lower respiratory diseases (J40-J47)	30	1.5	0.4
9	Certain conditions originating in the perinatal period (P00-P96)	25	1.3	0.3
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	1.2	0.3
10	Cerebrovascular diseases (I60-I69)	23	1.2	0.3
12	Acute bronchitis and bronchiolitis (J20-J21)	11	0.6	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.4	*
14	Meningitis (G00,G03)	7	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.3	*
...	All other causes (Residual)	499	25.5	6.0

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 5-14 years				
...	All causes	2,403	100.0	12.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	637	26.5	3.2
2	Malignant neoplasms (C00-C97)	408	17.0	2.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	158	6.6	0.8
4	Influenza and pneumonia (J09-J18)	114	4.7	0.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	109	4.5	0.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	105	4.4	0.5
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	90	3.7	0.5
8	Chronic lower respiratory diseases (J40-J47)	46	1.9	0.2
9	Septicemia (A40-A41)	43	1.8	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	34	1.4	0.2
11	Cerebrovascular diseases (I60-I69)	29	1.2	0.1
12	Anemias (D50-D64)	21	0.9	0.1
13	Diabetes mellitus (E10-E14)	17	0.7	*
14	Certain conditions originating in the perinatal period (P00-P96)	15	0.6	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	0.5	*
...	All other causes (Residual)	566	23.6	2.9
All races, female, 15-24 years				
...	All causes	8,104	100.0	38.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,219	39.7	15.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	776	9.6	3.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	657	8.1	3.1
4	Malignant neoplasms (C00-C97)	643	7.9	3.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	348	4.3	1.7
6	Pregnancy, childbirth and the puerperium (O00-O99)	227	2.8	1.1
7	Influenza and pneumonia (J09-J18)	194	2.4	0.9
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	194	2.4	0.9
9	Cerebrovascular diseases (I60-I69)	85	1.0	0.4
10	Chronic lower respiratory diseases (J40-J47)	79	1.0	0.4
11	Diabetes mellitus (E10-E14)	77	1.0	0.4
12	Septicemia (A40-A41)	63	0.8	0.3
13	Human immunodeficiency virus (HIV) disease (B20-B24)	59	0.7	0.3
14	Anemias (D50-D64)	41	0.5	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	41	0.5	0.2
...	All other causes (Residual)	1,401	17.3	6.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 25-34 years				
...	All causes	13,303	100.0	65.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,530	26.5	17.4
2	Malignant neoplasms (C00-C97)	1,852	13.9	9.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,027	7.7	5.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,010	7.6	5.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	741	5.6	3.6
6	Pregnancy, childbirth and the puerperium (O00-O99)	403	3.0	2.0
7	Influenza and pneumonia (J09-J18)	385	2.9	1.9
8	Human immunodeficiency virus (HIV) disease (B20-B24)	320	2.4	1.6
9	Diabetes mellitus (E10-E14)	251	1.9	1.2
10	Cerebrovascular diseases (I60-I69)	239	1.8	1.2
11	Septicemia (A40-A41)	186	1.4	0.9
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	171	1.3	0.8
13	Chronic liver disease and cirrhosis (K70,K73-K74)	154	1.2	0.8
14	Chronic lower respiratory diseases (J40-J47)	140	1.1	0.7
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	126	0.9	0.6
...	All other causes (Residual)	2,768	20.8	13.6
All races, female, 35-44 years				
...	All causes	28,110	100.0	136.0
1	Malignant neoplasms (C00-C97)	7,121	25.3	34.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,453	15.8	21.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,343	11.9	16.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,535	5.5	7.4
5	Cerebrovascular diseases (I60-I69)	872	3.1	4.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	833	3.0	4.0
7	Human immunodeficiency virus (HIV) disease (B20-B24)	736	2.6	3.6
8	Diabetes mellitus (E10-E14)	664	2.4	3.2
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	640	2.3	3.1
10	Influenza and pneumonia (J09-J18)	582	2.1	2.8
11	Septicemia (A40-A41)	467	1.7	2.3
12	Chronic lower respiratory diseases (J40-J47)	388	1.4	1.9
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	362	1.3	1.8
14	Pregnancy, childbirth and the puerperium (O00-O99)	228	0.8	1.1
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	185	0.7	0.9
...	All other causes (Residual)	5,701	20.3	27.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 45-54 years				
...	All causes	72,748	100.0	321.6
1	Malignant neoplasms (C00-C97)	25,169	34.6	111.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,495	14.4	46.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,520	9.0	28.8
4	Cerebrovascular diseases (I60-I69)	2,778	3.8	12.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,620	3.6	11.6
6	Chronic lower respiratory diseases (J40-J47)	2,394	3.3	10.6
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,129	2.9	9.4
8	Diabetes mellitus (E10-E14)	2,123	2.9	9.4
9	Influenza and pneumonia (J09-J18)	1,360	1.9	6.0
10	Septicemia (A40-A41)	1,119	1.5	4.9
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,005	1.4	4.4
12	Human immunodeficiency virus (HIV) disease (B20-B24)	908	1.2	4.0
13	Viral hepatitis (B15-B19)	797	1.1	3.5
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	558	0.8	2.5
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	498	0.7	2.2
...	All other causes (Residual)	12,275	16.9	54.3
All races, female, 55-64 years				
...	All causes	119,165	100.0	661.8
1	Malignant neoplasms (C00-C97)	47,771	40.1	265.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,459	17.2	113.6
3	Chronic lower respiratory diseases (J40-J47)	6,806	5.7	37.8
4	Cerebrovascular diseases (I60-I69)	4,546	3.8	25.2
5	Diabetes mellitus (E10-E14)	4,524	3.8	25.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,973	3.3	22.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,680	2.2	14.9
8	Septicemia (A40-A41)	2,183	1.8	12.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,085	1.7	11.6
10	Influenza and pneumonia (J09-J18)	1,809	1.5	10.0
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,351	1.1	7.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	976	0.8	5.4
13	Viral hepatitis (B15-B19)	835	0.7	4.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	627	0.5	3.5
15	Human immunodeficiency virus (HIV) disease (B20-B24)	410	0.3	2.3
...	All other causes (Residual)	18,130	15.2	100.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 65-74 years				
...	All causes	175,238	100.0	1,564.8
1	Malignant neoplasms (C00-C97)	63,951	36.5	571.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	33,463	19.1	298.8
3	Chronic lower respiratory diseases (J40-J47)	15,597	8.9	139.3
4	Cerebrovascular diseases (I60-I69)	8,295	4.7	74.1
5	Diabetes mellitus (E10-E14)	6,490	3.7	58.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,863	2.2	34.5
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,330	1.9	29.7
8	Septicemia (A40-A41)	3,311	1.9	29.6
9	Influenza and pneumonia (J09-J18)	2,794	1.6	24.9
10	Alzheimer's disease (G30)	2,290	1.3	20.4
11	Chronic liver disease and cirrhosis (K70,K73-K74)	2,053	1.2	18.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,661	0.9	14.8
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,041	0.6	9.3
14	Parkinson's disease (G20-G21)	778	0.4	6.9
15	Aortic aneurysm and dissection (I71)	753	0.4	6.7
...	All other causes (Residual)	25,568	14.6	228.3
All races, female, 75-84 years				
...	All causes	316,695	100.0	4,112.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	75,399	23.8	979.2
2	Malignant neoplasms (C00-C97)	74,397	23.5	966.1
3	Chronic lower respiratory diseases (J40-J47)	25,276	8.0	328.2
4	Cerebrovascular diseases (I60-I69)	22,060	7.0	286.5
5	Alzheimer's disease (G30)	14,665	4.6	190.4
6	Diabetes mellitus (E10-E14)	9,934	3.1	129.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,487	2.4	97.2
8	Influenza and pneumonia (J09-J18)	6,891	2.2	89.5
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,209	2.0	80.6
10	Septicemia (A40-A41)	5,529	1.7	71.8
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,875	1.2	50.3
12	Parkinson's disease (G20-G21)	3,566	1.1	46.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,080	0.7	27.0
14	Pneumonitis due to solids and liquids (J69)	2,059	0.7	26.7
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,738	0.5	22.6
...	All other causes (Residual)	55,530	17.5	721.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 85 years and over				
...	All causes	470,396	100.0	12,224.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	147,312	31.3	3,828.3
2	Malignant neoplasms (C00-C97)	49,344	10.5	1,282.3
3	Cerebrovascular diseases (I60-I69)	37,781	8.0	981.8
4	Alzheimer's disease (G30)	37,697	8.0	979.7
5	Chronic lower respiratory diseases (J40-J47)	21,467	4.6	557.9
6	Influenza and pneumonia (J09-J18)	14,252	3.0	370.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,095	2.1	262.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9,972	2.1	259.1
9	Diabetes mellitus (E10-E14)	9,571	2.0	248.7
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	8,038	1.7	208.9
11	Septicemia (A40-A41)	6,234	1.3	162.0
12	Pneumonitis due to solids and liquids (J69)	4,178	0.9	108.6
13	Parkinson's disease (G20-G21)	3,944	0.8	102.5
14	Atherosclerosis (I70)	2,860	0.6	74.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,644	0.6	68.7
...	All other causes (Residual)	105,007	22.3	2,728.9
White, both sexes, all ages				
...	All causes	2,086,355	100.0	844.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	515,587	24.7	208.8
2	Malignant neoplasms (C00-C97)	486,995	23.3	197.2
3	Chronic lower respiratory diseases (J40-J47)	126,733	6.1	51.3
4	Cerebrovascular diseases (I60-I69)	108,764	5.2	44.0
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	102,170	4.9	41.4
6	Alzheimer's disease (G30)	72,696	3.5	29.4
7	Diabetes mellitus (E10-E14)	54,113	2.6	21.9
8	Influenza and pneumonia (J09-J18)	46,432	2.2	18.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	38,758	1.9	15.7
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	33,425	1.6	13.5
11	Septicemia (A40-A41)	28,609	1.4	11.6
12	Chronic liver disease and cirrhosis (K70,K73-K74)	26,820	1.3	10.9
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	20,052	1.0	8.1
14	Parkinson's disease (G20-G21)	19,401	0.9	7.9
15	Pneumonitis due to solids and liquids (J69)	14,133	0.7	5.7
...	All other causes (Residual)	391,667	18.8	158.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 1-4 years				
...	All causes	3,141	100.0	24.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,082	34.4	8.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	350	11.1	2.7
3	Malignant neoplasms (C00-C97)	274	8.7	2.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	211	6.7	1.6
5	Influenza and pneumonia (J09-J18)	108	3.4	0.8
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	96	3.1	0.7
7	Septicemia (A40-A41)	47	1.5	0.4
8	Cerebrovascular diseases (I60-I69)	38	1.2	0.3
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	37	1.2	0.3
10	Certain conditions originating in the perinatal period (P00-P96)	36	1.1	0.3
11	Chronic lower respiratory diseases (J40-J47)	32	1.0	0.2
12	Meningitis (G00,G03)	15	0.5	*
13	Anemias (D50-D64)	12	0.4	*
14	Acute bronchitis and bronchiolitis (J20-J21)	9	0.3	*
15	Meningococcal infection (A39)	8	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.3	*
...	All other causes (Residual)	778	24.8	6.0
White, both sexes, 5-14 years				
...	All causes	4,005	100.0	12.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,219	30.4	3.9
2	Malignant neoplasms (C00-C97)	703	17.6	2.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	259	6.5	0.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	201	5.0	0.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	185	4.6	0.6
6	Influenza and pneumonia (J09-J18)	165	4.1	0.5
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	132	3.3	0.4
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	63	1.6	0.2
9	Cerebrovascular diseases (I60-I69)	53	1.3	0.2
10	Septicemia (A40-A41)	43	1.1	0.1
11	Chronic lower respiratory diseases (J40-J47)	40	1.0	0.1
12	Diabetes mellitus (E10-E14)	21	0.5	0.1
13	Certain conditions originating in the perinatal period (P00-P96)	20	0.5	0.1
14	Anemias (D50-D64)	16	0.4	*
15	Meningitis (G00,G03)	15	0.4	*
...	All other causes (Residual)	870	21.7	2.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 15-24 years				
...	All causes	22,241	100.0	66.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,526	47.3	31.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,619	16.3	10.8
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,902	8.6	5.7
4	Malignant neoplasms (C00-C97)	1,256	5.6	3.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	643	2.9	1.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	340	1.5	1.0
7	Influenza and pneumonia (J09-J18)	308	1.4	0.9
8	Cerebrovascular diseases (I60-I69)	137	0.6	0.4
9	Pregnancy, childbirth and the puerperium (O00-O99)	133	0.6	0.4
10	Diabetes mellitus (E10-E14)	113	0.5	0.3
11	Septicemia (A40-A41)	108	0.5	0.3
12	Chronic lower respiratory diseases (J40-J47)	92	0.4	0.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	52	0.2	0.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	52	0.2	0.2
15	Human immunodeficiency virus (HIV) disease (B20-B24)	43	0.2	0.1
...	All other causes (Residual)	2,917	13.1	8.7
White, both sexes, 25-34 years				
...	All causes	31,489	100.0	97.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,912	37.8	36.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,569	14.5	14.1
3	Malignant neoplasms (C00-C97)	2,822	9.0	8.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,072	6.6	6.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,765	5.6	5.4
6	Influenza and pneumonia (J09-J18)	610	1.9	1.9
7	Diabetes mellitus (E10-E14)	394	1.3	1.2
8	Chronic liver disease and cirrhosis (K70,K73-K74)	365	1.2	1.1
9	Cerebrovascular diseases (I60-I69)	360	1.1	1.1
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	313	1.0	1.0
11	Human immunodeficiency virus (HIV) disease (B20-B24)	261	0.8	0.8
12	Septicemia (A40-A41)	250	0.8	0.8
12	Pregnancy, childbirth and the puerperium (O00-O99)	250	0.8	0.8
14	Chronic lower respiratory diseases (J40-J47)	170	0.5	0.5
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	156	0.5	0.5
...	All other causes (Residual)	5,220	16.6	16.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 35-44 years				
...	All causes	56,337	100.0	171.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,774	22.7	38.8
2	Malignant neoplasms (C00-C97)	9,692	17.2	29.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,765	13.8	23.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,016	10.7	18.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,094	3.7	6.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,456	2.6	4.4
7	Diabetes mellitus (E10-E14)	1,260	2.2	3.8
8	Cerebrovascular diseases (I60-I69)	1,243	2.2	3.8
9	Human immunodeficiency virus (HIV) disease (B20-B24)	998	1.8	3.0
10	Influenza and pneumonia (J09-J18)	985	1.7	3.0
11	Septicemia (A40-A41)	613	1.1	1.9
12	Chronic lower respiratory diseases (J40-J47)	539	1.0	1.6
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	475	0.8	1.4
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	395	0.7	1.2
15	Viral hepatitis (B15-B19)	332	0.6	1.0
...	All other causes (Residual)	9,700	17.2	29.4
White, both sexes, 45-54 years				
...	All causes	146,108	100.0	399.8
1	Malignant neoplasms (C00-C97)	40,200	27.5	110.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	27,693	19.0	75.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17,002	11.6	46.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7,987	5.5	21.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	7,215	4.9	19.7
6	Diabetes mellitus (E10-E14)	4,166	2.9	11.4
7	Cerebrovascular diseases (I60-I69)	4,055	2.8	11.1
8	Chronic lower respiratory diseases (J40-J47)	3,870	2.6	10.6
9	Influenza and pneumonia (J09-J18)	2,268	1.6	6.2
10	Viral hepatitis (B15-B19)	2,150	1.5	5.9
11	Septicemia (A40-A41)	1,732	1.2	4.7
12	Human immunodeficiency virus (HIV) disease (B20-B24)	1,470	1.0	4.0
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,400	1.0	3.8
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,251	0.9	3.4
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	788	0.5	2.2
...	All other causes (Residual)	22,861	15.6	62.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 55-64 years				
...	All causes	243,809	100.0	832.4
1	Malignant neoplasms (C00-C97)	87,770	36.0	299.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	52,502	21.5	179.3
3	Chronic lower respiratory diseases (J40-J47)	12,601	5.2	43.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,949	4.5	37.4
5	Diabetes mellitus (E10-E14)	8,453	3.5	28.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	7,942	3.3	27.1
7	Cerebrovascular diseases (I60-I69)	7,424	3.0	25.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,464	2.2	18.7
9	Septicemia (A40-A41)	3,438	1.4	11.7
10	Influenza and pneumonia (J09-J18)	3,342	1.4	11.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,218	1.3	11.0
12	Viral hepatitis (B15-B19)	2,453	1.0	8.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,599	0.7	5.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,201	0.5	4.1
15	Aortic aneurysm and dissection (I71)	964	0.4	3.3
...	All other causes (Residual)	34,489	14.1	117.8
White, both sexes, 65-74 years				
...	All causes	338,048	100.0	1,892.9
1	Malignant neoplasms (C00-C97)	121,630	36.0	681.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	73,724	21.8	412.8
3	Chronic lower respiratory diseases (J40-J47)	28,834	8.5	161.5
4	Cerebrovascular diseases (I60-I69)	13,875	4.1	77.7
5	Diabetes mellitus (E10-E14)	11,407	3.4	63.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,827	2.3	43.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,002	1.8	33.6
8	Septicemia (A40-A41)	5,205	1.5	29.1
9	Influenza and pneumonia (J09-J18)	5,202	1.5	29.1
10	Chronic liver disease and cirrhosis (K70,K73-K74)	4,843	1.4	27.1
11	Alzheimer's disease (G30)	3,673	1.1	20.6
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,760	0.8	15.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,457	0.7	13.8
14	Parkinson's disease (G20-G21)	2,190	0.6	12.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,159	0.6	12.1
...	All other causes (Residual)	46,260	13.7	259.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 75-84 years				
...	All causes	555,168	100.0	4,818.6
1	Malignant neoplasms (C00-C97)	140,078	25.2	1,215.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	138,425	24.9	1,201.5
3	Chronic lower respiratory diseases (J40-J47)	45,982	8.3	399.1
4	Cerebrovascular diseases (I60-I69)	33,143	6.0	287.7
5	Alzheimer's disease (G30)	21,271	3.8	184.6
6	Diabetes mellitus (E10-E14)	15,475	2.8	134.3
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,376	2.2	107.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12,274	2.2	106.5
9	Influenza and pneumonia (J09-J18)	12,201	2.2	105.9
10	Parkinson's disease (G20-G21)	8,708	1.6	75.6
11	Septicemia (A40-A41)	8,630	1.6	74.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,329	1.0	46.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,155	0.7	36.1
14	Pneumonitis due to solids and liquids (J69)	4,132	0.7	35.9
15	Chronic liver disease and cirrhosis (K70,K73-K74)	3,282	0.6	28.5
...	All other causes (Residual)	89,707	16.2	778.6
White, both sexes, 85 years and over				
...	All causes	668,976	100.0	13,184.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	212,268	31.7	4,183.3
2	Malignant neoplasms (C00-C97)	82,506	12.3	1,626.0
3	Cerebrovascular diseases (I60-I69)	48,335	7.2	952.6
4	Alzheimer's disease (G30)	47,002	7.0	926.3
5	Chronic lower respiratory diseases (J40-J47)	34,555	5.2	681.0
6	Influenza and pneumonia (J09-J18)	21,092	3.2	415.7
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15,726	2.4	309.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15,085	2.3	297.3
9	Diabetes mellitus (E10-E14)	12,822	1.9	252.7
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9,595	1.4	189.1
11	Septicemia (A40-A41)	8,420	1.3	165.9
12	Parkinson's disease (G20-G21)	8,029	1.2	158.2
13	Pneumonitis due to solids and liquids (J69)	7,189	1.1	141.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,319	0.6	85.1
15	Atherosclerosis (I70)	3,895	0.6	76.8
...	All other causes (Residual)	138,138	20.6	2,722.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, male, all ages				
...	All causes	1,037,475	100.0	846.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	263,976	25.4	215.4
2	Malignant neoplasms (C00-C97)	255,042	24.6	208.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	64,501	6.2	52.6
4	Chronic lower respiratory diseases (J40-J47)	59,465	5.7	48.5
5	Cerebrovascular diseases (I60-I69)	43,190	4.2	35.2
6	Diabetes mellitus (E10-E14)	28,205	2.7	23.0
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	26,426	2.5	21.6
8	Alzheimer's disease (G30)	22,066	2.1	18.0
9	Influenza and pneumonia (J09-J18)	21,614	2.1	17.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19,312	1.9	15.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	17,475	1.7	14.3
12	Septicemia (A40-A41)	13,206	1.3	10.8
13	Parkinson's disease (G20-G21)	11,407	1.1	9.3
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	8,034	0.8	6.6
15	Pneumonitis due to solids and liquids (J69)	7,413	0.7	6.0
...	All other causes (Residual)	176,143	17.0	143.7
White, male, 1-4 years				
...	All causes	1,780	100.0	26.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	698	39.2	10.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	169	9.5	2.5
3	Malignant neoplasms (C00-C97)	147	8.3	2.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	100	5.6	1.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	52	2.9	0.8
6	Influenza and pneumonia (J09-J18)	47	2.6	0.7
7	Septicemia (A40-A41)	29	1.6	0.4
8	Certain conditions originating in the perinatal period (P00-P96)	21	1.2	0.3
9	Cerebrovascular diseases (I60-I69)	20	1.1	0.3
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	1.1	*
11	Chronic lower respiratory diseases (J40-J47)	18	1.0	*
12	Anemias (D50-D64)	9	0.5	*
12	Meningitis (G00,G03)	9	0.5	*
14	Meningococcal infection (A39)	7	0.4	*
15	Acute bronchitis and bronchiolitis (J20-J21)	4	0.2	*
...	All other causes (Residual)	431	24.2	6.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 5-14 years				
...	All causes	2,314	100.0	14.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	761	32.9	4.7
2	Malignant neoplasms (C00-C97)	390	16.9	2.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	147	6.4	0.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	131	5.7	0.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	115	5.0	0.7
6	Influenza and pneumonia (J09-J18)	84	3.6	0.5
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	66	2.9	0.4
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	36	1.6	0.2
9	Cerebrovascular diseases (I60-I69)	32	1.4	0.2
10	Chronic lower respiratory diseases (J40-J47)	28	1.2	0.2
11	Septicemia (A40-A41)	14	0.6	*
12	Diabetes mellitus (E10-E14)	10	0.4	*
13	Meningitis (G00,G03)	9	0.4	*
14	Certain conditions originating in the perinatal period (P00-P96)	8	0.3	*
15	Anemias (D50-D64)	7	0.3	*
...	All other causes (Residual)	476	20.6	2.9
White, male, 15-24 years				
...	All causes	16,243	100.0	94.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,862	48.4	45.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,012	18.5	17.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,563	9.6	9.1
4	Malignant neoplasms (C00-C97)	776	4.8	4.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	429	2.6	2.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	194	1.2	1.1
7	Influenza and pneumonia (J09-J18)	169	1.0	1.0
8	Cerebrovascular diseases (I60-I69)	79	0.5	0.5
9	Septicemia (A40-A41)	66	0.4	0.4
10	Diabetes mellitus (E10-E14)	62	0.4	0.4
11	Chronic lower respiratory diseases (J40-J47)	50	0.3	0.3
12	Legal intervention (Y35,Y89.0)	40	0.2	0.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	35	0.2	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	31	0.2	0.2
15	Pneumonitis due to solids and liquids (J69)	26	0.2	0.2
...	All other causes (Residual)	1,849	11.4	10.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 25-34 years				
...	All causes	21,780	100.0	130.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,930	41.0	53.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,715	17.1	22.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,466	6.7	8.7
4	Malignant neoplasms (C00-C97)	1,423	6.5	8.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,334	6.1	8.0
6	Influenza and pneumonia (J09-J18)	335	1.5	2.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	249	1.1	1.5
8	Diabetes mellitus (E10-E14)	228	1.0	1.4
9	Human immunodeficiency virus (HIV) disease (B20-B24)	203	0.9	1.2
9	Cerebrovascular diseases (I60-I69)	203	0.9	1.2
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	185	0.8	1.1
12	Septicemia (A40-A41)	131	0.6	0.8
13	Chronic lower respiratory diseases (J40-J47)	81	0.4	0.5
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	81	0.4	0.5
15	Legal intervention (Y35,Y89.0)	68	0.3	0.4
...	All other causes (Residual)	3,148	14.5	18.8
White, male, 35-44 years				
...	All causes	35,627	100.0	212.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,994	25.2	53.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,549	15.6	33.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,627	13.0	27.6
4	Malignant neoplasms (C00-C97)	4,260	12.0	25.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,410	4.0	8.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,037	2.9	6.2
7	Diabetes mellitus (E10-E14)	819	2.3	4.9
8	Human immunodeficiency virus (HIV) disease (B20-B24)	802	2.3	4.8
9	Cerebrovascular diseases (I60-I69)	684	1.9	4.1
10	Influenza and pneumonia (J09-J18)	565	1.6	3.4
11	Septicemia (A40-A41)	306	0.9	1.8
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	285	0.8	1.7
13	Chronic lower respiratory diseases (J40-J47)	235	0.7	1.4
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	234	0.7	1.4
15	Viral hepatitis (B15-B19)	218	0.6	1.3
...	All other causes (Residual)	5,602	15.7	33.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 45-54 years				
...	All causes	90,871	100.0	498.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,454	22.5	112.3
2	Malignant neoplasms (C00-C97)	20,344	22.4	111.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,434	12.6	62.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,031	6.6	33.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	5,013	5.5	27.5
6	Diabetes mellitus (E10-E14)	2,676	2.9	14.7
7	Cerebrovascular diseases (I60-I69)	2,291	2.5	12.6
8	Chronic lower respiratory diseases (J40-J47)	1,896	2.1	10.4
9	Viral hepatitis (B15-B19)	1,517	1.7	8.3
10	Influenza and pneumonia (J09-J18)	1,222	1.3	6.7
11	Human immunodeficiency virus (HIV) disease (B20-B24)	1,206	1.3	6.6
12	Septicemia (A40-A41)	961	1.1	5.3
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	879	1.0	4.8
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	835	0.9	4.6
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	531	0.6	2.9
...	All other causes (Residual)	13,581	14.9	74.6
White, male, 55-64 years				
...	All causes	149,005	100.0	1,042.0
1	Malignant neoplasms (C00-C97)	48,484	32.5	339.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	37,375	25.1	261.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,563	5.1	52.9
4	Chronic lower respiratory diseases (J40-J47)	6,495	4.4	45.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	5,651	3.8	39.5
6	Diabetes mellitus (E10-E14)	5,241	3.5	36.7
7	Cerebrovascular diseases (I60-I69)	4,229	2.8	29.6
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,191	2.8	29.3
9	Influenza and pneumonia (J09-J18)	1,878	1.3	13.1
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,864	1.3	13.0
11	Septicemia (A40-A41)	1,814	1.2	12.7
12	Viral hepatitis (B15-B19)	1,795	1.2	12.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,016	0.7	7.1
14	Aortic aneurysm and dissection (I71)	730	0.5	5.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	699	0.5	4.9
...	All other causes (Residual)	19,980	13.4	139.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 65-74 years				
...	All causes	191,494	100.0	2,299.3
1	Malignant neoplasms (C00-C97)	66,881	34.9	803.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	46,999	24.5	564.3
3	Chronic lower respiratory diseases (J40-J47)	14,343	7.5	172.2
4	Cerebrovascular diseases (I60-I69)	7,355	3.8	88.3
5	Diabetes mellitus (E10-E14)	6,599	3.4	79.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,905	2.6	58.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,286	1.7	39.5
8	Chronic liver disease and cirrhosis (K70,K73-K74)	3,012	1.6	36.2
9	Influenza and pneumonia (J09-J18)	2,865	1.5	34.4
10	Septicemia (A40-A41)	2,650	1.4	31.8
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,266	1.2	27.2
12	Alzheimer's disease (G30)	1,626	0.8	19.5
13	Parkinson's disease (G20-G21)	1,474	0.8	17.7
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,327	0.7	15.9
15	Aortic aneurysm and dissection (I71)	1,263	0.7	15.2
...	All other causes (Residual)	24,643	12.9	295.9
White, male, 75-84 years				
...	All causes	277,531	100.0	5,750.3
1	Malignant neoplasms (C00-C97)	74,413	26.8	1,541.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	73,348	26.4	1,519.7
3	Chronic lower respiratory diseases (J40-J47)	22,069	8.0	457.3
4	Cerebrovascular diseases (I60-I69)	14,168	5.1	293.6
5	Alzheimer's disease (G30)	7,929	2.9	164.3
6	Diabetes mellitus (E10-E14)	7,665	2.8	158.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,675	2.4	138.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,326	2.3	131.1
9	Influenza and pneumonia (J09-J18)	6,163	2.2	127.7
10	Parkinson's disease (G20-G21)	5,356	1.9	111.0
11	Septicemia (A40-A41)	4,075	1.5	84.4
12	Pneumonitis due to solids and liquids (J69)	2,333	0.8	48.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,288	0.8	47.4
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,257	0.8	46.8
15	Aortic aneurysm and dissection (I71)	1,750	0.6	36.3
...	All other causes (Residual)	40,716	14.7	843.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 85 years and over				
...	All causes	241,261	100.0	14,949.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	78,091	32.4	4,838.7
2	Malignant neoplasms (C00-C97)	37,888	15.7	2,347.6
3	Chronic lower respiratory diseases (J40-J47)	14,238	5.9	882.2
4	Cerebrovascular diseases (I60-I69)	14,072	5.8	871.9
5	Alzheimer's disease (G30)	12,175	5.0	754.4
6	Influenza and pneumonia (J09-J18)	8,190	3.4	507.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,551	2.7	405.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,211	2.6	384.8
9	Diabetes mellitus (E10-E14)	4,903	2.0	303.8
10	Parkinson's disease (G20-G21)	4,271	1.8	264.6
11	Pneumonitis due to solids and liquids (J69)	3,399	1.4	210.6
12	Septicemia (A40-A41)	3,091	1.3	191.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,707	1.1	167.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,905	0.8	118.0
15	Atherosclerosis (I70)	1,232	0.5	76.3
...	All other causes (Residual)	42,337	17.5	2,623.3
White, female, all ages				
...	All causes	1,048,880	100.0	843.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	251,611	24.0	202.2
2	Malignant neoplasms (C00-C97)	231,953	22.1	186.4
3	Chronic lower respiratory diseases (J40-J47)	67,268	6.4	54.1
4	Cerebrovascular diseases (I60-I69)	65,574	6.3	52.7
5	Alzheimer's disease (G30)	50,630	4.8	40.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	37,669	3.6	30.3
7	Diabetes mellitus (E10-E14)	25,908	2.5	20.8
8	Influenza and pneumonia (J09-J18)	24,818	2.4	19.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19,446	1.9	15.6
10	Septicemia (A40-A41)	15,403	1.5	12.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	12,018	1.1	9.7
12	Chronic liver disease and cirrhosis (K70,K73-K74)	9,345	0.9	7.5
13	Parkinson's disease (G20-G21)	7,994	0.8	6.4
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,999	0.7	5.6
15	Pneumonitis due to solids and liquids (J69)	6,720	0.6	5.4
...	All other causes (Residual)	215,524	20.5	173.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 1-4 years				
...	All causes	1,361	100.0	21.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	384	28.2	6.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	181	13.3	2.8
3	Malignant neoplasms (C00-C97)	127	9.3	2.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	111	8.2	1.7
5	Influenza and pneumonia (J09-J18)	61	4.5	1.0
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	44	3.2	0.7
7	Septicemia (A40-A41)	18	1.3	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	1.3	*
7	Cerebrovascular diseases (I60-I69)	18	1.3	*
10	Certain conditions originating in the perinatal period (P00-P96)	15	1.1	*
11	Chronic lower respiratory diseases (J40-J47)	14	1.0	*
12	Meningitis (G00,G03)	6	0.4	*
13	Acute bronchitis and bronchiolitis (J20-J21)	5	0.4	*
13	Pneumonitis due to solids and liquids (J69)	5	0.4	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.4	*
...	All other causes (Residual)	349	25.6	5.5
White, female, 5-14 years				
...	All causes	1,691	100.0	11.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	458	27.1	3.0
2	Malignant neoplasms (C00-C97)	313	18.5	2.0
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	112	6.6	0.7
4	Influenza and pneumonia (J09-J18)	81	4.8	0.5
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	70	4.1	0.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	70	4.1	0.5
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	66	3.9	0.4
8	Septicemia (A40-A41)	29	1.7	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	27	1.6	0.2
10	Cerebrovascular diseases (I60-I69)	21	1.2	0.1
11	Chronic lower respiratory diseases (J40-J47)	12	0.7	*
11	Certain conditions originating in the perinatal period (P00-P96)	12	0.7	*
13	Diabetes mellitus (E10-E14)	11	0.7	*
14	Anemias (D50-D64)	9	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.4	*
...	All other causes (Residual)	393	23.2	2.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 15-24 years				
...	All causes	5,998	100.0	37.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,664	44.4	16.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	607	10.1	3.7
3	Malignant neoplasms (C00-C97)	480	8.0	3.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	339	5.7	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	214	3.6	1.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	146	2.4	0.9
7	Influenza and pneumonia (J09-J18)	139	2.3	0.9
8	Pregnancy, childbirth and the puerperium (O00-O99)	133	2.2	0.8
9	Cerebrovascular diseases (I60-I69)	58	1.0	0.4
10	Diabetes mellitus (E10-E14)	51	0.9	0.3
11	Septicemia (A40-A41)	42	0.7	0.3
11	Chronic lower respiratory diseases (J40-J47)	42	0.7	0.3
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21	0.4	0.1
14	Human immunodeficiency virus (HIV) disease (B20-B24)	18	0.3	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	0.3	*
...	All other causes (Residual)	1,027	17.1	6.3
White, female, 25-34 years				
...	All causes	9,709	100.0	61.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,982	30.7	19.0
2	Malignant neoplasms (C00-C97)	1,399	14.4	8.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	854	8.8	5.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	606	6.2	3.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	431	4.4	2.7
6	Influenza and pneumonia (J09-J18)	275	2.8	1.7
7	Pregnancy, childbirth and the puerperium (O00-O99)	250	2.6	1.6
8	Diabetes mellitus (E10-E14)	166	1.7	1.1
9	Cerebrovascular diseases (I60-I69)	157	1.6	1.0
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	128	1.3	0.8
11	Septicemia (A40-A41)	119	1.2	0.8
12	Chronic liver disease and cirrhosis (K70,K73-K74)	116	1.2	0.7
13	Chronic lower respiratory diseases (J40-J47)	89	0.9	0.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	75	0.8	0.5
15	Human immunodeficiency virus (HIV) disease (B20-B24)	58	0.6	0.4
...	All other causes (Residual)	2,004	20.6	12.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 35-44 years				
...	All causes	20,710	100.0	127.9
1	Malignant neoplasms (C00-C97)	5,432	26.2	33.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,780	18.3	23.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,216	10.7	13.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,389	6.7	8.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	684	3.3	4.2
6	Cerebrovascular diseases (I60-I69)	559	2.7	3.5
7	Diabetes mellitus (E10-E14)	441	2.1	2.7
8	Influenza and pneumonia (J09-J18)	420	2.0	2.6
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	419	2.0	2.6
10	Septicemia (A40-A41)	307	1.5	1.9
11	Chronic lower respiratory diseases (J40-J47)	304	1.5	1.9
12	Human immunodeficiency virus (HIV) disease (B20-B24)	196	0.9	1.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	190	0.9	1.2
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	161	0.8	1.0
15	Pregnancy, childbirth and the puerperium (O00-O99)	149	0.7	0.9
...	All other causes (Residual)	4,063	19.6	25.1
White, female, 45-54 years				
...	All causes	55,237	100.0	301.4
1	Malignant neoplasms (C00-C97)	19,856	35.9	108.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,239	13.1	39.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,568	10.1	30.4
4	Chronic liver disease and cirrhosis (K70,K73-K74)	2,202	4.0	12.0
5	Chronic lower respiratory diseases (J40-J47)	1,974	3.6	10.8
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,956	3.5	10.7
7	Cerebrovascular diseases (I60-I69)	1,764	3.2	9.6
8	Diabetes mellitus (E10-E14)	1,490	2.7	8.1
9	Influenza and pneumonia (J09-J18)	1,046	1.9	5.7
10	Septicemia (A40-A41)	771	1.4	4.2
11	Viral hepatitis (B15-B19)	633	1.1	3.5
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	565	1.0	3.1
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	372	0.7	2.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	277	0.5	1.5
15	Human immunodeficiency virus (HIV) disease (B20-B24)	264	0.5	1.4
...	All other causes (Residual)	9,260	16.8	50.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 55-64 years				
...	All causes	94,804	100.0	632.5
1	Malignant neoplasms (C00-C97)	39,286	41.4	262.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,127	16.0	100.9
3	Chronic lower respiratory diseases (J40-J47)	6,106	6.4	40.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,386	3.6	22.6
5	Diabetes mellitus (E10-E14)	3,212	3.4	21.4
6	Cerebrovascular diseases (I60-I69)	3,195	3.4	21.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,291	2.4	15.3
8	Septicemia (A40-A41)	1,624	1.7	10.8
9	Influenza and pneumonia (J09-J18)	1,464	1.5	9.8
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,354	1.4	9.0
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,273	1.3	8.5
12	Viral hepatitis (B15-B19)	658	0.7	4.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	583	0.6	3.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	502	0.5	3.3
15	Alzheimer's disease (G30)	356	0.4	2.4
...	All other causes (Residual)	14,387	15.2	96.0
White, female, 65-74 years				
...	All causes	146,554	100.0	1,537.7
1	Malignant neoplasms (C00-C97)	54,749	37.4	574.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	26,725	18.2	280.4
3	Chronic lower respiratory diseases (J40-J47)	14,491	9.9	152.0
4	Cerebrovascular diseases (I60-I69)	6,520	4.4	68.4
5	Diabetes mellitus (E10-E14)	4,808	3.3	50.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,922	2.0	30.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,716	1.9	28.5
8	Septicemia (A40-A41)	2,555	1.7	26.8
9	Influenza and pneumonia (J09-J18)	2,337	1.6	24.5
10	Alzheimer's disease (G30)	2,047	1.4	21.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1,831	1.2	19.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,130	0.8	11.9
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	899	0.6	9.4
14	Parkinson's disease (G20-G21)	716	0.5	7.5
15	Aortic aneurysm and dissection (I71)	656	0.4	6.9
...	All other causes (Residual)	21,452	14.6	225.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 75-84 years				
...	All causes	277,637	100.0	4,146.9
1	Malignant neoplasms (C00-C97)	65,665	23.7	980.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	65,077	23.4	972.0
3	Chronic lower respiratory diseases (J40-J47)	23,913	8.6	357.2
4	Cerebrovascular diseases (I60-I69)	18,975	6.8	283.4
5	Alzheimer's disease (G30)	13,342	4.8	199.3
6	Diabetes mellitus (E10-E14)	7,810	2.8	116.7
7	Influenza and pneumonia (J09-J18)	6,038	2.2	90.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,948	2.1	88.8
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,701	2.1	85.2
10	Septicemia (A40-A41)	4,555	1.6	68.0
11	Parkinson's disease (G20-G21)	3,352	1.2	50.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,072	1.1	45.9
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,867	0.7	27.9
14	Pneumonitis due to solids and liquids (J69)	1,799	0.6	26.9
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,591	0.6	23.8
...	All other causes (Residual)	48,932	17.6	730.9
White, female, 85 years and over				
...	All causes	427,715	100.0	12,360.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	134,177	31.4	3,877.6
2	Malignant neoplasms (C00-C97)	44,618	10.4	1,289.4
3	Alzheimer's disease (G30)	34,827	8.1	1,006.5
4	Cerebrovascular diseases (I60-I69)	34,263	8.0	990.2
5	Chronic lower respiratory diseases (J40-J47)	20,317	4.8	587.2
6	Influenza and pneumonia (J09-J18)	12,902	3.0	372.9
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,515	2.2	275.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,534	2.0	246.6
9	Diabetes mellitus (E10-E14)	7,919	1.9	228.9
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	6,888	1.6	199.1
11	Septicemia (A40-A41)	5,329	1.2	154.0
12	Pneumonitis due to solids and liquids (J69)	3,790	0.9	109.5
13	Parkinson's disease (G20-G21)	3,758	0.9	108.6
14	Atherosclerosis (I70)	2,663	0.6	77.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,414	0.6	69.8
...	All other causes (Residual)	95,801	22.4	2,768.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, all ages				
...	All causes	286,623	100.0	699.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	69,687	24.3	170.0
2	Malignant neoplasms (C00-C97)	64,649	22.6	157.7
3	Cerebrovascular diseases (I60-I69)	15,878	5.5	38.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,074	4.2	29.4
5	Diabetes mellitus (E10-E14)	11,960	4.2	29.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,808	3.1	21.5
7	Chronic lower respiratory diseases (J40-J47)	8,530	3.0	20.8
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7,874	2.7	19.2
9	Septicemia (A40-A41)	6,179	2.2	15.1
10	Human immunodeficiency virus (HIV) disease (B20-B24)	5,377	1.9	13.1
11	Influenza and pneumonia (J09-J18)	5,290	1.8	12.9
12	Alzheimer's disease (G30)	5,090	1.8	12.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,868	1.7	11.9
14	Certain conditions originating in the perinatal period (P00-P96)	4,654	1.6	11.4
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,560	0.9	6.2
...	All other causes (Residual)	53,145	18.5	129.6
Black, both sexes, 1-4 years				
...	All causes	1,077	100.0	38.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	301	27.9	10.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	149	13.8	5.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	92	8.5	3.3
4	Malignant neoplasms (C00-C97)	59	5.5	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	50	4.6	1.8
6	Chronic lower respiratory diseases (J40-J47)	33	3.1	1.2
7	Influenza and pneumonia (J09-J18)	30	2.8	1.1
8	Certain conditions originating in the perinatal period (P00-P96)	20	1.9	0.7
9	Septicemia (A40-A41)	19	1.8	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	1.2	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	12	1.1	*
12	Cerebrovascular diseases (I60-I69)	9	0.8	*
12	Acute bronchitis and bronchiolitis (J20-J21)	9	0.8	*
14	Anemias (D50-D64)	8	0.7	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.6	*
...	All other causes (Residual)	267	24.8	9.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 5-14 years				
...	All causes	1,300	100.0	20.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	385	29.6	6.0
2	Malignant neoplasms (C00-C97)	133	10.2	2.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	108	8.3	1.7
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	81	6.2	1.3
5	Chronic lower respiratory diseases (J40-J47)	75	5.8	1.2
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	64	4.9	1.0
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	49	3.8	0.8
8	Influenza and pneumonia (J09-J18)	46	3.5	0.7
9	Septicemia (A40-A41)	22	1.7	0.3
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	1.5	0.3
11	Anemias (D50-D64)	19	1.5	*
12	Cerebrovascular diseases (I60-I69)	18	1.4	*
13	Diabetes mellitus (E10-E14)	8	0.6	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.5	*
15	Certain conditions originating in the perinatal period (P00-P96)	5	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.4	*
...	All other causes (Residual)	256	19.7	4.0
Black, both sexes, 15-24 years				
...	All causes	6,742	100.0	96.7
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,810	41.7	40.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,349	20.0	19.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	448	6.6	6.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	345	5.1	4.9
5	Malignant neoplasms (C00-C97)	291	4.3	4.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	106	1.6	1.5
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	91	1.3	1.3
8	Pregnancy, childbirth and the puerperium (O00-O99)	88	1.3	1.3
9	Influenza and pneumonia (J09-J18)	87	1.3	1.2
9	Chronic lower respiratory diseases (J40-J47)	87	1.3	1.2
11	Anemias (D50-D64)	66	1.0	0.9
12	Diabetes mellitus (E10-E14)	50	0.7	0.7
13	Cerebrovascular diseases (I60-I69)	41	0.6	0.6
14	Septicemia (A40-A41)	31	0.5	0.4
15	Legal intervention (Y35,Y89.0)	30	0.4	0.4
...	All other causes (Residual)	822	12.2	11.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 25-34 years				
...	All causes	9,196	100.0	154.1
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,329	25.3	39.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,616	17.6	27.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	963	10.5	16.1
4	Malignant neoplasms (C00-C97)	633	6.9	10.6
5	Human immunodeficiency virus (HIV) disease (B20-B24)	602	6.5	10.1
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	474	5.2	7.9
7	Diabetes mellitus (E10-E14)	190	2.1	3.2
8	Cerebrovascular diseases (I60-I69)	147	1.6	2.5
9	Influenza and pneumonia (J09-J18)	145	1.6	2.4
10	Pregnancy, childbirth and the puerperium (O00-O99)	130	1.4	2.2
11	Anemias (D50-D64)	114	1.2	1.9
12	Septicemia (A40-A41)	109	1.2	1.8
13	Chronic lower respiratory diseases (J40-J47)	105	1.1	1.8
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	99	1.1	1.7
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	79	0.9	1.3
...	All other causes (Residual)	1,461	15.9	24.5
Black, both sexes, 35-44 years				
...	All causes	15,153	100.0	273.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,889	19.1	52.2
2	Malignant neoplasms (C00-C97)	2,163	14.3	39.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,768	11.7	31.9
4	Human immunodeficiency virus (HIV) disease (B20-B24)	1,393	9.2	25.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,196	7.9	21.6
6	Cerebrovascular diseases (I60-I69)	548	3.6	9.9
7	Diabetes mellitus (E10-E14)	531	3.5	9.6
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	404	2.7	7.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	315	2.1	5.7
10	Septicemia (A40-A41)	274	1.8	4.9
11	Influenza and pneumonia (J09-J18)	254	1.7	4.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	203	1.3	3.7
13	Chronic liver disease and cirrhosis (K70,K73-K74)	187	1.2	3.4
14	Chronic lower respiratory diseases (J40-J47)	180	1.2	3.3
15	Anemias (D50-D64)	133	0.9	2.4
...	All other causes (Residual)	2,715	17.9	49.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 45-54 years				
...	All causes	35,447	100.0	646.3
1	Malignant neoplasms (C00-C97)	8,610	24.3	157.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,196	23.1	149.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,431	6.9	44.3
4	Human immunodeficiency virus (HIV) disease (B20-B24)	1,866	5.3	34.0
5	Cerebrovascular diseases (I60-I69)	1,793	5.1	32.7
6	Diabetes mellitus (E10-E14)	1,340	3.8	24.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	867	2.4	15.8
8	Chronic liver disease and cirrhosis (K70,K73-K74)	810	2.3	14.8
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	724	2.0	13.2
10	Chronic lower respiratory diseases (J40-J47)	706	2.0	12.9
11	Septicemia (A40-A41)	634	1.8	11.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	557	1.6	10.2
13	Influenza and pneumonia (J09-J18)	520	1.5	9.5
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	375	1.1	6.8
15	Viral hepatitis (B15-B19)	338	1.0	6.2
...	All other causes (Residual)	5,680	16.0	103.6
Black, both sexes, 55-64 years				
...	All causes	50,676	100.0	1,374.7
1	Malignant neoplasms (C00-C97)	15,703	31.0	426.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,940	25.5	351.0
3	Cerebrovascular diseases (I60-I69)	2,610	5.2	70.8
4	Diabetes mellitus (E10-E14)	2,461	4.9	66.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,639	3.2	44.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,428	2.8	38.7
7	Chronic lower respiratory diseases (J40-J47)	1,380	2.7	37.4
8	Septicemia (A40-A41)	1,082	2.1	29.4
9	Human immunodeficiency virus (HIV) disease (B20-B24)	1,048	2.1	28.4
10	Chronic liver disease and cirrhosis (K70,K73-K74)	941	1.9	25.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	833	1.6	22.6
12	Influenza and pneumonia (J09-J18)	650	1.3	17.6
13	Viral hepatitis (B15-B19)	519	1.0	14.1
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	275	0.5	7.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	226	0.4	6.1
...	All other causes (Residual)	6,941	13.7	188.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 65-74 years				
...	All causes	51,870	100.0	2,645.8
1	Malignant neoplasms (C00-C97)	16,411	31.6	837.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,528	26.1	690.0
3	Cerebrovascular diseases (I60-I69)	3,007	5.8	153.4
4	Diabetes mellitus (E10-E14)	2,767	5.3	141.1
5	Chronic lower respiratory diseases (J40-J47)	2,074	4.0	105.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,932	3.7	98.5
7	Septicemia (A40-A41)	1,270	2.4	64.8
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	877	1.7	44.7
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	829	1.6	42.3
10	Influenza and pneumonia (J09-J18)	823	1.6	42.0
11	Chronic liver disease and cirrhosis (K70,K73-K74)	373	0.7	19.0
12	Alzheimer's disease (G30)	364	0.7	18.6
13	Human immunodeficiency virus (HIV) disease (B20-B24)	277	0.5	14.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	223	0.4	11.4
15	Pneumonitis due to solids and liquids (J69)	220	0.4	11.2
...	All other causes (Residual)	6,895	13.3	351.7
Black, both sexes, 75-84 years				
...	All causes	57,464	100.0	5,180.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,566	27.1	1,403.2
2	Malignant neoplasms (C00-C97)	13,981	24.3	1,260.3
3	Cerebrovascular diseases (I60-I69)	4,007	7.0	361.2
4	Diabetes mellitus (E10-E14)	2,784	4.8	251.0
5	Chronic lower respiratory diseases (J40-J47)	2,339	4.1	210.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,307	4.0	208.0
7	Alzheimer's disease (G30)	1,670	2.9	150.5
8	Septicemia (A40-A41)	1,460	2.5	131.6
9	Influenza and pneumonia (J09-J18)	1,259	2.2	113.5
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,083	1.9	97.6
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	728	1.3	65.6
12	Pneumonitis due to solids and liquids (J69)	419	0.7	37.8
13	Parkinson's disease (G20-G21)	325	0.6	29.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	294	0.5	26.5
15	Aortic aneurysm and dissection (I71)	199	0.3	17.9
...	All other causes (Residual)	9,043	15.7	815.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 85 years and over				
...	All causes	49,356	100.0	13,463.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,020	30.4	4,097.1
2	Malignant neoplasms (C00-C97)	6,648	13.5	1,813.4
3	Cerebrovascular diseases (I60-I69)	3,652	7.4	996.2
4	Alzheimer's disease (G30)	2,975	6.0	811.5
5	Diabetes mellitus (E10-E14)	1,828	3.7	498.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,787	3.6	487.5
7	Chronic lower respiratory diseases (J40-J47)	1,542	3.1	420.6
8	Influenza and pneumonia (J09-J18)	1,394	2.8	380.3
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,244	2.5	339.3
10	Septicemia (A40-A41)	1,185	2.4	323.2
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	621	1.3	169.4
12	Pneumonitis due to solids and liquids (J69)	485	1.0	132.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	250	0.5	68.2
14	Parkinson's disease (G20-G21)	247	0.5	67.4
15	Atherosclerosis (I70)	216	0.4	58.9
...	All other causes (Residual)	10,262	20.8	2,799.2
Black, male, all ages				
...	All causes	146,239	100.0	745.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	35,420	24.2	180.6
2	Malignant neoplasms (C00-C97)	33,443	22.9	170.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,083	5.5	41.2
4	Cerebrovascular diseases (I60-I69)	6,962	4.8	35.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6,715	4.6	34.2
6	Diabetes mellitus (E10-E14)	5,488	3.8	28.0
7	Chronic lower respiratory diseases (J40-J47)	4,503	3.1	23.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,974	2.7	20.3
9	Human immunodeficiency virus (HIV) disease (B20-B24)	3,535	2.4	18.0
10	Septicemia (A40-A41)	2,745	1.9	14.0
11	Certain conditions originating in the perinatal period (P00-P96)	2,635	1.8	13.4
12	Influenza and pneumonia (J09-J18)	2,512	1.7	12.8
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,066	1.4	10.5
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,684	1.2	8.6
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,660	1.1	8.5
...	All other causes (Residual)	24,814	17.0	126.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 1-4 years				
...	All causes	576	100.0	40.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	175	30.4	12.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	80	13.9	5.7
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	50	8.7	3.5
4	Malignant neoplasms (C00-C97)	22	3.8	1.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	22	3.8	1.6
6	Chronic lower respiratory diseases (J40-J47)	18	3.1	*
7	Influenza and pneumonia (J09-J18)	14	2.4	*
8	Certain conditions originating in the perinatal period (P00-P96)	11	1.9	*
9	Complications of medical and surgical care (Y40-Y84,Y88)	10	1.7	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	1.6	*
11	Septicemia (A40-A41)	7	1.2	*
12	Anemias (D50-D64)	6	1.0	*
13	Cerebrovascular diseases (I60-I69)	4	0.7	*
14	Acute bronchitis and bronchiolitis (J20-J21)	3	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.5	*
...	All other causes (Residual)	142	24.7	10.0
Black, male, 5-14 years				
...	All causes	755	100.0	23.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	241	31.9	7.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	74	9.8	2.3
3	Malignant neoplasms (C00-C97)	64	8.5	2.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	48	6.4	1.5
5	Chronic lower respiratory diseases (J40-J47)	47	6.2	1.4
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	37	4.9	1.1
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	35	4.6	1.1
8	Influenza and pneumonia (J09-J18)	22	2.9	0.7
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	1.7	*
10	Cerebrovascular diseases (I60-I69)	11	1.5	*
11	Septicemia (A40-A41)	10	1.3	*
12	Anemias (D50-D64)	7	0.9	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.5	*
14	Diseases of appendix (K35-K38)	3	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.4	*
14	Certain conditions originating in the perinatal period (P00-P96)	3	0.4	*
...	All other causes (Residual)	133	17.6	4.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 15-24 years				
...	All causes	5,090	100.0	144.0
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,526	49.6	71.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	968	19.0	27.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	372	7.3	10.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	226	4.4	6.4
5	Malignant neoplasms (C00-C97)	162	3.2	4.6
6	Human immunodeficiency virus (HIV) disease (B20-B24)	65	1.3	1.8
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	56	1.1	1.6
8	Chronic lower respiratory diseases (J40-J47)	55	1.1	1.6
9	Influenza and pneumonia (J09-J18)	40	0.8	1.1
10	Anemias (D50-D64)	36	0.7	1.0
11	Legal intervention (Y35,Y89.0)	29	0.6	0.8
12	Diabetes mellitus (E10-E14)	25	0.5	0.7
13	Cerebrovascular diseases (I60-I69)	22	0.4	0.6
14	Septicemia (A40-A41)	12	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	0.2	*
...	All other causes (Residual)	484	9.5	13.7
Black, male, 25-34 years				
...	All causes	6,238	100.0	213.2
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,045	32.8	69.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,230	19.7	42.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	583	9.3	19.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	382	6.1	13.1
5	Human immunodeficiency virus (HIV) disease (B20-B24)	347	5.6	11.9
6	Malignant neoplasms (C00-C97)	286	4.6	9.8
7	Diabetes mellitus (E10-E14)	110	1.8	3.8
8	Cerebrovascular diseases (I60-I69)	80	1.3	2.7
9	Influenza and pneumonia (J09-J18)	64	1.0	2.2
10	Anemias (D50-D64)	61	1.0	2.1
11	Chronic lower respiratory diseases (J40-J47)	56	0.9	1.9
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	56	0.9	1.9
13	Septicemia (A40-A41)	52	0.8	1.8
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	42	0.7	1.4
15	Legal intervention (Y35,Y89.0)	37	0.6	1.3
...	All other causes (Residual)	807	12.9	27.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 35-44 years				
...	All causes	8,952	100.0	343.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,856	20.7	71.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,260	14.1	48.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,011	11.3	38.8
4	Human immunodeficiency virus (HIV) disease (B20-B24)	862	9.6	33.0
5	Malignant neoplasms (C00-C97)	833	9.3	31.9
6	Diabetes mellitus (E10-E14)	333	3.7	12.8
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	333	3.7	12.8
8	Cerebrovascular diseases (I60-I69)	289	3.2	11.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	161	1.8	6.2
10	Septicemia (A40-A41)	131	1.5	5.0
11	Influenza and pneumonia (J09-J18)	117	1.3	4.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	115	1.3	4.4
13	Chronic liver disease and cirrhosis (K70,K73-K74)	113	1.3	4.3
14	Chronic lower respiratory diseases (J40-J47)	106	1.2	4.1
15	Anemias (D50-D64)	64	0.7	2.5
...	All other causes (Residual)	1,368	15.3	52.4
Black, male, 45-54 years				
...	All causes	20,320	100.0	801.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,188	25.5	204.7
2	Malignant neoplasms (C00-C97)	4,206	20.7	166.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,634	8.0	64.5
4	Human immunodeficiency virus (HIV) disease (B20-B24)	1,233	6.1	48.7
5	Cerebrovascular diseases (I60-I69)	922	4.5	36.4
6	Diabetes mellitus (E10-E14)	780	3.8	30.8
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	567	2.8	22.4
8	Chronic liver disease and cirrhosis (K70,K73-K74)	537	2.6	21.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	462	2.3	18.2
10	Chronic lower respiratory diseases (J40-J47)	333	1.6	13.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	328	1.6	12.9
12	Septicemia (A40-A41)	322	1.6	12.7
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	281	1.4	11.1
14	Influenza and pneumonia (J09-J18)	276	1.4	10.9
15	Viral hepatitis (B15-B19)	216	1.1	8.5
...	All other causes (Residual)	3,035	14.9	119.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 55-64 years				
...	All causes	29,932	100.0	1,821.0
1	Malignant neoplasms (C00-C97)	8,802	29.4	535.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,121	27.1	494.1
3	Cerebrovascular diseases (I60-I69)	1,482	5.0	90.2
4	Diabetes mellitus (E10-E14)	1,312	4.4	79.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,162	3.9	70.7
6	Human immunodeficiency virus (HIV) disease (B20-B24)	771	2.6	46.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	764	2.6	46.5
8	Chronic lower respiratory diseases (J40-J47)	761	2.5	46.3
9	Chronic liver disease and cirrhosis (K70,K73-K74)	647	2.2	39.4
10	Septicemia (A40-A41)	570	1.9	34.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	466	1.6	28.3
12	Viral hepatitis (B15-B19)	376	1.3	22.9
13	Influenza and pneumonia (J09-J18)	362	1.2	22.0
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	228	0.8	13.9
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	144	0.5	8.8
...	All other causes (Residual)	3,964	13.2	241.2
Black, male, 65-74 years				
...	All causes	28,076	100.0	3,426.9
1	Malignant neoplasms (C00-C97)	8,972	32.0	1,095.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,717	27.5	941.9
3	Cerebrovascular diseases (I60-I69)	1,579	5.6	192.7
4	Diabetes mellitus (E10-E14)	1,341	4.8	163.7
5	Chronic lower respiratory diseases (J40-J47)	1,147	4.1	140.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	933	3.3	113.9
7	Septicemia (A40-A41)	602	2.1	73.5
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	548	2.0	66.9
9	Influenza and pneumonia (J09-J18)	474	1.7	57.9
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	410	1.5	50.0
11	Chronic liver disease and cirrhosis (K70,K73-K74)	239	0.9	29.2
12	Human immunodeficiency virus (HIV) disease (B20-B24)	198	0.7	24.2
13	Alzheimer's disease (G30)	164	0.6	20.0
14	Pneumonitis due to solids and liquids (J69)	115	0.4	14.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	112	0.4	13.7
...	All other causes (Residual)	3,525	12.6	430.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 75-84 years				
...	All causes	26,115	100.0	6,387.1
1	Malignant neoplasms (C00-C97)	7,124	27.3	1,742.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,071	27.1	1,729.4
3	Cerebrovascular diseases (I60-I69)	1,602	6.1	391.8
4	Chronic lower respiratory diseases (J40-J47)	1,282	4.9	313.5
5	Diabetes mellitus (E10-E14)	1,093	4.2	267.3
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	990	3.8	242.1
7	Influenza and pneumonia (J09-J18)	609	2.3	148.9
8	Septicemia (A40-A41)	603	2.3	147.5
9	Alzheimer's disease (G30)	595	2.3	145.5
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	410	1.6	100.3
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	405	1.6	99.1
12	Pneumonitis due to solids and liquids (J69)	210	0.8	51.4
13	Parkinson's disease (G20-G21)	186	0.7	45.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	133	0.5	32.5
15	Aortic aneurysm and dissection (I71)	88	0.3	21.5
...	All other causes (Residual)	3,714	14.2	908.4
Black, male, 85 years and over				
...	All causes	15,459	100.0	15,430.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,523	29.3	4,514.7
2	Malignant neoplasms (C00-C97)	2,963	19.2	2,957.6
3	Cerebrovascular diseases (I60-I69)	942	6.1	940.3
4	Chronic lower respiratory diseases (J40-J47)	693	4.5	691.7
5	Alzheimer's disease (G30)	646	4.2	644.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	570	3.7	569.0
7	Diabetes mellitus (E10-E14)	491	3.2	490.1
8	Influenza and pneumonia (J09-J18)	487	3.2	486.1
9	Septicemia (A40-A41)	387	2.5	386.3
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	310	2.0	309.4
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	218	1.4	217.6
12	Pneumonitis due to solids and liquids (J69)	188	1.2	187.7
13	Parkinson's disease (G20-G21)	127	0.8	126.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	75	0.5	74.9
15	Atherosclerosis (I70)	58	0.4	57.9
...	All other causes (Residual)	2,781	18.0	2,775.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, all ages				
...	All causes	140,384	100.0	656.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	34,267	24.4	160.2
2	Malignant neoplasms (C00-C97)	31,206	22.2	145.9
3	Cerebrovascular diseases (I60-I69)	8,916	6.4	41.7
4	Diabetes mellitus (E10-E14)	6,472	4.6	30.3
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,834	3.4	22.6
6	Chronic lower respiratory diseases (J40-J47)	4,027	2.9	18.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,991	2.8	18.7
8	Alzheimer's disease (G30)	3,639	2.6	17.0
9	Septicemia (A40-A41)	3,434	2.4	16.1
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,802	2.0	13.1
11	Influenza and pneumonia (J09-J18)	2,778	2.0	13.0
12	Certain conditions originating in the perinatal period (P00-P96)	2,019	1.4	9.4
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,842	1.3	8.6
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,159	0.8	5.4
15	Chronic liver disease and cirrhosis (K70,K73-K74)	900	0.6	4.2
...	All other causes (Residual)	28,098	20.0	131.4
Black, female, 1-4 years				
...	All causes	501	100.0	36.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	126	25.1	9.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	69	13.8	5.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	42	8.4	3.1
4	Malignant neoplasms (C00-C97)	37	7.4	2.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	28	5.6	2.1
6	Influenza and pneumonia (J09-J18)	16	3.2	*
7	Chronic lower respiratory diseases (J40-J47)	15	3.0	*
8	Septicemia (A40-A41)	12	2.4	*
9	Certain conditions originating in the perinatal period (P00-P96)	9	1.8	*
10	Acute bronchitis and bronchiolitis (J20-J21)	6	1.2	*
11	Cerebrovascular diseases (I60-I69)	5	1.0	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.8	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.6	*
14	Anemias (D50-D64)	2	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
...	All other causes (Residual)	125	25.0	9.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 5-14 years				
...	All causes	545	100.0	17.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	144	26.4	4.6
2	Malignant neoplasms (C00-C97)	69	12.7	2.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	34	6.2	1.1
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	33	6.1	1.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	29	5.3	0.9
6	Chronic lower respiratory diseases (J40-J47)	28	5.1	0.9
7	Influenza and pneumonia (J09-J18)	24	4.4	0.8
8	Septicemia (A40-A41)	12	2.2	*
8	Anemias (D50-D64)	12	2.2	*
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	12	2.2	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	1.3	*
11	Cerebrovascular diseases (I60-I69)	7	1.3	*
13	Diabetes mellitus (E10-E14)	6	1.1	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.6	*
14	Pregnancy, childbirth and the puerperium (O00-O99)	3	0.6	*
...	All other causes (Residual)	122	22.4	3.9
Black, female, 15-24 years				
...	All causes	1,652	100.0	48.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	381	23.1	11.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	284	17.2	8.3
3	Malignant neoplasms (C00-C97)	129	7.8	3.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	119	7.2	3.5
5	Pregnancy, childbirth and the puerperium (O00-O99)	88	5.3	2.6
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	76	4.6	2.2
7	Influenza and pneumonia (J09-J18)	47	2.8	1.4
8	Human immunodeficiency virus (HIV) disease (B20-B24)	41	2.5	1.2
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	35	2.1	1.0
10	Chronic lower respiratory diseases (J40-J47)	32	1.9	0.9
11	Anemias (D50-D64)	30	1.8	0.9
12	Diabetes mellitus (E10-E14)	25	1.5	0.7
13	Septicemia (A40-A41)	19	1.2	*
13	Cerebrovascular diseases (I60-I69)	19	1.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	1.0	*
...	All other causes (Residual)	310	18.8	9.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 25-34 years				
...	All causes	2,958	100.0	97.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	386	13.0	12.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	380	12.8	12.5
3	Malignant neoplasms (C00-C97)	347	11.7	11.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	284	9.6	9.3
5	Human immunodeficiency virus (HIV) disease (B20-B24)	255	8.6	8.4
6	Pregnancy, childbirth and the puerperium (O00-O99)	130	4.4	4.3
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	92	3.1	3.0
8	Influenza and pneumonia (J09-J18)	81	2.7	2.7
9	Diabetes mellitus (E10-E14)	80	2.7	2.6
10	Cerebrovascular diseases (I60-I69)	67	2.3	2.2
11	Septicemia (A40-A41)	57	1.9	1.9
12	Anemias (D50-D64)	53	1.8	1.7
13	Chronic lower respiratory diseases (J40-J47)	49	1.7	1.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	43	1.5	1.4
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	37	1.3	1.2
...	All other causes (Residual)	617	20.9	20.3
Black, female, 35-44 years				
...	All causes	6,201	100.0	211.8
1	Malignant neoplasms (C00-C97)	1,330	21.4	45.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,033	16.7	35.3
3	Human immunodeficiency virus (HIV) disease (B20-B24)	531	8.6	18.1
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	508	8.2	17.3
5	Cerebrovascular diseases (I60-I69)	259	4.2	8.8
6	Diabetes mellitus (E10-E14)	198	3.2	6.8
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	185	3.0	6.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	154	2.5	5.3
9	Septicemia (A40-A41)	143	2.3	4.9
10	Influenza and pneumonia (J09-J18)	137	2.2	4.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	88	1.4	3.0
12	Chronic lower respiratory diseases (J40-J47)	74	1.2	2.5
12	Chronic liver disease and cirrhosis (K70,K73-K74)	74	1.2	2.5
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	71	1.1	2.4
15	Anemias (D50-D64)	69	1.1	2.4
...	All other causes (Residual)	1,347	21.7	46.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 45-54 years				
...	All causes	15,127	100.0	512.6
1	Malignant neoplasms (C00-C97)	4,404	29.1	149.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,008	19.9	101.9
3	Cerebrovascular diseases (I60-I69)	871	5.8	29.5
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	797	5.3	27.0
5	Human immunodeficiency virus (HIV) disease (B20-B24)	633	4.2	21.4
6	Diabetes mellitus (E10-E14)	560	3.7	19.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	405	2.7	13.7
8	Chronic lower respiratory diseases (J40-J47)	373	2.5	12.6
9	Septicemia (A40-A41)	312	2.1	10.6
10	Chronic liver disease and cirrhosis (K70,K73-K74)	273	1.8	9.3
11	Influenza and pneumonia (J09-J18)	244	1.6	8.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	229	1.5	7.8
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	157	1.0	5.3
14	Viral hepatitis (B15-B19)	122	0.8	4.1
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	94	0.6	3.2
...	All other causes (Residual)	2,645	17.5	89.6
Black, female, 55-64 years				
...	All causes	20,744	100.0	1,015.5
1	Malignant neoplasms (C00-C97)	6,901	33.3	337.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,819	23.2	235.9
3	Diabetes mellitus (E10-E14)	1,149	5.5	56.2
4	Cerebrovascular diseases (I60-I69)	1,128	5.4	55.2
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	664	3.2	32.5
6	Chronic lower respiratory diseases (J40-J47)	619	3.0	30.3
7	Septicemia (A40-A41)	512	2.5	25.1
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	477	2.3	23.4
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	367	1.8	18.0
10	Chronic liver disease and cirrhosis (K70,K73-K74)	294	1.4	14.4
11	Influenza and pneumonia (J09-J18)	288	1.4	14.1
12	Human immunodeficiency virus (HIV) disease (B20-B24)	277	1.3	13.6
13	Viral hepatitis (B15-B19)	143	0.7	7.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	104	0.5	5.1
15	Pneumonitis due to solids and liquids (J69)	76	0.4	3.7
...	All other causes (Residual)	2,926	14.1	143.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 65-74 years				
...	All causes	23,794	100.0	2,085.0
1	Malignant neoplasms (C00-C97)	7,439	31.3	651.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,811	24.4	509.2
3	Cerebrovascular diseases (I60-I69)	1,428	6.0	125.1
4	Diabetes mellitus (E10-E14)	1,426	6.0	125.0
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	999	4.2	87.5
6	Chronic lower respiratory diseases (J40-J47)	927	3.9	81.2
7	Septicemia (A40-A41)	668	2.8	58.5
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	467	2.0	40.9
9	Influenza and pneumonia (J09-J18)	349	1.5	30.6
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	281	1.2	24.6
11	Alzheimer's disease (G30)	200	0.8	17.5
12	Chronic liver disease and cirrhosis (K70,K73-K74)	134	0.6	11.7
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	111	0.5	9.7
14	Pneumonitis due to solids and liquids (J69)	105	0.4	9.2
15	Human immunodeficiency virus (HIV) disease (B20-B24)	79	0.3	6.9
...	All other causes (Residual)	3,370	14.2	295.3
Black, female, 75-84 years				
...	All causes	31,349	100.0	4,475.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,495	27.1	1,212.8
2	Malignant neoplasms (C00-C97)	6,857	21.9	978.9
3	Cerebrovascular diseases (I60-I69)	2,405	7.7	343.3
4	Diabetes mellitus (E10-E14)	1,691	5.4	241.4
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,317	4.2	188.0
6	Alzheimer's disease (G30)	1,075	3.4	153.5
7	Chronic lower respiratory diseases (J40-J47)	1,057	3.4	150.9
8	Septicemia (A40-A41)	857	2.7	122.3
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	678	2.2	96.8
10	Influenza and pneumonia (J09-J18)	650	2.1	92.8
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	318	1.0	45.4
12	Pneumonitis due to solids and liquids (J69)	209	0.7	29.8
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	161	0.5	23.0
14	Parkinson's disease (G20-G21)	139	0.4	19.8
15	Atherosclerosis (I70)	113	0.4	16.1
...	All other causes (Residual)	5,327	17.0	760.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 85 years and over				
...	All causes	33,897	100.0	12,723.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,497	31.0	3,940.1
2	Malignant neoplasms (C00-C97)	3,685	10.9	1,383.2
3	Cerebrovascular diseases (I60-I69)	2,710	8.0	1,017.2
4	Alzheimer's disease (G30)	2,329	6.9	874.2
5	Diabetes mellitus (E10-E14)	1,337	3.9	501.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,217	3.6	456.8
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	934	2.8	350.6
8	Influenza and pneumonia (J09-J18)	907	2.7	340.4
9	Chronic lower respiratory diseases (J40-J47)	849	2.5	318.7
10	Septicemia (A40-A41)	798	2.4	299.5
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	403	1.2	151.3
12	Pneumonitis due to solids and liquids (J69)	297	0.9	111.5
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	175	0.5	65.7
14	Atherosclerosis (I70)	158	0.5	59.3
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	128	0.4	48.0
...	All other causes (Residual)	7,473	22.0	2,805.0
American Indian, both sexes, all ages				
...	All causes	14,960	100.0	427.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,764	18.5	79.0
2	Malignant neoplasms (C00-C97)	2,710	18.1	77.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,733	11.6	49.5
4	Diabetes mellitus (E10-E14)	786	5.3	22.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	714	4.8	20.4
6	Chronic lower respiratory diseases (J40-J47)	610	4.1	17.4
7	Cerebrovascular diseases (I60-I69)	561	3.8	16.0
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	429	2.9	12.3
9	Influenza and pneumonia (J09-J18)	402	2.7	11.5
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	307	2.1	8.8
11	Septicemia (A40-A41)	275	1.8	7.9
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	254	1.7	7.3
13	Alzheimer's disease (G30)	193	1.3	5.5
14	Certain conditions originating in the perinatal period (P00-P96)	148	1.0	4.2
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	125	0.8	3.6
...	All other causes (Residual)	2,949	19.7	84.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 1-4 years				
...	All causes	83	100.0	30.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	34	41.0	12.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	7.2	*
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	7.2	*
4	Influenza and pneumonia (J09-J18)	5	6.0	*
5	Malignant neoplasms (C00-C97)	4	4.8	*
6	Certain conditions originating in the perinatal period (P00-P96)	2	2.4	*
7	Meningococcal infection (A39)	1	1.2	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.2	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	1.2	*
...	All other causes (Residual)	23	27.7	8.5
American Indian, both sexes, 5-14 years				
...	All causes	115	100.0	20.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	38	33.0	6.7
2	Malignant neoplasms (C00-C97)	13	11.3	*
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	9	7.8	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	6	5.2	*
4	Influenza and pneumonia (J09-J18)	6	5.2	*
6	Septicemia (A40-A41)	5	4.3	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	3.5	*
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	3.5	*
9	Chronic lower respiratory diseases (J40-J47)	3	2.6	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.9	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.9	*
...	All other causes (Residual)	25	21.7	4.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 15-24 years				
...	All causes	670	100.0	109.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	327	48.8	53.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	152	22.7	24.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	78	11.6	12.8
4	Malignant neoplasms (C00-C97)	14	2.1	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	10	1.5	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	1.3	*
7	Influenza and pneumonia (J09-J18)	8	1.2	*
8	Chronic liver disease and cirrhosis (K70,K73-K74)	4	0.6	*
9	Cerebrovascular diseases (I60-I69)	3	0.4	*
9	Chronic lower respiratory diseases (J40-J47)	3	0.4	*
11	Diabetes mellitus (E10-E14)	2	0.3	*
12	Septicemia (A40-A41)	1	0.1	*
12	Anemias (D50-D64)	1	0.1	*
12	Pneumonitis due to solids and liquids (J69)	1	0.1	*
12	Pregnancy, childbirth and the puerperium (O00-O99)	1	0.1	*
12	Legal intervention (Y35,Y89.0)	1	0.1	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.1	*
...	All other causes (Residual)	54	8.1	8.9
American Indian, both sexes, 25-34 years				
...	All causes	752	100.0	141.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	283	37.6	53.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	91	12.1	17.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	61	8.1	11.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	50	6.6	9.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	43	5.7	8.1
6	Malignant neoplasms (C00-C97)	41	5.5	7.7
7	Influenza and pneumonia (J09-J18)	17	2.3	*
8	Septicemia (A40-A41)	13	1.7	*
8	Diabetes mellitus (E10-E14)	13	1.7	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	1.2	*
11	Pregnancy, childbirth and the puerperium (O00-O99)	8	1.1	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	7	0.9	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	0.8	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3	0.4	*
14	Cerebrovascular diseases (I60-I69)	3	0.4	*
14	Legal intervention (Y35,Y89.0)	3	0.4	*
...	All other causes (Residual)	101	13.4	19.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 35-44 years				
...	All causes	1,261	100.0	279.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	356	28.2	78.8
2	Chronic liver disease and cirrhosis (K70,K73-K74)	145	11.5	32.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	137	10.9	30.3
4	Malignant neoplasms (C00-C97)	92	7.3	20.4
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	77	6.1	17.0
6	Diabetes mellitus (E10-E14)	49	3.9	10.8
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	48	3.8	10.6
8	Influenza and pneumonia (J09-J18)	37	2.9	8.2
9	Septicemia (A40-A41)	20	1.6	4.4
10	Viral hepatitis (B15-B19)	18	1.4	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	15	1.2	*
12	Cerebrovascular diseases (I60-I69)	13	1.0	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	1.0	*
14	Chronic lower respiratory diseases (J40-J47)	9	0.7	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3	0.2	*
15	Pregnancy, childbirth and the puerperium (O00-O99)	3	0.2	*
...	All other causes (Residual)	226	17.9	50.0
American Indian, both sexes, 45-54 years				
...	All causes	2,174	100.0	486.2
1	Malignant neoplasms (C00-C97)	376	17.3	84.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	334	15.4	74.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	319	14.7	71.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	250	11.5	55.9
5	Diabetes mellitus (E10-E14)	98	4.5	21.9
6	Cerebrovascular diseases (I60-I69)	68	3.1	15.2
7	Influenza and pneumonia (J09-J18)	61	2.8	13.6
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	59	2.7	13.2
9	Viral hepatitis (B15-B19)	53	2.4	11.9
10	Chronic lower respiratory diseases (J40-J47)	46	2.1	10.3
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	37	1.7	8.3
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	33	1.5	7.4
13	Human immunodeficiency virus (HIV) disease (B20-B24)	31	1.4	6.9
14	Septicemia (A40-A41)	29	1.3	6.5
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	13	0.6	*
...	All other causes (Residual)	367	16.9	82.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 55-64 years				
...	All causes	2,525	100.0	827.9
1	Malignant neoplasms (C00-C97)	677	26.8	222.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	540	21.4	177.1
3	Diabetes mellitus (E10-E14)	165	6.5	54.1
4	Chronic liver disease and cirrhosis (K70,K73-K74)	158	6.3	51.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	155	6.1	50.8
6	Chronic lower respiratory diseases (J40-J47)	102	4.0	33.4
7	Cerebrovascular diseases (I60-I69)	82	3.2	26.9
8	Influenza and pneumonia (J09-J18)	52	2.1	17.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	52	2.1	17.0
10	Septicemia (A40-A41)	36	1.4	11.8
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	31	1.2	10.2
12	Viral hepatitis (B15-B19)	23	0.9	7.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	22	0.9	7.2
14	Pneumonitis due to solids and liquids (J69)	14	0.6	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	0.5	*
...	All other causes (Residual)	403	16.0	132.1
American Indian, both sexes, 65-74 years				
...	All causes	2,674	100.0	1,749.8
1	Malignant neoplasms (C00-C97)	743	27.8	486.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	599	22.4	392.0
3	Diabetes mellitus (E10-E14)	205	7.7	134.1
4	Chronic lower respiratory diseases (J40-J47)	170	6.4	111.2
5	Cerebrovascular diseases (I60-I69)	108	4.0	70.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	76	2.8	49.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	75	2.8	49.1
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	69	2.6	45.2
9	Septicemia (A40-A41)	63	2.4	41.2
9	Influenza and pneumonia (J09-J18)	63	2.4	41.2
11	Alzheimer's disease (G30)	22	0.8	14.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	21	0.8	13.7
12	Pneumonitis due to solids and liquids (J69)	21	0.8	13.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	0.5	*
15	Parkinson's disease (G20-G21)	13	0.5	*
...	All other causes (Residual)	412	15.4	269.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 75-84 years				
...	All causes	2,535	100.0	3,453.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	598	23.6	814.6
2	Malignant neoplasms (C00-C97)	538	21.2	732.9
3	Chronic lower respiratory diseases (J40-J47)	178	7.0	242.5
4	Diabetes mellitus (E10-E14)	177	7.0	241.1
5	Cerebrovascular diseases (I60-I69)	148	5.8	201.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	77	3.0	104.9
7	Influenza and pneumonia (J09-J18)	72	2.8	98.1
8	Alzheimer's disease (G30)	71	2.8	96.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	71	2.8	96.7
10	Septicemia (A40-A41)	68	2.7	92.6
11	Parkinson's disease (G20-G21)	25	1.0	34.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	25	1.0	34.1
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	0.7	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	16	0.6	*
15	Pneumonitis due to solids and liquids (J69)	14	0.6	*
...	All other causes (Residual)	438	17.3	596.7
American Indian, both sexes, 85 years and over				
...	All causes	1,785	100.0	6,718.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	491	27.5	1,848.2
2	Malignant neoplasms (C00-C97)	211	11.8	794.2
3	Cerebrovascular diseases (I60-I69)	136	7.6	511.9
4	Alzheimer's disease (G30)	97	5.4	365.1
4	Chronic lower respiratory diseases (J40-J47)	97	5.4	365.1
6	Diabetes mellitus (E10-E14)	77	4.3	289.8
7	Influenza and pneumonia (J09-J18)	73	4.1	274.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	60	3.4	225.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	42	2.4	158.1
10	Septicemia (A40-A41)	39	2.2	146.8
11	Pneumonitis due to solids and liquids (J69)	32	1.8	120.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	26	1.5	97.9
13	Parkinson's disease (G20-G21)	12	0.7	*
14	Aortic aneurysm and dissection (I71)	10	0.6	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.5	*
15	Nutritional deficiencies (E40-E64)	9	0.5	*
...	All other causes (Residual)	364	20.4	1,370.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, all ages				
...	All causes	8,105	100.0	462.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,591	19.6	90.9
2	Malignant neoplasms (C00-C97)	1,381	17.0	78.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,150	14.2	65.7
4	Diabetes mellitus (E10-E14)	418	5.2	23.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	380	4.7	21.7
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	313	3.9	17.9
7	Chronic lower respiratory diseases (J40-J47)	287	3.5	16.4
8	Cerebrovascular diseases (I60-I69)	263	3.2	15.0
9	Influenza and pneumonia (J09-J18)	193	2.4	11.0
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	193	2.4	11.0
11	Septicemia (A40-A41)	127	1.6	7.3
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	126	1.6	7.2
13	Certain conditions originating in the perinatal period (P00-P96)	92	1.1	5.3
14	Alzheimer's disease (G30)	65	0.8	3.7
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	62	0.8	3.5
...	All other causes (Residual)	1,464	18.1	83.6
American Indian, male, 1-4 years				
...	All causes	48	100.0	35.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	24	50.0	17.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	6.3	*
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	6.3	*
4	Malignant neoplasms (C00-C97)	2	4.2	*
4	Influenza and pneumonia (J09-J18)	2	4.2	*
6	Meningococcal infection (A39)	1	2.1	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	2.1	*
6	Certain conditions originating in the perinatal period (P00-P96)	1	2.1	*
...	All other causes (Residual)	11	22.9	*

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 5-14 years				
...	All causes	57	100.0	19.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	21	36.8	7.3
2	Malignant neoplasms (C00-C97)	6	10.5	*
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	8.8	*
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5	8.8	*
5	Influenza and pneumonia (J09-J18)	4	7.0	*
6	Septicemia (A40-A41)	3	5.3	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	3.5	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.8	*
8	Chronic lower respiratory diseases (J40-J47)	1	1.8	*
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1	1.8	*
...	All other causes (Residual)	8	14.0	*
American Indian, male, 15-24 years				
...	All causes	468	100.0	150.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	224	47.9	72.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	112	23.9	36.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	61	13.0	19.6
4	Malignant neoplasms (C00-C97)	10	2.1	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	6	1.3	*
5	Influenza and pneumonia (J09-J18)	6	1.3	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	1.3	*
8	Chronic liver disease and cirrhosis (K70,K73-K74)	4	0.9	*
9	Anemias (D50-D64)	1	0.2	*
9	Diabetes mellitus (E10-E14)	1	0.2	*
9	Cerebrovascular diseases (I60-I69)	1	0.2	*
9	Chronic lower respiratory diseases (J40-J47)	1	0.2	*
9	Pneumonitis due to solids and liquids (J69)	1	0.2	*
9	Legal intervention (Y35,Y89.0)	1	0.2	*
9	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
...	All other causes (Residual)	32	6.8	10.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 25-34 years				
...	All causes	502	100.0	182.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	193	38.4	70.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	66	13.1	24.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	54	10.8	19.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	34	6.8	12.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	32	6.4	11.6
6	Malignant neoplasms (C00-C97)	21	4.2	7.6
7	Diabetes mellitus (E10-E14)	10	2.0	*
8	Influenza and pneumonia (J09-J18)	9	1.8	*
9	Septicemia (A40-A41)	7	1.4	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.8	*
11	Cerebrovascular diseases (I60-I69)	3	0.6	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.6	*
11	Legal intervention (Y35,Y89.0)	3	0.6	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
15	Tuberculosis (A16-A19)	1	0.2	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.2	*
15	Chronic lower respiratory diseases (J40-J47)	1	0.2	*
15	Pneumonitis due to solids and liquids (J69)	1	0.2	*
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.2	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
...	All other causes (Residual)	55	11.0	20.0
American Indian, male, 35-44 years				
...	All causes	782	100.0	339.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	241	30.8	104.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	94	12.0	40.8
3	Chronic liver disease and cirrhosis (K70,K73-K74)	80	10.2	34.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	58	7.4	25.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	37	4.7	16.1
6	Malignant neoplasms (C00-C97)	35	4.5	15.2
7	Diabetes mellitus (E10-E14)	34	4.3	14.8
8	Influenza and pneumonia (J09-J18)	22	2.8	9.6
9	Human immunodeficiency virus (HIV) disease (B20-B24)	12	1.5	*
10	Septicemia (A40-A41)	10	1.3	*
11	Viral hepatitis (B15-B19)	9	1.2	*
12	Cerebrovascular diseases (I60-I69)	8	1.0	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.8	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3	0.4	*
14	Chronic lower respiratory diseases (J40-J47)	3	0.4	*
...	All other causes (Residual)	130	16.6	56.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 45-54 years				
...	All causes	1,250	100.0	573.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	222	17.8	101.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	215	17.2	98.6
3	Malignant neoplasms (C00-C97)	188	15.0	86.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	131	10.5	60.1
5	Diabetes mellitus (E10-E14)	53	4.2	24.3
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	42	3.4	19.3
7	Cerebrovascular diseases (I60-I69)	37	3.0	17.0
8	Viral hepatitis (B15-B19)	32	2.6	14.7
9	Influenza and pneumonia (J09-J18)	25	2.0	11.5
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	23	1.8	10.6
11	Human immunodeficiency virus (HIV) disease (B20-B24)	22	1.8	10.1
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20	1.6	9.2
13	Chronic lower respiratory diseases (J40-J47)	17	1.4	*
14	Septicemia (A40-A41)	14	1.1	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7	0.6	*
...	All other causes (Residual)	202	16.2	92.7
American Indian, male, 55-64 years				
...	All causes	1,496	100.0	1,024.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	366	24.5	250.5
2	Malignant neoplasms (C00-C97)	365	24.4	249.9
3	Diabetes mellitus (E10-E14)	114	7.6	78.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	113	7.6	77.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	91	6.1	62.3
6	Chronic lower respiratory diseases (J40-J47)	55	3.7	37.7
7	Cerebrovascular diseases (I60-I69)	40	2.7	27.4
8	Influenza and pneumonia (J09-J18)	29	1.9	19.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	1.5	15.7
10	Septicemia (A40-A41)	22	1.5	15.1
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	22	1.5	15.1
12	Viral hepatitis (B15-B19)	14	0.9	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	14	0.9	*
14	Pneumonitis due to solids and liquids (J69)	8	0.5	*
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8	0.5	*
...	All other causes (Residual)	212	14.2	145.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 65-74 years				
...	All causes	1,441	100.0	2,020.5
1	Malignant neoplasms (C00-C97)	396	27.5	555.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	379	26.3	531.4
3	Diabetes mellitus (E10-E14)	109	7.6	152.8
4	Chronic lower respiratory diseases (J40-J47)	79	5.5	110.8
5	Cerebrovascular diseases (I60-I69)	50	3.5	70.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	44	3.1	61.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	32	2.2	44.9
8	Septicemia (A40-A41)	31	2.2	43.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	29	2.0	40.7
10	Influenza and pneumonia (J09-J18)	28	1.9	39.3
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	11	0.8	*
12	Pneumonitis due to solids and liquids (J69)	9	0.6	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.6	*
13	Alzheimer's disease (G30)	8	0.6	*
15	Parkinson's disease (G20-G21)	6	0.4	*
15	Aortic aneurysm and dissection (I71)	6	0.4	*
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6	0.4	*
...	All other causes (Residual)	210	14.6	294.5
American Indian, male, 75-84 years				
...	All causes	1,219	100.0	3,907.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	308	25.3	987.3
2	Malignant neoplasms (C00-C97)	263	21.6	843.1
3	Chronic lower respiratory diseases (J40-J47)	87	7.1	278.9
4	Cerebrovascular diseases (I60-I69)	81	6.6	259.6
5	Diabetes mellitus (E10-E14)	67	5.5	214.8
6	Influenza and pneumonia (J09-J18)	42	3.4	134.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	42	3.4	134.6
8	Septicemia (A40-A41)	31	2.5	99.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	31	2.5	99.4
10	Alzheimer's disease (G30)	28	2.3	89.8
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	12	1.0	*
12	Parkinson's disease (G20-G21)	11	0.9	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	0.8	*
14	Chronic liver disease and cirrhosis (K70,K73-K74)	9	0.7	*
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	8	0.7	*
...	All other causes (Residual)	189	15.5	605.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 85 years and over				
...	All causes	627	100.0	7,013.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	175	27.9	1,957.5
2	Malignant neoplasms (C00-C97)	95	15.2	1,062.6
3	Cerebrovascular diseases (I60-I69)	43	6.9	481.0
4	Chronic lower respiratory diseases (J40-J47)	42	6.7	469.8
5	Diabetes mellitus (E10-E14)	30	4.8	335.6
6	Alzheimer's disease (G30)	28	4.5	313.2
7	Influenza and pneumonia (J09-J18)	22	3.5	246.1
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19	3.0	*
9	Pneumonitis due to solids and liquids (J69)	16	2.6	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	2.4	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	13	2.1	*
12	Septicemia (A40-A41)	9	1.4	*
13	Parkinson's disease (G20-G21)	7	1.1	*
14	Nutritional deficiencies (E40-E64)	5	0.8	*
15	Aortic aneurysm and dissection (I71)	4	0.6	*
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	4	0.6	*
...	All other causes (Residual)	100	15.9	1,118.6
American Indian, female, all ages				
...	All causes	6,855	100.0	391.9
1	Malignant neoplasms (C00-C97)	1,329	19.4	76.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,173	17.1	67.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	583	8.5	33.3
4	Diabetes mellitus (E10-E14)	368	5.4	21.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	334	4.9	19.1
6	Chronic lower respiratory diseases (J40-J47)	323	4.7	18.5
7	Cerebrovascular diseases (I60-I69)	298	4.3	17.0
8	Influenza and pneumonia (J09-J18)	209	3.0	11.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	181	2.6	10.3
10	Septicemia (A40-A41)	148	2.2	8.5
11	Alzheimer's disease (G30)	128	1.9	7.3
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	116	1.7	6.6
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	63	0.9	3.6
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	61	0.9	3.5
15	Certain conditions originating in the perinatal period (P00-P96)	56	0.8	3.2
...	All other causes (Residual)	1,485	21.7	84.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 1-4 years				
...	All causes	35	100.0	26.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10	28.6	*
2	Influenza and pneumonia (J09-J18)	3	8.6	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	8.6	*
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	8.6	*
5	Malignant neoplasms (C00-C97)	2	5.7	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	2.9	*
6	Certain conditions originating in the perinatal period (P00-P96)	1	2.9	*
...	All other causes (Residual)	12	34.3	*
American Indian, female, 5-14 years				
...	All causes	58	100.0	21.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17	29.3	*
2	Malignant neoplasms (C00-C97)	7	12.1	*
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4	6.9	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	5.2	*
5	Septicemia (A40-A41)	2	3.4	*
5	Influenza and pneumonia (J09-J18)	2	3.4	*
5	Chronic lower respiratory diseases (J40-J47)	2	3.4	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	3.4	*
9	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	1.7	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	1.7	*
...	All other causes (Residual)	17	29.3	*
American Indian, female, 15-24 years				
...	All causes	202	100.0	67.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	103	51.0	34.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	40	19.8	13.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	17	8.4	*
4	Malignant neoplasms (C00-C97)	4	2.0	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	2.0	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	1.5	*
7	Cerebrovascular diseases (I60-I69)	2	1.0	*
7	Influenza and pneumonia (J09-J18)	2	1.0	*
7	Chronic lower respiratory diseases (J40-J47)	2	1.0	*
10	Septicemia (A40-A41)	1	0.5	*
10	Diabetes mellitus (E10-E14)	1	0.5	*
10	Pregnancy, childbirth and the puerperium (O00-O99)	1	0.5	*
...	All other causes (Residual)	22	10.9	7.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 25-34 years				
...	All causes	250	100.0	98.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	90	36.0	35.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	25	10.0	9.8
3	Malignant neoplasms (C00-C97)	20	8.0	7.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	18	7.2	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	9	3.6	*
6	Influenza and pneumonia (J09-J18)	8	3.2	*
6	Pregnancy, childbirth and the puerperium (O00-O99)	8	3.2	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	2.8	*
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7	2.8	*
10	Septicemia (A40-A41)	6	2.4	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	3	1.2	*
11	Diabetes mellitus (E10-E14)	3	1.2	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	1.2	*
14	Viral hepatitis (B15-B19)	2	0.8	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2	0.8	*
...	All other causes (Residual)	39	15.6	15.3
American Indian, female, 35-44 years				
...	All causes	479	100.0	216.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	115	24.0	51.9
2	Chronic liver disease and cirrhosis (K70,K73-K74)	65	13.6	29.3
3	Malignant neoplasms (C00-C97)	57	11.9	25.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	43	9.0	19.4
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	19	4.0	*
6	Diabetes mellitus (E10-E14)	15	3.1	*
6	Influenza and pneumonia (J09-J18)	15	3.1	*
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	11	2.3	*
9	Septicemia (A40-A41)	10	2.1	*
10	Viral hepatitis (B15-B19)	9	1.9	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	1.5	*
12	Chronic lower respiratory diseases (J40-J47)	6	1.3	*
13	Cerebrovascular diseases (I60-I69)	5	1.0	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	3	0.6	*
14	Pregnancy, childbirth and the puerperium (O00-O99)	3	0.6	*
...	All other causes (Residual)	96	20.0	43.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 45-54 years				
...	All causes	924	100.0	403.2
1	Malignant neoplasms (C00-C97)	188	20.3	82.0
2	Chronic liver disease and cirrhosis (K70,K73-K74)	119	12.9	51.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	112	12.1	48.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	104	11.3	45.4
5	Diabetes mellitus (E10-E14)	45	4.9	19.6
6	Influenza and pneumonia (J09-J18)	36	3.9	15.7
7	Cerebrovascular diseases (I60-I69)	31	3.4	13.5
8	Chronic lower respiratory diseases (J40-J47)	29	3.1	12.7
9	Viral hepatitis (B15-B19)	21	2.3	9.2
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	1.8	*
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	17	1.8	*
12	Septicemia (A40-A41)	15	1.6	*
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	10	1.1	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	9	1.0	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	6	0.6	*
...	All other causes (Residual)	165	17.9	72.0
American Indian, female, 55-64 years				
...	All causes	1,029	100.0	647.5
1	Malignant neoplasms (C00-C97)	312	30.3	196.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	174	16.9	109.5
3	Chronic liver disease and cirrhosis (K70,K73-K74)	67	6.5	42.2
4	Diabetes mellitus (E10-E14)	51	5.0	32.1
5	Chronic lower respiratory diseases (J40-J47)	47	4.6	29.6
6	Cerebrovascular diseases (I60-I69)	42	4.1	26.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	42	4.1	26.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	29	2.8	18.2
9	Influenza and pneumonia (J09-J18)	23	2.2	14.5
10	Septicemia (A40-A41)	14	1.4	*
11	Viral hepatitis (B15-B19)	9	0.9	*
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	9	0.9	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	8	0.8	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.7	*
15	Pneumonitis due to solids and liquids (J69)	6	0.6	*
...	All other causes (Residual)	189	18.4	118.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 65-74 years				
...	All causes	1,233	100.0	1,512.9
1	Malignant neoplasms (C00-C97)	347	28.1	425.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	220	17.8	269.9
3	Diabetes mellitus (E10-E14)	96	7.8	117.8
4	Chronic lower respiratory diseases (J40-J47)	91	7.4	111.7
5	Cerebrovascular diseases (I60-I69)	58	4.7	71.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	47	3.8	57.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	43	3.5	52.8
8	Influenza and pneumonia (J09-J18)	35	2.8	42.9
9	Septicemia (A40-A41)	32	2.6	39.3
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	25	2.0	30.7
11	Alzheimer's disease (G30)	14	1.1	*
12	Pneumonitis due to solids and liquids (J69)	12	1.0	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10	0.8	*
14	Parkinson's disease (G20-G21)	7	0.6	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.5	*
...	All other causes (Residual)	190	15.4	233.1
American Indian, female, 75-84 years				
...	All causes	1,316	100.0	3,117.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	290	22.0	687.0
2	Malignant neoplasms (C00-C97)	275	20.9	651.5
3	Diabetes mellitus (E10-E14)	110	8.4	260.6
4	Chronic lower respiratory diseases (J40-J47)	91	6.9	215.6
5	Cerebrovascular diseases (I60-I69)	67	5.1	158.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	46	3.5	109.0
7	Alzheimer's disease (G30)	43	3.3	101.9
8	Septicemia (A40-A41)	37	2.8	87.7
9	Influenza and pneumonia (J09-J18)	30	2.3	71.1
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	29	2.2	68.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	16	1.2	*
12	Parkinson's disease (G20-G21)	14	1.1	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.7	*
14	Pneumonitis due to solids and liquids (J69)	7	0.5	*
15	Aortic aneurysm and dissection (I71)	6	0.5	*
...	All other causes (Residual)	246	18.7	582.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 85 years and over				
...	All causes	1,158	100.0	6,569.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	316	27.3	1,792.7
2	Malignant neoplasms (C00-C97)	116	10.0	658.1
3	Cerebrovascular diseases (I60-I69)	93	8.0	527.6
4	Alzheimer's disease (G30)	69	6.0	391.4
5	Chronic lower respiratory diseases (J40-J47)	55	4.7	312.0
6	Influenza and pneumonia (J09-J18)	51	4.4	289.3
7	Diabetes mellitus (E10-E14)	47	4.1	266.6
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	41	3.5	232.6
9	Septicemia (A40-A41)	30	2.6	170.2
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	27	2.3	153.2
11	Pneumonitis due to solids and liquids (J69)	16	1.4	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	13	1.1	*
13	Anemias (D50-D64)	7	0.6	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.5	*
14	Aortic aneurysm and dissection (I71)	6	0.5	*
14	Chronic liver disease and cirrhosis (K70,K73-K74)	6	0.5	*
...	All other causes (Residual)	259	22.4	1,469.3
Asian or Pacific Islander, both sexes, all ages				
...	All causes	49,225	100.0	317.0
1	Malignant neoplasms (C00-C97)	13,274	27.0	85.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,375	23.1	73.3
3	Cerebrovascular diseases (I60-I69)	3,639	7.4	23.4
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,044	4.2	13.2
5	Diabetes mellitus (E10-E14)	1,846	3.8	11.9
6	Influenza and pneumonia (J09-J18)	1,568	3.2	10.1
7	Chronic lower respiratory diseases (J40-J47)	1,480	3.0	9.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,062	2.2	6.8
9	Alzheimer's disease (G30)	1,024	2.1	6.6
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	971	2.0	6.3
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	710	1.4	4.6
12	Septicemia (A40-A41)	576	1.2	3.7
13	Chronic liver disease and cirrhosis (K70,K73-K74)	464	0.9	3.0
14	Certain conditions originating in the perinatal period (P00-P96)	455	0.9	2.9
15	Parkinson's disease (G20-G21)	367	0.7	2.4
...	All other causes (Residual)	8,370	17.0	53.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 1-4 years				
...	All causes	149	100.0	15.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	49	32.9	5.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	16	10.7	*
3	Malignant neoplasms (C00-C97)	13	8.7	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	10	6.7	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	7	4.7	*
6	Septicemia (A40-A41)	5	3.4	*
7	Influenza and pneumonia (J09-J18)	3	2.0	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	1.3	*
9	Nutritional deficiencies (E40-E64)	1	0.7	*
9	Meningitis (G00,G03)	1	0.7	*
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.7	*
9	Acute bronchitis and bronchiolitis (J20-J21)	1	0.7	*
9	Chronic lower respiratory diseases (J40-J47)	1	0.7	*
9	Hernia (K40-K46)	1	0.7	*
9	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.7	*
...	All other causes (Residual)	37	24.8	3.9
Asian or Pacific Islander, both sexes, 5-14 years				
...	All causes	231	100.0	11.1
1	Malignant neoplasms (C00-C97)	47	20.3	2.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	47	20.3	2.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	20	8.7	1.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	15	6.5	*
5	Influenza and pneumonia (J09-J18)	11	4.8	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8	3.5	*
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6	2.6	*
8	Chronic lower respiratory diseases (J40-J47)	5	2.2	*
9	Meningitis (G00,G03)	3	1.3	*
9	Cerebrovascular diseases (I60-I69)	3	1.3	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.4	*
11	Diseases of appendix (K35-K38)	1	0.4	*
11	Hernia (K40-K46)	1	0.4	*
11	Certain conditions originating in the perinatal period (P00-P96)	1	0.4	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.4	*
...	All other causes (Residual)	61	26.4	2.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 15-24 years				
...	All causes	763	100.0	37.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	256	33.6	12.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	152	19.9	7.5
3	Malignant neoplasms (C00-C97)	75	9.8	3.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	72	9.4	3.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	37	4.8	1.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	17	2.2	*
7	Influenza and pneumonia (J09-J18)	15	2.0	*
8	Cerebrovascular diseases (I60-I69)	12	1.6	*
9	Chronic lower respiratory diseases (J40-J47)	5	0.7	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.7	*
9	Pregnancy, childbirth and the puerperium (O00-O99)	5	0.7	*
12	Septicemia (A40-A41)	4	0.5	*
13	Tuberculosis (A16-A19)	3	0.4	*
13	Anemias (D50-D64)	3	0.4	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.3	*
15	Diabetes mellitus (E10-E14)	2	0.3	*
...	All other causes (Residual)	98	12.8	4.8
Asian or Pacific Islander, both sexes, 25-34 years				
...	All causes	1,065	100.0	41.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	251	23.6	9.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	186	17.5	7.2
3	Malignant neoplasms (C00-C97)	163	15.3	6.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	96	9.0	3.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	67	6.3	2.6
6	Influenza and pneumonia (J09-J18)	35	3.3	1.3
7	Cerebrovascular diseases (I60-I69)	27	2.5	1.0
8	Pregnancy, childbirth and the puerperium (O00-O99)	15	1.4	*
9	Septicemia (A40-A41)	11	1.0	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	11	1.0	*
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	11	1.0	*
12	Diabetes mellitus (E10-E14)	7	0.7	*
12	Chronic lower respiratory diseases (J40-J47)	7	0.7	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.6	*
15	Viral hepatitis (B15-B19)	4	0.4	*
15	Aortic aneurysm and dissection (I71)	4	0.4	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	4	0.4	*
15	Legal intervention (Y35,Y89.0)	4	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.4	*
...	All other causes (Residual)	152	14.3	5.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 35-44 years				
...	All causes	1,914	100.0	74.0
1	Malignant neoplasms (C00-C97)	572	29.9	22.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	290	15.2	11.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	204	10.7	7.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	180	9.4	7.0
5	Cerebrovascular diseases (I60-I69)	112	5.9	4.3
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	62	3.2	2.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	55	2.9	2.1
8	Influenza and pneumonia (J09-J18)	38	2.0	1.5
9	Diabetes mellitus (E10-E14)	32	1.7	1.2
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	26	1.4	1.0
11	Human immunodeficiency virus (HIV) disease (B20-B24)	19	1.0	*
12	Septicemia (A40-A41)	18	0.9	*
13	Chronic lower respiratory diseases (J40-J47)	15	0.8	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	0.7	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	14	0.7	*
...	All other causes (Residual)	263	13.7	10.2
Asian or Pacific Islander, both sexes, 45-54 years				
...	All causes	3,839	100.0	181.4
1	Malignant neoplasms (C00-C97)	1,430	37.2	67.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	704	18.3	33.3
3	Cerebrovascular diseases (I60-I69)	247	6.4	11.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	222	5.8	10.5
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	177	4.6	8.4
6	Diabetes mellitus (E10-E14)	121	3.2	5.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	102	2.7	4.8
8	Influenza and pneumonia (J09-J18)	69	1.8	3.3
9	Viral hepatitis (B15-B19)	61	1.6	2.9
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	58	1.5	2.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	48	1.3	2.3
12	Chronic lower respiratory diseases (J40-J47)	42	1.1	2.0
13	Septicemia (A40-A41)	38	1.0	1.8
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	32	0.8	1.5
15	Aortic aneurysm and dissection (I71)	26	0.7	1.2
...	All other causes (Residual)	462	12.0	21.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 55-64 years				
...	All causes	6,297	100.0	418.0
1	Malignant neoplasms (C00-C97)	2,679	42.5	177.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,279	20.3	84.9
3	Cerebrovascular diseases (I60-I69)	407	6.5	27.0
4	Diabetes mellitus (E10-E14)	282	4.5	18.7
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	190	3.0	12.6
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	135	2.1	9.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	113	1.8	7.5
8	Influenza and pneumonia (J09-J18)	95	1.5	6.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	94	1.5	6.2
10	Chronic lower respiratory diseases (J40-J47)	77	1.2	5.1
11	Septicemia (A40-A41)	72	1.1	4.8
12	Viral hepatitis (B15-B19)	68	1.1	4.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	49	0.8	3.3
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	40	0.6	2.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	36	0.6	2.4
...	All other causes (Residual)	681	10.8	45.2
Asian or Pacific Islander, both sexes, 65-74 years				
...	All causes	8,440	100.0	1,029.8
1	Malignant neoplasms (C00-C97)	3,087	36.6	376.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,914	22.7	233.5
3	Cerebrovascular diseases (I60-I69)	588	7.0	71.7
4	Diabetes mellitus (E10-E14)	406	4.8	49.5
5	Chronic lower respiratory diseases (J40-J47)	236	2.8	28.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	215	2.5	26.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	213	2.5	26.0
8	Influenza and pneumonia (J09-J18)	171	2.0	20.9
9	Septicemia (A40-A41)	124	1.5	15.1
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	100	1.2	12.2
11	Chronic liver disease and cirrhosis (K70,K73-K74)	92	1.1	11.2
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	67	0.8	8.2
13	Viral hepatitis (B15-B19)	59	0.7	7.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	56	0.7	6.8
15	Aortic aneurysm and dissection (I71)	53	0.6	6.5
...	All other causes (Residual)	1,059	12.5	129.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 75-84 years				
...	All causes	12,560	100.0	2,830.7
1	Malignant neoplasms (C00-C97)	3,363	26.8	757.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,097	24.7	698.0
3	Cerebrovascular diseases (I60-I69)	1,109	8.8	249.9
4	Diabetes mellitus (E10-E14)	555	4.4	125.1
5	Chronic lower respiratory diseases (J40-J47)	522	4.2	117.6
6	Influenza and pneumonia (J09-J18)	397	3.2	89.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	335	2.7	75.5
8	Alzheimer's disease (G30)	312	2.5	70.3
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	307	2.4	69.2
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	213	1.7	48.0
11	Parkinson's disease (G20-G21)	161	1.3	36.3
12	Septicemia (A40-A41)	155	1.2	34.9
13	Aortic aneurysm and dissection (I71)	101	0.8	22.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	93	0.7	21.0
15	Pneumonitis due to solids and liquids (J69)	86	0.7	19.4
...	All other causes (Residual)	1,754	14.0	395.3
Asian or Pacific Islander, both sexes, 85 years and over				
...	All causes	13,061	100.0	7,996.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,920	30.0	2,400.0
2	Malignant neoplasms (C00-C97)	1,839	14.1	1,125.9
3	Cerebrovascular diseases (I60-I69)	1,130	8.7	691.8
4	Influenza and pneumonia (J09-J18)	722	5.5	442.0
5	Alzheimer's disease (G30)	661	5.1	404.7
6	Chronic lower respiratory diseases (J40-J47)	569	4.4	348.4
7	Diabetes mellitus (E10-E14)	441	3.4	270.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	335	2.6	205.1
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	300	2.3	183.7
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	282	2.2	172.7
11	Pneumonitis due to solids and liquids (J69)	167	1.3	102.2
12	Septicemia (A40-A41)	144	1.1	88.2
13	Parkinson's disease (G20-G21)	141	1.1	86.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	78	0.6	47.8
15	Aortic aneurysm and dissection (I71)	54	0.4	33.1
...	All other causes (Residual)	2,278	17.4	1,394.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, all ages				
...	All causes	25,560	100.0	339.5
1	Malignant neoplasms (C00-C97)	6,897	27.0	91.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,238	24.4	82.9
3	Cerebrovascular diseases (I60-I69)	1,658	6.5	22.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,288	5.0	17.1
5	Diabetes mellitus (E10-E14)	943	3.7	12.5
6	Chronic lower respiratory diseases (J40-J47)	864	3.4	11.5
7	Influenza and pneumonia (J09-J18)	809	3.2	10.7
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	666	2.6	8.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	518	2.0	6.9
10	Alzheimer's disease (G30)	318	1.2	4.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	300	1.2	4.0
12	Septicemia (A40-A41)	293	1.1	3.9
13	Chronic liver disease and cirrhosis (K70,K73-K74)	289	1.1	3.8
14	Certain conditions originating in the perinatal period (P00-P96)	246	1.0	3.3
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	235	0.9	3.1
...	All other causes (Residual)	3,998	15.6	53.1
Asian or Pacific Islander, male, 1-4 years				
...	All causes	91	100.0	18.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	28	30.8	5.8
2	Malignant neoplasms (C00-C97)	9	9.9	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	8.8	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	5.5	*
5	Septicemia (A40-A41)	4	4.4	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	3.3	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.1	*
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	1.1	*
7	Influenza and pneumonia (J09-J18)	1	1.1	*
7	Acute bronchitis and bronchiolitis (J20-J21)	1	1.1	*
...	All other causes (Residual)	30	33.0	6.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 5-14 years				
...	All causes	122	100.0	11.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	29	23.8	2.7
2	Malignant neoplasms (C00-C97)	28	23.0	2.6
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	7.4	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	6	4.9	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	4.9	*
6	Influenza and pneumonia (J09-J18)	4	3.3	*
7	Cerebrovascular diseases (I60-I69)	2	1.6	*
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2	1.6	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.8	*
9	Meningitis (G00,G03)	1	0.8	*
9	Chronic lower respiratory diseases (J40-J47)	1	0.8	*
9	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.8	*
...	All other causes (Residual)	32	26.2	3.0
Asian or Pacific Islander, male, 15-24 years				
...	All causes	511	100.0	49.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	185	36.2	17.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	99	19.4	9.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	55	10.8	5.3
4	Malignant neoplasms (C00-C97)	45	8.8	4.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	26	5.1	2.5
6	Influenza and pneumonia (J09-J18)	9	1.8	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	1.4	*
8	Cerebrovascular diseases (I60-I69)	6	1.2	*
9	Septicemia (A40-A41)	3	0.6	*
10	Tuberculosis (A16-A19)	2	0.4	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.4	*
10	Anemias (D50-D64)	2	0.4	*
10	Diabetes mellitus (E10-E14)	2	0.4	*
10	Chronic lower respiratory diseases (J40-J47)	2	0.4	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
...	All other causes (Residual)	64	12.5	6.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 25-34 years				
...	All causes	679	100.0	53.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	179	26.4	14.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	130	19.1	10.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	81	11.9	6.4
4	Malignant neoplasms (C00-C97)	77	11.3	6.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	48	7.1	3.8
6	Influenza and pneumonia (J09-J18)	14	2.1	*
7	Cerebrovascular diseases (I60-I69)	12	1.8	*
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	1.2	*
9	Septicemia (A40-A41)	7	1.0	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	7	1.0	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.7	*
11	Diabetes mellitus (E10-E14)	5	0.7	*
11	Chronic lower respiratory diseases (J40-J47)	5	0.7	*
14	Aortic aneurysm and dissection (I71)	4	0.6	*
14	Legal intervention (Y35,Y89.0)	4	0.6	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.6	*
...	All other causes (Residual)	89	13.1	7.0
Asian or Pacific Islander, male, 35-44 years				
...	All causes	1,194	100.0	95.1
1	Malignant neoplasms (C00-C97)	270	22.6	21.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	239	20.0	19.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	154	12.9	12.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	124	10.4	9.9
5	Cerebrovascular diseases (I60-I69)	63	5.3	5.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	45	3.8	3.6
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	37	3.1	2.9
8	Influenza and pneumonia (J09-J18)	28	2.3	2.2
9	Diabetes mellitus (E10-E14)	22	1.8	1.8
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	1.3	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	13	1.1	*
12	Septicemia (A40-A41)	11	0.9	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.9	*
12	Chronic lower respiratory diseases (J40-J47)	11	0.9	*
15	Aortic aneurysm and dissection (I71)	8	0.7	*
...	All other causes (Residual)	143	12.0	11.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 45-54 years				
...	All causes	2,379	100.0	236.3
1	Malignant neoplasms (C00-C97)	709	29.8	70.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	568	23.9	56.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	171	7.2	17.0
4	Cerebrovascular diseases (I60-I69)	135	5.7	13.4
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	115	4.8	11.4
6	Diabetes mellitus (E10-E14)	93	3.9	9.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	76	3.2	7.5
8	Viral hepatitis (B15-B19)	40	1.7	4.0
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	39	1.6	3.9
10	Influenza and pneumonia (J09-J18)	35	1.5	3.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	30	1.3	3.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	26	1.1	2.6
13	Chronic lower respiratory diseases (J40-J47)	24	1.0	2.4
14	Human immunodeficiency virus (HIV) disease (B20-B24)	19	0.8	*
15	Septicemia (A40-A41)	17	0.7	*
15	Aortic aneurysm and dissection (I71)	17	0.7	*
...	All other causes (Residual)	265	11.1	26.3
Asian or Pacific Islander, male, 55-64 years				
...	All causes	3,709	100.0	536.0
1	Malignant neoplasms (C00-C97)	1,407	37.9	203.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	940	25.3	135.9
3	Cerebrovascular diseases (I60-I69)	226	6.1	32.7
4	Diabetes mellitus (E10-E14)	170	4.6	24.6
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	122	3.3	17.6
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	100	2.7	14.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	85	2.3	12.3
8	Influenza and pneumonia (J09-J18)	61	1.6	8.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	56	1.5	8.1
10	Viral hepatitis (B15-B19)	43	1.2	6.2
10	Chronic lower respiratory diseases (J40-J47)	43	1.2	6.2
12	Septicemia (A40-A41)	39	1.1	5.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	31	0.8	4.5
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	26	0.7	3.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	22	0.6	3.2
...	All other causes (Residual)	338	9.1	48.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 65-74 years				
...	All causes	4,783	100.0	1,277.0
1	Malignant neoplasms (C00-C97)	1,671	34.9	446.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,207	25.2	322.2
3	Cerebrovascular diseases (I60-I69)	299	6.3	79.8
4	Diabetes mellitus (E10-E14)	246	5.1	65.7
5	Chronic lower respiratory diseases (J40-J47)	148	3.1	39.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	113	2.4	30.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	112	2.3	29.9
8	Influenza and pneumonia (J09-J18)	98	2.0	26.2
9	Septicemia (A40-A41)	68	1.4	18.2
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	53	1.1	14.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	47	1.0	12.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	46	1.0	12.3
13	Parkinson's disease (G20-G21)	40	0.8	10.7
14	Aortic aneurysm and dissection (I71)	35	0.7	9.3
15	Viral hepatitis (B15-B19)	34	0.7	9.1
...	All other causes (Residual)	566	11.8	151.1
Asian or Pacific Islander, male, 75-84 years				
...	All causes	6,167	100.0	3,406.3
1	Malignant neoplasms (C00-C97)	1,763	28.6	973.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,560	25.3	861.7
3	Cerebrovascular diseases (I60-I69)	496	8.0	274.0
4	Chronic lower respiratory diseases (J40-J47)	307	5.0	169.6
5	Diabetes mellitus (E10-E14)	232	3.8	128.1
6	Influenza and pneumonia (J09-J18)	224	3.6	123.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	159	2.6	87.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	146	2.4	80.6
9	Alzheimer's disease (G30)	107	1.7	59.1
10	Parkinson's disease (G20-G21)	100	1.6	55.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	92	1.5	50.8
12	Septicemia (A40-A41)	75	1.2	41.4
13	Aortic aneurysm and dissection (I71)	53	0.9	29.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	50	0.8	27.6
15	Pneumonitis due to solids and liquids (J69)	42	0.7	23.2
...	All other causes (Residual)	761	12.3	420.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 85 years and over				
...	All causes	5,435	100.0	9,109.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,598	29.4	2,678.3
2	Malignant neoplasms (C00-C97)	914	16.8	1,531.9
3	Cerebrovascular diseases (I60-I69)	415	7.6	695.6
4	Influenza and pneumonia (J09-J18)	330	6.1	553.1
5	Chronic lower respiratory diseases (J40-J47)	323	5.9	541.4
6	Alzheimer's disease (G30)	189	3.5	316.8
7	Diabetes mellitus (E10-E14)	173	3.2	290.0
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	146	2.7	244.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	141	2.6	236.3
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	97	1.8	162.6
11	Pneumonitis due to solids and liquids (J69)	92	1.7	154.2
12	Parkinson's disease (G20-G21)	80	1.5	134.1
13	Septicemia (A40-A41)	67	1.2	112.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	29	0.5	48.6
15	Aortic aneurysm and dissection (I71)	27	0.5	45.3
...	All other causes (Residual)	814	15.0	1,364.3
Asian or Pacific Islander, female, all ages				
...	All causes	23,665	100.0	295.9
1	Malignant neoplasms (C00-C97)	6,377	26.9	79.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,137	21.7	64.2
3	Cerebrovascular diseases (I60-I69)	1,981	8.4	24.8
4	Diabetes mellitus (E10-E14)	903	3.8	11.3
5	Influenza and pneumonia (J09-J18)	759	3.2	9.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	756	3.2	9.5
7	Alzheimer's disease (G30)	706	3.0	8.8
8	Chronic lower respiratory diseases (J40-J47)	616	2.6	7.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	544	2.3	6.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	410	1.7	5.1
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	305	1.3	3.8
12	Septicemia (A40-A41)	283	1.2	3.5
13	Certain conditions originating in the perinatal period (P00-P96)	209	0.9	2.6
14	Chronic liver disease and cirrhosis (K70,K73-K74)	175	0.7	2.2
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	174	0.7	2.2
...	All other causes (Residual)	4,330	18.3	54.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 1-4 years				
...	All causes	58	100.0	12.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	21	36.2	4.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	13.8	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	8.6	*
4	Malignant neoplasms (C00-C97)	4	6.9	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	6.9	*
6	Influenza and pneumonia (J09-J18)	2	3.4	*
7	Septicemia (A40-A41)	1	1.7	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.7	*
7	Nutritional deficiencies (E40-E64)	1	1.7	*
7	Meningitis (G00,G03)	1	1.7	*
7	Chronic lower respiratory diseases (J40-J47)	1	1.7	*
7	Hernia (K40-K46)	1	1.7	*
7	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	1.7	*
...	All other causes (Residual)	7	12.1	*
Asian or Pacific Islander, female, 5-14 years				
...	All causes	109	100.0	10.6
1	Malignant neoplasms (C00-C97)	19	17.4	*
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18	16.5	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	11	10.1	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	9	8.3	*
5	Influenza and pneumonia (J09-J18)	7	6.4	*
6	Chronic lower respiratory diseases (J40-J47)	4	3.7	*
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4	3.7	*
8	Meningitis (G00,G03)	2	1.8	*
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	1.8	*
10	Cerebrovascular diseases (I60-I69)	1	0.9	*
10	Diseases of appendix (K35-K38)	1	0.9	*
10	Hernia (K40-K46)	1	0.9	*
10	Certain conditions originating in the perinatal period (P00-P96)	1	0.9	*
...	All other causes (Residual)	29	26.6	2.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 15-24 years				
...	All causes	252	100.0	25.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	71	28.2	7.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	53	21.0	5.3
3	Malignant neoplasms (C00-C97)	30	11.9	3.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	17	6.7	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	11	4.4	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	10	4.0	*
7	Cerebrovascular diseases (I60-I69)	6	2.4	*
7	Influenza and pneumonia (J09-J18)	6	2.4	*
9	Pregnancy, childbirth and the puerperium (O00-O99)	5	2.0	*
10	Chronic lower respiratory diseases (J40-J47)	3	1.2	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	1.2	*
12	Tuberculosis (A16-A19)	1	0.4	*
12	Septicemia (A40-A41)	1	0.4	*
12	Anemias (D50-D64)	1	0.4	*
...	All other causes (Residual)	34	13.5	3.4
Asian or Pacific Islander, female, 25-34 years				
...	All causes	386	100.0	29.0
1	Malignant neoplasms (C00-C97)	86	22.3	6.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	72	18.7	5.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	56	14.5	4.2
4	Influenza and pneumonia (J09-J18)	21	5.4	1.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	19	4.9	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	15	3.9	*
6	Cerebrovascular diseases (I60-I69)	15	3.9	*
6	Pregnancy, childbirth and the puerperium (O00-O99)	15	3.9	*
9	Septicemia (A40-A41)	4	1.0	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	4	1.0	*
11	Chronic liver disease and cirrhosis (K70,K73-K74)	3	0.8	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.8	*
13	Diabetes mellitus (E10-E14)	2	0.5	*
13	Chronic lower respiratory diseases (J40-J47)	2	0.5	*
15	Viral hepatitis (B15-B19)	1	0.3	*
15	Malaria (B50-B54)	1	0.3	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.3	*
15	Anemias (D50-D64)	1	0.3	*
15	Pneumonitis due to solids and liquids (J69)	1	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
...	All other causes (Residual)	63	16.3	4.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 35-44 years				
...	All causes	720	100.0	54.1
1	Malignant neoplasms (C00-C97)	302	41.9	22.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	56	7.8	4.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	51	7.1	3.8
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	50	6.9	3.8
5	Cerebrovascular diseases (I60-I69)	49	6.8	3.7
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	25	3.5	1.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	1.5	*
7	Pregnancy, childbirth and the puerperium (O00-O99)	11	1.5	*
9	Diabetes mellitus (E10-E14)	10	1.4	*
9	Influenza and pneumonia (J09-J18)	10	1.4	*
9	Chronic liver disease and cirrhosis (K70,K73-K74)	10	1.4	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	8	1.1	*
13	Septicemia (A40-A41)	7	1.0	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	6	0.8	*
15	Viral hepatitis (B15-B19)	4	0.6	*
15	Chronic lower respiratory diseases (J40-J47)	4	0.6	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	0.6	*
...	All other causes (Residual)	102	14.2	7.7
Asian or Pacific Islander, female, 45-54 years				
...	All causes	1,460	100.0	131.5
1	Malignant neoplasms (C00-C97)	721	49.4	65.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	136	9.3	12.3
3	Cerebrovascular diseases (I60-I69)	112	7.7	10.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	62	4.2	5.6
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	51	3.5	4.6
6	Influenza and pneumonia (J09-J18)	34	2.3	3.1
7	Diabetes mellitus (E10-E14)	28	1.9	2.5
8	Chronic liver disease and cirrhosis (K70,K73-K74)	26	1.8	2.3
9	Septicemia (A40-A41)	21	1.4	1.9
9	Viral hepatitis (B15-B19)	21	1.4	1.9
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	19	1.3	*
12	Chronic lower respiratory diseases (J40-J47)	18	1.2	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18	1.2	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	12	0.8	*
15	Aortic aneurysm and dissection (I71)	9	0.6	*
...	All other causes (Residual)	172	11.8	15.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 55-64 years				
...	All causes	2,588	100.0	317.8
1	Malignant neoplasms (C00-C97)	1,272	49.1	156.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	339	13.1	41.6
3	Cerebrovascular diseases (I60-I69)	181	7.0	22.2
4	Diabetes mellitus (E10-E14)	112	4.3	13.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	68	2.6	8.3
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	38	1.5	4.7
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	35	1.4	4.3
8	Influenza and pneumonia (J09-J18)	34	1.3	4.2
8	Chronic lower respiratory diseases (J40-J47)	34	1.3	4.2
10	Septicemia (A40-A41)	33	1.3	4.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	28	1.1	3.4
12	Viral hepatitis (B15-B19)	25	1.0	3.1
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	18	0.7	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	0.5	*
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	14	0.5	*
...	All other causes (Residual)	343	13.3	42.1
Asian or Pacific Islander, female, 65-74 years				
...	All causes	3,657	100.0	821.7
1	Malignant neoplasms (C00-C97)	1,416	38.7	318.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	707	19.3	158.9
3	Cerebrovascular diseases (I60-I69)	289	7.9	64.9
4	Diabetes mellitus (E10-E14)	160	4.4	36.0
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	102	2.8	22.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	101	2.8	22.7
7	Chronic lower respiratory diseases (J40-J47)	88	2.4	19.8
8	Influenza and pneumonia (J09-J18)	73	2.0	16.4
9	Septicemia (A40-A41)	56	1.5	12.6
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	54	1.5	12.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	45	1.2	10.1
12	Alzheimer's disease (G30)	29	0.8	6.5
13	Viral hepatitis (B15-B19)	25	0.7	5.6
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	25	0.7	5.6
15	Aortic aneurysm and dissection (I71)	18	0.5	*
...	All other causes (Residual)	469	12.8	105.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 75-84 years				
...	All causes	6,393	100.0	2,434.0
1	Malignant neoplasms (C00-C97)	1,600	25.0	609.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,537	24.0	585.2
3	Cerebrovascular diseases (I60-I69)	613	9.6	233.4
4	Diabetes mellitus (E10-E14)	323	5.1	123.0
5	Chronic lower respiratory diseases (J40-J47)	215	3.4	81.9
6	Alzheimer's disease (G30)	205	3.2	78.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	176	2.8	67.0
8	Influenza and pneumonia (J09-J18)	173	2.7	65.9
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	161	2.5	61.3
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	121	1.9	46.1
11	Septicemia (A40-A41)	80	1.3	30.5
12	Parkinson's disease (G20-G21)	61	1.0	23.2
13	Aortic aneurysm and dissection (I71)	48	0.8	18.3
14	Viral hepatitis (B15-B19)	44	0.7	16.8
14	Pneumonitis due to solids and liquids (J69)	44	0.7	16.8
...	All other causes (Residual)	992	15.5	377.7
Asian or Pacific Islander, female, 85 years and over				
...	All causes	7,626	100.0	7,356.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,322	30.4	2,239.8
2	Malignant neoplasms (C00-C97)	925	12.1	892.3
3	Cerebrovascular diseases (I60-I69)	715	9.4	689.7
4	Alzheimer's disease (G30)	472	6.2	455.3
5	Influenza and pneumonia (J09-J18)	392	5.1	378.1
6	Diabetes mellitus (E10-E14)	268	3.5	258.5
7	Chronic lower respiratory diseases (J40-J47)	246	3.2	237.3
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	203	2.7	195.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	194	2.5	187.1
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	136	1.8	131.2
11	Septicemia (A40-A41)	77	1.0	74.3
12	Pneumonitis due to solids and liquids (J69)	75	1.0	72.3
13	Parkinson's disease (G20-G21)	61	0.8	58.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	49	0.6	47.3
15	Atherosclerosis (I70)	34	0.4	32.8
...	All other causes (Residual)	1,457	19.1	1,405.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable