

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, all ages				
...	All causes	2,468,435	100.0	799.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	597,689	24.2	193.6
2	Malignant neoplasms (C00-C97)	574,743	23.3	186.2
3	Chronic lower respiratory diseases (J40-J47)	138,080	5.6	44.7
4	Cerebrovascular diseases (I60-I69)	129,476	5.2	41.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	120,859	4.9	39.1
6	Alzheimer's disease (G30)	83,494	3.4	27.0
7	Diabetes mellitus (E10-E14)	69,071	2.8	22.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	50,476	2.0	16.3
9	Influenza and pneumonia (J09-J18)	50,097	2.0	16.2
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	38,364	1.6	12.4
11	Septicemia (A40-A41)	34,812	1.4	11.3
12	Chronic liver disease and cirrhosis (K70,K73-K74)	31,903	1.3	10.3
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	26,634	1.1	8.6
14	Parkinson's disease (G20-G21)	22,032	0.9	7.1
15	Pneumonitis due to solids and liquids (J69)	17,011	0.7	5.5
...	All other causes (Residual)	483,694	19.6	156.7
All races, both sexes, 1-4 years				
...	All causes	4,316	100.0	26.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,394	32.3	8.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	507	11.7	3.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	385	8.9	2.4
4	Malignant neoplasms (C00-C97)	346	8.0	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	159	3.7	1.0
6	Influenza and pneumonia (J09-J18)	91	2.1	0.6
7	Septicemia (A40-A41)	62	1.4	0.4
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	59	1.4	0.4
9	Certain conditions originating in the perinatal period (P00-P96)	52	1.2	0.3
10	Chronic lower respiratory diseases (J40-J47)	51	1.2	0.3
11	Cerebrovascular diseases (I60-I69)	50	1.2	0.3
12	Anemias (D50-D64)	29	0.7	0.2
13	Meningitis (G00,G03)	22	0.5	0.1
14	Complications of medical and surgical care (Y40-Y84,Y88)	19	0.4	*
15	Pneumonitis due to solids and liquids (J69)	13	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	0.3	*
...	All other causes (Residual)	1,064	24.7	6.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 5-14 years				
...	All causes	5,279	100.0	12.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,643	31.1	4.0
2	Malignant neoplasms (C00-C97)	916	17.4	2.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	298	5.6	0.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	274	5.2	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	261	4.9	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	185	3.5	0.5
7	Chronic lower respiratory diseases (J40-J47)	133	2.5	0.3
8	Cerebrovascular diseases (I60-I69)	90	1.7	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	82	1.6	0.2
10	Influenza and pneumonia (J09-J18)	71	1.3	0.2
11	Septicemia (A40-A41)	67	1.3	0.2
12	Anemias (D50-D64)	26	0.5	0.1
12	Diabetes mellitus (E10-E14)	26	0.5	0.1
14	Certain conditions originating in the perinatal period (P00-P96)	25	0.5	0.1
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22	0.4	0.1
15	Complications of medical and surgical care (Y40-Y84,Y88)	22	0.4	0.1
...	All other causes (Residual)	1,138	21.6	2.8
All races, both sexes, 15-24 years				
...	All causes	29,551	100.0	67.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,341	41.8	28.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4,678	15.8	10.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,600	15.6	10.5
4	Malignant neoplasms (C00-C97)	1,604	5.4	3.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,028	3.5	2.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	412	1.4	0.9
7	Cerebrovascular diseases (I60-I69)	190	0.6	0.4
8	Influenza and pneumonia (J09-J18)	181	0.6	0.4
9	Diabetes mellitus (E10-E14)	165	0.6	0.4
10	Pregnancy, childbirth and the puerperium (O00-O99)	163	0.6	0.4
11	Chronic lower respiratory diseases (J40-J47)	149	0.5	0.3
12	Human immunodeficiency virus (HIV) disease (B20-B24)	147	0.5	0.3
13	Septicemia (A40-A41)	141	0.5	0.3
14	Anemias (D50-D64)	109	0.4	0.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	93	0.3	0.2
...	All other causes (Residual)	3,550	12.0	8.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 25-34 years				
...	All causes	42,259	100.0	102.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14,573	34.5	35.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,735	13.6	14.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4,258	10.1	10.4
4	Malignant neoplasms (C00-C97)	3,619	8.6	8.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,222	7.6	7.8
6	Human immunodeficiency virus (HIV) disease (B20-B24)	741	1.8	1.8
7	Diabetes mellitus (E10-E14)	606	1.4	1.5
8	Cerebrovascular diseases (I60-I69)	517	1.2	1.3
9	Chronic liver disease and cirrhosis (K70,K73-K74)	487	1.2	1.2
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	397	0.9	1.0
11	Influenza and pneumonia (J09-J18)	385	0.9	0.9
12	Pregnancy, childbirth and the puerperium (O00-O99)	367	0.9	0.9
13	Septicemia (A40-A41)	312	0.7	0.8
14	Chronic lower respiratory diseases (J40-J47)	272	0.6	0.7
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	243	0.6	0.6
...	All other causes (Residual)	6,525	15.4	15.9
All races, both sexes, 35-44 years				
...	All causes	70,033	100.0	170.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14,792	21.1	36.0
2	Malignant neoplasms (C00-C97)	11,809	16.9	28.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,594	15.1	25.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,571	9.4	16.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,473	3.5	6.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	2,423	3.5	5.9
7	Cerebrovascular diseases (I60-I69)	1,904	2.7	4.6
8	Human immunodeficiency virus (HIV) disease (B20-B24)	1,898	2.7	4.6
9	Diabetes mellitus (E10-E14)	1,789	2.6	4.4
10	Influenza and pneumonia (J09-J18)	773	1.1	1.9
11	Septicemia (A40-A41)	767	1.1	1.9
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	726	1.0	1.8
13	Chronic lower respiratory diseases (J40-J47)	709	1.0	1.7
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	449	0.6	1.1
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	394	0.6	1.0
...	All other causes (Residual)	11,962	17.1	29.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 45-54 years				
...	All causes	183,207	100.0	407.1
1	Malignant neoplasms (C00-C97)	50,211	27.4	111.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	36,729	20.0	81.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19,667	10.7	43.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8,799	4.8	19.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	8,651	4.7	19.2
6	Cerebrovascular diseases (I60-I69)	5,910	3.2	13.1
7	Diabetes mellitus (E10-E14)	5,610	3.1	12.5
8	Chronic lower respiratory diseases (J40-J47)	4,452	2.4	9.9
9	Human immunodeficiency virus (HIV) disease (B20-B24)	3,123	1.7	6.9
10	Viral hepatitis (B15-B19)	2,376	1.3	5.3
11	Septicemia (A40-A41)	2,333	1.3	5.2
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,222	1.2	4.9
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,997	1.1	4.4
14	Influenza and pneumonia (J09-J18)	1,926	1.1	4.3
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,384	0.8	3.1
...	All other causes (Residual)	27,817	15.2	61.8
All races, both sexes, 55-64 years				
...	All causes	310,802	100.0	851.9
1	Malignant neoplasms (C00-C97)	109,501	35.2	300.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	68,077	21.9	186.6
3	Chronic lower respiratory diseases (J40-J47)	14,242	4.6	39.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14,023	4.5	38.4
5	Diabetes mellitus (E10-E14)	11,677	3.8	32.0
6	Cerebrovascular diseases (I60-I69)	10,693	3.4	29.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	9,764	3.1	26.8
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,384	2.1	17.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,082	1.6	13.9
10	Septicemia (A40-A41)	4,604	1.5	12.6
11	Influenza and pneumonia (J09-J18)	3,627	1.2	9.9
12	Viral hepatitis (B15-B19)	3,218	1.0	8.8
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,672	0.9	7.3
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1,822	0.6	5.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,438	0.5	3.9
...	All other causes (Residual)	43,978	14.1	120.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 65-74 years				
...	All causes	407,151	100.0	1,875.1
1	Malignant neoplasms (C00-C97)	144,635	35.5	666.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	88,851	21.8	409.2
3	Chronic lower respiratory diseases (J40-J47)	31,777	7.8	146.3
4	Cerebrovascular diseases (I60-I69)	17,736	4.4	81.7
5	Diabetes mellitus (E10-E14)	14,687	3.6	67.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,407	2.3	43.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,541	2.1	39.3
8	Septicemia (A40-A41)	6,545	1.6	30.1
9	Influenza and pneumonia (J09-J18)	6,066	1.5	27.9
10	Chronic liver disease and cirrhosis (K70,K73-K74)	5,720	1.4	26.3
11	Alzheimer's disease (G30)	4,291	1.1	19.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,636	0.9	16.7
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,974	0.7	13.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,580	0.6	11.9
15	Parkinson's disease (G20-G21)	2,567	0.6	11.8
...	All other causes (Residual)	57,138	14.0	263.1
All races, both sexes, 75-84 years				
...	All causes	625,651	100.0	4,790.2
1	Malignant neoplasms (C00-C97)	157,025	25.1	1,202.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	153,080	24.5	1,172.0
3	Chronic lower respiratory diseases (J40-J47)	48,309	7.7	369.9
4	Cerebrovascular diseases (I60-I69)	37,659	6.0	288.3
5	Alzheimer's disease (G30)	24,099	3.9	184.5
6	Diabetes mellitus (E10-E14)	18,822	3.0	144.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15,118	2.4	115.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,853	2.2	106.1
9	Influenza and pneumonia (J09-J18)	13,369	2.1	102.4
10	Septicemia (A40-A41)	9,931	1.6	76.0
11	Parkinson's disease (G20-G21)	9,769	1.6	74.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	6,763	1.1	51.8
13	Pneumonitis due to solids and liquids (J69)	4,990	0.8	38.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,710	0.8	36.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	3,620	0.6	27.7
...	All other causes (Residual)	104,534	16.7	800.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 85 years and over				
...	All causes	765,474	100.0	13,934.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	235,407	30.8	4,285.2
2	Malignant neoplasms (C00-C97)	95,010	12.4	1,729.5
3	Cerebrovascular diseases (I60-I69)	54,595	7.1	993.8
4	Alzheimer's disease (G30)	54,226	7.1	987.1
5	Chronic lower respiratory diseases (J40-J47)	37,945	5.0	690.7
6	Influenza and pneumonia (J09-J18)	23,411	3.1	426.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18,335	2.4	333.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18,040	2.4	328.4
9	Diabetes mellitus (E10-E14)	15,682	2.0	285.5
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	11,648	1.5	212.0
11	Septicemia (A40-A41)	9,834	1.3	179.0
12	Parkinson's disease (G20-G21)	9,112	1.2	165.9
13	Pneumonitis due to solids and liquids (J69)	8,367	1.1	152.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,714	0.6	85.8
15	Atherosclerosis (I70)	4,162	0.5	75.8
...	All other causes (Residual)	164,986	21.6	3,003.3
All races, male, all ages				
...	All causes	1,232,432	100.0	812.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	307,384	24.9	202.5
2	Malignant neoplasms (C00-C97)	301,037	24.4	198.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	75,921	6.2	50.0
4	Chronic lower respiratory diseases (J40-J47)	65,423	5.3	43.1
5	Cerebrovascular diseases (I60-I69)	52,367	4.2	34.5
6	Diabetes mellitus (E10-E14)	35,490	2.9	23.4
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	30,277	2.5	19.9
8	Alzheimer's disease (G30)	25,364	2.1	16.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	24,865	2.0	16.4
10	Influenza and pneumonia (J09-J18)	23,615	1.9	15.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	20,798	1.7	13.7
12	Septicemia (A40-A41)	16,069	1.3	10.6
13	Parkinson's disease (G20-G21)	12,871	1.0	8.5
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	12,774	1.0	8.4
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10,846	0.9	7.1
...	All other causes (Residual)	217,331	17.6	143.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 1-4 years				
...	All causes	2,460	100.0	29.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	868	35.3	10.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	231	9.4	2.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	231	9.4	2.8
4	Malignant neoplasms (C00-C97)	197	8.0	2.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	90	3.7	1.1
6	Influenza and pneumonia (J09-J18)	50	2.0	0.6
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	32	1.3	0.4
8	Septicemia (A40-A41)	30	1.2	0.4
8	Certain conditions originating in the perinatal period (P00-P96)	30	1.2	0.4
10	Chronic lower respiratory diseases (J40-J47)	28	1.1	0.3
11	Cerebrovascular diseases (I60-I69)	26	1.1	0.3
12	Anemias (D50-D64)	17	0.7	*
13	Meningitis (G00,G03)	13	0.5	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	13	0.5	*
15	Pneumonitis due to solids and liquids (J69)	9	0.4	*
...	All other causes (Residual)	595	24.2	7.2
All races, male, 5-14 years				
...	All causes	3,054	100.0	14.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,052	34.4	5.0
2	Malignant neoplasms (C00-C97)	482	15.8	2.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	187	6.1	0.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	166	5.4	0.8
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	148	4.8	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	96	3.1	0.5
7	Chronic lower respiratory diseases (J40-J47)	90	2.9	0.4
8	Cerebrovascular diseases (I60-I69)	54	1.8	0.3
9	Septicemia (A40-A41)	38	1.2	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	35	1.1	0.2
11	Influenza and pneumonia (J09-J18)	33	1.1	0.2
12	Certain conditions originating in the perinatal period (P00-P96)	16	0.5	*
13	Diabetes mellitus (E10-E14)	14	0.5	*
14	Anemias (D50-D64)	11	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	0.4	*
...	All other causes (Residual)	621	20.3	3.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 15-24 years				
...	All causes	21,790	100.0	97.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,088	41.7	40.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4,067	18.7	18.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,771	17.3	16.9
4	Malignant neoplasms (C00-C97)	997	4.6	4.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	719	3.3	3.2
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	251	1.2	1.1
7	Cerebrovascular diseases (I60-I69)	107	0.5	0.5
8	Influenza and pneumonia (J09-J18)	105	0.5	0.5
9	Human immunodeficiency virus (HIV) disease (B20-B24)	100	0.5	0.4
10	Chronic lower respiratory diseases (J40-J47)	96	0.4	0.4
11	Diabetes mellitus (E10-E14)	90	0.4	0.4
12	Septicemia (A40-A41)	76	0.3	0.3
13	Legal intervention (Y35,Y89.0)	72	0.3	0.3
14	Anemias (D50-D64)	70	0.3	0.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	58	0.3	0.3
...	All other causes (Residual)	2,123	9.7	9.5
All races, male, 25-34 years				
...	All causes	29,192	100.0	141.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,803	37.0	52.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,643	15.9	22.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,574	12.2	17.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,212	7.6	10.7
5	Malignant neoplasms (C00-C97)	1,784	6.1	8.6
6	Human immunodeficiency virus (HIV) disease (B20-B24)	482	1.7	2.3
7	Diabetes mellitus (E10-E14)	344	1.2	1.7
8	Chronic liver disease and cirrhosis (K70,K73-K74)	307	1.1	1.5
9	Cerebrovascular diseases (I60-I69)	264	0.9	1.3
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	233	0.8	1.1
11	Influenza and pneumonia (J09-J18)	219	0.8	1.1
12	Chronic lower respiratory diseases (J40-J47)	170	0.6	0.8
13	Septicemia (A40-A41)	146	0.5	0.7
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	116	0.4	0.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	104	0.4	0.5
...	All other causes (Residual)	3,791	13.0	18.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 35-44 years				
...	All causes	43,434	100.0	212.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,196	23.5	49.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,366	17.0	36.0
3	Malignant neoplasms (C00-C97)	5,146	11.8	25.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,033	11.6	24.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,878	4.3	9.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1,612	3.7	7.9
7	Human immunodeficiency virus (HIV) disease (B20-B24)	1,294	3.0	6.3
8	Diabetes mellitus (E10-E14)	1,061	2.4	5.2
9	Cerebrovascular diseases (I60-I69)	1,029	2.4	5.0
10	Influenza and pneumonia (J09-J18)	439	1.0	2.1
11	Septicemia (A40-A41)	400	0.9	2.0
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	396	0.9	1.9
13	Chronic lower respiratory diseases (J40-J47)	334	0.8	1.6
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	251	0.6	1.2
15	Aortic aneurysm and dissection (I71)	248	0.6	1.2
...	All other causes (Residual)	6,751	15.5	33.0
All races, male, 45-54 years				
...	All causes	112,018	100.0	505.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	26,087	23.3	117.8
2	Malignant neoplasms (C00-C97)	25,188	22.5	113.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,178	11.8	59.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,733	6.0	30.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	5,896	5.3	26.6
6	Diabetes mellitus (E10-E14)	3,528	3.1	15.9
7	Cerebrovascular diseases (I60-I69)	3,299	2.9	14.9
8	Human immunodeficiency virus (HIV) disease (B20-B24)	2,342	2.1	10.6
9	Chronic lower respiratory diseases (J40-J47)	2,053	1.8	9.3
10	Viral hepatitis (B15-B19)	1,659	1.5	7.5
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,477	1.3	6.7
12	Septicemia (A40-A41)	1,254	1.1	5.7
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,246	1.1	5.6
14	Influenza and pneumonia (J09-J18)	1,087	1.0	4.9
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	848	0.8	3.8
...	All other causes (Residual)	16,143	14.4	72.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 55-64 years				
...	All causes	189,295	100.0	1,075.5
1	Malignant neoplasms (C00-C97)	60,700	32.1	344.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	47,433	25.1	269.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,609	5.1	54.6
4	Chronic lower respiratory diseases (J40-J47)	7,419	3.9	42.2
5	Diabetes mellitus (E10-E14)	7,122	3.8	40.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	6,947	3.7	39.5
7	Cerebrovascular diseases (I60-I69)	6,109	3.2	34.7
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,869	2.6	27.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,884	1.5	16.4
10	Septicemia (A40-A41)	2,445	1.3	13.9
11	Viral hepatitis (B15-B19)	2,306	1.2	13.1
12	Influenza and pneumonia (J09-J18)	2,119	1.1	12.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,659	0.9	9.4
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1,388	0.7	7.9
15	Aortic aneurysm and dissection (I71)	885	0.5	5.0
...	All other causes (Residual)	25,401	13.4	144.3
All races, male, 65-74 years				
...	All causes	229,704	100.0	2,275.1
1	Malignant neoplasms (C00-C97)	79,685	34.7	789.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	55,836	24.3	553.0
3	Chronic lower respiratory diseases (J40-J47)	16,218	7.1	160.6
4	Cerebrovascular diseases (I60-I69)	9,284	4.0	92.0
5	Diabetes mellitus (E10-E14)	8,168	3.6	80.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,938	2.6	58.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,579	2.0	45.4
8	Chronic liver disease and cirrhosis (K70,K73-K74)	3,561	1.6	35.3
9	Influenza and pneumonia (J09-J18)	3,425	1.5	33.9
10	Septicemia (A40-A41)	3,298	1.4	32.7
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,418	1.1	23.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,905	0.8	18.9
13	Alzheimer's disease (G30)	1,883	0.8	18.6
14	Parkinson's disease (G20-G21)	1,696	0.7	16.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,464	0.6	14.5
...	All other causes (Residual)	30,346	13.2	300.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 75-84 years				
...	All causes	311,830	100.0	5,693.7
1	Malignant neoplasms (C00-C97)	82,929	26.6	1,514.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	80,823	25.9	1,475.7
3	Chronic lower respiratory diseases (J40-J47)	23,285	7.5	425.2
4	Cerebrovascular diseases (I60-I69)	16,166	5.2	295.2
5	Diabetes mellitus (E10-E14)	9,267	3.0	169.2
6	Alzheimer's disease (G30)	9,065	2.9	165.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,758	2.5	141.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,359	2.4	134.4
9	Influenza and pneumonia (J09-J18)	6,820	2.2	124.5
10	Parkinson's disease (G20-G21)	6,118	2.0	111.7
11	Septicemia (A40-A41)	4,664	1.5	85.2
12	Pneumonitis due to solids and liquids (J69)	2,891	0.9	52.8
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,807	0.9	51.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,541	0.8	46.4
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,933	0.6	35.3
...	All other causes (Residual)	47,404	15.2	865.5
All races, male, 85 years and over				
...	All causes	275,866	100.0	15,414.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	86,506	31.4	4,833.6
2	Malignant neoplasms (C00-C97)	43,893	15.9	2,452.6
3	Cerebrovascular diseases (I60-I69)	15,964	5.8	892.0
4	Chronic lower respiratory diseases (J40-J47)	15,702	5.7	877.4
5	Alzheimer's disease (G30)	14,047	5.1	784.9
6	Influenza and pneumonia (J09-J18)	9,205	3.3	514.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,783	2.8	434.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,182	2.6	401.3
9	Diabetes mellitus (E10-E14)	5,892	2.1	329.2
10	Parkinson's disease (G20-G21)	4,666	1.7	260.7
11	Pneumonitis due to solids and liquids (J69)	4,081	1.5	228.0
12	Septicemia (A40-A41)	3,598	1.3	201.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,305	1.2	184.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,077	0.8	116.1
15	Atherosclerosis (I70)	1,329	0.5	74.3
...	All other causes (Residual)	50,636	18.4	2,829.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, all ages				
...	All causes	1,236,003	100.0	787.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	290,305	23.5	184.9
2	Malignant neoplasms (C00-C97)	273,706	22.1	174.4
3	Cerebrovascular diseases (I60-I69)	77,109	6.2	49.1
4	Chronic lower respiratory diseases (J40-J47)	72,657	5.9	46.3
5	Alzheimer's disease (G30)	58,130	4.7	37.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	44,938	3.6	28.6
7	Diabetes mellitus (E10-E14)	33,581	2.7	21.4
8	Influenza and pneumonia (J09-J18)	26,482	2.1	16.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	25,611	2.1	16.3
10	Septicemia (A40-A41)	18,743	1.5	11.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	15,788	1.3	10.1
12	Chronic liver disease and cirrhosis (K70,K73-K74)	11,105	0.9	7.1
13	Parkinson's disease (G20-G21)	9,161	0.7	5.8
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8,087	0.7	5.2
15	Pneumonitis due to solids and liquids (J69)	7,803	0.6	5.0
...	All other causes (Residual)	262,797	21.3	167.4
All races, female, 1-4 years				
...	All causes	1,856	100.0	23.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	526	28.3	6.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	276	14.9	3.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	154	8.3	1.9
4	Malignant neoplasms (C00-C97)	149	8.0	1.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	69	3.7	0.9
6	Influenza and pneumonia (J09-J18)	41	2.2	0.5
7	Septicemia (A40-A41)	32	1.7	0.4
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	27	1.5	0.3
9	Cerebrovascular diseases (I60-I69)	24	1.3	0.3
10	Chronic lower respiratory diseases (J40-J47)	23	1.2	0.3
11	Certain conditions originating in the perinatal period (P00-P96)	22	1.2	0.3
12	Anemias (D50-D64)	12	0.6	*
13	Meningitis (G00,G03)	9	0.5	*
14	Meningococcal infection (A39)	7	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.4	*
...	All other causes (Residual)	478	25.8	6.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 5-14 years				
...	All causes	2,225	100.0	11.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	591	26.6	2.9
2	Malignant neoplasms (C00-C97)	434	19.5	2.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	150	6.7	0.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	95	4.3	0.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	89	4.0	0.4
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	87	3.9	0.4
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	47	2.1	0.2
8	Chronic lower respiratory diseases (J40-J47)	43	1.9	0.2
9	Influenza and pneumonia (J09-J18)	38	1.7	0.2
10	Cerebrovascular diseases (I60-I69)	36	1.6	0.2
11	Septicemia (A40-A41)	29	1.3	0.1
12	Anemias (D50-D64)	15	0.7	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	14	0.6	*
14	Diabetes mellitus (E10-E14)	12	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	0.5	*
...	All other causes (Residual)	534	24.0	2.7
All races, female, 15-24 years				
...	All causes	7,761	100.0	36.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,253	41.9	15.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	829	10.7	3.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	611	7.9	2.9
4	Malignant neoplasms (C00-C97)	607	7.8	2.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	309	4.0	1.5
6	Pregnancy, childbirth and the puerperium (O00-O99)	163	2.1	0.8
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	161	2.1	0.8
8	Cerebrovascular diseases (I60-I69)	83	1.1	0.4
9	Influenza and pneumonia (J09-J18)	76	1.0	0.4
10	Diabetes mellitus (E10-E14)	75	1.0	0.4
11	Septicemia (A40-A41)	65	0.8	0.3
12	Chronic lower respiratory diseases (J40-J47)	53	0.7	0.2
13	Human immunodeficiency virus (HIV) disease (B20-B24)	47	0.6	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	42	0.5	0.2
15	Anemias (D50-D64)	39	0.5	0.2
...	All other causes (Residual)	1,348	17.4	6.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 25-34 years				
...	All causes	13,067	100.0	64.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,770	28.9	18.5
2	Malignant neoplasms (C00-C97)	1,835	14.0	9.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,092	8.4	5.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,010	7.7	4.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	684	5.2	3.3
6	Pregnancy, childbirth and the puerperium (O00-O99)	367	2.8	1.8
7	Diabetes mellitus (E10-E14)	262	2.0	1.3
8	Human immunodeficiency virus (HIV) disease (B20-B24)	259	2.0	1.3
9	Cerebrovascular diseases (I60-I69)	253	1.9	1.2
10	Chronic liver disease and cirrhosis (K70,K73-K74)	180	1.4	0.9
11	Septicemia (A40-A41)	166	1.3	0.8
11	Influenza and pneumonia (J09-J18)	166	1.3	0.8
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	164	1.3	0.8
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	127	1.0	0.6
15	Chronic lower respiratory diseases (J40-J47)	102	0.8	0.5
...	All other causes (Residual)	2,630	20.1	12.9
All races, female, 35-44 years				
...	All causes	26,599	100.0	128.9
1	Malignant neoplasms (C00-C97)	6,663	25.0	32.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,596	17.3	22.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,228	12.1	15.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,538	5.8	7.5
5	Cerebrovascular diseases (I60-I69)	875	3.3	4.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	811	3.0	3.9
7	Diabetes mellitus (E10-E14)	728	2.7	3.5
8	Human immunodeficiency virus (HIV) disease (B20-B24)	604	2.3	2.9
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	595	2.2	2.9
10	Chronic lower respiratory diseases (J40-J47)	375	1.4	1.8
11	Septicemia (A40-A41)	367	1.4	1.8
12	Influenza and pneumonia (J09-J18)	334	1.3	1.6
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	330	1.2	1.6
14	Pregnancy, childbirth and the puerperium (O00-O99)	223	0.8	1.1
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	198	0.7	1.0
...	All other causes (Residual)	5,134	19.3	24.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 45-54 years				
...	All causes	71,189	100.0	311.4
1	Malignant neoplasms (C00-C97)	25,023	35.2	109.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,642	14.9	46.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,489	9.1	28.4
4	Chronic liver disease and cirrhosis (K70,K73-K74)	2,755	3.9	12.0
5	Cerebrovascular diseases (I60-I69)	2,611	3.7	11.4
6	Chronic lower respiratory diseases (J40-J47)	2,399	3.4	10.5
7	Diabetes mellitus (E10-E14)	2,082	2.9	9.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,066	2.9	9.0
9	Septicemia (A40-A41)	1,079	1.5	4.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	976	1.4	4.3
11	Influenza and pneumonia (J09-J18)	839	1.2	3.7
12	Human immunodeficiency virus (HIV) disease (B20-B24)	781	1.1	3.4
13	Viral hepatitis (B15-B19)	717	1.0	3.1
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	536	0.8	2.3
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	520	0.7	2.3
...	All other causes (Residual)	11,674	16.4	51.1
All races, female, 55-64 years				
...	All causes	121,507	100.0	643.5
1	Malignant neoplasms (C00-C97)	48,801	40.2	258.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,644	17.0	109.3
3	Chronic lower respiratory diseases (J40-J47)	6,823	5.6	36.1
4	Cerebrovascular diseases (I60-I69)	4,584	3.8	24.3
5	Diabetes mellitus (E10-E14)	4,555	3.7	24.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,414	3.6	23.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,817	2.3	14.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,198	1.8	11.6
9	Septicemia (A40-A41)	2,159	1.8	11.4
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,515	1.2	8.0
11	Influenza and pneumonia (J09-J18)	1,508	1.2	8.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,013	0.8	5.4
13	Viral hepatitis (B15-B19)	912	0.8	4.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	598	0.5	3.2
15	Human immunodeficiency virus (HIV) disease (B20-B24)	434	0.4	2.3
...	All other causes (Residual)	18,532	15.3	98.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 65-74 years				
...	All causes	177,447	100.0	1,527.5
1	Malignant neoplasms (C00-C97)	64,950	36.6	559.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	33,015	18.6	284.2
3	Chronic lower respiratory diseases (J40-J47)	15,559	8.8	133.9
4	Cerebrovascular diseases (I60-I69)	8,452	4.8	72.8
5	Diabetes mellitus (E10-E14)	6,519	3.7	56.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,962	2.2	34.1
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,469	2.0	29.9
8	Septicemia (A40-A41)	3,247	1.8	28.0
9	Influenza and pneumonia (J09-J18)	2,641	1.5	22.7
10	Alzheimer's disease (G30)	2,408	1.4	20.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	2,159	1.2	18.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,731	1.0	14.9
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,116	0.6	9.6
14	Parkinson's disease (G20-G21)	871	0.5	7.5
15	Pneumonitis due to solids and liquids (J69)	752	0.4	6.5
...	All other causes (Residual)	26,596	15.0	228.9
All races, female, 75-84 years				
...	All causes	313,821	100.0	4,137.7
1	Malignant neoplasms (C00-C97)	74,096	23.6	977.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	72,257	23.0	952.7
3	Chronic lower respiratory diseases (J40-J47)	25,024	8.0	329.9
4	Cerebrovascular diseases (I60-I69)	21,493	6.8	283.4
5	Alzheimer's disease (G30)	15,034	4.8	198.2
6	Diabetes mellitus (E10-E14)	9,555	3.0	126.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,360	2.3	97.0
8	Influenza and pneumonia (J09-J18)	6,549	2.1	86.3
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,494	2.1	85.6
10	Septicemia (A40-A41)	5,267	1.7	69.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,956	1.3	52.2
12	Parkinson's disease (G20-G21)	3,651	1.2	48.1
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,169	0.7	28.6
14	Pneumonitis due to solids and liquids (J69)	2,099	0.7	27.7
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,687	0.5	22.2
...	All other causes (Residual)	57,130	18.2	753.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 85 years and over				
...	All causes	489,608	100.0	13,219.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	148,901	30.4	4,020.3
2	Malignant neoplasms (C00-C97)	51,117	10.4	1,380.1
3	Alzheimer's disease (G30)	40,179	8.2	1,084.8
4	Cerebrovascular diseases (I60-I69)	38,631	7.9	1,043.0
5	Chronic lower respiratory diseases (J40-J47)	22,243	4.5	600.6
6	Influenza and pneumonia (J09-J18)	14,206	2.9	383.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,858	2.2	293.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10,552	2.2	284.9
9	Diabetes mellitus (E10-E14)	9,790	2.0	264.3
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	8,343	1.7	225.3
11	Septicemia (A40-A41)	6,236	1.3	168.4
12	Parkinson's disease (G20-G21)	4,446	0.9	120.0
13	Pneumonitis due to solids and liquids (J69)	4,286	0.9	115.7
14	Atherosclerosis (I70)	2,833	0.6	76.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,637	0.5	71.2
...	All other causes (Residual)	114,350	23.4	3,087.4
White, both sexes, all ages				
...	All causes	2,114,749	100.0	861.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	514,323	24.3	209.6
2	Malignant neoplasms (C00-C97)	491,686	23.3	200.3
3	Chronic lower respiratory diseases (J40-J47)	127,176	6.0	51.8
4	Cerebrovascular diseases (I60-I69)	109,119	5.2	44.5
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	104,945	5.0	42.8
6	Alzheimer's disease (G30)	76,928	3.6	31.3
7	Diabetes mellitus (E10-E14)	54,250	2.6	22.1
8	Influenza and pneumonia (J09-J18)	43,296	2.0	17.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	40,205	1.9	16.4
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	34,690	1.6	14.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	28,014	1.3	11.4
12	Septicemia (A40-A41)	27,985	1.3	11.4
13	Parkinson's disease (G20-G21)	20,792	1.0	8.5
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	20,560	1.0	8.4
15	Pneumonitis due to solids and liquids (J69)	15,235	0.7	6.2
...	All other causes (Residual)	405,545	19.2	165.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 1-4 years				
...	All causes	3,015	100.0	24.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,054	35.0	8.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	348	11.5	2.8
3	Malignant neoplasms (C00-C97)	269	8.9	2.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	214	7.1	1.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	95	3.2	0.8
6	Influenza and pneumonia (J09-J18)	61	2.0	0.5
7	Septicemia (A40-A41)	47	1.6	0.4
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	43	1.4	0.4
9	Certain conditions originating in the perinatal period (P00-P96)	34	1.1	0.3
10	Cerebrovascular diseases (I60-I69)	30	1.0	0.2
11	Chronic lower respiratory diseases (J40-J47)	23	0.8	0.2
12	Meningitis (G00,G03)	16	0.5	*
13	Anemias (D50-D64)	14	0.5	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	13	0.4	*
15	Acute bronchitis and bronchiolitis (J20-J21)	10	0.3	*
...	All other causes (Residual)	744	24.7	6.1
White, both sexes, 5-14 years				
...	All causes	3,841	100.0	12.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,248	32.5	4.0
2	Malignant neoplasms (C00-C97)	707	18.4	2.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	214	5.6	0.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	209	5.4	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	148	3.9	0.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	116	3.0	0.4
7	Cerebrovascular diseases (I60-I69)	61	1.6	0.2
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	60	1.6	0.2
9	Influenza and pneumonia (J09-J18)	54	1.4	0.2
10	Chronic lower respiratory diseases (J40-J47)	53	1.4	0.2
11	Septicemia (A40-A41)	44	1.1	0.1
12	Diabetes mellitus (E10-E14)	19	0.5	*
13	Certain conditions originating in the perinatal period (P00-P96)	17	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	14	0.4	*
...	All other causes (Residual)	862	22.4	2.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 15-24 years				
...	All causes	21,509	100.0	64.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,358	48.2	31.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,792	17.6	11.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,724	8.0	5.2
4	Malignant neoplasms (C00-C97)	1,243	5.8	3.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	639	3.0	1.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	304	1.4	0.9
7	Influenza and pneumonia (J09-J18)	134	0.6	0.4
8	Cerebrovascular diseases (I60-I69)	129	0.6	0.4
9	Diabetes mellitus (E10-E14)	113	0.5	0.3
10	Pregnancy, childbirth and the puerperium (O00-O99)	106	0.5	0.3
11	Septicemia (A40-A41)	95	0.4	0.3
12	Chronic lower respiratory diseases (J40-J47)	82	0.4	0.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	68	0.3	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	44	0.2	0.1
15	Legal intervention (Y35,Y89.0)	41	0.2	0.1
...	All other causes (Residual)	2,637	12.3	7.9
White, both sexes, 25-34 years				
...	All causes	31,425	100.0	99.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,382	39.4	39.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,917	15.6	15.5
3	Malignant neoplasms (C00-C97)	2,790	8.9	8.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,173	6.9	6.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,720	5.5	5.4
6	Chronic liver disease and cirrhosis (K70,K73-K74)	391	1.2	1.2
7	Diabetes mellitus (E10-E14)	385	1.2	1.2
8	Cerebrovascular diseases (I60-I69)	349	1.1	1.1
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	300	1.0	0.9
10	Influenza and pneumonia (J09-J18)	282	0.9	0.9
11	Human immunodeficiency virus (HIV) disease (B20-B24)	256	0.8	0.8
12	Pregnancy, childbirth and the puerperium (O00-O99)	220	0.7	0.7
13	Septicemia (A40-A41)	210	0.7	0.7
14	Chronic lower respiratory diseases (J40-J47)	160	0.5	0.5
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	143	0.5	0.5
...	All other causes (Residual)	4,747	15.1	15.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 35-44 years				
...	All causes	53,060	100.0	165.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,701	23.9	39.5
2	Malignant neoplasms (C00-C97)	9,068	17.1	28.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,471	14.1	23.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,855	11.0	18.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,028	3.8	6.3
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,276	2.4	4.0
7	Cerebrovascular diseases (I60-I69)	1,217	2.3	3.8
8	Diabetes mellitus (E10-E14)	1,196	2.3	3.7
9	Human immunodeficiency virus (HIV) disease (B20-B24)	795	1.5	2.5
10	Influenza and pneumonia (J09-J18)	589	1.1	1.8
11	Septicemia (A40-A41)	526	1.0	1.6
12	Chronic lower respiratory diseases (J40-J47)	509	1.0	1.6
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	414	0.8	1.3
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	371	0.7	1.2
15	Viral hepatitis (B15-B19)	286	0.5	0.9
...	All other causes (Residual)	8,758	16.5	27.2
White, both sexes, 45-54 years				
...	All causes	143,049	100.0	392.9
1	Malignant neoplasms (C00-C97)	39,703	27.8	109.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	27,589	19.3	75.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	16,813	11.8	46.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8,198	5.7	22.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	7,473	5.2	20.5
6	Diabetes mellitus (E10-E14)	3,975	2.8	10.9
7	Cerebrovascular diseases (I60-I69)	3,869	2.7	10.6
8	Chronic lower respiratory diseases (J40-J47)	3,731	2.6	10.2
9	Viral hepatitis (B15-B19)	1,977	1.4	5.4
10	Septicemia (A40-A41)	1,658	1.2	4.6
11	Influenza and pneumonia (J09-J18)	1,468	1.0	4.0
12	Human immunodeficiency virus (HIV) disease (B20-B24)	1,430	1.0	3.9
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,334	0.9	3.7
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,230	0.9	3.4
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	806	0.6	2.2
...	All other causes (Residual)	21,795	15.2	59.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 55-64 years				
...	All causes	249,583	100.0	820.3
1	Malignant neoplasms (C00-C97)	89,487	35.9	294.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	53,448	21.4	175.7
3	Chronic lower respiratory diseases (J40-J47)	12,573	5.0	41.3
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,834	4.7	38.9
5	Diabetes mellitus (E10-E14)	8,706	3.5	28.6
6	Chronic liver disease and cirrhosis (K70,K73-K74)	8,440	3.4	27.7
7	Cerebrovascular diseases (I60-I69)	7,520	3.0	24.7
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,001	2.4	19.7
9	Septicemia (A40-A41)	3,384	1.4	11.1
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,375	1.4	11.1
11	Influenza and pneumonia (J09-J18)	2,899	1.2	9.5
12	Viral hepatitis (B15-B19)	2,582	1.0	8.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,700	0.7	5.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,204	0.5	4.0
15	Aortic aneurysm and dissection (I71)	995	0.4	3.3
...	All other causes (Residual)	35,435	14.2	116.5
White, both sexes, 65-74 years				
...	All causes	342,977	100.0	1,846.7
1	Malignant neoplasms (C00-C97)	123,807	36.1	666.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	73,227	21.4	394.3
3	Chronic lower respiratory diseases (J40-J47)	29,164	8.5	157.0
4	Cerebrovascular diseases (I60-I69)	13,842	4.0	74.5
5	Diabetes mellitus (E10-E14)	11,306	3.3	60.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,143	2.4	43.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,298	1.8	33.9
8	Chronic liver disease and cirrhosis (K70,K73-K74)	5,169	1.5	27.8
9	Septicemia (A40-A41)	5,124	1.5	27.6
10	Influenza and pneumonia (J09-J18)	4,965	1.4	26.7
11	Alzheimer's disease (G30)	3,799	1.1	20.5
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,823	0.8	15.2
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,549	0.7	13.7
14	Parkinson's disease (G20-G21)	2,349	0.7	12.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,267	0.7	12.2
...	All other causes (Residual)	48,145	14.0	259.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 75-84 years				
...	All causes	552,508	100.0	4,818.2
1	Malignant neoplasms (C00-C97)	138,816	25.1	1,210.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	133,933	24.2	1,168.0
3	Chronic lower respiratory diseases (J40-J47)	45,256	8.2	394.7
4	Cerebrovascular diseases (I60-I69)	32,524	5.9	283.6
5	Alzheimer's disease (G30)	22,018	4.0	192.0
6	Diabetes mellitus (E10-E14)	15,278	2.8	133.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,650	2.3	110.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12,465	2.3	108.7
9	Influenza and pneumonia (J09-J18)	11,619	2.1	101.3
10	Parkinson's disease (G20-G21)	9,258	1.7	80.7
11	Septicemia (A40-A41)	8,301	1.5	72.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,352	1.0	46.7
13	Pneumonitis due to solids and liquids (J69)	4,469	0.8	39.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,302	0.8	37.5
15	Chronic liver disease and cirrhosis (K70,K73-K74)	3,345	0.6	29.2
...	All other causes (Residual)	92,922	16.8	810.3
White, both sexes, 85 years and over				
...	All causes	697,733	100.0	14,147.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	215,399	30.9	4,367.6
2	Malignant neoplasms (C00-C97)	85,745	12.3	1,738.6
3	Alzheimer's disease (G30)	50,307	7.2	1,020.1
4	Cerebrovascular diseases (I60-I69)	49,502	7.1	1,003.7
5	Chronic lower respiratory diseases (J40-J47)	35,600	5.1	721.8
6	Influenza and pneumonia (J09-J18)	21,122	3.0	428.3
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17,046	2.4	345.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16,049	2.3	325.4
9	Diabetes mellitus (E10-E14)	13,269	1.9	269.0
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9,885	1.4	200.4
11	Parkinson's disease (G20-G21)	8,667	1.2	175.7
12	Septicemia (A40-A41)	8,481	1.2	172.0
13	Pneumonitis due to solids and liquids (J69)	7,734	1.1	156.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,360	0.6	88.4
15	Atherosclerosis (I70)	3,863	0.6	78.3
...	All other causes (Residual)	150,704	21.6	3,055.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, all ages				
...	All causes	1,051,514	100.0	866.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	264,425	25.1	217.8
2	Malignant neoplasms (C00-C97)	258,272	24.6	212.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	65,360	6.2	53.8
4	Chronic lower respiratory diseases (J40-J47)	59,632	5.7	49.1
5	Cerebrovascular diseases (I60-I69)	43,424	4.1	35.8
6	Diabetes mellitus (E10-E14)	28,486	2.7	23.5
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	27,422	2.6	22.6
8	Alzheimer's disease (G30)	23,442	2.2	19.3
9	Influenza and pneumonia (J09-J18)	20,238	1.9	16.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20,172	1.9	16.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	18,352	1.7	15.1
12	Septicemia (A40-A41)	12,976	1.2	10.7
13	Parkinson's disease (G20-G21)	12,173	1.2	10.0
14	Pneumonitis due to solids and liquids (J69)	8,264	0.8	6.8
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	8,229	0.8	6.8
...	All other causes (Residual)	180,647	17.2	148.8
White, male, 1-4 years				
...	All causes	1,718	100.0	27.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	653	38.0	10.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	160	9.3	2.5
3	Malignant neoplasms (C00-C97)	149	8.7	2.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	130	7.6	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	53	3.1	0.8
6	Influenza and pneumonia (J09-J18)	33	1.9	0.5
7	Septicemia (A40-A41)	23	1.3	0.4
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	1.3	0.4
9	Certain conditions originating in the perinatal period (P00-P96)	19	1.1	*
10	Cerebrovascular diseases (I60-I69)	18	1.0	*
11	Chronic lower respiratory diseases (J40-J47)	14	0.8	*
12	Meningitis (G00,G03)	11	0.6	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	9	0.5	*
14	Anemias (D50-D64)	8	0.5	*
15	Acute bronchitis and bronchiolitis (J20-J21)	7	0.4	*
...	All other causes (Residual)	408	23.7	6.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 5-14 years				
...	All causes	2,222	100.0	13.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	785	35.3	4.9
2	Malignant neoplasms (C00-C97)	386	17.4	2.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	151	6.8	0.9
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	104	4.7	0.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	90	4.1	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	54	2.4	0.3
7	Cerebrovascular diseases (I60-I69)	36	1.6	0.2
8	Chronic lower respiratory diseases (J40-J47)	31	1.4	0.2
9	Septicemia (A40-A41)	26	1.2	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	25	1.1	0.2
11	Influenza and pneumonia (J09-J18)	23	1.0	0.1
12	Certain conditions originating in the perinatal period (P00-P96)	13	0.6	*
13	Diabetes mellitus (E10-E14)	11	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.3	*
...	All other causes (Residual)	473	21.3	2.9
White, male, 15-24 years				
...	All causes	15,661	100.0	91.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,620	48.7	44.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,116	19.9	18.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,399	8.9	8.2
4	Malignant neoplasms (C00-C97)	788	5.0	4.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	471	3.0	2.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	184	1.2	1.1
7	Cerebrovascular diseases (I60-I69)	80	0.5	0.5
8	Influenza and pneumonia (J09-J18)	76	0.5	0.4
9	Diabetes mellitus (E10-E14)	63	0.4	0.4
10	Chronic lower respiratory diseases (J40-J47)	49	0.3	0.3
11	Septicemia (A40-A41)	46	0.3	0.3
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	41	0.3	0.2
13	Legal intervention (Y35,Y89.0)	40	0.3	0.2
14	Human immunodeficiency virus (HIV) disease (B20-B24)	27	0.2	0.2
14	Aortic aneurysm and dissection (I71)	27	0.2	0.2
...	All other causes (Residual)	1,634	10.4	9.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 25-34 years				
...	All causes	21,883	100.0	135.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,219	42.1	57.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,976	18.2	24.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,540	7.0	9.5
4	Malignant neoplasms (C00-C97)	1,414	6.5	8.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,335	6.1	8.3
6	Chronic liver disease and cirrhosis (K70,K73-K74)	264	1.2	1.6
7	Diabetes mellitus (E10-E14)	225	1.0	1.4
8	Human immunodeficiency virus (HIV) disease (B20-B24)	194	0.9	1.2
9	Cerebrovascular diseases (I60-I69)	173	0.8	1.1
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	173	0.8	1.1
11	Influenza and pneumonia (J09-J18)	158	0.7	1.0
12	Septicemia (A40-A41)	106	0.5	0.7
13	Chronic lower respiratory diseases (J40-J47)	92	0.4	0.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	85	0.4	0.5
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	71	0.3	0.4
...	All other causes (Residual)	2,858	13.1	17.7
White, male, 35-44 years				
...	All causes	33,486	100.0	206.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,741	26.1	53.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,375	16.1	33.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,500	13.4	27.8
4	Malignant neoplasms (C00-C97)	4,077	12.2	25.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,374	4.1	8.5
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	893	2.7	5.5
7	Diabetes mellitus (E10-E14)	714	2.1	4.4
8	Cerebrovascular diseases (I60-I69)	665	2.0	4.1
9	Human immunodeficiency virus (HIV) disease (B20-B24)	621	1.9	3.8
10	Influenza and pneumonia (J09-J18)	326	1.0	2.0
11	Septicemia (A40-A41)	278	0.8	1.7
12	Chronic lower respiratory diseases (J40-J47)	233	0.7	1.4
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	224	0.7	1.4
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	210	0.6	1.3
15	Viral hepatitis (B15-B19)	195	0.6	1.2
...	All other causes (Residual)	5,060	15.1	31.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 45-54 years				
...	All causes	89,017	100.0	491.9
1	Malignant neoplasms (C00-C97)	20,194	22.7	111.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,131	22.6	111.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,246	12.6	62.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,286	7.1	34.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	5,165	5.8	28.5
6	Diabetes mellitus (E10-E14)	2,579	2.9	14.3
7	Cerebrovascular diseases (I60-I69)	2,199	2.5	12.2
8	Chronic lower respiratory diseases (J40-J47)	1,731	1.9	9.6
9	Viral hepatitis (B15-B19)	1,386	1.6	7.7
10	Human immunodeficiency virus (HIV) disease (B20-B24)	1,207	1.4	6.7
11	Septicemia (A40-A41)	910	1.0	5.0
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	871	1.0	4.8
13	Influenza and pneumonia (J09-J18)	807	0.9	4.5
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	800	0.9	4.4
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	533	0.6	2.9
...	All other causes (Residual)	12,972	14.6	71.7
White, male, 55-64 years				
...	All causes	153,296	100.0	1,033.0
1	Malignant neoplasms (C00-C97)	49,696	32.4	334.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	38,139	24.9	257.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,073	5.3	54.4
4	Chronic lower respiratory diseases (J40-J47)	6,501	4.2	43.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	6,023	3.9	40.6
6	Diabetes mellitus (E10-E14)	5,461	3.6	36.8
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,572	3.0	30.8
8	Cerebrovascular diseases (I60-I69)	4,307	2.8	29.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,945	1.3	13.1
10	Viral hepatitis (B15-B19)	1,864	1.2	12.6
11	Septicemia (A40-A41)	1,807	1.2	12.2
12	Influenza and pneumonia (J09-J18)	1,696	1.1	11.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,108	0.7	7.5
14	Aortic aneurysm and dissection (I71)	749	0.5	5.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	714	0.5	4.8
...	All other causes (Residual)	20,641	13.5	139.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 65-74 years				
...	All causes	194,793	100.0	2,232.4
1	Malignant neoplasms (C00-C97)	68,301	35.1	782.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	46,798	24.0	536.3
3	Chronic lower respiratory diseases (J40-J47)	14,741	7.6	168.9
4	Cerebrovascular diseases (I60-I69)	7,268	3.7	83.3
5	Diabetes mellitus (E10-E14)	6,509	3.3	74.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,136	2.6	58.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,462	1.8	39.7
8	Chronic liver disease and cirrhosis (K70,K73-K74)	3,213	1.6	36.8
9	Influenza and pneumonia (J09-J18)	2,792	1.4	32.0
10	Septicemia (A40-A41)	2,630	1.4	30.1
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,293	1.2	26.3
12	Alzheimer's disease (G30)	1,672	0.9	19.2
13	Parkinson's disease (G20-G21)	1,565	0.8	17.9
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,370	0.7	15.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,320	0.7	15.1
...	All other causes (Residual)	25,723	13.2	294.8
White, male, 75-84 years				
...	All causes	277,543	100.0	5,703.6
1	Malignant neoplasms (C00-C97)	73,556	26.5	1,511.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	71,778	25.9	1,475.1
3	Chronic lower respiratory diseases (J40-J47)	21,631	7.8	444.5
4	Cerebrovascular diseases (I60-I69)	14,030	5.1	288.3
5	Alzheimer's disease (G30)	8,317	3.0	170.9
6	Diabetes mellitus (E10-E14)	7,777	2.8	159.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,713	2.4	138.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,603	2.4	135.7
9	Influenza and pneumonia (J09-J18)	5,963	2.1	122.5
10	Parkinson's disease (G20-G21)	5,813	2.1	119.5
11	Septicemia (A40-A41)	3,931	1.4	80.8
12	Pneumonitis due to solids and liquids (J69)	2,594	0.9	53.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,351	0.8	48.3
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,234	0.8	45.9
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,795	0.6	36.9
...	All other causes (Residual)	42,457	15.3	872.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 85 years and over				
...	All causes	252,958	100.0	15,640.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	79,948	31.6	4,943.1
2	Malignant neoplasms (C00-C97)	39,682	15.7	2,453.5
3	Cerebrovascular diseases (I60-I69)	14,608	5.8	903.2
4	Chronic lower respiratory diseases (J40-J47)	14,595	5.8	902.4
5	Alzheimer's disease (G30)	13,116	5.2	811.0
6	Influenza and pneumonia (J09-J18)	8,306	3.3	513.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,004	2.8	433.1
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,763	2.7	418.2
9	Diabetes mellitus (E10-E14)	5,146	2.0	318.2
10	Parkinson's disease (G20-G21)	4,451	1.8	275.2
11	Pneumonitis due to solids and liquids (J69)	3,821	1.5	236.3
12	Septicemia (A40-A41)	3,149	1.2	194.7
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,810	1.1	173.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,940	0.8	119.9
15	Atherosclerosis (I70)	1,240	0.5	76.7
...	All other causes (Residual)	46,379	18.3	2,867.6
White, female, all ages				
...	All causes	1,063,235	100.0	857.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	249,898	23.5	201.5
2	Malignant neoplasms (C00-C97)	233,414	22.0	188.2
3	Chronic lower respiratory diseases (J40-J47)	67,544	6.4	54.5
4	Cerebrovascular diseases (I60-I69)	65,695	6.2	53.0
5	Alzheimer's disease (G30)	53,486	5.0	43.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	39,585	3.7	31.9
7	Diabetes mellitus (E10-E14)	25,764	2.4	20.8
8	Influenza and pneumonia (J09-J18)	23,058	2.2	18.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20,033	1.9	16.2
10	Septicemia (A40-A41)	15,009	1.4	12.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	12,331	1.2	9.9
12	Chronic liver disease and cirrhosis (K70,K73-K74)	9,662	0.9	7.8
13	Parkinson's disease (G20-G21)	8,619	0.8	6.9
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7,268	0.7	5.9
15	Pneumonitis due to solids and liquids (J69)	6,971	0.7	5.6
...	All other causes (Residual)	224,898	21.2	181.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 1-4 years				
...	All causes	1,297	100.0	21.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	401	30.9	6.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	188	14.5	3.1
3	Malignant neoplasms (C00-C97)	120	9.3	2.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	84	6.5	1.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	42	3.2	0.7
6	Influenza and pneumonia (J09-J18)	28	2.2	0.5
7	Septicemia (A40-A41)	24	1.9	0.4
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	1.5	0.3
9	Certain conditions originating in the perinatal period (P00-P96)	15	1.2	*
10	Cerebrovascular diseases (I60-I69)	12	0.9	*
11	Chronic lower respiratory diseases (J40-J47)	9	0.7	*
12	Meningococcal infection (A39)	7	0.5	*
13	Anemias (D50-D64)	6	0.5	*
14	Meningitis (G00,G03)	5	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.3	*
...	All other causes (Residual)	328	25.3	5.5
White, female, 5-14 years				
...	All causes	1,619	100.0	10.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	463	28.6	3.0
2	Malignant neoplasms (C00-C97)	321	19.8	2.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	110	6.8	0.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	62	3.8	0.4
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	58	3.6	0.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	58	3.6	0.4
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	35	2.2	0.2
8	Influenza and pneumonia (J09-J18)	31	1.9	0.2
9	Cerebrovascular diseases (I60-I69)	25	1.5	0.2
10	Chronic lower respiratory diseases (J40-J47)	22	1.4	0.1
11	Septicemia (A40-A41)	18	1.1	*
12	Diabetes mellitus (E10-E14)	8	0.5	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.5	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.4	*
15	Anemias (D50-D64)	6	0.4	*
15	Meningitis (G00,G03)	6	0.4	*
15	Pneumonitis due to solids and liquids (J69)	6	0.4	*
...	All other causes (Residual)	375	23.2	2.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 15-24 years				
...	All causes	5,848	100.0	36.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,738	46.8	17.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	676	11.6	4.2
3	Malignant neoplasms (C00-C97)	455	7.8	2.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	325	5.6	2.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	168	2.9	1.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	120	2.1	0.7
7	Pregnancy, childbirth and the puerperium (O00-O99)	106	1.8	0.7
8	Influenza and pneumonia (J09-J18)	58	1.0	0.4
9	Diabetes mellitus (E10-E14)	50	0.9	0.3
10	Septicemia (A40-A41)	49	0.8	0.3
10	Cerebrovascular diseases (I60-I69)	49	0.8	0.3
12	Chronic lower respiratory diseases (J40-J47)	33	0.6	0.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	27	0.5	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	25	0.4	0.2
15	Complications of medical and surgical care (Y40-Y84,Y88)	14	0.2	*
...	All other causes (Residual)	955	16.3	5.9
White, female, 25-34 years				
...	All causes	9,542	100.0	61.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,163	33.1	20.3
2	Malignant neoplasms (C00-C97)	1,376	14.4	8.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	941	9.9	6.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	633	6.6	4.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	385	4.0	2.5
6	Pregnancy, childbirth and the puerperium (O00-O99)	220	2.3	1.4
7	Cerebrovascular diseases (I60-I69)	176	1.8	1.1
8	Diabetes mellitus (E10-E14)	160	1.7	1.0
9	Chronic liver disease and cirrhosis (K70,K73-K74)	127	1.3	0.8
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	127	1.3	0.8
11	Influenza and pneumonia (J09-J18)	124	1.3	0.8
12	Septicemia (A40-A41)	104	1.1	0.7
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	72	0.8	0.5
14	Chronic lower respiratory diseases (J40-J47)	68	0.7	0.4
15	Human immunodeficiency virus (HIV) disease (B20-B24)	62	0.6	0.4
...	All other causes (Residual)	1,804	18.9	11.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 35-44 years				
...	All causes	19,574	100.0	122.8
1	Malignant neoplasms (C00-C97)	4,991	25.5	31.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,960	20.2	24.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,096	10.7	13.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,355	6.9	8.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	654	3.3	4.1
6	Cerebrovascular diseases (I60-I69)	552	2.8	3.5
7	Diabetes mellitus (E10-E14)	482	2.5	3.0
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	383	2.0	2.4
9	Chronic lower respiratory diseases (J40-J47)	276	1.4	1.7
10	Influenza and pneumonia (J09-J18)	263	1.3	1.6
11	Septicemia (A40-A41)	248	1.3	1.6
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	190	1.0	1.2
13	Human immunodeficiency virus (HIV) disease (B20-B24)	174	0.9	1.1
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	161	0.8	1.0
15	Pregnancy, childbirth and the puerperium (O00-O99)	124	0.6	0.8
...	All other causes (Residual)	3,665	18.7	23.0
White, female, 45-54 years				
...	All causes	54,032	100.0	295.1
1	Malignant neoplasms (C00-C97)	19,509	36.1	106.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,458	13.8	40.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,567	10.3	30.4
4	Chronic liver disease and cirrhosis (K70,K73-K74)	2,308	4.3	12.6
5	Chronic lower respiratory diseases (J40-J47)	2,000	3.7	10.9
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,912	3.5	10.4
7	Cerebrovascular diseases (I60-I69)	1,670	3.1	9.1
8	Diabetes mellitus (E10-E14)	1,396	2.6	7.6
9	Septicemia (A40-A41)	748	1.4	4.1
10	Influenza and pneumonia (J09-J18)	661	1.2	3.6
11	Viral hepatitis (B15-B19)	591	1.1	3.2
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	534	1.0	2.9
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	359	0.7	2.0
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	273	0.5	1.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	272	0.5	1.5
...	All other causes (Residual)	8,774	16.2	47.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 55-64 years				
...	All causes	96,287	100.0	617.8
1	Malignant neoplasms (C00-C97)	39,791	41.3	255.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,309	15.9	98.2
3	Chronic lower respiratory diseases (J40-J47)	6,072	6.3	39.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,761	3.9	24.1
5	Diabetes mellitus (E10-E14)	3,245	3.4	20.8
6	Cerebrovascular diseases (I60-I69)	3,213	3.3	20.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,417	2.5	15.5
8	Septicemia (A40-A41)	1,577	1.6	10.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,430	1.5	9.2
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,429	1.5	9.2
11	Influenza and pneumonia (J09-J18)	1,203	1.2	7.7
12	Viral hepatitis (B15-B19)	718	0.7	4.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	592	0.6	3.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	490	0.5	3.1
15	Alzheimer's disease (G30)	391	0.4	2.5
...	All other causes (Residual)	14,649	15.2	94.0
White, female, 65-74 years				
...	All causes	148,184	100.0	1,504.9
1	Malignant neoplasms (C00-C97)	55,506	37.5	563.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	26,429	17.8	268.4
3	Chronic lower respiratory diseases (J40-J47)	14,423	9.7	146.5
4	Cerebrovascular diseases (I60-I69)	6,574	4.4	66.8
5	Diabetes mellitus (E10-E14)	4,797	3.2	48.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,007	2.0	30.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,836	1.9	28.8
8	Septicemia (A40-A41)	2,494	1.7	25.3
9	Influenza and pneumonia (J09-J18)	2,173	1.5	22.1
10	Alzheimer's disease (G30)	2,127	1.4	21.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1,956	1.3	19.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,179	0.8	12.0
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	947	0.6	9.6
14	Parkinson's disease (G20-G21)	784	0.5	8.0
15	Pneumonitis due to solids and liquids (J69)	645	0.4	6.6
...	All other causes (Residual)	22,307	15.1	226.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 75-84 years				
...	All causes	274,965	100.0	4,165.4
1	Malignant neoplasms (C00-C97)	65,260	23.7	988.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	62,155	22.6	941.6
3	Chronic lower respiratory diseases (J40-J47)	23,625	8.6	357.9
4	Cerebrovascular diseases (I60-I69)	18,494	6.7	280.2
5	Alzheimer's disease (G30)	13,701	5.0	207.6
6	Diabetes mellitus (E10-E14)	7,501	2.7	113.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,937	2.2	89.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,862	2.1	88.8
9	Influenza and pneumonia (J09-J18)	5,656	2.1	85.7
10	Septicemia (A40-A41)	4,370	1.6	66.2
11	Parkinson's disease (G20-G21)	3,445	1.3	52.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,118	1.1	47.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,951	0.7	29.6
14	Pneumonitis due to solids and liquids (J69)	1,875	0.7	28.4
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,550	0.6	23.5
...	All other causes (Residual)	50,465	18.4	764.5
White, female, 85 years and over				
...	All causes	444,775	100.0	13,419.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	135,451	30.5	4,086.7
2	Malignant neoplasms (C00-C97)	46,063	10.4	1,389.8
3	Alzheimer's disease (G30)	37,191	8.4	1,122.1
4	Cerebrovascular diseases (I60-I69)	34,894	7.8	1,052.8
5	Chronic lower respiratory diseases (J40-J47)	21,005	4.7	633.7
6	Influenza and pneumonia (J09-J18)	12,816	2.9	386.7
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,283	2.3	310.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9,045	2.0	272.9
9	Diabetes mellitus (E10-E14)	8,123	1.8	245.1
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7,075	1.6	213.5
11	Septicemia (A40-A41)	5,332	1.2	160.9
12	Parkinson's disease (G20-G21)	4,216	0.9	127.2
13	Pneumonitis due to solids and liquids (J69)	3,913	0.9	118.1
14	Atherosclerosis (I70)	2,623	0.6	79.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,420	0.5	73.0
...	All other causes (Residual)	104,325	23.5	3,147.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, all ages				
...	All causes	286,959	100.0	682.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	69,083	24.1	164.2
2	Malignant neoplasms (C00-C97)	65,930	23.0	156.7
3	Cerebrovascular diseases (I60-I69)	15,965	5.6	38.0
4	Diabetes mellitus (E10-E14)	12,126	4.2	28.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,069	4.2	28.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,841	3.1	21.0
7	Chronic lower respiratory diseases (J40-J47)	8,715	3.0	20.7
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7,818	2.7	18.6
9	Septicemia (A40-A41)	6,001	2.1	14.3
10	Alzheimer's disease (G30)	5,220	1.8	12.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,116	1.8	12.2
12	Influenza and pneumonia (J09-J18)	4,936	1.7	11.7
13	Human immunodeficiency virus (HIV) disease (B20-B24)	4,662	1.6	11.1
14	Certain conditions originating in the perinatal period (P00-P96)	4,031	1.4	9.6
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,635	0.9	6.3
...	All other causes (Residual)	53,811	18.8	127.9
Black, both sexes, 1-4 years				
...	All causes	1,041	100.0	38.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	267	25.6	9.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	156	15.0	5.7
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	127	12.2	4.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	51	4.9	1.9
5	Malignant neoplasms (C00-C97)	48	4.6	1.8
6	Influenza and pneumonia (J09-J18)	25	2.4	0.9
7	Chronic lower respiratory diseases (J40-J47)	21	2.0	0.8
8	Certain conditions originating in the perinatal period (P00-P96)	16	1.5	*
9	Cerebrovascular diseases (I60-I69)	15	1.4	*
10	Anemias (D50-D64)	14	1.3	*
11	Septicemia (A40-A41)	12	1.2	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	1.0	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.5	*
14	Meningitis (G00,G03)	4	0.4	*
14	Pneumonitis due to solids and liquids (J69)	4	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.4	*
...	All other causes (Residual)	262	25.2	9.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 5-14 years				
...	All causes	1,145	100.0	17.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	319	27.9	4.8
2	Malignant neoplasms (C00-C97)	146	12.8	2.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	101	8.8	1.5
4	Chronic lower respiratory diseases (J40-J47)	72	6.3	1.1
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	63	5.5	0.9
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	61	5.3	0.9
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	44	3.8	0.7
8	Cerebrovascular diseases (I60-I69)	22	1.9	0.3
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	1.5	*
10	Septicemia (A40-A41)	15	1.3	*
11	Anemias (D50-D64)	14	1.2	*
12	Influenza and pneumonia (J09-J18)	12	1.0	*
13	Diabetes mellitus (E10-E14)	7	0.6	*
14	Certain conditions originating in the perinatal period (P00-P96)	6	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.4	*
...	All other causes (Residual)	241	21.0	3.6
Black, both sexes, 15-24 years				
...	All causes	6,675	100.0	93.2
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,816	42.2	39.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,457	21.8	20.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	469	7.0	6.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	336	5.0	4.7
5	Malignant neoplasms (C00-C97)	267	4.0	3.7
6	Human immunodeficiency virus (HIV) disease (B20-B24)	111	1.7	1.6
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	91	1.4	1.3
8	Anemias (D50-D64)	79	1.2	1.1
9	Chronic lower respiratory diseases (J40-J47)	62	0.9	0.9
10	Diabetes mellitus (E10-E14)	51	0.8	0.7
10	Pregnancy, childbirth and the puerperium (O00-O99)	51	0.8	0.7
12	Cerebrovascular diseases (I60-I69)	49	0.7	0.7
13	Septicemia (A40-A41)	41	0.6	0.6
14	Influenza and pneumonia (J09-J18)	39	0.6	0.5
15	Legal intervention (Y35,Y89.0)	28	0.4	0.4
...	All other causes (Residual)	728	10.9	10.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 25-34 years				
...	All causes	8,920	100.0	152.0
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,401	26.9	40.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,612	18.1	27.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	914	10.2	15.6
4	Malignant neoplasms (C00-C97)	589	6.6	10.0
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	504	5.7	8.6
6	Human immunodeficiency virus (HIV) disease (B20-B24)	472	5.3	8.0
7	Diabetes mellitus (E10-E14)	209	2.3	3.6
8	Cerebrovascular diseases (I60-I69)	137	1.5	2.3
9	Pregnancy, childbirth and the puerperium (O00-O99)	118	1.3	2.0
10	Anemias (D50-D64)	109	1.2	1.9
11	Chronic lower respiratory diseases (J40-J47)	98	1.1	1.7
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	95	1.1	1.6
13	Septicemia (A40-A41)	86	1.0	1.5
14	Influenza and pneumonia (J09-J18)	85	1.0	1.4
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	70	0.8	1.2
...	All other causes (Residual)	1,421	15.9	24.2
Black, both sexes, 35-44 years				
...	All causes	14,016	100.0	250.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,726	19.4	48.7
2	Malignant neoplasms (C00-C97)	2,111	15.1	37.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,602	11.4	28.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,088	7.8	19.4
5	Human immunodeficiency virus (HIV) disease (B20-B24)	1,077	7.7	19.2
6	Cerebrovascular diseases (I60-I69)	559	4.0	10.0
7	Diabetes mellitus (E10-E14)	514	3.7	9.2
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	452	3.2	8.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	281	2.0	5.0
10	Septicemia (A40-A41)	212	1.5	3.8
11	Chronic lower respiratory diseases (J40-J47)	182	1.3	3.2
12	Chronic liver disease and cirrhosis (K70,K73-K74)	180	1.3	3.2
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	175	1.2	3.1
14	Influenza and pneumonia (J09-J18)	144	1.0	2.6
15	Anemias (D50-D64)	132	0.9	2.4
...	All other causes (Residual)	2,581	18.4	46.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 45-54 years				
...	All causes	34,093	100.0	591.5
1	Malignant neoplasms (C00-C97)	8,561	25.1	148.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,119	23.8	140.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,305	6.8	40.0
4	Cerebrovascular diseases (I60-I69)	1,726	5.1	29.9
5	Human immunodeficiency virus (HIV) disease (B20-B24)	1,635	4.8	28.4
6	Diabetes mellitus (E10-E14)	1,394	4.1	24.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	817	2.4	14.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	800	2.3	13.9
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	693	2.0	12.0
10	Chronic lower respiratory diseases (J40-J47)	649	1.9	11.3
11	Septicemia (A40-A41)	607	1.8	10.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	532	1.6	9.2
13	Influenza and pneumonia (J09-J18)	392	1.1	6.8
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	337	1.0	5.8
15	Viral hepatitis (B15-B19)	314	0.9	5.4
...	All other causes (Residual)	5,212	15.3	90.4
Black, both sexes, 55-64 years				
...	All causes	51,816	100.0	1,286.0
1	Malignant neoplasms (C00-C97)	16,375	31.6	406.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,826	24.8	318.3
3	Cerebrovascular diseases (I60-I69)	2,652	5.1	65.8
4	Diabetes mellitus (E10-E14)	2,489	4.8	61.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,769	3.4	43.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,538	3.0	38.2
7	Chronic lower respiratory diseases (J40-J47)	1,463	2.8	36.3
8	Septicemia (A40-A41)	1,105	2.1	27.4
9	Human immunodeficiency virus (HIV) disease (B20-B24)	1,029	2.0	25.5
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,026	2.0	25.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	884	1.7	21.9
12	Influenza and pneumonia (J09-J18)	599	1.2	14.9
13	Viral hepatitis (B15-B19)	502	1.0	12.5
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	323	0.6	8.0
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	206	0.4	5.1
...	All other causes (Residual)	7,030	13.6	174.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 65-74 years				
...	All causes	52,491	100.0	2,526.1
1	Malignant neoplasms (C00-C97)	16,627	31.7	800.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,184	25.1	634.5
3	Cerebrovascular diseases (I60-I69)	3,121	5.9	150.2
4	Diabetes mellitus (E10-E14)	2,781	5.3	133.8
5	Chronic lower respiratory diseases (J40-J47)	2,190	4.2	105.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,961	3.7	94.4
7	Septicemia (A40-A41)	1,269	2.4	61.1
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	951	1.8	45.8
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	915	1.7	44.0
10	Influenza and pneumonia (J09-J18)	870	1.7	41.9
11	Alzheimer's disease (G30)	405	0.8	19.5
12	Chronic liver disease and cirrhosis (K70,K73-K74)	385	0.7	18.5
13	Human immunodeficiency virus (HIV) disease (B20-B24)	261	0.5	12.6
14	Pneumonitis due to solids and liquids (J69)	234	0.4	11.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	233	0.4	11.2
...	All other causes (Residual)	7,104	13.5	341.9
Black, both sexes, 75-84 years				
...	All causes	57,543	100.0	5,371.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,441	26.8	1,441.5
2	Malignant neoplasms (C00-C97)	14,040	24.4	1,310.7
3	Cerebrovascular diseases (I60-I69)	3,920	6.8	365.9
4	Diabetes mellitus (E10-E14)	2,827	4.9	263.9
5	Chronic lower respiratory diseases (J40-J47)	2,330	4.0	217.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,222	3.9	207.4
7	Alzheimer's disease (G30)	1,664	2.9	155.3
8	Septicemia (A40-A41)	1,422	2.5	132.7
9	Influenza and pneumonia (J09-J18)	1,230	2.1	114.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,132	2.0	105.7
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	822	1.4	76.7
12	Pneumonitis due to solids and liquids (J69)	403	0.7	37.6
13	Parkinson's disease (G20-G21)	327	0.6	30.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	288	0.5	26.9
15	Aortic aneurysm and dissection (I71)	203	0.4	19.0
...	All other causes (Residual)	9,272	16.1	865.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 85 years and over				
...	All causes	51,795	100.0	13,187.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,324	29.6	3,901.5
2	Malignant neoplasms (C00-C97)	7,157	13.8	1,822.2
3	Cerebrovascular diseases (I60-I69)	3,716	7.2	946.1
4	Alzheimer's disease (G30)	3,090	6.0	786.7
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,878	3.6	478.1
6	Diabetes mellitus (E10-E14)	1,850	3.6	471.0
7	Chronic lower respiratory diseases (J40-J47)	1,639	3.2	417.3
8	Influenza and pneumonia (J09-J18)	1,467	2.8	373.5
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,374	2.7	349.8
10	Septicemia (A40-A41)	1,142	2.2	290.8
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	645	1.2	164.2
12	Pneumonitis due to solids and liquids (J69)	456	0.9	116.1
13	Parkinson's disease (G20-G21)	278	0.5	70.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	248	0.5	63.1
15	Atherosclerosis (I70)	230	0.4	58.6
...	All other causes (Residual)	11,301	21.8	2,877.3
Black, male, all ages				
...	All causes	145,802	100.0	725.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	35,089	24.1	174.6
2	Malignant neoplasms (C00-C97)	33,967	23.3	169.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,074	5.5	40.2
4	Cerebrovascular diseases (I60-I69)	6,938	4.8	34.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6,704	4.6	33.4
6	Diabetes mellitus (E10-E14)	5,640	3.9	28.1
7	Chronic lower respiratory diseases (J40-J47)	4,532	3.1	22.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,016	2.8	20.0
9	Human immunodeficiency virus (HIV) disease (B20-B24)	3,047	2.1	15.2
10	Septicemia (A40-A41)	2,691	1.8	13.4
11	Influenza and pneumonia (J09-J18)	2,380	1.6	11.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,218	1.5	11.0
13	Certain conditions originating in the perinatal period (P00-P96)	2,216	1.5	11.0
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,755	1.2	8.7
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,715	1.2	8.5
...	All other causes (Residual)	24,820	17.0	123.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 1-4 years				
...	All causes	595	100.0	42.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	170	28.6	12.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	93	15.6	6.7
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	57	9.6	4.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	29	4.9	2.1
5	Malignant neoplasms (C00-C97)	27	4.5	1.9
6	Influenza and pneumonia (J09-J18)	15	2.5	*
7	Chronic lower respiratory diseases (J40-J47)	11	1.8	*
8	Anemias (D50-D64)	9	1.5	*
8	Certain conditions originating in the perinatal period (P00-P96)	9	1.5	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.0	*
11	Septicemia (A40-A41)	5	0.8	*
11	Cerebrovascular diseases (I60-I69)	5	0.8	*
13	Meningococcal infection (A39)	3	0.5	*
13	Pneumonitis due to solids and liquids (J69)	3	0.5	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.5	*
...	All other causes (Residual)	150	25.2	10.8
Black, male, 5-14 years				
...	All causes	667	100.0	19.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	213	31.9	6.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	69	10.3	2.0
3	Malignant neoplasms (C00-C97)	65	9.7	1.9
4	Chronic lower respiratory diseases (J40-J47)	52	7.8	1.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	39	5.8	1.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	35	5.2	1.0
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	23	3.4	0.7
8	Cerebrovascular diseases (I60-I69)	14	2.1	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	1.3	*
10	Septicemia (A40-A41)	8	1.2	*
10	Influenza and pneumonia (J09-J18)	8	1.2	*
12	Anemias (D50-D64)	6	0.9	*
13	Diabetes mellitus (E10-E14)	3	0.4	*
13	Certain conditions originating in the perinatal period (P00-P96)	3	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.3	*
...	All other causes (Residual)	118	17.7	3.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 15-24 years				
...	All causes	5,129	100.0	142.8
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,548	49.7	71.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,071	20.9	29.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	397	7.7	11.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	211	4.1	5.9
5	Malignant neoplasms (C00-C97)	149	2.9	4.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	71	1.4	2.0
7	Anemias (D50-D64)	55	1.1	1.5
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	55	1.1	1.5
9	Chronic lower respiratory diseases (J40-J47)	43	0.8	1.2
10	Legal intervention (Y35,Y89.0)	27	0.5	0.8
11	Septicemia (A40-A41)	26	0.5	0.7
11	Diabetes mellitus (E10-E14)	26	0.5	0.7
13	Influenza and pneumonia (J09-J18)	24	0.5	0.7
14	Cerebrovascular diseases (I60-I69)	21	0.4	0.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	0.2	*
...	All other causes (Residual)	393	7.7	10.9
Black, male, 25-34 years				
...	All causes	6,071	100.0	216.7
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,132	35.1	76.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,182	19.5	42.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	576	9.5	20.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	433	7.1	15.5
5	Human immunodeficiency virus (HIV) disease (B20-B24)	275	4.5	9.8
6	Malignant neoplasms (C00-C97)	257	4.2	9.2
7	Diabetes mellitus (E10-E14)	113	1.9	4.0
8	Cerebrovascular diseases (I60-I69)	75	1.2	2.7
9	Chronic lower respiratory diseases (J40-J47)	69	1.1	2.5
10	Anemias (D50-D64)	52	0.9	1.9
11	Influenza and pneumonia (J09-J18)	50	0.8	1.8
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	43	0.7	1.5
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	42	0.7	1.5
14	Septicemia (A40-A41)	35	0.6	1.2
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	29	0.5	1.0
...	All other causes (Residual)	708	11.7	25.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 35-44 years				
...	All causes	8,115	100.0	307.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,675	20.6	63.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,109	13.7	42.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	911	11.2	34.5
4	Malignant neoplasms (C00-C97)	794	9.8	30.1
5	Human immunodeficiency virus (HIV) disease (B20-B24)	654	8.1	24.8
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	355	4.4	13.5
7	Diabetes mellitus (E10-E14)	304	3.7	11.5
8	Cerebrovascular diseases (I60-I69)	289	3.6	11.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	156	1.9	5.9
10	Chronic liver disease and cirrhosis (K70,K73-K74)	106	1.3	4.0
11	Septicemia (A40-A41)	105	1.3	4.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	98	1.2	3.7
13	Chronic lower respiratory diseases (J40-J47)	91	1.1	3.4
14	Influenza and pneumonia (J09-J18)	85	1.0	3.2
15	Aortic aneurysm and dissection (I71)	82	1.0	3.1
...	All other causes (Residual)	1,301	16.0	49.3
Black, male, 45-54 years				
...	All causes	19,393	100.0	716.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,168	26.6	190.9
2	Malignant neoplasms (C00-C97)	4,085	21.1	150.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,562	8.1	57.7
4	Human immunodeficiency virus (HIV) disease (B20-B24)	1,090	5.6	40.3
5	Cerebrovascular diseases (I60-I69)	911	4.7	33.6
6	Diabetes mellitus (E10-E14)	792	4.1	29.3
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	552	2.8	20.4
8	Chronic liver disease and cirrhosis (K70,K73-K74)	522	2.7	19.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	399	2.1	14.7
10	Septicemia (A40-A41)	315	1.6	11.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	288	1.5	10.6
12	Chronic lower respiratory diseases (J40-J47)	286	1.5	10.6
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	264	1.4	9.8
14	Influenza and pneumonia (J09-J18)	229	1.2	8.5
15	Viral hepatitis (B15-B19)	212	1.1	7.8
...	All other causes (Residual)	2,718	14.0	100.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 55-64 years				
...	All causes	30,458	100.0	1,662.1
1	Malignant neoplasms (C00-C97)	9,101	29.9	496.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,023	26.3	437.8
3	Cerebrovascular diseases (I60-I69)	1,524	5.0	83.2
4	Diabetes mellitus (E10-E14)	1,354	4.4	73.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,263	4.1	68.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	847	2.8	46.2
7	Chronic lower respiratory diseases (J40-J47)	796	2.6	43.4
8	Chronic liver disease and cirrhosis (K70,K73-K74)	727	2.4	39.7
9	Human immunodeficiency virus (HIV) disease (B20-B24)	717	2.4	39.1
10	Septicemia (A40-A41)	578	1.9	31.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	501	1.6	27.3
12	Viral hepatitis (B15-B19)	351	1.2	19.2
13	Influenza and pneumonia (J09-J18)	345	1.1	18.8
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	245	0.8	13.4
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	168	0.6	9.2
...	All other causes (Residual)	3,918	12.9	213.8
Black, male, 65-74 years				
...	All causes	28,395	100.0	3,205.6
1	Malignant neoplasms (C00-C97)	9,104	32.1	1,027.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,510	26.4	847.8
3	Cerebrovascular diseases (I60-I69)	1,617	5.7	182.6
4	Diabetes mellitus (E10-E14)	1,348	4.7	152.2
5	Chronic lower respiratory diseases (J40-J47)	1,221	4.3	137.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	974	3.4	110.0
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	601	2.1	67.8
8	Septicemia (A40-A41)	591	2.1	66.7
9	Influenza and pneumonia (J09-J18)	492	1.7	55.5
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	485	1.7	54.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	249	0.9	28.1
12	Human immunodeficiency virus (HIV) disease (B20-B24)	189	0.7	21.3
13	Alzheimer's disease (G30)	169	0.6	19.1
14	Pneumonitis due to solids and liquids (J69)	144	0.5	16.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	111	0.4	12.5
...	All other causes (Residual)	3,590	12.6	405.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 75-84 years				
...	All causes	26,602	100.0	6,721.5
1	Malignant neoplasms (C00-C97)	7,230	27.2	1,826.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,152	26.9	1,807.1
3	Cerebrovascular diseases (I60-I69)	1,575	5.9	398.0
4	Chronic lower respiratory diseases (J40-J47)	1,225	4.6	309.5
5	Diabetes mellitus (E10-E14)	1,179	4.4	297.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	956	3.6	241.6
7	Septicemia (A40-A41)	622	2.3	157.2
8	Alzheimer's disease (G30)	611	2.3	154.4
9	Influenza and pneumonia (J09-J18)	588	2.2	148.6
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	451	1.7	114.0
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	445	1.7	112.4
12	Pneumonitis due to solids and liquids (J69)	233	0.9	58.9
13	Parkinson's disease (G20-G21)	196	0.7	49.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	122	0.5	30.8
15	Chronic liver disease and cirrhosis (K70,K73-K74)	91	0.3	23.0
...	All other causes (Residual)	3,926	14.8	992.0
Black, male, 85 years and over				
...	All causes	16,243	100.0	14,715.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,639	28.6	4,202.7
2	Malignant neoplasms (C00-C97)	3,151	19.4	2,854.6
3	Cerebrovascular diseases (I60-I69)	888	5.5	804.5
4	Chronic lower respiratory diseases (J40-J47)	731	4.5	662.2
5	Alzheimer's disease (G30)	680	4.2	616.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	613	3.8	555.3
7	Diabetes mellitus (E10-E14)	518	3.2	469.3
8	Influenza and pneumonia (J09-J18)	501	3.1	453.9
9	Septicemia (A40-A41)	359	2.2	325.2
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	357	2.2	323.4
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	254	1.6	230.1
12	Pneumonitis due to solids and liquids (J69)	167	1.0	151.3
13	Parkinson's disease (G20-G21)	129	0.8	116.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	85	0.5	77.0
15	Atherosclerosis (I70)	60	0.4	54.4
...	All other causes (Residual)	3,111	19.2	2,818.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, all ages				
...	All causes	141,157	100.0	642.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	33,994	24.1	154.8
2	Malignant neoplasms (C00-C97)	31,963	22.6	145.5
3	Cerebrovascular diseases (I60-I69)	9,027	6.4	41.1
4	Diabetes mellitus (E10-E14)	6,486	4.6	29.5
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,825	3.4	22.0
6	Chronic lower respiratory diseases (J40-J47)	4,183	3.0	19.0
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,995	2.8	18.2
8	Alzheimer's disease (G30)	3,732	2.6	17.0
9	Septicemia (A40-A41)	3,310	2.3	15.1
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,898	2.1	13.2
11	Influenza and pneumonia (J09-J18)	2,556	1.8	11.6
12	Certain conditions originating in the perinatal period (P00-P96)	1,815	1.3	8.3
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,615	1.1	7.4
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,114	0.8	5.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	920	0.7	4.2
...	All other causes (Residual)	28,724	20.3	130.8
Black, female, 1-4 years				
...	All causes	446	100.0	33.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	97	21.7	7.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	70	15.7	5.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	63	14.1	4.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	22	4.9	1.6
5	Malignant neoplasms (C00-C97)	21	4.7	1.6
6	Cerebrovascular diseases (I60-I69)	10	2.2	*
6	Influenza and pneumonia (J09-J18)	10	2.2	*
6	Chronic lower respiratory diseases (J40-J47)	10	2.2	*
9	Septicemia (A40-A41)	7	1.6	*
9	Certain conditions originating in the perinatal period (P00-P96)	7	1.6	*
11	Anemias (D50-D64)	5	1.1	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.9	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.7	*
14	Meningitis (G00,G03)	2	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
...	All other causes (Residual)	113	25.3	8.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 5-14 years				
...	All causes	478	100.0	14.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	106	22.2	3.2
2	Malignant neoplasms (C00-C97)	81	16.9	2.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	32	6.7	1.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	28	5.9	0.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	22	4.6	0.7
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	21	4.4	0.6
7	Chronic lower respiratory diseases (J40-J47)	20	4.2	0.6
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	1.7	*
8	Anemias (D50-D64)	8	1.7	*
8	Cerebrovascular diseases (I60-I69)	8	1.7	*
11	Septicemia (A40-A41)	7	1.5	*
12	Diabetes mellitus (E10-E14)	4	0.8	*
12	Influenza and pneumonia (J09-J18)	4	0.8	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.6	*
14	Certain conditions originating in the perinatal period (P00-P96)	3	0.6	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.6	*
...	All other causes (Residual)	120	25.1	3.6
Black, female, 15-24 years				
...	All causes	1,546	100.0	43.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	386	25.0	10.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	268	17.3	7.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	125	8.1	3.5
4	Malignant neoplasms (C00-C97)	118	7.6	3.3
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	72	4.7	2.0
6	Pregnancy, childbirth and the puerperium (O00-O99)	51	3.3	1.4
7	Human immunodeficiency virus (HIV) disease (B20-B24)	40	2.6	1.1
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	36	2.3	1.0
9	Cerebrovascular diseases (I60-I69)	28	1.8	0.8
10	Diabetes mellitus (E10-E14)	25	1.6	0.7
11	Anemias (D50-D64)	24	1.6	0.7
12	Chronic lower respiratory diseases (J40-J47)	19	1.2	*
13	Septicemia (A40-A41)	15	1.0	*
13	Influenza and pneumonia (J09-J18)	15	1.0	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	1.0	*
...	All other causes (Residual)	309	20.0	8.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 25-34 years				
...	All causes	2,849	100.0	92.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	430	15.1	14.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	338	11.9	11.0
3	Malignant neoplasms (C00-C97)	332	11.7	10.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	269	9.4	8.8
5	Human immunodeficiency virus (HIV) disease (B20-B24)	197	6.9	6.4
6	Pregnancy, childbirth and the puerperium (O00-O99)	118	4.1	3.8
7	Diabetes mellitus (E10-E14)	96	3.4	3.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	71	2.5	2.3
9	Cerebrovascular diseases (I60-I69)	62	2.2	2.0
10	Anemias (D50-D64)	57	2.0	1.9
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	52	1.8	1.7
12	Septicemia (A40-A41)	51	1.8	1.7
13	Influenza and pneumonia (J09-J18)	35	1.2	1.1
14	Chronic lower respiratory diseases (J40-J47)	29	1.0	0.9
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	28	1.0	0.9
...	All other causes (Residual)	684	24.0	22.3
Black, female, 35-44 years				
...	All causes	5,901	100.0	199.3
1	Malignant neoplasms (C00-C97)	1,317	22.3	44.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,051	17.8	35.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	493	8.4	16.6
4	Human immunodeficiency virus (HIV) disease (B20-B24)	423	7.2	14.3
5	Cerebrovascular diseases (I60-I69)	270	4.6	9.1
6	Diabetes mellitus (E10-E14)	210	3.6	7.1
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	177	3.0	6.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	125	2.1	4.2
9	Septicemia (A40-A41)	107	1.8	3.6
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	97	1.6	3.3
11	Chronic lower respiratory diseases (J40-J47)	91	1.5	3.1
12	Pregnancy, childbirth and the puerperium (O00-O99)	80	1.4	2.7
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	77	1.3	2.6
14	Chronic liver disease and cirrhosis (K70,K73-K74)	74	1.3	2.5
15	Anemias (D50-D64)	66	1.1	2.2
...	All other causes (Residual)	1,243	21.1	42.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 45-54 years				
...	All causes	14,700	100.0	481.0
1	Malignant neoplasms (C00-C97)	4,476	30.4	146.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,951	20.1	96.6
3	Cerebrovascular diseases (I60-I69)	815	5.5	26.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	743	5.1	24.3
5	Diabetes mellitus (E10-E14)	602	4.1	19.7
6	Human immunodeficiency virus (HIV) disease (B20-B24)	545	3.7	17.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	401	2.7	13.1
8	Chronic lower respiratory diseases (J40-J47)	363	2.5	11.9
9	Chronic liver disease and cirrhosis (K70,K73-K74)	295	2.0	9.7
10	Septicemia (A40-A41)	292	2.0	9.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	244	1.7	8.0
12	Influenza and pneumonia (J09-J18)	163	1.1	5.3
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	141	1.0	4.6
14	Viral hepatitis (B15-B19)	102	0.7	3.3
15	Anemias (D50-D64)	75	0.5	2.5
...	All other causes (Residual)	2,492	17.0	81.5
Black, female, 55-64 years				
...	All causes	21,358	100.0	972.2
1	Malignant neoplasms (C00-C97)	7,274	34.1	331.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,803	22.5	218.6
3	Diabetes mellitus (E10-E14)	1,135	5.3	51.7
4	Cerebrovascular diseases (I60-I69)	1,128	5.3	51.3
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	691	3.2	31.5
6	Chronic lower respiratory diseases (J40-J47)	667	3.1	30.4
7	Septicemia (A40-A41)	527	2.5	24.0
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	506	2.4	23.0
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	383	1.8	17.4
10	Human immunodeficiency virus (HIV) disease (B20-B24)	312	1.5	14.2
11	Chronic liver disease and cirrhosis (K70,K73-K74)	299	1.4	13.6
12	Influenza and pneumonia (J09-J18)	254	1.2	11.6
13	Viral hepatitis (B15-B19)	151	0.7	6.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	91	0.4	4.1
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	78	0.4	3.6
...	All other causes (Residual)	3,059	14.3	139.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 65-74 years				
...	All causes	24,096	100.0	2,021.2
1	Malignant neoplasms (C00-C97)	7,523	31.2	631.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,674	23.5	475.9
3	Cerebrovascular diseases (I60-I69)	1,504	6.2	126.2
4	Diabetes mellitus (E10-E14)	1,433	5.9	120.2
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	987	4.1	82.8
6	Chronic lower respiratory diseases (J40-J47)	969	4.0	81.3
7	Septicemia (A40-A41)	678	2.8	56.9
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	466	1.9	39.1
9	Influenza and pneumonia (J09-J18)	378	1.6	31.7
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	314	1.3	26.3
11	Alzheimer's disease (G30)	236	1.0	19.8
12	Chronic liver disease and cirrhosis (K70,K73-K74)	136	0.6	11.4
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	122	0.5	10.2
14	Pneumonitis due to solids and liquids (J69)	90	0.4	7.5
15	Viral hepatitis (B15-B19)	73	0.3	6.1
...	All other causes (Residual)	3,513	14.6	294.7
Black, female, 75-84 years				
...	All causes	30,941	100.0	4,580.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,289	26.8	1,227.2
2	Malignant neoplasms (C00-C97)	6,810	22.0	1,008.2
3	Cerebrovascular diseases (I60-I69)	2,345	7.6	347.2
4	Diabetes mellitus (E10-E14)	1,648	5.3	244.0
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,266	4.1	187.4
6	Chronic lower respiratory diseases (J40-J47)	1,105	3.6	163.6
7	Alzheimer's disease (G30)	1,053	3.4	155.9
8	Septicemia (A40-A41)	800	2.6	118.4
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	681	2.2	100.8
10	Influenza and pneumonia (J09-J18)	642	2.1	95.1
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	377	1.2	55.8
12	Pneumonitis due to solids and liquids (J69)	170	0.5	25.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	166	0.5	24.6
14	Parkinson's disease (G20-G21)	131	0.4	19.4
15	Aortic aneurysm and dissection (I71)	120	0.4	17.8
...	All other causes (Residual)	5,338	17.3	790.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 85 years and over				
...	All causes	35,552	100.0	12,589.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,685	30.1	3,783.8
2	Malignant neoplasms (C00-C97)	4,006	11.3	1,418.6
3	Cerebrovascular diseases (I60-I69)	2,828	8.0	1,001.5
4	Alzheimer's disease (G30)	2,410	6.8	853.4
5	Diabetes mellitus (E10-E14)	1,332	3.7	471.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,265	3.6	448.0
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,017	2.9	360.1
8	Influenza and pneumonia (J09-J18)	966	2.7	342.1
9	Chronic lower respiratory diseases (J40-J47)	908	2.6	321.5
10	Septicemia (A40-A41)	783	2.2	277.3
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	391	1.1	138.5
12	Pneumonitis due to solids and liquids (J69)	289	0.8	102.3
13	Atherosclerosis (I70)	170	0.5	60.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	163	0.5	57.7
15	Parkinson's disease (G20-G21)	149	0.4	52.8
...	All other causes (Residual)	8,190	23.0	2,900.3
American Indian, both sexes, all ages				
...	All causes	15,565	100.0	365.1
1	Malignant neoplasms (C00-C97)	2,962	19.0	69.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,793	17.9	65.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,701	10.9	39.9
4	Diabetes mellitus (E10-E14)	857	5.5	20.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	787	5.1	18.5
6	Chronic lower respiratory diseases (J40-J47)	702	4.5	16.5
7	Cerebrovascular diseases (I60-I69)	559	3.6	13.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	469	3.0	11.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	339	2.2	8.0
10	Influenza and pneumonia (J09-J18)	326	2.1	7.6
11	Alzheimer's disease (G30)	264	1.7	6.2
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	257	1.7	6.0
13	Septicemia (A40-A41)	244	1.6	5.7
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	144	0.9	3.4
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	130	0.8	3.0
...	All other causes (Residual)	3,031	19.5	71.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 1-4 years				
...	All causes	93	100.0	29.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	31	33.3	9.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	10	10.8	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	9.7	*
4	Malignant neoplasms (C00-C97)	6	6.5	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	5.4	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	2.2	*
6	Chronic lower respiratory diseases (J40-J47)	2	2.2	*
8	Septicemia (A40-A41)	1	1.1	*
8	Meningitis (G00,G03)	1	1.1	*
8	Cerebrovascular diseases (I60-I69)	1	1.1	*
8	Influenza and pneumonia (J09-J18)	1	1.1	*
8	Acute bronchitis and bronchiolitis (J20-J21)	1	1.1	*
8	Pneumonitis due to solids and liquids (J69)	1	1.1	*
8	Certain conditions originating in the perinatal period (P00-P96)	1	1.1	*
8	Legal intervention (Y35,Y89.0)	1	1.1	*
8	Complications of medical and surgical care (Y40-Y84,Y88)	1	1.1	*
...	All other causes (Residual)	19	20.4	*
American Indian, both sexes, 5-14 years				
...	All causes	108	100.0	14.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	47	43.5	6.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	14	13.0	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	10	9.3	*
4	Malignant neoplasms (C00-C97)	8	7.4	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	3.7	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	2.8	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	1.9	*
7	Cerebrovascular diseases (I60-I69)	2	1.9	*
7	Influenza and pneumonia (J09-J18)	2	1.9	*
10	Septicemia (A40-A41)	1	0.9	*
10	Chronic lower respiratory diseases (J40-J47)	1	0.9	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.9	*
...	All other causes (Residual)	13	12.0	*

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 15-24 years				
...	All causes	615	100.0	81.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	270	43.9	35.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	158	25.7	20.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	75	12.2	9.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	16	2.6	*
5	Malignant neoplasms (C00-C97)	14	2.3	*
6	Pregnancy, childbirth and the puerperium (O00-O99)	5	0.8	*
7	Chronic liver disease and cirrhosis (K70,K73-K74)	4	0.7	*
7	Legal intervention (Y35,Y89.0)	4	0.7	*
9	Cerebrovascular diseases (I60-I69)	3	0.5	*
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.5	*
11	Septicemia (A40-A41)	2	0.3	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2	0.3	*
11	Influenza and pneumonia (J09-J18)	2	0.3	*
11	Chronic lower respiratory diseases (J40-J47)	2	0.3	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.3	*
...	All other causes (Residual)	53	8.6	7.0
American Indian, both sexes, 25-34 years				
...	All causes	795	100.0	122.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	310	39.0	47.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	100	12.6	15.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	73	9.2	11.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	48	6.0	7.4
5	Malignant neoplasms (C00-C97)	45	5.7	6.9
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	38	4.8	5.8
7	Human immunodeficiency virus (HIV) disease (B20-B24)	9	1.1	*
7	Diabetes mellitus (E10-E14)	9	1.1	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	1.1	*
10	Influenza and pneumonia (J09-J18)	7	0.9	*
11	Cerebrovascular diseases (I60-I69)	6	0.8	*
12	Septicemia (A40-A41)	5	0.6	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5	0.6	*
12	Chronic lower respiratory diseases (J40-J47)	5	0.6	*
15	Pregnancy, childbirth and the puerperium (O00-O99)	3	0.4	*
15	Legal intervention (Y35,Y89.0)	3	0.4	*
...	All other causes (Residual)	120	15.1	18.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 35-44 years				
...	All causes	1,163	100.0	203.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	302	26.0	52.8
2	Chronic liver disease and cirrhosis (K70,K73-K74)	171	14.7	29.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	145	12.5	25.3
4	Malignant neoplasms (C00-C97)	79	6.8	13.8
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	75	6.4	13.1
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	48	4.1	8.4
7	Diabetes mellitus (E10-E14)	40	3.4	7.0
8	Influenza and pneumonia (J09-J18)	19	1.6	*
9	Cerebrovascular diseases (I60-I69)	15	1.3	*
10	Septicemia (A40-A41)	14	1.2	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	12	1.0	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	0.9	*
13	Viral hepatitis (B15-B19)	10	0.9	*
14	Chronic lower respiratory diseases (J40-J47)	9	0.8	*
15	Pregnancy, childbirth and the puerperium (O00-O99)	5	0.4	*
...	All other causes (Residual)	208	17.9	36.3
American Indian, both sexes, 45-54 years				
...	All causes	2,203	100.0	409.7
1	Malignant neoplasms (C00-C97)	373	16.9	69.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	362	16.4	67.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	307	13.9	57.1
4	Chronic liver disease and cirrhosis (K70,K73-K74)	264	12.0	49.1
5	Diabetes mellitus (E10-E14)	122	5.5	22.7
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	77	3.5	14.3
7	Cerebrovascular diseases (I60-I69)	60	2.7	11.2
8	Chronic lower respiratory diseases (J40-J47)	43	2.0	8.0
9	Viral hepatitis (B15-B19)	42	1.9	7.8
10	Influenza and pneumonia (J09-J18)	36	1.6	6.7
11	Septicemia (A40-A41)	31	1.4	5.8
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	31	1.4	5.8
13	Human immunodeficiency virus (HIV) disease (B20-B24)	30	1.4	5.6
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	28	1.3	5.2
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	15	0.7	*
...	All other causes (Residual)	382	17.3	71.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 55-64 years				
...	All causes	2,704	100.0	782.1
1	Malignant neoplasms (C00-C97)	749	27.7	216.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	522	19.3	151.0
3	Diabetes mellitus (E10-E14)	202	7.5	58.4
4	Chronic liver disease and cirrhosis (K70,K73-K74)	181	6.7	52.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	179	6.6	51.8
6	Chronic lower respiratory diseases (J40-J47)	121	4.5	35.0
7	Cerebrovascular diseases (I60-I69)	77	2.8	22.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	62	2.3	17.9
9	Septicemia (A40-A41)	50	1.8	14.5
10	Influenza and pneumonia (J09-J18)	40	1.5	11.6
11	Viral hepatitis (B15-B19)	34	1.3	9.8
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	28	1.0	8.1
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	21	0.8	6.1
14	Pneumonitis due to solids and liquids (J69)	13	0.5	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	0.4	*
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	10	0.4	*
...	All other causes (Residual)	405	15.0	117.1
American Indian, both sexes, 65-74 years				
...	All causes	2,803	100.0	1,709.9
1	Malignant neoplasms (C00-C97)	835	29.8	509.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	542	19.3	330.6
3	Chronic lower respiratory diseases (J40-J47)	205	7.3	125.1
4	Diabetes mellitus (E10-E14)	201	7.2	122.6
5	Cerebrovascular diseases (I60-I69)	104	3.7	63.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	99	3.5	60.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	83	3.0	50.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	72	2.6	43.9
9	Septicemia (A40-A41)	55	2.0	33.6
10	Influenza and pneumonia (J09-J18)	50	1.8	30.5
11	Alzheimer's disease (G30)	32	1.1	19.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	32	1.1	19.5
13	Pneumonitis due to solids and liquids (J69)	20	0.7	12.2
14	Aortic aneurysm and dissection (I71)	13	0.5	*
15	Parkinson's disease (G20-G21)	12	0.4	*
...	All other causes (Residual)	448	16.0	273.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 75-84 years				
...	All causes	2,685	100.0	3,833.0
1	Malignant neoplasms (C00-C97)	601	22.4	858.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	584	21.8	833.7
3	Diabetes mellitus (E10-E14)	196	7.3	279.8
4	Chronic lower respiratory diseases (J40-J47)	194	7.2	276.9
5	Cerebrovascular diseases (I60-I69)	149	5.5	212.7
6	Alzheimer's disease (G30)	99	3.7	141.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	89	3.3	127.1
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	74	2.8	105.6
9	Influenza and pneumonia (J09-J18)	72	2.7	102.8
10	Septicemia (A40-A41)	44	1.6	62.8
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	36	1.3	51.4
12	Chronic liver disease and cirrhosis (K70,K73-K74)	33	1.2	47.1
13	Parkinson's disease (G20-G21)	30	1.1	42.8
14	Aortic aneurysm and dissection (I71)	21	0.8	30.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	0.6	*
...	All other causes (Residual)	447	16.6	638.1
American Indian, both sexes, 85 years and over				
...	All causes	2,042	100.0	9,615.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	571	28.0	2,688.7
2	Malignant neoplasms (C00-C97)	250	12.2	1,177.2
3	Cerebrovascular diseases (I60-I69)	140	6.9	659.2
4	Alzheimer's disease (G30)	130	6.4	612.1
5	Chronic lower respiratory diseases (J40-J47)	116	5.7	546.2
6	Influenza and pneumonia (J09-J18)	88	4.3	414.4
7	Diabetes mellitus (E10-E14)	86	4.2	405.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	69	3.4	324.9
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	56	2.7	263.7
10	Septicemia (A40-A41)	38	1.9	178.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	30	1.5	141.3
12	Pneumonitis due to solids and liquids (J69)	23	1.1	108.3
13	Parkinson's disease (G20-G21)	18	0.9	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.4	*
14	Nutritional deficiencies (E40-E64)	9	0.4	*
...	All other causes (Residual)	409	20.0	1,925.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, all ages				
...	All causes	8,516	100.0	397.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,608	18.9	75.0
2	Malignant neoplasms (C00-C97)	1,588	18.6	74.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,150	13.5	53.7
4	Diabetes mellitus (E10-E14)	432	5.1	20.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	429	5.0	20.0
6	Chronic lower respiratory diseases (J40-J47)	349	4.1	16.3
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	344	4.0	16.1
8	Cerebrovascular diseases (I60-I69)	258	3.0	12.0
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	204	2.4	9.5
10	Influenza and pneumonia (J09-J18)	172	2.0	8.0
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	153	1.8	7.1
12	Septicemia (A40-A41)	106	1.2	4.9
13	Alzheimer's disease (G30)	89	1.0	4.2
14	Certain conditions originating in the perinatal period (P00-P96)	76	0.9	3.5
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	71	0.8	3.3
...	All other causes (Residual)	1,487	17.5	69.4
American Indian, male, 1-4 years				
...	All causes	55	100.0	34.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19	34.5	*
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7	12.7	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	10.9	*
4	Malignant neoplasms (C00-C97)	4	7.3	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	5.5	*
6	Septicemia (A40-A41)	1	1.8	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.8	*
6	Chronic lower respiratory diseases (J40-J47)	1	1.8	*
6	Certain conditions originating in the perinatal period (P00-P96)	1	1.8	*
6	Complications of medical and surgical care (Y40-Y84,Y88)	1	1.8	*
...	All other causes (Residual)	11	20.0	*

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 5-14 years				
...	All causes	69	100.0	18.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	35	50.7	9.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8	11.6	*
3	Malignant neoplasms (C00-C97)	4	5.8	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	5.8	*
5	Cerebrovascular diseases (I60-I69)	2	2.9	*
6	Septicemia (A40-A41)	1	1.4	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.4	*
6	Influenza and pneumonia (J09-J18)	1	1.4	*
6	Chronic lower respiratory diseases (J40-J47)	1	1.4	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1	1.4	*
...	All other causes (Residual)	11	15.9	*
American Indian, male, 15-24 years				
...	All causes	456	100.0	116.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	201	44.1	51.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	120	26.3	30.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	69	15.1	17.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	11	2.4	*
5	Malignant neoplasms (C00-C97)	8	1.8	*
6	Legal intervention (Y35,Y89.0)	4	0.9	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.7	*
8	Cerebrovascular diseases (I60-I69)	2	0.4	*
8	Chronic liver disease and cirrhosis (K70,K73-K74)	2	0.4	*
10	Septicemia (A40-A41)	1	0.2	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.2	*
10	Diabetes mellitus (E10-E14)	1	0.2	*
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.2	*
10	Influenza and pneumonia (J09-J18)	1	0.2	*
10	Chronic lower respiratory diseases (J40-J47)	1	0.2	*
10	Pneumonitis due to solids and liquids (J69)	1	0.2	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
...	All other causes (Residual)	27	5.9	6.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 25-34 years				
...	All causes	525	100.0	156.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	199	37.9	59.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	80	15.2	23.8
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	59	11.2	17.6
4	Malignant neoplasms (C00-C97)	25	4.8	7.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	24	4.6	7.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	24	4.6	7.1
7	Human immunodeficiency virus (HIV) disease (B20-B24)	9	1.7	*
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	1.3	*
9	Diabetes mellitus (E10-E14)	4	0.8	*
9	Cerebrovascular diseases (I60-I69)	4	0.8	*
9	Chronic lower respiratory diseases (J40-J47)	4	0.8	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3	0.6	*
12	Influenza and pneumonia (J09-J18)	3	0.6	*
14	Septicemia (A40-A41)	2	0.4	*
14	Pneumonitis due to solids and liquids (J69)	2	0.4	*
...	All other causes (Residual)	76	14.5	22.6
American Indian, male, 35-44 years				
...	All causes	749	100.0	258.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	205	27.4	70.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	110	14.7	37.9
3	Chronic liver disease and cirrhosis (K70,K73-K74)	99	13.2	34.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	51	6.8	17.6
5	Malignant neoplasms (C00-C97)	39	5.2	13.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	34	4.5	11.7
7	Diabetes mellitus (E10-E14)	23	3.1	7.9
8	Influenza and pneumonia (J09-J18)	15	2.0	*
9	Septicemia (A40-A41)	9	1.2	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	1.2	*
11	Viral hepatitis (B15-B19)	8	1.1	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	8	1.1	*
11	Cerebrovascular diseases (I60-I69)	8	1.1	*
14	Chronic lower respiratory diseases (J40-J47)	6	0.8	*
15	Aortic aneurysm and dissection (I71)	4	0.5	*
...	All other causes (Residual)	121	16.2	41.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 45-54 years				
...	All causes	1,311	100.0	496.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	259	19.8	98.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	213	16.2	80.6
3	Malignant neoplasms (C00-C97)	185	14.1	70.0
4	Chronic liver disease and cirrhosis (K70,K73-K74)	137	10.5	51.8
5	Diabetes mellitus (E10-E14)	77	5.9	29.1
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	54	4.1	20.4
7	Cerebrovascular diseases (I60-I69)	31	2.4	11.7
8	Influenza and pneumonia (J09-J18)	28	2.1	10.6
9	Viral hepatitis (B15-B19)	27	2.1	10.2
10	Human immunodeficiency virus (HIV) disease (B20-B24)	24	1.8	9.1
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	22	1.7	8.3
12	Chronic lower respiratory diseases (J40-J47)	21	1.6	7.9
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	1.1	*
14	Septicemia (A40-A41)	13	1.0	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	11	0.8	*
...	All other causes (Residual)	195	14.9	73.8
American Indian, male, 55-64 years				
...	All causes	1,590	100.0	951.2
1	Malignant neoplasms (C00-C97)	417	26.2	249.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	363	22.8	217.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	128	8.1	76.6
4	Diabetes mellitus (E10-E14)	115	7.2	68.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	107	6.7	64.0
6	Chronic lower respiratory diseases (J40-J47)	63	4.0	37.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	39	2.5	23.3
8	Cerebrovascular diseases (I60-I69)	37	2.3	22.1
9	Influenza and pneumonia (J09-J18)	23	1.4	13.8
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	23	1.4	13.8
11	Septicemia (A40-A41)	21	1.3	12.6
12	Viral hepatitis (B15-B19)	20	1.3	12.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	11	0.7	*
14	Pneumonitis due to solids and liquids (J69)	10	0.6	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	6	0.4	*
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	0.4	*
...	All other causes (Residual)	201	12.6	120.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 65-74 years				
...	All causes	1,507	100.0	1,971.0
1	Malignant neoplasms (C00-C97)	457	30.3	597.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	325	21.6	425.1
3	Chronic lower respiratory diseases (J40-J47)	108	7.2	141.3
4	Diabetes mellitus (E10-E14)	102	6.8	133.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	65	4.3	85.0
6	Cerebrovascular diseases (I60-I69)	52	3.5	68.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	44	2.9	57.5
8	Influenza and pneumonia (J09-J18)	27	1.8	35.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	27	1.8	35.3
10	Septicemia (A40-A41)	26	1.7	34.0
11	Alzheimer's disease (G30)	17	1.1	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	12	0.8	*
12	Pneumonitis due to solids and liquids (J69)	12	0.8	*
14	Viral hepatitis (B15-B19)	7	0.5	*
14	Aortic aneurysm and dissection (I71)	7	0.5	*
...	All other causes (Residual)	219	14.5	286.4
American Indian, male, 75-84 years				
...	All causes	1,298	100.0	4,451.8
1	Malignant neoplasms (C00-C97)	322	24.8	1,104.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	304	23.4	1,042.6
3	Chronic lower respiratory diseases (J40-J47)	93	7.2	319.0
4	Cerebrovascular diseases (I60-I69)	78	6.0	267.5
5	Diabetes mellitus (E10-E14)	76	5.9	260.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	43	3.3	147.5
7	Alzheimer's disease (G30)	41	3.2	140.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	34	2.6	116.6
9	Influenza and pneumonia (J09-J18)	29	2.2	99.5
10	Septicemia (A40-A41)	23	1.8	78.9
10	Parkinson's disease (G20-G21)	23	1.8	78.9
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	1.1	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	14	1.1	*
12	Chronic liver disease and cirrhosis (K70,K73-K74)	14	1.1	*
15	Aortic aneurysm and dissection (I71)	9	0.7	*
...	All other causes (Residual)	181	13.9	620.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 85 years and over				
...	All causes	743	100.0	10,268.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	205	27.6	2,833.1
2	Malignant neoplasms (C00-C97)	126	17.0	1,741.3
3	Chronic lower respiratory diseases (J40-J47)	47	6.3	649.5
4	Cerebrovascular diseases (I60-I69)	42	5.7	580.4
5	Influenza and pneumonia (J09-J18)	40	5.4	552.8
6	Diabetes mellitus (E10-E14)	34	4.6	469.9
7	Alzheimer's disease (G30)	31	4.2	428.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	27	3.6	373.1
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	26	3.5	359.3
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10	1.3	*
11	Parkinson's disease (G20-G21)	9	1.2	*
12	Septicemia (A40-A41)	7	0.9	*
12	Pneumonitis due to solids and liquids (J69)	7	0.9	*
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	6	0.8	*
15	Nutritional deficiencies (E40-E64)	4	0.5	*
...	All other causes (Residual)	122	16.4	1,686.0
American Indian, female, all ages				
...	All causes	7,049	100.0	332.4
1	Malignant neoplasms (C00-C97)	1,374	19.5	64.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,185	16.8	55.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	551	7.8	26.0
4	Diabetes mellitus (E10-E14)	425	6.0	20.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	358	5.1	16.9
6	Chronic lower respiratory diseases (J40-J47)	353	5.0	16.6
7	Cerebrovascular diseases (I60-I69)	301	4.3	14.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	186	2.6	8.8
9	Alzheimer's disease (G30)	175	2.5	8.3
10	Influenza and pneumonia (J09-J18)	154	2.2	7.3
11	Septicemia (A40-A41)	138	2.0	6.5
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	125	1.8	5.9
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	82	1.2	3.9
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	59	0.8	2.8
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	53	0.8	2.5
...	All other causes (Residual)	1,530	21.7	72.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 1-4 years				
...	All causes	38	100.0	24.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12	31.6	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	7.9	*
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	7.9	*
4	Malignant neoplasms (C00-C97)	2	5.3	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	5.3	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	2.6	*
6	Meningitis (G00,G03)	1	2.6	*
6	Cerebrovascular diseases (I60-I69)	1	2.6	*
6	Influenza and pneumonia (J09-J18)	1	2.6	*
6	Acute bronchitis and bronchiolitis (J20-J21)	1	2.6	*
6	Chronic lower respiratory diseases (J40-J47)	1	2.6	*
6	Pneumonitis due to solids and liquids (J69)	1	2.6	*
6	Legal intervention (Y35,Y89.0)	1	2.6	*
...	All other causes (Residual)	8	21.1	*
American Indian, female, 5-14 years				
...	All causes	39	100.0	10.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12	30.8	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	15.4	*
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6	15.4	*
4	Malignant neoplasms (C00-C97)	4	10.3	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	7.7	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	5.1	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	5.1	*
8	Influenza and pneumonia (J09-J18)	1	2.6	*
8	Complications of medical and surgical care (Y40-Y84,Y88)	1	2.6	*
...	All other causes (Residual)	2	5.1	*

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 15-24 years				
...	All causes	159	100.0	43.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	69	43.4	18.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	38	23.9	10.4
3	Malignant neoplasms (C00-C97)	6	3.8	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	3.8	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	3.1	*
5	Pregnancy, childbirth and the puerperium (O00-O99)	5	3.1	*
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2	1.3	*
8	Septicemia (A40-A41)	1	0.6	*
8	Anemias (D50-D64)	1	0.6	*
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.6	*
8	Cerebrovascular diseases (I60-I69)	1	0.6	*
8	Influenza and pneumonia (J09-J18)	1	0.6	*
8	Chronic lower respiratory diseases (J40-J47)	1	0.6	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.6	*
...	All other causes (Residual)	21	13.2	5.8
American Indian, female, 25-34 years				
...	All causes	270	100.0	85.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	111	41.1	35.2
2	Chronic liver disease and cirrhosis (K70,K73-K74)	24	8.9	7.6
3	Malignant neoplasms (C00-C97)	20	7.4	6.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	20	7.4	6.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	14	5.2	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	14	5.2	*
7	Diabetes mellitus (E10-E14)	5	1.9	*
8	Influenza and pneumonia (J09-J18)	4	1.5	*
9	Septicemia (A40-A41)	3	1.1	*
9	Pregnancy, childbirth and the puerperium (O00-O99)	3	1.1	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2	0.7	*
11	Cerebrovascular diseases (I60-I69)	2	0.7	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	0.7	*
11	Legal intervention (Y35,Y89.0)	2	0.7	*
15	Meningitis (G00,G03)	1	0.4	*
15	Chronic lower respiratory diseases (J40-J47)	1	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.4	*
...	All other causes (Residual)	41	15.2	13.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 35-44 years				
...	All causes	414	100.0	146.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	97	23.4	34.4
2	Chronic liver disease and cirrhosis (K70,K73-K74)	72	17.4	25.5
3	Malignant neoplasms (C00-C97)	40	9.7	14.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	35	8.5	12.4
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	24	5.8	8.5
6	Diabetes mellitus (E10-E14)	17	4.1	*
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	14	3.4	*
8	Cerebrovascular diseases (I60-I69)	7	1.7	*
9	Septicemia (A40-A41)	5	1.2	*
9	Pregnancy, childbirth and the puerperium (O00-O99)	5	1.2	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	4	1.0	*
11	Influenza and pneumonia (J09-J18)	4	1.0	*
13	Anemias (D50-D64)	3	0.7	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3	0.7	*
13	Chronic lower respiratory diseases (J40-J47)	3	0.7	*
...	All other causes (Residual)	81	19.6	28.7
American Indian, female, 45-54 years				
...	All causes	892	100.0	326.2
1	Malignant neoplasms (C00-C97)	188	21.1	68.7
2	Chronic liver disease and cirrhosis (K70,K73-K74)	127	14.2	46.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	103	11.5	37.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	94	10.5	34.4
5	Diabetes mellitus (E10-E14)	45	5.0	16.5
6	Cerebrovascular diseases (I60-I69)	29	3.3	10.6
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	23	2.6	8.4
8	Chronic lower respiratory diseases (J40-J47)	22	2.5	8.0
9	Septicemia (A40-A41)	18	2.0	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	1.9	*
11	Viral hepatitis (B15-B19)	15	1.7	*
12	Influenza and pneumonia (J09-J18)	8	0.9	*
13	Pneumonitis due to solids and liquids (J69)	7	0.8	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	6	0.7	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	0.7	*
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	0.7	*
...	All other causes (Residual)	178	20.0	65.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 55-64 years				
...	All causes	1,114	100.0	623.8
1	Malignant neoplasms (C00-C97)	332	29.8	185.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	159	14.3	89.0
3	Diabetes mellitus (E10-E14)	87	7.8	48.7
4	Chronic liver disease and cirrhosis (K70,K73-K74)	74	6.6	41.4
5	Chronic lower respiratory diseases (J40-J47)	58	5.2	32.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	51	4.6	28.6
7	Cerebrovascular diseases (I60-I69)	40	3.6	22.4
8	Septicemia (A40-A41)	29	2.6	16.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	2.1	12.9
10	Influenza and pneumonia (J09-J18)	17	1.5	*
11	Viral hepatitis (B15-B19)	14	1.3	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10	0.9	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.4	*
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5	0.4	*
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	0.4	*
...	All other causes (Residual)	206	18.5	115.4
American Indian, female, 65-74 years				
...	All causes	1,296	100.0	1,481.7
1	Malignant neoplasms (C00-C97)	378	29.2	432.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	217	16.7	248.1
3	Diabetes mellitus (E10-E14)	99	7.6	113.2
4	Chronic lower respiratory diseases (J40-J47)	97	7.5	110.9
5	Cerebrovascular diseases (I60-I69)	52	4.0	59.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	45	3.5	51.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	39	3.0	44.6
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	34	2.6	38.9
9	Septicemia (A40-A41)	29	2.2	33.2
10	Influenza and pneumonia (J09-J18)	23	1.8	26.3
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	20	1.5	22.9
12	Alzheimer's disease (G30)	15	1.2	*
13	Pneumonitis due to solids and liquids (J69)	8	0.6	*
14	Parkinson's disease (G20-G21)	6	0.5	*
14	Aortic aneurysm and dissection (I71)	6	0.5	*
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6	0.5	*
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	6	0.5	*
...	All other causes (Residual)	216	16.7	246.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 75-84 years				
...	All causes	1,387	100.0	3,391.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	280	20.2	684.7
2	Malignant neoplasms (C00-C97)	279	20.1	682.3
3	Diabetes mellitus (E10-E14)	120	8.7	293.5
4	Chronic lower respiratory diseases (J40-J47)	101	7.3	247.0
5	Cerebrovascular diseases (I60-I69)	71	5.1	173.6
6	Alzheimer's disease (G30)	58	4.2	141.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	55	4.0	134.5
8	Influenza and pneumonia (J09-J18)	43	3.1	105.2
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	31	2.2	75.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	22	1.6	53.8
11	Septicemia (A40-A41)	21	1.5	51.4
12	Chronic liver disease and cirrhosis (K70,K73-K74)	19	1.4	*
13	Aortic aneurysm and dissection (I71)	12	0.9	*
14	Nutritional deficiencies (E40-E64)	7	0.5	*
14	Parkinson's disease (G20-G21)	7	0.5	*
14	Pneumonitis due to solids and liquids (J69)	7	0.5	*
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	7	0.5	*
...	All other causes (Residual)	247	17.8	604.0
American Indian, female, 85 years and over				
...	All causes	1,299	100.0	9,277.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	366	28.2	2,614.1
2	Malignant neoplasms (C00-C97)	124	9.5	885.7
3	Alzheimer's disease (G30)	99	7.6	707.1
4	Cerebrovascular diseases (I60-I69)	98	7.5	700.0
5	Chronic lower respiratory diseases (J40-J47)	69	5.3	492.8
6	Diabetes mellitus (E10-E14)	52	4.0	371.4
7	Influenza and pneumonia (J09-J18)	48	3.7	342.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	42	3.2	300.0
9	Septicemia (A40-A41)	31	2.4	221.4
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	30	2.3	214.3
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	20	1.5	142.8
12	Pneumonitis due to solids and liquids (J69)	16	1.2	*
13	Parkinson's disease (G20-G21)	9	0.7	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.6	*
15	Atherosclerosis (I70)	7	0.5	*
...	All other causes (Residual)	280	21.6	1,999.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, all ages				
...	All causes	51,162	100.0	301.1
1	Malignant neoplasms (C00-C97)	14,165	27.7	83.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,490	22.5	67.6
3	Cerebrovascular diseases (I60-I69)	3,833	7.5	22.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,144	4.2	12.6
5	Diabetes mellitus (E10-E14)	1,838	3.6	10.8
6	Influenza and pneumonia (J09-J18)	1,539	3.0	9.1
7	Chronic lower respiratory diseases (J40-J47)	1,487	2.9	8.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,091	2.1	6.4
9	Alzheimer's disease (G30)	1,082	2.1	6.4
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,061	2.1	6.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	814	1.6	4.8
12	Septicemia (A40-A41)	582	1.1	3.4
13	Chronic liver disease and cirrhosis (K70,K73-K74)	467	0.9	2.7
13	Certain conditions originating in the perinatal period (P00-P96)	467	0.9	2.7
15	Parkinson's disease (G20-G21)	374	0.7	2.2
...	All other causes (Residual)	8,728	17.1	51.4
Asian or Pacific Islander, both sexes, 1-4 years				
...	All causes	167	100.0	17.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	42	25.1	4.5
2	Malignant neoplasms (C00-C97)	23	13.8	2.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	23	13.8	2.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	8	4.8	*
5	Chronic lower respiratory diseases (J40-J47)	5	3.0	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	3.0	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	2.4	*
7	Cerebrovascular diseases (I60-I69)	4	2.4	*
7	Influenza and pneumonia (J09-J18)	4	2.4	*
10	Septicemia (A40-A41)	2	1.2	*
11	Anemias (D50-D64)	1	0.6	*
11	Meningitis (G00,G03)	1	0.6	*
11	Acute bronchitis and bronchiolitis (J20-J21)	1	0.6	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.6	*
11	Certain conditions originating in the perinatal period (P00-P96)	1	0.6	*
...	All other causes (Residual)	42	25.1	4.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 5-14 years				
...	All causes	185	100.0	8.2
1	Malignant neoplasms (C00-C97)	55	29.7	2.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	29	15.7	1.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	11	5.9	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8	4.3	*
5	Septicemia (A40-A41)	7	3.8	*
5	Chronic lower respiratory diseases (J40-J47)	7	3.8	*
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7	3.8	*
8	Diseases of heart (I00-I09,I11,I13,I20-I51)	6	3.2	*
9	Cerebrovascular diseases (I60-I69)	5	2.7	*
10	Influenza and pneumonia (J09-J18)	3	1.6	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	3	1.6	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	1.1	*
12	Anemias (D50-D64)	2	1.1	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	1.1	*
12	Certain conditions originating in the perinatal period (P00-P96)	2	1.1	*
...	All other causes (Residual)	36	19.5	1.6
Asian or Pacific Islander, both sexes, 15-24 years				
...	All causes	752	100.0	30.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	256	34.0	10.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	181	24.1	7.3
3	Malignant neoplasms (C00-C97)	80	10.6	3.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	63	8.4	2.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	37	4.9	1.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	14	1.9	*
7	Cerebrovascular diseases (I60-I69)	9	1.2	*
8	Influenza and pneumonia (J09-J18)	6	0.8	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.7	*
10	Septicemia (A40-A41)	3	0.4	*
10	Anemias (D50-D64)	3	0.4	*
10	Chronic lower respiratory diseases (J40-J47)	3	0.4	*
13	Tuberculosis (A16-A19)	2	0.3	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.3	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.3	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.3	*
...	All other causes (Residual)	84	11.2	3.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 25-34 years				
...	All causes	1,119	100.0	39.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	269	24.0	9.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	214	19.1	7.5
3	Malignant neoplasms (C00-C97)	195	17.4	6.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	97	8.7	3.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	64	5.7	2.2
6	Pregnancy, childbirth and the puerperium (O00-O99)	26	2.3	0.9
7	Cerebrovascular diseases (I60-I69)	25	2.2	0.9
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	18	1.6	*
9	Septicemia (A40-A41)	11	1.0	*
9	Influenza and pneumonia (J09-J18)	11	1.0	*
9	Chronic liver disease and cirrhosis (K70,K73-K74)	11	1.0	*
12	Chronic lower respiratory diseases (J40-J47)	9	0.8	*
13	Aortic aneurysm and dissection (I71)	7	0.6	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.4	*
15	Tuberculosis (A16-A19)	4	0.4	*
15	Viral hepatitis (B15-B19)	4	0.4	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.4	*
...	All other causes (Residual)	141	12.6	4.9
Asian or Pacific Islander, both sexes, 35-44 years				
...	All causes	1,794	100.0	65.3
1	Malignant neoplasms (C00-C97)	551	30.7	20.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	252	14.0	9.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	189	10.5	6.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	187	10.4	6.8
5	Cerebrovascular diseases (I60-I69)	113	6.3	4.1
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	61	3.4	2.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	44	2.5	1.6
8	Diabetes mellitus (E10-E14)	39	2.2	1.4
9	Influenza and pneumonia (J09-J18)	21	1.2	0.8
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20	1.1	0.7
11	Aortic aneurysm and dissection (I71)	16	0.9	*
12	Septicemia (A40-A41)	15	0.8	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	14	0.8	*
13	Pregnancy, childbirth and the puerperium (O00-O99)	14	0.8	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	12	0.7	*
...	All other causes (Residual)	246	13.7	9.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 45-54 years				
...	All causes	3,862	100.0	168.0
1	Malignant neoplasms (C00-C97)	1,574	40.8	68.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	659	17.1	28.7
3	Cerebrovascular diseases (I60-I69)	255	6.6	11.1
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	242	6.3	10.5
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	187	4.8	8.1
6	Diabetes mellitus (E10-E14)	119	3.1	5.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	97	2.5	4.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	57	1.5	2.5
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	46	1.2	2.0
10	Viral hepatitis (B15-B19)	43	1.1	1.9
11	Septicemia (A40-A41)	37	1.0	1.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	31	0.8	1.3
13	Influenza and pneumonia (J09-J18)	30	0.8	1.3
14	Chronic lower respiratory diseases (J40-J47)	29	0.8	1.3
15	Human immunodeficiency virus (HIV) disease (B20-B24)	28	0.7	1.2
...	All other causes (Residual)	428	11.1	18.6
Asian or Pacific Islander, both sexes, 55-64 years				
...	All causes	6,699	100.0	398.5
1	Malignant neoplasms (C00-C97)	2,890	43.1	171.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,281	19.1	76.2
3	Cerebrovascular diseases (I60-I69)	444	6.6	26.4
4	Diabetes mellitus (E10-E14)	280	4.2	16.7
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	241	3.6	14.3
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	149	2.2	8.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	117	1.7	7.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	107	1.6	6.4
9	Viral hepatitis (B15-B19)	100	1.5	5.9
10	Influenza and pneumonia (J09-J18)	89	1.3	5.3
11	Chronic lower respiratory diseases (J40-J47)	85	1.3	5.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	67	1.0	4.0
13	Septicemia (A40-A41)	65	1.0	3.9
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	41	0.6	2.4
15	Aortic aneurysm and dissection (I71)	30	0.4	1.8
...	All other causes (Residual)	713	10.6	42.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 65-74 years				
...	All causes	8,880	100.0	987.4
1	Malignant neoplasms (C00-C97)	3,366	37.9	374.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,898	21.4	211.0
3	Cerebrovascular diseases (I60-I69)	669	7.5	74.4
4	Diabetes mellitus (E10-E14)	399	4.5	44.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	250	2.8	27.8
6	Chronic lower respiratory diseases (J40-J47)	218	2.5	24.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	210	2.4	23.3
8	Influenza and pneumonia (J09-J18)	181	2.0	20.1
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	104	1.2	11.6
10	Septicemia (A40-A41)	97	1.1	10.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	83	0.9	9.2
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	70	0.8	7.8
13	Viral hepatitis (B15-B19)	69	0.8	7.7
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	69	0.8	7.7
15	Parkinson's disease (G20-G21)	55	0.6	6.1
15	Alzheimer's disease (G30)	55	0.6	6.1
...	All other causes (Residual)	1,087	12.2	120.9
Asian or Pacific Islander, both sexes, 75-84 years				
...	All causes	12,915	100.0	2,853.1
1	Malignant neoplasms (C00-C97)	3,568	27.6	788.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,122	24.2	689.7
3	Cerebrovascular diseases (I60-I69)	1,066	8.3	235.5
4	Chronic lower respiratory diseases (J40-J47)	529	4.1	116.9
5	Diabetes mellitus (E10-E14)	521	4.0	115.1
6	Influenza and pneumonia (J09-J18)	448	3.5	99.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	342	2.6	75.6
8	Alzheimer's disease (G30)	318	2.5	70.2
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	307	2.4	67.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	243	1.9	53.7
11	Septicemia (A40-A41)	164	1.3	36.2
12	Parkinson's disease (G20-G21)	154	1.2	34.0
13	Pneumonitis due to solids and liquids (J69)	105	0.8	23.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	104	0.8	23.0
15	Chronic liver disease and cirrhosis (K70,K73-K74)	82	0.6	18.1
...	All other causes (Residual)	1,842	14.3	406.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 85 years and over				
...	All causes	13,904	100.0	9,418.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,113	29.6	2,786.0
2	Malignant neoplasms (C00-C97)	1,858	13.4	1,258.6
3	Cerebrovascular diseases (I60-I69)	1,237	8.9	837.9
4	Influenza and pneumonia (J09-J18)	734	5.3	497.2
5	Alzheimer's disease (G30)	699	5.0	473.5
6	Chronic lower respiratory diseases (J40-J47)	590	4.2	399.6
7	Diabetes mellitus (E10-E14)	477	3.4	323.1
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	359	2.6	243.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	339	2.4	229.6
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	293	2.1	198.5
11	Septicemia (A40-A41)	173	1.2	117.2
12	Pneumonitis due to solids and liquids (J69)	154	1.1	104.3
13	Parkinson's disease (G20-G21)	149	1.1	100.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	97	0.7	65.7
15	Aortic aneurysm and dissection (I71)	72	0.5	48.8
...	All other causes (Residual)	2,560	18.4	1,734.1
Asian or Pacific Islander, male, all ages				
...	All causes	26,600	100.0	327.0
1	Malignant neoplasms (C00-C97)	7,210	27.1	88.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,262	23.5	77.0
3	Cerebrovascular diseases (I60-I69)	1,747	6.6	21.5
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,337	5.0	16.4
5	Diabetes mellitus (E10-E14)	932	3.5	11.5
6	Chronic lower respiratory diseases (J40-J47)	910	3.4	11.2
7	Influenza and pneumonia (J09-J18)	825	3.1	10.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	756	2.8	9.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	524	2.0	6.4
10	Alzheimer's disease (G30)	345	1.3	4.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	337	1.3	4.1
12	Chronic liver disease and cirrhosis (K70,K73-K74)	302	1.1	3.7
13	Septicemia (A40-A41)	296	1.1	3.6
14	Certain conditions originating in the perinatal period (P00-P96)	278	1.0	3.4
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	218	0.8	2.7
...	All other causes (Residual)	4,321	16.2	53.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 1-4 years				
...	All causes	92	100.0	19.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	26	28.3	5.5
2	Malignant neoplasms (C00-C97)	17	18.5	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	8.7	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	5.4	*
5	Cerebrovascular diseases (I60-I69)	3	3.3	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	2.2	*
6	Influenza and pneumonia (J09-J18)	2	2.2	*
6	Chronic lower respiratory diseases (J40-J47)	2	2.2	*
9	Septicemia (A40-A41)	1	1.1	*
9	Acute bronchitis and bronchiolitis (J20-J21)	1	1.1	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	1.1	*
9	Certain conditions originating in the perinatal period (P00-P96)	1	1.1	*
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1	1.1	*
...	All other causes (Residual)	22	23.9	4.6
Asian or Pacific Islander, male, 5-14 years				
...	All causes	96	100.0	8.4
1	Malignant neoplasms (C00-C97)	27	28.1	2.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19	19.8	*
3	Chronic lower respiratory diseases (J40-J47)	6	6.3	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	6.3	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	5.2	*
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5	5.2	*
7	Septicemia (A40-A41)	3	3.1	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	3.1	*
9	Cerebrovascular diseases (I60-I69)	2	2.1	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	2.1	*
11	Anemias (D50-D64)	1	1.0	*
11	Influenza and pneumonia (J09-J18)	1	1.0	*
...	All other causes (Residual)	16	16.7	*

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 15-24 years				
...	All causes	544	100.0	43.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	196	36.0	15.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	138	25.4	10.9
3	Malignant neoplasms (C00-C97)	52	9.6	4.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	51	9.4	4.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	26	4.8	2.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	1.7	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.7	*
7	Cerebrovascular diseases (I60-I69)	4	0.7	*
7	Influenza and pneumonia (J09-J18)	4	0.7	*
10	Septicemia (A40-A41)	3	0.6	*
10	Chronic lower respiratory diseases (J40-J47)	3	0.6	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.4	*
13	Tuberculosis (A16-A19)	1	0.2	*
13	Anemias (D50-D64)	1	0.2	*
13	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.2	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
13	Legal intervention (Y35,Y89.0)	1	0.2	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
...	All other causes (Residual)	46	8.5	3.6
Asian or Pacific Islander, male, 25-34 years				
...	All causes	713	100.0	52.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	203	28.5	15.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	154	21.6	11.4
3	Malignant neoplasms (C00-C97)	88	12.3	6.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	72	10.1	5.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	48	6.7	3.5
6	Cerebrovascular diseases (I60-I69)	12	1.7	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	11	1.5	*
8	Influenza and pneumonia (J09-J18)	8	1.1	*
9	Chronic liver disease and cirrhosis (K70,K73-K74)	7	1.0	*
10	Aortic aneurysm and dissection (I71)	6	0.8	*
11	Chronic lower respiratory diseases (J40-J47)	5	0.7	*
12	Viral hepatitis (B15-B19)	4	0.6	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.6	*
14	Septicemia (A40-A41)	3	0.4	*
14	Legal intervention (Y35,Y89.0)	3	0.4	*
...	All other causes (Residual)	85	11.9	6.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 35-44 years				
...	All causes	1,084	100.0	83.5
1	Malignant neoplasms (C00-C97)	236	21.8	18.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	206	19.0	15.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	141	13.0	10.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	127	11.7	9.8
5	Cerebrovascular diseases (I60-I69)	67	6.2	5.2
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	40	3.7	3.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	33	3.0	2.5
8	Diabetes mellitus (E10-E14)	20	1.8	1.5
9	Influenza and pneumonia (J09-J18)	13	1.2	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	11	1.0	*
11	Aortic aneurysm and dissection (I71)	10	0.9	*
12	Septicemia (A40-A41)	8	0.7	*
12	Viral hepatitis (B15-B19)	8	0.7	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.6	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	6	0.6	*
...	All other causes (Residual)	151	13.9	11.6
Asian or Pacific Islander, male, 45-54 years				
...	All causes	2,297	100.0	213.7
1	Malignant neoplasms (C00-C97)	724	31.5	67.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	529	23.0	49.2
3	Cerebrovascular diseases (I60-I69)	158	6.9	14.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	157	6.8	14.6
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	129	5.6	12.0
6	Diabetes mellitus (E10-E14)	80	3.5	7.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	72	3.1	6.7
8	Viral hepatitis (B15-B19)	34	1.5	3.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	33	1.4	3.1
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	32	1.4	3.0
11	Influenza and pneumonia (J09-J18)	23	1.0	2.1
12	Human immunodeficiency virus (HIV) disease (B20-B24)	21	0.9	2.0
13	Septicemia (A40-A41)	16	0.7	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	16	0.7	*
13	Aortic aneurysm and dissection (I71)	16	0.7	*
...	All other causes (Residual)	257	11.2	23.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 55-64 years				
...	All causes	3,951	100.0	519.0
1	Malignant neoplasms (C00-C97)	1,486	37.6	195.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	908	23.0	119.3
3	Cerebrovascular diseases (I60-I69)	241	6.1	31.7
4	Diabetes mellitus (E10-E14)	192	4.9	25.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	145	3.7	19.0
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	106	2.7	13.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	90	2.3	11.8
8	Viral hepatitis (B15-B19)	71	1.8	9.3
9	Chronic lower respiratory diseases (J40-J47)	59	1.5	7.8
10	Influenza and pneumonia (J09-J18)	55	1.4	7.2
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	53	1.3	7.0
12	Septicemia (A40-A41)	39	1.0	5.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	39	1.0	5.1
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	25	0.6	3.3
15	Aortic aneurysm and dissection (I71)	17	0.4	*
...	All other causes (Residual)	425	10.8	55.8
Asian or Pacific Islander, male, 65-74 years				
...	All causes	5,009	100.0	1,226.0
1	Malignant neoplasms (C00-C97)	1,823	36.4	446.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,203	24.0	294.4
3	Cerebrovascular diseases (I60-I69)	347	6.9	84.9
4	Diabetes mellitus (E10-E14)	209	4.2	51.2
5	Chronic lower respiratory diseases (J40-J47)	148	3.0	36.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	136	2.7	33.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	116	2.3	28.4
8	Influenza and pneumonia (J09-J18)	114	2.3	27.9
9	Chronic liver disease and cirrhosis (K70,K73-K74)	55	1.1	13.5
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	53	1.1	13.0
11	Septicemia (A40-A41)	51	1.0	12.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	38	0.8	9.3
13	Aortic aneurysm and dissection (I71)	33	0.7	8.1
14	Parkinson's disease (G20-G21)	30	0.6	7.3
15	Viral hepatitis (B15-B19)	28	0.6	6.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	28	0.6	6.9
...	All other causes (Residual)	597	11.9	146.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 75-84 years				
...	All causes	6,387	100.0	3,438.7
1	Malignant neoplasms (C00-C97)	1,821	28.5	980.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,589	24.9	855.5
3	Cerebrovascular diseases (I60-I69)	483	7.6	260.0
4	Chronic lower respiratory diseases (J40-J47)	336	5.3	180.9
5	Influenza and pneumonia (J09-J18)	240	3.8	129.2
6	Diabetes mellitus (E10-E14)	235	3.7	126.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	165	2.6	88.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	158	2.5	85.1
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	108	1.7	58.1
10	Alzheimer's disease (G30)	96	1.5	51.7
11	Septicemia (A40-A41)	88	1.4	47.4
12	Parkinson's disease (G20-G21)	86	1.3	46.3
13	Pneumonitis due to solids and liquids (J69)	58	0.9	31.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	54	0.8	29.1
15	Aortic aneurysm and dissection (I71)	44	0.7	23.7
...	All other causes (Residual)	826	12.9	444.7
Asian or Pacific Islander, male, 85 years and over				
...	All causes	5,922	100.0	10,824.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,714	28.9	3,132.9
2	Malignant neoplasms (C00-C97)	934	15.8	1,707.2
3	Cerebrovascular diseases (I60-I69)	426	7.2	778.7
4	Influenza and pneumonia (J09-J18)	358	6.0	654.4
5	Chronic lower respiratory diseases (J40-J47)	329	5.6	601.4
6	Alzheimer's disease (G30)	220	3.7	402.1
7	Diabetes mellitus (E10-E14)	194	3.3	354.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	139	2.3	254.1
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	139	2.3	254.1
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	128	2.2	234.0
11	Pneumonitis due to solids and liquids (J69)	86	1.5	157.2
12	Septicemia (A40-A41)	83	1.4	151.7
13	Parkinson's disease (G20-G21)	77	1.3	140.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	51	0.9	93.2
15	Aortic aneurysm and dissection (I71)	33	0.6	60.3
...	All other causes (Residual)	1,011	17.1	1,848.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, all ages				
...	All causes	24,562	100.0	277.3
1	Malignant neoplasms (C00-C97)	6,955	28.3	78.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,228	21.3	59.0
3	Cerebrovascular diseases (I60-I69)	2,086	8.5	23.5
4	Diabetes mellitus (E10-E14)	906	3.7	10.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	807	3.3	9.1
6	Alzheimer's disease (G30)	737	3.0	8.3
7	Influenza and pneumonia (J09-J18)	714	2.9	8.1
8	Chronic lower respiratory diseases (J40-J47)	577	2.3	6.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	567	2.3	6.4
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	477	1.9	5.4
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	305	1.2	3.4
12	Septicemia (A40-A41)	286	1.2	3.2
13	Certain conditions originating in the perinatal period (P00-P96)	189	0.8	2.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	178	0.7	2.0
15	Parkinson's disease (G20-G21)	168	0.7	1.9
...	All other causes (Residual)	4,382	17.8	49.5
Asian or Pacific Islander, female, 1-4 years				
...	All causes	75	100.0	16.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	16	21.3	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	15	20.0	*
3	Malignant neoplasms (C00-C97)	6	8.0	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	5.3	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	4.0	*
5	Chronic lower respiratory diseases (J40-J47)	3	4.0	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	2.7	*
7	Influenza and pneumonia (J09-J18)	2	2.7	*
9	Septicemia (A40-A41)	1	1.3	*
9	Anemias (D50-D64)	1	1.3	*
9	Meningitis (G00,G03)	1	1.3	*
9	Cerebrovascular diseases (I60-I69)	1	1.3	*
...	All other causes (Residual)	20	26.7	4.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 5-14 years				
...	All causes	89	100.0	7.9
1	Malignant neoplasms (C00-C97)	28	31.5	2.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10	11.2	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	6.7	*
4	Septicemia (A40-A41)	4	4.5	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	3.4	*
5	Cerebrovascular diseases (I60-I69)	3	3.4	*
5	Complications of medical and surgical care (Y40-Y84,Y88)	3	3.4	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	2.2	*
8	Influenza and pneumonia (J09-J18)	2	2.2	*
8	Certain conditions originating in the perinatal period (P00-P96)	2	2.2	*
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2	2.2	*
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	2.2	*
13	Meningococcal infection (A39)	1	1.1	*
13	Anemias (D50-D64)	1	1.1	*
13	Chronic lower respiratory diseases (J40-J47)	1	1.1	*
...	All other causes (Residual)	19	21.3	*
Asian or Pacific Islander, female, 15-24 years				
...	All causes	208	100.0	17.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	60	28.8	4.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	43	20.7	3.5
3	Malignant neoplasms (C00-C97)	28	13.5	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	12	5.8	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	11	5.3	*
6	Cerebrovascular diseases (I60-I69)	5	2.4	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	2.4	*
8	Anemias (D50-D64)	2	1.0	*
8	Influenza and pneumonia (J09-J18)	2	1.0	*
10	Tuberculosis (A16-A19)	1	0.5	*
10	Scarlet fever and erysipelas (A38,A46)	1	0.5	*
10	Meningococcal infection (A39)	1	0.5	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.5	*
10	Aortic aneurysm and dissection (I71)	1	0.5	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.5	*
10	Pregnancy, childbirth and the puerperium (O00-O99)	1	0.5	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.5	*
...	All other causes (Residual)	32	15.4	2.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 25-34 years				
...	All causes	406	100.0	27.1
1	Malignant neoplasms (C00-C97)	107	26.4	7.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	66	16.3	4.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	60	14.8	4.0
4	Pregnancy, childbirth and the puerperium (O00-O99)	26	6.4	1.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	25	6.2	1.7
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	16	3.9	*
7	Cerebrovascular diseases (I60-I69)	13	3.2	*
8	Septicemia (A40-A41)	8	2.0	*
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	1.7	*
10	Chronic lower respiratory diseases (J40-J47)	4	1.0	*
10	Chronic liver disease and cirrhosis (K70,K73-K74)	4	1.0	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.7	*
12	Influenza and pneumonia (J09-J18)	3	0.7	*
14	Tuberculosis (A16-A19)	2	0.5	*
14	Peptic ulcer (K25-K28)	2	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.5	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.5	*
...	All other causes (Residual)	56	13.8	3.7
Asian or Pacific Islander, female, 35-44 years				
...	All causes	710	100.0	49.0
1	Malignant neoplasms (C00-C97)	315	44.4	21.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	62	8.7	4.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	46	6.5	3.2
3	Cerebrovascular diseases (I60-I69)	46	6.5	3.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	46	6.5	3.2
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	21	3.0	1.4
7	Diabetes mellitus (E10-E14)	19	2.7	*
8	Pregnancy, childbirth and the puerperium (O00-O99)	14	2.0	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	1.8	*
10	Chronic liver disease and cirrhosis (K70,K73-K74)	11	1.5	*
11	Influenza and pneumonia (J09-J18)	8	1.1	*
12	Septicemia (A40-A41)	7	1.0	*
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	1.0	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.8	*
14	Aortic aneurysm and dissection (I71)	6	0.8	*
...	All other causes (Residual)	83	11.7	5.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 45-54 years				
...	All causes	1,565	100.0	127.9
1	Malignant neoplasms (C00-C97)	850	54.3	69.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	130	8.3	10.6
3	Cerebrovascular diseases (I60-I69)	97	6.2	7.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	85	5.4	6.9
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	58	3.7	4.7
6	Diabetes mellitus (E10-E14)	39	2.5	3.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	25	1.6	2.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	24	1.5	2.0
9	Septicemia (A40-A41)	21	1.3	1.7
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	15	1.0	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	0.9	*
11	Chronic lower respiratory diseases (J40-J47)	14	0.9	*
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	14	0.9	*
14	Viral hepatitis (B15-B19)	9	0.6	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	0.6	*
...	All other causes (Residual)	161	10.3	13.2
Asian or Pacific Islander, female, 55-64 years				
...	All causes	2,748	100.0	298.8
1	Malignant neoplasms (C00-C97)	1,404	51.1	152.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	373	13.6	40.6
3	Cerebrovascular diseases (I60-I69)	203	7.4	22.1
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	96	3.5	10.4
5	Diabetes mellitus (E10-E14)	88	3.2	9.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	54	2.0	5.9
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	43	1.6	4.7
8	Influenza and pneumonia (J09-J18)	34	1.2	3.7
9	Viral hepatitis (B15-B19)	29	1.1	3.2
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	28	1.0	3.0
11	Chronic liver disease and cirrhosis (K70,K73-K74)	27	1.0	2.9
12	Septicemia (A40-A41)	26	0.9	2.8
12	Chronic lower respiratory diseases (J40-J47)	26	0.9	2.8
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	16	0.6	*
15	Aortic aneurysm and dissection (I71)	13	0.5	*
...	All other causes (Residual)	288	10.5	31.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 65-74 years				
...	All causes	3,871	100.0	788.7
1	Malignant neoplasms (C00-C97)	1,543	39.9	314.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	695	18.0	141.6
3	Cerebrovascular diseases (I60-I69)	322	8.3	65.6
4	Diabetes mellitus (E10-E14)	190	4.9	38.7
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	114	2.9	23.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	94	2.4	19.2
7	Chronic lower respiratory diseases (J40-J47)	70	1.8	14.3
8	Influenza and pneumonia (J09-J18)	67	1.7	13.7
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	66	1.7	13.4
10	Septicemia (A40-A41)	46	1.2	9.4
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	42	1.1	8.6
12	Viral hepatitis (B15-B19)	41	1.1	8.4
13	Alzheimer's disease (G30)	30	0.8	6.1
14	Chronic liver disease and cirrhosis (K70,K73-K74)	28	0.7	5.7
15	Parkinson's disease (G20-G21)	25	0.6	5.1
...	All other causes (Residual)	498	12.9	101.5
Asian or Pacific Islander, female, 75-84 years				
...	All causes	6,528	100.0	2,445.5
1	Malignant neoplasms (C00-C97)	1,747	26.8	654.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,533	23.5	574.3
3	Cerebrovascular diseases (I60-I69)	583	8.9	218.4
4	Diabetes mellitus (E10-E14)	286	4.4	107.1
5	Alzheimer's disease (G30)	222	3.4	83.2
6	Influenza and pneumonia (J09-J18)	208	3.2	77.9
7	Chronic lower respiratory diseases (J40-J47)	193	3.0	72.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	177	2.7	66.3
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	149	2.3	55.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	135	2.1	50.6
11	Septicemia (A40-A41)	76	1.2	28.5
12	Parkinson's disease (G20-G21)	68	1.0	25.5
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	50	0.8	18.7
14	Chronic liver disease and cirrhosis (K70,K73-K74)	49	0.8	18.4
15	Pneumonitis due to solids and liquids (J69)	47	0.7	17.6
...	All other causes (Residual)	1,005	15.4	376.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 85 years and over				
...	All causes	7,982	100.0	8,590.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,399	30.1	2,581.8
2	Malignant neoplasms (C00-C97)	924	11.6	994.4
3	Cerebrovascular diseases (I60-I69)	811	10.2	872.8
4	Alzheimer's disease (G30)	479	6.0	515.5
5	Influenza and pneumonia (J09-J18)	376	4.7	404.6
6	Diabetes mellitus (E10-E14)	283	3.5	304.6
7	Chronic lower respiratory diseases (J40-J47)	261	3.3	280.9
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	231	2.9	248.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	200	2.5	215.2
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	154	1.9	165.7
11	Septicemia (A40-A41)	90	1.1	96.9
12	Parkinson's disease (G20-G21)	72	0.9	77.5
13	Pneumonitis due to solids and liquids (J69)	68	0.9	73.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	46	0.6	49.5
15	Aortic aneurysm and dissection (I71)	39	0.5	42.0
...	All other causes (Residual)	1,549	19.4	1,667.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable