

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, all ages				
...	All causes	2,596,993	100.0	821.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	611,105	23.5	193.3
2	Malignant neoplasms (C00-C97)	584,881	22.5	185.0
3	Chronic lower respiratory diseases (J40-J47)	149,205	5.7	47.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	130,557	5.0	41.3
5	Cerebrovascular diseases (I60-I69)	128,978	5.0	40.8
6	Alzheimer's disease (G30)	84,767	3.3	26.8
7	Diabetes mellitus (E10-E14)	75,578	2.9	23.9
8	Influenza and pneumonia (J09-J18)	56,979	2.2	18.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	47,112	1.8	14.9
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	41,149	1.6	13.0
11	Septicemia (A40-A41)	38,156	1.5	12.1
12	Chronic liver disease and cirrhosis (K70,K73-K74)	36,427	1.4	11.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	30,770	1.2	9.7
14	Parkinson's disease (G20-G21)	25,196	1.0	8.0
15	Pneumonitis due to solids and liquids (J69)	18,579	0.7	5.9
...	All other causes (Residual)	537,554	20.7	170.0
All races, both sexes, 1-4 years				
...	All causes	4,068	100.0	25.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,316	32.4	8.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	476	11.7	3.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	337	8.3	2.1
4	Malignant neoplasms (C00-C97)	328	8.1	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	169	4.2	1.1
6	Influenza and pneumonia (J09-J18)	102	2.5	0.6
7	Chronic lower respiratory diseases (J40-J47)	64	1.6	0.4
8	Septicemia (A40-A41)	53	1.3	0.3
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	47	1.2	0.3
10	Certain conditions originating in the perinatal period (P00-P96)	45	1.1	0.3
11	Cerebrovascular diseases (I60-I69)	38	0.9	0.2
12	Meningitis (G00,G03)	23	0.6	0.1
13	Anemias (D50-D64)	20	0.5	0.1
14	Complications of medical and surgical care (Y40-Y84,Y88)	15	0.4	*
15	Acute bronchitis and bronchiolitis (J20-J21)	14	0.3	*
...	All other causes (Residual)	1,021	25.1	6.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 5-14 years				
...	All causes	5,340	100.0	13.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,521	28.5	3.7
2	Malignant neoplasms (C00-C97)	895	16.8	2.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	395	7.4	1.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	340	6.4	0.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	277	5.2	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	173	3.2	0.4
7	Chronic lower respiratory diseases (J40-J47)	155	2.9	0.4
8	Influenza and pneumonia (J09-J18)	128	2.4	0.3
9	Cerebrovascular diseases (I60-I69)	89	1.7	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	65	1.2	0.2
11	Septicemia (A40-A41)	55	1.0	0.1
12	Certain conditions originating in the perinatal period (P00-P96)	33	0.6	0.1
13	Complications of medical and surgical care (Y40-Y84,Y88)	26	0.5	0.1
14	Diabetes mellitus (E10-E14)	24	0.4	0.1
15	Anemias (D50-D64)	17	0.3	*
...	All other causes (Residual)	1,147	21.5	2.8
All races, both sexes, 15-24 years				
...	All causes	28,486	100.0	64.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,619	40.8	26.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,878	17.1	11.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4,329	15.2	9.8
4	Malignant neoplasms (C00-C97)	1,496	5.3	3.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	941	3.3	2.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	362	1.3	0.8
7	Influenza and pneumonia (J09-J18)	197	0.7	0.4
8	Diabetes mellitus (E10-E14)	193	0.7	0.4
9	Pregnancy, childbirth and the puerperium (O00-O99)	178	0.6	0.4
10	Chronic lower respiratory diseases (J40-J47)	155	0.5	0.4
11	Cerebrovascular diseases (I60-I69)	153	0.5	0.3
12	Septicemia (A40-A41)	138	0.5	0.3
13	Human immunodeficiency virus (HIV) disease (B20-B24)	109	0.4	0.2
14	Legal intervention (Y35,Y89.0)	99	0.3	0.2
15	Anemias (D50-D64)	93	0.3	0.2
...	All other causes (Residual)	3,546	12.4	8.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 25-34 years				
...	All causes	45,463	100.0	106.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	16,209	35.7	37.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,348	14.0	14.8
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4,236	9.3	9.9
4	Malignant neoplasms (C00-C97)	3,673	8.1	8.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,258	7.2	7.6
6	Diabetes mellitus (E10-E14)	684	1.5	1.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	676	1.5	1.6
8	Human immunodeficiency virus (HIV) disease (B20-B24)	631	1.4	1.5
9	Cerebrovascular diseases (I60-I69)	508	1.1	1.2
10	Influenza and pneumonia (J09-J18)	449	1.0	1.0
11	Pregnancy, childbirth and the puerperium (O00-O99)	441	1.0	1.0
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	423	0.9	1.0
13	Septicemia (A40-A41)	333	0.7	0.8
14	Chronic lower respiratory diseases (J40-J47)	291	0.6	0.7
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	266	0.6	0.6
...	All other causes (Residual)	7,037	15.5	16.4
All races, both sexes, 35-44 years				
...	All causes	69,573	100.0	172.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15,354	22.1	38.0
2	Malignant neoplasms (C00-C97)	11,349	16.3	28.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,341	14.9	25.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,551	9.4	16.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,581	3.7	6.4
6	Chronic liver disease and cirrhosis (K70,K73-K74)	2,491	3.6	6.2
7	Diabetes mellitus (E10-E14)	1,952	2.8	4.8
8	Cerebrovascular diseases (I60-I69)	1,687	2.4	4.2
9	Human immunodeficiency virus (HIV) disease (B20-B24)	1,246	1.8	3.1
10	Influenza and pneumonia (J09-J18)	881	1.3	2.2
11	Septicemia (A40-A41)	820	1.2	2.0
12	Chronic lower respiratory diseases (J40-J47)	760	1.1	1.9
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	627	0.9	1.5
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	441	0.6	1.1
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	409	0.6	1.0
...	All other causes (Residual)	12,083	17.4	29.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 45-54 years				
...	All causes	177,724	100.0	406.1
1	Malignant neoplasms (C00-C97)	46,185	26.0	105.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	35,167	19.8	80.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	20,357	11.5	46.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	8,785	4.9	20.1
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8,621	4.9	19.7
6	Diabetes mellitus (E10-E14)	5,899	3.3	13.5
7	Cerebrovascular diseases (I60-I69)	5,425	3.1	12.4
8	Chronic lower respiratory diseases (J40-J47)	4,619	2.6	10.6
9	Septicemia (A40-A41)	2,445	1.4	5.6
10	Human immunodeficiency virus (HIV) disease (B20-B24)	2,378	1.3	5.4
11	Influenza and pneumonia (J09-J18)	2,233	1.3	5.1
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,025	1.1	4.6
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,989	1.1	4.5
14	Viral hepatitis (B15-B19)	1,911	1.1	4.4
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,513	0.9	3.5
...	All other causes (Residual)	28,172	15.9	64.4
All races, both sexes, 55-64 years				
...	All causes	338,127	100.0	860.0
1	Malignant neoplasms (C00-C97)	113,324	33.5	288.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	72,568	21.5	184.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17,057	5.0	43.4
4	Chronic lower respiratory diseases (J40-J47)	15,942	4.7	40.5
5	Diabetes mellitus (E10-E14)	13,061	3.9	33.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	11,951	3.5	30.4
7	Cerebrovascular diseases (I60-I69)	11,364	3.4	28.9
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7,135	2.1	18.1
9	Septicemia (A40-A41)	5,345	1.6	13.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,947	1.5	12.6
11	Influenza and pneumonia (J09-J18)	4,779	1.4	12.2
12	Viral hepatitis (B15-B19)	3,936	1.2	10.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,137	0.9	8.0
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1,820	0.5	4.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,502	0.4	3.8
...	All other causes (Residual)	50,259	14.9	127.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 65-74 years				
...	All causes	454,429	100.0	1,802.1
1	Malignant neoplasms (C00-C97)	155,552	34.2	616.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	98,432	21.7	390.3
3	Chronic lower respiratory diseases (J40-J47)	35,603	7.8	141.2
4	Cerebrovascular diseases (I60-I69)	18,722	4.1	74.2
5	Diabetes mellitus (E10-E14)	17,279	3.8	68.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,967	2.4	43.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,524	1.9	33.8
8	Septicemia (A40-A41)	7,693	1.7	30.5
9	Influenza and pneumonia (J09-J18)	7,441	1.6	29.5
10	Chronic liver disease and cirrhosis (K70,K73-K74)	7,087	1.6	28.1
11	Alzheimer's disease (G30)	4,561	1.0	18.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,357	1.0	17.3
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,794	0.8	15.0
14	Parkinson's disease (G20-G21)	3,215	0.7	12.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,773	0.6	11.0
...	All other causes (Residual)	68,429	15.1	271.4
All races, both sexes, 75-84 years				
...	All causes	625,013	100.0	4,648.1
1	Malignant neoplasms (C00-C97)	153,214	24.5	1,139.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	147,255	23.6	1,095.1
3	Chronic lower respiratory diseases (J40-J47)	49,346	7.9	367.0
4	Cerebrovascular diseases (I60-I69)	36,151	5.8	268.9
5	Alzheimer's disease (G30)	23,073	3.7	171.6
6	Diabetes mellitus (E10-E14)	19,587	3.1	145.7
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14,438	2.3	107.4
8	Influenza and pneumonia (J09-J18)	13,949	2.2	103.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13,317	2.1	99.0
10	Parkinson's disease (G20-G21)	10,549	1.7	78.5
11	Septicemia (A40-A41)	10,274	1.6	76.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7,216	1.2	53.7
13	Pneumonitis due to solids and liquids (J69)	4,933	0.8	36.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,554	0.7	33.9
15	Chronic liver disease and cirrhosis (K70,K73-K74)	4,014	0.6	29.9
...	All other causes (Residual)	113,143	18.1	841.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 85 years and over				
...	All causes	825,198	100.0	13,660.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	242,469	29.4	4,013.9
2	Malignant neoplasms (C00-C97)	98,792	12.0	1,635.4
3	Alzheimer's disease (G30)	56,152	6.8	929.5
4	Cerebrovascular diseases (I60-I69)	54,729	6.6	906.0
5	Chronic lower respiratory diseases (J40-J47)	42,245	5.1	699.3
6	Influenza and pneumonia (J09-J18)	26,641	3.2	441.0
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	20,537	2.5	340.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17,239	2.1	285.4
9	Diabetes mellitus (E10-E14)	16,885	2.0	279.5
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	13,991	1.7	231.6
11	Septicemia (A40-A41)	10,848	1.3	179.6
12	Parkinson's disease (G20-G21)	10,762	1.3	178.2
13	Pneumonitis due to solids and liquids (J69)	9,218	1.1	152.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5,106	0.6	84.5
15	Atherosclerosis (I70)	3,803	0.5	63.0
...	All other causes (Residual)	195,781	23.7	3,241.0
All races, male, all ages				
...	All causes	1,306,034	100.0	839.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	321,347	24.6	206.5
2	Malignant neoplasms (C00-C97)	307,559	23.5	197.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	81,916	6.3	52.6
4	Chronic lower respiratory diseases (J40-J47)	70,317	5.4	45.2
5	Cerebrovascular diseases (I60-I69)	53,691	4.1	34.5
6	Diabetes mellitus (E10-E14)	39,841	3.1	25.6
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	32,055	2.5	20.6
8	Influenza and pneumonia (J09-J18)	26,804	2.1	17.2
9	Alzheimer's disease (G30)	25,836	2.0	16.6
10	Chronic liver disease and cirrhosis (K70,K73-K74)	23,709	1.8	15.2
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23,493	1.8	15.1
12	Septicemia (A40-A41)	17,994	1.4	11.6
13	Parkinson's disease (G20-G21)	15,088	1.2	9.7
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	12,963	1.0	8.3
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	12,726	1.0	8.2
...	All other causes (Residual)	240,695	18.4	154.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 1-4 years				
...	All causes	2,323	100.0	28.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	831	35.8	10.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	232	10.0	2.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	184	7.9	2.3
4	Malignant neoplasms (C00-C97)	183	7.9	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	96	4.1	1.2
6	Influenza and pneumonia (J09-J18)	53	2.3	0.7
7	Chronic lower respiratory diseases (J40-J47)	38	1.6	0.5
8	Septicemia (A40-A41)	32	1.4	0.4
9	Certain conditions originating in the perinatal period (P00-P96)	29	1.2	0.4
10	Cerebrovascular diseases (I60-I69)	22	0.9	0.3
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	0.8	*
12	Meningitis (G00,G03)	13	0.6	*
13	Anemias (D50-D64)	11	0.5	*
14	Acute bronchitis and bronchiolitis (J20-J21)	7	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.3	*
...	All other causes (Residual)	566	24.4	7.0
All races, male, 5-14 years				
...	All causes	3,077	100.0	14.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	950	30.9	4.5
2	Malignant neoplasms (C00-C97)	471	15.3	2.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	252	8.2	1.2
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	176	5.7	0.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	168	5.5	0.8
6	Chronic lower respiratory diseases (J40-J47)	100	3.2	0.5
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	93	3.0	0.4
8	Influenza and pneumonia (J09-J18)	70	2.3	0.3
9	Cerebrovascular diseases (I60-I69)	44	1.4	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	40	1.3	0.2
11	Septicemia (A40-A41)	32	1.0	0.2
12	Certain conditions originating in the perinatal period (P00-P96)	17	0.6	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	15	0.5	*
14	Diabetes mellitus (E10-E14)	12	0.4	*
15	Meningitis (G00,G03)	11	0.4	*
...	All other causes (Residual)	626	20.3	3.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 15-24 years				
...	All causes	20,864	100.0	92.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,545	41.0	37.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,903	18.7	17.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,765	18.0	16.7
4	Malignant neoplasms (C00-C97)	859	4.1	3.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	628	3.0	2.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	223	1.1	1.0
7	Diabetes mellitus (E10-E14)	120	0.6	0.5
8	Influenza and pneumonia (J09-J18)	114	0.5	0.5
9	Chronic lower respiratory diseases (J40-J47)	101	0.5	0.4
10	Legal intervention (Y35,Y89.0)	95	0.5	0.4
11	Cerebrovascular diseases (I60-I69)	85	0.4	0.4
12	Human immunodeficiency virus (HIV) disease (B20-B24)	81	0.4	0.4
13	Septicemia (A40-A41)	74	0.4	0.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	57	0.3	0.3
15	Anemias (D50-D64)	54	0.3	0.2
...	All other causes (Residual)	2,160	10.4	9.6
All races, male, 25-34 years				
...	All causes	31,462	100.0	145.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,025	38.2	55.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,063	16.1	23.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,555	11.3	16.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,214	7.0	10.2
5	Malignant neoplasms (C00-C97)	1,857	5.9	8.6
6	Human immunodeficiency virus (HIV) disease (B20-B24)	457	1.5	2.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	435	1.4	2.0
8	Diabetes mellitus (E10-E14)	403	1.3	1.9
9	Cerebrovascular diseases (I60-I69)	279	0.9	1.3
10	Influenza and pneumonia (J09-J18)	253	0.8	1.2
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	240	0.8	1.1
12	Septicemia (A40-A41)	178	0.6	0.8
13	Chronic lower respiratory diseases (J40-J47)	160	0.5	0.7
14	Legal intervention (Y35,Y89.0)	141	0.4	0.7
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	140	0.4	0.6
...	All other causes (Residual)	4,062	12.9	18.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 35-44 years				
...	All causes	43,072	100.0	213.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,566	24.5	52.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,152	16.6	35.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,003	11.6	24.8
4	Malignant neoplasms (C00-C97)	4,827	11.2	24.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,025	4.7	10.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1,620	3.8	8.0
7	Diabetes mellitus (E10-E14)	1,211	2.8	6.0
8	Cerebrovascular diseases (I60-I69)	938	2.2	4.7
9	Human immunodeficiency virus (HIV) disease (B20-B24)	842	2.0	4.2
10	Influenza and pneumonia (J09-J18)	464	1.1	2.3
11	Septicemia (A40-A41)	397	0.9	2.0
12	Chronic lower respiratory diseases (J40-J47)	365	0.8	1.8
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	342	0.8	1.7
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	264	0.6	1.3
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	251	0.6	1.2
...	All other causes (Residual)	6,805	15.8	33.8
All races, male, 45-54 years				
...	All causes	107,997	100.0	500.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	24,830	23.0	115.1
2	Malignant neoplasms (C00-C97)	22,966	21.3	106.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,591	12.6	63.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,393	5.9	29.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	5,858	5.4	27.2
6	Diabetes mellitus (E10-E14)	3,633	3.4	16.8
7	Cerebrovascular diseases (I60-I69)	3,067	2.8	14.2
8	Chronic lower respiratory diseases (J40-J47)	2,142	2.0	9.9
9	Human immunodeficiency virus (HIV) disease (B20-B24)	1,760	1.6	8.2
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,479	1.4	6.9
11	Viral hepatitis (B15-B19)	1,328	1.2	6.2
12	Septicemia (A40-A41)	1,304	1.2	6.0
13	Influenza and pneumonia (J09-J18)	1,277	1.2	5.9
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,190	1.1	5.5
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	990	0.9	4.6
...	All other causes (Residual)	16,189	15.0	75.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 55-64 years				
...	All causes	206,325	100.0	1,088.4
1	Malignant neoplasms (C00-C97)	62,810	30.4	331.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	50,680	24.6	267.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,658	5.7	61.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	8,485	4.1	44.8
5	Chronic lower respiratory diseases (J40-J47)	8,287	4.0	43.7
6	Diabetes mellitus (E10-E14)	7,933	3.8	41.8
7	Cerebrovascular diseases (I60-I69)	6,651	3.2	35.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,373	2.6	28.3
9	Septicemia (A40-A41)	2,892	1.4	15.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,800	1.4	14.8
11	Viral hepatitis (B15-B19)	2,792	1.4	14.7
12	Influenza and pneumonia (J09-J18)	2,745	1.3	14.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,990	1.0	10.5
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1,369	0.7	7.2
15	Aortic aneurysm and dissection (I71)	934	0.5	4.9
...	All other causes (Residual)	28,926	14.0	152.6
All races, male, 65-74 years				
...	All causes	257,898	100.0	2,186.0
1	Malignant neoplasms (C00-C97)	85,669	33.2	726.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	62,634	24.3	530.9
3	Chronic lower respiratory diseases (J40-J47)	17,925	7.0	151.9
4	Cerebrovascular diseases (I60-I69)	10,033	3.9	85.0
5	Diabetes mellitus (E10-E14)	9,857	3.8	83.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,014	2.7	59.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	4,531	1.8	38.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,469	1.7	37.9
9	Influenza and pneumonia (J09-J18)	4,131	1.6	35.0
10	Septicemia (A40-A41)	4,002	1.6	33.9
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,072	1.2	26.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,359	0.9	20.0
13	Parkinson's disease (G20-G21)	2,160	0.8	18.3
14	Alzheimer's disease (G30)	1,984	0.8	16.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,657	0.6	14.0
...	All other causes (Residual)	36,401	14.1	308.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 75-84 years				
...	All causes	315,340	100.0	5,474.2
1	Malignant neoplasms (C00-C97)	81,480	25.8	1,414.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	79,634	25.3	1,382.4
3	Chronic lower respiratory diseases (J40-J47)	23,809	7.6	413.3
4	Cerebrovascular diseases (I60-I69)	16,006	5.1	277.9
5	Diabetes mellitus (E10-E14)	9,952	3.2	172.8
6	Alzheimer's disease (G30)	8,789	2.8	152.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,829	2.5	135.9
8	Influenza and pneumonia (J09-J18)	7,097	2.3	123.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,861	2.2	119.1
10	Parkinson's disease (G20-G21)	6,717	2.1	116.6
11	Septicemia (A40-A41)	4,830	1.5	83.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,152	1.0	54.7
13	Pneumonitis due to solids and liquids (J69)	2,856	0.9	49.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,535	0.8	44.0
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,132	0.7	37.0
...	All other causes (Residual)	51,661	16.4	896.8
All races, male, 85 years and over				
...	All causes	304,462	100.0	14,911.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	93,192	30.6	4,564.2
2	Malignant neoplasms (C00-C97)	46,401	15.2	2,272.6
3	Chronic lower respiratory diseases (J40-J47)	17,375	5.7	851.0
4	Cerebrovascular diseases (I60-I69)	16,505	5.4	808.4
5	Alzheimer's disease (G30)	14,647	4.8	717.4
6	Influenza and pneumonia (J09-J18)	10,489	3.4	513.7
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,226	2.7	402.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,595	2.5	372.0
9	Diabetes mellitus (E10-E14)	6,711	2.2	328.7
10	Parkinson's disease (G20-G21)	5,763	1.9	282.3
11	Pneumonitis due to solids and liquids (J69)	4,608	1.5	225.7
12	Septicemia (A40-A41)	4,165	1.4	204.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,114	1.4	201.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,313	0.8	113.3
15	Aortic aneurysm and dissection (I71)	1,217	0.4	59.6
...	All other causes (Residual)	61,141	20.1	2,994.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, all ages				
...	All causes	1,290,959	100.0	804.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	289,758	22.4	180.6
2	Malignant neoplasms (C00-C97)	277,322	21.5	172.8
3	Chronic lower respiratory diseases (J40-J47)	78,888	6.1	49.2
4	Cerebrovascular diseases (I60-I69)	75,287	5.8	46.9
5	Alzheimer's disease (G30)	58,931	4.6	36.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	48,641	3.8	30.3
7	Diabetes mellitus (E10-E14)	35,737	2.8	22.3
8	Influenza and pneumonia (J09-J18)	30,175	2.3	18.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23,619	1.8	14.7
10	Septicemia (A40-A41)	20,162	1.6	12.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	17,807	1.4	11.1
12	Chronic liver disease and cirrhosis (K70,K73-K74)	12,718	1.0	7.9
13	Parkinson's disease (G20-G21)	10,108	0.8	6.3
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	9,094	0.7	5.7
15	Pneumonitis due to solids and liquids (J69)	8,439	0.7	5.3
...	All other causes (Residual)	294,273	22.8	183.4
All races, female, 1-4 years				
...	All causes	1,745	100.0	22.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	485	27.8	6.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	244	14.0	3.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	153	8.8	2.0
4	Malignant neoplasms (C00-C97)	145	8.3	1.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	73	4.2	0.9
6	Influenza and pneumonia (J09-J18)	49	2.8	0.6
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	28	1.6	0.4
8	Chronic lower respiratory diseases (J40-J47)	26	1.5	0.3
9	Septicemia (A40-A41)	21	1.2	0.3
10	Cerebrovascular diseases (I60-I69)	16	0.9	*
10	Certain conditions originating in the perinatal period (P00-P96)	16	0.9	*
12	Meningitis (G00,G03)	10	0.6	*
13	Anemias (D50-D64)	9	0.5	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	8	0.5	*
15	Acute bronchitis and bronchiolitis (J20-J21)	7	0.4	*
...	All other causes (Residual)	455	26.1	5.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 5-14 years				
...	All causes	2,263	100.0	11.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	571	25.2	2.8
2	Malignant neoplasms (C00-C97)	424	18.7	2.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	164	7.2	0.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	143	6.3	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	109	4.8	0.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	80	3.5	0.4
7	Influenza and pneumonia (J09-J18)	58	2.6	0.3
8	Chronic lower respiratory diseases (J40-J47)	55	2.4	0.3
9	Cerebrovascular diseases (I60-I69)	45	2.0	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	25	1.1	0.1
11	Septicemia (A40-A41)	23	1.0	0.1
12	Certain conditions originating in the perinatal period (P00-P96)	16	0.7	*
13	Diabetes mellitus (E10-E14)	12	0.5	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	11	0.5	*
15	Anemias (D50-D64)	8	0.4	*
...	All other causes (Residual)	519	22.9	2.6
All races, female, 15-24 years				
...	All causes	7,622	100.0	35.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,074	40.3	14.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	975	12.8	4.5
3	Malignant neoplasms (C00-C97)	637	8.4	3.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	564	7.4	2.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	313	4.1	1.5
6	Pregnancy, childbirth and the puerperium (O00-O99)	178	2.3	0.8
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	139	1.8	0.6
8	Influenza and pneumonia (J09-J18)	83	1.1	0.4
9	Diabetes mellitus (E10-E14)	73	1.0	0.3
10	Cerebrovascular diseases (I60-I69)	68	0.9	0.3
11	Septicemia (A40-A41)	64	0.8	0.3
12	Chronic lower respiratory diseases (J40-J47)	54	0.7	0.3
13	Anemias (D50-D64)	39	0.5	0.2
14	Human immunodeficiency virus (HIV) disease (B20-B24)	28	0.4	0.1
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	26	0.3	0.1
...	All other causes (Residual)	1,307	17.1	6.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 25-34 years				
...	All causes	14,001	100.0	66.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,184	29.9	19.7
2	Malignant neoplasms (C00-C97)	1,816	13.0	8.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,285	9.2	6.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,044	7.5	4.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	681	4.9	3.2
6	Pregnancy, childbirth and the puerperium (O00-O99)	441	3.1	2.1
7	Diabetes mellitus (E10-E14)	281	2.0	1.3
8	Chronic liver disease and cirrhosis (K70,K73-K74)	241	1.7	1.1
9	Cerebrovascular diseases (I60-I69)	229	1.6	1.1
10	Influenza and pneumonia (J09-J18)	196	1.4	0.9
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	183	1.3	0.9
12	Human immunodeficiency virus (HIV) disease (B20-B24)	174	1.2	0.8
13	Septicemia (A40-A41)	155	1.1	0.7
14	Chronic lower respiratory diseases (J40-J47)	131	0.9	0.6
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	126	0.9	0.6
...	All other causes (Residual)	2,834	20.2	13.4
All races, female, 35-44 years				
...	All causes	26,501	100.0	130.5
1	Malignant neoplasms (C00-C97)	6,522	24.6	32.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,788	18.1	23.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,189	12.0	15.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,548	5.8	7.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	871	3.3	4.3
6	Cerebrovascular diseases (I60-I69)	749	2.8	3.7
7	Diabetes mellitus (E10-E14)	741	2.8	3.6
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	556	2.1	2.7
9	Septicemia (A40-A41)	423	1.6	2.1
10	Influenza and pneumonia (J09-J18)	417	1.6	2.1
11	Human immunodeficiency virus (HIV) disease (B20-B24)	404	1.5	2.0
12	Chronic lower respiratory diseases (J40-J47)	395	1.5	1.9
13	Pregnancy, childbirth and the puerperium (O00-O99)	301	1.1	1.5
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	285	1.1	1.4
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	190	0.7	0.9
...	All other causes (Residual)	5,122	19.3	25.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 45-54 years				
...	All causes	69,727	100.0	314.1
1	Malignant neoplasms (C00-C97)	23,219	33.3	104.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,337	14.8	46.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,766	9.7	30.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	2,927	4.2	13.2
5	Chronic lower respiratory diseases (J40-J47)	2,477	3.6	11.2
6	Cerebrovascular diseases (I60-I69)	2,358	3.4	10.6
7	Diabetes mellitus (E10-E14)	2,266	3.2	10.2
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,228	3.2	10.0
9	Septicemia (A40-A41)	1,141	1.6	5.1
10	Influenza and pneumonia (J09-J18)	956	1.4	4.3
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	835	1.2	3.8
12	Human immunodeficiency virus (HIV) disease (B20-B24)	618	0.9	2.8
13	Viral hepatitis (B15-B19)	583	0.8	2.6
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	523	0.8	2.4
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	510	0.7	2.3
...	All other causes (Residual)	11,983	17.2	54.0
All races, female, 55-64 years				
...	All causes	131,802	100.0	647.4
1	Malignant neoplasms (C00-C97)	50,514	38.3	248.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	21,888	16.6	107.5
3	Chronic lower respiratory diseases (J40-J47)	7,655	5.8	37.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,399	4.1	26.5
5	Diabetes mellitus (E10-E14)	5,128	3.9	25.2
6	Cerebrovascular diseases (I60-I69)	4,713	3.6	23.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	3,466	2.6	17.0
8	Septicemia (A40-A41)	2,453	1.9	12.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,147	1.6	10.5
10	Influenza and pneumonia (J09-J18)	2,034	1.5	10.0
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,762	1.3	8.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,147	0.9	5.6
13	Viral hepatitis (B15-B19)	1,144	0.9	5.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	641	0.5	3.1
15	Alzheimer's disease (G30)	509	0.4	2.5
...	All other causes (Residual)	21,202	16.1	104.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 65-74 years				
...	All causes	196,531	100.0	1,464.6
1	Malignant neoplasms (C00-C97)	69,883	35.6	520.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	35,798	18.2	266.8
3	Chronic lower respiratory diseases (J40-J47)	17,678	9.0	131.7
4	Cerebrovascular diseases (I60-I69)	8,689	4.4	64.8
5	Diabetes mellitus (E10-E14)	7,422	3.8	55.3
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,055	2.1	30.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,953	2.0	29.5
8	Septicemia (A40-A41)	3,691	1.9	27.5
9	Influenza and pneumonia (J09-J18)	3,310	1.7	24.7
10	Alzheimer's disease (G30)	2,577	1.3	19.2
11	Chronic liver disease and cirrhosis (K70,K73-K74)	2,556	1.3	19.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,998	1.0	14.9
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,116	0.6	8.3
14	Parkinson's disease (G20-G21)	1,055	0.5	7.9
15	Pneumonitis due to solids and liquids (J69)	883	0.4	6.6
...	All other causes (Residual)	31,867	16.2	237.5
All races, female, 75-84 years				
...	All causes	309,673	100.0	4,029.1
1	Malignant neoplasms (C00-C97)	71,734	23.2	933.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	67,621	21.8	879.8
3	Chronic lower respiratory diseases (J40-J47)	25,537	8.2	332.3
4	Cerebrovascular diseases (I60-I69)	20,145	6.5	262.1
5	Alzheimer's disease (G30)	14,284	4.6	185.8
6	Diabetes mellitus (E10-E14)	9,635	3.1	125.4
7	Influenza and pneumonia (J09-J18)	6,852	2.2	89.1
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,609	2.1	86.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,456	2.1	84.0
10	Septicemia (A40-A41)	5,444	1.8	70.8
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,064	1.3	52.9
12	Parkinson's disease (G20-G21)	3,832	1.2	49.9
13	Pneumonitis due to solids and liquids (J69)	2,077	0.7	27.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,019	0.7	26.3
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,882	0.6	24.5
...	All other causes (Residual)	61,482	19.9	799.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 85 years and over				
...	All causes	520,736	100.0	13,021.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	149,277	28.7	3,732.9
2	Malignant neoplasms (C00-C97)	52,391	10.1	1,310.1
3	Alzheimer's disease (G30)	41,505	8.0	1,037.9
4	Cerebrovascular diseases (I60-I69)	38,224	7.3	955.8
5	Chronic lower respiratory diseases (J40-J47)	24,870	4.8	621.9
6	Influenza and pneumonia (J09-J18)	16,152	3.1	403.9
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,311	2.4	307.9
8	Diabetes mellitus (E10-E14)	10,174	2.0	254.4
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9,877	1.9	247.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9,644	1.9	241.2
11	Septicemia (A40-A41)	6,683	1.3	167.1
12	Parkinson's disease (G20-G21)	4,999	1.0	125.0
13	Pneumonitis due to solids and liquids (J69)	4,610	0.9	115.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,793	0.5	69.8
15	Atherosclerosis (I70)	2,612	0.5	65.3
...	All other causes (Residual)	134,614	25.9	3,366.2
White, both sexes, all ages				
...	All causes	2,217,103	100.0	889.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	522,645	23.6	209.6
2	Malignant neoplasms (C00-C97)	498,116	22.5	199.8
3	Chronic lower respiratory diseases (J40-J47)	136,682	6.2	54.8
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	112,803	5.1	45.2
5	Cerebrovascular diseases (I60-I69)	107,909	4.9	43.3
6	Alzheimer's disease (G30)	77,387	3.5	31.0
7	Diabetes mellitus (E10-E14)	58,925	2.7	23.6
8	Influenza and pneumonia (J09-J18)	49,013	2.2	19.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	37,270	1.7	14.9
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	37,154	1.7	14.9
11	Chronic liver disease and cirrhosis (K70,K73-K74)	31,871	1.4	12.8
12	Septicemia (A40-A41)	30,889	1.4	12.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	24,149	1.1	9.7
14	Parkinson's disease (G20-G21)	23,553	1.1	9.4
15	Pneumonitis due to solids and liquids (J69)	16,458	0.7	6.6
...	All other causes (Residual)	452,279	20.4	181.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 1-4 years				
...	All causes	2,774	100.0	23.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	929	33.5	7.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	345	12.4	2.9
3	Malignant neoplasms (C00-C97)	246	8.9	2.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	179	6.5	1.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	103	3.7	0.9
6	Influenza and pneumonia (J09-J18)	70	2.5	0.6
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	37	1.3	0.3
8	Chronic lower respiratory diseases (J40-J47)	32	1.2	0.3
9	Certain conditions originating in the perinatal period (P00-P96)	30	1.1	0.3
10	Septicemia (A40-A41)	29	1.0	0.2
11	Cerebrovascular diseases (I60-I69)	20	0.7	0.2
12	Meningitis (G00,G03)	13	0.5	*
13	Anemias (D50-D64)	11	0.4	*
13	Acute bronchitis and bronchiolitis (J20-J21)	11	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	10	0.4	*
...	All other causes (Residual)	709	25.6	5.9
White, both sexes, 5-14 years				
...	All causes	3,841	100.0	12.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,105	28.8	3.5
2	Malignant neoplasms (C00-C97)	717	18.7	2.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	312	8.1	1.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	242	6.3	0.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	170	4.4	0.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	105	2.7	0.3
7	Influenza and pneumonia (J09-J18)	92	2.4	0.3
8	Cerebrovascular diseases (I60-I69)	66	1.7	0.2
9	Chronic lower respiratory diseases (J40-J47)	59	1.5	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	45	1.2	0.1
11	Septicemia (A40-A41)	38	1.0	0.1
12	Certain conditions originating in the perinatal period (P00-P96)	24	0.6	0.1
13	Diabetes mellitus (E10-E14)	20	0.5	0.1
14	Complications of medical and surgical care (Y40-Y84,Y88)	17	0.4	*
15	Meningitis (G00,G03)	9	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.2	*
...	All other causes (Residual)	811	21.1	2.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 15-24 years				
...	All causes	20,618	100.0	62.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,718	47.1	29.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,003	19.4	12.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,503	7.3	4.5
4	Malignant neoplasms (C00-C97)	1,138	5.5	3.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	602	2.9	1.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	281	1.4	0.8
7	Influenza and pneumonia (J09-J18)	159	0.8	0.5
8	Diabetes mellitus (E10-E14)	113	0.5	0.3
9	Cerebrovascular diseases (I60-I69)	107	0.5	0.3
10	Pregnancy, childbirth and the puerperium (O00-O99)	103	0.5	0.3
11	Septicemia (A40-A41)	92	0.4	0.3
12	Chronic lower respiratory diseases (J40-J47)	68	0.3	0.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	63	0.3	0.2
14	Legal intervention (Y35,Y89.0)	49	0.2	0.1
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	41	0.2	0.1
...	All other causes (Residual)	2,578	12.5	7.8
White, both sexes, 25-34 years				
...	All causes	33,777	100.0	103.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,758	40.7	42.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,394	16.0	16.5
3	Malignant neoplasms (C00-C97)	2,783	8.2	8.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,144	6.3	6.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,612	4.8	4.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	539	1.6	1.6
7	Diabetes mellitus (E10-E14)	429	1.3	1.3
8	Cerebrovascular diseases (I60-I69)	338	1.0	1.0
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	336	1.0	1.0
10	Influenza and pneumonia (J09-J18)	335	1.0	1.0
11	Pregnancy, childbirth and the puerperium (O00-O99)	262	0.8	0.8
12	Septicemia (A40-A41)	225	0.7	0.7
13	Human immunodeficiency virus (HIV) disease (B20-B24)	216	0.6	0.7
14	Chronic lower respiratory diseases (J40-J47)	184	0.5	0.6
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	153	0.5	0.5
...	All other causes (Residual)	5,069	15.0	15.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 35-44 years				
...	All causes	52,530	100.0	167.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,023	24.8	41.6
2	Malignant neoplasms (C00-C97)	8,704	16.6	27.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,110	13.5	22.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,824	11.1	18.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,074	3.9	6.6
6	Diabetes mellitus (E10-E14)	1,327	2.5	4.2
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,243	2.4	4.0
8	Cerebrovascular diseases (I60-I69)	1,057	2.0	3.4
9	Influenza and pneumonia (J09-J18)	676	1.3	2.2
10	Septicemia (A40-A41)	560	1.1	1.8
11	Human immunodeficiency virus (HIV) disease (B20-B24)	536	1.0	1.7
12	Chronic lower respiratory diseases (J40-J47)	526	1.0	1.7
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	354	0.7	1.1
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	353	0.7	1.1
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	211	0.4	0.7
...	All other causes (Residual)	8,952	17.0	28.6
White, both sexes, 45-54 years				
...	All causes	139,086	100.0	397.2
1	Malignant neoplasms (C00-C97)	36,544	26.3	104.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	26,367	19.0	75.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17,284	12.4	49.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7,984	5.7	22.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	7,632	5.5	21.8
6	Diabetes mellitus (E10-E14)	4,228	3.0	12.1
7	Chronic lower respiratory diseases (J40-J47)	3,872	2.8	11.1
8	Cerebrovascular diseases (I60-I69)	3,610	2.6	10.3
9	Septicemia (A40-A41)	1,713	1.2	4.9
10	Influenza and pneumonia (J09-J18)	1,703	1.2	4.9
11	Viral hepatitis (B15-B19)	1,602	1.2	4.6
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,272	0.9	3.6
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,149	0.8	3.3
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1,141	0.8	3.3
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	913	0.7	2.6
...	All other causes (Residual)	22,072	15.9	63.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 55-64 years				
...	All causes	269,285	100.0	831.8
1	Malignant neoplasms (C00-C97)	91,710	34.1	283.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	56,017	20.8	173.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14,220	5.3	43.9
4	Chronic lower respiratory diseases (J40-J47)	13,902	5.2	42.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	10,310	3.8	31.8
6	Diabetes mellitus (E10-E14)	9,597	3.6	29.6
7	Cerebrovascular diseases (I60-I69)	8,013	3.0	24.8
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,714	2.5	20.7
9	Septicemia (A40-A41)	4,037	1.5	12.5
10	Influenza and pneumonia (J09-J18)	3,728	1.4	11.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,225	1.2	10.0
12	Viral hepatitis (B15-B19)	3,151	1.2	9.7
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,123	0.8	6.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,239	0.5	3.8
15	Aortic aneurysm and dissection (I71)	1,014	0.4	3.1
...	All other causes (Residual)	40,285	15.0	124.4
White, both sexes, 65-74 years				
...	All causes	383,122	100.0	1,783.7
1	Malignant neoplasms (C00-C97)	133,064	34.7	619.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	81,003	21.1	377.1
3	Chronic lower respiratory diseases (J40-J47)	32,599	8.5	151.8
4	Cerebrovascular diseases (I60-I69)	14,672	3.8	68.3
5	Diabetes mellitus (E10-E14)	13,385	3.5	62.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,519	2.5	44.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,332	1.7	29.5
8	Chronic liver disease and cirrhosis (K70,K73-K74)	6,329	1.7	29.5
9	Septicemia (A40-A41)	6,125	1.6	28.5
10	Influenza and pneumonia (J09-J18)	6,117	1.6	28.5
11	Alzheimer's disease (G30)	4,056	1.1	18.9
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,616	0.9	16.8
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,095	0.8	14.4
14	Parkinson's disease (G20-G21)	2,942	0.8	13.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,436	0.6	11.3
...	All other causes (Residual)	57,832	15.1	269.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 75-84 years				
...	All causes	547,335	100.0	4,694.4
1	Malignant neoplasms (C00-C97)	134,311	24.5	1,152.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	127,797	23.3	1,096.1
3	Chronic lower respiratory diseases (J40-J47)	45,935	8.4	394.0
4	Cerebrovascular diseases (I60-I69)	30,722	5.6	263.5
5	Alzheimer's disease (G30)	20,866	3.8	179.0
6	Diabetes mellitus (E10-E14)	15,659	2.9	134.3
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,160	2.4	112.9
8	Influenza and pneumonia (J09-J18)	12,082	2.2	103.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10,798	2.0	92.6
10	Parkinson's disease (G20-G21)	9,873	1.8	84.7
11	Septicemia (A40-A41)	8,571	1.6	73.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,672	1.0	48.6
13	Pneumonitis due to solids and liquids (J69)	4,364	0.8	37.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,137	0.8	35.5
15	Chronic liver disease and cirrhosis (K70,K73-K74)	3,670	0.7	31.5
...	All other causes (Residual)	99,718	18.2	855.3
White, both sexes, 85 years and over				
...	All causes	749,509	100.0	13,965.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	221,185	29.5	4,121.3
2	Malignant neoplasms (C00-C97)	88,839	11.9	1,655.3
3	Alzheimer's disease (G30)	51,591	6.9	961.3
4	Cerebrovascular diseases (I60-I69)	49,239	6.6	917.5
5	Chronic lower respiratory diseases (J40-J47)	39,492	5.3	735.8
6	Influenza and pneumonia (J09-J18)	23,950	3.2	446.3
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19,374	2.6	361.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15,023	2.0	279.9
9	Diabetes mellitus (E10-E14)	14,159	1.9	263.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	12,057	1.6	224.7
11	Parkinson's disease (G20-G21)	10,151	1.4	189.1
12	Septicemia (A40-A41)	9,407	1.3	175.3
13	Pneumonitis due to solids and liquids (J69)	8,445	1.1	157.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,743	0.6	88.4
15	Atherosclerosis (I70)	3,525	0.5	65.7
...	All other causes (Residual)	178,329	23.8	3,322.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, all ages				
...	All causes	1,110,956	100.0	899.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	275,101	24.8	222.6
2	Malignant neoplasms (C00-C97)	263,167	23.7	213.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	70,161	6.3	56.8
4	Chronic lower respiratory diseases (J40-J47)	63,757	5.7	51.6
5	Cerebrovascular diseases (I60-I69)	44,203	4.0	35.8
6	Diabetes mellitus (E10-E14)	31,745	2.9	25.7
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	28,943	2.6	23.4
8	Alzheimer's disease (G30)	23,648	2.1	19.1
9	Influenza and pneumonia (J09-J18)	22,907	2.1	18.5
10	Chronic liver disease and cirrhosis (K70,K73-K74)	20,884	1.9	16.9
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18,800	1.7	15.2
12	Septicemia (A40-A41)	14,602	1.3	11.8
13	Parkinson's disease (G20-G21)	14,127	1.3	11.4
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9,940	0.9	8.0
15	Pneumonitis due to solids and liquids (J69)	8,989	0.8	7.3
...	All other causes (Residual)	199,982	18.0	161.9
White, male, 1-4 years				
...	All causes	1,601	100.0	26.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	604	37.7	9.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	162	10.1	2.7
3	Malignant neoplasms (C00-C97)	142	8.9	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	98	6.1	1.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	55	3.4	0.9
6	Influenza and pneumonia (J09-J18)	39	2.4	0.6
7	Certain conditions originating in the perinatal period (P00-P96)	22	1.4	0.4
8	Chronic lower respiratory diseases (J40-J47)	19	1.2	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	1.1	*
10	Septicemia (A40-A41)	16	1.0	*
11	Cerebrovascular diseases (I60-I69)	9	0.6	*
12	Meningitis (G00,G03)	7	0.4	*
13	Anemias (D50-D64)	6	0.4	*
13	Acute bronchitis and bronchiolitis (J20-J21)	6	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.3	*
...	All other causes (Residual)	393	24.5	6.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 5-14 years				
...	All causes	2,219	100.0	13.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	684	30.8	4.3
2	Malignant neoplasms (C00-C97)	371	16.7	2.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	203	9.1	1.3
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	130	5.9	0.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	103	4.6	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	53	2.4	0.3
7	Influenza and pneumonia (J09-J18)	48	2.2	0.3
8	Chronic lower respiratory diseases (J40-J47)	38	1.7	0.2
9	Cerebrovascular diseases (I60-I69)	31	1.4	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	26	1.2	0.2
11	Septicemia (A40-A41)	21	0.9	0.1
12	Certain conditions originating in the perinatal period (P00-P96)	12	0.5	*
13	Diabetes mellitus (E10-E14)	11	0.5	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	10	0.5	*
15	Meningitis (G00,G03)	8	0.4	*
...	All other causes (Residual)	470	21.2	2.9
White, male, 15-24 years				
...	All causes	14,880	100.0	87.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,172	48.2	42.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,203	21.5	18.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,212	8.1	7.1
4	Malignant neoplasms (C00-C97)	664	4.5	3.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	398	2.7	2.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	172	1.2	1.0
7	Influenza and pneumonia (J09-J18)	94	0.6	0.5
8	Diabetes mellitus (E10-E14)	67	0.5	0.4
9	Cerebrovascular diseases (I60-I69)	61	0.4	0.4
10	Septicemia (A40-A41)	50	0.3	0.3
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	47	0.3	0.3
12	Legal intervention (Y35,Y89.0)	45	0.3	0.3
13	Chronic lower respiratory diseases (J40-J47)	42	0.3	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	0.2	0.1
15	Human immunodeficiency virus (HIV) disease (B20-B24)	20	0.1	0.1
...	All other causes (Residual)	1,610	10.8	9.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 25-34 years				
...	All causes	23,435	100.0	139.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,236	43.7	61.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,293	18.3	25.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,481	6.3	8.8
4	Malignant neoplasms (C00-C97)	1,428	6.1	8.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,238	5.3	7.4
6	Chronic liver disease and cirrhosis (K70,K73-K74)	355	1.5	2.1
7	Diabetes mellitus (E10-E14)	270	1.2	1.6
8	Cerebrovascular diseases (I60-I69)	196	0.8	1.2
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	193	0.8	1.2
10	Influenza and pneumonia (J09-J18)	184	0.8	1.1
11	Human immunodeficiency virus (HIV) disease (B20-B24)	171	0.7	1.0
12	Septicemia (A40-A41)	129	0.6	0.8
13	Chronic lower respiratory diseases (J40-J47)	102	0.4	0.6
14	Legal intervention (Y35,Y89.0)	94	0.4	0.6
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	83	0.4	0.5
...	All other causes (Residual)	2,982	12.7	17.8
White, male, 35-44 years				
...	All causes	32,884	100.0	208.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,929	27.2	56.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,016	15.3	31.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,438	13.5	28.1
4	Malignant neoplasms (C00-C97)	3,786	11.5	23.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,389	4.2	8.8
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	890	2.7	5.6
7	Diabetes mellitus (E10-E14)	830	2.5	5.2
8	Cerebrovascular diseases (I60-I69)	603	1.8	3.8
9	Human immunodeficiency virus (HIV) disease (B20-B24)	432	1.3	2.7
10	Influenza and pneumonia (J09-J18)	362	1.1	2.3
11	Septicemia (A40-A41)	272	0.8	1.7
12	Chronic lower respiratory diseases (J40-J47)	228	0.7	1.4
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	200	0.6	1.3
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	190	0.6	1.2
15	Aortic aneurysm and dissection (I71)	147	0.4	0.9
...	All other causes (Residual)	5,172	15.7	32.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 45-54 years				
...	All causes	85,790	100.0	491.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	19,163	22.3	109.8
2	Malignant neoplasms (C00-C97)	18,450	21.5	105.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,522	13.4	66.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,916	6.9	33.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	5,156	6.0	29.6
6	Diabetes mellitus (E10-E14)	2,696	3.1	15.5
7	Cerebrovascular diseases (I60-I69)	2,033	2.4	11.7
8	Chronic lower respiratory diseases (J40-J47)	1,789	2.1	10.3
9	Viral hepatitis (B15-B19)	1,131	1.3	6.5
10	Influenza and pneumonia (J09-J18)	966	1.1	5.5
11	Human immunodeficiency virus (HIV) disease (B20-B24)	935	1.1	5.4
12	Septicemia (A40-A41)	914	1.1	5.2
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	820	1.0	4.7
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	745	0.9	4.3
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	616	0.7	3.5
...	All other causes (Residual)	12,938	15.1	74.2
White, male, 55-64 years				
...	All causes	165,870	100.0	1,050.7
1	Malignant neoplasms (C00-C97)	51,001	30.7	323.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	40,106	24.2	254.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,643	5.8	61.1
4	Chronic liver disease and cirrhosis (K70,K73-K74)	7,383	4.5	46.8
5	Chronic lower respiratory diseases (J40-J47)	7,242	4.4	45.9
6	Diabetes mellitus (E10-E14)	5,978	3.6	37.9
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,065	3.1	32.1
8	Cerebrovascular diseases (I60-I69)	4,726	2.8	29.9
9	Viral hepatitis (B15-B19)	2,269	1.4	14.4
10	Septicemia (A40-A41)	2,207	1.3	14.0
11	Influenza and pneumonia (J09-J18)	2,151	1.3	13.6
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,826	1.1	11.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,396	0.8	8.8
14	Aortic aneurysm and dissection (I71)	767	0.5	4.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	722	0.4	4.6
...	All other causes (Residual)	23,388	14.1	148.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 65-74 years				
...	All causes	218,640	100.0	2,152.0
1	Malignant neoplasms (C00-C97)	73,484	33.6	723.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	52,420	24.0	516.0
3	Chronic lower respiratory diseases (J40-J47)	16,242	7.4	159.9
4	Cerebrovascular diseases (I60-I69)	7,883	3.6	77.6
5	Diabetes mellitus (E10-E14)	7,850	3.6	77.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,079	2.8	59.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	4,031	1.8	39.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,404	1.6	33.5
9	Influenza and pneumonia (J09-J18)	3,396	1.6	33.4
10	Septicemia (A40-A41)	3,202	1.5	31.5
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,933	1.3	28.9
12	Parkinson's disease (G20-G21)	1,977	0.9	19.5
13	Alzheimer's disease (G30)	1,759	0.8	17.3
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,690	0.8	16.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,464	0.7	14.4
...	All other causes (Residual)	30,826	14.1	303.4
White, male, 75-84 years				
...	All causes	278,457	100.0	5,507.2
1	Malignant neoplasms (C00-C97)	71,883	25.8	1,421.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	70,230	25.2	1,389.0
3	Chronic lower respiratory diseases (J40-J47)	21,982	7.9	434.8
4	Cerebrovascular diseases (I60-I69)	13,707	4.9	271.1
5	Diabetes mellitus (E10-E14)	8,212	2.9	162.4
6	Alzheimer's disease (G30)	7,977	2.9	157.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,112	2.6	140.7
8	Parkinson's disease (G20-G21)	6,299	2.3	124.6
9	Influenza and pneumonia (J09-J18)	6,191	2.2	122.4
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,725	2.1	113.2
11	Septicemia (A40-A41)	4,086	1.5	80.8
12	Pneumonitis due to solids and liquids (J69)	2,530	0.9	50.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,474	0.9	48.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,330	0.8	46.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,969	0.7	38.9
...	All other causes (Residual)	45,750	16.4	904.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 85 years and over				
...	All causes	278,557	100.0	15,220.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	86,047	30.9	4,701.6
2	Malignant neoplasms (C00-C97)	41,924	15.1	2,290.7
3	Chronic lower respiratory diseases (J40-J47)	16,065	5.8	877.8
4	Cerebrovascular diseases (I60-I69)	14,919	5.4	815.2
5	Alzheimer's disease (G30)	13,557	4.9	740.8
6	Influenza and pneumonia (J09-J18)	9,411	3.4	514.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,746	2.8	423.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,765	2.4	369.6
9	Diabetes mellitus (E10-E14)	5,825	2.1	318.3
10	Parkinson's disease (G20-G21)	5,460	2.0	298.3
11	Pneumonitis due to solids and liquids (J69)	4,255	1.5	232.5
12	Septicemia (A40-A41)	3,652	1.3	199.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,565	1.3	194.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,164	0.8	118.2
15	Aortic aneurysm and dissection (I71)	1,141	0.4	62.3
...	All other causes (Residual)	56,061	20.1	3,063.2
White, female, all ages				
...	All causes	1,106,147	100.0	879.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	247,544	22.4	196.8
2	Malignant neoplasms (C00-C97)	234,949	21.2	186.8
3	Chronic lower respiratory diseases (J40-J47)	72,925	6.6	58.0
4	Cerebrovascular diseases (I60-I69)	63,706	5.8	50.6
5	Alzheimer's disease (G30)	53,739	4.9	42.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	42,642	3.9	33.9
7	Diabetes mellitus (E10-E14)	27,180	2.5	21.6
8	Influenza and pneumonia (J09-J18)	26,106	2.4	20.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18,470	1.7	14.7
10	Septicemia (A40-A41)	16,287	1.5	12.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	14,209	1.3	11.3
12	Chronic liver disease and cirrhosis (K70,K73-K74)	10,987	1.0	8.7
13	Parkinson's disease (G20-G21)	9,426	0.9	7.5
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8,211	0.7	6.5
15	Pneumonitis due to solids and liquids (J69)	7,469	0.7	5.9
...	All other causes (Residual)	252,297	22.8	200.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 1-4 years				
...	All causes	1,173	100.0	20.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	325	27.7	5.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	183	15.6	3.1
3	Malignant neoplasms (C00-C97)	104	8.9	1.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	81	6.9	1.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	48	4.1	0.8
6	Influenza and pneumonia (J09-J18)	31	2.6	0.5
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	1.6	*
8	Septicemia (A40-A41)	13	1.1	*
8	Chronic lower respiratory diseases (J40-J47)	13	1.1	*
10	Cerebrovascular diseases (I60-I69)	11	0.9	*
11	Certain conditions originating in the perinatal period (P00-P96)	8	0.7	*
12	Meningitis (G00,G03)	6	0.5	*
13	Anemias (D50-D64)	5	0.4	*
13	Acute bronchitis and bronchiolitis (J20-J21)	5	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.4	*
...	All other causes (Residual)	316	26.9	5.4
White, female, 5-14 years				
...	All causes	1,622	100.0	10.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	421	26.0	2.8
2	Malignant neoplasms (C00-C97)	346	21.3	2.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	112	6.9	0.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	109	6.7	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	67	4.1	0.4
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	52	3.2	0.3
7	Influenza and pneumonia (J09-J18)	44	2.7	0.3
8	Cerebrovascular diseases (I60-I69)	35	2.2	0.2
9	Chronic lower respiratory diseases (J40-J47)	21	1.3	0.1
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	1.2	*
11	Septicemia (A40-A41)	17	1.0	*
12	Certain conditions originating in the perinatal period (P00-P96)	12	0.7	*
13	Diabetes mellitus (E10-E14)	9	0.6	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.3	*
...	All other causes (Residual)	346	21.3	2.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 15-24 years				
...	All causes	5,738	100.0	35.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,546	44.4	15.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	800	13.9	5.0
3	Malignant neoplasms (C00-C97)	474	8.3	2.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	291	5.1	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	204	3.6	1.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	109	1.9	0.7
7	Pregnancy, childbirth and the puerperium (O00-O99)	103	1.8	0.6
8	Influenza and pneumonia (J09-J18)	65	1.1	0.4
9	Diabetes mellitus (E10-E14)	46	0.8	0.3
9	Cerebrovascular diseases (I60-I69)	46	0.8	0.3
11	Septicemia (A40-A41)	42	0.7	0.3
12	Chronic lower respiratory diseases (J40-J47)	26	0.5	0.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	17	0.3	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	0.3	*
...	All other causes (Residual)	935	16.3	5.8
White, female, 25-34 years				
...	All causes	10,342	100.0	64.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,522	34.1	22.0
2	Malignant neoplasms (C00-C97)	1,355	13.1	8.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,101	10.6	6.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	663	6.4	4.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	374	3.6	2.3
6	Pregnancy, childbirth and the puerperium (O00-O99)	262	2.5	1.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	184	1.8	1.1
8	Diabetes mellitus (E10-E14)	159	1.5	1.0
9	Influenza and pneumonia (J09-J18)	151	1.5	0.9
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	143	1.4	0.9
11	Cerebrovascular diseases (I60-I69)	142	1.4	0.9
12	Septicemia (A40-A41)	96	0.9	0.6
13	Chronic lower respiratory diseases (J40-J47)	82	0.8	0.5
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	70	0.7	0.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	54	0.5	0.3
...	All other causes (Residual)	1,984	19.2	12.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 35-44 years				
...	All causes	19,646	100.0	126.8
1	Malignant neoplasms (C00-C97)	4,918	25.0	31.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,094	20.8	26.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,094	10.7	13.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,386	7.1	8.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	685	3.5	4.4
6	Diabetes mellitus (E10-E14)	497	2.5	3.2
7	Cerebrovascular diseases (I60-I69)	454	2.3	2.9
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	353	1.8	2.3
9	Influenza and pneumonia (J09-J18)	314	1.6	2.0
10	Chronic lower respiratory diseases (J40-J47)	298	1.5	1.9
11	Septicemia (A40-A41)	288	1.5	1.9
12	Pregnancy, childbirth and the puerperium (O00-O99)	193	1.0	1.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	163	0.8	1.1
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	154	0.8	1.0
15	Human immunodeficiency virus (HIV) disease (B20-B24)	104	0.5	0.7
...	All other causes (Residual)	3,651	18.6	23.6
White, female, 45-54 years				
...	All causes	53,296	100.0	303.4
1	Malignant neoplasms (C00-C97)	18,094	34.0	103.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,204	13.5	41.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,762	10.8	32.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	2,476	4.6	14.1
5	Chronic lower respiratory diseases (J40-J47)	2,083	3.9	11.9
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,068	3.9	11.8
7	Cerebrovascular diseases (I60-I69)	1,577	3.0	9.0
8	Diabetes mellitus (E10-E14)	1,532	2.9	8.7
9	Septicemia (A40-A41)	799	1.5	4.5
10	Influenza and pneumonia (J09-J18)	737	1.4	4.2
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	527	1.0	3.0
12	Viral hepatitis (B15-B19)	471	0.9	2.7
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	329	0.6	1.9
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	297	0.6	1.7
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	282	0.5	1.6
...	All other causes (Residual)	9,058	17.0	51.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 55-64 years				
...	All causes	103,415	100.0	623.4
1	Malignant neoplasms (C00-C97)	40,709	39.4	245.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,911	15.4	95.9
3	Chronic lower respiratory diseases (J40-J47)	6,660	6.4	40.1
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,577	4.4	27.6
5	Diabetes mellitus (E10-E14)	3,619	3.5	21.8
6	Cerebrovascular diseases (I60-I69)	3,287	3.2	19.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,927	2.8	17.6
8	Septicemia (A40-A41)	1,830	1.8	11.0
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,649	1.6	9.9
10	Influenza and pneumonia (J09-J18)	1,577	1.5	9.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,399	1.4	8.4
12	Viral hepatitis (B15-B19)	882	0.9	5.3
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	727	0.7	4.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	517	0.5	3.1
15	Alzheimer's disease (G30)	466	0.5	2.8
...	All other causes (Residual)	16,678	16.1	100.5
White, female, 65-74 years				
...	All causes	164,482	100.0	1,453.2
1	Malignant neoplasms (C00-C97)	59,580	36.2	526.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	28,583	17.4	252.5
3	Chronic lower respiratory diseases (J40-J47)	16,357	9.9	144.5
4	Cerebrovascular diseases (I60-I69)	6,789	4.1	60.0
5	Diabetes mellitus (E10-E14)	5,535	3.4	48.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,440	2.1	30.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,928	1.8	25.9
8	Septicemia (A40-A41)	2,923	1.8	25.8
9	Influenza and pneumonia (J09-J18)	2,721	1.7	24.0
10	Chronic liver disease and cirrhosis (K70,K73-K74)	2,298	1.4	20.3
11	Alzheimer's disease (G30)	2,297	1.4	20.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,405	0.9	12.4
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	972	0.6	8.6
14	Parkinson's disease (G20-G21)	965	0.6	8.5
15	Pneumonitis due to solids and liquids (J69)	744	0.5	6.6
...	All other causes (Residual)	26,945	16.4	238.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 75-84 years				
...	All causes	268,878	100.0	4,072.0
1	Malignant neoplasms (C00-C97)	62,428	23.2	945.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	57,567	21.4	871.8
3	Chronic lower respiratory diseases (J40-J47)	23,953	8.9	362.8
4	Cerebrovascular diseases (I60-I69)	17,015	6.3	257.7
5	Alzheimer's disease (G30)	12,889	4.8	195.2
6	Diabetes mellitus (E10-E14)	7,447	2.8	112.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,048	2.2	91.6
8	Influenza and pneumonia (J09-J18)	5,891	2.2	89.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,073	1.9	76.8
10	Septicemia (A40-A41)	4,485	1.7	67.9
11	Parkinson's disease (G20-G21)	3,574	1.3	54.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,198	1.2	48.4
13	Pneumonitis due to solids and liquids (J69)	1,834	0.7	27.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,807	0.7	27.4
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,701	0.6	25.8
...	All other causes (Residual)	53,968	20.1	817.3
White, female, 85 years and over				
...	All causes	470,952	100.0	13,316.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	135,138	28.7	3,821.0
2	Malignant neoplasms (C00-C97)	46,915	10.0	1,326.5
3	Alzheimer's disease (G30)	38,034	8.1	1,075.4
4	Cerebrovascular diseases (I60-I69)	34,320	7.3	970.4
5	Chronic lower respiratory diseases (J40-J47)	23,427	5.0	662.4
6	Influenza and pneumonia (J09-J18)	14,539	3.1	411.1
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,628	2.5	328.8
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	8,492	1.8	240.1
9	Diabetes mellitus (E10-E14)	8,334	1.8	235.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,258	1.8	233.5
11	Septicemia (A40-A41)	5,755	1.2	162.7
12	Parkinson's disease (G20-G21)	4,691	1.0	132.6
13	Pneumonitis due to solids and liquids (J69)	4,190	0.9	118.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,579	0.5	72.9
15	Atherosclerosis (I70)	2,412	0.5	68.2
...	All other causes (Residual)	122,240	26.0	3,456.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, all ages				
...	All causes	302,969	100.0	693.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	72,010	23.8	164.8
2	Malignant neoplasms (C00-C97)	67,953	22.4	155.5
3	Cerebrovascular diseases (I60-I69)	16,269	5.4	37.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,413	4.4	30.7
5	Diabetes mellitus (E10-E14)	13,385	4.4	30.6
6	Chronic lower respiratory diseases (J40-J47)	9,918	3.3	22.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,393	2.8	19.2
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8,059	2.7	18.4
9	Septicemia (A40-A41)	6,250	2.1	14.3
10	Alzheimer's disease (G30)	5,714	1.9	13.1
11	Influenza and pneumonia (J09-J18)	5,567	1.8	12.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,455	1.8	12.5
13	Certain conditions originating in the perinatal period (P00-P96)	4,055	1.3	9.3
14	Human immunodeficiency virus (HIV) disease (B20-B24)	3,742	1.2	8.6
15	Chronic liver disease and cirrhosis (K70,K73-K74)	3,053	1.0	7.0
...	All other causes (Residual)	59,733	19.7	136.7
Black, both sexes, 1-4 years				
...	All causes	1,006	100.0	37.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	299	29.7	11.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	136	13.5	5.0
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	91	9.0	3.4
4	Malignant neoplasms (C00-C97)	64	6.4	2.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	59	5.9	2.2
6	Chronic lower respiratory diseases (J40-J47)	27	2.7	1.0
7	Influenza and pneumonia (J09-J18)	23	2.3	0.8
8	Septicemia (A40-A41)	20	2.0	0.7
9	Cerebrovascular diseases (I60-I69)	15	1.5	*
10	Certain conditions originating in the perinatal period (P00-P96)	13	1.3	*
11	Anemias (D50-D64)	8	0.8	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.7	*
13	Meningitis (G00,G03)	6	0.6	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.5	*
15	Acute bronchitis and bronchiolitis (J20-J21)	2	0.2	*
...	All other causes (Residual)	231	23.0	8.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 5-14 years				
...	All causes	1,169	100.0	17.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	330	28.2	4.9
2	Malignant neoplasms (C00-C97)	122	10.4	1.8
3	Chronic lower respiratory diseases (J40-J47)	87	7.4	1.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	87	7.4	1.3
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	84	7.2	1.2
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	58	5.0	0.9
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	51	4.4	0.8
8	Influenza and pneumonia (J09-J18)	28	2.4	0.4
9	Cerebrovascular diseases (I60-I69)	19	1.6	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	1.4	*
11	Septicemia (A40-A41)	13	1.1	*
12	Anemias (D50-D64)	11	0.9	*
13	Certain conditions originating in the perinatal period (P00-P96)	6	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.4	*
...	All other causes (Residual)	247	21.1	3.6
Black, both sexes, 15-24 years				
...	All causes	6,538	100.0	88.9
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,731	41.8	37.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,415	21.6	19.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	523	8.0	7.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	290	4.4	3.9
5	Malignant neoplasms (C00-C97)	241	3.7	3.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	85	1.3	1.2
7	Chronic lower respiratory diseases (J40-J47)	80	1.2	1.1
8	Diabetes mellitus (E10-E14)	73	1.1	1.0
9	Anemias (D50-D64)	70	1.1	1.0
10	Pregnancy, childbirth and the puerperium (O00-O99)	69	1.1	0.9
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	57	0.9	0.8
12	Legal intervention (Y35,Y89.0)	44	0.7	0.6
13	Cerebrovascular diseases (I60-I69)	38	0.6	0.5
14	Septicemia (A40-A41)	36	0.6	0.5
15	Influenza and pneumonia (J09-J18)	32	0.5	0.4
...	All other causes (Residual)	754	11.5	10.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 25-34 years				
...	All causes	9,531	100.0	152.7
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,483	26.1	39.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,801	18.9	28.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	974	10.2	15.6
4	Malignant neoplasms (C00-C97)	645	6.8	10.3
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	564	5.9	9.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	397	4.2	6.4
7	Diabetes mellitus (E10-E14)	226	2.4	3.6
8	Pregnancy, childbirth and the puerperium (O00-O99)	150	1.6	2.4
9	Cerebrovascular diseases (I60-I69)	145	1.5	2.3
10	Anemias (D50-D64)	114	1.2	1.8
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	100	1.0	1.6
12	Chronic lower respiratory diseases (J40-J47)	99	1.0	1.6
13	Influenza and pneumonia (J09-J18)	97	1.0	1.6
14	Septicemia (A40-A41)	90	0.9	1.4
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	74	0.8	1.2
...	All other causes (Residual)	1,572	16.5	25.2
Black, both sexes, 35-44 years				
...	All causes	13,759	100.0	246.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,799	20.3	50.2
2	Malignant neoplasms (C00-C97)	1,917	13.9	34.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,775	12.9	31.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,234	9.0	22.1
5	Human immunodeficiency virus (HIV) disease (B20-B24)	682	5.0	12.2
6	Diabetes mellitus (E10-E14)	534	3.9	9.6
7	Cerebrovascular diseases (I60-I69)	496	3.6	8.9
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	431	3.1	7.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	245	1.8	4.4
10	Septicemia (A40-A41)	228	1.7	4.1
11	Chronic lower respiratory diseases (J40-J47)	208	1.5	3.7
12	Chronic liver disease and cirrhosis (K70,K73-K74)	197	1.4	3.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	178	1.3	3.2
14	Influenza and pneumonia (J09-J18)	160	1.2	2.9
15	Aortic aneurysm and dissection (I71)	102	0.7	1.8
...	All other causes (Residual)	2,573	18.7	46.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 45-54 years				
...	All causes	32,111	100.0	559.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,668	23.9	133.7
2	Malignant neoplasms (C00-C97)	7,666	23.9	133.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,429	7.6	42.3
4	Cerebrovascular diseases (I60-I69)	1,470	4.6	25.6
5	Diabetes mellitus (E10-E14)	1,415	4.4	24.7
6	Human immunodeficiency virus (HIV) disease (B20-B24)	1,192	3.7	20.8
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	767	2.4	13.4
8	Chronic liver disease and cirrhosis (K70,K73-K74)	746	2.3	13.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	680	2.1	11.9
10	Chronic lower respiratory diseases (J40-J47)	659	2.1	11.5
11	Septicemia (A40-A41)	641	2.0	11.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	550	1.7	9.6
13	Influenza and pneumonia (J09-J18)	450	1.4	7.8
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	370	1.2	6.5
15	Viral hepatitis (B15-B19)	213	0.7	3.7
...	All other causes (Residual)	5,195	16.2	90.6
Black, both sexes, 55-64 years				
...	All causes	58,109	100.0	1,265.7
1	Malignant neoplasms (C00-C97)	17,705	30.5	385.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,320	24.6	311.9
3	Diabetes mellitus (E10-E14)	2,893	5.0	63.0
4	Cerebrovascular diseases (I60-I69)	2,826	4.9	61.6
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,345	4.0	51.1
6	Chronic lower respiratory diseases (J40-J47)	1,800	3.1	39.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,539	2.6	33.5
8	Chronic liver disease and cirrhosis (K70,K73-K74)	1,276	2.2	27.8
9	Septicemia (A40-A41)	1,176	2.0	25.6
10	Human immunodeficiency virus (HIV) disease (B20-B24)	982	1.7	21.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	908	1.6	19.8
12	Influenza and pneumonia (J09-J18)	865	1.5	18.8
13	Viral hepatitis (B15-B19)	644	1.1	14.0
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	352	0.6	7.7
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	255	0.4	5.6
...	All other causes (Residual)	8,223	14.2	179.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 65-74 years				
...	All causes	58,130	100.0	2,405.6
1	Malignant neoplasms (C00-C97)	17,953	30.9	743.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,597	25.1	604.1
3	Cerebrovascular diseases (I60-I69)	3,216	5.5	133.1
4	Diabetes mellitus (E10-E14)	3,175	5.5	131.4
5	Chronic lower respiratory diseases (J40-J47)	2,519	4.3	104.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,894	3.3	78.4
7	Septicemia (A40-A41)	1,343	2.3	55.6
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,093	1.9	45.2
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,085	1.9	44.9
10	Influenza and pneumonia (J09-J18)	1,049	1.8	43.4
11	Chronic liver disease and cirrhosis (K70,K73-K74)	522	0.9	21.6
12	Alzheimer's disease (G30)	419	0.7	17.3
13	Human immunodeficiency virus (HIV) disease (B20-B24)	308	0.5	12.7
14	Pneumonitis due to solids and liquids (J69)	306	0.5	12.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	250	0.4	10.3
...	All other causes (Residual)	8,401	14.5	347.7
Black, both sexes, 75-84 years				
...	All causes	59,873	100.0	5,153.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,401	25.7	1,325.6
2	Malignant neoplasms (C00-C97)	14,287	23.9	1,229.7
3	Cerebrovascular diseases (I60-I69)	4,108	6.9	353.6
4	Diabetes mellitus (E10-E14)	3,027	5.1	260.5
5	Chronic lower respiratory diseases (J40-J47)	2,573	4.3	221.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,132	3.6	183.5
7	Alzheimer's disease (G30)	1,762	2.9	151.7
8	Septicemia (A40-A41)	1,448	2.4	124.6
9	Influenza and pneumonia (J09-J18)	1,238	2.1	106.6
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,228	2.1	105.7
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	799	1.3	68.8
12	Parkinson's disease (G20-G21)	442	0.7	38.0
13	Pneumonitis due to solids and liquids (J69)	439	0.7	37.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	309	0.5	26.6
15	Chronic liver disease and cirrhosis (K70,K73-K74)	212	0.4	18.2
...	All other causes (Residual)	10,468	17.5	901.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 85 years and over				
...	All causes	55,594	100.0	12,428.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,747	28.3	3,520.5
2	Malignant neoplasms (C00-C97)	7,343	13.2	1,641.6
3	Cerebrovascular diseases (I60-I69)	3,896	7.0	871.0
4	Alzheimer's disease (G30)	3,445	6.2	770.2
5	Diabetes mellitus (E10-E14)	2,033	3.7	454.5
6	Chronic lower respiratory diseases (J40-J47)	1,855	3.3	414.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,758	3.2	393.0
8	Influenza and pneumonia (J09-J18)	1,555	2.8	347.6
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,434	2.6	320.6
10	Septicemia (A40-A41)	1,200	2.2	268.3
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	703	1.3	157.2
12	Pneumonitis due to solids and liquids (J69)	500	0.9	111.8
13	Parkinson's disease (G20-G21)	360	0.6	80.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	243	0.4	54.3
15	Atherosclerosis (I70)	202	0.4	45.2
...	All other causes (Residual)	13,320	24.0	2,977.9
Black, male, all ages				
...	All causes	154,767	100.0	739.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	37,096	24.0	177.2
2	Malignant neoplasms (C00-C97)	34,671	22.4	165.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,017	5.8	43.1
4	Cerebrovascular diseases (I60-I69)	7,338	4.7	35.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6,937	4.5	33.1
6	Diabetes mellitus (E10-E14)	6,378	4.1	30.5
7	Chronic lower respiratory diseases (J40-J47)	5,073	3.3	24.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,976	2.6	19.0
9	Septicemia (A40-A41)	2,882	1.9	13.8
10	Influenza and pneumonia (J09-J18)	2,696	1.7	12.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,496	1.6	11.9
12	Human immunodeficiency virus (HIV) disease (B20-B24)	2,479	1.6	11.8
13	Certain conditions originating in the perinatal period (P00-P96)	2,215	1.4	10.6
14	Chronic liver disease and cirrhosis (K70,K73-K74)	1,955	1.3	9.3
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,891	1.2	9.0
...	All other causes (Residual)	27,667	17.9	132.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 1-4 years				
...	All causes	561	100.0	40.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	177	31.6	12.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	70	12.5	5.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	50	8.9	3.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	36	6.4	2.6
5	Malignant neoplasms (C00-C97)	33	5.9	2.4
6	Chronic lower respiratory diseases (J40-J47)	15	2.7	*
7	Septicemia (A40-A41)	13	2.3	*
7	Influenza and pneumonia (J09-J18)	13	2.3	*
9	Cerebrovascular diseases (I60-I69)	12	2.1	*
10	Certain conditions originating in the perinatal period (P00-P96)	6	1.1	*
11	Anemias (D50-D64)	4	0.7	*
11	Meningitis (G00,G03)	4	0.7	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
14	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
14	Pneumoconioses and chemical effects (J60-J66,J68)	1	0.2	*
14	Pneumonitis due to solids and liquids (J69)	1	0.2	*
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.2	*
...	All other causes (Residual)	122	21.7	8.8
Black, male, 5-14 years				
...	All causes	675	100.0	19.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	215	31.9	6.2
2	Malignant neoplasms (C00-C97)	66	9.8	1.9
3	Chronic lower respiratory diseases (J40-J47)	55	8.1	1.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	55	8.1	1.6
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	40	5.9	1.2
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	36	5.3	1.0
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	31	4.6	0.9
8	Influenza and pneumonia (J09-J18)	18	2.7	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	1.6	*
10	Cerebrovascular diseases (I60-I69)	10	1.5	*
11	Septicemia (A40-A41)	8	1.2	*
12	Anemias (D50-D64)	5	0.7	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.4	*
13	Certain conditions originating in the perinatal period (P00-P96)	3	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.4	*
...	All other causes (Residual)	116	17.2	3.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 15-24 years				
...	All causes	5,041	100.0	135.5
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,479	49.2	66.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,023	20.3	27.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	429	8.5	11.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	194	3.8	5.2
5	Malignant neoplasms (C00-C97)	129	2.6	3.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	61	1.2	1.6
7	Chronic lower respiratory diseases (J40-J47)	55	1.1	1.5
8	Diabetes mellitus (E10-E14)	48	1.0	1.3
9	Anemias (D50-D64)	44	0.9	1.2
9	Legal intervention (Y35,Y89.0)	44	0.9	1.2
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	35	0.7	0.9
12	Cerebrovascular diseases (I60-I69)	21	0.4	0.6
13	Septicemia (A40-A41)	19	0.4	*
14	Influenza and pneumonia (J09-J18)	16	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.2	*
...	All other causes (Residual)	435	8.6	11.7
Black, male, 25-34 years				
...	All causes	6,598	100.0	218.6
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,209	33.5	73.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,335	20.2	44.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	631	9.6	20.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	466	7.1	15.4
5	Malignant neoplasms (C00-C97)	309	4.7	10.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	273	4.1	9.0
7	Diabetes mellitus (E10-E14)	121	1.8	4.0
8	Cerebrovascular diseases (I60-I69)	74	1.1	2.5
9	Anemias (D50-D64)	59	0.9	2.0
10	Influenza and pneumonia (J09-J18)	55	0.8	1.8
11	Chronic lower respiratory diseases (J40-J47)	53	0.8	1.8
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	49	0.7	1.6
13	Legal intervention (Y35,Y89.0)	43	0.7	1.4
14	Septicemia (A40-A41)	42	0.6	1.4
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	39	0.6	1.3
...	All other causes (Residual)	840	12.7	27.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 35-44 years				
...	All causes	8,214	100.0	312.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,813	22.1	69.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,246	15.2	47.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,063	12.9	40.4
4	Malignant neoplasms (C00-C97)	717	8.7	27.3
5	Human immunodeficiency virus (HIV) disease (B20-B24)	390	4.7	14.8
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	348	4.2	13.2
7	Diabetes mellitus (E10-E14)	321	3.9	12.2
8	Cerebrovascular diseases (I60-I69)	269	3.3	10.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	141	1.7	5.4
10	Chronic lower respiratory diseases (J40-J47)	119	1.4	4.5
11	Septicemia (A40-A41)	112	1.4	4.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	110	1.3	4.2
13	Chronic liver disease and cirrhosis (K70,K73-K74)	102	1.2	3.9
14	Influenza and pneumonia (J09-J18)	82	1.0	3.1
15	Aortic aneurysm and dissection (I71)	77	0.9	2.9
...	All other causes (Residual)	1,304	15.9	49.6
Black, male, 45-54 years				
...	All causes	18,281	100.0	678.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,802	26.3	178.3
2	Malignant neoplasms (C00-C97)	3,565	19.5	132.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,641	9.0	60.9
4	Cerebrovascular diseases (I60-I69)	827	4.5	30.7
5	Human immunodeficiency virus (HIV) disease (B20-B24)	787	4.3	29.2
6	Diabetes mellitus (E10-E14)	776	4.2	28.8
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	604	3.3	22.4
8	Chronic liver disease and cirrhosis (K70,K73-K74)	456	2.5	16.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	408	2.2	15.2
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	342	1.9	12.7
11	Septicemia (A40-A41)	338	1.8	12.6
12	Chronic lower respiratory diseases (J40-J47)	309	1.7	11.5
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	284	1.6	10.5
14	Influenza and pneumonia (J09-J18)	267	1.5	9.9
15	Viral hepatitis (B15-B19)	140	0.8	5.2
...	All other causes (Residual)	2,735	15.0	101.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 55-64 years				
...	All causes	34,066	100.0	1,628.0
1	Malignant neoplasms (C00-C97)	9,710	28.5	464.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,997	26.4	430.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,702	5.0	81.3
4	Cerebrovascular diseases (I60-I69)	1,634	4.8	78.1
5	Diabetes mellitus (E10-E14)	1,615	4.7	77.2
6	Chronic lower respiratory diseases (J40-J47)	914	2.7	43.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	879	2.6	42.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	874	2.6	41.8
9	Human immunodeficiency virus (HIV) disease (B20-B24)	689	2.0	32.9
10	Septicemia (A40-A41)	603	1.8	28.8
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	522	1.5	24.9
12	Influenza and pneumonia (J09-J18)	485	1.4	23.2
13	Viral hepatitis (B15-B19)	437	1.3	20.9
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	278	0.8	13.3
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	193	0.6	9.2
...	All other causes (Residual)	4,534	13.3	216.7
Black, male, 65-74 years				
...	All causes	31,820	100.0	3,064.4
1	Malignant neoplasms (C00-C97)	9,776	30.7	941.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,385	26.4	807.5
3	Cerebrovascular diseases (I60-I69)	1,713	5.4	165.0
4	Diabetes mellitus (E10-E14)	1,588	5.0	152.9
5	Chronic lower respiratory diseases (J40-J47)	1,410	4.4	135.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	911	2.9	87.7
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	729	2.3	70.2
8	Septicemia (A40-A41)	674	2.1	64.9
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	583	1.8	56.1
10	Influenza and pneumonia (J09-J18)	578	1.8	55.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	363	1.1	35.0
12	Human immunodeficiency virus (HIV) disease (B20-B24)	209	0.7	20.1
13	Pneumonitis due to solids and liquids (J69)	188	0.6	18.1
14	Alzheimer's disease (G30)	186	0.6	17.9
15	Viral hepatitis (B15-B19)	151	0.5	14.5
...	All other causes (Residual)	4,376	13.8	421.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 75-84 years				
...	All causes	27,952	100.0	6,362.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,293	26.1	1,660.0
2	Malignant neoplasms (C00-C97)	7,175	25.7	1,633.2
3	Cerebrovascular diseases (I60-I69)	1,702	6.1	387.4
4	Chronic lower respiratory diseases (J40-J47)	1,332	4.8	303.2
5	Diabetes mellitus (E10-E14)	1,295	4.6	294.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	949	3.4	216.0
7	Alzheimer's disease (G30)	654	2.3	148.9
8	Septicemia (A40-A41)	626	2.2	142.5
9	Influenza and pneumonia (J09-J18)	581	2.1	132.2
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	536	1.9	122.0
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	453	1.6	103.1
12	Parkinson's disease (G20-G21)	284	1.0	64.6
13	Pneumonitis due to solids and liquids (J69)	242	0.9	55.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	150	0.5	34.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	101	0.4	23.0
...	All other causes (Residual)	4,579	16.4	1,042.3
Black, male, 85 years and over				
...	All causes	17,664	100.0	13,657.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,856	27.5	3,754.5
2	Malignant neoplasms (C00-C97)	3,189	18.1	2,465.6
3	Cerebrovascular diseases (I60-I69)	1,053	6.0	814.1
4	Chronic lower respiratory diseases (J40-J47)	805	4.6	622.4
5	Alzheimer's disease (G30)	772	4.4	596.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	617	3.5	477.0
7	Diabetes mellitus (E10-E14)	610	3.5	471.6
8	Influenza and pneumonia (J09-J18)	561	3.2	433.7
9	Septicemia (A40-A41)	416	2.4	321.6
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	367	2.1	283.8
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	269	1.5	208.0
12	Pneumonitis due to solids and liquids (J69)	216	1.2	167.0
13	Parkinson's disease (G20-G21)	168	1.0	129.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	94	0.5	72.7
15	Atherosclerosis (I70)	54	0.3	41.8
...	All other causes (Residual)	3,617	20.5	2,796.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, all ages				
...	All causes	148,202	100.0	651.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	34,914	23.6	153.4
2	Malignant neoplasms (C00-C97)	33,282	22.5	146.2
3	Cerebrovascular diseases (I60-I69)	8,931	6.0	39.2
4	Diabetes mellitus (E10-E14)	7,007	4.7	30.8
5	Chronic lower respiratory diseases (J40-J47)	4,845	3.3	21.3
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,417	3.0	19.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,396	3.0	19.3
8	Alzheimer's disease (G30)	4,050	2.7	17.8
9	Septicemia (A40-A41)	3,368	2.3	14.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,959	2.0	13.0
11	Influenza and pneumonia (J09-J18)	2,871	1.9	12.6
12	Certain conditions originating in the perinatal period (P00-P96)	1,840	1.2	8.1
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,263	0.9	5.5
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,122	0.8	4.9
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,098	0.7	4.8
...	All other causes (Residual)	31,839	21.5	139.9
Black, female, 1-4 years				
...	All causes	445	100.0	33.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	122	27.4	9.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	66	14.8	5.0
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	41	9.2	3.1
4	Malignant neoplasms (C00-C97)	31	7.0	2.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	23	5.2	1.7
6	Chronic lower respiratory diseases (J40-J47)	12	2.7	*
7	Influenza and pneumonia (J09-J18)	10	2.2	*
8	Septicemia (A40-A41)	7	1.6	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	1.6	*
8	Certain conditions originating in the perinatal period (P00-P96)	7	1.6	*
11	Anemias (D50-D64)	4	0.9	*
12	Cerebrovascular diseases (I60-I69)	3	0.7	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.7	*
14	Meningitis (G00,G03)	2	0.4	*
15	Diabetes mellitus (E10-E14)	1	0.2	*
15	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
15	Diseases of appendix (K35-K38)	1	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
...	All other causes (Residual)	103	23.1	7.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 5-14 years				
...	All causes	494	100.0	14.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	115	23.3	3.5
2	Malignant neoplasms (C00-C97)	56	11.3	1.7
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	44	8.9	1.3
4	Chronic lower respiratory diseases (J40-J47)	32	6.5	1.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	32	6.5	1.0
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	22	4.5	0.7
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	20	4.0	0.6
8	Influenza and pneumonia (J09-J18)	10	2.0	*
9	Cerebrovascular diseases (I60-I69)	9	1.8	*
10	Anemias (D50-D64)	6	1.2	*
11	Septicemia (A40-A41)	5	1.0	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	1.0	*
13	Diabetes mellitus (E10-E14)	3	0.6	*
13	Certain conditions originating in the perinatal period (P00-P96)	3	0.6	*
15	Meningitis (G00,G03)	2	0.4	*
15	Pneumonitis due to solids and liquids (J69)	2	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
...	All other causes (Residual)	124	25.1	3.7
Black, female, 15-24 years				
...	All causes	1,497	100.0	41.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	392	26.2	10.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	252	16.8	6.9
3	Malignant neoplasms (C00-C97)	112	7.5	3.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	96	6.4	2.6
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	94	6.3	2.6
6	Pregnancy, childbirth and the puerperium (O00-O99)	69	4.6	1.9
7	Anemias (D50-D64)	26	1.7	0.7
8	Diabetes mellitus (E10-E14)	25	1.7	0.7
8	Chronic lower respiratory diseases (J40-J47)	25	1.7	0.7
10	Human immunodeficiency virus (HIV) disease (B20-B24)	24	1.6	0.7
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	22	1.5	0.6
12	Septicemia (A40-A41)	17	1.1	*
12	Cerebrovascular diseases (I60-I69)	17	1.1	*
14	Influenza and pneumonia (J09-J18)	16	1.1	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.5	*
...	All other causes (Residual)	303	20.2	8.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 25-34 years				
...	All causes	2,933	100.0	91.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	466	15.9	14.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	343	11.7	10.6
3	Malignant neoplasms (C00-C97)	336	11.5	10.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	274	9.3	8.5
5	Pregnancy, childbirth and the puerperium (O00-O99)	150	5.1	4.7
6	Human immunodeficiency virus (HIV) disease (B20-B24)	124	4.2	3.8
7	Diabetes mellitus (E10-E14)	105	3.6	3.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	98	3.3	3.0
9	Cerebrovascular diseases (I60-I69)	71	2.4	2.2
10	Anemias (D50-D64)	55	1.9	1.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	51	1.7	1.6
12	Septicemia (A40-A41)	48	1.6	1.5
13	Chronic lower respiratory diseases (J40-J47)	46	1.6	1.4
14	Influenza and pneumonia (J09-J18)	42	1.4	1.3
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	35	1.2	1.1
...	All other causes (Residual)	689	23.5	21.4
Black, female, 35-44 years				
...	All causes	5,545	100.0	187.9
1	Malignant neoplasms (C00-C97)	1,200	21.6	40.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	986	17.8	33.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	529	9.5	17.9
4	Human immunodeficiency virus (HIV) disease (B20-B24)	292	5.3	9.9
5	Cerebrovascular diseases (I60-I69)	227	4.1	7.7
6	Diabetes mellitus (E10-E14)	213	3.8	7.2
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	171	3.1	5.8
8	Septicemia (A40-A41)	116	2.1	3.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	104	1.9	3.5
10	Chronic liver disease and cirrhosis (K70,K73-K74)	95	1.7	3.2
11	Chronic lower respiratory diseases (J40-J47)	89	1.6	3.0
12	Pregnancy, childbirth and the puerperium (O00-O99)	87	1.6	2.9
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	83	1.5	2.8
14	Influenza and pneumonia (J09-J18)	78	1.4	2.6
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	68	1.2	2.3
...	All other causes (Residual)	1,207	21.8	40.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 45-54 years				
...	All causes	13,830	100.0	454.4
1	Malignant neoplasms (C00-C97)	4,101	29.7	134.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,866	20.7	94.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	788	5.7	25.9
4	Cerebrovascular diseases (I60-I69)	643	4.6	21.1
5	Diabetes mellitus (E10-E14)	639	4.6	21.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	405	2.9	13.3
7	Chronic lower respiratory diseases (J40-J47)	350	2.5	11.5
8	Septicemia (A40-A41)	303	2.2	10.0
9	Chronic liver disease and cirrhosis (K70,K73-K74)	290	2.1	9.5
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	272	2.0	8.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	208	1.5	6.8
12	Influenza and pneumonia (J09-J18)	183	1.3	6.0
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	163	1.2	5.4
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	86	0.6	2.8
15	Viral hepatitis (B15-B19)	73	0.5	2.4
...	All other causes (Residual)	2,460	17.8	80.8
Black, female, 55-64 years				
...	All causes	24,043	100.0	962.2
1	Malignant neoplasms (C00-C97)	7,995	33.3	320.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,323	22.1	213.0
3	Diabetes mellitus (E10-E14)	1,278	5.3	51.1
4	Cerebrovascular diseases (I60-I69)	1,192	5.0	47.7
5	Chronic lower respiratory diseases (J40-J47)	886	3.7	35.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	665	2.8	26.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	643	2.7	25.7
8	Septicemia (A40-A41)	573	2.4	22.9
9	Chronic liver disease and cirrhosis (K70,K73-K74)	397	1.7	15.9
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	386	1.6	15.4
11	Influenza and pneumonia (J09-J18)	380	1.6	15.2
12	Human immunodeficiency virus (HIV) disease (B20-B24)	293	1.2	11.7
13	Viral hepatitis (B15-B19)	207	0.9	8.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	94	0.4	3.8
15	Pneumonitis due to solids and liquids (J69)	84	0.3	3.4
...	All other causes (Residual)	3,647	15.2	146.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 65-74 years				
...	All causes	26,310	100.0	1,909.2
1	Malignant neoplasms (C00-C97)	8,177	31.1	593.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,212	23.6	450.8
3	Diabetes mellitus (E10-E14)	1,587	6.0	115.2
4	Cerebrovascular diseases (I60-I69)	1,503	5.7	109.1
5	Chronic lower respiratory diseases (J40-J47)	1,109	4.2	80.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	983	3.7	71.3
7	Septicemia (A40-A41)	669	2.5	48.5
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	510	1.9	37.0
9	Influenza and pneumonia (J09-J18)	471	1.8	34.2
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	356	1.4	25.8
11	Alzheimer's disease (G30)	233	0.9	16.9
12	Chronic liver disease and cirrhosis (K70,K73-K74)	159	0.6	11.5
13	Pneumonitis due to solids and liquids (J69)	118	0.4	8.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	105	0.4	7.6
15	Human immunodeficiency virus (HIV) disease (B20-B24)	99	0.4	7.2
...	All other causes (Residual)	4,019	15.3	291.6
Black, female, 75-84 years				
...	All causes	31,921	100.0	4,418.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,108	25.4	1,122.2
2	Malignant neoplasms (C00-C97)	7,112	22.3	984.3
3	Cerebrovascular diseases (I60-I69)	2,406	7.5	333.0
4	Diabetes mellitus (E10-E14)	1,732	5.4	239.7
5	Chronic lower respiratory diseases (J40-J47)	1,241	3.9	171.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,183	3.7	163.7
7	Alzheimer's disease (G30)	1,108	3.5	153.4
8	Septicemia (A40-A41)	822	2.6	113.8
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	692	2.2	95.8
10	Influenza and pneumonia (J09-J18)	657	2.1	90.9
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	346	1.1	47.9
12	Pneumonitis due to solids and liquids (J69)	197	0.6	27.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	159	0.5	22.0
14	Parkinson's disease (G20-G21)	158	0.5	21.9
15	Chronic liver disease and cirrhosis (K70,K73-K74)	111	0.3	15.4
...	All other causes (Residual)	5,889	18.4	815.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 85 years and over				
...	All causes	37,930	100.0	11,929.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,891	28.7	3,425.3
2	Malignant neoplasms (C00-C97)	4,154	11.0	1,306.5
3	Cerebrovascular diseases (I60-I69)	2,843	7.5	894.1
4	Alzheimer's disease (G30)	2,673	7.0	840.7
5	Diabetes mellitus (E10-E14)	1,423	3.8	447.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,141	3.0	358.9
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,067	2.8	335.6
8	Chronic lower respiratory diseases (J40-J47)	1,050	2.8	330.2
9	Influenza and pneumonia (J09-J18)	994	2.6	312.6
10	Septicemia (A40-A41)	784	2.1	246.6
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	434	1.1	136.5
12	Pneumonitis due to solids and liquids (J69)	284	0.7	89.3
13	Parkinson's disease (G20-G21)	192	0.5	60.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	149	0.4	46.9
15	Atherosclerosis (I70)	148	0.4	46.5
...	All other causes (Residual)	9,703	25.6	3,051.6
American Indian, both sexes, all ages				
...	All causes	17,052	100.0	382.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,139	18.4	70.4
2	Malignant neoplasms (C00-C97)	3,109	18.2	69.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,833	10.7	41.1
4	Diabetes mellitus (E10-E14)	959	5.6	21.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	944	5.5	21.2
6	Chronic lower respiratory diseases (J40-J47)	757	4.4	17.0
7	Cerebrovascular diseases (I60-I69)	595	3.5	13.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	521	3.1	11.7
9	Influenza and pneumonia (J09-J18)	375	2.2	8.4
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	302	1.8	6.8
11	Septicemia (A40-A41)	288	1.7	6.5
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	241	1.4	5.4
13	Alzheimer's disease (G30)	238	1.4	5.3
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	183	1.1	4.1
15	Viral hepatitis (B15-B19)	133	0.8	3.0
...	All other causes (Residual)	3,435	20.1	77.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 1-4 years				
...	All causes	104	100.0	33.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	43	41.3	13.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	16	15.4	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	11	10.6	*
4	Malignant neoplasms (C00-C97)	3	2.9	*
4	Influenza and pneumonia (J09-J18)	3	2.9	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	1.9	*
7	Nutritional deficiencies (E40-E64)	1	1.0	*
7	Cerebrovascular diseases (I60-I69)	1	1.0	*
7	Acute bronchitis and bronchiolitis (J20-J21)	1	1.0	*
7	Chronic lower respiratory diseases (J40-J47)	1	1.0	*
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	1.0	*
...	All other causes (Residual)	21	20.2	6.8
American Indian, both sexes, 5-14 years				
...	All causes	88	100.0	11.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	35	39.8	4.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11	12.5	*
3	Malignant neoplasms (C00-C97)	10	11.4	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	6.8	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	3.4	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	2.3	*
6	Chronic lower respiratory diseases (J40-J47)	2	2.3	*
8	Viral hepatitis (B15-B19)	1	1.1	*
8	Influenza and pneumonia (J09-J18)	1	1.1	*
8	Pneumonitis due to solids and liquids (J69)	1	1.1	*
8	Certain conditions originating in the perinatal period (P00-P96)	1	1.1	*
...	All other causes (Residual)	15	17.0	*

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 15-24 years				
...	All causes	548	100.0	71.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	245	44.7	31.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	150	27.4	19.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	40	7.3	5.2
4	Malignant neoplasms (C00-C97)	15	2.7	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	14	2.6	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	1.3	*
7	Diabetes mellitus (E10-E14)	5	0.9	*
8	Septicemia (A40-A41)	3	0.5	*
8	Chronic lower respiratory diseases (J40-J47)	3	0.5	*
8	Chronic liver disease and cirrhosis (K70,K73-K74)	3	0.5	*
8	Pregnancy, childbirth and the puerperium (O00-O99)	3	0.5	*
8	Legal intervention (Y35,Y89.0)	3	0.5	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.2	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.2	*
13	Influenza and pneumonia (J09-J18)	1	0.2	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
13	Certain conditions originating in the perinatal period (P00-P96)	1	0.2	*
...	All other causes (Residual)	52	9.5	6.8
American Indian, both sexes, 25-34 years				
...	All causes	923	100.0	134.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	353	38.2	51.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	132	14.3	19.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	74	8.0	10.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	72	7.8	10.5
5	Malignant neoplasms (C00-C97)	43	4.7	6.3
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	40	4.3	5.8
7	Diabetes mellitus (E10-E14)	18	2.0	*
8	Influenza and pneumonia (J09-J18)	9	1.0	*
9	Septicemia (A40-A41)	8	0.9	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.8	*
10	Pregnancy, childbirth and the puerperium (O00-O99)	7	0.8	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	5	0.5	*
12	Cerebrovascular diseases (I60-I69)	5	0.5	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	0.4	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.2	*
...	All other causes (Residual)	144	15.6	21.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 35-44 years				
...	All causes	1,226	100.0	209.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	304	24.8	51.8
2	Chronic liver disease and cirrhosis (K70,K73-K74)	166	13.5	28.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	139	11.3	23.7
4	Malignant neoplasms (C00-C97)	93	7.6	15.9
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	88	7.2	15.0
6	Diabetes mellitus (E10-E14)	54	4.4	9.2
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	50	4.1	8.5
8	Influenza and pneumonia (J09-J18)	28	2.3	4.8
9	Cerebrovascular diseases (I60-I69)	23	1.9	3.9
10	Septicemia (A40-A41)	20	1.6	3.4
11	Viral hepatitis (B15-B19)	9	0.7	*
11	Chronic lower respiratory diseases (J40-J47)	9	0.7	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.7	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	8	0.7	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	0.4	*
...	All other causes (Residual)	221	18.0	37.7
American Indian, both sexes, 45-54 years				
...	All causes	2,300	100.0	419.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	379	16.5	69.2
2	Malignant neoplasms (C00-C97)	359	15.6	65.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	359	15.6	65.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	316	13.7	57.7
5	Diabetes mellitus (E10-E14)	107	4.7	19.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	83	3.6	15.1
7	Viral hepatitis (B15-B19)	51	2.2	9.3
8	Cerebrovascular diseases (I60-I69)	50	2.2	9.1
9	Septicemia (A40-A41)	47	2.0	8.6
10	Chronic lower respiratory diseases (J40-J47)	46	2.0	8.4
11	Influenza and pneumonia (J09-J18)	36	1.6	6.6
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	29	1.3	5.3
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	26	1.1	4.7
14	Human immunodeficiency virus (HIV) disease (B20-B24)	23	1.0	4.2
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	18	0.8	*
...	All other causes (Residual)	371	16.1	67.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 55-64 years				
...	All causes	3,103	100.0	771.0
1	Malignant neoplasms (C00-C97)	776	25.0	192.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	609	19.6	151.3
3	Diabetes mellitus (E10-E14)	238	7.7	59.1
4	Chronic liver disease and cirrhosis (K70,K73-K74)	236	7.6	58.6
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	214	6.9	53.2
6	Chronic lower respiratory diseases (J40-J47)	113	3.6	28.1
7	Cerebrovascular diseases (I60-I69)	86	2.8	21.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	63	2.0	15.7
9	Septicemia (A40-A41)	56	1.8	13.9
10	Viral hepatitis (B15-B19)	54	1.7	13.4
11	Influenza and pneumonia (J09-J18)	53	1.7	13.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	38	1.2	9.4
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	34	1.1	8.4
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	15	0.5	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	0.5	*
...	All other causes (Residual)	504	16.2	125.2
American Indian, both sexes, 65-74 years				
...	All causes	3,168	100.0	1,564.6
1	Malignant neoplasms (C00-C97)	855	27.0	422.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	679	21.4	335.3
3	Chronic lower respiratory diseases (J40-J47)	238	7.5	117.5
4	Diabetes mellitus (E10-E14)	234	7.4	115.6
5	Cerebrovascular diseases (I60-I69)	134	4.2	66.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	109	3.4	53.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	94	3.0	46.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	80	2.5	39.5
9	Septicemia (A40-A41)	62	2.0	30.6
9	Influenza and pneumonia (J09-J18)	62	2.0	30.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	29	0.9	14.3
12	Alzheimer's disease (G30)	17	0.5	*
13	Viral hepatitis (B15-B19)	15	0.5	*
13	Aortic aneurysm and dissection (I71)	15	0.5	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	0.4	*
15	Pneumonitis due to solids and liquids (J69)	13	0.4	*
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	13	0.4	*
...	All other causes (Residual)	506	16.0	249.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 75-84 years				
...	All causes	3,014	100.0	3,647.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	692	23.0	837.5
2	Malignant neoplasms (C00-C97)	666	22.1	806.0
3	Chronic lower respiratory diseases (J40-J47)	215	7.1	260.2
4	Diabetes mellitus (E10-E14)	195	6.5	236.0
5	Cerebrovascular diseases (I60-I69)	156	5.2	188.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	95	3.2	115.0
7	Alzheimer's disease (G30)	94	3.1	113.8
8	Influenza and pneumonia (J09-J18)	74	2.5	89.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	69	2.3	83.5
10	Septicemia (A40-A41)	51	1.7	61.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	44	1.5	53.2
12	Chronic liver disease and cirrhosis (K70,K73-K74)	38	1.3	46.0
13	Parkinson's disease (G20-G21)	24	0.8	29.0
14	Pneumonitis due to solids and liquids (J69)	21	0.7	25.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	0.5	*
...	All other causes (Residual)	566	18.8	685.0
American Indian, both sexes, 85 years and over				
...	All causes	2,259	100.0	8,368.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	581	25.7	2,152.3
2	Malignant neoplasms (C00-C97)	289	12.8	1,070.6
3	Cerebrovascular diseases (I60-I69)	137	6.1	507.5
4	Chronic lower respiratory diseases (J40-J47)	128	5.7	474.2
5	Alzheimer's disease (G30)	121	5.4	448.2
6	Diabetes mellitus (E10-E14)	107	4.7	396.4
7	Influenza and pneumonia (J09-J18)	101	4.5	374.2
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	66	2.9	244.5
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	48	2.1	177.8
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	46	2.0	170.4
11	Septicemia (A40-A41)	39	1.7	144.5
12	Parkinson's disease (G20-G21)	29	1.3	107.4
13	Pneumonitis due to solids and liquids (J69)	25	1.1	92.6
14	Nutritional deficiencies (E40-E64)	12	0.5	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.4	*
15	Atherosclerosis (I70)	9	0.4	*
...	All other causes (Residual)	512	22.7	1,896.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, all ages				
...	All causes	9,331	100.0	416.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,843	19.8	82.3
2	Malignant neoplasms (C00-C97)	1,650	17.7	73.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,177	12.6	52.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	511	5.5	22.8
5	Diabetes mellitus (E10-E14)	490	5.3	21.9
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	401	4.3	17.9
7	Chronic lower respiratory diseases (J40-J47)	370	4.0	16.5
8	Cerebrovascular diseases (I60-I69)	253	2.7	11.3
9	Influenza and pneumonia (J09-J18)	190	2.0	8.5
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	187	2.0	8.3
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	143	1.5	6.4
12	Septicemia (A40-A41)	139	1.5	6.2
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	94	1.0	4.2
14	Viral hepatitis (B15-B19)	80	0.9	3.6
15	Alzheimer's disease (G30)	77	0.8	3.4
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	77	0.8	3.4
...	All other causes (Residual)	1,649	17.7	73.6
American Indian, male, 1-4 years				
...	All causes	65	100.0	41.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	24	36.9	15.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	11	16.9	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	10.8	*
4	Malignant neoplasms (C00-C97)	2	3.1	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	3.1	*
6	Nutritional deficiencies (E40-E64)	1	1.5	*
6	Cerebrovascular diseases (I60-I69)	1	1.5	*
6	Influenza and pneumonia (J09-J18)	1	1.5	*
6	Chronic lower respiratory diseases (J40-J47)	1	1.5	*
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	1.5	*
...	All other causes (Residual)	14	21.5	*

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 5-14 years				
...	All causes	47	100.0	12.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18	38.3	*
2	Malignant neoplasms (C00-C97)	8	17.0	*
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7	14.9	*
4	Chronic lower respiratory diseases (J40-J47)	2	4.3	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	4.3	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	4.3	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	2.1	*
7	Pneumonitis due to solids and liquids (J69)	1	2.1	*
7	Certain conditions originating in the perinatal period (P00-P96)	1	2.1	*
...	All other causes (Residual)	5	10.6	*
American Indian, male, 15-24 years				
...	All causes	388	100.0	98.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	172	44.3	43.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	115	29.6	29.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	28	7.2	7.1
4	Malignant neoplasms (C00-C97)	9	2.3	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	9	2.3	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	1.5	*
7	Diabetes mellitus (E10-E14)	4	1.0	*
8	Legal intervention (Y35,Y89.0)	3	0.8	*
9	Septicemia (A40-A41)	2	0.5	*
9	Chronic lower respiratory diseases (J40-J47)	2	0.5	*
9	Chronic liver disease and cirrhosis (K70,K73-K74)	2	0.5	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.3	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.3	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
12	Certain conditions originating in the perinatal period (P00-P96)	1	0.3	*
...	All other causes (Residual)	32	8.2	8.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 25-34 years				
...	All causes	615	100.0	172.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	241	39.2	67.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	105	17.1	29.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	61	9.9	17.1
4	Chronic liver disease and cirrhosis (K70,K73-K74)	39	6.3	10.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	28	4.6	7.9
6	Malignant neoplasms (C00-C97)	22	3.6	6.2
7	Diabetes mellitus (E10-E14)	7	1.1	*
7	Influenza and pneumonia (J09-J18)	7	1.1	*
9	Septicemia (A40-A41)	3	0.5	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.5	*
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.5	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.3	*
12	Cerebrovascular diseases (I60-I69)	2	0.3	*
14	Tuberculosis (A16-A19)	1	0.2	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.2	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.2	*
14	Hernia (K40-K46)	1	0.2	*
14	Legal intervention (Y35,Y89.0)	1	0.2	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.2	*
...	All other causes (Residual)	85	13.8	23.9
American Indian, male, 35-44 years				
...	All causes	730	100.0	244.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	190	26.0	63.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	94	12.9	31.5
3	Chronic liver disease and cirrhosis (K70,K73-K74)	92	12.6	30.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	69	9.5	23.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	40	5.5	13.4
6	Malignant neoplasms (C00-C97)	39	5.3	13.1
7	Diabetes mellitus (E10-E14)	34	4.7	11.4
8	Septicemia (A40-A41)	10	1.4	*
9	Cerebrovascular diseases (I60-I69)	9	1.2	*
9	Influenza and pneumonia (J09-J18)	9	1.2	*
11	Viral hepatitis (B15-B19)	6	0.8	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	6	0.8	*
11	Chronic lower respiratory diseases (J40-J47)	6	0.8	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.5	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	0.5	*
...	All other causes (Residual)	118	16.2	39.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 45-54 years				
...	All causes	1,375	100.0	506.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	276	20.1	101.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	242	17.6	89.1
3	Chronic liver disease and cirrhosis (K70,K73-K74)	169	12.3	62.2
4	Malignant neoplasms (C00-C97)	168	12.2	61.9
5	Diabetes mellitus (E10-E14)	66	4.8	24.3
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	66	4.8	24.3
7	Viral hepatitis (B15-B19)	29	2.1	10.7
8	Cerebrovascular diseases (I60-I69)	25	1.8	9.2
9	Influenza and pneumonia (J09-J18)	24	1.7	8.8
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	24	1.7	8.8
11	Septicemia (A40-A41)	22	1.6	8.1
12	Human immunodeficiency virus (HIV) disease (B20-B24)	19	1.4	*
13	Chronic lower respiratory diseases (J40-J47)	16	1.2	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	12	0.9	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	0.7	*
...	All other causes (Residual)	207	15.1	76.2
American Indian, male, 55-64 years				
...	All causes	1,815	100.0	937.8
1	Malignant neoplasms (C00-C97)	452	24.9	233.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	399	22.0	206.2
3	Chronic liver disease and cirrhosis (K70,K73-K74)	136	7.5	70.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	136	7.5	70.3
5	Diabetes mellitus (E10-E14)	127	7.0	65.6
6	Chronic lower respiratory diseases (J40-J47)	51	2.8	26.4
7	Cerebrovascular diseases (I60-I69)	46	2.5	23.8
8	Septicemia (A40-A41)	34	1.9	17.6
8	Viral hepatitis (B15-B19)	34	1.9	17.6
10	Influenza and pneumonia (J09-J18)	30	1.7	15.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	29	1.6	15.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	28	1.5	14.5
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	19	1.0	*
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	13	0.7	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	9	0.5	*
...	All other causes (Residual)	272	15.0	140.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 65-74 years				
...	All causes	1,758	100.0	1,845.9
1	Malignant neoplasms (C00-C97)	455	25.9	477.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	443	25.2	465.2
3	Chronic lower respiratory diseases (J40-J47)	120	6.8	126.0
4	Diabetes mellitus (E10-E14)	118	6.7	123.9
5	Cerebrovascular diseases (I60-I69)	69	3.9	72.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	63	3.6	66.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	55	3.1	57.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	42	2.4	44.1
9	Influenza and pneumonia (J09-J18)	34	1.9	35.7
10	Septicemia (A40-A41)	31	1.8	32.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	18	1.0	*
12	Viral hepatitis (B15-B19)	11	0.6	*
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11	0.6	*
14	Parkinson's disease (G20-G21)	9	0.5	*
14	Alzheimer's disease (G30)	9	0.5	*
14	Aortic aneurysm and dissection (I71)	9	0.5	*
...	All other causes (Residual)	261	14.8	274.1
American Indian, male, 75-84 years				
...	All causes	1,482	100.0	4,224.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	376	25.4	1,071.8
2	Malignant neoplasms (C00-C97)	354	23.9	1,009.1
3	Chronic lower respiratory diseases (J40-J47)	103	7.0	293.6
4	Diabetes mellitus (E10-E14)	97	6.5	276.5
5	Cerebrovascular diseases (I60-I69)	67	4.5	191.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	48	3.2	136.8
7	Influenza and pneumonia (J09-J18)	37	2.5	105.5
8	Alzheimer's disease (G30)	35	2.4	99.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	29	2.0	82.7
10	Septicemia (A40-A41)	23	1.6	65.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	18	1.2	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	16	1.1	*
13	Parkinson's disease (G20-G21)	14	0.9	*
14	Pneumonitis due to solids and liquids (J69)	13	0.9	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.5	*
...	All other causes (Residual)	244	16.5	695.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 85 years and over				
...	All causes	856	100.0	9,034.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	215	25.1	2,269.1
2	Malignant neoplasms (C00-C97)	141	16.5	1,488.1
3	Chronic lower respiratory diseases (J40-J47)	68	7.9	717.7
4	Influenza and pneumonia (J09-J18)	43	5.0	453.8
5	Diabetes mellitus (E10-E14)	37	4.3	390.5
6	Cerebrovascular diseases (I60-I69)	33	3.9	348.3
7	Alzheimer's disease (G30)	29	3.4	306.1
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	28	3.3	295.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	24	2.8	253.3
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	16	1.9	*
11	Parkinson's disease (G20-G21)	15	1.8	*
12	Septicemia (A40-A41)	13	1.5	*
13	Pneumonitis due to solids and liquids (J69)	11	1.3	*
14	Nutritional deficiencies (E40-E64)	7	0.8	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.7	*
...	All other causes (Residual)	170	19.9	1,794.2
American Indian, female, all ages				
...	All causes	7,721	100.0	348.2
1	Malignant neoplasms (C00-C97)	1,459	18.9	65.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,296	16.8	58.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	656	8.5	29.6
4	Diabetes mellitus (E10-E14)	469	6.1	21.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	433	5.6	19.5
6	Chronic lower respiratory diseases (J40-J47)	387	5.0	17.5
7	Cerebrovascular diseases (I60-I69)	342	4.4	15.4
8	Influenza and pneumonia (J09-J18)	185	2.4	8.3
9	Alzheimer's disease (G30)	161	2.1	7.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	159	2.1	7.2
11	Septicemia (A40-A41)	149	1.9	6.7
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	120	1.6	5.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	89	1.2	4.0
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	54	0.7	2.4
15	Viral hepatitis (B15-B19)	53	0.7	2.4
...	All other causes (Residual)	1,709	22.1	77.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 1-4 years				
...	All causes	39	100.0	25.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19	48.7	*
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	12.8	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	10.3	*
4	Influenza and pneumonia (J09-J18)	2	5.1	*
5	Malignant neoplasms (C00-C97)	1	2.6	*
5	Acute bronchitis and bronchiolitis (J20-J21)	1	2.6	*
...	All other causes (Residual)	7	17.9	*
American Indian, female, 5-14 years				
...	All causes	41	100.0	10.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17	41.5	*
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4	9.8	*
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	9.8	*
4	Malignant neoplasms (C00-C97)	2	4.9	*
5	Viral hepatitis (B15-B19)	1	2.4	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	2.4	*
5	Influenza and pneumonia (J09-J18)	1	2.4	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	2.4	*
...	All other causes (Residual)	10	24.4	*
American Indian, female, 15-24 years				
...	All causes	160	100.0	42.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	73	45.6	19.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	35	21.9	9.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	12	7.5	*
4	Malignant neoplasms (C00-C97)	6	3.8	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	3.1	*
6	Pregnancy, childbirth and the puerperium (O00-O99)	3	1.9	*
7	Septicemia (A40-A41)	1	0.6	*
7	Diabetes mellitus (E10-E14)	1	0.6	*
7	Influenza and pneumonia (J09-J18)	1	0.6	*
7	Chronic lower respiratory diseases (J40-J47)	1	0.6	*
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.6	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.6	*
...	All other causes (Residual)	20	12.5	5.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 25-34 years				
...	All causes	308	100.0	93.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	112	36.4	34.0
2	Chronic liver disease and cirrhosis (K70,K73-K74)	33	10.7	10.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	27	8.8	8.2
4	Malignant neoplasms (C00-C97)	21	6.8	6.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	13	4.2	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	12	3.9	*
7	Diabetes mellitus (E10-E14)	11	3.6	*
8	Pregnancy, childbirth and the puerperium (O00-O99)	7	2.3	*
9	Septicemia (A40-A41)	5	1.6	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	4	1.3	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	1.3	*
12	Cerebrovascular diseases (I60-I69)	3	1.0	*
13	Influenza and pneumonia (J09-J18)	2	0.6	*
14	Chronic lower respiratory diseases (J40-J47)	1	0.3	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.3	*
...	All other causes (Residual)	52	16.9	15.8
American Indian, female, 35-44 years				
...	All causes	496	100.0	172.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	114	23.0	39.6
2	Chronic liver disease and cirrhosis (K70,K73-K74)	74	14.9	25.7
3	Malignant neoplasms (C00-C97)	54	10.9	18.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	45	9.1	15.6
5	Diabetes mellitus (E10-E14)	20	4.0	6.9
6	Influenza and pneumonia (J09-J18)	19	3.8	*
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	19	3.8	*
8	Cerebrovascular diseases (I60-I69)	14	2.8	*
9	Septicemia (A40-A41)	10	2.0	*
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	10	2.0	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	1.0	*
12	Pregnancy, childbirth and the puerperium (O00-O99)	4	0.8	*
13	Viral hepatitis (B15-B19)	3	0.6	*
13	Chronic lower respiratory diseases (J40-J47)	3	0.6	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.4	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2	0.4	*
...	All other causes (Residual)	98	19.8	34.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 45-54 years				
...	All causes	925	100.0	334.5
1	Malignant neoplasms (C00-C97)	191	20.6	69.1
2	Chronic liver disease and cirrhosis (K70,K73-K74)	147	15.9	53.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	117	12.6	42.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	103	11.1	37.3
5	Diabetes mellitus (E10-E14)	41	4.4	14.8
6	Chronic lower respiratory diseases (J40-J47)	30	3.2	10.8
7	Septicemia (A40-A41)	25	2.7	9.0
7	Cerebrovascular diseases (I60-I69)	25	2.7	9.0
9	Viral hepatitis (B15-B19)	22	2.4	8.0
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	17	1.8	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	1.7	*
12	Influenza and pneumonia (J09-J18)	12	1.3	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	6	0.6	*
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	0.5	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.4	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.4	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	0.4	*
...	All other causes (Residual)	156	16.9	56.4
American Indian, female, 55-64 years				
...	All causes	1,288	100.0	616.6
1	Malignant neoplasms (C00-C97)	324	25.2	155.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	210	16.3	100.5
3	Diabetes mellitus (E10-E14)	111	8.6	53.1
4	Chronic liver disease and cirrhosis (K70,K73-K74)	100	7.8	47.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	78	6.1	37.3
6	Chronic lower respiratory diseases (J40-J47)	62	4.8	29.7
7	Cerebrovascular diseases (I60-I69)	40	3.1	19.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	34	2.6	16.3
9	Influenza and pneumonia (J09-J18)	23	1.8	11.0
10	Septicemia (A40-A41)	22	1.7	10.5
11	Viral hepatitis (B15-B19)	20	1.6	9.6
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	15	1.2	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10	0.8	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.5	*
15	Infections of kidney (N10-N12,N13.6,N15.1)	4	0.3	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	0.3	*
...	All other causes (Residual)	225	17.5	107.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 65-74 years				
...	All causes	1,410	100.0	1,314.7
1	Malignant neoplasms (C00-C97)	400	28.4	373.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	236	16.7	220.1
3	Chronic lower respiratory diseases (J40-J47)	118	8.4	110.0
4	Diabetes mellitus (E10-E14)	116	8.2	108.2
5	Cerebrovascular diseases (I60-I69)	65	4.6	60.6
6	Chronic liver disease and cirrhosis (K70,K73-K74)	54	3.8	50.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	38	2.7	35.4
8	Septicemia (A40-A41)	31	2.2	28.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	31	2.2	28.9
10	Influenza and pneumonia (J09-J18)	28	2.0	26.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	11	0.8	*
12	Alzheimer's disease (G30)	8	0.6	*
13	Aortic aneurysm and dissection (I71)	6	0.4	*
13	Pneumonitis due to solids and liquids (J69)	6	0.4	*
13	Cholelithiasis and other disorders of gallbladder (K80-K82)	6	0.4	*
...	All other causes (Residual)	256	18.2	238.7
American Indian, female, 75-84 years				
...	All causes	1,532	100.0	3,221.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	316	20.6	664.6
2	Malignant neoplasms (C00-C97)	312	20.4	656.2
3	Chronic lower respiratory diseases (J40-J47)	112	7.3	235.5
4	Diabetes mellitus (E10-E14)	98	6.4	206.1
5	Cerebrovascular diseases (I60-I69)	89	5.8	187.2
6	Alzheimer's disease (G30)	59	3.9	124.1
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	47	3.1	98.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	40	2.6	84.1
9	Influenza and pneumonia (J09-J18)	37	2.4	77.8
10	Septicemia (A40-A41)	28	1.8	58.9
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	28	1.8	58.9
12	Chronic liver disease and cirrhosis (K70,K73-K74)	20	1.3	42.1
13	Parkinson's disease (G20-G21)	10	0.7	*
14	Aortic aneurysm and dissection (I71)	8	0.5	*
14	Pneumonitis due to solids and liquids (J69)	8	0.5	*
...	All other causes (Residual)	320	20.9	673.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 85 years and over				
...	All causes	1,403	100.0	8,008.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	366	26.1	2,089.2
2	Malignant neoplasms (C00-C97)	148	10.5	844.8
3	Cerebrovascular diseases (I60-I69)	104	7.4	593.6
4	Alzheimer's disease (G30)	92	6.6	525.1
5	Diabetes mellitus (E10-E14)	70	5.0	399.6
6	Chronic lower respiratory diseases (J40-J47)	60	4.3	342.5
7	Influenza and pneumonia (J09-J18)	58	4.1	331.1
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	38	2.7	216.9
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	32	2.3	182.7
10	Septicemia (A40-A41)	26	1.9	148.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22	1.6	125.6
12	Parkinson's disease (G20-G21)	14	1.0	*
12	Pneumonitis due to solids and liquids (J69)	14	1.0	*
14	Atherosclerosis (I70)	7	0.5	*
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	6	0.4	*
...	All other causes (Residual)	346	24.7	1,975.0
Asian or Pacific Islander, both sexes, all ages				
...	All causes	59,869	100.0	321.4
1	Malignant neoplasms (C00-C97)	15,703	26.2	84.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,311	22.2	71.4
3	Cerebrovascular diseases (I60-I69)	4,205	7.0	22.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,508	4.2	13.5
5	Diabetes mellitus (E10-E14)	2,309	3.9	12.4
6	Influenza and pneumonia (J09-J18)	2,024	3.4	10.9
7	Chronic lower respiratory diseases (J40-J47)	1,848	3.1	9.9
8	Alzheimer's disease (G30)	1,428	2.4	7.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,147	1.9	6.2
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,121	1.9	6.0
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	983	1.6	5.3
12	Septicemia (A40-A41)	729	1.2	3.9
13	Chronic liver disease and cirrhosis (K70,K73-K74)	559	0.9	3.0
14	Parkinson's disease (G20-G21)	528	0.9	2.8
15	Certain conditions originating in the perinatal period (P00-P96)	487	0.8	2.6
...	All other causes (Residual)	10,979	18.3	58.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 1-4 years				
...	All causes	184	100.0	18.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	45	24.5	4.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	29	15.8	3.0
3	Malignant neoplasms (C00-C97)	15	8.2	*
4	Influenza and pneumonia (J09-J18)	6	3.3	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	3.3	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	2.7	*
7	Septicemia (A40-A41)	4	2.2	*
7	Meningitis (G00,G03)	4	2.2	*
7	Chronic lower respiratory diseases (J40-J47)	4	2.2	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	1.6	*
11	Cerebrovascular diseases (I60-I69)	2	1.1	*
11	Certain conditions originating in the perinatal period (P00-P96)	2	1.1	*
13	Shigellosis and amebiasis (A03,A06)	1	0.5	*
13	Anemias (D50-D64)	1	0.5	*
13	Hernia (K40-K46)	1	0.5	*
13	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.5	*
...	All other causes (Residual)	55	29.9	5.6
Asian or Pacific Islander, both sexes, 5-14 years				
...	All causes	242	100.0	10.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	51	21.1	2.1
2	Malignant neoplasms (C00-C97)	46	19.0	1.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	15	6.2	*
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	14	5.8	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	14	5.8	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	11	4.5	*
7	Influenza and pneumonia (J09-J18)	7	2.9	*
7	Chronic lower respiratory diseases (J40-J47)	7	2.9	*
9	Septicemia (A40-A41)	4	1.7	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	1.7	*
9	Cerebrovascular diseases (I60-I69)	4	1.7	*
9	Complications of medical and surgical care (Y40-Y84,Y88)	4	1.7	*
13	Meningitis (G00,G03)	2	0.8	*
13	Certain conditions originating in the perinatal period (P00-P96)	2	0.8	*
15	Hernia (K40-K46)	1	0.4	*
...	All other causes (Residual)	56	23.1	2.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 15-24 years				
...	All causes	782	100.0	30.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	241	30.8	9.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	202	25.8	7.8
3	Malignant neoplasms (C00-C97)	102	13.0	3.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	55	7.0	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	35	4.5	1.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	17	2.2	*
7	Cerebrovascular diseases (I60-I69)	8	1.0	*
8	Septicemia (A40-A41)	7	0.9	*
9	Influenza and pneumonia (J09-J18)	5	0.6	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.5	*
10	Chronic lower respiratory diseases (J40-J47)	4	0.5	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.4	*
12	Pregnancy, childbirth and the puerperium (O00-O99)	3	0.4	*
12	Legal intervention (Y35,Y89.0)	3	0.4	*
15	Tuberculosis (A16-A19)	2	0.3	*
15	Diabetes mellitus (E10-E14)	2	0.3	*
...	All other causes (Residual)	89	11.4	3.4
Asian or Pacific Islander, both sexes, 25-34 years				
...	All causes	1,232	100.0	39.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	297	24.1	9.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	258	20.9	8.2
3	Malignant neoplasms (C00-C97)	202	16.4	6.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	100	8.1	3.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	67	5.4	2.1
6	Pregnancy, childbirth and the puerperium (O00-O99)	22	1.8	0.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	21	1.7	0.7
8	Cerebrovascular diseases (I60-I69)	20	1.6	0.6
9	Human immunodeficiency virus (HIV) disease (B20-B24)	13	1.1	*
10	Diabetes mellitus (E10-E14)	11	0.9	*
11	Septicemia (A40-A41)	10	0.8	*
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	0.7	*
13	Influenza and pneumonia (J09-J18)	8	0.6	*
14	Chronic lower respiratory diseases (J40-J47)	7	0.6	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.5	*
...	All other causes (Residual)	175	14.2	5.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 35-44 years				
...	All causes	2,058	100.0	68.9
1	Malignant neoplasms (C00-C97)	635	30.9	21.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	293	14.2	9.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	252	12.2	8.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	208	10.1	7.0
5	Cerebrovascular diseases (I60-I69)	111	5.4	3.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	54	2.6	1.8
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	54	2.6	1.8
8	Diabetes mellitus (E10-E14)	37	1.8	1.2
9	Human immunodeficiency virus (HIV) disease (B20-B24)	20	1.0	0.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20	1.0	0.7
11	Influenza and pneumonia (J09-J18)	17	0.8	*
11	Chronic lower respiratory diseases (J40-J47)	17	0.8	*
11	Pregnancy, childbirth and the puerperium (O00-O99)	17	0.8	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	16	0.8	*
15	Viral hepatitis (B15-B19)	13	0.6	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	0.6	*
...	All other causes (Residual)	281	13.7	9.4
Asian or Pacific Islander, both sexes, 45-54 years				
...	All causes	4,227	100.0	171.3
1	Malignant neoplasms (C00-C97)	1,616	38.2	65.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	753	17.8	30.5
3	Cerebrovascular diseases (I60-I69)	295	7.0	12.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	285	6.7	11.6
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	184	4.4	7.5
6	Diabetes mellitus (E10-E14)	149	3.5	6.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	91	2.2	3.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	47	1.1	1.9
9	Viral hepatitis (B15-B19)	45	1.1	1.8
10	Septicemia (A40-A41)	44	1.0	1.8
10	Influenza and pneumonia (J09-J18)	44	1.0	1.8
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	44	1.0	1.8
13	Chronic lower respiratory diseases (J40-J47)	42	1.0	1.7
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	32	0.8	1.3
15	Aortic aneurysm and dissection (I71)	29	0.7	1.2
...	All other causes (Residual)	527	12.5	21.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 55-64 years				
...	All causes	7,630	100.0	391.7
1	Malignant neoplasms (C00-C97)	3,133	41.1	160.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,622	21.3	83.3
3	Cerebrovascular diseases (I60-I69)	439	5.8	22.5
4	Diabetes mellitus (E10-E14)	333	4.4	17.1
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	278	3.6	14.3
6	Influenza and pneumonia (J09-J18)	133	1.7	6.8
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	132	1.7	6.8
8	Chronic liver disease and cirrhosis (K70,K73-K74)	129	1.7	6.6
9	Chronic lower respiratory diseases (J40-J47)	127	1.7	6.5
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	120	1.6	6.2
11	Viral hepatitis (B15-B19)	87	1.1	4.5
12	Septicemia (A40-A41)	76	1.0	3.9
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	68	0.9	3.5
14	Aortic aneurysm and dissection (I71)	50	0.7	2.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	41	0.5	2.1
...	All other causes (Residual)	862	11.3	44.3
Asian or Pacific Islander, both sexes, 65-74 years				
...	All causes	10,009	100.0	894.3
1	Malignant neoplasms (C00-C97)	3,680	36.8	328.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,153	21.5	192.4
3	Cerebrovascular diseases (I60-I69)	700	7.0	62.5
4	Diabetes mellitus (E10-E14)	485	4.8	43.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	269	2.7	24.0
6	Chronic lower respiratory diseases (J40-J47)	247	2.5	22.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	218	2.2	19.5
8	Influenza and pneumonia (J09-J18)	213	2.1	19.0
9	Septicemia (A40-A41)	163	1.6	14.6
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	140	1.4	12.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	127	1.3	11.3
12	Parkinson's disease (G20-G21)	79	0.8	7.1
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	74	0.7	6.6
14	Viral hepatitis (B15-B19)	69	0.7	6.2
14	Alzheimer's disease (G30)	69	0.7	6.2
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	69	0.7	6.2
...	All other causes (Residual)	1,254	12.5	112.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 75-84 years				
...	All causes	14,791	100.0	2,725.0
1	Malignant neoplasms (C00-C97)	3,950	26.7	727.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,365	22.8	620.0
3	Cerebrovascular diseases (I60-I69)	1,165	7.9	214.6
4	Diabetes mellitus (E10-E14)	706	4.8	130.1
5	Chronic lower respiratory diseases (J40-J47)	623	4.2	114.8
6	Influenza and pneumonia (J09-J18)	555	3.8	102.3
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	384	2.6	70.7
8	Alzheimer's disease (G30)	351	2.4	64.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	318	2.1	58.6
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	272	1.8	50.1
11	Parkinson's disease (G20-G21)	210	1.4	38.7
12	Septicemia (A40-A41)	204	1.4	37.6
13	Pneumonitis due to solids and liquids (J69)	109	0.7	20.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	94	0.6	17.3
14	Chronic liver disease and cirrhosis (K70,K73-K74)	94	0.6	17.3
...	All other causes (Residual)	2,391	16.2	440.5
Asian or Pacific Islander, both sexes, 85 years and over				
...	All causes	17,836	100.0	8,934.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,956	27.8	2,482.5
2	Malignant neoplasms (C00-C97)	2,321	13.0	1,162.6
3	Cerebrovascular diseases (I60-I69)	1,457	8.2	729.8
4	Influenza and pneumonia (J09-J18)	1,035	5.8	518.4
5	Alzheimer's disease (G30)	995	5.6	498.4
6	Chronic lower respiratory diseases (J40-J47)	770	4.3	385.7
7	Diabetes mellitus (E10-E14)	586	3.3	293.5
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	452	2.5	226.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	412	2.3	206.4
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	394	2.2	197.4
11	Pneumonitis due to solids and liquids (J69)	248	1.4	124.2
12	Parkinson's disease (G20-G21)	222	1.2	111.2
13	Septicemia (A40-A41)	202	1.1	101.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	111	0.6	55.6
15	Aortic aneurysm and dissection (I71)	77	0.4	38.6
...	All other causes (Residual)	3,598	20.2	1,802.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, all ages				
...	All causes	30,980	100.0	347.4
1	Malignant neoplasms (C00-C97)	8,071	26.1	90.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,307	23.6	81.9
3	Cerebrovascular diseases (I60-I69)	1,897	6.1	21.3
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,561	5.0	17.5
5	Diabetes mellitus (E10-E14)	1,228	4.0	13.8
6	Chronic lower respiratory diseases (J40-J47)	1,117	3.6	12.5
7	Influenza and pneumonia (J09-J18)	1,011	3.3	11.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	820	2.6	9.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	574	1.9	6.4
10	Alzheimer's disease (G30)	447	1.4	5.0
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	433	1.4	4.9
12	Septicemia (A40-A41)	371	1.2	4.2
13	Chronic liver disease and cirrhosis (K70,K73-K74)	359	1.2	4.0
14	Parkinson's disease (G20-G21)	299	1.0	3.4
15	Certain conditions originating in the perinatal period (P00-P96)	277	0.9	3.1
...	All other causes (Residual)	5,208	16.8	58.4
Asian or Pacific Islander, male, 1-4 years				
...	All causes	96	100.0	19.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	26	27.1	5.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	13	13.5	*
3	Malignant neoplasms (C00-C97)	6	6.3	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	5.2	*
5	Septicemia (A40-A41)	3	3.1	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	3.1	*
5	Chronic lower respiratory diseases (J40-J47)	3	3.1	*
8	Meningitis (G00,G03)	2	2.1	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.0	*
9	Anemias (D50-D64)	1	1.0	*
9	Hernia (K40-K46)	1	1.0	*
9	Infections of kidney (N10-N12,N13.6,N15.1)	1	1.0	*
9	Certain conditions originating in the perinatal period (P00-P96)	1	1.0	*
...	All other causes (Residual)	30	31.3	6.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 5-14 years				
...	All causes	136	100.0	11.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	33	24.3	2.7
2	Malignant neoplasms (C00-C97)	26	19.1	2.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	8	5.9	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8	5.9	*
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6	4.4	*
6	Chronic lower respiratory diseases (J40-J47)	5	3.7	*
7	Influenza and pneumonia (J09-J18)	4	2.9	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	2.9	*
9	Septicemia (A40-A41)	3	2.2	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	2.2	*
9	Cerebrovascular diseases (I60-I69)	3	2.2	*
12	Meningitis (G00,G03)	2	1.5	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	2	1.5	*
14	Hernia (K40-K46)	1	0.7	*
14	Certain conditions originating in the perinatal period (P00-P96)	1	0.7	*
...	All other causes (Residual)	27	19.9	2.2
Asian or Pacific Islander, male, 15-24 years				
...	All causes	555	100.0	42.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	178	32.1	13.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	156	28.1	11.9
3	Malignant neoplasms (C00-C97)	57	10.3	4.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	46	8.3	3.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	27	4.9	2.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	10	1.8	*
7	Influenza and pneumonia (J09-J18)	4	0.7	*
8	Septicemia (A40-A41)	3	0.5	*
8	Cerebrovascular diseases (I60-I69)	3	0.5	*
8	Legal intervention (Y35,Y89.0)	3	0.5	*
11	Tuberculosis (A16-A19)	2	0.4	*
11	Chronic lower respiratory diseases (J40-J47)	2	0.4	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.2	*
14	Diabetes mellitus (E10-E14)	1	0.2	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.2	*
14	Certain conditions originating in the perinatal period (P00-P96)	1	0.2	*
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.2	*
...	All other causes (Residual)	57	10.3	4.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 25-34 years				
...	All causes	814	100.0	54.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	213	26.2	14.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	199	24.4	13.2
3	Malignant neoplasms (C00-C97)	98	12.0	6.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	74	9.1	4.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	47	5.8	3.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	14	1.7	*
7	Human immunodeficiency virus (HIV) disease (B20-B24)	12	1.5	*
8	Cerebrovascular diseases (I60-I69)	7	0.9	*
8	Influenza and pneumonia (J09-J18)	7	0.9	*
10	Diabetes mellitus (E10-E14)	5	0.6	*
10	Chronic lower respiratory diseases (J40-J47)	5	0.6	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.6	*
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	0.6	*
14	Septicemia (A40-A41)	4	0.5	*
14	Viral hepatitis (B15-B19)	4	0.5	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.5	*
14	Aortic aneurysm and dissection (I71)	4	0.5	*
...	All other causes (Residual)	107	13.1	7.1
Asian or Pacific Islander, male, 35-44 years				
...	All causes	1,244	100.0	88.5
1	Malignant neoplasms (C00-C97)	285	22.9	20.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	229	18.4	16.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	201	16.2	14.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	148	11.9	10.5
5	Cerebrovascular diseases (I60-I69)	57	4.6	4.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	37	3.0	2.6
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	32	2.6	2.3
8	Diabetes mellitus (E10-E14)	26	2.1	1.9
9	Human immunodeficiency virus (HIV) disease (B20-B24)	14	1.1	*
10	Chronic lower respiratory diseases (J40-J47)	12	1.0	*
11	Viral hepatitis (B15-B19)	11	0.9	*
11	Influenza and pneumonia (J09-J18)	11	0.9	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	0.7	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7	0.6	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.6	*
...	All other causes (Residual)	158	12.7	11.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 45-54 years				
...	All causes	2,551	100.0	220.4
1	Malignant neoplasms (C00-C97)	783	30.7	67.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	589	23.1	50.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	186	7.3	16.1
4	Cerebrovascular diseases (I60-I69)	182	7.1	15.7
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	127	5.0	11.0
6	Diabetes mellitus (E10-E14)	95	3.7	8.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	77	3.0	6.7
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	31	1.2	2.7
9	Septicemia (A40-A41)	30	1.2	2.6
10	Viral hepatitis (B15-B19)	28	1.1	2.4
10	Chronic lower respiratory diseases (J40-J47)	28	1.1	2.4
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	27	1.1	2.3
13	Aortic aneurysm and dissection (I71)	22	0.9	1.9
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	20	0.8	1.7
14	Influenza and pneumonia (J09-J18)	20	0.8	1.7
...	All other causes (Residual)	306	12.0	26.4
Asian or Pacific Islander, male, 55-64 years				
...	All causes	4,574	100.0	517.6
1	Malignant neoplasms (C00-C97)	1,647	36.0	186.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,178	25.8	133.3
3	Cerebrovascular diseases (I60-I69)	245	5.4	27.7
4	Diabetes mellitus (E10-E14)	213	4.7	24.1
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	177	3.9	20.0
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	96	2.1	10.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	87	1.9	9.8
8	Chronic lower respiratory diseases (J40-J47)	80	1.7	9.1
9	Influenza and pneumonia (J09-J18)	79	1.7	8.9
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	71	1.6	8.0
11	Viral hepatitis (B15-B19)	52	1.1	5.9
12	Septicemia (A40-A41)	48	1.0	5.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	44	1.0	5.0
14	Aortic aneurysm and dissection (I71)	35	0.8	4.0
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	21	0.5	2.4
...	All other causes (Residual)	501	11.0	56.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 65-74 years				
...	All causes	5,680	100.0	1,126.4
1	Malignant neoplasms (C00-C97)	1,954	34.4	387.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,386	24.4	274.9
3	Cerebrovascular diseases (I60-I69)	368	6.5	73.0
4	Diabetes mellitus (E10-E14)	301	5.3	59.7
5	Chronic lower respiratory diseases (J40-J47)	153	2.7	30.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	143	2.5	28.4
7	Influenza and pneumonia (J09-J18)	123	2.2	24.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	112	2.0	22.2
9	Septicemia (A40-A41)	95	1.7	18.8
10	Chronic liver disease and cirrhosis (K70,K73-K74)	82	1.4	16.3
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	68	1.2	13.5
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	47	0.8	9.3
13	Parkinson's disease (G20-G21)	45	0.8	8.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	40	0.7	7.9
15	Viral hepatitis (B15-B19)	38	0.7	7.5
...	All other causes (Residual)	725	12.8	143.8
Asian or Pacific Islander, male, 75-84 years				
...	All causes	7,449	100.0	3,239.9
1	Malignant neoplasms (C00-C97)	2,068	27.8	899.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,735	23.3	754.6
3	Cerebrovascular diseases (I60-I69)	530	7.1	230.5
4	Chronic lower respiratory diseases (J40-J47)	392	5.3	170.5
5	Diabetes mellitus (E10-E14)	348	4.7	151.4
6	Influenza and pneumonia (J09-J18)	288	3.9	125.3
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	216	2.9	93.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	158	2.1	68.7
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	126	1.7	54.8
10	Alzheimer's disease (G30)	123	1.7	53.5
11	Parkinson's disease (G20-G21)	120	1.6	52.2
12	Septicemia (A40-A41)	95	1.3	41.3
13	Pneumonitis due to solids and liquids (J69)	71	1.0	30.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	47	0.6	20.4
15	Chronic liver disease and cirrhosis (K70,K73-K74)	44	0.6	19.1
...	All other causes (Residual)	1,088	14.6	473.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 85 years and over				
...	All causes	7,385	100.0	10,142.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,074	28.1	2,848.5
2	Malignant neoplasms (C00-C97)	1,147	15.5	1,575.3
3	Cerebrovascular diseases (I60-I69)	500	6.8	686.7
4	Influenza and pneumonia (J09-J18)	474	6.4	651.0
5	Chronic lower respiratory diseases (J40-J47)	437	5.9	600.2
6	Alzheimer's disease (G30)	289	3.9	396.9
7	Diabetes mellitus (E10-E14)	239	3.2	328.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	189	2.6	259.6
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	183	2.5	251.3
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	166	2.2	228.0
11	Pneumonitis due to solids and liquids (J69)	126	1.7	173.1
12	Parkinson's disease (G20-G21)	120	1.6	164.8
13	Septicemia (A40-A41)	84	1.1	115.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	49	0.7	67.3
15	Aortic aneurysm and dissection (I71)	31	0.4	42.6
...	All other causes (Residual)	1,277	17.3	1,753.9
Asian or Pacific Islander, female, all ages				
...	All causes	28,889	100.0	297.4
1	Malignant neoplasms (C00-C97)	7,632	26.4	78.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,004	20.8	61.8
3	Cerebrovascular diseases (I60-I69)	2,308	8.0	23.8
4	Diabetes mellitus (E10-E14)	1,081	3.7	11.1
5	Influenza and pneumonia (J09-J18)	1,013	3.5	10.4
6	Alzheimer's disease (G30)	981	3.4	10.1
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	947	3.3	9.7
8	Chronic lower respiratory diseases (J40-J47)	731	2.5	7.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	573	2.0	5.9
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	550	1.9	5.7
11	Septicemia (A40-A41)	358	1.2	3.7
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	301	1.0	3.1
13	Parkinson's disease (G20-G21)	229	0.8	2.4
14	Certain conditions originating in the perinatal period (P00-P96)	210	0.7	2.2
15	Chronic liver disease and cirrhosis (K70,K73-K74)	200	0.7	2.1
...	All other causes (Residual)	5,771	20.0	59.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 1-4 years				
...	All causes	88	100.0	18.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19	21.6	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	16	18.2	*
3	Malignant neoplasms (C00-C97)	9	10.2	*
4	Influenza and pneumonia (J09-J18)	6	6.8	*
5	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	2.3	*
5	Meningitis (G00,G03)	2	2.3	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	2.3	*
5	Cerebrovascular diseases (I60-I69)	2	2.3	*
9	Shigellosis and amebiasis (A03,A06)	1	1.1	*
9	Septicemia (A40-A41)	1	1.1	*
9	Chronic lower respiratory diseases (J40-J47)	1	1.1	*
9	Certain conditions originating in the perinatal period (P00-P96)	1	1.1	*
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1	1.1	*
...	All other causes (Residual)	25	28.4	5.2
Asian or Pacific Islander, female, 5-14 years				
...	All causes	106	100.0	8.8
1	Malignant neoplasms (C00-C97)	20	18.9	1.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18	17.0	*
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8	7.5	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	7	6.6	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	6.6	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	5.7	*
7	Influenza and pneumonia (J09-J18)	3	2.8	*
8	Chronic lower respiratory diseases (J40-J47)	2	1.9	*
8	Complications of medical and surgical care (Y40-Y84,Y88)	2	1.9	*
10	Septicemia (A40-A41)	1	0.9	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.9	*
10	Cerebrovascular diseases (I60-I69)	1	0.9	*
10	Certain conditions originating in the perinatal period (P00-P96)	1	0.9	*
...	All other causes (Residual)	29	27.4	2.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 15-24 years				
...	All causes	227	100.0	17.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	63	27.8	4.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	46	20.3	3.6
3	Malignant neoplasms (C00-C97)	45	19.8	3.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	9	4.0	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	8	3.5	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	3.1	*
7	Cerebrovascular diseases (I60-I69)	5	2.2	*
8	Septicemia (A40-A41)	4	1.8	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	1.3	*
9	Pregnancy, childbirth and the puerperium (O00-O99)	3	1.3	*
11	Chronic lower respiratory diseases (J40-J47)	2	0.9	*
12	Diabetes mellitus (E10-E14)	1	0.4	*
12	Influenza and pneumonia (J09-J18)	1	0.4	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.4	*
...	All other causes (Residual)	29	12.8	2.3
Asian or Pacific Islander, female, 25-34 years				
...	All causes	418	100.0	25.7
1	Malignant neoplasms (C00-C97)	104	24.9	6.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	84	20.1	5.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	59	14.1	3.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	26	6.2	1.6
5	Pregnancy, childbirth and the puerperium (O00-O99)	22	5.3	1.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	20	4.8	1.2
7	Cerebrovascular diseases (I60-I69)	13	3.1	*
8	Chronic liver disease and cirrhosis (K70,K73-K74)	7	1.7	*
9	Septicemia (A40-A41)	6	1.4	*
9	Diabetes mellitus (E10-E14)	6	1.4	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	1.0	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.5	*
12	Chronic lower respiratory diseases (J40-J47)	2	0.5	*
14	Tuberculosis (A16-A19)	1	0.2	*
14	Viral hepatitis (B15-B19)	1	0.2	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.2	*
14	Anemias (D50-D64)	1	0.2	*
14	Nutritional deficiencies (E40-E64)	1	0.2	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.2	*
14	Aortic aneurysm and dissection (I71)	1	0.2	*
14	Influenza and pneumonia (J09-J18)	1	0.2	*
14	Pneumonitis due to solids and liquids (J69)	1	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
...	All other causes (Residual)	52	12.4	3.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 35-44 years				
...	All causes	814	100.0	51.5
1	Malignant neoplasms (C00-C97)	350	43.0	22.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	64	7.9	4.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	60	7.4	3.8
4	Cerebrovascular diseases (I60-I69)	54	6.6	3.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	51	6.3	3.2
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	22	2.7	1.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	17	2.1	*
7	Pregnancy, childbirth and the puerperium (O00-O99)	17	2.1	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	1.6	*
10	Diabetes mellitus (E10-E14)	11	1.4	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	1.2	*
12	Septicemia (A40-A41)	9	1.1	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9	1.1	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	6	0.7	*
14	Influenza and pneumonia (J09-J18)	6	0.7	*
...	All other causes (Residual)	115	14.1	7.3
Asian or Pacific Islander, female, 45-54 years				
...	All causes	1,676	100.0	128.0
1	Malignant neoplasms (C00-C97)	833	49.7	63.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	164	9.8	12.5
3	Cerebrovascular diseases (I60-I69)	113	6.7	8.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	99	5.9	7.6
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	57	3.4	4.4
6	Diabetes mellitus (E10-E14)	54	3.2	4.1
7	Influenza and pneumonia (J09-J18)	24	1.4	1.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20	1.2	1.5
9	Viral hepatitis (B15-B19)	17	1.0	*
10	Septicemia (A40-A41)	14	0.8	*
10	Chronic lower respiratory diseases (J40-J47)	14	0.8	*
10	Chronic liver disease and cirrhosis (K70,K73-K74)	14	0.8	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	0.8	*
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	13	0.8	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	12	0.7	*
...	All other causes (Residual)	215	12.8	16.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 55-64 years				
...	All causes	3,056	100.0	287.2
1	Malignant neoplasms (C00-C97)	1,486	48.6	139.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	444	14.5	41.7
3	Cerebrovascular diseases (I60-I69)	194	6.3	18.2
4	Diabetes mellitus (E10-E14)	120	3.9	11.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	101	3.3	9.5
6	Influenza and pneumonia (J09-J18)	54	1.8	5.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	49	1.6	4.6
8	Chronic lower respiratory diseases (J40-J47)	47	1.5	4.4
9	Chronic liver disease and cirrhosis (K70,K73-K74)	42	1.4	3.9
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	36	1.2	3.4
11	Viral hepatitis (B15-B19)	35	1.1	3.3
12	Septicemia (A40-A41)	28	0.9	2.6
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	24	0.8	2.3
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	24	0.8	2.3
15	Aortic aneurysm and dissection (I71)	15	0.5	*
...	All other causes (Residual)	357	11.7	33.6
Asian or Pacific Islander, female, 65-74 years				
...	All causes	4,329	100.0	703.9
1	Malignant neoplasms (C00-C97)	1,726	39.9	280.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	767	17.7	124.7
3	Cerebrovascular diseases (I60-I69)	332	7.7	54.0
4	Diabetes mellitus (E10-E14)	184	4.3	29.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	126	2.9	20.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	106	2.4	17.2
7	Chronic lower respiratory diseases (J40-J47)	94	2.2	15.3
8	Influenza and pneumonia (J09-J18)	90	2.1	14.6
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	72	1.7	11.7
10	Septicemia (A40-A41)	68	1.6	11.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	45	1.0	7.3
12	Alzheimer's disease (G30)	39	0.9	6.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	34	0.8	5.5
13	Parkinson's disease (G20-G21)	34	0.8	5.5
15	Viral hepatitis (B15-B19)	31	0.7	5.0
...	All other causes (Residual)	581	13.4	94.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 75-84 years				
...	All causes	7,342	100.0	2,346.7
1	Malignant neoplasms (C00-C97)	1,882	25.6	601.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,630	22.2	521.0
3	Cerebrovascular diseases (I60-I69)	635	8.6	203.0
4	Diabetes mellitus (E10-E14)	358	4.9	114.4
5	Influenza and pneumonia (J09-J18)	267	3.6	85.3
6	Chronic lower respiratory diseases (J40-J47)	231	3.1	73.8
7	Alzheimer's disease (G30)	228	3.1	72.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	168	2.3	53.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	160	2.2	51.1
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	146	2.0	46.7
11	Septicemia (A40-A41)	109	1.5	34.8
12	Parkinson's disease (G20-G21)	90	1.2	28.8
13	Chronic liver disease and cirrhosis (K70,K73-K74)	50	0.7	16.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	47	0.6	15.0
15	Viral hepatitis (B15-B19)	42	0.6	13.4
...	All other causes (Residual)	1,299	17.7	415.2
Asian or Pacific Islander, female, 85 years and over				
...	All causes	10,451	100.0	8,240.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,882	27.6	2,272.4
2	Malignant neoplasms (C00-C97)	1,174	11.2	925.7
3	Cerebrovascular diseases (I60-I69)	957	9.2	754.6
4	Alzheimer's disease (G30)	706	6.8	556.7
5	Influenza and pneumonia (J09-J18)	561	5.4	442.3
6	Diabetes mellitus (E10-E14)	347	3.3	273.6
7	Chronic lower respiratory diseases (J40-J47)	333	3.2	262.6
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	286	2.7	225.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	223	2.1	175.8
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	211	2.0	166.4
11	Pneumonitis due to solids and liquids (J69)	122	1.2	96.2
12	Septicemia (A40-A41)	118	1.1	93.0
13	Parkinson's disease (G20-G21)	102	1.0	80.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	62	0.6	48.9
15	Aortic aneurysm and dissection (I71)	46	0.4	36.3
...	All other causes (Residual)	2,321	22.2	1,830.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013