

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, all ages				
...	All causes	2,397,615	100.0	816.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	652,486	27.2	222.2
2	Malignant neoplasms (C00-C97)	553,888	23.1	188.6
3	Cerebrovascular diseases (I60-I69)	150,074	6.3	51.1
4	Chronic lower respiratory diseases (J40-J47)	121,987	5.1	41.5
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	112,012	4.7	38.1
6	Diabetes mellitus (E10-E14)	73,138	3.1	24.9
7	Alzheimer's disease (G30)	65,965	2.8	22.5
8	Influenza and pneumonia (J10-J18)	59,664	2.5	20.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	42,480	1.8	14.5
10	Septicemia (A40-A41)	33,373	1.4	11.4
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	32,439	1.4	11.0
12	Chronic liver disease and cirrhosis (K70,K73-K74)	27,013	1.1	9.2
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	23,076	1.0	7.9
14	Parkinson's disease (G20-G21)	17,989	0.8	6.1
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	17,357	0.7	5.9
...	All other causes (Residual)	414,674	17.3	141.2
All races, both sexes, 1-4 years				
...	All causes	4,785	100.0	29.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,641	34.3	10.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	569	11.9	3.6
3	Malignant neoplasms (C00-C97)	399	8.3	2.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	377	7.9	2.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	187	3.9	1.2
6	Influenza and pneumonia (J10-J18)	119	2.5	0.7
7	Septicemia (A40-A41)	84	1.8	0.5
8	Certain conditions originating in the perinatal period (P00-P96)	61	1.3	0.4
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	53	1.1	0.3
10	Chronic lower respiratory diseases (J40-J47)	48	1.0	0.3
11	Cerebrovascular diseases (I60-I69)	41	0.9	0.3
12	Anemias (D50-D64)	33	0.7	0.2
13	Meningitis (G00,G03)	25	0.5	0.2
14	Complications of medical and surgical care (Y40-Y84,Y88)	19	0.4	*
15	Meningococcal infection (A39)	18	0.4	*
...	All other causes (Residual)	1,111	23.2	6.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 5-14 years				
...	All causes	6,834	100.0	16.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,666	39.0	6.5
2	Malignant neoplasms (C00-C97)	1,019	14.9	2.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	389	5.7	1.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	329	4.8	0.8
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	285	4.2	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	245	3.6	0.6
7	Chronic lower respiratory diseases (J40-J47)	120	1.8	0.3
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	84	1.2	0.2
9	Influenza and pneumonia (J10-J18)	82	1.2	0.2
10	Cerebrovascular diseases (I60-I69)	77	1.1	0.2
11	Septicemia (A40-A41)	66	1.0	0.2
12	Anemias (D50-D64)	46	0.7	0.1
13	Diabetes mellitus (E10-E14)	37	0.5	0.1
13	Certain conditions originating in the perinatal period (P00-P96)	37	0.5	0.1
15	Human immunodeficiency virus (HIV) disease (B20-B24)	25	0.4	0.1
15	Meningitis (G00,G03)	25	0.4	0.1
...	All other causes (Residual)	1,302	19.1	3.2
All races, both sexes, 15-24 years				
...	All causes	33,421	100.0	80.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15,449	46.2	37.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5,085	15.2	12.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,316	12.9	10.3
4	Malignant neoplasms (C00-C97)	1,709	5.1	4.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,038	3.1	2.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	483	1.4	1.2
7	Cerebrovascular diseases (I60-I69)	211	0.6	0.5
8	Human immunodeficiency virus (HIV) disease (B20-B24)	191	0.6	0.5
9	Influenza and pneumonia (J10-J18)	185	0.6	0.4
10	Chronic lower respiratory diseases (J40-J47)	179	0.5	0.4
11	Pregnancy, childbirth and the puerperium (O00-099)	168	0.5	0.4
12	Diabetes mellitus (E10-E14)	161	0.5	0.4
13	Septicemia (A40-A41)	135	0.4	0.3
14	Anemias (D50-D64)	128	0.4	0.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	110	0.3	0.3
...	All other causes (Residual)	3,873	11.6	9.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 25-34 years				
...	All causes	40,868	100.0	102.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,032	31.9	32.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,074	12.4	12.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4,495	11.0	11.2
4	Malignant neoplasms (C00-C97)	3,633	8.9	9.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,163	7.7	7.9
6	Human immunodeficiency virus (HIV) disease (B20-B24)	1,468	3.6	3.7
7	Diabetes mellitus (E10-E14)	599	1.5	1.5
8	Cerebrovascular diseases (I60-I69)	567	1.4	1.4
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	420	1.0	1.0
10	Septicemia (A40-A41)	328	0.8	0.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	309	0.8	0.8
12	Influenza and pneumonia (J10-J18)	303	0.7	0.8
13	Pregnancy, childbirth and the puerperium (O00-099)	287	0.7	0.7
14	Chronic lower respiratory diseases (J40-J47)	255	0.6	0.6
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	246	0.6	0.6
...	All other causes (Residual)	6,689	16.4	16.7
All races, both sexes, 35-44 years				
...	All causes	85,362	100.0	193.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	16,471	19.3	37.3
2	Malignant neoplasms (C00-C97)	14,723	17.2	33.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,925	15.1	29.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,638	7.8	15.0
5	Human immunodeficiency virus (HIV) disease (B20-B24)	4,826	5.7	10.9
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,984	3.5	6.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,799	3.3	6.3
8	Cerebrovascular diseases (I60-I69)	2,361	2.8	5.4
9	Diabetes mellitus (E10-E14)	2,026	2.4	4.6
10	Influenza and pneumonia (J10-J18)	891	1.0	2.0
11	Chronic lower respiratory diseases (J40-J47)	887	1.0	2.0
12	Septicemia (A40-A41)	846	1.0	1.9
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	801	0.9	1.8
14	Viral hepatitis (B15-B19)	610	0.7	1.4
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	493	0.6	1.1
...	All other causes (Residual)	15,081	17.7	34.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 45-54 years				
...	All causes	177,697	100.0	427.0
1	Malignant neoplasms (C00-C97)	49,520	27.9	119.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	37,556	21.1	90.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	16,942	9.5	40.7
4	Chronic liver disease and cirrhosis (K70,K73-K74)	7,496	4.2	18.0
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,906	3.9	16.6
6	Cerebrovascular diseases (I60-I69)	6,181	3.5	14.9
7	Diabetes mellitus (E10-E14)	5,567	3.1	13.4
8	Human immunodeficiency virus (HIV) disease (B20-B24)	4,422	2.5	10.6
9	Chronic lower respiratory diseases (J40-J47)	3,511	2.0	8.4
10	Septicemia (A40-A41)	2,251	1.3	5.4
11	Viral hepatitis (B15-B19)	2,226	1.3	5.3
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,067	1.2	5.0
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,008	1.1	4.8
14	Influenza and pneumonia (J10-J18)	1,897	1.1	4.6
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,138	0.6	2.7
...	All other causes (Residual)	28,009	15.8	67.3
All races, both sexes, 55-64 years				
...	All causes	264,697	100.0	910.3
1	Malignant neoplasms (C00-C97)	96,956	36.6	333.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	63,613	24.0	218.8
3	Chronic lower respiratory diseases (J40-J47)	11,754	4.4	40.4
4	Diabetes mellitus (E10-E14)	10,780	4.1	37.1
5	Cerebrovascular diseases (I60-I69)	9,966	3.8	34.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,651	3.6	33.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	6,569	2.5	22.6
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,011	1.5	13.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,963	1.5	13.6
10	Septicemia (A40-A41)	3,745	1.4	12.9
11	Influenza and pneumonia (J10-J18)	3,154	1.2	10.8
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,839	0.7	6.3
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,562	0.6	5.4
14	Aortic aneurysm and dissection (I71)	1,318	0.5	4.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,310	0.5	4.5
...	All other causes (Residual)	34,506	13.0	118.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 65-74 years				
...	All causes	399,666	100.0	2,164.6
1	Malignant neoplasms (C00-C97)	139,417	34.9	755.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	99,999	25.0	541.6
3	Chronic lower respiratory diseases (J40-J47)	28,390	7.1	153.8
4	Cerebrovascular diseases (I60-I69)	19,901	5.0	107.8
5	Diabetes mellitus (E10-E14)	16,093	4.0	87.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,116	2.0	44.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,119	1.8	38.6
8	Influenza and pneumonia (J10-J18)	6,382	1.6	34.6
9	Septicemia (A40-A41)	5,983	1.5	32.4
10	Chronic liver disease and cirrhosis (K70,K73-K74)	5,119	1.3	27.7
11	Alzheimer's disease (G30)	3,641	0.9	19.7
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	3,152	0.8	17.1
13	Aortic aneurysm and dissection (I71)	2,932	0.7	15.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,343	0.6	12.7
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,279	0.6	12.3
...	All other causes (Residual)	48,800	12.2	264.3
All races, both sexes, 75-84 years				
...	All causes	684,230	100.0	5,275.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	195,379	28.6	1,506.3
2	Malignant neoplasms (C00-C97)	166,085	24.3	1,280.4
3	Cerebrovascular diseases (I60-I69)	50,092	7.3	386.2
4	Chronic lower respiratory diseases (J40-J47)	47,568	7.0	366.7
5	Diabetes mellitus (E10-E14)	22,945	3.4	176.9
6	Alzheimer's disease (G30)	21,882	3.2	168.7
7	Influenza and pneumonia (J10-J18)	18,066	2.6	139.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14,058	2.1	108.4
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,457	2.0	103.7
10	Septicemia (A40-A41)	10,586	1.5	81.6
11	Parkinson's disease (G20-G21)	8,752	1.3	67.5
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	6,819	1.0	52.6
13	Pneumonitis due to solids and liquids (J69)	5,564	0.8	42.9
14	Aortic aneurysm and dissection (I71)	5,131	0.7	39.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,644	0.7	35.8
...	All other causes (Residual)	93,202	13.6	718.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 85 years and over				
...	All causes	671,773	100.0	13,823.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	237,924	35.4	4,895.9
2	Malignant neoplasms (C00-C97)	80,345	12.0	1,653.3
3	Cerebrovascular diseases (I60-I69)	60,545	9.0	1,245.9
4	Alzheimer's disease (G30)	39,790	5.9	818.8
5	Chronic lower respiratory diseases (J40-J47)	29,239	4.4	601.7
6	Influenza and pneumonia (J10-J18)	28,312	4.2	582.6
7	Diabetes mellitus (E10-E14)	14,918	2.2	307.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13,928	2.1	286.6
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,447	2.0	276.7
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	9,648	1.4	198.5
11	Septicemia (A40-A41)	9,075	1.4	186.7
12	Pneumonitis due to solids and liquids (J69)	8,014	1.2	164.9
13	Atherosclerosis (I70)	6,659	1.0	137.0
14	Parkinson's disease (G20-G21)	6,598	1.0	135.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3,774	0.6	77.7
...	All other causes (Residual)	109,557	16.3	2,254.4
All races, male, all ages				
...	All causes	1,181,668	100.0	817.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	321,973	27.2	222.8
2	Malignant neoplasms (C00-C97)	286,830	24.3	198.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	72,050	6.1	49.8
4	Cerebrovascular diseases (I60-I69)	58,800	5.0	40.7
5	Chronic lower respiratory diseases (J40-J47)	58,646	5.0	40.6
6	Diabetes mellitus (E10-E14)	35,267	3.0	24.4
7	Influenza and pneumonia (J10-J18)	26,861	2.3	18.6
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	25,566	2.2	17.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20,370	1.7	14.1
10	Alzheimer's disease (G30)	18,974	1.6	13.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	17,545	1.5	12.1
12	Septicemia (A40-A41)	15,011	1.3	10.4
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	13,578	1.1	9.4
14	Parkinson's disease (G20-G21)	10,366	0.9	7.2
15	Human immunodeficiency virus (HIV) disease (B20-B24)	9,506	0.8	6.6
...	All other causes (Residual)	190,325	16.1	131.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 1-4 years				
...	All causes	2,649	100.0	32.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	964	36.4	11.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	318	12.0	3.9
3	Malignant neoplasms (C00-C97)	212	8.0	2.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	204	7.7	2.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	94	3.5	1.1
6	Influenza and pneumonia (J10-J18)	47	1.8	0.6
7	Septicemia (A40-A41)	41	1.5	0.5
8	Certain conditions originating in the perinatal period (P00-P96)	38	1.4	0.5
9	Chronic lower respiratory diseases (J40-J47)	30	1.1	0.4
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	28	1.1	0.3
11	Cerebrovascular diseases (I60-I69)	23	0.9	0.3
12	Anemias (D50-D64)	18	0.7	*
13	Meningitis (G00,G03)	15	0.6	*
14	Meningococcal infection (A39)	11	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	10	0.4	*
...	All other causes (Residual)	586	22.1	7.2
All races, male, 5-14 years				
...	All causes	3,999	100.0	19.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,661	41.5	8.0
2	Malignant neoplasms (C00-C97)	572	14.3	2.7
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	211	5.3	1.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	200	5.0	1.0
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	187	4.7	0.9
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	122	3.1	0.6
7	Chronic lower respiratory diseases (J40-J47)	75	1.9	0.4
8	Influenza and pneumonia (J10-J18)	47	1.2	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	46	1.2	0.2
10	Cerebrovascular diseases (I60-I69)	40	1.0	0.2
11	Septicemia (A40-A41)	31	0.8	0.1
12	Certain conditions originating in the perinatal period (P00-P96)	23	0.6	0.1
13	Anemias (D50-D64)	21	0.5	0.1
14	Diabetes mellitus (E10-E14)	16	0.4	*
15	Meningitis (G00,G03)	13	0.3	*
...	All other causes (Residual)	734	18.4	3.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 15-24 years				
...	All causes	24,587	100.0	114.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,423	46.5	53.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4,380	17.8	20.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,596	14.6	16.8
4	Malignant neoplasms (C00-C97)	1,027	4.2	4.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	691	2.8	3.2
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	282	1.1	1.3
7	Influenza and pneumonia (J10-J18)	118	0.5	0.6
8	Cerebrovascular diseases (I60-I69)	116	0.5	0.5
9	Chronic lower respiratory diseases (J40-J47)	110	0.4	0.5
10	Human immunodeficiency virus (HIV) disease (B20-B24)	107	0.4	0.5
11	Diabetes mellitus (E10-E14)	83	0.3	0.4
12	Legal intervention (Y35,Y89.0)	79	0.3	0.4
13	Septicemia (A40-A41)	71	0.3	0.3
14	Anemias (D50-D64)	69	0.3	0.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	63	0.3	0.3
...	All other causes (Residual)	2,372	9.6	11.1
All races, male, 25-34 years				
...	All causes	28,359	100.0	139.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,871	34.8	48.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,142	14.6	20.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,764	13.3	18.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,143	7.6	10.5
5	Malignant neoplasms (C00-C97)	1,750	6.2	8.6
6	Human immunodeficiency virus (HIV) disease (B20-B24)	917	3.2	4.5
7	Diabetes mellitus (E10-E14)	340	1.2	1.7
8	Cerebrovascular diseases (I60-I69)	284	1.0	1.4
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	235	0.8	1.2
10	Chronic liver disease and cirrhosis (K70,K73-K74)	221	0.8	1.1
11	Influenza and pneumonia (J10-J18)	194	0.7	1.0
12	Septicemia (A40-A41)	165	0.6	0.8
13	Chronic lower respiratory diseases (J40-J47)	151	0.5	0.7
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	113	0.4	0.6
15	Legal intervention (Y35,Y89.0)	104	0.4	0.5
...	All other causes (Residual)	3,965	14.0	19.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 35-44 years				
...	All causes	53,677	100.0	243.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,671	21.7	53.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,011	16.8	40.9
3	Malignant neoplasms (C00-C97)	6,408	11.9	29.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,067	9.4	23.0
5	Human immunodeficiency virus (HIV) disease (B20-B24)	3,466	6.5	15.7
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,224	4.1	10.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,916	3.6	8.7
8	Diabetes mellitus (E10-E14)	1,258	2.3	5.7
9	Cerebrovascular diseases (I60-I69)	1,240	2.3	5.6
10	Influenza and pneumonia (J10-J18)	494	0.9	2.2
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	451	0.8	2.0
12	Septicemia (A40-A41)	433	0.8	2.0
13	Chronic lower respiratory diseases (J40-J47)	431	0.8	2.0
14	Viral hepatitis (B15-B19)	399	0.7	1.8
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	279	0.5	1.3
...	All other causes (Residual)	8,929	16.6	40.5
All races, male, 45-54 years				
...	All causes	111,163	100.0	543.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	27,050	24.3	132.3
2	Malignant neoplasms (C00-C97)	25,432	22.9	124.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,988	10.8	58.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	5,382	4.8	26.3
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,078	4.6	24.8
6	Cerebrovascular diseases (I60-I69)	3,409	3.1	16.7
7	Diabetes mellitus (E10-E14)	3,376	3.0	16.5
8	Human immunodeficiency virus (HIV) disease (B20-B24)	3,327	3.0	16.3
9	Chronic lower respiratory diseases (J40-J47)	1,806	1.6	8.8
10	Viral hepatitis (B15-B19)	1,645	1.5	8.0
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,492	1.3	7.3
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,208	1.1	5.9
13	Septicemia (A40-A41)	1,207	1.1	5.9
14	Influenza and pneumonia (J10-J18)	1,142	1.0	5.6
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	714	0.6	3.5
...	All other causes (Residual)	16,907	15.2	82.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 55-64 years				
...	All causes	158,032	100.0	1,128.8
1	Malignant neoplasms (C00-C97)	52,738	33.4	376.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	43,788	27.7	312.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,562	4.2	46.9
4	Diabetes mellitus (E10-E14)	6,153	3.9	44.0
5	Chronic lower respiratory diseases (J40-J47)	6,029	3.8	43.1
6	Cerebrovascular diseases (I60-I69)	5,523	3.5	39.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	4,587	2.9	32.8
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,088	2.0	22.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,120	1.3	15.1
10	Septicemia (A40-A41)	1,934	1.2	13.8
11	Influenza and pneumonia (J10-J18)	1,823	1.2	13.0
12	Human immunodeficiency virus (HIV) disease (B20-B24)	1,262	0.8	9.0
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,036	0.7	7.4
14	Aortic aneurysm and dissection (I71)	972	0.6	6.9
15	Viral hepatitis (B15-B19)	860	0.5	6.1
...	All other causes (Residual)	19,557	12.4	139.7
All races, male, 65-74 years				
...	All causes	222,891	100.0	2,644.8
1	Malignant neoplasms (C00-C97)	76,487	34.3	907.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	61,000	27.4	723.8
3	Chronic lower respiratory diseases (J40-J47)	14,503	6.5	172.1
4	Cerebrovascular diseases (I60-I69)	10,207	4.6	121.1
5	Diabetes mellitus (E10-E14)	8,430	3.8	100.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,992	2.2	59.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,677	1.6	43.6
8	Influenza and pneumonia (J10-J18)	3,605	1.6	42.8
9	Chronic liver disease and cirrhosis (K70,K73-K74)	3,122	1.4	37.0
10	Septicemia (A40-A41)	3,017	1.4	35.8
11	Aortic aneurysm and dissection (I71)	1,946	0.9	23.1
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,901	0.9	22.6
13	Alzheimer's disease (G30)	1,558	0.7	18.5
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,535	0.7	18.2
15	Parkinson's disease (G20-G21)	1,457	0.7	17.3
...	All other causes (Residual)	25,454	11.4	302.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 75-84 years				
...	All causes	333,669	100.0	6,394.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	98,812	29.6	1,893.6
2	Malignant neoplasms (C00-C97)	86,734	26.0	1,662.1
3	Chronic lower respiratory diseases (J40-J47)	23,255	7.0	445.6
4	Cerebrovascular diseases (I60-I69)	21,025	6.3	402.9
5	Diabetes mellitus (E10-E14)	10,569	3.2	202.5
6	Influenza and pneumonia (J10-J18)	9,013	2.7	172.7
7	Alzheimer's disease (G30)	7,777	2.3	149.0
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,030	2.1	134.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,007	2.1	134.3
10	Parkinson's disease (G20-G21)	5,306	1.6	101.7
11	Septicemia (A40-A41)	4,870	1.5	93.3
12	Pneumonitis due to solids and liquids (J69)	3,059	0.9	58.6
13	Aortic aneurysm and dissection (I71)	2,859	0.9	54.8
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,782	0.8	53.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,378	0.7	45.6
...	All other causes (Residual)	41,193	12.3	789.4
All races, male, 85 years and over				
...	All causes	226,650	100.0	15,031.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	79,002	34.9	5,239.3
2	Malignant neoplasms (C00-C97)	35,427	15.6	2,349.5
3	Cerebrovascular diseases (I60-I69)	16,859	7.4	1,118.1
4	Chronic lower respiratory diseases (J40-J47)	12,231	5.4	811.1
5	Influenza and pneumonia (J10-J18)	10,212	4.5	677.2
6	Alzheimer's disease (G30)	9,343	4.1	619.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,612	2.5	372.2
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,210	2.3	345.5
9	Diabetes mellitus (E10-E14)	5,035	2.2	333.9
10	Pneumonitis due to solids and liquids (J69)	3,576	1.6	237.2
11	Parkinson's disease (G20-G21)	3,318	1.5	220.0
12	Septicemia (A40-A41)	3,074	1.4	203.9
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,503	1.1	166.0
14	Atherosclerosis (I70)	1,933	0.9	128.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,492	0.7	98.9
...	All other causes (Residual)	31,823	14.0	2,110.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, all ages				
...	All causes	1,215,947	100.0	815.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	330,513	27.2	221.6
2	Malignant neoplasms (C00-C97)	267,058	22.0	179.1
3	Cerebrovascular diseases (I60-I69)	91,274	7.5	61.2
4	Chronic lower respiratory diseases (J40-J47)	63,341	5.2	42.5
5	Alzheimer's disease (G30)	46,991	3.9	31.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	39,962	3.3	26.8
7	Diabetes mellitus (E10-E14)	37,871	3.1	25.4
8	Influenza and pneumonia (J10-J18)	32,803	2.7	22.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22,110	1.8	14.8
10	Septicemia (A40-A41)	18,362	1.5	12.3
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	14,207	1.2	9.5
12	Chronic liver disease and cirrhosis (K70,K73-K74)	9,468	0.8	6.3
13	Pneumonitis due to solids and liquids (J69)	8,187	0.7	5.5
14	Parkinson's disease (G20-G21)	7,623	0.6	5.1
15	Atherosclerosis (I70)	7,351	0.6	4.9
...	All other causes (Residual)	218,826	18.0	146.7
All races, female, 1-4 years				
...	All causes	2,136	100.0	27.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	677	31.7	8.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	251	11.8	3.2
3	Malignant neoplasms (C00-C97)	187	8.8	2.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	173	8.1	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	93	4.4	1.2
6	Influenza and pneumonia (J10-J18)	72	3.4	0.9
7	Septicemia (A40-A41)	43	2.0	0.6
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	25	1.2	0.3
9	Certain conditions originating in the perinatal period (P00-P96)	23	1.1	0.3
10	Cerebrovascular diseases (I60-I69)	18	0.8	*
10	Chronic lower respiratory diseases (J40-J47)	18	0.8	*
12	Anemias (D50-D64)	15	0.7	*
13	Meningitis (G00,G03)	10	0.5	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	9	0.4	*
15	Meningococcal infection (A39)	7	0.3	*
15	Acute bronchitis and bronchiolitis (J20-J21)	7	0.3	*
...	All other causes (Residual)	508	23.8	6.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 5-14 years				
...	All causes	2,835	100.0	14.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,005	35.4	5.1
2	Malignant neoplasms (C00-C97)	447	15.8	2.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	178	6.3	0.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	129	4.6	0.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	123	4.3	0.6
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	98	3.5	0.5
7	Chronic lower respiratory diseases (J40-J47)	45	1.6	0.2
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	38	1.3	0.2
9	Cerebrovascular diseases (I60-I69)	37	1.3	0.2
10	Septicemia (A40-A41)	35	1.2	0.2
10	Influenza and pneumonia (J10-J18)	35	1.2	0.2
12	Anemias (D50-D64)	25	0.9	0.1
13	Diabetes mellitus (E10-E14)	21	0.7	0.1
14	Certain conditions originating in the perinatal period (P00-P96)	14	0.5	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	13	0.5	*
...	All other causes (Residual)	592	20.9	3.0
All races, female, 15-24 years				
...	All causes	8,834	100.0	43.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,026	45.6	19.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	720	8.2	3.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	705	8.0	3.5
4	Malignant neoplasms (C00-C97)	682	7.7	3.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	347	3.9	1.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	201	2.3	1.0
7	Pregnancy, childbirth and the puerperium (O00-099)	168	1.9	0.8
8	Cerebrovascular diseases (I60-I69)	95	1.1	0.5
9	Human immunodeficiency virus (HIV) disease (B20-B24)	84	1.0	0.4
10	Diabetes mellitus (E10-E14)	78	0.9	0.4
11	Chronic lower respiratory diseases (J40-J47)	69	0.8	0.3
12	Influenza and pneumonia (J10-J18)	67	0.8	0.3
13	Septicemia (A40-A41)	64	0.7	0.3
14	Anemias (D50-D64)	59	0.7	0.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	47	0.5	0.2
...	All other causes (Residual)	1,422	16.1	7.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 25-34 years				
...	All causes	12,509	100.0	63.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,161	25.3	16.0
2	Malignant neoplasms (C00-C97)	1,883	15.1	9.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,020	8.2	5.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	932	7.5	4.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	731	5.8	3.7
6	Human immunodeficiency virus (HIV) disease (B20-B24)	551	4.4	2.8
7	Pregnancy, childbirth and the puerperium (O00-099)	287	2.3	1.5
8	Cerebrovascular diseases (I60-I69)	283	2.3	1.4
9	Diabetes mellitus (E10-E14)	259	2.1	1.3
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	185	1.5	0.9
11	Septicemia (A40-A41)	163	1.3	0.8
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	133	1.1	0.7
13	Influenza and pneumonia (J10-J18)	109	0.9	0.6
14	Chronic lower respiratory diseases (J40-J47)	104	0.8	0.5
15	Chronic liver disease and cirrhosis (K70,K73-K74)	88	0.7	0.4
...	All other causes (Residual)	2,620	20.9	13.3
All races, female, 35-44 years				
...	All causes	31,685	100.0	143.5
1	Malignant neoplasms (C00-C97)	8,315	26.2	37.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,800	15.1	21.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,914	12.4	17.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,571	5.0	7.1
5	Human immunodeficiency virus (HIV) disease (B20-B24)	1,360	4.3	6.2
6	Cerebrovascular diseases (I60-I69)	1,121	3.5	5.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	883	2.8	4.0
8	Diabetes mellitus (E10-E14)	768	2.4	3.5
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	760	2.4	3.4
10	Chronic lower respiratory diseases (J40-J47)	456	1.4	2.1
11	Septicemia (A40-A41)	413	1.3	1.9
12	Influenza and pneumonia (J10-J18)	397	1.3	1.8
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	350	1.1	1.6
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	214	0.7	1.0
15	Viral hepatitis (B15-B19)	211	0.7	1.0
...	All other causes (Residual)	6,152	19.4	27.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 45-54 years				
...	All causes	66,534	100.0	314.3
1	Malignant neoplasms (C00-C97)	24,088	36.2	113.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,506	15.8	49.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,954	7.4	23.4
4	Cerebrovascular diseases (I60-I69)	2,772	4.2	13.1
5	Diabetes mellitus (E10-E14)	2,191	3.3	10.4
6	Chronic liver disease and cirrhosis (K70,K73-K74)	2,114	3.2	10.0
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,828	2.7	8.6
8	Chronic lower respiratory diseases (J40-J47)	1,705	2.6	8.1
9	Human immunodeficiency virus (HIV) disease (B20-B24)	1,095	1.6	5.2
10	Septicemia (A40-A41)	1,044	1.6	4.9
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	859	1.3	4.1
12	Influenza and pneumonia (J10-J18)	755	1.1	3.6
13	Viral hepatitis (B15-B19)	581	0.9	2.7
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	516	0.8	2.4
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	424	0.6	2.0
...	All other causes (Residual)	11,102	16.7	52.5
All races, female, 55-64 years				
...	All causes	106,665	100.0	707.4
1	Malignant neoplasms (C00-C97)	44,218	41.5	293.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	19,825	18.6	131.5
3	Chronic lower respiratory diseases (J40-J47)	5,725	5.4	38.0
4	Diabetes mellitus (E10-E14)	4,627	4.3	30.7
5	Cerebrovascular diseases (I60-I69)	4,443	4.2	29.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,089	2.9	20.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,982	1.9	13.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,843	1.7	12.2
9	Septicemia (A40-A41)	1,811	1.7	12.0
10	Influenza and pneumonia (J10-J18)	1,331	1.2	8.8
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	923	0.9	6.1
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	803	0.8	5.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	560	0.5	3.7
14	Viral hepatitis (B15-B19)	384	0.4	2.5
15	Aortic aneurysm and dissection (I71)	346	0.3	2.3
...	All other causes (Residual)	14,755	13.8	97.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 65-74 years				
...	All causes	176,775	100.0	1,761.4
1	Malignant neoplasms (C00-C97)	62,930	35.6	627.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	38,999	22.1	388.6
3	Chronic lower respiratory diseases (J40-J47)	13,887	7.9	138.4
4	Cerebrovascular diseases (I60-I69)	9,694	5.5	96.6
5	Diabetes mellitus (E10-E14)	7,663	4.3	76.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,442	1.9	34.3
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,124	1.8	31.1
8	Septicemia (A40-A41)	2,966	1.7	29.6
9	Influenza and pneumonia (J10-J18)	2,777	1.6	27.7
10	Alzheimer's disease (G30)	2,083	1.2	20.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1,997	1.1	19.9
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,617	0.9	16.1
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,021	0.6	10.2
14	Aortic aneurysm and dissection (I71)	986	0.6	9.8
15	Parkinson's disease (G20-G21)	757	0.4	7.5
...	All other causes (Residual)	22,832	12.9	227.5
All races, female, 75-84 years				
...	All causes	350,561	100.0	4,521.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	96,567	27.5	1,245.6
2	Malignant neoplasms (C00-C97)	79,351	22.6	1,023.5
3	Cerebrovascular diseases (I60-I69)	29,067	8.3	374.9
4	Chronic lower respiratory diseases (J40-J47)	24,313	6.9	313.6
5	Alzheimer's disease (G30)	14,105	4.0	181.9
6	Diabetes mellitus (E10-E14)	12,376	3.5	159.6
7	Influenza and pneumonia (J10-J18)	9,053	2.6	116.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,051	2.0	90.9
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,427	1.8	82.9
10	Septicemia (A40-A41)	5,716	1.6	73.7
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	4,037	1.2	52.1
12	Parkinson's disease (G20-G21)	3,446	1.0	44.4
13	Pneumonitis due to solids and liquids (J69)	2,505	0.7	32.3
14	Aortic aneurysm and dissection (I71)	2,272	0.6	29.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,266	0.6	29.2
...	All other causes (Residual)	52,009	14.8	670.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 85 years and over				
...	All causes	445,123	100.0	13,280.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	158,922	35.7	4,741.5
2	Malignant neoplasms (C00-C97)	44,918	10.1	1,340.1
3	Cerebrovascular diseases (I60-I69)	43,686	9.8	1,303.4
4	Alzheimer's disease (G30)	30,447	6.8	908.4
5	Influenza and pneumonia (J10-J18)	18,100	4.1	540.0
6	Chronic lower respiratory diseases (J40-J47)	17,008	3.8	507.4
7	Diabetes mellitus (E10-E14)	9,883	2.2	294.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,316	1.9	248.1
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,237	1.9	245.8
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	7,145	1.6	213.2
11	Septicemia (A40-A41)	6,001	1.3	179.0
12	Atherosclerosis (I70)	4,726	1.1	141.0
13	Pneumonitis due to solids and liquids (J69)	4,438	1.0	132.4
14	Parkinson's disease (G20-G21)	3,280	0.7	97.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,282	0.5	68.1
...	All other causes (Residual)	77,734	17.5	2,319.2
White, both sexes, all ages				
...	All causes	2,056,643	100.0	863.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	565,703	27.5	237.4
2	Malignant neoplasms (C00-C97)	478,134	23.2	200.7
3	Cerebrovascular diseases (I60-I69)	127,868	6.2	53.7
4	Chronic lower respiratory diseases (J40-J47)	112,914	5.5	47.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	95,890	4.7	40.2
6	Alzheimer's disease (G30)	61,087	3.0	25.6
7	Diabetes mellitus (E10-E14)	58,087	2.8	24.4
8	Influenza and pneumonia (J10-J18)	52,430	2.5	22.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	33,691	1.6	14.1
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	29,251	1.4	12.3
11	Septicemia (A40-A41)	26,678	1.3	11.2
12	Chronic liver disease and cirrhosis (K70,K73-K74)	23,562	1.1	9.9
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	17,693	0.9	7.4
14	Parkinson's disease (G20-G21)	17,041	0.8	7.2
15	Pneumonitis due to solids and liquids (J69)	14,968	0.7	6.3
...	All other causes (Residual)	341,646	16.6	143.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 1-4 years				
...	All causes	3,363	100.0	27.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,242	36.9	10.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	400	11.9	3.2
3	Malignant neoplasms (C00-C97)	311	9.2	2.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	201	6.0	1.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	116	3.4	0.9
6	Influenza and pneumonia (J10-J18)	77	2.3	0.6
7	Septicemia (A40-A41)	60	1.8	0.5
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	43	1.3	0.3
9	Certain conditions originating in the perinatal period (P00-P96)	33	1.0	0.3
10	Cerebrovascular diseases (I60-I69)	28	0.8	0.2
11	Chronic lower respiratory diseases (J40-J47)	22	0.7	0.2
12	Meningitis (G00,G03)	19	0.6	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	16	0.5	*
14	Meningococcal infection (A39)	15	0.4	*
15	Anemias (D50-D64)	13	0.4	*
...	All other causes (Residual)	767	22.8	6.2
White, both sexes, 5-14 years				
...	All causes	4,900	100.0	15.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,953	39.9	6.2
2	Malignant neoplasms (C00-C97)	793	16.2	2.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	287	5.9	0.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	218	4.4	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	184	3.8	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	159	3.2	0.5
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	59	1.2	0.2
8	Cerebrovascular diseases (I60-I69)	58	1.2	0.2
9	Influenza and pneumonia (J10-J18)	57	1.2	0.2
10	Chronic lower respiratory diseases (J40-J47)	53	1.1	0.2
11	Septicemia (A40-A41)	50	1.0	0.2
12	Certain conditions originating in the perinatal period (P00-P96)	26	0.5	0.1
13	Anemias (D50-D64)	20	0.4	0.1
14	Diabetes mellitus (E10-E14)	19	0.4	*
14	Meningitis (G00,G03)	19	0.4	*
...	All other causes (Residual)	945	19.3	3.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 15-24 years				
...	All causes	25,013	100.0	76.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,174	52.7	40.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,610	14.4	11.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,126	8.5	6.5
4	Malignant neoplasms (C00-C97)	1,323	5.3	4.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	678	2.7	2.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	352	1.4	1.1
7	Cerebrovascular diseases (I60-I69)	152	0.6	0.5
8	Influenza and pneumonia (J10-J18)	142	0.6	0.4
9	Diabetes mellitus (E10-E14)	99	0.4	0.3
10	Chronic lower respiratory diseases (J40-J47)	91	0.4	0.3
11	Pregnancy, childbirth and the puerperium (O00-099)	88	0.4	0.3
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	86	0.3	0.3
13	Septicemia (A40-A41)	82	0.3	0.3
14	Legal intervention (Y35,Y89.0)	55	0.2	0.2
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	53	0.2	0.2
...	All other causes (Residual)	2,902	11.6	8.9
White, both sexes, 25-34 years				
...	All causes	29,275	100.0	92.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,787	36.8	34.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,288	14.6	13.6
3	Malignant neoplasms (C00-C97)	2,783	9.5	8.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,040	7.0	6.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,920	6.6	6.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	503	1.7	1.6
7	Diabetes mellitus (E10-E14)	396	1.4	1.3
8	Cerebrovascular diseases (I60-I69)	373	1.3	1.2
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	345	1.2	1.1
10	Chronic liver disease and cirrhosis (K70,K73-K74)	251	0.9	0.8
11	Influenza and pneumonia (J10-J18)	228	0.8	0.7
12	Septicemia (A40-A41)	206	0.7	0.7
13	Pregnancy, childbirth and the puerperium (O00-099)	168	0.6	0.5
14	Chronic lower respiratory diseases (J40-J47)	150	0.5	0.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	129	0.4	0.4
...	All other causes (Residual)	4,708	16.1	14.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 35-44 years				
...	All causes	64,040	100.0	179.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,841	21.6	38.9
2	Malignant neoplasms (C00-C97)	11,386	17.8	32.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,351	14.6	26.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,031	9.4	16.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,327	3.6	6.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	2,196	3.4	6.2
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,592	2.5	4.5
8	Cerebrovascular diseases (I60-I69)	1,557	2.4	4.4
9	Diabetes mellitus (E10-E14)	1,421	2.2	4.0
10	Chronic lower respiratory diseases (J40-J47)	644	1.0	1.8
11	Influenza and pneumonia (J10-J18)	635	1.0	1.8
12	Septicemia (A40-A41)	559	0.9	1.6
13	Viral hepatitis (B15-B19)	500	0.8	1.4
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	465	0.7	1.3
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	409	0.6	1.1
...	All other causes (Residual)	11,126	17.4	31.2
White, both sexes, 45-54 years				
...	All causes	135,719	100.0	394.0
1	Malignant neoplasms (C00-C97)	39,028	28.8	113.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	27,969	20.6	81.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14,030	10.3	40.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,420	4.7	18.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	6,348	4.7	18.4
6	Diabetes mellitus (E10-E14)	3,993	2.9	11.6
7	Cerebrovascular diseases (I60-I69)	3,943	2.9	11.4
8	Chronic lower respiratory diseases (J40-J47)	2,830	2.1	8.2
9	Human immunodeficiency virus (HIV) disease (B20-B24)	1,900	1.4	5.5
10	Viral hepatitis (B15-B19)	1,789	1.3	5.2
11	Septicemia (A40-A41)	1,515	1.1	4.4
12	Influenza and pneumonia (J10-J18)	1,399	1.0	4.1
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,186	0.9	3.4
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,170	0.9	3.4
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	599	0.4	1.7
...	All other causes (Residual)	21,600	15.9	62.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 55-64 years				
...	All causes	214,171	100.0	863.5
1	Malignant neoplasms (C00-C97)	81,073	37.9	326.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	50,449	23.6	203.4
3	Chronic lower respiratory diseases (J40-J47)	10,483	4.9	42.3
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,078	3.8	32.6
5	Diabetes mellitus (E10-E14)	8,030	3.7	32.4
6	Cerebrovascular diseases (I60-I69)	7,103	3.3	28.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	5,619	2.6	22.7
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,751	1.8	15.1
9	Septicemia (A40-A41)	2,727	1.3	11.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,552	1.2	10.3
11	Influenza and pneumonia (J10-J18)	2,510	1.2	10.1
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,093	0.5	4.4
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,079	0.5	4.4
14	Aortic aneurysm and dissection (I71)	1,060	0.5	4.3
15	Viral hepatitis (B15-B19)	934	0.4	3.8
...	All other causes (Residual)	27,630	12.9	111.4
White, both sexes, 65-74 years				
...	All causes	338,262	100.0	2,120.4
1	Malignant neoplasms (C00-C97)	120,167	35.5	753.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	83,145	24.6	521.2
3	Chronic lower respiratory diseases (J40-J47)	26,273	7.8	164.7
4	Cerebrovascular diseases (I60-I69)	15,799	4.7	99.0
5	Diabetes mellitus (E10-E14)	12,456	3.7	78.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,006	2.1	43.9
7	Influenza and pneumonia (J10-J18)	5,330	1.6	33.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,254	1.6	32.9
9	Septicemia (A40-A41)	4,649	1.4	29.1
10	Chronic liver disease and cirrhosis (K70,K73-K74)	4,613	1.4	28.9
11	Alzheimer's disease (G30)	3,263	1.0	20.5
12	Aortic aneurysm and dissection (I71)	2,637	0.8	16.5
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,148	0.6	13.5
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,137	0.6	13.4
15	Parkinson's disease (G20-G21)	2,075	0.6	13.0
...	All other causes (Residual)	41,310	12.2	259.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 75-84 years				
...	All causes	609,660	100.0	5,272.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	173,433	28.4	1,500.0
2	Malignant neoplasms (C00-C97)	148,497	24.4	1,284.3
3	Chronic lower respiratory diseases (J40-J47)	44,742	7.3	387.0
4	Cerebrovascular diseases (I60-I69)	43,672	7.2	377.7
5	Alzheimer's disease (G30)	20,130	3.3	174.1
6	Diabetes mellitus (E10-E14)	18,988	3.1	164.2
7	Influenza and pneumonia (J10-J18)	16,014	2.6	138.5
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,294	2.0	106.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11,739	1.9	101.5
10	Septicemia (A40-A41)	8,859	1.5	76.6
11	Parkinson's disease (G20-G21)	8,313	1.4	71.9
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	5,405	0.9	46.7
13	Pneumonitis due to solids and liquids (J69)	4,994	0.8	43.2
14	Aortic aneurysm and dissection (I71)	4,698	0.8	40.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,244	0.7	36.7
...	All other causes (Residual)	83,638	13.7	723.4
White, both sexes, 85 years and over				
...	All causes	613,740	100.0	14,010.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	218,052	35.5	4,977.6
2	Malignant neoplasms (C00-C97)	72,704	11.8	1,659.7
3	Cerebrovascular diseases (I60-I69)	55,101	9.0	1,257.8
4	Alzheimer's disease (G30)	37,094	6.0	846.8
5	Chronic lower respiratory diseases (J40-J47)	27,602	4.5	630.1
6	Influenza and pneumonia (J10-J18)	25,877	4.2	590.7
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,700	2.1	289.9
8	Diabetes mellitus (E10-E14)	12,677	2.1	289.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12,188	2.0	278.2
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	8,242	1.3	188.1
11	Septicemia (A40-A41)	7,804	1.3	178.1
12	Pneumonitis due to solids and liquids (J69)	7,352	1.2	167.8
13	Parkinson's disease (G20-G21)	6,267	1.0	143.1
14	Atherosclerosis (I70)	6,219	1.0	142.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3,515	0.6	80.2
...	All other causes (Residual)	100,346	16.3	2,290.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, male, all ages				
...	All causes	1,007,266	100.0	854.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	278,914	27.7	236.5
2	Malignant neoplasms (C00-C97)	247,450	24.6	209.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	61,267	6.1	52.0
4	Chronic lower respiratory diseases (J40-J47)	53,622	5.3	45.5
5	Cerebrovascular diseases (I60-I69)	49,296	4.9	41.8
6	Diabetes mellitus (E10-E14)	28,629	2.8	24.3
7	Influenza and pneumonia (J10-J18)	23,348	2.3	19.8
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	23,081	2.3	19.6
9	Alzheimer's disease (G30)	17,605	1.7	14.9
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16,349	1.6	13.9
11	Chronic liver disease and cirrhosis (K70,K73-K74)	15,334	1.5	13.0
12	Septicemia (A40-A41)	11,986	1.2	10.2
13	Parkinson's disease (G20-G21)	9,829	1.0	8.3
14	Pneumonitis due to solids and liquids (J69)	7,644	0.8	6.5
15	Aortic aneurysm and dissection (I71)	7,226	0.7	6.1
...	All other causes (Residual)	155,686	15.5	132.0
White, male, 1-4 years				
...	All causes	1,878	100.0	29.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	734	39.1	11.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	222	11.8	3.5
3	Malignant neoplasms (C00-C97)	161	8.6	2.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	113	6.0	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	61	3.2	1.0
6	Influenza and pneumonia (J10-J18)	32	1.7	0.5
7	Septicemia (A40-A41)	29	1.5	0.5
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	22	1.2	0.3
8	Certain conditions originating in the perinatal period (P00-P96)	22	1.2	0.3
10	Cerebrovascular diseases (I60-I69)	18	1.0	*
11	Chronic lower respiratory diseases (J40-J47)	15	0.8	*
12	Meningitis (G00,G03)	12	0.6	*
13	Meningococcal infection (A39)	10	0.5	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	8	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.4	*
...	All other causes (Residual)	412	21.9	6.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 5-14 years				
...	All causes	2,905	100.0	17.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,223	42.1	7.5
2	Malignant neoplasms (C00-C97)	453	15.6	2.8
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	162	5.6	1.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	143	4.9	0.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	109	3.8	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	82	2.8	0.5
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	33	1.1	0.2
8	Influenza and pneumonia (J10-J18)	32	1.1	0.2
9	Chronic lower respiratory diseases (J40-J47)	30	1.0	0.2
10	Cerebrovascular diseases (I60-I69)	29	1.0	0.2
11	Septicemia (A40-A41)	26	0.9	0.2
12	Certain conditions originating in the perinatal period (P00-P96)	16	0.6	*
13	Meningitis (G00,G03)	11	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	10	0.3	*
15	Meningococcal infection (A39)	8	0.3	*
15	Anemias (D50-D64)	8	0.3	*
15	Diseases of appendix (K35-K38)	8	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.3	*
...	All other causes (Residual)	514	17.7	3.2
White, male, 15-24 years				
...	All causes	18,306	100.0	108.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,770	53.4	57.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,016	16.5	17.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,727	9.4	10.2
4	Malignant neoplasms (C00-C97)	801	4.4	4.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	459	2.5	2.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	196	1.1	1.2
7	Influenza and pneumonia (J10-J18)	98	0.5	0.6
8	Cerebrovascular diseases (I60-I69)	82	0.4	0.5
9	Chronic lower respiratory diseases (J40-J47)	55	0.3	0.3
10	Legal intervention (Y35,Y89.0)	54	0.3	0.3
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	52	0.3	0.3
11	Diabetes mellitus (E10-E14)	52	0.3	0.3
13	Septicemia (A40-A41)	46	0.3	0.3
14	Human immunodeficiency virus (HIV) disease (B20-B24)	33	0.2	0.2
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	29	0.2	0.2
...	All other causes (Residual)	1,836	10.0	10.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 25-34 years				
...	All causes	20,584	100.0	127.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,180	39.7	50.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,526	17.1	21.8
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,501	7.3	9.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,431	7.0	8.8
5	Malignant neoplasms (C00-C97)	1,401	6.8	8.6
6	Human immunodeficiency virus (HIV) disease (B20-B24)	392	1.9	2.4
7	Diabetes mellitus (E10-E14)	227	1.1	1.4
8	Cerebrovascular diseases (I60-I69)	198	1.0	1.2
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	189	0.9	1.2
10	Chronic liver disease and cirrhosis (K70,K73-K74)	182	0.9	1.1
11	Influenza and pneumonia (J10-J18)	143	0.7	0.9
12	Septicemia (A40-A41)	101	0.5	0.6
13	Chronic lower respiratory diseases (J40-J47)	82	0.4	0.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	74	0.4	0.5
15	Legal intervention (Y35,Y89.0)	65	0.3	0.4
...	All other causes (Residual)	2,892	14.0	17.8
White, male, 35-44 years				
...	All causes	41,215	100.0	229.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,865	23.9	54.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,796	16.5	37.8
3	Malignant neoplasms (C00-C97)	5,068	12.3	28.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,606	11.2	25.6
5	Human immunodeficiency virus (HIV) disease (B20-B24)	1,777	4.3	9.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1,614	3.9	9.0
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,130	2.7	6.3
8	Diabetes mellitus (E10-E14)	905	2.2	5.0
9	Cerebrovascular diseases (I60-I69)	842	2.0	4.7
10	Influenza and pneumonia (J10-J18)	348	0.8	1.9
11	Viral hepatitis (B15-B19)	338	0.8	1.9
12	Chronic lower respiratory diseases (J40-J47)	312	0.8	1.7
13	Septicemia (A40-A41)	294	0.7	1.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	272	0.7	1.5
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	227	0.6	1.3
...	All other causes (Residual)	6,821	16.5	37.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 45-54 years				
...	All causes	86,266	100.0	504.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,911	24.2	122.2
2	Malignant neoplasms (C00-C97)	20,108	23.3	117.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,933	11.5	58.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,716	5.5	27.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	4,604	5.3	26.9
6	Diabetes mellitus (E10-E14)	2,508	2.9	14.7
7	Cerebrovascular diseases (I60-I69)	2,195	2.5	12.8
8	Human immunodeficiency virus (HIV) disease (B20-B24)	1,578	1.8	9.2
9	Chronic lower respiratory diseases (J40-J47)	1,462	1.7	8.5
10	Viral hepatitis (B15-B19)	1,336	1.5	7.8
11	Influenza and pneumonia (J10-J18)	846	1.0	4.9
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	839	1.0	4.9
13	Septicemia (A40-A41)	825	1.0	4.8
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	692	0.8	4.0
15	Aortic aneurysm and dissection (I71)	409	0.5	2.4
...	All other causes (Residual)	13,304	15.4	77.7
White, male, 55-64 years				
...	All causes	128,575	100.0	1,065.9
1	Malignant neoplasms (C00-C97)	44,020	34.2	364.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	35,512	27.6	294.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,438	4.2	45.1
4	Chronic lower respiratory diseases (J40-J47)	5,284	4.1	43.8
5	Diabetes mellitus (E10-E14)	4,728	3.7	39.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	3,950	3.1	32.7
7	Cerebrovascular diseases (I60-I69)	3,906	3.0	32.4
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,885	2.2	23.9
9	Influenza and pneumonia (J10-J18)	1,454	1.1	12.1
10	Septicemia (A40-A41)	1,411	1.1	11.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,411	1.1	11.7
12	Aortic aneurysm and dissection (I71)	789	0.6	6.5
13	Viral hepatitis (B15-B19)	644	0.5	5.3
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	640	0.5	5.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	631	0.5	5.2
...	All other causes (Residual)	15,872	12.3	131.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 65-74 years				
...	All causes	190,123	100.0	2,584.0
1	Malignant neoplasms (C00-C97)	65,946	34.7	896.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	51,740	27.2	703.2
3	Chronic lower respiratory diseases (J40-J47)	13,285	7.0	180.6
4	Cerebrovascular diseases (I60-I69)	8,149	4.3	110.8
5	Diabetes mellitus (E10-E14)	6,741	3.5	91.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,327	2.3	58.8
7	Influenza and pneumonia (J10-J18)	2,985	1.6	40.6
8	Chronic liver disease and cirrhosis (K70,K73-K74)	2,807	1.5	38.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,792	1.5	37.9
10	Septicemia (A40-A41)	2,370	1.2	32.2
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,783	0.9	24.2
12	Aortic aneurysm and dissection (I71)	1,770	0.9	24.1
13	Alzheimer's disease (G30)	1,400	0.7	19.0
14	Parkinson's disease (G20-G21)	1,367	0.7	18.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,186	0.6	16.1
...	All other causes (Residual)	21,475	11.3	291.9
White, male, 75-84 years				
...	All causes	299,342	100.0	6,384.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	88,944	29.7	1,897.1
2	Malignant neoplasms (C00-C97)	77,484	25.9	1,652.7
3	Chronic lower respiratory diseases (J40-J47)	21,681	7.2	462.4
4	Cerebrovascular diseases (I60-I69)	18,457	6.2	393.7
5	Diabetes mellitus (E10-E14)	9,061	3.0	193.3
6	Influenza and pneumonia (J10-J18)	7,979	2.7	170.2
7	Alzheimer's disease (G30)	7,208	2.4	153.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,423	2.1	137.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,029	2.0	128.6
10	Parkinson's disease (G20-G21)	5,058	1.7	107.9
11	Septicemia (A40-A41)	4,105	1.4	87.6
12	Pneumonitis due to solids and liquids (J69)	2,755	0.9	58.8
13	Aortic aneurysm and dissection (I71)	2,652	0.9	56.6
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,210	0.7	47.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,192	0.7	46.8
...	All other causes (Residual)	37,104	12.4	791.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 85 years and over				
...	All causes	207,589	100.0	15,250.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	72,801	35.1	5,348.4
2	Malignant neoplasms (C00-C97)	31,973	15.4	2,348.9
3	Cerebrovascular diseases (I60-I69)	15,372	7.4	1,129.3
4	Chronic lower respiratory diseases (J40-J47)	11,402	5.5	837.7
5	Influenza and pneumonia (J10-J18)	9,331	4.5	685.5
6	Alzheimer's disease (G30)	8,724	4.2	640.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,992	2.4	366.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,892	2.4	359.4
9	Diabetes mellitus (E10-E14)	4,396	2.1	323.0
10	Pneumonitis due to solids and liquids (J69)	3,290	1.6	241.7
11	Parkinson's disease (G20-G21)	3,148	1.5	231.3
12	Septicemia (A40-A41)	2,675	1.3	196.5
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,170	1.0	159.4
14	Atherosclerosis (I70)	1,829	0.9	134.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,405	0.7	103.2
...	All other causes (Residual)	29,189	14.1	2,144.4
White, female, all ages				
...	All causes	1,049,377	100.0	871.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	286,789	27.3	238.3
2	Malignant neoplasms (C00-C97)	230,684	22.0	191.7
3	Cerebrovascular diseases (I60-I69)	78,572	7.5	65.3
4	Chronic lower respiratory diseases (J40-J47)	59,292	5.7	49.3
5	Alzheimer's disease (G30)	43,482	4.1	36.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	34,623	3.3	28.8
7	Diabetes mellitus (E10-E14)	29,458	2.8	24.5
8	Influenza and pneumonia (J10-J18)	29,082	2.8	24.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17,342	1.7	14.4
10	Septicemia (A40-A41)	14,692	1.4	12.2
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	11,053	1.1	9.2
12	Chronic liver disease and cirrhosis (K70,K73-K74)	8,228	0.8	6.8
13	Pneumonitis due to solids and liquids (J69)	7,324	0.7	6.1
14	Parkinson's disease (G20-G21)	7,212	0.7	6.0
15	Atherosclerosis (I70)	6,731	0.6	5.6
...	All other causes (Residual)	184,813	17.6	153.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 1-4 years				
...	All causes	1,485	100.0	24.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	508	34.2	8.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	178	12.0	2.9
3	Malignant neoplasms (C00-C97)	150	10.1	2.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	88	5.9	1.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	55	3.7	0.9
6	Influenza and pneumonia (J10-J18)	45	3.0	0.7
7	Septicemia (A40-A41)	31	2.1	0.5
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	21	1.4	0.3
9	Certain conditions originating in the perinatal period (P00-P96)	11	0.7	*
10	Cerebrovascular diseases (I60-I69)	10	0.7	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	8	0.5	*
12	Anemias (D50-D64)	7	0.5	*
12	Meningitis (G00,G03)	7	0.5	*
12	Chronic lower respiratory diseases (J40-J47)	7	0.5	*
15	Meningococcal infection (A39)	5	0.3	*
15	Acute bronchitis and bronchiolitis (J20-J21)	5	0.3	*
...	All other causes (Residual)	349	23.5	5.7
White, female, 5-14 years				
...	All causes	1,995	100.0	13.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	730	36.6	4.7
2	Malignant neoplasms (C00-C97)	340	17.0	2.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	125	6.3	0.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	77	3.9	0.5
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	75	3.8	0.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	75	3.8	0.5
7	Cerebrovascular diseases (I60-I69)	29	1.5	0.2
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	26	1.3	0.2
9	Influenza and pneumonia (J10-J18)	25	1.3	0.2
10	Septicemia (A40-A41)	24	1.2	0.2
11	Chronic lower respiratory diseases (J40-J47)	23	1.2	0.1
12	Anemias (D50-D64)	12	0.6	*
12	Diabetes mellitus (E10-E14)	12	0.6	*
14	Certain conditions originating in the perinatal period (P00-P96)	10	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.5	*
...	All other causes (Residual)	403	20.2	2.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 15-24 years				
...	All causes	6,707	100.0	42.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,404	50.8	21.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	594	8.9	3.8
3	Malignant neoplasms (C00-C97)	522	7.8	3.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	399	5.9	2.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	219	3.3	1.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	156	2.3	1.0
7	Pregnancy, childbirth and the puerperium (O00-099)	88	1.3	0.6
8	Cerebrovascular diseases (I60-I69)	70	1.0	0.4
9	Diabetes mellitus (E10-E14)	47	0.7	0.3
10	Influenza and pneumonia (J10-J18)	44	0.7	0.3
11	Septicemia (A40-A41)	36	0.5	0.2
11	Chronic lower respiratory diseases (J40-J47)	36	0.5	0.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	34	0.5	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	24	0.4	0.2
15	Complications of medical and surgical care (Y40-Y84,Y88)	17	0.3	*
...	All other causes (Residual)	1,017	15.2	6.4
White, female, 25-34 years				
...	All causes	8,691	100.0	56.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,607	30.0	17.0
2	Malignant neoplasms (C00-C97)	1,382	15.9	9.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	762	8.8	5.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	609	7.0	4.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	419	4.8	2.7
6	Cerebrovascular diseases (I60-I69)	175	2.0	1.1
7	Diabetes mellitus (E10-E14)	169	1.9	1.1
8	Pregnancy, childbirth and the puerperium (O00-099)	168	1.9	1.1
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	156	1.8	1.0
10	Human immunodeficiency virus (HIV) disease (B20-B24)	111	1.3	0.7
11	Septicemia (A40-A41)	105	1.2	0.7
12	Influenza and pneumonia (J10-J18)	85	1.0	0.6
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	73	0.8	0.5
14	Chronic liver disease and cirrhosis (K70,K73-K74)	69	0.8	0.5
15	Chronic lower respiratory diseases (J40-J47)	68	0.8	0.4
...	All other causes (Residual)	1,733	19.9	11.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 35-44 years				
...	All causes	22,825	100.0	129.5
1	Malignant neoplasms (C00-C97)	6,318	27.7	35.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,976	17.4	22.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,555	11.2	14.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,425	6.2	8.1
5	Cerebrovascular diseases (I60-I69)	715	3.1	4.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	713	3.1	4.0
7	Diabetes mellitus (E10-E14)	516	2.3	2.9
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	462	2.0	2.6
9	Human immunodeficiency virus (HIV) disease (B20-B24)	419	1.8	2.4
10	Chronic lower respiratory diseases (J40-J47)	332	1.5	1.9
11	Influenza and pneumonia (J10-J18)	287	1.3	1.6
12	Septicemia (A40-A41)	265	1.2	1.5
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	193	0.8	1.1
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	182	0.8	1.0
15	Viral hepatitis (B15-B19)	162	0.7	0.9
...	All other causes (Residual)	4,305	18.9	24.4
White, female, 45-54 years				
...	All causes	49,453	100.0	285.4
1	Malignant neoplasms (C00-C97)	18,920	38.3	109.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,058	14.3	40.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,097	8.3	23.6
4	Cerebrovascular diseases (I60-I69)	1,748	3.5	10.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,744	3.5	10.1
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,704	3.4	9.8
7	Diabetes mellitus (E10-E14)	1,485	3.0	8.6
8	Chronic lower respiratory diseases (J40-J47)	1,368	2.8	7.9
9	Septicemia (A40-A41)	690	1.4	4.0
10	Influenza and pneumonia (J10-J18)	553	1.1	3.2
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	494	1.0	2.9
12	Viral hepatitis (B15-B19)	453	0.9	2.6
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	331	0.7	1.9
14	Human immunodeficiency virus (HIV) disease (B20-B24)	322	0.7	1.9
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	269	0.5	1.6
...	All other causes (Residual)	8,217	16.6	47.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 55-64 years				
...	All causes	85,596	100.0	671.8
1	Malignant neoplasms (C00-C97)	37,053	43.3	290.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,937	17.5	117.2
3	Chronic lower respiratory diseases (J40-J47)	5,199	6.1	40.8
4	Diabetes mellitus (E10-E14)	3,302	3.9	25.9
5	Cerebrovascular diseases (I60-I69)	3,197	3.7	25.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,640	3.1	20.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,669	1.9	13.1
8	Septicemia (A40-A41)	1,316	1.5	10.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,141	1.3	9.0
10	Influenza and pneumonia (J10-J18)	1,056	1.2	8.3
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	866	1.0	6.8
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	462	0.5	3.6
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	439	0.5	3.4
14	Viral hepatitis (B15-B19)	290	0.3	2.3
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	286	0.3	2.2
...	All other causes (Residual)	11,743	13.7	92.2
White, female, 65-74 years				
...	All causes	148,139	100.0	1,723.6
1	Malignant neoplasms (C00-C97)	54,221	36.6	630.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	31,405	21.2	365.4
3	Chronic lower respiratory diseases (J40-J47)	12,988	8.8	151.1
4	Cerebrovascular diseases (I60-I69)	7,650	5.2	89.0
5	Diabetes mellitus (E10-E14)	5,715	3.9	66.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,679	1.8	31.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,462	1.7	28.6
8	Influenza and pneumonia (J10-J18)	2,345	1.6	27.3
9	Septicemia (A40-A41)	2,279	1.5	26.5
10	Alzheimer's disease (G30)	1,863	1.3	21.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1,806	1.2	21.0
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,075	0.7	12.5
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	879	0.6	10.2
14	Aortic aneurysm and dissection (I71)	867	0.6	10.1
15	Parkinson's disease (G20-G21)	708	0.5	8.2
...	All other causes (Residual)	19,197	13.0	223.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 75-84 years				
...	All causes	310,318	100.0	4,514.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	84,489	27.2	1,229.1
2	Malignant neoplasms (C00-C97)	71,013	22.9	1,033.1
3	Cerebrovascular diseases (I60-I69)	25,215	8.1	366.8
4	Chronic lower respiratory diseases (J40-J47)	23,061	7.4	335.5
5	Alzheimer's disease (G30)	12,922	4.2	188.0
6	Diabetes mellitus (E10-E14)	9,927	3.2	144.4
7	Influenza and pneumonia (J10-J18)	8,035	2.6	116.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,871	1.9	85.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,710	1.8	83.1
10	Septicemia (A40-A41)	4,754	1.5	69.2
11	Parkinson's disease (G20-G21)	3,255	1.0	47.4
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	3,195	1.0	46.5
13	Pneumonitis due to solids and liquids (J69)	2,239	0.7	32.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,052	0.7	29.9
15	Aortic aneurysm and dissection (I71)	2,046	0.7	29.8
...	All other causes (Residual)	46,534	15.0	677.0
White, female, 85 years and over				
...	All causes	406,151	100.0	13,450.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	145,251	35.8	4,810.4
2	Malignant neoplasms (C00-C97)	40,731	10.0	1,348.9
3	Cerebrovascular diseases (I60-I69)	39,729	9.8	1,315.7
4	Alzheimer's disease (G30)	28,370	7.0	939.6
5	Influenza and pneumonia (J10-J18)	16,546	4.1	548.0
6	Chronic lower respiratory diseases (J40-J47)	16,200	4.0	536.5
7	Diabetes mellitus (E10-E14)	8,281	2.0	274.3
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,808	1.9	258.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,196	1.8	238.3
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	6,072	1.5	201.1
11	Septicemia (A40-A41)	5,129	1.3	169.9
12	Atherosclerosis (I70)	4,390	1.1	145.4
13	Pneumonitis due to solids and liquids (J69)	4,062	1.0	134.5
14	Parkinson's disease (G20-G21)	3,119	0.8	103.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,110	0.5	69.9
...	All other causes (Residual)	71,157	17.5	2,356.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, all ages				
...	All causes	287,315	100.0	744.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	74,225	25.8	192.3
2	Malignant neoplasms (C00-C97)	62,499	21.8	161.9
3	Cerebrovascular diseases (I60-I69)	18,118	6.3	46.9
4	Diabetes mellitus (E10-E14)	12,834	4.5	33.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,670	4.4	32.8
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8,135	2.8	21.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,834	2.7	20.3
8	Chronic lower respiratory diseases (J40-J47)	7,400	2.6	19.2
9	Human immunodeficiency virus (HIV) disease (B20-B24)	7,271	2.5	18.8
10	Septicemia (A40-A41)	6,010	2.1	15.6
11	Influenza and pneumonia (J10-J18)	5,700	2.0	14.8
12	Certain conditions originating in the perinatal period (P00-P96)	4,920	1.7	12.7
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	4,747	1.7	12.3
14	Alzheimer's disease (G30)	4,232	1.5	11.0
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,534	0.9	6.6
...	All other causes (Residual)	48,186	16.8	124.8
Black, both sexes, 1-4 years				
...	All causes	1,164	100.0	44.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	324	27.8	12.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	163	14.0	6.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	132	11.3	5.1
4	Malignant neoplasms (C00-C97)	65	5.6	2.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	62	5.3	2.4
6	Influenza and pneumonia (J10-J18)	35	3.0	1.3
7	Certain conditions originating in the perinatal period (P00-P96)	23	2.0	0.9
8	Chronic lower respiratory diseases (J40-J47)	22	1.9	0.8
9	Septicemia (A40-A41)	19	1.6	*
9	Anemias (D50-D64)	19	1.6	*
11	Cerebrovascular diseases (I60-I69)	12	1.0	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.7	*
13	Acute bronchitis and bronchiolitis (J20-J21)	6	0.5	*
14	Meningitis (G00,G03)	5	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.3	*
...	All other causes (Residual)	265	22.8	10.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 5-14 years				
...	All causes	1,584	100.0	23.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	570	36.0	8.5
2	Malignant neoplasms (C00-C97)	170	10.7	2.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	131	8.3	1.9
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	85	5.4	1.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	80	5.1	1.2
6	Chronic lower respiratory diseases (J40-J47)	64	4.0	1.0
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	49	3.1	0.7
8	Anemias (D50-D64)	23	1.5	0.3
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	21	1.3	0.3
10	Influenza and pneumonia (J10-J18)	20	1.3	0.3
11	Human immunodeficiency virus (HIV) disease (B20-B24)	18	1.1	*
12	Cerebrovascular diseases (I60-I69)	17	1.1	*
13	Diabetes mellitus (E10-E14)	16	1.0	*
14	Septicemia (A40-A41)	12	0.8	*
15	Certain conditions originating in the perinatal period (P00-P96)	9	0.6	*
...	All other causes (Residual)	299	18.9	4.4
Black, both sexes, 15-24 years				
...	All causes	7,049	100.0	109.4
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,803	39.8	43.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,661	23.6	25.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	465	6.6	7.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	316	4.5	4.9
5	Malignant neoplasms (C00-C97)	311	4.4	4.8
6	Human immunodeficiency virus (HIV) disease (B20-B24)	142	2.0	2.2
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	111	1.6	1.7
8	Anemias (D50-D64)	93	1.3	1.4
9	Chronic lower respiratory diseases (J40-J47)	79	1.1	1.2
10	Pregnancy, childbirth and the puerperium (O00-099)	74	1.0	1.1
11	Diabetes mellitus (E10-E14)	61	0.9	0.9
12	Cerebrovascular diseases (I60-I69)	49	0.7	0.8
13	Septicemia (A40-A41)	46	0.7	0.7
14	Influenza and pneumonia (J10-J18)	40	0.6	0.6
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	31	0.4	0.5
...	All other causes (Residual)	767	10.9	11.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 25-34 years				
...	All causes	9,925	100.0	179.3
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,445	24.6	44.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,740	17.5	31.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	980	9.9	17.7
4	Human immunodeficiency virus (HIV) disease (B20-B24)	944	9.5	17.1
5	Malignant neoplasms (C00-C97)	667	6.7	12.0
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	521	5.2	9.4
7	Diabetes mellitus (E10-E14)	183	1.8	3.3
8	Cerebrovascular diseases (I60-I69)	173	1.7	3.1
9	Anemias (D50-D64)	129	1.3	2.3
10	Pregnancy, childbirth and the puerperium (O00-099)	109	1.1	2.0
11	Septicemia (A40-A41)	104	1.0	1.9
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	103	1.0	1.9
13	Chronic lower respiratory diseases (J40-J47)	101	1.0	1.8
14	Influenza and pneumonia (J10-J18)	66	0.7	1.2
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	60	0.6	1.1
...	All other causes (Residual)	1,600	16.1	28.9
Black, both sexes, 35-44 years				
...	All causes	18,474	100.0	322.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,236	17.5	56.5
2	Malignant neoplasms (C00-C97)	2,682	14.5	46.8
3	Human immunodeficiency virus (HIV) disease (B20-B24)	2,554	13.8	44.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,104	11.4	36.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,297	7.0	22.6
6	Cerebrovascular diseases (I60-I69)	703	3.8	12.3
7	Diabetes mellitus (E10-E14)	547	3.0	9.5
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	381	2.1	6.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	317	1.7	5.5
10	Chronic liver disease and cirrhosis (K70,K73-K74)	310	1.7	5.4
11	Septicemia (A40-A41)	257	1.4	4.5
12	Influenza and pneumonia (J10-J18)	229	1.2	4.0
13	Chronic lower respiratory diseases (J40-J47)	225	1.2	3.9
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	178	1.0	3.1
15	Anemias (D50-D64)	140	0.8	2.4
...	All other causes (Residual)	3,314	17.9	57.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 45-54 years				
...	All causes	36,710	100.0	744.8
1	Malignant neoplasms (C00-C97)	8,753	23.8	177.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,680	23.6	176.1
3	Human immunodeficiency virus (HIV) disease (B20-B24)	2,464	6.7	50.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,455	6.7	49.8
5	Cerebrovascular diseases (I60-I69)	1,920	5.2	39.0
6	Diabetes mellitus (E10-E14)	1,385	3.8	28.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	903	2.5	18.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	820	2.2	16.6
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	752	2.0	15.3
10	Septicemia (A40-A41)	664	1.8	13.5
11	Chronic lower respiratory diseases (J40-J47)	622	1.7	12.6
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	501	1.4	10.2
13	Influenza and pneumonia (J10-J18)	447	1.2	9.1
14	Viral hepatitis (B15-B19)	369	1.0	7.5
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	288	0.8	5.8
...	All other causes (Residual)	5,687	15.5	115.4
Black, both sexes, 55-64 years				
...	All causes	43,490	100.0	1,499.6
1	Malignant neoplasms (C00-C97)	13,296	30.6	458.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,679	26.9	402.7
3	Cerebrovascular diseases (I60-I69)	2,433	5.6	83.9
4	Diabetes mellitus (E10-E14)	2,334	5.4	80.5
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,291	3.0	44.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,268	2.9	43.7
7	Chronic lower respiratory diseases (J40-J47)	1,106	2.5	38.1
8	Septicemia (A40-A41)	916	2.1	31.6
9	Human immunodeficiency virus (HIV) disease (B20-B24)	860	2.0	29.7
10	Chronic liver disease and cirrhosis (K70,K73-K74)	733	1.7	25.3
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	693	1.6	23.9
12	Influenza and pneumonia (J10-J18)	552	1.3	19.0
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	254	0.6	8.8
14	Viral hepatitis (B15-B19)	251	0.6	8.7
15	Aortic aneurysm and dissection (I71)	219	0.5	7.6
...	All other causes (Residual)	5,605	12.9	193.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 65-74 years				
...	All causes	51,682	100.0	2,982.6
1	Malignant neoplasms (C00-C97)	15,871	30.7	915.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,552	28.2	839.8
3	Cerebrovascular diseases (I60-I69)	3,384	6.5	195.3
4	Diabetes mellitus (E10-E14)	3,107	6.0	179.3
5	Chronic lower respiratory diseases (J40-J47)	1,770	3.4	102.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,653	3.2	95.4
7	Septicemia (A40-A41)	1,211	2.3	69.9
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	899	1.7	51.9
9	Influenza and pneumonia (J10-J18)	863	1.7	49.8
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	835	1.6	48.2
11	Chronic liver disease and cirrhosis (K70,K73-K74)	374	0.7	21.6
12	Alzheimer's disease (G30)	334	0.6	19.3
13	Pneumonitis due to solids and liquids (J69)	241	0.5	13.9
14	Human immunodeficiency virus (HIV) disease (B20-B24)	235	0.5	13.6
15	Aortic aneurysm and dissection (I71)	226	0.4	13.0
...	All other causes (Residual)	6,127	11.9	353.6
Black, both sexes, 75-84 years				
...	All causes	61,379	100.0	6,170.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	18,344	29.9	1,844.1
2	Malignant neoplasms (C00-C97)	14,384	23.4	1,446.0
3	Cerebrovascular diseases (I60-I69)	5,090	8.3	511.7
4	Diabetes mellitus (E10-E14)	3,310	5.4	332.8
5	Chronic lower respiratory diseases (J40-J47)	2,216	3.6	222.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,017	3.3	202.8
7	Influenza and pneumonia (J10-J18)	1,564	2.5	157.2
8	Septicemia (A40-A41)	1,548	2.5	155.6
9	Alzheimer's disease (G30)	1,523	2.5	153.1
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,220	2.0	122.6
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	866	1.4	87.1
12	Pneumonitis due to solids and liquids (J69)	467	0.8	46.9
13	Aortic aneurysm and dissection (I71)	318	0.5	32.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	313	0.5	31.5
15	Parkinson's disease (G20-G21)	290	0.5	29.2
...	All other causes (Residual)	7,909	12.9	795.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 85 years and over				
...	All causes	47,300	100.0	13,338.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	16,184	34.2	4,563.8
2	Malignant neoplasms (C00-C97)	6,291	13.3	1,774.0
3	Cerebrovascular diseases (I60-I69)	4,295	9.1	1,211.2
4	Alzheimer's disease (G30)	2,333	4.9	657.9
5	Diabetes mellitus (E10-E14)	1,888	4.0	532.4
6	Influenza and pneumonia (J10-J18)	1,791	3.8	505.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,528	3.2	430.9
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,190	2.5	335.6
9	Chronic lower respiratory diseases (J40-J47)	1,184	2.5	333.9
10	Septicemia (A40-A41)	1,136	2.4	320.3
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	540	1.1	152.3
12	Pneumonitis due to solids and liquids (J69)	513	1.1	144.7
13	Atherosclerosis (I70)	365	0.8	102.9
14	Parkinson's disease (G20-G21)	217	0.5	61.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	202	0.4	57.0
...	All other causes (Residual)	7,643	16.2	2,155.3
Black, male, all ages				
...	All causes	145,970	100.0	792.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	36,222	24.8	196.7
2	Malignant neoplasms (C00-C97)	32,443	22.2	176.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,603	5.9	46.7
4	Cerebrovascular diseases (I60-I69)	7,644	5.2	41.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6,839	4.7	37.1
6	Diabetes mellitus (E10-E14)	5,565	3.8	30.2
7	Human immunodeficiency virus (HIV) disease (B20-B24)	4,769	3.3	25.9
8	Chronic lower respiratory diseases (J40-J47)	4,046	2.8	22.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,537	2.4	19.2
10	Influenza and pneumonia (J10-J18)	2,775	1.9	15.1
11	Certain conditions originating in the perinatal period (P00-P96)	2,753	1.9	14.9
12	Septicemia (A40-A41)	2,690	1.8	14.6
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,964	1.3	10.7
14	Chronic liver disease and cirrhosis (K70,K73-K74)	1,670	1.1	9.1
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,655	1.1	9.0
...	All other causes (Residual)	22,795	15.6	123.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 1-4 years				
...	All causes	641	100.0	48.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	188	29.3	14.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	83	12.9	6.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	77	12.0	5.8
4	Malignant neoplasms (C00-C97)	37	5.8	2.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	32	5.0	2.4
6	Chronic lower respiratory diseases (J40-J47)	14	2.2	*
7	Certain conditions originating in the perinatal period (P00-P96)	13	2.0	*
8	Anemias (D50-D64)	12	1.9	*
8	Influenza and pneumonia (J10-J18)	12	1.9	*
10	Septicemia (A40-A41)	9	1.4	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.8	*
11	Cerebrovascular diseases (I60-I69)	5	0.8	*
13	Acute bronchitis and bronchiolitis (J20-J21)	4	0.6	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.5	*
15	Meningitis (G00,G03)	2	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.3	*
...	All other causes (Residual)	143	22.3	10.8
Black, male, 5-14 years				
...	All causes	888	100.0	26.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	351	39.5	10.3
2	Malignant neoplasms (C00-C97)	84	9.5	2.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	82	9.2	2.4
4	Chronic lower respiratory diseases (J40-J47)	44	5.0	1.3
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	39	4.4	1.1
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	37	4.2	1.1
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	30	3.4	0.9
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	1.4	*
8	Anemias (D50-D64)	12	1.4	*
8	Influenza and pneumonia (J10-J18)	12	1.4	*
11	Cerebrovascular diseases (I60-I69)	10	1.1	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	9	1.0	*
13	Diabetes mellitus (E10-E14)	8	0.9	*
14	Certain conditions originating in the perinatal period (P00-P96)	5	0.6	*
15	Septicemia (A40-A41)	4	0.5	*
...	All other causes (Residual)	149	16.8	4.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 15-24 years				
...	All causes	5,336	100.0	164.3
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,522	47.3	77.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,228	23.0	37.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	396	7.4	12.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	207	3.9	6.4
5	Malignant neoplasms (C00-C97)	176	3.3	5.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	73	1.4	2.2
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	73	1.4	2.2
8	Anemias (D50-D64)	50	0.9	1.5
9	Chronic lower respiratory diseases (J40-J47)	48	0.9	1.5
10	Diabetes mellitus (E10-E14)	31	0.6	1.0
10	Cerebrovascular diseases (I60-I69)	31	0.6	1.0
12	Septicemia (A40-A41)	22	0.4	0.7
13	Legal intervention (Y35,Y89.0)	20	0.4	0.6
14	Influenza and pneumonia (J10-J18)	19	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19	0.4	*
...	All other causes (Residual)	421	7.9	13.0
Black, male, 25-34 years				
...	All causes	6,684	100.0	252.3
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,163	32.4	81.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,333	19.9	50.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	597	8.9	22.5
4	Human immunodeficiency virus (HIV) disease (B20-B24)	508	7.6	19.2
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	429	6.4	16.2
6	Malignant neoplasms (C00-C97)	265	4.0	10.0
7	Diabetes mellitus (E10-E14)	104	1.6	3.9
8	Cerebrovascular diseases (I60-I69)	77	1.2	2.9
9	Anemias (D50-D64)	71	1.1	2.7
10	Chronic lower respiratory diseases (J40-J47)	66	1.0	2.5
11	Septicemia (A40-A41)	53	0.8	2.0
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	51	0.8	1.9
13	Influenza and pneumonia (J10-J18)	44	0.7	1.7
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	37	0.6	1.4
15	Legal intervention (Y35,Y89.0)	34	0.5	1.3
...	All other causes (Residual)	852	12.7	32.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 35-44 years				
...	All causes	10,700	100.0	397.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,961	18.3	72.8
2	Human immunodeficiency virus (HIV) disease (B20-B24)	1,624	15.2	60.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,457	13.6	54.1
4	Malignant neoplasms (C00-C97)	1,035	9.7	38.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,022	9.6	37.9
6	Cerebrovascular diseases (I60-I69)	345	3.2	12.8
7	Diabetes mellitus (E10-E14)	317	3.0	11.8
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	304	2.8	11.3
9	Chronic liver disease and cirrhosis (K70,K73-K74)	199	1.9	7.4
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	167	1.6	6.2
11	Influenza and pneumonia (J10-J18)	130	1.2	4.8
12	Septicemia (A40-A41)	126	1.2	4.7
13	Chronic lower respiratory diseases (J40-J47)	108	1.0	4.0
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	104	1.0	3.9
15	Anemias (D50-D64)	74	0.7	2.7
...	All other causes (Residual)	1,727	16.1	64.1
Black, male, 45-54 years				
...	All causes	21,756	100.0	954.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,468	25.1	240.0
2	Malignant neoplasms (C00-C97)	4,489	20.6	197.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,747	8.0	76.7
4	Human immunodeficiency virus (HIV) disease (B20-B24)	1,706	7.8	74.9
5	Cerebrovascular diseases (I60-I69)	1,021	4.7	44.8
6	Diabetes mellitus (E10-E14)	754	3.5	33.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	618	2.8	27.1
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	592	2.7	26.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	478	2.2	21.0
10	Septicemia (A40-A41)	345	1.6	15.1
11	Chronic lower respiratory diseases (J40-J47)	317	1.5	13.9
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	293	1.3	12.9
13	Influenza and pneumonia (J10-J18)	268	1.2	11.8
14	Viral hepatitis (B15-B19)	267	1.2	11.7
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	221	1.0	9.7
...	All other causes (Residual)	3,172	14.6	139.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 55-64 years				
...	All causes	25,352	100.0	1,960.8
1	Malignant neoplasms (C00-C97)	7,359	29.0	569.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,243	28.6	560.2
3	Cerebrovascular diseases (I60-I69)	1,388	5.5	107.4
4	Diabetes mellitus (E10-E14)	1,184	4.7	91.6
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	912	3.6	70.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	650	2.6	50.3
7	Chronic lower respiratory diseases (J40-J47)	648	2.6	50.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	644	2.5	49.8
9	Chronic liver disease and cirrhosis (K70,K73-K74)	505	2.0	39.1
10	Septicemia (A40-A41)	468	1.8	36.2
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	366	1.4	28.3
12	Influenza and pneumonia (J10-J18)	322	1.3	24.9
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	195	0.8	15.1
14	Viral hepatitis (B15-B19)	179	0.7	13.8
15	Aortic aneurysm and dissection (I71)	152	0.6	11.8
...	All other causes (Residual)	3,137	12.4	242.6
Black, male, 65-74 years				
...	All causes	27,555	100.0	3,818.3
1	Malignant neoplasms (C00-C97)	8,730	31.7	1,209.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,914	28.7	1,096.6
3	Cerebrovascular diseases (I60-I69)	1,697	6.2	235.2
4	Diabetes mellitus (E10-E14)	1,432	5.2	198.4
5	Chronic lower respiratory diseases (J40-J47)	1,013	3.7	140.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	780	2.8	108.1
7	Septicemia (A40-A41)	585	2.1	81.1
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	532	1.9	73.7
9	Influenza and pneumonia (J10-J18)	523	1.9	72.5
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	410	1.5	56.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	244	0.9	33.8
12	Human immunodeficiency virus (HIV) disease (B20-B24)	169	0.6	23.4
13	Pneumonitis due to solids and liquids (J69)	153	0.6	21.2
14	Alzheimer's disease (G30)	138	0.5	19.1
15	Aortic aneurysm and dissection (I71)	131	0.5	18.2
...	All other causes (Residual)	3,104	11.3	430.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 75-84 years				
...	All causes	27,695	100.0	7,710.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,030	29.0	2,235.5
2	Malignant neoplasms (C00-C97)	7,497	27.1	2,087.2
3	Cerebrovascular diseases (I60-I69)	1,979	7.1	551.0
4	Diabetes mellitus (E10-E14)	1,224	4.4	340.8
5	Chronic lower respiratory diseases (J40-J47)	1,208	4.4	336.3
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	836	3.0	232.7
7	Influenza and pneumonia (J10-J18)	788	2.8	219.4
8	Septicemia (A40-A41)	679	2.5	189.0
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	492	1.8	137.0
10	Alzheimer's disease (G30)	491	1.8	136.7
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	474	1.7	132.0
12	Pneumonitis due to solids and liquids (J69)	243	0.9	67.7
13	Parkinson's disease (G20-G21)	156	0.6	43.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	141	0.5	39.3
15	Aortic aneurysm and dissection (I71)	132	0.5	36.7
...	All other causes (Residual)	3,325	12.0	925.7
Black, male, 85 years and over				
...	All causes	14,548	100.0	14,452.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,668	32.1	4,637.3
2	Malignant neoplasms (C00-C97)	2,767	19.0	2,748.8
3	Cerebrovascular diseases (I60-I69)	1,068	7.3	1,061.0
4	Influenza and pneumonia (J10-J18)	602	4.1	598.0
5	Chronic lower respiratory diseases (J40-J47)	570	3.9	566.3
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	516	3.5	512.6
7	Diabetes mellitus (E10-E14)	509	3.5	505.7
8	Alzheimer's disease (G30)	501	3.4	497.7
9	Septicemia (A40-A41)	344	2.4	341.7
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	265	1.8	263.3
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	215	1.5	213.6
12	Pneumonitis due to solids and liquids (J69)	204	1.4	202.7
13	Parkinson's disease (G20-G21)	107	0.7	106.3
14	Atherosclerosis (I70)	79	0.5	78.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	62	0.4	61.6
...	All other causes (Residual)	2,071	14.2	2,057.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, all ages				
...	All causes	141,345	100.0	700.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	38,003	26.9	188.3
2	Malignant neoplasms (C00-C97)	30,056	21.3	148.9
3	Cerebrovascular diseases (I60-I69)	10,474	7.4	51.9
4	Diabetes mellitus (E10-E14)	7,269	5.1	36.0
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,297	3.0	21.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,067	2.9	20.1
7	Chronic lower respiratory diseases (J40-J47)	3,354	2.4	16.6
8	Septicemia (A40-A41)	3,320	2.3	16.4
9	Alzheimer's disease (G30)	3,085	2.2	15.3
10	Influenza and pneumonia (J10-J18)	2,925	2.1	14.5
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,783	2.0	13.8
12	Human immunodeficiency virus (HIV) disease (B20-B24)	2,502	1.8	12.4
13	Certain conditions originating in the perinatal period (P00-P96)	2,167	1.5	10.7
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,296	0.9	6.4
15	Chronic liver disease and cirrhosis (K70,K73-K74)	864	0.6	4.3
...	All other causes (Residual)	24,883	17.6	123.3
Black, female, 1-4 years				
...	All causes	523	100.0	40.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	136	26.0	10.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	80	15.3	6.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	55	10.5	4.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	30	5.7	2.3
5	Malignant neoplasms (C00-C97)	28	5.4	2.2
6	Influenza and pneumonia (J10-J18)	23	4.4	1.8
7	Septicemia (A40-A41)	10	1.9	*
7	Certain conditions originating in the perinatal period (P00-P96)	10	1.9	*
9	Chronic lower respiratory diseases (J40-J47)	8	1.5	*
10	Anemias (D50-D64)	7	1.3	*
10	Cerebrovascular diseases (I60-I69)	7	1.3	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.6	*
12	Meningitis (G00,G03)	3	0.6	*
14	Meningococcal infection (A39)	2	0.4	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.4	*
14	Acute bronchitis and bronchiolitis (J20-J21)	2	0.4	*
...	All other causes (Residual)	117	22.4	9.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 5-14 years				
...	All causes	696	100.0	21.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	219	31.5	6.6
2	Malignant neoplasms (C00-C97)	86	12.4	2.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	49	7.0	1.5
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	46	6.6	1.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	43	6.2	1.3
6	Chronic lower respiratory diseases (J40-J47)	20	2.9	0.6
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	19	2.7	*
8	Anemias (D50-D64)	11	1.6	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	9	1.3	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	1.3	*
11	Septicemia (A40-A41)	8	1.1	*
11	Diabetes mellitus (E10-E14)	8	1.1	*
11	Influenza and pneumonia (J10-J18)	8	1.1	*
14	Cerebrovascular diseases (I60-I69)	7	1.0	*
15	Meningitis (G00,G03)	4	0.6	*
15	Certain conditions originating in the perinatal period (P00-P96)	4	0.6	*
...	All other causes (Residual)	146	21.0	4.4
Black, female, 15-24 years				
...	All causes	1,713	100.0	53.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	433	25.3	13.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	281	16.4	8.8
3	Malignant neoplasms (C00-C97)	135	7.9	4.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	109	6.4	3.4
5	Pregnancy, childbirth and the puerperium (O00-099)	74	4.3	2.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	69	4.0	2.2
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	69	4.0	2.2
8	Anemias (D50-D64)	43	2.5	1.3
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	38	2.2	1.2
10	Chronic lower respiratory diseases (J40-J47)	31	1.8	1.0
11	Diabetes mellitus (E10-E14)	30	1.8	0.9
12	Septicemia (A40-A41)	24	1.4	0.8
13	Influenza and pneumonia (J10-J18)	21	1.2	0.7
14	Cerebrovascular diseases (I60-I69)	18	1.1	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	0.7	*
...	All other causes (Residual)	326	19.0	10.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 25-34 years				
...	All causes	3,241	100.0	112.3
1	Human immunodeficiency virus (HIV) disease (B20-B24)	436	13.5	15.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	407	12.6	14.1
3	Malignant neoplasms (C00-C97)	402	12.4	13.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	383	11.8	13.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	282	8.7	9.8
6	Pregnancy, childbirth and the puerperium (O00-099)	109	3.4	3.8
7	Cerebrovascular diseases (I60-I69)	96	3.0	3.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	92	2.8	3.2
9	Diabetes mellitus (E10-E14)	79	2.4	2.7
10	Anemias (D50-D64)	58	1.8	2.0
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	52	1.6	1.8
12	Septicemia (A40-A41)	51	1.6	1.8
13	Chronic lower respiratory diseases (J40-J47)	35	1.1	1.2
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	26	0.8	0.9
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	23	0.7	0.8
...	All other causes (Residual)	710	21.9	24.6
Black, female, 35-44 years				
...	All causes	7,774	100.0	256.0
1	Malignant neoplasms (C00-C97)	1,647	21.2	54.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,275	16.4	42.0
3	Human immunodeficiency virus (HIV) disease (B20-B24)	930	12.0	30.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	647	8.3	21.3
5	Cerebrovascular diseases (I60-I69)	358	4.6	11.8
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	275	3.5	9.1
7	Diabetes mellitus (E10-E14)	230	3.0	7.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	150	1.9	4.9
9	Septicemia (A40-A41)	131	1.7	4.3
10	Chronic lower respiratory diseases (J40-J47)	117	1.5	3.9
11	Chronic liver disease and cirrhosis (K70,K73-K74)	111	1.4	3.7
12	Influenza and pneumonia (J10-J18)	99	1.3	3.3
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	77	1.0	2.5
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	74	1.0	2.4
15	Anemias (D50-D64)	66	0.8	2.2
...	All other causes (Residual)	1,587	20.4	52.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 45-54 years				
...	All causes	14,954	100.0	564.1
1	Malignant neoplasms (C00-C97)	4,264	28.5	160.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,212	21.5	121.2
3	Cerebrovascular diseases (I60-I69)	899	6.0	33.9
4	Human immunodeficiency virus (HIV) disease (B20-B24)	758	5.1	28.6
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	708	4.7	26.7
6	Diabetes mellitus (E10-E14)	631	4.2	23.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	342	2.3	12.9
8	Septicemia (A40-A41)	319	2.1	12.0
9	Chronic lower respiratory diseases (J40-J47)	305	2.0	11.5
10	Chronic liver disease and cirrhosis (K70,K73-K74)	285	1.9	10.8
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	208	1.4	7.8
12	Influenza and pneumonia (J10-J18)	179	1.2	6.8
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	160	1.1	6.0
14	Viral hepatitis (B15-B19)	102	0.7	3.8
15	Anemias (D50-D64)	77	0.5	2.9
...	All other causes (Residual)	2,505	16.8	94.5
Black, female, 55-64 years				
...	All causes	18,138	100.0	1,128.6
1	Malignant neoplasms (C00-C97)	5,937	32.7	369.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,436	24.5	276.0
3	Diabetes mellitus (E10-E14)	1,150	6.3	71.6
4	Cerebrovascular diseases (I60-I69)	1,045	5.8	65.0
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	647	3.6	40.3
6	Chronic lower respiratory diseases (J40-J47)	458	2.5	28.5
7	Septicemia (A40-A41)	448	2.5	27.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	356	2.0	22.2
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	327	1.8	20.3
10	Influenza and pneumonia (J10-J18)	230	1.3	14.3
11	Chronic liver disease and cirrhosis (K70,K73-K74)	228	1.3	14.2
12	Human immunodeficiency virus (HIV) disease (B20-B24)	210	1.2	13.1
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	80	0.4	5.0
14	Viral hepatitis (B15-B19)	72	0.4	4.5
15	Aortic aneurysm and dissection (I71)	67	0.4	4.2
...	All other causes (Residual)	2,447	13.5	152.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 65-74 years				
...	All causes	24,127	100.0	2,386.1
1	Malignant neoplasms (C00-C97)	7,141	29.6	706.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,638	27.5	656.5
3	Cerebrovascular diseases (I60-I69)	1,687	7.0	166.8
4	Diabetes mellitus (E10-E14)	1,675	6.9	165.7
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	873	3.6	86.3
6	Chronic lower respiratory diseases (J40-J47)	757	3.1	74.9
7	Septicemia (A40-A41)	626	2.6	61.9
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	489	2.0	48.4
9	Influenza and pneumonia (J10-J18)	340	1.4	33.6
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	303	1.3	30.0
11	Alzheimer's disease (G30)	196	0.8	19.4
12	Chronic liver disease and cirrhosis (K70,K73-K74)	130	0.5	12.9
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	111	0.5	11.0
14	Aortic aneurysm and dissection (I71)	95	0.4	9.4
15	Pneumonitis due to solids and liquids (J69)	88	0.4	8.7
...	All other causes (Residual)	2,978	12.3	294.5
Black, female, 75-84 years				
...	All causes	33,684	100.0	5,300.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,314	30.6	1,622.9
2	Malignant neoplasms (C00-C97)	6,887	20.4	1,083.6
3	Cerebrovascular diseases (I60-I69)	3,111	9.2	489.5
4	Diabetes mellitus (E10-E14)	2,086	6.2	328.2
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,181	3.5	185.8
6	Alzheimer's disease (G30)	1,032	3.1	162.4
7	Chronic lower respiratory diseases (J40-J47)	1,008	3.0	158.6
8	Septicemia (A40-A41)	869	2.6	136.7
9	Influenza and pneumonia (J10-J18)	776	2.3	122.1
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	728	2.2	114.5
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	392	1.2	61.7
12	Pneumonitis due to solids and liquids (J69)	224	0.7	35.2
13	Aortic aneurysm and dissection (I71)	186	0.6	29.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	172	0.5	27.1
15	Parkinson's disease (G20-G21)	134	0.4	21.1
...	All other causes (Residual)	4,584	13.6	721.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 85 years and over				
...	All causes	32,752	100.0	12,896.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,516	35.2	4,534.7
2	Malignant neoplasms (C00-C97)	3,524	10.8	1,387.7
3	Cerebrovascular diseases (I60-I69)	3,227	9.9	1,270.7
4	Alzheimer's disease (G30)	1,832	5.6	721.4
5	Diabetes mellitus (E10-E14)	1,379	4.2	543.0
6	Influenza and pneumonia (J10-J18)	1,189	3.6	468.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,012	3.1	398.5
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	925	2.8	364.2
9	Septicemia (A40-A41)	792	2.4	311.9
10	Chronic lower respiratory diseases (J40-J47)	614	1.9	241.8
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	325	1.0	128.0
12	Pneumonitis due to solids and liquids (J69)	309	0.9	121.7
13	Atherosclerosis (I70)	286	0.9	112.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	140	0.4	55.1
15	Anemias (D50-D64)	128	0.4	50.4
...	All other causes (Residual)	5,554	17.0	2,187.0
American Indian, both sexes, all ages				
...	All causes	13,124	100.0	416.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,598	19.8	82.5
2	Malignant neoplasms (C00-C97)	2,392	18.2	76.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,520	11.6	48.3
4	Diabetes mellitus (E10-E14)	746	5.7	23.7
5	Cerebrovascular diseases (I60-I69)	581	4.4	18.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	577	4.4	18.3
7	Chronic lower respiratory diseases (J40-J47)	486	3.7	15.4
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	404	3.1	12.8
9	Influenza and pneumonia (J10-J18)	291	2.2	9.2
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	247	1.9	7.8
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	228	1.7	7.2
12	Septicemia (A40-A41)	183	1.4	5.8
13	Certain conditions originating in the perinatal period (P00-P96)	137	1.0	4.4
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	125	1.0	4.0
15	Alzheimer's disease (G30)	112	0.9	3.6
...	All other causes (Residual)	2,497	19.0	79.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 1-4 years				
...	All causes	97	100.0	54.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	42	43.3	23.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	8.2	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7	7.2	*
4	Influenza and pneumonia (J10-J18)	4	4.1	*
5	Malignant neoplasms (C00-C97)	3	3.1	*
6	Septicemia (A40-A41)	2	2.1	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	2.1	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.0	*
8	Chronic lower respiratory diseases (J40-J47)	1	1.0	*
8	Certain conditions originating in the perinatal period (P00-P96)	1	1.0	*
...	All other causes (Residual)	26	26.8	14.6
American Indian, both sexes, 5-14 years				
...	All causes	127	100.0	22.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	61	48.0	10.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	10	7.9	*
3	Malignant neoplasms (C00-C97)	8	6.3	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	5.5	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	4.7	*
6	Septicemia (A40-A41)	4	3.1	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	1.6	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.8	*
8	Anemias (D50-D64)	1	0.8	*
8	Cerebrovascular diseases (I60-I69)	1	0.8	*
8	Influenza and pneumonia (J10-J18)	1	0.8	*
8	Chronic lower respiratory diseases (J40-J47)	1	0.8	*
8	Pneumonitis due to solids and liquids (J69)	1	0.8	*
8	Diseases of appendix (K35-K38)	1	0.8	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.8	*
...	All other causes (Residual)	21	16.5	3.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 15-24 years				
...	All causes	593	100.0	101.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	295	49.7	50.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	122	20.6	20.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	67	11.3	11.5
4	Malignant neoplasms (C00-C97)	16	2.7	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	13	2.2	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	0.8	*
7	Chronic lower respiratory diseases (J40-J47)	3	0.5	*
7	Chronic liver disease and cirrhosis (K70,K73-K74)	3	0.5	*
9	Septicemia (A40-A41)	2	0.3	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.3	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.2	*
11	Diabetes mellitus (E10-E14)	1	0.2	*
11	Parkinson's disease (G20-G21)	1	0.2	*
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1	0.2	*
11	Cerebrovascular diseases (I60-I69)	1	0.2	*
11	Diseases of appendix (K35-K38)	1	0.2	*
11	Pregnancy, childbirth and the puerperium (O00-099)	1	0.2	*
11	Legal intervention (Y35,Y89.0)	1	0.2	*
...	All other causes (Residual)	57	9.6	9.8
American Indian, both sexes, 25-34 years				
...	All causes	695	100.0	147.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	262	37.7	55.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	102	14.7	21.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	54	7.8	11.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	48	6.9	10.2
5	Malignant neoplasms (C00-C97)	25	3.6	5.3
6	Chronic liver disease and cirrhosis (K70,K73-K74)	21	3.0	4.5
7	Diabetes mellitus (E10-E14)	11	1.6	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	1.4	*
9	Septicemia (A40-A41)	9	1.3	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	9	1.3	*
11	Influenza and pneumonia (J10-J18)	7	1.0	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	1.0	*
13	Cerebrovascular diseases (I60-I69)	5	0.7	*
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2	0.3	*
14	Aortic aneurysm and dissection (I71)	2	0.3	*
14	Chronic lower respiratory diseases (J40-J47)	2	0.3	*
...	All other causes (Residual)	119	17.1	25.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 35-44 years				
...	All causes	1,194	100.0	256.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	290	24.3	62.3
2	Chronic liver disease and cirrhosis (K70,K73-K74)	128	10.7	27.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	118	9.9	25.3
4	Malignant neoplasms (C00-C97)	106	8.9	22.8
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	90	7.5	19.3
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	46	3.9	9.9
7	Human immunodeficiency virus (HIV) disease (B20-B24)	39	3.3	8.4
8	Diabetes mellitus (E10-E14)	36	3.0	7.7
9	Cerebrovascular diseases (I60-I69)	24	2.0	5.2
10	Septicemia (A40-A41)	16	1.3	*
11	Viral hepatitis (B15-B19)	13	1.1	*
11	Influenza and pneumonia (J10-J18)	13	1.1	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.8	*
14	Chronic lower respiratory diseases (J40-J47)	8	0.7	*
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	6	0.5	*
...	All other causes (Residual)	252	21.1	54.1
American Indian, both sexes, 45-54 years				
...	All causes	1,739	100.0	438.7
1	Malignant neoplasms (C00-C97)	313	18.0	79.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	282	16.2	71.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	234	13.5	59.0
4	Chronic liver disease and cirrhosis (K70,K73-K74)	184	10.6	46.4
5	Diabetes mellitus (E10-E14)	85	4.9	21.4
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	54	3.1	13.6
7	Cerebrovascular diseases (I60-I69)	49	2.8	12.4
8	Septicemia (A40-A41)	31	1.8	7.8
8	Viral hepatitis (B15-B19)	31	1.8	7.8
10	Chronic lower respiratory diseases (J40-J47)	29	1.7	7.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	29	1.7	7.3
12	Human immunodeficiency virus (HIV) disease (B20-B24)	26	1.5	6.6
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	22	1.3	5.6
14	Influenza and pneumonia (J10-J18)	19	1.1	*
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	12	0.7	*
...	All other causes (Residual)	339	19.5	85.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 55-64 years				
...	All causes	2,073	100.0	878.9
1	Malignant neoplasms (C00-C97)	564	27.2	239.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	439	21.2	186.1
3	Diabetes mellitus (E10-E14)	193	9.3	81.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	144	6.9	61.1
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	117	5.6	49.6
6	Chronic lower respiratory diseases (J40-J47)	89	4.3	37.7
7	Cerebrovascular diseases (I60-I69)	64	3.1	27.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	39	1.9	16.5
9	Septicemia (A40-A41)	33	1.6	14.0
10	Influenza and pneumonia (J10-J18)	32	1.5	13.6
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	18	0.9	*
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	12	0.6	*
13	Viral hepatitis (B15-B19)	11	0.5	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	9	0.4	*
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8	0.4	*
...	All other causes (Residual)	301	14.5	127.6
American Indian, both sexes, 65-74 years				
...	All causes	2,329	100.0	1,930.1
1	Malignant neoplasms (C00-C97)	670	28.8	555.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	535	23.0	443.4
3	Diabetes mellitus (E10-E14)	197	8.5	163.3
4	Chronic lower respiratory diseases (J40-J47)	136	5.8	112.7
5	Cerebrovascular diseases (I60-I69)	133	5.7	110.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	89	3.8	73.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	64	2.7	53.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	60	2.6	49.7
9	Influenza and pneumonia (J10-J18)	38	1.6	31.5
10	Septicemia (A40-A41)	25	1.1	20.7
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	23	1.0	19.1
12	Alzheimer's disease (G30)	11	0.5	*
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11	0.5	*
14	Atherosclerosis (I70)	10	0.4	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.4	*
15	Pneumonitis due to solids and liquids (J69)	9	0.4	*
...	All other causes (Residual)	309	13.3	256.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 75-84 years				
...	All causes	2,375	100.0	3,972.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	624	26.3	1,043.8
2	Malignant neoplasms (C00-C97)	511	21.5	854.8
3	Diabetes mellitus (E10-E14)	173	7.3	289.4
4	Cerebrovascular diseases (I60-I69)	168	7.1	281.0
5	Chronic lower respiratory diseases (J40-J47)	161	6.8	269.3
6	Influenza and pneumonia (J10-J18)	91	3.8	152.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	65	2.7	108.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	59	2.5	98.7
9	Alzheimer's disease (G30)	43	1.8	71.9
10	Septicemia (A40-A41)	31	1.3	51.9
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	29	1.2	48.5
12	Chronic liver disease and cirrhosis (K70,K73-K74)	28	1.2	46.8
13	Parkinson's disease (G20-G21)	25	1.1	41.8
14	Pneumonitis due to solids and liquids (J69)	21	0.9	35.1
15	Aortic aneurysm and dissection (I71)	15	0.6	*
...	All other causes (Residual)	331	13.9	553.7
American Indian, both sexes, 85 years and over				
...	All causes	1,503	100.0	7,362.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	524	34.9	2,566.9
2	Malignant neoplasms (C00-C97)	175	11.6	857.3
3	Cerebrovascular diseases (I60-I69)	135	9.0	661.3
4	Influenza and pneumonia (J10-J18)	74	4.9	362.5
5	Alzheimer's disease (G30)	56	3.7	274.3
6	Chronic lower respiratory diseases (J40-J47)	55	3.7	269.4
7	Diabetes mellitus (E10-E14)	50	3.3	244.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	37	2.5	181.2
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	37	2.5	181.2
10	Septicemia (A40-A41)	25	1.7	122.5
11	Pneumonitis due to solids and liquids (J69)	17	1.1	*
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	16	1.1	*
13	Parkinson's disease (G20-G21)	14	0.9	*
14	Atherosclerosis (I70)	12	0.8	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.7	*
...	All other causes (Residual)	265	17.6	1,298.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, all ages				
...	All causes	7,134	100.0	453.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,437	20.1	91.4
2	Malignant neoplasms (C00-C97)	1,241	17.4	78.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,010	14.2	64.2
4	Diabetes mellitus (E10-E14)	365	5.1	23.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	323	4.5	20.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	306	4.3	19.5
7	Cerebrovascular diseases (I60-I69)	246	3.4	15.6
8	Chronic lower respiratory diseases (J40-J47)	235	3.3	14.9
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	181	2.5	11.5
10	Influenza and pneumonia (J10-J18)	142	2.0	9.0
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	112	1.6	7.1
12	Septicemia (A40-A41)	85	1.2	5.4
13	Certain conditions originating in the perinatal period (P00-P96)	79	1.1	5.0
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	70	1.0	4.5
15	Human immunodeficiency virus (HIV) disease (B20-B24)	63	0.9	4.0
...	All other causes (Residual)	1,239	17.4	78.8
American Indian, male, 1-4 years				
...	All causes	50	100.0	55.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	26	52.0	28.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	10.0	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	6.0	*
4	Septicemia (A40-A41)	2	4.0	*
4	Malignant neoplasms (C00-C97)	2	4.0	*
6	Influenza and pneumonia (J10-J18)	1	2.0	*
...	All other causes (Residual)	11	22.0	*

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 5-14 years				
...	All causes	70	100.0	24.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	33	47.1	11.4
2	Malignant neoplasms (C00-C97)	7	10.0	*
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6	8.6	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	7.1	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	5.7	*
6	Septicemia (A40-A41)	1	1.4	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.4	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	1.4	*
6	Cerebrovascular diseases (I60-I69)	1	1.4	*
6	Pneumonitis due to solids and liquids (J69)	1	1.4	*
6	Diseases of appendix (K35-K38)	1	1.4	*
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	1.4	*
...	All other causes (Residual)	8	11.4	*
American Indian, male, 15-24 years				
...	All causes	409	100.0	136.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	203	49.6	67.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	92	22.5	30.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	53	13.0	17.7
4	Malignant neoplasms (C00-C97)	9	2.2	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	1.2	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.7	*
7	Chronic lower respiratory diseases (J40-J47)	2	0.5	*
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2	0.5	*
9	Septicemia (A40-A41)	1	0.2	*
9	Parkinson's disease (G20-G21)	1	0.2	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
9	Legal intervention (Y35,Y89.0)	1	0.2	*
...	All other causes (Residual)	36	8.8	12.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 25-34 years				
...	All causes	461	100.0	188.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	184	39.9	75.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	76	16.5	31.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	47	10.2	19.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	37	8.0	15.2
5	Malignant neoplasms (C00-C97)	11	2.4	*
5	Chronic liver disease and cirrhosis (K70,K73-K74)	11	2.4	*
7	Human immunodeficiency virus (HIV) disease (B20-B24)	8	1.7	*
8	Diabetes mellitus (E10-E14)	5	1.1	*
8	Influenza and pneumonia (J10-J18)	5	1.1	*
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	1.1	*
11	Septicemia (A40-A41)	4	0.9	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.7	*
13	Cerebrovascular diseases (I60-I69)	2	0.4	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.2	*
14	Meningitis (G00,G03)	1	0.2	*
14	Aortic aneurysm and dissection (I71)	1	0.2	*
14	Chronic lower respiratory diseases (J40-J47)	1	0.2	*
14	Legal intervention (Y35,Y89.0)	1	0.2	*
...	All other causes (Residual)	58	12.6	23.8
American Indian, male, 35-44 years				
...	All causes	747	100.0	321.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	191	25.6	82.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	83	11.1	35.7
3	Chronic liver disease and cirrhosis (K70,K73-K74)	75	10.0	32.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	71	9.5	30.5
5	Malignant neoplasms (C00-C97)	42	5.6	18.1
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	38	5.1	16.3
7	Human immunodeficiency virus (HIV) disease (B20-B24)	33	4.4	14.2
8	Diabetes mellitus (E10-E14)	25	3.3	10.8
9	Septicemia (A40-A41)	8	1.1	*
9	Viral hepatitis (B15-B19)	8	1.1	*
11	Influenza and pneumonia (J10-J18)	7	0.9	*
12	Cerebrovascular diseases (I60-I69)	6	0.8	*
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	5	0.7	*
13	Chronic lower respiratory diseases (J40-J47)	5	0.7	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.5	*
...	All other causes (Residual)	146	19.5	62.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 45-54 years				
...	All causes	1,045	100.0	543.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	181	17.3	94.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	169	16.2	87.9
3	Malignant neoplasms (C00-C97)	166	15.9	86.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	105	10.0	54.6
5	Diabetes mellitus (E10-E14)	54	5.2	28.1
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	40	3.8	20.8
7	Cerebrovascular diseases (I60-I69)	27	2.6	14.0
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	18	1.7	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	17	1.6	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	1.6	*
11	Septicemia (A40-A41)	15	1.4	*
12	Viral hepatitis (B15-B19)	14	1.3	*
13	Chronic lower respiratory diseases (J40-J47)	13	1.2	*
14	Influenza and pneumonia (J10-J18)	12	1.1	*
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	8	0.8	*
...	All other causes (Residual)	189	18.1	98.3
American Indian, male, 55-64 years				
...	All causes	1,212	100.0	1,067.3
1	Malignant neoplasms (C00-C97)	305	25.2	268.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	296	24.4	260.7
3	Diabetes mellitus (E10-E14)	106	8.7	93.3
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	94	7.8	82.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	84	6.9	74.0
6	Chronic lower respiratory diseases (J40-J47)	46	3.8	40.5
7	Cerebrovascular diseases (I60-I69)	34	2.8	29.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19	1.6	*
9	Septicemia (A40-A41)	14	1.2	*
10	Influenza and pneumonia (J10-J18)	13	1.1	*
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	9	0.7	*
12	Viral hepatitis (B15-B19)	7	0.6	*
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	7	0.6	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	5	0.4	*
14	Aortic aneurysm and dissection (I71)	5	0.4	*
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	0.4	*
...	All other causes (Residual)	163	13.4	143.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 65-74 years				
...	All causes	1,225	100.0	2,196.8
1	Malignant neoplasms (C00-C97)	358	29.2	642.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	329	26.9	590.0
3	Diabetes mellitus (E10-E14)	81	6.6	145.3
4	Chronic lower respiratory diseases (J40-J47)	68	5.6	121.9
5	Cerebrovascular diseases (I60-I69)	61	5.0	109.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	46	3.8	82.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	30	2.4	53.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	28	2.3	50.2
9	Influenza and pneumonia (J10-J18)	18	1.5	*
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	15	1.2	*
11	Septicemia (A40-A41)	12	1.0	*
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	9	0.7	*
13	Alzheimer's disease (G30)	6	0.5	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.4	*
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	0.4	*
...	All other causes (Residual)	154	12.6	276.2
American Indian, male, 75-84 years				
...	All causes	1,146	100.0	4,584.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	313	27.3	1,252.1
2	Malignant neoplasms (C00-C97)	265	23.1	1,060.0
3	Diabetes mellitus (E10-E14)	79	6.9	316.0
4	Cerebrovascular diseases (I60-I69)	78	6.8	312.0
5	Chronic lower respiratory diseases (J40-J47)	75	6.5	300.0
6	Influenza and pneumonia (J10-J18)	46	4.0	184.0
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	27	2.4	108.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22	1.9	88.0
9	Parkinson's disease (G20-G21)	17	1.5	*
10	Chronic liver disease and cirrhosis (K70,K73-K74)	15	1.3	*
11	Alzheimer's disease (G30)	14	1.2	*
12	Septicemia (A40-A41)	13	1.1	*
13	Pneumonitis due to solids and liquids (J69)	12	1.0	*
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	9	0.8	*
14	Aortic aneurysm and dissection (I71)	9	0.8	*
...	All other causes (Residual)	152	13.3	608.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 85 years and over				
...	All causes	524	100.0	7,923.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	186	35.5	2,812.6
2	Malignant neoplasms (C00-C97)	75	14.3	1,134.1
3	Cerebrovascular diseases (I60-I69)	37	7.1	559.5
4	Influenza and pneumonia (J10-J18)	33	6.3	499.0
5	Chronic lower respiratory diseases (J40-J47)	24	4.6	362.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19	3.6	*
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	3.1	*
8	Diabetes mellitus (E10-E14)	15	2.9	*
9	Septicemia (A40-A41)	10	1.9	*
9	Alzheimer's disease (G30)	10	1.9	*
11	Pneumonitis due to solids and liquids (J69)	7	1.3	*
12	Parkinson's disease (G20-G21)	6	1.1	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	1.0	*
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	4	0.8	*
15	Atherosclerosis (I70)	3	0.6	*
15	Aortic aneurysm and dissection (I71)	3	0.6	*
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	3	0.6	*
...	All other causes (Residual)	68	13.0	1,028.3
American Indian, female, all ages				
...	All causes	5,990	100.0	380.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,161	19.4	73.6
2	Malignant neoplasms (C00-C97)	1,151	19.2	73.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	510	8.5	32.4
4	Diabetes mellitus (E10-E14)	381	6.4	24.2
5	Cerebrovascular diseases (I60-I69)	335	5.6	21.3
6	Chronic liver disease and cirrhosis (K70,K73-K74)	254	4.2	16.1
7	Chronic lower respiratory diseases (J40-J47)	251	4.2	15.9
8	Influenza and pneumonia (J10-J18)	149	2.5	9.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	135	2.3	8.6
10	Septicemia (A40-A41)	98	1.6	6.2
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	98	1.6	6.2
12	Alzheimer's disease (G30)	80	1.3	5.1
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	59	1.0	3.7
14	Certain conditions originating in the perinatal period (P00-P96)	58	1.0	3.7
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	55	0.9	3.5
...	All other causes (Residual)	1,215	20.3	77.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 1-4 years				
...	All causes	47	100.0	53.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	16	34.0	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	10.6	*
3	Influenza and pneumonia (J10-J18)	3	6.4	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	4.3	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	4.3	*
6	Malignant neoplasms (C00-C97)	1	2.1	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	2.1	*
6	Chronic lower respiratory diseases (J40-J47)	1	2.1	*
6	Certain conditions originating in the perinatal period (P00-P96)	1	2.1	*
...	All other causes (Residual)	15	31.9	*
American Indian, female, 5-14 years				
...	All causes	57	100.0	20.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	28	49.1	9.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4	7.0	*
3	Septicemia (A40-A41)	3	5.3	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	3.5	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	3.5	*
6	Malignant neoplasms (C00-C97)	1	1.8	*
6	Anemias (D50-D64)	1	1.8	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	1.8	*
6	Influenza and pneumonia (J10-J18)	1	1.8	*
6	Chronic lower respiratory diseases (J40-J47)	1	1.8	*
...	All other causes (Residual)	13	22.8	*
American Indian, female, 15-24 years				
...	All causes	184	100.0	64.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	92	50.0	32.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	30	16.3	10.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	14	7.6	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	8	4.3	*
5	Malignant neoplasms (C00-C97)	7	3.8	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	1.1	*
7	Septicemia (A40-A41)	1	0.5	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.5	*
7	Diabetes mellitus (E10-E14)	1	0.5	*
7	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1	0.5	*
7	Cerebrovascular diseases (I60-I69)	1	0.5	*
7	Chronic lower respiratory diseases (J40-J47)	1	0.5	*
7	Diseases of appendix (K35-K38)	1	0.5	*
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.5	*
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.5	*
7	Pregnancy, childbirth and the puerperium (O00-099)	1	0.5	*
...	All other causes (Residual)	21	11.4	7.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 25-34 years				
...	All causes	234	100.0	102.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	78	33.3	34.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	26	11.1	11.4
3	Malignant neoplasms (C00-C97)	14	6.0	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	11	4.7	*
5	Chronic liver disease and cirrhosis (K70,K73-K74)	10	4.3	*
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	3.0	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7	3.0	*
8	Diabetes mellitus (E10-E14)	6	2.6	*
9	Septicemia (A40-A41)	5	2.1	*
10	Cerebrovascular diseases (I60-I69)	3	1.3	*
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2	0.9	*
11	Influenza and pneumonia (J10-J18)	2	0.9	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	0.9	*
14	Meningococcal infection (A39)	1	0.4	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.4	*
14	Aortic aneurysm and dissection (I71)	1	0.4	*
14	Chronic lower respiratory diseases (J40-J47)	1	0.4	*
14	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.4	*
14	Pregnancy, childbirth and the puerperium (O00-099)	1	0.4	*
...	All other causes (Residual)	55	23.5	24.2
American Indian, female, 35-44 years				
...	All causes	447	100.0	191.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	99	22.1	42.4
2	Malignant neoplasms (C00-C97)	64	14.3	27.4
3	Chronic liver disease and cirrhosis (K70,K73-K74)	53	11.9	22.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	35	7.8	15.0
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	19	4.3	*
6	Cerebrovascular diseases (I60-I69)	18	4.0	*
7	Diabetes mellitus (E10-E14)	11	2.5	*
8	Septicemia (A40-A41)	8	1.8	*
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8	1.8	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	6	1.3	*
10	Influenza and pneumonia (J10-J18)	6	1.3	*
12	Viral hepatitis (B15-B19)	5	1.1	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	1.1	*
14	Chronic lower respiratory diseases (J40-J47)	3	0.7	*
14	Pregnancy, childbirth and the puerperium (O00-099)	3	0.7	*
...	All other causes (Residual)	104	23.3	44.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 45-54 years				
...	All causes	694	100.0	340.0
1	Malignant neoplasms (C00-C97)	147	21.2	72.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	101	14.6	49.5
3	Chronic liver disease and cirrhosis (K70,K73-K74)	79	11.4	38.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	65	9.4	31.8
5	Diabetes mellitus (E10-E14)	31	4.5	15.2
6	Cerebrovascular diseases (I60-I69)	22	3.2	10.8
7	Viral hepatitis (B15-B19)	17	2.4	*
8	Septicemia (A40-A41)	16	2.3	*
8	Chronic lower respiratory diseases (J40-J47)	16	2.3	*
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	14	2.0	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	1.7	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	9	1.3	*
13	Influenza and pneumonia (J10-J18)	7	1.0	*
14	Pneumonitis due to solids and liquids (J69)	6	0.9	*
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	4	0.6	*
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	0.6	*
...	All other causes (Residual)	144	20.7	70.6
American Indian, female, 55-64 years				
...	All causes	861	100.0	704.0
1	Malignant neoplasms (C00-C97)	259	30.1	211.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	143	16.6	116.9
3	Diabetes mellitus (E10-E14)	87	10.1	71.1
4	Chronic liver disease and cirrhosis (K70,K73-K74)	60	7.0	49.1
5	Chronic lower respiratory diseases (J40-J47)	43	5.0	35.2
6	Cerebrovascular diseases (I60-I69)	30	3.5	24.5
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	23	2.7	18.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20	2.3	16.4
9	Septicemia (A40-A41)	19	2.2	*
9	Influenza and pneumonia (J10-J18)	19	2.2	*
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	11	1.3	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.6	*
13	Viral hepatitis (B15-B19)	4	0.5	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.5	*
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3	0.3	*
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	0.3	*
...	All other causes (Residual)	128	14.9	104.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 65-74 years				
...	All causes	1,104	100.0	1,700.9
1	Malignant neoplasms (C00-C97)	312	28.3	480.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	206	18.7	317.4
3	Diabetes mellitus (E10-E14)	116	10.5	178.7
4	Cerebrovascular diseases (I60-I69)	72	6.5	110.9
5	Chronic lower respiratory diseases (J40-J47)	68	6.2	104.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	43	3.9	66.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	34	3.1	52.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	32	2.9	49.3
9	Influenza and pneumonia (J10-J18)	20	1.8	30.8
10	Septicemia (A40-A41)	13	1.2	*
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	8	0.7	*
12	Atherosclerosis (I70)	7	0.6	*
13	Alzheimer's disease (G30)	5	0.5	*
13	Pneumonitis due to solids and liquids (J69)	5	0.5	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.4	*
15	Aortic aneurysm and dissection (I71)	4	0.4	*
...	All other causes (Residual)	155	14.0	238.8
American Indian, female, 75-84 years				
...	All causes	1,229	100.0	3,533.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	311	25.3	894.1
2	Malignant neoplasms (C00-C97)	246	20.0	707.3
3	Diabetes mellitus (E10-E14)	94	7.6	270.3
4	Cerebrovascular diseases (I60-I69)	90	7.3	258.8
5	Chronic lower respiratory diseases (J40-J47)	86	7.0	247.3
6	Influenza and pneumonia (J10-J18)	45	3.7	129.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	38	3.1	109.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	37	3.0	106.4
9	Alzheimer's disease (G30)	29	2.4	83.4
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	20	1.6	57.5
11	Septicemia (A40-A41)	18	1.5	*
12	Chronic liver disease and cirrhosis (K70,K73-K74)	13	1.1	*
13	Pneumonitis due to solids and liquids (J69)	9	0.7	*
14	Parkinson's disease (G20-G21)	8	0.7	*
15	Aortic aneurysm and dissection (I71)	6	0.5	*
...	All other causes (Residual)	179	14.6	514.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 85 years and over				
...	All causes	979	100.0	7,093.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	338	34.5	2,449.1
2	Malignant neoplasms (C00-C97)	100	10.2	724.6
3	Cerebrovascular diseases (I60-I69)	98	10.0	710.1
4	Alzheimer's disease (G30)	46	4.7	333.3
5	Influenza and pneumonia (J10-J18)	41	4.2	297.1
6	Diabetes mellitus (E10-E14)	35	3.6	253.6
7	Chronic lower respiratory diseases (J40-J47)	31	3.2	224.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21	2.1	152.2
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18	1.8	*
10	Septicemia (A40-A41)	15	1.5	*
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	12	1.2	*
12	Pneumonitis due to solids and liquids (J69)	10	1.0	*
13	Atherosclerosis (I70)	9	0.9	*
14	Parkinson's disease (G20-G21)	8	0.8	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.6	*
15	Nutritional deficiencies (E40-E64)	6	0.6	*
...	All other causes (Residual)	185	18.9	1,340.5
Asian or Pacific Islander, both sexes, all ages				
...	All causes	40,533	100.0	297.2
1	Malignant neoplasms (C00-C97)	10,863	26.8	79.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,960	24.6	73.0
3	Cerebrovascular diseases (I60-I69)	3,507	8.7	25.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,932	4.8	14.2
5	Diabetes mellitus (E10-E14)	1,471	3.6	10.8
6	Influenza and pneumonia (J10-J18)	1,243	3.1	9.1
7	Chronic lower respiratory diseases (J40-J47)	1,187	2.9	8.7
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	765	1.9	5.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	708	1.7	5.2
10	Alzheimer's disease (G30)	534	1.3	3.9
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	529	1.3	3.9
12	Septicemia (A40-A41)	502	1.2	3.7
13	Certain conditions originating in the perinatal period (P00-P96)	424	1.0	3.1
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	351	0.9	2.6
15	Chronic liver disease and cirrhosis (K70,K73-K74)	340	0.8	2.5
...	All other causes (Residual)	6,217	15.3	45.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 1-4 years				
...	All causes	161	100.0	21.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	33	20.5	4.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	29	18.0	3.8
3	Malignant neoplasms (C00-C97)	20	12.4	2.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	7	4.3	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	3.7	*
6	Certain conditions originating in the perinatal period (P00-P96)	4	2.5	*
7	Septicemia (A40-A41)	3	1.9	*
7	Influenza and pneumonia (J10-J18)	3	1.9	*
7	Chronic lower respiratory diseases (J40-J47)	3	1.9	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.6	*
10	Anemias (D50-D64)	1	0.6	*
10	Diabetes mellitus (E10-E14)	1	0.6	*
10	Meningitis (G00,G03)	1	0.6	*
10	Cerebrovascular diseases (I60-I69)	1	0.6	*
10	Acute bronchitis and bronchiolitis (J20-J21)	1	0.6	*
10	Peptic ulcer (K25-K28)	1	0.6	*
...	All other causes (Residual)	46	28.6	6.1
Asian or Pacific Islander, both sexes, 5-14 years				
...	All causes	223	100.0	12.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	82	36.8	4.5
2	Malignant neoplasms (C00-C97)	48	21.5	2.6
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	10	4.5	*
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8	3.6	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8	3.6	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	1.8	*
6	Influenza and pneumonia (J10-J18)	4	1.8	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	1.3	*
9	Anemias (D50-D64)	2	0.9	*
9	Diabetes mellitus (E10-E14)	2	0.9	*
9	Chronic lower respiratory diseases (J40-J47)	2	0.9	*
9	Pregnancy, childbirth and the puerperium (O00-O99)	2	0.9	*
9	Certain conditions originating in the perinatal period (P00-P96)	2	0.9	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.4	*
14	Cerebrovascular diseases (I60-I69)	1	0.4	*
14	Diseases of appendix (K35-K38)	1	0.4	*
...	All other causes (Residual)	43	19.3	2.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 15-24 years				
...	All causes	766	100.0	39.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	319	41.6	16.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	119	15.5	6.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	89	11.6	4.6
4	Malignant neoplasms (C00-C97)	59	7.7	3.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	31	4.0	1.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	15	2.0	*
7	Cerebrovascular diseases (I60-I69)	9	1.2	*
8	Chronic lower respiratory diseases (J40-J47)	6	0.8	*
9	Septicemia (A40-A41)	5	0.7	*
9	Pregnancy, childbirth and the puerperium (O00-099)	5	0.7	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.5	*
11	Aortic aneurysm and dissection (I71)	4	0.5	*
11	Legal intervention (Y35,Y89.0)	4	0.5	*
14	Influenza and pneumonia (J10-J18)	3	0.4	*
15	Meningococcal infection (A39)	2	0.3	*
15	Viral hepatitis (B15-B19)	2	0.3	*
15	Meningitis (G00,G03)	2	0.3	*
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	2	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.3	*
...	All other causes (Residual)	84	11.0	4.3
Asian or Pacific Islander, both sexes, 25-34 years				
...	All causes	973	100.0	38.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	243	25.0	9.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	163	16.8	6.5
3	Malignant neoplasms (C00-C97)	158	16.2	6.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	95	9.8	3.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	76	7.8	3.0
6	Cerebrovascular diseases (I60-I69)	16	1.6	*
7	Chronic liver disease and cirrhosis (K70,K73-K74)	15	1.5	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	12	1.2	*
9	Septicemia (A40-A41)	9	0.9	*
9	Diabetes mellitus (E10-E14)	9	0.9	*
9	Pregnancy, childbirth and the puerperium (O00-099)	9	0.9	*
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	0.8	*
13	Viral hepatitis (B15-B19)	7	0.7	*
14	Aortic aneurysm and dissection (I71)	6	0.6	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.5	*
15	Legal intervention (Y35,Y89.0)	5	0.5	*
...	All other causes (Residual)	132	13.6	5.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 35-44 years				
...	All causes	1,654	100.0	72.2
1	Malignant neoplasms (C00-C97)	549	33.2	24.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	236	14.3	10.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	220	13.3	9.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	136	8.2	5.9
5	Cerebrovascular diseases (I60-I69)	77	4.7	3.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	49	3.0	2.1
7	Human immunodeficiency virus (HIV) disease (B20-B24)	37	2.2	1.6
8	Chronic liver disease and cirrhosis (K70,K73-K74)	34	2.1	1.5
9	Diabetes mellitus (E10-E14)	22	1.3	1.0
10	Viral hepatitis (B15-B19)	15	0.9	*
11	Septicemia (A40-A41)	14	0.8	*
11	Influenza and pneumonia (J10-J18)	14	0.8	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	11	0.7	*
14	Chronic lower respiratory diseases (J40-J47)	10	0.6	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	0.6	*
14	Pregnancy, childbirth and the puerperium (O00-099)	10	0.6	*
...	All other causes (Residual)	210	12.7	9.2
Asian or Pacific Islander, both sexes, 45-54 years				
...	All causes	3,529	100.0	190.9
1	Malignant neoplasms (C00-C97)	1,426	40.4	77.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	625	17.7	33.8
3	Cerebrovascular diseases (I60-I69)	269	7.6	14.5
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	223	6.3	12.1
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	144	4.1	7.8
6	Diabetes mellitus (E10-E14)	104	2.9	5.6
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	64	1.8	3.5
8	Chronic liver disease and cirrhosis (K70,K73-K74)	61	1.7	3.3
9	Septicemia (A40-A41)	41	1.2	2.2
10	Viral hepatitis (B15-B19)	37	1.0	2.0
11	Human immunodeficiency virus (HIV) disease (B20-B24)	32	0.9	1.7
11	Influenza and pneumonia (J10-J18)	32	0.9	1.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	32	0.9	1.7
14	Chronic lower respiratory diseases (J40-J47)	30	0.9	1.6
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	28	0.8	1.5
...	All other causes (Residual)	381	10.8	20.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 55-64 years				
...	All causes	4,963	100.0	435.5
1	Malignant neoplasms (C00-C97)	2,023	40.8	177.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,046	21.1	91.8
3	Cerebrovascular diseases (I60-I69)	366	7.4	32.1
4	Diabetes mellitus (E10-E14)	223	4.5	19.6
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	188	3.8	16.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	82	1.7	7.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	81	1.6	7.1
8	Chronic lower respiratory diseases (J40-J47)	76	1.5	6.7
9	Chronic liver disease and cirrhosis (K70,K73-K74)	73	1.5	6.4
10	Septicemia (A40-A41)	69	1.4	6.1
11	Influenza and pneumonia (J10-J18)	60	1.2	5.3
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	49	1.0	4.3
13	Viral hepatitis (B15-B19)	48	1.0	4.2
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	35	0.7	3.1
15	Aortic aneurysm and dissection (I71)	34	0.7	3.0
...	All other causes (Residual)	510	10.3	44.8
Asian or Pacific Islander, both sexes, 65-74 years				
...	All causes	7,393	100.0	1,124.7
1	Malignant neoplasms (C00-C97)	2,709	36.6	412.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,767	23.9	268.8
3	Cerebrovascular diseases (I60-I69)	585	7.9	89.0
4	Diabetes mellitus (E10-E14)	333	4.5	50.7
5	Chronic lower respiratory diseases (J40-J47)	211	2.9	32.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	186	2.5	28.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	152	2.1	23.1
8	Influenza and pneumonia (J10-J18)	151	2.0	23.0
9	Septicemia (A40-A41)	98	1.3	14.9
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	82	1.1	12.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	68	0.9	10.3
12	Aortic aneurysm and dissection (I71)	63	0.9	9.6
13	Viral hepatitis (B15-B19)	58	0.8	8.8
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	55	0.7	8.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	47	0.6	7.1
...	All other causes (Residual)	828	11.2	126.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 75-84 years				
...	All causes	10,816	100.0	3,055.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,978	27.5	841.2
2	Malignant neoplasms (C00-C97)	2,693	24.9	760.7
3	Cerebrovascular diseases (I60-I69)	1,162	10.7	328.2
4	Diabetes mellitus (E10-E14)	474	4.4	133.9
5	Chronic lower respiratory diseases (J40-J47)	449	4.2	126.8
6	Influenza and pneumonia (J10-J18)	397	3.7	112.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	243	2.2	68.6
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	232	2.1	65.5
9	Alzheimer's disease (G30)	186	1.7	52.5
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	165	1.5	46.6
11	Septicemia (A40-A41)	148	1.4	41.8
12	Parkinson's disease (G20-G21)	124	1.1	35.0
13	Aortic aneurysm and dissection (I71)	100	0.9	28.2
14	Pneumonitis due to solids and liquids (J69)	82	0.8	23.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	78	0.7	22.0
...	All other causes (Residual)	1,305	12.1	368.6
Asian or Pacific Islander, both sexes, 85 years and over				
...	All causes	9,230	100.0	8,881.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,164	34.3	3,044.6
2	Malignant neoplasms (C00-C97)	1,175	12.7	1,130.7
3	Cerebrovascular diseases (I60-I69)	1,014	11.0	975.7
4	Influenza and pneumonia (J10-J18)	570	6.2	548.5
5	Chronic lower respiratory diseases (J40-J47)	398	4.3	383.0
6	Alzheimer's disease (G30)	307	3.3	295.4
7	Diabetes mellitus (E10-E14)	303	3.3	291.6
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	200	2.2	192.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	175	1.9	168.4
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	170	1.8	163.6
11	Pneumonitis due to solids and liquids (J69)	132	1.4	127.0
12	Septicemia (A40-A41)	110	1.2	105.8
13	Parkinson's disease (G20-G21)	100	1.1	96.2
14	Atherosclerosis (I70)	63	0.7	60.6
15	Aortic aneurysm and dissection (I71)	60	0.7	57.7
...	All other causes (Residual)	1,289	14.0	1,240.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, all ages				
...	All causes	21,298	100.0	321.1
1	Malignant neoplasms (C00-C97)	5,696	26.7	85.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,400	25.4	81.4
3	Cerebrovascular diseases (I60-I69)	1,614	7.6	24.3
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,170	5.5	17.6
5	Chronic lower respiratory diseases (J40-J47)	743	3.5	11.2
6	Diabetes mellitus (E10-E14)	708	3.3	10.7
7	Influenza and pneumonia (J10-J18)	596	2.8	9.0
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	524	2.5	7.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	372	1.7	5.6
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	256	1.2	3.9
11	Septicemia (A40-A41)	250	1.2	3.8
12	Certain conditions originating in the perinatal period (P00-P96)	236	1.1	3.6
13	Chronic liver disease and cirrhosis (K70,K73-K74)	218	1.0	3.3
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	217	1.0	3.3
15	Aortic aneurysm and dissection (I71)	194	0.9	2.9
...	All other causes (Residual)	3,104	14.6	46.8
Asian or Pacific Islander, male, 1-4 years				
...	All causes	80	100.0	20.7
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	16	20.0	*
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	16	20.0	*
3	Malignant neoplasms (C00-C97)	12	15.0	*
4	Certain conditions originating in the perinatal period (P00-P96)	3	3.8	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	3.8	*
6	Influenza and pneumonia (J10-J18)	2	2.5	*
7	Septicemia (A40-A41)	1	1.3	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.3	*
7	Diabetes mellitus (E10-E14)	1	1.3	*
7	Meningitis (G00,G03)	1	1.3	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	1.3	*
7	Acute bronchitis and bronchiolitis (J20-J21)	1	1.3	*
7	Chronic lower respiratory diseases (J40-J47)	1	1.3	*
...	All other causes (Residual)	21	26.3	5.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 5-14 years				
...	All causes	136	100.0	14.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	54	39.7	5.8
2	Malignant neoplasms (C00-C97)	28	20.6	3.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8	5.9	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	3.7	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	3.7	*
6	Influenza and pneumonia (J10-J18)	3	2.2	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	1.5	*
7	Certain conditions originating in the perinatal period (P00-P96)	2	1.5	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.7	*
9	Anemias (D50-D64)	1	0.7	*
9	Diabetes mellitus (E10-E14)	1	0.7	*
9	Chronic lower respiratory diseases (J40-J47)	1	0.7	*
...	All other causes (Residual)	25	18.4	2.7
Asian or Pacific Islander, male, 15-24 years				
...	All causes	536	100.0	53.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	222	41.4	22.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	92	17.2	9.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	78	14.6	7.8
4	Malignant neoplasms (C00-C97)	41	7.6	4.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	20	3.7	2.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	10	1.9	*
7	Chronic lower respiratory diseases (J40-J47)	5	0.9	*
8	Legal intervention (Y35,Y89.0)	4	0.7	*
9	Cerebrovascular diseases (I60-I69)	3	0.6	*
10	Meningococcal infection (A39)	2	0.4	*
10	Septicemia (A40-A41)	2	0.4	*
10	Aortic aneurysm and dissection (I71)	2	0.4	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
14	Viral hepatitis (B15-B19)	1	0.2	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.2	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.2	*
14	Meningitis (G00,G03)	1	0.2	*
14	Influenza and pneumonia (J10-J18)	1	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
...	All other causes (Residual)	47	8.8	4.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 25-34 years				
...	All causes	630	100.0	50.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	174	27.6	14.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	111	17.6	9.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	78	12.4	6.3
4	Malignant neoplasms (C00-C97)	73	11.6	5.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	53	8.4	4.3
6	Chronic liver disease and cirrhosis (K70,K73-K74)	14	2.2	*
7	Human immunodeficiency virus (HIV) disease (B20-B24)	9	1.4	*
8	Septicemia (A40-A41)	7	1.1	*
8	Cerebrovascular diseases (I60-I69)	7	1.1	*
10	Viral hepatitis (B15-B19)	4	0.6	*
10	Diabetes mellitus (E10-E14)	4	0.6	*
10	Aortic aneurysm and dissection (I71)	4	0.6	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.6	*
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	0.6	*
10	Legal intervention (Y35,Y89.0)	4	0.6	*
...	All other causes (Residual)	80	12.7	6.5
Asian or Pacific Islander, male, 35-44 years				
...	All causes	1,015	100.0	91.3
1	Malignant neoplasms (C00-C97)	263	25.9	23.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	171	16.8	15.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	158	15.6	14.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	86	8.5	7.7
5	Cerebrovascular diseases (I60-I69)	47	4.6	4.2
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	34	3.3	3.1
7	Human immunodeficiency virus (HIV) disease (B20-B24)	32	3.2	2.9
8	Chronic liver disease and cirrhosis (K70,K73-K74)	28	2.8	2.5
9	Viral hepatitis (B15-B19)	13	1.3	*
10	Diabetes mellitus (E10-E14)	11	1.1	*
11	Influenza and pneumonia (J10-J18)	9	0.9	*
12	Aortic aneurysm and dissection (I71)	8	0.8	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.8	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	0.7	*
15	Chronic lower respiratory diseases (J40-J47)	6	0.6	*
...	All other causes (Residual)	134	13.2	12.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 45-54 years				
...	All causes	2,096	100.0	241.9
1	Malignant neoplasms (C00-C97)	669	31.9	77.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	490	23.4	56.6
3	Cerebrovascular diseases (I60-I69)	166	7.9	19.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	139	6.6	16.0
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	101	4.8	11.7
6	Diabetes mellitus (E10-E14)	60	2.9	6.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	55	2.6	6.3
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	43	2.1	5.0
9	Viral hepatitis (B15-B19)	28	1.3	3.2
10	Human immunodeficiency virus (HIV) disease (B20-B24)	26	1.2	3.0
11	Septicemia (A40-A41)	22	1.0	2.5
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21	1.0	2.4
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	19	0.9	*
14	Aortic aneurysm and dissection (I71)	16	0.8	*
14	Influenza and pneumonia (J10-J18)	16	0.8	*
...	All other causes (Residual)	225	10.7	26.0
Asian or Pacific Islander, male, 55-64 years				
...	All causes	2,893	100.0	545.3
1	Malignant neoplasms (C00-C97)	1,054	36.4	198.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	737	25.5	138.9
3	Cerebrovascular diseases (I60-I69)	195	6.7	36.8
4	Diabetes mellitus (E10-E14)	135	4.7	25.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	118	4.1	22.2
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	54	1.9	10.2
7	Chronic lower respiratory diseases (J40-J47)	51	1.8	9.6
8	Chronic liver disease and cirrhosis (K70,K73-K74)	48	1.7	9.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	46	1.6	8.7
10	Septicemia (A40-A41)	41	1.4	7.7
11	Influenza and pneumonia (J10-J18)	34	1.2	6.4
12	Viral hepatitis (B15-B19)	30	1.0	5.7
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	28	1.0	5.3
14	Aortic aneurysm and dissection (I71)	26	0.9	4.9
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	23	0.8	4.3
...	All other causes (Residual)	273	9.4	51.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 65-74 years				
...	All causes	3,988	100.0	1,363.4
1	Malignant neoplasms (C00-C97)	1,453	36.4	496.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,017	25.5	347.7
3	Cerebrovascular diseases (I60-I69)	300	7.5	102.6
4	Diabetes mellitus (E10-E14)	176	4.4	60.2
5	Chronic lower respiratory diseases (J40-J47)	137	3.4	46.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	87	2.2	29.7
7	Influenza and pneumonia (J10-J18)	79	2.0	27.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	77	1.9	26.3
9	Septicemia (A40-A41)	50	1.3	17.1
10	Aortic aneurysm and dissection (I71)	43	1.1	14.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	41	1.0	14.0
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	38	1.0	13.0
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	37	0.9	12.6
14	Parkinson's disease (G20-G21)	27	0.7	9.2
15	Viral hepatitis (B15-B19)	22	0.6	7.5
...	All other causes (Residual)	404	10.1	138.1
Asian or Pacific Islander, male, 75-84 years				
...	All causes	5,486	100.0	3,766.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,525	27.8	1,047.0
2	Malignant neoplasms (C00-C97)	1,488	27.1	1,021.6
3	Cerebrovascular diseases (I60-I69)	511	9.3	350.8
4	Chronic lower respiratory diseases (J40-J47)	291	5.3	199.8
5	Diabetes mellitus (E10-E14)	205	3.7	140.7
6	Influenza and pneumonia (J10-J18)	200	3.6	137.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	120	2.2	82.4
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	106	1.9	72.8
9	Parkinson's disease (G20-G21)	75	1.4	51.5
10	Septicemia (A40-A41)	73	1.3	50.1
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	71	1.3	48.7
12	Aortic aneurysm and dissection (I71)	66	1.2	45.3
13	Alzheimer's disease (G30)	64	1.2	43.9
14	Pneumonitis due to solids and liquids (J69)	49	0.9	33.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	40	0.7	27.5
...	All other causes (Residual)	602	11.0	413.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 85 years and over				
...	All causes	3,989	100.0	10,118.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,347	33.8	3,416.7
2	Malignant neoplasms (C00-C97)	612	15.3	1,552.4
3	Cerebrovascular diseases (I60-I69)	382	9.6	969.0
4	Influenza and pneumonia (J10-J18)	246	6.2	624.0
5	Chronic lower respiratory diseases (J40-J47)	235	5.9	596.1
6	Diabetes mellitus (E10-E14)	115	2.9	291.7
7	Alzheimer's disease (G30)	108	2.7	273.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	88	2.2	223.2
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	84	2.1	213.1
10	Pneumonitis due to solids and liquids (J69)	75	1.9	190.2
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	64	1.6	162.3
12	Parkinson's disease (G20-G21)	57	1.4	144.6
13	Septicemia (A40-A41)	45	1.1	114.1
14	Aortic aneurysm and dissection (I71)	29	0.7	73.6
15	Atherosclerosis (I70)	22	0.6	55.8
...	All other causes (Residual)	480	12.0	1,217.5
Asian or Pacific Islander, female, all ages				
...	All causes	19,235	100.0	274.6
1	Malignant neoplasms (C00-C97)	5,167	26.9	73.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,560	23.7	65.1
3	Cerebrovascular diseases (I60-I69)	1,893	9.8	27.0
4	Diabetes mellitus (E10-E14)	763	4.0	10.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	762	4.0	10.9
6	Influenza and pneumonia (J10-J18)	647	3.4	9.2
7	Chronic lower respiratory diseases (J40-J47)	444	2.3	6.3
8	Alzheimer's disease (G30)	344	1.8	4.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	336	1.7	4.8
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	312	1.6	4.5
11	Septicemia (A40-A41)	252	1.3	3.6
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	241	1.3	3.4
13	Certain conditions originating in the perinatal period (P00-P96)	188	1.0	2.7
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	146	0.8	2.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	129	0.7	1.8
...	All other causes (Residual)	3,051	15.9	43.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 1-4 years				
...	All causes	81	100.0	22.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17	21.0	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	13	16.0	*
3	Malignant neoplasms (C00-C97)	8	9.9	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	6	7.4	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	3.7	*
6	Septicemia (A40-A41)	2	2.5	*
6	Chronic lower respiratory diseases (J40-J47)	2	2.5	*
8	Anemias (D50-D64)	1	1.2	*
8	Cerebrovascular diseases (I60-I69)	1	1.2	*
8	Influenza and pneumonia (J10-J18)	1	1.2	*
8	Peptic ulcer (K25-K28)	1	1.2	*
8	Certain conditions originating in the perinatal period (P00-P96)	1	1.2	*
...	All other causes (Residual)	25	30.9	6.8
Asian or Pacific Islander, female, 5-14 years				
...	All causes	87	100.0	9.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	28	32.2	3.1
2	Malignant neoplasms (C00-C97)	20	23.0	2.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	5.7	*
4	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	3.4	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	3.4	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	2.3	*
6	Pregnancy, childbirth and the puerperium (O00-099)	2	2.3	*
8	Anemias (D50-D64)	1	1.1	*
8	Diabetes mellitus (E10-E14)	1	1.1	*
8	Cerebrovascular diseases (I60-I69)	1	1.1	*
8	Influenza and pneumonia (J10-J18)	1	1.1	*
8	Chronic lower respiratory diseases (J40-J47)	1	1.1	*
8	Diseases of appendix (K35-K38)	1	1.1	*
...	All other causes (Residual)	18	20.7	*

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 15-24 years				
...	All causes	230	100.0	24.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	97	42.2	10.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	27	11.7	2.8
3	Malignant neoplasms (C00-C97)	18	7.8	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	11	4.8	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	11	4.8	*
6	Cerebrovascular diseases (I60-I69)	6	2.6	*
7	Pregnancy, childbirth and the puerperium (O00-099)	5	2.2	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	2.2	*
9	Septicemia (A40-A41)	3	1.3	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	1.3	*
11	Aortic aneurysm and dissection (I71)	2	0.9	*
11	Influenza and pneumonia (J10-J18)	2	0.9	*
11	Cholelithiasis and other disorders of gallbladder (K80-K82)	2	0.9	*
14	Tuberculosis (A16-A19)	1	0.4	*
14	Viral hepatitis (B15-B19)	1	0.4	*
14	Nutritional deficiencies (E40-E64)	1	0.4	*
14	Meningitis (G00,G03)	1	0.4	*
14	Chronic lower respiratory diseases (J40-J47)	1	0.4	*
...	All other causes (Residual)	33	14.3	3.5
Asian or Pacific Islander, female, 25-34 years				
...	All causes	343	100.0	26.8
1	Malignant neoplasms (C00-C97)	85	24.8	6.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	69	20.1	5.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	52	15.2	4.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	23	6.7	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	17	5.0	*
6	Cerebrovascular diseases (I60-I69)	9	2.6	*
6	Pregnancy, childbirth and the puerperium (O00-099)	9	2.6	*
8	Diabetes mellitus (E10-E14)	5	1.5	*
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	1.2	*
10	Viral hepatitis (B15-B19)	3	0.9	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	3	0.9	*
12	Septicemia (A40-A41)	2	0.6	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.6	*
12	Aortic aneurysm and dissection (I71)	2	0.6	*
12	Pneumonitis due to solids and liquids (J69)	2	0.6	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.6	*
...	All other causes (Residual)	54	15.7	4.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 35-44 years				
...	All causes	639	100.0	54.2
1	Malignant neoplasms (C00-C97)	286	44.8	24.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	78	12.2	6.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	50	7.8	4.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	49	7.7	4.2
5	Cerebrovascular diseases (I60-I69)	30	4.7	2.5
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	15	2.3	*
7	Diabetes mellitus (E10-E14)	11	1.7	*
8	Pregnancy, childbirth and the puerperium (O00-099)	10	1.6	*
9	Septicemia (A40-A41)	9	1.4	*
10	Chronic liver disease and cirrhosis (K70,K73-K74)	6	0.9	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	5	0.8	*
11	Influenza and pneumonia (J10-J18)	5	0.8	*
13	Anemias (D50-D64)	4	0.6	*
13	Chronic lower respiratory diseases (J40-J47)	4	0.6	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	0.6	*
...	All other causes (Residual)	73	11.4	6.2
Asian or Pacific Islander, female, 45-54 years				
...	All causes	1,433	100.0	145.8
1	Malignant neoplasms (C00-C97)	757	52.8	77.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	135	9.4	13.7
3	Cerebrovascular diseases (I60-I69)	103	7.2	10.5
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	84	5.9	8.5
5	Diabetes mellitus (E10-E14)	44	3.1	4.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	43	3.0	4.4
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	21	1.5	2.1
8	Septicemia (A40-A41)	19	1.3	*
9	Influenza and pneumonia (J10-J18)	16	1.1	*
9	Chronic lower respiratory diseases (J40-J47)	16	1.1	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	0.9	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	0.8	*
13	Viral hepatitis (B15-B19)	9	0.6	*
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	9	0.6	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	6	0.4	*
15	Aortic aneurysm and dissection (I71)	6	0.4	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	6	0.4	*
...	All other causes (Residual)	135	9.4	13.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 55-64 years				
...	All causes	2,070	100.0	339.9
1	Malignant neoplasms (C00-C97)	969	46.8	159.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	309	14.9	50.7
3	Cerebrovascular diseases (I60-I69)	171	8.3	28.1
4	Diabetes mellitus (E10-E14)	88	4.3	14.5
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	70	3.4	11.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	35	1.7	5.7
7	Septicemia (A40-A41)	28	1.4	4.6
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	28	1.4	4.6
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	26	1.3	4.3
9	Influenza and pneumonia (J10-J18)	26	1.3	4.3
11	Chronic lower respiratory diseases (J40-J47)	25	1.2	4.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	25	1.2	4.1
13	Viral hepatitis (B15-B19)	18	0.9	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	0.6	*
15	Aortic aneurysm and dissection (I71)	8	0.4	*
...	All other causes (Residual)	231	11.2	37.9
Asian or Pacific Islander, female, 65-74 years				
...	All causes	3,405	100.0	933.2
1	Malignant neoplasms (C00-C97)	1,256	36.9	344.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	750	22.0	205.6
3	Cerebrovascular diseases (I60-I69)	285	8.4	78.1
4	Diabetes mellitus (E10-E14)	157	4.6	43.0
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	99	2.9	27.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	75	2.2	20.6
7	Chronic lower respiratory diseases (J40-J47)	74	2.2	20.3
8	Influenza and pneumonia (J10-J18)	72	2.1	19.7
9	Septicemia (A40-A41)	48	1.4	13.2
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	45	1.3	12.3
11	Viral hepatitis (B15-B19)	36	1.1	9.9
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	27	0.8	7.4
12	Chronic liver disease and cirrhosis (K70,K73-K74)	27	0.8	7.4
14	Aortic aneurysm and dissection (I71)	20	0.6	5.5
15	Alzheimer's disease (G30)	19	0.6	*
...	All other causes (Residual)	415	12.2	113.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2004

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 75-84 years				
...	All causes	5,330	100.0	2,558.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,453	27.3	697.4
2	Malignant neoplasms (C00-C97)	1,205	22.6	578.4
3	Cerebrovascular diseases (I60-I69)	651	12.2	312.5
4	Diabetes mellitus (E10-E14)	269	5.0	129.1
5	Influenza and pneumonia (J10-J18)	197	3.7	94.6
6	Chronic lower respiratory diseases (J40-J47)	158	3.0	75.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	126	2.4	60.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	123	2.3	59.0
9	Alzheimer's disease (G30)	122	2.3	58.6
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	94	1.8	45.1
11	Septicemia (A40-A41)	75	1.4	36.0
12	Parkinson's disease (G20-G21)	49	0.9	23.5
13	Chronic liver disease and cirrhosis (K70,K73-K74)	42	0.8	20.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	38	0.7	18.2
15	Aortic aneurysm and dissection (I71)	34	0.6	16.3
...	All other causes (Residual)	694	13.0	333.1
Asian or Pacific Islander, female, 85 years and over				
...	All causes	5,241	100.0	8,125.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,817	34.7	2,817.1
2	Cerebrovascular diseases (I60-I69)	632	12.1	979.9
3	Malignant neoplasms (C00-C97)	563	10.7	872.9
4	Influenza and pneumonia (J10-J18)	324	6.2	502.3
5	Alzheimer's disease (G30)	199	3.8	308.5
6	Diabetes mellitus (E10-E14)	188	3.6	291.5
7	Chronic lower respiratory diseases (J40-J47)	163	3.1	252.7
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	136	2.6	210.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	87	1.7	134.9
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	86	1.6	133.3
11	Septicemia (A40-A41)	65	1.2	100.8
12	Pneumonitis due to solids and liquids (J69)	57	1.1	88.4
13	Parkinson's disease (G20-G21)	43	0.8	66.7
14	Atherosclerosis (I70)	41	0.8	63.6
15	Aortic aneurysm and dissection (I71)	31	0.6	48.1
...	All other causes (Residual)	809	15.4	1,254.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable