

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, all ages				
...	All causes	2,448,017	100.0	825.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	652,091	26.6	220.0
2	Malignant neoplasms (C00-C97)	559,312	22.8	188.7
3	Cerebrovascular diseases (I60-I69)	143,579	5.9	48.4
4	Chronic lower respiratory diseases (J40-J47)	130,933	5.3	44.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	117,809	4.8	39.7
6	Diabetes mellitus (E10-E14)	75,119	3.1	25.3
7	Alzheimer's disease (G30)	71,599	2.9	24.2
8	Influenza and pneumonia (J10-J18)	63,001	2.6	21.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	43,901	1.8	14.8
10	Septicemia (A40-A41)	34,136	1.4	11.5
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	32,637	1.3	11.0
12	Chronic liver disease and cirrhosis (K70,K73-K74)	27,530	1.1	9.3
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	24,902	1.0	8.4
14	Parkinson's disease (G20-G21)	19,544	0.8	6.6
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	18,124	0.7	6.1
...	All other causes (Residual)	433,800	17.7	146.4
All races, both sexes, 1-4 years				
...	All causes	4,756	100.0	29.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,664	35.0	10.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	522	11.0	3.2
3	Malignant neoplasms (C00-C97)	377	7.9	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	375	7.9	2.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	151	3.2	0.9
6	Influenza and pneumonia (J10-J18)	110	2.3	0.7
7	Septicemia (A40-A41)	85	1.8	0.5
8	Cerebrovascular diseases (I60-I69)	62	1.3	0.4
9	Certain conditions originating in the perinatal period (P00-P96)	58	1.2	0.4
10	Chronic lower respiratory diseases (J40-J47)	56	1.2	0.3
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	52	1.1	0.3
12	Meningitis (G00,G03)	32	0.7	0.2
13	Anemias (D50-D64)	23	0.5	0.1
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	16	0.3	*
...	All other causes (Residual)	1,157	24.3	7.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 5-14 years				
...	All causes	6,602	100.0	16.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,415	36.6	6.0
2	Malignant neoplasms (C00-C97)	1,000	15.1	2.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	396	6.0	1.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	341	5.2	0.8
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	272	4.1	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	252	3.8	0.6
7	Influenza and pneumonia (J10-J18)	106	1.6	0.3
8	Chronic lower respiratory diseases (J40-J47)	104	1.6	0.3
9	Cerebrovascular diseases (I60-I69)	95	1.4	0.2
10	Septicemia (A40-A41)	81	1.2	0.2
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	76	1.2	0.2
12	Diabetes mellitus (E10-E14)	38	0.6	0.1
13	Anemias (D50-D64)	26	0.4	0.1
13	Meningitis (G00,G03)	26	0.4	0.1
13	Certain conditions originating in the perinatal period (P00-P96)	26	0.4	0.1
...	All other causes (Residual)	1,348	20.4	3.3
All races, both sexes, 15-24 years				
...	All causes	34,234	100.0	81.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15,753	46.0	37.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5,466	16.0	13.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,212	12.3	10.0
4	Malignant neoplasms (C00-C97)	1,717	5.0	4.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,119	3.3	2.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	504	1.5	1.2
7	Diabetes mellitus (E10-E14)	202	0.6	0.5
8	Cerebrovascular diseases (I60-I69)	196	0.6	0.5
9	Pregnancy, childbirth and the puerperium (O00-099)	183	0.5	0.4
10	Influenza and pneumonia (J10-J18)	172	0.5	0.4
11	Human immunodeficiency virus (HIV) disease (B20-B24)	159	0.5	0.4
12	Septicemia (A40-A41)	148	0.4	0.4
12	Chronic lower respiratory diseases (J40-J47)	148	0.4	0.4
14	Anemias (D50-D64)	103	0.3	0.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	93	0.3	0.2
...	All other causes (Residual)	4,059	11.9	9.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 25-44 years				
...	All causes	126,710	100.0	150.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	30,916	24.4	36.8
2	Malignant neoplasms (C00-C97)	18,167	14.3	21.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,937	12.6	19.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11,540	9.1	13.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7,861	6.2	9.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	5,681	4.5	6.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,999	2.4	3.6
8	Cerebrovascular diseases (I60-I69)	2,806	2.2	3.3
9	Diabetes mellitus (E10-E14)	2,662	2.1	3.2
10	Influenza and pneumonia (J10-J18)	1,288	1.0	1.5
11	Septicemia (A40-A41)	1,151	0.9	1.4
12	Chronic lower respiratory diseases (J40-J47)	1,148	0.9	1.4
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,027	0.8	1.2
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	953	0.8	1.1
15	Viral hepatitis (B15-B19)	570	0.4	0.7
...	All other causes (Residual)	22,004	17.4	26.2
All races, both sexes, 45-64 years				
...	All causes	458,831	100.0	629.9
1	Malignant neoplasms (C00-C97)	149,645	32.6	205.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	103,311	22.5	141.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	29,192	6.4	40.1
4	Diabetes mellitus (E10-E14)	16,992	3.7	23.3
5	Chronic lower respiratory diseases (J40-J47)	16,724	3.6	23.0
6	Cerebrovascular diseases (I60-I69)	16,409	3.6	22.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	14,643	3.2	20.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11,201	2.4	15.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,169	1.3	8.5
10	Human immunodeficiency virus (HIV) disease (B20-B24)	6,128	1.3	8.4
11	Septicemia (A40-A41)	6,123	1.3	8.4
12	Influenza and pneumonia (J10-J18)	5,605	1.2	7.7
13	Viral hepatitis (B15-B19)	3,691	0.8	5.1
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	3,120	0.7	4.3
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,922	0.6	4.0
...	All other causes (Residual)	66,956	14.6	91.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 65 years and over				
...	All causes	1,788,189	100.0	4,860.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	530,926	29.7	1,443.1
2	Malignant neoplasms (C00-C97)	388,322	21.7	1,055.5
3	Cerebrovascular diseases (I60-I69)	123,881	6.9	336.7
4	Chronic lower respiratory diseases (J40-J47)	112,716	6.3	306.4
5	Alzheimer's disease (G30)	70,858	4.0	192.6
6	Influenza and pneumonia (J10-J18)	55,453	3.1	150.7
7	Diabetes mellitus (E10-E14)	55,222	3.1	150.1
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	36,729	2.1	99.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	36,416	2.0	99.0
10	Septicemia (A40-A41)	26,243	1.5	71.3
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	21,265	1.2	57.8
12	Parkinson's disease (G20-G21)	19,030	1.1	51.7
13	Pneumonitis due to solids and liquids (J69)	15,651	0.9	42.5
14	Atherosclerosis (I70)	11,168	0.6	30.4
15	Aortic aneurysm and dissection (I71)	11,147	0.6	30.3
...	All other causes (Residual)	273,162	15.3	742.5
All races, male, all ages				
...	All causes	1,207,675	100.0	827.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	322,841	26.7	221.1
2	Malignant neoplasms (C00-C97)	290,422	24.0	198.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	76,375	6.3	52.3
4	Chronic lower respiratory diseases (J40-J47)	62,435	5.2	42.8
5	Cerebrovascular diseases (I60-I69)	56,586	4.7	38.8
6	Diabetes mellitus (E10-E14)	36,538	3.0	25.0
7	Influenza and pneumonia (J10-J18)	28,052	2.3	19.2
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	25,907	2.1	17.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21,268	1.8	14.6
10	Alzheimer's disease (G30)	20,559	1.7	14.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	17,937	1.5	12.3
12	Septicemia (A40-A41)	15,322	1.3	10.5
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	14,376	1.2	9.8
14	Parkinson's disease (G20-G21)	11,247	0.9	7.7
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	9,458	0.8	6.5
...	All other causes (Residual)	198,352	16.4	135.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 1-4 years				
...	All causes	2,765	100.0	33.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,027	37.1	12.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	264	9.5	3.2
3	Malignant neoplasms (C00-C97)	217	7.8	2.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	214	7.7	2.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	81	2.9	1.0
6	Influenza and pneumonia (J10-J18)	56	2.0	0.7
7	Septicemia (A40-A41)	47	1.7	0.6
8	Cerebrovascular diseases (I60-I69)	41	1.5	0.5
9	Certain conditions originating in the perinatal period (P00-P96)	36	1.3	0.4
10	Chronic lower respiratory diseases (J40-J47)	33	1.2	0.4
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	31	1.1	0.4
12	Meningitis (G00,G03)	21	0.8	0.3
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	10	0.4	*
15	Anemias (D50-D64)	9	0.3	*
...	All other causes (Residual)	666	24.1	8.0
All races, male, 5-14 years				
...	All causes	3,853	100.0	18.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,493	38.7	7.2
2	Malignant neoplasms (C00-C97)	557	14.5	2.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	204	5.3	1.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	204	5.3	1.0
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	195	5.1	0.9
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	131	3.4	0.6
7	Chronic lower respiratory diseases (J40-J47)	63	1.6	0.3
8	Cerebrovascular diseases (I60-I69)	55	1.4	0.3
9	Influenza and pneumonia (J10-J18)	53	1.4	0.3
10	Septicemia (A40-A41)	49	1.3	0.2
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	36	0.9	0.2
12	Diabetes mellitus (E10-E14)	15	0.4	*
12	Meningitis (G00,G03)	15	0.4	*
14	Anemias (D50-D64)	14	0.4	*
15	Certain conditions originating in the perinatal period (P00-P96)	11	0.3	*
...	All other causes (Residual)	758	19.7	3.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 15-24 years				
...	All causes	25,509	100.0	117.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,827	46.4	54.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4,765	18.7	22.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,498	13.7	16.2
4	Malignant neoplasms (C00-C97)	1,035	4.1	4.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	770	3.0	3.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	326	1.3	1.5
7	Influenza and pneumonia (J10-J18)	117	0.5	0.5
8	Diabetes mellitus (E10-E14)	115	0.5	0.5
9	Chronic lower respiratory diseases (J40-J47)	95	0.4	0.4
10	Cerebrovascular diseases (I60-I69)	94	0.4	0.4
11	Human immunodeficiency virus (HIV) disease (B20-B24)	89	0.3	0.4
12	Septicemia (A40-A41)	82	0.3	0.4
12	Legal intervention (Y35,Y89.0)	82	0.3	0.4
14	Anemias (D50-D64)	57	0.2	0.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	54	0.2	0.2
...	All other causes (Residual)	2,503	9.8	11.6
All races, male, 25-44 years				
...	All causes	82,592	100.0	195.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	22,691	27.5	53.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,137	13.5	26.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	9,129	11.1	21.6
4	Malignant neoplasms (C00-C97)	8,143	9.9	19.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6,325	7.7	14.9
6	Human immunodeficiency virus (HIV) disease (B20-B24)	3,955	4.8	9.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,982	2.4	4.7
8	Diabetes mellitus (E10-E14)	1,642	2.0	3.9
9	Cerebrovascular diseases (I60-I69)	1,449	1.8	3.4
10	Influenza and pneumonia (J10-J18)	736	0.9	1.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	576	0.7	1.4
12	Septicemia (A40-A41)	566	0.7	1.3
13	Chronic lower respiratory diseases (J40-J47)	556	0.7	1.3
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	516	0.6	1.2
15	Viral hepatitis (B15-B19)	383	0.5	0.9
...	All other causes (Residual)	12,806	15.5	30.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 45-64 years				
...	All causes	279,901	100.0	788.0
1	Malignant neoplasms (C00-C97)	79,442	28.4	223.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	72,381	25.9	203.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	20,509	7.3	57.7
4	Chronic liver disease and cirrhosis (K70,K73-K74)	10,517	3.8	29.6
5	Diabetes mellitus (E10-E14)	10,037	3.6	28.3
6	Cerebrovascular diseases (I60-I69)	9,082	3.2	25.6
7	Chronic lower respiratory diseases (J40-J47)	8,623	3.1	24.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8,519	3.0	24.0
9	Human immunodeficiency virus (HIV) disease (B20-B24)	4,709	1.7	13.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,589	1.3	10.1
11	Influenza and pneumonia (J10-J18)	3,287	1.2	9.3
12	Septicemia (A40-A41)	3,249	1.2	9.1
13	Viral hepatitis (B15-B19)	2,674	1.0	7.5
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,215	0.8	6.2
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,884	0.7	5.3
...	All other causes (Residual)	39,184	14.0	110.3
All races, male, 65 years and over				
...	All causes	796,838	100.0	5,170.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	238,119	29.9	1,545.0
2	Malignant neoplasms (C00-C97)	200,978	25.2	1,304.0
3	Chronic lower respiratory diseases (J40-J47)	53,043	6.7	344.2
4	Cerebrovascular diseases (I60-I69)	45,788	5.7	297.1
5	Diabetes mellitus (E10-E14)	24,729	3.1	160.4
6	Influenza and pneumonia (J10-J18)	23,653	3.0	153.5
7	Alzheimer's disease (G30)	20,236	2.5	131.3
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18,201	2.3	118.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16,948	2.1	110.0
10	Septicemia (A40-A41)	11,156	1.4	72.4
11	Parkinson's disease (G20-G21)	10,920	1.4	70.9
12	Pneumonitis due to solids and liquids (J69)	7,799	1.0	50.6
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	7,272	0.9	47.2
14	Aortic aneurysm and dissection (I71)	6,149	0.8	39.9
15	Chronic liver disease and cirrhosis (K70,K73-K74)	5,414	0.7	35.1
...	All other causes (Residual)	106,433	13.4	690.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, all ages				
...	All causes	1,240,342	100.0	824.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	329,250	26.5	218.9
2	Malignant neoplasms (C00-C97)	268,890	21.7	178.8
3	Cerebrovascular diseases (I60-I69)	86,993	7.0	57.8
4	Chronic lower respiratory diseases (J40-J47)	68,498	5.5	45.5
5	Alzheimer's disease (G30)	51,040	4.1	33.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	41,434	3.3	27.5
7	Diabetes mellitus (E10-E14)	38,581	3.1	25.7
8	Influenza and pneumonia (J10-J18)	34,949	2.8	23.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22,633	1.8	15.0
10	Septicemia (A40-A41)	18,814	1.5	12.5
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	15,444	1.2	10.3
12	Chronic liver disease and cirrhosis (K70,K73-K74)	9,593	0.8	6.4
13	Pneumonitis due to solids and liquids (J69)	8,519	0.7	5.7
14	Parkinson's disease (G20-G21)	8,297	0.7	5.5
15	Atherosclerosis (I70)	7,369	0.6	4.9
...	All other causes (Residual)	230,038	18.5	152.9
All races, female, 1-4 years				
...	All causes	1,991	100.0	25.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	637	32.0	8.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	258	13.0	3.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	161	8.1	2.0
4	Malignant neoplasms (C00-C97)	160	8.0	2.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	70	3.5	0.9
6	Influenza and pneumonia (J10-J18)	54	2.7	0.7
7	Septicemia (A40-A41)	38	1.9	0.5
8	Chronic lower respiratory diseases (J40-J47)	23	1.2	0.3
9	Certain conditions originating in the perinatal period (P00-P96)	22	1.1	0.3
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	21	1.1	0.3
10	Cerebrovascular diseases (I60-I69)	21	1.1	0.3
12	Anemias (D50-D64)	14	0.7	*
13	Meningitis (G00,G03)	11	0.6	*
14	Acute bronchitis and bronchiolitis (J20-J21)	6	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.3	*
...	All other causes (Residual)	489	24.6	6.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 5-14 years				
...	All causes	2,749	100.0	13.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	922	33.5	4.7
2	Malignant neoplasms (C00-C97)	443	16.1	2.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	201	7.3	1.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	137	5.0	0.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	121	4.4	0.6
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	68	2.5	0.3
7	Influenza and pneumonia (J10-J18)	53	1.9	0.3
8	Chronic lower respiratory diseases (J40-J47)	41	1.5	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	40	1.5	0.2
9	Cerebrovascular diseases (I60-I69)	40	1.5	0.2
11	Septicemia (A40-A41)	32	1.2	0.2
12	Diabetes mellitus (E10-E14)	23	0.8	0.1
13	Certain conditions originating in the perinatal period (P00-P96)	15	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	0.5	*
15	Anemias (D50-D64)	12	0.4	*
...	All other causes (Residual)	588	21.4	3.0
All races, female, 15-24 years				
...	All causes	8,725	100.0	42.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,926	45.0	19.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	714	8.2	3.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	701	8.0	3.4
4	Malignant neoplasms (C00-C97)	682	7.8	3.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	349	4.0	1.7
6	Pregnancy, childbirth and the puerperium (O00-099)	183	2.1	0.9
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	178	2.0	0.9
8	Cerebrovascular diseases (I60-I69)	102	1.2	0.5
9	Diabetes mellitus (E10-E14)	87	1.0	0.4
10	Human immunodeficiency virus (HIV) disease (B20-B24)	70	0.8	0.3
11	Septicemia (A40-A41)	66	0.8	0.3
12	Influenza and pneumonia (J10-J18)	55	0.6	0.3
13	Chronic lower respiratory diseases (J40-J47)	53	0.6	0.3
14	Anemias (D50-D64)	46	0.5	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	46	0.5	0.2
...	All other causes (Residual)	1,467	16.8	7.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 25-44 years				
...	All causes	44,118	100.0	105.9
1	Malignant neoplasms (C00-C97)	10,024	22.7	24.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,225	18.6	19.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,800	10.9	11.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,411	5.5	5.8
5	Human immunodeficiency virus (HIV) disease (B20-B24)	1,726	3.9	4.1
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,536	3.5	3.7
7	Cerebrovascular diseases (I60-I69)	1,357	3.1	3.3
8	Diabetes mellitus (E10-E14)	1,020	2.3	2.4
9	Chronic liver disease and cirrhosis (K70,K73-K74)	1,017	2.3	2.4
10	Chronic lower respiratory diseases (J40-J47)	592	1.3	1.4
11	Septicemia (A40-A41)	585	1.3	1.4
12	Influenza and pneumonia (J10-J18)	552	1.3	1.3
13	Pregnancy, childbirth and the puerperium (O00-099)	519	1.2	1.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	451	1.0	1.1
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	437	1.0	1.0
...	All other causes (Residual)	8,866	20.1	21.3
All races, female, 45-64 years				
...	All causes	178,930	100.0	479.5
1	Malignant neoplasms (C00-C97)	70,203	39.2	188.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	30,930	17.3	82.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,683	4.9	23.3
4	Chronic lower respiratory diseases (J40-J47)	8,101	4.5	21.7
5	Cerebrovascular diseases (I60-I69)	7,327	4.1	19.6
6	Diabetes mellitus (E10-E14)	6,955	3.9	18.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	4,126	2.3	11.1
8	Septicemia (A40-A41)	2,874	1.6	7.7
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,682	1.5	7.2
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,580	1.4	6.9
11	Influenza and pneumonia (J10-J18)	2,318	1.3	6.2
12	Human immunodeficiency virus (HIV) disease (B20-B24)	1,419	0.8	3.8
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,236	0.7	3.3
14	Viral hepatitis (B15-B19)	1,017	0.6	2.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	889	0.5	2.4
...	All other causes (Residual)	27,590	15.4	73.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 65 years and over				
...	All causes	991,351	100.0	4,637.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	292,807	29.5	1,369.7
2	Malignant neoplasms (C00-C97)	187,344	18.9	876.4
3	Cerebrovascular diseases (I60-I69)	78,093	7.9	365.3
4	Chronic lower respiratory diseases (J40-J47)	59,673	6.0	279.1
5	Alzheimer's disease (G30)	50,622	5.1	236.8
6	Influenza and pneumonia (J10-J18)	31,800	3.2	148.8
7	Diabetes mellitus (E10-E14)	30,493	3.1	142.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19,468	2.0	91.1
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18,528	1.9	86.7
10	Septicemia (A40-A41)	15,087	1.5	70.6
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	13,993	1.4	65.5
12	Parkinson's disease (G20-G21)	8,110	0.8	37.9
13	Pneumonitis due to solids and liquids (J69)	7,852	0.8	36.7
14	Atherosclerosis (I70)	7,124	0.7	33.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5,606	0.6	26.2
...	All other causes (Residual)	164,751	16.6	770.7
White, both sexes, all ages				
...	All causes	2,098,097	100.0	873.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	564,796	26.9	235.2
2	Malignant neoplasms (C00-C97)	482,132	23.0	200.8
3	Cerebrovascular diseases (I60-I69)	121,868	5.8	50.7
4	Chronic lower respiratory diseases (J40-J47)	120,884	5.8	50.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	100,406	4.8	41.8
6	Alzheimer's disease (G30)	66,191	3.2	27.6
7	Diabetes mellitus (E10-E14)	59,755	2.8	24.9
8	Influenza and pneumonia (J10-J18)	55,540	2.6	23.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	34,806	1.7	14.5
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	29,527	1.4	12.3
11	Septicemia (A40-A41)	27,194	1.3	11.3
12	Chronic liver disease and cirrhosis (K70,K73-K74)	24,003	1.1	10.0
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	19,254	0.9	8.0
14	Parkinson's disease (G20-G21)	18,496	0.9	7.7
15	Pneumonitis due to solids and liquids (J69)	15,338	0.7	6.4
...	All other causes (Residual)	357,907	17.1	149.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 1-4 years				
...	All causes	3,408	100.0	27.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,262	37.0	10.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	385	11.3	3.1
3	Malignant neoplasms (C00-C97)	293	8.6	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	217	6.4	1.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	93	2.7	0.7
6	Influenza and pneumonia (J10-J18)	76	2.2	0.6
7	Septicemia (A40-A41)	59	1.7	0.5
8	Cerebrovascular diseases (I60-I69)	45	1.3	0.4
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	42	1.2	0.3
10	Certain conditions originating in the perinatal period (P00-P96)	35	1.0	0.3
11	Chronic lower respiratory diseases (J40-J47)	24	0.7	0.2
12	Meningitis (G00,G03)	20	0.6	0.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	13	0.4	*
15	Acute bronchitis and bronchiolitis (J20-J21)	12	0.4	*
...	All other causes (Residual)	819	24.0	6.5
White, both sexes, 5-14 years				
...	All causes	4,710	100.0	15.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,785	37.9	5.7
2	Malignant neoplasms (C00-C97)	787	16.7	2.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	285	6.1	0.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	209	4.4	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	168	3.6	0.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	159	3.4	0.5
7	Influenza and pneumonia (J10-J18)	77	1.6	0.2
8	Cerebrovascular diseases (I60-I69)	64	1.4	0.2
9	Septicemia (A40-A41)	57	1.2	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	57	1.2	0.2
11	Chronic lower respiratory diseases (J40-J47)	45	1.0	0.1
12	Diabetes mellitus (E10-E14)	25	0.5	0.1
13	Certain conditions originating in the perinatal period (P00-P96)	22	0.5	0.1
14	Meningitis (G00,G03)	18	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18	0.4	*
...	All other causes (Residual)	934	19.8	3.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 15-24 years				
...	All causes	25,410	100.0	77.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,325	52.4	40.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,540	13.9	10.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,172	8.5	6.6
4	Malignant neoplasms (C00-C97)	1,365	5.4	4.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	736	2.9	2.2
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	386	1.5	1.2
7	Cerebrovascular diseases (I60-I69)	136	0.5	0.4
8	Influenza and pneumonia (J10-J18)	134	0.5	0.4
9	Diabetes mellitus (E10-E14)	124	0.5	0.4
10	Pregnancy, childbirth and the puerperium (O00-099)	110	0.4	0.3
11	Septicemia (A40-A41)	95	0.4	0.3
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	76	0.3	0.2
13	Chronic lower respiratory diseases (J40-J47)	71	0.3	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	60	0.2	0.2
15	Legal intervention (Y35,Y89.0)	59	0.2	0.2
...	All other causes (Residual)	3,021	11.9	9.2
White, both sexes, 25-44 years				
...	All causes	93,687	100.0	140.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	25,800	27.5	38.6
2	Malignant neoplasms (C00-C97)	14,046	15.0	21.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,185	11.9	16.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	10,146	10.8	15.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,637	3.9	5.4
6	Chronic liver disease and cirrhosis (K70,K73-K74)	2,515	2.7	3.8
7	Human immunodeficiency virus (HIV) disease (B20-B24)	2,422	2.6	3.6
8	Cerebrovascular diseases (I60-I69)	1,806	1.9	2.7
9	Diabetes mellitus (E10-E14)	1,800	1.9	2.7
10	Influenza and pneumonia (J10-J18)	945	1.0	1.4
11	Chronic lower respiratory diseases (J40-J47)	799	0.9	1.2
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	766	0.8	1.1
13	Septicemia (A40-A41)	743	0.8	1.1
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	553	0.6	0.8
15	Viral hepatitis (B15-B19)	461	0.5	0.7
...	All other causes (Residual)	16,063	17.1	24.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 45-64 years				
...	All causes	362,442	100.0	595.1
1	Malignant neoplasms (C00-C97)	122,156	33.7	200.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	79,781	22.0	131.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	24,167	6.7	39.7
4	Chronic lower respiratory diseases (J40-J47)	14,580	4.0	23.9
5	Diabetes mellitus (E10-E14)	12,537	3.5	20.6
6	Chronic liver disease and cirrhosis (K70,K73-K74)	12,480	3.4	20.5
7	Cerebrovascular diseases (I60-I69)	11,223	3.1	18.4
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	10,484	2.9	17.2
9	Septicemia (A40-A41)	4,376	1.2	7.2
10	Influenza and pneumonia (J10-J18)	4,323	1.2	7.1
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,884	1.1	6.4
12	Viral hepatitis (B15-B19)	2,954	0.8	4.8
13	Human immunodeficiency virus (HIV) disease (B20-B24)	2,630	0.7	4.3
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,813	0.5	3.0
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,755	0.5	2.9
...	All other causes (Residual)	53,299	14.7	87.5
White, both sexes, 65 years and over				
...	All causes	1,589,721	100.0	4,929.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	472,587	29.7	1,465.5
2	Malignant neoplasms (C00-C97)	343,416	21.6	1,064.9
3	Cerebrovascular diseases (I60-I69)	108,515	6.8	336.5
4	Chronic lower respiratory diseases (J40-J47)	105,342	6.6	326.7
5	Alzheimer's disease (G30)	65,515	4.1	203.2
6	Influenza and pneumonia (J10-J18)	49,825	3.1	154.5
7	Diabetes mellitus (E10-E14)	45,267	2.8	140.4
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	33,281	2.1	103.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	30,182	1.9	93.6
10	Septicemia (A40-A41)	21,703	1.4	67.3
11	Parkinson's disease (G20-G21)	18,027	1.1	55.9
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	17,205	1.1	53.4
13	Pneumonitis due to solids and liquids (J69)	14,022	0.9	43.5
14	Atherosclerosis (I70)	10,299	0.6	31.9
15	Aortic aneurysm and dissection (I71)	10,106	0.6	31.3
...	All other causes (Residual)	244,429	15.4	758.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, male, all ages				
...	All causes	1,028,152	100.0	864.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	279,324	27.2	234.9
2	Malignant neoplasms (C00-C97)	250,478	24.4	210.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	64,600	6.3	54.3
4	Chronic lower respiratory diseases (J40-J47)	56,911	5.5	47.9
5	Cerebrovascular diseases (I60-I69)	47,194	4.6	39.7
6	Diabetes mellitus (E10-E14)	29,628	2.9	24.9
7	Influenza and pneumonia (J10-J18)	24,425	2.4	20.5
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	23,478	2.3	19.7
9	Alzheimer's disease (G30)	18,990	1.8	16.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17,137	1.7	14.4
11	Chronic liver disease and cirrhosis (K70,K73-K74)	15,679	1.5	13.2
12	Septicemia (A40-A41)	12,172	1.2	10.2
13	Parkinson's disease (G20-G21)	10,649	1.0	9.0
14	Pneumonitis due to solids and liquids (J69)	7,759	0.8	6.5
15	Aortic aneurysm and dissection (I71)	7,203	0.7	6.1
...	All other causes (Residual)	162,525	15.8	136.7
White, male, 1-4 years				
...	All causes	1,996	100.0	30.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	793	39.7	12.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	201	10.1	3.1
3	Malignant neoplasms (C00-C97)	168	8.4	2.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	119	6.0	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	55	2.8	0.9
6	Influenza and pneumonia (J10-J18)	36	1.8	0.6
7	Septicemia (A40-A41)	33	1.7	0.5
8	Cerebrovascular diseases (I60-I69)	28	1.4	0.4
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	26	1.3	0.4
10	Certain conditions originating in the perinatal period (P00-P96)	20	1.0	0.3
11	Meningitis (G00,G03)	13	0.7	*
12	Chronic lower respiratory diseases (J40-J47)	11	0.6	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.5	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	9	0.5	*
15	Acute bronchitis and bronchiolitis (J20-J21)	8	0.4	*
...	All other causes (Residual)	467	23.4	7.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 5-14 years				
...	All causes	2,749	100.0	17.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,101	40.1	6.8
2	Malignant neoplasms (C00-C97)	429	15.6	2.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	160	5.8	1.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	145	5.3	0.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	96	3.5	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	90	3.3	0.6
7	Influenza and pneumonia (J10-J18)	36	1.3	0.2
8	Cerebrovascular diseases (I60-I69)	35	1.3	0.2
9	Septicemia (A40-A41)	33	1.2	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	29	1.1	0.2
11	Chronic lower respiratory diseases (J40-J47)	26	0.9	0.2
12	Diabetes mellitus (E10-E14)	11	0.4	*
12	Certain conditions originating in the perinatal period (P00-P96)	11	0.4	*
14	Meningitis (G00,G03)	9	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	8	0.3	*
...	All other causes (Residual)	530	19.3	3.3
White, male, 15-24 years				
...	All causes	18,792	100.0	110.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,035	53.4	58.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,945	15.7	17.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,800	9.6	10.6
4	Malignant neoplasms (C00-C97)	826	4.4	4.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	509	2.7	3.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	252	1.3	1.5
7	Influenza and pneumonia (J10-J18)	90	0.5	0.5
8	Diabetes mellitus (E10-E14)	73	0.4	0.4
9	Cerebrovascular diseases (I60-I69)	65	0.3	0.4
10	Legal intervention (Y35,Y89.0)	58	0.3	0.3
11	Septicemia (A40-A41)	56	0.3	0.3
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	47	0.3	0.3
13	Chronic lower respiratory diseases (J40-J47)	46	0.2	0.3
14	Human immunodeficiency virus (HIV) disease (B20-B24)	32	0.2	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	32	0.2	0.2
...	All other causes (Residual)	1,926	10.2	11.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 25-44 years				
...	All causes	62,056	100.0	182.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18,972	30.6	55.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,113	13.1	23.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8,019	12.9	23.5
4	Malignant neoplasms (C00-C97)	6,450	10.4	18.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,721	4.4	8.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	1,945	3.1	5.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,691	2.7	5.0
8	Diabetes mellitus (E10-E14)	1,139	1.8	3.3
9	Cerebrovascular diseases (I60-I69)	951	1.5	2.8
10	Influenza and pneumonia (J10-J18)	533	0.9	1.6
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	413	0.7	1.2
12	Chronic lower respiratory diseases (J40-J47)	383	0.6	1.1
13	Septicemia (A40-A41)	373	0.6	1.1
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	320	0.5	0.9
15	Viral hepatitis (B15-B19)	319	0.5	0.9
...	All other causes (Residual)	9,714	15.7	28.5
White, male, 45-64 years				
...	All causes	223,154	100.0	743.2
1	Malignant neoplasms (C00-C97)	65,010	29.1	216.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	57,408	25.7	191.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	16,899	7.6	56.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	9,040	4.1	30.1
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7,986	3.6	26.6
6	Diabetes mellitus (E10-E14)	7,600	3.4	25.3
7	Chronic lower respiratory diseases (J40-J47)	7,462	3.3	24.9
8	Cerebrovascular diseases (I60-I69)	6,225	2.8	20.7
9	Influenza and pneumonia (J10-J18)	2,504	1.1	8.3
10	Septicemia (A40-A41)	2,344	1.1	7.8
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,313	1.0	7.7
12	Human immunodeficiency virus (HIV) disease (B20-B24)	2,196	1.0	7.3
13	Viral hepatitis (B15-B19)	2,169	1.0	7.2
14	Aortic aneurysm and dissection (I71)	1,308	0.6	4.4
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,291	0.6	4.3
...	All other causes (Residual)	31,399	14.1	104.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 65 years and over				
...	All causes	708,768	100.0	5,210.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	213,005	30.1	1,565.8
2	Malignant neoplasms (C00-C97)	177,554	25.1	1,305.2
3	Chronic lower respiratory diseases (J40-J47)	48,968	6.9	360.0
4	Cerebrovascular diseases (I60-I69)	39,844	5.6	292.9
5	Influenza and pneumonia (J10-J18)	21,128	3.0	155.3
6	Diabetes mellitus (E10-E14)	20,805	2.9	152.9
7	Alzheimer's disease (G30)	18,693	2.6	137.4
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	16,366	2.3	120.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14,405	2.0	105.9
10	Parkinson's disease (G20-G21)	10,352	1.5	76.1
11	Septicemia (A40-A41)	9,237	1.3	67.9
12	Pneumonitis due to solids and liquids (J69)	6,985	1.0	51.3
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	5,741	0.8	42.2
14	Aortic aneurysm and dissection (I71)	5,647	0.8	41.5
15	Chronic liver disease and cirrhosis (K70,K73-K74)	4,932	0.7	36.3
...	All other causes (Residual)	95,106	13.4	699.1
White, female, all ages				
...	All causes	1,069,945	100.0	882.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	285,472	26.7	235.5
2	Malignant neoplasms (C00-C97)	231,654	21.7	191.1
3	Cerebrovascular diseases (I60-I69)	74,674	7.0	61.6
4	Chronic lower respiratory diseases (J40-J47)	63,973	6.0	52.8
5	Alzheimer's disease (G30)	47,201	4.4	38.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	35,806	3.3	29.5
7	Influenza and pneumonia (J10-J18)	31,115	2.9	25.7
8	Diabetes mellitus (E10-E14)	30,127	2.8	24.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17,669	1.7	14.6
10	Septicemia (A40-A41)	15,022	1.4	12.4
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	12,236	1.1	10.1
12	Chronic liver disease and cirrhosis (K70,K73-K74)	8,324	0.8	6.9
13	Parkinson's disease (G20-G21)	7,847	0.7	6.5
14	Pneumonitis due to solids and liquids (J69)	7,579	0.7	6.3
15	Atherosclerosis (I70)	6,752	0.6	5.6
...	All other causes (Residual)	194,494	18.2	160.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 1-4 years				
...	All causes	1,412	100.0	22.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	469	33.2	7.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	184	13.0	3.0
3	Malignant neoplasms (C00-C97)	125	8.9	2.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	98	6.9	1.6
5	Influenza and pneumonia (J10-J18)	40	2.8	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	38	2.7	0.6
7	Septicemia (A40-A41)	26	1.8	0.4
8	Cerebrovascular diseases (I60-I69)	17	1.2	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	1.1	*
10	Certain conditions originating in the perinatal period (P00-P96)	15	1.1	*
11	Chronic lower respiratory diseases (J40-J47)	13	0.9	*
12	Anemias (D50-D64)	7	0.5	*
12	Meningitis (G00,G03)	7	0.5	*
14	Meningococcal infection (A39)	5	0.4	*
15	Acute bronchitis and bronchiolitis (J20-J21)	4	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.3	*
...	All other causes (Residual)	340	24.1	5.5
White, female, 5-14 years				
...	All causes	1,961	100.0	12.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	684	34.9	4.5
2	Malignant neoplasms (C00-C97)	358	18.3	2.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	140	7.1	0.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	72	3.7	0.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	69	3.5	0.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	49	2.5	0.3
7	Influenza and pneumonia (J10-J18)	41	2.1	0.3
8	Cerebrovascular diseases (I60-I69)	29	1.5	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	28	1.4	0.2
10	Septicemia (A40-A41)	24	1.2	0.2
11	Chronic lower respiratory diseases (J40-J47)	19	1.0	*
12	Diabetes mellitus (E10-E14)	14	0.7	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	0.6	*
13	Certain conditions originating in the perinatal period (P00-P96)	11	0.6	*
15	Meningitis (G00,G03)	9	0.5	*
...	All other causes (Residual)	403	20.6	2.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 15-24 years				
...	All causes	6,618	100.0	41.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,290	49.7	20.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	595	9.0	3.7
3	Malignant neoplasms (C00-C97)	539	8.1	3.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	372	5.6	2.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	227	3.4	1.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	134	2.0	0.8
7	Pregnancy, childbirth and the puerperium (O00-099)	110	1.7	0.7
8	Cerebrovascular diseases (I60-I69)	71	1.1	0.4
9	Diabetes mellitus (E10-E14)	51	0.8	0.3
10	Influenza and pneumonia (J10-J18)	44	0.7	0.3
11	Septicemia (A40-A41)	39	0.6	0.2
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	29	0.4	0.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	28	0.4	0.2
14	Chronic lower respiratory diseases (J40-J47)	25	0.4	0.2
15	Human immunodeficiency virus (HIV) disease (B20-B24)	18	0.3	*
...	All other causes (Residual)	1,046	15.8	6.6
White, female, 25-44 years				
...	All causes	31,631	100.0	96.6
1	Malignant neoplasms (C00-C97)	7,596	24.0	23.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,828	21.6	20.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,072	9.7	9.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,127	6.7	6.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	916	2.9	2.8
6	Cerebrovascular diseases (I60-I69)	855	2.7	2.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	824	2.6	2.5
8	Diabetes mellitus (E10-E14)	661	2.1	2.0
9	Human immunodeficiency virus (HIV) disease (B20-B24)	477	1.5	1.5
10	Chronic lower respiratory diseases (J40-J47)	416	1.3	1.3
11	Influenza and pneumonia (J10-J18)	412	1.3	1.3
12	Septicemia (A40-A41)	370	1.2	1.1
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	353	1.1	1.1
14	Pregnancy, childbirth and the puerperium (O00-099)	305	1.0	0.9
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	233	0.7	0.7
...	All other causes (Residual)	6,186	19.6	18.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 45-64 years				
...	All causes	139,288	100.0	451.0
1	Malignant neoplasms (C00-C97)	57,146	41.0	185.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	22,373	16.1	72.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,268	5.2	23.5
4	Chronic lower respiratory diseases (J40-J47)	7,118	5.1	23.0
5	Cerebrovascular diseases (I60-I69)	4,998	3.6	16.2
6	Diabetes mellitus (E10-E14)	4,937	3.5	16.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	3,440	2.5	11.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,498	1.8	8.1
9	Septicemia (A40-A41)	2,032	1.5	6.6
10	Influenza and pneumonia (J10-J18)	1,819	1.3	5.9
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,571	1.1	5.1
12	Viral hepatitis (B15-B19)	785	0.6	2.5
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	716	0.5	2.3
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	689	0.5	2.2
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	606	0.4	2.0
...	All other causes (Residual)	21,292	15.3	68.9
White, female, 65 years and over				
...	All causes	880,953	100.0	4,725.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	259,582	29.5	1,392.3
2	Malignant neoplasms (C00-C97)	165,862	18.8	889.6
3	Cerebrovascular diseases (I60-I69)	68,671	7.8	368.3
4	Chronic lower respiratory diseases (J40-J47)	56,374	6.4	302.4
5	Alzheimer's disease (G30)	46,822	5.3	251.1
6	Influenza and pneumonia (J10-J18)	28,697	3.3	153.9
7	Diabetes mellitus (E10-E14)	24,462	2.8	131.2
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	16,915	1.9	90.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15,777	1.8	84.6
10	Septicemia (A40-A41)	12,466	1.4	66.9
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	11,464	1.3	61.5
12	Parkinson's disease (G20-G21)	7,675	0.9	41.2
13	Pneumonitis due to solids and liquids (J69)	7,037	0.8	37.7
14	Atherosclerosis (I70)	6,557	0.7	35.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5,077	0.6	27.2
...	All other causes (Residual)	147,515	16.7	791.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, all ages				
...	All causes	292,808	100.0	749.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	74,159	25.3	189.8
2	Malignant neoplasms (C00-C97)	63,165	21.6	161.7
3	Cerebrovascular diseases (I60-I69)	17,541	6.0	44.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,652	4.7	34.9
5	Diabetes mellitus (E10-E14)	12,970	4.4	33.2
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8,669	3.0	22.2
7	Chronic lower respiratory diseases (J40-J47)	8,229	2.8	21.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,075	2.8	20.7
9	Human immunodeficiency virus (HIV) disease (B20-B24)	7,022	2.4	18.0
10	Septicemia (A40-A41)	6,221	2.1	15.9
11	Influenza and pneumonia (J10-J18)	5,780	2.0	14.8
12	Certain conditions originating in the perinatal period (P00-P96)	5,050	1.7	12.9
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	4,953	1.7	12.7
14	Alzheimer's disease (G30)	4,620	1.6	11.8
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,530	0.9	6.5
...	All other causes (Residual)	50,172	17.1	128.4
Black, both sexes, 1-4 years				
...	All causes	1,097	100.0	41.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	327	29.8	12.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	140	12.8	5.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	106	9.7	4.0
4	Malignant neoplasms (C00-C97)	64	5.8	2.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	45	4.1	1.7
6	Chronic lower respiratory diseases (J40-J47)	31	2.8	1.2
7	Influenza and pneumonia (J10-J18)	29	2.6	1.1
8	Certain conditions originating in the perinatal period (P00-P96)	23	2.1	0.9
9	Septicemia (A40-A41)	20	1.8	0.8
10	Cerebrovascular diseases (I60-I69)	14	1.3	*
11	Anemias (D50-D64)	11	1.0	*
12	Meningitis (G00,G03)	6	0.5	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.5	*
13	Pneumonitis due to solids and liquids (J69)	5	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.3	*
...	All other causes (Residual)	268	24.4	10.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 5-14 years				
...	All causes	1,543	100.0	23.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	501	32.5	7.6
2	Malignant neoplasms (C00-C97)	172	11.1	2.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	150	9.7	2.3
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	88	5.7	1.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	77	5.0	1.2
6	Chronic lower respiratory diseases (J40-J47)	53	3.4	0.8
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	41	2.7	0.6
8	Cerebrovascular diseases (I60-I69)	24	1.6	0.4
9	Influenza and pneumonia (J10-J18)	22	1.4	0.3
10	Septicemia (A40-A41)	21	1.4	0.3
11	Anemias (D50-D64)	15	1.0	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	0.9	*
13	Diabetes mellitus (E10-E14)	12	0.8	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	9	0.6	*
15	Meningitis (G00,G03)	7	0.5	*
...	All other causes (Residual)	337	21.8	5.1
Black, both sexes, 15-24 years				
...	All causes	7,371	100.0	112.3
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,077	41.7	46.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,780	24.1	27.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	437	5.9	6.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	334	4.5	5.1
5	Malignant neoplasms (C00-C97)	264	3.6	4.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	109	1.5	1.7
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	100	1.4	1.5
8	Anemias (D50-D64)	77	1.0	1.2
9	Chronic lower respiratory diseases (J40-J47)	75	1.0	1.1
10	Diabetes mellitus (E10-E14)	73	1.0	1.1
11	Pregnancy, childbirth and the puerperium (O00-099)	69	0.9	1.1
12	Cerebrovascular diseases (I60-I69)	51	0.7	0.8
13	Septicemia (A40-A41)	44	0.6	0.7
14	Influenza and pneumonia (J10-J18)	31	0.4	0.5
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	29	0.4	0.4
...	All other causes (Residual)	821	11.1	12.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 25-44 years				
...	All causes	28,222	100.0	249.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,211	14.9	37.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,004	14.2	35.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,960	14.0	35.0
4	Malignant neoplasms (C00-C97)	3,248	11.5	28.7
5	Human immunodeficiency virus (HIV) disease (B20-B24)	3,173	11.2	28.1
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	904	3.2	8.0
7	Cerebrovascular diseases (I60-I69)	852	3.0	7.5
8	Diabetes mellitus (E10-E14)	760	2.7	6.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	434	1.5	3.8
10	Septicemia (A40-A41)	370	1.3	3.3
11	Chronic lower respiratory diseases (J40-J47)	316	1.1	2.8
12	Influenza and pneumonia (J10-J18)	295	1.0	2.6
13	Anemias (D50-D64)	278	1.0	2.5
14	Chronic liver disease and cirrhosis (K70,K73-K74)	272	1.0	2.4
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	249	0.9	2.2
...	All other causes (Residual)	4,896	17.3	43.3
Black, both sexes, 45-64 years				
...	All causes	83,165	100.0	1,022.4
1	Malignant neoplasms (C00-C97)	22,861	27.5	281.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,831	25.0	256.1
3	Cerebrovascular diseases (I60-I69)	4,434	5.3	54.5
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,160	5.0	51.1
5	Diabetes mellitus (E10-E14)	3,850	4.6	47.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	3,431	4.1	42.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,100	2.5	25.8
8	Chronic lower respiratory diseases (J40-J47)	1,910	2.3	23.5
9	Chronic liver disease and cirrhosis (K70,K73-K74)	1,655	2.0	20.3
10	Septicemia (A40-A41)	1,600	1.9	19.7
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,223	1.5	15.0
12	Influenza and pneumonia (J10-J18)	1,110	1.3	13.6
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,045	1.3	12.8
14	Viral hepatitis (B15-B19)	612	0.7	7.5
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	458	0.6	5.6
...	All other causes (Residual)	11,885	14.3	146.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 65 years and over				
...	All causes	162,668	100.0	5,162.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	48,545	29.8	1,540.6
2	Malignant neoplasms (C00-C97)	36,542	22.5	1,159.6
3	Cerebrovascular diseases (I60-I69)	12,122	7.5	384.7
4	Diabetes mellitus (E10-E14)	8,274	5.1	262.6
5	Chronic lower respiratory diseases (J40-J47)	5,831	3.6	185.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,447	3.3	172.9
7	Alzheimer's disease (G30)	4,566	2.8	144.9
8	Influenza and pneumonia (J10-J18)	4,195	2.6	133.1
9	Septicemia (A40-A41)	4,033	2.5	128.0
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	3,468	2.1	110.1
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,561	1.6	81.3
12	Pneumonitis due to solids and liquids (J69)	1,338	0.8	42.5
13	Aortic aneurysm and dissection (I71)	755	0.5	24.0
14	Atherosclerosis (I70)	735	0.5	23.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	712	0.4	22.6
...	All other causes (Residual)	23,544	14.5	747.2
Black, male, all ages				
...	All causes	149,108	100.0	799.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	36,343	24.4	194.8
2	Malignant neoplasms (C00-C97)	32,726	21.9	175.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,329	6.3	50.0
4	Cerebrovascular diseases (I60-I69)	7,519	5.0	40.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7,412	5.0	39.7
6	Diabetes mellitus (E10-E14)	5,730	3.8	30.7
7	Human immunodeficiency virus (HIV) disease (B20-B24)	4,684	3.1	25.1
8	Chronic lower respiratory diseases (J40-J47)	4,464	3.0	23.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,645	2.4	19.5
10	Certain conditions originating in the perinatal period (P00-P96)	2,828	1.9	15.2
11	Septicemia (A40-A41)	2,786	1.9	14.9
12	Influenza and pneumonia (J10-J18)	2,729	1.8	14.6
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,128	1.4	11.4
14	Chronic liver disease and cirrhosis (K70,K73-K74)	1,656	1.1	8.9
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,621	1.1	8.7
...	All other causes (Residual)	23,508	15.8	126.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 1-4 years				
...	All causes	623	100.0	46.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	184	29.5	13.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	88	14.1	6.6
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	48	7.7	3.6
4	Malignant neoplasms (C00-C97)	37	5.9	2.8
5	Chronic lower respiratory diseases (J40-J47)	22	3.5	1.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	19	3.0	*
7	Influenza and pneumonia (J10-J18)	17	2.7	*
8	Certain conditions originating in the perinatal period (P00-P96)	16	2.6	*
9	Septicemia (A40-A41)	13	2.1	*
10	Cerebrovascular diseases (I60-I69)	10	1.6	*
11	Anemias (D50-D64)	5	0.8	*
12	Meningitis (G00,G03)	3	0.5	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.5	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.3	*
14	Pneumonitis due to solids and liquids (J69)	2	0.3	*
...	All other causes (Residual)	154	24.7	11.5
Black, male, 5-14 years				
...	All causes	909	100.0	27.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	314	34.5	9.3
2	Malignant neoplasms (C00-C97)	103	11.3	3.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	97	10.7	2.9
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	41	4.5	1.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	36	4.0	1.1
6	Chronic lower respiratory diseases (J40-J47)	34	3.7	1.0
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	27	3.0	0.8
8	Cerebrovascular diseases (I60-I69)	15	1.7	*
9	Septicemia (A40-A41)	14	1.5	*
10	Influenza and pneumonia (J10-J18)	13	1.4	*
11	Anemias (D50-D64)	9	1.0	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	6	0.7	*
12	Meningitis (G00,G03)	6	0.7	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.4	*
15	Diabetes mellitus (E10-E14)	3	0.3	*
15	Diseases of appendix (K35-K38)	3	0.3	*
...	All other causes (Residual)	184	20.2	5.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 15-24 years				
...	All causes	5,709	100.0	172.1
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,790	48.9	84.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,334	23.4	40.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	382	6.7	11.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	229	4.0	6.9
5	Malignant neoplasms (C00-C97)	155	2.7	4.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	64	1.1	1.9
7	Human immunodeficiency virus (HIV) disease (B20-B24)	57	1.0	1.7
8	Chronic lower respiratory diseases (J40-J47)	48	0.8	1.4
9	Anemias (D50-D64)	45	0.8	1.4
10	Diabetes mellitus (E10-E14)	39	0.7	1.2
11	Septicemia (A40-A41)	24	0.4	0.7
11	Cerebrovascular diseases (I60-I69)	24	0.4	0.7
11	Legal intervention (Y35,Y89.0)	24	0.4	0.7
14	Influenza and pneumonia (J10-J18)	23	0.4	0.7
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	0.2	*
...	All other causes (Residual)	460	8.1	13.9
Black, male, 25-44 years				
...	All causes	17,469	100.0	324.9
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,410	19.5	63.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,926	16.7	54.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,622	15.0	48.8
4	Human immunodeficiency virus (HIV) disease (B20-B24)	1,947	11.1	36.2
5	Malignant neoplasms (C00-C97)	1,294	7.4	24.1
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	739	4.2	13.7
7	Diabetes mellitus (E10-E14)	438	2.5	8.1
8	Cerebrovascular diseases (I60-I69)	421	2.4	7.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	226	1.3	4.2
10	Influenza and pneumonia (J10-J18)	174	1.0	3.2
11	Septicemia (A40-A41)	172	1.0	3.2
12	Chronic liver disease and cirrhosis (K70,K73-K74)	158	0.9	2.9
13	Chronic lower respiratory diseases (J40-J47)	156	0.9	2.9
14	Anemias (D50-D64)	148	0.8	2.8
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	132	0.8	2.5
...	All other causes (Residual)	2,506	14.3	46.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 45-64 years				
...	All causes	48,959	100.0	1,318.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,074	26.7	352.2
2	Malignant neoplasms (C00-C97)	12,132	24.8	326.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,030	6.2	81.6
4	Cerebrovascular diseases (I60-I69)	2,467	5.0	66.5
5	Human immunodeficiency virus (HIV) disease (B20-B24)	2,456	5.0	66.2
6	Diabetes mellitus (E10-E14)	2,067	4.2	55.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,193	2.4	32.1
8	Chronic liver disease and cirrhosis (K70,K73-K74)	1,139	2.3	30.7
9	Chronic lower respiratory diseases (J40-J47)	1,023	2.1	27.6
10	Septicemia (A40-A41)	834	1.7	22.5
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	829	1.7	22.3
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	710	1.5	19.1
13	Influenza and pneumonia (J10-J18)	679	1.4	18.3
14	Viral hepatitis (B15-B19)	415	0.8	11.2
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	349	0.7	9.4
...	All other causes (Residual)	6,562	13.4	176.8
Black, male, 65 years and over				
...	All causes	70,541	100.0	5,826.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,296	28.8	1,676.5
2	Malignant neoplasms (C00-C97)	18,997	26.9	1,569.2
3	Cerebrovascular diseases (I60-I69)	4,555	6.5	376.3
4	Diabetes mellitus (E10-E14)	3,183	4.5	262.9
5	Chronic lower respiratory diseases (J40-J47)	3,174	4.5	262.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,174	3.1	179.6
7	Influenza and pneumonia (J10-J18)	1,776	2.5	146.7
8	Septicemia (A40-A41)	1,658	2.4	137.0
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,366	1.9	112.8
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,280	1.8	105.7
11	Alzheimer's disease (G30)	1,270	1.8	104.9
12	Pneumonitis due to solids and liquids (J69)	653	0.9	53.9
13	Parkinson's disease (G20-G21)	393	0.6	32.5
14	Chronic liver disease and cirrhosis (K70,K73-K74)	353	0.5	29.2
15	Aortic aneurysm and dissection (I71)	331	0.5	27.3
...	All other causes (Residual)	9,082	12.9	750.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, all ages				
...	All causes	143,700	100.0	703.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	37,816	26.3	185.2
2	Malignant neoplasms (C00-C97)	30,439	21.2	149.1
3	Cerebrovascular diseases (I60-I69)	10,022	7.0	49.1
4	Diabetes mellitus (E10-E14)	7,240	5.0	35.5
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,430	3.1	21.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,323	3.0	21.2
7	Chronic lower respiratory diseases (J40-J47)	3,765	2.6	18.4
8	Septicemia (A40-A41)	3,435	2.4	16.8
9	Alzheimer's disease (G30)	3,328	2.3	16.3
10	Influenza and pneumonia (J10-J18)	3,051	2.1	14.9
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,825	2.0	13.8
12	Human immunodeficiency virus (HIV) disease (B20-B24)	2,338	1.6	11.5
13	Certain conditions originating in the perinatal period (P00-P96)	2,222	1.5	10.9
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,257	0.9	6.2
15	Chronic liver disease and cirrhosis (K70,K73-K74)	874	0.6	4.3
...	All other causes (Residual)	26,335	18.3	129.0
Black, female, 1-4 years				
...	All causes	474	100.0	36.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	143	30.2	11.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	58	12.2	4.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	52	11.0	4.0
4	Malignant neoplasms (C00-C97)	27	5.7	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	26	5.5	2.0
6	Influenza and pneumonia (J10-J18)	12	2.5	*
7	Chronic lower respiratory diseases (J40-J47)	9	1.9	*
8	Septicemia (A40-A41)	7	1.5	*
8	Certain conditions originating in the perinatal period (P00-P96)	7	1.5	*
10	Anemias (D50-D64)	6	1.3	*
11	Cerebrovascular diseases (I60-I69)	4	0.8	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.6	*
12	Meningitis (G00,G03)	3	0.6	*
12	Pneumonitis due to solids and liquids (J69)	3	0.6	*
15	Acute bronchitis and bronchiolitis (J20-J21)	2	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
...	All other causes (Residual)	110	23.2	8.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 5-14 years				
...	All causes	634	100.0	19.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	187	29.5	5.7
2	Malignant neoplasms (C00-C97)	69	10.9	2.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	53	8.4	1.6
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	47	7.4	1.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	41	6.5	1.3
6	Chronic lower respiratory diseases (J40-J47)	19	3.0	*
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	14	2.2	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	1.6	*
9	Diabetes mellitus (E10-E14)	9	1.4	*
9	Cerebrovascular diseases (I60-I69)	9	1.4	*
9	Influenza and pneumonia (J10-J18)	9	1.4	*
12	Septicemia (A40-A41)	7	1.1	*
13	Anemias (D50-D64)	6	0.9	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	3	0.5	*
14	Certain conditions originating in the perinatal period (P00-P96)	3	0.5	*
...	All other causes (Residual)	148	23.3	4.5
Black, female, 15-24 years				
...	All causes	1,662	100.0	51.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	446	26.8	13.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	287	17.3	8.8
3	Malignant neoplasms (C00-C97)	109	6.6	3.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	105	6.3	3.2
5	Pregnancy, childbirth and the puerperium (O00-099)	69	4.2	2.1
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	55	3.3	1.7
7	Human immunodeficiency virus (HIV) disease (B20-B24)	52	3.1	1.6
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	36	2.2	1.1
9	Diabetes mellitus (E10-E14)	34	2.0	1.0
10	Anemias (D50-D64)	32	1.9	1.0
11	Cerebrovascular diseases (I60-I69)	27	1.6	0.8
11	Chronic lower respiratory diseases (J40-J47)	27	1.6	0.8
13	Septicemia (A40-A41)	20	1.2	0.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18	1.1	*
15	Influenza and pneumonia (J10-J18)	8	0.5	*
...	All other causes (Residual)	337	20.3	10.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 25-44 years				
...	All causes	10,753	100.0	181.2
1	Malignant neoplasms (C00-C97)	1,954	18.2	32.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,589	14.8	26.8
3	Human immunodeficiency virus (HIV) disease (B20-B24)	1,226	11.4	20.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,078	10.0	18.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	550	5.1	9.3
6	Cerebrovascular diseases (I60-I69)	431	4.0	7.3
7	Diabetes mellitus (E10-E14)	322	3.0	5.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	208	1.9	3.5
9	Septicemia (A40-A41)	198	1.8	3.3
10	Pregnancy, childbirth and the puerperium (O00-099)	182	1.7	3.1
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	165	1.5	2.8
12	Chronic lower respiratory diseases (J40-J47)	160	1.5	2.7
13	Anemias (D50-D64)	130	1.2	2.2
14	Influenza and pneumonia (J10-J18)	121	1.1	2.0
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	117	1.1	2.0
...	All other causes (Residual)	2,322	21.6	39.1
Black, female, 45-64 years				
...	All causes	34,206	100.0	773.5
1	Malignant neoplasms (C00-C97)	10,729	31.4	242.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,757	22.7	175.4
3	Cerebrovascular diseases (I60-I69)	1,967	5.8	44.5
4	Diabetes mellitus (E10-E14)	1,783	5.2	40.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,130	3.3	25.6
6	Human immunodeficiency virus (HIV) disease (B20-B24)	975	2.9	22.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	907	2.7	20.5
8	Chronic lower respiratory diseases (J40-J47)	887	2.6	20.1
9	Septicemia (A40-A41)	766	2.2	17.3
10	Chronic liver disease and cirrhosis (K70,K73-K74)	516	1.5	11.7
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	513	1.5	11.6
12	Influenza and pneumonia (J10-J18)	431	1.3	9.7
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	216	0.6	4.9
14	Viral hepatitis (B15-B19)	197	0.6	4.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	142	0.4	3.2
...	All other causes (Residual)	5,290	15.5	119.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 65 years and over				
...	All causes	92,127	100.0	4,747.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	28,249	30.7	1,455.8
2	Malignant neoplasms (C00-C97)	17,545	19.0	904.1
3	Cerebrovascular diseases (I60-I69)	7,567	8.2	390.0
4	Diabetes mellitus (E10-E14)	5,091	5.5	262.4
5	Alzheimer's disease (G30)	3,296	3.6	169.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,273	3.6	168.7
7	Chronic lower respiratory diseases (J40-J47)	2,657	2.9	136.9
8	Influenza and pneumonia (J10-J18)	2,419	2.6	124.7
9	Septicemia (A40-A41)	2,375	2.6	122.4
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,188	2.4	112.8
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,195	1.3	61.6
12	Pneumonitis due to solids and liquids (J69)	685	0.7	35.3
13	Atherosclerosis (I70)	488	0.5	25.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	426	0.5	22.0
15	Aortic aneurysm and dissection (I71)	424	0.5	21.8
...	All other causes (Residual)	14,249	15.5	734.3
American Indian, both sexes, all ages				
...	All causes	13,918	100.0	440.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,659	19.1	84.1
2	Malignant neoplasms (C00-C97)	2,465	17.7	78.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,626	11.7	51.4
4	Diabetes mellitus (E10-E14)	818	5.9	25.9
5	Cerebrovascular diseases (I60-I69)	627	4.5	19.8
6	Chronic liver disease and cirrhosis (K70,K73-K74)	596	4.3	18.9
7	Chronic lower respiratory diseases (J40-J47)	518	3.7	16.4
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	392	2.8	12.4
9	Influenza and pneumonia (J10-J18)	354	2.5	11.2
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	265	1.9	8.4
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	257	1.8	8.1
12	Septicemia (A40-A41)	217	1.6	6.9
13	Alzheimer's disease (G30)	173	1.2	5.5
14	Certain conditions originating in the perinatal period (P00-P96)	130	0.9	4.1
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	117	0.8	3.7
...	All other causes (Residual)	2,704	19.4	85.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 1-4 years				
...	All causes	100	100.0	59.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	44	44.0	26.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	11	11.0	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	10	10.0	*
4	Malignant neoplasms (C00-C97)	4	4.0	*
5	Meningitis (G00,G03)	3	3.0	*
6	Septicemia (A40-A41)	1	1.0	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	1.0	*
6	Influenza and pneumonia (J10-J18)	1	1.0	*
6	Chronic lower respiratory diseases (J40-J47)	1	1.0	*
...	All other causes (Residual)	24	24.0	14.2
American Indian, both sexes, 5-14 years				
...	All causes	110	100.0	19.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	55	50.0	9.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	14	12.7	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7	6.4	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	5.5	*
5	Malignant neoplasms (C00-C97)	4	3.6	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	1.8	*
6	Chronic lower respiratory diseases (J40-J47)	2	1.8	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.9	*
8	Meningitis (G00,G03)	1	0.9	*
8	Influenza and pneumonia (J10-J18)	1	0.9	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.9	*
...	All other causes (Residual)	16	14.5	*
American Indian, both sexes, 15-24 years				
...	All causes	635	100.0	107.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	321	50.6	54.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	128	20.2	21.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	82	12.9	13.9
4	Malignant neoplasms (C00-C97)	15	2.4	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	13	2.0	*
6	Diabetes mellitus (E10-E14)	4	0.6	*
7	Septicemia (A40-A41)	3	0.5	*
7	Chronic liver disease and cirrhosis (K70,K73-K74)	3	0.5	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.3	*
9	Cerebrovascular diseases (I60-I69)	2	0.3	*
9	Pregnancy, childbirth and the puerperium (O00-099)	2	0.3	*
12	Salmonella infections (A01-A02)	1	0.2	*
12	Anemias (D50-D64)	1	0.2	*
12	Influenza and pneumonia (J10-J18)	1	0.2	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.2	*
...	All other causes (Residual)	55	8.7	9.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 25-44 years				
...	All causes	1,943	100.0	207.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	629	32.4	67.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	175	9.0	18.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	166	8.5	17.7
4	Chronic liver disease and cirrhosis (K70,K73-K74)	161	8.3	17.2
5	Malignant neoplasms (C00-C97)	145	7.5	15.5
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	105	5.4	11.2
7	Diabetes mellitus (E10-E14)	58	3.0	6.2
8	Human immunodeficiency virus (HIV) disease (B20-B24)	45	2.3	4.8
9	Cerebrovascular diseases (I60-I69)	36	1.9	3.8
10	Influenza and pneumonia (J10-J18)	23	1.2	2.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21	1.1	2.2
12	Septicemia (A40-A41)	20	1.0	2.1
13	Chronic lower respiratory diseases (J40-J47)	15	0.8	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	11	0.6	*
15	Viral hepatitis (B15-B19)	10	0.5	*
...	All other causes (Residual)	323	16.6	34.4
American Indian, both sexes, 45-64 years				
...	All causes	4,167	100.0	636.9
1	Malignant neoplasms (C00-C97)	920	22.1	140.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	815	19.6	124.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	393	9.4	60.1
4	Chronic liver disease and cirrhosis (K70,K73-K74)	339	8.1	51.8
5	Diabetes mellitus (E10-E14)	275	6.6	42.0
6	Cerebrovascular diseases (I60-I69)	153	3.7	23.4
7	Chronic lower respiratory diseases (J40-J47)	124	3.0	19.0
8	Influenza and pneumonia (J10-J18)	70	1.7	10.7
9	Septicemia (A40-A41)	69	1.7	10.5
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	67	1.6	10.2
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	65	1.6	9.9
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	40	1.0	6.1
13	Viral hepatitis (B15-B19)	37	0.9	5.7
14	Human immunodeficiency virus (HIV) disease (B20-B24)	32	0.8	4.9
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	17	0.4	*
...	All other causes (Residual)	751	18.0	114.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 65 years and over				
...	All causes	6,604	100.0	3,163.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,657	25.1	793.8
2	Malignant neoplasms (C00-C97)	1,377	20.9	659.6
3	Diabetes mellitus (E10-E14)	481	7.3	230.4
4	Cerebrovascular diseases (I60-I69)	434	6.6	207.9
5	Chronic lower respiratory diseases (J40-J47)	375	5.7	179.6
6	Influenza and pneumonia (J10-J18)	253	3.8	121.2
7	Alzheimer's disease (G30)	173	2.6	82.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	173	2.6	82.9
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	165	2.5	79.0
10	Septicemia (A40-A41)	122	1.8	58.4
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	93	1.4	44.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	93	1.4	44.6
13	Parkinson's disease (G20-G21)	59	0.9	28.3
14	Pneumonitis due to solids and liquids (J69)	53	0.8	25.4
15	Atherosclerosis (I70)	38	0.6	18.2
...	All other causes (Residual)	1,058	16.0	506.8
American Indian, male, all ages				
...	All causes	7,607	100.0	481.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,471	19.3	93.2
2	Malignant neoplasms (C00-C97)	1,297	17.1	82.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,118	14.7	70.8
4	Diabetes mellitus (E10-E14)	412	5.4	26.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	348	4.6	22.0
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	313	4.1	19.8
7	Chronic lower respiratory diseases (J40-J47)	257	3.4	16.3
8	Cerebrovascular diseases (I60-I69)	249	3.3	15.8
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	193	2.5	12.2
10	Influenza and pneumonia (J10-J18)	180	2.4	11.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	105	1.4	6.7
12	Septicemia (A40-A41)	104	1.4	6.6
13	Certain conditions originating in the perinatal period (P00-P96)	75	1.0	4.8
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	61	0.8	3.9
15	Human immunodeficiency virus (HIV) disease (B20-B24)	59	0.8	3.7
...	All other causes (Residual)	1,365	17.9	86.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 1-4 years				
...	All causes	62	100.0	72.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	31	50.0	36.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	6.5	*
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	6.5	*
4	Malignant neoplasms (C00-C97)	3	4.8	*
5	Meningitis (G00,G03)	2	3.2	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	1.6	*
...	All other causes (Residual)	17	27.4	*
American Indian, male, 5-14 years				
...	All causes	64	100.0	22.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	36	56.3	12.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11	17.2	*
3	Malignant neoplasms (C00-C97)	4	6.3	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	4.7	*
5	Human immunodeficiency virus (HIV) disease (B20-B24)	1	1.6	*
5	Influenza and pneumonia (J10-J18)	1	1.6	*
5	Chronic lower respiratory diseases (J40-J47)	1	1.6	*
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	1.6	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	1.6	*
...	All other causes (Residual)	5	7.8	*
American Indian, male, 15-24 years				
...	All causes	439	100.0	145.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	223	50.8	73.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	99	22.6	32.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	67	15.3	22.1
4	Malignant neoplasms (C00-C97)	10	2.3	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	7	1.6	*
6	Diabetes mellitus (E10-E14)	2	0.5	*
6	Chronic liver disease and cirrhosis (K70,K73-K74)	2	0.5	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.2	*
8	Cerebrovascular diseases (I60-I69)	1	0.2	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
...	All other causes (Residual)	26	5.9	8.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 25-44 years				
...	All causes	1,289	100.0	269.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	453	35.1	94.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	141	10.9	29.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	117	9.1	24.4
4	Chronic liver disease and cirrhosis (K70,K73-K74)	92	7.1	19.2
5	Malignant neoplasms (C00-C97)	78	6.1	16.3
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	77	6.0	16.1
7	Diabetes mellitus (E10-E14)	38	2.9	7.9
8	Human immunodeficiency virus (HIV) disease (B20-B24)	29	2.2	6.1
9	Cerebrovascular diseases (I60-I69)	18	1.4	*
10	Influenza and pneumonia (J10-J18)	16	1.2	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	1.1	*
12	Septicemia (A40-A41)	11	0.9	*
13	Viral hepatitis (B15-B19)	7	0.5	*
14	Legal intervention (Y35,Y89.0)	6	0.5	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	0.4	*
...	All other causes (Residual)	187	14.5	39.0
American Indian, male, 45-64 years				
...	All causes	2,501	100.0	790.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	549	22.0	173.6
2	Malignant neoplasms (C00-C97)	482	19.3	152.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	280	11.2	88.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	211	8.4	66.7
5	Diabetes mellitus (E10-E14)	157	6.3	49.6
6	Cerebrovascular diseases (I60-I69)	70	2.8	22.1
7	Chronic lower respiratory diseases (J40-J47)	63	2.5	19.9
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	53	2.1	16.8
9	Influenza and pneumonia (J10-J18)	40	1.6	12.6
10	Septicemia (A40-A41)	34	1.4	10.8
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	34	1.4	10.8
12	Viral hepatitis (B15-B19)	28	1.1	8.9
12	Human immunodeficiency virus (HIV) disease (B20-B24)	28	1.1	8.9
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	0.9	7.3
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	13	0.5	*
...	All other causes (Residual)	436	17.4	137.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 65 years and over				
...	All causes	3,055	100.0	3,357.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	796	26.1	874.8
2	Malignant neoplasms (C00-C97)	720	23.6	791.3
3	Diabetes mellitus (E10-E14)	215	7.0	236.3
4	Chronic lower respiratory diseases (J40-J47)	189	6.2	207.7
5	Cerebrovascular diseases (I60-I69)	159	5.2	174.7
6	Influenza and pneumonia (J10-J18)	121	4.0	133.0
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	85	2.8	93.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	65	2.1	71.4
9	Septicemia (A40-A41)	59	1.9	64.8
10	Alzheimer's disease (G30)	48	1.6	52.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	43	1.4	47.3
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	42	1.4	46.2
13	Parkinson's disease (G20-G21)	30	1.0	33.0
14	Pneumonitis due to solids and liquids (J69)	29	0.9	31.9
15	Aortic aneurysm and dissection (I71)	17	0.6	*
...	All other causes (Residual)	437	14.3	480.3
American Indian, female, all ages				
...	All causes	6,311	100.0	398.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,188	18.8	75.1
2	Malignant neoplasms (C00-C97)	1,168	18.5	73.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	508	8.0	32.1
4	Diabetes mellitus (E10-E14)	406	6.4	25.7
5	Cerebrovascular diseases (I60-I69)	378	6.0	23.9
6	Chronic lower respiratory diseases (J40-J47)	261	4.1	16.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	248	3.9	15.7
8	Influenza and pneumonia (J10-J18)	174	2.8	11.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	160	2.5	10.1
10	Alzheimer's disease (G30)	125	2.0	7.9
11	Septicemia (A40-A41)	113	1.8	7.1
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	79	1.3	5.0
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	64	1.0	4.0
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	57	0.9	3.6
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	56	0.9	3.5
...	All other causes (Residual)	1,326	21.0	83.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 1-4 years				
...	All causes	38	100.0	45.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13	34.2	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	18.4	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	15.8	*
4	Septicemia (A40-A41)	1	2.6	*
4	Malignant neoplasms (C00-C97)	1	2.6	*
4	Meningitis (G00,G03)	1	2.6	*
4	Influenza and pneumonia (J10-J18)	1	2.6	*
4	Chronic lower respiratory diseases (J40-J47)	1	2.6	*
...	All other causes (Residual)	7	18.4	*
American Indian, female, 5-14 years				
...	All causes	46	100.0	16.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19	41.3	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	10.9	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	8.7	*
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3	6.5	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	4.3	*
6	Meningitis (G00,G03)	1	2.2	*
6	Chronic lower respiratory diseases (J40-J47)	1	2.2	*
...	All other causes (Residual)	11	23.9	*
American Indian, female, 15-24 years				
...	All causes	196	100.0	67.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	98	50.0	34.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	29	14.8	10.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	15	7.7	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	6	3.1	*
5	Malignant neoplasms (C00-C97)	5	2.6	*
6	Septicemia (A40-A41)	3	1.5	*
7	Diabetes mellitus (E10-E14)	2	1.0	*
7	Pregnancy, childbirth and the puerperium (O00-099)	2	1.0	*
9	Salmonella infections (A01-A02)	1	0.5	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.5	*
9	Anemias (D50-D64)	1	0.5	*
9	Cerebrovascular diseases (I60-I69)	1	0.5	*
9	Influenza and pneumonia (J10-J18)	1	0.5	*
9	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.5	*
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.5	*
...	All other causes (Residual)	29	14.8	10.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 25-44 years				
...	All causes	654	100.0	142.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	176	26.9	38.3
2	Chronic liver disease and cirrhosis (K70,K73-K74)	69	10.6	15.0
3	Malignant neoplasms (C00-C97)	67	10.2	14.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	49	7.5	10.7
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	34	5.2	7.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	28	4.3	6.1
7	Diabetes mellitus (E10-E14)	20	3.1	4.4
8	Cerebrovascular diseases (I60-I69)	18	2.8	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	16	2.4	*
10	Chronic lower respiratory diseases (J40-J47)	11	1.7	*
11	Septicemia (A40-A41)	9	1.4	*
12	Influenza and pneumonia (J10-J18)	7	1.1	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	1.1	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	0.9	*
15	Pregnancy, childbirth and the puerperium (O00-099)	5	0.8	*
...	All other causes (Residual)	132	20.2	28.7
American Indian, female, 45-64 years				
...	All causes	1,666	100.0	492.9
1	Malignant neoplasms (C00-C97)	438	26.3	129.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	266	16.0	78.7
3	Chronic liver disease and cirrhosis (K70,K73-K74)	128	7.7	37.9
4	Diabetes mellitus (E10-E14)	118	7.1	34.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	113	6.8	33.4
6	Cerebrovascular diseases (I60-I69)	83	5.0	24.6
7	Chronic lower respiratory diseases (J40-J47)	61	3.7	18.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	44	2.6	13.0
9	Septicemia (A40-A41)	35	2.1	10.4
10	Influenza and pneumonia (J10-J18)	30	1.8	8.9
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	12	0.7	*
12	Viral hepatitis (B15-B19)	9	0.5	*
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	0.4	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.3	*
14	Atherosclerosis (I70)	5	0.3	*
...	All other causes (Residual)	313	18.8	92.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 65 years and over				
...	All causes	3,549	100.0	3,013.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	861	24.3	731.2
2	Malignant neoplasms (C00-C97)	657	18.5	557.9
3	Cerebrovascular diseases (I60-I69)	275	7.7	233.5
4	Diabetes mellitus (E10-E14)	266	7.5	225.9
5	Chronic lower respiratory diseases (J40-J47)	186	5.2	157.9
6	Influenza and pneumonia (J10-J18)	132	3.7	112.1
7	Alzheimer's disease (G30)	125	3.5	106.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	108	3.0	91.7
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	80	2.3	67.9
10	Septicemia (A40-A41)	63	1.8	53.5
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	51	1.4	43.3
12	Chronic liver disease and cirrhosis (K70,K73-K74)	50	1.4	42.5
13	Parkinson's disease (G20-G21)	29	0.8	24.6
14	Pneumonitis due to solids and liquids (J69)	24	0.7	20.4
15	Atherosclerosis (I70)	22	0.6	18.7
...	All other causes (Residual)	620	17.5	526.5
Asian or Pacific Islander, both sexes, all ages				
...	All causes	43,194	100.0	307.7
1	Malignant neoplasms (C00-C97)	11,550	26.7	82.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,477	24.3	74.6
3	Cerebrovascular diseases (I60-I69)	3,543	8.2	25.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,125	4.9	15.1
5	Diabetes mellitus (E10-E14)	1,576	3.6	11.2
6	Influenza and pneumonia (J10-J18)	1,327	3.1	9.5
7	Chronic lower respiratory diseases (J40-J47)	1,302	3.0	9.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	755	1.7	5.4
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	726	1.7	5.2
10	Alzheimer's disease (G30)	615	1.4	4.4
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	582	1.3	4.1
12	Septicemia (A40-A41)	504	1.2	3.6
13	Certain conditions originating in the perinatal period (P00-P96)	463	1.1	3.3
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	428	1.0	3.0
15	Chronic liver disease and cirrhosis (K70,K73-K74)	401	0.9	2.9
...	All other causes (Residual)	6,820	15.8	48.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 1-4 years				
...	All causes	151	100.0	19.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	31	20.5	3.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	20	13.2	2.5
3	Malignant neoplasms (C00-C97)	16	10.6	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	12	7.9	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8	5.3	*
6	Septicemia (A40-A41)	5	3.3	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	3.3	*
8	Influenza and pneumonia (J10-J18)	4	2.6	*
9	Meningitis (G00,G03)	3	2.0	*
9	Cerebrovascular diseases (I60-I69)	3	2.0	*
11	Anemias (D50-D64)	1	0.7	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.7	*
...	All other causes (Residual)	42	27.8	5.3
Asian or Pacific Islander, both sexes, 5-14 years				
...	All causes	239	100.0	12.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	74	31.0	4.0
2	Malignant neoplasms (C00-C97)	37	15.5	2.0
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	17	7.1	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	16	6.7	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	14	5.9	*
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8	3.3	*
7	Cerebrovascular diseases (I60-I69)	7	2.9	*
8	Influenza and pneumonia (J10-J18)	6	2.5	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	2.1	*
10	Chronic lower respiratory diseases (J40-J47)	4	1.7	*
11	Septicemia (A40-A41)	3	1.3	*
12	Meningococcal infection (A39)	1	0.4	*
12	Diabetes mellitus (E10-E14)	1	0.4	*
12	Pneumonitis due to solids and liquids (J69)	1	0.4	*
12	Diseases of appendix (K35-K38)	1	0.4	*
12	Certain conditions originating in the perinatal period (P00-P96)	1	0.4	*
...	All other causes (Residual)	43	18.0	2.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 15-24 years				
...	All causes	818	100.0	41.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	327	40.0	16.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	135	16.5	6.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	107	13.1	5.5
4	Malignant neoplasms (C00-C97)	73	8.9	3.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	36	4.4	1.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	17	2.1	*
7	Cerebrovascular diseases (I60-I69)	7	0.9	*
8	Septicemia (A40-A41)	6	0.7	*
8	Influenza and pneumonia (J10-J18)	6	0.7	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.5	*
11	Tuberculosis (A16-A19)	3	0.4	*
11	Anemias (D50-D64)	3	0.4	*
13	Viral hepatitis (B15-B19)	2	0.2	*
13	Meningitis (G00,G03)	2	0.2	*
13	Chronic lower respiratory diseases (J40-J47)	2	0.2	*
13	Pregnancy, childbirth and the puerperium (O00-099)	2	0.2	*
...	All other causes (Residual)	86	10.5	4.4
Asian or Pacific Islander, both sexes, 25-44 years				
...	All causes	2,858	100.0	58.0
1	Malignant neoplasms (C00-C97)	728	25.5	14.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	483	16.9	9.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	375	13.1	7.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	315	11.0	6.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	159	5.6	3.2
6	Cerebrovascular diseases (I60-I69)	112	3.9	2.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	51	1.8	1.0
8	Diabetes mellitus (E10-E14)	44	1.5	0.9
9	Human immunodeficiency virus (HIV) disease (B20-B24)	41	1.4	0.8
10	Pregnancy, childbirth and the puerperium (O00-099)	27	0.9	0.5
11	Aortic aneurysm and dissection (I71)	25	0.9	0.5
11	Influenza and pneumonia (J10-J18)	25	0.9	0.5
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	20	0.7	0.4
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19	0.7	*
15	Septicemia (A40-A41)	18	0.6	*
15	Chronic lower respiratory diseases (J40-J47)	18	0.6	*
...	All other causes (Residual)	398	13.9	8.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 45-64 years				
...	All causes	9,057	100.0	288.4
1	Malignant neoplasms (C00-C97)	3,708	40.9	118.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,884	20.8	60.0
3	Cerebrovascular diseases (I60-I69)	599	6.6	19.1
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	472	5.2	15.0
5	Diabetes mellitus (E10-E14)	330	3.6	10.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	194	2.1	6.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	169	1.9	5.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	118	1.3	3.8
9	Chronic lower respiratory diseases (J40-J47)	110	1.2	3.5
10	Influenza and pneumonia (J10-J18)	102	1.1	3.2
11	Viral hepatitis (B15-B19)	88	1.0	2.8
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	82	0.9	2.6
13	Septicemia (A40-A41)	78	0.9	2.5
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	67	0.7	2.1
15	Aortic aneurysm and dissection (I71)	64	0.7	2.0
...	All other causes (Residual)	992	11.0	31.6
Asian or Pacific Islander, both sexes, 65 years and over				
...	All causes	29,196	100.0	2,468.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,137	27.9	688.0
2	Malignant neoplasms (C00-C97)	6,987	23.9	590.8
3	Cerebrovascular diseases (I60-I69)	2,810	9.6	237.6
4	Diabetes mellitus (E10-E14)	1,200	4.1	101.5
5	Influenza and pneumonia (J10-J18)	1,180	4.0	99.8
6	Chronic lower respiratory diseases (J40-J47)	1,168	4.0	98.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	722	2.5	61.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	614	2.1	51.9
9	Alzheimer's disease (G30)	604	2.1	51.1
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	499	1.7	42.2
11	Septicemia (A40-A41)	385	1.3	32.6
12	Parkinson's disease (G20-G21)	263	0.9	22.2
13	Aortic aneurysm and dissection (I71)	251	0.9	21.2
14	Pneumonitis due to solids and liquids (J69)	238	0.8	20.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	180	0.6	15.2
...	All other causes (Residual)	3,958	13.6	334.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, all ages				
...	All causes	22,808	100.0	333.9
1	Malignant neoplasms (C00-C97)	5,921	26.0	86.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,703	25.0	83.5
3	Cerebrovascular diseases (I60-I69)	1,624	7.1	23.8
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,328	5.8	19.4
5	Chronic lower respiratory diseases (J40-J47)	803	3.5	11.8
6	Diabetes mellitus (E10-E14)	768	3.4	11.2
7	Influenza and pneumonia (J10-J18)	718	3.1	10.5
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	495	2.2	7.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	381	1.7	5.6
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	314	1.4	4.6
11	Septicemia (A40-A41)	260	1.1	3.8
12	Certain conditions originating in the perinatal period (P00-P96)	258	1.1	3.8
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	256	1.1	3.7
14	Chronic liver disease and cirrhosis (K70,K73-K74)	254	1.1	3.7
15	Alzheimer's disease (G30)	229	1.0	3.4
...	All other causes (Residual)	3,496	15.3	51.2
Asian or Pacific Islander, male, 1-4 years				
...	All causes	84	100.0	20.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19	22.6	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	11	13.1	*
3	Malignant neoplasms (C00-C97)	9	10.7	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	6	7.1	*
5	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	3.6	*
5	Meningitis (G00,G03)	3	3.6	*
5	Cerebrovascular diseases (I60-I69)	3	3.6	*
5	Influenza and pneumonia (J10-J18)	3	3.6	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	3.6	*
10	Septicemia (A40-A41)	1	1.2	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	1	1.2	*
...	All other causes (Residual)	22	26.2	5.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 5-14 years				
...	All causes	131	100.0	14.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	42	32.1	4.5
2	Malignant neoplasms (C00-C97)	21	16.0	2.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	6.1	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8	6.1	*
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6	4.6	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	3.8	*
6	Cerebrovascular diseases (I60-I69)	5	3.8	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	2.3	*
8	Influenza and pneumonia (J10-J18)	3	2.3	*
10	Septicemia (A40-A41)	2	1.5	*
10	Chronic lower respiratory diseases (J40-J47)	2	1.5	*
12	Meningococcal infection (A39)	1	0.8	*
12	Diabetes mellitus (E10-E14)	1	0.8	*
...	All other causes (Residual)	24	18.3	2.6
Asian or Pacific Islander, male, 15-24 years				
...	All causes	569	100.0	56.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	235	41.3	23.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	108	19.0	10.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	72	12.7	7.2
4	Malignant neoplasms (C00-C97)	44	7.7	4.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	25	4.4	2.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	10	1.8	*
7	Cerebrovascular diseases (I60-I69)	4	0.7	*
7	Influenza and pneumonia (J10-J18)	4	0.7	*
9	Tuberculosis (A16-A19)	2	0.4	*
9	Septicemia (A40-A41)	2	0.4	*
9	Anemias (D50-D64)	2	0.4	*
12	Viral hepatitis (B15-B19)	1	0.2	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.2	*
12	Diabetes mellitus (E10-E14)	1	0.2	*
12	Meningitis (G00,G03)	1	0.2	*
12	Chronic lower respiratory diseases (J40-J47)	1	0.2	*
12	Pneumonitis due to solids and liquids (J69)	1	0.2	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
...	All other causes (Residual)	53	9.3	5.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 25-44 years				
...	All causes	1,778	100.0	73.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	340	19.1	14.1
2	Malignant neoplasms (C00-C97)	321	18.1	13.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	285	16.0	11.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	230	12.9	9.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	117	6.6	4.9
6	Cerebrovascular diseases (I60-I69)	59	3.3	2.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	41	2.3	1.7
8	Human immunodeficiency virus (HIV) disease (B20-B24)	34	1.9	1.4
9	Diabetes mellitus (E10-E14)	27	1.5	1.1
10	Aortic aneurysm and dissection (I71)	23	1.3	1.0
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	0.9	*
12	Influenza and pneumonia (J10-J18)	13	0.7	*
12	Chronic lower respiratory diseases (J40-J47)	13	0.7	*
14	Viral hepatitis (B15-B19)	11	0.6	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	11	0.6	*
...	All other causes (Residual)	237	13.3	9.8
Asian or Pacific Islander, male, 45-64 years				
...	All causes	5,287	100.0	359.8
1	Malignant neoplasms (C00-C97)	1,818	34.4	123.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,350	25.5	91.9
3	Cerebrovascular diseases (I60-I69)	320	6.1	21.8
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	300	5.7	20.4
5	Diabetes mellitus (E10-E14)	213	4.0	14.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	131	2.5	8.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	127	2.4	8.6
8	Chronic lower respiratory diseases (J40-J47)	75	1.4	5.1
9	Influenza and pneumonia (J10-J18)	64	1.2	4.4
10	Viral hepatitis (B15-B19)	62	1.2	4.2
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	61	1.2	4.2
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	60	1.1	4.1
13	Aortic aneurysm and dissection (I71)	48	0.9	3.3
14	Septicemia (A40-A41)	37	0.7	2.5
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	37	0.7	2.5
...	All other causes (Residual)	584	11.0	39.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 65 years and over				
...	All causes	14,474	100.0	2,851.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,022	27.8	792.3
2	Malignant neoplasms (C00-C97)	3,707	25.6	730.2
3	Cerebrovascular diseases (I60-I69)	1,230	8.5	242.3
4	Chronic lower respiratory diseases (J40-J47)	712	4.9	140.3
5	Influenza and pneumonia (J10-J18)	628	4.3	123.7
6	Diabetes mellitus (E10-E14)	526	3.6	103.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	384	2.7	75.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	304	2.1	59.9
9	Alzheimer's disease (G30)	225	1.6	44.3
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	209	1.4	41.2
11	Septicemia (A40-A41)	202	1.4	39.8
12	Aortic aneurysm and dissection (I71)	154	1.1	30.3
13	Parkinson's disease (G20-G21)	145	1.0	28.6
14	Pneumonitis due to solids and liquids (J69)	132	0.9	26.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	92	0.6	18.1
...	All other causes (Residual)	1,802	12.4	355.0
Asian or Pacific Islander, female, all ages				
...	All causes	20,386	100.0	282.8
1	Malignant neoplasms (C00-C97)	5,629	27.6	78.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,774	23.4	66.2
3	Cerebrovascular diseases (I60-I69)	1,919	9.4	26.6
4	Diabetes mellitus (E10-E14)	808	4.0	11.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	797	3.9	11.1
6	Influenza and pneumonia (J10-J18)	609	3.0	8.4
7	Chronic lower respiratory diseases (J40-J47)	499	2.4	6.9
8	Alzheimer's disease (G30)	386	1.9	5.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	374	1.8	5.2
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	326	1.6	4.5
11	Septicemia (A40-A41)	244	1.2	3.4
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	231	1.1	3.2
13	Certain conditions originating in the perinatal period (P00-P96)	205	1.0	2.8
14	Chronic liver disease and cirrhosis (K70,K73-K74)	147	0.7	2.0
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	146	0.7	2.0
...	All other causes (Residual)	3,292	16.1	45.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 1-4 years				
...	All causes	67	100.0	17.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12	17.9	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	13.4	*
3	Malignant neoplasms (C00-C97)	7	10.4	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	6	9.0	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	7.5	*
6	Septicemia (A40-A41)	4	6.0	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	3.0	*
8	Anemias (D50-D64)	1	1.5	*
8	Influenza and pneumonia (J10-J18)	1	1.5	*
...	All other causes (Residual)	20	29.9	5.2
Asian or Pacific Islander, female, 5-14 years				
...	All causes	108	100.0	11.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	32	29.6	3.5
2	Malignant neoplasms (C00-C97)	16	14.8	*
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	9	8.3	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	8.3	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8	7.4	*
6	Influenza and pneumonia (J10-J18)	3	2.8	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	1.9	*
7	Cerebrovascular diseases (I60-I69)	2	1.9	*
7	Chronic lower respiratory diseases (J40-J47)	2	1.9	*
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2	1.9	*
11	Septicemia (A40-A41)	1	0.9	*
11	Pneumonitis due to solids and liquids (J69)	1	0.9	*
11	Diseases of appendix (K35-K38)	1	0.9	*
11	Certain conditions originating in the perinatal period (P00-P96)	1	0.9	*
...	All other causes (Residual)	19	17.6	*

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 15-24 years				
...	All causes	249	100.0	26.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	92	36.9	9.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	35	14.1	3.7
3	Malignant neoplasms (C00-C97)	29	11.6	3.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	27	10.8	2.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	11	4.4	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	2.8	*
7	Septicemia (A40-A41)	4	1.6	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	1.2	*
8	Cerebrovascular diseases (I60-I69)	3	1.2	*
10	Influenza and pneumonia (J10-J18)	2	0.8	*
10	Pregnancy, childbirth and the puerperium (O00-099)	2	0.8	*
12	Tuberculosis (A16-A19)	1	0.4	*
12	Viral hepatitis (B15-B19)	1	0.4	*
12	Anemias (D50-D64)	1	0.4	*
12	Nutritional deficiencies (E40-E64)	1	0.4	*
12	Meningitis (G00,G03)	1	0.4	*
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1	0.4	*
12	Chronic lower respiratory diseases (J40-J47)	1	0.4	*
12	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.4	*
...	All other causes (Residual)	26	10.4	2.7
Asian or Pacific Islander, female, 25-44 years				
...	All causes	1,080	100.0	42.9
1	Malignant neoplasms (C00-C97)	407	37.7	16.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	143	13.2	5.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	90	8.3	3.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	85	7.9	3.4
5	Cerebrovascular diseases (I60-I69)	53	4.9	2.1
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	42	3.9	1.7
7	Pregnancy, childbirth and the puerperium (O00-099)	27	2.5	1.1
8	Diabetes mellitus (E10-E14)	17	1.6	*
9	Influenza and pneumonia (J10-J18)	12	1.1	*
10	Chronic liver disease and cirrhosis (K70,K73-K74)	10	0.9	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	0.8	*
12	Septicemia (A40-A41)	8	0.7	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	7	0.6	*
14	Viral hepatitis (B15-B19)	5	0.5	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.5	*
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	5	0.5	*
14	Chronic lower respiratory diseases (J40-J47)	5	0.5	*
...	All other causes (Residual)	150	13.9	6.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2005

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 45-64 years				
...	All causes	3,770	100.0	225.6
1	Malignant neoplasms (C00-C97)	1,890	50.1	113.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	534	14.2	32.0
3	Cerebrovascular diseases (I60-I69)	279	7.4	16.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	172	4.6	10.3
5	Diabetes mellitus (E10-E14)	117	3.1	7.0
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	63	1.7	3.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	58	1.5	3.5
8	Chronic liver disease and cirrhosis (K70,K73-K74)	42	1.1	2.5
9	Septicemia (A40-A41)	41	1.1	2.5
10	Influenza and pneumonia (J10-J18)	38	1.0	2.3
11	Chronic lower respiratory diseases (J40-J47)	35	0.9	2.1
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	30	0.8	1.8
13	Viral hepatitis (B15-B19)	26	0.7	1.6
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	26	0.7	1.6
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	21	0.6	1.3
...	All other causes (Residual)	398	10.6	23.8
Asian or Pacific Islander, female, 65 years and over				
...	All causes	14,722	100.0	2,180.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,115	28.0	609.6
2	Malignant neoplasms (C00-C97)	3,280	22.3	485.9
3	Cerebrovascular diseases (I60-I69)	1,580	10.7	234.0
4	Diabetes mellitus (E10-E14)	674	4.6	99.8
5	Influenza and pneumonia (J10-J18)	552	3.7	81.8
6	Chronic lower respiratory diseases (J40-J47)	456	3.1	67.5
7	Alzheimer's disease (G30)	379	2.6	56.1
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	338	2.3	50.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	310	2.1	45.9
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	290	2.0	43.0
11	Septicemia (A40-A41)	183	1.2	27.1
12	Parkinson's disease (G20-G21)	118	0.8	17.5
13	Pneumonitis due to solids and liquids (J69)	106	0.7	15.7
14	Aortic aneurysm and dissection (I71)	97	0.7	14.4
15	Chronic liver disease and cirrhosis (K70,K73-K74)	94	0.6	13.9
...	All other causes (Residual)	2,150	14.6	318.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable