

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, all ages				
...	All causes	2,437,163	100.0	793.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	599,413	24.6	195.2
2	Malignant neoplasms (C00-C97)	567,628	23.3	184.9
3	Chronic lower respiratory diseases (J40-J47)	137,353	5.6	44.7
4	Cerebrovascular diseases (I60-I69)	128,842	5.3	42.0
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	118,021	4.8	38.4
6	Alzheimer's disease (G30)	79,003	3.2	25.7
7	Diabetes mellitus (E10-E14)	68,705	2.8	22.4
8	Influenza and pneumonia (J09-J18)	53,692	2.2	17.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	48,935	2.0	15.9
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	36,909	1.5	12.0
11	Septicemia (A40-A41)	35,639	1.5	11.6
12	Chronic liver disease and cirrhosis (K70,K73-K74)	30,558	1.3	10.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	25,734	1.1	8.4
14	Parkinson's disease (G20-G21)	20,565	0.8	6.7
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	16,799	0.7	5.5
...	All other causes (Residual)	469,367	19.3	152.9
All races, both sexes, 1-4 years				
...	All causes	4,450	100.0	26.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,466	32.9	8.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	464	10.4	2.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	376	8.4	2.2
4	Malignant neoplasms (C00-C97)	350	7.9	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	154	3.5	0.9
6	Influenza and pneumonia (J09-J18)	146	3.3	0.9
7	Septicemia (A40-A41)	71	1.6	0.4
8	Chronic lower respiratory diseases (J40-J47)	66	1.5	0.4
9	Certain conditions originating in the perinatal period (P00-P96)	58	1.3	0.3
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	53	1.2	0.3
11	Cerebrovascular diseases (I60-I69)	47	1.1	0.3
12	Anemias (D50-D64)	20	0.4	0.1
13	Acute bronchitis and bronchiolitis (J20-J21)	19	0.4	*
14	Meningitis (G00,G03)	18	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	18	0.4	*
...	All other causes (Residual)	1,124	25.3	6.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 5-14 years				
...	All causes	5,651	100.0	13.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,689	29.9	4.2
2	Malignant neoplasms (C00-C97)	896	15.9	2.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	364	6.4	0.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	305	5.4	0.8
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	265	4.7	0.7
6	Influenza and pneumonia (J09-J18)	228	4.0	0.6
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	217	3.8	0.5
8	Chronic lower respiratory diseases (J40-J47)	123	2.2	0.3
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	85	1.5	0.2
10	Cerebrovascular diseases (I60-I69)	74	1.3	0.2
11	Septicemia (A40-A41)	70	1.2	0.2
12	Anemias (D50-D64)	35	0.6	0.1
13	Diabetes mellitus (E10-E14)	29	0.5	0.1
14	Certain conditions originating in the perinatal period (P00-P96)	26	0.5	0.1
15	Meningitis (G00,G03)	22	0.4	0.1
...	All other causes (Residual)	1,223	21.6	3.0
All races, both sexes, 15-24 years				
...	All causes	30,416	100.0	70.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,458	41.0	28.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4,862	16.0	11.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,371	14.4	10.1
4	Malignant neoplasms (C00-C97)	1,636	5.4	3.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,035	3.4	2.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	457	1.5	1.1
7	Influenza and pneumonia (J09-J18)	418	1.4	1.0
8	Pregnancy, childbirth and the puerperium (O00-O99)	227	0.7	0.5
9	Cerebrovascular diseases (I60-I69)	193	0.6	0.4
10	Chronic lower respiratory diseases (J40-J47)	187	0.6	0.4
11	Diabetes mellitus (E10-E14)	167	0.5	0.4
12	Human immunodeficiency virus (HIV) disease (B20-B24)	149	0.5	0.3
13	Septicemia (A40-A41)	144	0.5	0.3
14	Anemias (D50-D64)	93	0.3	0.2
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	86	0.3	0.2
...	All other causes (Residual)	3,933	12.9	9.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 25-44 years				
...	All causes	117,167	100.0	141.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	29,164	24.9	35.1
2	Malignant neoplasms (C00-C97)	16,178	13.8	19.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,255	12.2	17.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11,997	10.2	14.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6,984	6.0	8.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	3,306	2.8	4.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,940	2.5	3.5
8	Diabetes mellitus (E10-E14)	2,476	2.1	3.0
9	Cerebrovascular diseases (I60-I69)	2,453	2.1	3.0
10	Influenza and pneumonia (J09-J18)	2,121	1.8	2.6
11	Septicemia (A40-A41)	1,308	1.1	1.6
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,096	0.9	1.3
13	Chronic lower respiratory diseases (J40-J47)	1,026	0.9	1.2
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	871	0.7	1.0
15	Pregnancy, childbirth and the puerperium (O00-O99)	631	0.5	0.8
...	All other causes (Residual)	20,361	17.4	24.5
All races, both sexes, 45-64 years				
...	All causes	490,875	100.0	618.4
1	Malignant neoplasms (C00-C97)	157,445	32.1	198.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	104,188	21.2	131.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	32,907	6.7	41.5
4	Chronic lower respiratory diseases (J40-J47)	18,824	3.8	23.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	17,531	3.6	22.1
6	Diabetes mellitus (E10-E14)	17,086	3.5	21.5
7	Cerebrovascular diseases (I60-I69)	16,686	3.4	21.0
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	14,406	2.9	18.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,144	1.5	9.0
10	Septicemia (A40-A41)	7,061	1.4	8.9
11	Influenza and pneumonia (J09-J18)	7,057	1.4	8.9
12	Viral hepatitis (B15-B19)	5,665	1.2	7.1
13	Human immunodeficiency virus (HIV) disease (B20-B24)	5,242	1.1	6.6
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,893	0.8	4.9
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,077	0.6	3.9
...	All other causes (Residual)	72,663	14.8	91.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 65 years and over				
...	All causes	1,761,937	100.0	4,452.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	479,150	27.2	1,210.9
2	Malignant neoplasms (C00-C97)	391,035	22.2	988.2
3	Chronic lower respiratory diseases (J40-J47)	117,098	6.6	295.9
4	Cerebrovascular diseases (I60-I69)	109,238	6.2	276.1
5	Alzheimer's disease (G30)	78,168	4.4	197.5
6	Diabetes mellitus (E10-E14)	48,944	2.8	123.7
7	Influenza and pneumonia (J09-J18)	43,469	2.5	109.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	40,465	2.3	102.3
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	39,111	2.2	98.8
10	Septicemia (A40-A41)	26,763	1.5	67.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	21,261	1.2	53.7
12	Parkinson's disease (G20-G21)	20,028	1.1	50.6
13	Pneumonitis due to solids and liquids (J69)	14,150	0.8	35.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11,669	0.7	29.5
15	Chronic liver disease and cirrhosis (K70,K73-K74)	10,055	0.6	25.4
...	All other causes (Residual)	311,333	17.7	786.8
All races, male, all ages ³				
...	All causes	1,217,379	100.0	803.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	307,225	25.2	202.9
2	Malignant neoplasms (C00-C97)	296,763	24.4	195.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	75,022	6.2	49.5
4	Chronic lower respiratory diseases (J40-J47)	65,119	5.3	43.0
5	Cerebrovascular diseases (I60-I69)	52,073	4.3	34.4
6	Diabetes mellitus (E10-E14)	35,054	2.9	23.1
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	29,089	2.4	19.2
8	Influenza and pneumonia (J09-J18)	25,128	2.1	16.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23,930	2.0	15.8
10	Alzheimer's disease (G30)	23,900	2.0	15.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	19,804	1.6	13.1
12	Septicemia (A40-A41)	16,371	1.3	10.8
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	13,126	1.1	8.7
14	Parkinson's disease (G20-G21)	12,088	1.0	8.0
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10,461	0.9	6.9
...	All other causes (Residual)	212,226	17.4	140.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 1-4 years				
...	All causes	2,495	100.0	28.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	925	37.1	10.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	230	9.2	2.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	188	7.5	2.2
4	Malignant neoplasms (C00-C97)	180	7.2	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	77	3.1	0.9
6	Influenza and pneumonia (J09-J18)	64	2.6	0.7
7	Septicemia (A40-A41)	40	1.6	0.5
8	Chronic lower respiratory diseases (J40-J47)	36	1.4	0.4
9	Certain conditions originating in the perinatal period (P00-P96)	33	1.3	0.4
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	30	1.2	0.3
11	Cerebrovascular diseases (I60-I69)	24	1.0	0.3
12	Anemias (D50-D64)	15	0.6	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	12	0.5	*
14	Meningitis (G00,G03)	11	0.4	*
15	Meningococcal infection (A39)	9	0.4	*
...	All other causes (Residual)	621	24.9	7.1
All races, male, 5-14 years				
...	All causes	3,248	100.0	15.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,052	32.4	5.1
2	Malignant neoplasms (C00-C97)	488	15.0	2.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	206	6.3	1.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	196	6.0	0.9
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	175	5.4	0.8
6	Influenza and pneumonia (J09-J18)	114	3.5	0.5
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	112	3.4	0.5
8	Chronic lower respiratory diseases (J40-J47)	77	2.4	0.4
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	51	1.6	0.2
10	Cerebrovascular diseases (I60-I69)	45	1.4	0.2
11	Septicemia (A40-A41)	27	0.8	0.1
12	Anemias (D50-D64)	14	0.4	*
13	Diabetes mellitus (E10-E14)	12	0.4	*
13	Meningitis (G00,G03)	12	0.4	*
15	Certain conditions originating in the perinatal period (P00-P96)	11	0.3	*
...	All other causes (Residual)	656	20.2	3.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 15-24 years				
...	All causes	22,312	100.0	100.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,239	41.4	41.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4,205	18.8	19.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,595	16.1	16.2
4	Malignant neoplasms (C00-C97)	993	4.5	4.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	687	3.1	3.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	263	1.2	1.2
7	Influenza and pneumonia (J09-J18)	224	1.0	1.0
8	Cerebrovascular diseases (I60-I69)	108	0.5	0.5
8	Chronic lower respiratory diseases (J40-J47)	108	0.5	0.5
10	Human immunodeficiency virus (HIV) disease (B20-B24)	90	0.4	0.4
10	Diabetes mellitus (E10-E14)	90	0.4	0.4
12	Septicemia (A40-A41)	81	0.4	0.4
13	Legal intervention (Y35,Y89.0)	71	0.3	0.3
14	Anemias (D50-D64)	52	0.2	0.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	47	0.2	0.2
...	All other causes (Residual)	2,459	11.0	11.1
All races, male, 25-44 years				
...	All causes	75,754	100.0	180.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	21,181	28.0	50.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,902	13.1	23.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	9,435	12.5	22.4
4	Malignant neoplasms (C00-C97)	7,205	9.5	17.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5,603	7.4	13.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	2,250	3.0	5.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,953	2.6	4.6
8	Diabetes mellitus (E10-E14)	1,561	2.1	3.7
9	Cerebrovascular diseases (I60-I69)	1,342	1.8	3.2
10	Influenza and pneumonia (J09-J18)	1,154	1.5	2.7
11	Septicemia (A40-A41)	655	0.9	1.6
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	608	0.8	1.4
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	515	0.7	1.2
14	Chronic lower respiratory diseases (J40-J47)	498	0.7	1.2
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	343	0.5	0.8
...	All other causes (Residual)	11,549	15.2	27.4

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 45-64 years				
...	All causes	298,962	100.0	771.4
1	Malignant neoplasms (C00-C97)	84,505	28.3	218.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	73,234	24.5	189.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	22,414	7.5	57.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	12,231	4.1	31.6
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	10,926	3.7	28.2
6	Diabetes mellitus (E10-E14)	10,439	3.5	26.9
7	Chronic lower respiratory diseases (J40-J47)	9,624	3.2	24.8
8	Cerebrovascular diseases (I60-I69)	9,362	3.1	24.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,054	1.4	10.5
10	Viral hepatitis (B15-B19)	4,033	1.3	10.4
11	Human immunodeficiency virus (HIV) disease (B20-B24)	3,924	1.3	10.1
12	Influenza and pneumonia (J09-J18)	3,888	1.3	10.0
13	Septicemia (A40-A41)	3,759	1.3	9.7
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,419	0.8	6.2
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,231	0.7	5.8
...	All other causes (Residual)	41,919	14.0	108.2
All races, male, 65 years and over				
...	All causes	799,608	100.0	4,752.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	222,976	27.9	1,325.4
2	Malignant neoplasms (C00-C97)	203,343	25.4	1,208.7
3	Chronic lower respiratory diseases (J40-J47)	54,758	6.8	325.5
4	Cerebrovascular diseases (I60-I69)	41,102	5.1	244.3
5	Alzheimer's disease (G30)	23,516	2.9	139.8
6	Diabetes mellitus (E10-E14)	22,949	2.9	136.4
7	Influenza and pneumonia (J09-J18)	19,532	2.4	116.1
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19,477	2.4	115.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19,143	2.4	113.8
10	Parkinson's disease (G20-G21)	11,740	1.5	69.8
11	Septicemia (A40-A41)	11,689	1.5	69.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7,687	1.0	45.7
13	Pneumonitis due to solids and liquids (J69)	7,216	0.9	42.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5,904	0.7	35.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	5,595	0.7	33.3
...	All other causes (Residual)	122,981	15.4	731.0

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, all ages				
...	All causes	1,219,784	100.0	784.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	292,188	24.0	187.8
2	Malignant neoplasms (C00-C97)	270,865	22.2	174.1
3	Cerebrovascular diseases (I60-I69)	76,769	6.3	49.4
4	Chronic lower respiratory diseases (J40-J47)	72,234	5.9	46.4
5	Alzheimer's disease (G30)	55,103	4.5	35.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	42,999	3.5	27.6
7	Diabetes mellitus (E10-E14)	33,651	2.8	21.6
8	Influenza and pneumonia (J09-J18)	28,564	2.3	18.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	25,005	2.0	16.1
10	Septicemia (A40-A41)	19,268	1.6	12.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	15,273	1.3	9.8
12	Chronic liver disease and cirrhosis (K70,K73-K74)	10,754	0.9	6.9
13	Parkinson's disease (G20-G21)	8,477	0.7	5.4
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7,820	0.6	5.0
15	Pneumonitis due to solids and liquids (J69)	7,649	0.6	4.9
...	All other causes (Residual)	253,165	20.8	162.7
All races, female, 1-4 years				
...	All causes	1,955	100.0	23.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	541	27.7	6.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	234	12.0	2.8
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	188	9.6	2.3
4	Malignant neoplasms (C00-C97)	170	8.7	2.0
5	Influenza and pneumonia (J09-J18)	82	4.2	1.0
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	77	3.9	0.9
7	Septicemia (A40-A41)	31	1.6	0.4
8	Chronic lower respiratory diseases (J40-J47)	30	1.5	0.4
9	Certain conditions originating in the perinatal period (P00-P96)	25	1.3	0.3
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	1.2	0.3
10	Cerebrovascular diseases (I60-I69)	23	1.2	0.3
12	Acute bronchitis and bronchiolitis (J20-J21)	11	0.6	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.4	*
14	Meningitis (G00,G03)	7	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.3	*
...	All other causes (Residual)	499	25.5	6.0

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 5-14 years				
...	All causes	2,403	100.0	12.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	637	26.5	3.2
2	Malignant neoplasms (C00-C97)	408	17.0	2.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	158	6.6	0.8
4	Influenza and pneumonia (J09-J18)	114	4.7	0.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	109	4.5	0.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	105	4.4	0.5
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	90	3.7	0.5
8	Chronic lower respiratory diseases (J40-J47)	46	1.9	0.2
9	Septicemia (A40-A41)	43	1.8	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	34	1.4	0.2
11	Cerebrovascular diseases (I60-I69)	29	1.2	0.1
12	Anemias (D50-D64)	21	0.9	0.1
13	Diabetes mellitus (E10-E14)	17	0.7	*
14	Certain conditions originating in the perinatal period (P00-P96)	15	0.6	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	0.5	*
...	All other causes (Residual)	566	23.6	2.9
All races, female, 15-24 years				
...	All causes	8,104	100.0	38.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,219	39.7	15.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	776	9.6	3.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	657	8.1	3.1
4	Malignant neoplasms (C00-C97)	643	7.9	3.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	348	4.3	1.7
6	Pregnancy, childbirth and the puerperium (O00-O99)	227	2.8	1.1
7	Influenza and pneumonia (J09-J18)	194	2.4	0.9
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	194	2.4	0.9
9	Cerebrovascular diseases (I60-I69)	85	1.0	0.4
10	Chronic lower respiratory diseases (J40-J47)	79	1.0	0.4
11	Diabetes mellitus (E10-E14)	77	1.0	0.4
12	Septicemia (A40-A41)	63	0.8	0.3
13	Human immunodeficiency virus (HIV) disease (B20-B24)	59	0.7	0.3
14	Anemias (D50-D64)	41	0.5	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	41	0.5	0.2
...	All other causes (Residual)	1,401	17.3	6.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 25-44 years				
...	All causes	41,413	100.0	101.0
1	Malignant neoplasms (C00-C97)	8,973	21.7	21.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,983	19.3	19.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,353	10.5	10.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,562	6.2	6.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,381	3.3	3.4
6	Cerebrovascular diseases (I60-I69)	1,111	2.7	2.7
7	Human immunodeficiency virus (HIV) disease (B20-B24)	1,056	2.5	2.6
8	Chronic liver disease and cirrhosis (K70,K73-K74)	987	2.4	2.4
9	Influenza and pneumonia (J09-J18)	967	2.3	2.4
10	Diabetes mellitus (E10-E14)	915	2.2	2.2
11	Septicemia (A40-A41)	653	1.6	1.6
12	Pregnancy, childbirth and the puerperium (O00-O99)	631	1.5	1.5
13	Chronic lower respiratory diseases (J40-J47)	528	1.3	1.3
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	488	1.2	1.2
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	356	0.9	0.9
...	All other causes (Residual)	8,469	20.5	20.6
All races, female, 45-64 years				
...	All causes	191,913	100.0	472.4
1	Malignant neoplasms (C00-C97)	72,940	38.0	179.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	30,954	16.1	76.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,493	5.5	25.8
4	Chronic lower respiratory diseases (J40-J47)	9,200	4.8	22.6
5	Cerebrovascular diseases (I60-I69)	7,324	3.8	18.0
6	Diabetes mellitus (E10-E14)	6,647	3.5	16.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	5,300	2.8	13.0
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,480	1.8	8.6
9	Septicemia (A40-A41)	3,302	1.7	8.1
10	Influenza and pneumonia (J09-J18)	3,169	1.7	7.8
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,090	1.6	7.6
12	Viral hepatitis (B15-B19)	1,632	0.9	4.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,474	0.8	3.6
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1,318	0.7	3.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	982	0.5	2.4
...	All other causes (Residual)	30,608	15.9	75.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 65 years and over				
...	All causes	962,329	100.0	4,230.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	256,174	26.6	1,126.2
2	Malignant neoplasms (C00-C97)	187,692	19.5	825.1
3	Cerebrovascular diseases (I60-I69)	68,136	7.1	299.5
4	Chronic lower respiratory diseases (J40-J47)	62,340	6.5	274.1
5	Alzheimer's disease (G30)	54,652	5.7	240.3
6	Diabetes mellitus (E10-E14)	25,995	2.7	114.3
7	Influenza and pneumonia (J09-J18)	23,937	2.5	105.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21,322	2.2	93.7
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19,634	2.0	86.3
10	Septicemia (A40-A41)	15,074	1.6	66.3
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	13,574	1.4	59.7
12	Parkinson's disease (G20-G21)	8,288	0.9	36.4
13	Pneumonitis due to solids and liquids (J69)	6,934	0.7	30.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5,765	0.6	25.3
15	Chronic liver disease and cirrhosis (K70,K73-K74)	4,460	0.5	19.6
...	All other causes (Residual)	188,352	19.6	828.0
White, both sexes, all ages				
...	All causes	2,086,355	100.0	844.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	515,587	24.7	208.8
2	Malignant neoplasms (C00-C97)	486,995	23.3	197.2
3	Chronic lower respiratory diseases (J40-J47)	126,733	6.1	51.3
4	Cerebrovascular diseases (I60-I69)	108,764	5.2	44.0
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	102,170	4.9	41.4
6	Alzheimer's disease (G30)	72,696	3.5	29.4
7	Diabetes mellitus (E10-E14)	54,113	2.6	21.9
8	Influenza and pneumonia (J09-J18)	46,432	2.2	18.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	38,758	1.9	15.7
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	33,425	1.6	13.5
11	Septicemia (A40-A41)	28,609	1.4	11.6
12	Chronic liver disease and cirrhosis (K70,K73-K74)	26,820	1.3	10.9
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	20,052	1.0	8.1
14	Parkinson's disease (G20-G21)	19,401	0.9	7.9
15	Pneumonitis due to solids and liquids (J69)	14,133	0.7	5.7
...	All other causes (Residual)	391,667	18.8	158.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 1-4 years				
...	All causes	3,141	100.0	24.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,082	34.4	8.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	350	11.1	2.7
3	Malignant neoplasms (C00-C97)	274	8.7	2.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	211	6.7	1.6
5	Influenza and pneumonia (J09-J18)	108	3.4	0.8
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	96	3.1	0.7
7	Septicemia (A40-A41)	47	1.5	0.4
8	Cerebrovascular diseases (I60-I69)	38	1.2	0.3
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	37	1.2	0.3
10	Certain conditions originating in the perinatal period (P00-P96)	36	1.1	0.3
11	Chronic lower respiratory diseases (J40-J47)	32	1.0	0.2
12	Meningitis (G00,G03)	15	0.5	*
13	Anemias (D50-D64)	12	0.4	*
14	Acute bronchitis and bronchiolitis (J20-J21)	9	0.3	*
15	Meningococcal infection (A39)	8	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.3	*
...	All other causes (Residual)	778	24.8	6.0
White, both sexes, 5-14 years				
...	All causes	4,005	100.0	12.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,219	30.4	3.9
2	Malignant neoplasms (C00-C97)	703	17.6	2.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	259	6.5	0.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	201	5.0	0.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	185	4.6	0.6
6	Influenza and pneumonia (J09-J18)	165	4.1	0.5
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	132	3.3	0.4
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	63	1.6	0.2
9	Cerebrovascular diseases (I60-I69)	53	1.3	0.2
10	Septicemia (A40-A41)	43	1.1	0.1
11	Chronic lower respiratory diseases (J40-J47)	40	1.0	0.1
12	Diabetes mellitus (E10-E14)	21	0.5	0.1
13	Certain conditions originating in the perinatal period (P00-P96)	20	0.5	0.1
14	Anemias (D50-D64)	16	0.4	*
15	Meningitis (G00,G03)	15	0.4	*
...	All other causes (Residual)	870	21.7	2.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 15-24 years				
...	All causes	22,241	100.0	66.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,526	47.3	31.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,619	16.3	10.8
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,902	8.6	5.7
4	Malignant neoplasms (C00-C97)	1,256	5.6	3.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	643	2.9	1.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	340	1.5	1.0
7	Influenza and pneumonia (J09-J18)	308	1.4	0.9
8	Cerebrovascular diseases (I60-I69)	137	0.6	0.4
9	Pregnancy, childbirth and the puerperium (O00-O99)	133	0.6	0.4
10	Diabetes mellitus (E10-E14)	113	0.5	0.3
11	Septicemia (A40-A41)	108	0.5	0.3
12	Chronic lower respiratory diseases (J40-J47)	92	0.4	0.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	52	0.2	0.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	52	0.2	0.2
15	Human immunodeficiency virus (HIV) disease (B20-B24)	43	0.2	0.1
...	All other causes (Residual)	2,917	13.1	8.7
White, both sexes, 25-44 years				
...	All causes	87,826	100.0	134.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	24,686	28.1	37.7
2	Malignant neoplasms (C00-C97)	12,514	14.2	19.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	10,585	12.1	16.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,837	11.2	15.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,221	3.7	4.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	2,459	2.8	3.8
7	Diabetes mellitus (E10-E14)	1,654	1.9	2.5
8	Cerebrovascular diseases (I60-I69)	1,603	1.8	2.4
9	Influenza and pneumonia (J09-J18)	1,595	1.8	2.4
10	Human immunodeficiency virus (HIV) disease (B20-B24)	1,259	1.4	1.9
11	Septicemia (A40-A41)	863	1.0	1.3
12	Chronic lower respiratory diseases (J40-J47)	709	0.8	1.1
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	708	0.8	1.1
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	631	0.7	1.0
15	Pregnancy, childbirth and the puerperium (O00-O99)	399	0.5	0.6
...	All other causes (Residual)	15,103	17.2	23.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 45-64 years				
...	All causes	389,917	100.0	592.3
1	Malignant neoplasms (C00-C97)	127,970	32.8	194.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	80,195	20.6	121.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	27,951	7.2	42.5
4	Chronic lower respiratory diseases (J40-J47)	16,471	4.2	25.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	15,157	3.9	23.0
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	13,451	3.4	20.4
7	Diabetes mellitus (E10-E14)	12,619	3.2	19.2
8	Cerebrovascular diseases (I60-I69)	11,479	2.9	17.4
9	Influenza and pneumonia (J09-J18)	5,610	1.4	8.5
10	Septicemia (A40-A41)	5,170	1.3	7.9
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,618	1.2	7.0
12	Viral hepatitis (B15-B19)	4,603	1.2	7.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,387	0.6	3.6
14	Human immunodeficiency virus (HIV) disease (B20-B24)	2,255	0.6	3.4
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,936	0.5	2.9
...	All other causes (Residual)	58,045	14.9	88.2
White, both sexes, 65 years and over				
...	All causes	1,562,192	100.0	4,534.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	424,417	27.2	1,231.8
2	Malignant neoplasms (C00-C97)	344,214	22.0	999.0
3	Chronic lower respiratory diseases (J40-J47)	109,371	7.0	317.4
4	Cerebrovascular diseases (I60-I69)	95,353	6.1	276.7
5	Alzheimer's disease (G30)	71,946	4.6	208.8
6	Diabetes mellitus (E10-E14)	39,704	2.5	115.2
7	Influenza and pneumonia (J09-J18)	38,495	2.5	111.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	35,929	2.3	104.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	33,361	2.1	96.8
10	Septicemia (A40-A41)	22,255	1.4	64.6
11	Parkinson's disease (G20-G21)	18,927	1.2	54.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	17,381	1.1	50.4
13	Pneumonitis due to solids and liquids (J69)	12,677	0.8	36.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10,633	0.7	30.9
15	Chronic liver disease and cirrhosis (K70,K73-K74)	9,183	0.6	26.7
...	All other causes (Residual)	278,346	17.8	807.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, male, all ages				
...	All causes	1,037,475	100.0	846.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	263,976	25.4	215.4
2	Malignant neoplasms (C00-C97)	255,042	24.6	208.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	64,501	6.2	52.6
4	Chronic lower respiratory diseases (J40-J47)	59,465	5.7	48.5
5	Cerebrovascular diseases (I60-I69)	43,190	4.2	35.2
6	Diabetes mellitus (E10-E14)	28,205	2.7	23.0
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	26,426	2.5	21.6
8	Alzheimer's disease (G30)	22,066	2.1	18.0
9	Influenza and pneumonia (J09-J18)	21,614	2.1	17.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19,312	1.9	15.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	17,475	1.7	14.3
12	Septicemia (A40-A41)	13,206	1.3	10.8
13	Parkinson's disease (G20-G21)	11,407	1.1	9.3
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	8,034	0.8	6.6
15	Pneumonitis due to solids and liquids (J69)	7,413	0.7	6.0
...	All other causes (Residual)	176,143	17.0	143.7
White, male, 1-4 years				
...	All causes	1,780	100.0	26.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	698	39.2	10.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	169	9.5	2.5
3	Malignant neoplasms (C00-C97)	147	8.3	2.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	100	5.6	1.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	52	2.9	0.8
6	Influenza and pneumonia (J09-J18)	47	2.6	0.7
7	Septicemia (A40-A41)	29	1.6	0.4
8	Certain conditions originating in the perinatal period (P00-P96)	21	1.2	0.3
9	Cerebrovascular diseases (I60-I69)	20	1.1	0.3
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	1.1	*
11	Chronic lower respiratory diseases (J40-J47)	18	1.0	*
12	Anemias (D50-D64)	9	0.5	*
12	Meningitis (G00,G03)	9	0.5	*
14	Meningococcal infection (A39)	7	0.4	*
15	Acute bronchitis and bronchiolitis (J20-J21)	4	0.2	*
...	All other causes (Residual)	431	24.2	6.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 5-14 years				
...	All causes	2,314	100.0	14.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	761	32.9	4.7
2	Malignant neoplasms (C00-C97)	390	16.9	2.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	147	6.4	0.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	131	5.7	0.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	115	5.0	0.7
6	Influenza and pneumonia (J09-J18)	84	3.6	0.5
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	66	2.9	0.4
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	36	1.6	0.2
9	Cerebrovascular diseases (I60-I69)	32	1.4	0.2
10	Chronic lower respiratory diseases (J40-J47)	28	1.2	0.2
11	Septicemia (A40-A41)	14	0.6	*
12	Diabetes mellitus (E10-E14)	10	0.4	*
13	Meningitis (G00,G03)	9	0.4	*
14	Certain conditions originating in the perinatal period (P00-P96)	8	0.3	*
15	Anemias (D50-D64)	7	0.3	*
...	All other causes (Residual)	476	20.6	2.9
White, male, 15-24 years				
...	All causes	16,243	100.0	94.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,862	48.4	45.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,012	18.5	17.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,563	9.6	9.1
4	Malignant neoplasms (C00-C97)	776	4.8	4.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	429	2.6	2.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	194	1.2	1.1
7	Influenza and pneumonia (J09-J18)	169	1.0	1.0
8	Cerebrovascular diseases (I60-I69)	79	0.5	0.5
9	Septicemia (A40-A41)	66	0.4	0.4
10	Diabetes mellitus (E10-E14)	62	0.4	0.4
11	Chronic lower respiratory diseases (J40-J47)	50	0.3	0.3
12	Legal intervention (Y35,Y89.0)	40	0.2	0.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	35	0.2	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	31	0.2	0.2
15	Pneumonitis due to solids and liquids (J69)	26	0.2	0.2
...	All other causes (Residual)	1,849	11.4	10.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 25-44 years				
...	All causes	57,407	100.0	171.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17,924	31.2	53.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8,342	14.5	24.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,015	12.2	20.9
4	Malignant neoplasms (C00-C97)	5,683	9.9	17.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,371	4.1	7.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1,659	2.9	4.9
7	Diabetes mellitus (E10-E14)	1,047	1.8	3.1
8	Human immunodeficiency virus (HIV) disease (B20-B24)	1,005	1.8	3.0
9	Influenza and pneumonia (J09-J18)	900	1.6	2.7
10	Cerebrovascular diseases (I60-I69)	887	1.5	2.6
11	Septicemia (A40-A41)	437	0.8	1.3
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	419	0.7	1.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	366	0.6	1.1
14	Chronic lower respiratory diseases (J40-J47)	316	0.6	0.9
15	Viral hepatitis (B15-B19)	238	0.4	0.7
...	All other causes (Residual)	8,798	15.3	26.2
White, male, 45-64 years				
...	All causes	239,876	100.0	737.8
1	Malignant neoplasms (C00-C97)	68,828	28.7	211.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	57,829	24.1	177.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18,997	7.9	58.4
4	Chronic liver disease and cirrhosis (K70,K73-K74)	10,664	4.4	32.8
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	10,222	4.3	31.4
6	Chronic lower respiratory diseases (J40-J47)	8,391	3.5	25.8
7	Diabetes mellitus (E10-E14)	7,917	3.3	24.3
8	Cerebrovascular diseases (I60-I69)	6,520	2.7	20.1
9	Viral hepatitis (B15-B19)	3,312	1.4	10.2
10	Influenza and pneumonia (J09-J18)	3,100	1.3	9.5
11	Septicemia (A40-A41)	2,775	1.2	8.5
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,699	1.1	8.3
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,863	0.8	5.7
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,547	0.6	4.8
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,340	0.6	4.1
...	All other causes (Residual)	33,872	14.1	104.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 65 years and over				
...	All causes	710,286	100.0	4,809.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	198,438	27.9	1,343.7
2	Malignant neoplasms (C00-C97)	179,182	25.2	1,213.3
3	Chronic lower respiratory diseases (J40-J47)	50,650	7.1	343.0
4	Cerebrovascular diseases (I60-I69)	35,595	5.0	241.0
5	Alzheimer's disease (G30)	21,730	3.1	147.1
6	Diabetes mellitus (E10-E14)	19,167	2.7	129.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17,791	2.5	120.5
8	Influenza and pneumonia (J09-J18)	17,218	2.4	116.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16,163	2.3	109.4
10	Parkinson's disease (G20-G21)	11,101	1.6	75.2
11	Septicemia (A40-A41)	9,816	1.4	66.5
12	Pneumonitis due to solids and liquids (J69)	6,518	0.9	44.1
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	6,291	0.9	42.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5,453	0.8	36.9
15	Chronic liver disease and cirrhosis (K70,K73-K74)	5,138	0.7	34.8
...	All other causes (Residual)	110,035	15.5	745.1
White, female, all ages				
...	All causes	1,048,880	100.0	843.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	251,611	24.0	202.2
2	Malignant neoplasms (C00-C97)	231,953	22.1	186.4
3	Chronic lower respiratory diseases (J40-J47)	67,268	6.4	54.1
4	Cerebrovascular diseases (I60-I69)	65,574	6.3	52.7
5	Alzheimer's disease (G30)	50,630	4.8	40.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	37,669	3.6	30.3
7	Diabetes mellitus (E10-E14)	25,908	2.5	20.8
8	Influenza and pneumonia (J09-J18)	24,818	2.4	19.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19,446	1.9	15.6
10	Septicemia (A40-A41)	15,403	1.5	12.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	12,018	1.1	9.7
12	Chronic liver disease and cirrhosis (K70,K73-K74)	9,345	0.9	7.5
13	Parkinson's disease (G20-G21)	7,994	0.8	6.4
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,999	0.7	5.6
15	Pneumonitis due to solids and liquids (J69)	6,720	0.6	5.4
...	All other causes (Residual)	215,524	20.5	173.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 1-4 years				
...	All causes	1,361	100.0	21.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	384	28.2	6.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	181	13.3	2.8
3	Malignant neoplasms (C00-C97)	127	9.3	2.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	111	8.2	1.7
5	Influenza and pneumonia (J09-J18)	61	4.5	1.0
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	44	3.2	0.7
7	Septicemia (A40-A41)	18	1.3	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	1.3	*
7	Cerebrovascular diseases (I60-I69)	18	1.3	*
10	Certain conditions originating in the perinatal period (P00-P96)	15	1.1	*
11	Chronic lower respiratory diseases (J40-J47)	14	1.0	*
12	Meningitis (G00,G03)	6	0.4	*
13	Acute bronchitis and bronchiolitis (J20-J21)	5	0.4	*
13	Pneumonitis due to solids and liquids (J69)	5	0.4	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.4	*
...	All other causes (Residual)	349	25.6	5.5
White, female, 5-14 years				
...	All causes	1,691	100.0	11.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	458	27.1	3.0
2	Malignant neoplasms (C00-C97)	313	18.5	2.0
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	112	6.6	0.7
4	Influenza and pneumonia (J09-J18)	81	4.8	0.5
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	70	4.1	0.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	70	4.1	0.5
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	66	3.9	0.4
8	Septicemia (A40-A41)	29	1.7	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	27	1.6	0.2
10	Cerebrovascular diseases (I60-I69)	21	1.2	0.1
11	Chronic lower respiratory diseases (J40-J47)	12	0.7	*
11	Certain conditions originating in the perinatal period (P00-P96)	12	0.7	*
13	Diabetes mellitus (E10-E14)	11	0.7	*
14	Anemias (D50-D64)	9	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.4	*
...	All other causes (Residual)	393	23.2	2.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 15-24 years				
...	All causes	5,998	100.0	37.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,664	44.4	16.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	607	10.1	3.7
3	Malignant neoplasms (C00-C97)	480	8.0	3.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	339	5.7	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	214	3.6	1.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	146	2.4	0.9
7	Influenza and pneumonia (J09-J18)	139	2.3	0.9
8	Pregnancy, childbirth and the puerperium (O00-O99)	133	2.2	0.8
9	Cerebrovascular diseases (I60-I69)	58	1.0	0.4
10	Diabetes mellitus (E10-E14)	51	0.9	0.3
11	Septicemia (A40-A41)	42	0.7	0.3
11	Chronic lower respiratory diseases (J40-J47)	42	0.7	0.3
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21	0.4	0.1
14	Human immunodeficiency virus (HIV) disease (B20-B24)	18	0.3	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	0.3	*
...	All other causes (Residual)	1,027	17.1	6.3
White, female, 25-44 years				
...	All causes	30,419	100.0	95.3
1	Malignant neoplasms (C00-C97)	6,831	22.5	21.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,762	22.2	21.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,822	9.3	8.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,243	7.4	7.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	850	2.8	2.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	800	2.6	2.5
7	Cerebrovascular diseases (I60-I69)	716	2.4	2.2
8	Influenza and pneumonia (J09-J18)	695	2.3	2.2
9	Diabetes mellitus (E10-E14)	607	2.0	1.9
10	Septicemia (A40-A41)	426	1.4	1.3
11	Pregnancy, childbirth and the puerperium (O00-O99)	399	1.3	1.3
12	Chronic lower respiratory diseases (J40-J47)	393	1.3	1.2
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	289	1.0	0.9
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	265	0.9	0.8
15	Human immunodeficiency virus (HIV) disease (B20-B24)	254	0.8	0.8
...	All other causes (Residual)	6,067	19.9	19.0

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 45-64 years				
...	All causes	150,041	100.0	450.3
1	Malignant neoplasms (C00-C97)	59,142	39.4	177.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	22,366	14.9	67.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,954	6.0	26.9
4	Chronic lower respiratory diseases (J40-J47)	8,080	5.4	24.3
5	Cerebrovascular diseases (I60-I69)	4,959	3.3	14.9
6	Diabetes mellitus (E10-E14)	4,702	3.1	14.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	4,493	3.0	13.5
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,229	2.2	9.7
9	Influenza and pneumonia (J09-J18)	2,510	1.7	7.5
10	Septicemia (A40-A41)	2,395	1.6	7.2
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,919	1.3	5.8
12	Viral hepatitis (B15-B19)	1,291	0.9	3.9
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	840	0.6	2.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	779	0.5	2.3
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	596	0.4	1.8
...	All other causes (Residual)	23,786	15.9	71.4
White, female, 65 years and over				
...	All causes	851,906	100.0	4,327.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	225,979	26.5	1,147.9
2	Malignant neoplasms (C00-C97)	165,032	19.4	838.3
3	Cerebrovascular diseases (I60-I69)	59,758	7.0	303.6
4	Chronic lower respiratory diseases (J40-J47)	58,721	6.9	298.3
5	Alzheimer's disease (G30)	50,216	5.9	255.1
6	Influenza and pneumonia (J09-J18)	21,277	2.5	108.1
7	Diabetes mellitus (E10-E14)	20,537	2.4	104.3
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18,138	2.1	92.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17,198	2.0	87.4
10	Septicemia (A40-A41)	12,439	1.5	63.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	11,090	1.3	56.3
12	Parkinson's disease (G20-G21)	7,826	0.9	39.8
13	Pneumonitis due to solids and liquids (J69)	6,159	0.7	31.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5,180	0.6	26.3
15	Chronic liver disease and cirrhosis (K70,K73-K74)	4,045	0.5	20.5
...	All other causes (Residual)	168,311	19.8	855.0

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, all ages				
...	All causes	286,623	100.0	699.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	69,687	24.3	170.0
2	Malignant neoplasms (C00-C97)	64,649	22.6	157.7
3	Cerebrovascular diseases (I60-I69)	15,878	5.5	38.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,074	4.2	29.4
5	Diabetes mellitus (E10-E14)	11,960	4.2	29.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,808	3.1	21.5
7	Chronic lower respiratory diseases (J40-J47)	8,530	3.0	20.8
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7,874	2.7	19.2
9	Septicemia (A40-A41)	6,179	2.2	15.1
10	Human immunodeficiency virus (HIV) disease (B20-B24)	5,377	1.9	13.1
11	Influenza and pneumonia (J09-J18)	5,290	1.8	12.9
12	Alzheimer's disease (G30)	5,090	1.8	12.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,868	1.7	11.9
14	Certain conditions originating in the perinatal period (P00-P96)	4,654	1.6	11.4
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,560	0.9	6.2
...	All other causes (Residual)	53,145	18.5	129.6
Black, both sexes, 1-4 years				
...	All causes	1,077	100.0	38.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	301	27.9	10.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	149	13.8	5.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	92	8.5	3.3
4	Malignant neoplasms (C00-C97)	59	5.5	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	50	4.6	1.8
6	Chronic lower respiratory diseases (J40-J47)	33	3.1	1.2
7	Influenza and pneumonia (J09-J18)	30	2.8	1.1
8	Certain conditions originating in the perinatal period (P00-P96)	20	1.9	0.7
9	Septicemia (A40-A41)	19	1.8	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	1.2	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	12	1.1	*
12	Cerebrovascular diseases (I60-I69)	9	0.8	*
12	Acute bronchitis and bronchiolitis (J20-J21)	9	0.8	*
14	Anemias (D50-D64)	8	0.7	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.6	*
...	All other causes (Residual)	267	24.8	9.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 5-14 years				
...	All causes	1,300	100.0	20.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	385	29.6	6.0
2	Malignant neoplasms (C00-C97)	133	10.2	2.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	108	8.3	1.7
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	81	6.2	1.3
5	Chronic lower respiratory diseases (J40-J47)	75	5.8	1.2
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	64	4.9	1.0
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	49	3.8	0.8
8	Influenza and pneumonia (J09-J18)	46	3.5	0.7
9	Septicemia (A40-A41)	22	1.7	0.3
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	1.5	0.3
11	Anemias (D50-D64)	19	1.5	*
12	Cerebrovascular diseases (I60-I69)	18	1.4	*
13	Diabetes mellitus (E10-E14)	8	0.6	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.5	*
15	Certain conditions originating in the perinatal period (P00-P96)	5	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.4	*
...	All other causes (Residual)	256	19.7	4.0
Black, both sexes, 15-24 years				
...	All causes	6,742	100.0	96.7
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,810	41.7	40.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,349	20.0	19.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	448	6.6	6.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	345	5.1	4.9
5	Malignant neoplasms (C00-C97)	291	4.3	4.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	106	1.6	1.5
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	91	1.3	1.3
8	Pregnancy, childbirth and the puerperium (O00-O99)	88	1.3	1.3
9	Influenza and pneumonia (J09-J18)	87	1.3	1.2
9	Chronic lower respiratory diseases (J40-J47)	87	1.3	1.2
11	Anemias (D50-D64)	66	1.0	0.9
12	Diabetes mellitus (E10-E14)	50	0.7	0.7
13	Cerebrovascular diseases (I60-I69)	41	0.6	0.6
14	Septicemia (A40-A41)	31	0.5	0.4
15	Legal intervention (Y35,Y89.0)	30	0.4	0.4
...	All other causes (Residual)	822	12.2	11.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 25-44 years				
...	All causes	24,349	100.0	211.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,852	15.8	33.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,525	14.5	30.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,384	13.9	29.4
4	Malignant neoplasms (C00-C97)	2,796	11.5	24.3
5	Human immunodeficiency virus (HIV) disease (B20-B24)	1,995	8.2	17.3
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	878	3.6	7.6
7	Diabetes mellitus (E10-E14)	721	3.0	6.3
8	Cerebrovascular diseases (I60-I69)	695	2.9	6.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	414	1.7	3.6
10	Influenza and pneumonia (J09-J18)	399	1.6	3.5
11	Septicemia (A40-A41)	383	1.6	3.3
12	Chronic lower respiratory diseases (J40-J47)	285	1.2	2.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	265	1.1	2.3
14	Anemias (D50-D64)	247	1.0	2.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	227	0.9	2.0
...	All other causes (Residual)	4,283	17.6	37.2
Black, both sexes, 45-64 years				
...	All causes	86,123	100.0	939.0
1	Malignant neoplasms (C00-C97)	24,313	28.2	265.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	21,136	24.5	230.5
3	Cerebrovascular diseases (I60-I69)	4,403	5.1	48.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,070	4.7	44.4
5	Diabetes mellitus (E10-E14)	3,801	4.4	41.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	2,914	3.4	31.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,295	2.7	25.0
8	Chronic lower respiratory diseases (J40-J47)	2,086	2.4	22.7
9	Chronic liver disease and cirrhosis (K70,K73-K74)	1,751	2.0	19.1
10	Septicemia (A40-A41)	1,716	2.0	18.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,390	1.6	15.2
12	Influenza and pneumonia (J09-J18)	1,170	1.4	12.8
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	999	1.2	10.9
14	Viral hepatitis (B15-B19)	857	1.0	9.3
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	553	0.6	6.0
...	All other causes (Residual)	12,669	14.7	138.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 65 years and over				
...	All causes	158,690	100.0	4,617.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	44,114	27.8	1,283.7
2	Malignant neoplasms (C00-C97)	37,040	23.3	1,077.9
3	Cerebrovascular diseases (I60-I69)	10,666	6.7	310.4
4	Diabetes mellitus (E10-E14)	7,379	4.6	214.7
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,026	3.8	175.4
6	Chronic lower respiratory diseases (J40-J47)	5,955	3.8	173.3
7	Alzheimer's disease (G30)	5,009	3.2	145.8
8	Septicemia (A40-A41)	3,915	2.5	113.9
9	Influenza and pneumonia (J09-J18)	3,476	2.2	101.2
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,204	2.0	93.2
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,178	1.4	63.4
12	Pneumonitis due to solids and liquids (J69)	1,124	0.7	32.7
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	767	0.5	22.3
14	Parkinson's disease (G20-G21)	698	0.4	20.3
15	Chronic liver disease and cirrhosis (K70,K73-K74)	577	0.4	16.8
...	All other causes (Residual)	26,562	16.7	773.0
Black, male, all ages				
...	All causes	146,239	100.0	745.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	35,420	24.2	180.6
2	Malignant neoplasms (C00-C97)	33,443	22.9	170.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,083	5.5	41.2
4	Cerebrovascular diseases (I60-I69)	6,962	4.8	35.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6,715	4.6	34.2
6	Diabetes mellitus (E10-E14)	5,488	3.8	28.0
7	Chronic lower respiratory diseases (J40-J47)	4,503	3.1	23.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,974	2.7	20.3
9	Human immunodeficiency virus (HIV) disease (B20-B24)	3,535	2.4	18.0
10	Septicemia (A40-A41)	2,745	1.9	14.0
11	Certain conditions originating in the perinatal period (P00-P96)	2,635	1.8	13.4
12	Influenza and pneumonia (J09-J18)	2,512	1.7	12.8
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,066	1.4	10.5
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,684	1.2	8.6
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,660	1.1	8.5
...	All other causes (Residual)	24,814	17.0	126.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 1-4 years				
...	All causes	576	100.0	40.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	175	30.4	12.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	80	13.9	5.7
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	50	8.7	3.5
4	Malignant neoplasms (C00-C97)	22	3.8	1.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	22	3.8	1.6
6	Chronic lower respiratory diseases (J40-J47)	18	3.1	*
7	Influenza and pneumonia (J09-J18)	14	2.4	*
8	Certain conditions originating in the perinatal period (P00-P96)	11	1.9	*
9	Complications of medical and surgical care (Y40-Y84,Y88)	10	1.7	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	1.6	*
11	Septicemia (A40-A41)	7	1.2	*
12	Anemias (D50-D64)	6	1.0	*
13	Cerebrovascular diseases (I60-I69)	4	0.7	*
14	Acute bronchitis and bronchiolitis (J20-J21)	3	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.5	*
...	All other causes (Residual)	142	24.7	10.0
Black, male, 5-14 years				
...	All causes	755	100.0	23.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	241	31.9	7.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	74	9.8	2.3
3	Malignant neoplasms (C00-C97)	64	8.5	2.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	48	6.4	1.5
5	Chronic lower respiratory diseases (J40-J47)	47	6.2	1.4
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	37	4.9	1.1
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	35	4.6	1.1
8	Influenza and pneumonia (J09-J18)	22	2.9	0.7
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	1.7	*
10	Cerebrovascular diseases (I60-I69)	11	1.5	*
11	Septicemia (A40-A41)	10	1.3	*
12	Anemias (D50-D64)	7	0.9	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.5	*
14	Diseases of appendix (K35-K38)	3	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.4	*
14	Certain conditions originating in the perinatal period (P00-P96)	3	0.4	*
...	All other causes (Residual)	133	17.6	4.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 15-24 years				
...	All causes	5,090	100.0	144.0
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,526	49.6	71.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	968	19.0	27.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	372	7.3	10.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	226	4.4	6.4
5	Malignant neoplasms (C00-C97)	162	3.2	4.6
6	Human immunodeficiency virus (HIV) disease (B20-B24)	65	1.3	1.8
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	56	1.1	1.6
8	Chronic lower respiratory diseases (J40-J47)	55	1.1	1.6
9	Influenza and pneumonia (J09-J18)	40	0.8	1.1
10	Anemias (D50-D64)	36	0.7	1.0
11	Legal intervention (Y35,Y89.0)	29	0.6	0.8
12	Diabetes mellitus (E10-E14)	25	0.5	0.7
13	Cerebrovascular diseases (I60-I69)	22	0.4	0.6
14	Septicemia (A40-A41)	12	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	0.2	*
...	All other causes (Residual)	484	9.5	13.7
Black, male, 25-44 years				
...	All causes	15,190	100.0	274.5
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,056	20.1	55.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,490	16.4	45.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,439	16.1	44.1
4	Human immunodeficiency virus (HIV) disease (B20-B24)	1,209	8.0	21.8
5	Malignant neoplasms (C00-C97)	1,119	7.4	20.2
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	715	4.7	12.9
7	Diabetes mellitus (E10-E14)	443	2.9	8.0
8	Cerebrovascular diseases (I60-I69)	369	2.4	6.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	217	1.4	3.9
10	Septicemia (A40-A41)	183	1.2	3.3
11	Influenza and pneumonia (J09-J18)	181	1.2	3.3
12	Chronic lower respiratory diseases (J40-J47)	162	1.1	2.9
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	144	0.9	2.6
14	Chronic liver disease and cirrhosis (K70,K73-K74)	136	0.9	2.5
15	Anemias (D50-D64)	125	0.8	2.3
...	All other causes (Residual)	2,202	14.5	39.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 45-64 years				
...	All causes	50,252	100.0	1,202.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,309	26.5	318.6
2	Malignant neoplasms (C00-C97)	13,008	25.9	311.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,796	5.6	66.9
4	Cerebrovascular diseases (I60-I69)	2,404	4.8	57.5
5	Diabetes mellitus (E10-E14)	2,092	4.2	50.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	2,004	4.0	48.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,226	2.4	29.3
8	Chronic liver disease and cirrhosis (K70,K73-K74)	1,184	2.4	28.3
9	Chronic lower respiratory diseases (J40-J47)	1,094	2.2	26.2
10	Septicemia (A40-A41)	892	1.8	21.4
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	795	1.6	19.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	794	1.6	19.0
13	Influenza and pneumonia (J09-J18)	638	1.3	15.3
14	Viral hepatitis (B15-B19)	592	1.2	14.2
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	425	0.8	10.2
...	All other causes (Residual)	6,999	13.9	167.5
Black, male, 65 years and over				
...	All causes	69,650	100.0	5,243.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	19,311	27.7	1,453.8
2	Malignant neoplasms (C00-C97)	19,059	27.4	1,434.8
3	Cerebrovascular diseases (I60-I69)	4,123	5.9	310.4
4	Chronic lower respiratory diseases (J40-J47)	3,122	4.5	235.0
5	Diabetes mellitus (E10-E14)	2,925	4.2	220.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,493	3.6	187.7
7	Septicemia (A40-A41)	1,592	2.3	119.8
8	Influenza and pneumonia (J09-J18)	1,570	2.3	118.2
9	Alzheimer's disease (G30)	1,405	2.0	105.8
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,176	1.7	88.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,125	1.6	84.7
12	Pneumonitis due to solids and liquids (J69)	513	0.7	38.6
13	Parkinson's disease (G20-G21)	395	0.6	29.7
14	Chronic liver disease and cirrhosis (K70,K73-K74)	335	0.5	25.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	320	0.5	24.1
...	All other causes (Residual)	10,186	14.6	766.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, all ages				
...	All causes	140,384	100.0	656.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	34,267	24.4	160.2
2	Malignant neoplasms (C00-C97)	31,206	22.2	145.9
3	Cerebrovascular diseases (I60-I69)	8,916	6.4	41.7
4	Diabetes mellitus (E10-E14)	6,472	4.6	30.3
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,834	3.4	22.6
6	Chronic lower respiratory diseases (J40-J47)	4,027	2.9	18.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,991	2.8	18.7
8	Alzheimer's disease (G30)	3,639	2.6	17.0
9	Septicemia (A40-A41)	3,434	2.4	16.1
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,802	2.0	13.1
11	Influenza and pneumonia (J09-J18)	2,778	2.0	13.0
12	Certain conditions originating in the perinatal period (P00-P96)	2,019	1.4	9.4
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,842	1.3	8.6
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,159	0.8	5.4
15	Chronic liver disease and cirrhosis (K70,K73-K74)	900	0.6	4.2
...	All other causes (Residual)	28,098	20.0	131.4
Black, female, 1-4 years				
...	All causes	501	100.0	36.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	126	25.1	9.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	69	13.8	5.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	42	8.4	3.1
4	Malignant neoplasms (C00-C97)	37	7.4	2.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	28	5.6	2.1
6	Influenza and pneumonia (J09-J18)	16	3.2	*
7	Chronic lower respiratory diseases (J40-J47)	15	3.0	*
8	Septicemia (A40-A41)	12	2.4	*
9	Certain conditions originating in the perinatal period (P00-P96)	9	1.8	*
10	Acute bronchitis and bronchiolitis (J20-J21)	6	1.2	*
11	Cerebrovascular diseases (I60-I69)	5	1.0	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.8	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.6	*
14	Anemias (D50-D64)	2	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
...	All other causes (Residual)	125	25.0	9.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 5-14 years				
...	All causes	545	100.0	17.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	144	26.4	4.6
2	Malignant neoplasms (C00-C97)	69	12.7	2.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	34	6.2	1.1
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	33	6.1	1.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	29	5.3	0.9
6	Chronic lower respiratory diseases (J40-J47)	28	5.1	0.9
7	Influenza and pneumonia (J09-J18)	24	4.4	0.8
8	Septicemia (A40-A41)	12	2.2	*
8	Anemias (D50-D64)	12	2.2	*
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	12	2.2	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	1.3	*
11	Cerebrovascular diseases (I60-I69)	7	1.3	*
13	Diabetes mellitus (E10-E14)	6	1.1	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.6	*
14	Pregnancy, childbirth and the puerperium (O00-O99)	3	0.6	*
...	All other causes (Residual)	122	22.4	3.9
Black, female, 15-24 years				
...	All causes	1,652	100.0	48.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	381	23.1	11.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	284	17.2	8.3
3	Malignant neoplasms (C00-C97)	129	7.8	3.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	119	7.2	3.5
5	Pregnancy, childbirth and the puerperium (O00-O99)	88	5.3	2.6
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	76	4.6	2.2
7	Influenza and pneumonia (J09-J18)	47	2.8	1.4
8	Human immunodeficiency virus (HIV) disease (B20-B24)	41	2.5	1.2
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	35	2.1	1.0
10	Chronic lower respiratory diseases (J40-J47)	32	1.9	0.9
11	Anemias (D50-D64)	30	1.8	0.9
12	Diabetes mellitus (E10-E14)	25	1.5	0.7
13	Septicemia (A40-A41)	19	1.2	*
13	Cerebrovascular diseases (I60-I69)	19	1.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	1.0	*
...	All other causes (Residual)	310	18.8	9.0

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 25-44 years				
...	All causes	9,159	100.0	153.4
1	Malignant neoplasms (C00-C97)	1,677	18.3	28.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,413	15.4	23.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	894	9.8	15.0
4	Human immunodeficiency virus (HIV) disease (B20-B24)	786	8.6	13.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	469	5.1	7.9
6	Cerebrovascular diseases (I60-I69)	326	3.6	5.5
7	Diabetes mellitus (E10-E14)	278	3.0	4.7
8	Influenza and pneumonia (J09-J18)	218	2.4	3.7
9	Septicemia (A40-A41)	200	2.2	3.4
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	197	2.2	3.3
11	Pregnancy, childbirth and the puerperium (O00-O99)	195	2.1	3.3
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	163	1.8	2.7
13	Chronic lower respiratory diseases (J40-J47)	123	1.3	2.1
14	Anemias (D50-D64)	122	1.3	2.0
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	121	1.3	2.0
...	All other causes (Residual)	1,977	21.6	33.1
Black, female, 45-64 years				
...	All causes	35,871	100.0	718.3
1	Malignant neoplasms (C00-C97)	11,305	31.5	226.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,827	21.8	156.7
3	Cerebrovascular diseases (I60-I69)	1,999	5.6	40.0
4	Diabetes mellitus (E10-E14)	1,709	4.8	34.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,274	3.6	25.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,069	3.0	21.4
7	Chronic lower respiratory diseases (J40-J47)	992	2.8	19.9
8	Human immunodeficiency virus (HIV) disease (B20-B24)	910	2.5	18.2
9	Septicemia (A40-A41)	824	2.3	16.5
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	596	1.7	11.9
11	Chronic liver disease and cirrhosis (K70,K73-K74)	567	1.6	11.4
12	Influenza and pneumonia (J09-J18)	532	1.5	10.7
13	Viral hepatitis (B15-B19)	265	0.7	5.3
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	204	0.6	4.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	173	0.5	3.5
...	All other causes (Residual)	5,625	15.7	112.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 65 years and over				
...	All causes	89,040	100.0	4,223.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	24,803	27.9	1,176.6
2	Malignant neoplasms (C00-C97)	17,981	20.2	853.0
3	Cerebrovascular diseases (I60-I69)	6,543	7.3	310.4
4	Diabetes mellitus (E10-E14)	4,454	5.0	211.3
5	Alzheimer's disease (G30)	3,604	4.0	171.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,533	4.0	167.6
7	Chronic lower respiratory diseases (J40-J47)	2,833	3.2	134.4
8	Septicemia (A40-A41)	2,323	2.6	110.2
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,079	2.3	98.6
10	Influenza and pneumonia (J09-J18)	1,906	2.1	90.4
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,002	1.1	47.5
12	Pneumonitis due to solids and liquids (J69)	611	0.7	29.0
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	447	0.5	21.2
14	Atherosclerosis (I70)	311	0.3	14.8
15	Parkinson's disease (G20-G21)	303	0.3	14.4
...	All other causes (Residual)	16,307	18.3	773.6
American Indian, both sexes, all ages				
...	All causes	14,960	100.0	427.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,764	18.5	79.0
2	Malignant neoplasms (C00-C97)	2,710	18.1	77.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,733	11.6	49.5
4	Diabetes mellitus (E10-E14)	786	5.3	22.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	714	4.8	20.4
6	Chronic lower respiratory diseases (J40-J47)	610	4.1	17.4
7	Cerebrovascular diseases (I60-I69)	561	3.8	16.0
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	429	2.9	12.3
9	Influenza and pneumonia (J09-J18)	402	2.7	11.5
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	307	2.1	8.8
11	Septicemia (A40-A41)	275	1.8	7.9
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	254	1.7	7.3
13	Alzheimer's disease (G30)	193	1.3	5.5
14	Certain conditions originating in the perinatal period (P00-P96)	148	1.0	4.2
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	125	0.8	3.6
...	All other causes (Residual)	2,949	19.7	84.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 1-4 years				
...	All causes	83	100.0	30.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	34	41.0	12.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	7.2	*
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	7.2	*
4	Influenza and pneumonia (J09-J18)	5	6.0	*
5	Malignant neoplasms (C00-C97)	4	4.8	*
6	Certain conditions originating in the perinatal period (P00-P96)	2	2.4	*
7	Meningococcal infection (A39)	1	1.2	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.2	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	1.2	*
...	All other causes (Residual)	23	27.7	8.5
American Indian, both sexes, 5-14 years				
...	All causes	115	100.0	20.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	38	33.0	6.7
2	Malignant neoplasms (C00-C97)	13	11.3	*
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	9	7.8	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	6	5.2	*
4	Influenza and pneumonia (J09-J18)	6	5.2	*
6	Septicemia (A40-A41)	5	4.3	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	3.5	*
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	3.5	*
9	Chronic lower respiratory diseases (J40-J47)	3	2.6	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.9	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.9	*
...	All other causes (Residual)	25	21.7	4.4

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 15-24 years				
...	All causes	670	100.0	109.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	327	48.8	53.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	152	22.7	24.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	78	11.6	12.8
4	Malignant neoplasms (C00-C97)	14	2.1	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	10	1.5	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	1.3	*
7	Influenza and pneumonia (J09-J18)	8	1.2	*
8	Chronic liver disease and cirrhosis (K70,K73-K74)	4	0.6	*
9	Cerebrovascular diseases (I60-I69)	3	0.4	*
9	Chronic lower respiratory diseases (J40-J47)	3	0.4	*
11	Diabetes mellitus (E10-E14)	2	0.3	*
12	Septicemia (A40-A41)	1	0.1	*
12	Anemias (D50-D64)	1	0.1	*
12	Pneumonitis due to solids and liquids (J69)	1	0.1	*
12	Pregnancy, childbirth and the puerperium (O00-O99)	1	0.1	*
12	Legal intervention (Y35,Y89.0)	1	0.1	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.1	*
...	All other causes (Residual)	54	8.1	8.9
American Indian, both sexes, 25-44 years				
...	All causes	2,013	100.0	205.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	639	31.7	65.1
2	Chronic liver disease and cirrhosis (K70,K73-K74)	195	9.7	19.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	180	8.9	18.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	168	8.3	17.1
5	Malignant neoplasms (C00-C97)	133	6.6	13.5
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	109	5.4	11.1
7	Diabetes mellitus (E10-E14)	62	3.1	6.3
8	Influenza and pneumonia (J09-J18)	54	2.7	5.5
9	Septicemia (A40-A41)	33	1.6	3.4
10	Human immunodeficiency virus (HIV) disease (B20-B24)	22	1.1	2.2
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22	1.1	2.2
12	Viral hepatitis (B15-B19)	20	1.0	2.0
13	Cerebrovascular diseases (I60-I69)	16	0.8	*
14	Pregnancy, childbirth and the puerperium (O00-O99)	11	0.5	*
15	Chronic lower respiratory diseases (J40-J47)	10	0.5	*
...	All other causes (Residual)	339	16.8	34.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 45-64 years				
...	All causes	4,699	100.0	624.8
1	Malignant neoplasms (C00-C97)	1,053	22.4	140.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	874	18.6	116.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	474	10.1	63.0
4	Chronic liver disease and cirrhosis (K70,K73-K74)	408	8.7	54.2
5	Diabetes mellitus (E10-E14)	263	5.6	35.0
6	Cerebrovascular diseases (I60-I69)	150	3.2	19.9
7	Chronic lower respiratory diseases (J40-J47)	148	3.1	19.7
8	Influenza and pneumonia (J09-J18)	113	2.4	15.0
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	90	1.9	12.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	89	1.9	11.8
11	Viral hepatitis (B15-B19)	76	1.6	10.1
12	Septicemia (A40-A41)	65	1.4	8.6
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	44	0.9	5.9
14	Human immunodeficiency virus (HIV) disease (B20-B24)	40	0.9	5.3
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	35	0.7	4.7
...	All other causes (Residual)	777	16.5	103.3
American Indian, both sexes, 65 years and over				
...	All causes	6,994	100.0	2,766.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,688	24.1	667.7
2	Malignant neoplasms (C00-C97)	1,492	21.3	590.2
3	Diabetes mellitus (E10-E14)	459	6.6	181.6
4	Chronic lower respiratory diseases (J40-J47)	445	6.4	176.0
5	Cerebrovascular diseases (I60-I69)	392	5.6	155.1
6	Influenza and pneumonia (J09-J18)	208	3.0	82.3
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	200	2.9	79.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	195	2.8	77.1
9	Alzheimer's disease (G30)	190	2.7	75.2
10	Septicemia (A40-A41)	170	2.4	67.2
11	Chronic liver disease and cirrhosis (K70,K73-K74)	107	1.5	42.3
12	Pneumonitis due to solids and liquids (J69)	67	1.0	26.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	63	0.9	24.9
14	Parkinson's disease (G20-G21)	50	0.7	19.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	42	0.6	16.6
...	All other causes (Residual)	1,226	17.5	485.0

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, all ages				
...	All causes	8,105	100.0	462.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,591	19.6	90.9
2	Malignant neoplasms (C00-C97)	1,381	17.0	78.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,150	14.2	65.7
4	Diabetes mellitus (E10-E14)	418	5.2	23.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	380	4.7	21.7
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	313	3.9	17.9
7	Chronic lower respiratory diseases (J40-J47)	287	3.5	16.4
8	Cerebrovascular diseases (I60-I69)	263	3.2	15.0
9	Influenza and pneumonia (J09-J18)	193	2.4	11.0
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	193	2.4	11.0
11	Septicemia (A40-A41)	127	1.6	7.3
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	126	1.6	7.2
13	Certain conditions originating in the perinatal period (P00-P96)	92	1.1	5.3
14	Alzheimer's disease (G30)	65	0.8	3.7
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	62	0.8	3.5
...	All other causes (Residual)	1,464	18.1	83.6
American Indian, male, 1-4 years				
...	All causes	48	100.0	35.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	24	50.0	17.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	6.3	*
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	6.3	*
4	Malignant neoplasms (C00-C97)	2	4.2	*
4	Influenza and pneumonia (J09-J18)	2	4.2	*
6	Meningococcal infection (A39)	1	2.1	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	2.1	*
6	Certain conditions originating in the perinatal period (P00-P96)	1	2.1	*
...	All other causes (Residual)	11	22.9	*

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 5-14 years				
...	All causes	57	100.0	19.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	21	36.8	7.3
2	Malignant neoplasms (C00-C97)	6	10.5	*
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	8.8	*
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5	8.8	*
5	Influenza and pneumonia (J09-J18)	4	7.0	*
6	Septicemia (A40-A41)	3	5.3	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	3.5	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.8	*
8	Chronic lower respiratory diseases (J40-J47)	1	1.8	*
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1	1.8	*
...	All other causes (Residual)	8	14.0	*
American Indian, male, 15-24 years				
...	All causes	468	100.0	150.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	224	47.9	72.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	112	23.9	36.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	61	13.0	19.6
4	Malignant neoplasms (C00-C97)	10	2.1	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	6	1.3	*
5	Influenza and pneumonia (J09-J18)	6	1.3	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	1.3	*
8	Chronic liver disease and cirrhosis (K70,K73-K74)	4	0.9	*
9	Anemias (D50-D64)	1	0.2	*
9	Diabetes mellitus (E10-E14)	1	0.2	*
9	Cerebrovascular diseases (I60-I69)	1	0.2	*
9	Chronic lower respiratory diseases (J40-J47)	1	0.2	*
9	Pneumonitis due to solids and liquids (J69)	1	0.2	*
9	Legal intervention (Y35,Y89.0)	1	0.2	*
9	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
...	All other causes (Residual)	32	6.8	10.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 25-44 years				
...	All causes	1,284	100.0	254.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	434	33.8	85.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	128	10.0	25.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	124	9.7	24.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	112	8.7	22.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	91	7.1	18.0
6	Malignant neoplasms (C00-C97)	56	4.4	11.1
7	Diabetes mellitus (E10-E14)	44	3.4	8.7
8	Influenza and pneumonia (J09-J18)	31	2.4	6.1
9	Septicemia (A40-A41)	17	1.3	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	16	1.2	*
11	Cerebrovascular diseases (I60-I69)	11	0.9	*
12	Viral hepatitis (B15-B19)	9	0.7	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.6	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4	0.3	*
14	Chronic lower respiratory diseases (J40-J47)	4	0.3	*
14	Legal intervention (Y35,Y89.0)	4	0.3	*
...	All other causes (Residual)	191	14.9	37.8
American Indian, male, 45-64 years				
...	All causes	2,746	100.0	754.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	588	21.4	161.5
2	Malignant neoplasms (C00-C97)	553	20.1	151.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	328	11.9	90.1
4	Chronic liver disease and cirrhosis (K70,K73-K74)	222	8.1	61.0
5	Diabetes mellitus (E10-E14)	167	6.1	45.9
6	Cerebrovascular diseases (I60-I69)	77	2.8	21.2
7	Chronic lower respiratory diseases (J40-J47)	72	2.6	19.8
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	64	2.3	17.6
9	Influenza and pneumonia (J09-J18)	54	2.0	14.8
10	Viral hepatitis (B15-B19)	46	1.7	12.6
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	43	1.6	11.8
12	Septicemia (A40-A41)	36	1.3	9.9
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	31	1.1	8.5
14	Human immunodeficiency virus (HIV) disease (B20-B24)	28	1.0	7.7
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	21	0.8	5.8
...	All other causes (Residual)	416	15.1	114.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 65 years and over				
...	All causes	3,287	100.0	2,949.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	862	26.2	773.4
2	Malignant neoplasms (C00-C97)	754	22.9	676.5
3	Chronic lower respiratory diseases (J40-J47)	208	6.3	186.6
4	Diabetes mellitus (E10-E14)	206	6.3	184.8
5	Cerebrovascular diseases (I60-I69)	174	5.3	156.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	105	3.2	94.2
7	Influenza and pneumonia (J09-J18)	92	2.8	82.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	75	2.3	67.3
9	Septicemia (A40-A41)	71	2.2	63.7
10	Alzheimer's disease (G30)	64	1.9	57.4
11	Chronic liver disease and cirrhosis (K70,K73-K74)	42	1.3	37.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	36	1.1	32.3
13	Pneumonitis due to solids and liquids (J69)	32	1.0	28.7
14	Parkinson's disease (G20-G21)	24	0.7	21.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	21	0.6	18.8
...	All other causes (Residual)	521	15.9	467.5
American Indian, female, all ages				
...	All causes	6,855	100.0	391.9
1	Malignant neoplasms (C00-C97)	1,329	19.4	76.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,173	17.1	67.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	583	8.5	33.3
4	Diabetes mellitus (E10-E14)	368	5.4	21.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	334	4.9	19.1
6	Chronic lower respiratory diseases (J40-J47)	323	4.7	18.5
7	Cerebrovascular diseases (I60-I69)	298	4.3	17.0
8	Influenza and pneumonia (J09-J18)	209	3.0	11.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	181	2.6	10.3
10	Septicemia (A40-A41)	148	2.2	8.5
11	Alzheimer's disease (G30)	128	1.9	7.3
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	116	1.7	6.6
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	63	0.9	3.6
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	61	0.9	3.5
15	Certain conditions originating in the perinatal period (P00-P96)	56	0.8	3.2
...	All other causes (Residual)	1,485	21.7	84.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 1-4 years				
...	All causes	35	100.0	26.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10	28.6	*
2	Influenza and pneumonia (J09-J18)	3	8.6	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	8.6	*
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	8.6	*
5	Malignant neoplasms (C00-C97)	2	5.7	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	2.9	*
6	Certain conditions originating in the perinatal period (P00-P96)	1	2.9	*
...	All other causes (Residual)	12	34.3	*
American Indian, female, 5-14 years				
...	All causes	58	100.0	21.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17	29.3	*
2	Malignant neoplasms (C00-C97)	7	12.1	*
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4	6.9	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	5.2	*
5	Septicemia (A40-A41)	2	3.4	*
5	Influenza and pneumonia (J09-J18)	2	3.4	*
5	Chronic lower respiratory diseases (J40-J47)	2	3.4	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	3.4	*
9	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	1.7	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	1.7	*
...	All other causes (Residual)	17	29.3	*
American Indian, female, 15-24 years				
...	All causes	202	100.0	67.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	103	51.0	34.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	40	19.8	13.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	17	8.4	*
4	Malignant neoplasms (C00-C97)	4	2.0	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	2.0	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	1.5	*
7	Cerebrovascular diseases (I60-I69)	2	1.0	*
7	Influenza and pneumonia (J09-J18)	2	1.0	*
7	Chronic lower respiratory diseases (J40-J47)	2	1.0	*
10	Septicemia (A40-A41)	1	0.5	*
10	Diabetes mellitus (E10-E14)	1	0.5	*
10	Pregnancy, childbirth and the puerperium (O00-O99)	1	0.5	*
...	All other causes (Residual)	22	10.9	7.4

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 25-44 years				
...	All causes	729	100.0	152.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	205	28.1	43.0
2	Chronic liver disease and cirrhosis (K70,K73-K74)	83	11.4	17.4
3	Malignant neoplasms (C00-C97)	77	10.6	16.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	52	7.1	10.9
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	44	6.0	9.2
6	Influenza and pneumonia (J09-J18)	23	3.2	4.8
7	Diabetes mellitus (E10-E14)	18	2.5	*
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	18	2.5	*
9	Septicemia (A40-A41)	16	2.2	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	1.9	*
11	Viral hepatitis (B15-B19)	11	1.5	*
11	Pregnancy, childbirth and the puerperium (O00-O99)	11	1.5	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	6	0.8	*
13	Chronic lower respiratory diseases (J40-J47)	6	0.8	*
15	Cerebrovascular diseases (I60-I69)	5	0.7	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	0.7	*
...	All other causes (Residual)	135	18.5	28.3
American Indian, female, 45-64 years				
...	All causes	1,953	100.0	503.3
1	Malignant neoplasms (C00-C97)	500	25.6	128.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	286	14.6	73.7
3	Chronic liver disease and cirrhosis (K70,K73-K74)	186	9.5	47.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	146	7.5	37.6
5	Diabetes mellitus (E10-E14)	96	4.9	24.7
6	Chronic lower respiratory diseases (J40-J47)	76	3.9	19.6
7	Cerebrovascular diseases (I60-I69)	73	3.7	18.8
8	Influenza and pneumonia (J09-J18)	59	3.0	15.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	46	2.4	11.9
10	Viral hepatitis (B15-B19)	30	1.5	7.7
11	Septicemia (A40-A41)	29	1.5	7.5
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	26	1.3	6.7
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	14	0.7	*
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	13	0.7	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	12	0.6	*
...	All other causes (Residual)	361	18.5	93.0

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 65 years and over				
...	All causes	3,707	100.0	2,622.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	826	22.3	584.4
2	Malignant neoplasms (C00-C97)	738	19.9	522.1
3	Diabetes mellitus (E10-E14)	253	6.8	179.0
4	Chronic lower respiratory diseases (J40-J47)	237	6.4	167.7
5	Cerebrovascular diseases (I60-I69)	218	5.9	154.2
6	Alzheimer's disease (G30)	126	3.4	89.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	120	3.2	84.9
8	Influenza and pneumonia (J09-J18)	116	3.1	82.1
9	Septicemia (A40-A41)	99	2.7	70.0
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	95	2.6	67.2
11	Chronic liver disease and cirrhosis (K70,K73-K74)	65	1.8	46.0
12	Pneumonitis due to solids and liquids (J69)	35	0.9	24.8
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	27	0.7	19.1
14	Parkinson's disease (G20-G21)	26	0.7	18.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	21	0.6	14.9
...	All other causes (Residual)	705	19.0	498.8
Asian or Pacific Islander, both sexes, all ages				
...	All causes	49,225	100.0	317.0
1	Malignant neoplasms (C00-C97)	13,274	27.0	85.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,375	23.1	73.3
3	Cerebrovascular diseases (I60-I69)	3,639	7.4	23.4
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,044	4.2	13.2
5	Diabetes mellitus (E10-E14)	1,846	3.8	11.9
6	Influenza and pneumonia (J09-J18)	1,568	3.2	10.1
7	Chronic lower respiratory diseases (J40-J47)	1,480	3.0	9.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,062	2.2	6.8
9	Alzheimer's disease (G30)	1,024	2.1	6.6
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	971	2.0	6.3
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	710	1.4	4.6
12	Septicemia (A40-A41)	576	1.2	3.7
13	Chronic liver disease and cirrhosis (K70,K73-K74)	464	0.9	3.0
14	Certain conditions originating in the perinatal period (P00-P96)	455	0.9	2.9
15	Parkinson's disease (G20-G21)	367	0.7	2.4
...	All other causes (Residual)	8,370	17.0	53.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 1-4 years				
...	All causes	149	100.0	15.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	49	32.9	5.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	16	10.7	*
3	Malignant neoplasms (C00-C97)	13	8.7	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	10	6.7	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	7	4.7	*
6	Septicemia (A40-A41)	5	3.4	*
7	Influenza and pneumonia (J09-J18)	3	2.0	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	1.3	*
9	Nutritional deficiencies (E40-E64)	1	0.7	*
9	Meningitis (G00,G03)	1	0.7	*
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.7	*
9	Acute bronchitis and bronchiolitis (J20-J21)	1	0.7	*
9	Chronic lower respiratory diseases (J40-J47)	1	0.7	*
9	Hernia (K40-K46)	1	0.7	*
9	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.7	*
...	All other causes (Residual)	37	24.8	3.9
Asian or Pacific Islander, both sexes, 5-14 years				
...	All causes	231	100.0	11.1
1	Malignant neoplasms (C00-C97)	47	20.3	2.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	47	20.3	2.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	20	8.7	1.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	15	6.5	*
5	Influenza and pneumonia (J09-J18)	11	4.8	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8	3.5	*
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6	2.6	*
8	Chronic lower respiratory diseases (J40-J47)	5	2.2	*
9	Meningitis (G00,G03)	3	1.3	*
9	Cerebrovascular diseases (I60-I69)	3	1.3	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.4	*
11	Diseases of appendix (K35-K38)	1	0.4	*
11	Hernia (K40-K46)	1	0.4	*
11	Certain conditions originating in the perinatal period (P00-P96)	1	0.4	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.4	*
...	All other causes (Residual)	61	26.4	2.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 15-24 years				
...	All causes	763	100.0	37.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	256	33.6	12.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	152	19.9	7.5
3	Malignant neoplasms (C00-C97)	75	9.8	3.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	72	9.4	3.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	37	4.8	1.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	17	2.2	*
7	Influenza and pneumonia (J09-J18)	15	2.0	*
8	Cerebrovascular diseases (I60-I69)	12	1.6	*
9	Chronic lower respiratory diseases (J40-J47)	5	0.7	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.7	*
9	Pregnancy, childbirth and the puerperium (O00-O99)	5	0.7	*
12	Septicemia (A40-A41)	4	0.5	*
13	Tuberculosis (A16-A19)	3	0.4	*
13	Anemias (D50-D64)	3	0.4	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.3	*
15	Diabetes mellitus (E10-E14)	2	0.3	*
...	All other causes (Residual)	98	12.8	4.8
Asian or Pacific Islander, both sexes, 25-44 years				
...	All causes	2,979	100.0	57.5
1	Malignant neoplasms (C00-C97)	735	24.7	14.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	455	15.3	8.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	386	13.0	7.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	366	12.3	7.1
5	Cerebrovascular diseases (I60-I69)	139	4.7	2.7
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	129	4.3	2.5
7	Influenza and pneumonia (J09-J18)	73	2.5	1.4
8	Chronic liver disease and cirrhosis (K70,K73-K74)	59	2.0	1.1
9	Diabetes mellitus (E10-E14)	39	1.3	0.8
10	Human immunodeficiency virus (HIV) disease (B20-B24)	30	1.0	0.6
11	Septicemia (A40-A41)	29	1.0	0.6
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	29	1.0	0.6
13	Pregnancy, childbirth and the puerperium (O00-O99)	26	0.9	0.5
14	Chronic lower respiratory diseases (J40-J47)	22	0.7	0.4
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	21	0.7	0.4
...	All other causes (Residual)	441	14.8	8.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 45-64 years				
...	All causes	10,136	100.0	279.7
1	Malignant neoplasms (C00-C97)	4,109	40.5	113.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,983	19.6	54.7
3	Cerebrovascular diseases (I60-I69)	654	6.5	18.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	412	4.1	11.4
5	Diabetes mellitus (E10-E14)	403	4.0	11.1
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	312	3.1	8.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	215	2.1	5.9
8	Influenza and pneumonia (J09-J18)	164	1.6	4.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	142	1.4	3.9
10	Viral hepatitis (B15-B19)	129	1.3	3.6
11	Chronic lower respiratory diseases (J40-J47)	119	1.2	3.3
12	Septicemia (A40-A41)	110	1.1	3.0
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	98	1.0	2.7
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	81	0.8	2.2
15	Aortic aneurysm and dissection (I71)	59	0.6	1.6
...	All other causes (Residual)	1,146	11.3	31.6
Asian or Pacific Islander, both sexes, 65 years and over				
...	All causes	34,061	100.0	2,387.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,931	26.2	626.0
2	Malignant neoplasms (C00-C97)	8,289	24.3	581.0
3	Cerebrovascular diseases (I60-I69)	2,827	8.3	198.2
4	Diabetes mellitus (E10-E14)	1,402	4.1	98.3
5	Chronic lower respiratory diseases (J40-J47)	1,327	3.9	93.0
6	Influenza and pneumonia (J09-J18)	1,290	3.8	90.4
7	Alzheimer's disease (G30)	1,023	3.0	71.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	883	2.6	61.9
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	804	2.4	56.4
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	613	1.8	43.0
11	Septicemia (A40-A41)	423	1.2	29.7
12	Parkinson's disease (G20-G21)	353	1.0	24.7
13	Pneumonitis due to solids and liquids (J69)	282	0.8	19.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	227	0.7	15.9
15	Aortic aneurysm and dissection (I71)	208	0.6	14.6
...	All other causes (Residual)	5,179	15.2	363.0

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, all ages				
...	All causes	25,560	100.0	339.5
1	Malignant neoplasms (C00-C97)	6,897	27.0	91.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,238	24.4	82.9
3	Cerebrovascular diseases (I60-I69)	1,658	6.5	22.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,288	5.0	17.1
5	Diabetes mellitus (E10-E14)	943	3.7	12.5
6	Chronic lower respiratory diseases (J40-J47)	864	3.4	11.5
7	Influenza and pneumonia (J09-J18)	809	3.2	10.7
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	666	2.6	8.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	518	2.0	6.9
10	Alzheimer's disease (G30)	318	1.2	4.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	300	1.2	4.0
12	Septicemia (A40-A41)	293	1.1	3.9
13	Chronic liver disease and cirrhosis (K70,K73-K74)	289	1.1	3.8
14	Certain conditions originating in the perinatal period (P00-P96)	246	1.0	3.3
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	235	0.9	3.1
...	All other causes (Residual)	3,998	15.6	53.1
Asian or Pacific Islander, male, 1-4 years				
...	All causes	91	100.0	18.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	28	30.8	5.8
2	Malignant neoplasms (C00-C97)	9	9.9	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	8.8	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	5.5	*
5	Septicemia (A40-A41)	4	4.4	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	3.3	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.1	*
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	1.1	*
7	Influenza and pneumonia (J09-J18)	1	1.1	*
7	Acute bronchitis and bronchiolitis (J20-J21)	1	1.1	*
...	All other causes (Residual)	30	33.0	6.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 5-14 years				
...	All causes	122	100.0	11.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	29	23.8	2.7
2	Malignant neoplasms (C00-C97)	28	23.0	2.6
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	7.4	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	6	4.9	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	4.9	*
6	Influenza and pneumonia (J09-J18)	4	3.3	*
7	Cerebrovascular diseases (I60-I69)	2	1.6	*
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2	1.6	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.8	*
9	Meningitis (G00,G03)	1	0.8	*
9	Chronic lower respiratory diseases (J40-J47)	1	0.8	*
9	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.8	*
...	All other causes (Residual)	32	26.2	3.0
Asian or Pacific Islander, male, 15-24 years				
...	All causes	511	100.0	49.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	185	36.2	17.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	99	19.4	9.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	55	10.8	5.3
4	Malignant neoplasms (C00-C97)	45	8.8	4.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	26	5.1	2.5
6	Influenza and pneumonia (J09-J18)	9	1.8	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	1.4	*
8	Cerebrovascular diseases (I60-I69)	6	1.2	*
9	Septicemia (A40-A41)	3	0.6	*
10	Tuberculosis (A16-A19)	2	0.4	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.4	*
10	Anemias (D50-D64)	2	0.4	*
10	Diabetes mellitus (E10-E14)	2	0.4	*
10	Chronic lower respiratory diseases (J40-J47)	2	0.4	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
...	All other causes (Residual)	64	12.5	6.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 25-44 years				
...	All causes	1,873	100.0	74.4
1	Malignant neoplasms (C00-C97)	347	18.5	13.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	333	17.8	13.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	320	17.1	12.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	254	13.6	10.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	85	4.5	3.4
6	Cerebrovascular diseases (I60-I69)	75	4.0	3.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	46	2.5	1.8
8	Influenza and pneumonia (J09-J18)	42	2.2	1.7
9	Diabetes mellitus (E10-E14)	27	1.4	1.1
10	Human immunodeficiency virus (HIV) disease (B20-B24)	20	1.1	0.8
11	Septicemia (A40-A41)	18	1.0	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	0.9	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	0.9	*
13	Chronic lower respiratory diseases (J40-J47)	16	0.9	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	14	0.7	*
...	All other causes (Residual)	243	13.0	9.6
Asian or Pacific Islander, male, 45-64 years				
...	All causes	6,088	100.0	358.4
1	Malignant neoplasms (C00-C97)	2,116	34.8	124.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,508	24.8	88.8
3	Cerebrovascular diseases (I60-I69)	361	5.9	21.3
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	293	4.8	17.2
5	Diabetes mellitus (E10-E14)	263	4.3	15.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	215	3.5	12.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	161	2.6	9.5
8	Influenza and pneumonia (J09-J18)	96	1.6	5.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	86	1.4	5.1
10	Viral hepatitis (B15-B19)	83	1.4	4.9
11	Chronic lower respiratory diseases (J40-J47)	67	1.1	3.9
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	65	1.1	3.8
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	57	0.9	3.4
14	Septicemia (A40-A41)	56	0.9	3.3
15	Aortic aneurysm and dissection (I71)	37	0.6	2.2
...	All other causes (Residual)	624	10.2	36.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 65 years and over				
...	All causes	16,385	100.0	2,663.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,365	26.6	709.4
2	Malignant neoplasms (C00-C97)	4,348	26.5	706.7
3	Cerebrovascular diseases (I60-I69)	1,210	7.4	196.7
4	Chronic lower respiratory diseases (J40-J47)	778	4.7	126.4
5	Influenza and pneumonia (J09-J18)	652	4.0	106.0
6	Diabetes mellitus (E10-E14)	651	4.0	105.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	412	2.5	67.0
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	405	2.5	65.8
9	Alzheimer's disease (G30)	317	1.9	51.5
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	235	1.4	38.2
11	Parkinson's disease (G20-G21)	220	1.3	35.8
12	Septicemia (A40-A41)	210	1.3	34.1
13	Pneumonitis due to solids and liquids (J69)	153	0.9	24.9
14	Aortic aneurysm and dissection (I71)	115	0.7	18.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	110	0.7	17.9
...	All other causes (Residual)	2,204	13.5	358.2
Asian or Pacific Islander, female, all ages				
...	All causes	23,665	100.0	295.9
1	Malignant neoplasms (C00-C97)	6,377	26.9	79.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,137	21.7	64.2
3	Cerebrovascular diseases (I60-I69)	1,981	8.4	24.8
4	Diabetes mellitus (E10-E14)	903	3.8	11.3
5	Influenza and pneumonia (J09-J18)	759	3.2	9.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	756	3.2	9.5
7	Alzheimer's disease (G30)	706	3.0	8.8
8	Chronic lower respiratory diseases (J40-J47)	616	2.6	7.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	544	2.3	6.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	410	1.7	5.1
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	305	1.3	3.8
12	Septicemia (A40-A41)	283	1.2	3.5
13	Certain conditions originating in the perinatal period (P00-P96)	209	0.9	2.6
14	Chronic liver disease and cirrhosis (K70,K73-K74)	175	0.7	2.2
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	174	0.7	2.2
...	All other causes (Residual)	4,330	18.3	54.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 1-4 years				
...	All causes	58	100.0	12.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	21	36.2	4.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	13.8	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	8.6	*
4	Malignant neoplasms (C00-C97)	4	6.9	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	6.9	*
6	Influenza and pneumonia (J09-J18)	2	3.4	*
7	Septicemia (A40-A41)	1	1.7	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.7	*
7	Nutritional deficiencies (E40-E64)	1	1.7	*
7	Meningitis (G00,G03)	1	1.7	*
7	Chronic lower respiratory diseases (J40-J47)	1	1.7	*
7	Hernia (K40-K46)	1	1.7	*
7	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	1.7	*
...	All other causes (Residual)	7	12.1	*
Asian or Pacific Islander, female, 5-14 years				
...	All causes	109	100.0	10.6
1	Malignant neoplasms (C00-C97)	19	17.4	*
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18	16.5	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	11	10.1	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	9	8.3	*
5	Influenza and pneumonia (J09-J18)	7	6.4	*
6	Chronic lower respiratory diseases (J40-J47)	4	3.7	*
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4	3.7	*
8	Meningitis (G00,G03)	2	1.8	*
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	1.8	*
10	Cerebrovascular diseases (I60-I69)	1	0.9	*
10	Diseases of appendix (K35-K38)	1	0.9	*
10	Hernia (K40-K46)	1	0.9	*
10	Certain conditions originating in the perinatal period (P00-P96)	1	0.9	*
...	All other causes (Residual)	29	26.6	2.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 15-24 years				
...	All causes	252	100.0	25.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	71	28.2	7.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	53	21.0	5.3
3	Malignant neoplasms (C00-C97)	30	11.9	3.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	17	6.7	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	11	4.4	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	10	4.0	*
7	Cerebrovascular diseases (I60-I69)	6	2.4	*
7	Influenza and pneumonia (J09-J18)	6	2.4	*
9	Pregnancy, childbirth and the puerperium (O00-O99)	5	2.0	*
10	Chronic lower respiratory diseases (J40-J47)	3	1.2	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	1.2	*
12	Tuberculosis (A16-A19)	1	0.4	*
12	Septicemia (A40-A41)	1	0.4	*
12	Anemias (D50-D64)	1	0.4	*
...	All other causes (Residual)	34	13.5	3.4
Asian or Pacific Islander, female, 25-44 years				
...	All causes	1,106	100.0	41.6
1	Malignant neoplasms (C00-C97)	388	35.1	14.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	122	11.0	4.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	112	10.1	4.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	66	6.0	2.5
5	Cerebrovascular diseases (I60-I69)	64	5.8	2.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	44	4.0	1.7
7	Influenza and pneumonia (J09-J18)	31	2.8	1.2
8	Pregnancy, childbirth and the puerperium (O00-O99)	26	2.4	1.0
9	Chronic liver disease and cirrhosis (K70,K73-K74)	13	1.2	*
10	Diabetes mellitus (E10-E14)	12	1.1	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	1.1	*
12	Septicemia (A40-A41)	11	1.0	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	10	0.9	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	8	0.7	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	0.6	*
...	All other causes (Residual)	180	16.3	6.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 45-64 years				
...	All causes	4,048	100.0	210.3
1	Malignant neoplasms (C00-C97)	1,993	49.2	103.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	475	11.7	24.7
3	Cerebrovascular diseases (I60-I69)	293	7.2	15.2
4	Diabetes mellitus (E10-E14)	140	3.5	7.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	119	2.9	6.2
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	97	2.4	5.0
7	Influenza and pneumonia (J09-J18)	68	1.7	3.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	56	1.4	2.9
9	Septicemia (A40-A41)	54	1.3	2.8
9	Chronic liver disease and cirrhosis (K70,K73-K74)	54	1.3	2.8
11	Chronic lower respiratory diseases (J40-J47)	52	1.3	2.7
12	Viral hepatitis (B15-B19)	46	1.1	2.4
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	33	0.8	1.7
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	24	0.6	1.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	22	0.5	1.1
15	Aortic aneurysm and dissection (I71)	22	0.5	1.1
...	All other causes (Residual)	500	12.4	26.0
Asian or Pacific Islander, female, 65 years and over				
...	All causes	17,676	100.0	2,178.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,566	25.8	562.8
2	Malignant neoplasms (C00-C97)	3,941	22.3	485.7
3	Cerebrovascular diseases (I60-I69)	1,617	9.1	199.3
4	Diabetes mellitus (E10-E14)	751	4.2	92.6
5	Alzheimer's disease (G30)	706	4.0	87.0
6	Influenza and pneumonia (J09-J18)	638	3.6	78.6
7	Chronic lower respiratory diseases (J40-J47)	549	3.1	67.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	471	2.7	58.1
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	399	2.3	49.2
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	378	2.1	46.6
11	Septicemia (A40-A41)	213	1.2	26.3
12	Parkinson's disease (G20-G21)	133	0.8	16.4
13	Pneumonitis due to solids and liquids (J69)	129	0.7	15.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	117	0.7	14.4
15	Chronic liver disease and cirrhosis (K70,K73-K74)	108	0.6	13.3
...	All other causes (Residual)	2,960	16.7	364.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable