

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total deaths		
		Number	Rate	Rate
Hispanic, both sexes, all ages				
...	All causes	107,254	100.0	330.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	25,819	24.1	79.5
2	Malignant neoplasms (C00-C97)	21,160	19.7	65.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,830	8.2	27.2
4	Cerebrovascular diseases (I60-I69)	6,187	5.8	19.1
5	Diabetes mellitus (E10-E14)	5,328	5.0	16.4
6	Chronic liver disease and cirrhosis (K70,K73-K74)	3,187	3.0	9.8
7	Assault (homicide) (X85-Y09,Y87.1)	2,917	2.7	9.0
8	Chronic lower respiratory diseases (J40-J47)	2,689	2.5	8.3
9	Influenza and pneumonia (J10-J18)	2,625	2.4	8.1
10	Certain conditions originating in the perinatal period (P00-P96)	2,145	2.0	6.6
11	Human immunodeficiency virus (HIV) disease (B20-B24)	1,908	1.8	5.9
12	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,787	1.7	5.5
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,646	1.5	5.1
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,612	1.5	5.0
15	Septicemia (A40-A41)	1,226	1.1	3.8
...	All other causes (Residual)	18,188	17.0	56.0
Hispanic, both sexes, 1-4 years				
...	All causes	872	100.0	31.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	306	35.1	10.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	111	12.7	3.9
3	Malignant neoplasms (C00-C97)	75	8.6	2.7
4	Assault (homicide) (X85-Y09,Y87.1)	57	6.5	2.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	29	3.3	1.0
6	Influenza and pneumonia (J10-J18)	17	1.9	*
7	Septicemia (A40-A41)	15	1.7	*
8	Certain conditions originating in the perinatal period (P00-P96)	12	1.4	*
9	Chronic lower respiratory diseases (J40-J47)	9	1.0	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.8	*
11	Cerebrovascular diseases (I60-I69)	6	0.7	*
12	Meningitis (G00,G03)	4	0.5	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.5	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.5	*
15	Meningococcal infection (A39)	3	0.3	*
...	All other causes (Residual)	213	24.4	7.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total deaths		
		Number	Rate	Rate
Hispanic, both sexes, 5-9 years				
...	All causes	510	100.0	15.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	205	40.2	6.1
2	Malignant neoplasms (C00-C97)	91	17.8	2.7
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	33	6.5	1.0
4	Assault (homicide) (X85-Y09,Y87.1)	24	4.7	0.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	17	3.3	*
6	Influenza and pneumonia (J10-J18)	13	2.5	*
7	Certain conditions originating in the perinatal period (P00-P96)	7	1.4	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	5	1.0	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	1.0	*
8	Chronic lower respiratory diseases (J40-J47)	5	1.0	*
11	Cerebrovascular diseases (I60-I69)	4	0.8	*
12	Septicemia (A40-A41)	3	0.6	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.6	*
14	Meningococcal infection (A39)	2	0.4	*
14	Anemias (D50-D64)	2	0.4	*
14	Meningitis (G00,G03)	2	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
14	Intentional self-harm (suicide) (X60-X84,Y87.0)	2	0.4	*
...	All other causes (Residual)	85	16.7	2.5
Hispanic, both sexes, 10-14 years				
...	All causes	555	100.0	19.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	178	32.1	6.2
2	Malignant neoplasms (C00-C97)	107	19.3	3.7
3	Assault (homicide) (X85-Y09,Y87.1)	52	9.4	1.8
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	27	4.9	0.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	21	3.8	0.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	19	3.4	*
7	Chronic lower respiratory diseases (J40-J47)	10	1.8	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	1.4	*
8	Influenza and pneumonia (J10-J18)	8	1.4	*
10	Anemias (D50-D64)	5	0.9	*
11	Septicemia (A40-A41)	4	0.7	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.7	*
11	Cerebrovascular diseases (I60-I69)	4	0.7	*
11	Diseases of appendix (K35-K38)	4	0.7	*
15	Diabetes mellitus (E10-E14)	3	0.5	*
15	Certain conditions originating in the perinatal period (P00-P96)	3	0.5	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.5	*
...	All other causes (Residual)	95	17.1	3.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total deaths		
		Number	Rate	Rate
Hispanic, both sexes, 15-19 years				
...	All causes	1,953	100.0	68.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	884	45.3	31.0
2	Assault (homicide) (X85-Y09,Y87.1)	476	24.4	16.7
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	178	9.1	6.2
4	Malignant neoplasms (C00-C97)	122	6.2	4.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	49	2.5	1.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	24	1.2	0.8
7	Human immunodeficiency virus (HIV) disease (B20-B24)	7	0.4	*
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.4	*
9	Chronic lower respiratory diseases (J40-J47)	6	0.3	*
10	Diabetes mellitus (E10-E14)	5	0.3	*
10	Cerebrovascular diseases (I60-I69)	5	0.3	*
10	Aortic aneurysm and dissection (I71)	5	0.3	*
10	Legal intervention (Y35,Y89.0)	5	0.3	*
14	Influenza and pneumonia (J10-J18)	4	0.2	*
15	Meningococcal infection (A39)	3	0.2	*
15	Septicemia (A40-A41)	3	0.2	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.2	*
15	Diseases of appendix (K35-K38)	3	0.2	*
...	All other causes (Residual)	164	8.4	5.8
Hispanic, both sexes, 20-24 years				
...	All causes	2,840	100.0	101.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,211	42.6	43.5
2	Assault (homicide) (X85-Y09,Y87.1)	650	22.9	23.3
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	269	9.5	9.7
4	Malignant neoplasms (C00-C97)	153	5.4	5.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	76	2.7	2.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	30	1.1	1.1
7	Cerebrovascular diseases (I60-I69)	26	0.9	0.9
8	Pregnancy, childbirth and the puerperium (O00-099)	22	0.8	0.8
9	Influenza and pneumonia (J10-J18)	17	0.6	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	16	0.6	*
11	Legal intervention (Y35,Y89.0)	13	0.5	*
12	Chronic lower respiratory diseases (J40-J47)	10	0.4	*
13	Septicemia (A40-A41)	9	0.3	*
14	Diabetes mellitus (E10-E14)	8	0.3	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.2	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	7	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.2	*
...	All other causes (Residual)	309	10.9	11.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total deaths		
		Number	Rate	Rate
Hispanic, both sexes, 25-29 years				
...	All causes	2,598	100.0	99.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	944	36.3	36.2
2	Assault (homicide) (X85-Y09,Y87.1)	463	17.8	17.8
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	227	8.7	8.7
4	Malignant neoplasms (C00-C97)	196	7.5	7.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	112	4.3	4.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	98	3.8	3.8
7	Cerebrovascular diseases (I60-I69)	37	1.4	1.4
8	Diabetes mellitus (E10-E14)	21	0.8	0.8
9	Pregnancy, childbirth and the puerperium (O00-099)	20	0.8	0.8
10	Influenza and pneumonia (J10-J18)	19	0.7	*
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	19	0.7	*
12	Septicemia (A40-A41)	17	0.7	*
13	Chronic liver disease and cirrhosis (K70,K73-K74)	16	0.6	*
14	Anemias (D50-D64)	9	0.3	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.3	*
15	Aortic aneurysm and dissection (I71)	8	0.3	*
...	All other causes (Residual)	384	14.8	14.7
Hispanic, both sexes, 30-34 years				
...	All causes	2,911	100.0	108.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	832	28.6	31.1
2	Assault (homicide) (X85-Y09,Y87.1)	347	11.9	13.0
3	Malignant neoplasms (C00-C97)	280	9.6	10.5
4	Human immunodeficiency virus (HIV) disease (B20-B24)	267	9.2	10.0
5	Intentional self-harm (suicide) (X60-X84,Y87.0)	193	6.6	7.2
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	168	5.8	6.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	76	2.6	2.8
8	Cerebrovascular diseases (I60-I69)	71	2.4	2.7
9	Diabetes mellitus (E10-E14)	43	1.5	1.6
10	Influenza and pneumonia (J10-J18)	29	1.0	1.1
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	27	0.9	1.0
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20	0.7	0.7
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	0.6	*
14	Pregnancy, childbirth and the puerperium (O00-099)	16	0.5	*
15	Septicemia (A40-A41)	14	0.5	*
...	All other causes (Residual)	511	17.6	19.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total		
		Number	deaths	Rate
Hispanic, both sexes, 35-39 years				
...	All causes	3,656	100.0	137.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	856	23.4	32.1
2	Malignant neoplasms (C00-C97)	515	14.1	19.3
3	Human immunodeficiency virus (HIV) disease (B20-B24)	382	10.4	14.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	284	7.8	10.7
5	Assault (homicide) (X85-Y09,Y87.1)	248	6.8	9.3
6	Intentional self-harm (suicide) (X60-X84,Y87.0)	194	5.3	7.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	170	4.6	6.4
8	Cerebrovascular diseases (I60-I69)	95	2.6	3.6
9	Diabetes mellitus (E10-E14)	49	1.3	1.8
10	Viral hepatitis (B15-B19)	39	1.1	1.5
11	Influenza and pneumonia (J10-J18)	33	0.9	1.2
12	Septicemia (A40-A41)	28	0.8	1.1
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	24	0.7	0.9
14	Chronic lower respiratory diseases (J40-J47)	21	0.6	0.8
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19	0.5	*
15	Pregnancy, childbirth and the puerperium (O00-099)	19	0.5	*
...	All other causes (Residual)	680	18.6	25.5
Hispanic, both sexes, 40-44 years				
...	All causes	4,429	100.0	194.7
1	Malignant neoplasms (C00-C97)	780	17.6	34.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	762	17.2	33.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	476	10.7	20.9
4	Human immunodeficiency virus (HIV) disease (B20-B24)	380	8.6	16.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	331	7.5	14.6
6	Assault (homicide) (X85-Y09,Y87.1)	199	4.5	8.7
7	Intentional self-harm (suicide) (X60-X84,Y87.0)	191	4.3	8.4
8	Cerebrovascular diseases (I60-I69)	172	3.9	7.6
9	Diabetes mellitus (E10-E14)	105	2.4	4.6
10	Viral hepatitis (B15-B19)	87	2.0	3.8
11	Septicemia (A40-A41)	51	1.2	2.2
12	Influenza and pneumonia (J10-J18)	49	1.1	2.2
13	Chronic lower respiratory diseases (J40-J47)	34	0.8	1.5
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	34	0.8	1.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	0.5	1.0
...	All other causes (Residual)	755	17.0	33.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total		
		Number	deaths	Rate
Hispanic, both sexes, 45-49 years				
...	All causes	4,990	100.0	283.2
1	Malignant neoplasms (C00-C97)	1,141	22.9	64.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	709	14.2	40.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	542	10.9	30.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	440	8.8	25.0
5	Human immunodeficiency virus (HIV) disease (B20-B24)	322	6.5	18.3
6	Cerebrovascular diseases (I60-I69)	245	4.9	13.9
7	Diabetes mellitus (E10-E14)	174	3.5	9.9
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	141	2.8	8.0
9	Assault (homicide) (X85-Y09,Y87.1)	135	2.7	7.7
10	Viral hepatitis (B15-B19)	119	2.4	6.8
11	Influenza and pneumonia (J10-J18)	54	1.1	3.1
12	Septicemia (A40-A41)	51	1.0	2.9
13	Chronic lower respiratory diseases (J40-J47)	45	0.9	2.6
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	45	0.9	2.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	0.4	1.1
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	20	0.4	1.1
...	All other causes (Residual)	787	15.8	44.7
Hispanic, both sexes, 50-54 years				
...	All causes	5,366	100.0	398.4
1	Malignant neoplasms (C00-C97)	1,434	26.7	106.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,045	19.5	77.6
3	Chronic liver disease and cirrhosis (K70,K73-K74)	414	7.7	30.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	411	7.7	30.5
5	Diabetes mellitus (E10-E14)	265	4.9	19.7
6	Cerebrovascular diseases (I60-I69)	232	4.3	17.2
7	Human immunodeficiency virus (HIV) disease (B20-B24)	209	3.9	15.5
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	103	1.9	7.6
9	Viral hepatitis (B15-B19)	99	1.8	7.3
10	Assault (homicide) (X85-Y09,Y87.1)	76	1.4	5.6
11	Septicemia (A40-A41)	71	1.3	5.3
12	Influenza and pneumonia (J10-J18)	66	1.2	4.9
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	64	1.2	4.8
14	Chronic lower respiratory diseases (J40-J47)	60	1.1	4.5
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	37	0.7	2.7
...	All other causes (Residual)	780	14.5	57.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total		
		Number	deaths	Rate
Hispanic, both sexes, 55-59 years				
...	All causes	5,876	100.0	595.0
1	Malignant neoplasms (C00-C97)	1,735	29.5	175.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,355	23.1	137.2
3	Diabetes mellitus (E10-E14)	436	7.4	44.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	380	6.5	38.5
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	272	4.6	27.5
6	Cerebrovascular diseases (I60-I69)	271	4.6	27.4
7	Human immunodeficiency virus (HIV) disease (B20-B24)	96	1.6	9.7
8	Chronic lower respiratory diseases (J40-J47)	87	1.5	8.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	87	1.5	8.8
10	Influenza and pneumonia (J10-J18)	82	1.4	8.3
11	Viral hepatitis (B15-B19)	67	1.1	6.8
12	Intentional self-harm (suicide) (X60-X84,Y87.0)	62	1.1	6.3
13	Septicemia (A40-A41)	61	1.0	6.2
14	Assault (homicide) (X85-Y09,Y87.1)	50	0.9	5.1
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	34	0.6	3.4
...	All other causes (Residual)	801	13.6	81.1
Hispanic, both sexes, 60-64 years				
...	All causes	6,857	100.0	881.6
1	Malignant neoplasms (C00-C97)	2,045	29.8	262.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,781	26.0	229.0
3	Diabetes mellitus (E10-E14)	514	7.5	66.1
4	Cerebrovascular diseases (I60-I69)	410	6.0	52.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	328	4.8	42.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	220	3.2	28.3
7	Chronic lower respiratory diseases (J40-J47)	157	2.3	20.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	145	2.1	18.6
9	Influenza and pneumonia (J10-J18)	109	1.6	14.0
10	Septicemia (A40-A41)	84	1.2	10.8
11	Viral hepatitis (B15-B19)	80	1.2	10.3
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	75	1.1	9.6
13	Human immunodeficiency virus (HIV) disease (B20-B24)	67	1.0	8.6
14	Intentional self-harm (suicide) (X60-X84,Y87.0)	40	0.6	5.1
15	Aortic aneurysm and dissection (I71)	32	0.5	4.1
...	All other causes (Residual)	770	11.2	99.0

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total		
		Number	deaths	Rate
Hispanic, both sexes, 65-69 years				
...	All causes	8,546	100.0	1,340.7
1	Malignant neoplasms (C00-C97)	2,532	29.6	397.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,328	27.2	365.2
3	Diabetes mellitus (E10-E14)	684	8.0	107.3
4	Cerebrovascular diseases (I60-I69)	498	5.8	78.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	333	3.9	52.2
6	Chronic lower respiratory diseases (J40-J47)	240	2.8	37.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	195	2.3	30.6
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	187	2.2	29.3
9	Influenza and pneumonia (J10-J18)	138	1.6	21.6
10	Septicemia (A40-A41)	97	1.1	15.2
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	65	0.8	10.2
12	Viral hepatitis (B15-B19)	62	0.7	9.7
13	Aortic aneurysm and dissection (I71)	46	0.5	7.2
14	Alzheimer's disease (G30)	43	0.5	6.7
14	Intentional self-harm (suicide) (X60-X84,Y87.0)	43	0.5	6.7
...	All other causes (Residual)	1,055	12.3	165.5
Hispanic, both sexes, 70-74 years				
...	All causes	10,874	100.0	2,106.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,125	28.7	605.4
2	Malignant neoplasms (C00-C97)	3,025	27.8	586.1
3	Diabetes mellitus (E10-E14)	908	8.4	175.9
4	Cerebrovascular diseases (I60-I69)	721	6.6	139.7
5	Chronic lower respiratory diseases (J40-J47)	332	3.1	64.3
6	Chronic liver disease and cirrhosis (K70,K73-K74)	295	2.7	57.2
7	Influenza and pneumonia (J10-J18)	267	2.5	51.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	231	2.1	44.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	209	1.9	40.5
10	Septicemia (A40-A41)	138	1.3	26.7
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	97	0.9	18.8
12	Alzheimer's disease (G30)	85	0.8	16.5
13	Aortic aneurysm and dissection (I71)	75	0.7	14.5
14	Viral hepatitis (B15-B19)	58	0.5	11.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	58	0.5	11.2
...	All other causes (Residual)	1,250	11.5	242.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total		
		Number	deaths	Rate
Hispanic, both sexes, 75-79 years				
...	All causes	11,578	100.0	3,085.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,583	30.9	954.8
2	Malignant neoplasms (C00-C97)	2,783	24.0	741.6
3	Cerebrovascular diseases (I60-I69)	859	7.4	228.9
4	Diabetes mellitus (E10-E14)	817	7.1	217.7
5	Chronic lower respiratory diseases (J40-J47)	468	4.0	124.7
6	Influenza and pneumonia (J10-J18)	313	2.7	83.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	235	2.0	62.6
8	Chronic liver disease and cirrhosis (K70,K73-K74)	215	1.9	57.3
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	193	1.7	51.4
10	Alzheimer's disease (G30)	180	1.6	48.0
11	Septicemia (A40-A41)	157	1.4	41.8
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	118	1.0	31.4
13	Parkinson's disease (G20-G21)	89	0.8	23.7
14	Aortic aneurysm and dissection (I71)	83	0.7	22.1
15	Pneumonitis due to solids and liquids (J69)	68	0.6	18.1
...	All other causes (Residual)	1,417	12.2	377.6
Hispanic, both sexes, 80-84 years				
...	All causes	10,373	100.0	4,772.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,484	33.6	1,602.9
2	Malignant neoplasms (C00-C97)	2,044	19.7	940.4
3	Cerebrovascular diseases (I60-I69)	883	8.5	406.2
4	Diabetes mellitus (E10-E14)	593	5.7	272.8
5	Chronic lower respiratory diseases (J40-J47)	453	4.4	208.4
6	Influenza and pneumonia (J10-J18)	377	3.6	173.4
7	Alzheimer's disease (G30)	221	2.1	101.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	194	1.9	89.3
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	172	1.7	79.1
10	Septicemia (A40-A41)	143	1.4	65.8
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	114	1.1	52.4
12	Chronic liver disease and cirrhosis (K70,K73-K74)	108	1.0	49.7
13	Parkinson's disease (G20-G21)	102	1.0	46.9
14	Pneumonitis due to solids and liquids (J69)	73	0.7	33.6
15	Atherosclerosis (I70)	71	0.7	32.7
...	All other causes (Residual)	1,341	12.9	616.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total		
		Number	deaths	Rate
Hispanic, both sexes, 85-89 years				
...	All causes	9,170	100.0	7,336.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,438	37.5	2,750.6
2	Malignant neoplasms (C00-C97)	1,292	14.1	1,033.7
3	Cerebrovascular diseases (I60-I69)	852	9.3	681.6
4	Influenza and pneumonia (J10-J18)	440	4.8	352.0
5	Diabetes mellitus (E10-E14)	412	4.5	329.6
6	Chronic lower respiratory diseases (J40-J47)	400	4.4	320.0
7	Alzheimer's disease (G30)	288	3.1	230.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	182	2.0	145.6
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	146	1.6	116.8
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	122	1.3	97.6
11	Septicemia (A40-A41)	119	1.3	95.2
12	Pneumonitis due to solids and liquids (J69)	87	0.9	69.6
13	Atherosclerosis (I70)	83	0.9	66.4
14	Parkinson's disease (G20-G21)	81	0.9	64.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	63	0.7	50.4
...	All other causes (Residual)	1,165	12.7	932.1
Hispanic, both sexes, 90-94 years				
...	All causes	5,853	100.0	10,270.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,438	41.7	4,278.2
2	Malignant neoplasms (C00-C97)	586	10.0	1,028.3
3	Cerebrovascular diseases (I60-I69)	539	9.2	945.8
4	Influenza and pneumonia (J10-J18)	339	5.8	594.9
5	Chronic lower respiratory diseases (J40-J47)	217	3.7	380.8
6	Diabetes mellitus (E10-E14)	213	3.6	373.8
7	Alzheimer's disease (G30)	201	3.4	352.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	127	2.2	222.9
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	94	1.6	164.9
10	Pneumonitis due to solids and liquids (J69)	78	1.3	136.9
11	Atherosclerosis (I70)	77	1.3	135.1
12	Septicemia (A40-A41)	70	1.2	122.8
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	55	0.9	96.5
14	Parkinson's disease (G20-G21)	42	0.7	73.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	29	0.5	50.9
15	Nutritional deficiencies (E40-E64)	29	0.5	50.9
...	All other causes (Residual)	719	12.3	1,261.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total deaths		
		Number	Rate	Rate
Hispanic, both sexes, 95-99 years				
...	All causes	2,334	100.0	12,226.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	995	42.6	5,212.2
2	Cerebrovascular diseases (I60-I69)	209	9.0	1,094.8
3	Malignant neoplasms (C00-C97)	183	7.8	958.6
4	Influenza and pneumonia (J10-J18)	161	6.9	843.4
5	Chronic lower respiratory diseases (J40-J47)	94	4.0	492.4
6	Alzheimer's disease (G30)	76	3.3	398.1
7	Diabetes mellitus (E10-E14)	66	2.8	345.7
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	38	1.6	199.1
8	Pneumonitis due to solids and liquids (J69)	38	1.6	199.1
10	Atherosclerosis (I70)	32	1.4	167.6
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	31	1.3	162.4
12	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	30	1.3	157.2
13	Septicemia (A40-A41)	24	1.0	125.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	0.5	*
15	Nutritional deficiencies (E40-E64)	8	0.3	*
...	All other causes (Residual)	337	14.4	1,765.3
Hispanic, both sexes, 100 years and over				
...	All causes	461	100.0	13,627.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	215	46.6	6,355.3
2	Cerebrovascular diseases (I60-I69)	32	6.9	945.9
3	Influenza and pneumonia (J10-J18)	30	6.5	886.8
4	Malignant neoplasms (C00-C97)	24	5.2	709.4
5	Chronic lower respiratory diseases (J40-J47)	14	3.0	*
6	Diabetes mellitus (E10-E14)	10	2.2	*
7	Alzheimer's disease (G30)	8	1.7	*
7	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	8	1.7	*
7	Atherosclerosis (I70)	8	1.7	*
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8	1.7	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	1.3	*
12	Septicemia (A40-A41)	5	1.1	*
13	Pneumonitis due to solids and liquids (J69)	4	0.9	*
14	Nutritional deficiencies (E40-E64)	3	0.7	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.4	*
15	Parkinson's disease (G20-G21)	2	0.4	*
15	Peptic ulcer (K25-K28)	2	0.4	*
...	All other causes (Residual)	80	17.4	2,364.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total deaths		
		Number	Rate	Rate
Hispanic, male, all ages				
...	All causes	60,172	100.0	368.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,566	22.5	83.2
2	Malignant neoplasms (C00-C97)	11,138	18.5	68.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,696	11.1	41.1
4	Cerebrovascular diseases (I60-I69)	2,865	4.8	17.6
5	Diabetes mellitus (E10-E14)	2,507	4.2	15.4
6	Assault (homicide) (X85-Y09,Y87.1)	2,431	4.0	14.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,312	3.8	14.2
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,525	2.5	9.3
9	Human immunodeficiency virus (HIV) disease (B20-B24)	1,493	2.5	9.2
10	Chronic lower respiratory diseases (J40-J47)	1,451	2.4	8.9
11	Influenza and pneumonia (J10-J18)	1,303	2.2	8.0
12	Certain conditions originating in the perinatal period (P00-P96)	1,194	2.0	7.3
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	857	1.4	5.3
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	805	1.3	4.9
15	Septicemia (A40-A41)	625	1.0	3.8
...	All other causes (Residual)	9,404	15.6	57.7
Hispanic, male, 1-4 years				
...	All causes	475	100.0	33.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	174	36.6	12.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	60	12.6	4.2
3	Malignant neoplasms (C00-C97)	43	9.1	3.0
4	Assault (homicide) (X85-Y09,Y87.1)	30	6.3	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	21	4.4	1.5
6	Septicemia (A40-A41)	9	1.9	*
7	Certain conditions originating in the perinatal period (P00-P96)	7	1.5	*
8	Influenza and pneumonia (J10-J18)	5	1.1	*
8	Chronic lower respiratory diseases (J40-J47)	5	1.1	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.6	*
11	Meningococcal infection (A39)	2	0.4	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.4	*
11	Meningitis (G00,G03)	2	0.4	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
15	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
15	Diseases of appendix (K35-K38)	1	0.2	*
...	All other causes (Residual)	108	22.7	7.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total deaths		
		Number	Rate	Rate
Hispanic, male, 5-9 years				
...	All causes	277	100.0	16.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	105	37.9	6.1
2	Malignant neoplasms (C00-C97)	55	19.9	3.2
3	Assault (homicide) (X85-Y09,Y87.1)	19	6.9	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	16	5.8	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	9	3.2	*
6	Influenza and pneumonia (J10-J18)	7	2.5	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	1.4	*
7	Certain conditions originating in the perinatal period (P00-P96)	4	1.4	*
9	Septicemia (A40-A41)	3	1.1	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	3	1.1	*
11	Chronic lower respiratory diseases (J40-J47)	2	0.7	*
11	Intentional self-harm (suicide) (X60-X84,Y87.0)	2	0.7	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.7	*
14	Anemias (D50-D64)	1	0.4	*
14	Meningitis (G00,G03)	1	0.4	*
14	Cerebrovascular diseases (I60-I69)	1	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.4	*
...	All other causes (Residual)	42	15.2	2.5
Hispanic, male, 10-14 years				
...	All causes	345	100.0	23.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	105	30.4	7.1
2	Malignant neoplasms (C00-C97)	67	19.4	4.6
3	Assault (homicide) (X85-Y09,Y87.1)	37	10.7	2.5
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	20	5.8	1.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	17	4.9	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	12	3.5	*
7	Chronic lower respiratory diseases (J40-J47)	7	2.0	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	1.2	*
9	Influenza and pneumonia (J10-J18)	3	0.9	*
9	Diseases of appendix (K35-K38)	3	0.9	*
11	Septicemia (A40-A41)	2	0.6	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.6	*
11	Anemias (D50-D64)	2	0.6	*
11	Diabetes mellitus (E10-E14)	2	0.6	*
11	Cerebrovascular diseases (I60-I69)	2	0.6	*
11	Certain conditions originating in the perinatal period (P00-P96)	2	0.6	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.6	*
...	All other causes (Residual)	56	16.2	3.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total		
		Number	deaths	Rate
Hispanic, male, 15-19 years				
...	All causes	1,528	100.0	103.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	689	45.1	46.6
2	Assault (homicide) (X85-Y09,Y87.1)	434	28.4	29.3
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	143	9.4	9.7
4	Malignant neoplasms (C00-C97)	72	4.7	4.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	36	2.4	2.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	19	1.2	*
7	Chronic lower respiratory diseases (J40-J47)	5	0.3	*
7	Legal intervention (Y35,Y89.0)	5	0.3	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.3	*
9	Diabetes mellitus (E10-E14)	4	0.3	*
11	Aortic aneurysm and dissection (I71)	3	0.2	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.2	*
13	Meningococcal infection (A39)	2	0.1	*
13	Septicemia (A40-A41)	2	0.1	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.1	*
13	Cerebrovascular diseases (I60-I69)	2	0.1	*
...	All other causes (Residual)	103	6.7	7.0
Hispanic, male, 20-24 years				
...	All causes	2,309	100.0	159.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,028	44.5	71.1
2	Assault (homicide) (X85-Y09,Y87.1)	580	25.1	40.1
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	244	10.6	16.9
4	Malignant neoplasms (C00-C97)	90	3.9	6.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	51	2.2	3.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	20	0.9	1.4
7	Cerebrovascular diseases (I60-I69)	15	0.6	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	14	0.6	*
9	Legal intervention (Y35,Y89.0)	12	0.5	*
10	Influenza and pneumonia (J10-J18)	9	0.4	*
11	Chronic lower respiratory diseases (J40-J47)	8	0.3	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.3	*
13	Chronic liver disease and cirrhosis (K70,K73-K74)	5	0.2	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.2	*
15	Meningococcal infection (A39)	4	0.2	*
15	Septicemia (A40-A41)	4	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.2	*
...	All other causes (Residual)	210	9.1	14.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total deaths		
		Number	Rate	Rate
Hispanic, male, 25-29 years				
...	All causes	2,014	100.0	151.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	804	39.9	60.6
2	Assault (homicide) (X85-Y09,Y87.1)	405	20.1	30.5
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	203	10.1	15.3
4	Malignant neoplasms (C00-C97)	105	5.2	7.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	79	3.9	6.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	64	3.2	4.8
7	Cerebrovascular diseases (I60-I69)	22	1.1	1.7
8	Diabetes mellitus (E10-E14)	16	0.8	*
9	Chronic liver disease and cirrhosis (K70,K73-K74)	14	0.7	*
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	11	0.5	*
11	Influenza and pneumonia (J10-J18)	10	0.5	*
12	Septicemia (A40-A41)	9	0.4	*
13	Aortic aneurysm and dissection (I71)	6	0.3	*
13	Legal intervention (Y35,Y89.0)	6	0.3	*
15	Anemias (D50-D64)	4	0.2	*
15	Chronic lower respiratory diseases (J40-J47)	4	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.2	*
...	All other causes (Residual)	248	12.3	18.7
Hispanic, male, 30-34 years				
...	All causes	2,186	100.0	158.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	695	31.8	50.3
2	Assault (homicide) (X85-Y09,Y87.1)	292	13.4	21.1
3	Human immunodeficiency virus (HIV) disease (B20-B24)	210	9.6	15.2
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	167	7.6	12.1
5	Malignant neoplasms (C00-C97)	136	6.2	9.8
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	120	5.5	8.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	68	3.1	4.9
8	Cerebrovascular diseases (I60-I69)	43	2.0	3.1
9	Diabetes mellitus (E10-E14)	32	1.5	2.3
10	Influenza and pneumonia (J10-J18)	17	0.8	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	15	0.7	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	0.6	*
13	Septicemia (A40-A41)	12	0.5	*
14	Viral hepatitis (B15-B19)	9	0.4	*
14	Legal intervention (Y35,Y89.0)	9	0.4	*
...	All other causes (Residual)	348	15.9	25.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total		
		Number	deaths	Rate
Hispanic, male, 35-39 years				
...	All causes	2,543	100.0	185.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	682	26.8	49.8
2	Human immunodeficiency virus (HIV) disease (B20-B24)	309	12.2	22.6
3	Malignant neoplasms (C00-C97)	204	8.0	14.9
4	Assault (homicide) (X85-Y09,Y87.1)	199	7.8	14.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	198	7.8	14.5
6	Intentional self-harm (suicide) (X60-X84,Y87.0)	158	6.2	11.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	134	5.3	9.8
8	Cerebrovascular diseases (I60-I69)	53	2.1	3.9
9	Viral hepatitis (B15-B19)	30	1.2	2.2
10	Diabetes mellitus (E10-E14)	28	1.1	2.0
11	Influenza and pneumonia (J10-J18)	21	0.8	1.5
12	Septicemia (A40-A41)	16	0.6	*
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	16	0.6	*
14	Chronic lower respiratory diseases (J40-J47)	11	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.4	*
...	All other causes (Residual)	475	18.7	34.7
Hispanic, male, 40-44 years				
...	All causes	3,054	100.0	265.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	635	20.8	55.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	352	11.5	30.6
3	Malignant neoplasms (C00-C97)	330	10.8	28.6
4	Human immunodeficiency virus (HIV) disease (B20-B24)	294	9.6	25.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	265	8.7	23.0
6	Intentional self-harm (suicide) (X60-X84,Y87.0)	162	5.3	14.1
7	Assault (homicide) (X85-Y09,Y87.1)	160	5.2	13.9
8	Cerebrovascular diseases (I60-I69)	92	3.0	8.0
9	Viral hepatitis (B15-B19)	70	2.3	6.1
10	Diabetes mellitus (E10-E14)	67	2.2	5.8
11	Influenza and pneumonia (J10-J18)	35	1.1	3.0
12	Septicemia (A40-A41)	30	1.0	2.6
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	27	0.9	2.3
14	Chronic lower respiratory diseases (J40-J47)	13	0.4	*
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	12	0.4	*
...	All other causes (Residual)	510	16.7	44.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total deaths		
		Number	Rate	Rate
Hispanic, male, 45-49 years				
...	All causes	3,357	100.0	384.7
1	Malignant neoplasms (C00-C97)	540	16.1	61.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	539	16.1	61.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	435	13.0	49.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	365	10.9	41.8
5	Human immunodeficiency virus (HIV) disease (B20-B24)	252	7.5	28.9
6	Cerebrovascular diseases (I60-I69)	144	4.3	16.5
7	Intentional self-harm (suicide) (X60-X84,Y87.0)	112	3.3	12.8
8	Diabetes mellitus (E10-E14)	98	2.9	11.2
8	Assault (homicide) (X85-Y09,Y87.1)	98	2.9	11.2
10	Viral hepatitis (B15-B19)	91	2.7	10.4
11	Influenza and pneumonia (J10-J18)	41	1.2	4.7
12	Septicemia (A40-A41)	36	1.1	4.1
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	32	1.0	3.7
14	Chronic lower respiratory diseases (J40-J47)	21	0.6	2.4
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	16	0.5	*
...	All other causes (Residual)	537	16.0	61.5
Hispanic, male, 50-54 years				
...	All causes	3,451	100.0	531.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	768	22.3	118.3
2	Malignant neoplasms (C00-C97)	692	20.1	106.6
3	Chronic liver disease and cirrhosis (K70,K73-K74)	342	9.9	52.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	315	9.1	48.5
5	Human immunodeficiency virus (HIV) disease (B20-B24)	174	5.0	26.8
6	Diabetes mellitus (E10-E14)	142	4.1	21.9
7	Cerebrovascular diseases (I60-I69)	136	3.9	20.9
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	82	2.4	12.6
9	Viral hepatitis (B15-B19)	74	2.1	11.4
10	Assault (homicide) (X85-Y09,Y87.1)	60	1.7	9.2
11	Septicemia (A40-A41)	42	1.2	6.5
12	Influenza and pneumonia (J10-J18)	41	1.2	6.3
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	37	1.1	5.7
14	Chronic lower respiratory diseases (J40-J47)	32	0.9	4.9
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	28	0.8	4.3
...	All other causes (Residual)	486	14.1	74.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total deaths		
		Number	Rate	Rate
Hispanic, male, 55-59 years				
...	All causes	3,656	100.0	790.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	952	26.0	205.9
2	Malignant neoplasms (C00-C97)	911	24.9	197.0
3	Chronic liver disease and cirrhosis (K70,K73-K74)	295	8.1	63.8
4	Diabetes mellitus (E10-E14)	241	6.6	52.1
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	193	5.3	41.7
6	Cerebrovascular diseases (I60-I69)	172	4.7	37.2
7	Human immunodeficiency virus (HIV) disease (B20-B24)	79	2.2	17.1
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	54	1.5	11.7
9	Influenza and pneumonia (J10-J18)	50	1.4	10.8
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	45	1.2	9.7
11	Viral hepatitis (B15-B19)	43	1.2	9.3
12	Chronic lower respiratory diseases (J40-J47)	40	1.1	8.6
12	Assault (homicide) (X85-Y09,Y87.1)	40	1.1	8.6
14	Septicemia (A40-A41)	37	1.0	8.0
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	20	0.5	4.3
...	All other causes (Residual)	484	13.2	104.7
Hispanic, male, 60-64 years				
...	All causes	4,104	100.0	1,154.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,176	28.7	330.8
2	Malignant neoplasms (C00-C97)	1,123	27.4	315.9
3	Diabetes mellitus (E10-E14)	279	6.8	78.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	236	5.8	66.4
5	Cerebrovascular diseases (I60-I69)	220	5.4	61.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	154	3.8	43.3
7	Chronic lower respiratory diseases (J40-J47)	106	2.6	29.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	70	1.7	19.7
9	Influenza and pneumonia (J10-J18)	69	1.7	19.4
10	Viral hepatitis (B15-B19)	51	1.2	14.3
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	45	1.1	12.7
12	Human immunodeficiency virus (HIV) disease (B20-B24)	44	1.1	12.4
13	Septicemia (A40-A41)	43	1.0	12.1
14	Intentional self-harm (suicide) (X60-X84,Y87.0)	35	0.9	9.8
15	Aortic aneurysm and dissection (I71)	23	0.6	6.5
...	All other causes (Residual)	430	10.5	120.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total		
		Number	deaths	Rate
Hispanic, male, 65-69 years				
...	All causes	4,838	100.0	1,702.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,409	29.1	495.8
2	Malignant neoplasms (C00-C97)	1,398	28.9	491.9
3	Diabetes mellitus (E10-E14)	338	7.0	118.9
4	Cerebrovascular diseases (I60-I69)	267	5.5	94.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	233	4.8	82.0
6	Chronic lower respiratory diseases (J40-J47)	136	2.8	47.9
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	126	2.6	44.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	90	1.9	31.7
9	Influenza and pneumonia (J10-J18)	75	1.6	26.4
10	Septicemia (A40-A41)	60	1.2	21.1
11	Intentional self-harm (suicide) (X60-X84,Y87.0)	39	0.8	13.7
12	Aortic aneurysm and dissection (I71)	33	0.7	11.6
13	Viral hepatitis (B15-B19)	28	0.6	9.9
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	26	0.5	9.1
15	Human immunodeficiency virus (HIV) disease (B20-B24)	22	0.5	7.7
...	All other causes (Residual)	558	11.5	196.4
Hispanic, male, 70-74 years				
...	All causes	6,004	100.0	2,663.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,836	30.6	814.6
2	Malignant neoplasms (C00-C97)	1,688	28.1	748.9
3	Diabetes mellitus (E10-E14)	424	7.1	188.1
4	Cerebrovascular diseases (I60-I69)	378	6.3	167.7
5	Chronic lower respiratory diseases (J40-J47)	207	3.4	91.8
6	Chronic liver disease and cirrhosis (K70,K73-K74)	162	2.7	71.9
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	145	2.4	64.3
8	Influenza and pneumonia (J10-J18)	143	2.4	63.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	104	1.7	46.1
10	Septicemia (A40-A41)	66	1.1	29.3
11	Alzheimer's disease (G30)	47	0.8	20.9
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	46	0.8	20.4
13	Aortic aneurysm and dissection (I71)	43	0.7	19.1
14	Intentional self-harm (suicide) (X60-X84,Y87.0)	35	0.6	15.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	30	0.5	13.3
...	All other causes (Residual)	650	10.8	288.4

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total		
		Number	deaths	Rate
Hispanic, male, 75-79 years				
...	All causes	6,069	100.0	3,782.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,912	31.5	1,191.5
2	Malignant neoplasms (C00-C97)	1,545	25.5	962.8
3	Cerebrovascular diseases (I60-I69)	421	6.9	262.4
4	Diabetes mellitus (E10-E14)	362	6.0	225.6
5	Chronic lower respiratory diseases (J40-J47)	260	4.3	162.0
6	Influenza and pneumonia (J10-J18)	166	2.7	103.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	121	2.0	75.4
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	117	1.9	72.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	116	1.9	72.3
10	Septicemia (A40-A41)	79	1.3	49.2
11	Alzheimer's disease (G30)	75	1.2	46.7
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	57	0.9	35.5
12	Aortic aneurysm and dissection (I71)	57	0.9	35.5
14	Parkinson's disease (G20-G21)	53	0.9	33.0
15	Pneumonitis due to solids and liquids (J69)	44	0.7	27.4
...	All other causes (Residual)	684	11.3	426.2
Hispanic, male, 80-84 years				
...	All causes	4,904	100.0	5,769.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,613	32.9	1,897.7
2	Malignant neoplasms (C00-C97)	1,112	22.7	1,308.3
3	Cerebrovascular diseases (I60-I69)	378	7.7	444.7
4	Chronic lower respiratory diseases (J40-J47)	249	5.1	293.0
5	Diabetes mellitus (E10-E14)	247	5.0	290.6
6	Influenza and pneumonia (J10-J18)	201	4.1	236.5
7	Alzheimer's disease (G30)	91	1.9	107.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	90	1.8	105.9
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	74	1.5	87.1
10	Septicemia (A40-A41)	64	1.3	75.3
11	Parkinson's disease (G20-G21)	59	1.2	69.4
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	50	1.0	58.8
13	Pneumonitis due to solids and liquids (J69)	43	0.9	50.6
14	Chronic liver disease and cirrhosis (K70,K73-K74)	42	0.9	49.4
15	Atherosclerosis (I70)	31	0.6	36.5
...	All other causes (Residual)	560	11.4	658.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 85-89 years				
...	All causes	3,617	100.0	8,222.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,312	36.3	2,982.6
2	Malignant neoplasms (C00-C97)	632	17.5	1,436.8
3	Cerebrovascular diseases (I60-I69)	282	7.8	641.1
4	Chronic lower respiratory diseases (J40-J47)	190	5.3	431.9
5	Influenza and pneumonia (J10-J18)	164	4.5	372.8
6	Diabetes mellitus (E10-E14)	146	4.0	331.9
7	Alzheimer's disease (G30)	83	2.3	188.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	79	2.2	179.6
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	69	1.9	156.9
10	Parkinson's disease (G20-G21)	49	1.4	111.4
11	Septicemia (A40-A41)	41	1.1	93.2
11	Pneumonitis due to solids and liquids (J69)	41	1.1	93.2
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	38	1.1	86.4
14	Atherosclerosis (I70)	27	0.7	61.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	26	0.7	59.1
...	All other causes (Residual)	438	12.1	995.7
Hispanic, male, 90-94 years				
...	All causes	2,035	100.0	11,011.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	793	39.0	4,290.9
2	Malignant neoplasms (C00-C97)	292	14.3	1,580.0
3	Cerebrovascular diseases (I60-I69)	159	7.8	860.3
4	Influenza and pneumonia (J10-J18)	144	7.1	779.2
5	Chronic lower respiratory diseases (J40-J47)	95	4.7	514.0
6	Diabetes mellitus (E10-E14)	60	2.9	324.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	52	2.6	281.4
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	42	2.1	227.3
9	Alzheimer's disease (G30)	40	2.0	216.4
10	Pneumonitis due to solids and liquids (J69)	33	1.6	178.6
11	Septicemia (A40-A41)	24	1.2	129.9
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	21	1.0	113.6
13	Parkinson's disease (G20-G21)	18	0.9	*
14	Atherosclerosis (I70)	17	0.8	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	0.6	*
15	Aortic aneurysm and dissection (I71)	12	0.6	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	12	0.6	*
...	All other causes (Residual)	209	10.3	1,130.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total deaths		
		Number	Rate	Rate
Hispanic, male, 95-99 years				
...	All causes	725	100.0	12,569.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	275	37.9	4,767.7
2	Malignant neoplasms (C00-C97)	88	12.1	1,525.7
3	Cerebrovascular diseases (I60-I69)	63	8.7	1,092.2
4	Influenza and pneumonia (J10-J18)	62	8.6	1,074.9
5	Chronic lower respiratory diseases (J40-J47)	45	6.2	780.2
6	Diabetes mellitus (E10-E14)	17	2.3	*
7	Alzheimer's disease (G30)	16	2.2	*
8	Septicemia (A40-A41)	11	1.5	*
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11	1.5	*
10	Pneumonitis due to solids and liquids (J69)	10	1.4	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	1.4	*
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	7	1.0	*
12	Atherosclerosis (I70)	7	1.0	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.7	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.4	*
...	All other causes (Residual)	95	13.1	1,647.0
Hispanic, male, 100 years and over				
...	All causes	116	100.0	12,262.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	50	43.1	5,285.4
2	Influenza and pneumonia (J10-J18)	9	7.8	*
3	Cerebrovascular diseases (I60-I69)	7	6.0	*
4	Malignant neoplasms (C00-C97)	5	4.3	*
5	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	4	3.4	*
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	3.4	*
7	Chronic lower respiratory diseases (J40-J47)	3	2.6	*
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3	2.6	*
9	Septicemia (A40-A41)	2	1.7	*
9	Parkinson's disease (G20-G21)	2	1.7	*
9	Alzheimer's disease (G30)	2	1.7	*
12	Nutritional deficiencies (E40-E64)	1	0.9	*
12	Atherosclerosis (I70)	1	0.9	*
...	All other causes (Residual)	23	19.8	2,431.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, all ages				
...	All causes	47,082	100.0	291.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,253	26.0	75.9
2	Malignant neoplasms (C00-C97)	10,022	21.3	62.0
3	Cerebrovascular diseases (I60-I69)	3,322	7.1	20.6
4	Diabetes mellitus (E10-E14)	2,821	6.0	17.5
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,134	4.5	13.2
6	Influenza and pneumonia (J10-J18)	1,322	2.8	8.2
7	Chronic lower respiratory diseases (J40-J47)	1,238	2.6	7.7
8	Certain conditions originating in the perinatal period (P00-P96)	951	2.0	5.9
9	Chronic liver disease and cirrhosis (K70,K73-K74)	875	1.9	5.4
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	841	1.8	5.2
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	755	1.6	4.7
12	Alzheimer's disease (G30)	744	1.6	4.6
13	Septicemia (A40-A41)	601	1.3	3.7
14	Assault (homicide) (X85-Y09,Y87.1)	486	1.0	3.0
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	435	0.9	2.7
...	All other causes (Residual)	8,282	17.6	51.3
Hispanic, female, 1-4 years				
...	All causes	397	100.0	28.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	132	33.2	9.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	51	12.8	3.7
3	Malignant neoplasms (C00-C97)	32	8.1	2.3
4	Assault (homicide) (X85-Y09,Y87.1)	27	6.8	2.0
5	Influenza and pneumonia (J10-J18)	12	3.0	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	8	2.0	*
7	Septicemia (A40-A41)	6	1.5	*
7	Cerebrovascular diseases (I60-I69)	6	1.5	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	1.3	*
9	Certain conditions originating in the perinatal period (P00-P96)	5	1.3	*
11	Chronic lower respiratory diseases (J40-J47)	4	1.0	*
12	Tuberculosis (A16-A19)	2	0.5	*
12	Meningitis (G00,G03)	2	0.5	*
12	Pneumonitis due to solids and liquids (J69)	2	0.5	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.5	*
...	All other causes (Residual)	101	25.4	7.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	
			Rate	
Hispanic, female, 5-9 years				
...	All causes	233	100.0	14.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	100	42.9	6.1
2	Malignant neoplasms (C00-C97)	36	15.5	2.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	17	7.3	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	8	3.4	*
5	Influenza and pneumonia (J10-J18)	6	2.6	*
6	Assault (homicide) (X85-Y09,Y87.1)	5	2.1	*
7	Cerebrovascular diseases (I60-I69)	3	1.3	*
7	Chronic lower respiratory diseases (J40-J47)	3	1.3	*
7	Certain conditions originating in the perinatal period (P00-P96)	3	1.3	*
10	Meningococcal infection (A39)	2	0.9	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.9	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.4	*
12	Anemias (D50-D64)	1	0.4	*
12	Diabetes mellitus (E10-E14)	1	0.4	*
12	Meningitis (G00,G03)	1	0.4	*
12	Pneumonitis due to solids and liquids (J69)	1	0.4	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.4	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.4	*
...	All other causes (Residual)	41	17.6	2.5
Hispanic, female, 10-14 years				
...	All causes	210	100.0	14.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	73	34.8	5.2
2	Malignant neoplasms (C00-C97)	40	19.0	2.8
3	Assault (homicide) (X85-Y09,Y87.1)	15	7.1	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	3.3	*
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	7	3.3	*
6	Influenza and pneumonia (J10-J18)	5	2.4	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	1.9	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	1.9	*
9	Anemias (D50-D64)	3	1.4	*
9	Chronic lower respiratory diseases (J40-J47)	3	1.4	*
11	Septicemia (A40-A41)	2	1.0	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	2	1.0	*
11	Cerebrovascular diseases (I60-I69)	2	1.0	*
14	Tuberculosis (A16-A19)	1	0.5	*
14	Diabetes mellitus (E10-E14)	1	0.5	*
14	Meningitis (G00,G03)	1	0.5	*
14	Diseases of appendix (K35-K38)	1	0.5	*
14	Certain conditions originating in the perinatal period (P00-P96)	1	0.5	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.5	*
...	All other causes (Residual)	37	17.6	2.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total deaths		
		Number	Rate	Rate
	Hispanic, female, 15-19 years			
...	All causes	425	100.0	31.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	195	45.9	14.2
2	Malignant neoplasms (C00-C97)	50	11.8	3.6
3	Assault (homicide) (X85-Y09,Y87.1)	42	9.9	3.1
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	35	8.2	2.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	13	3.1	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	1.2	*
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.9	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	3	0.7	*
8	Cerebrovascular diseases (I60-I69)	3	0.7	*
8	Influenza and pneumonia (J10-J18)	3	0.7	*
11	Aortic aneurysm and dissection (I71)	2	0.5	*
11	Diseases of appendix (K35-K38)	2	0.5	*
11	Pregnancy, childbirth and the puerperium (O00-099)	2	0.5	*
14	Meningococcal infection (A39)	1	0.2	*
14	Septicemia (A40-A41)	1	0.2	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.2	*
14	Anemias (D50-D64)	1	0.2	*
14	Diabetes mellitus (E10-E14)	1	0.2	*
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1	0.2	*
14	Chronic lower respiratory diseases (J40-J47)	1	0.2	*
14	Hernia (K40-K46)	1	0.2	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
...	All other causes (Residual)	57	13.4	4.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 20-24 years				
...	All causes	531	100.0	39.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	183	34.5	13.6
2	Assault (homicide) (X85-Y09,Y87.1)	70	13.2	5.2
3	Malignant neoplasms (C00-C97)	63	11.9	4.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	25	4.7	1.9
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	25	4.7	1.9
6	Pregnancy, childbirth and the puerperium (O00-099)	22	4.1	1.6
7	Cerebrovascular diseases (I60-I69)	11	2.1	*
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	10	1.9	*
9	Influenza and pneumonia (J10-J18)	8	1.5	*
10	Septicemia (A40-A41)	5	0.9	*
10	Diabetes mellitus (E10-E14)	5	0.9	*
12	Anemias (D50-D64)	3	0.6	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.6	*
14	Tuberculosis (A16-A19)	2	0.4	*
14	Meningococcal infection (A39)	2	0.4	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.4	*
14	Aortic aneurysm and dissection (I71)	2	0.4	*
14	Chronic lower respiratory diseases (J40-J47)	2	0.4	*
14	Chronic liver disease and cirrhosis (K70,K73-K74)	2	0.4	*
14	Cholelithiasis and other disorders of gallbladder (K80-K82)	2	0.4	*
...	All other causes (Residual)	84	15.8	6.3
Hispanic, female, 25-29 years				
...	All causes	584	100.0	45.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	140	24.0	11.0
2	Malignant neoplasms (C00-C97)	91	15.6	7.1
3	Assault (homicide) (X85-Y09,Y87.1)	58	9.9	4.5
4	Human immunodeficiency virus (HIV) disease (B20-B24)	34	5.8	2.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	33	5.7	2.6
6	Intentional self-harm (suicide) (X60-X84,Y87.0)	24	4.1	1.9
7	Pregnancy, childbirth and the puerperium (O00-099)	20	3.4	1.6
8	Cerebrovascular diseases (I60-I69)	15	2.6	*
9	Influenza and pneumonia (J10-J18)	9	1.5	*
10	Septicemia (A40-A41)	8	1.4	*
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	1.4	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.0	*
13	Anemias (D50-D64)	5	0.9	*
13	Diabetes mellitus (E10-E14)	5	0.9	*
15	Chronic lower respiratory diseases (J40-J47)	3	0.5	*
...	All other causes (Residual)	125	21.4	9.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	
			Rate	
Hispanic, female, 30-34 years				
...	All causes	725	100.0	55.9
1	Malignant neoplasms (C00-C97)	144	19.9	11.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	137	18.9	10.6
3	Human immunodeficiency virus (HIV) disease (B20-B24)	57	7.9	4.4
4	Assault (homicide) (X85-Y09,Y87.1)	55	7.6	4.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	48	6.6	3.7
6	Cerebrovascular diseases (I60-I69)	28	3.9	2.2
7	Intentional self-harm (suicide) (X60-X84,Y87.0)	26	3.6	2.0
8	Pregnancy, childbirth and the puerperium (O00-099)	16	2.2	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	1.8	*
10	Influenza and pneumonia (J10-J18)	12	1.7	*
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	12	1.7	*
12	Diabetes mellitus (E10-E14)	11	1.5	*
13	Chronic liver disease and cirrhosis (K70,K73-K74)	8	1.1	*
14	Chronic lower respiratory diseases (J40-J47)	5	0.7	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.6	*
...	All other causes (Residual)	149	20.6	11.5
Hispanic, female, 35-39 years				
...	All causes	1,113	100.0	86.0
1	Malignant neoplasms (C00-C97)	311	27.9	24.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	174	15.6	13.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	86	7.7	6.6
4	Human immunodeficiency virus (HIV) disease (B20-B24)	73	6.6	5.6
5	Assault (homicide) (X85-Y09,Y87.1)	49	4.4	3.8
6	Cerebrovascular diseases (I60-I69)	42	3.8	3.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	36	3.2	2.8
7	Intentional self-harm (suicide) (X60-X84,Y87.0)	36	3.2	2.8
9	Diabetes mellitus (E10-E14)	21	1.9	1.6
10	Pregnancy, childbirth and the puerperium (O00-099)	19	1.7	*
11	Septicemia (A40-A41)	12	1.1	*
11	Influenza and pneumonia (J10-J18)	12	1.1	*
13	Chronic lower respiratory diseases (J40-J47)	10	0.9	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	0.9	*
15	Viral hepatitis (B15-B19)	9	0.8	*
...	All other causes (Residual)	213	19.1	16.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total deaths		
		Number	Rate	Rate
Hispanic, female, 40-44 years				
...	All causes	1,375	100.0	122.5
1	Malignant neoplasms (C00-C97)	450	32.7	40.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	127	9.2	11.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	124	9.0	11.0
4	Human immunodeficiency virus (HIV) disease (B20-B24)	86	6.3	7.7
5	Cerebrovascular diseases (I60-I69)	80	5.8	7.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	66	4.8	5.9
7	Assault (homicide) (X85-Y09,Y87.1)	39	2.8	3.5
8	Diabetes mellitus (E10-E14)	38	2.8	3.4
9	Intentional self-harm (suicide) (X60-X84,Y87.0)	29	2.1	2.6
10	Septicemia (A40-A41)	21	1.5	1.9
10	Chronic lower respiratory diseases (J40-J47)	21	1.5	1.9
12	Viral hepatitis (B15-B19)	17	1.2	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	1.2	*
14	Influenza and pneumonia (J10-J18)	14	1.0	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	0.6	*
...	All other causes (Residual)	239	17.4	21.3
Hispanic, female, 45-49 years				
...	All causes	1,633	100.0	183.6
1	Malignant neoplasms (C00-C97)	601	36.8	67.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	170	10.4	19.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	107	6.6	12.0
4	Cerebrovascular diseases (I60-I69)	101	6.2	11.4
5	Diabetes mellitus (E10-E14)	76	4.7	8.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	75	4.6	8.4
7	Human immunodeficiency virus (HIV) disease (B20-B24)	70	4.3	7.9
8	Assault (homicide) (X85-Y09,Y87.1)	37	2.3	4.2
9	Intentional self-harm (suicide) (X60-X84,Y87.0)	29	1.8	3.3
10	Viral hepatitis (B15-B19)	28	1.7	3.1
11	Chronic lower respiratory diseases (J40-J47)	24	1.5	2.7
12	Septicemia (A40-A41)	15	0.9	*
13	Influenza and pneumonia (J10-J18)	13	0.8	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	0.8	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.7	*
...	All other causes (Residual)	263	16.1	29.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total deaths		
		Number	Rate	Rate
Hispanic, female, 50-54 years				
...	All causes	1,915	100.0	274.5
1	Malignant neoplasms (C00-C97)	742	38.7	106.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	277	14.5	39.7
3	Diabetes mellitus (E10-E14)	123	6.4	17.6
4	Cerebrovascular diseases (I60-I69)	96	5.0	13.8
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	96	5.0	13.8
6	Chronic liver disease and cirrhosis (K70,K73-K74)	72	3.8	10.3
7	Human immunodeficiency virus (HIV) disease (B20-B24)	35	1.8	5.0
8	Septicemia (A40-A41)	29	1.5	4.2
9	Chronic lower respiratory diseases (J40-J47)	28	1.5	4.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	27	1.4	3.9
11	Viral hepatitis (B15-B19)	25	1.3	3.6
11	Influenza and pneumonia (J10-J18)	25	1.3	3.6
13	Intentional self-harm (suicide) (X60-X84,Y87.0)	21	1.1	3.0
14	Assault (homicide) (X85-Y09,Y87.1)	16	0.8	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	0.7	*
...	All other causes (Residual)	289	15.1	41.4
Hispanic, female, 55-59 years				
...	All causes	2,220	100.0	422.8
1	Malignant neoplasms (C00-C97)	824	37.1	156.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	403	18.2	76.8
3	Diabetes mellitus (E10-E14)	195	8.8	37.1
4	Cerebrovascular diseases (I60-I69)	99	4.5	18.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	85	3.8	16.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	79	3.6	15.0
7	Chronic lower respiratory diseases (J40-J47)	47	2.1	9.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	42	1.9	8.0
9	Influenza and pneumonia (J10-J18)	32	1.4	6.1
10	Septicemia (A40-A41)	24	1.1	4.6
10	Viral hepatitis (B15-B19)	24	1.1	4.6
12	Human immunodeficiency virus (HIV) disease (B20-B24)	17	0.8	*
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	14	0.6	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	0.6	*
15	Assault (homicide) (X85-Y09,Y87.1)	10	0.5	*
...	All other causes (Residual)	312	14.1	59.4

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 60-64 years				
...	All causes	2,753	100.0	652.1
1	Malignant neoplasms (C00-C97)	922	33.5	218.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	605	22.0	143.3
3	Diabetes mellitus (E10-E14)	235	8.5	55.7
4	Cerebrovascular diseases (I60-I69)	190	6.9	45.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	92	3.3	21.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	75	2.7	17.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	66	2.4	15.6
8	Chronic lower respiratory diseases (J40-J47)	51	1.9	12.1
9	Septicemia (A40-A41)	41	1.5	9.7
10	Influenza and pneumonia (J10-J18)	40	1.5	9.5
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	30	1.1	7.1
12	Viral hepatitis (B15-B19)	29	1.1	6.9
13	Human immunodeficiency virus (HIV) disease (B20-B24)	23	0.8	5.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	15	0.5	*
15	Assault (homicide) (X85-Y09,Y87.1)	10	0.4	*
...	All other causes (Residual)	329	12.0	77.9
Hispanic, female, 65-69 years				
...	All causes	3,708	100.0	1,049.7
1	Malignant neoplasms (C00-C97)	1,134	30.6	321.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	919	24.8	260.2
3	Diabetes mellitus (E10-E14)	346	9.3	97.9
4	Cerebrovascular diseases (I60-I69)	231	6.2	65.4
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	105	2.8	29.7
6	Chronic lower respiratory diseases (J40-J47)	104	2.8	29.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	100	2.7	28.3
8	Influenza and pneumonia (J10-J18)	63	1.7	17.8
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	61	1.6	17.3
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	39	1.1	11.0
11	Septicemia (A40-A41)	37	1.0	10.5
12	Viral hepatitis (B15-B19)	34	0.9	9.6
13	Alzheimer's disease (G30)	27	0.7	7.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	24	0.6	6.8
15	Parkinson's disease (G20-G21)	15	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	15	0.4	*
...	All other causes (Residual)	454	12.2	128.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total deaths		
		Number	Rate	Rate
Hispanic, female, 70-74 years				
...	All causes	4,870	100.0	1,674.8
1	Malignant neoplasms (C00-C97)	1,337	27.5	459.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,289	26.5	443.3
3	Diabetes mellitus (E10-E14)	484	9.9	166.5
4	Cerebrovascular diseases (I60-I69)	343	7.0	118.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	133	2.7	45.7
6	Chronic lower respiratory diseases (J40-J47)	125	2.6	43.0
7	Influenza and pneumonia (J10-J18)	124	2.5	42.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	105	2.2	36.1
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	86	1.8	29.6
10	Septicemia (A40-A41)	72	1.5	24.8
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	51	1.0	17.5
12	Alzheimer's disease (G30)	38	0.8	13.1
13	Viral hepatitis (B15-B19)	34	0.7	11.7
14	Aortic aneurysm and dissection (I71)	32	0.7	11.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	28	0.6	9.6
...	All other causes (Residual)	589	12.1	202.6
Hispanic, female, 75-79 years				
...	All causes	5,509	100.0	2,564.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,671	30.3	778.0
2	Malignant neoplasms (C00-C97)	1,238	22.5	576.4
3	Diabetes mellitus (E10-E14)	455	8.3	211.8
4	Cerebrovascular diseases (I60-I69)	438	8.0	203.9
5	Chronic lower respiratory diseases (J40-J47)	208	3.8	96.8
6	Influenza and pneumonia (J10-J18)	147	2.7	68.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	119	2.2	55.4
8	Alzheimer's disease (G30)	105	1.9	48.9
9	Chronic liver disease and cirrhosis (K70,K73-K74)	94	1.7	43.8
10	Septicemia (A40-A41)	78	1.4	36.3
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	76	1.4	35.4
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	61	1.1	28.4
13	Parkinson's disease (G20-G21)	36	0.7	16.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	31	0.6	14.4
15	Aortic aneurysm and dissection (I71)	26	0.5	12.1
...	All other causes (Residual)	726	13.2	338.0

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total deaths		
		Number	Rate	Rate
Hispanic, female, 80-84 years				
...	All causes	5,469	100.0	4,131.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,871	34.2	1,413.5
2	Malignant neoplasms (C00-C97)	932	17.0	704.1
3	Cerebrovascular diseases (I60-I69)	505	9.2	381.5
4	Diabetes mellitus (E10-E14)	346	6.3	261.4
5	Chronic lower respiratory diseases (J40-J47)	204	3.7	154.1
6	Influenza and pneumonia (J10-J18)	176	3.2	133.0
7	Alzheimer's disease (G30)	130	2.4	98.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	104	1.9	78.6
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	98	1.8	74.0
10	Septicemia (A40-A41)	79	1.4	59.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	66	1.2	49.9
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	64	1.2	48.4
13	Parkinson's disease (G20-G21)	43	0.8	32.5
14	Atherosclerosis (I70)	40	0.7	30.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	38	0.7	28.7
...	All other causes (Residual)	773	14.1	584.0
Hispanic, female, 85-89 years				
...	All causes	5,553	100.0	6,855.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,126	38.3	2,624.5
2	Malignant neoplasms (C00-C97)	660	11.9	814.8
3	Cerebrovascular diseases (I60-I69)	570	10.3	703.7
4	Influenza and pneumonia (J10-J18)	276	5.0	340.7
5	Diabetes mellitus (E10-E14)	266	4.8	328.4
6	Chronic lower respiratory diseases (J40-J47)	210	3.8	259.2
7	Alzheimer's disease (G30)	205	3.7	253.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	103	1.9	127.2
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	84	1.5	103.7
10	Septicemia (A40-A41)	78	1.4	96.3
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	77	1.4	95.1
12	Atherosclerosis (I70)	56	1.0	69.1
13	Pneumonitis due to solids and liquids (J69)	46	0.8	56.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	37	0.7	45.7
15	Chronic liver disease and cirrhosis (K70,K73-K74)	34	0.6	42.0
...	All other causes (Residual)	725	13.1	895.0

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Hispanic, female, 90-94 years				
...	All causes	3,818	100.0	9,915.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,645	43.1	4,272.1
2	Cerebrovascular diseases (I60-I69)	380	10.0	986.9
3	Malignant neoplasms (C00-C97)	294	7.7	763.5
4	Influenza and pneumonia (J10-J18)	195	5.1	506.4
5	Alzheimer's disease (G30)	161	4.2	418.1
6	Diabetes mellitus (E10-E14)	153	4.0	397.3
7	Chronic lower respiratory diseases (J40-J47)	122	3.2	316.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	75	2.0	194.8
9	Atherosclerosis (I70)	60	1.6	155.8
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	52	1.4	135.0
11	Septicemia (A40-A41)	46	1.2	119.5
12	Pneumonitis due to solids and liquids (J69)	45	1.2	116.9
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	34	0.9	88.3
14	Parkinson's disease (G20-G21)	24	0.6	62.3
15	Nutritional deficiencies (E40-E64)	20	0.5	51.9
...	All other causes (Residual)	512	13.4	1,329.7
Hispanic, female, 95-99 years				
...	All causes	1,609	100.0	12,077.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	720	44.7	5,404.6
2	Cerebrovascular diseases (I60-I69)	146	9.1	1,095.9
3	Influenza and pneumonia (J10-J18)	99	6.2	743.1
4	Malignant neoplasms (C00-C97)	95	5.9	713.1
5	Alzheimer's disease (G30)	60	3.7	450.4
6	Diabetes mellitus (E10-E14)	49	3.0	367.8
6	Chronic lower respiratory diseases (J40-J47)	49	3.0	367.8
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	31	1.9	232.7
9	Pneumonitis due to solids and liquids (J69)	28	1.7	210.2
10	Atherosclerosis (I70)	25	1.6	187.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21	1.3	157.6
12	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19	1.2	*
13	Septicemia (A40-A41)	13	0.8	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.4	*
14	Nutritional deficiencies (E40-E64)	7	0.4	*
...	All other causes (Residual)	240	14.9	1,801.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total deaths		
		Number	Rate	Rate
Hispanic, female, 100 years and over				
...	All causes	345	100.0	14,156.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	165	47.8	6,770.6
2	Cerebrovascular diseases (I60-I69)	25	7.2	1,025.9
3	Influenza and pneumonia (J10-J18)	21	6.1	861.7
4	Malignant neoplasms (C00-C97)	19	5.5	*
5	Chronic lower respiratory diseases (J40-J47)	11	3.2	*
6	Diabetes mellitus (E10-E14)	10	2.9	*
7	Atherosclerosis (I70)	7	2.0	*
8	Alzheimer's disease (G30)	6	1.7	*
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5	1.4	*
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	4	1.2	*
10	Pneumonitis due to solids and liquids (J69)	4	1.2	*
12	Septicemia (A40-A41)	3	0.9	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.6	*
13	Nutritional deficiencies (E40-E64)	2	0.6	*
13	Peptic ulcer (K25-K28)	2	0.6	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.6	*
...	All other causes (Residual)	57	16.5	2,338.9
White, non-Hispanic, both sexes, all ages				
...	All causes	1,959,919	100.0	996.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	594,465	30.3	302.3
2	Malignant neoplasms (C00-C97)	458,200	23.4	233.0
3	Cerebrovascular diseases (I60-I69)	138,210	7.1	70.3
4	Chronic lower respiratory diseases (J40-J47)	109,901	5.6	55.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	73,536	3.8	37.4
6	Influenza and pneumonia (J10-J18)	55,135	2.8	28.0
7	Diabetes mellitus (E10-E14)	50,163	2.6	25.5
8	Alzheimer's disease (G30)	45,246	2.3	23.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	27,924	1.4	14.2
10	Intentional self-harm (suicide) (X60-X84,Y87.0)	24,616	1.3	12.5
11	Septicemia (A40-A41)	23,517	1.2	12.0
12	Chronic liver disease and cirrhosis (K70,K73-K74)	19,739	1.0	10.0
13	Parkinson's disease (G20-G21)	14,497	0.7	7.4
14	Pneumonitis due to solids and liquids (J69)	14,343	0.7	7.3
15	Aortic aneurysm and dissection (I71)	13,722	0.7	7.0
...	All other causes (Residual)	296,705	15.1	150.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total deaths		
		Number	Rate	Rate
White, non-Hispanic, both sexes, 1-4 years				
...	All causes	2,641	100.0	27.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,033	39.1	10.9
2	Malignant neoplasms (C00-C97)	256	9.7	2.7
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	240	9.1	2.5
4	Assault (homicide) (X85-Y09,Y87.1)	144	5.5	1.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	98	3.7	1.0
6	Septicemia (A40-A41)	51	1.9	0.5
6	Influenza and pneumonia (J10-J18)	51	1.9	0.5
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	37	1.4	0.4
9	Certain conditions originating in the perinatal period (P00-P96)	34	1.3	0.4
10	Cerebrovascular diseases (I60-I69)	25	0.9	0.3
11	Meningitis (G00,G03)	19	0.7	*
11	Chronic lower respiratory diseases (J40-J47)	19	0.7	*
13	Anemias (D50-D64)	14	0.5	*
14	Meningococcal infection (A39)	11	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	8	0.3	*
...	All other causes (Residual)	601	22.8	6.4
White, non-Hispanic, both sexes, 5-9 years				
...	All causes	1,850	100.0	14.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	796	43.0	6.4
2	Malignant neoplasms (C00-C97)	307	16.6	2.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	113	6.1	0.9
4	Assault (homicide) (X85-Y09,Y87.1)	72	3.9	0.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	54	2.9	0.4
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	41	2.2	0.3
7	Septicemia (A40-A41)	25	1.4	0.2
8	Influenza and pneumonia (J10-J18)	23	1.2	0.2
9	Cerebrovascular diseases (I60-I69)	17	0.9	*
9	Chronic lower respiratory diseases (J40-J47)	17	0.9	*
11	Certain conditions originating in the perinatal period (P00-P96)	11	0.6	*
12	Meningitis (G00,G03)	9	0.5	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	9	0.5	*
14	Anemias (D50-D64)	7	0.4	*
14	Pneumonitis due to solids and liquids (J69)	7	0.4	*
...	All other causes (Residual)	342	18.5	2.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total		
		Number	deaths	Rate
White, non-Hispanic, both sexes, 10-14 years				
...	All causes	2,547	100.0	19.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,047	41.1	8.0
2	Malignant neoplasms (C00-C97)	320	12.6	2.5
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	220	8.6	1.7
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	127	5.0	1.0
5	Assault (homicide) (X85-Y09,Y87.1)	92	3.6	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	86	3.4	0.7
7	Chronic lower respiratory diseases (J40-J47)	30	1.2	0.2
8	Cerebrovascular diseases (I60-I69)	29	1.1	0.2
9	Influenza and pneumonia (J10-J18)	26	1.0	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	22	0.9	0.2
11	Septicemia (A40-A41)	17	0.7	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	0.4	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	10	0.4	*
14	Diabetes mellitus (E10-E14)	9	0.4	*
14	Meningitis (G00,G03)	9	0.4	*
14	Certain conditions originating in the perinatal period (P00-P96)	9	0.4	*
...	All other causes (Residual)	483	19.0	3.7
White, non-Hispanic, both sexes, 15-19 years				
...	All causes	8,327	100.0	63.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,852	58.3	36.9
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,175	14.1	8.9
3	Malignant neoplasms (C00-C97)	452	5.4	3.4
4	Assault (homicide) (X85-Y09,Y87.1)	352	4.2	2.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	216	2.6	1.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	134	1.6	1.0
7	Cerebrovascular diseases (I60-I69)	46	0.6	0.3
8	Influenza and pneumonia (J10-J18)	43	0.5	0.3
9	Chronic lower respiratory diseases (J40-J47)	34	0.4	0.3
10	Diabetes mellitus (E10-E14)	28	0.3	0.2
11	Meningococcal infection (A39)	27	0.3	0.2
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	26	0.3	0.2
13	Septicemia (A40-A41)	21	0.3	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	0.1	*
15	Anemias (D50-D64)	9	0.1	*
15	Pneumonitis due to solids and liquids (J69)	9	0.1	*
15	Pregnancy, childbirth and the puerperium (O00-099)	9	0.1	*
...	All other causes (Residual)	882	10.6	6.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total deaths		
		Number	Rate	Rate
White, non-Hispanic, both sexes, 20-24 years				
...	All causes	9,934	100.0	81.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,883	49.2	40.2
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,693	17.0	13.9
3	Malignant neoplasms (C00-C97)	607	6.1	5.0
4	Assault (homicide) (X85-Y09,Y87.1)	522	5.3	4.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	315	3.2	2.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	145	1.5	1.2
7	Influenza and pneumonia (J10-J18)	73	0.7	0.6
8	Cerebrovascular diseases (I60-I69)	72	0.7	0.6
9	Diabetes mellitus (E10-E14)	64	0.6	0.5
10	Chronic lower respiratory diseases (J40-J47)	53	0.5	0.4
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	37	0.4	0.3
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	36	0.4	0.3
13	Septicemia (A40-A41)	27	0.3	0.2
14	Human immunodeficiency virus (HIV) disease (B20-B24)	24	0.2	0.2
14	Pregnancy, childbirth and the puerperium (O00-099)	24	0.2	0.2
...	All other causes (Residual)	1,359	13.7	11.2
White, non-Hispanic, both sexes, 25-29 years				
...	All causes	9,822	100.0	83.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,788	38.6	32.2
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,709	17.4	14.5
3	Malignant neoplasms (C00-C97)	884	9.0	7.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	523	5.3	4.4
5	Assault (homicide) (X85-Y09,Y87.1)	490	5.0	4.2
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	144	1.5	1.2
7	Human immunodeficiency virus (HIV) disease (B20-B24)	139	1.4	1.2
8	Cerebrovascular diseases (I60-I69)	123	1.3	1.0
9	Diabetes mellitus (E10-E14)	107	1.1	0.9
10	Influenza and pneumonia (J10-J18)	79	0.8	0.7
11	Chronic lower respiratory diseases (J40-J47)	59	0.6	0.5
12	Septicemia (A40-A41)	44	0.4	0.4
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	43	0.4	0.4
14	Chronic liver disease and cirrhosis (K70,K73-K74)	42	0.4	0.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	40	0.4	0.3
...	All other causes (Residual)	1,608	16.4	13.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total deaths		
		Number	Rate	Rate
White, non-Hispanic, both sexes, 30-34 years				
...	All causes	13,369	100.0	100.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,942	29.5	29.7
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,949	14.6	14.7
3	Malignant neoplasms (C00-C97)	1,715	12.8	12.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,130	8.5	8.5
5	Assault (homicide) (X85-Y09,Y87.1)	533	4.0	4.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	480	3.6	3.6
7	Diabetes mellitus (E10-E14)	250	1.9	1.9
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	200	1.5	1.5
9	Cerebrovascular diseases (I60-I69)	190	1.4	1.4
10	Chronic liver disease and cirrhosis (K70,K73-K74)	175	1.3	1.3
11	Influenza and pneumonia (J10-J18)	134	1.0	1.0
12	Septicemia (A40-A41)	90	0.7	0.7
13	Chronic lower respiratory diseases (J40-J47)	85	0.6	0.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	67	0.5	0.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	65	0.5	0.5
...	All other causes (Residual)	2,364	17.7	17.8
White, non-Hispanic, both sexes, 35-39 years				
...	All causes	22,912	100.0	145.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,302	23.1	33.6
2	Malignant neoplasms (C00-C97)	3,802	16.6	24.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,789	12.2	17.7
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	2,599	11.3	16.5
5	Human immunodeficiency virus (HIV) disease (B20-B24)	1,007	4.4	6.4
6	Chronic liver disease and cirrhosis (K70,K73-K74)	702	3.1	4.4
7	Assault (homicide) (X85-Y09,Y87.1)	656	2.9	4.2
8	Cerebrovascular diseases (I60-I69)	464	2.0	2.9
9	Diabetes mellitus (E10-E14)	441	1.9	2.8
10	Influenza and pneumonia (J10-J18)	268	1.2	1.7
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	201	0.9	1.3
12	Septicemia (A40-A41)	175	0.8	1.1
13	Chronic lower respiratory diseases (J40-J47)	174	0.8	1.1
14	Viral hepatitis (B15-B19)	142	0.6	0.9
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	126	0.5	0.8
...	All other causes (Residual)	4,064	17.7	25.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total deaths		
		Number	Rate	Rate
White, non-Hispanic, both sexes, 40-44 years				
...	All causes	35,319	100.0	212.4
1	Malignant neoplasms (C00-C97)	7,656	21.7	46.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,954	16.9	35.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,706	16.2	34.3
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	2,941	8.3	17.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,538	4.4	9.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	993	2.8	6.0
7	Cerebrovascular diseases (I60-I69)	914	2.6	5.5
8	Diabetes mellitus (E10-E14)	819	2.3	4.9
9	Assault (homicide) (X85-Y09,Y87.1)	617	1.7	3.7
10	Influenza and pneumonia (J10-J18)	410	1.2	2.5
11	Chronic lower respiratory diseases (J40-J47)	396	1.1	2.4
12	Viral hepatitis (B15-B19)	359	1.0	2.2
13	Septicemia (A40-A41)	323	0.9	1.9
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	233	0.7	1.4
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	218	0.6	1.3
...	All other causes (Residual)	6,242	17.7	37.5
White, non-Hispanic, both sexes, 45-49 years				
...	All causes	48,005	100.0	319.5
1	Malignant neoplasms (C00-C97)	13,598	28.3	90.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,905	20.6	65.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,978	10.4	33.1
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	2,611	5.4	17.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,365	4.9	15.7
6	Cerebrovascular diseases (I60-I69)	1,380	2.9	9.2
7	Diabetes mellitus (E10-E14)	1,175	2.4	7.8
8	Chronic lower respiratory diseases (J40-J47)	830	1.7	5.5
9	Human immunodeficiency virus (HIV) disease (B20-B24)	826	1.7	5.5
10	Viral hepatitis (B15-B19)	707	1.5	4.7
11	Influenza and pneumonia (J10-J18)	501	1.0	3.3
12	Assault (homicide) (X85-Y09,Y87.1)	479	1.0	3.2
13	Septicemia (A40-A41)	445	0.9	3.0
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	334	0.7	2.2
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	224	0.5	1.5
...	All other causes (Residual)	7,647	15.9	50.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total deaths		
		Number	Rate	Rate
White, non-Hispanic, both sexes, 50-54 years				
...	All causes	63,959	100.0	475.1
1	Malignant neoplasms (C00-C97)	22,263	34.8	165.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,248	23.8	113.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,904	6.1	29.0
4	Chronic liver disease and cirrhosis (K70,K73-K74)	2,272	3.6	16.9
5	Intentional self-harm (suicide) (X60-X84,Y87.0)	2,163	3.4	16.1
6	Cerebrovascular diseases (I60-I69)	2,053	3.2	15.3
7	Diabetes mellitus (E10-E14)	1,903	3.0	14.1
8	Chronic lower respiratory diseases (J40-J47)	1,633	2.6	12.1
9	Influenza and pneumonia (J10-J18)	664	1.0	4.9
10	Septicemia (A40-A41)	647	1.0	4.8
11	Viral hepatitis (B15-B19)	536	0.8	4.0
12	Human immunodeficiency virus (HIV) disease (B20-B24)	499	0.8	3.7
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	495	0.8	3.7
14	Assault (homicide) (X85-Y09,Y87.1)	355	0.6	2.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	310	0.5	2.3
...	All other causes (Residual)	9,014	14.1	67.0
White, non-Hispanic, both sexes, 55-59 years				
...	All causes	79,771	100.0	756.4
1	Malignant neoplasms (C00-C97)	30,664	38.4	290.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,515	25.7	194.5
3	Chronic lower respiratory diseases (J40-J47)	3,369	4.2	31.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,044	3.8	28.9
5	Cerebrovascular diseases (I60-I69)	2,657	3.3	25.2
6	Diabetes mellitus (E10-E14)	2,526	3.2	24.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,121	2.7	20.1
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,536	1.9	14.6
9	Influenza and pneumonia (J10-J18)	844	1.1	8.0
10	Septicemia (A40-A41)	836	1.0	7.9
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	704	0.9	6.7
12	Aortic aneurysm and dissection (I71)	451	0.6	4.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	423	0.5	4.0
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	307	0.4	2.9
15	Human immunodeficiency virus (HIV) disease (B20-B24)	291	0.4	2.8
...	All other causes (Residual)	9,483	11.9	89.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total deaths		
		Number	Rate	Rate
White, non-Hispanic, both sexes, 60-64 years				
...	All causes	102,577	100.0	1,214.2
1	Malignant neoplasms (C00-C97)	40,201	39.2	475.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	26,777	26.1	317.0
3	Chronic lower respiratory diseases (J40-J47)	5,884	5.7	69.6
4	Cerebrovascular diseases (I60-I69)	3,898	3.8	46.1
5	Diabetes mellitus (E10-E14)	3,397	3.3	40.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,709	2.6	32.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,114	2.1	25.0
8	Influenza and pneumonia (J10-J18)	1,217	1.2	14.4
9	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,123	1.1	13.3
10	Septicemia (A40-A41)	1,083	1.1	12.8
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,072	1.0	12.7
12	Aortic aneurysm and dissection (I71)	750	0.7	8.9
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	581	0.6	6.9
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	429	0.4	5.1
15	Alzheimer's disease (G30)	301	0.3	3.6
...	All other causes (Residual)	11,041	10.8	130.7
White, non-Hispanic, both sexes, 65-69 years				
...	All causes	143,765	100.0	1,899.4
1	Malignant neoplasms (C00-C97)	54,052	37.6	714.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	38,325	26.7	506.3
3	Chronic lower respiratory diseases (J40-J47)	10,523	7.3	139.0
4	Cerebrovascular diseases (I60-I69)	6,277	4.4	82.9
5	Diabetes mellitus (E10-E14)	4,714	3.3	62.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,659	1.8	35.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,093	1.5	27.7
8	Influenza and pneumonia (J10-J18)	1,884	1.3	24.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,776	1.2	23.5
10	Septicemia (A40-A41)	1,610	1.1	21.3
11	Aortic aneurysm and dissection (I71)	1,335	0.9	17.6
12	Intentional self-harm (suicide) (X60-X84,Y87.0)	969	0.7	12.8
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	847	0.6	11.2
14	Alzheimer's disease (G30)	751	0.5	9.9
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	644	0.4	8.5
...	All other causes (Residual)	15,306	10.6	202.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total deaths		
		Number	Rate	Rate
White, non-Hispanic, both sexes, 70-74 years				
...	All causes	213,104	100.0	2,942.3
1	Malignant neoplasms (C00-C97)	70,790	33.2	977.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	59,423	27.9	820.5
3	Chronic lower respiratory diseases (J40-J47)	17,599	8.3	243.0
4	Cerebrovascular diseases (I60-I69)	11,712	5.5	161.7
5	Diabetes mellitus (E10-E14)	6,761	3.2	93.3
6	Influenza and pneumonia (J10-J18)	3,662	1.7	50.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,522	1.7	48.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,965	1.4	40.9
9	Septicemia (A40-A41)	2,563	1.2	35.4
10	Alzheimer's disease (G30)	2,253	1.1	31.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	2,192	1.0	30.3
12	Aortic aneurysm and dissection (I71)	2,191	1.0	30.3
13	Parkinson's disease (G20-G21)	1,345	0.6	18.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,297	0.6	17.9
15	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,106	0.5	15.3
...	All other causes (Residual)	23,723	11.1	327.5
White, non-Hispanic, both sexes, 75-79 years				
...	All causes	287,117	100.0	4,553.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	85,862	29.9	1,361.6
2	Malignant neoplasms (C00-C97)	77,899	27.1	1,235.3
3	Chronic lower respiratory diseases (J40-J47)	22,319	7.8	353.9
4	Cerebrovascular diseases (I60-I69)	20,501	7.1	325.1
5	Diabetes mellitus (E10-E14)	8,725	3.0	138.4
6	Influenza and pneumonia (J10-J18)	6,696	2.3	106.2
7	Alzheimer's disease (G30)	5,676	2.0	90.0
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,834	1.7	76.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,616	1.6	73.2
10	Septicemia (A40-A41)	3,690	1.3	58.5
11	Parkinson's disease (G20-G21)	3,070	1.1	48.7
12	Aortic aneurysm and dissection (I71)	2,705	0.9	42.9
13	Chronic liver disease and cirrhosis (K70,K73-K74)	2,008	0.7	31.8
14	Pneumonitis due to solids and liquids (J69)	1,934	0.7	30.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,894	0.7	30.0
...	All other causes (Residual)	34,688	12.1	550.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total		
		Number	deaths	Rate
White, non-Hispanic, both sexes, 80-84 years				
...	All causes	317,541	100.0	7,473.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	103,504	32.6	2,436.0
2	Malignant neoplasms (C00-C97)	65,180	20.5	1,534.1
3	Cerebrovascular diseases (I60-I69)	28,055	8.8	660.3
4	Chronic lower respiratory diseases (J40-J47)	21,705	6.8	510.8
5	Influenza and pneumonia (J10-J18)	10,372	3.3	244.1
6	Alzheimer's disease (G30)	10,028	3.2	236.0
7	Diabetes mellitus (E10-E14)	8,315	2.6	195.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,522	1.7	130.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,410	1.7	127.3
10	Septicemia (A40-A41)	4,211	1.3	99.1
11	Parkinson's disease (G20-G21)	4,036	1.3	95.0
12	Pneumonitis due to solids and liquids (J69)	2,859	0.9	67.3
13	Aortic aneurysm and dissection (I71)	2,545	0.8	59.9
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,454	0.8	57.8
15	Atherosclerosis (I70)	2,303	0.7	54.2
...	All other causes (Residual)	41,042	12.9	966.0
White, non-Hispanic, both sexes, 85-89 years				
...	All causes	297,730	100.0	12,734.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	107,375	36.1	4,592.5
2	Malignant neoplasms (C00-C97)	43,210	14.5	1,848.1
3	Cerebrovascular diseases (I60-I69)	30,109	10.1	1,287.8
4	Chronic lower respiratory diseases (J40-J47)	15,765	5.3	674.3
5	Alzheimer's disease (G30)	12,242	4.1	523.6
6	Influenza and pneumonia (J10-J18)	12,218	4.1	522.6
7	Diabetes mellitus (E10-E14)	6,472	2.2	276.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,410	1.8	231.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,202	1.7	222.5
10	Septicemia (A40-A41)	4,075	1.4	174.3
11	Pneumonitis due to solids and liquids (J69)	3,437	1.2	147.0
12	Parkinson's disease (G20-G21)	3,332	1.1	142.5
13	Atherosclerosis (I70)	2,892	1.0	123.7
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,644	0.9	113.1
15	Aortic aneurysm and dissection (I71)	1,948	0.7	83.3
...	All other causes (Residual)	41,399	13.9	1,770.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total deaths		
		Number	Rate	Rate
White, non-Hispanic, both sexes, 90-94 years				
...	All causes	195,687	100.0	19,501.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	77,430	39.6	7,716.6
2	Cerebrovascular diseases (I60-I69)	20,776	10.6	2,070.5
3	Malignant neoplasms (C00-C97)	18,871	9.6	1,880.7
4	Influenza and pneumonia (J10-J18)	10,117	5.2	1,008.2
5	Alzheimer's disease (G30)	9,507	4.9	947.5
6	Chronic lower respiratory diseases (J40-J47)	7,125	3.6	710.1
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,594	1.8	358.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,318	1.7	330.7
9	Diabetes mellitus (E10-E14)	3,309	1.7	329.8
10	Atherosclerosis (I70)	2,610	1.3	260.1
11	Pneumonitis due to solids and liquids (J69)	2,482	1.3	247.4
12	Septicemia (A40-A41)	2,468	1.3	246.0
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,992	1.0	198.5
14	Parkinson's disease (G20-G21)	1,515	0.8	151.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,033	0.5	102.9
...	All other causes (Residual)	29,540	15.1	2,943.9
White, non-Hispanic, both sexes, 95-99 years				
...	All causes	74,671	100.0	25,154.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	31,597	42.3	10,644.0
2	Cerebrovascular diseases (I60-I69)	7,491	10.0	2,523.5
3	Malignant neoplasms (C00-C97)	4,800	6.4	1,617.0
4	Influenza and pneumonia (J10-J18)	4,554	6.1	1,534.1
5	Alzheimer's disease (G30)	3,681	4.9	1,240.0
6	Chronic lower respiratory diseases (J40-J47)	1,970	2.6	663.6
7	Atherosclerosis (I70)	1,394	1.9	469.6
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,284	1.7	432.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,223	1.6	412.0
10	Diabetes mellitus (E10-E14)	998	1.3	336.2
11	Pneumonitis due to solids and liquids (J69)	985	1.3	331.8
12	Septicemia (A40-A41)	848	1.1	285.7
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	818	1.1	275.6
14	Parkinson's disease (G20-G21)	345	0.5	116.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	279	0.4	94.0
...	All other causes (Residual)	12,404	16.6	4,178.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total deaths		
		Number	Rate	Rate
White, non-Hispanic, both sexes, 100 years and over				
...	All causes	15,669	100.0	31,226.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,052	45.0	14,053.7
2	Cerebrovascular diseases (I60-I69)	1,362	8.7	2,714.3
3	Influenza and pneumonia (J10-J18)	1,184	7.6	2,359.6
4	Alzheimer's disease (G30)	622	4.0	1,239.6
5	Malignant neoplasms (C00-C97)	606	3.9	1,207.7
6	Atherosclerosis (I70)	374	2.4	745.3
7	Chronic lower respiratory diseases (J40-J47)	295	1.9	587.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	264	1.7	526.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	210	1.3	418.5
10	Pneumonitis due to solids and liquids (J69)	207	1.3	412.5
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	187	1.2	372.7
12	Septicemia (A40-A41)	159	1.0	316.9
13	Diabetes mellitus (E10-E14)	138	0.9	275.0
14	Nutritional deficiencies (E40-E64)	68	0.4	135.5
15	Anemias (D50-D64)	67	0.4	133.5
...	All other causes (Residual)	2,874	18.3	5,727.5
White, non-Hispanic, male, all ages				
...	All causes	944,781	100.0	980.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	287,210	30.4	298.2
2	Malignant neoplasms (C00-C97)	235,932	25.0	245.0
3	Chronic lower respiratory diseases (J40-J47)	53,231	5.6	55.3
4	Cerebrovascular diseases (I60-I69)	52,000	5.5	54.0
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	46,470	4.9	48.2
6	Influenza and pneumonia (J10-J18)	23,609	2.5	24.5
7	Diabetes mellitus (E10-E14)	23,465	2.5	24.4
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	19,711	2.1	20.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13,567	1.4	14.1
10	Alzheimer's disease (G30)	13,123	1.4	13.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	12,640	1.3	13.1
12	Septicemia (A40-A41)	10,056	1.1	10.4
13	Aortic aneurysm and dissection (I71)	8,488	0.9	8.8
14	Parkinson's disease (G20-G21)	8,116	0.9	8.4
15	Pneumonitis due to solids and liquids (J69)	7,214	0.8	7.5
...	All other causes (Residual)	129,949	13.8	134.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total deaths		
		Number	Rate	Rate
White, non-Hispanic, male, 1-4 years				
...	All causes	1,541	100.0	31.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	638	41.4	13.2
2	Malignant neoplasms (C00-C97)	146	9.5	3.0
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	124	8.0	2.6
4	Assault (homicide) (X85-Y09,Y87.1)	86	5.6	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	59	3.8	1.2
6	Septicemia (A40-A41)	26	1.7	0.5
7	Influenza and pneumonia (J10-J18)	24	1.6	0.5
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	1.5	0.5
8	Certain conditions originating in the perinatal period (P00-P96)	23	1.5	0.5
10	Cerebrovascular diseases (I60-I69)	14	0.9	*
11	Meningitis (G00,G03)	12	0.8	*
12	Anemias (D50-D64)	8	0.5	*
12	Chronic lower respiratory diseases (J40-J47)	8	0.5	*
14	Meningococcal infection (A39)	6	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.4	*
...	All other causes (Residual)	338	21.9	7.0
White, non-Hispanic, male, 5-9 years				
...	All causes	1,077	100.0	16.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	487	45.2	7.6
2	Malignant neoplasms (C00-C97)	161	14.9	2.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	55	5.1	0.9
4	Assault (homicide) (X85-Y09,Y87.1)	39	3.6	0.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	34	3.2	0.5
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	1.9	0.3
7	Septicemia (A40-A41)	16	1.5	*
8	Influenza and pneumonia (J10-J18)	15	1.4	*
9	Cerebrovascular diseases (I60-I69)	11	1.0	*
10	Chronic lower respiratory diseases (J40-J47)	10	0.9	*
11	Anemias (D50-D64)	6	0.6	*
11	Certain conditions originating in the perinatal period (P00-P96)	6	0.6	*
13	Meningitis (G00,G03)	5	0.5	*
13	Pneumonitis due to solids and liquids (J69)	5	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.5	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.5	*
...	All other causes (Residual)	197	18.3	3.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total		
		Number	deaths	Rate
White, non-Hispanic, male, 10-14 years				
...	All causes	1,575	100.0	23.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	689	43.7	10.3
2	Malignant neoplasms (C00-C97)	177	11.2	2.6
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	175	11.1	2.6
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	64	4.1	1.0
5	Assault (homicide) (X85-Y09,Y87.1)	54	3.4	0.8
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	53	3.4	0.8
7	Influenza and pneumonia (J10-J18)	14	0.9	*
7	Chronic lower respiratory diseases (J40-J47)	14	0.9	*
9	Cerebrovascular diseases (I60-I69)	12	0.8	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.7	*
11	Septicemia (A40-A41)	9	0.6	*
12	Certain conditions originating in the perinatal period (P00-P96)	8	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.4	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.3	*
14	Diabetes mellitus (E10-E14)	4	0.3	*
14	Meningitis (G00,G03)	4	0.3	*
...	All other causes (Residual)	276	17.5	4.1
White, non-Hispanic, male, 15-19 years				
...	All causes	5,726	100.0	84.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,326	58.1	49.1
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	987	17.2	14.6
3	Malignant neoplasms (C00-C97)	270	4.7	4.0
4	Assault (homicide) (X85-Y09,Y87.1)	231	4.0	3.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	126	2.2	1.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	74	1.3	1.1
7	Cerebrovascular diseases (I60-I69)	31	0.5	0.5
8	Influenza and pneumonia (J10-J18)	25	0.4	0.4
9	Chronic lower respiratory diseases (J40-J47)	21	0.4	0.3
10	Meningococcal infection (A39)	18	0.3	*
10	Diabetes mellitus (E10-E14)	18	0.3	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	0.3	*
13	Septicemia (A40-A41)	13	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.1	*
14	Certain conditions originating in the perinatal period (P00-P96)	7	0.1	*
...	All other causes (Residual)	556	9.7	8.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total deaths		
		Number	Rate	Rate
White, non-Hispanic, male, 20-24 years				
...	All causes	7,340	100.0	117.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,739	50.9	60.1
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,480	20.2	23.8
3	Malignant neoplasms (C00-C97)	366	5.0	5.9
4	Assault (homicide) (X85-Y09,Y87.1)	357	4.9	5.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	200	2.7	3.2
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	89	1.2	1.4
7	Influenza and pneumonia (J10-J18)	43	0.6	0.7
8	Diabetes mellitus (E10-E14)	39	0.5	0.6
9	Cerebrovascular diseases (I60-I69)	36	0.5	0.6
10	Chronic lower respiratory diseases (J40-J47)	31	0.4	0.5
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	24	0.3	0.4
12	Septicemia (A40-A41)	20	0.3	0.3
13	Legal intervention (Y35,Y89.0)	19	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18	0.2	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	16	0.2	*
...	All other causes (Residual)	863	11.8	13.9
White, non-Hispanic, male, 25-29 years				
...	All causes	6,920	100.0	117.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,899	41.9	49.2
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,424	20.6	24.2
3	Malignant neoplasms (C00-C97)	452	6.5	7.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	336	4.9	5.7
5	Assault (homicide) (X85-Y09,Y87.1)	321	4.6	5.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	95	1.4	1.6
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	82	1.2	1.4
8	Cerebrovascular diseases (I60-I69)	67	1.0	1.1
9	Diabetes mellitus (E10-E14)	60	0.9	1.0
10	Influenza and pneumonia (J10-J18)	41	0.6	0.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	33	0.5	0.6
12	Chronic lower respiratory diseases (J40-J47)	31	0.4	0.5
13	Legal intervention (Y35,Y89.0)	29	0.4	0.5
14	Septicemia (A40-A41)	25	0.4	0.4
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	24	0.3	0.4
...	All other causes (Residual)	1,001	14.5	17.0

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total deaths		
		Number	Rate	Rate
White, non-Hispanic, male, 30-34 years				
...	All causes	9,014	100.0	136.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,962	32.9	44.7
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,565	17.4	23.6
3	Malignant neoplasms (C00-C97)	807	9.0	12.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	779	8.6	11.8
5	Human immunodeficiency virus (HIV) disease (B20-B24)	380	4.2	5.7
6	Assault (homicide) (X85-Y09,Y87.1)	356	3.9	5.4
7	Diabetes mellitus (E10-E14)	160	1.8	2.4
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	121	1.3	1.8
9	Chronic liver disease and cirrhosis (K70,K73-K74)	107	1.2	1.6
10	Cerebrovascular diseases (I60-I69)	94	1.0	1.4
11	Influenza and pneumonia (J10-J18)	70	0.8	1.1
12	Septicemia (A40-A41)	49	0.5	0.7
13	Chronic lower respiratory diseases (J40-J47)	40	0.4	0.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	38	0.4	0.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	33	0.4	0.5
...	All other causes (Residual)	1,453	16.1	22.0
White, non-Hispanic, male, 35-39 years				
...	All causes	14,922	100.0	188.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,914	26.2	49.5
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	2,038	13.7	25.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,979	13.3	25.0
4	Malignant neoplasms (C00-C97)	1,705	11.4	21.6
5	Human immunodeficiency virus (HIV) disease (B20-B24)	844	5.7	10.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	453	3.0	5.7
7	Assault (homicide) (X85-Y09,Y87.1)	407	2.7	5.2
8	Diabetes mellitus (E10-E14)	253	1.7	3.2
9	Cerebrovascular diseases (I60-I69)	229	1.5	2.9
10	Influenza and pneumonia (J10-J18)	137	0.9	1.7
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	112	0.8	1.4
12	Viral hepatitis (B15-B19)	94	0.6	1.2
13	Septicemia (A40-A41)	93	0.6	1.2
14	Chronic lower respiratory diseases (J40-J47)	74	0.5	0.9
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	71	0.5	0.9
...	All other causes (Residual)	2,519	16.9	31.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total deaths		
		Number	Rate	Rate
White, non-Hispanic, male, 40-44 years				
...	All causes	22,700	100.0	272.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,467	19.7	53.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,199	18.5	50.4
3	Malignant neoplasms (C00-C97)	3,496	15.4	42.0
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	2,249	9.9	27.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,065	4.7	12.8
6	Human immunodeficiency virus (HIV) disease (B20-B24)	834	3.7	10.0
7	Diabetes mellitus (E10-E14)	513	2.3	6.2
8	Cerebrovascular diseases (I60-I69)	465	2.0	5.6
9	Assault (homicide) (X85-Y09,Y87.1)	429	1.9	5.2
10	Influenza and pneumonia (J10-J18)	257	1.1	3.1
11	Viral hepatitis (B15-B19)	248	1.1	3.0
12	Chronic lower respiratory diseases (J40-J47)	193	0.9	2.3
13	Septicemia (A40-A41)	169	0.7	2.0
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	135	0.6	1.6
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	121	0.5	1.5
...	All other causes (Residual)	3,860	17.0	46.4
White, non-Hispanic, male, 45-49 years				
...	All causes	31,133	100.0	416.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,699	24.7	103.1
2	Malignant neoplasms (C00-C97)	6,745	21.7	90.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,680	11.8	49.3
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	2,008	6.4	26.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,785	5.7	23.9
6	Cerebrovascular diseases (I60-I69)	752	2.4	10.1
7	Human immunodeficiency virus (HIV) disease (B20-B24)	720	2.3	9.6
8	Diabetes mellitus (E10-E14)	715	2.3	9.6
9	Viral hepatitis (B15-B19)	545	1.8	7.3
10	Chronic lower respiratory diseases (J40-J47)	445	1.4	6.0
11	Assault (homicide) (X85-Y09,Y87.1)	325	1.0	4.4
12	Influenza and pneumonia (J10-J18)	280	0.9	3.7
13	Septicemia (A40-A41)	254	0.8	3.4
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	208	0.7	2.8
15	Aortic aneurysm and dissection (I71)	139	0.4	1.9
...	All other causes (Residual)	4,833	15.5	64.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
White, non-Hispanic, male, 50-54 years				
...	All causes	39,655	100.0	597.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,402	28.8	171.8
2	Malignant neoplasms (C00-C97)	11,347	28.6	170.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,841	7.2	42.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	1,692	4.3	25.5
5	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,619	4.1	24.4
6	Diabetes mellitus (E10-E14)	1,127	2.8	17.0
7	Cerebrovascular diseases (I60-I69)	1,104	2.8	16.6
8	Chronic lower respiratory diseases (J40-J47)	814	2.1	12.3
9	Human immunodeficiency virus (HIV) disease (B20-B24)	441	1.1	6.6
10	Viral hepatitis (B15-B19)	413	1.0	6.2
11	Influenza and pneumonia (J10-J18)	387	1.0	5.8
12	Septicemia (A40-A41)	359	0.9	5.4
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	283	0.7	4.3
14	Assault (homicide) (X85-Y09,Y87.1)	235	0.6	3.5
15	Aortic aneurysm and dissection (I71)	216	0.5	3.3
...	All other causes (Residual)	5,375	13.6	81.0
White, non-Hispanic, male, 55-59 years				
...	All causes	48,255	100.0	939.2
1	Malignant neoplasms (C00-C97)	16,345	33.9	318.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,800	30.7	288.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,089	4.3	40.7
4	Chronic lower respiratory diseases (J40-J47)	1,744	3.6	33.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,480	3.1	28.8
6	Cerebrovascular diseases (I60-I69)	1,431	3.0	27.9
7	Diabetes mellitus (E10-E14)	1,417	2.9	27.6
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,165	2.4	22.7
9	Influenza and pneumonia (J10-J18)	480	1.0	9.3
10	Septicemia (A40-A41)	437	0.9	8.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	400	0.8	7.8
12	Aortic aneurysm and dissection (I71)	346	0.7	6.7
13	Human immunodeficiency virus (HIV) disease (B20-B24)	253	0.5	4.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	229	0.5	4.5
15	Viral hepatitis (B15-B19)	205	0.4	4.0
...	All other causes (Residual)	5,434	11.3	105.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	
			Rate	
White, non-Hispanic, male, 60-64 years				
...	All causes	60,776	100.0	1,497.8
1	Malignant neoplasms (C00-C97)	22,023	36.2	542.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	18,379	30.2	453.0
3	Chronic lower respiratory diseases (J40-J47)	2,984	4.9	73.5
4	Cerebrovascular diseases (I60-I69)	2,174	3.6	53.6
5	Diabetes mellitus (E10-E14)	1,881	3.1	46.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,782	2.9	43.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,455	2.4	35.9
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	866	1.4	21.3
9	Influenza and pneumonia (J10-J18)	696	1.1	17.2
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	590	1.0	14.5
11	Septicemia (A40-A41)	578	1.0	14.2
12	Aortic aneurysm and dissection (I71)	568	0.9	14.0
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	310	0.5	7.6
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	244	0.4	6.0
15	Pneumonitis due to solids and liquids (J69)	150	0.2	3.7
...	All other causes (Residual)	6,096	10.0	150.2
White, non-Hispanic, male, 65-69 years				
...	All causes	83,126	100.0	2,357.5
1	Malignant neoplasms (C00-C97)	30,179	36.3	855.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	24,751	29.8	701.9
3	Chronic lower respiratory diseases (J40-J47)	5,367	6.5	152.2
4	Cerebrovascular diseases (I60-I69)	3,274	3.9	92.9
5	Diabetes mellitus (E10-E14)	2,510	3.0	71.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,636	2.0	46.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,314	1.6	37.3
8	Influenza and pneumonia (J10-J18)	1,065	1.3	30.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	983	1.2	27.9
10	Aortic aneurysm and dissection (I71)	974	1.2	27.6
11	Septicemia (A40-A41)	826	1.0	23.4
12	Intentional self-harm (suicide) (X60-X84,Y87.0)	779	0.9	22.1
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	500	0.6	14.2
14	Alzheimer's disease (G30)	373	0.4	10.6
15	Parkinson's disease (G20-G21)	352	0.4	10.0
...	All other causes (Residual)	8,243	9.9	233.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total deaths		
		Number	Rate	Rate
White, non-Hispanic, male, 70-74 years				
...	All causes	118,912	100.0	3,671.9
1	Malignant neoplasms (C00-C97)	38,906	32.7	1,201.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	36,273	30.5	1,120.1
3	Chronic lower respiratory diseases (J40-J47)	9,031	7.6	278.9
4	Cerebrovascular diseases (I60-I69)	5,902	5.0	182.3
5	Diabetes mellitus (E10-E14)	3,457	2.9	106.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,056	1.7	63.5
7	Influenza and pneumonia (J10-J18)	2,019	1.7	62.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,591	1.3	49.1
9	Aortic aneurysm and dissection (I71)	1,479	1.2	45.7
10	Septicemia (A40-A41)	1,305	1.1	40.3
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1,288	1.1	39.8
12	Alzheimer's disease (G30)	994	0.8	30.7
13	Intentional self-harm (suicide) (X60-X84,Y87.0)	922	0.8	28.5
14	Parkinson's disease (G20-G21)	862	0.7	26.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	702	0.6	21.7
...	All other causes (Residual)	12,125	10.2	374.4
White, non-Hispanic, male, 75-79 years				
...	All causes	148,070	100.0	5,591.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	46,643	31.5	1,761.5
2	Malignant neoplasms (C00-C97)	41,247	27.9	1,557.7
3	Chronic lower respiratory diseases (J40-J47)	11,134	7.5	420.5
4	Cerebrovascular diseases (I60-I69)	9,308	6.3	351.5
5	Diabetes mellitus (E10-E14)	4,135	2.8	156.2
6	Influenza and pneumonia (J10-J18)	3,446	2.3	130.1
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,589	1.7	97.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,449	1.7	92.5
9	Alzheimer's disease (G30)	2,195	1.5	82.9
10	Parkinson's disease (G20-G21)	1,887	1.3	71.3
11	Septicemia (A40-A41)	1,748	1.2	66.0
12	Aortic aneurysm and dissection (I71)	1,685	1.1	63.6
13	Pneumonitis due to solids and liquids (J69)	1,091	0.7	41.2
14	Chronic liver disease and cirrhosis (K70,K73-K74)	1,040	0.7	39.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	994	0.7	37.5
...	All other causes (Residual)	16,479	11.1	622.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total		
		Number	deaths	Rate
White, non-Hispanic, male, 80-84 years				
...	All causes	146,772	100.0	9,077.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	48,933	33.3	3,026.2
2	Malignant neoplasms (C00-C97)	32,705	22.3	2,022.6
3	Cerebrovascular diseases (I60-I69)	11,071	7.5	684.7
4	Chronic lower respiratory diseases (J40-J47)	10,569	7.2	653.6
5	Influenza and pneumonia (J10-J18)	4,900	3.3	303.0
6	Diabetes mellitus (E10-E14)	3,661	2.5	226.4
7	Alzheimer's disease (G30)	3,491	2.4	215.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,703	1.8	167.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,692	1.8	166.5
10	Parkinson's disease (G20-G21)	2,304	1.6	142.5
11	Septicemia (A40-A41)	1,831	1.2	113.2
12	Pneumonitis due to solids and liquids (J69)	1,557	1.1	96.3
13	Aortic aneurysm and dissection (I71)	1,451	1.0	89.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,114	0.8	68.9
15	Atherosclerosis (I70)	976	0.7	60.4
...	All other causes (Residual)	16,814	11.5	1,039.8
White, non-Hispanic, male, 85-89 years				
...	All causes	114,484	100.0	15,056.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	40,903	35.7	5,379.3
2	Malignant neoplasms (C00-C97)	19,854	17.3	2,611.1
3	Cerebrovascular diseases (I60-I69)	9,594	8.4	1,261.7
4	Chronic lower respiratory diseases (J40-J47)	7,141	6.2	939.1
5	Influenza and pneumonia (J10-J18)	4,952	4.3	651.3
6	Alzheimer's disease (G30)	3,376	2.9	444.0
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,326	2.0	305.9
8	Diabetes mellitus (E10-E14)	2,320	2.0	305.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,318	2.0	304.8
10	Parkinson's disease (G20-G21)	1,752	1.5	230.4
11	Pneumonitis due to solids and liquids (J69)	1,676	1.5	220.4
12	Septicemia (A40-A41)	1,401	1.2	184.2
13	Aortic aneurysm and dissection (I71)	994	0.9	130.7
14	Atherosclerosis (I70)	983	0.9	129.3
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	801	0.7	105.3
...	All other causes (Residual)	14,093	12.3	1,853.4

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total deaths		
		Number	Rate	Rate
White, non-Hispanic, male, 90-94 years				
...	All causes	56,864	100.0	21,399.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	21,817	38.4	8,210.4
2	Malignant neoplasms (C00-C97)	7,309	12.9	2,750.6
3	Cerebrovascular diseases (I60-I69)	4,928	8.7	1,854.6
4	Influenza and pneumonia (J10-J18)	3,263	5.7	1,228.0
5	Chronic lower respiratory diseases (J40-J47)	2,836	5.0	1,067.3
6	Alzheimer's disease (G30)	1,930	3.4	726.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,268	2.2	477.2
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,246	2.2	468.9
9	Pneumonitis due to solids and liquids (J69)	995	1.7	374.5
10	Diabetes mellitus (E10-E14)	962	1.7	362.0
11	Parkinson's disease (G20-G21)	669	1.2	251.8
12	Septicemia (A40-A41)	668	1.2	251.4
13	Atherosclerosis (I70)	666	1.2	250.6
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	462	0.8	173.9
15	Aortic aneurysm and dissection (I71)	375	0.7	141.1
...	All other causes (Residual)	7,470	13.1	2,811.2
White, non-Hispanic, male, 95-99 years				
...	All causes	15,857	100.0	25,162.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,412	40.4	10,174.9
2	Malignant neoplasms (C00-C97)	1,507	9.5	2,391.4
3	Cerebrovascular diseases (I60-I69)	1,301	8.2	2,064.5
4	Influenza and pneumonia (J10-J18)	1,194	7.5	1,894.7
5	Chronic lower respiratory diseases (J40-J47)	657	4.1	1,042.6
6	Alzheimer's disease (G30)	509	3.2	807.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	375	2.4	595.1
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	345	2.2	547.5
9	Pneumonitis due to solids and liquids (J69)	328	2.1	520.5
10	Atherosclerosis (I70)	254	1.6	403.1
11	Diabetes mellitus (E10-E14)	210	1.3	333.2
12	Septicemia (A40-A41)	150	0.9	238.0
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	116	0.7	184.1
14	Parkinson's disease (G20-G21)	107	0.7	169.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	77	0.5	122.2
...	All other causes (Residual)	2,315	14.6	3,673.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total		
		Number	deaths	Rate
White, non-Hispanic, male, 100 years and over				
...	All causes	2,332	100.0	27,643.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,006	43.1	11,925.1
2	Influenza and pneumonia (J10-J18)	232	9.9	2,750.1
3	Cerebrovascular diseases (I60-I69)	164	7.0	1,944.0
4	Malignant neoplasms (C00-C97)	144	6.2	1,707.0
5	Chronic lower respiratory diseases (J40-J47)	74	3.2	877.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	62	2.7	734.9
7	Pneumonitis due to solids and liquids (J69)	60	2.6	711.2
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	51	2.2	604.6
9	Atherosclerosis (I70)	46	2.0	545.3
10	Alzheimer's disease (G30)	42	1.8	497.9
11	Septicemia (A40-A41)	23	1.0	272.6
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	20	0.9	237.1
13	Diabetes mellitus (E10-E14)	16	0.7	*
14	Anemias (D50-D64)	13	0.6	*
15	Nutritional deficiencies (E40-E64)	10	0.4	*
...	All other causes (Residual)	369	15.8	4,374.1
White, non-Hispanic, female, all ages				
...	All causes	1,015,138	100.0	1,011.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	307,255	30.3	306.2
2	Malignant neoplasms (C00-C97)	222,268	21.9	221.5
3	Cerebrovascular diseases (I60-I69)	86,210	8.5	85.9
4	Chronic lower respiratory diseases (J40-J47)	56,670	5.6	56.5
5	Alzheimer's disease (G30)	32,123	3.2	32.0
6	Influenza and pneumonia (J10-J18)	31,526	3.1	31.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	27,066	2.7	27.0
8	Diabetes mellitus (E10-E14)	26,698	2.6	26.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14,357	1.4	14.3
10	Septicemia (A40-A41)	13,461	1.3	13.4
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	8,196	0.8	8.2
12	Atherosclerosis (I70)	8,052	0.8	8.0
13	Pneumonitis due to solids and liquids (J69)	7,129	0.7	7.1
14	Chronic liver disease and cirrhosis (K70,K73-K74)	7,099	0.7	7.1
15	Parkinson's disease (G20-G21)	6,381	0.6	6.4
...	All other causes (Residual)	160,647	15.8	160.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total deaths		
		Number	Rate	Rate
White, non-Hispanic, female, 1-4 years				
...	All causes	1,100	100.0	23.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	395	35.9	8.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	116	10.5	2.5
3	Malignant neoplasms (C00-C97)	110	10.0	2.4
4	Assault (homicide) (X85-Y09,Y87.1)	58	5.3	1.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	39	3.5	0.8
6	Influenza and pneumonia (J10-J18)	27	2.5	0.6
7	Septicemia (A40-A41)	25	2.3	0.5
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	1.3	*
9	Cerebrovascular diseases (I60-I69)	11	1.0	*
9	Chronic lower respiratory diseases (J40-J47)	11	1.0	*
9	Certain conditions originating in the perinatal period (P00-P96)	11	1.0	*
12	Meningitis (G00,G03)	7	0.6	*
13	Anemias (D50-D64)	6	0.5	*
14	Meningococcal infection (A39)	5	0.5	*
15	Diabetes mellitus (E10-E14)	4	0.4	*
...	All other causes (Residual)	261	23.7	5.7
White, non-Hispanic, female, 5-9 years				
...	All causes	773	100.0	12.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	309	40.0	5.1
2	Malignant neoplasms (C00-C97)	146	18.9	2.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	58	7.5	1.0
4	Assault (homicide) (X85-Y09,Y87.1)	33	4.3	0.5
5	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	21	2.7	0.3
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	20	2.6	0.3
7	Septicemia (A40-A41)	9	1.2	*
8	Influenza and pneumonia (J10-J18)	8	1.0	*
9	Chronic lower respiratory diseases (J40-J47)	7	0.9	*
10	Cerebrovascular diseases (I60-I69)	6	0.8	*
11	Certain conditions originating in the perinatal period (P00-P96)	5	0.6	*
12	Meningitis (G00,G03)	4	0.5	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.5	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.3	*
14	Pneumonitis due to solids and liquids (J69)	2	0.3	*
14	Diseases of appendix (K35-K38)	2	0.3	*
...	All other causes (Residual)	137	17.7	2.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total	
			deaths	Rate
White, non-Hispanic, female, 10-14 years				
...	All causes	972	100.0	15.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	358	36.8	5.6
2	Malignant neoplasms (C00-C97)	143	14.7	2.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	63	6.5	1.0
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	45	4.6	0.7
5	Assault (homicide) (X85-Y09,Y87.1)	38	3.9	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	33	3.4	0.5
7	Cerebrovascular diseases (I60-I69)	17	1.7	*
8	Chronic lower respiratory diseases (J40-J47)	16	1.6	*
9	Influenza and pneumonia (J10-J18)	12	1.2	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	1.1	*
11	Septicemia (A40-A41)	8	0.8	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	6	0.6	*
12	Diseases of appendix (K35-K38)	6	0.6	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.6	*
15	Diabetes mellitus (E10-E14)	5	0.5	*
15	Meningitis (G00,G03)	5	0.5	*
...	All other causes (Residual)	200	20.6	3.2
White, non-Hispanic, female, 15-19 years				
...	All causes	2,601	100.0	40.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,526	58.7	23.9
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	188	7.2	2.9
3	Malignant neoplasms (C00-C97)	182	7.0	2.9
4	Assault (homicide) (X85-Y09,Y87.1)	121	4.7	1.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	90	3.5	1.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	60	2.3	0.9
7	Influenza and pneumonia (J10-J18)	18	0.7	*
8	Cerebrovascular diseases (I60-I69)	15	0.6	*
9	Chronic lower respiratory diseases (J40-J47)	13	0.5	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	0.4	*
10	Diabetes mellitus (E10-E14)	10	0.4	*
12	Meningococcal infection (A39)	9	0.3	*
12	Pregnancy, childbirth and the puerperium (O00-099)	9	0.3	*
14	Septicemia (A40-A41)	8	0.3	*
15	Anemias (D50-D64)	7	0.3	*
...	All other causes (Residual)	335	12.9	5.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
White, non-Hispanic, female, 20-24 years				
...	All causes	2,594	100.0	43.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,144	44.1	19.3
2	Malignant neoplasms (C00-C97)	241	9.3	4.1
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	213	8.2	3.6
4	Assault (homicide) (X85-Y09,Y87.1)	165	6.4	2.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	115	4.4	1.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	56	2.2	0.9
7	Cerebrovascular diseases (I60-I69)	36	1.4	0.6
8	Influenza and pneumonia (J10-J18)	30	1.2	0.5
9	Diabetes mellitus (E10-E14)	25	1.0	0.4
10	Pregnancy, childbirth and the puerperium (O00-099)	24	0.9	0.4
11	Chronic lower respiratory diseases (J40-J47)	22	0.8	0.4
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18	0.7	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	0.5	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	8	0.3	*
15	Septicemia (A40-A41)	7	0.3	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	7	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.3	*
...	All other causes (Residual)	463	17.8	7.8
White, non-Hispanic, female, 25-29 years				
...	All causes	2,902	100.0	49.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	889	30.6	15.2
2	Malignant neoplasms (C00-C97)	432	14.9	7.4
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	285	9.8	4.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	187	6.4	3.2
5	Assault (homicide) (X85-Y09,Y87.1)	169	5.8	2.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	62	2.1	1.1
7	Cerebrovascular diseases (I60-I69)	56	1.9	1.0
8	Diabetes mellitus (E10-E14)	47	1.6	0.8
9	Human immunodeficiency virus (HIV) disease (B20-B24)	44	1.5	0.8
10	Influenza and pneumonia (J10-J18)	38	1.3	0.6
11	Pregnancy, childbirth and the puerperium (O00-099)	33	1.1	0.6
12	Chronic lower respiratory diseases (J40-J47)	28	1.0	0.5
13	Septicemia (A40-A41)	19	0.7	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19	0.7	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	0.6	*
...	All other causes (Residual)	576	19.8	9.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total deaths		
		Number	Rate	Rate
White, non-Hispanic, female, 30-34 years				
...	All causes	4,355	100.0	65.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	980	22.5	14.7
2	Malignant neoplasms (C00-C97)	908	20.8	13.6
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	384	8.8	5.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	351	8.1	5.3
5	Assault (homicide) (X85-Y09,Y87.1)	177	4.1	2.7
6	Human immunodeficiency virus (HIV) disease (B20-B24)	100	2.3	1.5
7	Cerebrovascular diseases (I60-I69)	96	2.2	1.4
8	Diabetes mellitus (E10-E14)	90	2.1	1.3
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	79	1.8	1.2
10	Chronic liver disease and cirrhosis (K70,K73-K74)	68	1.6	1.0
11	Influenza and pneumonia (J10-J18)	64	1.5	1.0
12	Chronic lower respiratory diseases (J40-J47)	45	1.0	0.7
13	Septicemia (A40-A41)	41	0.9	0.6
14	Pregnancy, childbirth and the puerperium (O00-099)	39	0.9	0.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	32	0.7	0.5
...	All other causes (Residual)	901	20.7	13.5
White, non-Hispanic, female, 35-39 years				
...	All causes	7,990	100.0	101.4
1	Malignant neoplasms (C00-C97)	2,097	26.2	26.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,388	17.4	17.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	810	10.1	10.3
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	561	7.0	7.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	249	3.1	3.2
5	Assault (homicide) (X85-Y09,Y87.1)	249	3.1	3.2
7	Cerebrovascular diseases (I60-I69)	235	2.9	3.0
8	Diabetes mellitus (E10-E14)	188	2.4	2.4
9	Human immunodeficiency virus (HIV) disease (B20-B24)	163	2.0	2.1
10	Influenza and pneumonia (J10-J18)	131	1.6	1.7
11	Chronic lower respiratory diseases (J40-J47)	100	1.3	1.3
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	89	1.1	1.1
13	Septicemia (A40-A41)	82	1.0	1.0
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	55	0.7	0.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	49	0.6	0.6
...	All other causes (Residual)	1,544	19.3	19.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total deaths		
		Number	Rate	Rate
White, non-Hispanic, female, 40-44 years				
...	All causes	12,619	100.0	152.0
1	Malignant neoplasms (C00-C97)	4,160	33.0	50.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,507	11.9	18.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,487	11.8	17.9
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	692	5.5	8.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	473	3.7	5.7
6	Cerebrovascular diseases (I60-I69)	449	3.6	5.4
7	Diabetes mellitus (E10-E14)	306	2.4	3.7
8	Chronic lower respiratory diseases (J40-J47)	203	1.6	2.4
9	Assault (homicide) (X85-Y09,Y87.1)	188	1.5	2.3
10	Human immunodeficiency virus (HIV) disease (B20-B24)	159	1.3	1.9
11	Septicemia (A40-A41)	154	1.2	1.9
12	Influenza and pneumonia (J10-J18)	153	1.2	1.8
13	Viral hepatitis (B15-B19)	111	0.9	1.3
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	98	0.8	1.2
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	97	0.8	1.2
...	All other causes (Residual)	2,382	18.9	28.7
White, non-Hispanic, female, 45-49 years				
...	All causes	16,872	100.0	223.3
1	Malignant neoplasms (C00-C97)	6,853	40.6	90.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,206	13.1	29.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,298	7.7	17.2
4	Cerebrovascular diseases (I60-I69)	628	3.7	8.3
5	Intentional self-harm (suicide) (X60-X84,Y87.0)	603	3.6	8.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	580	3.4	7.7
7	Diabetes mellitus (E10-E14)	460	2.7	6.1
8	Chronic lower respiratory diseases (J40-J47)	385	2.3	5.1
9	Influenza and pneumonia (J10-J18)	221	1.3	2.9
10	Septicemia (A40-A41)	191	1.1	2.5
11	Viral hepatitis (B15-B19)	162	1.0	2.1
12	Assault (homicide) (X85-Y09,Y87.1)	154	0.9	2.0
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	126	0.7	1.7
14	Human immunodeficiency virus (HIV) disease (B20-B24)	106	0.6	1.4
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	104	0.6	1.4
...	All other causes (Residual)	2,795	16.6	37.0

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total deaths		
		Number	Rate	Rate
White, non-Hispanic, female, 50-54 years				
...	All causes	24,304	100.0	356.2
1	Malignant neoplasms (C00-C97)	10,916	44.9	160.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,846	15.8	56.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,063	4.4	15.6
4	Cerebrovascular diseases (I60-I69)	949	3.9	13.9
5	Chronic lower respiratory diseases (J40-J47)	819	3.4	12.0
6	Diabetes mellitus (E10-E14)	776	3.2	11.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	580	2.4	8.5
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	544	2.2	8.0
9	Septicemia (A40-A41)	288	1.2	4.2
10	Influenza and pneumonia (J10-J18)	277	1.1	4.1
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	212	0.9	3.1
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	127	0.5	1.9
13	Viral hepatitis (B15-B19)	123	0.5	1.8
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	121	0.5	1.8
15	Assault (homicide) (X85-Y09,Y87.1)	120	0.5	1.8
...	All other causes (Residual)	3,543	14.6	51.9
White, non-Hispanic, female, 55-59 years				
...	All causes	31,516	100.0	582.8
1	Malignant neoplasms (C00-C97)	14,319	45.4	264.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,715	18.1	105.7
3	Chronic lower respiratory diseases (J40-J47)	1,625	5.2	30.0
4	Cerebrovascular diseases (I60-I69)	1,226	3.9	22.7
5	Diabetes mellitus (E10-E14)	1,109	3.5	20.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	955	3.0	17.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	641	2.0	11.9
8	Septicemia (A40-A41)	399	1.3	7.4
9	Intentional self-harm (suicide) (X60-X84,Y87.0)	371	1.2	6.9
10	Influenza and pneumonia (J10-J18)	364	1.2	6.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	304	1.0	5.6
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	194	0.6	3.6
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	133	0.4	2.5
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	117	0.4	2.2
15	Aortic aneurysm and dissection (I71)	105	0.3	1.9
...	All other causes (Residual)	3,939	12.5	72.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total deaths		
		Number	Rate	Rate
White, non-Hispanic, female, 60-64 years				
...	All causes	41,801	100.0	952.1
1	Malignant neoplasms (C00-C97)	18,178	43.5	414.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,398	20.1	191.3
3	Chronic lower respiratory diseases (J40-J47)	2,900	6.9	66.1
4	Cerebrovascular diseases (I60-I69)	1,724	4.1	39.3
5	Diabetes mellitus (E10-E14)	1,516	3.6	34.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	927	2.2	21.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	659	1.6	15.0
8	Influenza and pneumonia (J10-J18)	521	1.2	11.9
9	Septicemia (A40-A41)	505	1.2	11.5
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	482	1.2	11.0
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	271	0.6	6.2
12	Intentional self-harm (suicide) (X60-X84,Y87.0)	257	0.6	5.9
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	185	0.4	4.2
14	Aortic aneurysm and dissection (I71)	182	0.4	4.1
15	Alzheimer's disease (G30)	171	0.4	3.9
...	All other causes (Residual)	4,925	11.8	112.2
White, non-Hispanic, female, 65-69 years				
...	All causes	60,639	100.0	1,499.8
1	Malignant neoplasms (C00-C97)	23,873	39.4	590.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,574	22.4	335.7
3	Chronic lower respiratory diseases (J40-J47)	5,156	8.5	127.5
4	Cerebrovascular diseases (I60-I69)	3,003	5.0	74.3
5	Diabetes mellitus (E10-E14)	2,204	3.6	54.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,023	1.7	25.3
7	Influenza and pneumonia (J10-J18)	819	1.4	20.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	793	1.3	19.6
9	Septicemia (A40-A41)	784	1.3	19.4
10	Chronic liver disease and cirrhosis (K70,K73-K74)	779	1.3	19.3
11	Alzheimer's disease (G30)	378	0.6	9.3
12	Aortic aneurysm and dissection (I71)	361	0.6	8.9
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	347	0.6	8.6
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	329	0.5	8.1
15	Parkinson's disease (G20-G21)	198	0.3	4.9
...	All other causes (Residual)	7,018	11.6	173.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total deaths		
		Number	Rate	Rate
White, non-Hispanic, female, 70-74 years				
...	All causes	94,192	100.0	2,352.3
1	Malignant neoplasms (C00-C97)	31,884	33.9	796.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	23,150	24.6	578.1
3	Chronic lower respiratory diseases (J40-J47)	8,568	9.1	214.0
4	Cerebrovascular diseases (I60-I69)	5,810	6.2	145.1
5	Diabetes mellitus (E10-E14)	3,304	3.5	82.5
6	Influenza and pneumonia (J10-J18)	1,643	1.7	41.0
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,466	1.6	36.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,374	1.5	34.3
9	Alzheimer's disease (G30)	1,259	1.3	31.4
10	Septicemia (A40-A41)	1,258	1.3	31.4
11	Chronic liver disease and cirrhosis (K70,K73-K74)	904	1.0	22.6
12	Aortic aneurysm and dissection (I71)	712	0.8	17.8
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	595	0.6	14.9
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	563	0.6	14.1
15	Parkinson's disease (G20-G21)	483	0.5	12.1
...	All other causes (Residual)	11,219	11.9	280.2
White, non-Hispanic, female, 75-79 years				
...	All causes	139,047	100.0	3,801.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	39,219	28.2	1,072.1
2	Malignant neoplasms (C00-C97)	36,652	26.4	1,001.9
3	Cerebrovascular diseases (I60-I69)	11,193	8.0	306.0
4	Chronic lower respiratory diseases (J40-J47)	11,185	8.0	305.8
5	Diabetes mellitus (E10-E14)	4,590	3.3	125.5
6	Alzheimer's disease (G30)	3,481	2.5	95.2
7	Influenza and pneumonia (J10-J18)	3,250	2.3	88.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,245	1.6	61.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,167	1.6	59.2
10	Septicemia (A40-A41)	1,942	1.4	53.1
11	Parkinson's disease (G20-G21)	1,183	0.9	32.3
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,035	0.7	28.3
13	Aortic aneurysm and dissection (I71)	1,020	0.7	27.9
14	Chronic liver disease and cirrhosis (K70,K73-K74)	968	0.7	26.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	900	0.6	24.6
...	All other causes (Residual)	18,017	13.0	492.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total		
		Number	deaths	Rate
White, non-Hispanic, female, 80-84 years				
...	All causes	170,769	100.0	6,488.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	54,571	32.0	2,073.5
2	Malignant neoplasms (C00-C97)	32,475	19.0	1,233.9
3	Cerebrovascular diseases (I60-I69)	16,984	9.9	645.3
4	Chronic lower respiratory diseases (J40-J47)	11,136	6.5	423.1
5	Alzheimer's disease (G30)	6,537	3.8	248.4
6	Influenza and pneumonia (J10-J18)	5,472	3.2	207.9
7	Diabetes mellitus (E10-E14)	4,654	2.7	176.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,819	1.7	107.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,718	1.6	103.3
10	Septicemia (A40-A41)	2,380	1.4	90.4
11	Parkinson's disease (G20-G21)	1,732	1.0	65.8
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,546	0.9	58.7
13	Atherosclerosis (I70)	1,327	0.8	50.4
14	Pneumonitis due to solids and liquids (J69)	1,302	0.8	49.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,163	0.7	44.2
...	All other causes (Residual)	23,953	14.0	910.1
White, non-Hispanic, female, 85-89 years				
...	All causes	183,246	100.0	11,615.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	66,472	36.3	4,213.3
2	Malignant neoplasms (C00-C97)	23,356	12.7	1,480.4
3	Cerebrovascular diseases (I60-I69)	20,515	11.2	1,300.3
4	Alzheimer's disease (G30)	8,866	4.8	562.0
5	Chronic lower respiratory diseases (J40-J47)	8,624	4.7	546.6
6	Influenza and pneumonia (J10-J18)	7,266	4.0	460.6
7	Diabetes mellitus (E10-E14)	4,152	2.3	263.2
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,084	1.7	195.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,884	1.6	182.8
10	Septicemia (A40-A41)	2,674	1.5	169.5
11	Atherosclerosis (I70)	1,909	1.0	121.0
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,843	1.0	116.8
13	Pneumonitis due to solids and liquids (J69)	1,761	1.0	111.6
14	Parkinson's disease (G20-G21)	1,580	0.9	100.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,103	0.6	69.9
...	All other causes (Residual)	27,157	14.8	1,721.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total		
		Number	deaths	Rate
White, non-Hispanic, female, 90-94 years				
...	All causes	138,823	100.0	18,818.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	55,613	40.1	7,538.7
2	Cerebrovascular diseases (I60-I69)	15,848	11.4	2,148.3
3	Malignant neoplasms (C00-C97)	11,562	8.3	1,567.3
4	Alzheimer's disease (G30)	7,577	5.5	1,027.1
5	Influenza and pneumonia (J10-J18)	6,854	4.9	929.1
6	Chronic lower respiratory diseases (J40-J47)	4,289	3.1	581.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,348	1.7	318.3
8	Diabetes mellitus (E10-E14)	2,347	1.7	318.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,050	1.5	277.9
10	Atherosclerosis (I70)	1,944	1.4	263.5
11	Septicemia (A40-A41)	1,800	1.3	244.0
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,530	1.1	207.4
13	Pneumonitis due to solids and liquids (J69)	1,487	1.1	201.6
14	Parkinson's disease (G20-G21)	846	0.6	114.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	705	0.5	95.6
...	All other causes (Residual)	22,023	15.9	2,985.4
White, non-Hispanic, female, 95-99 years				
...	All causes	58,814	100.0	25,151.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	25,185	42.8	10,770.4
2	Cerebrovascular diseases (I60-I69)	6,190	10.5	2,647.2
3	Influenza and pneumonia (J10-J18)	3,360	5.7	1,436.9
4	Malignant neoplasms (C00-C97)	3,293	5.6	1,408.3
5	Alzheimer's disease (G30)	3,172	5.4	1,356.5
6	Chronic lower respiratory diseases (J40-J47)	1,313	2.2	561.5
7	Atherosclerosis (I70)	1,140	1.9	487.5
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	939	1.6	401.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	848	1.4	362.6
10	Diabetes mellitus (E10-E14)	788	1.3	337.0
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	702	1.2	300.2
12	Septicemia (A40-A41)	698	1.2	298.5
13	Pneumonitis due to solids and liquids (J69)	657	1.1	281.0
14	Parkinson's disease (G20-G21)	238	0.4	101.8
15	Nutritional deficiencies (E40-E64)	207	0.4	88.5
...	All other causes (Residual)	10,084	17.1	4,312.4

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total deaths		
		Number	Rate	Rate
White, non-Hispanic, female, 100 years and over				
...	All causes	13,337	100.0	31,950.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,046	45.3	14,483.9
2	Cerebrovascular diseases (I60-I69)	1,198	9.0	2,869.9
3	Influenza and pneumonia (J10-J18)	952	7.1	2,280.6
4	Alzheimer's disease (G30)	580	4.3	1,389.5
5	Malignant neoplasms (C00-C97)	462	3.5	1,106.8
6	Atherosclerosis (I70)	328	2.5	785.8
7	Chronic lower respiratory diseases (J40-J47)	221	1.7	529.4
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	213	1.6	510.3
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	167	1.3	400.1
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	148	1.1	354.6
11	Pneumonitis due to solids and liquids (J69)	147	1.1	352.2
12	Septicemia (A40-A41)	136	1.0	325.8
13	Diabetes mellitus (E10-E14)	122	0.9	292.3
14	Nutritional deficiencies (E40-E64)	58	0.4	138.9
15	Anemias (D50-D64)	54	0.4	129.4
...	All other causes (Residual)	2,505	18.8	6,001.0
Black, non-Hispanic, both sexes, all ages				
...	All causes	282,676	100.0	844.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	76,706	27.1	229.1
2	Malignant neoplasms (C00-C97)	61,390	21.7	183.4
3	Cerebrovascular diseases (I60-I69)	19,053	6.7	56.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,087	4.3	36.1
5	Diabetes mellitus (E10-E14)	11,918	4.2	35.6
6	Assault (homicide) (X85-Y09,Y87.1)	7,748	2.7	23.1
7	Human immunodeficiency virus (HIV) disease (B20-B24)	7,708	2.7	23.0
8	Chronic lower respiratory diseases (J40-J47)	7,528	2.7	22.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,853	2.4	20.5
10	Influenza and pneumonia (J10-J18)	5,920	2.1	17.7
11	Septicemia (A40-A41)	5,863	2.1	17.5
12	Certain conditions originating in the perinatal period (P00-P96)	4,878	1.7	14.6
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	3,862	1.4	11.5
14	Alzheimer's disease (G30)	2,719	1.0	8.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,705	1.0	8.1
...	All other causes (Residual)	45,738	16.2	136.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 1-4 years				
...	All causes	1,217	100.0	58.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	396	32.5	19.1
2	Assault (homicide) (X85-Y09,Y87.1)	143	11.8	6.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	123	10.1	5.9
4	Malignant neoplasms (C00-C97)	66	5.4	3.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	47	3.9	2.3
6	Influenza and pneumonia (J10-J18)	29	2.4	1.4
7	Certain conditions originating in the perinatal period (P00-P96)	27	2.2	1.3
8	Septicemia (A40-A41)	25	2.1	1.2
9	Chronic lower respiratory diseases (J40-J47)	21	1.7	1.0
10	Anemias (D50-D64)	19	1.6	*
11	Meningitis (G00,G03)	13	1.1	*
11	Cerebrovascular diseases (I60-I69)	13	1.1	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	12	1.0	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.7	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	8	0.7	*
...	All other causes (Residual)	266	21.9	12.8
Black, non-Hispanic, both sexes, 5-9 years				
...	All causes	745	100.0	25.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	330	44.3	11.4
2	Malignant neoplasms (C00-C97)	71	9.5	2.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	44	5.9	1.5
4	Assault (homicide) (X85-Y09,Y87.1)	40	5.4	1.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	29	3.9	1.0
6	Chronic lower respiratory diseases (J40-J47)	24	3.2	0.8
7	Anemias (D50-D64)	14	1.9	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	1.7	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	9	1.2	*
10	Septicemia (A40-A41)	8	1.1	*
11	Meningococcal infection (A39)	7	0.9	*
11	Influenza and pneumonia (J10-J18)	7	0.9	*
13	Certain conditions originating in the perinatal period (P00-P96)	6	0.8	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.5	*
15	Diabetes mellitus (E10-E14)	3	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.4	*
...	All other causes (Residual)	133	17.9	4.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total deaths		
		Number	Rate	Rate
Black, non-Hispanic, both sexes, 10-14 years				
...	All causes	875	100.0	29.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	289	33.0	9.6
2	Assault (homicide) (X85-Y09,Y87.1)	82	9.4	2.7
3	Malignant neoplasms (C00-C97)	76	8.7	2.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	46	5.3	1.5
4	Chronic lower respiratory diseases (J40-J47)	46	5.3	1.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	45	5.1	1.5
7	Intentional self-harm (suicide) (X60-X84,Y87.0)	39	4.5	1.3
8	Anemias (D50-D64)	16	1.8	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	14	1.6	*
10	Cerebrovascular diseases (I60-I69)	13	1.5	*
11	Meningitis (G00,G03)	8	0.9	*
12	Diabetes mellitus (E10-E14)	6	0.7	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.6	*
14	Influenza and pneumonia (J10-J18)	4	0.5	*
14	Certain conditions originating in the perinatal period (P00-P96)	4	0.5	*
...	All other causes (Residual)	182	20.8	6.1
Black, non-Hispanic, both sexes, 15-19 years				
...	All causes	2,672	100.0	92.3
1	Assault (homicide) (X85-Y09,Y87.1)	1,012	37.9	35.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	721	27.0	24.9
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	170	6.4	5.9
4	Malignant neoplasms (C00-C97)	135	5.1	4.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	124	4.6	4.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	56	2.1	1.9
7	Chronic lower respiratory diseases (J40-J47)	44	1.6	1.5
8	Human immunodeficiency virus (HIV) disease (B20-B24)	22	0.8	0.8
9	Anemias (D50-D64)	16	0.6	*
10	Legal intervention (Y35,Y89.0)	15	0.6	*
11	Diabetes mellitus (E10-E14)	14	0.5	*
11	Influenza and pneumonia (J10-J18)	14	0.5	*
13	Cerebrovascular diseases (I60-I69)	13	0.5	*
14	Septicemia (A40-A41)	12	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	0.4	*
...	All other causes (Residual)	292	10.9	10.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total deaths		
		Number	Rate	Rate
Black, non-Hispanic, both sexes, 20-24 years				
...	All causes	4,254	100.0	161.4
1	Assault (homicide) (X85-Y09,Y87.1)	1,759	41.3	66.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	941	22.1	35.7
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	301	7.1	11.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	207	4.9	7.9
5	Malignant neoplasms (C00-C97)	162	3.8	6.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	103	2.4	3.9
7	Anemias (D50-D64)	45	1.1	1.7
8	Diabetes mellitus (E10-E14)	41	1.0	1.6
9	Chronic lower respiratory diseases (J40-J47)	36	0.8	1.4
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	35	0.8	1.3
11	Influenza and pneumonia (J10-J18)	30	0.7	1.1
11	Pregnancy, childbirth and the puerperium (O00-099)	30	0.7	1.1
13	Cerebrovascular diseases (I60-I69)	29	0.7	1.1
14	Septicemia (A40-A41)	21	0.5	0.8
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	0.3	*
15	Legal intervention (Y35,Y89.0)	12	0.3	*
...	All other causes (Residual)	490	11.5	18.6
Black, non-Hispanic, both sexes, 25-29 years				
...	All causes	4,489	100.0	183.2
1	Assault (homicide) (X85-Y09,Y87.1)	1,291	28.8	52.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	877	19.5	35.8
3	Human immunodeficiency virus (HIV) disease (B20-B24)	405	9.0	16.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	343	7.6	14.0
5	Intentional self-harm (suicide) (X60-X84,Y87.0)	262	5.8	10.7
6	Malignant neoplasms (C00-C97)	213	4.7	8.7
7	Diabetes mellitus (E10-E14)	75	1.7	3.1
8	Chronic lower respiratory diseases (J40-J47)	49	1.1	2.0
9	Anemias (D50-D64)	47	1.0	1.9
10	Cerebrovascular diseases (I60-I69)	41	0.9	1.7
11	Septicemia (A40-A41)	39	0.9	1.6
12	Influenza and pneumonia (J10-J18)	37	0.8	1.5
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	36	0.8	1.5
14	Pregnancy, childbirth and the puerperium (O00-099)	31	0.7	1.3
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	28	0.6	1.1
...	All other causes (Residual)	715	15.9	29.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total deaths		
		Number	Rate	Rate
Black, non-Hispanic, both sexes, 30-34 years				
...	All causes	5,614	100.0	224.7
1	Human immunodeficiency virus (HIV) disease (B20-B24)	1,009	18.0	40.4
2	Assault (homicide) (X85-Y09,Y87.1)	902	16.1	36.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	863	15.4	34.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	586	10.4	23.5
5	Malignant neoplasms (C00-C97)	428	7.6	17.1
6	Intentional self-harm (suicide) (X60-X84,Y87.0)	207	3.7	8.3
7	Cerebrovascular diseases (I60-I69)	117	2.1	4.7
8	Diabetes mellitus (E10-E14)	112	2.0	4.5
9	Anemias (D50-D64)	77	1.4	3.1
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	71	1.3	2.8
11	Septicemia (A40-A41)	68	1.2	2.7
12	Chronic lower respiratory diseases (J40-J47)	65	1.2	2.6
13	Influenza and pneumonia (J10-J18)	55	1.0	2.2
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	38	0.7	1.5
15	Chronic liver disease and cirrhosis (K70,K73-K74)	36	0.6	1.4
...	All other causes (Residual)	980	17.5	39.2
Black, non-Hispanic, both sexes, 35-39 years				
...	All causes	8,247	100.0	301.4
1	Human immunodeficiency virus (HIV) disease (B20-B24)	1,447	17.5	52.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,118	13.6	40.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,042	12.6	38.1
4	Malignant neoplasms (C00-C97)	1,006	12.2	36.8
5	Assault (homicide) (X85-Y09,Y87.1)	754	9.1	27.6
6	Cerebrovascular diseases (I60-I69)	281	3.4	10.3
7	Intentional self-harm (suicide) (X60-X84,Y87.0)	205	2.5	7.5
8	Diabetes mellitus (E10-E14)	176	2.1	6.4
9	Chronic liver disease and cirrhosis (K70,K73-K74)	142	1.7	5.2
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	129	1.6	4.7
11	Septicemia (A40-A41)	109	1.3	4.0
12	Chronic lower respiratory diseases (J40-J47)	102	1.2	3.7
13	Influenza and pneumonia (J10-J18)	83	1.0	3.0
14	Anemias (D50-D64)	65	0.8	2.4
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	61	0.7	2.2
...	All other causes (Residual)	1,527	18.5	55.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total deaths		
		Number	Rate	Rate
Black, non-Hispanic, both sexes, 40-44 years				
...	All causes	12,032	100.0	450.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,141	17.8	80.1
2	Malignant neoplasms (C00-C97)	2,066	17.2	77.3
3	Human immunodeficiency virus (HIV) disease (B20-B24)	1,581	13.1	59.1
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,211	10.1	45.3
5	Assault (homicide) (X85-Y09,Y87.1)	605	5.0	22.6
6	Cerebrovascular diseases (I60-I69)	548	4.6	20.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	304	2.5	11.4
8	Diabetes mellitus (E10-E14)	282	2.3	10.5
9	Intentional self-harm (suicide) (X60-X84,Y87.0)	222	1.8	8.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	176	1.5	6.6
11	Influenza and pneumonia (J10-J18)	171	1.4	6.4
12	Chronic lower respiratory diseases (J40-J47)	170	1.4	6.4
13	Septicemia (A40-A41)	160	1.3	6.0
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	115	1.0	4.3
15	Viral hepatitis (B15-B19)	88	0.7	3.3
...	All other causes (Residual)	2,192	18.2	82.0
Black, non-Hispanic, both sexes, 45-49 years				
...	All causes	15,545	100.0	701.4
1	Malignant neoplasms (C00-C97)	3,398	21.9	153.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,320	21.4	149.8
3	Human immunodeficiency virus (HIV) disease (B20-B24)	1,360	8.7	61.4
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,096	7.1	49.4
5	Cerebrovascular diseases (I60-I69)	868	5.6	39.2
6	Diabetes mellitus (E10-E14)	513	3.3	23.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	452	2.9	20.4
8	Assault (homicide) (X85-Y09,Y87.1)	390	2.5	17.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	300	1.9	13.5
10	Septicemia (A40-A41)	258	1.7	11.6
10	Chronic lower respiratory diseases (J40-J47)	258	1.7	11.6
12	Influenza and pneumonia (J10-J18)	197	1.3	8.9
13	Viral hepatitis (B15-B19)	170	1.1	7.7
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	168	1.1	7.6
15	Intentional self-harm (suicide) (X60-X84,Y87.0)	150	1.0	6.8
...	All other causes (Residual)	2,647	17.0	119.4

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Black, non-Hispanic, both sexes, 50-54 years				
...	All causes	17,366	100.0	1,004.4
1	Malignant neoplasms (C00-C97)	4,670	26.9	270.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,286	24.7	247.9
3	Cerebrovascular diseases (I60-I69)	951	5.5	55.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	865	5.0	50.0
5	Human immunodeficiency virus (HIV) disease (B20-B24)	851	4.9	49.2
6	Diabetes mellitus (E10-E14)	757	4.4	43.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	477	2.7	27.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	364	2.1	21.1
9	Chronic lower respiratory diseases (J40-J47)	335	1.9	19.4
10	Septicemia (A40-A41)	314	1.8	18.2
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	237	1.4	13.7
12	Influenza and pneumonia (J10-J18)	226	1.3	13.1
13	Assault (homicide) (X85-Y09,Y87.1)	225	1.3	13.0
14	Viral hepatitis (B15-B19)	159	0.9	9.2
15	Intentional self-harm (suicide) (X60-X84,Y87.0)	116	0.7	6.7
...	All other causes (Residual)	2,533	14.6	146.5
Black, non-Hispanic, both sexes, 55-59 years				
...	All causes	17,965	100.0	1,411.2
1	Malignant neoplasms (C00-C97)	5,447	30.3	427.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,076	28.3	398.7
3	Cerebrovascular diseases (I60-I69)	1,032	5.7	81.1
4	Diabetes mellitus (E10-E14)	889	4.9	69.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	561	3.1	44.1
6	Chronic lower respiratory diseases (J40-J47)	443	2.5	34.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	433	2.4	34.0
8	Human immunodeficiency virus (HIV) disease (B20-B24)	388	2.2	30.5
9	Chronic liver disease and cirrhosis (K70,K73-K74)	341	1.9	26.8
10	Septicemia (A40-A41)	324	1.8	25.5
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	240	1.3	18.9
12	Influenza and pneumonia (J10-J18)	237	1.3	18.6
13	Assault (homicide) (X85-Y09,Y87.1)	121	0.7	9.5
14	Aortic aneurysm and dissection (I71)	95	0.5	7.5
15	Viral hepatitis (B15-B19)	90	0.5	7.1
...	All other causes (Residual)	2,248	12.5	176.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total		
		Number	deaths	Rate
Black, non-Hispanic, both sexes, 60-64 years				
...	All causes	20,902	100.0	2,013.6
1	Malignant neoplasms (C00-C97)	6,572	31.4	633.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,181	29.6	595.4
3	Cerebrovascular diseases (I60-I69)	1,233	5.9	118.8
4	Diabetes mellitus (E10-E14)	1,106	5.3	106.5
5	Chronic lower respiratory diseases (J40-J47)	620	3.0	59.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	547	2.6	52.7
7	Septicemia (A40-A41)	432	2.1	41.6
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	422	2.0	40.7
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	315	1.5	30.3
10	Influenza and pneumonia (J10-J18)	294	1.4	28.3
11	Chronic liver disease and cirrhosis (K70,K73-K74)	281	1.3	27.1
12	Human immunodeficiency virus (HIV) disease (B20-B24)	237	1.1	22.8
13	Aortic aneurysm and dissection (I71)	101	0.5	9.7
14	Assault (homicide) (X85-Y09,Y87.1)	80	0.4	7.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	78	0.4	7.5
...	All other causes (Residual)	2,403	11.5	231.5
Black, non-Hispanic, both sexes, 65-69 years				
...	All causes	24,831	100.0	2,745.7
1	Malignant neoplasms (C00-C97)	7,733	31.1	855.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,302	29.4	807.4
3	Cerebrovascular diseases (I60-I69)	1,573	6.3	173.9
4	Diabetes mellitus (E10-E14)	1,420	5.7	157.0
5	Chronic lower respiratory diseases (J40-J47)	872	3.5	96.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	748	3.0	82.7
7	Septicemia (A40-A41)	537	2.2	59.4
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	414	1.7	45.8
9	Influenza and pneumonia (J10-J18)	408	1.6	45.1
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	362	1.5	40.0
11	Chronic liver disease and cirrhosis (K70,K73-K74)	246	1.0	27.2
12	Human immunodeficiency virus (HIV) disease (B20-B24)	130	0.5	14.4
13	Aortic aneurysm and dissection (I71)	129	0.5	14.3
14	Pneumonitis due to solids and liquids (J69)	102	0.4	11.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	96	0.4	10.6
...	All other causes (Residual)	2,759	11.1	305.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total		
		Number	deaths	Rate
Black, non-Hispanic, both sexes, 70-74 years				
...	All causes	29,814	100.0	4,089.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,111	30.6	1,249.6
2	Malignant neoplasms (C00-C97)	8,752	29.4	1,200.4
3	Cerebrovascular diseases (I60-I69)	2,087	7.0	286.2
4	Diabetes mellitus (E10-E14)	1,663	5.6	228.1
5	Chronic lower respiratory diseases (J40-J47)	1,121	3.8	153.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	889	3.0	121.9
7	Septicemia (A40-A41)	636	2.1	87.2
8	Influenza and pneumonia (J10-J18)	576	1.9	79.0
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	443	1.5	60.8
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	439	1.5	60.2
11	Alzheimer's disease (G30)	194	0.7	26.6
12	Pneumonitis due to solids and liquids (J69)	187	0.6	25.6
13	Chronic liver disease and cirrhosis (K70,K73-K74)	183	0.6	25.1
14	Aortic aneurysm and dissection (I71)	150	0.5	20.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	149	0.5	20.4
...	All other causes (Residual)	3,234	10.8	443.6
Black, non-Hispanic, both sexes, 75-79 years				
...	All causes	32,219	100.0	5,939.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,227	31.7	1,885.3
2	Malignant neoplasms (C00-C97)	8,264	25.6	1,523.5
3	Cerebrovascular diseases (I60-I69)	2,644	8.2	487.4
4	Diabetes mellitus (E10-E14)	1,769	5.5	326.1
5	Chronic lower respiratory diseases (J40-J47)	1,176	3.7	216.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	923	2.9	170.2
7	Septicemia (A40-A41)	817	2.5	150.6
8	Influenza and pneumonia (J10-J18)	689	2.1	127.0
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	518	1.6	95.5
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	419	1.3	77.2
11	Alzheimer's disease (G30)	399	1.2	73.6
12	Pneumonitis due to solids and liquids (J69)	216	0.7	39.8
13	Aortic aneurysm and dissection (I71)	188	0.6	34.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	149	0.5	27.5
15	Atherosclerosis (I70)	130	0.4	24.0
...	All other causes (Residual)	3,691	11.5	680.4

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total deaths		
		Number	Rate	
Black, non-Hispanic, both sexes, 80-84 years				
...	All causes	28,655	100.0	8,704.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,373	32.7	2,847.3
2	Malignant neoplasms (C00-C97)	5,980	20.9	1,816.6
3	Cerebrovascular diseases (I60-I69)	2,700	9.4	820.2
4	Diabetes mellitus (E10-E14)	1,424	5.0	432.6
5	Chronic lower respiratory diseases (J40-J47)	952	3.3	289.2
6	Influenza and pneumonia (J10-J18)	861	3.0	261.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	853	3.0	259.1
8	Septicemia (A40-A41)	759	2.6	230.6
9	Alzheimer's disease (G30)	605	2.1	183.8
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	462	1.6	140.3
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	349	1.2	106.0
12	Pneumonitis due to solids and liquids (J69)	252	0.9	76.6
13	Atherosclerosis (I70)	190	0.7	57.7
14	Aortic aneurysm and dissection (I71)	143	0.5	43.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	133	0.5	40.4
...	All other causes (Residual)	3,619	12.6	1,099.4
Black, non-Hispanic, both sexes, 85-89 years				
...	All causes	23,696	100.0	13,331.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,381	35.4	4,715.1
2	Malignant neoplasms (C00-C97)	3,860	16.3	2,171.6
3	Cerebrovascular diseases (I60-I69)	2,472	10.4	1,390.7
4	Diabetes mellitus (E10-E14)	953	4.0	536.1
5	Influenza and pneumonia (J10-J18)	815	3.4	458.5
6	Alzheimer's disease (G30)	680	2.9	382.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	667	2.8	375.2
8	Chronic lower respiratory diseases (J40-J47)	663	2.8	373.0
9	Septicemia (A40-A41)	631	2.7	355.0
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	435	1.8	244.7
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	279	1.2	157.0
12	Pneumonitis due to solids and liquids (J69)	232	1.0	130.5
13	Atherosclerosis (I70)	172	0.7	96.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	116	0.5	65.3
15	Aortic aneurysm and dissection (I71)	113	0.5	63.6
...	All other causes (Residual)	3,227	13.6	1,815.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total		
		Number	deaths	Rate
Black, non-Hispanic, both sexes, 90-94 years				
...	All causes	14,579	100.0	16,424.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,367	36.8	6,046.5
2	Malignant neoplasms (C00-C97)	1,769	12.1	1,993.0
3	Cerebrovascular diseases (I60-I69)	1,507	10.3	1,697.8
4	Influenza and pneumonia (J10-J18)	653	4.5	735.7
5	Diabetes mellitus (E10-E14)	505	3.5	568.9
6	Alzheimer's disease (G30)	463	3.2	521.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	411	2.8	463.0
8	Septicemia (A40-A41)	378	2.6	425.9
9	Chronic lower respiratory diseases (J40-J47)	358	2.5	403.3
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	281	1.9	316.6
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	189	1.3	212.9
12	Pneumonitis due to solids and liquids (J69)	178	1.2	200.5
13	Atherosclerosis (I70)	153	1.0	172.4
14	Aortic aneurysm and dissection (I71)	64	0.4	72.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	62	0.4	69.8
...	All other causes (Residual)	2,241	15.4	2,524.7
Black, non-Hispanic, both sexes, 95-99 years				
...	All causes	6,362	100.0	17,651.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,479	39.0	6,877.9
2	Cerebrovascular diseases (I60-I69)	685	10.8	1,900.5
3	Malignant neoplasms (C00-C97)	591	9.3	1,639.7
4	Influenza and pneumonia (J10-J18)	311	4.9	862.9
5	Alzheimer's disease (G30)	218	3.4	604.8
6	Septicemia (A40-A41)	185	2.9	513.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	168	2.6	466.1
8	Diabetes mellitus (E10-E14)	160	2.5	443.9
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	142	2.2	394.0
10	Chronic lower respiratory diseases (J40-J47)	131	2.1	363.5
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	80	1.3	222.0
12	Atherosclerosis (I70)	76	1.2	210.9
13	Pneumonitis due to solids and liquids (J69)	74	1.2	205.3
14	Nutritional deficiencies (E40-E64)	40	0.6	111.0
15	Anemias (D50-D64)	28	0.4	77.7
...	All other causes (Residual)	994	15.6	2,757.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total		
		Number	deaths	Rate
Black, non-Hispanic, both sexes, 100 years and over				
...	All causes	2,023	100.0	24,402.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	832	41.1	10,036.2
2	Cerebrovascular diseases (I60-I69)	197	9.7	2,376.4
3	Influenza and pneumonia (J10-J18)	122	6.0	1,471.7
4	Malignant neoplasms (C00-C97)	117	5.8	1,411.3
5	Alzheimer's disease (G30)	57	2.8	687.6
6	Septicemia (A40-A41)	51	2.5	615.2
7	Diabetes mellitus (E10-E14)	49	2.4	591.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	44	2.2	530.8
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	41	2.0	494.6
10	Atherosclerosis (I70)	40	2.0	482.5
11	Chronic lower respiratory diseases (J40-J47)	31	1.5	373.9
12	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	28	1.4	337.8
13	Pneumonitis due to solids and liquids (J69)	21	1.0	253.3
14	Nutritional deficiencies (E40-E64)	16	0.8	*
15	Anemias (D50-D64)	10	0.5	*
...	All other causes (Residual)	367	18.1	4,427.0
Black, non-Hispanic, male, all ages				
...	All causes	143,297	100.0	903.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	36,282	25.3	228.7
2	Malignant neoplasms (C00-C97)	32,493	22.7	204.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,380	5.8	52.8
4	Cerebrovascular diseases (I60-I69)	7,945	5.5	50.1
5	Assault (homicide) (X85-Y09,Y87.1)	6,382	4.5	40.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	5,291	3.7	33.4
7	Diabetes mellitus (E10-E14)	4,720	3.3	29.8
8	Chronic lower respiratory diseases (J40-J47)	4,184	2.9	26.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,045	2.1	19.2
10	Influenza and pneumonia (J10-J18)	2,880	2.0	18.2
11	Certain conditions originating in the perinatal period (P00-P96)	2,760	1.9	17.4
12	Septicemia (A40-A41)	2,559	1.8	16.1
13	Chronic liver disease and cirrhosis (K70,K73-K74)	1,709	1.2	10.8
14	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,601	1.1	10.1
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,576	1.1	9.9
...	All other causes (Residual)	21,490	15.0	135.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 1-4 years				
...	All causes	672	100.0	63.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	232	34.5	22.1
2	Assault (homicide) (X85-Y09,Y87.1)	73	10.9	6.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	67	10.0	6.4
4	Malignant neoplasms (C00-C97)	31	4.6	2.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	24	3.6	2.3
6	Influenza and pneumonia (J10-J18)	20	3.0	1.9
7	Certain conditions originating in the perinatal period (P00-P96)	18	2.7	*
8	Chronic lower respiratory diseases (J40-J47)	14	2.1	*
9	Septicemia (A40-A41)	13	1.9	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	1.2	*
10	Anemias (D50-D64)	8	1.2	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	6	0.9	*
12	Meningitis (G00,G03)	6	0.9	*
14	Meningococcal infection (A39)	4	0.6	*
15	Cerebrovascular diseases (I60-I69)	3	0.4	*
...	All other causes (Residual)	145	21.6	13.8
Black, non-Hispanic, male, 5-9 years				
...	All causes	425	100.0	28.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	204	48.0	13.9
2	Malignant neoplasms (C00-C97)	42	9.9	2.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	29	6.8	2.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	17	4.0	*
5	Assault (homicide) (X85-Y09,Y87.1)	16	3.8	*
6	Chronic lower respiratory diseases (J40-J47)	14	3.3	*
7	Anemias (D50-D64)	10	2.4	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.4	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	5	1.2	*
10	Septicemia (A40-A41)	4	0.9	*
10	Influenza and pneumonia (J10-J18)	4	0.9	*
12	Meningococcal infection (A39)	3	0.7	*
12	Certain conditions originating in the perinatal period (P00-P96)	3	0.7	*
14	Diabetes mellitus (E10-E14)	2	0.5	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.5	*
...	All other causes (Residual)	64	15.1	4.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total deaths		
		Number	Rate	Rate
Black, non-Hispanic, male, 10-14 years				
...	All causes	534	100.0	35.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	202	37.8	13.3
2	Assault (homicide) (X85-Y09,Y87.1)	54	10.1	3.5
3	Malignant neoplasms (C00-C97)	44	8.2	2.9
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	34	6.4	2.2
5	Chronic lower respiratory diseases (J40-J47)	29	5.4	1.9
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	24	4.5	1.6
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	21	3.9	1.4
8	Human immunodeficiency virus (HIV) disease (B20-B24)	7	1.3	*
8	Anemias (D50-D64)	7	1.3	*
10	Meningitis (G00,G03)	6	1.1	*
11	Cerebrovascular diseases (I60-I69)	5	0.9	*
12	Influenza and pneumonia (J10-J18)	3	0.6	*
13	Septicemia (A40-A41)	2	0.4	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.4	*
13	Diabetes mellitus (E10-E14)	2	0.4	*
13	Legal intervention (Y35,Y89.0)	2	0.4	*
...	All other causes (Residual)	90	16.9	5.9
Black, non-Hispanic, male, 15-19 years				
...	All causes	2,012	100.0	136.6
1	Assault (homicide) (X85-Y09,Y87.1)	886	44.0	60.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	530	26.3	36.0
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	148	7.4	10.0
4	Malignant neoplasms (C00-C97)	82	4.1	5.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	74	3.7	5.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	39	1.9	2.6
7	Chronic lower respiratory diseases (J40-J47)	31	1.5	2.1
8	Legal intervention (Y35,Y89.0)	15	0.7	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	11	0.5	*
10	Diabetes mellitus (E10-E14)	8	0.4	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.3	*
11	Anemias (D50-D64)	7	0.3	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.3	*
14	Septicemia (A40-A41)	6	0.3	*
14	Influenza and pneumonia (J10-J18)	6	0.3	*
...	All other causes (Residual)	155	7.7	10.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total deaths		
		Number	Rate	Rate
Black, non-Hispanic, male, 20-24 years				
...	All causes	3,212	100.0	246.4
1	Assault (homicide) (X85-Y09,Y87.1)	1,573	49.0	120.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	703	21.9	53.9
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	259	8.1	19.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	118	3.7	9.1
5	Malignant neoplasms (C00-C97)	96	3.0	7.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	49	1.5	3.8
7	Anemias (D50-D64)	22	0.7	1.7
8	Diabetes mellitus (E10-E14)	20	0.6	1.5
8	Chronic lower respiratory diseases (J40-J47)	20	0.6	1.5
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	19	0.6	*
11	Cerebrovascular diseases (I60-I69)	15	0.5	*
11	Influenza and pneumonia (J10-J18)	15	0.5	*
13	Septicemia (A40-A41)	12	0.4	*
13	Legal intervention (Y35,Y89.0)	12	0.4	*
15	Viral hepatitis (B15-B19)	5	0.2	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	5	0.2	*
...	All other causes (Residual)	269	8.4	20.6
Black, non-Hispanic, male, 25-29 years				
...	All causes	3,127	100.0	266.9
1	Assault (homicide) (X85-Y09,Y87.1)	1,138	36.4	97.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	681	21.8	58.1
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	222	7.1	18.9
4	Human immunodeficiency virus (HIV) disease (B20-B24)	208	6.7	17.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	200	6.4	17.1
6	Malignant neoplasms (C00-C97)	93	3.0	7.9
7	Diabetes mellitus (E10-E14)	40	1.3	3.4
8	Chronic lower respiratory diseases (J40-J47)	29	0.9	2.5
9	Septicemia (A40-A41)	27	0.9	2.3
10	Influenza and pneumonia (J10-J18)	26	0.8	2.2
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	22	0.7	1.9
12	Anemias (D50-D64)	20	0.6	1.7
13	Legal intervention (Y35,Y89.0)	19	0.6	*
14	Cerebrovascular diseases (I60-I69)	18	0.6	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	0.4	*
...	All other causes (Residual)	371	11.9	31.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total		
		Number	deaths	Rate
Black, non-Hispanic, male, 30-34 years				
...	All causes	3,522	100.0	301.2
1	Assault (homicide) (X85-Y09,Y87.1)	745	21.2	63.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	635	18.0	54.3
3	Human immunodeficiency virus (HIV) disease (B20-B24)	614	17.4	52.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	343	9.7	29.3
5	Intentional self-harm (suicide) (X60-X84,Y87.0)	175	5.0	15.0
6	Malignant neoplasms (C00-C97)	166	4.7	14.2
7	Diabetes mellitus (E10-E14)	58	1.6	5.0
8	Cerebrovascular diseases (I60-I69)	52	1.5	4.4
9	Chronic lower respiratory diseases (J40-J47)	41	1.2	3.5
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	37	1.1	3.2
11	Anemias (D50-D64)	35	1.0	3.0
12	Septicemia (A40-A41)	32	0.9	2.7
12	Influenza and pneumonia (J10-J18)	32	0.9	2.7
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	21	0.6	1.8
15	Chronic liver disease and cirrhosis (K70,K73-K74)	15	0.4	*
...	All other causes (Residual)	521	14.8	44.6
Black, non-Hispanic, male, 35-39 years				
...	All causes	4,947	100.0	385.9
1	Human immunodeficiency virus (HIV) disease (B20-B24)	973	19.7	75.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	759	15.3	59.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	667	13.5	52.0
4	Assault (homicide) (X85-Y09,Y87.1)	584	11.8	45.6
5	Malignant neoplasms (C00-C97)	377	7.6	29.4
6	Intentional self-harm (suicide) (X60-X84,Y87.0)	172	3.5	13.4
7	Cerebrovascular diseases (I60-I69)	127	2.6	9.9
8	Diabetes mellitus (E10-E14)	99	2.0	7.7
9	Chronic liver disease and cirrhosis (K70,K73-K74)	90	1.8	7.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	71	1.4	5.5
11	Chronic lower respiratory diseases (J40-J47)	51	1.0	4.0
12	Septicemia (A40-A41)	48	1.0	3.7
13	Influenza and pneumonia (J10-J18)	42	0.8	3.3
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	29	0.6	2.3
15	Anemias (D50-D64)	26	0.5	2.0
...	All other causes (Residual)	832	16.8	64.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total deaths		
		Number	Rate	Rate
Black, non-Hispanic, male, 40-44 years				
...	All causes	7,114	100.0	569.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,305	18.3	104.4
2	Human immunodeficiency virus (HIV) disease (B20-B24)	1,087	15.3	87.0
3	Malignant neoplasms (C00-C97)	915	12.9	73.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	890	12.5	71.2
5	Assault (homicide) (X85-Y09,Y87.1)	443	6.2	35.4
6	Cerebrovascular diseases (I60-I69)	267	3.8	21.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	189	2.7	15.1
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	174	2.4	13.9
9	Diabetes mellitus (E10-E14)	153	2.2	12.2
10	Influenza and pneumonia (J10-J18)	109	1.5	8.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	104	1.5	8.3
12	Chronic lower respiratory diseases (J40-J47)	78	1.1	6.2
13	Septicemia (A40-A41)	77	1.1	6.2
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	69	1.0	5.5
15	Viral hepatitis (B15-B19)	56	0.8	4.5
...	All other causes (Residual)	1,198	16.8	95.9
Black, non-Hispanic, male, 45-49 years				
...	All causes	9,225	100.0	910.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,069	22.4	204.2
2	Malignant neoplasms (C00-C97)	1,648	17.9	162.6
3	Human immunodeficiency virus (HIV) disease (B20-B24)	995	10.8	98.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	808	8.8	79.7
5	Cerebrovascular diseases (I60-I69)	440	4.8	43.4
6	Assault (homicide) (X85-Y09,Y87.1)	310	3.4	30.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	289	3.1	28.5
8	Diabetes mellitus (E10-E14)	277	3.0	27.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	191	2.1	18.9
10	Septicemia (A40-A41)	136	1.5	13.4
11	Chronic lower respiratory diseases (J40-J47)	125	1.4	12.3
12	Influenza and pneumonia (J10-J18)	124	1.3	12.2
13	Intentional self-harm (suicide) (X60-X84,Y87.0)	123	1.3	12.1
14	Viral hepatitis (B15-B19)	119	1.3	11.7
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	96	1.0	9.5
...	All other causes (Residual)	1,475	16.0	145.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total deaths		
		Number	Rate	Rate
Black, non-Hispanic, male, 50-54 years				
...	All causes	10,422	100.0	1,346.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,718	26.1	351.0
2	Malignant neoplasms (C00-C97)	2,504	24.0	323.4
3	Human immunodeficiency virus (HIV) disease (B20-B24)	647	6.2	83.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	646	6.2	83.4
5	Cerebrovascular diseases (I60-I69)	515	4.9	66.5
6	Diabetes mellitus (E10-E14)	371	3.6	47.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	344	3.3	44.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	203	1.9	26.2
9	Assault (homicide) (X85-Y09,Y87.1)	188	1.8	24.3
10	Septicemia (A40-A41)	183	1.8	23.6
11	Chronic lower respiratory diseases (J40-J47)	164	1.6	21.2
12	Influenza and pneumonia (J10-J18)	147	1.4	19.0
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	133	1.3	17.2
14	Viral hepatitis (B15-B19)	115	1.1	14.9
15	Intentional self-harm (suicide) (X60-X84,Y87.0)	87	0.8	11.2
...	All other causes (Residual)	1,457	14.0	188.2
Black, non-Hispanic, male, 55-59 years				
...	All causes	10,649	100.0	1,916.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,193	30.0	574.7
2	Malignant neoplasms (C00-C97)	3,079	28.9	554.2
3	Cerebrovascular diseases (I60-I69)	587	5.5	105.7
4	Diabetes mellitus (E10-E14)	434	4.1	78.1
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	429	4.0	77.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	303	2.8	54.5
7	Chronic lower respiratory diseases (J40-J47)	238	2.2	42.8
8	Chronic liver disease and cirrhosis (K70,K73-K74)	230	2.2	41.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	226	2.1	40.7
10	Septicemia (A40-A41)	165	1.5	29.7
11	Influenza and pneumonia (J10-J18)	155	1.5	27.9
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	135	1.3	24.3
13	Assault (homicide) (X85-Y09,Y87.1)	96	0.9	17.3
14	Viral hepatitis (B15-B19)	61	0.6	11.0
14	Aortic aneurysm and dissection (I71)	61	0.6	11.0
...	All other causes (Residual)	1,257	11.8	226.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 60-64 years				
...	All causes	11,753	100.0	2,658.1
1	Malignant neoplasms (C00-C97)	3,684	31.3	833.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,593	30.6	812.6
3	Cerebrovascular diseases (I60-I69)	653	5.6	147.7
4	Diabetes mellitus (E10-E14)	518	4.4	117.2
5	Chronic lower respiratory diseases (J40-J47)	349	3.0	78.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	307	2.6	69.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	258	2.2	58.3
8	Septicemia (A40-A41)	216	1.8	48.9
9	Human immunodeficiency virus (HIV) disease (B20-B24)	189	1.6	42.7
10	Chronic liver disease and cirrhosis (K70,K73-K74)	178	1.5	40.3
11	Influenza and pneumonia (J10-J18)	167	1.4	37.8
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	166	1.4	37.5
13	Assault (homicide) (X85-Y09,Y87.1)	63	0.5	14.2
14	Aortic aneurysm and dissection (I71)	62	0.5	14.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	49	0.4	11.1
...	All other causes (Residual)	1,301	11.1	294.2
Black, non-Hispanic, male, 65-69 years				
...	All causes	13,355	100.0	3,467.8
1	Malignant neoplasms (C00-C97)	4,332	32.4	1,124.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,929	29.4	1,020.2
3	Cerebrovascular diseases (I60-I69)	812	6.1	210.8
4	Diabetes mellitus (E10-E14)	583	4.4	151.4
5	Chronic lower respiratory diseases (J40-J47)	526	3.9	136.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	330	2.5	85.7
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	282	2.1	73.2
8	Septicemia (A40-A41)	262	2.0	68.0
9	Influenza and pneumonia (J10-J18)	244	1.8	63.4
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	153	1.1	39.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	144	1.1	37.4
12	Human immunodeficiency virus (HIV) disease (B20-B24)	95	0.7	24.7
13	Aortic aneurysm and dissection (I71)	68	0.5	17.7
14	Pneumonitis due to solids and liquids (J69)	67	0.5	17.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	60	0.4	15.6
...	All other causes (Residual)	1,468	11.0	381.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total		
		Number	deaths	Rate
Black, non-Hispanic, male, 70-74 years				
...	All causes	15,390	100.0	5,092.5
1	Malignant neoplasms (C00-C97)	4,903	31.9	1,622.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,631	30.1	1,532.4
3	Cerebrovascular diseases (I60-I69)	1,005	6.5	332.6
4	Chronic lower respiratory diseases (J40-J47)	669	4.3	221.4
5	Diabetes mellitus (E10-E14)	665	4.3	220.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	376	2.4	124.4
7	Influenza and pneumonia (J10-J18)	318	2.1	105.2
8	Septicemia (A40-A41)	283	1.8	93.6
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	255	1.7	84.4
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	191	1.2	63.2
11	Pneumonitis due to solids and liquids (J69)	123	0.8	40.7
12	Chronic liver disease and cirrhosis (K70,K73-K74)	111	0.7	36.7
13	Alzheimer's disease (G30)	83	0.5	27.5
13	Aortic aneurysm and dissection (I71)	83	0.5	27.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	70	0.5	23.2
...	All other causes (Residual)	1,624	10.6	537.4
Black, non-Hispanic, male, 75-79 years				
...	All causes	15,491	100.0	7,297.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,692	30.3	2,210.4
2	Malignant neoplasms (C00-C97)	4,532	29.3	2,135.0
3	Cerebrovascular diseases (I60-I69)	1,138	7.3	536.1
4	Chronic lower respiratory diseases (J40-J47)	681	4.4	320.8
5	Diabetes mellitus (E10-E14)	652	4.2	307.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	418	2.7	196.9
7	Septicemia (A40-A41)	384	2.5	180.9
8	Influenza and pneumonia (J10-J18)	360	2.3	169.6
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	250	1.6	117.8
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	195	1.3	91.9
11	Alzheimer's disease (G30)	136	0.9	64.1
12	Pneumonitis due to solids and liquids (J69)	129	0.8	60.8
13	Aortic aneurysm and dissection (I71)	99	0.6	46.6
14	Parkinson's disease (G20-G21)	78	0.5	36.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	72	0.5	33.9
...	All other causes (Residual)	1,675	10.8	789.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total deaths		
		Number	Rate	Rate
Black, non-Hispanic, male, 80-84 years				
...	All causes	12,224	100.0	10,482.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,784	31.0	3,244.8
2	Malignant neoplasms (C00-C97)	3,157	25.8	2,707.1
3	Cerebrovascular diseases (I60-I69)	1,017	8.3	872.1
4	Chronic lower respiratory diseases (J40-J47)	526	4.3	451.0
5	Diabetes mellitus (E10-E14)	448	3.7	384.2
6	Influenza and pneumonia (J10-J18)	403	3.3	345.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	341	2.8	292.4
8	Septicemia (A40-A41)	293	2.4	251.2
9	Alzheimer's disease (G30)	184	1.5	157.8
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	182	1.5	156.1
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	163	1.3	139.8
12	Pneumonitis due to solids and liquids (J69)	114	0.9	97.8
13	Atherosclerosis (I70)	79	0.6	67.7
14	Parkinson's disease (G20-G21)	76	0.6	65.2
15	Aortic aneurysm and dissection (I71)	54	0.4	46.3
...	All other causes (Residual)	1,403	11.5	1,203.1
Black, non-Hispanic, male, 85-89 years				
...	All causes	8,341	100.0	14,907.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,740	32.8	4,897.1
2	Malignant neoplasms (C00-C97)	1,811	21.7	3,236.8
3	Cerebrovascular diseases (I60-I69)	727	8.7	1,299.4
4	Chronic lower respiratory diseases (J40-J47)	356	4.3	636.3
5	Influenza and pneumonia (J10-J18)	321	3.8	573.7
6	Diabetes mellitus (E10-E14)	258	3.1	461.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	233	2.8	416.4
8	Septicemia (A40-A41)	216	2.6	386.1
9	Alzheimer's disease (G30)	181	2.2	323.5
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	123	1.5	219.8
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	120	1.4	214.5
12	Pneumonitis due to solids and liquids (J69)	88	1.1	157.3
13	Atherosclerosis (I70)	63	0.8	112.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	44	0.5	78.6
14	Parkinson's disease (G20-G21)	44	0.5	78.6
...	All other causes (Residual)	1,016	12.2	1,815.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Black, non-Hispanic, male, 90-94 years				
...	All causes	4,218	100.0	16,960.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,398	33.1	5,621.2
2	Malignant neoplasms (C00-C97)	734	17.4	2,951.3
3	Cerebrovascular diseases (I60-I69)	367	8.7	1,475.7
4	Influenza and pneumonia (J10-J18)	218	5.2	876.6
5	Chronic lower respiratory diseases (J40-J47)	171	4.1	687.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	126	3.0	506.6
7	Septicemia (A40-A41)	109	2.6	438.3
8	Diabetes mellitus (E10-E14)	108	2.6	434.3
9	Alzheimer's disease (G30)	99	2.3	398.1
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	77	1.8	309.6
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	74	1.8	297.5
12	Pneumonitis due to solids and liquids (J69)	66	1.6	265.4
13	Atherosclerosis (I70)	41	1.0	164.9
14	Parkinson's disease (G20-G21)	24	0.6	96.5
15	Anemias (D50-D64)	22	0.5	88.5
...	All other causes (Residual)	584	13.8	2,348.2
Black, non-Hispanic, male, 95-99 years				
...	All causes	1,523	100.0	17,253.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	544	35.7	6,162.9
2	Malignant neoplasms (C00-C97)	219	14.4	2,481.0
3	Cerebrovascular diseases (I60-I69)	145	9.5	1,642.7
4	Influenza and pneumonia (J10-J18)	86	5.6	974.3
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	59	3.9	668.4
6	Chronic lower respiratory diseases (J40-J47)	58	3.8	657.1
7	Alzheimer's disease (G30)	36	2.4	407.8
8	Septicemia (A40-A41)	34	2.2	385.2
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	32	2.1	362.5
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	27	1.8	305.9
11	Diabetes mellitus (E10-E14)	21	1.4	237.9
11	Pneumonitis due to solids and liquids (J69)	21	1.4	237.9
13	Atherosclerosis (I70)	19	1.2	*
14	Nutritional deficiencies (E40-E64)	11	0.7	*
15	Parkinson's disease (G20-G21)	6	0.4	*
...	All other causes (Residual)	205	13.5	2,322.4

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 100 years and over				
...	All causes	349	100.0	20,553.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	153	43.8	9,010.6
2	Malignant neoplasms (C00-C97)	38	10.9	2,237.9
3	Influenza and pneumonia (J10-J18)	25	7.2	1,472.3
4	Cerebrovascular diseases (I60-I69)	21	6.0	1,236.7
5	Septicemia (A40-A41)	9	2.6	*
6	Chronic lower respiratory diseases (J40-J47)	8	2.3	*
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7	2.0	*
8	Alzheimer's disease (G30)	6	1.7	*
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	5	1.4	*
9	Atherosclerosis (I70)	5	1.4	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	1.4	*
12	Pneumonitis due to solids and liquids (J69)	4	1.1	*
13	Diabetes mellitus (E10-E14)	2	0.6	*
14	Anemias (D50-D64)	1	0.3	*
14	Nutritional deficiencies (E40-E64)	1	0.3	*
14	Parkinson's disease (G20-G21)	1	0.3	*
14	Peptic ulcer (K25-K28)	1	0.3	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.3	*
14	Intentional self-harm (suicide) (X60-X84,Y87.0)	1	0.3	*
...	All other causes (Residual)	55	15.8	3,239.1
Black, non-Hispanic, female, all ages				
...	All causes	139,379	100.0	791.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	40,424	29.0	229.5
2	Malignant neoplasms (C00-C97)	28,897	20.7	164.1
3	Cerebrovascular diseases (I60-I69)	11,108	8.0	63.1
4	Diabetes mellitus (E10-E14)	7,198	5.2	40.9
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,808	2.7	21.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,707	2.7	21.0
7	Chronic lower respiratory diseases (J40-J47)	3,344	2.4	19.0
8	Septicemia (A40-A41)	3,304	2.4	18.8
9	Influenza and pneumonia (J10-J18)	3,040	2.2	17.3
10	Human immunodeficiency virus (HIV) disease (B20-B24)	2,417	1.7	13.7
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,286	1.6	13.0
12	Certain conditions originating in the perinatal period (P00-P96)	2,118	1.5	12.0
13	Alzheimer's disease (G30)	1,945	1.4	11.0
14	Assault (homicide) (X85-Y09,Y87.1)	1,366	1.0	7.8
15	Chronic liver disease and cirrhosis (K70,K73-K74)	996	0.7	5.7
...	All other causes (Residual)	23,421	16.8	133.0

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 1-4 years				
...	All causes	545	100.0	53.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	164	30.1	16.1
2	Assault (homicide) (X85-Y09,Y87.1)	70	12.8	6.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	56	10.3	5.5
4	Malignant neoplasms (C00-C97)	35	6.4	3.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	23	4.2	2.3
6	Septicemia (A40-A41)	12	2.2	*
7	Anemias (D50-D64)	11	2.0	*
8	Cerebrovascular diseases (I60-I69)	10	1.8	*
9	Influenza and pneumonia (J10-J18)	9	1.7	*
9	Certain conditions originating in the perinatal period (P00-P96)	9	1.7	*
11	Meningitis (G00,G03)	7	1.3	*
11	Chronic lower respiratory diseases (J40-J47)	7	1.3	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	6	1.1	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	6	1.1	*
15	Acute bronchitis and bronchiolitis (J20-J21)	3	0.6	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.6	*
...	All other causes (Residual)	114	20.9	11.2
Black, non-Hispanic, female, 5-9 years				
...	All causes	320	100.0	22.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	126	39.4	8.8
2	Malignant neoplasms (C00-C97)	29	9.1	2.0
3	Assault (homicide) (X85-Y09,Y87.1)	24	7.5	1.7
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	15	4.7	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	12	3.8	*
6	Chronic lower respiratory diseases (J40-J47)	10	3.1	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	2.2	*
8	Meningococcal infection (A39)	4	1.3	*
8	Septicemia (A40-A41)	4	1.3	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	4	1.3	*
8	Anemias (D50-D64)	4	1.3	*
12	Influenza and pneumonia (J10-J18)	3	0.9	*
12	Certain conditions originating in the perinatal period (P00-P96)	3	0.9	*
14	Cerebrovascular diseases (I60-I69)	2	0.6	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.6	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.6	*
...	All other causes (Residual)	69	21.6	4.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total deaths		
		Number	Rate	Rate
Black, non-Hispanic, female, 10-14 years				
...	All causes	341	100.0	23.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	87	25.5	5.9
2	Malignant neoplasms (C00-C97)	32	9.4	2.2
3	Assault (homicide) (X85-Y09,Y87.1)	28	8.2	1.9
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	24	7.0	1.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	22	6.5	1.5
6	Chronic lower respiratory diseases (J40-J47)	17	5.0	*
7	Anemias (D50-D64)	9	2.6	*
8	Cerebrovascular diseases (I60-I69)	8	2.3	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	7	2.1	*
10	Intentional self-harm (suicide) (X60-X84,Y87.0)	5	1.5	*
11	Diabetes mellitus (E10-E14)	4	1.2	*
11	Certain conditions originating in the perinatal period (P00-P96)	4	1.2	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.9	*
14	Nutritional deficiencies (E40-E64)	2	0.6	*
14	Meningitis (G00,G03)	2	0.6	*
14	Aortic aneurysm and dissection (I71)	2	0.6	*
14	Acute bronchitis and bronchiolitis (J20-J21)	2	0.6	*
...	All other causes (Residual)	83	24.3	5.6
Black, non-Hispanic, female, 15-19 years				
...	All causes	660	100.0	46.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	191	28.9	13.4
2	Assault (homicide) (X85-Y09,Y87.1)	126	19.1	8.9
3	Malignant neoplasms (C00-C97)	53	8.0	3.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	50	7.6	3.5
5	Intentional self-harm (suicide) (X60-X84,Y87.0)	22	3.3	1.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	17	2.6	*
7	Chronic lower respiratory diseases (J40-J47)	13	2.0	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	11	1.7	*
9	Anemias (D50-D64)	9	1.4	*
9	Cerebrovascular diseases (I60-I69)	9	1.4	*
11	Influenza and pneumonia (J10-J18)	8	1.2	*
12	Septicemia (A40-A41)	6	0.9	*
12	Diabetes mellitus (E10-E14)	6	0.9	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.8	*
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	4	0.6	*
15	Pregnancy, childbirth and the puerperium (O00-099)	4	0.6	*
...	All other causes (Residual)	126	19.1	8.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total		
		Number	deaths	Rate
Black, non-Hispanic, female, 20-24 years				
...	All causes	1,042	100.0	78.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	238	22.8	17.9
2	Assault (homicide) (X85-Y09,Y87.1)	186	17.9	14.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	89	8.5	6.7
4	Malignant neoplasms (C00-C97)	66	6.3	5.0
5	Human immunodeficiency virus (HIV) disease (B20-B24)	54	5.2	4.1
6	Intentional self-harm (suicide) (X60-X84,Y87.0)	42	4.0	3.2
7	Pregnancy, childbirth and the puerperium (O00-099)	30	2.9	2.3
8	Anemias (D50-D64)	23	2.2	1.7
9	Diabetes mellitus (E10-E14)	21	2.0	1.6
10	Chronic lower respiratory diseases (J40-J47)	16	1.5	*
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	16	1.5	*
12	Influenza and pneumonia (J10-J18)	15	1.4	*
13	Cerebrovascular diseases (I60-I69)	14	1.3	*
14	Septicemia (A40-A41)	9	0.9	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.9	*
...	All other causes (Residual)	214	20.5	16.1
Black, non-Hispanic, female, 25-29 years				
...	All causes	1,362	100.0	106.6
1	Human immunodeficiency virus (HIV) disease (B20-B24)	197	14.5	15.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	196	14.4	15.3
3	Assault (homicide) (X85-Y09,Y87.1)	153	11.2	12.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	143	10.5	11.2
5	Malignant neoplasms (C00-C97)	120	8.8	9.4
6	Intentional self-harm (suicide) (X60-X84,Y87.0)	40	2.9	3.1
7	Diabetes mellitus (E10-E14)	35	2.6	2.7
8	Pregnancy, childbirth and the puerperium (O00-099)	31	2.3	2.4
9	Anemias (D50-D64)	27	2.0	2.1
10	Cerebrovascular diseases (I60-I69)	23	1.7	1.8
11	Chronic lower respiratory diseases (J40-J47)	20	1.5	1.6
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	1.1	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	14	1.0	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	1.0	*
15	Septicemia (A40-A41)	12	0.9	*
...	All other causes (Residual)	323	23.7	25.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total deaths		
		Number	Rate	Rate
Black, non-Hispanic, female, 30-34 years				
...	All causes	2,092	100.0	157.4
1	Human immunodeficiency virus (HIV) disease (B20-B24)	395	18.9	29.7
2	Malignant neoplasms (C00-C97)	262	12.5	19.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	243	11.6	18.3
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	228	10.9	17.2
5	Assault (homicide) (X85-Y09,Y87.1)	157	7.5	11.8
6	Cerebrovascular diseases (I60-I69)	65	3.1	4.9
7	Diabetes mellitus (E10-E14)	54	2.6	4.1
8	Anemias (D50-D64)	42	2.0	3.2
9	Septicemia (A40-A41)	36	1.7	2.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	34	1.6	2.6
10	Pregnancy, childbirth and the puerperium (O00-099)	34	1.6	2.6
12	Intentional self-harm (suicide) (X60-X84,Y87.0)	32	1.5	2.4
13	Chronic lower respiratory diseases (J40-J47)	24	1.1	1.8
14	Influenza and pneumonia (J10-J18)	23	1.1	1.7
15	Chronic liver disease and cirrhosis (K70,K73-K74)	21	1.0	1.6
...	All other causes (Residual)	442	21.1	33.3
Black, non-Hispanic, female, 35-39 years				
...	All causes	3,300	100.0	226.9
1	Malignant neoplasms (C00-C97)	629	19.1	43.2
2	Human immunodeficiency virus (HIV) disease (B20-B24)	474	14.4	32.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	451	13.7	31.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	283	8.6	19.5
5	Assault (homicide) (X85-Y09,Y87.1)	170	5.2	11.7
6	Cerebrovascular diseases (I60-I69)	154	4.7	10.6
7	Diabetes mellitus (E10-E14)	77	2.3	5.3
8	Septicemia (A40-A41)	61	1.8	4.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	58	1.8	4.0
10	Chronic liver disease and cirrhosis (K70,K73-K74)	52	1.6	3.6
11	Chronic lower respiratory diseases (J40-J47)	51	1.5	3.5
12	Influenza and pneumonia (J10-J18)	41	1.2	2.8
13	Anemias (D50-D64)	39	1.2	2.7
14	Intentional self-harm (suicide) (X60-X84,Y87.0)	33	1.0	2.3
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	32	1.0	2.2
...	All other causes (Residual)	695	21.1	47.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total		
		Number	deaths	Rate
Black, non-Hispanic, female, 40-44 years				
...	All causes	4,918	100.0	345.4
1	Malignant neoplasms (C00-C97)	1,151	23.4	80.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	836	17.0	58.7
3	Human immunodeficiency virus (HIV) disease (B20-B24)	494	10.0	34.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	321	6.5	22.5
5	Cerebrovascular diseases (I60-I69)	281	5.7	19.7
6	Assault (homicide) (X85-Y09,Y87.1)	162	3.3	11.4
7	Diabetes mellitus (E10-E14)	129	2.6	9.1
8	Chronic liver disease and cirrhosis (K70,K73-K74)	115	2.3	8.1
9	Chronic lower respiratory diseases (J40-J47)	92	1.9	6.5
10	Septicemia (A40-A41)	83	1.7	5.8
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	72	1.5	5.1
12	Influenza and pneumonia (J10-J18)	62	1.3	4.4
13	Intentional self-harm (suicide) (X60-X84,Y87.0)	48	1.0	3.4
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	46	0.9	3.2
15	Anemias (D50-D64)	45	0.9	3.2
...	All other causes (Residual)	981	19.9	68.9
Black, non-Hispanic, female, 45-49 years				
...	All causes	6,320	100.0	525.3
1	Malignant neoplasms (C00-C97)	1,750	27.7	145.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,251	19.8	104.0
3	Cerebrovascular diseases (I60-I69)	428	6.8	35.6
4	Human immunodeficiency virus (HIV) disease (B20-B24)	365	5.8	30.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	288	4.6	23.9
6	Diabetes mellitus (E10-E14)	236	3.7	19.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	163	2.6	13.5
8	Chronic lower respiratory diseases (J40-J47)	133	2.1	11.1
9	Septicemia (A40-A41)	122	1.9	10.1
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	109	1.7	9.1
11	Assault (homicide) (X85-Y09,Y87.1)	80	1.3	6.6
12	Influenza and pneumonia (J10-J18)	73	1.2	6.1
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	72	1.1	6.0
14	Viral hepatitis (B15-B19)	51	0.8	4.2
15	Anemias (D50-D64)	34	0.5	2.8
...	All other causes (Residual)	1,165	18.4	96.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total deaths		
		Number	Rate	Rate
Black, non-Hispanic, female, 50-54 years				
...	All causes	6,944	100.0	727.3
1	Malignant neoplasms (C00-C97)	2,166	31.2	226.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,568	22.6	164.2
3	Cerebrovascular diseases (I60-I69)	436	6.3	45.7
4	Diabetes mellitus (E10-E14)	386	5.6	40.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	219	3.2	22.9
6	Human immunodeficiency virus (HIV) disease (B20-B24)	204	2.9	21.4
7	Chronic lower respiratory diseases (J40-J47)	171	2.5	17.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	161	2.3	16.9
9	Chronic liver disease and cirrhosis (K70,K73-K74)	133	1.9	13.9
10	Septicemia (A40-A41)	131	1.9	13.7
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	104	1.5	10.9
12	Influenza and pneumonia (J10-J18)	79	1.1	8.3
13	Viral hepatitis (B15-B19)	44	0.6	4.6
14	Assault (homicide) (X85-Y09,Y87.1)	37	0.5	3.9
15	Anemias (D50-D64)	30	0.4	3.1
...	All other causes (Residual)	1,075	15.5	112.6
Black, non-Hispanic, female, 55-59 years				
...	All causes	7,316	100.0	1,019.7
1	Malignant neoplasms (C00-C97)	2,368	32.4	330.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,883	25.7	262.5
3	Diabetes mellitus (E10-E14)	455	6.2	63.4
4	Cerebrovascular diseases (I60-I69)	445	6.1	62.0
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	207	2.8	28.9
6	Chronic lower respiratory diseases (J40-J47)	205	2.8	28.6
7	Septicemia (A40-A41)	159	2.2	22.2
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	132	1.8	18.4
9	Chronic liver disease and cirrhosis (K70,K73-K74)	111	1.5	15.5
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	105	1.4	14.6
11	Human immunodeficiency virus (HIV) disease (B20-B24)	85	1.2	11.8
12	Influenza and pneumonia (J10-J18)	82	1.1	11.4
13	Aortic aneurysm and dissection (I71)	34	0.5	4.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	30	0.4	4.2
15	Viral hepatitis (B15-B19)	29	0.4	4.0
...	All other causes (Residual)	986	13.5	137.4

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total deaths		
		Number	Rate	Rate
Black, non-Hispanic, female, 60-64 years				
...	All causes	9,149	100.0	1,535.3
1	Malignant neoplasms (C00-C97)	2,888	31.6	484.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,588	28.3	434.3
3	Diabetes mellitus (E10-E14)	588	6.4	98.7
4	Cerebrovascular diseases (I60-I69)	580	6.3	97.3
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	289	3.2	48.5
6	Chronic lower respiratory diseases (J40-J47)	271	3.0	45.5
7	Septicemia (A40-A41)	216	2.4	36.2
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	149	1.6	25.0
9	Influenza and pneumonia (J10-J18)	127	1.4	21.3
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	115	1.3	19.3
11	Chronic liver disease and cirrhosis (K70,K73-K74)	103	1.1	17.3
12	Human immunodeficiency virus (HIV) disease (B20-B24)	48	0.5	8.1
13	Aortic aneurysm and dissection (I71)	39	0.4	6.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	29	0.3	4.9
14	Complications of medical and surgical care (Y40-Y84,Y88)	29	0.3	4.9
...	All other causes (Residual)	1,090	11.9	182.9
Black, non-Hispanic, female, 65-69 years				
...	All causes	11,476	100.0	2,210.1
1	Malignant neoplasms (C00-C97)	3,401	29.6	655.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,373	29.4	649.6
3	Diabetes mellitus (E10-E14)	837	7.3	161.2
4	Cerebrovascular diseases (I60-I69)	761	6.6	146.6
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	418	3.6	80.5
6	Chronic lower respiratory diseases (J40-J47)	346	3.0	66.6
7	Septicemia (A40-A41)	275	2.4	53.0
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	209	1.8	40.2
9	Influenza and pneumonia (J10-J18)	164	1.4	31.6
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	132	1.2	25.4
11	Chronic liver disease and cirrhosis (K70,K73-K74)	102	0.9	19.6
12	Aortic aneurysm and dissection (I71)	61	0.5	11.7
13	Alzheimer's disease (G30)	37	0.3	7.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	36	0.3	6.9
15	Human immunodeficiency virus (HIV) disease (B20-B24)	35	0.3	6.7
15	Pneumonitis due to solids and liquids (J69)	35	0.3	6.7
...	All other causes (Residual)	1,254	10.9	241.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total deaths		
		Number	Rate	Rate
Black, non-Hispanic, female, 70-74 years				
...	All causes	14,424	100.0	3,378.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,480	31.1	1,049.5
2	Malignant neoplasms (C00-C97)	3,849	26.7	901.6
3	Cerebrovascular diseases (I60-I69)	1,082	7.5	253.5
4	Diabetes mellitus (E10-E14)	998	6.9	233.8
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	513	3.6	120.2
6	Chronic lower respiratory diseases (J40-J47)	452	3.1	105.9
7	Septicemia (A40-A41)	353	2.4	82.7
8	Influenza and pneumonia (J10-J18)	258	1.8	60.4
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	248	1.7	58.1
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	188	1.3	44.0
11	Alzheimer's disease (G30)	111	0.8	26.0
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	79	0.5	18.5
13	Chronic liver disease and cirrhosis (K70,K73-K74)	72	0.5	16.9
14	Aortic aneurysm and dissection (I71)	67	0.5	15.7
15	Pneumonitis due to solids and liquids (J69)	64	0.4	15.0
...	All other causes (Residual)	1,610	11.2	377.2
Black, non-Hispanic, female, 75-79 years				
...	All causes	16,728	100.0	5,066.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,535	33.1	1,676.4
2	Malignant neoplasms (C00-C97)	3,732	22.3	1,130.3
3	Cerebrovascular diseases (I60-I69)	1,506	9.0	456.1
4	Diabetes mellitus (E10-E14)	1,117	6.7	338.3
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	505	3.0	152.9
6	Chronic lower respiratory diseases (J40-J47)	495	3.0	149.9
7	Septicemia (A40-A41)	433	2.6	131.1
8	Influenza and pneumonia (J10-J18)	329	2.0	99.6
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	323	1.9	97.8
10	Alzheimer's disease (G30)	263	1.6	79.7
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	169	1.0	51.2
12	Aortic aneurysm and dissection (I71)	89	0.5	27.0
13	Pneumonitis due to solids and liquids (J69)	87	0.5	26.3
14	Atherosclerosis (I70)	78	0.5	23.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	77	0.5	23.3
...	All other causes (Residual)	1,990	11.9	602.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total		
		Number	deaths	Rate
Black, non-Hispanic, female, 80-84 years				
...	All causes	16,431	100.0	7,729.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,589	34.0	2,629.2
2	Malignant neoplasms (C00-C97)	2,823	17.2	1,328.0
3	Cerebrovascular diseases (I60-I69)	1,683	10.2	791.7
4	Diabetes mellitus (E10-E14)	976	5.9	459.1
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	512	3.1	240.9
6	Septicemia (A40-A41)	466	2.8	219.2
7	Influenza and pneumonia (J10-J18)	458	2.8	215.5
8	Chronic lower respiratory diseases (J40-J47)	426	2.6	200.4
9	Alzheimer's disease (G30)	421	2.6	198.1
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	299	1.8	140.7
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	167	1.0	78.6
12	Pneumonitis due to solids and liquids (J69)	138	0.8	64.9
13	Atherosclerosis (I70)	111	0.7	52.2
14	Aortic aneurysm and dissection (I71)	89	0.5	41.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	87	0.5	40.9
...	All other causes (Residual)	2,186	13.3	1,028.4
Black, non-Hispanic, female, 85-89 years				
...	All causes	15,355	100.0	12,606.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,641	36.7	4,631.4
2	Malignant neoplasms (C00-C97)	2,049	13.3	1,682.3
3	Cerebrovascular diseases (I60-I69)	1,745	11.4	1,432.7
4	Diabetes mellitus (E10-E14)	695	4.5	570.6
5	Alzheimer's disease (G30)	499	3.2	409.7
6	Influenza and pneumonia (J10-J18)	494	3.2	405.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	434	2.8	356.3
8	Septicemia (A40-A41)	415	2.7	340.7
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	312	2.0	256.2
10	Chronic lower respiratory diseases (J40-J47)	307	2.0	252.1
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	159	1.0	130.5
12	Pneumonitis due to solids and liquids (J69)	144	0.9	118.2
13	Atherosclerosis (I70)	109	0.7	89.5
14	Aortic aneurysm and dissection (I71)	78	0.5	64.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	72	0.5	59.1
...	All other causes (Residual)	2,202	14.3	1,807.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total deaths		
		Number	Rate	Rate
Black, non-Hispanic, female, 90-94 years				
...	All causes	10,361	100.0	16,216.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,969	38.3	6,212.0
2	Cerebrovascular diseases (I60-I69)	1,140	11.0	1,784.3
3	Malignant neoplasms (C00-C97)	1,035	10.0	1,619.9
4	Influenza and pneumonia (J10-J18)	435	4.2	680.8
5	Diabetes mellitus (E10-E14)	397	3.8	621.4
6	Alzheimer's disease (G30)	364	3.5	569.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	285	2.8	446.1
8	Septicemia (A40-A41)	269	2.6	421.0
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	207	2.0	324.0
10	Chronic lower respiratory diseases (J40-J47)	187	1.8	292.7
11	Atherosclerosis (I70)	112	1.1	175.3
11	Pneumonitis due to solids and liquids (J69)	112	1.1	175.3
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	112	1.1	175.3
14	Nutritional deficiencies (E40-E64)	48	0.5	75.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	44	0.4	68.9
...	All other causes (Residual)	1,645	15.9	2,574.7
Black, non-Hispanic, female, 95-99 years				
...	All causes	4,839	100.0	17,780.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,935	40.0	7,109.8
2	Cerebrovascular diseases (I60-I69)	540	11.2	1,984.1
3	Malignant neoplasms (C00-C97)	372	7.7	1,366.8
4	Influenza and pneumonia (J10-J18)	225	4.6	826.7
5	Alzheimer's disease (G30)	182	3.8	668.7
6	Septicemia (A40-A41)	151	3.1	554.8
7	Diabetes mellitus (E10-E14)	139	2.9	510.7
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	115	2.4	422.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	109	2.3	400.5
10	Chronic lower respiratory diseases (J40-J47)	73	1.5	268.2
11	Atherosclerosis (I70)	57	1.2	209.4
12	Pneumonitis due to solids and liquids (J69)	53	1.1	194.7
13	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	48	1.0	176.4
14	Nutritional deficiencies (E40-E64)	29	0.6	106.6
15	Anemias (D50-D64)	23	0.5	84.5
...	All other causes (Residual)	788	16.3	2,895.4

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups,
by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
	Black, non-Hispanic, female, 100 years and over			
...	All causes	1,674	100.0	25,394.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	679	40.6	10,300.4
2	Cerebrovascular diseases (I60-I69)	176	10.5	2,669.9
3	Influenza and pneumonia (J10-J18)	97	5.8	1,471.5
4	Malignant neoplasms (C00-C97)	79	4.7	1,198.4
5	Alzheimer's disease (G30)	51	3.0	773.7
6	Diabetes mellitus (E10-E14)	47	2.8	713.0
7	Septicemia (A40-A41)	42	2.5	637.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	39	2.3	591.6
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	36	2.2	546.1
10	Atherosclerosis (I70)	35	2.1	530.9
11	Chronic lower respiratory diseases (J40-J47)	23	1.4	348.9
12	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	21	1.3	318.6
13	Pneumonitis due to solids and liquids (J69)	17	1.0	*
14	Nutritional deficiencies (E40-E64)	15	0.9	*
15	Anemias (D50-D64)	9	0.5	*
...	All other causes (Residual)	308	18.4	4,672.3
14	Nutritional deficiencies (E40-E64)	12	0.8	*
...	All other causes (Residual)	279	18.2	4,788.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable