

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|---------------------------------|---|---------|-------------------------|-------|
| Hispanic, both sexes, all ages | | | | |
| ... | All causes | 113,413 | 100.0 | 306.8 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 27,090 | 23.9 | 73.3 |
| 2 | Malignant neoplasms (C00-C97) | 22,371 | 19.7 | 60.5 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 9,523 | 8.4 | 25.8 |
| 4 | Cerebrovascular diseases (I60-I69) | 6,416 | 5.7 | 17.4 |
| 5 | Diabetes mellitus (E10-E14) | 5,663 | 5.0 | 15.3 |
| 6 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 3,331 | 2.9 | 9.0 |
| 7 | Chronic liver disease and cirrhosis (K70,K73-K74) | 3,301 | 2.9 | 8.9 |
| 8 | Chronic lower respiratory diseases (J40-J47) | 2,832 | 2.5 | 7.7 |
| 9 | Influenza and pneumonia (J10-J18) | 2,722 | 2.4 | 7.4 |
| 10 | Certain conditions originating in the perinatal period (P00-P96) | 2,227 | 2.0 | 6.0 |
| 11 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1,857 | 1.6 | 5.0 |
| 12 | Human immunodeficiency virus (HIV) disease (B20-B24) | 1,853 | 1.6 | 5.0 |
| 13 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 1,850 | 1.6 | 5.0 |
| 14 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 1,632 | 1.4 | 4.4 |
| 15 | Septicemia (A40-A41) | 1,386 | 1.2 | 3.7 |
| ... | All other causes (Residual) | 19,359 | 17.1 | 52.4 |
| Hispanic, both sexes, 1-4 years | | | | |
| ... | All causes | 917 | 100.0 | 30.6 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 310 | 33.8 | 10.3 |
| 2 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 103 | 11.2 | 3.4 |
| 3 | Malignant neoplasms (C00-C97) | 84 | 9.2 | 2.8 |
| 4 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 69 | 7.5 | 2.3 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 33 | 3.6 | 1.1 |
| 6 | Influenza and pneumonia (J10-J18) | 32 | 3.5 | 1.1 |
| 7 | Septicemia (A40-A41) | 18 | 2.0 | * |
| 8 | Certain conditions originating in the perinatal period (P00-P96) | 14 | 1.5 | * |
| 9 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 13 | 1.4 | * |
| 10 | Cerebrovascular diseases (I60-I69) | 10 | 1.1 | * |
| 11 | Chronic lower respiratory diseases (J40-J47) | 8 | 0.9 | * |
| 12 | Complications of medical and surgical care (Y40-Y84,Y88) | 6 | 0.7 | * |
| 13 | Meningococcal infection (A39) | 5 | 0.5 | * |
| 14 | Meningitis (G00,G03) | 4 | 0.4 | * |
| 14 | Diseases of appendix (K35-K38) | 4 | 0.4 | * |
| ... | All other causes (Residual) | 204 | 22.2 | 6.8 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|-----------------------------------|---|--------|-------------------------|------|
| Hispanic, both sexes, 5-9 years | | | | |
| ... | All causes | 480 | 100.0 | 13.0 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 186 | 38.8 | 5.0 |
| 2 | Malignant neoplasms (C00-C97) | 87 | 18.1 | 2.4 |
| 3 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 28 | 5.8 | 0.8 |
| 4 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 20 | 4.2 | 0.5 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 17 | 3.5 | * |
| 6 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 14 | 2.9 | * |
| 7 | Septicemia (A40-A41) | 8 | 1.7 | * |
| 8 | Cerebrovascular diseases (I60-I69) | 7 | 1.5 | * |
| 9 | Influenza and pneumonia (J10-J18) | 5 | 1.0 | * |
| 10 | Certain conditions originating in the perinatal period (P00-P96) | 3 | 0.6 | * |
| 11 | Meningococcal infection (A39) | 2 | 0.4 | * |
| 11 | Viral hepatitis (B15-B19) | 2 | 0.4 | * |
| 11 | Anemias (D50-D64) | 2 | 0.4 | * |
| 11 | Meningitis (G00,G03) | 2 | 0.4 | * |
| 11 | Chronic lower respiratory diseases (J40-J47) | 2 | 0.4 | * |
| 11 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 2 | 0.4 | * |
| ... | All other causes (Residual) | 93 | 19.4 | 2.5 |
| Hispanic, both sexes, 10-14 years | | | | |
| ... | All causes | 558 | 100.0 | 16.6 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 193 | 34.6 | 5.7 |
| 2 | Malignant neoplasms (C00-C97) | 90 | 16.1 | 2.7 |
| 3 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 37 | 6.6 | 1.1 |
| 4 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 36 | 6.5 | 1.1 |
| 5 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 30 | 5.4 | 0.9 |
| 6 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 21 | 3.8 | 0.6 |
| 7 | Septicemia (A40-A41) | 10 | 1.8 | * |
| 8 | Cerebrovascular diseases (I60-I69) | 9 | 1.6 | * |
| 9 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 7 | 1.3 | * |
| 9 | Chronic lower respiratory diseases (J40-J47) | 7 | 1.3 | * |
| 11 | Influenza and pneumonia (J10-J18) | 5 | 0.9 | * |
| 12 | Pneumonitis due to solids and liquids (J69) | 4 | 0.7 | * |
| 13 | Anemias (D50-D64) | 3 | 0.5 | * |
| 13 | Complications of medical and surgical care (Y40-Y84,Y88) | 3 | 0.5 | * |
| 15 | Human immunodeficiency virus (HIV) disease (B20-B24) | 2 | 0.4 | * |
| 15 | Diabetes mellitus (E10-E14) | 2 | 0.4 | * |
| 15 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 2 | 0.4 | * |
| ... | All other causes (Residual) | 97 | 17.4 | 2.9 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|-----------------------------------|---|--------|-------------------------|------|
| Hispanic, both sexes, 15-19 years | | | | |
| ... | All causes | 1,998 | 100.0 | 63.1 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 879 | 44.0 | 27.8 |
| 2 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 436 | 21.8 | 13.8 |
| 3 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 166 | 8.3 | 5.2 |
| 4 | Malignant neoplasms (C00-C97) | 143 | 7.2 | 4.5 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 45 | 2.3 | 1.4 |
| 6 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 32 | 1.6 | 1.0 |
| 7 | Cerebrovascular diseases (I60-I69) | 13 | 0.7 | * |
| 8 | Pregnancy, childbirth and the puerperium (O00-099) | 9 | 0.5 | * |
| 9 | Diabetes mellitus (E10-E14) | 8 | 0.4 | * |
| 10 | Influenza and pneumonia (J10-J18) | 7 | 0.4 | * |
| 10 | Chronic lower respiratory diseases (J40-J47) | 7 | 0.4 | * |
| 10 | Legal intervention (Y35,Y89.0) | 7 | 0.4 | * |
| 13 | Anemias (D50-D64) | 6 | 0.3 | * |
| 14 | Septicemia (A40-A41) | 5 | 0.3 | * |
| 15 | Tuberculosis (A16-A19) | 4 | 0.2 | * |
| 15 | Human immunodeficiency virus (HIV) disease (B20-B24) | 4 | 0.2 | * |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 4 | 0.2 | * |
| 15 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 4 | 0.2 | * |
| 15 | Complications of medical and surgical care (Y40-Y84,Y88) | 4 | 0.2 | * |
| ... | All other causes (Residual) | 215 | 10.8 | 6.8 |
| Hispanic, both sexes, 20-24 years | | | | |
| ... | All causes | 3,060 | 100.0 | 86.9 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,303 | 42.6 | 37.0 |
| 2 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 786 | 25.7 | 22.3 |
| 3 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 248 | 8.1 | 7.0 |
| 4 | Malignant neoplasms (C00-C97) | 146 | 4.8 | 4.1 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 67 | 2.2 | 1.9 |
| 6 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 41 | 1.3 | 1.2 |
| 7 | Human immunodeficiency virus (HIV) disease (B20-B24) | 28 | 0.9 | 0.8 |
| 8 | Cerebrovascular diseases (I60-I69) | 27 | 0.9 | 0.8 |
| 9 | Pregnancy, childbirth and the puerperium (O00-099) | 17 | 0.6 | * |
| 10 | Legal intervention (Y35,Y89.0) | 16 | 0.5 | * |
| 11 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 12 | 0.4 | * |
| 12 | Diabetes mellitus (E10-E14) | 11 | 0.4 | * |
| 12 | Influenza and pneumonia (J10-J18) | 11 | 0.4 | * |
| 14 | Septicemia (A40-A41) | 8 | 0.3 | * |
| 15 | Anemias (D50-D64) | 5 | 0.2 | * |
| ... | All other causes (Residual) | 334 | 10.9 | 9.5 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|-----------------------------------|---|--------|-------------------------|------|
| Hispanic, both sexes, 25-29 years | | | | |
| ... | All causes | 2,863 | 100.0 | 80.6 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,073 | 37.5 | 30.2 |
| 2 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 573 | 20.0 | 16.1 |
| 3 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 259 | 9.0 | 7.3 |
| 4 | Malignant neoplasms (C00-C97) | 204 | 7.1 | 5.7 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 115 | 4.0 | 3.2 |
| 6 | Human immunodeficiency virus (HIV) disease (B20-B24) | 80 | 2.8 | 2.3 |
| 7 | Cerebrovascular diseases (I60-I69) | 34 | 1.2 | 1.0 |
| 8 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 26 | 0.9 | 0.7 |
| 9 | Pregnancy, childbirth and the puerperium (O00-099) | 21 | 0.7 | 0.6 |
| 10 | Diabetes mellitus (E10-E14) | 20 | 0.7 | 0.6 |
| 11 | Chronic liver disease and cirrhosis (K70,K73-K74) | 16 | 0.6 | * |
| 12 | Legal intervention (Y35,Y89.0) | 15 | 0.5 | * |
| 13 | Influenza and pneumonia (J10-J18) | 10 | 0.3 | * |
| 13 | Chronic lower respiratory diseases (J40-J47) | 10 | 0.3 | * |
| 13 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 10 | 0.3 | * |
| ... | All other causes (Residual) | 397 | 13.9 | 11.2 |
| Hispanic, both sexes, 30-34 years | | | | |
| ... | All causes | 2,934 | 100.0 | 88.7 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 865 | 29.5 | 26.1 |
| 2 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 387 | 13.2 | 11.7 |
| 3 | Malignant neoplasms (C00-C97) | 293 | 10.0 | 8.9 |
| 4 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 231 | 7.9 | 7.0 |
| 5 | Human immunodeficiency virus (HIV) disease (B20-B24) | 220 | 7.5 | 6.6 |
| 6 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 199 | 6.8 | 6.0 |
| 7 | Cerebrovascular diseases (I60-I69) | 65 | 2.2 | 2.0 |
| 7 | Chronic liver disease and cirrhosis (K70,K73-K74) | 65 | 2.2 | 2.0 |
| 9 | Diabetes mellitus (E10-E14) | 29 | 1.0 | 0.9 |
| 10 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 19 | 0.6 | * |
| 11 | Pregnancy, childbirth and the puerperium (O00-099) | 17 | 0.6 | * |
| 12 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 15 | 0.5 | * |
| 13 | Septicemia (A40-A41) | 14 | 0.5 | * |
| 14 | Influenza and pneumonia (J10-J18) | 13 | 0.4 | * |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 12 | 0.4 | * |
| 15 | Chronic lower respiratory diseases (J40-J47) | 12 | 0.4 | * |
| ... | All other causes (Residual) | 478 | 16.3 | 14.4 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|-----------------------------------|---|--------|-------------------------|-------|
| Hispanic, both sexes, 35-39 years | | | | |
| ... | All causes | 3,859 | 100.0 | 129.5 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 942 | 24.4 | 31.6 |
| 2 | Malignant neoplasms (C00-C97) | 545 | 14.1 | 18.3 |
| 3 | Human immunodeficiency virus (HIV) disease (B20-B24) | 355 | 9.2 | 11.9 |
| 4 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 329 | 8.5 | 11.0 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 302 | 7.8 | 10.1 |
| 6 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 224 | 5.8 | 7.5 |
| 7 | Chronic liver disease and cirrhosis (K70,K73-K74) | 187 | 4.8 | 6.3 |
| 8 | Cerebrovascular diseases (I60-I69) | 102 | 2.6 | 3.4 |
| 9 | Diabetes mellitus (E10-E14) | 69 | 1.8 | 2.3 |
| 10 | Viral hepatitis (B15-B19) | 35 | 0.9 | 1.2 |
| 11 | Influenza and pneumonia (J10-J18) | 32 | 0.8 | 1.1 |
| 12 | Septicemia (A40-A41) | 28 | 0.7 | 0.9 |
| 13 | Chronic lower respiratory diseases (J40-J47) | 25 | 0.6 | 0.8 |
| 14 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 24 | 0.6 | 0.8 |
| 15 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 19 | 0.5 | * |
| ... | All other causes (Residual) | 641 | 16.6 | 21.5 |
| Hispanic, both sexes, 40-44 years | | | | |
| ... | All causes | 4,582 | 100.0 | 185.3 |
| 1 | Malignant neoplasms (C00-C97) | 851 | 18.6 | 34.4 |
| 2 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 818 | 17.9 | 33.1 |
| 3 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 448 | 9.8 | 18.1 |
| 4 | Human immunodeficiency virus (HIV) disease (B20-B24) | 404 | 8.8 | 16.3 |
| 5 | Chronic liver disease and cirrhosis (K70,K73-K74) | 343 | 7.5 | 13.9 |
| 6 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 238 | 5.2 | 9.6 |
| 7 | Cerebrovascular diseases (I60-I69) | 173 | 3.8 | 7.0 |
| 7 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 173 | 3.8 | 7.0 |
| 9 | Diabetes mellitus (E10-E14) | 98 | 2.1 | 4.0 |
| 10 | Viral hepatitis (B15-B19) | 91 | 2.0 | 3.7 |
| 11 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 50 | 1.1 | 2.0 |
| 12 | Septicemia (A40-A41) | 44 | 1.0 | 1.8 |
| 13 | Influenza and pneumonia (J10-J18) | 41 | 0.9 | 1.7 |
| 14 | Chronic lower respiratory diseases (J40-J47) | 27 | 0.6 | 1.1 |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 25 | 0.5 | 1.0 |
| ... | All other causes (Residual) | 758 | 16.5 | 30.7 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|-----------------------------------|---|--------|-------------------------|-------|
| Hispanic, both sexes, 45-49 years | | | | |
| ... | All causes | 5,397 | 100.0 | 280.9 |
| 1 | Malignant neoplasms (C00-C97) | 1,186 | 22.0 | 61.7 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 736 | 13.6 | 38.3 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 627 | 11.6 | 32.6 |
| 4 | Chronic liver disease and cirrhosis (K70,K73-K74) | 480 | 8.9 | 25.0 |
| 5 | Human immunodeficiency virus (HIV) disease (B20-B24) | 316 | 5.9 | 16.4 |
| 6 | Cerebrovascular diseases (I60-I69) | 239 | 4.4 | 12.4 |
| 7 | Diabetes mellitus (E10-E14) | 229 | 4.2 | 11.9 |
| 8 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 156 | 2.9 | 8.1 |
| 9 | Viral hepatitis (B15-B19) | 155 | 2.9 | 8.1 |
| 10 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 130 | 2.4 | 6.8 |
| 11 | Septicemia (A40-A41) | 70 | 1.3 | 3.6 |
| 12 | Influenza and pneumonia (J10-J18) | 56 | 1.0 | 2.9 |
| 13 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 51 | 0.9 | 2.7 |
| 14 | Chronic lower respiratory diseases (J40-J47) | 47 | 0.9 | 2.4 |
| 15 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 28 | 0.5 | 1.5 |
| ... | All other causes (Residual) | 891 | 16.5 | 46.4 |
| Hispanic, both sexes, 50-54 years | | | | |
| ... | All causes | 5,965 | 100.0 | 403.9 |
| 1 | Malignant neoplasms (C00-C97) | 1,629 | 27.3 | 110.3 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 1,139 | 19.1 | 77.1 |
| 3 | Chronic liver disease and cirrhosis (K70,K73-K74) | 471 | 7.9 | 31.9 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 451 | 7.6 | 30.5 |
| 5 | Cerebrovascular diseases (I60-I69) | 300 | 5.0 | 20.3 |
| 6 | Diabetes mellitus (E10-E14) | 278 | 4.7 | 18.8 |
| 7 | Human immunodeficiency virus (HIV) disease (B20-B24) | 210 | 3.5 | 14.2 |
| 8 | Viral hepatitis (B15-B19) | 111 | 1.9 | 7.5 |
| 9 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 109 | 1.8 | 7.4 |
| 10 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 92 | 1.5 | 6.2 |
| 11 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 81 | 1.4 | 5.5 |
| 12 | Chronic lower respiratory diseases (J40-J47) | 77 | 1.3 | 5.2 |
| 13 | Influenza and pneumonia (J10-J18) | 65 | 1.1 | 4.4 |
| 14 | Septicemia (A40-A41) | 63 | 1.1 | 4.3 |
| 15 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 32 | 0.5 | 2.2 |
| ... | All other causes (Residual) | 857 | 14.4 | 58.0 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|-----------------------------------|---|--------|-------------------------|-------|
| Hispanic, both sexes, 55-59 years | | | | |
| ... | All causes | 6,114 | 100.0 | 580.6 |
| 1 | Malignant neoplasms (C00-C97) | 1,814 | 29.7 | 172.3 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 1,362 | 22.3 | 129.3 |
| 3 | Diabetes mellitus (E10-E14) | 404 | 6.6 | 38.4 |
| 4 | Chronic liver disease and cirrhosis (K70,K73-K74) | 378 | 6.2 | 35.9 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 313 | 5.1 | 29.7 |
| 6 | Cerebrovascular diseases (I60-I69) | 301 | 4.9 | 28.6 |
| 7 | Viral hepatitis (B15-B19) | 117 | 1.9 | 11.1 |
| 7 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 117 | 1.9 | 11.1 |
| 9 | Human immunodeficiency virus (HIV) disease (B20-B24) | 111 | 1.8 | 10.5 |
| 10 | Chronic lower respiratory diseases (J40-J47) | 100 | 1.6 | 9.5 |
| 11 | Influenza and pneumonia (J10-J18) | 72 | 1.2 | 6.8 |
| 12 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 65 | 1.1 | 6.2 |
| 13 | Septicemia (A40-A41) | 57 | 0.9 | 5.4 |
| 14 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 46 | 0.8 | 4.4 |
| 15 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 31 | 0.5 | 2.9 |
| ... | All other causes (Residual) | 826 | 13.5 | 78.4 |
| Hispanic, both sexes, 60-64 years | | | | |
| ... | All causes | 7,223 | 100.0 | 910.1 |
| 1 | Malignant neoplasms (C00-C97) | 2,312 | 32.0 | 291.3 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 1,790 | 24.8 | 225.5 |
| 3 | Diabetes mellitus (E10-E14) | 578 | 8.0 | 72.8 |
| 4 | Cerebrovascular diseases (I60-I69) | 353 | 4.9 | 44.5 |
| 5 | Chronic liver disease and cirrhosis (K70,K73-K74) | 343 | 4.7 | 43.2 |
| 6 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 251 | 3.5 | 31.6 |
| 7 | Chronic lower respiratory diseases (J40-J47) | 158 | 2.2 | 19.9 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 120 | 1.7 | 15.1 |
| 9 | Influenza and pneumonia (J10-J18) | 96 | 1.3 | 12.1 |
| 10 | Septicemia (A40-A41) | 94 | 1.3 | 11.8 |
| 11 | Viral hepatitis (B15-B19) | 75 | 1.0 | 9.5 |
| 12 | Human immunodeficiency virus (HIV) disease (B20-B24) | 58 | 0.8 | 7.3 |
| 13 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 52 | 0.7 | 6.6 |
| 14 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 49 | 0.7 | 6.2 |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 42 | 0.6 | 5.3 |
| ... | All other causes (Residual) | 852 | 11.8 | 107.4 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|-----------------------------------|---|--------|-------------------------|---------|
| Hispanic, both sexes, 65-69 years | | | | |
| ... | All causes | 8,910 | 100.0 | 1,412.8 |
| 1 | Malignant neoplasms (C00-C97) | 2,608 | 29.3 | 413.5 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 2,466 | 27.7 | 391.0 |
| 3 | Diabetes mellitus (E10-E14) | 764 | 8.6 | 121.1 |
| 4 | Cerebrovascular diseases (I60-I69) | 471 | 5.3 | 74.7 |
| 5 | Chronic liver disease and cirrhosis (K70,K73-K74) | 322 | 3.6 | 51.1 |
| 6 | Chronic lower respiratory diseases (J40-J47) | 265 | 3.0 | 42.0 |
| 7 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 218 | 2.4 | 34.6 |
| 8 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 205 | 2.3 | 32.5 |
| 9 | Influenza and pneumonia (J10-J18) | 140 | 1.6 | 22.2 |
| 10 | Septicemia (A40-A41) | 129 | 1.4 | 20.5 |
| 11 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 72 | 0.8 | 11.4 |
| 12 | Viral hepatitis (B15-B19) | 52 | 0.6 | 8.2 |
| 13 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 47 | 0.5 | 7.5 |
| 14 | Aortic aneurysm and dissection (I71) | 45 | 0.5 | 7.1 |
| 15 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 42 | 0.5 | 6.7 |
| ... | All other causes (Residual) | 1,064 | 11.9 | 168.7 |
| Hispanic, both sexes, 70-74 years | | | | |
| ... | All causes | 11,262 | 100.0 | 2,229.5 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 3,264 | 29.0 | 646.2 |
| 2 | Malignant neoplasms (C00-C97) | 3,116 | 27.7 | 616.9 |
| 3 | Diabetes mellitus (E10-E14) | 840 | 7.5 | 166.3 |
| 4 | Cerebrovascular diseases (I60-I69) | 735 | 6.5 | 145.5 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 369 | 3.3 | 73.1 |
| 6 | Chronic liver disease and cirrhosis (K70,K73-K74) | 296 | 2.6 | 58.6 |
| 7 | Influenza and pneumonia (J10-J18) | 262 | 2.3 | 51.9 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 252 | 2.2 | 49.9 |
| 9 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 249 | 2.2 | 49.3 |
| 10 | Septicemia (A40-A41) | 159 | 1.4 | 31.5 |
| 11 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 115 | 1.0 | 22.8 |
| 12 | Alzheimer's disease (G30) | 80 | 0.7 | 15.8 |
| 13 | Parkinson's disease (G20-G21) | 68 | 0.6 | 13.5 |
| 14 | Viral hepatitis (B15-B19) | 61 | 0.5 | 12.1 |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 53 | 0.5 | 10.5 |
| ... | All other causes (Residual) | 1,343 | 11.9 | 265.9 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|-----------------------------------|---|--------|-------------------------|---------|
| Hispanic, both sexes, 75-79 years | | | | |
| ... | All causes | 12,560 | 100.0 | 3,556.3 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 3,875 | 30.9 | 1,097.2 |
| 2 | Malignant neoplasms (C00-C97) | 2,988 | 23.8 | 846.0 |
| 3 | Cerebrovascular diseases (I60-I69) | 959 | 7.6 | 271.5 |
| 4 | Diabetes mellitus (E10-E14) | 899 | 7.2 | 254.5 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 513 | 4.1 | 145.3 |
| 6 | Influenza and pneumonia (J10-J18) | 346 | 2.8 | 98.0 |
| 7 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 296 | 2.4 | 83.8 |
| 8 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 222 | 1.8 | 62.9 |
| 9 | Chronic liver disease and cirrhosis (K70,K73-K74) | 219 | 1.7 | 62.0 |
| 10 | Alzheimer's disease (G30) | 194 | 1.5 | 54.9 |
| 11 | Septicemia (A40-A41) | 191 | 1.5 | 54.1 |
| 12 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 135 | 1.1 | 38.2 |
| 13 | Parkinson's disease (G20-G21) | 108 | 0.9 | 30.6 |
| 14 | Aortic aneurysm and dissection (I71) | 75 | 0.6 | 21.2 |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 67 | 0.5 | 19.0 |
| 15 | Pneumonitis due to solids and liquids (J69) | 67 | 0.5 | 19.0 |
| ... | All other causes (Residual) | 1,406 | 11.2 | 398.1 |
| Hispanic, both sexes, 80-84 years | | | | |
| ... | All causes | 11,077 | 100.0 | 5,501.5 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 3,757 | 33.9 | 1,866.0 |
| 2 | Malignant neoplasms (C00-C97) | 2,080 | 18.8 | 1,033.1 |
| 3 | Cerebrovascular diseases (I60-I69) | 933 | 8.4 | 463.4 |
| 4 | Diabetes mellitus (E10-E14) | 649 | 5.9 | 322.3 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 427 | 3.9 | 212.1 |
| 6 | Influenza and pneumonia (J10-J18) | 415 | 3.7 | 206.1 |
| 7 | Alzheimer's disease (G30) | 271 | 2.4 | 134.6 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 228 | 2.1 | 113.2 |
| 9 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 187 | 1.7 | 92.9 |
| 10 | Septicemia (A40-A41) | 155 | 1.4 | 77.0 |
| 11 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 136 | 1.2 | 67.5 |
| 12 | Parkinson's disease (G20-G21) | 122 | 1.1 | 60.6 |
| 13 | Chronic liver disease and cirrhosis (K70,K73-K74) | 116 | 1.0 | 57.6 |
| 14 | Pneumonitis due to solids and liquids (J69) | 85 | 0.8 | 42.2 |
| 15 | Atherosclerosis (I70) | 68 | 0.6 | 33.8 |
| ... | All other causes (Residual) | 1,448 | 13.1 | 719.2 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|-----------------------------------|---|--------|-------------------------|------|
| Hispanic, both sexes, 85-89 years | | | | |
| ... | All causes | 9,601 | 100.0 | --- |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 3,506 | 36.5 | --- |
| 2 | Malignant neoplasms (C00-C97) | 1,322 | 13.8 | --- |
| 3 | Cerebrovascular diseases (I60-I69) | 868 | 9.0 | --- |
| 4 | Diabetes mellitus (E10-E14) | 477 | 5.0 | --- |
| 5 | Influenza and pneumonia (J10-J18) | 465 | 4.8 | --- |
| 6 | Chronic lower respiratory diseases (J40-J47) | 399 | 4.2 | --- |
| 7 | Alzheimer's disease (G30) | 313 | 3.3 | --- |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 180 | 1.9 | --- |
| 9 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 140 | 1.5 | --- |
| 10 | Septicemia (A40-A41) | 136 | 1.4 | --- |
| 11 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 129 | 1.3 | --- |
| 12 | Pneumonitis due to solids and liquids (J69) | 112 | 1.2 | --- |
| 13 | Atherosclerosis (I70) | 99 | 1.0 | --- |
| 14 | Parkinson's disease (G20-G21) | 94 | 1.0 | --- |
| 15 | Chronic liver disease and cirrhosis (K70,K73-K74) | 46 | 0.5 | --- |
| ... | All other causes (Residual) | 1,315 | 13.7 | --- |
| Hispanic, both sexes, 90-94 years | | | | |
| ... | All causes | 6,177 | 100.0 | --- |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 2,531 | 41.0 | --- |
| 2 | Malignant neoplasms (C00-C97) | 615 | 10.0 | --- |
| 3 | Cerebrovascular diseases (I60-I69) | 532 | 8.6 | --- |
| 4 | Influenza and pneumonia (J10-J18) | 382 | 6.2 | --- |
| 5 | Chronic lower respiratory diseases (J40-J47) | 271 | 4.4 | --- |
| 6 | Diabetes mellitus (E10-E14) | 212 | 3.4 | --- |
| 7 | Alzheimer's disease (G30) | 197 | 3.2 | --- |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 121 | 2.0 | --- |
| 9 | Pneumonitis due to solids and liquids (J69) | 97 | 1.6 | --- |
| 10 | Septicemia (A40-A41) | 93 | 1.5 | --- |
| 11 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 84 | 1.4 | --- |
| 12 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 82 | 1.3 | --- |
| 13 | Atherosclerosis (I70) | 59 | 1.0 | --- |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 34 | 0.6 | --- |
| 14 | Parkinson's disease (G20-G21) | 34 | 0.6 | --- |
| ... | All other causes (Residual) | 833 | 13.5 | --- |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|------|
| Hispanic, both sexes, 95-99 years | | | | |
| ... | All causes | 2,582 | 100.0 | --- |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 1,080 | 41.8 | --- |
| 2 | Malignant neoplasms (C00-C97) | 222 | 8.6 | --- |
| 3 | Cerebrovascular diseases (I60-I69) | 221 | 8.6 | --- |
| 4 | Influenza and pneumonia (J10-J18) | 173 | 6.7 | --- |
| 5 | Alzheimer's disease (G30) | 106 | 4.1 | --- |
| 6 | Diabetes mellitus (E10-E14) | 88 | 3.4 | --- |
| 7 | Chronic lower respiratory diseases (J40-J47) | 80 | 3.1 | --- |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 44 | 1.7 | --- |
| 9 | Pneumonitis due to solids and liquids (J69) | 41 | 1.6 | --- |
| 10 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 40 | 1.5 | --- |
| 11 | Atherosclerosis (I70) | 39 | 1.5 | --- |
| 12 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 36 | 1.4 | --- |
| 13 | Septicemia (A40-A41) | 35 | 1.4 | --- |
| 14 | Nutritional deficiencies (E40-E64) | 12 | 0.5 | --- |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 11 | 0.4 | --- |
| ... | All other causes (Residual) | 354 | 13.7 | --- |
| Hispanic, both sexes, 100 years and over | | | | |
| ... | All causes | 547 | 100.0 | --- |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 252 | 46.1 | --- |
| 2 | Cerebrovascular diseases (I60-I69) | 47 | 8.6 | --- |
| 3 | Influenza and pneumonia (J10-J18) | 36 | 6.6 | --- |
| 4 | Alzheimer's disease (G30) | 23 | 4.2 | --- |
| 5 | Malignant neoplasms (C00-C97) | 22 | 4.0 | --- |
| 6 | Chronic lower respiratory diseases (J40-J47) | 18 | 3.3 | --- |
| 7 | Atherosclerosis (I70) | 12 | 2.2 | --- |
| 8 | Septicemia (A40-A41) | 10 | 1.8 | --- |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 10 | 1.8 | --- |
| 10 | Pneumonitis due to solids and liquids (J69) | 8 | 1.5 | --- |
| 11 | Diabetes mellitus (E10-E14) | 6 | 1.1 | --- |
| 11 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 6 | 1.1 | --- |
| 13 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 5 | 0.9 | --- |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 3 | 0.5 | --- |
| 14 | Cholelithiasis and other disorders of gallbladder (K80-K82) | 3 | 0.5 | --- |
| ... | All other causes (Residual) | 86 | 15.7 | --- |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|---------------------------|---|--------|-------------------------|-------|
| Hispanic, male, all ages | | | | |
| ... | All causes | 63,317 | 100.0 | 332.9 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 14,195 | 22.4 | 74.6 |
| 2 | Malignant neoplasms (C00-C97) | 11,825 | 18.7 | 62.2 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 7,157 | 11.3 | 37.6 |
| 4 | Cerebrovascular diseases (I60-I69) | 2,982 | 4.7 | 15.7 |
| 5 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 2,756 | 4.4 | 14.5 |
| 6 | Diabetes mellitus (E10-E14) | 2,590 | 4.1 | 13.6 |
| 7 | Chronic liver disease and cirrhosis (K70,K73-K74) | 2,410 | 3.8 | 12.7 |
| 8 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 1,576 | 2.5 | 8.3 |
| 9 | Chronic lower respiratory diseases (J40-J47) | 1,482 | 2.3 | 7.8 |
| 10 | Human immunodeficiency virus (HIV) disease (B20-B24) | 1,437 | 2.3 | 7.6 |
| 11 | Influenza and pneumonia (J10-J18) | 1,309 | 2.1 | 6.9 |
| 12 | Certain conditions originating in the perinatal period (P00-P96) | 1,268 | 2.0 | 6.7 |
| 13 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 892 | 1.4 | 4.7 |
| 13 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 892 | 1.4 | 4.7 |
| 15 | Septicemia (A40-A41) | 684 | 1.1 | 3.6 |
| ... | All other causes (Residual) | 9,862 | 15.6 | 51.9 |
| Hispanic, male, 1-4 years | | | | |
| ... | All causes | 518 | 100.0 | 33.8 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 195 | 37.6 | 12.7 |
| 2 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 54 | 10.4 | 3.5 |
| 3 | Malignant neoplasms (C00-C97) | 44 | 8.5 | 2.9 |
| 4 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 38 | 7.3 | 2.5 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 16 | 3.1 | * |
| 5 | Influenza and pneumonia (J10-J18) | 16 | 3.1 | * |
| 7 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 7 | 1.4 | * |
| 8 | Certain conditions originating in the perinatal period (P00-P96) | 6 | 1.2 | * |
| 9 | Septicemia (A40-A41) | 5 | 1.0 | * |
| 10 | Cerebrovascular diseases (I60-I69) | 4 | 0.8 | * |
| 10 | Chronic lower respiratory diseases (J40-J47) | 4 | 0.8 | * |
| 12 | Meningococcal infection (A39) | 3 | 0.6 | * |
| 12 | Meningitis (G00,G03) | 3 | 0.6 | * |
| 12 | Diseases of appendix (K35-K38) | 3 | 0.6 | * |
| 12 | Complications of medical and surgical care (Y40-Y84,Y88) | 3 | 0.6 | * |
| ... | All other causes (Residual) | 117 | 22.6 | 7.6 |

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|-------------------|---|--------|-------------------------|------|
| | Hispanic, male, 5-9 years | | | |
| ... | All causes | 281 | 100.0 | 14.9 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 114 | 40.6 | 6.0 |
| 2 | Malignant neoplasms (C00-C97) | 49 | 17.4 | 2.6 |
| 3 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 17 | 6.0 | * |
| 4 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 12 | 4.3 | * |
| 5 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 9 | 3.2 | * |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 9 | 3.2 | * |
| 7 | Septicemia (A40-A41) | 7 | 2.5 | * |
| 8 | Cerebrovascular diseases (I60-I69) | 5 | 1.8 | * |
| 9 | Influenza and pneumonia (J10-J18) | 3 | 1.1 | * |
| 10 | Meningococcal infection (A39) | 2 | 0.7 | * |
| 10 | Viral hepatitis (B15-B19) | 2 | 0.7 | * |
| 10 | Anemias (D50-D64) | 2 | 0.7 | * |
| 10 | Certain conditions originating in the perinatal period (P00-P96) | 2 | 0.7 | * |
| 14 | Shigellosis and amebiasis (A03,A06) | 1 | 0.4 | * |
| 14 | Human immunodeficiency virus (HIV) disease (B20-B24) | 1 | 0.4 | * |
| 14 | Meningitis (G00,G03) | 1 | 0.4 | * |
| 14 | Chronic lower respiratory diseases (J40-J47) | 1 | 0.4 | * |
| 14 | Pneumonitis due to solids and liquids (J69) | 1 | 0.4 | * |
| 14 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1 | 0.4 | * |
| 14 | Complications of medical and surgical care (Y40-Y84,Y88) | 1 | 0.4 | * |
| ... | All other causes (Residual) | 41 | 14.6 | 2.2 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|-------------------|---|--------|-------------------------|------|
| | Hispanic, male, 10-14 years | | | |
| ... | All causes | 319 | 100.0 | 18.6 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 110 | 34.5 | 6.4 |
| 2 | Malignant neoplasms (C00-C97) | 41 | 12.9 | 2.4 |
| 3 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 26 | 8.2 | 1.5 |
| 3 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 26 | 8.2 | 1.5 |
| 5 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 19 | 6.0 | * |
| 6 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 10 | 3.1 | * |
| 7 | Septicemia (A40-A41) | 7 | 2.2 | * |
| 8 | Cerebrovascular diseases (I60-I69) | 5 | 1.6 | * |
| 8 | Chronic lower respiratory diseases (J40-J47) | 5 | 1.6 | * |
| 10 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 3 | 0.9 | * |
| 10 | Anemias (D50-D64) | 3 | 0.9 | * |
| 10 | Influenza and pneumonia (J10-J18) | 3 | 0.9 | * |
| 13 | Diabetes mellitus (E10-E14) | 2 | 0.6 | * |
| 13 | Complications of medical and surgical care (Y40-Y84,Y88) | 2 | 0.6 | * |
| 15 | Human immunodeficiency virus (HIV) disease (B20-B24) | 1 | 0.3 | * |
| 15 | Nutritional deficiencies (E40-E64) | 1 | 0.3 | * |
| 15 | Meningitis (G00,G03) | 1 | 0.3 | * |
| 15 | Pneumonitis due to solids and liquids (J69) | 1 | 0.3 | * |
| 15 | Diseases of appendix (K35-K38) | 1 | 0.3 | * |
| 15 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1 | 0.3 | * |
| ... | All other causes (Residual) | 51 | 16.0 | 3.0 |

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|-----------------------------|---|--------|-------------------------|-------|
| Hispanic, male, 15-19 years | | | | |
| ... | All causes | 1,530 | 100.0 | 92.0 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 669 | 43.7 | 40.2 |
| 2 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 397 | 25.9 | 23.9 |
| 3 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 129 | 8.4 | 7.8 |
| 4 | Malignant neoplasms (C00-C97) | 95 | 6.2 | 5.7 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 27 | 1.8 | 1.6 |
| 6 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 17 | 1.1 | * |
| 7 | Cerebrovascular diseases (I60-I69) | 8 | 0.5 | * |
| 8 | Legal intervention (Y35,Y89.0) | 7 | 0.5 | * |
| 9 | Influenza and pneumonia (J10-J18) | 6 | 0.4 | * |
| 10 | Diabetes mellitus (E10-E14) | 5 | 0.3 | * |
| 11 | Tuberculosis (A16-A19) | 4 | 0.3 | * |
| 11 | Septicemia (A40-A41) | 4 | 0.3 | * |
| 11 | Chronic lower respiratory diseases (J40-J47) | 4 | 0.3 | * |
| 11 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 4 | 0.3 | * |
| 15 | Meningococcal infection (A39) | 2 | 0.1 | * |
| 15 | Human immunodeficiency virus (HIV) disease (B20-B24) | 2 | 0.1 | * |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 2 | 0.1 | * |
| 15 | Anemias (D50-D64) | 2 | 0.1 | * |
| 15 | Aortic aneurysm and dissection (I71) | 2 | 0.1 | * |
| ... | All other causes (Residual) | 144 | 9.4 | 8.7 |
| Hispanic, male, 20-24 years | | | | |
| ... | All causes | 2,489 | 100.0 | 128.2 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,104 | 44.4 | 56.9 |
| 2 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 702 | 28.2 | 36.1 |
| 3 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 214 | 8.6 | 11.0 |
| 4 | Malignant neoplasms (C00-C97) | 104 | 4.2 | 5.4 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 49 | 2.0 | 2.5 |
| 6 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 22 | 0.9 | 1.1 |
| 7 | Human immunodeficiency virus (HIV) disease (B20-B24) | 19 | 0.8 | * |
| 8 | Cerebrovascular diseases (I60-I69) | 18 | 0.7 | * |
| 9 | Legal intervention (Y35,Y89.0) | 16 | 0.6 | * |
| 10 | Influenza and pneumonia (J10-J18) | 9 | 0.4 | * |
| 11 | Diabetes mellitus (E10-E14) | 6 | 0.2 | * |
| 12 | Septicemia (A40-A41) | 5 | 0.2 | * |
| 12 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 5 | 0.2 | * |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 4 | 0.2 | * |
| 14 | Anemias (D50-D64) | 4 | 0.2 | * |
| ... | All other causes (Residual) | 208 | 8.4 | 10.7 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|-----------------------------|---|--------|-------------------------|-------|
| Hispanic, male, 25-29 years | | | | |
| ... | All causes | 2,223 | 100.0 | 115.2 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 908 | 40.8 | 47.0 |
| 2 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 488 | 22.0 | 25.3 |
| 3 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 223 | 10.0 | 11.6 |
| 4 | Malignant neoplasms (C00-C97) | 100 | 4.5 | 5.2 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 84 | 3.8 | 4.4 |
| 6 | Human immunodeficiency virus (HIV) disease (B20-B24) | 56 | 2.5 | 2.9 |
| 7 | Cerebrovascular diseases (I60-I69) | 20 | 0.9 | 1.0 |
| 8 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 17 | 0.8 | * |
| 9 | Chronic liver disease and cirrhosis (K70,K73-K74) | 15 | 0.7 | * |
| 9 | Legal intervention (Y35,Y89.0) | 15 | 0.7 | * |
| 11 | Diabetes mellitus (E10-E14) | 9 | 0.4 | * |
| 12 | Influenza and pneumonia (J10-J18) | 7 | 0.3 | * |
| 13 | Chronic lower respiratory diseases (J40-J47) | 6 | 0.3 | * |
| 14 | Tuberculosis (A16-A19) | 5 | 0.2 | * |
| 14 | Septicemia (A40-A41) | 5 | 0.2 | * |
| ... | All other causes (Residual) | 265 | 11.9 | 13.7 |
| Hispanic, male, 30-34 years | | | | |
| ... | All causes | 2,143 | 100.0 | 121.1 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 711 | 33.2 | 40.2 |
| 2 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 312 | 14.6 | 17.6 |
| 3 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 196 | 9.1 | 11.1 |
| 4 | Human immunodeficiency virus (HIV) disease (B20-B24) | 166 | 7.7 | 9.4 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 142 | 6.6 | 8.0 |
| 6 | Malignant neoplasms (C00-C97) | 133 | 6.2 | 7.5 |
| 7 | Chronic liver disease and cirrhosis (K70,K73-K74) | 52 | 2.4 | 2.9 |
| 8 | Cerebrovascular diseases (I60-I69) | 40 | 1.9 | 2.3 |
| 9 | Diabetes mellitus (E10-E14) | 19 | 0.9 | * |
| 10 | Septicemia (A40-A41) | 10 | 0.5 | * |
| 10 | Aortic aneurysm and dissection (I71) | 10 | 0.5 | * |
| 12 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 9 | 0.4 | * |
| 12 | Legal intervention (Y35,Y89.0) | 9 | 0.4 | * |
| 14 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 7 | 0.3 | * |
| 14 | Influenza and pneumonia (J10-J18) | 7 | 0.3 | * |
| 14 | Chronic lower respiratory diseases (J40-J47) | 7 | 0.3 | * |
| ... | All other causes (Residual) | 313 | 14.6 | 17.7 |

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|-----------------------------|---|--------|-------------------------|-------|
| Hispanic, male, 35-39 years | | | | |
| ... | All causes | 2,739 | 100.0 | 175.5 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 763 | 27.9 | 48.9 |
| 2 | Human immunodeficiency virus (HIV) disease (B20-B24) | 273 | 10.0 | 17.5 |
| 3 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 258 | 9.4 | 16.5 |
| 4 | Malignant neoplasms (C00-C97) | 236 | 8.6 | 15.1 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 228 | 8.3 | 14.6 |
| 6 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 200 | 7.3 | 12.8 |
| 7 | Chronic liver disease and cirrhosis (K70,K73-K74) | 150 | 5.5 | 9.6 |
| 8 | Cerebrovascular diseases (I60-I69) | 58 | 2.1 | 3.7 |
| 9 | Diabetes mellitus (E10-E14) | 40 | 1.5 | 2.6 |
| 10 | Viral hepatitis (B15-B19) | 23 | 0.8 | 1.5 |
| 11 | Septicemia (A40-A41) | 20 | 0.7 | 1.3 |
| 11 | Influenza and pneumonia (J10-J18) | 20 | 0.7 | 1.3 |
| 13 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 14 | 0.5 | * |
| 14 | Chronic lower respiratory diseases (J40-J47) | 12 | 0.4 | * |
| 15 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 11 | 0.4 | * |
| ... | All other causes (Residual) | 433 | 15.8 | 27.8 |
| Hispanic, male, 40-44 years | | | | |
| ... | All causes | 3,156 | 100.0 | 249.0 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 655 | 20.8 | 51.7 |
| 2 | Malignant neoplasms (C00-C97) | 379 | 12.0 | 29.9 |
| 3 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 337 | 10.7 | 26.6 |
| 4 | Human immunodeficiency virus (HIV) disease (B20-B24) | 316 | 10.0 | 24.9 |
| 5 | Chronic liver disease and cirrhosis (K70,K73-K74) | 275 | 8.7 | 21.7 |
| 6 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 193 | 6.1 | 15.2 |
| 7 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 150 | 4.8 | 11.8 |
| 8 | Cerebrovascular diseases (I60-I69) | 93 | 2.9 | 7.3 |
| 9 | Diabetes mellitus (E10-E14) | 69 | 2.2 | 5.4 |
| 10 | Viral hepatitis (B15-B19) | 66 | 2.1 | 5.2 |
| 11 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 32 | 1.0 | 2.5 |
| 12 | Influenza and pneumonia (J10-J18) | 28 | 0.9 | 2.2 |
| 13 | Septicemia (A40-A41) | 23 | 0.7 | 1.8 |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 18 | 0.6 | * |
| 15 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 16 | 0.5 | * |
| ... | All other causes (Residual) | 506 | 16.0 | 39.9 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|-----------------------------|---|--------|-------------------------|-------|
| Hispanic, male, 45-49 years | | | | |
| ... | All causes | 3,655 | 100.0 | 379.7 |
| 1 | Malignant neoplasms (C00-C97) | 579 | 15.8 | 60.1 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 565 | 15.5 | 58.7 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 481 | 13.2 | 50.0 |
| 4 | Chronic liver disease and cirrhosis (K70,K73-K74) | 403 | 11.0 | 41.9 |
| 5 | Human immunodeficiency virus (HIV) disease (B20-B24) | 241 | 6.6 | 25.0 |
| 6 | Cerebrovascular diseases (I60-I69) | 141 | 3.9 | 14.6 |
| 7 | Diabetes mellitus (E10-E14) | 139 | 3.8 | 14.4 |
| 8 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 121 | 3.3 | 12.6 |
| 9 | Viral hepatitis (B15-B19) | 119 | 3.3 | 12.4 |
| 10 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 106 | 2.9 | 11.0 |
| 11 | Influenza and pneumonia (J10-J18) | 47 | 1.3 | 4.9 |
| 12 | Septicemia (A40-A41) | 37 | 1.0 | 3.8 |
| 13 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 27 | 0.7 | 2.8 |
| 14 | Chronic lower respiratory diseases (J40-J47) | 22 | 0.6 | 2.3 |
| 15 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 21 | 0.6 | 2.2 |
| ... | All other causes (Residual) | 606 | 16.6 | 63.0 |
| Hispanic, male, 50-54 years | | | | |
| ... | All causes | 3,822 | 100.0 | 529.6 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 832 | 21.8 | 115.3 |
| 2 | Malignant neoplasms (C00-C97) | 793 | 20.7 | 109.9 |
| 3 | Chronic liver disease and cirrhosis (K70,K73-K74) | 379 | 9.9 | 52.5 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 341 | 8.9 | 47.3 |
| 5 | Human immunodeficiency virus (HIV) disease (B20-B24) | 179 | 4.7 | 24.8 |
| 6 | Cerebrovascular diseases (I60-I69) | 174 | 4.6 | 24.1 |
| 7 | Diabetes mellitus (E10-E14) | 172 | 4.5 | 23.8 |
| 8 | Viral hepatitis (B15-B19) | 87 | 2.3 | 12.1 |
| 9 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 86 | 2.3 | 11.9 |
| 10 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 69 | 1.8 | 9.6 |
| 11 | Septicemia (A40-A41) | 43 | 1.1 | 6.0 |
| 12 | Influenza and pneumonia (J10-J18) | 38 | 1.0 | 5.3 |
| 13 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 37 | 1.0 | 5.1 |
| 14 | Chronic lower respiratory diseases (J40-J47) | 32 | 0.8 | 4.4 |
| 15 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 18 | 0.5 | * |
| ... | All other causes (Residual) | 542 | 14.2 | 75.1 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|-----------------------------|---|--------|-------------------------|---------|
| Hispanic, male, 55-59 years | | | | |
| ... | All causes | 3,760 | 100.0 | 751.2 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 956 | 25.4 | 191.0 |
| 2 | Malignant neoplasms (C00-C97) | 948 | 25.2 | 189.4 |
| 3 | Chronic liver disease and cirrhosis (K70,K73-K74) | 295 | 7.8 | 58.9 |
| 4 | Diabetes mellitus (E10-E14) | 217 | 5.8 | 43.4 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 213 | 5.7 | 42.6 |
| 6 | Cerebrovascular diseases (I60-I69) | 171 | 4.5 | 34.2 |
| 7 | Human immunodeficiency virus (HIV) disease (B20-B24) | 88 | 2.3 | 17.6 |
| 8 | Viral hepatitis (B15-B19) | 85 | 2.3 | 17.0 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 63 | 1.7 | 12.6 |
| 10 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 56 | 1.5 | 11.2 |
| 11 | Chronic lower respiratory diseases (J40-J47) | 53 | 1.4 | 10.6 |
| 12 | Influenza and pneumonia (J10-J18) | 50 | 1.3 | 10.0 |
| 13 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 33 | 0.9 | 6.6 |
| 14 | Septicemia (A40-A41) | 32 | 0.9 | 6.4 |
| 15 | Pneumonitis due to solids and liquids (J69) | 14 | 0.4 | * |
| ... | All other causes (Residual) | 486 | 12.9 | 97.1 |
| Hispanic, male, 60-64 years | | | | |
| ... | All causes | 4,266 | 100.0 | 1,158.6 |
| 1 | Malignant neoplasms (C00-C97) | 1,254 | 29.4 | 340.6 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 1,148 | 26.9 | 311.8 |
| 3 | Diabetes mellitus (E10-E14) | 293 | 6.9 | 79.6 |
| 4 | Chronic liver disease and cirrhosis (K70,K73-K74) | 260 | 6.1 | 70.6 |
| 5 | Cerebrovascular diseases (I60-I69) | 207 | 4.9 | 56.2 |
| 6 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 178 | 4.2 | 48.3 |
| 7 | Chronic lower respiratory diseases (J40-J47) | 87 | 2.0 | 23.6 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 69 | 1.6 | 18.7 |
| 9 | Septicemia (A40-A41) | 54 | 1.3 | 14.7 |
| 10 | Influenza and pneumonia (J10-J18) | 49 | 1.1 | 13.3 |
| 11 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 44 | 1.0 | 11.9 |
| 12 | Human immunodeficiency virus (HIV) disease (B20-B24) | 41 | 1.0 | 11.1 |
| 13 | Viral hepatitis (B15-B19) | 38 | 0.9 | 10.3 |
| 14 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 26 | 0.6 | 7.1 |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 25 | 0.6 | 6.8 |
| ... | All other causes (Residual) | 493 | 11.6 | 133.9 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|-----------------------------|---|--------|-------------------------|---------|
| Hispanic, male, 65-69 years | | | | |
| ... | All causes | 5,129 | 100.0 | 1,817.5 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 1,544 | 30.1 | 547.1 |
| 2 | Malignant neoplasms (C00-C97) | 1,465 | 28.6 | 519.1 |
| 3 | Diabetes mellitus (E10-E14) | 374 | 7.3 | 132.5 |
| 4 | Cerebrovascular diseases (I60-I69) | 269 | 5.2 | 95.3 |
| 5 | Chronic liver disease and cirrhosis (K70,K73-K74) | 220 | 4.3 | 78.0 |
| 6 | Chronic lower respiratory diseases (J40-J47) | 161 | 3.1 | 57.1 |
| 7 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 130 | 2.5 | 46.1 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 100 | 1.9 | 35.4 |
| 9 | Influenza and pneumonia (J10-J18) | 71 | 1.4 | 25.2 |
| 10 | Septicemia (A40-A41) | 69 | 1.3 | 24.5 |
| 11 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 39 | 0.8 | 13.8 |
| 12 | Aortic aneurysm and dissection (I71) | 35 | 0.7 | 12.4 |
| 13 | Human immunodeficiency virus (HIV) disease (B20-B24) | 32 | 0.6 | 11.3 |
| 14 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 30 | 0.6 | 10.6 |
| 15 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 24 | 0.5 | 8.5 |
| ... | All other causes (Residual) | 566 | 11.0 | 200.6 |
| Hispanic, male, 70-74 years | | | | |
| ... | All causes | 6,102 | 100.0 | 2,791.5 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 1,880 | 30.8 | 860.1 |
| 2 | Malignant neoplasms (C00-C97) | 1,754 | 28.7 | 802.4 |
| 3 | Diabetes mellitus (E10-E14) | 384 | 6.3 | 175.7 |
| 4 | Cerebrovascular diseases (I60-I69) | 368 | 6.0 | 168.4 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 217 | 3.6 | 99.3 |
| 6 | Chronic liver disease and cirrhosis (K70,K73-K74) | 166 | 2.7 | 75.9 |
| 7 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 145 | 2.4 | 66.3 |
| 8 | Influenza and pneumonia (J10-J18) | 133 | 2.2 | 60.8 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 117 | 1.9 | 53.5 |
| 10 | Septicemia (A40-A41) | 80 | 1.3 | 36.6 |
| 11 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 54 | 0.9 | 24.7 |
| 12 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 38 | 0.6 | 17.4 |
| 13 | Parkinson's disease (G20-G21) | 37 | 0.6 | 16.9 |
| 14 | Aortic aneurysm and dissection (I71) | 36 | 0.6 | 16.5 |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 29 | 0.5 | 13.3 |
| ... | All other causes (Residual) | 664 | 10.9 | 303.8 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|-----------------------------|---|--------|-------------------------|---------|
| Hispanic, male, 75-79 years | | | | |
| ... | All causes | 6,535 | 100.0 | 4,456.8 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 2,031 | 31.1 | 1,385.1 |
| 2 | Malignant neoplasms (C00-C97) | 1,718 | 26.3 | 1,171.7 |
| 3 | Cerebrovascular diseases (I60-I69) | 456 | 7.0 | 311.0 |
| 4 | Diabetes mellitus (E10-E14) | 385 | 5.9 | 262.6 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 286 | 4.4 | 195.1 |
| 6 | Influenza and pneumonia (J10-J18) | 180 | 2.8 | 122.8 |
| 7 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 150 | 2.3 | 102.3 |
| 8 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 136 | 2.1 | 92.8 |
| 9 | Chronic liver disease and cirrhosis (K70,K73-K74) | 120 | 1.8 | 81.8 |
| 10 | Septicemia (A40-A41) | 101 | 1.5 | 68.9 |
| 11 | Alzheimer's disease (G30) | 89 | 1.4 | 60.7 |
| 12 | Parkinson's disease (G20-G21) | 63 | 1.0 | 43.0 |
| 13 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 58 | 0.9 | 39.6 |
| 14 | Aortic aneurysm and dissection (I71) | 52 | 0.8 | 35.5 |
| 15 | Pneumonitis due to solids and liquids (J69) | 37 | 0.6 | 25.2 |
| ... | All other causes (Residual) | 673 | 10.3 | 459.0 |
| Hispanic, male, 80-84 years | | | | |
| ... | All causes | 5,224 | 100.0 | 6,783.5 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 1,777 | 34.0 | 2,307.5 |
| 2 | Malignant neoplasms (C00-C97) | 1,096 | 21.0 | 1,423.2 |
| 3 | Cerebrovascular diseases (I60-I69) | 408 | 7.8 | 529.8 |
| 4 | Diabetes mellitus (E10-E14) | 253 | 4.8 | 328.5 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 215 | 4.1 | 279.2 |
| 6 | Influenza and pneumonia (J10-J18) | 208 | 4.0 | 270.1 |
| 7 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 101 | 1.9 | 131.2 |
| 8 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 99 | 1.9 | 128.6 |
| 9 | Alzheimer's disease (G30) | 97 | 1.9 | 126.0 |
| 10 | Parkinson's disease (G20-G21) | 78 | 1.5 | 101.3 |
| 11 | Septicemia (A40-A41) | 69 | 1.3 | 89.6 |
| 12 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 56 | 1.1 | 72.7 |
| 13 | Chronic liver disease and cirrhosis (K70,K73-K74) | 50 | 1.0 | 64.9 |
| 14 | Pneumonitis due to solids and liquids (J69) | 44 | 0.8 | 57.1 |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 38 | 0.7 | 49.3 |
| ... | All other causes (Residual) | 635 | 12.2 | 824.6 |

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|-----------------------------|---|--------|-------------------------|------|
| Hispanic, male, 85-89 years | | | | |
| ... | All causes | 3,821 | 100.0 | --- |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 1,385 | 36.2 | --- |
| 2 | Malignant neoplasms (C00-C97) | 642 | 16.8 | --- |
| 3 | Cerebrovascular diseases (I60-I69) | 300 | 7.9 | --- |
| 4 | Chronic lower respiratory diseases (J40-J47) | 203 | 5.3 | --- |
| 5 | Influenza and pneumonia (J10-J18) | 177 | 4.6 | --- |
| 6 | Diabetes mellitus (E10-E14) | 146 | 3.8 | --- |
| 7 | Alzheimer's disease (G30) | 88 | 2.3 | --- |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 82 | 2.1 | --- |
| 9 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 59 | 1.5 | --- |
| 10 | Pneumonitis due to solids and liquids (J69) | 56 | 1.5 | --- |
| 11 | Parkinson's disease (G20-G21) | 50 | 1.3 | --- |
| 12 | Septicemia (A40-A41) | 49 | 1.3 | --- |
| 13 | Atherosclerosis (I70) | 39 | 1.0 | --- |
| 14 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 37 | 1.0 | --- |
| 15 | Aortic aneurysm and dissection (I71) | 23 | 0.6 | --- |
| ... | All other causes (Residual) | 485 | 12.7 | --- |
| Hispanic, male, 90-94 years | | | | |
| ... | All causes | 2,051 | 100.0 | --- |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 808 | 39.4 | --- |
| 2 | Malignant neoplasms (C00-C97) | 271 | 13.2 | --- |
| 3 | Cerebrovascular diseases (I60-I69) | 161 | 7.8 | --- |
| 4 | Influenza and pneumonia (J10-J18) | 145 | 7.1 | --- |
| 5 | Chronic lower respiratory diseases (J40-J47) | 112 | 5.5 | --- |
| 6 | Diabetes mellitus (E10-E14) | 56 | 2.7 | --- |
| 7 | Alzheimer's disease (G30) | 50 | 2.4 | --- |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 48 | 2.3 | --- |
| 9 | Pneumonitis due to solids and liquids (J69) | 41 | 2.0 | --- |
| 10 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 35 | 1.7 | --- |
| 11 | Septicemia (A40-A41) | 25 | 1.2 | --- |
| 12 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 24 | 1.2 | --- |
| 13 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 15 | 0.7 | --- |
| 14 | Parkinson's disease (G20-G21) | 14 | 0.7 | --- |
| 15 | Atherosclerosis (I70) | 13 | 0.6 | --- |
| ... | All other causes (Residual) | 233 | 11.4 | --- |

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|------------------------------------|---|--------|-------------------------|------|
| Hispanic, male, 95-99 years | | | | |
| ... | All causes | 754 | 100.0 | --- |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 265 | 35.1 | --- |
| 2 | Malignant neoplasms (C00-C97) | 109 | 14.5 | --- |
| 3 | Influenza and pneumonia (J10-J18) | 69 | 9.2 | --- |
| 4 | Cerebrovascular diseases (I60-I69) | 55 | 7.3 | --- |
| 5 | Chronic lower respiratory diseases (J40-J47) | 34 | 4.5 | --- |
| 6 | Alzheimer's disease (G30) | 20 | 2.7 | --- |
| 7 | Diabetes mellitus (E10-E14) | 19 | 2.5 | --- |
| 8 | Pneumonitis due to solids and liquids (J69) | 16 | 2.1 | --- |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 15 | 2.0 | --- |
| 10 | Atherosclerosis (I70) | 11 | 1.5 | --- |
| 11 | Septicemia (A40-A41) | 9 | 1.2 | --- |
| 12 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 8 | 1.1 | --- |
| 13 | Nutritional deficiencies (E40-E64) | 7 | 0.9 | --- |
| 14 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 6 | 0.8 | --- |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 3 | 0.4 | --- |
| 15 | Parkinson's disease (G20-G21) | 3 | 0.4 | --- |
| ... | All other causes (Residual) | 105 | 13.9 | --- |
| Hispanic, male, 100 years and over | | | | |
| ... | All causes | 141 | 100.0 | --- |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 54 | 38.3 | --- |
| 2 | Cerebrovascular diseases (I60-I69) | 12 | 8.5 | --- |
| 2 | Malignant neoplasms (C00-C97) | 11 | 7.8 | --- |
| 3 | Influenza and pneumonia (J10-J18) | 11 | 7.8 | --- |
| 5 | Chronic lower respiratory diseases (J40-J47) | 6 | 4.3 | --- |
| 6 | Atherosclerosis (I70) | 4 | 2.8 | --- |
| 7 | Alzheimer's disease (G30) | 3 | 2.1 | --- |
| 7 | Pneumonitis due to solids and liquids (J69) | 3 | 2.1 | --- |
| 7 | Cholelithiasis and other disorders of gallbladder (K80-K82) | 3 | 2.1 | --- |
| 7 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 3 | 2.1 | --- |
| 11 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 2 | 1.4 | --- |
| 12 | Septicemia (A40-A41) | 1 | 0.7 | --- |
| 12 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 1 | 0.7 | --- |
| 12 | Anemias (D50-D64) | 1 | 0.7 | --- |
| 12 | Diabetes mellitus (E10-E14) | 1 | 0.7 | --- |
| 12 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 1 | 0.7 | --- |
| 12 | Pneumoconioses and chemical effects (J60-J66,J68) | 1 | 0.7 | --- |
| ... | All other causes (Residual) | 23 | 16.3 | --- |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|-----------------------------|---|--------|-------------------------|-------|
| Hispanic, female, all ages | | | | |
| ... | All causes | 50,096 | 100.0 | 279.0 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 12,895 | 25.7 | 71.8 |
| 2 | Malignant neoplasms (C00-C97) | 10,546 | 21.1 | 58.7 |
| 3 | Cerebrovascular diseases (I60-I69) | 3,434 | 6.9 | 19.1 |
| 4 | Diabetes mellitus (E10-E14) | 3,073 | 6.1 | 17.1 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 2,366 | 4.7 | 13.2 |
| 6 | Influenza and pneumonia (J10-J18) | 1,413 | 2.8 | 7.9 |
| 7 | Chronic lower respiratory diseases (J40-J47) | 1,350 | 2.7 | 7.5 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 965 | 1.9 | 5.4 |
| 9 | Certain conditions originating in the perinatal period (P00-P96) | 959 | 1.9 | 5.3 |
| 10 | Chronic liver disease and cirrhosis (K70,K73-K74) | 891 | 1.8 | 5.0 |
| 11 | Alzheimer's disease (G30) | 838 | 1.7 | 4.7 |
| 12 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 740 | 1.5 | 4.1 |
| 13 | Septicemia (A40-A41) | 702 | 1.4 | 3.9 |
| 14 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 575 | 1.1 | 3.2 |
| 15 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 524 | 1.0 | 2.9 |
| ... | All other causes (Residual) | 8,825 | 17.6 | 49.2 |
| Hispanic, female, 1-4 years | | | | |
| ... | All causes | 399 | 100.0 | 27.2 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 115 | 28.8 | 7.8 |
| 2 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 49 | 12.3 | 3.3 |
| 3 | Malignant neoplasms (C00-C97) | 40 | 10.0 | 2.7 |
| 4 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 31 | 7.8 | 2.1 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 17 | 4.3 | * |
| 6 | Influenza and pneumonia (J10-J18) | 16 | 4.0 | * |
| 7 | Septicemia (A40-A41) | 13 | 3.3 | * |
| 8 | Certain conditions originating in the perinatal period (P00-P96) | 8 | 2.0 | * |
| 9 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 6 | 1.5 | * |
| 9 | Cerebrovascular diseases (I60-I69) | 6 | 1.5 | * |
| 11 | Chronic lower respiratory diseases (J40-J47) | 4 | 1.0 | * |
| 12 | Complications of medical and surgical care (Y40-Y84,Y88) | 3 | 0.8 | * |
| 13 | Meningococcal infection (A39) | 2 | 0.5 | * |
| 13 | Pneumonitis due to solids and liquids (J69) | 2 | 0.5 | * |
| 13 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 2 | 0.5 | * |
| ... | All other causes (Residual) | 85 | 21.3 | 5.8 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|-------------------------------|---|--------|-------------------------|------|
| Hispanic, female, 5-9 years | | | | |
| ... | All causes | 199 | 100.0 | 11.0 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 72 | 36.2 | 4.0 |
| 2 | Malignant neoplasms (C00-C97) | 38 | 19.1 | 2.1 |
| 3 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 11 | 5.5 | * |
| 4 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 8 | 4.0 | * |
| 4 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 8 | 4.0 | * |
| 6 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 5 | 2.5 | * |
| 7 | Cerebrovascular diseases (I60-I69) | 2 | 1.0 | * |
| 7 | Influenza and pneumonia (J10-J18) | 2 | 1.0 | * |
| 9 | Septicemia (A40-A41) | 1 | 0.5 | * |
| 9 | Meningitis (G00,G03) | 1 | 0.5 | * |
| 9 | Chronic lower respiratory diseases (J40-J47) | 1 | 0.5 | * |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1 | 0.5 | * |
| 9 | Certain conditions originating in the perinatal period (P00-P96) | 1 | 0.5 | * |
| ... | All other causes (Residual) | 48 | 24.1 | 2.7 |
| Hispanic, female, 10-14 years | | | | |
| ... | All causes | 239 | 100.0 | 14.5 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 83 | 34.7 | 5.0 |
| 2 | Malignant neoplasms (C00-C97) | 49 | 20.5 | 3.0 |
| 3 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 11 | 4.6 | * |
| 3 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 11 | 4.6 | * |
| 3 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 11 | 4.6 | * |
| 6 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 10 | 4.2 | * |
| 7 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 4 | 1.7 | * |
| 7 | Cerebrovascular diseases (I60-I69) | 4 | 1.7 | * |
| 9 | Septicemia (A40-A41) | 3 | 1.3 | * |
| 9 | Pneumonitis due to solids and liquids (J69) | 3 | 1.3 | * |
| 11 | Influenza and pneumonia (J10-J18) | 2 | 0.8 | * |
| 11 | Chronic lower respiratory diseases (J40-J47) | 2 | 0.8 | * |
| 13 | Human immunodeficiency virus (HIV) disease (B20-B24) | 1 | 0.4 | * |
| 13 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1 | 0.4 | * |
| 13 | Pregnancy, childbirth and the puerperium (O00-099) | 1 | 0.4 | * |
| 13 | Complications of medical and surgical care (Y40-Y84,Y88) | 1 | 0.4 | * |
| ... | All other causes (Residual) | 42 | 17.6 | 2.6 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|-------------------------------|---|--------|-------------------------|------|
| Hispanic, female, 15-19 years | | | | |
| ... | All causes | 468 | 100.0 | 31.1 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 210 | 44.9 | 14.0 |
| 2 | Malignant neoplasms (C00-C97) | 48 | 10.3 | 3.2 |
| 3 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 39 | 8.3 | 2.6 |
| 4 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 37 | 7.9 | 2.5 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 18 | 3.8 | * |
| 6 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 15 | 3.2 | * |
| 7 | Pregnancy, childbirth and the puerperium (O00-099) | 9 | 1.9 | * |
| 8 | Cerebrovascular diseases (I60-I69) | 5 | 1.1 | * |
| 9 | Anemias (D50-D64) | 4 | 0.9 | * |
| 10 | Diabetes mellitus (E10-E14) | 3 | 0.6 | * |
| 10 | Chronic lower respiratory diseases (J40-J47) | 3 | 0.6 | * |
| 10 | Complications of medical and surgical care (Y40-Y84,Y88) | 3 | 0.6 | * |
| 13 | Viral hepatitis (B15-B19) | 2 | 0.4 | * |
| 13 | Human immunodeficiency virus (HIV) disease (B20-B24) | 2 | 0.4 | * |
| 13 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 2 | 0.4 | * |
| 13 | Pneumonitis due to solids and liquids (J69) | 2 | 0.4 | * |
| ... | All other causes (Residual) | 66 | 14.1 | 4.4 |
| Hispanic, female, 20-24 years | | | | |
| ... | All causes | 571 | 100.0 | 36.1 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 199 | 34.9 | 12.6 |
| 2 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 84 | 14.7 | 5.3 |
| 3 | Malignant neoplasms (C00-C97) | 42 | 7.4 | 2.7 |
| 4 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 34 | 6.0 | 2.2 |
| 5 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 19 | 3.3 | * |
| 6 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 18 | 3.2 | * |
| 7 | Pregnancy, childbirth and the puerperium (O00-099) | 17 | 3.0 | * |
| 8 | Human immunodeficiency virus (HIV) disease (B20-B24) | 9 | 1.6 | * |
| 8 | Cerebrovascular diseases (I60-I69) | 9 | 1.6 | * |
| 10 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 7 | 1.2 | * |
| 11 | Diabetes mellitus (E10-E14) | 5 | 0.9 | * |
| 12 | Septicemia (A40-A41) | 3 | 0.5 | * |
| 13 | Viral hepatitis (B15-B19) | 2 | 0.4 | * |
| 13 | Meningitis (G00,G03) | 2 | 0.4 | * |
| 13 | Influenza and pneumonia (J10-J18) | 2 | 0.4 | * |
| 13 | Chronic liver disease and cirrhosis (K70,K73-K74) | 2 | 0.4 | * |
| 13 | Certain conditions originating in the perinatal period (P00-P96) | 2 | 0.4 | * |
| 13 | Complications of medical and surgical care (Y40-Y84,Y88) | 2 | 0.4 | * |
| ... | All other causes (Residual) | 113 | 19.8 | 7.1 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|-------------------------------|---|--------|-------------------------|------|
| Hispanic, female, 25-29 years | | | | |
| ... | All causes | 640 | 100.0 | 39.4 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 165 | 25.8 | 10.2 |
| 2 | Malignant neoplasms (C00-C97) | 104 | 16.3 | 6.4 |
| 3 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 85 | 13.3 | 5.2 |
| 4 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 36 | 5.6 | 2.2 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 31 | 4.8 | 1.9 |
| 6 | Human immunodeficiency virus (HIV) disease (B20-B24) | 24 | 3.8 | 1.5 |
| 7 | Pregnancy, childbirth and the puerperium (O00-099) | 21 | 3.3 | 1.3 |
| 8 | Cerebrovascular diseases (I60-I69) | 14 | 2.2 | * |
| 9 | Diabetes mellitus (E10-E14) | 11 | 1.7 | * |
| 10 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 9 | 1.4 | * |
| 11 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 6 | 0.9 | * |
| 12 | Septicemia (A40-A41) | 4 | 0.6 | * |
| 12 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 4 | 0.6 | * |
| 12 | Chronic lower respiratory diseases (J40-J47) | 4 | 0.6 | * |
| 15 | Influenza and pneumonia (J10-J18) | 3 | 0.5 | * |
| 15 | Certain conditions originating in the perinatal period (P00-P96) | 3 | 0.5 | * |
| 15 | Complications of medical and surgical care (Y40-Y84,Y88) | 3 | 0.5 | * |
| ... | All other causes (Residual) | 113 | 17.7 | 7.0 |
| Hispanic, female, 30-34 years | | | | |
| ... | All causes | 791 | 100.0 | 51.3 |
| 1 | Malignant neoplasms (C00-C97) | 160 | 20.2 | 10.4 |
| 2 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 154 | 19.5 | 10.0 |
| 3 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 75 | 9.5 | 4.9 |
| 4 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 57 | 7.2 | 3.7 |
| 5 | Human immunodeficiency virus (HIV) disease (B20-B24) | 54 | 6.8 | 3.5 |
| 6 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 35 | 4.4 | 2.3 |
| 7 | Cerebrovascular diseases (I60-I69) | 25 | 3.2 | 1.6 |
| 8 | Pregnancy, childbirth and the puerperium (O00-099) | 17 | 2.1 | * |
| 9 | Chronic liver disease and cirrhosis (K70,K73-K74) | 13 | 1.6 | * |
| 10 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 10 | 1.3 | * |
| 10 | Diabetes mellitus (E10-E14) | 10 | 1.3 | * |
| 10 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 10 | 1.3 | * |
| 13 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 9 | 1.1 | * |
| 14 | Viral hepatitis (B15-B19) | 7 | 0.9 | * |
| 15 | Influenza and pneumonia (J10-J18) | 6 | 0.8 | * |
| ... | All other causes (Residual) | 149 | 18.8 | 9.7 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|-------------------------------|---|--------|-------------------------|-------|
| Hispanic, female, 35-39 years | | | | |
| ... | All causes | 1,120 | 100.0 | 79.0 |
| 1 | Malignant neoplasms (C00-C97) | 309 | 27.6 | 21.8 |
| 2 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 179 | 16.0 | 12.6 |
| 3 | Human immunodeficiency virus (HIV) disease (B20-B24) | 82 | 7.3 | 5.8 |
| 4 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 74 | 6.6 | 5.2 |
| 5 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 71 | 6.3 | 5.0 |
| 6 | Cerebrovascular diseases (I60-I69) | 44 | 3.9 | 3.1 |
| 7 | Chronic liver disease and cirrhosis (K70,K73-K74) | 37 | 3.3 | 2.6 |
| 8 | Diabetes mellitus (E10-E14) | 29 | 2.6 | 2.0 |
| 9 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 24 | 2.1 | 1.7 |
| 10 | Chronic lower respiratory diseases (J40-J47) | 13 | 1.2 | * |
| 11 | Viral hepatitis (B15-B19) | 12 | 1.1 | * |
| 11 | Influenza and pneumonia (J10-J18) | 12 | 1.1 | * |
| 13 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 10 | 0.9 | * |
| 13 | Pregnancy, childbirth and the puerperium (O00-099) | 10 | 0.9 | * |
| 15 | Septicemia (A40-A41) | 8 | 0.7 | * |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 8 | 0.7 | * |
| 15 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 8 | 0.7 | * |
| ... | All other causes (Residual) | 190 | 17.0 | 13.4 |
| Hispanic, female, 40-44 years | | | | |
| ... | All causes | 1,426 | 100.0 | 118.3 |
| 1 | Malignant neoplasms (C00-C97) | 472 | 33.1 | 39.2 |
| 2 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 163 | 11.4 | 13.5 |
| 3 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 111 | 7.8 | 9.2 |
| 4 | Human immunodeficiency virus (HIV) disease (B20-B24) | 88 | 6.2 | 7.3 |
| 5 | Cerebrovascular diseases (I60-I69) | 80 | 5.6 | 6.6 |
| 6 | Chronic liver disease and cirrhosis (K70,K73-K74) | 68 | 4.8 | 5.6 |
| 7 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 45 | 3.2 | 3.7 |
| 8 | Diabetes mellitus (E10-E14) | 29 | 2.0 | 2.4 |
| 9 | Viral hepatitis (B15-B19) | 25 | 1.8 | 2.1 |
| 10 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 23 | 1.6 | 1.9 |
| 11 | Septicemia (A40-A41) | 21 | 1.5 | 1.7 |
| 12 | Chronic lower respiratory diseases (J40-J47) | 18 | 1.3 | * |
| 12 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 18 | 1.3 | * |
| 14 | Influenza and pneumonia (J10-J18) | 13 | 0.9 | * |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 7 | 0.5 | * |
| 15 | Pregnancy, childbirth and the puerperium (O00-099) | 7 | 0.5 | * |
| 15 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 7 | 0.5 | * |
| ... | All other causes (Residual) | 231 | 16.2 | 19.2 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|-------------------------------|---|--------|-------------------------|-------|
| Hispanic, female, 45-49 years | | | | |
| ... | All causes | 1,742 | 100.0 | 181.7 |
| 1 | Malignant neoplasms (C00-C97) | 607 | 34.8 | 63.3 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 171 | 9.8 | 17.8 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 146 | 8.4 | 15.2 |
| 4 | Cerebrovascular diseases (I60-I69) | 98 | 5.6 | 10.2 |
| 5 | Diabetes mellitus (E10-E14) | 90 | 5.2 | 9.4 |
| 6 | Chronic liver disease and cirrhosis (K70,K73-K74) | 77 | 4.4 | 8.0 |
| 7 | Human immunodeficiency virus (HIV) disease (B20-B24) | 75 | 4.3 | 7.8 |
| 8 | Viral hepatitis (B15-B19) | 36 | 2.1 | 3.8 |
| 9 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 35 | 2.0 | 3.7 |
| 10 | Septicemia (A40-A41) | 33 | 1.9 | 3.4 |
| 11 | Chronic lower respiratory diseases (J40-J47) | 25 | 1.4 | 2.6 |
| 12 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 24 | 1.4 | 2.5 |
| 12 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 24 | 1.4 | 2.5 |
| 14 | Influenza and pneumonia (J10-J18) | 9 | 0.5 | * |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 8 | 0.5 | * |
| ... | All other causes (Residual) | 284 | 16.3 | 29.6 |
| Hispanic, female, 50-54 years | | | | |
| ... | All causes | 2,143 | 100.0 | 283.8 |
| 1 | Malignant neoplasms (C00-C97) | 836 | 39.0 | 110.7 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 307 | 14.3 | 40.7 |
| 3 | Cerebrovascular diseases (I60-I69) | 126 | 5.9 | 16.7 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 110 | 5.1 | 14.6 |
| 5 | Diabetes mellitus (E10-E14) | 106 | 4.9 | 14.0 |
| 6 | Chronic liver disease and cirrhosis (K70,K73-K74) | 92 | 4.3 | 12.2 |
| 7 | Chronic lower respiratory diseases (J40-J47) | 45 | 2.1 | 6.0 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 44 | 2.1 | 5.8 |
| 9 | Human immunodeficiency virus (HIV) disease (B20-B24) | 31 | 1.4 | 4.1 |
| 10 | Influenza and pneumonia (J10-J18) | 27 | 1.3 | 3.6 |
| 11 | Viral hepatitis (B15-B19) | 24 | 1.1 | 3.2 |
| 12 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 23 | 1.1 | 3.0 |
| 12 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 23 | 1.1 | 3.0 |
| 14 | Septicemia (A40-A41) | 20 | 0.9 | 2.6 |
| 15 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 14 | 0.7 | * |
| ... | All other causes (Residual) | 315 | 14.7 | 41.7 |

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|-------------------------------|---|--------|-------------------------|-------|
| Hispanic, female, 55-59 years | | | | |
| ... | All causes | 2,354 | 100.0 | 426.0 |
| 1 | Malignant neoplasms (C00-C97) | 866 | 36.8 | 156.7 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 406 | 17.2 | 73.5 |
| 3 | Diabetes mellitus (E10-E14) | 187 | 7.9 | 33.8 |
| 4 | Cerebrovascular diseases (I60-I69) | 130 | 5.5 | 23.5 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 100 | 4.2 | 18.1 |
| 6 | Chronic liver disease and cirrhosis (K70,K73-K74) | 83 | 3.5 | 15.0 |
| 7 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 54 | 2.3 | 9.8 |
| 8 | Chronic lower respiratory diseases (J40-J47) | 47 | 2.0 | 8.5 |
| 9 | Viral hepatitis (B15-B19) | 32 | 1.4 | 5.8 |
| 10 | Septicemia (A40-A41) | 25 | 1.1 | 4.5 |
| 11 | Human immunodeficiency virus (HIV) disease (B20-B24) | 23 | 1.0 | 4.2 |
| 12 | Influenza and pneumonia (J10-J18) | 22 | 0.9 | 4.0 |
| 13 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 18 | 0.8 | * |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 15 | 0.6 | * |
| 15 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 13 | 0.6 | * |
| ... | All other causes (Residual) | 333 | 14.1 | 60.3 |
| Hispanic, female, 60-64 years | | | | |
| ... | All causes | 2,957 | 100.0 | 695.1 |
| 1 | Malignant neoplasms (C00-C97) | 1,058 | 35.8 | 248.7 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 642 | 21.7 | 150.9 |
| 3 | Diabetes mellitus (E10-E14) | 285 | 9.6 | 67.0 |
| 4 | Cerebrovascular diseases (I60-I69) | 146 | 4.9 | 34.3 |
| 5 | Chronic liver disease and cirrhosis (K70,K73-K74) | 83 | 2.8 | 19.5 |
| 6 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 73 | 2.5 | 17.2 |
| 7 | Chronic lower respiratory diseases (J40-J47) | 71 | 2.4 | 16.7 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 51 | 1.7 | 12.0 |
| 9 | Influenza and pneumonia (J10-J18) | 47 | 1.6 | 11.0 |
| 10 | Septicemia (A40-A41) | 40 | 1.4 | 9.4 |
| 11 | Viral hepatitis (B15-B19) | 37 | 1.3 | 8.7 |
| 12 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 31 | 1.0 | 7.3 |
| 13 | Human immunodeficiency virus (HIV) disease (B20-B24) | 17 | 0.6 | * |
| 13 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 17 | 0.6 | * |
| 15 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 10 | 0.3 | * |
| ... | All other causes (Residual) | 349 | 11.8 | 82.0 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|-------------------------------|---|--------|-------------------------|---------|
| Hispanic, female, 65-69 years | | | | |
| ... | All causes | 3,781 | 100.0 | 1,085.0 |
| 1 | Malignant neoplasms (C00-C97) | 1,143 | 30.2 | 328.0 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 922 | 24.4 | 264.6 |
| 3 | Diabetes mellitus (E10-E14) | 390 | 10.3 | 111.9 |
| 4 | Cerebrovascular diseases (I60-I69) | 202 | 5.3 | 58.0 |
| 5 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 118 | 3.1 | 33.9 |
| 6 | Chronic lower respiratory diseases (J40-J47) | 104 | 2.8 | 29.8 |
| 7 | Chronic liver disease and cirrhosis (K70,K73-K74) | 102 | 2.7 | 29.3 |
| 8 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 75 | 2.0 | 21.5 |
| 9 | Influenza and pneumonia (J10-J18) | 69 | 1.8 | 19.8 |
| 10 | Septicemia (A40-A41) | 60 | 1.6 | 17.2 |
| 11 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 42 | 1.1 | 12.1 |
| 12 | Viral hepatitis (B15-B19) | 35 | 0.9 | 10.0 |
| 13 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 26 | 0.7 | 7.5 |
| 14 | Alzheimer's disease (G30) | 16 | 0.4 | * |
| 15 | Parkinson's disease (G20-G21) | 11 | 0.3 | * |
| 15 | Complications of medical and surgical care (Y40-Y84,Y88) | 11 | 0.3 | * |
| ... | All other causes (Residual) | 455 | 12.0 | 130.6 |
| Hispanic, female, 70-74 years | | | | |
| ... | All causes | 5,160 | 100.0 | 1,800.8 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 1,384 | 26.8 | 483.0 |
| 2 | Malignant neoplasms (C00-C97) | 1,362 | 26.4 | 475.3 |
| 3 | Diabetes mellitus (E10-E14) | 456 | 8.8 | 159.1 |
| 4 | Cerebrovascular diseases (I60-I69) | 367 | 7.1 | 128.1 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 152 | 2.9 | 53.0 |
| 6 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 135 | 2.6 | 47.1 |
| 7 | Chronic liver disease and cirrhosis (K70,K73-K74) | 130 | 2.5 | 45.4 |
| 8 | Influenza and pneumonia (J10-J18) | 129 | 2.5 | 45.0 |
| 9 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 104 | 2.0 | 36.3 |
| 10 | Septicemia (A40-A41) | 79 | 1.5 | 27.6 |
| 11 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 61 | 1.2 | 21.3 |
| 12 | Alzheimer's disease (G30) | 52 | 1.0 | 18.1 |
| 13 | Viral hepatitis (B15-B19) | 43 | 0.8 | 15.0 |
| 14 | Parkinson's disease (G20-G21) | 31 | 0.6 | 10.8 |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 24 | 0.5 | 8.4 |
| ... | All other causes (Residual) | 651 | 12.6 | 227.2 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|-------------------------------|---|--------|-------------------------|---------|
| Hispanic, female, 75-79 years | | | | |
| ... | All causes | 6,025 | 100.0 | 2,917.0 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 1,844 | 30.6 | 892.8 |
| 2 | Malignant neoplasms (C00-C97) | 1,270 | 21.1 | 614.9 |
| 3 | Diabetes mellitus (E10-E14) | 514 | 8.5 | 248.9 |
| 4 | Cerebrovascular diseases (I60-I69) | 503 | 8.3 | 243.5 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 227 | 3.8 | 109.9 |
| 6 | Influenza and pneumonia (J10-J18) | 166 | 2.8 | 80.4 |
| 7 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 146 | 2.4 | 70.7 |
| 8 | Alzheimer's disease (G30) | 105 | 1.7 | 50.8 |
| 9 | Chronic liver disease and cirrhosis (K70,K73-K74) | 99 | 1.6 | 47.9 |
| 10 | Septicemia (A40-A41) | 90 | 1.5 | 43.6 |
| 11 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 86 | 1.4 | 41.6 |
| 12 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 77 | 1.3 | 37.3 |
| 13 | Parkinson's disease (G20-G21) | 45 | 0.7 | 21.8 |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 38 | 0.6 | 18.4 |
| 15 | Pneumonitis due to solids and liquids (J69) | 30 | 0.5 | 14.5 |
| ... | All other causes (Residual) | 785 | 13.0 | 380.1 |
| Hispanic, female, 80-84 years | | | | |
| ... | All causes | 5,853 | 100.0 | 4,707.4 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 1,980 | 33.8 | 1,592.5 |
| 2 | Malignant neoplasms (C00-C97) | 984 | 16.8 | 791.4 |
| 3 | Cerebrovascular diseases (I60-I69) | 525 | 9.0 | 422.2 |
| 4 | Diabetes mellitus (E10-E14) | 396 | 6.8 | 318.5 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 212 | 3.6 | 170.5 |
| 6 | Influenza and pneumonia (J10-J18) | 207 | 3.5 | 166.5 |
| 7 | Alzheimer's disease (G30) | 174 | 3.0 | 139.9 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 127 | 2.2 | 102.1 |
| 9 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 88 | 1.5 | 70.8 |
| 10 | Septicemia (A40-A41) | 86 | 1.5 | 69.2 |
| 11 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 80 | 1.4 | 64.3 |
| 12 | Chronic liver disease and cirrhosis (K70,K73-K74) | 66 | 1.1 | 53.1 |
| 13 | Parkinson's disease (G20-G21) | 44 | 0.8 | 35.4 |
| 14 | Pneumonitis due to solids and liquids (J69) | 41 | 0.7 | 33.0 |
| 15 | Atherosclerosis (I70) | 38 | 0.6 | 30.6 |
| ... | All other causes (Residual) | 805 | 13.8 | 647.4 |

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|-------------------------------|---|--------|-------------------------|------|
| Hispanic, female, 85-89 years | | | | |
| ... | All causes | 5,780 | 100.0 | --- |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 2,121 | 36.7 | --- |
| 2 | Malignant neoplasms (C00-C97) | 680 | 11.8 | --- |
| 3 | Cerebrovascular diseases (I60-I69) | 568 | 9.8 | --- |
| 4 | Diabetes mellitus (E10-E14) | 331 | 5.7 | --- |
| 5 | Influenza and pneumonia (J10-J18) | 288 | 5.0 | --- |
| 6 | Alzheimer's disease (G30) | 225 | 3.9 | --- |
| 7 | Chronic lower respiratory diseases (J40-J47) | 196 | 3.4 | --- |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 98 | 1.7 | --- |
| 9 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 92 | 1.6 | --- |
| 10 | Septicemia (A40-A41) | 87 | 1.5 | --- |
| 11 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 81 | 1.4 | --- |
| 12 | Atherosclerosis (I70) | 60 | 1.0 | --- |
| 13 | Pneumonitis due to solids and liquids (J69) | 56 | 1.0 | --- |
| 14 | Parkinson's disease (G20-G21) | 44 | 0.8 | --- |
| 15 | Chronic liver disease and cirrhosis (K70,K73-K74) | 30 | 0.5 | --- |
| ... | All other causes (Residual) | 823 | 14.2 | --- |
| Hispanic, female, 90-94 years | | | | |
| ... | All causes | 4,126 | 100.0 | --- |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 1,723 | 41.8 | --- |
| 2 | Cerebrovascular diseases (I60-I69) | 371 | 9.0 | --- |
| 3 | Malignant neoplasms (C00-C97) | 344 | 8.3 | --- |
| 4 | Influenza and pneumonia (J10-J18) | 237 | 5.7 | --- |
| 5 | Chronic lower respiratory diseases (J40-J47) | 159 | 3.9 | --- |
| 6 | Diabetes mellitus (E10-E14) | 156 | 3.8 | --- |
| 7 | Alzheimer's disease (G30) | 147 | 3.6 | --- |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 73 | 1.8 | --- |
| 9 | Septicemia (A40-A41) | 68 | 1.6 | --- |
| 10 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 58 | 1.4 | --- |
| 11 | Pneumonitis due to solids and liquids (J69) | 56 | 1.4 | --- |
| 12 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 49 | 1.2 | --- |
| 13 | Atherosclerosis (I70) | 46 | 1.1 | --- |
| 14 | Parkinson's disease (G20-G21) | 20 | 0.5 | --- |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 19 | 0.5 | --- |
| ... | All other causes (Residual) | 600 | 14.5 | --- |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|--------------------------------------|---|--------|-------------------------|------|
| Hispanic, female, 95-99 years | | | | |
| ... | All causes | 1,828 | 100.0 | --- |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 815 | 44.6 | --- |
| 2 | Cerebrovascular diseases (I60-I69) | 166 | 9.1 | --- |
| 3 | Malignant neoplasms (C00-C97) | 113 | 6.2 | --- |
| 4 | Influenza and pneumonia (J10-J18) | 104 | 5.7 | --- |
| 5 | Alzheimer's disease (G30) | 86 | 4.7 | --- |
| 6 | Diabetes mellitus (E10-E14) | 69 | 3.8 | --- |
| 7 | Chronic lower respiratory diseases (J40-J47) | 46 | 2.5 | --- |
| 8 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 34 | 1.9 | --- |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 29 | 1.6 | --- |
| 10 | Atherosclerosis (I70) | 28 | 1.5 | --- |
| 10 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 28 | 1.5 | --- |
| 12 | Septicemia (A40-A41) | 26 | 1.4 | --- |
| 13 | Pneumonitis due to solids and liquids (J69) | 25 | 1.4 | --- |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 8 | 0.4 | --- |
| 15 | Anemias (D50-D64) | 5 | 0.3 | --- |
| 15 | Nutritional deficiencies (E40-E64) | 5 | 0.3 | --- |
| 15 | Parkinson's disease (G20-G21) | 5 | 0.3 | --- |
| ... | All other causes (Residual) | 236 | 12.9 | --- |
| Hispanic, female, 100 years and over | | | | |
| ... | All causes | 406 | 100.0 | --- |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 198 | 48.8 | --- |
| 2 | Cerebrovascular diseases (I60-I69) | 35 | 8.6 | --- |
| 3 | Influenza and pneumonia (J10-J18) | 25 | 6.2 | --- |
| 4 | Alzheimer's disease (G30) | 20 | 4.9 | --- |
| 5 | Chronic lower respiratory diseases (J40-J47) | 12 | 3.0 | --- |
| 6 | Malignant neoplasms (C00-C97) | 11 | 2.7 | --- |
| 7 | Septicemia (A40-A41) | 9 | 2.2 | --- |
| 8 | Atherosclerosis (I70) | 8 | 2.0 | --- |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 7 | 1.7 | --- |
| 10 | Diabetes mellitus (E10-E14) | 5 | 1.2 | --- |
| 10 | Pneumonitis due to solids and liquids (J69) | 5 | 1.2 | --- |
| 12 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 4 | 1.0 | --- |
| 12 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 4 | 1.0 | --- |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 2 | 0.5 | --- |
| 14 | Parkinson's disease (G20-G21) | 2 | 0.5 | --- |
| ... | All other causes (Residual) | 59 | 14.5 | --- |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|--|---|-----------|-------------------------|-------|
| White, non-Hispanic, both sexes, all ages | | | | |
| ... | All causes | 1,962,810 | 100.0 | 991.1 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 582,349 | 29.7 | 294.1 |
| 2 | Malignant neoplasms (C00-C97) | 456,709 | 23.3 | 230.6 |
| 3 | Cerebrovascular diseases (I60-I69) | 133,879 | 6.8 | 67.6 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 110,753 | 5.6 | 55.9 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 76,262 | 3.9 | 38.5 |
| 6 | Influenza and pneumonia (J10-J18) | 51,952 | 2.6 | 26.2 |
| 7 | Diabetes mellitus (E10-E14) | 51,482 | 2.6 | 26.0 |
| 8 | Alzheimer's disease (G30) | 49,030 | 2.5 | 24.8 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 29,449 | 1.5 | 14.9 |
| 10 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 25,813 | 1.3 | 13.0 |
| 11 | Septicemia (A40-A41) | 24,389 | 1.2 | 12.3 |
| 12 | Chronic liver disease and cirrhosis (K70,K73-K74) | 20,065 | 1.0 | 10.1 |
| 13 | Parkinson's disease (G20-G21) | 15,239 | 0.8 | 7.7 |
| 14 | Pneumonitis due to solids and liquids (J69) | 14,843 | 0.8 | 7.5 |
| 15 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 13,837 | 0.7 | 7.0 |
| ... | All other causes (Residual) | 306,759 | 15.6 | 154.9 |
| White, non-Hispanic, both sexes, 1-4 years | | | | |
| ... | All causes | 2,772 | 100.0 | 30.1 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,006 | 36.3 | 10.9 |
| 2 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 287 | 10.4 | 3.1 |
| 3 | Malignant neoplasms (C00-C97) | 268 | 9.7 | 2.9 |
| 4 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 160 | 5.8 | 1.7 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 108 | 3.9 | 1.2 |
| 6 | Septicemia (A40-A41) | 63 | 2.3 | 0.7 |
| 7 | Influenza and pneumonia (J10-J18) | 48 | 1.7 | 0.5 |
| 8 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 36 | 1.3 | 0.4 |
| 9 | Certain conditions originating in the perinatal period (P00-P96) | 34 | 1.2 | 0.4 |
| 10 | Cerebrovascular diseases (I60-I69) | 28 | 1.0 | 0.3 |
| 11 | Meningitis (G00,G03) | 15 | 0.5 | * |
| 12 | Chronic lower respiratory diseases (J40-J47) | 13 | 0.5 | * |
| 13 | Acute bronchitis and bronchiolitis (J20-J21) | 12 | 0.4 | * |
| 14 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 11 | 0.4 | * |
| 15 | Meningococcal infection (A39) | 10 | 0.4 | * |
| 15 | Complications of medical and surgical care (Y40-Y84,Y88) | 10 | 0.4 | * |
| ... | All other causes (Residual) | 663 | 23.9 | 7.2 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|------|
| White, non-Hispanic, both sexes, 5-9 years | | | | |
| ... | All causes | 1,777 | 100.0 | 14.5 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 722 | 40.6 | 5.9 |
| 2 | Malignant neoplasms (C00-C97) | 302 | 17.0 | 2.5 |
| 3 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 111 | 6.2 | 0.9 |
| 4 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 82 | 4.6 | 0.7 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 52 | 2.9 | 0.4 |
| 6 | Influenza and pneumonia (J10-J18) | 28 | 1.6 | 0.2 |
| 7 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 24 | 1.4 | 0.2 |
| 8 | Cerebrovascular diseases (I60-I69) | 21 | 1.2 | 0.2 |
| 9 | Septicemia (A40-A41) | 17 | 1.0 | * |
| 10 | Chronic lower respiratory diseases (J40-J47) | 16 | 0.9 | * |
| 11 | Certain conditions originating in the perinatal period (P00-P96) | 11 | 0.6 | * |
| 12 | Anemias (D50-D64) | 9 | 0.5 | * |
| 13 | Meningitis (G00,G03) | 8 | 0.5 | * |
| 14 | Meningococcal infection (A39) | 6 | 0.3 | * |
| 14 | Human immunodeficiency virus (HIV) disease (B20-B24) | 6 | 0.3 | * |
| ... | All other causes (Residual) | 362 | 20.4 | 2.9 |
| White, non-Hispanic, both sexes, 10-14 years | | | | |
| ... | All causes | 2,374 | 100.0 | 18.0 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 966 | 40.7 | 7.3 |
| 2 | Malignant neoplasms (C00-C97) | 305 | 12.8 | 2.3 |
| 3 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 178 | 7.5 | 1.3 |
| 4 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 115 | 4.8 | 0.9 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 100 | 4.2 | 0.8 |
| 6 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 82 | 3.5 | 0.6 |
| 7 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 32 | 1.3 | 0.2 |
| 8 | Influenza and pneumonia (J10-J18) | 29 | 1.2 | 0.2 |
| 9 | Chronic lower respiratory diseases (J40-J47) | 23 | 1.0 | 0.2 |
| 10 | Cerebrovascular diseases (I60-I69) | 21 | 0.9 | 0.2 |
| 11 | Septicemia (A40-A41) | 18 | 0.8 | * |
| 12 | Diabetes mellitus (E10-E14) | 16 | 0.7 | * |
| 13 | Anemias (D50-D64) | 13 | 0.5 | * |
| 14 | Meningococcal infection (A39) | 12 | 0.5 | * |
| 15 | Meningitis (G00,G03) | 7 | 0.3 | * |
| 15 | Certain conditions originating in the perinatal period (P00-P96) | 7 | 0.3 | * |
| ... | All other causes (Residual) | 450 | 19.0 | 3.4 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|------|
| White, non-Hispanic, both sexes, 15-19 years | | | | |
| ... | All causes | 8,264 | 100.0 | 63.4 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 4,692 | 56.8 | 36.0 |
| 2 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 1,217 | 14.7 | 9.3 |
| 3 | Malignant neoplasms (C00-C97) | 448 | 5.4 | 3.4 |
| 4 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 382 | 4.6 | 2.9 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 186 | 2.3 | 1.4 |
| 6 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 157 | 1.9 | 1.2 |
| 7 | Cerebrovascular diseases (I60-I69) | 40 | 0.5 | 0.3 |
| 8 | Influenza and pneumonia (J10-J18) | 39 | 0.5 | 0.3 |
| 9 | Chronic lower respiratory diseases (J40-J47) | 33 | 0.4 | 0.3 |
| 10 | Septicemia (A40-A41) | 32 | 0.4 | 0.2 |
| 11 | Meningococcal infection (A39) | 29 | 0.4 | 0.2 |
| 11 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 29 | 0.4 | 0.2 |
| 13 | Diabetes mellitus (E10-E14) | 20 | 0.2 | 0.2 |
| 14 | Meningitis (G00,G03) | 14 | 0.2 | * |
| 15 | Pregnancy, childbirth and the puerperium (O00-099) | 13 | 0.2 | * |
| ... | All other causes (Residual) | 933 | 11.3 | 7.2 |
| White, non-Hispanic, both sexes, 20-24 years | | | | |
| ... | All causes | 10,466 | 100.0 | 85.4 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 5,175 | 49.4 | 42.2 |
| 2 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 1,709 | 16.3 | 13.9 |
| 3 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 676 | 6.5 | 5.5 |
| 4 | Malignant neoplasms (C00-C97) | 628 | 6.0 | 5.1 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 336 | 3.2 | 2.7 |
| 6 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 145 | 1.4 | 1.2 |
| 7 | Influenza and pneumonia (J10-J18) | 69 | 0.7 | 0.6 |
| 8 | Cerebrovascular diseases (I60-I69) | 58 | 0.6 | 0.5 |
| 9 | Diabetes mellitus (E10-E14) | 46 | 0.4 | 0.4 |
| 10 | Chronic lower respiratory diseases (J40-J47) | 42 | 0.4 | 0.3 |
| 11 | Septicemia (A40-A41) | 37 | 0.4 | 0.3 |
| 11 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 37 | 0.4 | 0.3 |
| 13 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 34 | 0.3 | 0.3 |
| 14 | Legal intervention (Y35,Y89.0) | 29 | 0.3 | 0.2 |
| 15 | Human immunodeficiency virus (HIV) disease (B20-B24) | 28 | 0.3 | 0.2 |
| ... | All other causes (Residual) | 1,417 | 13.5 | 11.6 |

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|-------|
| White, non-Hispanic, both sexes, 25-29 years | | | | |
| ... | All causes | 10,130 | 100.0 | 87.2 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 3,819 | 37.7 | 32.9 |
| 2 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 1,755 | 17.3 | 15.1 |
| 3 | Malignant neoplasms (C00-C97) | 865 | 8.5 | 7.4 |
| 4 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 750 | 7.4 | 6.5 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 563 | 5.6 | 4.8 |
| 6 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 152 | 1.5 | 1.3 |
| 7 | Human immunodeficiency virus (HIV) disease (B20-B24) | 110 | 1.1 | 0.9 |
| 8 | Diabetes mellitus (E10-E14) | 105 | 1.0 | 0.9 |
| 9 | Cerebrovascular diseases (I60-I69) | 101 | 1.0 | 0.9 |
| 10 | Influenza and pneumonia (J10-J18) | 72 | 0.7 | 0.6 |
| 11 | Septicemia (A40-A41) | 52 | 0.5 | 0.4 |
| 12 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 45 | 0.4 | 0.4 |
| 13 | Chronic lower respiratory diseases (J40-J47) | 43 | 0.4 | 0.4 |
| 14 | Chronic liver disease and cirrhosis (K70,K73-K74) | 35 | 0.3 | 0.3 |
| 14 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 35 | 0.3 | 0.3 |
| ... | All other causes (Residual) | 1,628 | 16.1 | 14.0 |
| White, non-Hispanic, both sexes, 30-34 years | | | | |
| ... | All causes | 14,061 | 100.0 | 104.4 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 3,979 | 28.3 | 29.6 |
| 2 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 2,116 | 15.0 | 15.7 |
| 3 | Malignant neoplasms (C00-C97) | 1,729 | 12.3 | 12.8 |
| 4 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 1,212 | 8.6 | 9.0 |
| 5 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 892 | 6.3 | 6.6 |
| 6 | Human immunodeficiency virus (HIV) disease (B20-B24) | 409 | 2.9 | 3.0 |
| 7 | Diabetes mellitus (E10-E14) | 230 | 1.6 | 1.7 |
| 8 | Cerebrovascular diseases (I60-I69) | 219 | 1.6 | 1.6 |
| 9 | Chronic liver disease and cirrhosis (K70,K73-K74) | 192 | 1.4 | 1.4 |
| 10 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 178 | 1.3 | 1.3 |
| 11 | Influenza and pneumonia (J10-J18) | 135 | 1.0 | 1.0 |
| 12 | Septicemia (A40-A41) | 93 | 0.7 | 0.7 |
| 13 | Chronic lower respiratory diseases (J40-J47) | 91 | 0.6 | 0.7 |
| 14 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 65 | 0.5 | 0.5 |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 61 | 0.4 | 0.5 |
| ... | All other causes (Residual) | 2,460 | 17.5 | 18.3 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|-------|
| White, non-Hispanic, both sexes, 35-39 years | | | | |
| ... | All causes | 23,334 | 100.0 | 153.1 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 5,314 | 22.8 | 34.9 |
| 2 | Malignant neoplasms (C00-C97) | 3,728 | 16.0 | 24.5 |
| 3 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 2,803 | 12.0 | 18.4 |
| 4 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 2,621 | 11.2 | 17.2 |
| 5 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 1,066 | 4.6 | 7.0 |
| 6 | Human immunodeficiency virus (HIV) disease (B20-B24) | 914 | 3.9 | 6.0 |
| 7 | Chronic liver disease and cirrhosis (K70,K73-K74) | 733 | 3.1 | 4.8 |
| 8 | Cerebrovascular diseases (I60-I69) | 452 | 1.9 | 3.0 |
| 9 | Diabetes mellitus (E10-E14) | 442 | 1.9 | 2.9 |
| 10 | Influenza and pneumonia (J10-J18) | 243 | 1.0 | 1.6 |
| 11 | Chronic lower respiratory diseases (J40-J47) | 187 | 0.8 | 1.2 |
| 11 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 187 | 0.8 | 1.2 |
| 13 | Septicemia (A40-A41) | 162 | 0.7 | 1.1 |
| 14 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 142 | 0.6 | 0.9 |
| 15 | Viral hepatitis (B15-B19) | 131 | 0.6 | 0.9 |
| ... | All other causes (Residual) | 4,209 | 18.0 | 27.6 |
| White, non-Hispanic, both sexes, 40-44 years | | | | |
| ... | All causes | 36,212 | 100.0 | 220.9 |
| 1 | Malignant neoplasms (C00-C97) | 7,823 | 21.6 | 47.7 |
| 2 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 6,026 | 16.6 | 36.8 |
| 3 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 5,931 | 16.4 | 36.2 |
| 4 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 2,983 | 8.2 | 18.2 |
| 5 | Chronic liver disease and cirrhosis (K70,K73-K74) | 1,479 | 4.1 | 9.0 |
| 6 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 1,015 | 2.8 | 6.2 |
| 7 | Human immunodeficiency virus (HIV) disease (B20-B24) | 997 | 2.8 | 6.1 |
| 8 | Cerebrovascular diseases (I60-I69) | 881 | 2.4 | 5.4 |
| 9 | Diabetes mellitus (E10-E14) | 791 | 2.2 | 4.8 |
| 10 | Chronic lower respiratory diseases (J40-J47) | 426 | 1.2 | 2.6 |
| 11 | Viral hepatitis (B15-B19) | 391 | 1.1 | 2.4 |
| 12 | Influenza and pneumonia (J10-J18) | 362 | 1.0 | 2.2 |
| 13 | Septicemia (A40-A41) | 296 | 0.8 | 1.8 |
| 14 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 252 | 0.7 | 1.5 |
| 15 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 217 | 0.6 | 1.3 |
| ... | All other causes (Residual) | 6,342 | 17.5 | 38.7 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|-------|
| White, non-Hispanic, both sexes, 45-49 years | | | | |
| ... | All causes | 50,151 | 100.0 | 324.7 |
| 1 | Malignant neoplasms (C00-C97) | 13,929 | 27.8 | 90.2 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 9,888 | 19.7 | 64.0 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 5,361 | 10.7 | 34.7 |
| 4 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 2,897 | 5.8 | 18.8 |
| 5 | Chronic liver disease and cirrhosis (K70,K73-K74) | 2,478 | 4.9 | 16.0 |
| 6 | Cerebrovascular diseases (I60-I69) | 1,363 | 2.7 | 8.8 |
| 7 | Diabetes mellitus (E10-E14) | 1,250 | 2.5 | 8.1 |
| 8 | Chronic lower respiratory diseases (J40-J47) | 831 | 1.7 | 5.4 |
| 9 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 792 | 1.6 | 5.1 |
| 10 | Human immunodeficiency virus (HIV) disease (B20-B24) | 771 | 1.5 | 5.0 |
| 11 | Viral hepatitis (B15-B19) | 742 | 1.5 | 4.8 |
| 12 | Influenza and pneumonia (J10-J18) | 516 | 1.0 | 3.3 |
| 13 | Septicemia (A40-A41) | 478 | 1.0 | 3.1 |
| 14 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 352 | 0.7 | 2.3 |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 235 | 0.5 | 1.5 |
| ... | All other causes (Residual) | 8,268 | 16.5 | 53.5 |
| White, non-Hispanic, both sexes, 50-54 years | | | | |
| ... | All causes | 66,935 | 100.0 | 474.0 |
| 1 | Malignant neoplasms (C00-C97) | 22,907 | 34.2 | 162.2 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 15,656 | 23.4 | 110.9 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 4,343 | 6.5 | 30.8 |
| 4 | Chronic liver disease and cirrhosis (K70,K73-K74) | 2,557 | 3.8 | 18.1 |
| 5 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 2,395 | 3.6 | 17.0 |
| 6 | Diabetes mellitus (E10-E14) | 2,082 | 3.1 | 14.7 |
| 7 | Cerebrovascular diseases (I60-I69) | 1,837 | 2.7 | 13.0 |
| 8 | Chronic lower respiratory diseases (J40-J47) | 1,739 | 2.6 | 12.3 |
| 9 | Septicemia (A40-A41) | 703 | 1.1 | 5.0 |
| 10 | Influenza and pneumonia (J10-J18) | 677 | 1.0 | 4.8 |
| 11 | Viral hepatitis (B15-B19) | 622 | 0.9 | 4.4 |
| 12 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 581 | 0.9 | 4.1 |
| 13 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 552 | 0.8 | 3.9 |
| 14 | Human immunodeficiency virus (HIV) disease (B20-B24) | 477 | 0.7 | 3.4 |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 326 | 0.5 | 2.3 |
| ... | All other causes (Residual) | 9,481 | 14.2 | 67.1 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|--|---|---------|-------------------------|---------|
| White, non-Hispanic, both sexes, 55-59 years | | | | |
| ... | All causes | 81,505 | 100.0 | 730.8 |
| 1 | Malignant neoplasms (C00-C97) | 31,194 | 38.3 | 279.7 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 20,256 | 24.9 | 181.6 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 3,548 | 4.4 | 31.8 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 3,132 | 3.8 | 28.1 |
| 5 | Diabetes mellitus (E10-E14) | 2,664 | 3.3 | 23.9 |
| 6 | Cerebrovascular diseases (I60-I69) | 2,606 | 3.2 | 23.4 |
| 7 | Chronic liver disease and cirrhosis (K70,K73-K74) | 2,127 | 2.6 | 19.1 |
| 8 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 1,787 | 2.2 | 16.0 |
| 9 | Septicemia (A40-A41) | 919 | 1.1 | 8.2 |
| 10 | Influenza and pneumonia (J10-J18) | 831 | 1.0 | 7.5 |
| 11 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 763 | 0.9 | 6.8 |
| 12 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 451 | 0.6 | 4.0 |
| 13 | Aortic aneurysm and dissection (I71) | 410 | 0.5 | 3.7 |
| 14 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 408 | 0.5 | 3.7 |
| 15 | Viral hepatitis (B15-B19) | 330 | 0.4 | 3.0 |
| ... | All other causes (Residual) | 10,079 | 12.4 | 90.4 |
| White, non-Hispanic, both sexes, 60-64 years | | | | |
| ... | All causes | 102,604 | 100.0 | 1,169.7 |
| 1 | Malignant neoplasms (C00-C97) | 40,195 | 39.2 | 458.2 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 26,177 | 25.5 | 298.4 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 6,130 | 6.0 | 69.9 |
| 4 | Cerebrovascular diseases (I60-I69) | 3,602 | 3.5 | 41.1 |
| 5 | Diabetes mellitus (E10-E14) | 3,454 | 3.4 | 39.4 |
| 6 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 2,644 | 2.6 | 30.1 |
| 7 | Chronic liver disease and cirrhosis (K70,K73-K74) | 2,062 | 2.0 | 23.5 |
| 8 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 1,192 | 1.2 | 13.6 |
| 9 | Septicemia (A40-A41) | 1,162 | 1.1 | 13.2 |
| 10 | Influenza and pneumonia (J10-J18) | 1,132 | 1.1 | 12.9 |
| 11 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1,098 | 1.1 | 12.5 |
| 12 | Aortic aneurysm and dissection (I71) | 707 | 0.7 | 8.1 |
| 13 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 555 | 0.5 | 6.3 |
| 14 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 453 | 0.4 | 5.2 |
| 15 | Alzheimer's disease (G30) | 328 | 0.3 | 3.7 |
| ... | All other causes (Residual) | 11,713 | 11.4 | 133.5 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|-------------------|---|---------|-------------------------|---------|
| | White, non-Hispanic, both sexes, 65-69 years | | | |
| ... | All causes | 139,503 | 100.0 | 1,827.4 |
| 1 | Malignant neoplasms (C00-C97) | 52,117 | 37.4 | 682.7 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 36,014 | 25.8 | 471.8 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 10,528 | 7.5 | 137.9 |
| 4 | Cerebrovascular diseases (I60-I69) | 5,715 | 4.1 | 74.9 |
| 5 | Diabetes mellitus (E10-E14) | 4,736 | 3.4 | 62.0 |
| 6 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 2,784 | 2.0 | 36.5 |
| 7 | Chronic liver disease and cirrhosis (K70,K73-K74) | 2,112 | 1.5 | 27.7 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1,907 | 1.4 | 25.0 |
| 9 | Influenza and pneumonia (J10-J18) | 1,773 | 1.3 | 23.2 |
| 10 | Septicemia (A40-A41) | 1,704 | 1.2 | 22.3 |
| 11 | Aortic aneurysm and dissection (I71) | 1,228 | 0.9 | 16.1 |
| 12 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 1,085 | 0.8 | 14.2 |
| 13 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 837 | 0.6 | 11.0 |
| 14 | Alzheimer's disease (G30) | 736 | 0.5 | 9.6 |
| 15 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 679 | 0.5 | 8.9 |
| ... | All other causes (Residual) | 15,548 | 11.1 | 203.7 |
| | White, non-Hispanic, both sexes, 70-74 years | | | |
| ... | All causes | 206,886 | 100.0 | 2,849.2 |
| 1 | Malignant neoplasms (C00-C97) | 69,412 | 33.6 | 955.9 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 55,841 | 27.0 | 769.0 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 17,165 | 8.3 | 236.4 |
| 4 | Cerebrovascular diseases (I60-I69) | 11,123 | 5.4 | 153.2 |
| 5 | Diabetes mellitus (E10-E14) | 6,728 | 3.3 | 92.7 |
| 6 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 3,525 | 1.7 | 48.5 |
| 7 | Influenza and pneumonia (J10-J18) | 3,441 | 1.7 | 47.4 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 3,077 | 1.5 | 42.4 |
| 9 | Septicemia (A40-A41) | 2,681 | 1.3 | 36.9 |
| 10 | Alzheimer's disease (G30) | 2,284 | 1.1 | 31.5 |
| 11 | Chronic liver disease and cirrhosis (K70,K73-K74) | 2,195 | 1.1 | 30.2 |
| 12 | Aortic aneurysm and dissection (I71) | 1,851 | 0.9 | 25.5 |
| 13 | Parkinson's disease (G20-G21) | 1,428 | 0.7 | 19.7 |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 1,348 | 0.7 | 18.6 |
| 15 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 1,102 | 0.5 | 15.2 |
| ... | All other causes (Residual) | 23,685 | 11.4 | 326.2 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|--|---|---------|-------------------------|---------|
| White, non-Hispanic, both sexes, 75-79 years | | | | |
| ... | All causes | 282,018 | 100.0 | 4,476.8 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 82,060 | 29.1 | 1,302.6 |
| 2 | Malignant neoplasms (C00-C97) | 77,026 | 27.3 | 1,222.7 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 22,403 | 7.9 | 355.6 |
| 4 | Cerebrovascular diseases (I60-I69) | 19,553 | 6.9 | 310.4 |
| 5 | Diabetes mellitus (E10-E14) | 8,585 | 3.0 | 136.3 |
| 6 | Influenza and pneumonia (J10-J18) | 6,188 | 2.2 | 98.2 |
| 7 | Alzheimer's disease (G30) | 5,829 | 2.1 | 92.5 |
| 8 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 5,160 | 1.8 | 81.9 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 4,686 | 1.7 | 74.4 |
| 10 | Septicemia (A40-A41) | 3,928 | 1.4 | 62.4 |
| 11 | Parkinson's disease (G20-G21) | 3,172 | 1.1 | 50.4 |
| 12 | Aortic aneurysm and dissection (I71) | 2,561 | 0.9 | 40.7 |
| 13 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 1,945 | 0.7 | 30.9 |
| 13 | Chronic liver disease and cirrhosis (K70,K73-K74) | 1,945 | 0.7 | 30.9 |
| 15 | Pneumonitis due to solids and liquids (J69) | 1,933 | 0.7 | 30.7 |
| ... | All other causes (Residual) | 35,044 | 12.4 | 556.3 |
| White, non-Hispanic, both sexes, 80-84 years | | | | |
| ... | All causes | 321,125 | 100.0 | 7,196.1 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 103,476 | 32.2 | 2,318.8 |
| 2 | Malignant neoplasms (C00-C97) | 65,734 | 20.5 | 1,473.0 |
| 3 | Cerebrovascular diseases (I60-I69) | 27,751 | 8.6 | 621.9 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 21,569 | 6.7 | 483.3 |
| 5 | Alzheimer's disease (G30) | 10,913 | 3.4 | 244.5 |
| 6 | Influenza and pneumonia (J10-J18) | 9,649 | 3.0 | 216.2 |
| 7 | Diabetes mellitus (E10-E14) | 8,922 | 2.8 | 199.9 |
| 8 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 5,997 | 1.9 | 134.4 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 5,712 | 1.8 | 128.0 |
| 10 | Septicemia (A40-A41) | 4,424 | 1.4 | 99.1 |
| 11 | Parkinson's disease (G20-G21) | 4,349 | 1.4 | 97.5 |
| 12 | Pneumonitis due to solids and liquids (J69) | 3,079 | 1.0 | 69.0 |
| 13 | Aortic aneurysm and dissection (I71) | 2,594 | 0.8 | 58.1 |
| 14 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 2,591 | 0.8 | 58.1 |
| 15 | Atherosclerosis (I70) | 2,267 | 0.7 | 50.8 |
| ... | All other causes (Residual) | 42,098 | 13.1 | 943.4 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|--|---|---------|-------------------------|------|
| White, non-Hispanic, both sexes, 85-89 years | | | | |
| ... | All causes | 301,010 | 100.0 | --- |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 106,333 | 35.3 | --- |
| 2 | Malignant neoplasms (C00-C97) | 43,739 | 14.5 | --- |
| 3 | Cerebrovascular diseases (I60-I69) | 29,529 | 9.8 | --- |
| 4 | Chronic lower respiratory diseases (J40-J47) | 16,195 | 5.4 | --- |
| 5 | Alzheimer's disease (G30) | 13,619 | 4.5 | --- |
| 6 | Influenza and pneumonia (J10-J18) | 11,821 | 3.9 | --- |
| 7 | Diabetes mellitus (E10-E14) | 6,742 | 2.2 | --- |
| 8 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 5,711 | 1.9 | --- |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 5,617 | 1.9 | --- |
| 10 | Septicemia (A40-A41) | 4,040 | 1.3 | --- |
| 11 | Parkinson's disease (G20-G21) | 3,528 | 1.2 | --- |
| 12 | Pneumonitis due to solids and liquids (J69) | 3,503 | 1.2 | --- |
| 13 | Atherosclerosis (I70) | 2,926 | 1.0 | --- |
| 14 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 2,922 | 1.0 | --- |
| 15 | Aortic aneurysm and dissection (I71) | 1,951 | 0.6 | --- |
| ... | All other causes (Residual) | 42,834 | 14.2 | --- |
| White, non-Hispanic, both sexes, 90-94 years | | | | |
| ... | All causes | 197,069 | 100.0 | --- |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 76,775 | 39.0 | --- |
| 2 | Cerebrovascular diseases (I60-I69) | 20,201 | 10.3 | --- |
| 3 | Malignant neoplasms (C00-C97) | 18,926 | 9.6 | --- |
| 4 | Alzheimer's disease (G30) | 10,349 | 5.3 | --- |
| 5 | Influenza and pneumonia (J10-J18) | 9,463 | 4.8 | --- |
| 6 | Chronic lower respiratory diseases (J40-J47) | 7,328 | 3.7 | --- |
| 7 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 3,729 | 1.9 | --- |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 3,572 | 1.8 | --- |
| 9 | Diabetes mellitus (E10-E14) | 3,467 | 1.8 | --- |
| 10 | Pneumonitis due to solids and liquids (J69) | 2,601 | 1.3 | --- |
| 11 | Atherosclerosis (I70) | 2,561 | 1.3 | --- |
| 12 | Septicemia (A40-A41) | 2,499 | 1.3 | --- |
| 13 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 2,156 | 1.1 | --- |
| 14 | Parkinson's disease (G20-G21) | 1,547 | 0.8 | --- |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 1,015 | 0.5 | --- |
| ... | All other causes (Residual) | 30,880 | 15.7 | --- |

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|---|---|--------|-------------------------|------|
| White, non-Hispanic, both sexes, 95-99 years | | | | |
| ... | All causes | 75,705 | 100.0 | --- |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 31,359 | 41.4 | --- |
| 2 | Cerebrovascular diseases (I60-I69) | 7,436 | 9.8 | --- |
| 3 | Malignant neoplasms (C00-C97) | 4,811 | 6.4 | --- |
| 4 | Influenza and pneumonia (J10-J18) | 4,242 | 5.6 | --- |
| 5 | Alzheimer's disease (G30) | 4,070 | 5.4 | --- |
| 6 | Chronic lower respiratory diseases (J40-J47) | 2,108 | 2.8 | --- |
| 7 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,376 | 1.8 | --- |
| 8 | Atherosclerosis (I70) | 1,333 | 1.8 | --- |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1,293 | 1.7 | --- |
| 10 | Diabetes mellitus (E10-E14) | 1,057 | 1.4 | --- |
| 11 | Pneumonitis due to solids and liquids (J69) | 1,011 | 1.3 | --- |
| 12 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 958 | 1.3 | --- |
| 13 | Septicemia (A40-A41) | 825 | 1.1 | --- |
| 14 | Parkinson's disease (G20-G21) | 392 | 0.5 | --- |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 303 | 0.4 | --- |
| ... | All other causes (Residual) | 13,131 | 17.3 | --- |
| White, non-Hispanic, both sexes, 100 years and over | | | | |
| ... | All causes | 15,485 | 100.0 | --- |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 6,963 | 45.0 | --- |
| 2 | Cerebrovascular diseases (I60-I69) | 1,295 | 8.4 | --- |
| 3 | Influenza and pneumonia (J10-J18) | 1,076 | 6.9 | --- |
| 4 | Alzheimer's disease (G30) | 682 | 4.4 | --- |
| 5 | Malignant neoplasms (C00-C97) | 577 | 3.7 | --- |
| 6 | Atherosclerosis (I70) | 366 | 2.4 | --- |
| 7 | Chronic lower respiratory diseases (J40-J47) | 319 | 2.1 | --- |
| 8 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 276 | 1.8 | --- |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 241 | 1.6 | --- |
| 10 | Pneumonitis due to solids and liquids (J69) | 214 | 1.4 | --- |
| 11 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 206 | 1.3 | --- |
| 12 | Diabetes mellitus (E10-E14) | 136 | 0.9 | --- |
| 13 | Septicemia (A40-A41) | 134 | 0.9 | --- |
| 14 | Nutritional deficiencies (E40-E64) | 76 | 0.5 | --- |
| 15 | Anemias (D50-D64) | 66 | 0.4 | --- |
| ... | All other causes (Residual) | 2,858 | 18.5 | --- |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|--------------------------------------|---|---------|-------------------------|-------|
| White, non-Hispanic, male, all ages | | | | |
| ... | All causes | 945,967 | 100.0 | 975.6 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 280,713 | 29.7 | 289.5 |
| 2 | Malignant neoplasms (C00-C97) | 236,004 | 24.9 | 243.4 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 52,952 | 5.6 | 54.6 |
| 4 | Cerebrovascular diseases (I60-I69) | 50,382 | 5.3 | 52.0 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 48,180 | 5.1 | 49.7 |
| 6 | Diabetes mellitus (E10-E14) | 24,302 | 2.6 | 25.1 |
| 7 | Influenza and pneumonia (J10-J18) | 22,389 | 2.4 | 23.1 |
| 8 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 20,710 | 2.2 | 21.4 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 14,331 | 1.5 | 14.8 |
| 10 | Alzheimer's disease (G30) | 14,295 | 1.5 | 14.7 |
| 11 | Chronic liver disease and cirrhosis (K70,K73-K74) | 12,609 | 1.3 | 13.0 |
| 12 | Septicemia (A40-A41) | 10,668 | 1.1 | 11.0 |
| 13 | Parkinson's disease (G20-G21) | 8,685 | 0.9 | 9.0 |
| 14 | Aortic aneurysm and dissection (I71) | 7,924 | 0.8 | 8.2 |
| 15 | Pneumonitis due to solids and liquids (J69) | 7,411 | 0.8 | 7.6 |
| ... | All other causes (Residual) | 134,412 | 14.2 | 138.6 |
| White, non-Hispanic, male, 1-4 years | | | | |
| ... | All causes | 1,592 | 100.0 | 33.7 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 624 | 39.2 | 13.2 |
| 2 | Malignant neoplasms (C00-C97) | 144 | 9.0 | 3.1 |
| 2 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 144 | 9.0 | 3.1 |
| 4 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 95 | 6.0 | 2.0 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 59 | 3.7 | 1.3 |
| 6 | Septicemia (A40-A41) | 41 | 2.6 | 0.9 |
| 7 | Influenza and pneumonia (J10-J18) | 26 | 1.6 | 0.6 |
| 8 | Certain conditions originating in the perinatal period (P00-P96) | 22 | 1.4 | 0.5 |
| 9 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 18 | 1.1 | * |
| 10 | Cerebrovascular diseases (I60-I69) | 14 | 0.9 | * |
| 11 | Meningitis (G00,G03) | 10 | 0.6 | * |
| 12 | Chronic lower respiratory diseases (J40-J47) | 9 | 0.6 | * |
| 13 | Complications of medical and surgical care (Y40-Y84,Y88) | 7 | 0.4 | * |
| 14 | Meningococcal infection (A39) | 5 | 0.3 | * |
| 14 | Acute bronchitis and bronchiolitis (J20-J21) | 5 | 0.3 | * |
| ... | All other causes (Residual) | 369 | 23.2 | 7.8 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|------|
| White, non-Hispanic, male, 5-9 years | | | | |
| ... | All causes | 957 | 100.0 | 15.2 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 424 | 44.3 | 6.7 |
| 2 | Malignant neoplasms (C00-C97) | 146 | 15.3 | 2.3 |
| 3 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 64 | 6.7 | 1.0 |
| 4 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 40 | 4.2 | 0.6 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 23 | 2.4 | 0.4 |
| 6 | Influenza and pneumonia (J10-J18) | 14 | 1.5 | * |
| 7 | Cerebrovascular diseases (I60-I69) | 12 | 1.3 | * |
| 8 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 10 | 1.0 | * |
| 9 | Chronic lower respiratory diseases (J40-J47) | 9 | 0.9 | * |
| 9 | Certain conditions originating in the perinatal period (P00-P96) | 9 | 0.9 | * |
| 11 | Septicemia (A40-A41) | 7 | 0.7 | * |
| 12 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 5 | 0.5 | * |
| 13 | Human immunodeficiency virus (HIV) disease (B20-B24) | 4 | 0.4 | * |
| 13 | Anemias (D50-D64) | 4 | 0.4 | * |
| 13 | Meningitis (G00,G03) | 4 | 0.4 | * |
| ... | All other causes (Residual) | 182 | 19.0 | 2.9 |
| White, non-Hispanic, male, 10-14 years | | | | |
| ... | All causes | 1,471 | 100.0 | 21.7 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 642 | 43.6 | 9.5 |
| 2 | Malignant neoplasms (C00-C97) | 164 | 11.1 | 2.4 |
| 3 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 139 | 9.4 | 2.1 |
| 4 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 71 | 4.8 | 1.0 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 57 | 3.9 | 0.8 |
| 6 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 43 | 2.9 | 0.6 |
| 7 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 15 | 1.0 | * |
| 8 | Influenza and pneumonia (J10-J18) | 14 | 1.0 | * |
| 9 | Chronic lower respiratory diseases (J40-J47) | 13 | 0.9 | * |
| 10 | Diabetes mellitus (E10-E14) | 12 | 0.8 | * |
| 10 | Cerebrovascular diseases (I60-I69) | 12 | 0.8 | * |
| 12 | Septicemia (A40-A41) | 11 | 0.7 | * |
| 13 | Meningococcal infection (A39) | 8 | 0.5 | * |
| 14 | Anemias (D50-D64) | 7 | 0.5 | * |
| 15 | Certain conditions originating in the perinatal period (P00-P96) | 5 | 0.3 | * |
| ... | All other causes (Residual) | 258 | 17.5 | 3.8 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|-------|
| White, non-Hispanic, male, 15-19 years | | | | |
| ... | All causes | 5,756 | 100.0 | 86.1 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 3,266 | 56.7 | 48.8 |
| 2 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 1,030 | 17.9 | 15.4 |
| 3 | Malignant neoplasms (C00-C97) | 270 | 4.7 | 4.0 |
| 4 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 263 | 4.6 | 3.9 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 117 | 2.0 | 1.7 |
| 6 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 100 | 1.7 | 1.5 |
| 7 | Cerebrovascular diseases (I60-I69) | 20 | 0.3 | 0.3 |
| 8 | Septicemia (A40-A41) | 18 | 0.3 | * |
| 8 | Influenza and pneumonia (J10-J18) | 18 | 0.3 | * |
| 10 | Chronic lower respiratory diseases (J40-J47) | 15 | 0.3 | * |
| 11 | Diabetes mellitus (E10-E14) | 14 | 0.2 | * |
| 12 | Meningococcal infection (A39) | 13 | 0.2 | * |
| 13 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 10 | 0.2 | * |
| 14 | Meningitis (G00,G03) | 7 | 0.1 | * |
| 14 | Aortic aneurysm and dissection (I71) | 7 | 0.1 | * |
| 14 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 7 | 0.1 | * |
| 14 | Legal intervention (Y35,Y89.0) | 7 | 0.1 | * |
| ... | All other causes (Residual) | 574 | 10.0 | 8.6 |
| White, non-Hispanic, male, 20-24 years | | | | |
| ... | All causes | 7,786 | 100.0 | 125.6 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 4,031 | 51.8 | 65.0 |
| 2 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 1,492 | 19.2 | 24.1 |
| 3 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 466 | 6.0 | 7.5 |
| 4 | Malignant neoplasms (C00-C97) | 362 | 4.6 | 5.8 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 219 | 2.8 | 3.5 |
| 6 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 83 | 1.1 | 1.3 |
| 7 | Influenza and pneumonia (J10-J18) | 46 | 0.6 | 0.7 |
| 8 | Cerebrovascular diseases (I60-I69) | 35 | 0.4 | 0.6 |
| 9 | Diabetes mellitus (E10-E14) | 28 | 0.4 | 0.5 |
| 9 | Legal intervention (Y35,Y89.0) | 28 | 0.4 | 0.5 |
| 11 | Chronic lower respiratory diseases (J40-J47) | 24 | 0.3 | 0.4 |
| 12 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 21 | 0.3 | 0.3 |
| 13 | Septicemia (A40-A41) | 19 | 0.2 | * |
| 14 | Human immunodeficiency virus (HIV) disease (B20-B24) | 15 | 0.2 | * |
| 15 | Aortic aneurysm and dissection (I71) | 14 | 0.2 | * |
| 15 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 14 | 0.2 | * |
| ... | All other causes (Residual) | 889 | 11.4 | 14.3 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|-------|
| White, non-Hispanic, male, 25-29 years | | | | |
| ... | All causes | 7,168 | 100.0 | 122.4 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 2,908 | 40.6 | 49.7 |
| 2 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 1,479 | 20.6 | 25.3 |
| 3 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 542 | 7.6 | 9.3 |
| 4 | Malignant neoplasms (C00-C97) | 449 | 6.3 | 7.7 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 367 | 5.1 | 6.3 |
| 6 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 87 | 1.2 | 1.5 |
| 7 | Human immunodeficiency virus (HIV) disease (B20-B24) | 79 | 1.1 | 1.3 |
| 8 | Diabetes mellitus (E10-E14) | 54 | 0.8 | 0.9 |
| 9 | Cerebrovascular diseases (I60-I69) | 49 | 0.7 | 0.8 |
| 10 | Influenza and pneumonia (J10-J18) | 33 | 0.5 | 0.6 |
| 11 | Septicemia (A40-A41) | 28 | 0.4 | 0.5 |
| 12 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 25 | 0.3 | 0.4 |
| 13 | Aortic aneurysm and dissection (I71) | 24 | 0.3 | 0.4 |
| 14 | Chronic lower respiratory diseases (J40-J47) | 23 | 0.3 | 0.4 |
| 14 | Chronic liver disease and cirrhosis (K70,K73-K74) | 23 | 0.3 | 0.4 |
| ... | All other causes (Residual) | 998 | 13.9 | 17.0 |
| White, non-Hispanic, male, 30-34 years | | | | |
| ... | All causes | 9,445 | 100.0 | 139.3 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 2,998 | 31.7 | 44.2 |
| 2 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 1,711 | 18.1 | 25.2 |
| 3 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 817 | 8.7 | 12.1 |
| 4 | Malignant neoplasms (C00-C97) | 811 | 8.6 | 12.0 |
| 5 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 635 | 6.7 | 9.4 |
| 6 | Human immunodeficiency virus (HIV) disease (B20-B24) | 331 | 3.5 | 4.9 |
| 7 | Diabetes mellitus (E10-E14) | 130 | 1.4 | 1.9 |
| 8 | Chronic liver disease and cirrhosis (K70,K73-K74) | 122 | 1.3 | 1.8 |
| 9 | Cerebrovascular diseases (I60-I69) | 112 | 1.2 | 1.7 |
| 10 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 99 | 1.0 | 1.5 |
| 11 | Influenza and pneumonia (J10-J18) | 79 | 0.8 | 1.2 |
| 12 | Septicemia (A40-A41) | 41 | 0.4 | 0.6 |
| 13 | Chronic lower respiratory diseases (J40-J47) | 39 | 0.4 | 0.6 |
| 14 | Legal intervention (Y35,Y89.0) | 36 | 0.4 | 0.5 |
| 15 | Aortic aneurysm and dissection (I71) | 34 | 0.4 | 0.5 |
| ... | All other causes (Residual) | 1,450 | 15.4 | 21.4 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|-------|
| White, non-Hispanic, male, 35-39 years | | | | |
| ... | All causes | 15,102 | 100.0 | 197.9 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 3,794 | 25.1 | 49.7 |
| 2 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 2,046 | 13.5 | 26.8 |
| 3 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 1,984 | 13.1 | 26.0 |
| 4 | Malignant neoplasms (C00-C97) | 1,621 | 10.7 | 21.2 |
| 5 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 788 | 5.2 | 10.3 |
| 6 | Human immunodeficiency virus (HIV) disease (B20-B24) | 781 | 5.2 | 10.2 |
| 7 | Chronic liver disease and cirrhosis (K70,K73-K74) | 443 | 2.9 | 5.8 |
| 8 | Diabetes mellitus (E10-E14) | 262 | 1.7 | 3.4 |
| 9 | Cerebrovascular diseases (I60-I69) | 223 | 1.5 | 2.9 |
| 10 | Influenza and pneumonia (J10-J18) | 131 | 0.9 | 1.7 |
| 11 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 102 | 0.7 | 1.3 |
| 12 | Chronic lower respiratory diseases (J40-J47) | 97 | 0.6 | 1.3 |
| 13 | Septicemia (A40-A41) | 95 | 0.6 | 1.2 |
| 14 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 89 | 0.6 | 1.2 |
| 15 | Viral hepatitis (B15-B19) | 88 | 0.6 | 1.2 |
| ... | All other causes (Residual) | 2,558 | 16.9 | 33.5 |
| White, non-Hispanic, male, 40-44 years | | | | |
| ... | All causes | 23,111 | 100.0 | 282.4 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 4,411 | 19.1 | 53.9 |
| 2 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 4,394 | 19.0 | 53.7 |
| 3 | Malignant neoplasms (C00-C97) | 3,539 | 15.3 | 43.2 |
| 4 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 2,283 | 9.9 | 27.9 |
| 5 | Chronic liver disease and cirrhosis (K70,K73-K74) | 960 | 4.2 | 11.7 |
| 6 | Human immunodeficiency virus (HIV) disease (B20-B24) | 858 | 3.7 | 10.5 |
| 7 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 710 | 3.1 | 8.7 |
| 8 | Diabetes mellitus (E10-E14) | 505 | 2.2 | 6.2 |
| 9 | Cerebrovascular diseases (I60-I69) | 461 | 2.0 | 5.6 |
| 10 | Viral hepatitis (B15-B19) | 280 | 1.2 | 3.4 |
| 11 | Influenza and pneumonia (J10-J18) | 202 | 0.9 | 2.5 |
| 12 | Chronic lower respiratory diseases (J40-J47) | 196 | 0.8 | 2.4 |
| 13 | Septicemia (A40-A41) | 171 | 0.7 | 2.1 |
| 14 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 134 | 0.6 | 1.6 |
| 15 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 122 | 0.5 | 1.5 |
| ... | All other causes (Residual) | 3,885 | 16.8 | 47.5 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|-------|
| White, non-Hispanic, male, 45-49 years | | | | |
| ... | All causes | 32,045 | 100.0 | 417.3 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 7,516 | 23.5 | 97.9 |
| 2 | Malignant neoplasms (C00-C97) | 6,840 | 21.3 | 89.1 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 3,941 | 12.3 | 51.3 |
| 4 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 2,215 | 6.9 | 28.8 |
| 5 | Chronic liver disease and cirrhosis (K70,K73-K74) | 1,780 | 5.6 | 23.2 |
| 6 | Diabetes mellitus (E10-E14) | 761 | 2.4 | 9.9 |
| 7 | Cerebrovascular diseases (I60-I69) | 739 | 2.3 | 9.6 |
| 8 | Human immunodeficiency virus (HIV) disease (B20-B24) | 674 | 2.1 | 8.8 |
| 9 | Viral hepatitis (B15-B19) | 570 | 1.8 | 7.4 |
| 9 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 570 | 1.8 | 7.4 |
| 11 | Chronic lower respiratory diseases (J40-J47) | 456 | 1.4 | 5.9 |
| 12 | Influenza and pneumonia (J10-J18) | 311 | 1.0 | 4.0 |
| 13 | Septicemia (A40-A41) | 250 | 0.8 | 3.3 |
| 14 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 209 | 0.7 | 2.7 |
| 15 | Aortic aneurysm and dissection (I71) | 148 | 0.5 | 1.9 |
| ... | All other causes (Residual) | 5,065 | 15.8 | 66.0 |
| White, non-Hispanic, male, 50-54 years | | | | |
| ... | All causes | 41,686 | 100.0 | 596.5 |
| 1 | Malignant neoplasms (C00-C97) | 11,839 | 28.4 | 169.4 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 11,662 | 28.0 | 166.9 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 3,093 | 7.4 | 44.3 |
| 4 | Chronic liver disease and cirrhosis (K70,K73-K74) | 1,865 | 4.5 | 26.7 |
| 5 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 1,801 | 4.3 | 25.8 |
| 6 | Diabetes mellitus (E10-E14) | 1,253 | 3.0 | 17.9 |
| 7 | Cerebrovascular diseases (I60-I69) | 1,006 | 2.4 | 14.4 |
| 8 | Chronic lower respiratory diseases (J40-J47) | 875 | 2.1 | 12.5 |
| 9 | Viral hepatitis (B15-B19) | 492 | 1.2 | 7.0 |
| 10 | Human immunodeficiency virus (HIV) disease (B20-B24) | 432 | 1.0 | 6.2 |
| 11 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 420 | 1.0 | 6.0 |
| 12 | Influenza and pneumonia (J10-J18) | 396 | 0.9 | 5.7 |
| 13 | Septicemia (A40-A41) | 365 | 0.9 | 5.2 |
| 14 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 334 | 0.8 | 4.8 |
| 15 | Aortic aneurysm and dissection (I71) | 194 | 0.5 | 2.8 |
| ... | All other causes (Residual) | 5,659 | 13.6 | 81.0 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|---------|
| White, non-Hispanic, male, 55-59 years | | | | |
| ... | All causes | 49,212 | 100.0 | 901.6 |
| 1 | Malignant neoplasms (C00-C97) | 16,696 | 33.9 | 305.9 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 14,516 | 29.5 | 265.9 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 2,142 | 4.4 | 39.2 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 1,823 | 3.7 | 33.4 |
| 5 | Diabetes mellitus (E10-E14) | 1,585 | 3.2 | 29.0 |
| 6 | Chronic liver disease and cirrhosis (K70,K73-K74) | 1,511 | 3.1 | 27.7 |
| 7 | Cerebrovascular diseases (I60-I69) | 1,421 | 2.9 | 26.0 |
| 8 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 1,375 | 2.8 | 25.2 |
| 9 | Influenza and pneumonia (J10-J18) | 484 | 1.0 | 8.9 |
| 10 | Septicemia (A40-A41) | 470 | 1.0 | 8.6 |
| 11 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 425 | 0.9 | 7.8 |
| 12 | Aortic aneurysm and dissection (I71) | 320 | 0.7 | 5.9 |
| 13 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 292 | 0.6 | 5.3 |
| 14 | Human immunodeficiency virus (HIV) disease (B20-B24) | 262 | 0.5 | 4.8 |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 234 | 0.5 | 4.3 |
| ... | All other causes (Residual) | 5,656 | 11.5 | 103.6 |
| White, non-Hispanic, male, 60-64 years | | | | |
| ... | All causes | 60,781 | 100.0 | 1,439.8 |
| 1 | Malignant neoplasms (C00-C97) | 22,188 | 36.5 | 525.6 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 17,988 | 29.6 | 426.1 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 3,062 | 5.0 | 72.5 |
| 4 | Cerebrovascular diseases (I60-I69) | 1,979 | 3.3 | 46.9 |
| 5 | Diabetes mellitus (E10-E14) | 1,939 | 3.2 | 45.9 |
| 6 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,763 | 2.9 | 41.8 |
| 7 | Chronic liver disease and cirrhosis (K70,K73-K74) | 1,364 | 2.2 | 32.3 |
| 8 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 909 | 1.5 | 21.5 |
| 9 | Influenza and pneumonia (J10-J18) | 648 | 1.1 | 15.3 |
| 10 | Septicemia (A40-A41) | 601 | 1.0 | 14.2 |
| 11 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 587 | 1.0 | 13.9 |
| 12 | Aortic aneurysm and dissection (I71) | 521 | 0.9 | 12.3 |
| 13 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 313 | 0.5 | 7.4 |
| 14 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 247 | 0.4 | 5.9 |
| 15 | Pneumonitis due to solids and liquids (J69) | 173 | 0.3 | 4.1 |
| ... | All other causes (Residual) | 6,499 | 10.7 | 153.9 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|--|---|---------|-------------------------|---------|
| White, non-Hispanic, male, 65-69 years | | | | |
| ... | All causes | 80,805 | 100.0 | 2,256.8 |
| 1 | Malignant neoplasms (C00-C97) | 29,183 | 36.1 | 815.0 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 23,298 | 28.8 | 650.7 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 5,295 | 6.6 | 147.9 |
| 4 | Cerebrovascular diseases (I60-I69) | 3,066 | 3.8 | 85.6 |
| 5 | Diabetes mellitus (E10-E14) | 2,607 | 3.2 | 72.8 |
| 6 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,748 | 2.2 | 48.8 |
| 7 | Chronic liver disease and cirrhosis (K70,K73-K74) | 1,318 | 1.6 | 36.8 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1,072 | 1.3 | 29.9 |
| 9 | Influenza and pneumonia (J10-J18) | 1,013 | 1.3 | 28.3 |
| 10 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 899 | 1.1 | 25.1 |
| 11 | Septicemia (A40-A41) | 868 | 1.1 | 24.2 |
| 12 | Aortic aneurysm and dissection (I71) | 856 | 1.1 | 23.9 |
| 13 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 483 | 0.6 | 13.5 |
| 14 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 364 | 0.5 | 10.2 |
| 15 | Alzheimer's disease (G30) | 338 | 0.4 | 9.4 |
| ... | All other causes (Residual) | 8,397 | 10.4 | 234.5 |
| White, non-Hispanic, male, 70-74 years | | | | |
| ... | All causes | 115,353 | 100.0 | 3,543.1 |
| 1 | Malignant neoplasms (C00-C97) | 38,371 | 33.3 | 1,178.6 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 33,761 | 29.3 | 1,037.0 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 8,794 | 7.6 | 270.1 |
| 4 | Cerebrovascular diseases (I60-I69) | 5,687 | 4.9 | 174.7 |
| 5 | Diabetes mellitus (E10-E14) | 3,535 | 3.1 | 108.6 |
| 6 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 2,089 | 1.8 | 64.2 |
| 7 | Influenza and pneumonia (J10-J18) | 1,906 | 1.7 | 58.5 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1,657 | 1.4 | 50.9 |
| 9 | Septicemia (A40-A41) | 1,355 | 1.2 | 41.6 |
| 10 | Chronic liver disease and cirrhosis (K70,K73-K74) | 1,288 | 1.1 | 39.6 |
| 11 | Aortic aneurysm and dissection (I71) | 1,253 | 1.1 | 38.5 |
| 12 | Alzheimer's disease (G30) | 967 | 0.8 | 29.7 |
| 13 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 937 | 0.8 | 28.8 |
| 14 | Parkinson's disease (G20-G21) | 933 | 0.8 | 28.7 |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 767 | 0.7 | 23.6 |
| ... | All other causes (Residual) | 12,053 | 10.4 | 370.2 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|--|---|---------|-------------------------|---------|
| White, non-Hispanic, male, 75-79 years | | | | |
| ... | All causes | 145,725 | 100.0 | 5,580.2 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 45,093 | 30.9 | 1,726.7 |
| 2 | Malignant neoplasms (C00-C97) | 40,816 | 28.0 | 1,562.9 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 11,220 | 7.7 | 429.6 |
| 4 | Cerebrovascular diseases (I60-I69) | 8,818 | 6.1 | 337.7 |
| 5 | Diabetes mellitus (E10-E14) | 4,063 | 2.8 | 155.6 |
| 6 | Influenza and pneumonia (J10-J18) | 3,268 | 2.2 | 125.1 |
| 7 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 2,838 | 1.9 | 108.7 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 2,454 | 1.7 | 94.0 |
| 9 | Alzheimer's disease (G30) | 2,329 | 1.6 | 89.2 |
| 10 | Parkinson's disease (G20-G21) | 1,950 | 1.3 | 74.7 |
| 11 | Septicemia (A40-A41) | 1,892 | 1.3 | 72.4 |
| 12 | Aortic aneurysm and dissection (I71) | 1,534 | 1.1 | 58.7 |
| 13 | Pneumonitis due to solids and liquids (J69) | 1,079 | 0.7 | 41.3 |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 1,078 | 0.7 | 41.3 |
| 15 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 966 | 0.7 | 37.0 |
| ... | All other causes (Residual) | 16,327 | 11.2 | 625.2 |
| White, non-Hispanic, male, 80-84 years | | | | |
| ... | All causes | 148,288 | 100.0 | 8,828.0 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 48,903 | 33.0 | 2,911.3 |
| 2 | Malignant neoplasms (C00-C97) | 33,069 | 22.3 | 1,968.7 |
| 3 | Cerebrovascular diseases (I60-I69) | 10,944 | 7.4 | 651.5 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 10,276 | 6.9 | 611.8 |
| 5 | Influenza and pneumonia (J10-J18) | 4,562 | 3.1 | 271.6 |
| 6 | Diabetes mellitus (E10-E14) | 3,811 | 2.6 | 226.9 |
| 7 | Alzheimer's disease (G30) | 3,808 | 2.6 | 226.7 |
| 8 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 2,962 | 2.0 | 176.3 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 2,856 | 1.9 | 170.0 |
| 10 | Parkinson's disease (G20-G21) | 2,536 | 1.7 | 151.0 |
| 11 | Septicemia (A40-A41) | 1,959 | 1.3 | 116.6 |
| 12 | Pneumonitis due to solids and liquids (J69) | 1,686 | 1.1 | 100.4 |
| 13 | Aortic aneurysm and dissection (I71) | 1,468 | 1.0 | 87.4 |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 1,074 | 0.7 | 63.9 |
| 15 | Atherosclerosis (I70) | 914 | 0.6 | 54.4 |
| ... | All other causes (Residual) | 17,460 | 11.8 | 1,039.4 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|--|---|---------|-------------------------|------|
| White, non-Hispanic, male, 85-89 years | | | | |
| ... | All causes | 115,799 | 100.0 | --- |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 40,491 | 35.0 | --- |
| 2 | Malignant neoplasms (C00-C97) | 20,248 | 17.5 | --- |
| 3 | Cerebrovascular diseases (I60-I69) | 9,439 | 8.2 | --- |
| 4 | Chronic lower respiratory diseases (J40-J47) | 7,101 | 6.1 | --- |
| 5 | Influenza and pneumonia (J10-J18) | 4,746 | 4.1 | --- |
| 6 | Alzheimer's disease (G30) | 3,868 | 3.3 | --- |
| 7 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 2,598 | 2.2 | --- |
| 8 | Diabetes mellitus (E10-E14) | 2,515 | 2.2 | --- |
| 9 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 2,482 | 2.1 | --- |
| 10 | Parkinson's disease (G20-G21) | 1,881 | 1.6 | --- |
| 11 | Pneumonitis due to solids and liquids (J69) | 1,688 | 1.5 | --- |
| 12 | Septicemia (A40-A41) | 1,492 | 1.3 | --- |
| 13 | Atherosclerosis (I70) | 1,025 | 0.9 | --- |
| 14 | Aortic aneurysm and dissection (I71) | 927 | 0.8 | --- |
| 15 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 917 | 0.8 | --- |
| ... | All other causes (Residual) | 14,381 | 12.4 | --- |
| White, non-Hispanic, male, 90-94 years | | | | |
| ... | All causes | 57,884 | 100.0 | --- |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 21,898 | 37.8 | --- |
| 2 | Malignant neoplasms (C00-C97) | 7,441 | 12.9 | --- |
| 3 | Cerebrovascular diseases (I60-I69) | 4,851 | 8.4 | --- |
| 4 | Influenza and pneumonia (J10-J18) | 3,119 | 5.4 | --- |
| 5 | Chronic lower respiratory diseases (J40-J47) | 2,916 | 5.0 | --- |
| 6 | Alzheimer's disease (G30) | 2,125 | 3.7 | --- |
| 7 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1,399 | 2.4 | --- |
| 8 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,348 | 2.3 | --- |
| 9 | Pneumonitis due to solids and liquids (J69) | 1,048 | 1.8 | --- |
| 10 | Diabetes mellitus (E10-E14) | 990 | 1.7 | --- |
| 11 | Septicemia (A40-A41) | 712 | 1.2 | --- |
| 12 | Parkinson's disease (G20-G21) | 709 | 1.2 | --- |
| 13 | Atherosclerosis (I70) | 652 | 1.1 | --- |
| 14 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 508 | 0.9 | --- |
| 15 | Aortic aneurysm and dissection (I71) | 379 | 0.7 | --- |
| ... | All other causes (Residual) | 7,789 | 13.5 | --- |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|---|---|--------|-------------------------|------|
| White, non-Hispanic, male, 95-99 years | | | | |
| ... | All causes | 16,125 | 100.0 | --- |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 6,442 | 40.0 | --- |
| 2 | Malignant neoplasms (C00-C97) | 1,599 | 9.9 | --- |
| 3 | Cerebrovascular diseases (I60-I69) | 1,294 | 8.0 | --- |
| 4 | Influenza and pneumonia (J10-J18) | 1,089 | 6.8 | --- |
| 5 | Chronic lower respiratory diseases (J40-J47) | 629 | 3.9 | --- |
| 6 | Alzheimer's disease (G30) | 557 | 3.5 | --- |
| 7 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 372 | 2.3 | --- |
| 8 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 339 | 2.1 | --- |
| 9 | Pneumonitis due to solids and liquids (J69) | 333 | 2.1 | --- |
| 10 | Atherosclerosis (I70) | 256 | 1.6 | --- |
| 11 | Diabetes mellitus (E10-E14) | 210 | 1.3 | --- |
| 12 | Septicemia (A40-A41) | 184 | 1.1 | --- |
| 13 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 155 | 1.0 | --- |
| 14 | Parkinson's disease (G20-G21) | 123 | 0.8 | --- |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 91 | 0.6 | --- |
| ... | All other causes (Residual) | 2,452 | 15.2 | --- |
| White, non-Hispanic, male, 100 years and over | | | | |
| ... | All causes | 2,276 | 100.0 | --- |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 944 | 41.5 | --- |
| 2 | Influenza and pneumonia (J10-J18) | 212 | 9.3 | --- |
| 3 | Malignant neoplasms (C00-C97) | 183 | 8.0 | --- |
| 4 | Cerebrovascular diseases (I60-I69) | 172 | 7.6 | --- |
| 5 | Chronic lower respiratory diseases (J40-J47) | 70 | 3.1 | --- |
| 6 | Alzheimer's disease (G30) | 63 | 2.8 | --- |
| 7 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 53 | 2.3 | --- |
| 8 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 51 | 2.2 | --- |
| 9 | Pneumonitis due to solids and liquids (J69) | 50 | 2.2 | --- |
| 10 | Atherosclerosis (I70) | 40 | 1.8 | --- |
| 11 | Diabetes mellitus (E10-E14) | 25 | 1.1 | --- |
| 12 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 20 | 0.9 | --- |
| 13 | Septicemia (A40-A41) | 18 | 0.8 | --- |
| 14 | Anemias (D50-D64) | 10 | 0.4 | --- |
| 15 | Parkinson's disease (G20-G21) | 9 | 0.4 | --- |
| ... | All other causes (Residual) | 356 | 15.6 | --- |

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|--|---|-----------|-------------------------|---------|
| White, non-Hispanic, female, all ages | | | | |
| ... | All causes | 1,016,843 | 100.0 | 1,006.1 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 301,636 | 29.7 | 298.4 |
| 2 | Malignant neoplasms (C00-C97) | 220,705 | 21.7 | 218.4 |
| 3 | Cerebrovascular diseases (I60-I69) | 83,497 | 8.2 | 82.6 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 57,801 | 5.7 | 57.2 |
| 5 | Alzheimer's disease (G30) | 34,735 | 3.4 | 34.4 |
| 6 | Influenza and pneumonia (J10-J18) | 29,563 | 2.9 | 29.2 |
| 7 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 28,082 | 2.8 | 27.8 |
| 8 | Diabetes mellitus (E10-E14) | 27,180 | 2.7 | 26.9 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 15,118 | 1.5 | 15.0 |
| 10 | Septicemia (A40-A41) | 13,721 | 1.3 | 13.6 |
| 11 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 8,867 | 0.9 | 8.8 |
| 12 | Atherosclerosis (I70) | 8,006 | 0.8 | 7.9 |
| 13 | Chronic liver disease and cirrhosis (K70,K73-K74) | 7,456 | 0.7 | 7.4 |
| 14 | Pneumonitis due to solids and liquids (J69) | 7,432 | 0.7 | 7.4 |
| 15 | Parkinson's disease (G20-G21) | 6,554 | 0.6 | 6.5 |
| ... | All other causes (Residual) | 166,490 | 16.4 | 164.7 |
| White, non-Hispanic, female, 1-4 years | | | | |
| ... | All causes | 1,180 | 100.0 | 26.3 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 382 | 32.4 | 8.5 |
| 2 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 143 | 12.1 | 3.2 |
| 3 | Malignant neoplasms (C00-C97) | 124 | 10.5 | 2.8 |
| 4 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 65 | 5.5 | 1.5 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 49 | 4.2 | 1.1 |
| 6 | Septicemia (A40-A41) | 22 | 1.9 | 0.5 |
| 6 | Influenza and pneumonia (J10-J18) | 22 | 1.9 | 0.5 |
| 8 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 18 | 1.5 | * |
| 9 | Cerebrovascular diseases (I60-I69) | 14 | 1.2 | * |
| 10 | Certain conditions originating in the perinatal period (P00-P96) | 12 | 1.0 | * |
| 11 | Acute bronchitis and bronchiolitis (J20-J21) | 7 | 0.6 | * |
| 11 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 7 | 0.6 | * |
| 13 | Pneumonitis due to solids and liquids (J69) | 6 | 0.5 | * |
| 14 | Meningococcal infection (A39) | 5 | 0.4 | * |
| 14 | Anemias (D50-D64) | 5 | 0.4 | * |
| 14 | Meningitis (G00,G03) | 5 | 0.4 | * |
| ... | All other causes (Residual) | 294 | 24.9 | 6.6 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|------|
| White, non-Hispanic, female, 5-9 years | | | | |
| ... | All causes | 820 | 100.0 | 13.7 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 298 | 36.3 | 5.0 |
| 2 | Malignant neoplasms (C00-C97) | 156 | 19.0 | 2.6 |
| 3 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 47 | 5.7 | 0.8 |
| 4 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 42 | 5.1 | 0.7 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 29 | 3.5 | 0.5 |
| 6 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 14 | 1.7 | * |
| 6 | Influenza and pneumonia (J10-J18) | 14 | 1.7 | * |
| 8 | Septicemia (A40-A41) | 10 | 1.2 | * |
| 9 | Cerebrovascular diseases (I60-I69) | 9 | 1.1 | * |
| 10 | Chronic lower respiratory diseases (J40-J47) | 7 | 0.9 | * |
| 11 | Anemias (D50-D64) | 5 | 0.6 | * |
| 12 | Meningococcal infection (A39) | 4 | 0.5 | * |
| 12 | Meningitis (G00,G03) | 4 | 0.5 | * |
| 14 | Human immunodeficiency virus (HIV) disease (B20-B24) | 2 | 0.2 | * |
| 14 | Diabetes mellitus (E10-E14) | 2 | 0.2 | * |
| 14 | Certain conditions originating in the perinatal period (P00-P96) | 2 | 0.2 | * |
| 14 | Complications of medical and surgical care (Y40-Y84,Y88) | 2 | 0.2 | * |
| ... | All other causes (Residual) | 173 | 21.1 | 2.9 |
| White, non-Hispanic, female, 10-14 years | | | | |
| ... | All causes | 903 | 100.0 | 14.1 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 324 | 35.9 | 5.1 |
| 2 | Malignant neoplasms (C00-C97) | 141 | 15.6 | 2.2 |
| 3 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 44 | 4.9 | 0.7 |
| 4 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 43 | 4.8 | 0.7 |
| 5 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 39 | 4.3 | 0.6 |
| 5 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 39 | 4.3 | 0.6 |
| 7 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 17 | 1.9 | * |
| 8 | Influenza and pneumonia (J10-J18) | 15 | 1.7 | * |
| 9 | Chronic lower respiratory diseases (J40-J47) | 10 | 1.1 | * |
| 10 | Cerebrovascular diseases (I60-I69) | 9 | 1.0 | * |
| 11 | Septicemia (A40-A41) | 7 | 0.8 | * |
| 12 | Anemias (D50-D64) | 6 | 0.7 | * |
| 13 | Meningococcal infection (A39) | 4 | 0.4 | * |
| 13 | Diabetes mellitus (E10-E14) | 4 | 0.4 | * |
| 15 | Meningitis (G00,G03) | 3 | 0.3 | * |
| ... | All other causes (Residual) | 198 | 21.9 | 3.1 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|------|
| White, non-Hispanic, female, 15-19 years | | | | |
| ... | All causes | 2,508 | 100.0 | 39.6 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,426 | 56.9 | 22.5 |
| 2 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 187 | 7.5 | 3.0 |
| 3 | Malignant neoplasms (C00-C97) | 178 | 7.1 | 2.8 |
| 4 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 119 | 4.7 | 1.9 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 69 | 2.8 | 1.1 |
| 6 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 57 | 2.3 | 0.9 |
| 7 | Influenza and pneumonia (J10-J18) | 21 | 0.8 | 0.3 |
| 8 | Cerebrovascular diseases (I60-I69) | 20 | 0.8 | 0.3 |
| 9 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 19 | 0.8 | * |
| 10 | Chronic lower respiratory diseases (J40-J47) | 18 | 0.7 | * |
| 11 | Meningococcal infection (A39) | 16 | 0.6 | * |
| 12 | Septicemia (A40-A41) | 14 | 0.6 | * |
| 13 | Pregnancy, childbirth and the puerperium (O00-099) | 13 | 0.5 | * |
| 14 | Meningitis (G00,G03) | 7 | 0.3 | * |
| 15 | Diabetes mellitus (E10-E14) | 6 | 0.2 | * |
| ... | All other causes (Residual) | 338 | 13.5 | 5.3 |
| White, non-Hispanic, female, 20-24 years | | | | |
| ... | All causes | 2,680 | 100.0 | 44.3 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,144 | 42.7 | 18.9 |
| 2 | Malignant neoplasms (C00-C97) | 266 | 9.9 | 4.4 |
| 3 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 217 | 8.1 | 3.6 |
| 4 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 210 | 7.8 | 3.5 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 117 | 4.4 | 1.9 |
| 6 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 62 | 2.3 | 1.0 |
| 7 | Pregnancy, childbirth and the puerperium (O00-099) | 25 | 0.9 | 0.4 |
| 8 | Cerebrovascular diseases (I60-I69) | 23 | 0.9 | 0.4 |
| 8 | Influenza and pneumonia (J10-J18) | 23 | 0.9 | 0.4 |
| 10 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 20 | 0.7 | 0.3 |
| 11 | Septicemia (A40-A41) | 18 | 0.7 | * |
| 11 | Diabetes mellitus (E10-E14) | 18 | 0.7 | * |
| 11 | Chronic lower respiratory diseases (J40-J47) | 18 | 0.7 | * |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 16 | 0.6 | * |
| 15 | Human immunodeficiency virus (HIV) disease (B20-B24) | 13 | 0.5 | * |
| ... | All other causes (Residual) | 490 | 18.3 | 8.1 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|------|
| White, non-Hispanic, female, 25-29 years | | | | |
| ... | All causes | 2,962 | 100.0 | 51.4 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 911 | 30.8 | 15.8 |
| 2 | Malignant neoplasms (C00-C97) | 416 | 14.0 | 7.2 |
| 3 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 276 | 9.3 | 4.8 |
| 4 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 208 | 7.0 | 3.6 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 196 | 6.6 | 3.4 |
| 6 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 65 | 2.2 | 1.1 |
| 7 | Cerebrovascular diseases (I60-I69) | 52 | 1.8 | 0.9 |
| 8 | Diabetes mellitus (E10-E14) | 51 | 1.7 | 0.9 |
| 9 | Influenza and pneumonia (J10-J18) | 39 | 1.3 | 0.7 |
| 10 | Human immunodeficiency virus (HIV) disease (B20-B24) | 31 | 1.0 | 0.5 |
| 10 | Pregnancy, childbirth and the puerperium (O00-099) | 31 | 1.0 | 0.5 |
| 12 | Septicemia (A40-A41) | 24 | 0.8 | 0.4 |
| 13 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 20 | 0.7 | 0.3 |
| 13 | Chronic lower respiratory diseases (J40-J47) | 20 | 0.7 | 0.3 |
| 15 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 17 | 0.6 | * |
| ... | All other causes (Residual) | 605 | 20.4 | 10.5 |
| White, non-Hispanic, female, 30-34 years | | | | |
| ... | All causes | 4,616 | 100.0 | 69.0 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 981 | 21.3 | 14.7 |
| 2 | Malignant neoplasms (C00-C97) | 918 | 19.9 | 13.7 |
| 3 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 405 | 8.8 | 6.1 |
| 4 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 395 | 8.6 | 5.9 |
| 5 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 257 | 5.6 | 3.8 |
| 6 | Cerebrovascular diseases (I60-I69) | 107 | 2.3 | 1.6 |
| 7 | Diabetes mellitus (E10-E14) | 100 | 2.2 | 1.5 |
| 8 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 79 | 1.7 | 1.2 |
| 9 | Human immunodeficiency virus (HIV) disease (B20-B24) | 78 | 1.7 | 1.2 |
| 10 | Chronic liver disease and cirrhosis (K70,K73-K74) | 70 | 1.5 | 1.0 |
| 11 | Influenza and pneumonia (J10-J18) | 56 | 1.2 | 0.8 |
| 12 | Septicemia (A40-A41) | 52 | 1.1 | 0.8 |
| 12 | Chronic lower respiratory diseases (J40-J47) | 52 | 1.1 | 0.8 |
| 14 | Pregnancy, childbirth and the puerperium (O00-099) | 46 | 1.0 | 0.7 |
| 15 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 38 | 0.8 | 0.6 |
| ... | All other causes (Residual) | 982 | 21.3 | 14.7 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|-------|
| White, non-Hispanic, female, 35-39 years | | | | |
| ... | All causes | 8,232 | 100.0 | 108.2 |
| 1 | Malignant neoplasms (C00-C97) | 2,107 | 25.6 | 27.7 |
| 2 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,520 | 18.5 | 20.0 |
| 3 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 819 | 9.9 | 10.8 |
| 4 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 575 | 7.0 | 7.6 |
| 5 | Chronic liver disease and cirrhosis (K70,K73-K74) | 290 | 3.5 | 3.8 |
| 6 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 278 | 3.4 | 3.7 |
| 7 | Cerebrovascular diseases (I60-I69) | 229 | 2.8 | 3.0 |
| 8 | Diabetes mellitus (E10-E14) | 180 | 2.2 | 2.4 |
| 9 | Human immunodeficiency virus (HIV) disease (B20-B24) | 133 | 1.6 | 1.7 |
| 10 | Influenza and pneumonia (J10-J18) | 112 | 1.4 | 1.5 |
| 11 | Chronic lower respiratory diseases (J40-J47) | 90 | 1.1 | 1.2 |
| 12 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 85 | 1.0 | 1.1 |
| 13 | Septicemia (A40-A41) | 67 | 0.8 | 0.9 |
| 14 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 53 | 0.6 | 0.7 |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 52 | 0.6 | 0.7 |
| ... | All other causes (Residual) | 1,642 | 19.9 | 21.6 |
| White, non-Hispanic, female, 40-44 years | | | | |
| ... | All causes | 13,101 | 100.0 | 159.7 |
| 1 | Malignant neoplasms (C00-C97) | 4,284 | 32.7 | 52.2 |
| 2 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,632 | 12.5 | 19.9 |
| 3 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 1,520 | 11.6 | 18.5 |
| 4 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 700 | 5.3 | 8.5 |
| 5 | Chronic liver disease and cirrhosis (K70,K73-K74) | 519 | 4.0 | 6.3 |
| 6 | Cerebrovascular diseases (I60-I69) | 420 | 3.2 | 5.1 |
| 7 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 305 | 2.3 | 3.7 |
| 8 | Diabetes mellitus (E10-E14) | 286 | 2.2 | 3.5 |
| 9 | Chronic lower respiratory diseases (J40-J47) | 230 | 1.8 | 2.8 |
| 10 | Influenza and pneumonia (J10-J18) | 160 | 1.2 | 2.0 |
| 11 | Human immunodeficiency virus (HIV) disease (B20-B24) | 139 | 1.1 | 1.7 |
| 12 | Septicemia (A40-A41) | 125 | 1.0 | 1.5 |
| 13 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 118 | 0.9 | 1.4 |
| 14 | Viral hepatitis (B15-B19) | 111 | 0.8 | 1.4 |
| 15 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 95 | 0.7 | 1.2 |
| ... | All other causes (Residual) | 2,457 | 18.8 | 29.9 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|-------|
| White, non-Hispanic, female, 45-49 years | | | | |
| ... | All causes | 18,106 | 100.0 | 233.1 |
| 1 | Malignant neoplasms (C00-C97) | 7,089 | 39.2 | 91.3 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 2,372 | 13.1 | 30.5 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,420 | 7.8 | 18.3 |
| 4 | Chronic liver disease and cirrhosis (K70,K73-K74) | 698 | 3.9 | 9.0 |
| 5 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 682 | 3.8 | 8.8 |
| 6 | Cerebrovascular diseases (I60-I69) | 624 | 3.4 | 8.0 |
| 7 | Diabetes mellitus (E10-E14) | 489 | 2.7 | 6.3 |
| 8 | Chronic lower respiratory diseases (J40-J47) | 375 | 2.1 | 4.8 |
| 9 | Septicemia (A40-A41) | 228 | 1.3 | 2.9 |
| 10 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 222 | 1.2 | 2.9 |
| 11 | Influenza and pneumonia (J10-J18) | 205 | 1.1 | 2.6 |
| 12 | Viral hepatitis (B15-B19) | 172 | 0.9 | 2.2 |
| 13 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 143 | 0.8 | 1.8 |
| 14 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 113 | 0.6 | 1.5 |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 100 | 0.6 | 1.3 |
| ... | All other causes (Residual) | 3,174 | 17.5 | 40.9 |
| White, non-Hispanic, female, 50-54 years | | | | |
| ... | All causes | 25,249 | 100.0 | 353.9 |
| 1 | Malignant neoplasms (C00-C97) | 11,068 | 43.8 | 155.1 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 3,994 | 15.8 | 56.0 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,250 | 5.0 | 17.5 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 864 | 3.4 | 12.1 |
| 5 | Cerebrovascular diseases (I60-I69) | 831 | 3.3 | 11.6 |
| 6 | Diabetes mellitus (E10-E14) | 829 | 3.3 | 11.6 |
| 7 | Chronic liver disease and cirrhosis (K70,K73-K74) | 692 | 2.7 | 9.7 |
| 8 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 594 | 2.4 | 8.3 |
| 9 | Septicemia (A40-A41) | 338 | 1.3 | 4.7 |
| 10 | Influenza and pneumonia (J10-J18) | 281 | 1.1 | 3.9 |
| 11 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 218 | 0.9 | 3.1 |
| 12 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 161 | 0.6 | 2.3 |
| 13 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 149 | 0.6 | 2.1 |
| 14 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 139 | 0.6 | 1.9 |
| 15 | Viral hepatitis (B15-B19) | 130 | 0.5 | 1.8 |
| ... | All other causes (Residual) | 3,711 | 14.7 | 52.0 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|-------|
| White, non-Hispanic, female, 55-59 years | | | | |
| ... | All causes | 32,293 | 100.0 | 567.1 |
| 1 | Malignant neoplasms (C00-C97) | 14,498 | 44.9 | 254.6 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 5,740 | 17.8 | 100.8 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 1,725 | 5.3 | 30.3 |
| 4 | Cerebrovascular diseases (I60-I69) | 1,185 | 3.7 | 20.8 |
| 5 | Diabetes mellitus (E10-E14) | 1,079 | 3.3 | 18.9 |
| 6 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 990 | 3.1 | 17.4 |
| 7 | Chronic liver disease and cirrhosis (K70,K73-K74) | 616 | 1.9 | 10.8 |
| 8 | Septicemia (A40-A41) | 449 | 1.4 | 7.9 |
| 9 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 412 | 1.3 | 7.2 |
| 10 | Influenza and pneumonia (J10-J18) | 347 | 1.1 | 6.1 |
| 11 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 338 | 1.0 | 5.9 |
| 12 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 217 | 0.7 | 3.8 |
| 13 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 131 | 0.4 | 2.3 |
| 13 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 131 | 0.4 | 2.3 |
| 15 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 116 | 0.4 | 2.0 |
| ... | All other causes (Residual) | 4,319 | 13.4 | 75.9 |
| White, non-Hispanic, female, 60-64 years | | | | |
| ... | All causes | 41,823 | 100.0 | 919.2 |
| 1 | Malignant neoplasms (C00-C97) | 18,007 | 43.1 | 395.8 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 8,189 | 19.6 | 180.0 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 3,068 | 7.3 | 67.4 |
| 4 | Cerebrovascular diseases (I60-I69) | 1,623 | 3.9 | 35.7 |
| 5 | Diabetes mellitus (E10-E14) | 1,515 | 3.6 | 33.3 |
| 6 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 881 | 2.1 | 19.4 |
| 7 | Chronic liver disease and cirrhosis (K70,K73-K74) | 698 | 1.7 | 15.3 |
| 8 | Septicemia (A40-A41) | 561 | 1.3 | 12.3 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 511 | 1.2 | 11.2 |
| 10 | Influenza and pneumonia (J10-J18) | 484 | 1.2 | 10.6 |
| 11 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 283 | 0.7 | 6.2 |
| 12 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 242 | 0.6 | 5.3 |
| 13 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 206 | 0.5 | 4.5 |
| 14 | Aortic aneurysm and dissection (I71) | 186 | 0.4 | 4.1 |
| 15 | Alzheimer's disease (G30) | 179 | 0.4 | 3.9 |
| ... | All other causes (Residual) | 5,190 | 12.4 | 114.1 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|---------|
| White, non-Hispanic, female, 65-69 years | | | | |
| ... | All causes | 58,698 | 100.0 | 1,448.1 |
| 1 | Malignant neoplasms (C00-C97) | 22,934 | 39.1 | 565.8 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 12,716 | 21.7 | 313.7 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 5,233 | 8.9 | 129.1 |
| 4 | Cerebrovascular diseases (I60-I69) | 2,649 | 4.5 | 65.4 |
| 5 | Diabetes mellitus (E10-E14) | 2,129 | 3.6 | 52.5 |
| 6 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,036 | 1.8 | 25.6 |
| 7 | Septicemia (A40-A41) | 836 | 1.4 | 20.6 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 835 | 1.4 | 20.6 |
| 9 | Chronic liver disease and cirrhosis (K70,K73-K74) | 794 | 1.4 | 19.6 |
| 10 | Influenza and pneumonia (J10-J18) | 760 | 1.3 | 18.7 |
| 11 | Alzheimer's disease (G30) | 398 | 0.7 | 9.8 |
| 12 | Aortic aneurysm and dissection (I71) | 372 | 0.6 | 9.2 |
| 13 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 354 | 0.6 | 8.7 |
| 14 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 315 | 0.5 | 7.8 |
| 15 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 186 | 0.3 | 4.6 |
| ... | All other causes (Residual) | 7,151 | 12.2 | 176.4 |
| White, non-Hispanic, female, 70-74 years | | | | |
| ... | All causes | 91,533 | 100.0 | 2,285.2 |
| 1 | Malignant neoplasms (C00-C97) | 31,041 | 33.9 | 775.0 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 22,080 | 24.1 | 551.2 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 8,371 | 9.1 | 209.0 |
| 4 | Cerebrovascular diseases (I60-I69) | 5,436 | 5.9 | 135.7 |
| 5 | Diabetes mellitus (E10-E14) | 3,193 | 3.5 | 79.7 |
| 6 | Influenza and pneumonia (J10-J18) | 1,535 | 1.7 | 38.3 |
| 7 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,436 | 1.6 | 35.9 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1,420 | 1.6 | 35.5 |
| 9 | Septicemia (A40-A41) | 1,326 | 1.4 | 33.1 |
| 10 | Alzheimer's disease (G30) | 1,317 | 1.4 | 32.9 |
| 11 | Chronic liver disease and cirrhosis (K70,K73-K74) | 907 | 1.0 | 22.6 |
| 12 | Aortic aneurysm and dissection (I71) | 598 | 0.7 | 14.9 |
| 13 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 581 | 0.6 | 14.5 |
| 14 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 559 | 0.6 | 14.0 |
| 15 | Parkinson's disease (G20-G21) | 495 | 0.5 | 12.4 |
| ... | All other causes (Residual) | 11,238 | 12.3 | 280.6 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|--|---|---------|-------------------------|---------|
| White, non-Hispanic, female, 75-79 years | | | | |
| ... | All causes | 136,293 | 100.0 | 3,695.5 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 36,967 | 27.1 | 1,002.4 |
| 2 | Malignant neoplasms (C00-C97) | 36,210 | 26.6 | 981.8 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 11,183 | 8.2 | 303.2 |
| 4 | Cerebrovascular diseases (I60-I69) | 10,735 | 7.9 | 291.1 |
| 5 | Diabetes mellitus (E10-E14) | 4,522 | 3.3 | 122.6 |
| 6 | Alzheimer's disease (G30) | 3,500 | 2.6 | 94.9 |
| 7 | Influenza and pneumonia (J10-J18) | 2,920 | 2.1 | 79.2 |
| 8 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 2,322 | 1.7 | 63.0 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 2,232 | 1.6 | 60.5 |
| 10 | Septicemia (A40-A41) | 2,036 | 1.5 | 55.2 |
| 11 | Parkinson's disease (G20-G21) | 1,222 | 0.9 | 33.1 |
| 12 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 1,104 | 0.8 | 29.9 |
| 13 | Aortic aneurysm and dissection (I71) | 1,027 | 0.8 | 27.8 |
| 14 | Chronic liver disease and cirrhosis (K70,K73-K74) | 989 | 0.7 | 26.8 |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 867 | 0.6 | 23.5 |
| ... | All other causes (Residual) | 18,457 | 13.5 | 500.5 |
| White, non-Hispanic, female, 80-84 years | | | | |
| ... | All causes | 172,837 | 100.0 | 6,211.0 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 54,573 | 31.6 | 1,961.1 |
| 2 | Malignant neoplasms (C00-C97) | 32,665 | 18.9 | 1,173.8 |
| 3 | Cerebrovascular diseases (I60-I69) | 16,807 | 9.7 | 604.0 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 11,293 | 6.5 | 405.8 |
| 5 | Alzheimer's disease (G30) | 7,105 | 4.1 | 255.3 |
| 6 | Diabetes mellitus (E10-E14) | 5,111 | 3.0 | 183.7 |
| 7 | Influenza and pneumonia (J10-J18) | 5,087 | 2.9 | 182.8 |
| 8 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 3,035 | 1.8 | 109.1 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 2,856 | 1.7 | 102.6 |
| 10 | Septicemia (A40-A41) | 2,465 | 1.4 | 88.6 |
| 11 | Parkinson's disease (G20-G21) | 1,813 | 1.0 | 65.2 |
| 12 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 1,694 | 1.0 | 60.9 |
| 13 | Pneumonitis due to solids and liquids (J69) | 1,393 | 0.8 | 50.1 |
| 14 | Atherosclerosis (I70) | 1,353 | 0.8 | 48.6 |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 1,131 | 0.7 | 40.6 |
| ... | All other causes (Residual) | 24,456 | 14.1 | 878.8 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|--|---|---------|-------------------------|------|
| White, non-Hispanic, female, 85-89 years | | | | |
| ... | All causes | 185,211 | 100.0 | --- |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 65,842 | 35.5 | --- |
| 2 | Malignant neoplasms (C00-C97) | 23,491 | 12.7 | --- |
| 3 | Cerebrovascular diseases (I60-I69) | 20,090 | 10.8 | --- |
| 4 | Alzheimer's disease (G30) | 9,751 | 5.3 | --- |
| 5 | Chronic lower respiratory diseases (J40-J47) | 9,094 | 4.9 | --- |
| 6 | Influenza and pneumonia (J10-J18) | 7,075 | 3.8 | --- |
| 7 | Diabetes mellitus (E10-E14) | 4,227 | 2.3 | --- |
| 8 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 3,229 | 1.7 | --- |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 3,019 | 1.6 | --- |
| 10 | Septicemia (A40-A41) | 2,548 | 1.4 | --- |
| 11 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 2,005 | 1.1 | --- |
| 12 | Atherosclerosis (I70) | 1,901 | 1.0 | --- |
| 13 | Pneumonitis due to solids and liquids (J69) | 1,815 | 1.0 | --- |
| 14 | Parkinson's disease (G20-G21) | 1,647 | 0.9 | --- |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 1,101 | 0.6 | --- |
| ... | All other causes (Residual) | 28,376 | 15.3 | --- |
| White, non-Hispanic, female, 90-94 years | | | | |
| ... | All causes | 139,185 | 100.0 | --- |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 54,877 | 39.4 | --- |
| 2 | Cerebrovascular diseases (I60-I69) | 15,350 | 11.0 | --- |
| 3 | Malignant neoplasms (C00-C97) | 11,485 | 8.3 | --- |
| 4 | Alzheimer's disease (G30) | 8,224 | 5.9 | --- |
| 5 | Influenza and pneumonia (J10-J18) | 6,344 | 4.6 | --- |
| 6 | Chronic lower respiratory diseases (J40-J47) | 4,412 | 3.2 | --- |
| 7 | Diabetes mellitus (E10-E14) | 2,477 | 1.8 | --- |
| 8 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 2,381 | 1.7 | --- |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 2,173 | 1.6 | --- |
| 10 | Atherosclerosis (I70) | 1,909 | 1.4 | --- |
| 11 | Septicemia (A40-A41) | 1,787 | 1.3 | --- |
| 12 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 1,648 | 1.2 | --- |
| 13 | Pneumonitis due to solids and liquids (J69) | 1,553 | 1.1 | --- |
| 14 | Parkinson's disease (G20-G21) | 838 | 0.6 | --- |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 654 | 0.5 | --- |
| ... | All other causes (Residual) | 23,073 | 16.6 | --- |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|---|---|--------|-------------------------|------|
| White, non-Hispanic, female, 95-99 years | | | | |
| ... | All causes | 59,580 | 100.0 | --- |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 24,917 | 41.8 | --- |
| 2 | Cerebrovascular diseases (I60-I69) | 6,142 | 10.3 | --- |
| 3 | Alzheimer's disease (G30) | 3,513 | 5.9 | --- |
| 4 | Malignant neoplasms (C00-C97) | 3,212 | 5.4 | --- |
| 5 | Influenza and pneumonia (J10-J18) | 3,153 | 5.3 | --- |
| 6 | Chronic lower respiratory diseases (J40-J47) | 1,479 | 2.5 | --- |
| 7 | Atherosclerosis (I70) | 1,077 | 1.8 | --- |
| 8 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,037 | 1.7 | --- |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 921 | 1.5 | --- |
| 10 | Diabetes mellitus (E10-E14) | 847 | 1.4 | --- |
| 11 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 803 | 1.3 | --- |
| 12 | Pneumonitis due to solids and liquids (J69) | 678 | 1.1 | --- |
| 13 | Septicemia (A40-A41) | 641 | 1.1 | --- |
| 14 | Parkinson's disease (G20-G21) | 269 | 0.5 | --- |
| 15 | Nutritional deficiencies (E40-E64) | 233 | 0.4 | --- |
| ... | All other causes (Residual) | 10,658 | 17.9 | --- |
| White, non-Hispanic, female, 100 years and over | | | | |
| ... | All causes | 13,209 | 100.0 | --- |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 6,019 | 45.6 | --- |
| 2 | Cerebrovascular diseases (I60-I69) | 1,123 | 8.5 | --- |
| 3 | Influenza and pneumonia (J10-J18) | 864 | 6.5 | --- |
| 4 | Alzheimer's disease (G30) | 619 | 4.7 | --- |
| 5 | Malignant neoplasms (C00-C97) | 394 | 3.0 | --- |
| 6 | Atherosclerosis (I70) | 326 | 2.5 | --- |
| 7 | Chronic lower respiratory diseases (J40-J47) | 249 | 1.9 | --- |
| 8 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 225 | 1.7 | --- |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 188 | 1.4 | --- |
| 10 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 186 | 1.4 | --- |
| 11 | Pneumonitis due to solids and liquids (J69) | 164 | 1.2 | --- |
| 12 | Septicemia (A40-A41) | 116 | 0.9 | --- |
| 13 | Diabetes mellitus (E10-E14) | 111 | 0.8 | --- |
| 14 | Nutritional deficiencies (E40-E64) | 74 | 0.6 | --- |
| 15 | Anemias (D50-D64) | 56 | 0.4 | --- |
| ... | All other causes (Residual) | 2,495 | 18.9 | --- |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|--|---|---------|-------------------------|-------|
| Black, non-Hispanic, both sexes, all ages | | | | |
| ... | All causes | 284,343 | 100.0 | 798.1 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 76,794 | 27.0 | 215.5 |
| 2 | Malignant neoplasms (C00-C97) | 61,580 | 21.7 | 172.8 |
| 3 | Cerebrovascular diseases (I60-I69) | 18,844 | 6.6 | 52.9 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 12,239 | 4.3 | 34.4 |
| 5 | Diabetes mellitus (E10-E14) | 12,167 | 4.3 | 34.1 |
| 6 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 8,077 | 2.8 | 22.7 |
| 7 | Human immunodeficiency virus (HIV) disease (B20-B24) | 7,710 | 2.7 | 21.6 |
| 8 | Chronic lower respiratory diseases (J40-J47) | 7,504 | 2.6 | 21.1 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 7,214 | 2.5 | 20.2 |
| 10 | Septicemia (A40-A41) | 5,829 | 2.0 | 16.4 |
| 11 | Influenza and pneumonia (J10-J18) | 5,686 | 2.0 | 16.0 |
| 12 | Certain conditions originating in the perinatal period (P00-P96) | 4,713 | 1.7 | 13.2 |
| 13 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 4,012 | 1.4 | 11.3 |
| 14 | Alzheimer's disease (G30) | 3,104 | 1.1 | 8.7 |
| 15 | Chronic liver disease and cirrhosis (K70,K73-K74) | 2,733 | 1.0 | 7.7 |
| ... | All other causes (Residual) | 46,137 | 16.2 | 129.5 |
| Black, non-Hispanic, both sexes, 1-4 years | | | | |
| ... | All causes | 1,156 | 100.0 | 49.5 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 331 | 28.6 | 14.2 |
| 2 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 166 | 14.4 | 7.1 |
| 3 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 127 | 11.0 | 5.4 |
| 4 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 73 | 6.3 | 3.1 |
| 5 | Malignant neoplasms (C00-C97) | 46 | 4.0 | 2.0 |
| 6 | Influenza and pneumonia (J10-J18) | 24 | 2.1 | 1.0 |
| 7 | Chronic lower respiratory diseases (J40-J47) | 22 | 1.9 | 0.9 |
| 7 | Certain conditions originating in the perinatal period (P00-P96) | 22 | 1.9 | 0.9 |
| 9 | Septicemia (A40-A41) | 21 | 1.8 | 0.9 |
| 10 | Cerebrovascular diseases (I60-I69) | 12 | 1.0 | * |
| 11 | Anemias (D50-D64) | 11 | 1.0 | * |
| 12 | Human immunodeficiency virus (HIV) disease (B20-B24) | 10 | 0.9 | * |
| 13 | Complications of medical and surgical care (Y40-Y84,Y88) | 7 | 0.6 | * |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 5 | 0.4 | * |
| 14 | Meningitis (G00,G03) | 5 | 0.4 | * |
| ... | All other causes (Residual) | 274 | 23.7 | 11.7 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|------|
| Black, non-Hispanic, both sexes, 5-9 years | | | | |
| ... | All causes | 685 | 100.0 | 21.4 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 313 | 45.7 | 9.8 |
| 2 | Malignant neoplasms (C00-C97) | 78 | 11.4 | 2.4 |
| 3 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 32 | 4.7 | 1.0 |
| 4 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 28 | 4.1 | 0.9 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 27 | 3.9 | 0.8 |
| 6 | Chronic lower respiratory diseases (J40-J47) | 23 | 3.4 | 0.7 |
| 7 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 13 | 1.9 | * |
| 7 | Anemias (D50-D64) | 13 | 1.9 | * |
| 9 | Human immunodeficiency virus (HIV) disease (B20-B24) | 9 | 1.3 | * |
| 9 | Influenza and pneumonia (J10-J18) | 9 | 1.3 | * |
| 11 | Cerebrovascular diseases (I60-I69) | 6 | 0.9 | * |
| 11 | Certain conditions originating in the perinatal period (P00-P96) | 6 | 0.9 | * |
| 13 | Meningococcal infection (A39) | 4 | 0.6 | * |
| 13 | Diabetes mellitus (E10-E14) | 4 | 0.6 | * |
| 13 | Meningitis (G00,G03) | 4 | 0.6 | * |
| 13 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 4 | 0.6 | * |
| ... | All other causes (Residual) | 112 | 16.4 | 3.5 |
| Black, non-Hispanic, both sexes, 10-14 years | | | | |
| ... | All causes | 872 | 100.0 | 26.6 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 317 | 36.4 | 9.7 |
| 2 | Malignant neoplasms (C00-C97) | 94 | 10.8 | 2.9 |
| 3 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 64 | 7.3 | 1.9 |
| 4 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 43 | 4.9 | 1.3 |
| 5 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 41 | 4.7 | 1.2 |
| 6 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 38 | 4.4 | 1.2 |
| 7 | Chronic lower respiratory diseases (J40-J47) | 30 | 3.4 | 0.9 |
| 8 | Human immunodeficiency virus (HIV) disease (B20-B24) | 17 | 1.9 | * |
| 9 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 13 | 1.5 | * |
| 9 | Anemias (D50-D64) | 13 | 1.5 | * |
| 11 | Diabetes mellitus (E10-E14) | 11 | 1.3 | * |
| 12 | Septicemia (A40-A41) | 8 | 0.9 | * |
| 12 | Cerebrovascular diseases (I60-I69) | 8 | 0.9 | * |
| 12 | Influenza and pneumonia (J10-J18) | 8 | 0.9 | * |
| 15 | Certain conditions originating in the perinatal period (P00-P96) | 5 | 0.6 | * |
| ... | All other causes (Residual) | 162 | 18.6 | 4.9 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|-------|
| Black, non-Hispanic, both sexes, 15-19 years | | | | |
| ... | All causes | 2,662 | 100.0 | 88.7 |
| 1 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 1,009 | 37.9 | 33.6 |
| 2 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 760 | 28.5 | 25.3 |
| 3 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 134 | 5.0 | 4.5 |
| 4 | Malignant neoplasms (C00-C97) | 110 | 4.1 | 3.7 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 104 | 3.9 | 3.5 |
| 6 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 61 | 2.3 | 2.0 |
| 7 | Human immunodeficiency virus (HIV) disease (B20-B24) | 34 | 1.3 | 1.1 |
| 8 | Chronic lower respiratory diseases (J40-J47) | 30 | 1.1 | 1.0 |
| 9 | Anemias (D50-D64) | 27 | 1.0 | 0.9 |
| 10 | Septicemia (A40-A41) | 19 | 0.7 | * |
| 11 | Diabetes mellitus (E10-E14) | 17 | 0.6 | * |
| 12 | Influenza and pneumonia (J10-J18) | 16 | 0.6 | * |
| 13 | Pregnancy, childbirth and the puerperium (O00-099) | 14 | 0.5 | * |
| 14 | Cerebrovascular diseases (I60-I69) | 13 | 0.5 | * |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 12 | 0.5 | * |
| ... | All other causes (Residual) | 302 | 11.3 | 10.1 |
| Black, non-Hispanic, both sexes, 20-24 years | | | | |
| ... | All causes | 4,362 | 100.0 | 156.7 |
| 1 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 1,808 | 41.4 | 64.9 |
| 2 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 953 | 21.8 | 34.2 |
| 3 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 291 | 6.7 | 10.5 |
| 4 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 218 | 5.0 | 7.8 |
| 5 | Malignant neoplasms (C00-C97) | 147 | 3.4 | 5.3 |
| 6 | Human immunodeficiency virus (HIV) disease (B20-B24) | 127 | 2.9 | 4.6 |
| 7 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 52 | 1.2 | 1.9 |
| 8 | Chronic lower respiratory diseases (J40-J47) | 47 | 1.1 | 1.7 |
| 9 | Anemias (D50-D64) | 45 | 1.0 | 1.6 |
| 10 | Diabetes mellitus (E10-E14) | 39 | 0.9 | 1.4 |
| 11 | Cerebrovascular diseases (I60-I69) | 38 | 0.9 | 1.4 |
| 12 | Pregnancy, childbirth and the puerperium (O00-099) | 31 | 0.7 | 1.1 |
| 13 | Influenza and pneumonia (J10-J18) | 29 | 0.7 | 1.0 |
| 14 | Septicemia (A40-A41) | 22 | 0.5 | 0.8 |
| 14 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 22 | 0.5 | 0.8 |
| ... | All other causes (Residual) | 493 | 11.3 | 17.7 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|-------|
| Black, non-Hispanic, both sexes, 25-29 years | | | | |
| ... | All causes | 4,388 | 100.0 | 174.9 |
| 1 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 1,370 | 31.2 | 54.6 |
| 2 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 784 | 17.9 | 31.3 |
| 3 | Human immunodeficiency virus (HIV) disease (B20-B24) | 381 | 8.7 | 15.2 |
| 4 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 339 | 7.7 | 13.5 |
| 5 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 243 | 5.5 | 9.7 |
| 6 | Malignant neoplasms (C00-C97) | 241 | 5.5 | 9.6 |
| 7 | Diabetes mellitus (E10-E14) | 83 | 1.9 | 3.3 |
| 8 | Cerebrovascular diseases (I60-I69) | 57 | 1.3 | 2.3 |
| 9 | Anemias (D50-D64) | 50 | 1.1 | 2.0 |
| 10 | Chronic lower respiratory diseases (J40-J47) | 47 | 1.1 | 1.9 |
| 11 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 35 | 0.8 | 1.4 |
| 11 | Pregnancy, childbirth and the puerperium (O00-099) | 35 | 0.8 | 1.4 |
| 11 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 35 | 0.8 | 1.4 |
| 14 | Influenza and pneumonia (J10-J18) | 31 | 0.7 | 1.2 |
| 15 | Septicemia (A40-A41) | 28 | 0.6 | 1.1 |
| ... | All other causes (Residual) | 629 | 14.3 | 25.1 |
| Black, non-Hispanic, both sexes, 30-34 years | | | | |
| ... | All causes | 5,471 | 100.0 | 207.1 |
| 1 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 965 | 17.6 | 36.5 |
| 2 | Human immunodeficiency virus (HIV) disease (B20-B24) | 853 | 15.6 | 32.3 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 816 | 14.9 | 30.9 |
| 4 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 622 | 11.4 | 23.5 |
| 5 | Malignant neoplasms (C00-C97) | 450 | 8.2 | 17.0 |
| 6 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 229 | 4.2 | 8.7 |
| 7 | Diabetes mellitus (E10-E14) | 115 | 2.1 | 4.4 |
| 8 | Cerebrovascular diseases (I60-I69) | 96 | 1.8 | 3.6 |
| 9 | Chronic lower respiratory diseases (J40-J47) | 81 | 1.5 | 3.1 |
| 10 | Septicemia (A40-A41) | 70 | 1.3 | 2.7 |
| 11 | Anemias (D50-D64) | 67 | 1.2 | 2.5 |
| 12 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 58 | 1.1 | 2.2 |
| 13 | Influenza and pneumonia (J10-J18) | 56 | 1.0 | 2.1 |
| 14 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 37 | 0.7 | 1.4 |
| 15 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 32 | 0.6 | 1.2 |
| ... | All other causes (Residual) | 924 | 16.9 | 35.0 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|-------|
| Black, non-Hispanic, both sexes, 35-39 years | | | | |
| ... | All causes | 8,287 | 100.0 | 294.6 |
| 1 | Human immunodeficiency virus (HIV) disease (B20-B24) | 1,417 | 17.1 | 50.4 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 1,162 | 14.0 | 41.3 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,026 | 12.4 | 36.5 |
| 4 | Malignant neoplasms (C00-C97) | 949 | 11.5 | 33.7 |
| 5 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 817 | 9.9 | 29.0 |
| 6 | Cerebrovascular diseases (I60-I69) | 233 | 2.8 | 8.3 |
| 7 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 227 | 2.7 | 8.1 |
| 8 | Diabetes mellitus (E10-E14) | 202 | 2.4 | 7.2 |
| 9 | Chronic lower respiratory diseases (J40-J47) | 131 | 1.6 | 4.7 |
| 10 | Chronic liver disease and cirrhosis (K70,K73-K74) | 117 | 1.4 | 4.2 |
| 11 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 114 | 1.4 | 4.1 |
| 12 | Septicemia (A40-A41) | 109 | 1.3 | 3.9 |
| 12 | Influenza and pneumonia (J10-J18) | 109 | 1.3 | 3.9 |
| 14 | Anemias (D50-D64) | 62 | 0.7 | 2.2 |
| 15 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 53 | 0.6 | 1.9 |
| ... | All other causes (Residual) | 1,559 | 18.8 | 55.4 |
| Black, non-Hispanic, both sexes, 40-44 years | | | | |
| ... | All causes | 12,149 | 100.0 | 438.1 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 2,262 | 18.6 | 81.6 |
| 2 | Malignant neoplasms (C00-C97) | 2,001 | 16.5 | 72.2 |
| 3 | Human immunodeficiency virus (HIV) disease (B20-B24) | 1,655 | 13.6 | 59.7 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,256 | 10.3 | 45.3 |
| 5 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 610 | 5.0 | 22.0 |
| 6 | Cerebrovascular diseases (I60-I69) | 525 | 4.3 | 18.9 |
| 7 | Diabetes mellitus (E10-E14) | 304 | 2.5 | 11.0 |
| 8 | Chronic liver disease and cirrhosis (K70,K73-K74) | 303 | 2.5 | 10.9 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 192 | 1.6 | 6.9 |
| 10 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 191 | 1.6 | 6.9 |
| 11 | Influenza and pneumonia (J10-J18) | 165 | 1.4 | 6.0 |
| 12 | Septicemia (A40-A41) | 159 | 1.3 | 5.7 |
| 13 | Chronic lower respiratory diseases (J40-J47) | 142 | 1.2 | 5.1 |
| 14 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 124 | 1.0 | 4.5 |
| 15 | Viral hepatitis (B15-B19) | 94 | 0.8 | 3.4 |
| ... | All other causes (Residual) | 2,166 | 17.8 | 78.1 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|-------|
| Black, non-Hispanic, both sexes, 45-49 years | | | | |
| ... | All causes | 15,657 | 100.0 | 656.8 |
| 1 | Malignant neoplasms (C00-C97) | 3,370 | 21.5 | 141.4 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 3,345 | 21.4 | 140.3 |
| 3 | Human immunodeficiency virus (HIV) disease (B20-B24) | 1,372 | 8.8 | 57.6 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,212 | 7.7 | 50.8 |
| 5 | Cerebrovascular diseases (I60-I69) | 790 | 5.0 | 33.1 |
| 6 | Diabetes mellitus (E10-E14) | 507 | 3.2 | 21.3 |
| 7 | Chronic liver disease and cirrhosis (K70,K73-K74) | 463 | 3.0 | 19.4 |
| 8 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 433 | 2.8 | 18.2 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 307 | 2.0 | 12.9 |
| 10 | Septicemia (A40-A41) | 261 | 1.7 | 10.9 |
| 11 | Chronic lower respiratory diseases (J40-J47) | 222 | 1.4 | 9.3 |
| 12 | Influenza and pneumonia (J10-J18) | 219 | 1.4 | 9.2 |
| 13 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 188 | 1.2 | 7.9 |
| 14 | Viral hepatitis (B15-B19) | 162 | 1.0 | 6.8 |
| 15 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 152 | 1.0 | 6.4 |
| ... | All other causes (Residual) | 2,654 | 17.0 | 111.3 |
| Black, non-Hispanic, both sexes, 50-54 years | | | | |
| ... | All causes | 18,441 | 100.0 | 946.1 |
| 1 | Malignant neoplasms (C00-C97) | 4,805 | 26.1 | 246.5 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 4,611 | 25.0 | 236.6 |
| 3 | Cerebrovascular diseases (I60-I69) | 1,096 | 5.9 | 56.2 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 891 | 4.8 | 45.7 |
| 5 | Human immunodeficiency virus (HIV) disease (B20-B24) | 877 | 4.8 | 45.0 |
| 6 | Diabetes mellitus (E10-E14) | 799 | 4.3 | 41.0 |
| 7 | Chronic liver disease and cirrhosis (K70,K73-K74) | 530 | 2.9 | 27.2 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 395 | 2.1 | 20.3 |
| 9 | Septicemia (A40-A41) | 334 | 1.8 | 17.1 |
| 10 | Chronic lower respiratory diseases (J40-J47) | 327 | 1.8 | 16.8 |
| 11 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 271 | 1.5 | 13.9 |
| 12 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 251 | 1.4 | 12.9 |
| 13 | Influenza and pneumonia (J10-J18) | 211 | 1.1 | 10.8 |
| 14 | Viral hepatitis (B15-B19) | 187 | 1.0 | 9.6 |
| 15 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 116 | 0.6 | 6.0 |
| ... | All other causes (Residual) | 2,740 | 14.9 | 140.6 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|---------|
| Black, non-Hispanic, both sexes, 55-59 years | | | | |
| ... | All causes | 18,646 | 100.0 | 1,351.9 |
| 1 | Malignant neoplasms (C00-C97) | 5,694 | 30.5 | 412.8 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 5,178 | 27.8 | 375.4 |
| 3 | Cerebrovascular diseases (I60-I69) | 992 | 5.3 | 71.9 |
| 4 | Diabetes mellitus (E10-E14) | 974 | 5.2 | 70.6 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 568 | 3.0 | 41.2 |
| 6 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 459 | 2.5 | 33.3 |
| 7 | Chronic lower respiratory diseases (J40-J47) | 446 | 2.4 | 32.3 |
| 8 | Human immunodeficiency virus (HIV) disease (B20-B24) | 404 | 2.2 | 29.3 |
| 9 | Chronic liver disease and cirrhosis (K70,K73-K74) | 390 | 2.1 | 28.3 |
| 10 | Septicemia (A40-A41) | 381 | 2.0 | 27.6 |
| 11 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 253 | 1.4 | 18.3 |
| 12 | Influenza and pneumonia (J10-J18) | 213 | 1.1 | 15.4 |
| 13 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 138 | 0.7 | 10.0 |
| 14 | Viral hepatitis (B15-B19) | 100 | 0.5 | 7.3 |
| 15 | Aortic aneurysm and dissection (I71) | 86 | 0.5 | 6.2 |
| ... | All other causes (Residual) | 2,370 | 12.7 | 171.8 |
| Black, non-Hispanic, both sexes, 60-64 years | | | | |
| ... | All causes | 20,877 | 100.0 | 1,920.5 |
| 1 | Malignant neoplasms (C00-C97) | 6,502 | 31.1 | 598.1 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 6,031 | 28.9 | 554.8 |
| 3 | Cerebrovascular diseases (I60-I69) | 1,297 | 6.2 | 119.3 |
| 4 | Diabetes mellitus (E10-E14) | 1,159 | 5.6 | 106.6 |
| 5 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 589 | 2.8 | 54.2 |
| 6 | Chronic lower respiratory diseases (J40-J47) | 573 | 2.7 | 52.7 |
| 7 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 477 | 2.3 | 43.9 |
| 8 | Septicemia (A40-A41) | 423 | 2.0 | 38.9 |
| 9 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 300 | 1.4 | 27.6 |
| 10 | Human immunodeficiency virus (HIV) disease (B20-B24) | 286 | 1.4 | 26.3 |
| 11 | Chronic liver disease and cirrhosis (K70,K73-K74) | 259 | 1.2 | 23.8 |
| 12 | Influenza and pneumonia (J10-J18) | 258 | 1.2 | 23.7 |
| 13 | Aortic aneurysm and dissection (I71) | 103 | 0.5 | 9.5 |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 84 | 0.4 | 7.7 |
| 15 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 81 | 0.4 | 7.5 |
| ... | All other causes (Residual) | 2,455 | 11.8 | 225.8 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|---------|
| Black, non-Hispanic, both sexes, 65-69 years | | | | |
| ... | All causes | 25,103 | 100.0 | 2,777.9 |
| 1 | Malignant neoplasms (C00-C97) | 7,877 | 31.4 | 871.7 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 7,418 | 29.6 | 820.9 |
| 3 | Cerebrovascular diseases (I60-I69) | 1,640 | 6.5 | 181.5 |
| 4 | Diabetes mellitus (E10-E14) | 1,440 | 5.7 | 159.4 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 867 | 3.5 | 95.9 |
| 6 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 817 | 3.3 | 90.4 |
| 7 | Septicemia (A40-A41) | 526 | 2.1 | 58.2 |
| 8 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 415 | 1.7 | 45.9 |
| 9 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 372 | 1.5 | 41.2 |
| 10 | Influenza and pneumonia (J10-J18) | 315 | 1.3 | 34.9 |
| 11 | Chronic liver disease and cirrhosis (K70,K73-K74) | 245 | 1.0 | 27.1 |
| 12 | Human immunodeficiency virus (HIV) disease (B20-B24) | 146 | 0.6 | 16.2 |
| 13 | Aortic aneurysm and dissection (I71) | 105 | 0.4 | 11.6 |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 99 | 0.4 | 11.0 |
| 15 | Pneumonitis due to solids and liquids (J69) | 86 | 0.3 | 9.5 |
| ... | All other causes (Residual) | 2,735 | 10.9 | 302.7 |
| Black, non-Hispanic, both sexes, 70-74 years | | | | |
| ... | All causes | 28,700 | 100.0 | 3,951.1 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 8,522 | 29.7 | 1,173.2 |
| 2 | Malignant neoplasms (C00-C97) | 8,515 | 29.7 | 1,172.3 |
| 3 | Cerebrovascular diseases (I60-I69) | 2,044 | 7.1 | 281.4 |
| 4 | Diabetes mellitus (E10-E14) | 1,670 | 5.8 | 229.9 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 1,095 | 3.8 | 150.7 |
| 6 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 881 | 3.1 | 121.3 |
| 7 | Septicemia (A40-A41) | 660 | 2.3 | 90.9 |
| 8 | Influenza and pneumonia (J10-J18) | 481 | 1.7 | 66.2 |
| 9 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 421 | 1.5 | 58.0 |
| 10 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 417 | 1.5 | 57.4 |
| 11 | Alzheimer's disease (G30) | 173 | 0.6 | 23.8 |
| 12 | Chronic liver disease and cirrhosis (K70,K73-K74) | 167 | 0.6 | 23.0 |
| 13 | Pneumonitis due to solids and liquids (J69) | 165 | 0.6 | 22.7 |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 137 | 0.5 | 18.9 |
| 15 | Aortic aneurysm and dissection (I71) | 135 | 0.5 | 18.6 |
| ... | All other causes (Residual) | 3,217 | 11.2 | 442.9 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|---------|
| Black, non-Hispanic, both sexes, 75-79 years | | | | |
| ... | All causes | 32,243 | 100.0 | 5,712.5 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 10,132 | 31.4 | 1,795.1 |
| 2 | Malignant neoplasms (C00-C97) | 8,163 | 25.3 | 1,446.2 |
| 3 | Cerebrovascular diseases (I60-I69) | 2,627 | 8.1 | 465.4 |
| 4 | Diabetes mellitus (E10-E14) | 1,735 | 5.4 | 307.4 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 1,178 | 3.7 | 208.7 |
| 6 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 982 | 3.0 | 174.0 |
| 7 | Septicemia (A40-A41) | 797 | 2.5 | 141.2 |
| 8 | Influenza and pneumonia (J10-J18) | 756 | 2.3 | 133.9 |
| 9 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 512 | 1.6 | 90.7 |
| 10 | Alzheimer's disease (G30) | 499 | 1.5 | 88.4 |
| 11 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 447 | 1.4 | 79.2 |
| 12 | Pneumonitis due to solids and liquids (J69) | 221 | 0.7 | 39.2 |
| 13 | Aortic aneurysm and dissection (I71) | 195 | 0.6 | 34.5 |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 141 | 0.4 | 25.0 |
| 15 | Atherosclerosis (I70) | 131 | 0.4 | 23.2 |
| ... | All other causes (Residual) | 3,727 | 11.6 | 660.3 |
| Black, non-Hispanic, both sexes, 80-84 years | | | | |
| ... | All causes | 29,349 | 100.0 | 8,098.0 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 9,549 | 32.5 | 2,634.8 |
| 2 | Malignant neoplasms (C00-C97) | 6,291 | 21.4 | 1,735.8 |
| 3 | Cerebrovascular diseases (I60-I69) | 2,630 | 9.0 | 725.7 |
| 4 | Diabetes mellitus (E10-E14) | 1,397 | 4.8 | 385.5 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 1,012 | 3.4 | 279.2 |
| 6 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 909 | 3.1 | 250.8 |
| 7 | Influenza and pneumonia (J10-J18) | 815 | 2.8 | 224.9 |
| 8 | Septicemia (A40-A41) | 704 | 2.4 | 194.2 |
| 9 | Alzheimer's disease (G30) | 677 | 2.3 | 186.8 |
| 10 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 544 | 1.9 | 150.1 |
| 11 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 383 | 1.3 | 105.7 |
| 12 | Pneumonitis due to solids and liquids (J69) | 276 | 0.9 | 76.2 |
| 13 | Aortic aneurysm and dissection (I71) | 157 | 0.5 | 43.3 |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 155 | 0.5 | 42.8 |
| 15 | Atherosclerosis (I70) | 154 | 0.5 | 42.5 |
| ... | All other causes (Residual) | 3,696 | 12.6 | 1,019.8 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|------|
| Black, non-Hispanic, both sexes, 85-89 years | | | | |
| ... | All causes | 23,592 | 100.0 | --- |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 8,157 | 34.6 | --- |
| 2 | Malignant neoplasms (C00-C97) | 3,776 | 16.0 | --- |
| 3 | Cerebrovascular diseases (I60-I69) | 2,391 | 10.1 | --- |
| 4 | Diabetes mellitus (E10-E14) | 975 | 4.1 | --- |
| 5 | Influenza and pneumonia (J10-J18) | 784 | 3.3 | --- |
| 6 | Alzheimer's disease (G30) | 782 | 3.3 | --- |
| 7 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 707 | 3.0 | --- |
| 8 | Chronic lower respiratory diseases (J40-J47) | 697 | 3.0 | --- |
| 9 | Septicemia (A40-A41) | 626 | 2.7 | --- |
| 10 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 478 | 2.0 | --- |
| 11 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 295 | 1.3 | --- |
| 12 | Pneumonitis due to solids and liquids (J69) | 271 | 1.1 | --- |
| 13 | Atherosclerosis (I70) | 149 | 0.6 | --- |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 116 | 0.5 | --- |
| 15 | Parkinson's disease (G20-G21) | 113 | 0.5 | --- |
| ... | All other causes (Residual) | 3,275 | 13.9 | --- |
| Black, non-Hispanic, both sexes, 90-94 years | | | | |
| ... | All causes | 14,846 | 100.0 | --- |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 5,431 | 36.6 | --- |
| 2 | Malignant neoplasms (C00-C97) | 1,780 | 12.0 | --- |
| 3 | Cerebrovascular diseases (I60-I69) | 1,530 | 10.3 | --- |
| 4 | Influenza and pneumonia (J10-J18) | 629 | 4.2 | --- |
| 5 | Alzheimer's disease (G30) | 522 | 3.5 | --- |
| 6 | Diabetes mellitus (E10-E14) | 507 | 3.4 | --- |
| 7 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 454 | 3.1 | --- |
| 8 | Chronic lower respiratory diseases (J40-J47) | 372 | 2.5 | --- |
| 9 | Septicemia (A40-A41) | 364 | 2.5 | --- |
| 10 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 287 | 1.9 | --- |
| 11 | Pneumonitis due to solids and liquids (J69) | 203 | 1.4 | --- |
| 12 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 196 | 1.3 | --- |
| 13 | Atherosclerosis (I70) | 137 | 0.9 | --- |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 75 | 0.5 | --- |
| 15 | Nutritional deficiencies (E40-E64) | 68 | 0.5 | --- |
| ... | All other causes (Residual) | 2,291 | 15.4 | --- |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|---|---|--------|-------------------------|------|
| Black, non-Hispanic, both sexes, 95-99 years | | | | |
| ... | All causes | 6,423 | 100.0 | --- |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 2,542 | 39.6 | --- |
| 2 | Cerebrovascular diseases (I60-I69) | 604 | 9.4 | --- |
| 3 | Malignant neoplasms (C00-C97) | 567 | 8.8 | --- |
| 4 | Influenza and pneumonia (J10-J18) | 320 | 5.0 | --- |
| 5 | Alzheimer's disease (G30) | 260 | 4.0 | --- |
| 6 | Diabetes mellitus (E10-E14) | 179 | 2.8 | --- |
| 7 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 176 | 2.7 | --- |
| 8 | Septicemia (A40-A41) | 148 | 2.3 | --- |
| 9 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 126 | 2.0 | --- |
| 10 | Chronic lower respiratory diseases (J40-J47) | 117 | 1.8 | --- |
| 11 | Pneumonitis due to solids and liquids (J69) | 92 | 1.4 | --- |
| 12 | Atherosclerosis (I70) | 86 | 1.3 | --- |
| 13 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 73 | 1.1 | --- |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 33 | 0.5 | --- |
| 15 | Nutritional deficiencies (E40-E64) | 30 | 0.5 | --- |
| ... | All other causes (Residual) | 1,070 | 16.7 | --- |
| Black, non-Hispanic, both sexes, 100 years and over | | | | |
| ... | All causes | 2,107 | 100.0 | --- |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 874 | 41.5 | --- |
| 2 | Cerebrovascular diseases (I60-I69) | 170 | 8.1 | --- |
| 3 | Influenza and pneumonia (J10-J18) | 135 | 6.4 | --- |
| 4 | Malignant neoplasms (C00-C97) | 116 | 5.5 | --- |
| 5 | Alzheimer's disease (G30) | 80 | 3.8 | --- |
| 6 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 50 | 2.4 | --- |
| 7 | Diabetes mellitus (E10-E14) | 49 | 2.3 | --- |
| 8 | Septicemia (A40-A41) | 43 | 2.0 | --- |
| 8 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 43 | 2.0 | --- |
| 10 | Pneumonitis due to solids and liquids (J69) | 34 | 1.6 | --- |
| 11 | Atherosclerosis (I70) | 32 | 1.5 | --- |
| 12 | Chronic lower respiratory diseases (J40-J47) | 29 | 1.4 | --- |
| 13 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 25 | 1.2 | --- |
| 14 | Anemias (D50-D64) | 18 | 0.9 | --- |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 9 | 0.4 | --- |
| ... | All other causes (Residual) | 400 | 19.0 | --- |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|--------------------------------------|---|---------|-------------------------|-------|
| Black, non-Hispanic, male, all ages | | | | |
| ... | All causes | 143,971 | 100.0 | 849.7 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 36,541 | 25.4 | 215.7 |
| 2 | Malignant neoplasms (C00-C97) | 32,335 | 22.5 | 190.8 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 8,381 | 5.8 | 49.5 |
| 4 | Cerebrovascular diseases (I60-I69) | 7,828 | 5.4 | 46.2 |
| 5 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 6,658 | 4.6 | 39.3 |
| 6 | Human immunodeficiency virus (HIV) disease (B20-B24) | 5,227 | 3.6 | 30.8 |
| 7 | Diabetes mellitus (E10-E14) | 4,983 | 3.5 | 29.4 |
| 8 | Chronic lower respiratory diseases (J40-J47) | 4,143 | 2.9 | 24.5 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 3,158 | 2.2 | 18.6 |
| 10 | Influenza and pneumonia (J10-J18) | 2,769 | 1.9 | 16.3 |
| 11 | Certain conditions originating in the perinatal period (P00-P96) | 2,674 | 1.9 | 15.8 |
| 12 | Septicemia (A40-A41) | 2,611 | 1.8 | 15.4 |
| 13 | Chronic liver disease and cirrhosis (K70,K73-K74) | 1,819 | 1.3 | 10.7 |
| 14 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 1,698 | 1.2 | 10.0 |
| 15 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 1,578 | 1.1 | 9.3 |
| ... | All other causes (Residual) | 21,568 | 15.0 | 127.3 |
| Black, non-Hispanic, male, 1-4 years | | | | |
| ... | All causes | 655 | 100.0 | 55.1 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 188 | 28.7 | 15.8 |
| 2 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 100 | 15.3 | 8.4 |
| 3 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 65 | 9.9 | 5.5 |
| 4 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 40 | 6.1 | 3.4 |
| 5 | Malignant neoplasms (C00-C97) | 29 | 4.4 | 2.4 |
| 6 | Chronic lower respiratory diseases (J40-J47) | 16 | 2.4 | * |
| 7 | Certain conditions originating in the perinatal period (P00-P96) | 14 | 2.1 | * |
| 8 | Influenza and pneumonia (J10-J18) | 13 | 2.0 | * |
| 9 | Septicemia (A40-A41) | 12 | 1.8 | * |
| 10 | Human immunodeficiency virus (HIV) disease (B20-B24) | 8 | 1.2 | * |
| 11 | Anemias (D50-D64) | 7 | 1.1 | * |
| 12 | Cerebrovascular diseases (I60-I69) | 5 | 0.8 | * |
| 13 | Complications of medical and surgical care (Y40-Y84,Y88) | 4 | 0.6 | * |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 3 | 0.5 | * |
| 14 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 3 | 0.5 | * |
| ... | All other causes (Residual) | 148 | 22.6 | 12.5 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|------|
| Black, non-Hispanic, male, 5-9 years | | | | |
| ... | All causes | 400 | 100.0 | 24.6 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 212 | 53.0 | 13.0 |
| 2 | Malignant neoplasms (C00-C97) | 43 | 10.8 | 2.6 |
| 3 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 19 | 4.8 | * |
| 4 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 15 | 3.8 | * |
| 5 | Chronic lower respiratory diseases (J40-J47) | 14 | 3.5 | * |
| 6 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 10 | 2.5 | * |
| 7 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 6 | 1.5 | * |
| 8 | Human immunodeficiency virus (HIV) disease (B20-B24) | 5 | 1.3 | * |
| 8 | Anemias (D50-D64) | 5 | 1.3 | * |
| 10 | Influenza and pneumonia (J10-J18) | 4 | 1.0 | * |
| 11 | Meningitis (G00,G03) | 3 | 0.8 | * |
| 11 | Certain conditions originating in the perinatal period (P00-P96) | 3 | 0.8 | * |
| 13 | Meningococcal infection (A39) | 2 | 0.5 | * |
| 13 | Cerebrovascular diseases (I60-I69) | 2 | 0.5 | * |
| 13 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 2 | 0.5 | * |
| ... | All other causes (Residual) | 55 | 13.8 | 3.4 |
| Black, non-Hispanic, male, 10-14 years | | | | |
| ... | All causes | 541 | 100.0 | 32.5 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 220 | 40.7 | 13.2 |
| 2 | Malignant neoplasms (C00-C97) | 57 | 10.5 | 3.4 |
| 3 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 47 | 8.7 | 2.8 |
| 4 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 29 | 5.4 | 1.7 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 24 | 4.4 | 1.4 |
| 6 | Chronic lower respiratory diseases (J40-J47) | 18 | 3.3 | * |
| 6 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 18 | 3.3 | * |
| 8 | Human immunodeficiency virus (HIV) disease (B20-B24) | 10 | 1.8 | * |
| 9 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 7 | 1.3 | * |
| 10 | Anemias (D50-D64) | 5 | 0.9 | * |
| 10 | Diabetes mellitus (E10-E14) | 5 | 0.9 | * |
| 10 | Influenza and pneumonia (J10-J18) | 5 | 0.9 | * |
| 13 | Cerebrovascular diseases (I60-I69) | 3 | 0.6 | * |
| 14 | Septicemia (A40-A41) | 2 | 0.4 | * |
| 14 | Pneumonitis due to solids and liquids (J69) | 2 | 0.4 | * |
| 14 | Certain conditions originating in the perinatal period (P00-P96) | 2 | 0.4 | * |
| 14 | Complications of medical and surgical care (Y40-Y84,Y88) | 2 | 0.4 | * |
| ... | All other causes (Residual) | 85 | 15.7 | 5.1 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|-------|
| Black, non-Hispanic, male, 15-19 years | | | | |
| ... | All causes | 2,041 | 100.0 | 133.9 |
| 1 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 911 | 44.6 | 59.7 |
| 2 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 564 | 27.6 | 37.0 |
| 3 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 115 | 5.6 | 7.5 |
| 4 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 69 | 3.4 | 4.5 |
| 5 | Malignant neoplasms (C00-C97) | 65 | 3.2 | 4.3 |
| 6 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 41 | 2.0 | 2.7 |
| 7 | Chronic lower respiratory diseases (J40-J47) | 22 | 1.1 | 1.4 |
| 8 | Human immunodeficiency virus (HIV) disease (B20-B24) | 18 | 0.9 | * |
| 8 | Anemias (D50-D64) | 18 | 0.9 | * |
| 10 | Influenza and pneumonia (J10-J18) | 11 | 0.5 | * |
| 11 | Legal intervention (Y35,Y89.0) | 10 | 0.5 | * |
| 12 | Septicemia (A40-A41) | 9 | 0.4 | * |
| 13 | Cerebrovascular diseases (I60-I69) | 8 | 0.4 | * |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 7 | 0.3 | * |
| 14 | Diabetes mellitus (E10-E14) | 7 | 0.3 | * |
| ... | All other causes (Residual) | 166 | 8.1 | 10.9 |
| Black, non-Hispanic, male, 20-24 years | | | | |
| ... | All causes | 3,336 | 100.0 | 244.7 |
| 1 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 1,640 | 49.2 | 120.3 |
| 2 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 721 | 21.6 | 52.9 |
| 3 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 273 | 8.2 | 20.0 |
| 4 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 136 | 4.1 | 10.0 |
| 5 | Malignant neoplasms (C00-C97) | 83 | 2.5 | 6.1 |
| 6 | Human immunodeficiency virus (HIV) disease (B20-B24) | 50 | 1.5 | 3.7 |
| 7 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 31 | 0.9 | 2.3 |
| 8 | Chronic lower respiratory diseases (J40-J47) | 26 | 0.8 | 1.9 |
| 9 | Anemias (D50-D64) | 23 | 0.7 | 1.7 |
| 10 | Diabetes mellitus (E10-E14) | 20 | 0.6 | 1.5 |
| 11 | Septicemia (A40-A41) | 16 | 0.5 | * |
| 12 | Legal intervention (Y35,Y89.0) | 14 | 0.4 | * |
| 13 | Cerebrovascular diseases (I60-I69) | 12 | 0.4 | * |
| 13 | Influenza and pneumonia (J10-J18) | 12 | 0.4 | * |
| 15 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 10 | 0.3 | * |
| ... | All other causes (Residual) | 269 | 8.1 | 19.7 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|-------|
| Black, non-Hispanic, male, 25-29 years | | | | |
| ... | All causes | 3,056 | 100.0 | 255.7 |
| 1 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 1,192 | 39.0 | 99.7 |
| 2 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 592 | 19.4 | 49.5 |
| 3 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 212 | 6.9 | 17.7 |
| 4 | Human immunodeficiency virus (HIV) disease (B20-B24) | 210 | 6.9 | 17.6 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 199 | 6.5 | 16.7 |
| 6 | Malignant neoplasms (C00-C97) | 106 | 3.5 | 8.9 |
| 7 | Diabetes mellitus (E10-E14) | 49 | 1.6 | 4.1 |
| 8 | Cerebrovascular diseases (I60-I69) | 33 | 1.1 | 2.8 |
| 9 | Anemias (D50-D64) | 31 | 1.0 | 2.6 |
| 10 | Chronic lower respiratory diseases (J40-J47) | 26 | 0.9 | 2.2 |
| 11 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 22 | 0.7 | 1.8 |
| 12 | Influenza and pneumonia (J10-J18) | 17 | 0.6 | * |
| 13 | Septicemia (A40-A41) | 13 | 0.4 | * |
| 14 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 12 | 0.4 | * |
| 15 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 11 | 0.4 | * |
| ... | All other causes (Residual) | 331 | 10.8 | 27.7 |
| Black, non-Hispanic, male, 30-34 years | | | | |
| ... | All causes | 3,492 | 100.0 | 279.6 |
| 1 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 806 | 23.1 | 64.5 |
| 2 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 600 | 17.2 | 48.0 |
| 3 | Human immunodeficiency virus (HIV) disease (B20-B24) | 543 | 15.5 | 43.5 |
| 4 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 379 | 10.9 | 30.3 |
| 5 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 194 | 5.6 | 15.5 |
| 6 | Malignant neoplasms (C00-C97) | 161 | 4.6 | 12.9 |
| 7 | Diabetes mellitus (E10-E14) | 73 | 2.1 | 5.8 |
| 8 | Chronic lower respiratory diseases (J40-J47) | 43 | 1.2 | 3.4 |
| 9 | Anemias (D50-D64) | 42 | 1.2 | 3.4 |
| 9 | Cerebrovascular diseases (I60-I69) | 42 | 1.2 | 3.4 |
| 11 | Influenza and pneumonia (J10-J18) | 35 | 1.0 | 2.8 |
| 12 | Septicemia (A40-A41) | 33 | 0.9 | 2.6 |
| 13 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 32 | 0.9 | 2.6 |
| 14 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 19 | 0.5 | * |
| 15 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 17 | 0.5 | * |
| ... | All other causes (Residual) | 473 | 13.5 | 37.9 |

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|-------|
| Black, non-Hispanic, male, 35-39 years | | | | |
| ... | All causes | 4,959 | 100.0 | 375.1 |
| 1 | Human immunodeficiency virus (HIV) disease (B20-B24) | 947 | 19.1 | 71.6 |
| 2 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 718 | 14.5 | 54.3 |
| 3 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 715 | 14.4 | 54.1 |
| 4 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 584 | 11.8 | 44.2 |
| 5 | Malignant neoplasms (C00-C97) | 365 | 7.4 | 27.6 |
| 6 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 188 | 3.8 | 14.2 |
| 7 | Diabetes mellitus (E10-E14) | 119 | 2.4 | 9.0 |
| 8 | Cerebrovascular diseases (I60-I69) | 118 | 2.4 | 8.9 |
| 9 | Chronic liver disease and cirrhosis (K70,K73-K74) | 77 | 1.6 | 5.8 |
| 10 | Influenza and pneumonia (J10-J18) | 62 | 1.3 | 4.7 |
| 11 | Chronic lower respiratory diseases (J40-J47) | 58 | 1.2 | 4.4 |
| 12 | Septicemia (A40-A41) | 51 | 1.0 | 3.9 |
| 13 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 45 | 0.9 | 3.4 |
| 14 | Anemias (D50-D64) | 30 | 0.6 | 2.3 |
| 14 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 30 | 0.6 | 2.3 |
| ... | All other causes (Residual) | 852 | 17.2 | 64.4 |
| Black, non-Hispanic, male, 40-44 years | | | | |
| ... | All causes | 7,012 | 100.0 | 538.9 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 1,356 | 19.3 | 104.2 |
| 2 | Human immunodeficiency virus (HIV) disease (B20-B24) | 1,093 | 15.6 | 84.0 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 890 | 12.7 | 68.4 |
| 4 | Malignant neoplasms (C00-C97) | 836 | 11.9 | 64.3 |
| 5 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 452 | 6.4 | 34.7 |
| 6 | Cerebrovascular diseases (I60-I69) | 237 | 3.4 | 18.2 |
| 7 | Chronic liver disease and cirrhosis (K70,K73-K74) | 196 | 2.8 | 15.1 |
| 8 | Diabetes mellitus (E10-E14) | 156 | 2.2 | 12.0 |
| 9 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 141 | 2.0 | 10.8 |
| 10 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 94 | 1.3 | 7.2 |
| 11 | Influenza and pneumonia (J10-J18) | 93 | 1.3 | 7.1 |
| 12 | Septicemia (A40-A41) | 77 | 1.1 | 5.9 |
| 13 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 72 | 1.0 | 5.5 |
| 14 | Chronic lower respiratory diseases (J40-J47) | 66 | 0.9 | 5.1 |
| 15 | Viral hepatitis (B15-B19) | 64 | 0.9 | 4.9 |
| ... | All other causes (Residual) | 1,189 | 17.0 | 91.4 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|---------|
| Black, non-Hispanic, male, 45-49 years | | | | |
| ... | All causes | 9,354 | 100.0 | 846.0 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 2,087 | 22.3 | 188.8 |
| 2 | Malignant neoplasms (C00-C97) | 1,628 | 17.4 | 147.2 |
| 3 | Human immunodeficiency virus (HIV) disease (B20-B24) | 981 | 10.5 | 88.7 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 918 | 9.8 | 83.0 |
| 5 | Cerebrovascular diseases (I60-I69) | 412 | 4.4 | 37.3 |
| 6 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 333 | 3.6 | 30.1 |
| 7 | Chronic liver disease and cirrhosis (K70,K73-K74) | 319 | 3.4 | 28.9 |
| 8 | Diabetes mellitus (E10-E14) | 285 | 3.0 | 25.8 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 188 | 2.0 | 17.0 |
| 10 | Septicemia (A40-A41) | 133 | 1.4 | 12.0 |
| 11 | Influenza and pneumonia (J10-J18) | 128 | 1.4 | 11.6 |
| 12 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 113 | 1.2 | 10.2 |
| 13 | Viral hepatitis (B15-B19) | 112 | 1.2 | 10.1 |
| 14 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 109 | 1.2 | 9.9 |
| 15 | Chronic lower respiratory diseases (J40-J47) | 101 | 1.1 | 9.1 |
| ... | All other causes (Residual) | 1,507 | 16.1 | 136.3 |
| Black, non-Hispanic, male, 50-54 years | | | | |
| ... | All causes | 10,953 | 100.0 | 1,222.7 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 2,868 | 26.2 | 320.2 |
| 2 | Malignant neoplasms (C00-C97) | 2,568 | 23.4 | 286.7 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 656 | 6.0 | 73.2 |
| 4 | Human immunodeficiency virus (HIV) disease (B20-B24) | 648 | 5.9 | 72.3 |
| 5 | Cerebrovascular diseases (I60-I69) | 586 | 5.4 | 65.4 |
| 6 | Diabetes mellitus (E10-E14) | 416 | 3.8 | 46.4 |
| 7 | Chronic liver disease and cirrhosis (K70,K73-K74) | 383 | 3.5 | 42.8 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 211 | 1.9 | 23.6 |
| 8 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 211 | 1.9 | 23.6 |
| 10 | Septicemia (A40-A41) | 171 | 1.6 | 19.1 |
| 11 | Chronic lower respiratory diseases (J40-J47) | 170 | 1.6 | 19.0 |
| 12 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 150 | 1.4 | 16.7 |
| 13 | Influenza and pneumonia (J10-J18) | 142 | 1.3 | 15.9 |
| 14 | Viral hepatitis (B15-B19) | 128 | 1.2 | 14.3 |
| 15 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 87 | 0.8 | 9.7 |
| ... | All other causes (Residual) | 1,558 | 14.2 | 173.9 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|---------|
| Black, non-Hispanic, male, 55-59 years | | | | |
| ... | All causes | 10,845 | 100.0 | 1,746.2 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 3,170 | 29.2 | 510.4 |
| 2 | Malignant neoplasms (C00-C97) | 3,133 | 28.9 | 504.5 |
| 3 | Cerebrovascular diseases (I60-I69) | 554 | 5.1 | 89.2 |
| 4 | Diabetes mellitus (E10-E14) | 483 | 4.5 | 77.8 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 416 | 3.8 | 67.0 |
| 6 | Human immunodeficiency virus (HIV) disease (B20-B24) | 315 | 2.9 | 50.7 |
| 7 | Chronic liver disease and cirrhosis (K70,K73-K74) | 278 | 2.6 | 44.8 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 240 | 2.2 | 38.6 |
| 9 | Chronic lower respiratory diseases (J40-J47) | 235 | 2.2 | 37.8 |
| 10 | Septicemia (A40-A41) | 211 | 1.9 | 34.0 |
| 11 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 136 | 1.3 | 21.9 |
| 12 | Influenza and pneumonia (J10-J18) | 129 | 1.2 | 20.8 |
| 13 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 109 | 1.0 | 17.6 |
| 14 | Viral hepatitis (B15-B19) | 63 | 0.6 | 10.1 |
| 15 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 62 | 0.6 | 10.0 |
| ... | All other causes (Residual) | 1,311 | 12.1 | 211.1 |
| Black, non-Hispanic, male, 60-64 years | | | | |
| ... | All causes | 11,755 | 100.0 | 2,456.8 |
| 1 | Malignant neoplasms (C00-C97) | 3,619 | 30.8 | 756.4 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 3,543 | 30.1 | 740.5 |
| 3 | Cerebrovascular diseases (I60-I69) | 698 | 5.9 | 145.9 |
| 4 | Diabetes mellitus (E10-E14) | 548 | 4.7 | 114.5 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 334 | 2.8 | 69.8 |
| 6 | Chronic lower respiratory diseases (J40-J47) | 319 | 2.7 | 66.7 |
| 7 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 269 | 2.3 | 56.2 |
| 8 | Human immunodeficiency virus (HIV) disease (B20-B24) | 209 | 1.8 | 43.7 |
| 9 | Septicemia (A40-A41) | 200 | 1.7 | 41.8 |
| 10 | Chronic liver disease and cirrhosis (K70,K73-K74) | 184 | 1.6 | 38.5 |
| 11 | Influenza and pneumonia (J10-J18) | 164 | 1.4 | 34.3 |
| 12 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 148 | 1.3 | 30.9 |
| 13 | Aortic aneurysm and dissection (I71) | 71 | 0.6 | 14.8 |
| 14 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 56 | 0.5 | 11.7 |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 45 | 0.4 | 9.4 |
| ... | All other causes (Residual) | 1,348 | 11.5 | 281.7 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|---------|
| Black, non-Hispanic, male, 65-69 years | | | | |
| ... | All causes | 13,617 | 100.0 | 3,547.9 |
| 1 | Malignant neoplasms (C00-C97) | 4,430 | 32.5 | 1,154.2 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 4,044 | 29.7 | 1,053.7 |
| 3 | Cerebrovascular diseases (I60-I69) | 854 | 6.3 | 222.5 |
| 4 | Diabetes mellitus (E10-E14) | 629 | 4.6 | 163.9 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 496 | 3.6 | 129.2 |
| 6 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 397 | 2.9 | 103.4 |
| 7 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 268 | 2.0 | 69.8 |
| 8 | Septicemia (A40-A41) | 259 | 1.9 | 67.5 |
| 9 | Influenza and pneumonia (J10-J18) | 195 | 1.4 | 50.8 |
| 10 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 190 | 1.4 | 49.5 |
| 11 | Chronic liver disease and cirrhosis (K70,K73-K74) | 168 | 1.2 | 43.8 |
| 12 | Human immunodeficiency virus (HIV) disease (B20-B24) | 99 | 0.7 | 25.8 |
| 13 | Aortic aneurysm and dissection (I71) | 57 | 0.4 | 14.9 |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 51 | 0.4 | 13.3 |
| 15 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 50 | 0.4 | 13.0 |
| ... | All other causes (Residual) | 1,430 | 10.5 | 372.6 |
| Black, non-Hispanic, male, 70-74 years | | | | |
| ... | All causes | 14,829 | 100.0 | 5,105.6 |
| 1 | Malignant neoplasms (C00-C97) | 4,704 | 31.7 | 1,619.6 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 4,447 | 30.0 | 1,531.1 |
| 3 | Cerebrovascular diseases (I60-I69) | 990 | 6.7 | 340.9 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 651 | 4.4 | 224.1 |
| 5 | Diabetes mellitus (E10-E14) | 649 | 4.4 | 223.4 |
| 6 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 385 | 2.6 | 132.6 |
| 7 | Septicemia (A40-A41) | 327 | 2.2 | 112.6 |
| 8 | Influenza and pneumonia (J10-J18) | 277 | 1.9 | 95.4 |
| 9 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 237 | 1.6 | 81.6 |
| 10 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 184 | 1.2 | 63.4 |
| 11 | Chronic liver disease and cirrhosis (K70,K73-K74) | 100 | 0.7 | 34.4 |
| 12 | Pneumonitis due to solids and liquids (J69) | 91 | 0.6 | 31.3 |
| 13 | Aortic aneurysm and dissection (I71) | 77 | 0.5 | 26.5 |
| 14 | Alzheimer's disease (G30) | 72 | 0.5 | 24.8 |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 68 | 0.5 | 23.4 |
| ... | All other causes (Residual) | 1,570 | 10.6 | 540.5 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|----------|
| Black, non-Hispanic, male, 75-79 years | | | | |
| ... | All causes | 15,429 | 100.0 | 7,313.8 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 4,705 | 30.5 | 2,230.3 |
| 2 | Malignant neoplasms (C00-C97) | 4,432 | 28.7 | 2,100.9 |
| 3 | Cerebrovascular diseases (I60-I69) | 1,119 | 7.3 | 530.4 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 703 | 4.6 | 333.2 |
| 5 | Diabetes mellitus (E10-E14) | 655 | 4.2 | 310.5 |
| 6 | Influenza and pneumonia (J10-J18) | 429 | 2.8 | 203.4 |
| 7 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 402 | 2.6 | 190.6 |
| 8 | Septicemia (A40-A41) | 365 | 2.4 | 173.0 |
| 9 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 256 | 1.7 | 121.4 |
| 10 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 202 | 1.3 | 95.8 |
| 11 | Alzheimer's disease (G30) | 182 | 1.2 | 86.3 |
| 12 | Pneumonitis due to solids and liquids (J69) | 120 | 0.8 | 56.9 |
| 13 | Aortic aneurysm and dissection (I71) | 90 | 0.6 | 42.7 |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 68 | 0.4 | 32.2 |
| 15 | Atherosclerosis (I70) | 65 | 0.4 | 30.8 |
| ... | All other causes (Residual) | 1,636 | 10.6 | 775.5 |
| Black, non-Hispanic, male, 80-84 years | | | | |
| ... | All causes | 12,654 | 100.0 | 10,346.7 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 3,916 | 30.9 | 3,202.0 |
| 2 | Malignant neoplasms (C00-C97) | 3,268 | 25.8 | 2,672.1 |
| 3 | Cerebrovascular diseases (I60-I69) | 949 | 7.5 | 776.0 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 582 | 4.6 | 475.9 |
| 5 | Diabetes mellitus (E10-E14) | 495 | 3.9 | 404.7 |
| 6 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 382 | 3.0 | 312.3 |
| 7 | Influenza and pneumonia (J10-J18) | 379 | 3.0 | 309.9 |
| 8 | Septicemia (A40-A41) | 291 | 2.3 | 237.9 |
| 9 | Alzheimer's disease (G30) | 230 | 1.8 | 188.1 |
| 10 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 205 | 1.6 | 167.6 |
| 11 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 197 | 1.6 | 161.1 |
| 12 | Pneumonitis due to solids and liquids (J69) | 132 | 1.0 | 107.9 |
| 13 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 73 | 0.6 | 59.7 |
| 14 | Parkinson's disease (G20-G21) | 72 | 0.6 | 58.9 |
| 15 | Aortic aneurysm and dissection (I71) | 68 | 0.5 | 55.6 |
| ... | All other causes (Residual) | 1,415 | 11.2 | 1,157.0 |

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|------|
| Black, non-Hispanic, male, 85-89 years | | | | |
| ... | All causes | 8,313 | 100.0 | --- |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 2,630 | 31.6 | --- |
| 2 | Malignant neoplasms (C00-C97) | 1,830 | 22.0 | --- |
| 3 | Cerebrovascular diseases (I60-I69) | 726 | 8.7 | --- |
| 4 | Chronic lower respiratory diseases (J40-J47) | 362 | 4.4 | --- |
| 5 | Influenza and pneumonia (J10-J18) | 289 | 3.5 | --- |
| 6 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 260 | 3.1 | --- |
| 7 | Diabetes mellitus (E10-E14) | 236 | 2.8 | --- |
| 8 | Septicemia (A40-A41) | 225 | 2.7 | --- |
| 9 | Alzheimer's disease (G30) | 219 | 2.6 | --- |
| 10 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 146 | 1.8 | --- |
| 11 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 122 | 1.5 | --- |
| 12 | Pneumonitis due to solids and liquids (J69) | 115 | 1.4 | --- |
| 13 | Parkinson's disease (G20-G21) | 53 | 0.6 | --- |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 45 | 0.5 | --- |
| 15 | Atherosclerosis (I70) | 44 | 0.5 | --- |
| ... | All other causes (Residual) | 1,011 | 12.2 | --- |
| Black, non-Hispanic, male, 90-94 years | | | | |
| ... | All causes | 4,249 | 100.0 | --- |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 1,451 | 34.1 | --- |
| 2 | Malignant neoplasms (C00-C97) | 734 | 17.3 | --- |
| 3 | Cerebrovascular diseases (I60-I69) | 319 | 7.5 | --- |
| 4 | Influenza and pneumonia (J10-J18) | 224 | 5.3 | --- |
| 5 | Chronic lower respiratory diseases (J40-J47) | 166 | 3.9 | --- |
| 6 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 142 | 3.3 | --- |
| 7 | Diabetes mellitus (E10-E14) | 120 | 2.8 | --- |
| 8 | Septicemia (A40-A41) | 110 | 2.6 | --- |
| 9 | Alzheimer's disease (G30) | 109 | 2.6 | --- |
| 10 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 73 | 1.7 | --- |
| 11 | Pneumonitis due to solids and liquids (J69) | 69 | 1.6 | --- |
| 12 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 60 | 1.4 | --- |
| 13 | Atherosclerosis (I70) | 35 | 0.8 | --- |
| 14 | Parkinson's disease (G20-G21) | 29 | 0.7 | --- |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 22 | 0.5 | --- |
| ... | All other causes (Residual) | 586 | 13.8 | --- |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|---|---|--------|-------------------------|------|
| Black, non-Hispanic, male, 95-99 years | | | | |
| ... | All causes | 1,415 | 100.0 | --- |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 528 | 37.3 | --- |
| 2 | Malignant neoplasms (C00-C97) | 203 | 14.3 | --- |
| 3 | Cerebrovascular diseases (I60-I69) | 114 | 8.1 | --- |
| 4 | Influenza and pneumonia (J10-J18) | 82 | 5.8 | --- |
| 5 | Chronic lower respiratory diseases (J40-J47) | 51 | 3.6 | --- |
| 6 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 39 | 2.8 | --- |
| 7 | Diabetes mellitus (E10-E14) | 29 | 2.0 | --- |
| 8 | Septicemia (A40-A41) | 27 | 1.9 | --- |
| 8 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 27 | 1.9 | --- |
| 10 | Alzheimer's disease (G30) | 25 | 1.8 | --- |
| 11 | Pneumonitis due to solids and liquids (J69) | 22 | 1.6 | --- |
| 12 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 21 | 1.5 | --- |
| 13 | Atherosclerosis (I70) | 12 | 0.8 | --- |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 7 | 0.5 | --- |
| 15 | Anemias (D50-D64) | 6 | 0.4 | --- |
| ... | All other causes (Residual) | 222 | 15.7 | --- |
| Black, non-Hispanic, male, 100 years and over | | | | |
| ... | All causes | 375 | 100.0 | --- |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 149 | 39.7 | --- |
| 2 | Malignant neoplasms (C00-C97) | 37 | 9.9 | --- |
| 3 | Influenza and pneumonia (J10-J18) | 27 | 7.2 | --- |
| 4 | Cerebrovascular diseases (I60-I69) | 18 | 4.8 | --- |
| 5 | Chronic lower respiratory diseases (J40-J47) | 12 | 3.2 | --- |
| 6 | Septicemia (A40-A41) | 11 | 2.9 | --- |
| 6 | Alzheimer's disease (G30) | 11 | 2.9 | --- |
| 8 | Diabetes mellitus (E10-E14) | 7 | 1.9 | --- |
| 8 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 7 | 1.9 | --- |
| 8 | Atherosclerosis (I70) | 7 | 1.9 | --- |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 7 | 1.9 | --- |
| 12 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 6 | 1.6 | --- |
| 13 | Pneumonitis due to solids and liquids (J69) | 5 | 1.3 | --- |
| 14 | Anemias (D50-D64) | 3 | 0.8 | --- |
| 14 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 3 | 0.8 | --- |
| ... | All other causes (Residual) | 65 | 17.3 | --- |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|--|---|---------|-------------------------|-------|
| Black, non-Hispanic, female, all ages | | | | |
| ... | All causes | 140,372 | 100.0 | 751.2 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 40,253 | 28.7 | 215.4 |
| 2 | Malignant neoplasms (C00-C97) | 29,245 | 20.8 | 156.5 |
| 3 | Cerebrovascular diseases (I60-I69) | 11,016 | 7.8 | 59.0 |
| 4 | Diabetes mellitus (E10-E14) | 7,184 | 5.1 | 38.4 |
| 5 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 4,056 | 2.9 | 21.7 |
| 6 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 3,858 | 2.7 | 20.6 |
| 7 | Chronic lower respiratory diseases (J40-J47) | 3,361 | 2.4 | 18.0 |
| 8 | Septicemia (A40-A41) | 3,218 | 2.3 | 17.2 |
| 9 | Influenza and pneumonia (J10-J18) | 2,917 | 2.1 | 15.6 |
| 10 | Human immunodeficiency virus (HIV) disease (B20-B24) | 2,483 | 1.8 | 13.3 |
| 11 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 2,314 | 1.6 | 12.4 |
| 12 | Alzheimer's disease (G30) | 2,198 | 1.6 | 11.8 |
| 13 | Certain conditions originating in the perinatal period (P00-P96) | 2,039 | 1.5 | 10.9 |
| 14 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 1,419 | 1.0 | 7.6 |
| 15 | Chronic liver disease and cirrhosis (K70,K73-K74) | 914 | 0.7 | 4.9 |
| ... | All other causes (Residual) | 23,897 | 17.0 | 127.9 |
| Black, non-Hispanic, female, 1-4 years | | | | |
| ... | All causes | 501 | 100.0 | 43.6 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 143 | 28.5 | 12.4 |
| 2 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 66 | 13.2 | 5.7 |
| 3 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 62 | 12.4 | 5.4 |
| 4 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 33 | 6.6 | 2.9 |
| 5 | Malignant neoplasms (C00-C97) | 17 | 3.4 | * |
| 6 | Influenza and pneumonia (J10-J18) | 11 | 2.2 | * |
| 7 | Septicemia (A40-A41) | 9 | 1.8 | * |
| 8 | Certain conditions originating in the perinatal period (P00-P96) | 8 | 1.6 | * |
| 9 | Cerebrovascular diseases (I60-I69) | 7 | 1.4 | * |
| 10 | Chronic lower respiratory diseases (J40-J47) | 6 | 1.2 | * |
| 11 | Anemias (D50-D64) | 4 | 0.8 | * |
| 12 | Meningococcal infection (A39) | 3 | 0.6 | * |
| 12 | Meningitis (G00,G03) | 3 | 0.6 | * |
| 12 | Complications of medical and surgical care (Y40-Y84,Y88) | 3 | 0.6 | * |
| 15 | Human immunodeficiency virus (HIV) disease (B20-B24) | 2 | 0.4 | * |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 2 | 0.4 | * |
| ... | All other causes (Residual) | 122 | 24.4 | 10.6 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|------|
| Black, non-Hispanic, female, 5-9 years | | | | |
| ... | All causes | 285 | 100.0 | 18.1 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 101 | 35.4 | 6.4 |
| 2 | Malignant neoplasms (C00-C97) | 35 | 12.3 | 2.2 |
| 3 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 17 | 6.0 | * |
| 3 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 17 | 6.0 | * |
| 5 | Chronic lower respiratory diseases (J40-J47) | 9 | 3.2 | * |
| 5 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 9 | 3.2 | * |
| 7 | Anemias (D50-D64) | 8 | 2.8 | * |
| 8 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 7 | 2.5 | * |
| 9 | Influenza and pneumonia (J10-J18) | 5 | 1.8 | * |
| 10 | Human immunodeficiency virus (HIV) disease (B20-B24) | 4 | 1.4 | * |
| 10 | Cerebrovascular diseases (I60-I69) | 4 | 1.4 | * |
| 10 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 4 | 1.4 | * |
| 13 | Diabetes mellitus (E10-E14) | 3 | 1.1 | * |
| 13 | Certain conditions originating in the perinatal period (P00-P96) | 3 | 1.1 | * |
| 15 | Meningococcal infection (A39) | 2 | 0.7 | * |
| ... | All other causes (Residual) | 57 | 20.0 | 3.6 |
| Black, non-Hispanic, female, 10-14 years | | | | |
| ... | All causes | 331 | 100.0 | 20.5 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 97 | 29.3 | 6.0 |
| 2 | Malignant neoplasms (C00-C97) | 37 | 11.2 | 2.3 |
| 3 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 23 | 6.9 | 1.4 |
| 4 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 19 | 5.7 | * |
| 5 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 17 | 5.1 | * |
| 6 | Chronic lower respiratory diseases (J40-J47) | 12 | 3.6 | * |
| 7 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 9 | 2.7 | * |
| 8 | Anemias (D50-D64) | 8 | 2.4 | * |
| 9 | Human immunodeficiency virus (HIV) disease (B20-B24) | 7 | 2.1 | * |
| 10 | Septicemia (A40-A41) | 6 | 1.8 | * |
| 10 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 6 | 1.8 | * |
| 10 | Diabetes mellitus (E10-E14) | 6 | 1.8 | * |
| 13 | Cerebrovascular diseases (I60-I69) | 5 | 1.5 | * |
| 14 | Influenza and pneumonia (J10-J18) | 3 | 0.9 | * |
| 14 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 3 | 0.9 | * |
| 14 | Certain conditions originating in the perinatal period (P00-P96) | 3 | 0.9 | * |
| ... | All other causes (Residual) | 70 | 21.1 | 4.3 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|------|
| Black, non-Hispanic, female, 15-19 years | | | | |
| ... | All causes | 621 | 100.0 | 42.0 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 196 | 31.6 | 13.3 |
| 2 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 98 | 15.8 | 6.6 |
| 3 | Malignant neoplasms (C00-C97) | 45 | 7.2 | 3.0 |
| 4 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 35 | 5.6 | 2.4 |
| 5 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 20 | 3.2 | 1.4 |
| 6 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 19 | 3.1 | * |
| 7 | Human immunodeficiency virus (HIV) disease (B20-B24) | 16 | 2.6 | * |
| 8 | Pregnancy, childbirth and the puerperium (O00-099) | 14 | 2.3 | * |
| 9 | Septicemia (A40-A41) | 10 | 1.6 | * |
| 9 | Diabetes mellitus (E10-E14) | 10 | 1.6 | * |
| 11 | Anemias (D50-D64) | 9 | 1.4 | * |
| 12 | Chronic lower respiratory diseases (J40-J47) | 8 | 1.3 | * |
| 13 | Meningococcal infection (A39) | 6 | 1.0 | * |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 5 | 0.8 | * |
| 14 | Cerebrovascular diseases (I60-I69) | 5 | 0.8 | * |
| 14 | Influenza and pneumonia (J10-J18) | 5 | 0.8 | * |
| ... | All other causes (Residual) | 120 | 19.3 | 8.1 |
| Black, non-Hispanic, female, 20-24 years | | | | |
| ... | All causes | 1,026 | 100.0 | 72.2 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 232 | 22.6 | 16.3 |
| 2 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 168 | 16.4 | 11.8 |
| 3 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 82 | 8.0 | 5.8 |
| 4 | Human immunodeficiency virus (HIV) disease (B20-B24) | 77 | 7.5 | 5.4 |
| 5 | Malignant neoplasms (C00-C97) | 64 | 6.2 | 4.5 |
| 6 | Pregnancy, childbirth and the puerperium (O00-099) | 31 | 3.0 | 2.2 |
| 7 | Cerebrovascular diseases (I60-I69) | 26 | 2.5 | 1.8 |
| 8 | Anemias (D50-D64) | 22 | 2.1 | 1.5 |
| 9 | Chronic lower respiratory diseases (J40-J47) | 21 | 2.0 | 1.5 |
| 9 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 21 | 2.0 | 1.5 |
| 11 | Diabetes mellitus (E10-E14) | 19 | 1.9 | * |
| 12 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 18 | 1.8 | * |
| 13 | Influenza and pneumonia (J10-J18) | 17 | 1.7 | * |
| 14 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 12 | 1.2 | * |
| 15 | Chronic liver disease and cirrhosis (K70,K73-K74) | 7 | 0.7 | * |
| ... | All other causes (Residual) | 209 | 20.4 | 14.7 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|-------|
| Black, non-Hispanic, female, 25-29 years | | | | |
| ... | All causes | 1,332 | 100.0 | 101.4 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 192 | 14.4 | 14.6 |
| 2 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 178 | 13.4 | 13.6 |
| 3 | Human immunodeficiency virus (HIV) disease (B20-B24) | 171 | 12.8 | 13.0 |
| 4 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 140 | 10.5 | 10.7 |
| 5 | Malignant neoplasms (C00-C97) | 135 | 10.1 | 10.3 |
| 6 | Pregnancy, childbirth and the puerperium (O00-099) | 35 | 2.6 | 2.7 |
| 7 | Diabetes mellitus (E10-E14) | 34 | 2.6 | 2.6 |
| 8 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 31 | 2.3 | 2.4 |
| 9 | Cerebrovascular diseases (I60-I69) | 24 | 1.8 | 1.8 |
| 9 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 24 | 1.8 | 1.8 |
| 11 | Chronic lower respiratory diseases (J40-J47) | 21 | 1.6 | 1.6 |
| 12 | Anemias (D50-D64) | 19 | 1.4 | * |
| 13 | Septicemia (A40-A41) | 15 | 1.1 | * |
| 14 | Influenza and pneumonia (J10-J18) | 14 | 1.1 | * |
| 15 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 13 | 1.0 | * |
| ... | All other causes (Residual) | 286 | 21.5 | 21.8 |
| Black, non-Hispanic, female, 30-34 years | | | | |
| ... | All causes | 1,979 | 100.0 | 142.2 |
| 1 | Human immunodeficiency virus (HIV) disease (B20-B24) | 310 | 15.7 | 22.3 |
| 2 | Malignant neoplasms (C00-C97) | 289 | 14.6 | 20.8 |
| 3 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 243 | 12.3 | 17.5 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 216 | 10.9 | 15.5 |
| 5 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 159 | 8.0 | 11.4 |
| 6 | Cerebrovascular diseases (I60-I69) | 54 | 2.7 | 3.9 |
| 7 | Diabetes mellitus (E10-E14) | 42 | 2.1 | 3.0 |
| 8 | Chronic lower respiratory diseases (J40-J47) | 38 | 1.9 | 2.7 |
| 9 | Septicemia (A40-A41) | 37 | 1.9 | 2.7 |
| 10 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 35 | 1.8 | 2.5 |
| 11 | Pregnancy, childbirth and the puerperium (O00-099) | 30 | 1.5 | 2.2 |
| 12 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 26 | 1.3 | 1.9 |
| 13 | Anemias (D50-D64) | 25 | 1.3 | 1.8 |
| 14 | Influenza and pneumonia (J10-J18) | 21 | 1.1 | 1.5 |
| 15 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 20 | 1.0 | 1.4 |
| ... | All other causes (Residual) | 434 | 21.9 | 31.2 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|-------|
| Black, non-Hispanic, female, 35-39 years | | | | |
| ... | All causes | 3,328 | 100.0 | 223.2 |
| 1 | Malignant neoplasms (C00-C97) | 584 | 17.5 | 39.2 |
| 2 | Human immunodeficiency virus (HIV) disease (B20-B24) | 470 | 14.1 | 31.5 |
| 3 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 447 | 13.4 | 30.0 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 308 | 9.3 | 20.7 |
| 5 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 233 | 7.0 | 15.6 |
| 6 | Cerebrovascular diseases (I60-I69) | 115 | 3.5 | 7.7 |
| 7 | Diabetes mellitus (E10-E14) | 83 | 2.5 | 5.6 |
| 8 | Chronic lower respiratory diseases (J40-J47) | 73 | 2.2 | 4.9 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 69 | 2.1 | 4.6 |
| 10 | Septicemia (A40-A41) | 58 | 1.7 | 3.9 |
| 11 | Influenza and pneumonia (J10-J18) | 47 | 1.4 | 3.2 |
| 12 | Chronic liver disease and cirrhosis (K70,K73-K74) | 40 | 1.2 | 2.7 |
| 13 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 39 | 1.2 | 2.6 |
| 14 | Anemias (D50-D64) | 32 | 1.0 | 2.1 |
| 15 | Pregnancy, childbirth and the puerperium (O00-099) | 28 | 0.8 | 1.9 |
| ... | All other causes (Residual) | 702 | 21.1 | 47.1 |
| Black, non-Hispanic, female, 40-44 years | | | | |
| ... | All causes | 5,137 | 100.0 | 349.0 |
| 1 | Malignant neoplasms (C00-C97) | 1,165 | 22.7 | 79.2 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 906 | 17.6 | 61.6 |
| 3 | Human immunodeficiency virus (HIV) disease (B20-B24) | 562 | 10.9 | 38.2 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 366 | 7.1 | 24.9 |
| 5 | Cerebrovascular diseases (I60-I69) | 288 | 5.6 | 19.6 |
| 6 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 158 | 3.1 | 10.7 |
| 7 | Diabetes mellitus (E10-E14) | 148 | 2.9 | 10.1 |
| 8 | Chronic liver disease and cirrhosis (K70,K73-K74) | 107 | 2.1 | 7.3 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 98 | 1.9 | 6.7 |
| 10 | Septicemia (A40-A41) | 82 | 1.6 | 5.6 |
| 11 | Chronic lower respiratory diseases (J40-J47) | 76 | 1.5 | 5.2 |
| 12 | Influenza and pneumonia (J10-J18) | 72 | 1.4 | 4.9 |
| 13 | Anemias (D50-D64) | 56 | 1.1 | 3.8 |
| 14 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 52 | 1.0 | 3.5 |
| 15 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 50 | 1.0 | 3.4 |
| ... | All other causes (Residual) | 951 | 18.5 | 64.6 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|-------------------|---|--------|-------------------------|-------|
| | Black, non-Hispanic, female, 45-49 years | | | |
| ... | All causes | 6,303 | 100.0 | 493.1 |
| 1 | Malignant neoplasms (C00-C97) | 1,742 | 27.6 | 136.3 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 1,258 | 20.0 | 98.4 |
| 3 | Human immunodeficiency virus (HIV) disease (B20-B24) | 391 | 6.2 | 30.6 |
| 4 | Cerebrovascular diseases (I60-I69) | 378 | 6.0 | 29.6 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 294 | 4.7 | 23.0 |
| 6 | Diabetes mellitus (E10-E14) | 222 | 3.5 | 17.4 |
| 7 | Chronic liver disease and cirrhosis (K70,K73-K74) | 144 | 2.3 | 11.3 |
| 8 | Septicemia (A40-A41) | 128 | 2.0 | 10.0 |
| 9 | Chronic lower respiratory diseases (J40-J47) | 121 | 1.9 | 9.5 |
| 10 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 119 | 1.9 | 9.3 |
| 11 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 100 | 1.6 | 7.8 |
| 12 | Influenza and pneumonia (J10-J18) | 91 | 1.4 | 7.1 |
| 13 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 75 | 1.2 | 5.9 |
| 14 | Viral hepatitis (B15-B19) | 50 | 0.8 | 3.9 |
| 15 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 43 | 0.7 | 3.4 |
| ... | All other causes (Residual) | 1,147 | 18.2 | 89.7 |
| | Black, non-Hispanic, female, 50-54 years | | | |
| ... | All causes | 7,488 | 100.0 | 710.9 |
| 1 | Malignant neoplasms (C00-C97) | 2,237 | 29.9 | 212.4 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 1,743 | 23.3 | 165.5 |
| 3 | Cerebrovascular diseases (I60-I69) | 510 | 6.8 | 48.4 |
| 4 | Diabetes mellitus (E10-E14) | 383 | 5.1 | 36.4 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 235 | 3.1 | 22.3 |
| 6 | Human immunodeficiency virus (HIV) disease (B20-B24) | 229 | 3.1 | 21.7 |
| 7 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 184 | 2.5 | 17.5 |
| 8 | Septicemia (A40-A41) | 163 | 2.2 | 15.5 |
| 9 | Chronic lower respiratory diseases (J40-J47) | 157 | 2.1 | 14.9 |
| 10 | Chronic liver disease and cirrhosis (K70,K73-K74) | 147 | 2.0 | 14.0 |
| 11 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 101 | 1.3 | 9.6 |
| 12 | Influenza and pneumonia (J10-J18) | 69 | 0.9 | 6.6 |
| 13 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 60 | 0.8 | 5.7 |
| 14 | Viral hepatitis (B15-B19) | 59 | 0.8 | 5.6 |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 32 | 0.4 | 3.0 |
| ... | All other causes (Residual) | 1,179 | 15.7 | 111.9 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|---------|
| Black, non-Hispanic, female, 55-59 years | | | | |
| ... | All causes | 7,801 | 100.0 | 1,028.9 |
| 1 | Malignant neoplasms (C00-C97) | 2,561 | 32.8 | 337.8 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 2,008 | 25.7 | 264.8 |
| 3 | Diabetes mellitus (E10-E14) | 491 | 6.3 | 64.8 |
| 4 | Cerebrovascular diseases (I60-I69) | 438 | 5.6 | 57.8 |
| 5 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 219 | 2.8 | 28.9 |
| 6 | Chronic lower respiratory diseases (J40-J47) | 211 | 2.7 | 27.8 |
| 7 | Septicemia (A40-A41) | 170 | 2.2 | 22.4 |
| 8 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 152 | 1.9 | 20.0 |
| 9 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 117 | 1.5 | 15.4 |
| 10 | Chronic liver disease and cirrhosis (K70,K73-K74) | 112 | 1.4 | 14.8 |
| 11 | Human immunodeficiency virus (HIV) disease (B20-B24) | 89 | 1.1 | 11.7 |
| 12 | Influenza and pneumonia (J10-J18) | 84 | 1.1 | 11.1 |
| 13 | Viral hepatitis (B15-B19) | 37 | 0.5 | 4.9 |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 35 | 0.4 | 4.6 |
| 15 | Aortic aneurysm and dissection (I71) | 33 | 0.4 | 4.4 |
| ... | All other causes (Residual) | 1,044 | 13.4 | 137.7 |
| Black, non-Hispanic, female, 60-64 years | | | | |
| ... | All causes | 9,122 | 100.0 | 1,498.8 |
| 1 | Malignant neoplasms (C00-C97) | 2,883 | 31.6 | 473.7 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 2,488 | 27.3 | 408.8 |
| 3 | Diabetes mellitus (E10-E14) | 611 | 6.7 | 100.4 |
| 4 | Cerebrovascular diseases (I60-I69) | 599 | 6.6 | 98.4 |
| 5 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 320 | 3.5 | 52.6 |
| 6 | Chronic lower respiratory diseases (J40-J47) | 254 | 2.8 | 41.7 |
| 7 | Septicemia (A40-A41) | 223 | 2.4 | 36.6 |
| 8 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 152 | 1.7 | 25.0 |
| 9 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 143 | 1.6 | 23.5 |
| 10 | Influenza and pneumonia (J10-J18) | 94 | 1.0 | 15.4 |
| 11 | Human immunodeficiency virus (HIV) disease (B20-B24) | 77 | 0.8 | 12.7 |
| 12 | Chronic liver disease and cirrhosis (K70,K73-K74) | 75 | 0.8 | 12.3 |
| 13 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 39 | 0.4 | 6.4 |
| 14 | Aortic aneurysm and dissection (I71) | 32 | 0.4 | 5.3 |
| 15 | Viral hepatitis (B15-B19) | 31 | 0.3 | 5.1 |
| ... | All other causes (Residual) | 1,101 | 12.1 | 180.9 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|---------|
| Black, non-Hispanic, female, 65-69 years | | | | |
| ... | All causes | 11,486 | 100.0 | 2,209.4 |
| 1 | Malignant neoplasms (C00-C97) | 3,447 | 30.0 | 663.1 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 3,374 | 29.4 | 649.0 |
| 3 | Diabetes mellitus (E10-E14) | 811 | 7.1 | 156.0 |
| 4 | Cerebrovascular diseases (I60-I69) | 786 | 6.8 | 151.2 |
| 5 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 420 | 3.7 | 80.8 |
| 6 | Chronic lower respiratory diseases (J40-J47) | 371 | 3.2 | 71.4 |
| 7 | Septicemia (A40-A41) | 267 | 2.3 | 51.4 |
| 8 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 182 | 1.6 | 35.0 |
| 9 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 147 | 1.3 | 28.3 |
| 10 | Influenza and pneumonia (J10-J18) | 120 | 1.0 | 23.1 |
| 11 | Chronic liver disease and cirrhosis (K70,K73-K74) | 77 | 0.7 | 14.8 |
| 12 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 48 | 0.4 | 9.2 |
| 12 | Aortic aneurysm and dissection (I71) | 48 | 0.4 | 9.2 |
| 14 | Human immunodeficiency virus (HIV) disease (B20-B24) | 47 | 0.4 | 9.0 |
| 15 | Alzheimer's disease (G30) | 42 | 0.4 | 8.1 |
| ... | All other causes (Residual) | 1,299 | 11.3 | 249.9 |
| Black, non-Hispanic, female, 70-74 years | | | | |
| ... | All causes | 13,871 | 100.0 | 3,182.0 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 4,075 | 29.4 | 934.8 |
| 2 | Malignant neoplasms (C00-C97) | 3,811 | 27.5 | 874.2 |
| 3 | Cerebrovascular diseases (I60-I69) | 1,054 | 7.6 | 241.8 |
| 4 | Diabetes mellitus (E10-E14) | 1,021 | 7.4 | 234.2 |
| 5 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 496 | 3.6 | 113.8 |
| 6 | Chronic lower respiratory diseases (J40-J47) | 444 | 3.2 | 101.9 |
| 7 | Septicemia (A40-A41) | 333 | 2.4 | 76.4 |
| 8 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 233 | 1.7 | 53.4 |
| 9 | Influenza and pneumonia (J10-J18) | 204 | 1.5 | 46.8 |
| 10 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 184 | 1.3 | 42.2 |
| 11 | Alzheimer's disease (G30) | 101 | 0.7 | 23.2 |
| 12 | Pneumonitis due to solids and liquids (J69) | 74 | 0.5 | 17.0 |
| 13 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 69 | 0.5 | 15.8 |
| 14 | Chronic liver disease and cirrhosis (K70,K73-K74) | 67 | 0.5 | 15.4 |
| 15 | Aortic aneurysm and dissection (I71) | 58 | 0.4 | 13.3 |
| ... | All other causes (Residual) | 1,647 | 11.9 | 377.8 |

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|---------|
| Black, non-Hispanic, female, 75-79 years | | | | |
| ... | All causes | 16,814 | 100.0 | 4,756.8 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 5,427 | 32.3 | 1,535.3 |
| 2 | Malignant neoplasms (C00-C97) | 3,731 | 22.2 | 1,055.5 |
| 3 | Cerebrovascular diseases (I60-I69) | 1,508 | 9.0 | 426.6 |
| 4 | Diabetes mellitus (E10-E14) | 1,080 | 6.4 | 305.5 |
| 5 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 580 | 3.4 | 164.1 |
| 6 | Chronic lower respiratory diseases (J40-J47) | 475 | 2.8 | 134.4 |
| 7 | Septicemia (A40-A41) | 432 | 2.6 | 122.2 |
| 8 | Influenza and pneumonia (J10-J18) | 327 | 1.9 | 92.5 |
| 9 | Alzheimer's disease (G30) | 317 | 1.9 | 89.7 |
| 10 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 310 | 1.8 | 87.7 |
| 11 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 191 | 1.1 | 54.0 |
| 12 | Aortic aneurysm and dissection (I71) | 105 | 0.6 | 29.7 |
| 13 | Pneumonitis due to solids and liquids (J69) | 101 | 0.6 | 28.6 |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 73 | 0.4 | 20.7 |
| 15 | Atherosclerosis (I70) | 66 | 0.4 | 18.7 |
| ... | All other causes (Residual) | 2,091 | 12.4 | 591.6 |
| Black, non-Hispanic, female, 80-84 years | | | | |
| ... | All causes | 16,695 | 100.0 | 6,952.7 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 5,633 | 33.7 | 2,345.9 |
| 2 | Malignant neoplasms (C00-C97) | 3,023 | 18.1 | 1,258.9 |
| 3 | Cerebrovascular diseases (I60-I69) | 1,681 | 10.1 | 700.1 |
| 4 | Diabetes mellitus (E10-E14) | 902 | 5.4 | 375.6 |
| 5 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 527 | 3.2 | 219.5 |
| 6 | Alzheimer's disease (G30) | 447 | 2.7 | 186.2 |
| 7 | Influenza and pneumonia (J10-J18) | 436 | 2.6 | 181.6 |
| 8 | Chronic lower respiratory diseases (J40-J47) | 430 | 2.6 | 179.1 |
| 9 | Septicemia (A40-A41) | 413 | 2.5 | 172.0 |
| 10 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 347 | 2.1 | 144.5 |
| 11 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 178 | 1.1 | 74.1 |
| 12 | Pneumonitis due to solids and liquids (J69) | 144 | 0.9 | 60.0 |
| 13 | Atherosclerosis (I70) | 95 | 0.6 | 39.6 |
| 14 | Aortic aneurysm and dissection (I71) | 89 | 0.5 | 37.1 |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 82 | 0.5 | 34.1 |
| ... | All other causes (Residual) | 2,268 | 13.6 | 944.5 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|------|
| Black, non-Hispanic, female, 85-89 years | | | | |
| ... | All causes | 15,279 | 100.0 | --- |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 5,527 | 36.2 | --- |
| 2 | Malignant neoplasms (C00-C97) | 1,946 | 12.7 | --- |
| 3 | Cerebrovascular diseases (I60-I69) | 1,665 | 10.9 | --- |
| 4 | Diabetes mellitus (E10-E14) | 739 | 4.8 | --- |
| 5 | Alzheimer's disease (G30) | 563 | 3.7 | --- |
| 6 | Influenza and pneumonia (J10-J18) | 495 | 3.2 | --- |
| 7 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 447 | 2.9 | --- |
| 8 | Septicemia (A40-A41) | 401 | 2.6 | --- |
| 9 | Chronic lower respiratory diseases (J40-J47) | 335 | 2.2 | --- |
| 10 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 332 | 2.2 | --- |
| 11 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 173 | 1.1 | --- |
| 12 | Pneumonitis due to solids and liquids (J69) | 156 | 1.0 | --- |
| 13 | Atherosclerosis (I70) | 105 | 0.7 | --- |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 71 | 0.5 | --- |
| 15 | Aortic aneurysm and dissection (I71) | 69 | 0.5 | --- |
| ... | All other causes (Residual) | 2,255 | 14.8 | --- |
| Black, non-Hispanic, female, 90-94 years | | | | |
| ... | All causes | 10,597 | 100.0 | --- |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 3,980 | 37.6 | --- |
| 2 | Cerebrovascular diseases (I60-I69) | 1,211 | 11.4 | --- |
| 3 | Malignant neoplasms (C00-C97) | 1,046 | 9.9 | --- |
| 4 | Alzheimer's disease (G30) | 413 | 3.9 | --- |
| 5 | Influenza and pneumonia (J10-J18) | 405 | 3.8 | --- |
| 6 | Diabetes mellitus (E10-E14) | 387 | 3.7 | --- |
| 7 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 312 | 2.9 | --- |
| 8 | Septicemia (A40-A41) | 254 | 2.4 | --- |
| 9 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 227 | 2.1 | --- |
| 10 | Chronic lower respiratory diseases (J40-J47) | 206 | 1.9 | --- |
| 11 | Pneumonitis due to solids and liquids (J69) | 134 | 1.3 | --- |
| 12 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 123 | 1.2 | --- |
| 13 | Atherosclerosis (I70) | 102 | 1.0 | --- |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 53 | 0.5 | --- |
| 15 | Aortic aneurysm and dissection (I71) | 51 | 0.5 | --- |
| ... | All other causes (Residual) | 1,693 | 16.0 | --- |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|---|---|--------|-------------------------|------|
| Black, non-Hispanic, female, 95-99 years | | | | |
| ... | All causes | 5,008 | 100.0 | --- |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 2,014 | 40.2 | --- |
| 2 | Cerebrovascular diseases (I60-I69) | 490 | 9.8 | --- |
| 3 | Malignant neoplasms (C00-C97) | 364 | 7.3 | --- |
| 4 | Influenza and pneumonia (J10-J18) | 238 | 4.8 | --- |
| 5 | Alzheimer's disease (G30) | 235 | 4.7 | --- |
| 6 | Diabetes mellitus (E10-E14) | 150 | 3.0 | --- |
| 7 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 137 | 2.7 | --- |
| 8 | Septicemia (A40-A41) | 121 | 2.4 | --- |
| 9 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 99 | 2.0 | --- |
| 10 | Atherosclerosis (I70) | 74 | 1.5 | --- |
| 11 | Pneumonitis due to solids and liquids (J69) | 70 | 1.4 | --- |
| 12 | Chronic lower respiratory diseases (J40-J47) | 66 | 1.3 | --- |
| 13 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 52 | 1.0 | --- |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 26 | 0.5 | --- |
| 15 | Nutritional deficiencies (E40-E64) | 25 | 0.5 | --- |
| ... | All other causes (Residual) | 847 | 16.9 | --- |
| Black, non-Hispanic, female, 100 years and over | | | | |
| ... | All causes | 1,732 | 100.0 | --- |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 725 | 41.9 | --- |
| 2 | Cerebrovascular diseases (I60-I69) | 152 | 8.8 | --- |
| 3 | Influenza and pneumonia (J10-J18) | 108 | 6.2 | --- |
| 4 | Malignant neoplasms (C00-C97) | 79 | 4.6 | --- |
| 5 | Alzheimer's disease (G30) | 69 | 4.0 | --- |
| 6 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 43 | 2.5 | --- |
| 7 | Diabetes mellitus (E10-E14) | 42 | 2.4 | --- |
| 8 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 36 | 2.1 | --- |
| 9 | Septicemia (A40-A41) | 32 | 1.8 | --- |
| 10 | Pneumonitis due to solids and liquids (J69) | 29 | 1.7 | --- |
| 11 | Atherosclerosis (I70) | 25 | 1.4 | --- |
| 12 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 19 | 1.1 | --- |
| 13 | Chronic lower respiratory diseases (J40-J47) | 17 | 1.0 | --- |
| 14 | Anemias (D50-D64) | 15 | 0.9 | --- |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 7 | 0.4 | --- |
| ... | All other causes (Residual) | 334 | 19.3 | --- |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available