

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, all ages				
...	All causes	117,135	100.0	302.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	27,887	23.8	71.9
2	Malignant neoplasms (C00-C97)	23,141	19.8	59.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,106	8.6	26.1
4	Cerebrovascular diseases (I60-I69)	6,451	5.5	16.6
5	Diabetes mellitus (E10-E14)	5,912	5.0	15.3
6	Chronic liver disease and cirrhosis (K70,K73-K74)	3,409	2.9	8.8
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,129	2.7	8.1
8	Chronic lower respiratory diseases (J40-J47)	3,058	2.6	7.9
9	Influenza and pneumonia (J10-J18)	2,824	2.4	7.3
10	Certain conditions originating in the perinatal period (P00-P96)	2,402	2.1	6.2
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,954	1.7	5.0
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,898	1.6	4.9
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,855	1.6	4.8
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,715	1.5	4.4
15	Septicemia (A40-A41)	1,521	1.3	3.9
...	All other causes (Residual)	19,873	17.0	51.3
Hispanic, both sexes, 1-4 years				
...	All causes	933	100.0	29.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	341	36.5	10.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	111	11.9	3.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	81	8.7	2.6
4	Malignant neoplasms (C00-C97)	72	7.7	2.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	24	2.6	0.8
6	Influenza and pneumonia (J10-J18)	17	1.8	*
7	Septicemia (A40-A41)	16	1.7	*
8	Certain conditions originating in the perinatal period (P00-P96)	14	1.5	*
9	Cerebrovascular diseases (I60-I69)	12	1.3	*
10	Chronic lower respiratory diseases (J40-J47)	9	1.0	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	7	0.8	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.8	*
13	Meningitis (G00,G03)	5	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.5	*
15	Anemias (D50-D64)	3	0.3	*
...	All other causes (Residual)	209	22.4	6.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 5-9 years				
...	All causes	538	100.0	14.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	199	37.0	5.3
2	Malignant neoplasms (C00-C97)	103	19.1	2.8
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	40	7.4	1.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	28	5.2	0.8
5	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	16	3.0	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	12	2.2	*
7	Septicemia (A40-A41)	7	1.3	*
8	Influenza and pneumonia (J10-J18)	6	1.1	*
8	Chronic lower respiratory diseases (J40-J47)	6	1.1	*
10	Cerebrovascular diseases (I60-I69)	5	0.9	*
11	Certain conditions originating in the perinatal period (P00-P96)	4	0.7	*
12	Anemias (D50-D64)	3	0.6	*
13	Pneumonitis due to solids and liquids (J69)	2	0.4	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2	0.4	*
...	All other causes (Residual)	103	19.1	2.8
Hispanic, both sexes, 10-14 years				
...	All causes	586	100.0	16.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	190	32.4	5.4
2	Malignant neoplasms (C00-C97)	107	18.3	3.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	36	6.1	1.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	35	6.0	1.0
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	30	5.1	0.8
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	18	3.1	*
7	Cerebrovascular diseases (I60-I69)	15	2.6	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	10	1.7	*
9	Influenza and pneumonia (J10-J18)	8	1.4	*
10	Septicemia (A40-A41)	7	1.2	*
10	Chronic lower respiratory diseases (J40-J47)	7	1.2	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.7	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	3	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.5	*
15	Infections of kidney (N10-N12,N13.6,N15.1)	2	0.3	*
15	Certain conditions originating in the perinatal period (P00-P96)	2	0.3	*
...	All other causes (Residual)	109	18.6	3.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 15-19 years				
...	All causes	2,086	100.0	65.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	984	47.2	30.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	472	22.6	14.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	181	8.7	5.7
4	Malignant neoplasms (C00-C97)	121	5.8	3.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	39	1.9	1.2
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	37	1.8	1.2
7	Cerebrovascular diseases (I60-I69)	10	0.5	*
8	Influenza and pneumonia (J10-J18)	9	0.4	*
9	Septicemia (A40-A41)	7	0.3	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	6	0.3	*
10	Chronic lower respiratory diseases (J40-J47)	6	0.3	*
10	Pregnancy, childbirth and the puerperium (O00-099)	6	0.3	*
13	Diabetes mellitus (E10-E14)	5	0.2	*
13	Legal intervention (Y35,Y89.0)	5	0.2	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.2	*
...	All other causes (Residual)	193	9.3	6.0
Hispanic, both sexes, 20-24 years				
...	All causes	3,169	100.0	87.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,429	45.1	39.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	728	23.0	20.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	274	8.6	7.6
4	Malignant neoplasms (C00-C97)	183	5.8	5.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	89	2.8	2.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	30	0.9	0.8
7	Human immunodeficiency virus (HIV) disease (B20-B24)	23	0.7	0.6
8	Cerebrovascular diseases (I60-I69)	19	0.6	*
9	Legal intervention (Y35,Y89.0)	17	0.5	*
10	Pregnancy, childbirth and the puerperium (O00-099)	13	0.4	*
11	Diabetes mellitus (E10-E14)	10	0.3	*
11	Influenza and pneumonia (J10-J18)	10	0.3	*
13	Chronic lower respiratory diseases (J40-J47)	9	0.3	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.3	*
15	Septicemia (A40-A41)	8	0.3	*
...	All other causes (Residual)	318	10.0	8.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 25-29 years				
...	All causes	2,803	100.0	74.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,082	38.6	28.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	554	19.8	14.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	248	8.8	6.6
4	Malignant neoplasms (C00-C97)	177	6.3	4.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	126	4.5	3.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	80	2.9	2.1
7	Cerebrovascular diseases (I60-I69)	40	1.4	1.1
8	Diabetes mellitus (E10-E14)	32	1.1	0.9
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	23	0.8	0.6
10	Influenza and pneumonia (J10-J18)	18	0.6	*
10	Chronic liver disease and cirrhosis (K70,K73-K74)	18	0.6	*
12	Legal intervention (Y35,Y89.0)	13	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	0.4	*
14	Septicemia (A40-A41)	11	0.4	*
15	Pregnancy, childbirth and the puerperium (O00-099)	10	0.4	*
...	All other causes (Residual)	359	12.8	9.6
Hispanic, both sexes, 30-34 years				
...	All causes	3,013	100.0	84.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	945	31.4	26.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	349	11.6	9.8
3	Malignant neoplasms (C00-C97)	325	10.8	9.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	236	7.8	6.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	195	6.5	5.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	182	6.0	5.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	63	2.1	1.8
8	Cerebrovascular diseases (I60-I69)	53	1.8	1.5
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	30	1.0	0.8
10	Septicemia (A40-A41)	27	0.9	0.8
11	Influenza and pneumonia (J10-J18)	25	0.8	0.7
12	Diabetes mellitus (E10-E14)	24	0.8	0.7
13	Chronic lower respiratory diseases (J40-J47)	22	0.7	0.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21	0.7	0.6
15	Pregnancy, childbirth and the puerperium (O00-099)	18	0.6	*
...	All other causes (Residual)	498	16.5	13.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 35-39 years				
...	All causes	3,644	100.0	115.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	898	24.6	28.5
2	Malignant neoplasms (C00-C97)	500	13.7	15.8
3	Human immunodeficiency virus (HIV) disease (B20-B24)	353	9.7	11.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	313	8.6	9.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	273	7.5	8.7
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	222	6.1	7.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	162	4.4	5.1
8	Cerebrovascular diseases (I60-I69)	94	2.6	3.0
9	Diabetes mellitus (E10-E14)	73	2.0	2.3
10	Influenza and pneumonia (J10-J18)	37	1.0	1.2
11	Viral hepatitis (B15-B19)	30	0.8	1.0
12	Septicemia (A40-A41)	27	0.7	0.9
13	Chronic lower respiratory diseases (J40-J47)	25	0.7	0.8
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	0.6	0.7
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	23	0.6	0.7
...	All other causes (Residual)	591	16.2	18.7
Hispanic, both sexes, 40-44 years				
...	All causes	4,823	100.0	181.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	943	19.6	35.6
2	Malignant neoplasms (C00-C97)	846	17.5	31.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	539	11.2	20.3
4	Human immunodeficiency virus (HIV) disease (B20-B24)	426	8.8	16.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	345	7.2	13.0
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	191	4.0	7.2
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	185	3.8	7.0
8	Cerebrovascular diseases (I60-I69)	162	3.4	6.1
9	Diabetes mellitus (E10-E14)	132	2.7	5.0
10	Viral hepatitis (B15-B19)	71	1.5	2.7
11	Chronic lower respiratory diseases (J40-J47)	45	0.9	1.7
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	42	0.9	1.6
13	Septicemia (A40-A41)	40	0.8	1.5
13	Influenza and pneumonia (J10-J18)	40	0.8	1.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	20	0.4	0.8
...	All other causes (Residual)	796	16.5	30.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 45-49 years				
...	All causes	5,527	100.0	266.6
1	Malignant neoplasms (C00-C97)	1,190	21.5	57.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	851	15.4	41.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	685	12.4	33.0
4	Chronic liver disease and cirrhosis (K70,K73-K74)	478	8.6	23.1
5	Human immunodeficiency virus (HIV) disease (B20-B24)	320	5.8	15.4
6	Cerebrovascular diseases (I60-I69)	269	4.9	13.0
7	Diabetes mellitus (E10-E14)	191	3.5	9.2
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	149	2.7	7.2
9	Viral hepatitis (B15-B19)	142	2.6	6.8
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	129	2.3	6.2
11	Influenza and pneumonia (J10-J18)	61	1.1	2.9
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	57	1.0	2.7
13	Septicemia (A40-A41)	56	1.0	2.7
14	Chronic lower respiratory diseases (J40-J47)	50	0.9	2.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	30	0.5	1.4
...	All other causes (Residual)	869	15.7	41.9
Hispanic, both sexes, 50-54 years				
...	All causes	6,218	100.0	393.2
1	Malignant neoplasms (C00-C97)	1,669	26.8	105.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,169	18.8	73.9
3	Chronic liver disease and cirrhosis (K70,K73-K74)	505	8.1	31.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	483	7.8	30.5
5	Diabetes mellitus (E10-E14)	322	5.2	20.4
6	Cerebrovascular diseases (I60-I69)	291	4.7	18.4
7	Human immunodeficiency virus (HIV) disease (B20-B24)	217	3.5	13.7
8	Viral hepatitis (B15-B19)	149	2.4	9.4
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	111	1.8	7.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	100	1.6	6.3
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	87	1.4	5.5
12	Septicemia (A40-A41)	80	1.3	5.1
13	Chronic lower respiratory diseases (J40-J47)	77	1.2	4.9
14	Influenza and pneumonia (J10-J18)	63	1.0	4.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	24	0.4	1.5
...	All other causes (Residual)	871	14.0	55.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 55-59 years				
...	All causes	6,634	100.0	583.0
1	Malignant neoplasms (C00-C97)	1,971	29.7	173.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,528	23.0	134.3
3	Diabetes mellitus (E10-E14)	432	6.5	38.0
4	Chronic liver disease and cirrhosis (K70,K73-K74)	422	6.4	37.1
5	Cerebrovascular diseases (I60-I69)	321	4.8	28.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	311	4.7	27.3
7	Human immunodeficiency virus (HIV) disease (B20-B24)	123	1.9	10.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	119	1.8	10.5
9	Chronic lower respiratory diseases (J40-J47)	102	1.5	9.0
10	Influenza and pneumonia (J10-J18)	95	1.4	8.3
11	Viral hepatitis (B15-B19)	92	1.4	8.1
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	90	1.4	7.9
13	Septicemia (A40-A41)	71	1.1	6.2
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	56	0.8	4.9
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	50	0.8	4.4
...	All other causes (Residual)	851	12.8	74.8
Hispanic, both sexes, 60-64 years				
...	All causes	7,581	100.0	905.8
1	Malignant neoplasms (C00-C97)	2,276	30.0	271.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,964	25.9	234.7
3	Diabetes mellitus (E10-E14)	597	7.9	71.3
4	Cerebrovascular diseases (I60-I69)	387	5.1	46.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	336	4.4	40.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	255	3.4	30.5
7	Chronic lower respiratory diseases (J40-J47)	172	2.3	20.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	148	2.0	17.7
9	Influenza and pneumonia (J10-J18)	112	1.5	13.4
10	Septicemia (A40-A41)	94	1.2	11.2
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	73	1.0	8.7
12	Viral hepatitis (B15-B19)	72	0.9	8.6
13	Human immunodeficiency virus (HIV) disease (B20-B24)	65	0.9	7.8
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	52	0.7	6.2
15	Aortic aneurysm and dissection (I71)	44	0.6	5.3
...	All other causes (Residual)	934	12.3	111.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 65-69 years				
...	All causes	9,060	100.0	1,378.9
1	Malignant neoplasms (C00-C97)	2,712	29.9	412.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,472	27.3	376.2
3	Diabetes mellitus (E10-E14)	711	7.8	108.2
4	Cerebrovascular diseases (I60-I69)	462	5.1	70.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	332	3.7	50.5
6	Chronic lower respiratory diseases (J40-J47)	252	2.8	38.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	228	2.5	34.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	216	2.4	32.9
9	Influenza and pneumonia (J10-J18)	150	1.7	22.8
10	Septicemia (A40-A41)	132	1.5	20.1
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	66	0.7	10.0
12	Viral hepatitis (B15-B19)	58	0.6	8.8
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	45	0.5	6.8
14	Aortic aneurysm and dissection (I71)	42	0.5	6.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	41	0.5	6.2
...	All other causes (Residual)	1,141	12.6	173.7
Hispanic, both sexes, 70-74 years				
...	All causes	11,416	100.0	2,179.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,247	28.4	620.0
2	Malignant neoplasms (C00-C97)	3,145	27.5	600.5
3	Diabetes mellitus (E10-E14)	880	7.7	168.0
4	Cerebrovascular diseases (I60-I69)	753	6.6	143.8
5	Chronic lower respiratory diseases (J40-J47)	383	3.4	73.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	331	2.9	63.2
7	Influenza and pneumonia (J10-J18)	249	2.2	47.5
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	238	2.1	45.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	211	1.8	40.3
10	Septicemia (A40-A41)	209	1.8	39.9
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	117	1.0	22.3
12	Alzheimer's disease (G30)	88	0.8	16.8
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	64	0.6	12.2
14	Parkinson's disease (G20-G21)	63	0.6	12.0
15	Aortic aneurysm and dissection (I71)	47	0.4	9.0
...	All other causes (Residual)	1,391	12.2	265.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 75-79 years				
...	All causes	12,797	100.0	3,411.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,859	30.2	1,028.8
2	Malignant neoplasms (C00-C97)	3,120	24.4	831.8
3	Diabetes mellitus (E10-E14)	964	7.5	257.0
4	Cerebrovascular diseases (I60-I69)	901	7.0	240.2
5	Chronic lower respiratory diseases (J40-J47)	510	4.0	136.0
6	Influenza and pneumonia (J10-J18)	385	3.0	102.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	281	2.2	74.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	262	2.0	69.9
9	Alzheimer's disease (G30)	211	1.6	56.3
10	Septicemia (A40-A41)	206	1.6	54.9
11	Chronic liver disease and cirrhosis (K70,K73-K74)	198	1.5	52.8
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	130	1.0	34.7
13	Parkinson's disease (G20-G21)	104	0.8	27.7
14	Pneumonitis due to solids and liquids (J69)	73	0.6	19.5
15	Aortic aneurysm and dissection (I71)	72	0.6	19.2
...	All other causes (Residual)	1,521	11.9	405.5
Hispanic, both sexes, 80-84 years				
...	All causes	12,041	100.0	5,365.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,920	32.6	1,746.8
2	Malignant neoplasms (C00-C97)	2,245	18.6	1,000.4
3	Cerebrovascular diseases (I60-I69)	1,017	8.4	453.2
4	Diabetes mellitus (E10-E14)	748	6.2	333.3
5	Chronic lower respiratory diseases (J40-J47)	545	4.5	242.9
6	Influenza and pneumonia (J10-J18)	470	3.9	209.4
7	Alzheimer's disease (G30)	324	2.7	144.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	251	2.1	111.8
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	202	1.7	90.0
10	Septicemia (A40-A41)	176	1.5	78.4
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	153	1.3	68.2
12	Chronic liver disease and cirrhosis (K70,K73-K74)	123	1.0	54.8
13	Parkinson's disease (G20-G21)	119	1.0	53.0
14	Pneumonitis due to solids and liquids (J69)	101	0.8	45.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	83	0.7	37.0
...	All other causes (Residual)	1,564	13.0	696.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 85-89 years				
...	All causes	9,591	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,441	35.9	...
2	Malignant neoplasms (C00-C97)	1,446	15.1	...
3	Cerebrovascular diseases (I60-I69)	838	8.7	...
4	Diabetes mellitus (E10-E14)	457	4.8	...
5	Influenza and pneumonia (J10-J18)	443	4.6	...
6	Chronic lower respiratory diseases (J40-J47)	439	4.6	...
7	Alzheimer's disease (G30)	359	3.7	...
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	179	1.9	...
9	Septicemia (A40-A41)	144	1.5	...
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	130	1.4	...
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	116	1.2	...
12	Pneumonitis due to solids and liquids (J69)	110	1.1	...
13	Parkinson's disease (G20-G21)	96	1.0	...
14	Atherosclerosis (I70)	78	0.8	...
15	Chronic liver disease and cirrhosis (K70,K73-K74)	58	0.6	...
...	All other causes (Residual)	1,257	13.1	...
Hispanic, both sexes, 90-94 years				
...	All causes	6,460	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,548	39.4	...
2	Malignant neoplasms (C00-C97)	707	10.9	...
3	Cerebrovascular diseases (I60-I69)	535	8.3	...
4	Influenza and pneumonia (J10-J18)	358	5.5	...
5	Alzheimer's disease (G30)	297	4.6	...
6	Chronic lower respiratory diseases (J40-J47)	280	4.3	...
7	Diabetes mellitus (E10-E14)	249	3.9	...
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	120	1.9	...
9	Septicemia (A40-A41)	96	1.5	...
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	96	1.5	...
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	92	1.4	...
12	Pneumonitis due to solids and liquids (J69)	86	1.3	...
13	Atherosclerosis (I70)	74	1.1	...
14	Parkinson's disease (G20-G21)	33	0.5	...
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	31	0.5	...
...	All other causes (Residual)	858	13.3	...

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 95-99 years				
...	All causes	2,668	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,197	44.9	...
2	Cerebrovascular diseases (I60-I69)	202	7.6	...
3	Malignant neoplasms (C00-C97)	188	7.0	...
4	Influenza and pneumonia (J10-J18)	177	6.6	...
5	Alzheimer's disease (G30)	118	4.4	...
6	Chronic lower respiratory diseases (J40-J47)	89	3.3	...
7	Diabetes mellitus (E10-E14)	72	2.7	...
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	47	1.8	...
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	45	1.7	...
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	43	1.6	...
11	Septicemia (A40-A41)	41	1.5	...
12	Pneumonitis due to solids and liquids (J69)	35	1.3	...
13	Atherosclerosis (I70)	30	1.1	...
14	Anemias (D50-D64)	14	0.5	...
15	Nutritional deficiencies (E40-E64)	8	0.3	...
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	8	0.3	...
...	All other causes (Residual)	354	13.3	...
Hispanic, both sexes, 100 years and over				
...	All causes	546	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	250	45.8	...
2	Cerebrovascular diseases (I60-I69)	45	8.2	...
3	Influenza and pneumonia (J10-J18)	43	7.9	...
4	Malignant neoplasms (C00-C97)	20	3.7	...
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19	3.5	...
6	Alzheimer's disease (G30)	17	3.1	...
6	Chronic lower respiratory diseases (J40-J47)	17	3.1	...
8	Atherosclerosis (I70)	14	2.6	...
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	13	2.4	...
10	Diabetes mellitus (E10-E14)	12	2.2	...
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7	1.3	...
12	Pneumonitis due to solids and liquids (J69)	6	1.1	...
13	Nutritional deficiencies (E40-E64)	4	0.7	...
14	Parkinson's disease (G20-G21)	2	0.4	...
15	Septicemia (A40-A41)	1	0.2	...
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	1	0.2	...
15	Anemias (D50-D64)	1	0.2	...
15	Aortic aneurysm and dissection (I71)	1	0.2	...
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.2	...
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.2	...
15	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	...
...	All other causes (Residual)	70	12.8	...

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, all ages				
...	All causes	65,703	100.0	328.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,798	22.5	74.0
2	Malignant neoplasms (C00-C97)	12,235	18.6	61.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,698	11.7	38.5
4	Cerebrovascular diseases (I60-I69)	3,003	4.6	15.0
5	Diabetes mellitus (E10-E14)	2,779	4.2	13.9
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,635	4.0	13.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,437	3.7	12.2
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,651	2.5	8.3
9	Chronic lower respiratory diseases (J40-J47)	1,625	2.5	8.1
10	Human immunodeficiency virus (HIV) disease (B20-B24)	1,440	2.2	7.2
11	Influenza and pneumonia (J10-J18)	1,398	2.1	7.0
12	Certain conditions originating in the perinatal period (P00-P96)	1,352	2.1	6.8
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	937	1.4	4.7
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	903	1.4	4.5
15	Septicemia (A40-A41)	744	1.1	3.7
...	All other causes (Residual)	10,068	15.3	50.4
Hispanic, male, 1-4 years				
...	All causes	546	100.0	34.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	215	39.4	13.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	60	11.0	3.8
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	49	9.0	3.1
4	Malignant neoplasms (C00-C97)	35	6.4	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	11	2.0	*
5	Influenza and pneumonia (J10-J18)	11	2.0	*
7	Septicemia (A40-A41)	9	1.6	*
7	Certain conditions originating in the perinatal period (P00-P96)	9	1.6	*
9	Cerebrovascular diseases (I60-I69)	5	0.9	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	4	0.7	*
10	Meningitis (G00,G03)	4	0.7	*
10	Chronic lower respiratory diseases (J40-J47)	4	0.7	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.7	*
14	Anemias (D50-D64)	1	0.2	*
14	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
...	All other causes (Residual)	123	22.5	7.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 5-9 years				
...	All causes	303	100.0	15.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	120	39.6	6.3
2	Malignant neoplasms (C00-C97)	51	16.8	2.7
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	18	5.9	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	15	5.0	*
5	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	7	2.3	*
6	Septicemia (A40-A41)	6	2.0	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	6	2.0	*
8	Chronic lower respiratory diseases (J40-J47)	4	1.3	*
9	Certain conditions originating in the perinatal period (P00-P96)	3	1.0	*
10	Anemias (D50-D64)	2	0.7	*
10	Influenza and pneumonia (J10-J18)	2	0.7	*
12	Meningococcal infection (A39)	1	0.3	*
12	Cerebrovascular diseases (I60-I69)	1	0.3	*
12	Acute bronchitis and bronchiolitis (J20-J21)	1	0.3	*
12	Pneumonitis due to solids and liquids (J69)	1	0.3	*
12	Diseases of appendix (K35-K38)	1	0.3	*
12	Hernia (K40-K46)	1	0.3	*
12	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.3	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1	0.3	*
...	All other causes (Residual)	60	19.8	3.1
Hispanic, male, 10-14 years				
...	All causes	344	100.0	19.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	124	36.0	6.8
2	Malignant neoplasms (C00-C97)	57	16.6	3.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	22	6.4	1.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	20	5.8	1.1
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	17	4.9	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	8	2.3	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	8	2.3	*
8	Cerebrovascular diseases (I60-I69)	6	1.7	*
9	Influenza and pneumonia (J10-J18)	4	1.2	*
9	Chronic lower respiratory diseases (J40-J47)	4	1.2	*
9	Complications of medical and surgical care (Y40-Y84,Y88)	4	1.2	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	3	0.9	*
13	Septicemia (A40-A41)	2	0.6	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.6	*
13	Certain conditions originating in the perinatal period (P00-P96)	2	0.6	*
...	All other causes (Residual)	61	17.7	3.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 15-19 years				
...	All causes	1,614	100.0	97.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	763	47.3	45.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	425	26.3	25.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	151	9.4	9.1
4	Malignant neoplasms (C00-C97)	81	5.0	4.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	26	1.6	1.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	22	1.4	1.3
7	Cerebrovascular diseases (I60-I69)	6	0.4	*
7	Influenza and pneumonia (J10-J18)	6	0.4	*
9	Septicemia (A40-A41)	5	0.3	*
9	Chronic lower respiratory diseases (J40-J47)	5	0.3	*
9	Legal intervention (Y35,Y89.0)	5	0.3	*
12	Diabetes mellitus (E10-E14)	3	0.2	*
12	Meningitis (G00,G03)	3	0.2	*
12	Aortic aneurysm and dissection (I71)	3	0.2	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.2	*
...	All other causes (Residual)	107	6.6	6.4
Hispanic, male, 20-24 years				
...	All causes	2,568	100.0	128.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,195	46.5	60.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	656	25.5	32.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	237	9.2	11.9
4	Malignant neoplasms (C00-C97)	110	4.3	5.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	68	2.6	3.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	18	0.7	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	17	0.7	*
7	Legal intervention (Y35,Y89.0)	17	0.7	*
9	Cerebrovascular diseases (I60-I69)	11	0.4	*
10	Diabetes mellitus (E10-E14)	7	0.3	*
10	Chronic lower respiratory diseases (J40-J47)	7	0.3	*
12	Influenza and pneumonia (J10-J18)	5	0.2	*
12	Chronic liver disease and cirrhosis (K70,K73-K74)	5	0.2	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.2	*
15	Anemias (D50-D64)	4	0.2	*
...	All other causes (Residual)	206	8.0	10.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 25-29 years				
...	All causes	2,239	100.0	108.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	917	41.0	44.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	493	22.0	23.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	222	9.9	10.8
4	Malignant neoplasms (C00-C97)	103	4.6	5.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	98	4.4	4.8
6	Human immunodeficiency virus (HIV) disease (B20-B24)	59	2.6	2.9
7	Cerebrovascular diseases (I60-I69)	29	1.3	1.4
8	Diabetes mellitus (E10-E14)	19	0.8	*
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	14	0.6	*
10	Chronic liver disease and cirrhosis (K70,K73-K74)	13	0.6	*
10	Legal intervention (Y35,Y89.0)	13	0.6	*
12	Influenza and pneumonia (J10-J18)	12	0.5	*
13	Septicemia (A40-A41)	8	0.4	*
13	Chronic lower respiratory diseases (J40-J47)	8	0.4	*
15	Viral hepatitis (B15-B19)	4	0.2	*
...	All other causes (Residual)	227	10.1	11.0
Hispanic, male, 30-34 years				
...	All causes	2,235	100.0	116.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	792	35.4	41.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	296	13.2	15.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	204	9.1	10.6
4	Malignant neoplasms (C00-C97)	149	6.7	7.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	144	6.4	7.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	142	6.4	7.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	49	2.2	2.6
8	Cerebrovascular diseases (I60-I69)	32	1.4	1.7
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	22	1.0	1.1
10	Septicemia (A40-A41)	17	0.8	*
11	Diabetes mellitus (E10-E14)	15	0.7	*
12	Influenza and pneumonia (J10-J18)	14	0.6	*
13	Chronic lower respiratory diseases (J40-J47)	11	0.5	*
14	Viral hepatitis (B15-B19)	10	0.4	*
15	Aortic aneurysm and dissection (I71)	9	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.4	*
...	All other causes (Residual)	320	14.3	16.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 35-39 years				
...	All causes	2,514	100.0	151.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	712	28.3	42.9
2	Human immunodeficiency virus (HIV) disease (B20-B24)	281	11.2	16.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	216	8.6	13.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	215	8.6	12.9
5	Malignant neoplasms (C00-C97)	198	7.9	11.9
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	188	7.5	11.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	127	5.1	7.6
8	Cerebrovascular diseases (I60-I69)	57	2.3	3.4
9	Diabetes mellitus (E10-E14)	41	1.6	2.5
10	Influenza and pneumonia (J10-J18)	23	0.9	1.4
11	Viral hepatitis (B15-B19)	22	0.9	1.3
12	Septicemia (A40-A41)	17	0.7	*
13	Chronic lower respiratory diseases (J40-J47)	12	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	0.5	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	11	0.4	*
...	All other causes (Residual)	382	15.2	23.0
Hispanic, male, 40-44 years				
...	All causes	3,314	100.0	242.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	756	22.8	55.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	410	12.4	30.0
3	Malignant neoplasms (C00-C97)	359	10.8	26.3
4	Human immunodeficiency virus (HIV) disease (B20-B24)	326	9.8	23.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	283	8.5	20.7
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	152	4.6	11.1
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	151	4.6	11.1
8	Cerebrovascular diseases (I60-I69)	97	2.9	7.1
9	Diabetes mellitus (E10-E14)	83	2.5	6.1
10	Viral hepatitis (B15-B19)	53	1.6	3.9
11	Septicemia (A40-A41)	30	0.9	2.2
12	Influenza and pneumonia (J10-J18)	29	0.9	2.1
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	26	0.8	1.9
14	Chronic lower respiratory diseases (J40-J47)	19	0.6	*
15	Aortic aneurysm and dissection (I71)	12	0.4	*
...	All other causes (Residual)	528	15.9	38.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 45-49 years				
...	All causes	3,695	100.0	353.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	621	16.8	59.4
2	Malignant neoplasms (C00-C97)	566	15.3	54.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	522	14.1	50.0
4	Chronic liver disease and cirrhosis (K70,K73-K74)	397	10.7	38.0
5	Human immunodeficiency virus (HIV) disease (B20-B24)	251	6.8	24.0
6	Cerebrovascular diseases (I60-I69)	169	4.6	16.2
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	115	3.1	11.0
8	Viral hepatitis (B15-B19)	113	3.1	10.8
9	Diabetes mellitus (E10-E14)	107	2.9	10.2
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	104	2.8	10.0
11	Influenza and pneumonia (J10-J18)	41	1.1	3.9
12	Septicemia (A40-A41)	40	1.1	3.8
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	29	0.8	2.8
14	Chronic lower respiratory diseases (J40-J47)	28	0.8	2.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	16	0.4	*
...	All other causes (Residual)	576	15.6	55.1
Hispanic, male, 50-54 years				
...	All causes	4,024	100.0	516.8
1	Malignant neoplasms (C00-C97)	864	21.5	111.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	846	21.0	108.7
3	Chronic liver disease and cirrhosis (K70,K73-K74)	408	10.1	52.4
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	366	9.1	47.0
5	Diabetes mellitus (E10-E14)	198	4.9	25.4
6	Cerebrovascular diseases (I60-I69)	171	4.2	22.0
7	Human immunodeficiency virus (HIV) disease (B20-B24)	170	4.2	21.8
8	Viral hepatitis (B15-B19)	105	2.6	13.5
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	91	2.3	11.7
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	62	1.5	8.0
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	59	1.5	7.6
12	Septicemia (A40-A41)	54	1.3	6.9
13	Influenza and pneumonia (J10-J18)	45	1.1	5.8
14	Chronic lower respiratory diseases (J40-J47)	42	1.0	5.4
15	Aortic aneurysm and dissection (I71)	14	0.3	*
...	All other causes (Residual)	529	13.1	67.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 55-59 years				
...	All causes	4,247	100.0	779.4
1	Malignant neoplasms (C00-C97)	1,116	26.3	204.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,087	25.6	199.5
3	Chronic liver disease and cirrhosis (K70,K73-K74)	333	7.8	61.1
4	Diabetes mellitus (E10-E14)	236	5.6	43.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	235	5.5	43.1
6	Cerebrovascular diseases (I60-I69)	196	4.6	36.0
7	Human immunodeficiency virus (HIV) disease (B20-B24)	95	2.2	17.4
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	79	1.9	14.5
9	Viral hepatitis (B15-B19)	65	1.5	11.9
10	Chronic lower respiratory diseases (J40-J47)	64	1.5	11.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	64	1.5	11.7
12	Influenza and pneumonia (J10-J18)	60	1.4	11.0
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	47	1.1	8.6
14	Septicemia (A40-A41)	35	0.8	6.4
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	30	0.7	5.5
...	All other causes (Residual)	505	11.9	92.7
Hispanic, male, 60-64 years				
...	All causes	4,515	100.0	1,158.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,306	28.9	335.0
2	Malignant neoplasms (C00-C97)	1,261	27.9	323.5
3	Diabetes mellitus (E10-E14)	315	7.0	80.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	228	5.0	58.5
5	Cerebrovascular diseases (I60-I69)	225	5.0	57.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	172	3.8	44.1
7	Chronic lower respiratory diseases (J40-J47)	100	2.2	25.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	77	1.7	19.8
9	Influenza and pneumonia (J10-J18)	70	1.6	18.0
10	Septicemia (A40-A41)	49	1.1	12.6
11	Human immunodeficiency virus (HIV) disease (B20-B24)	45	1.0	11.5
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	44	1.0	11.3
13	Viral hepatitis (B15-B19)	43	1.0	11.0
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	36	0.8	9.2
15	Aortic aneurysm and dissection (I71)	30	0.7	7.7
...	All other causes (Residual)	514	11.4	131.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 65-69 years				
...	All causes	5,214	100.0	1,765.5
1	Malignant neoplasms (C00-C97)	1,532	29.4	518.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,511	29.0	511.6
3	Diabetes mellitus (E10-E14)	327	6.3	110.7
4	Cerebrovascular diseases (I60-I69)	256	4.9	86.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	226	4.3	76.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	166	3.2	56.2
7	Chronic lower respiratory diseases (J40-J47)	152	2.9	51.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	115	2.2	38.9
9	Influenza and pneumonia (J10-J18)	88	1.7	29.8
10	Septicemia (A40-A41)	61	1.2	20.7
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	39	0.7	13.2
12	Aortic aneurysm and dissection (I71)	28	0.5	9.5
13	Parkinson's disease (G20-G21)	26	0.5	8.8
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	26	0.5	8.8
15	Human immunodeficiency virus (HIV) disease (B20-B24)	25	0.5	8.5
...	All other causes (Residual)	636	12.2	215.4
Hispanic, male, 70-74 years				
...	All causes	6,262	100.0	2,747.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,930	30.8	846.9
2	Malignant neoplasms (C00-C97)	1,724	27.5	756.5
3	Diabetes mellitus (E10-E14)	425	6.8	186.5
4	Cerebrovascular diseases (I60-I69)	396	6.3	173.8
5	Chronic lower respiratory diseases (J40-J47)	210	3.4	92.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	181	2.9	79.4
7	Influenza and pneumonia (J10-J18)	141	2.3	61.9
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	141	2.3	61.9
9	Septicemia (A40-A41)	100	1.6	43.9
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	99	1.6	43.4
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	54	0.9	23.7
12	Parkinson's disease (G20-G21)	45	0.7	19.7
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	38	0.6	16.7
14	Alzheimer's disease (G30)	37	0.6	16.2
15	Aortic aneurysm and dissection (I71)	35	0.6	15.4
...	All other causes (Residual)	706	11.3	309.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 75-79 years				
...	All causes	6,654	100.0	4,258.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,054	30.9	1,314.4
2	Malignant neoplasms (C00-C97)	1,737	26.1	1,111.5
3	Diabetes mellitus (E10-E14)	442	6.6	282.8
4	Cerebrovascular diseases (I60-I69)	421	6.3	269.4
5	Chronic lower respiratory diseases (J40-J47)	312	4.7	199.7
6	Influenza and pneumonia (J10-J18)	189	2.8	120.9
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	152	2.3	97.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	144	2.2	92.1
9	Septicemia (A40-A41)	96	1.4	61.4
10	Chronic liver disease and cirrhosis (K70,K73-K74)	95	1.4	60.8
11	Alzheimer's disease (G30)	85	1.3	54.4
12	Parkinson's disease (G20-G21)	63	0.9	40.3
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	61	0.9	39.0
14	Pneumonitis due to solids and liquids (J69)	42	0.6	26.9
15	Aortic aneurysm and dissection (I71)	40	0.6	25.6
...	All other causes (Residual)	721	10.8	461.4
Hispanic, male, 80-84 years				
...	All causes	5,637	100.0	6,447.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,844	32.7	2,109.2
2	Malignant neoplasms (C00-C97)	1,165	20.7	1,332.6
3	Cerebrovascular diseases (I60-I69)	404	7.2	462.1
4	Diabetes mellitus (E10-E14)	306	5.4	350.0
5	Chronic lower respiratory diseases (J40-J47)	281	5.0	321.4
6	Influenza and pneumonia (J10-J18)	233	4.1	266.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	129	2.3	147.6
8	Alzheimer's disease (G30)	111	2.0	127.0
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	106	1.9	121.2
10	Septicemia (A40-A41)	80	1.4	91.5
11	Pneumonitis due to solids and liquids (J69)	63	1.1	72.1
12	Chronic liver disease and cirrhosis (K70,K73-K74)	62	1.1	70.9
13	Parkinson's disease (G20-G21)	59	1.0	67.5
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	52	0.9	59.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	46	0.8	52.6
...	All other causes (Residual)	696	12.3	796.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 85-89 years				
...	All causes	3,846	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,372	35.7	...
2	Malignant neoplasms (C00-C97)	700	18.2	...
3	Cerebrovascular diseases (I60-I69)	282	7.3	...
4	Chronic lower respiratory diseases (J40-J47)	201	5.2	...
5	Influenza and pneumonia (J10-J18)	186	4.8	...
6	Diabetes mellitus (E10-E14)	162	4.2	...
7	Alzheimer's disease (G30)	102	2.7	...
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	72	1.9	...
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	66	1.7	...
10	Parkinson's disease (G20-G21)	53	1.4	...
11	Septicemia (A40-A41)	52	1.4	...
12	Pneumonitis due to solids and liquids (J69)	46	1.2	...
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	33	0.9	...
14	Atherosclerosis (I70)	29	0.8	...
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	24	0.6	...
...	All other causes (Residual)	466	12.1	...
Hispanic, male, 90-94 years				
...	All causes	2,226	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	820	36.8	...
2	Malignant neoplasms (C00-C97)	319	14.3	...
3	Cerebrovascular diseases (I60-I69)	177	8.0	...
4	Influenza and pneumonia (J10-J18)	137	6.2	...
5	Chronic lower respiratory diseases (J40-J47)	122	5.5	...
6	Alzheimer's disease (G30)	72	3.2	...
7	Diabetes mellitus (E10-E14)	71	3.2	...
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	43	1.9	...
9	Pneumonitis due to solids and liquids (J69)	41	1.8	...
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	41	1.8	...
11	Septicemia (A40-A41)	32	1.4	...
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	32	1.4	...
13	Atherosclerosis (I70)	23	1.0	...
14	Anemias (D50-D64)	14	0.6	...
14	Parkinson's disease (G20-G21)	14	0.6	...
...	All other causes (Residual)	268	12.0	...

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 95-99 years				
...	All causes	768	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	327	42.6	...
2	Malignant neoplasms (C00-C97)	92	12.0	...
3	Influenza and pneumonia (J10-J18)	55	7.2	...
4	Cerebrovascular diseases (I60-I69)	42	5.5	...
5	Chronic lower respiratory diseases (J40-J47)	34	4.4	...
6	Alzheimer's disease (G30)	25	3.3	...
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21	2.7	...
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	21	2.7	...
9	Diabetes mellitus (E10-E14)	16	2.1	...
10	Septicemia (A40-A41)	13	1.7	...
11	Pneumonitis due to solids and liquids (J69)	11	1.4	...
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	9	1.2	...
13	Anemias (D50-D64)	7	0.9	...
14	Atherosclerosis (I70)	6	0.8	...
15	Nutritional deficiencies (E40-E64)	3	0.4	...
15	Peptic ulcer (K25-K28)	3	0.4	...
...	All other causes (Residual)	83	10.8	...
Hispanic, male, 100 years and over				
...	All causes	137	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	52	38.0	...
2	Influenza and pneumonia (J10-J18)	14	10.2	...
3	Cerebrovascular diseases (I60-I69)	11	8.0	...
4	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	5.1	...
5	Malignant neoplasms (C00-C97)	6	4.4	...
6	Diabetes mellitus (E10-E14)	5	3.6	...
6	Alzheimer's disease (G30)	5	3.6	...
6	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	5	3.6	...
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4	2.9	...
10	Atherosclerosis (I70)	3	2.2	...
10	Chronic lower respiratory diseases (J40-J47)	3	2.2	...
12	Pneumonitis due to solids and liquids (J69)	2	1.5	...
13	Septicemia (A40-A41)	1	0.7	...
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	1	0.7	...
13	Aortic aneurysm and dissection (I71)	1	0.7	...
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.7	...
...	All other causes (Residual)	16	11.7	...

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, all ages				
...	All causes	51,432	100.0	274.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,089	25.4	69.7
2	Malignant neoplasms (C00-C97)	10,906	21.2	58.1
3	Cerebrovascular diseases (I60-I69)	3,448	6.7	18.4
4	Diabetes mellitus (E10-E14)	3,133	6.1	16.7
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,408	4.7	12.8
6	Chronic lower respiratory diseases (J40-J47)	1,433	2.8	7.6
7	Influenza and pneumonia (J10-J18)	1,426	2.8	7.6
8	Certain conditions originating in the perinatal period (P00-P96)	1,050	2.0	5.6
9	Alzheimer's disease (G30)	1,010	2.0	5.4
10	Chronic liver disease and cirrhosis (K70,K73-K74)	972	1.9	5.2
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	961	1.9	5.1
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	812	1.6	4.3
13	Septicemia (A40-A41)	777	1.5	4.1
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	557	1.1	3.0
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	494	1.0	2.6
...	All other causes (Residual)	8,956	17.4	47.7
Hispanic, female, 1-4 years				
...	All causes	387	100.0	25.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	126	32.6	8.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	51	13.2	3.3
3	Malignant neoplasms (C00-C97)	37	9.6	2.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	32	8.3	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	13	3.4	*
6	Septicemia (A40-A41)	7	1.8	*
6	Cerebrovascular diseases (I60-I69)	7	1.8	*
8	Influenza and pneumonia (J10-J18)	6	1.6	*
9	Chronic lower respiratory diseases (J40-J47)	5	1.3	*
9	Certain conditions originating in the perinatal period (P00-P96)	5	1.3	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	1.0	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	3	0.8	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.8	*
14	Anemias (D50-D64)	2	0.5	*
15	Meningitis (G00,G03)	1	0.3	*
15	Acute bronchitis and bronchiolitis (J20-J21)	1	0.3	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.3	*
...	All other causes (Residual)	83	21.4	5.4

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 5-9 years				
...	All causes	235	100.0	12.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	79	33.6	4.3
2	Malignant neoplasms (C00-C97)	52	22.1	2.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	22	9.4	1.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	13	5.5	*
5	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	9	3.8	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	6	2.6	*
7	Cerebrovascular diseases (I60-I69)	4	1.7	*
7	Influenza and pneumonia (J10-J18)	4	1.7	*
9	Chronic lower respiratory diseases (J40-J47)	2	0.9	*
10	Septicemia (A40-A41)	1	0.4	*
10	Anemias (D50-D64)	1	0.4	*
10	Pneumonitis due to solids and liquids (J69)	1	0.4	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.4	*
10	Certain conditions originating in the perinatal period (P00-P96)	1	0.4	*
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1	0.4	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.4	*
...	All other causes (Residual)	37	15.7	2.0
Hispanic, female, 10-14 years				
...	All causes	242	100.0	14.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	66	27.3	3.8
2	Malignant neoplasms (C00-C97)	50	20.7	2.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	15	6.2	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	14	5.8	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	13	5.4	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	10	4.1	*
7	Cerebrovascular diseases (I60-I69)	9	3.7	*
8	Septicemia (A40-A41)	5	2.1	*
9	Influenza and pneumonia (J10-J18)	4	1.7	*
10	Chronic lower respiratory diseases (J40-J47)	3	1.2	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	2	0.8	*
11	Infections of kidney (N10-N12,N13.6,N15.1)	2	0.8	*
13	Acute bronchitis and bronchiolitis (J20-J21)	1	0.4	*
13	Pneumonitis due to solids and liquids (J69)	1	0.4	*
13	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.4	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.4	*
...	All other causes (Residual)	45	18.6	2.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 15-19 years				
...	All causes	472	100.0	30.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	221	46.8	14.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	47	10.0	3.1
3	Malignant neoplasms (C00-C97)	40	8.5	2.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	30	6.4	2.0
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	15	3.2	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	13	2.8	*
7	Pregnancy, childbirth and the puerperium (O00-099)	6	1.3	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	5	1.1	*
9	Cerebrovascular diseases (I60-I69)	4	0.8	*
10	Influenza and pneumonia (J10-J18)	3	0.6	*
11	Meningococcal infection (A39)	2	0.4	*
11	Septicemia (A40-A41)	2	0.4	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	2	0.4	*
11	Anemias (D50-D64)	2	0.4	*
11	Diabetes mellitus (E10-E14)	2	0.4	*
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2	0.4	*
11	Chronic liver disease and cirrhosis (K70,K73-K74)	2	0.4	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
...	All other causes (Residual)	72	15.3	4.7
Hispanic, female, 20-24 years				
...	All causes	601	100.0	37.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	234	38.9	14.5
2	Malignant neoplasms (C00-C97)	73	12.1	4.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	72	12.0	4.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	37	6.2	2.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	21	3.5	1.3
6	Pregnancy, childbirth and the puerperium (O00-099)	13	2.2	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	13	2.2	*
8	Cerebrovascular diseases (I60-I69)	8	1.3	*
9	Septicemia (A40-A41)	5	0.8	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	5	0.8	*
9	Influenza and pneumonia (J10-J18)	5	0.8	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.7	*
13	Diabetes mellitus (E10-E14)	3	0.5	*
14	Nutritional deficiencies (E40-E64)	2	0.3	*
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2	0.3	*
14	Aortic aneurysm and dissection (I71)	2	0.3	*
14	Chronic lower respiratory diseases (J40-J47)	2	0.3	*
14	Diseases of appendix (K35-K38)	2	0.3	*
14	Chronic liver disease and cirrhosis (K70,K73-K74)	2	0.3	*
14	Cholelithiasis and other disorders of gallbladder (K80-K82)	2	0.3	*
...	All other causes (Residual)	94	15.6	5.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 25-29 years				
...	All causes	564	100.0	33.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	165	29.3	9.7
2	Malignant neoplasms (C00-C97)	74	13.1	4.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	61	10.8	3.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	28	5.0	1.7
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	26	4.6	1.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	21	3.7	1.2
7	Diabetes mellitus (E10-E14)	13	2.3	*
8	Cerebrovascular diseases (I60-I69)	11	2.0	*
9	Pregnancy, childbirth and the puerperium (O00-099)	10	1.8	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	1.6	*
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	1.6	*
12	Influenza and pneumonia (J10-J18)	6	1.1	*
13	Chronic liver disease and cirrhosis (K70,K73-K74)	5	0.9	*
14	Septicemia (A40-A41)	3	0.5	*
14	Anemias (D50-D64)	3	0.5	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.5	*
...	All other causes (Residual)	117	20.7	6.9
Hispanic, female, 30-34 years				
...	All causes	778	100.0	46.9
1	Malignant neoplasms (C00-C97)	176	22.6	10.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	153	19.7	9.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	53	6.8	3.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	51	6.6	3.1
5	Human immunodeficiency virus (HIV) disease (B20-B24)	40	5.1	2.4
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	32	4.1	1.9
7	Cerebrovascular diseases (I60-I69)	21	2.7	1.3
8	Pregnancy, childbirth and the puerperium (O00-099)	18	2.3	*
9	Chronic liver disease and cirrhosis (K70,K73-K74)	14	1.8	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	1.5	*
11	Influenza and pneumonia (J10-J18)	11	1.4	*
11	Chronic lower respiratory diseases (J40-J47)	11	1.4	*
13	Septicemia (A40-A41)	10	1.3	*
14	Diabetes mellitus (E10-E14)	9	1.2	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	1.0	*
...	All other causes (Residual)	159	20.4	9.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 35-39 years				
...	All causes	1,130	100.0	75.6
1	Malignant neoplasms (C00-C97)	302	26.7	20.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	186	16.5	12.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	98	8.7	6.6
4	Human immunodeficiency virus (HIV) disease (B20-B24)	72	6.4	4.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	57	5.0	3.8
6	Cerebrovascular diseases (I60-I69)	37	3.3	2.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	35	3.1	2.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	34	3.0	2.3
9	Diabetes mellitus (E10-E14)	32	2.8	2.1
10	Pregnancy, childbirth and the puerperium (O00-099)	16	1.4	*
11	Influenza and pneumonia (J10-J18)	14	1.2	*
12	Chronic lower respiratory diseases (J40-J47)	13	1.2	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	12	1.1	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	1.0	*
15	Septicemia (A40-A41)	10	0.9	*
...	All other causes (Residual)	201	17.8	13.4
Hispanic, female, 40-44 years				
...	All causes	1,509	100.0	117.3
1	Malignant neoplasms (C00-C97)	487	32.3	37.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	187	12.4	14.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	129	8.5	10.0
4	Human immunodeficiency virus (HIV) disease (B20-B24)	100	6.6	7.8
5	Cerebrovascular diseases (I60-I69)	65	4.3	5.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	62	4.1	4.8
7	Diabetes mellitus (E10-E14)	49	3.2	3.8
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	40	2.7	3.1
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	33	2.2	2.6
10	Chronic lower respiratory diseases (J40-J47)	26	1.7	2.0
11	Viral hepatitis (B15-B19)	18	1.2	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	1.1	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	12	0.8	*
14	Influenza and pneumonia (J10-J18)	11	0.7	*
15	Septicemia (A40-A41)	10	0.7	*
...	All other causes (Residual)	264	17.5	20.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 45-49 years				
...	All causes	1,832	100.0	178.1
1	Malignant neoplasms (C00-C97)	624	34.1	60.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	230	12.6	22.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	163	8.9	15.8
4	Cerebrovascular diseases (I60-I69)	100	5.5	9.7
5	Diabetes mellitus (E10-E14)	84	4.6	8.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	81	4.4	7.9
7	Human immunodeficiency virus (HIV) disease (B20-B24)	69	3.8	6.7
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	34	1.9	3.3
9	Viral hepatitis (B15-B19)	29	1.6	2.8
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	28	1.5	2.7
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	25	1.4	2.4
12	Chronic lower respiratory diseases (J40-J47)	22	1.2	2.1
13	Influenza and pneumonia (J10-J18)	20	1.1	1.9
14	Septicemia (A40-A41)	16	0.9	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	14	0.8	*
...	All other causes (Residual)	293	16.0	28.5
Hispanic, female, 50-54 years				
...	All causes	2,194	100.0	273.2
1	Malignant neoplasms (C00-C97)	805	36.7	100.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	323	14.7	40.2
3	Diabetes mellitus (E10-E14)	124	5.7	15.4
4	Cerebrovascular diseases (I60-I69)	120	5.5	14.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	117	5.3	14.6
6	Chronic liver disease and cirrhosis (K70,K73-K74)	97	4.4	12.1
7	Human immunodeficiency virus (HIV) disease (B20-B24)	47	2.1	5.9
8	Viral hepatitis (B15-B19)	44	2.0	5.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	41	1.9	5.1
10	Chronic lower respiratory diseases (J40-J47)	35	1.6	4.4
11	Septicemia (A40-A41)	26	1.2	3.2
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	25	1.1	3.1
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	20	0.9	2.5
14	Influenza and pneumonia (J10-J18)	18	0.8	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	14	0.6	*
...	All other causes (Residual)	338	15.4	42.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 55-59 years				
...	All causes	2,387	100.0	402.6
1	Malignant neoplasms (C00-C97)	855	35.8	144.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	441	18.5	74.4
3	Diabetes mellitus (E10-E14)	196	8.2	33.1
4	Cerebrovascular diseases (I60-I69)	125	5.2	21.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	89	3.7	15.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	76	3.2	12.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	55	2.3	9.3
8	Chronic lower respiratory diseases (J40-J47)	38	1.6	6.4
9	Septicemia (A40-A41)	36	1.5	6.1
10	Influenza and pneumonia (J10-J18)	35	1.5	5.9
11	Human immunodeficiency virus (HIV) disease (B20-B24)	28	1.2	4.7
12	Viral hepatitis (B15-B19)	27	1.1	4.6
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	20	0.8	3.4
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11	0.5	*
15	Aortic aneurysm and dissection (I71)	10	0.4	*
...	All other causes (Residual)	345	14.5	58.2
Hispanic, female, 60-64 years				
...	All causes	3,066	100.0	685.7
1	Malignant neoplasms (C00-C97)	1,015	33.1	227.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	658	21.5	147.2
3	Diabetes mellitus (E10-E14)	282	9.2	63.1
4	Cerebrovascular diseases (I60-I69)	162	5.3	36.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	108	3.5	24.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	83	2.7	18.6
7	Chronic lower respiratory diseases (J40-J47)	72	2.3	16.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	71	2.3	15.9
9	Septicemia (A40-A41)	45	1.5	10.1
10	Influenza and pneumonia (J10-J18)	42	1.4	9.4
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	37	1.2	8.3
12	Viral hepatitis (B15-B19)	29	0.9	6.5
13	Human immunodeficiency virus (HIV) disease (B20-B24)	20	0.7	4.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	19	0.6	*
15	Aortic aneurysm and dissection (I71)	14	0.5	*
...	All other causes (Residual)	409	13.3	91.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 65-69 years				
...	All causes	3,846	100.0	1,063.3
1	Malignant neoplasms (C00-C97)	1,180	30.7	326.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	961	25.0	265.7
3	Diabetes mellitus (E10-E14)	384	10.0	106.2
4	Cerebrovascular diseases (I60-I69)	206	5.4	57.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	106	2.8	29.3
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	101	2.6	27.9
7	Chronic lower respiratory diseases (J40-J47)	100	2.6	27.6
8	Septicemia (A40-A41)	71	1.8	19.6
9	Influenza and pneumonia (J10-J18)	62	1.6	17.1
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	62	1.6	17.1
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	40	1.0	11.1
12	Viral hepatitis (B15-B19)	36	0.9	10.0
13	Alzheimer's disease (G30)	21	0.5	5.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	18	0.5	*
15	Aortic aneurysm and dissection (I71)	14	0.4	*
...	All other causes (Residual)	484	12.6	133.8
Hispanic, female, 70-74 years				
...	All causes	5,154	100.0	1,742.1
1	Malignant neoplasms (C00-C97)	1,421	27.6	480.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,317	25.6	445.1
3	Diabetes mellitus (E10-E14)	455	8.8	153.8
4	Cerebrovascular diseases (I60-I69)	357	6.9	120.7
5	Chronic lower respiratory diseases (J40-J47)	173	3.4	58.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	150	2.9	50.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	112	2.2	37.9
8	Septicemia (A40-A41)	109	2.1	36.8
9	Influenza and pneumonia (J10-J18)	108	2.1	36.5
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	97	1.9	32.8
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	63	1.2	21.3
12	Alzheimer's disease (G30)	51	1.0	17.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	33	0.6	11.2
14	Viral hepatitis (B15-B19)	26	0.5	8.8
15	Pneumonitis due to solids and liquids (J69)	20	0.4	6.8
...	All other causes (Residual)	662	12.8	223.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 75-79 years				
...	All causes	6,143	100.0	2,807.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,805	29.4	824.9
2	Malignant neoplasms (C00-C97)	1,383	22.5	632.0
3	Diabetes mellitus (E10-E14)	522	8.5	238.6
4	Cerebrovascular diseases (I60-I69)	480	7.8	219.4
5	Chronic lower respiratory diseases (J40-J47)	198	3.2	90.5
6	Influenza and pneumonia (J10-J18)	196	3.2	89.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	137	2.2	62.6
8	Alzheimer's disease (G30)	126	2.1	57.6
9	Septicemia (A40-A41)	110	1.8	50.3
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	110	1.8	50.3
11	Chronic liver disease and cirrhosis (K70,K73-K74)	103	1.7	47.1
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	69	1.1	31.5
13	Parkinson's disease (G20-G21)	41	0.7	18.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	34	0.6	15.5
15	Aortic aneurysm and dissection (I71)	32	0.5	14.6
...	All other causes (Residual)	797	13.0	364.2
Hispanic, female, 80-84 years				
...	All causes	6,404	100.0	4,674.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,076	32.4	1,515.4
2	Malignant neoplasms (C00-C97)	1,080	16.9	788.4
3	Cerebrovascular diseases (I60-I69)	613	9.6	447.5
4	Diabetes mellitus (E10-E14)	442	6.9	322.7
5	Chronic lower respiratory diseases (J40-J47)	264	4.1	192.7
6	Influenza and pneumonia (J10-J18)	237	3.7	173.0
7	Alzheimer's disease (G30)	213	3.3	155.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	122	1.9	89.1
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	101	1.6	73.7
10	Septicemia (A40-A41)	96	1.5	70.1
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	96	1.5	70.1
12	Chronic liver disease and cirrhosis (K70,K73-K74)	61	1.0	44.5
13	Parkinson's disease (G20-G21)	60	0.9	43.8
14	Atherosclerosis (I70)	42	0.7	30.7
15	Pneumonitis due to solids and liquids (J69)	38	0.6	27.7
...	All other causes (Residual)	863	13.5	630.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 85-89 years				
...	All causes	5,745	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,069	36.0	...
2	Malignant neoplasms (C00-C97)	746	13.0	...
3	Cerebrovascular diseases (I60-I69)	556	9.7	...
4	Diabetes mellitus (E10-E14)	295	5.1	...
5	Alzheimer's disease (G30)	257	4.5	...
5	Influenza and pneumonia (J10-J18)	257	4.5	...
7	Chronic lower respiratory diseases (J40-J47)	238	4.1	...
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	107	1.9	...
9	Septicemia (A40-A41)	92	1.6	...
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	83	1.4	...
11	Pneumonitis due to solids and liquids (J69)	64	1.1	...
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	64	1.1	...
13	Atherosclerosis (I70)	49	0.9	...
14	Parkinson's disease (G20-G21)	43	0.7	...
15	Chronic liver disease and cirrhosis (K70,K73-K74)	40	0.7	...
...	All other causes (Residual)	785	13.7	...
Hispanic, female, 90-94 years				
...	All causes	4,234	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,728	40.8	...
2	Malignant neoplasms (C00-C97)	388	9.2	...
3	Cerebrovascular diseases (I60-I69)	358	8.5	...
4	Alzheimer's disease (G30)	225	5.3	...
5	Influenza and pneumonia (J10-J18)	221	5.2	...
6	Diabetes mellitus (E10-E14)	178	4.2	...
7	Chronic lower respiratory diseases (J40-J47)	158	3.7	...
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	79	1.9	...
9	Septicemia (A40-A41)	64	1.5	...
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	64	1.5	...
11	Atherosclerosis (I70)	51	1.2	...
12	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	49	1.2	...
13	Pneumonitis due to solids and liquids (J69)	45	1.1	...
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	22	0.5	...
15	Parkinson's disease (G20-G21)	19	0.4	...
...	All other causes (Residual)	585	13.8	...

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 95-99 years				
...	All causes	1,900	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	870	45.8	...
2	Cerebrovascular diseases (I60-I69)	160	8.4	...
3	Influenza and pneumonia (J10-J18)	122	6.4	...
4	Malignant neoplasms (C00-C97)	96	5.1	...
5	Alzheimer's disease (G30)	93	4.9	...
6	Diabetes mellitus (E10-E14)	56	2.9	...
7	Chronic lower respiratory diseases (J40-J47)	55	2.9	...
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	34	1.8	...
9	Septicemia (A40-A41)	28	1.5	...
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	26	1.4	...
11	Atherosclerosis (I70)	24	1.3	...
11	Pneumonitis due to solids and liquids (J69)	24	1.3	...
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	24	1.3	...
14	Anemias (D50-D64)	7	0.4	...
15	Parkinson's disease (G20-G21)	6	0.3	...
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	6	0.3	...
...	All other causes (Residual)	269	14.2	...
Hispanic, female, 100 years and over				
...	All causes	409	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	198	48.4	...
2	Cerebrovascular diseases (I60-I69)	34	8.3	...
3	Influenza and pneumonia (J10-J18)	29	7.1	...
4	Malignant neoplasms (C00-C97)	14	3.4	...
4	Chronic lower respiratory diseases (J40-J47)	14	3.4	...
6	Alzheimer's disease (G30)	12	2.9	...
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	2.9	...
8	Atherosclerosis (I70)	11	2.7	...
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	8	2.0	...
10	Diabetes mellitus (E10-E14)	7	1.7	...
11	Nutritional deficiencies (E40-E64)	4	1.0	...
11	Pneumonitis due to solids and liquids (J69)	4	1.0	...
13	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3	0.7	...
14	Parkinson's disease (G20-G21)	2	0.5	...
15	Anemias (D50-D64)	1	0.2	...
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.2	...
15	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	...
...	All other causes (Residual)	54	13.2	...

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, all ages				
...	All causes	1,981,973	100.0	997.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	577,761	29.2	290.8
2	Malignant neoplasms (C00-C97)	458,754	23.1	230.9
3	Cerebrovascular diseases (I60-I69)	133,118	6.7	67.0
4	Chronic lower respiratory diseases (J40-J47)	112,128	5.7	56.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	80,605	4.1	40.6
6	Influenza and pneumonia (J10-J18)	55,419	2.8	27.9
7	Alzheimer's disease (G30)	53,486	2.7	26.9
8	Diabetes mellitus (E10-E14)	52,463	2.6	26.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	30,669	1.5	15.4
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	26,691	1.3	13.4
11	Septicemia (A40-A41)	25,545	1.3	12.9
12	Chronic liver disease and cirrhosis (K70,K73-K74)	20,353	1.0	10.2
13	Parkinson's disease (G20-G21)	15,626	0.8	7.9
14	Pneumonitis due to solids and liquids (J69)	15,203	0.8	7.7
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	14,617	0.7	7.4
...	All other causes (Residual)	309,535	15.6	155.8
White, non-Hispanic, both sexes, 1-4 years				
...	All causes	2,502	100.0	27.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	865	34.6	9.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	281	11.2	3.0
3	Malignant neoplasms (C00-C97)	236	9.4	2.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	169	6.8	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	91	3.6	1.0
6	Influenza and pneumonia (J10-J18)	46	1.8	0.5
7	Septicemia (A40-A41)	42	1.7	0.5
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	36	1.4	0.4
9	Cerebrovascular diseases (I60-I69)	29	1.2	0.3
10	Chronic lower respiratory diseases (J40-J47)	23	0.9	0.2
11	Certain conditions originating in the perinatal period (P00-P96)	22	0.9	0.2
12	Acute bronchitis and bronchiolitis (J20-J21)	18	0.7	*
13	Anemias (D50-D64)	10	0.4	*
13	Meningitis (G00,G03)	10	0.4	*
15	Meningococcal infection (A39)	8	0.3	*
...	All other causes (Residual)	616	24.6	6.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 5-9 years				
...	All causes	1,643	100.0	13.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	616	37.5	5.1
2	Malignant neoplasms (C00-C97)	331	20.1	2.8
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	117	7.1	1.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	56	3.4	0.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	53	3.2	0.4
6	Septicemia (A40-A41)	24	1.5	0.2
7	Influenza and pneumonia (J10-J18)	23	1.4	0.2
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	20	1.2	0.2
8	Cerebrovascular diseases (I60-I69)	20	1.2	0.2
10	Anemias (D50-D64)	12	0.7	*
11	Chronic lower respiratory diseases (J40-J47)	11	0.7	*
12	Meningitis (G00,G03)	9	0.5	*
13	Acute bronchitis and bronchiolitis (J20-J21)	6	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.3	*
15	Certain conditions originating in the perinatal period (P00-P96)	5	0.3	*
...	All other causes (Residual)	329	20.0	2.7
White, non-Hispanic, both sexes, 10-14 years				
...	All causes	2,388	100.0	18.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	965	40.4	7.3
2	Malignant neoplasms (C00-C97)	311	13.0	2.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	176	7.4	1.3
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	136	5.7	1.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	93	3.9	0.7
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	77	3.2	0.6
7	Influenza and pneumonia (J10-J18)	32	1.3	0.2
8	Septicemia (A40-A41)	31	1.3	0.2
8	Chronic lower respiratory diseases (J40-J47)	31	1.3	0.2
10	Cerebrovascular diseases (I60-I69)	24	1.0	0.2
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	21	0.9	0.2
12	Diabetes mellitus (E10-E14)	15	0.6	*
13	Meningitis (G00,G03)	11	0.5	*
14	Certain conditions originating in the perinatal period (P00-P96)	10	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	10	0.4	*
...	All other causes (Residual)	445	18.6	3.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 15-19 years				
...	All causes	8,552	100.0	65.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,099	59.6	39.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,124	13.1	8.6
3	Malignant neoplasms (C00-C97)	454	5.3	3.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	366	4.3	2.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	240	2.8	1.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	153	1.8	1.2
7	Influenza and pneumonia (J10-J18)	42	0.5	0.3
8	Chronic lower respiratory diseases (J40-J47)	40	0.5	0.3
9	Diabetes mellitus (E10-E14)	32	0.4	0.2
10	Septicemia (A40-A41)	31	0.4	0.2
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	30	0.4	0.2
12	Cerebrovascular diseases (I60-I69)	28	0.3	0.2
13	Meningococcal infection (A39)	14	0.2	*
13	Aortic aneurysm and dissection (I71)	14	0.2	*
15	Certain conditions originating in the perinatal period (P00-P96)	11	0.1	*
...	All other causes (Residual)	874	10.2	6.7
White, non-Hispanic, both sexes, 20-24 years				
...	All causes	10,906	100.0	86.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,565	51.0	44.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,827	16.8	14.5
3	Malignant neoplasms (C00-C97)	643	5.9	5.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	592	5.4	4.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	307	2.8	2.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	162	1.5	1.3
7	Cerebrovascular diseases (I60-I69)	73	0.7	0.6
8	Diabetes mellitus (E10-E14)	63	0.6	0.5
9	Influenza and pneumonia (J10-J18)	61	0.6	0.5
10	Chronic lower respiratory diseases (J40-J47)	54	0.5	0.4
11	Septicemia (A40-A41)	41	0.4	0.3
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	29	0.3	0.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	0.2	0.2
14	Legal intervention (Y35,Y89.0)	19	0.2	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	17	0.2	*
15	Pneumonitis due to solids and liquids (J69)	17	0.2	*
...	All other causes (Residual)	1,413	13.0	11.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 25-29 years				
...	All causes	9,900	100.0	86.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,921	39.6	34.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,771	17.9	15.5
3	Malignant neoplasms (C00-C97)	795	8.0	6.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	554	5.6	4.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	464	4.7	4.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	146	1.5	1.3
7	Diabetes mellitus (E10-E14)	120	1.2	1.0
8	Human immunodeficiency virus (HIV) disease (B20-B24)	105	1.1	0.9
9	Cerebrovascular diseases (I60-I69)	85	0.9	0.7
10	Influenza and pneumonia (J10-J18)	83	0.8	0.7
11	Septicemia (A40-A41)	63	0.6	0.6
12	Chronic lower respiratory diseases (J40-J47)	59	0.6	0.5
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	44	0.4	0.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	37	0.4	0.3
14	Chronic liver disease and cirrhosis (K70,K73-K74)	37	0.4	0.3
...	All other causes (Residual)	1,616	16.3	14.1
White, non-Hispanic, both sexes, 30-34 years				
...	All causes	13,858	100.0	103.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,423	31.9	33.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,038	14.7	15.3
3	Malignant neoplasms (C00-C97)	1,688	12.2	12.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,164	8.4	8.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	513	3.7	3.8
6	Human immunodeficiency virus (HIV) disease (B20-B24)	324	2.3	2.4
7	Diabetes mellitus (E10-E14)	262	1.9	2.0
8	Cerebrovascular diseases (I60-I69)	200	1.4	1.5
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	185	1.3	1.4
10	Chronic liver disease and cirrhosis (K70,K73-K74)	167	1.2	1.3
11	Influenza and pneumonia (J10-J18)	124	0.9	0.9
12	Septicemia (A40-A41)	102	0.7	0.8
13	Chronic lower respiratory diseases (J40-J47)	98	0.7	0.7
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	75	0.5	0.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	68	0.5	0.5
...	All other causes (Residual)	2,427	17.5	18.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 35-39 years				
...	All causes	22,230	100.0	151.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,358	24.1	36.4
2	Malignant neoplasms (C00-C97)	3,509	15.8	23.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,818	12.7	19.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,570	11.6	17.5
5	Human immunodeficiency virus (HIV) disease (B20-B24)	842	3.8	5.7
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	618	2.8	4.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	603	2.7	4.1
8	Diabetes mellitus (E10-E14)	474	2.1	3.2
9	Cerebrovascular diseases (I60-I69)	412	1.9	2.8
10	Influenza and pneumonia (J10-J18)	219	1.0	1.5
11	Chronic lower respiratory diseases (J40-J47)	201	0.9	1.4
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	191	0.9	1.3
13	Septicemia (A40-A41)	184	0.8	1.3
14	Viral hepatitis (B15-B19)	147	0.7	1.0
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	127	0.6	0.9
...	All other causes (Residual)	3,957	17.8	26.9
White, non-Hispanic, both sexes, 40-44 years				
...	All causes	37,299	100.0	228.1
1	Malignant neoplasms (C00-C97)	7,725	20.7	47.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,677	17.9	40.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,296	16.9	38.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,225	8.6	19.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,545	4.1	9.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	948	2.5	5.8
7	Cerebrovascular diseases (I60-I69)	876	2.3	5.4
8	Diabetes mellitus (E10-E14)	871	2.3	5.3
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	650	1.7	4.0
10	Chronic lower respiratory diseases (J40-J47)	423	1.1	2.6
11	Influenza and pneumonia (J10-J18)	398	1.1	2.4
12	Viral hepatitis (B15-B19)	349	0.9	2.1
13	Septicemia (A40-A41)	326	0.9	2.0
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	232	0.6	1.4
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	226	0.6	1.4
...	All other causes (Residual)	6,532	17.5	39.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 45-49 years				
...	All causes	52,044	100.0	331.7
1	Malignant neoplasms (C00-C97)	13,970	26.8	89.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,407	20.0	66.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,080	11.7	38.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,084	5.9	19.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,353	4.5	15.0
6	Cerebrovascular diseases (I60-I69)	1,407	2.7	9.0
7	Diabetes mellitus (E10-E14)	1,289	2.5	8.2
8	Chronic lower respiratory diseases (J40-J47)	917	1.8	5.8
9	Human immunodeficiency virus (HIV) disease (B20-B24)	864	1.7	5.5
10	Viral hepatitis (B15-B19)	820	1.6	5.2
11	Influenza and pneumonia (J10-J18)	611	1.2	3.9
12	Septicemia (A40-A41)	572	1.1	3.6
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	533	1.0	3.4
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	403	0.8	2.6
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	270	0.5	1.7
...	All other causes (Residual)	8,464	16.3	53.9
White, non-Hispanic, both sexes, 50-54 years				
...	All causes	67,768	100.0	475.5
1	Malignant neoplasms (C00-C97)	22,639	33.4	158.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,799	23.3	110.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,779	7.1	33.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	2,670	3.9	18.7
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,519	3.7	17.7
6	Diabetes mellitus (E10-E14)	2,096	3.1	14.7
7	Cerebrovascular diseases (I60-I69)	1,913	2.8	13.4
8	Chronic lower respiratory diseases (J40-J47)	1,769	2.6	12.4
9	Viral hepatitis (B15-B19)	770	1.1	5.4
10	Influenza and pneumonia (J10-J18)	718	1.1	5.0
11	Septicemia (A40-A41)	702	1.0	4.9
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	519	0.8	3.6
13	Human immunodeficiency virus (HIV) disease (B20-B24)	502	0.7	3.5
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	365	0.5	2.6
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	292	0.4	2.0
...	All other causes (Residual)	9,716	14.3	68.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 55-59 years				
...	All causes	85,916	100.0	731.3
1	Malignant neoplasms (C00-C97)	32,418	37.7	276.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	21,225	24.7	180.7
3	Chronic lower respiratory diseases (J40-J47)	3,600	4.2	30.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,459	4.0	29.4
5	Diabetes mellitus (E10-E14)	2,896	3.4	24.7
6	Cerebrovascular diseases (I60-I69)	2,733	3.2	23.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,364	2.8	20.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,965	2.3	16.7
9	Septicemia (A40-A41)	996	1.2	8.5
10	Influenza and pneumonia (J10-J18)	887	1.0	7.6
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	822	1.0	7.0
12	Aortic aneurysm and dissection (I71)	451	0.5	3.8
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	449	0.5	3.8
14	Viral hepatitis (B15-B19)	388	0.5	3.3
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	339	0.4	2.9
...	All other causes (Residual)	10,924	12.7	93.0
White, non-Hispanic, both sexes, 60-64 years				
...	All causes	104,927	100.0	1,145.4
1	Malignant neoplasms (C00-C97)	41,391	39.4	451.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	26,269	25.0	286.8
3	Chronic lower respiratory diseases (J40-J47)	6,157	5.9	67.2
4	Cerebrovascular diseases (I60-I69)	3,699	3.5	40.4
5	Diabetes mellitus (E10-E14)	3,507	3.3	38.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,934	2.8	32.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,119	2.0	23.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,289	1.2	14.1
9	Influenza and pneumonia (J10-J18)	1,265	1.2	13.8
10	Septicemia (A40-A41)	1,245	1.2	13.6
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,156	1.1	12.6
12	Aortic aneurysm and dissection (I71)	705	0.7	7.7
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	513	0.5	5.6
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	466	0.4	5.1
15	Alzheimer's disease (G30)	324	0.3	3.5
...	All other causes (Residual)	11,888	11.3	129.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 65-69 years				
...	All causes	137,191	100.0	1,800.0
1	Malignant neoplasms (C00-C97)	51,562	37.6	676.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	35,099	25.6	460.5
3	Chronic lower respiratory diseases (J40-J47)	10,170	7.4	133.4
4	Cerebrovascular diseases (I60-I69)	5,654	4.1	74.2
5	Diabetes mellitus (E10-E14)	4,717	3.4	61.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,840	2.1	37.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,047	1.5	26.9
8	Influenza and pneumonia (J10-J18)	1,866	1.4	24.5
9	Septicemia (A40-A41)	1,805	1.3	23.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,794	1.3	23.5
11	Aortic aneurysm and dissection (I71)	1,216	0.9	16.0
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,074	0.8	14.1
13	Alzheimer's disease (G30)	804	0.6	10.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	772	0.6	10.1
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	675	0.5	8.9
...	All other causes (Residual)	15,096	11.0	198.1
White, non-Hispanic, both sexes, 70-74 years				
...	All causes	201,770	100.0	2,825.5
1	Malignant neoplasms (C00-C97)	67,902	33.7	950.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	53,429	26.5	748.2
3	Chronic lower respiratory diseases (J40-J47)	16,638	8.2	233.0
4	Cerebrovascular diseases (I60-I69)	10,701	5.3	149.9
5	Diabetes mellitus (E10-E14)	6,620	3.3	92.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,641	1.8	51.0
7	Influenza and pneumonia (J10-J18)	3,553	1.8	49.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,076	1.5	43.1
9	Septicemia (A40-A41)	2,778	1.4	38.9
10	Alzheimer's disease (G30)	2,345	1.2	32.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	2,158	1.1	30.2
12	Aortic aneurysm and dissection (I71)	1,841	0.9	25.8
13	Parkinson's disease (G20-G21)	1,429	0.7	20.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	1,213	0.6	17.0
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,163	0.6	16.3
...	All other causes (Residual)	23,283	11.5	326.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 75-79 years				
...	All causes	278,144	100.0	4,446.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	78,991	28.4	1,262.7
2	Malignant neoplasms (C00-C97)	76,462	27.5	1,222.3
3	Chronic lower respiratory diseases (J40-J47)	22,378	8.0	357.7
4	Cerebrovascular diseases (I60-I69)	18,821	6.8	300.9
5	Diabetes mellitus (E10-E14)	8,632	3.1	138.0
6	Influenza and pneumonia (J10-J18)	6,568	2.4	105.0
7	Alzheimer's disease (G30)	6,332	2.3	101.2
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,082	1.8	81.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,879	1.8	78.0
10	Septicemia (A40-A41)	4,043	1.5	64.6
11	Parkinson's disease (G20-G21)	3,084	1.1	49.3
12	Aortic aneurysm and dissection (I71)	2,499	0.9	39.9
13	Chronic liver disease and cirrhosis (K70,K73-K74)	2,061	0.7	32.9
14	Pneumonitis due to solids and liquids (J69)	1,969	0.7	31.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	1,890	0.7	30.2
...	All other causes (Residual)	34,453	12.4	550.7
White, non-Hispanic, both sexes, 80-84 years				
...	All causes	328,287	100.0	7,166.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	102,984	31.4	2,248.0
2	Malignant neoplasms (C00-C97)	67,495	20.6	1,473.3
3	Cerebrovascular diseases (I60-I69)	27,506	8.4	600.4
4	Chronic lower respiratory diseases (J40-J47)	22,739	6.9	496.4
5	Alzheimer's disease (G30)	11,895	3.6	259.6
6	Influenza and pneumonia (J10-J18)	10,433	3.2	227.7
7	Diabetes mellitus (E10-E14)	9,060	2.8	197.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,234	1.9	136.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,231	1.9	136.0
10	Septicemia (A40-A41)	4,763	1.5	104.0
11	Parkinson's disease (G20-G21)	4,461	1.4	97.4
12	Pneumonitis due to solids and liquids (J69)	3,162	1.0	69.0
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,735	0.8	59.7
14	Aortic aneurysm and dissection (I71)	2,489	0.8	54.3
15	Atherosclerosis (I70)	2,236	0.7	48.8
...	All other causes (Residual)	43,864	13.4	957.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 85-89 years				
...	All causes	306,287	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	106,117	34.6	...
2	Malignant neoplasms (C00-C97)	44,205	14.4	...
3	Cerebrovascular diseases (I60-I69)	29,412	9.6	...
4	Chronic lower respiratory diseases (J40-J47)	16,549	5.4	...
5	Alzheimer's disease (G30)	14,914	4.9	...
6	Influenza and pneumonia (J10-J18)	12,381	4.0	...
7	Diabetes mellitus (E10-E14)	7,066	2.3	...
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,977	2.0	...
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,901	1.9	...
10	Septicemia (A40-A41)	4,168	1.4	...
11	Parkinson's disease (G20-G21)	3,695	1.2	...
12	Pneumonitis due to solids and liquids (J69)	3,664	1.2	...
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	3,077	1.0	...
14	Atherosclerosis (I70)	2,828	0.9	...
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	1,869	0.6	...
...	All other causes (Residual)	44,464	14.5	...
White, non-Hispanic, both sexes, 90-94 years				
...	All causes	203,219	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	77,340	38.1	...
2	Cerebrovascular diseases (I60-I69)	20,578	10.1	...
3	Malignant neoplasms (C00-C97)	19,422	9.6	...
4	Alzheimer's disease (G30)	11,241	5.5	...
5	Influenza and pneumonia (J10-J18)	10,228	5.0	...
6	Chronic lower respiratory diseases (J40-J47)	7,760	3.8	...
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,894	1.9	...
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,787	1.9	...
9	Diabetes mellitus (E10-E14)	3,536	1.7	...
10	Pneumonitis due to solids and liquids (J69)	2,686	1.3	...
11	Atherosclerosis (I70)	2,560	1.3	...
12	Septicemia (A40-A41)	2,466	1.2	...
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,404	1.2	...
14	Parkinson's disease (G20-G21)	1,672	0.8	...
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	1,043	0.5	...
...	All other causes (Residual)	32,602	16.0	...

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 95-99 years				
...	All causes	77,698	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	31,405	40.4	...
2	Cerebrovascular diseases (I60-I69)	7,553	9.7	...
3	Malignant neoplasms (C00-C97)	4,915	6.3	...
4	Alzheimer's disease (G30)	4,684	6.0	...
5	Influenza and pneumonia (J10-J18)	4,585	5.9	...
6	Chronic lower respiratory diseases (J40-J47)	2,169	2.8	...
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,448	1.9	...
8	Atherosclerosis (I70)	1,301	1.7	...
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,280	1.6	...
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,075	1.4	...
11	Pneumonitis due to solids and liquids (J69)	1,067	1.4	...
12	Diabetes mellitus (E10-E14)	1,064	1.4	...
13	Septicemia (A40-A41)	883	1.1	...
14	Parkinson's disease (G20-G21)	408	0.5	...
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	319	0.4	...
...	All other causes (Residual)	13,542	17.4	...
White, non-Hispanic, both sexes, 100 years and over				
...	All causes	15,859	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,808	42.9	...
2	Cerebrovascular diseases (I60-I69)	1,341	8.5	...
3	Influenza and pneumonia (J10-J18)	1,179	7.4	...
4	Alzheimer's disease (G30)	766	4.8	...
5	Malignant neoplasms (C00-C97)	642	4.0	...
6	Atherosclerosis (I70)	357	2.3	...
7	Chronic lower respiratory diseases (J40-J47)	318	2.0	...
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	260	1.6	...
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	234	1.5	...
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	224	1.4	...
11	Pneumonitis due to solids and liquids (J69)	199	1.3	...
12	Septicemia (A40-A41)	149	0.9	...
13	Diabetes mellitus (E10-E14)	135	0.9	...
14	Nutritional deficiencies (E40-E64)	73	0.5	...
15	Anemias (D50-D64)	72	0.5	...
...	All other causes (Residual)	3,102	19.6	...

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, all ages				
...	All causes	957,645	100.0	983.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	281,442	29.4	289.2
2	Malignant neoplasms (C00-C97)	237,323	24.8	243.8
3	Chronic lower respiratory diseases (J40-J47)	53,662	5.6	55.1
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	50,652	5.3	52.0
5	Cerebrovascular diseases (I60-I69)	49,908	5.2	51.3
6	Diabetes mellitus (E10-E14)	25,273	2.6	26.0
7	Influenza and pneumonia (J10-J18)	23,934	2.5	24.6
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	21,323	2.2	21.9
9	Alzheimer's disease (G30)	15,399	1.6	15.8
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14,883	1.6	15.3
11	Chronic liver disease and cirrhosis (K70,K73-K74)	12,800	1.3	13.2
12	Septicemia (A40-A41)	11,148	1.2	11.5
13	Parkinson's disease (G20-G21)	8,838	0.9	9.1
14	Aortic aneurysm and dissection (I71)	7,834	0.8	8.0
15	Pneumonitis due to solids and liquids (J69)	7,654	0.8	7.9
...	All other causes (Residual)	135,572	14.2	139.3
White, non-Hispanic, male, 1-4 years				
...	All causes	1,431	100.0	30.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	527	36.8	11.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	151	10.6	3.2
3	Malignant neoplasms (C00-C97)	134	9.4	2.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	89	6.2	1.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	54	3.8	1.1
6	Septicemia (A40-A41)	25	1.7	0.5
7	Influenza and pneumonia (J10-J18)	24	1.7	0.5
8	Cerebrovascular diseases (I60-I69)	18	1.3	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	17	1.2	*
9	Chronic lower respiratory diseases (J40-J47)	17	1.2	*
11	Certain conditions originating in the perinatal period (P00-P96)	10	0.7	*
12	Meningococcal infection (A39)	6	0.4	*
12	Meningitis (G00,G03)	6	0.4	*
12	Acute bronchitis and bronchiolitis (J20-J21)	6	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.3	*
...	All other causes (Residual)	342	23.9	7.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 5-9 years				
...	All causes	923	100.0	14.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	378	41.0	6.1
2	Malignant neoplasms (C00-C97)	182	19.7	2.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	58	6.3	0.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	24	2.6	0.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	21	2.3	0.3
6	Septicemia (A40-A41)	15	1.6	*
7	Cerebrovascular diseases (I60-I69)	10	1.1	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	9	1.0	*
9	Influenza and pneumonia (J10-J18)	8	0.9	*
9	Chronic lower respiratory diseases (J40-J47)	8	0.9	*
11	Acute bronchitis and bronchiolitis (J20-J21)	6	0.7	*
12	Meningitis (G00,G03)	5	0.5	*
13	Meningococcal infection (A39)	4	0.4	*
13	Anemias (D50-D64)	4	0.4	*
13	Diabetes mellitus (E10-E14)	4	0.4	*
13	Certain conditions originating in the perinatal period (P00-P96)	4	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.4	*
...	All other causes (Residual)	179	19.4	2.9
White, non-Hispanic, male, 10-14 years				
...	All causes	1,442	100.0	21.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	619	42.9	9.2
2	Malignant neoplasms (C00-C97)	173	12.0	2.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	137	9.5	2.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	67	4.6	1.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	61	4.2	0.9
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	40	2.8	0.6
7	Septicemia (A40-A41)	19	1.3	*
8	Chronic lower respiratory diseases (J40-J47)	15	1.0	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	13	0.9	*
9	Cerebrovascular diseases (I60-I69)	13	0.9	*
11	Diabetes mellitus (E10-E14)	12	0.8	*
11	Influenza and pneumonia (J10-J18)	12	0.8	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	8	0.6	*
14	Meningitis (G00,G03)	6	0.4	*
15	Pneumonitis due to solids and liquids (J69)	5	0.3	*
...	All other causes (Residual)	242	16.8	3.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 15-19 years				
...	All causes	5,884	100.0	87.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,466	58.9	51.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	954	16.2	14.2
3	Malignant neoplasms (C00-C97)	264	4.5	3.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	263	4.5	3.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	163	2.8	2.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	90	1.5	1.3
7	Chronic lower respiratory diseases (J40-J47)	25	0.4	0.4
8	Influenza and pneumonia (J10-J18)	24	0.4	0.4
9	Cerebrovascular diseases (I60-I69)	17	0.3	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	16	0.3	*
10	Diabetes mellitus (E10-E14)	16	0.3	*
12	Aortic aneurysm and dissection (I71)	12	0.2	*
13	Septicemia (A40-A41)	11	0.2	*
14	Legal intervention (Y35,Y89.0)	8	0.1	*
15	Meningococcal infection (A39)	7	0.1	*
...	All other causes (Residual)	548	9.3	8.2
White, non-Hispanic, male, 20-24 years				
...	All causes	8,073	100.0	126.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,280	53.0	67.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,573	19.5	24.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	422	5.2	6.6
4	Malignant neoplasms (C00-C97)	375	4.6	5.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	207	2.6	3.2
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	88	1.1	1.4
7	Cerebrovascular diseases (I60-I69)	45	0.6	0.7
8	Influenza and pneumonia (J10-J18)	38	0.5	0.6
9	Diabetes mellitus (E10-E14)	35	0.4	0.5
10	Septicemia (A40-A41)	26	0.3	0.4
11	Chronic lower respiratory diseases (J40-J47)	23	0.3	0.4
12	Legal intervention (Y35,Y89.0)	17	0.2	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	16	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	0.1	*
15	Pneumonitis due to solids and liquids (J69)	9	0.1	*
...	All other causes (Residual)	907	11.2	14.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 25-29 years				
...	All causes	6,988	100.0	121.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,992	42.8	51.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,427	20.4	24.7
3	Malignant neoplasms (C00-C97)	427	6.1	7.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	369	5.3	6.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	323	4.6	5.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	90	1.3	1.6
7	Human immunodeficiency virus (HIV) disease (B20-B24)	77	1.1	1.3
8	Diabetes mellitus (E10-E14)	69	1.0	1.2
9	Cerebrovascular diseases (I60-I69)	47	0.7	0.8
10	Influenza and pneumonia (J10-J18)	38	0.5	0.7
11	Septicemia (A40-A41)	37	0.5	0.6
12	Chronic lower respiratory diseases (J40-J47)	29	0.4	0.5
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	21	0.3	0.4
13	Chronic liver disease and cirrhosis (K70,K73-K74)	21	0.3	0.4
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20	0.3	0.3
...	All other causes (Residual)	1,001	14.3	17.4
White, non-Hispanic, male, 30-34 years				
...	All causes	9,351	100.0	139.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,281	35.1	48.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,638	17.5	24.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	808	8.6	12.0
4	Malignant neoplasms (C00-C97)	791	8.5	11.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	337	3.6	5.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	258	2.8	3.8
7	Diabetes mellitus (E10-E14)	162	1.7	2.4
8	Chronic liver disease and cirrhosis (K70,K73-K74)	109	1.2	1.6
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	105	1.1	1.6
10	Cerebrovascular diseases (I60-I69)	89	1.0	1.3
11	Influenza and pneumonia (J10-J18)	73	0.8	1.1
12	Chronic lower respiratory diseases (J40-J47)	52	0.6	0.8
13	Septicemia (A40-A41)	49	0.5	0.7
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	44	0.5	0.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	41	0.4	0.6
...	All other causes (Residual)	1,514	16.2	22.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 35-39 years				
...	All causes	14,299	100.0	194.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,822	26.7	51.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,013	14.1	27.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,984	13.9	26.9
4	Malignant neoplasms (C00-C97)	1,511	10.6	20.5
5	Human immunodeficiency virus (HIV) disease (B20-B24)	710	5.0	9.6
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	411	2.9	5.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	376	2.6	5.1
8	Diabetes mellitus (E10-E14)	284	2.0	3.9
9	Cerebrovascular diseases (I60-I69)	195	1.4	2.6
10	Influenza and pneumonia (J10-J18)	113	0.8	1.5
11	Viral hepatitis (B15-B19)	110	0.8	1.5
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	96	0.7	1.3
13	Septicemia (A40-A41)	87	0.6	1.2
14	Chronic lower respiratory diseases (J40-J47)	84	0.6	1.1
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	70	0.5	0.9
...	All other causes (Residual)	2,433	17.0	33.0
White, non-Hispanic, male, 40-44 years				
...	All causes	23,750	100.0	290.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,728	19.9	57.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,673	19.7	57.2
3	Malignant neoplasms (C00-C97)	3,530	14.9	43.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,449	10.3	30.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	995	4.2	12.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	807	3.4	9.9
7	Diabetes mellitus (E10-E14)	565	2.4	6.9
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	463	1.9	5.7
9	Cerebrovascular diseases (I60-I69)	422	1.8	5.2
10	Viral hepatitis (B15-B19)	241	1.0	3.0
11	Influenza and pneumonia (J10-J18)	227	1.0	2.8
12	Chronic lower respiratory diseases (J40-J47)	220	0.9	2.7
13	Septicemia (A40-A41)	162	0.7	2.0
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	127	0.5	1.6
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	121	0.5	1.5
...	All other causes (Residual)	4,020	16.9	49.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 45-49 years				
...	All causes	33,102	100.0	424.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,866	23.8	100.8
2	Malignant neoplasms (C00-C97)	6,912	20.9	88.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,412	13.3	56.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,296	6.9	29.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,682	5.1	21.6
6	Diabetes mellitus (E10-E14)	792	2.4	10.2
7	Cerebrovascular diseases (I60-I69)	765	2.3	9.8
8	Human immunodeficiency virus (HIV) disease (B20-B24)	739	2.2	9.5
9	Viral hepatitis (B15-B19)	613	1.9	7.9
10	Chronic lower respiratory diseases (J40-J47)	459	1.4	5.9
11	Influenza and pneumonia (J10-J18)	344	1.0	4.4
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	339	1.0	4.3
13	Septicemia (A40-A41)	312	0.9	4.0
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	234	0.7	3.0
15	Aortic aneurysm and dissection (I71)	158	0.5	2.0
...	All other causes (Residual)	5,179	15.6	66.4
White, non-Hispanic, male, 50-54 years				
...	All causes	42,622	100.0	604.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,908	27.9	168.9
2	Malignant neoplasms (C00-C97)	11,787	27.7	167.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,333	7.8	47.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	1,988	4.7	28.2
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,956	4.6	27.7
6	Diabetes mellitus (E10-E14)	1,262	3.0	17.9
7	Cerebrovascular diseases (I60-I69)	1,032	2.4	14.6
8	Chronic lower respiratory diseases (J40-J47)	924	2.2	13.1
9	Viral hepatitis (B15-B19)	599	1.4	8.5
10	Human immunodeficiency virus (HIV) disease (B20-B24)	444	1.0	6.3
11	Influenza and pneumonia (J10-J18)	424	1.0	6.0
12	Septicemia (A40-A41)	368	0.9	5.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	299	0.7	4.2
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	247	0.6	3.5
15	Aortic aneurysm and dissection (I71)	197	0.5	2.8
...	All other causes (Residual)	5,854	13.7	83.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 55-59 years				
...	All causes	52,169	100.0	906.2
1	Malignant neoplasms (C00-C97)	17,454	33.5	303.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,237	29.2	264.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,427	4.7	42.2
4	Chronic lower respiratory diseases (J40-J47)	1,793	3.4	31.1
5	Diabetes mellitus (E10-E14)	1,767	3.4	30.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1,674	3.2	29.1
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,524	2.9	26.5
8	Cerebrovascular diseases (I60-I69)	1,490	2.9	25.9
9	Septicemia (A40-A41)	524	1.0	9.1
10	Influenza and pneumonia (J10-J18)	508	1.0	8.8
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	438	0.8	7.6
12	Aortic aneurysm and dissection (I71)	349	0.7	6.1
13	Viral hepatitis (B15-B19)	271	0.5	4.7
14	Human immunodeficiency virus (HIV) disease (B20-B24)	269	0.5	4.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	261	0.5	4.5
...	All other causes (Residual)	6,183	11.9	107.4
White, non-Hispanic, male, 60-64 years				
...	All causes	62,366	100.0	1,413.9
1	Malignant neoplasms (C00-C97)	22,756	36.5	515.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	18,076	29.0	409.8
3	Chronic lower respiratory diseases (J40-J47)	3,116	5.0	70.6
4	Diabetes mellitus (E10-E14)	2,048	3.3	46.4
5	Cerebrovascular diseases (I60-I69)	2,021	3.2	45.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,947	3.1	44.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,423	2.3	32.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,022	1.6	23.2
9	Influenza and pneumonia (J10-J18)	720	1.2	16.3
10	Septicemia (A40-A41)	680	1.1	15.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	607	1.0	13.8
12	Aortic aneurysm and dissection (I71)	526	0.8	11.9
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	286	0.5	6.5
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	267	0.4	6.1
15	Pneumonitis due to solids and liquids (J69)	180	0.3	4.1
...	All other causes (Residual)	6,691	10.7	151.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 65-69 years				
...	All causes	79,363	100.0	2,216.9
1	Malignant neoplasms (C00-C97)	28,750	36.2	803.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	22,786	28.7	636.5
3	Chronic lower respiratory diseases (J40-J47)	5,124	6.5	143.1
4	Cerebrovascular diseases (I60-I69)	3,032	3.8	84.7
5	Diabetes mellitus (E10-E14)	2,595	3.3	72.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,795	2.3	50.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,246	1.6	34.8
8	Influenza and pneumonia (J10-J18)	1,059	1.3	29.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	983	1.2	27.5
10	Aortic aneurysm and dissection (I71)	904	1.1	25.3
11	Septicemia (A40-A41)	903	1.1	25.2
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	870	1.1	24.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	457	0.6	12.8
14	Parkinson's disease (G20-G21)	369	0.5	10.3
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	350	0.4	9.8
...	All other causes (Residual)	8,140	10.3	227.4
White, non-Hispanic, male, 70-74 years				
...	All causes	112,481	100.0	3,500.6
1	Malignant neoplasms (C00-C97)	37,500	33.3	1,167.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	32,479	28.9	1,010.8
3	Chronic lower respiratory diseases (J40-J47)	8,491	7.5	264.3
4	Cerebrovascular diseases (I60-I69)	5,336	4.7	166.1
5	Diabetes mellitus (E10-E14)	3,446	3.1	107.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,196	2.0	68.3
7	Influenza and pneumonia (J10-J18)	1,991	1.8	62.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,711	1.5	53.2
9	Septicemia (A40-A41)	1,407	1.3	43.8
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,243	1.1	38.7
11	Aortic aneurysm and dissection (I71)	1,201	1.1	37.4
12	Alzheimer's disease (G30)	1,008	0.9	31.4
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	995	0.9	31.0
14	Parkinson's disease (G20-G21)	943	0.8	29.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	678	0.6	21.1
...	All other causes (Residual)	11,856	10.5	369.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 75-79 years				
...	All causes	143,909	100.0	5,514.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	43,461	30.2	1,665.4
2	Malignant neoplasms (C00-C97)	40,610	28.2	1,556.2
3	Chronic lower respiratory diseases (J40-J47)	11,168	7.8	428.0
4	Cerebrovascular diseases (I60-I69)	8,484	5.9	325.1
5	Diabetes mellitus (E10-E14)	4,179	2.9	160.1
6	Influenza and pneumonia (J10-J18)	3,427	2.4	131.3
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,778	1.9	106.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,583	1.8	99.0
9	Alzheimer's disease (G30)	2,510	1.7	96.2
10	Septicemia (A40-A41)	1,921	1.3	73.6
11	Parkinson's disease (G20-G21)	1,896	1.3	72.7
12	Aortic aneurysm and dissection (I71)	1,542	1.1	59.1
13	Pneumonitis due to solids and liquids (J69)	1,131	0.8	43.3
14	Chronic liver disease and cirrhosis (K70,K73-K74)	1,050	0.7	40.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	1,023	0.7	39.2
...	All other causes (Residual)	16,146	11.2	618.7
White, non-Hispanic, male, 80-84 years				
...	All causes	151,854	100.0	8,733.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	49,155	32.4	2,827.0
2	Malignant neoplasms (C00-C97)	34,029	22.4	1,957.1
3	Chronic lower respiratory diseases (J40-J47)	10,866	7.2	624.9
4	Cerebrovascular diseases (I60-I69)	10,697	7.0	615.2
5	Influenza and pneumonia (J10-J18)	4,975	3.3	286.1
6	Alzheimer's disease (G30)	4,034	2.7	232.0
7	Diabetes mellitus (E10-E14)	3,972	2.6	228.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,104	2.0	178.5
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,027	2.0	174.1
10	Parkinson's disease (G20-G21)	2,577	1.7	148.2
11	Septicemia (A40-A41)	2,085	1.4	119.9
12	Pneumonitis due to solids and liquids (J69)	1,691	1.1	97.3
13	Aortic aneurysm and dissection (I71)	1,375	0.9	79.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	1,046	0.7	60.2
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	997	0.7	57.3
...	All other causes (Residual)	18,224	12.0	1,048.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 85-89 years				
...	All causes	119,953	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	41,657	34.7	...
2	Malignant neoplasms (C00-C97)	20,477	17.1	...
3	Cerebrovascular diseases (I60-I69)	9,680	8.1	...
4	Chronic lower respiratory diseases (J40-J47)	7,394	6.2	...
5	Influenza and pneumonia (J10-J18)	5,057	4.2	...
6	Alzheimer's disease (G30)	4,217	3.5	...
7	Diabetes mellitus (E10-E14)	2,720	2.3	...
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,679	2.2	...
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,544	2.1	...
10	Parkinson's disease (G20-G21)	1,969	1.6	...
11	Pneumonitis due to solids and liquids (J69)	1,800	1.5	...
12	Septicemia (A40-A41)	1,557	1.3	...
13	Atherosclerosis (I70)	999	0.8	...
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	977	0.8	...
15	Aortic aneurysm and dissection (I71)	968	0.8	...
...	All other causes (Residual)	15,258	12.7	...
White, non-Hispanic, male, 90-94 years				
...	All causes	60,743	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	22,700	37.4	...
2	Malignant neoplasms (C00-C97)	7,837	12.9	...
3	Cerebrovascular diseases (I60-I69)	5,016	8.3	...
4	Influenza and pneumonia (J10-J18)	3,416	5.6	...
5	Chronic lower respiratory diseases (J40-J47)	3,109	5.1	...
6	Alzheimer's disease (G30)	2,343	3.9	...
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,454	2.4	...
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,332	2.2	...
9	Pneumonitis due to solids and liquids (J69)	1,124	1.9	...
10	Diabetes mellitus (E10-E14)	1,071	1.8	...
11	Parkinson's disease (G20-G21)	741	1.2	...
12	Atherosclerosis (I70)	688	1.1	...
13	Septicemia (A40-A41)	676	1.1	...
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	576	0.9	...
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	357	0.6	...
...	All other causes (Residual)	8,303	13.7	...

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 95-99 years				
...	All causes	16,825	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,647	39.5	...
2	Malignant neoplasms (C00-C97)	1,629	9.7	...
3	Cerebrovascular diseases (I60-I69)	1,313	7.8	...
4	Influenza and pneumonia (J10-J18)	1,164	6.9	...
5	Chronic lower respiratory diseases (J40-J47)	659	3.9	...
6	Alzheimer's disease (G30)	651	3.9	...
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	425	2.5	...
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	406	2.4	...
9	Pneumonitis due to solids and liquids (J69)	367	2.2	...
10	Atherosclerosis (I70)	260	1.5	...
11	Diabetes mellitus (E10-E14)	255	1.5	...
12	Septicemia (A40-A41)	192	1.1	...
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	176	1.0	...
14	Parkinson's disease (G20-G21)	136	0.8	...
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	85	0.5	...
...	All other causes (Residual)	2,460	14.6	...
White, non-Hispanic, male, 100 years and over				
...	All causes	2,313	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	972	42.0	...
2	Influenza and pneumonia (J10-J18)	217	9.4	...
3	Malignant neoplasms (C00-C97)	169	7.3	...
4	Cerebrovascular diseases (I60-I69)	153	6.6	...
5	Chronic lower respiratory diseases (J40-J47)	70	3.0	...
6	Alzheimer's disease (G30)	59	2.6	...
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	55	2.4	...
8	Pneumonitis due to solids and liquids (J69)	52	2.2	...
9	Atherosclerosis (I70)	43	1.9	...
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	41	1.8	...
11	Septicemia (A40-A41)	23	1.0	...
12	Diabetes mellitus (E10-E14)	18	0.8	...
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	17	0.7	...
14	Anemias (D50-D64)	12	0.5	...
15	Nutritional deficiencies (E40-E64)	10	0.4	...
15	Parkinson's disease (G20-G21)	10	0.4	...
15	Hyperplasia of prostate (N40)	10	0.4	...
...	All other causes (Residual)	382	16.5	...

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, all ages				
...	All causes	1,024,328	100.0	1,010.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	296,319	28.9	292.3
2	Malignant neoplasms (C00-C97)	221,431	21.6	218.5
3	Cerebrovascular diseases (I60-I69)	83,210	8.1	82.1
4	Chronic lower respiratory diseases (J40-J47)	58,466	5.7	57.7
5	Alzheimer's disease (G30)	38,087	3.7	37.6
6	Influenza and pneumonia (J10-J18)	31,485	3.1	31.1
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	29,953	2.9	29.6
8	Diabetes mellitus (E10-E14)	27,190	2.7	26.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15,786	1.5	15.6
10	Septicemia (A40-A41)	14,397	1.4	14.2
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	9,292	0.9	9.2
12	Atherosclerosis (I70)	7,683	0.8	7.6
13	Chronic liver disease and cirrhosis (K70,K73-K74)	7,553	0.7	7.5
14	Pneumonitis due to solids and liquids (J69)	7,549	0.7	7.4
15	Parkinson's disease (G20-G21)	6,788	0.7	6.7
...	All other causes (Residual)	169,139	16.5	166.9
White, non-Hispanic, female, 1-4 years				
...	All causes	1,071	100.0	23.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	338	31.6	7.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	130	12.1	2.9
3	Malignant neoplasms (C00-C97)	102	9.5	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	80	7.5	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	37	3.5	0.8
6	Influenza and pneumonia (J10-J18)	22	2.1	0.5
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	19	1.8	*
8	Septicemia (A40-A41)	17	1.6	*
9	Acute bronchitis and bronchiolitis (J20-J21)	12	1.1	*
9	Certain conditions originating in the perinatal period (P00-P96)	12	1.1	*
11	Cerebrovascular diseases (I60-I69)	11	1.0	*
12	Anemias (D50-D64)	6	0.6	*
12	Chronic lower respiratory diseases (J40-J47)	6	0.6	*
14	Meningitis (G00,G03)	4	0.4	*
15	Meningococcal infection (A39)	2	0.2	*
15	Diabetes mellitus (E10-E14)	2	0.2	*
15	Pneumonitis due to solids and liquids (J69)	2	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.2	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.2	*
...	All other causes (Residual)	265	24.7	5.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 5-9 years				
...	All causes	720	100.0	12.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	238	33.1	4.1
2	Malignant neoplasms (C00-C97)	149	20.7	2.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	59	8.2	1.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	35	4.9	0.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	29	4.0	0.5
6	Influenza and pneumonia (J10-J18)	15	2.1	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	11	1.5	*
8	Cerebrovascular diseases (I60-I69)	10	1.4	*
9	Septicemia (A40-A41)	9	1.3	*
10	Anemias (D50-D64)	8	1.1	*
11	Meningitis (G00,G03)	4	0.6	*
12	Chronic lower respiratory diseases (J40-J47)	3	0.4	*
13	Pneumonitis due to solids and liquids (J69)	2	0.3	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.3	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.3	*
...	All other causes (Residual)	144	20.0	2.5
White, non-Hispanic, female, 10-14 years				
...	All causes	946	100.0	14.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	346	36.6	5.4
2	Malignant neoplasms (C00-C97)	138	14.6	2.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	69	7.3	1.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	39	4.1	0.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	37	3.9	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	32	3.4	0.5
7	Influenza and pneumonia (J10-J18)	20	2.1	0.3
8	Chronic lower respiratory diseases (J40-J47)	16	1.7	*
9	Septicemia (A40-A41)	12	1.3	*
10	Cerebrovascular diseases (I60-I69)	11	1.2	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	8	0.8	*
12	Certain conditions originating in the perinatal period (P00-P96)	7	0.7	*
13	Meningitis (G00,G03)	5	0.5	*
14	Anemias (D50-D64)	3	0.3	*
14	Diabetes mellitus (E10-E14)	3	0.3	*
...	All other causes (Residual)	200	21.1	3.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 15-19 years				
...	All causes	2,668	100.0	42.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,633	61.2	25.7
2	Malignant neoplasms (C00-C97)	190	7.1	3.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	170	6.4	2.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	103	3.9	1.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	77	2.9	1.2
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	63	2.4	1.0
7	Septicemia (A40-A41)	20	0.7	0.3
8	Influenza and pneumonia (J10-J18)	18	0.7	*
9	Diabetes mellitus (E10-E14)	16	0.6	*
10	Chronic lower respiratory diseases (J40-J47)	15	0.6	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	14	0.5	*
12	Cerebrovascular diseases (I60-I69)	11	0.4	*
13	Meningococcal infection (A39)	7	0.3	*
13	Viral hepatitis (B15-B19)	7	0.3	*
15	Pregnancy, childbirth and the puerperium (O00-099)	6	0.2	*
...	All other causes (Residual)	318	11.9	5.0
White, non-Hispanic, female, 20-24 years				
...	All causes	2,833	100.0	45.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,285	45.4	20.7
2	Malignant neoplasms (C00-C97)	268	9.5	4.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	254	9.0	4.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	170	6.0	2.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	100	3.5	1.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	74	2.6	1.2
7	Chronic lower respiratory diseases (J40-J47)	31	1.1	0.5
8	Diabetes mellitus (E10-E14)	28	1.0	0.5
8	Cerebrovascular diseases (I60-I69)	28	1.0	0.5
10	Influenza and pneumonia (J10-J18)	23	0.8	0.4
11	Septicemia (A40-A41)	15	0.5	*
11	Pregnancy, childbirth and the puerperium (O00-099)	15	0.5	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	13	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	0.4	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	9	0.3	*
...	All other causes (Residual)	509	18.0	8.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 25-29 years				
...	All causes	2,912	100.0	51.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	929	31.9	16.4
2	Malignant neoplasms (C00-C97)	368	12.6	6.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	344	11.8	6.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	185	6.4	3.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	141	4.8	2.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	56	1.9	1.0
7	Diabetes mellitus (E10-E14)	51	1.8	0.9
8	Influenza and pneumonia (J10-J18)	45	1.5	0.8
9	Cerebrovascular diseases (I60-I69)	38	1.3	0.7
10	Pregnancy, childbirth and the puerperium (O00-099)	31	1.1	0.5
11	Chronic lower respiratory diseases (J40-J47)	30	1.0	0.5
12	Human immunodeficiency virus (HIV) disease (B20-B24)	28	1.0	0.5
13	Septicemia (A40-A41)	26	0.9	0.5
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	24	0.8	0.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	16	0.5	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	16	0.5	*
...	All other causes (Residual)	584	20.1	10.3
White, non-Hispanic, female, 30-34 years				
...	All causes	4,507	100.0	68.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,142	25.3	17.2
2	Malignant neoplasms (C00-C97)	897	19.9	13.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	400	8.9	6.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	356	7.9	5.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	176	3.9	2.7
6	Cerebrovascular diseases (I60-I69)	111	2.5	1.7
7	Diabetes mellitus (E10-E14)	100	2.2	1.5
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	80	1.8	1.2
9	Human immunodeficiency virus (HIV) disease (B20-B24)	66	1.5	1.0
10	Chronic liver disease and cirrhosis (K70,K73-K74)	58	1.3	0.9
11	Septicemia (A40-A41)	53	1.2	0.8
12	Influenza and pneumonia (J10-J18)	51	1.1	0.8
13	Chronic lower respiratory diseases (J40-J47)	46	1.0	0.7
14	Pregnancy, childbirth and the puerperium (O00-099)	38	0.8	0.6
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	31	0.7	0.5
...	All other causes (Residual)	902	20.0	13.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 35-39 years				
...	All causes	7,931	100.0	108.0
1	Malignant neoplasms (C00-C97)	1,998	25.2	27.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,536	19.4	20.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	834	10.5	11.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	557	7.0	7.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	227	2.9	3.1
6	Cerebrovascular diseases (I60-I69)	217	2.7	3.0
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	207	2.6	2.8
8	Diabetes mellitus (E10-E14)	190	2.4	2.6
9	Human immunodeficiency virus (HIV) disease (B20-B24)	132	1.7	1.8
10	Chronic lower respiratory diseases (J40-J47)	117	1.5	1.6
11	Influenza and pneumonia (J10-J18)	106	1.3	1.4
12	Septicemia (A40-A41)	97	1.2	1.3
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	95	1.2	1.3
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	57	0.7	0.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	40	0.5	0.5
...	All other causes (Residual)	1,521	19.2	20.7
White, non-Hispanic, female, 40-44 years				
...	All causes	13,549	100.0	165.4
1	Malignant neoplasms (C00-C97)	4,195	31.0	51.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,949	14.4	23.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,623	12.0	19.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	776	5.7	9.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	550	4.1	6.7
6	Cerebrovascular diseases (I60-I69)	454	3.4	5.5
7	Diabetes mellitus (E10-E14)	306	2.3	3.7
8	Chronic lower respiratory diseases (J40-J47)	203	1.5	2.5
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	187	1.4	2.3
10	Influenza and pneumonia (J10-J18)	171	1.3	2.1
11	Septicemia (A40-A41)	164	1.2	2.0
12	Human immunodeficiency virus (HIV) disease (B20-B24)	141	1.0	1.7
13	Viral hepatitis (B15-B19)	108	0.8	1.3
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	105	0.8	1.3
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	105	0.8	1.3
...	All other causes (Residual)	2,512	18.5	30.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 45-49 years				
...	All causes	18,942	100.0	240.1
1	Malignant neoplasms (C00-C97)	7,058	37.3	89.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,541	13.4	32.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,668	8.8	21.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	788	4.2	10.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	671	3.5	8.5
6	Cerebrovascular diseases (I60-I69)	642	3.4	8.1
7	Diabetes mellitus (E10-E14)	497	2.6	6.3
8	Chronic lower respiratory diseases (J40-J47)	458	2.4	5.8
9	Influenza and pneumonia (J10-J18)	267	1.4	3.4
10	Septicemia (A40-A41)	260	1.4	3.3
11	Viral hepatitis (B15-B19)	207	1.1	2.6
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	194	1.0	2.5
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	169	0.9	2.1
14	Human immunodeficiency virus (HIV) disease (B20-B24)	125	0.7	1.6
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	120	0.6	1.5
...	All other causes (Residual)	3,277	17.3	41.5
White, non-Hispanic, female, 50-54 years				
...	All causes	25,146	100.0	349.1
1	Malignant neoplasms (C00-C97)	10,852	43.2	150.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,891	15.5	54.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,446	5.8	20.1
4	Cerebrovascular diseases (I60-I69)	881	3.5	12.2
5	Chronic lower respiratory diseases (J40-J47)	845	3.4	11.7
6	Diabetes mellitus (E10-E14)	834	3.3	11.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	682	2.7	9.5
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	563	2.2	7.8
9	Septicemia (A40-A41)	334	1.3	4.6
10	Influenza and pneumonia (J10-J18)	294	1.2	4.1
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	220	0.9	3.1
12	Viral hepatitis (B15-B19)	171	0.7	2.4
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	147	0.6	2.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	133	0.5	1.8
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	118	0.5	1.6
...	All other causes (Residual)	3,735	14.9	51.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 55-59 years				
...	All causes	33,747	100.0	563.3
1	Malignant neoplasms (C00-C97)	14,964	44.3	249.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,988	17.7	99.9
3	Chronic lower respiratory diseases (J40-J47)	1,807	5.4	30.2
4	Cerebrovascular diseases (I60-I69)	1,243	3.7	20.7
5	Diabetes mellitus (E10-E14)	1,129	3.3	18.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,032	3.1	17.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	690	2.0	11.5
8	Septicemia (A40-A41)	472	1.4	7.9
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	441	1.3	7.4
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	384	1.1	6.4
11	Influenza and pneumonia (J10-J18)	379	1.1	6.3
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	188	0.6	3.1
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	151	0.4	2.5
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	129	0.4	2.2
15	Viral hepatitis (B15-B19)	117	0.3	2.0
...	All other causes (Residual)	4,633	13.7	77.3
White, non-Hispanic, female, 60-64 years				
...	All causes	42,561	100.0	896.1
1	Malignant neoplasms (C00-C97)	18,635	43.8	392.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,193	19.3	172.5
3	Chronic lower respiratory diseases (J40-J47)	3,041	7.1	64.0
4	Cerebrovascular diseases (I60-I69)	1,678	3.9	35.3
5	Diabetes mellitus (E10-E14)	1,459	3.4	30.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	987	2.3	20.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	696	1.6	14.7
8	Septicemia (A40-A41)	565	1.3	11.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	549	1.3	11.6
10	Influenza and pneumonia (J10-J18)	545	1.3	11.5
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	267	0.6	5.6
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	227	0.5	4.8
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	199	0.5	4.2
14	Aortic aneurysm and dissection (I71)	179	0.4	3.8
15	Alzheimer's disease (G30)	172	0.4	3.6
...	All other causes (Residual)	5,169	12.1	108.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 65-69 years				
...	All causes	57,828	100.0	1,430.8
1	Malignant neoplasms (C00-C97)	22,812	39.4	564.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,313	21.3	304.7
3	Chronic lower respiratory diseases (J40-J47)	5,046	8.7	124.8
4	Cerebrovascular diseases (I60-I69)	2,622	4.5	64.9
5	Diabetes mellitus (E10-E14)	2,122	3.7	52.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,045	1.8	25.9
7	Septicemia (A40-A41)	902	1.6	22.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	811	1.4	20.1
9	Influenza and pneumonia (J10-J18)	807	1.4	20.0
10	Chronic liver disease and cirrhosis (K70,K73-K74)	801	1.4	19.8
11	Alzheimer's disease (G30)	459	0.8	11.4
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	325	0.6	8.0
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	315	0.5	7.8
14	Aortic aneurysm and dissection (I71)	312	0.5	7.7
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	204	0.4	5.0
...	All other causes (Residual)	6,932	12.0	171.5
White, non-Hispanic, female, 70-74 years				
...	All causes	89,289	100.0	2,273.2
1	Malignant neoplasms (C00-C97)	30,402	34.0	774.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,950	23.5	533.4
3	Chronic lower respiratory diseases (J40-J47)	8,147	9.1	207.4
4	Cerebrovascular diseases (I60-I69)	5,365	6.0	136.6
5	Diabetes mellitus (E10-E14)	3,174	3.6	80.8
6	Influenza and pneumonia (J10-J18)	1,562	1.7	39.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,445	1.6	36.8
8	Septicemia (A40-A41)	1,371	1.5	34.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,365	1.5	34.8
10	Alzheimer's disease (G30)	1,337	1.5	34.0
11	Chronic liver disease and cirrhosis (K70,K73-K74)	915	1.0	23.3
12	Aortic aneurysm and dissection (I71)	640	0.7	16.3
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	593	0.7	15.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	535	0.6	13.6
15	Parkinson's disease (G20-G21)	486	0.5	12.4
...	All other causes (Residual)	11,002	12.3	280.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 75-79 years				
...	All causes	134,235	100.0	3,681.6
1	Malignant neoplasms (C00-C97)	35,852	26.7	983.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	35,530	26.5	974.5
3	Chronic lower respiratory diseases (J40-J47)	11,210	8.4	307.4
4	Cerebrovascular diseases (I60-I69)	10,337	7.7	283.5
5	Diabetes mellitus (E10-E14)	4,453	3.3	122.1
6	Alzheimer's disease (G30)	3,822	2.8	104.8
7	Influenza and pneumonia (J10-J18)	3,141	2.3	86.1
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,304	1.7	63.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,296	1.7	63.0
10	Septicemia (A40-A41)	2,122	1.6	58.2
11	Parkinson's disease (G20-G21)	1,188	0.9	32.6
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,048	0.8	28.7
13	Chronic liver disease and cirrhosis (K70,K73-K74)	1,011	0.8	27.7
14	Aortic aneurysm and dissection (I71)	957	0.7	26.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	867	0.6	23.8
...	All other causes (Residual)	18,097	13.5	496.3
White, non-Hispanic, female, 80-84 years				
...	All causes	176,433	100.0	6,207.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	53,829	30.5	1,893.8
2	Malignant neoplasms (C00-C97)	33,466	19.0	1,177.4
3	Cerebrovascular diseases (I60-I69)	16,809	9.5	591.4
4	Chronic lower respiratory diseases (J40-J47)	11,873	6.7	417.7
5	Alzheimer's disease (G30)	7,861	4.5	276.6
6	Influenza and pneumonia (J10-J18)	5,458	3.1	192.0
7	Diabetes mellitus (E10-E14)	5,088	2.9	179.0
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,207	1.8	112.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,127	1.8	110.0
10	Septicemia (A40-A41)	2,678	1.5	94.2
11	Parkinson's disease (G20-G21)	1,884	1.1	66.3
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,738	1.0	61.1
13	Pneumonitis due to solids and liquids (J69)	1,471	0.8	51.8
14	Atherosclerosis (I70)	1,306	0.7	45.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	1,127	0.6	39.6
...	All other causes (Residual)	25,511	14.5	897.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 85-89 years				
...	All causes	186,334	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	64,460	34.6	...
2	Malignant neoplasms (C00-C97)	23,728	12.7	...
3	Cerebrovascular diseases (I60-I69)	19,732	10.6	...
4	Alzheimer's disease (G30)	10,697	5.7	...
5	Chronic lower respiratory diseases (J40-J47)	9,155	4.9	...
6	Influenza and pneumonia (J10-J18)	7,324	3.9	...
7	Diabetes mellitus (E10-E14)	4,346	2.3	...
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,433	1.8	...
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,222	1.7	...
10	Septicemia (A40-A41)	2,611	1.4	...
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,100	1.1	...
12	Pneumonitis due to solids and liquids (J69)	1,864	1.0	...
13	Atherosclerosis (I70)	1,829	1.0	...
14	Parkinson's disease (G20-G21)	1,726	0.9	...
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	1,045	0.6	...
...	All other causes (Residual)	29,062	15.6	...
White, non-Hispanic, female, 90-94 years				
...	All causes	142,476	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	54,640	38.4	...
2	Cerebrovascular diseases (I60-I69)	15,562	10.9	...
3	Malignant neoplasms (C00-C97)	11,585	8.1	...
4	Alzheimer's disease (G30)	8,898	6.2	...
5	Influenza and pneumonia (J10-J18)	6,812	4.8	...
6	Chronic lower respiratory diseases (J40-J47)	4,651	3.3	...
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,562	1.8	...
8	Diabetes mellitus (E10-E14)	2,465	1.7	...
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,333	1.6	...
10	Atherosclerosis (I70)	1,872	1.3	...
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,828	1.3	...
12	Septicemia (A40-A41)	1,790	1.3	...
13	Pneumonitis due to solids and liquids (J69)	1,562	1.1	...
14	Parkinson's disease (G20-G21)	931	0.7	...
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	686	0.5	...
...	All other causes (Residual)	24,299	17.1	...

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 95-99 years				
...	All causes	60,873	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	24,758	40.7	...
2	Cerebrovascular diseases (I60-I69)	6,240	10.3	...
3	Alzheimer's disease (G30)	4,033	6.6	...
4	Influenza and pneumonia (J10-J18)	3,421	5.6	...
5	Malignant neoplasms (C00-C97)	3,286	5.4	...
6	Chronic lower respiratory diseases (J40-J47)	1,510	2.5	...
7	Atherosclerosis (I70)	1,041	1.7	...
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,023	1.7	...
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	899	1.5	...
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	874	1.4	...
11	Diabetes mellitus (E10-E14)	809	1.3	...
12	Pneumonitis due to solids and liquids (J69)	700	1.1	...
13	Septicemia (A40-A41)	691	1.1	...
14	Parkinson's disease (G20-G21)	272	0.4	...
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	234	0.4	...
...	All other causes (Residual)	11,082	18.2	...
White, non-Hispanic, female, 100 years and over				
...	All causes	13,546	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,836	43.1	...
2	Cerebrovascular diseases (I60-I69)	1,188	8.8	...
3	Influenza and pneumonia (J10-J18)	962	7.1	...
4	Alzheimer's disease (G30)	707	5.2	...
5	Malignant neoplasms (C00-C97)	473	3.5	...
6	Atherosclerosis (I70)	314	2.3	...
7	Chronic lower respiratory diseases (J40-J47)	248	1.8	...
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	219	1.6	...
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	217	1.6	...
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	169	1.2	...
11	Pneumonitis due to solids and liquids (J69)	147	1.1	...
12	Septicemia (A40-A41)	126	0.9	...
13	Diabetes mellitus (E10-E14)	117	0.9	...
14	Nutritional deficiencies (E40-E64)	63	0.5	...
15	Anemias (D50-D64)	60	0.4	...
...	All other causes (Residual)	2,700	19.9	...

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, all ages				
...	All causes	286,573	100.0	792.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	76,694	26.8	212.2
2	Malignant neoplasms (C00-C97)	61,996	21.6	171.5
3	Cerebrovascular diseases (I60-I69)	18,691	6.5	51.7
4	Diabetes mellitus (E10-E14)	12,583	4.4	34.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,285	4.3	34.0
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8,147	2.8	22.5
7	Chronic lower respiratory diseases (J40-J47)	7,730	2.7	21.4
8	Human immunodeficiency virus (HIV) disease (B20-B24)	7,714	2.7	21.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,410	2.6	20.5
10	Septicemia (A40-A41)	6,074	2.1	16.8
11	Influenza and pneumonia (J10-J18)	5,803	2.0	16.1
12	Certain conditions originating in the perinatal period (P00-P96)	4,777	1.7	13.2
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	4,165	1.5	11.5
14	Alzheimer's disease (G30)	3,303	1.2	9.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,545	0.9	7.0
...	All other causes (Residual)	46,656	16.3	129.1
Black, non-Hispanic, both sexes, 1-4 years				
...	All causes	1,163	100.0	48.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	350	30.1	14.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	156	13.4	6.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	113	9.7	4.7
4	Malignant neoplasms (C00-C97)	71	6.1	3.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	40	3.4	1.7
5	Influenza and pneumonia (J10-J18)	40	3.4	1.7
7	Chronic lower respiratory diseases (J40-J47)	29	2.5	1.2
8	Certain conditions originating in the perinatal period (P00-P96)	26	2.2	1.1
9	Septicemia (A40-A41)	17	1.5	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	14	1.2	*
11	Cerebrovascular diseases (I60-I69)	10	0.9	*
12	Anemias (D50-D64)	7	0.6	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	5	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.4	*
15	Meningitis (G00,G03)	4	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.3	*
...	All other causes (Residual)	272	23.4	11.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 5-9 years				
...	All causes	686	100.0	22.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	298	43.4	9.6
2	Malignant neoplasms (C00-C97)	81	11.8	2.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	45	6.6	1.4
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	38	5.5	1.2
5	Chronic lower respiratory diseases (J40-J47)	21	3.1	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	19	2.8	*
7	Anemias (D50-D64)	12	1.7	*
8	Septicemia (A40-A41)	7	1.0	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	7	1.0	*
8	Cerebrovascular diseases (I60-I69)	7	1.0	*
8	Influenza and pneumonia (J10-J18)	7	1.0	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.9	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.6	*
13	Certain conditions originating in the perinatal period (P00-P96)	4	0.6	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.3	*
...	All other causes (Residual)	128	18.7	4.1
Black, non-Hispanic, both sexes, 10-14 years				
...	All causes	948	100.0	28.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	309	32.6	9.2
2	Malignant neoplasms (C00-C97)	87	9.2	2.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	87	9.2	2.6
4	Chronic lower respiratory diseases (J40-J47)	53	5.6	1.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	46	4.9	1.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	43	4.5	1.3
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	34	3.6	1.0
8	Cerebrovascular diseases (I60-I69)	16	1.7	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	15	1.6	*
10	Anemias (D50-D64)	14	1.5	*
11	Septicemia (A40-A41)	13	1.4	*
11	Diabetes mellitus (E10-E14)	13	1.4	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	11	1.2	*
13	Influenza and pneumonia (J10-J18)	11	1.2	*
15	Meningitis (G00,G03)	9	0.9	*
...	All other causes (Residual)	187	19.7	5.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 15-19 years				
...	All causes	2,542	100.0	83.8
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	967	38.0	31.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	740	29.1	24.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	122	4.8	4.0
4	Malignant neoplasms (C00-C97)	110	4.3	3.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	105	4.1	3.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	52	2.0	1.7
7	Chronic lower respiratory diseases (J40-J47)	43	1.7	1.4
8	Anemias (D50-D64)	27	1.1	0.9
9	Human immunodeficiency virus (HIV) disease (B20-B24)	25	1.0	0.8
10	Influenza and pneumonia (J10-J18)	18	0.7	*
11	Diabetes mellitus (E10-E14)	16	0.6	*
11	Pregnancy, childbirth and the puerperium (O00-099)	16	0.6	*
13	Cerebrovascular diseases (I60-I69)	13	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	0.4	*
14	Legal intervention (Y35,Y89.0)	10	0.4	*
...	All other causes (Residual)	268	10.5	8.8
Black, non-Hispanic, both sexes, 20-24 years				
...	All causes	4,362	100.0	151.7
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,887	43.3	65.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	940	21.5	32.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	270	6.2	9.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	194	4.4	6.7
5	Malignant neoplasms (C00-C97)	132	3.0	4.6
6	Human immunodeficiency virus (HIV) disease (B20-B24)	99	2.3	3.4
7	Anemias (D50-D64)	51	1.2	1.8
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	46	1.1	1.6
9	Diabetes mellitus (E10-E14)	41	0.9	1.4
10	Chronic lower respiratory diseases (J40-J47)	35	0.8	1.2
11	Pregnancy, childbirth and the puerperium (O00-099)	29	0.7	1.0
12	Legal intervention (Y35,Y89.0)	27	0.6	0.9
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	26	0.6	0.9
14	Cerebrovascular diseases (I60-I69)	25	0.6	0.9
15	Influenza and pneumonia (J10-J18)	20	0.5	0.7
...	All other causes (Residual)	540	12.4	18.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 25-29 years				
...	All causes	4,487	100.0	179.6
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,412	31.5	56.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	853	19.0	34.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	352	7.8	14.1
4	Human immunodeficiency virus (HIV) disease (B20-B24)	306	6.8	12.2
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	271	6.0	10.8
6	Malignant neoplasms (C00-C97)	211	4.7	8.4
7	Diabetes mellitus (E10-E14)	73	1.6	2.9
8	Anemias (D50-D64)	64	1.4	2.6
9	Cerebrovascular diseases (I60-I69)	50	1.1	2.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	48	1.1	1.9
11	Chronic lower respiratory diseases (J40-J47)	43	1.0	1.7
12	Septicemia (A40-A41)	41	0.9	1.6
13	Pregnancy, childbirth and the puerperium (O00-099)	37	0.8	1.5
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	35	0.8	1.4
15	Influenza and pneumonia (J10-J18)	27	0.6	1.1
...	All other causes (Residual)	664	14.8	26.6
Black, non-Hispanic, both sexes, 30-34 years				
...	All causes	5,522	100.0	207.4
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,013	18.3	38.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	823	14.9	30.9
3	Human immunodeficiency virus (HIV) disease (B20-B24)	804	14.6	30.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	641	11.6	24.1
5	Malignant neoplasms (C00-C97)	452	8.2	17.0
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	226	4.1	8.5
7	Diabetes mellitus (E10-E14)	121	2.2	4.5
8	Cerebrovascular diseases (I60-I69)	116	2.1	4.4
9	Anemias (D50-D64)	63	1.1	2.4
10	Septicemia (A40-A41)	61	1.1	2.3
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	60	1.1	2.3
12	Chronic lower respiratory diseases (J40-J47)	59	1.1	2.2
13	Influenza and pneumonia (J10-J18)	53	1.0	2.0
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	40	0.7	1.5
15	Chronic liver disease and cirrhosis (K70,K73-K74)	37	0.7	1.4
...	All other causes (Residual)	953	17.3	35.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 35-39 years				
...	All causes	8,102	100.0	292.2
1	Human immunodeficiency virus (HIV) disease (B20-B24)	1,368	16.9	49.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,167	14.4	42.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,084	13.4	39.1
4	Malignant neoplasms (C00-C97)	907	11.2	32.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	770	9.5	27.8
6	Cerebrovascular diseases (I60-I69)	237	2.9	8.5
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	229	2.8	8.3
8	Diabetes mellitus (E10-E14)	208	2.6	7.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	114	1.4	4.1
10	Chronic lower respiratory diseases (J40-J47)	109	1.3	3.9
11	Influenza and pneumonia (J10-J18)	101	1.2	3.6
12	Septicemia (A40-A41)	96	1.2	3.5
13	Chronic liver disease and cirrhosis (K70,K73-K74)	84	1.0	3.0
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	79	1.0	2.8
15	Anemias (D50-D64)	72	0.9	2.6
...	All other causes (Residual)	1,477	18.2	53.3
Black, non-Hispanic, both sexes, 40-44 years				
...	All causes	11,853	100.0	422.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,145	18.1	76.5
2	Malignant neoplasms (C00-C97)	1,963	16.6	70.0
3	Human immunodeficiency virus (HIV) disease (B20-B24)	1,643	13.9	58.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,143	9.6	40.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	587	5.0	20.9
6	Cerebrovascular diseases (I60-I69)	504	4.3	18.0
7	Diabetes mellitus (E10-E14)	346	2.9	12.3
8	Chronic liver disease and cirrhosis (K70,K73-K74)	229	1.9	8.2
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	203	1.7	7.2
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	192	1.6	6.8
11	Chronic lower respiratory diseases (J40-J47)	183	1.5	6.5
12	Septicemia (A40-A41)	158	1.3	5.6
13	Influenza and pneumonia (J10-J18)	149	1.3	5.3
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	110	0.9	3.9
15	Viral hepatitis (B15-B19)	80	0.7	2.9
...	All other causes (Residual)	2,218	18.7	79.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 45-49 years				
...	All causes	16,002	100.0	648.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,506	21.9	142.1
2	Malignant neoplasms (C00-C97)	3,420	21.4	138.6
3	Human immunodeficiency virus (HIV) disease (B20-B24)	1,441	9.0	58.4
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,211	7.6	49.1
5	Cerebrovascular diseases (I60-I69)	807	5.0	32.7
6	Diabetes mellitus (E10-E14)	543	3.4	22.0
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	442	2.8	17.9
8	Chronic liver disease and cirrhosis (K70,K73-K74)	440	2.7	17.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	296	1.8	12.0
10	Septicemia (A40-A41)	238	1.5	9.6
11	Chronic lower respiratory diseases (J40-J47)	230	1.4	9.3
12	Influenza and pneumonia (J10-J18)	189	1.2	7.7
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	188	1.2	7.6
14	Viral hepatitis (B15-B19)	180	1.1	7.3
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	149	0.9	6.0
...	All other causes (Residual)	2,722	17.0	110.3
Black, non-Hispanic, both sexes, 50-54 years				
...	All causes	19,209	100.0	943.7
1	Malignant neoplasms (C00-C97)	4,956	25.8	243.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,812	25.1	236.4
3	Cerebrovascular diseases (I60-I69)	1,067	5.6	52.4
4	Human immunodeficiency virus (HIV) disease (B20-B24)	1,028	5.4	50.5
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	934	4.9	45.9
6	Diabetes mellitus (E10-E14)	850	4.4	41.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	516	2.7	25.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	449	2.3	22.1
9	Chronic lower respiratory diseases (J40-J47)	359	1.9	17.6
10	Septicemia (A40-A41)	355	1.8	17.4
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	268	1.4	13.2
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	225	1.2	11.1
13	Influenza and pneumonia (J10-J18)	210	1.1	10.3
14	Viral hepatitis (B15-B19)	188	1.0	9.2
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	116	0.6	5.7
...	All other causes (Residual)	2,876	15.0	141.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 55-59 years				
...	All causes	19,483	100.0	1,346.7
1	Malignant neoplasms (C00-C97)	5,899	30.3	407.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,389	27.7	372.5
3	Cerebrovascular diseases (I60-I69)	1,048	5.4	72.4
4	Diabetes mellitus (E10-E14)	1,003	5.1	69.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	618	3.2	42.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	497	2.6	34.4
7	Chronic lower respiratory diseases (J40-J47)	454	2.3	31.4
8	Human immunodeficiency virus (HIV) disease (B20-B24)	445	2.3	30.8
9	Septicemia (A40-A41)	400	2.1	27.6
10	Chronic liver disease and cirrhosis (K70,K73-K74)	360	1.8	24.9
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	272	1.4	18.8
12	Influenza and pneumonia (J10-J18)	254	1.3	17.6
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	130	0.7	9.0
14	Viral hepatitis (B15-B19)	113	0.6	7.8
15	Aortic aneurysm and dissection (I71)	99	0.5	6.8
...	All other causes (Residual)	2,502	12.8	172.9
Black, non-Hispanic, both sexes, 60-64 years				
...	All causes	21,395	100.0	1,911.2
1	Malignant neoplasms (C00-C97)	6,794	31.8	606.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,165	28.8	550.7
3	Cerebrovascular diseases (I60-I69)	1,255	5.9	112.1
4	Diabetes mellitus (E10-E14)	1,201	5.6	107.3
5	Chronic lower respiratory diseases (J40-J47)	613	2.9	54.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	575	2.7	51.4
7	Septicemia (A40-A41)	468	2.2	41.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	457	2.1	40.8
9	Influenza and pneumonia (J10-J18)	300	1.4	26.8
10	Chronic liver disease and cirrhosis (K70,K73-K74)	289	1.4	25.8
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	278	1.3	24.8
12	Human immunodeficiency virus (HIV) disease (B20-B24)	230	1.1	20.5
13	Aortic aneurysm and dissection (I71)	93	0.4	8.3
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	86	0.4	7.7
15	Viral hepatitis (B15-B19)	71	0.3	6.3
...	All other causes (Residual)	2,520	11.8	225.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 65-69 years				
...	All causes	24,508	100.0	2,670.7
1	Malignant neoplasms (C00-C97)	7,759	31.7	845.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,109	29.0	774.7
3	Cerebrovascular diseases (I60-I69)	1,584	6.5	172.6
4	Diabetes mellitus (E10-E14)	1,513	6.2	164.9
5	Chronic lower respiratory diseases (J40-J47)	802	3.3	87.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	737	3.0	80.3
7	Septicemia (A40-A41)	581	2.4	63.3
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	435	1.8	47.4
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	339	1.4	36.9
10	Influenza and pneumonia (J10-J18)	327	1.3	35.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	213	0.9	23.2
12	Human immunodeficiency virus (HIV) disease (B20-B24)	163	0.7	17.8
13	Pneumonitis due to solids and liquids (J69)	110	0.4	12.0
14	Aortic aneurysm and dissection (I71)	107	0.4	11.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	99	0.4	10.8
...	All other causes (Residual)	2,630	10.7	286.6
Black, non-Hispanic, both sexes, 70-74 years				
...	All causes	28,240	100.0	3,877.8
1	Malignant neoplasms (C00-C97)	8,325	29.5	1,143.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,302	29.4	1,140.0
3	Cerebrovascular diseases (I60-I69)	2,016	7.1	276.8
4	Diabetes mellitus (E10-E14)	1,639	5.8	225.1
5	Chronic lower respiratory diseases (J40-J47)	1,090	3.9	149.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	933	3.3	128.1
7	Septicemia (A40-A41)	670	2.4	92.0
8	Influenza and pneumonia (J10-J18)	510	1.8	70.0
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	456	1.6	62.6
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	383	1.4	52.6
11	Alzheimer's disease (G30)	200	0.7	27.5
12	Chronic liver disease and cirrhosis (K70,K73-K74)	145	0.5	19.9
13	Pneumonitis due to solids and liquids (J69)	140	0.5	19.2
14	Aortic aneurysm and dissection (I71)	137	0.5	18.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	112	0.4	15.4
...	All other causes (Residual)	3,182	11.3	436.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 75-79 years				
...	All causes	32,076	100.0	5,630.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,863	30.7	1,731.2
2	Malignant neoplasms (C00-C97)	8,221	25.6	1,443.0
3	Cerebrovascular diseases (I60-I69)	2,576	8.0	452.2
4	Diabetes mellitus (E10-E14)	1,728	5.4	303.3
5	Chronic lower respiratory diseases (J40-J47)	1,240	3.9	217.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,024	3.2	179.7
7	Septicemia (A40-A41)	845	2.6	148.3
8	Influenza and pneumonia (J10-J18)	708	2.2	124.3
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	559	1.7	98.1
10	Alzheimer's disease (G30)	468	1.5	82.1
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	433	1.3	76.0
12	Pneumonitis due to solids and liquids (J69)	243	0.8	42.7
13	Atherosclerosis (I70)	154	0.5	27.0
14	Aortic aneurysm and dissection (I71)	143	0.4	25.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	141	0.4	24.7
...	All other causes (Residual)	3,730	11.6	654.7
Black, non-Hispanic, both sexes, 80-84 years				
...	All causes	30,107	100.0	8,036.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,657	32.1	2,577.8
2	Malignant neoplasms (C00-C97)	6,297	20.9	1,680.9
3	Cerebrovascular diseases (I60-I69)	2,701	9.0	721.0
4	Diabetes mellitus (E10-E14)	1,485	4.9	396.4
5	Chronic lower respiratory diseases (J40-J47)	1,126	3.7	300.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	917	3.0	244.8
7	Influenza and pneumonia (J10-J18)	876	2.9	233.8
8	Septicemia (A40-A41)	788	2.6	210.3
9	Alzheimer's disease (G30)	705	2.3	188.2
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	574	1.9	153.2
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	366	1.2	97.7
12	Pneumonitis due to solids and liquids (J69)	256	0.9	68.3
13	Atherosclerosis (I70)	195	0.6	52.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	159	0.5	42.4
15	Aortic aneurysm and dissection (I71)	142	0.5	37.9
...	All other causes (Residual)	3,863	12.8	1,031.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 85-89 years				
...	All causes	23,674	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,003	33.8	...
2	Malignant neoplasms (C00-C97)	3,818	16.1	...
3	Cerebrovascular diseases (I60-I69)	2,298	9.7	...
4	Diabetes mellitus (E10-E14)	1,045	4.4	...
5	Influenza and pneumonia (J10-J18)	849	3.6	...
6	Alzheimer's disease (G30)	834	3.5	...
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	759	3.2	...
8	Chronic lower respiratory diseases (J40-J47)	710	3.0	...
9	Septicemia (A40-A41)	632	2.7	...
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	512	2.2	...
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	289	1.2	...
12	Pneumonitis due to solids and liquids (J69)	259	1.1	...
13	Atherosclerosis (I70)	150	0.6	...
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	108	0.5	...
15	Parkinson's disease (G20-G21)	103	0.4	...
...	All other causes (Residual)	3,305	14.0	...
Black, non-Hispanic, both sexes, 90-94 years				
...	All causes	15,316	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,623	36.7	...
2	Malignant neoplasms (C00-C97)	1,819	11.9	...
3	Cerebrovascular diseases (I60-I69)	1,546	10.1	...
4	Influenza and pneumonia (J10-J18)	651	4.3	...
5	Alzheimer's disease (G30)	611	4.0	...
6	Diabetes mellitus (E10-E14)	516	3.4	...
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	462	3.0	...
8	Septicemia (A40-A41)	381	2.5	...
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	351	2.3	...
10	Chronic lower respiratory diseases (J40-J47)	347	2.3	...
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	195	1.3	...
12	Pneumonitis due to solids and liquids (J69)	184	1.2	...
13	Atherosclerosis (I70)	135	0.9	...
14	Anemias (D50-D64)	68	0.4	...
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	67	0.4	...
...	All other causes (Residual)	2,360	15.4	...

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 95-99 years				
...	All causes	6,521	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,529	38.8	...
2	Cerebrovascular diseases (I60-I69)	610	9.4	...
3	Malignant neoplasms (C00-C97)	550	8.4	...
4	Influenza and pneumonia (J10-J18)	310	4.8	...
5	Alzheimer's disease (G30)	288	4.4	...
6	Diabetes mellitus (E10-E14)	195	3.0	...
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	195	3.0	...
8	Septicemia (A40-A41)	166	2.5	...
9	Chronic lower respiratory diseases (J40-J47)	138	2.1	...
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	130	2.0	...
11	Pneumonitis due to solids and liquids (J69)	93	1.4	...
12	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	76	1.2	...
13	Atherosclerosis (I70)	68	1.0	...
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	34	0.5	...
15	Anemias (D50-D64)	33	0.5	...
...	All other causes (Residual)	1,106	17.0	...
Black, non-Hispanic, both sexes, 100 years and over				
...	All causes	2,054	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	875	42.6	...
2	Cerebrovascular diseases (I60-I69)	168	8.2	...
3	Malignant neoplasms (C00-C97)	109	5.3	...
4	Influenza and pneumonia (J10-J18)	107	5.2	...
5	Alzheimer's disease (G30)	81	3.9	...
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	52	2.5	...
7	Septicemia (A40-A41)	47	2.3	...
8	Diabetes mellitus (E10-E14)	45	2.2	...
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	40	1.9	...
10	Atherosclerosis (I70)	37	1.8	...
11	Chronic lower respiratory diseases (J40-J47)	31	1.5	...
12	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	20	1.0	...
13	Pneumonitis due to solids and liquids (J69)	17	0.8	...
14	Nutritional deficiencies (E40-E64)	13	0.6	...
15	Aortic aneurysm and dissection (I71)	5	0.2	...
...	All other causes (Residual)	407	19.8	...

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, all ages				
...	All causes	144,802	100.0	842.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	36,596	25.3	212.9
2	Malignant neoplasms (C00-C97)	32,287	22.3	187.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,428	5.8	49.0
4	Cerebrovascular diseases (I60-I69)	7,754	5.4	45.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6,780	4.7	39.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	5,203	3.6	30.3
7	Diabetes mellitus (E10-E14)	5,158	3.6	30.0
8	Chronic lower respiratory diseases (J40-J47)	4,282	3.0	24.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,389	2.3	19.7
10	Influenza and pneumonia (J10-J18)	2,734	1.9	15.9
11	Septicemia (A40-A41)	2,668	1.8	15.5
12	Certain conditions originating in the perinatal period (P00-P96)	2,640	1.8	15.4
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,701	1.2	9.9
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,595	1.1	9.3
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,592	1.1	9.3
...	All other causes (Residual)	21,995	15.2	127.9
Black, non-Hispanic, male, 1-4 years				
...	All causes	681	100.0	55.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	231	33.9	18.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	89	13.1	7.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	58	8.5	4.8
4	Malignant neoplasms (C00-C97)	33	4.8	2.7
5	Influenza and pneumonia (J10-J18)	22	3.2	1.8
6	Chronic lower respiratory diseases (J40-J47)	20	2.9	1.6
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	18	2.6	*
7	Certain conditions originating in the perinatal period (P00-P96)	18	2.6	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	8	1.2	*
10	Cerebrovascular diseases (I60-I69)	6	0.9	*
11	Septicemia (A40-A41)	5	0.7	*
12	Anemias (D50-D64)	4	0.6	*
13	Meningitis (G00,G03)	3	0.4	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.4	*
15	Meningococcal infection (A39)	2	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.3	*
...	All other causes (Residual)	159	23.3	13.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 5-9 years				
...	All causes	392	100.0	24.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	180	45.9	11.4
2	Malignant neoplasms (C00-C97)	52	13.3	3.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	24	6.1	1.5
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	22	5.6	1.4
5	Chronic lower respiratory diseases (J40-J47)	11	2.8	*
6	Anemias (D50-D64)	7	1.8	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	6	1.5	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	6	1.5	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	4	1.0	*
9	Cerebrovascular diseases (I60-I69)	4	1.0	*
9	Influenza and pneumonia (J10-J18)	4	1.0	*
9	Complications of medical and surgical care (Y40-Y84,Y88)	4	1.0	*
13	Septicemia (A40-A41)	3	0.8	*
14	Certain conditions originating in the perinatal period (P00-P96)	2	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1	0.3	*
...	All other causes (Residual)	61	15.6	3.9
Black, non-Hispanic, male, 10-14 years				
...	All causes	585	100.0	34.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	211	36.1	12.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	60	10.3	3.5
3	Malignant neoplasms (C00-C97)	51	8.7	3.0
4	Chronic lower respiratory diseases (J40-J47)	28	4.8	1.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	28	4.8	1.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	27	4.6	1.6
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	25	4.3	1.5
8	Cerebrovascular diseases (I60-I69)	11	1.9	*
9	Anemias (D50-D64)	7	1.2	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	6	1.0	*
10	Influenza and pneumonia (J10-J18)	6	1.0	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	6	1.0	*
13	Diabetes mellitus (E10-E14)	5	0.9	*
13	Meningitis (G00,G03)	5	0.9	*
15	Meningococcal infection (A39)	4	0.7	*
15	Septicemia (A40-A41)	4	0.7	*
15	Certain conditions originating in the perinatal period (P00-P96)	4	0.7	*
...	All other causes (Residual)	97	16.6	5.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 15-19 years				
...	All causes	1,914	100.0	124.1
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	845	44.1	54.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	541	28.3	35.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	107	5.6	6.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	68	3.6	4.4
5	Malignant neoplasms (C00-C97)	64	3.3	4.2
6	Chronic lower respiratory diseases (J40-J47)	36	1.9	2.3
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	33	1.7	2.1
8	Anemias (D50-D64)	16	0.8	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	11	0.6	*
10	Cerebrovascular diseases (I60-I69)	10	0.5	*
10	Legal intervention (Y35,Y89.0)	10	0.5	*
12	Diabetes mellitus (E10-E14)	9	0.5	*
13	Influenza and pneumonia (J10-J18)	8	0.4	*
14	Septicemia (A40-A41)	5	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.2	*
...	All other causes (Residual)	147	7.7	9.5
Black, non-Hispanic, male, 20-24 years				
...	All causes	3,334	100.0	235.3
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,695	50.8	119.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	719	21.6	50.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	235	7.0	16.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	117	3.5	8.3
5	Malignant neoplasms (C00-C97)	73	2.2	5.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	49	1.5	3.5
7	Anemias (D50-D64)	33	1.0	2.3
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	26	0.8	1.8
8	Legal intervention (Y35,Y89.0)	26	0.8	1.8
10	Diabetes mellitus (E10-E14)	25	0.7	1.8
11	Chronic lower respiratory diseases (J40-J47)	22	0.7	1.6
12	Cerebrovascular diseases (I60-I69)	12	0.4	*
12	Influenza and pneumonia (J10-J18)	12	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	0.3	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	6	0.2	*
...	All other causes (Residual)	273	8.2	19.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 25-29 years				
...	All causes	3,179	100.0	266.6
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,246	39.2	104.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	665	20.9	55.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	244	7.7	20.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	220	6.9	18.5
5	Human immunodeficiency virus (HIV) disease (B20-B24)	145	4.6	12.2
6	Malignant neoplasms (C00-C97)	102	3.2	8.6
7	Anemias (D50-D64)	39	1.2	3.3
8	Diabetes mellitus (E10-E14)	37	1.2	3.1
9	Chronic lower respiratory diseases (J40-J47)	28	0.9	2.3
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	22	0.7	1.8
11	Cerebrovascular diseases (I60-I69)	20	0.6	1.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20	0.6	1.7
13	Legal intervention (Y35,Y89.0)	19	0.6	*
14	Septicemia (A40-A41)	15	0.5	*
15	Influenza and pneumonia (J10-J18)	13	0.4	*
...	All other causes (Residual)	344	10.8	28.9
Black, non-Hispanic, male, 30-34 years				
...	All causes	3,544	100.0	281.5
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	846	23.9	67.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	614	17.3	48.8
3	Human immunodeficiency virus (HIV) disease (B20-B24)	479	13.5	38.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	392	11.1	31.1
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	197	5.6	15.6
6	Malignant neoplasms (C00-C97)	185	5.2	14.7
7	Diabetes mellitus (E10-E14)	75	2.1	6.0
8	Cerebrovascular diseases (I60-I69)	52	1.5	4.1
9	Anemias (D50-D64)	33	0.9	2.6
10	Septicemia (A40-A41)	32	0.9	2.5
11	Influenza and pneumonia (J10-J18)	30	0.8	2.4
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	29	0.8	2.3
13	Chronic lower respiratory diseases (J40-J47)	28	0.8	2.2
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	21	0.6	1.7
15	Chronic liver disease and cirrhosis (K70,K73-K74)	19	0.5	*
...	All other causes (Residual)	512	14.4	40.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 35-39 years				
...	All causes	4,906	100.0	376.4
1	Human immunodeficiency virus (HIV) disease (B20-B24)	911	18.6	69.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	786	16.0	60.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	730	14.9	56.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	607	12.4	46.6
5	Malignant neoplasms (C00-C97)	345	7.0	26.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	191	3.9	14.7
7	Diabetes mellitus (E10-E14)	114	2.3	8.7
8	Cerebrovascular diseases (I60-I69)	102	2.1	7.8
9	Chronic lower respiratory diseases (J40-J47)	60	1.2	4.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	59	1.2	4.5
11	Influenza and pneumonia (J10-J18)	54	1.1	4.1
12	Septicemia (A40-A41)	50	1.0	3.8
13	Chronic liver disease and cirrhosis (K70,K73-K74)	49	1.0	3.8
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	48	1.0	3.7
15	Anemias (D50-D64)	35	0.7	2.7
...	All other causes (Residual)	765	15.6	58.7
Black, non-Hispanic, male, 40-44 years				
...	All causes	6,755	100.0	514.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,301	19.3	99.1
2	Human immunodeficiency virus (HIV) disease (B20-B24)	1,067	15.8	81.3
3	Malignant neoplasms (C00-C97)	813	12.0	61.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	776	11.5	59.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	435	6.4	33.1
6	Cerebrovascular diseases (I60-I69)	229	3.4	17.4
7	Diabetes mellitus (E10-E14)	202	3.0	15.4
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	154	2.3	11.7
9	Chronic liver disease and cirrhosis (K70,K73-K74)	131	1.9	10.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	102	1.5	7.8
11	Chronic lower respiratory diseases (J40-J47)	90	1.3	6.9
12	Influenza and pneumonia (J10-J18)	82	1.2	6.2
13	Septicemia (A40-A41)	80	1.2	6.1
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	64	0.9	4.9
15	Viral hepatitis (B15-B19)	51	0.8	3.9
...	All other causes (Residual)	1,178	17.4	89.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 45-49 years				
...	All causes	9,372	100.0	818.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,184	23.3	190.7
2	Malignant neoplasms (C00-C97)	1,612	17.2	140.8
3	Human immunodeficiency virus (HIV) disease (B20-B24)	999	10.7	87.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	836	8.9	73.0
5	Cerebrovascular diseases (I60-I69)	398	4.2	34.8
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	359	3.8	31.4
7	Diabetes mellitus (E10-E14)	299	3.2	26.1
8	Chronic liver disease and cirrhosis (K70,K73-K74)	274	2.9	23.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	172	1.8	15.0
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	129	1.4	11.3
11	Viral hepatitis (B15-B19)	121	1.3	10.6
12	Septicemia (A40-A41)	120	1.3	10.5
13	Influenza and pneumonia (J10-J18)	116	1.2	10.1
14	Chronic lower respiratory diseases (J40-J47)	115	1.2	10.0
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	109	1.2	9.5
...	All other causes (Residual)	1,529	16.3	133.5
Black, non-Hispanic, male, 50-54 years				
...	All causes	11,467	100.0	1,229.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,030	26.4	324.8
2	Malignant neoplasms (C00-C97)	2,591	22.6	277.8
3	Human immunodeficiency virus (HIV) disease (B20-B24)	792	6.9	84.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	687	6.0	73.6
5	Cerebrovascular diseases (I60-I69)	588	5.1	63.0
6	Diabetes mellitus (E10-E14)	445	3.9	47.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	363	3.2	38.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	256	2.2	27.4
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	213	1.9	22.8
10	Septicemia (A40-A41)	192	1.7	20.6
11	Chronic lower respiratory diseases (J40-J47)	189	1.6	20.3
12	Viral hepatitis (B15-B19)	141	1.2	15.1
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	125	1.1	13.4
14	Influenza and pneumonia (J10-J18)	123	1.1	13.2
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	83	0.7	8.9
...	All other causes (Residual)	1,649	14.4	176.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 55-59 years				
...	All causes	11,414	100.0	1,751.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,338	29.2	512.2
2	Malignant neoplasms (C00-C97)	3,223	28.2	494.5
3	Cerebrovascular diseases (I60-I69)	592	5.2	90.8
4	Diabetes mellitus (E10-E14)	512	4.5	78.6
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	470	4.1	72.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	332	2.9	50.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	267	2.3	41.0
8	Chronic lower respiratory diseases (J40-J47)	250	2.2	38.4
9	Chronic liver disease and cirrhosis (K70,K73-K74)	243	2.1	37.3
10	Septicemia (A40-A41)	191	1.7	29.3
11	Influenza and pneumonia (J10-J18)	158	1.4	24.2
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	145	1.3	22.2
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	112	1.0	17.2
14	Viral hepatitis (B15-B19)	82	0.7	12.6
15	Aortic aneurysm and dissection (I71)	73	0.6	11.2
...	All other causes (Residual)	1,426	12.5	218.8
Black, non-Hispanic, male, 60-64 years				
...	All causes	12,292	100.0	2,503.4
1	Malignant neoplasms (C00-C97)	3,883	31.6	790.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,690	30.0	751.5
3	Cerebrovascular diseases (I60-I69)	696	5.7	141.7
4	Diabetes mellitus (E10-E14)	592	4.8	120.6
5	Chronic lower respiratory diseases (J40-J47)	357	2.9	72.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	321	2.6	65.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	306	2.5	62.3
8	Septicemia (A40-A41)	227	1.8	46.2
9	Chronic liver disease and cirrhosis (K70,K73-K74)	193	1.6	39.3
10	Human immunodeficiency virus (HIV) disease (B20-B24)	180	1.5	36.7
10	Influenza and pneumonia (J10-J18)	180	1.5	36.7
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	129	1.0	26.3
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	68	0.6	13.8
14	Aortic aneurysm and dissection (I71)	65	0.5	13.2
15	Viral hepatitis (B15-B19)	40	0.3	8.1
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	40	0.3	8.1
...	All other causes (Residual)	1,325	10.8	269.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 65-69 years				
...	All causes	13,292	100.0	3,402.3
1	Malignant neoplasms (C00-C97)	4,276	32.2	1,094.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,965	29.8	1,014.9
3	Cerebrovascular diseases (I60-I69)	848	6.4	217.1
4	Diabetes mellitus (E10-E14)	658	5.0	168.4
5	Chronic lower respiratory diseases (J40-J47)	461	3.5	118.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	353	2.7	90.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	295	2.2	75.5
8	Septicemia (A40-A41)	286	2.2	73.2
9	Influenza and pneumonia (J10-J18)	188	1.4	48.1
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	159	1.2	40.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	137	1.0	35.1
12	Human immunodeficiency virus (HIV) disease (B20-B24)	125	0.9	32.0
13	Pneumonitis due to solids and liquids (J69)	67	0.5	17.1
14	Aortic aneurysm and dissection (I71)	58	0.4	14.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	50	0.4	12.8
...	All other causes (Residual)	1,366	10.3	349.7
Black, non-Hispanic, male, 70-74 years				
...	All causes	14,567	100.0	4,983.1
1	Malignant neoplasms (C00-C97)	4,559	31.3	1,559.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,282	29.4	1,464.8
3	Cerebrovascular diseases (I60-I69)	978	6.7	334.6
4	Diabetes mellitus (E10-E14)	662	4.5	226.5
5	Chronic lower respiratory diseases (J40-J47)	632	4.3	216.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	441	3.0	150.9
7	Septicemia (A40-A41)	312	2.1	106.7
8	Influenza and pneumonia (J10-J18)	266	1.8	91.0
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	232	1.6	79.4
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	200	1.4	68.4
11	Alzheimer's disease (G30)	86	0.6	29.4
12	Pneumonitis due to solids and liquids (J69)	85	0.6	29.1
13	Chronic liver disease and cirrhosis (K70,K73-K74)	84	0.6	28.7
14	Aortic aneurysm and dissection (I71)	74	0.5	25.3
15	Human immunodeficiency virus (HIV) disease (B20-B24)	57	0.4	19.5
...	All other causes (Residual)	1,617	11.1	553.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 75-79 years				
...	All causes	15,246	100.0	7,157.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,503	29.5	2,114.1
2	Malignant neoplasms (C00-C97)	4,413	28.9	2,071.8
3	Cerebrovascular diseases (I60-I69)	1,084	7.1	508.9
4	Chronic lower respiratory diseases (J40-J47)	735	4.8	345.1
5	Diabetes mellitus (E10-E14)	634	4.2	297.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	444	2.9	208.4
7	Septicemia (A40-A41)	395	2.6	185.4
8	Influenza and pneumonia (J10-J18)	373	2.4	175.1
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	254	1.7	119.2
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	228	1.5	107.0
11	Alzheimer's disease (G30)	175	1.1	82.2
12	Pneumonitis due to solids and liquids (J69)	135	0.9	63.4
13	Atherosclerosis (I70)	76	0.5	35.7
14	Parkinson's disease (G20-G21)	66	0.4	31.0
14	Aortic aneurysm and dissection (I71)	66	0.4	31.0
...	All other causes (Residual)	1,665	10.9	781.7
Black, non-Hispanic, male, 80-84 years				
...	All causes	12,891	100.0	10,054.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,925	30.4	3,061.4
2	Malignant neoplasms (C00-C97)	3,256	25.3	2,539.6
3	Cerebrovascular diseases (I60-I69)	947	7.3	738.6
4	Chronic lower respiratory diseases (J40-J47)	615	4.8	479.7
5	Diabetes mellitus (E10-E14)	501	3.9	390.8
6	Influenza and pneumonia (J10-J18)	416	3.2	324.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	365	2.8	284.7
8	Septicemia (A40-A41)	334	2.6	260.5
9	Alzheimer's disease (G30)	243	1.9	189.5
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	214	1.7	166.9
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	194	1.5	151.3
12	Pneumonitis due to solids and liquids (J69)	118	0.9	92.0
13	Atherosclerosis (I70)	77	0.6	60.1
14	Parkinson's disease (G20-G21)	72	0.6	56.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	70	0.5	54.6
...	All other causes (Residual)	1,544	12.0	1,204.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 85-89 years				
...	All causes	8,195	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,580	31.5	...
2	Malignant neoplasms (C00-C97)	1,755	21.4	...
3	Cerebrovascular diseases (I60-I69)	671	8.2	...
4	Chronic lower respiratory diseases (J40-J47)	378	4.6	...
5	Influenza and pneumonia (J10-J18)	318	3.9	...
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	282	3.4	...
7	Diabetes mellitus (E10-E14)	259	3.2	...
8	Septicemia (A40-A41)	212	2.6	...
9	Alzheimer's disease (G30)	199	2.4	...
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	138	1.7	...
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	129	1.6	...
12	Pneumonitis due to solids and liquids (J69)	105	1.3	...
13	Parkinson's disease (G20-G21)	56	0.7	...
14	Atherosclerosis (I70)	48	0.6	...
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	41	0.5	...
...	All other causes (Residual)	1,024	12.5	...
Black, non-Hispanic, male, 90-94 years				
...	All causes	4,407	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,496	33.9	...
2	Malignant neoplasms (C00-C97)	770	17.5	...
3	Cerebrovascular diseases (I60-I69)	349	7.9	...
4	Influenza and pneumonia (J10-J18)	208	4.7	...
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	165	3.7	...
6	Chronic lower respiratory diseases (J40-J47)	155	3.5	...
7	Alzheimer's disease (G30)	118	2.7	...
8	Septicemia (A40-A41)	108	2.5	...
9	Diabetes mellitus (E10-E14)	90	2.0	...
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	83	1.9	...
11	Pneumonitis due to solids and liquids (J69)	81	1.8	...
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	81	1.8	...
13	Atherosclerosis (I70)	35	0.8	...
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	22	0.5	...
14	Anemias (D50-D64)	22	0.5	...
...	All other causes (Residual)	624	14.2	...

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 95-99 years				
...	All causes	1,441	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	507	35.2	...
2	Malignant neoplasms (C00-C97)	187	13.0	...
3	Cerebrovascular diseases (I60-I69)	100	6.9	...
4	Influenza and pneumonia (J10-J18)	92	6.4	...
5	Chronic lower respiratory diseases (J40-J47)	54	3.7	...
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	54	3.7	...
7	Alzheimer's disease (G30)	50	3.5	...
8	Septicemia (A40-A41)	35	2.4	...
9	Diabetes mellitus (E10-E14)	33	2.3	...
10	Pneumonitis due to solids and liquids (J69)	30	2.1	...
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	26	1.8	...
12	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	25	1.7	...
13	Atherosclerosis (I70)	13	0.9	...
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	11	0.8	...
15	Anemias (D50-D64)	9	0.6	...
...	All other causes (Residual)	215	14.9	...
Black, non-Hispanic, male, 100 years and over				
...	All causes	379	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	135	35.6	...
2	Malignant neoplasms (C00-C97)	38	10.0	...
2	Cerebrovascular diseases (I60-I69)	38	10.0	...
4	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19	5.0	...
5	Influenza and pneumonia (J10-J18)	18	4.7	...
6	Alzheimer's disease (G30)	12	3.2	...
7	Septicemia (A40-A41)	10	2.6	...
7	Chronic lower respiratory diseases (J40-J47)	10	2.6	...
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	6	1.6	...
10	Diabetes mellitus (E10-E14)	5	1.3	...
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5	1.3	...
12	Atherosclerosis (I70)	4	1.1	...
13	Nutritional deficiencies (E40-E64)	2	0.5	...
14	Anemias (D50-D64)	1	0.3	...
14	Aortic aneurysm and dissection (I71)	1	0.3	...
14	Pneumoconioses and chemical effects (J60-J66,J68)	1	0.3	...
14	Pneumonitis due to solids and liquids (J69)	1	0.3	...
14	Diseases of appendix (K35-K38)	1	0.3	...
14	Hyperplasia of prostate (N40)	1	0.3	...
...	All other causes (Residual)	71	18.7	...

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, all ages				
...	All causes	141,771	100.0	748.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	40,098	28.3	211.6
2	Malignant neoplasms (C00-C97)	29,709	21.0	156.7
3	Cerebrovascular diseases (I60-I69)	10,937	7.7	57.7
4	Diabetes mellitus (E10-E14)	7,425	5.2	39.2
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,021	2.8	21.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,857	2.7	20.3
7	Chronic lower respiratory diseases (J40-J47)	3,448	2.4	18.2
8	Septicemia (A40-A41)	3,406	2.4	18.0
9	Influenza and pneumonia (J10-J18)	3,069	2.2	16.2
10	Human immunodeficiency virus (HIV) disease (B20-B24)	2,511	1.8	13.2
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,464	1.7	13.0
12	Alzheimer's disease (G30)	2,363	1.7	12.5
13	Certain conditions originating in the perinatal period (P00-P96)	2,137	1.5	11.3
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,367	1.0	7.2
15	Chronic liver disease and cirrhosis (K70,K73-K74)	953	0.7	5.0
...	All other causes (Residual)	24,006	16.9	126.7
Black, non-Hispanic, female, 1-4 years				
...	All causes	482	100.0	40.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	119	24.7	10.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	67	13.9	5.7
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	55	11.4	4.7
4	Malignant neoplasms (C00-C97)	38	7.9	3.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	22	4.6	1.9
6	Influenza and pneumonia (J10-J18)	18	3.7	*
7	Septicemia (A40-A41)	12	2.5	*
8	Chronic lower respiratory diseases (J40-J47)	9	1.9	*
9	Certain conditions originating in the perinatal period (P00-P96)	8	1.7	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	6	1.2	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.8	*
11	Cerebrovascular diseases (I60-I69)	4	0.8	*
13	Anemias (D50-D64)	3	0.6	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.6	*
15	Pneumonitis due to solids and liquids (J69)	2	0.4	*
...	All other causes (Residual)	112	23.2	9.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 5-9 years				
...	All causes	294	100.0	19.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	118	40.1	7.7
2	Malignant neoplasms (C00-C97)	29	9.9	1.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	21	7.1	1.4
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	16	5.4	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	13	4.4	*
6	Chronic lower respiratory diseases (J40-J47)	10	3.4	*
7	Anemias (D50-D64)	5	1.7	*
8	Septicemia (A40-A41)	4	1.4	*
9	Cerebrovascular diseases (I60-I69)	3	1.0	*
9	Influenza and pneumonia (J10-J18)	3	1.0	*
11	Certain conditions originating in the perinatal period (P00-P96)	2	0.7	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.7	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	1	0.3	*
13	Diabetes mellitus (E10-E14)	1	0.3	*
13	Meningitis (G00,G03)	1	0.3	*
13	Pneumonitis due to solids and liquids (J69)	1	0.3	*
13	Diseases of appendix (K35-K38)	1	0.3	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
...	All other causes (Residual)	62	21.1	4.0
Black, non-Hispanic, female, 10-14 years				
...	All causes	363	100.0	22.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	98	27.0	5.9
2	Malignant neoplasms (C00-C97)	36	9.9	2.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	27	7.4	1.6
4	Chronic lower respiratory diseases (J40-J47)	25	6.9	1.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	19	5.2	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	18	5.0	*
7	Septicemia (A40-A41)	9	2.5	*
7	Human immunodeficiency virus (HIV) disease (B20-B24)	9	2.5	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	8	2.2	*
9	Diabetes mellitus (E10-E14)	8	2.2	*
11	Anemias (D50-D64)	7	1.9	*
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6	1.7	*
13	Cerebrovascular diseases (I60-I69)	5	1.4	*
13	Influenza and pneumonia (J10-J18)	5	1.4	*
15	Meningococcal infection (A39)	4	1.1	*
15	Meningitis (G00,G03)	4	1.1	*
...	All other causes (Residual)	75	20.7	4.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 15-19 years				
...	All causes	628	100.0	42.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	199	31.7	13.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	122	19.4	8.2
3	Malignant neoplasms (C00-C97)	46	7.3	3.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	37	5.9	2.5
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	19	3.0	*
6	Pregnancy, childbirth and the puerperium (O00-099)	16	2.5	*
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	15	2.4	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	14	2.2	*
9	Anemias (D50-D64)	11	1.8	*
10	Influenza and pneumonia (J10-J18)	10	1.6	*
11	Diabetes mellitus (E10-E14)	7	1.1	*
11	Chronic lower respiratory diseases (J40-J47)	7	1.1	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	1.0	*
14	Septicemia (A40-A41)	4	0.6	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	3	0.5	*
15	Cerebrovascular diseases (I60-I69)	3	0.5	*
...	All other causes (Residual)	109	17.4	7.3
Black, non-Hispanic, female, 20-24 years				
...	All causes	1,028	100.0	70.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	221	21.5	15.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	192	18.7	13.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	77	7.5	5.3
4	Malignant neoplasms (C00-C97)	59	5.7	4.0
5	Human immunodeficiency virus (HIV) disease (B20-B24)	50	4.9	3.4
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	35	3.4	2.4
7	Pregnancy, childbirth and the puerperium (O00-099)	29	2.8	2.0
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	20	1.9	1.4
9	Anemias (D50-D64)	18	1.8	*
10	Diabetes mellitus (E10-E14)	16	1.6	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	1.5	*
12	Cerebrovascular diseases (I60-I69)	13	1.3	*
12	Chronic lower respiratory diseases (J40-J47)	13	1.3	*
14	Septicemia (A40-A41)	10	1.0	*
15	Influenza and pneumonia (J10-J18)	8	0.8	*
...	All other causes (Residual)	252	24.5	17.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 25-29 years				
...	All causes	1,308	100.0	100.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	188	14.4	14.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	166	12.7	12.7
3	Human immunodeficiency virus (HIV) disease (B20-B24)	161	12.3	12.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	132	10.1	10.1
5	Malignant neoplasms (C00-C97)	109	8.3	8.3
6	Pregnancy, childbirth and the puerperium (O00-099)	37	2.8	2.8
7	Diabetes mellitus (E10-E14)	36	2.8	2.8
8	Cerebrovascular diseases (I60-I69)	30	2.3	2.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	28	2.1	2.1
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	27	2.1	2.1
11	Septicemia (A40-A41)	26	2.0	2.0
12	Anemias (D50-D64)	25	1.9	1.9
13	Chronic lower respiratory diseases (J40-J47)	15	1.1	*
14	Influenza and pneumonia (J10-J18)	14	1.1	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	13	1.0	*
...	All other causes (Residual)	301	23.0	23.0
Black, non-Hispanic, female, 30-34 years				
...	All causes	1,978	100.0	141.0
1	Human immunodeficiency virus (HIV) disease (B20-B24)	325	16.4	23.2
2	Malignant neoplasms (C00-C97)	267	13.5	19.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	249	12.6	17.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	209	10.6	14.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	167	8.4	11.9
6	Cerebrovascular diseases (I60-I69)	64	3.2	4.6
7	Diabetes mellitus (E10-E14)	46	2.3	3.3
8	Chronic lower respiratory diseases (J40-J47)	31	1.6	2.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	31	1.6	2.2
10	Anemias (D50-D64)	30	1.5	2.1
11	Septicemia (A40-A41)	29	1.5	2.1
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	29	1.5	2.1
13	Pregnancy, childbirth and the puerperium (O00-099)	27	1.4	1.9
14	Influenza and pneumonia (J10-J18)	23	1.2	1.6
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	19	1.0	*
...	All other causes (Residual)	432	21.8	30.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 35-39 years				
...	All causes	3,196	100.0	217.4
1	Malignant neoplasms (C00-C97)	562	17.6	38.2
2	Human immunodeficiency virus (HIV) disease (B20-B24)	457	14.3	31.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	437	13.7	29.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	298	9.3	20.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	163	5.1	11.1
6	Cerebrovascular diseases (I60-I69)	135	4.2	9.2
7	Diabetes mellitus (E10-E14)	94	2.9	6.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	55	1.7	3.7
9	Chronic lower respiratory diseases (J40-J47)	49	1.5	3.3
10	Influenza and pneumonia (J10-J18)	47	1.5	3.2
11	Septicemia (A40-A41)	46	1.4	3.1
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	38	1.2	2.6
13	Anemias (D50-D64)	37	1.2	2.5
14	Chronic liver disease and cirrhosis (K70,K73-K74)	35	1.1	2.4
15	Pregnancy, childbirth and the puerperium (O00-099)	34	1.1	2.3
...	All other causes (Residual)	709	22.2	48.2
Black, non-Hispanic, female, 40-44 years				
...	All causes	5,098	100.0	342.0
1	Malignant neoplasms (C00-C97)	1,150	22.6	77.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	844	16.6	56.6
3	Human immunodeficiency virus (HIV) disease (B20-B24)	576	11.3	38.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	367	7.2	24.6
5	Cerebrovascular diseases (I60-I69)	275	5.4	18.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	152	3.0	10.2
7	Diabetes mellitus (E10-E14)	144	2.8	9.7
8	Chronic liver disease and cirrhosis (K70,K73-K74)	98	1.9	6.6
9	Chronic lower respiratory diseases (J40-J47)	93	1.8	6.2
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	90	1.8	6.0
11	Septicemia (A40-A41)	78	1.5	5.2
12	Influenza and pneumonia (J10-J18)	67	1.3	4.5
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	49	1.0	3.3
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	46	0.9	3.1
15	Anemias (D50-D64)	39	0.8	2.6
...	All other causes (Residual)	1,030	20.2	69.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 45-49 years				
...	All causes	6,630	100.0	501.4
1	Malignant neoplasms (C00-C97)	1,808	27.3	136.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,322	19.9	100.0
3	Human immunodeficiency virus (HIV) disease (B20-B24)	442	6.7	33.4
4	Cerebrovascular diseases (I60-I69)	409	6.2	30.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	375	5.7	28.4
6	Diabetes mellitus (E10-E14)	244	3.7	18.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	166	2.5	12.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	124	1.9	9.4
9	Septicemia (A40-A41)	118	1.8	8.9
10	Chronic lower respiratory diseases (J40-J47)	115	1.7	8.7
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	83	1.3	6.3
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	79	1.2	6.0
13	Influenza and pneumonia (J10-J18)	73	1.1	5.5
14	Viral hepatitis (B15-B19)	59	0.9	4.5
15	Anemias (D50-D64)	47	0.7	3.6
...	All other causes (Residual)	1,166	17.6	88.2
Black, non-Hispanic, female, 50-54 years				
...	All causes	7,742	100.0	702.1
1	Malignant neoplasms (C00-C97)	2,365	30.5	214.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,782	23.0	161.6
3	Cerebrovascular diseases (I60-I69)	479	6.2	43.4
4	Diabetes mellitus (E10-E14)	405	5.2	36.7
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	247	3.2	22.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	236	3.0	21.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	193	2.5	17.5
8	Chronic lower respiratory diseases (J40-J47)	170	2.2	15.4
9	Septicemia (A40-A41)	163	2.1	14.8
10	Chronic liver disease and cirrhosis (K70,K73-K74)	153	2.0	13.9
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	100	1.3	9.1
12	Influenza and pneumonia (J10-J18)	87	1.1	7.9
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	55	0.7	5.0
14	Viral hepatitis (B15-B19)	47	0.6	4.3
15	Anemias (D50-D64)	39	0.5	3.5
...	All other causes (Residual)	1,221	15.8	110.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 55-59 years				
...	All causes	8,069	100.0	1,014.9
1	Malignant neoplasms (C00-C97)	2,676	33.2	336.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,051	25.4	258.0
3	Diabetes mellitus (E10-E14)	491	6.1	61.8
4	Cerebrovascular diseases (I60-I69)	456	5.7	57.4
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	230	2.9	28.9
6	Septicemia (A40-A41)	209	2.6	26.3
7	Chronic lower respiratory diseases (J40-J47)	204	2.5	25.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	148	1.8	18.6
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	127	1.6	16.0
10	Chronic liver disease and cirrhosis (K70,K73-K74)	117	1.4	14.7
11	Human immunodeficiency virus (HIV) disease (B20-B24)	113	1.4	14.2
12	Influenza and pneumonia (J10-J18)	96	1.2	12.1
13	Viral hepatitis (B15-B19)	31	0.4	3.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	27	0.3	3.4
14	Anemias (D50-D64)	27	0.3	3.4
...	All other causes (Residual)	1,066	13.2	134.1
Black, non-Hispanic, female, 60-64 years				
...	All causes	9,103	100.0	1,448.4
1	Malignant neoplasms (C00-C97)	2,911	32.0	463.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,475	27.2	393.8
3	Diabetes mellitus (E10-E14)	609	6.7	96.9
4	Cerebrovascular diseases (I60-I69)	559	6.1	88.9
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	269	3.0	42.8
6	Chronic lower respiratory diseases (J40-J47)	256	2.8	40.7
7	Septicemia (A40-A41)	241	2.6	38.3
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	149	1.6	23.7
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	136	1.5	21.6
10	Influenza and pneumonia (J10-J18)	120	1.3	19.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	96	1.1	15.3
12	Human immunodeficiency virus (HIV) disease (B20-B24)	50	0.5	8.0
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	34	0.4	5.4
14	Viral hepatitis (B15-B19)	31	0.3	4.9
14	Pneumonitis due to solids and liquids (J69)	31	0.3	4.9
...	All other causes (Residual)	1,136	12.5	180.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 65-69 years				
...	All causes	11,216	100.0	2,128.3
1	Malignant neoplasms (C00-C97)	3,483	31.1	660.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,144	28.0	596.6
3	Diabetes mellitus (E10-E14)	855	7.6	162.2
4	Cerebrovascular diseases (I60-I69)	736	6.6	139.7
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	384	3.4	72.9
6	Chronic lower respiratory diseases (J40-J47)	341	3.0	64.7
7	Septicemia (A40-A41)	295	2.6	56.0
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	180	1.6	34.2
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	140	1.2	26.6
10	Influenza and pneumonia (J10-J18)	139	1.2	26.4
11	Chronic liver disease and cirrhosis (K70,K73-K74)	76	0.7	14.4
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	49	0.4	9.3
12	Aortic aneurysm and dissection (I71)	49	0.4	9.3
14	Pneumonitis due to solids and liquids (J69)	43	0.4	8.2
15	Alzheimer's disease (G30)	41	0.4	7.8
...	All other causes (Residual)	1,261	11.2	239.3
Black, non-Hispanic, female, 70-74 years				
...	All causes	13,673	100.0	3,136.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,020	29.4	922.2
2	Malignant neoplasms (C00-C97)	3,766	27.5	863.9
3	Cerebrovascular diseases (I60-I69)	1,038	7.6	238.1
4	Diabetes mellitus (E10-E14)	977	7.1	224.1
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	492	3.6	112.9
6	Chronic lower respiratory diseases (J40-J47)	458	3.3	105.1
7	Septicemia (A40-A41)	358	2.6	82.1
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	256	1.9	58.7
9	Influenza and pneumonia (J10-J18)	244	1.8	56.0
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	151	1.1	34.6
11	Alzheimer's disease (G30)	114	0.8	26.2
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	64	0.5	14.7
13	Aortic aneurysm and dissection (I71)	63	0.5	14.5
14	Chronic liver disease and cirrhosis (K70,K73-K74)	61	0.4	14.0
15	Pneumonitis due to solids and liquids (J69)	55	0.4	12.6
...	All other causes (Residual)	1,556	11.4	357.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 75-79 years				
...	All causes	16,830	100.0	4,718.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,360	31.8	1,502.6
2	Malignant neoplasms (C00-C97)	3,808	22.6	1,067.5
3	Cerebrovascular diseases (I60-I69)	1,492	8.9	418.3
4	Diabetes mellitus (E10-E14)	1,094	6.5	306.7
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	580	3.4	162.6
6	Chronic lower respiratory diseases (J40-J47)	505	3.0	141.6
7	Septicemia (A40-A41)	450	2.7	126.1
8	Influenza and pneumonia (J10-J18)	335	2.0	93.9
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	331	2.0	92.8
10	Alzheimer's disease (G30)	293	1.7	82.1
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	179	1.1	50.2
12	Pneumonitis due to solids and liquids (J69)	108	0.6	30.3
13	Atherosclerosis (I70)	78	0.5	21.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	77	0.5	21.6
14	Aortic aneurysm and dissection (I71)	77	0.5	21.6
...	All other causes (Residual)	2,063	12.3	578.3
Black, non-Hispanic, female, 80-84 years				
...	All causes	17,216	100.0	6,986.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,732	33.3	2,326.2
2	Malignant neoplasms (C00-C97)	3,041	17.7	1,234.1
3	Cerebrovascular diseases (I60-I69)	1,754	10.2	711.8
4	Diabetes mellitus (E10-E14)	984	5.7	399.3
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	552	3.2	224.0
6	Chronic lower respiratory diseases (J40-J47)	511	3.0	207.4
7	Alzheimer's disease (G30)	462	2.7	187.5
8	Influenza and pneumonia (J10-J18)	460	2.7	186.7
9	Septicemia (A40-A41)	454	2.6	184.2
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	360	2.1	146.1
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	172	1.0	69.8
12	Pneumonitis due to solids and liquids (J69)	138	0.8	56.0
13	Atherosclerosis (I70)	118	0.7	47.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	89	0.5	36.1
14	Aortic aneurysm and dissection (I71)	89	0.5	36.1
...	All other causes (Residual)	2,300	13.4	933.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 85-89 years				
...	All causes	15,479	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,423	35.0	...
2	Malignant neoplasms (C00-C97)	2,063	13.3	...
3	Cerebrovascular diseases (I60-I69)	1,627	10.5	...
4	Diabetes mellitus (E10-E14)	786	5.1	...
5	Alzheimer's disease (G30)	635	4.1	...
6	Influenza and pneumonia (J10-J18)	531	3.4	...
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	477	3.1	...
8	Septicemia (A40-A41)	420	2.7	...
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	374	2.4	...
10	Chronic lower respiratory diseases (J40-J47)	332	2.1	...
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	160	1.0	...
12	Pneumonitis due to solids and liquids (J69)	154	1.0	...
13	Atherosclerosis (I70)	102	0.7	...
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	67	0.4	...
14	Aortic aneurysm and dissection (I71)	67	0.4	...
...	All other causes (Residual)	2,261	14.6	...
Black, non-Hispanic, female, 90-94 years				
...	All causes	10,909	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,127	37.8	...
2	Cerebrovascular diseases (I60-I69)	1,197	11.0	...
3	Malignant neoplasms (C00-C97)	1,049	9.6	...
4	Alzheimer's disease (G30)	493	4.5	...
5	Influenza and pneumonia (J10-J18)	443	4.1	...
6	Diabetes mellitus (E10-E14)	426	3.9	...
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	297	2.7	...
8	Septicemia (A40-A41)	273	2.5	...
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	268	2.5	...
10	Chronic lower respiratory diseases (J40-J47)	192	1.8	...
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	114	1.0	...
12	Pneumonitis due to solids and liquids (J69)	103	0.9	...
13	Atherosclerosis (I70)	100	0.9	...
14	Nutritional deficiencies (E40-E64)	54	0.5	...
15	Anemias (D50-D64)	46	0.4	...
...	All other causes (Residual)	1,727	15.8	...

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 95-99 years				
...	All causes	5,080	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,022	39.8	...
2	Cerebrovascular diseases (I60-I69)	510	10.0	...
3	Malignant neoplasms (C00-C97)	363	7.1	...
4	Alzheimer's disease (G30)	238	4.7	...
5	Influenza and pneumonia (J10-J18)	218	4.3	...
6	Diabetes mellitus (E10-E14)	162	3.2	...
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	141	2.8	...
8	Septicemia (A40-A41)	131	2.6	...
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	104	2.0	...
10	Chronic lower respiratory diseases (J40-J47)	84	1.7	...
11	Pneumonitis due to solids and liquids (J69)	63	1.2	...
12	Atherosclerosis (I70)	55	1.1	...
13	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	51	1.0	...
14	Nutritional deficiencies (E40-E64)	30	0.6	...
15	Anemias (D50-D64)	24	0.5	...
...	All other causes (Residual)	884	17.4	...
Black, non-Hispanic, female, 100 years and over				
...	All causes	1,675	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	740	44.2	...
2	Cerebrovascular diseases (I60-I69)	130	7.8	...
3	Influenza and pneumonia (J10-J18)	89	5.3	...
4	Malignant neoplasms (C00-C97)	71	4.2	...
5	Alzheimer's disease (G30)	69	4.1	...
6	Diabetes mellitus (E10-E14)	40	2.4	...
7	Septicemia (A40-A41)	37	2.2	...
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	34	2.0	...
9	Atherosclerosis (I70)	33	2.0	...
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	33	2.0	...
11	Chronic lower respiratory diseases (J40-J47)	21	1.3	...
12	Pneumonitis due to solids and liquids (J69)	16	1.0	...
13	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15	0.9	...
14	Nutritional deficiencies (E40-E64)	11	0.7	...
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	4	0.2	...
15	Aortic aneurysm and dissection (I71)	4	0.2	...
15	Peptic ulcer (K25-K28)	4	0.2	...
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	4	0.2	...
...	All other causes (Residual)	320	19.1	...

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available