

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, all ages				
...	All causes	122,026	100.0	305.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	28,298	23.2	70.9
2	Malignant neoplasms (C00-C97)	24,070	19.7	60.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,418	8.5	26.1
4	Cerebrovascular diseases (I60-I69)	6,658	5.5	16.7
5	Diabetes mellitus (E10-E14)	6,179	5.1	15.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	3,382	2.8	8.5
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,355	2.7	8.4
8	Chronic lower respiratory diseases (J40-J47)	3,174	2.6	8.0
9	Influenza and pneumonia (J10-J18)	2,948	2.4	7.4
10	Certain conditions originating in the perinatal period (P00-P96)	2,628	2.2	6.6
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,170	1.8	5.4
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,007	1.6	5.0
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,890	1.5	4.7
14	Alzheimer's disease (G30)	1,835	1.5	4.6
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,787	1.5	4.5
...	All other causes (Residual)	21,227	17.4	53.2
Hispanic, both sexes, 1-4 years				
...	All causes	993	100.0	30.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	325	32.7	9.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	122	12.3	3.7
3	Malignant neoplasms (C00-C97)	98	9.9	3.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	82	8.3	2.5
5	Influenza and pneumonia (J10-J18)	37	3.7	1.1
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	35	3.5	1.1
7	Certain conditions originating in the perinatal period (P00-P96)	15	1.5	*
8	Chronic lower respiratory diseases (J40-J47)	11	1.1	*
9	Septicemia (A40-A41)	10	1.0	*
10	Cerebrovascular diseases (I60-I69)	8	0.8	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.7	*
12	Acute bronchitis and bronchiolitis (J20-J21)	6	0.6	*
13	Anemias (D50-D64)	5	0.5	*
14	Meningitis (G00,G03)	4	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.4	*
...	All other causes (Residual)	224	22.6	6.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 5-9 years				
...	All causes	538	100.0	14.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	218	40.5	5.7
2	Malignant neoplasms (C00-C97)	111	20.6	2.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	31	5.8	0.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	16	3.0	*
5	Influenza and pneumonia (J10-J18)	13	2.4	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	9	1.7	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	1.5	*
8	Septicemia (A40-A41)	5	0.9	*
8	Cerebrovascular diseases (I60-I69)	5	0.9	*
10	Chronic lower respiratory diseases (J40-J47)	4	0.7	*
10	Diseases of appendix (K35-K38)	4	0.7	*
10	Certain conditions originating in the perinatal period (P00-P96)	4	0.7	*
13	Anemias (D50-D64)	2	0.4	*
14	Salmonella infections (A01-A02)	1	0.2	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.2	*
14	Diabetes mellitus (E10-E14)	1	0.2	*
14	Meningitis (G00,G03)	1	0.2	*
14	Pneumonitis due to solids and liquids (J69)	1	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
...	All other causes (Residual)	102	19.0	2.7
Hispanic, both sexes, 10-14 years				
...	All causes	635	100.0	17.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	213	33.5	5.8
2	Malignant neoplasms (C00-C97)	106	16.7	2.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	43	6.8	1.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	37	5.8	1.0
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	34	5.4	0.9
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	22	3.5	0.6
7	Chronic lower respiratory diseases (J40-J47)	11	1.7	*
8	Cerebrovascular diseases (I60-I69)	9	1.4	*
8	Influenza and pneumonia (J10-J18)	9	1.4	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.8	*
11	Certain conditions originating in the perinatal period (P00-P96)	4	0.6	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.5	*
13	Septicemia (A40-A41)	2	0.3	*
13	Anemias (D50-D64)	2	0.3	*
13	Diabetes mellitus (E10-E14)	2	0.3	*
13	Meningitis (G00,G03)	2	0.3	*
...	All other causes (Residual)	131	20.6	3.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Hispanic, both sexes, 15-19 years			
...	All causes	2,183	100.0	67.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,005	46.0	31.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	471	21.6	14.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	192	8.8	5.9
4	Malignant neoplasms (C00-C97)	124	5.7	3.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	46	2.1	1.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	27	1.2	0.8
7	Chronic lower respiratory diseases (J40-J47)	12	0.5	*
8	Influenza and pneumonia (J10-J18)	11	0.5	*
9	Cerebrovascular diseases (I60-I69)	9	0.4	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.4	*
10	Pregnancy, childbirth and the puerperium (O00-099)	8	0.4	*
12	Septicemia (A40-A41)	7	0.3	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	7	0.3	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.3	*
14	Diabetes mellitus (E10-E14)	6	0.3	*
14	Legal intervention (Y35,Y89.0)	6	0.3	*
...	All other causes (Residual)	238	10.9	7.3
	Hispanic, both sexes, 20-24 years			
...	All causes	3,211	100.0	85.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,361	42.4	36.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	815	25.4	21.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	299	9.3	8.0
4	Malignant neoplasms (C00-C97)	164	5.1	4.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	89	2.8	2.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	30	0.9	0.8
7	Influenza and pneumonia (J10-J18)	25	0.8	0.7
8	Human immunodeficiency virus (HIV) disease (B20-B24)	24	0.7	0.6
9	Legal intervention (Y35,Y89.0)	22	0.7	0.6
10	Cerebrovascular diseases (I60-I69)	19	0.6	*
11	Pregnancy, childbirth and the puerperium (O00-099)	16	0.5	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	15	0.5	*
13	Diabetes mellitus (E10-E14)	13	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	0.3	*
15	Septicemia (A40-A41)	10	0.3	*
...	All other causes (Residual)	298	9.3	8.0

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total	
			deaths	Rate
	Hispanic, both sexes, 25-29 years			
...	All causes	3,088	100.0	81.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,208	39.1	31.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	592	19.2	15.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	251	8.1	6.6
4	Malignant neoplasms (C00-C97)	208	6.7	5.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	134	4.3	3.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	75	2.4	2.0
7	Cerebrovascular diseases (I60-I69)	35	1.1	0.9
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	35	1.1	0.9
9	Pregnancy, childbirth and the puerperium (O00-099)	31	1.0	0.8
10	Diabetes mellitus (E10-E14)	28	0.9	0.7
11	Influenza and pneumonia (J10-J18)	24	0.8	0.6
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	0.5	*
13	Chronic lower respiratory diseases (J40-J47)	14	0.5	*
13	Legal intervention (Y35,Y89.0)	14	0.5	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	0.4	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	13	0.4	*
...	All other causes (Residual)	398	12.9	10.5
	Hispanic, both sexes, 30-34 years			
...	All causes	3,101	100.0	86.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	959	30.9	26.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	379	12.2	10.6
3	Malignant neoplasms (C00-C97)	314	10.1	8.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	248	8.0	6.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	213	6.9	6.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	169	5.4	4.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	87	2.8	2.4
8	Cerebrovascular diseases (I60-I69)	54	1.7	1.5
9	Diabetes mellitus (E10-E14)	36	1.2	1.0
10	Influenza and pneumonia (J10-J18)	30	1.0	0.8
11	Septicemia (A40-A41)	23	0.7	0.6
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	23	0.7	0.6
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	0.6	*
14	Legal intervention (Y35,Y89.0)	18	0.6	*
15	Chronic lower respiratory diseases (J40-J47)	15	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	0.5	*
15	Pregnancy, childbirth and the puerperium (O00-099)	15	0.5	*
...	All other causes (Residual)	484	15.6	13.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 35-39 years				
...	All causes	3,683	100.0	116.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	948	25.7	29.9
2	Malignant neoplasms (C00-C97)	513	13.9	16.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	327	8.9	10.3
4	Human immunodeficiency virus (HIV) disease (B20-B24)	311	8.4	9.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	296	8.0	9.3
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	203	5.5	6.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	156	4.2	4.9
8	Cerebrovascular diseases (I60-I69)	80	2.2	2.5
9	Diabetes mellitus (E10-E14)	59	1.6	1.9
10	Septicemia (A40-A41)	31	0.8	1.0
11	Viral hepatitis (B15-B19)	29	0.8	0.9
12	Influenza and pneumonia (J10-J18)	27	0.7	0.9
13	Chronic lower respiratory diseases (J40-J47)	25	0.7	0.8
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	25	0.7	0.8
15	Pregnancy, childbirth and the puerperium (O00-099)	17	0.5	*
...	All other causes (Residual)	636	17.3	20.0
Hispanic, both sexes, 40-44 years				
...	All causes	4,893	100.0	178.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	929	19.0	33.9
2	Malignant neoplasms (C00-C97)	842	17.2	30.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	552	11.3	20.1
4	Human immunodeficiency virus (HIV) disease (B20-B24)	416	8.5	15.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	313	6.4	11.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	206	4.2	7.5
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	199	4.1	7.3
8	Cerebrovascular diseases (I60-I69)	187	3.8	6.8
9	Diabetes mellitus (E10-E14)	138	2.8	5.0
10	Viral hepatitis (B15-B19)	78	1.6	2.8
11	Septicemia (A40-A41)	55	1.1	2.0
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	51	1.0	1.9
13	Influenza and pneumonia (J10-J18)	47	1.0	1.7
14	Chronic lower respiratory diseases (J40-J47)	33	0.7	1.2
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	28	0.6	1.0
...	All other causes (Residual)	819	16.7	29.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total	
			deaths	Rate
Hispanic, both sexes, 45-49 years				
...	All causes	5,917	100.0	272.7
1	Malignant neoplasms (C00-C97)	1,316	22.2	60.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	865	14.6	39.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	735	12.4	33.9
4	Chronic liver disease and cirrhosis (K70,K73-K74)	464	7.8	21.4
5	Human immunodeficiency virus (HIV) disease (B20-B24)	368	6.2	17.0
6	Cerebrovascular diseases (I60-I69)	251	4.2	11.6
7	Diabetes mellitus (E10-E14)	219	3.7	10.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	161	2.7	7.4
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	158	2.7	7.3
10	Viral hepatitis (B15-B19)	124	2.1	5.7
11	Septicemia (A40-A41)	74	1.3	3.4
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	70	1.2	3.2
13	Influenza and pneumonia (J10-J18)	63	1.1	2.9
14	Chronic lower respiratory diseases (J40-J47)	61	1.0	2.8
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	37	0.6	1.7
...	All other causes (Residual)	951	16.1	43.8
Hispanic, both sexes, 50-54 years				
...	All causes	6,599	100.0	400.3
1	Malignant neoplasms (C00-C97)	1,765	26.7	107.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,170	17.7	71.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	519	7.9	31.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	463	7.0	28.1
5	Diabetes mellitus (E10-E14)	374	5.7	22.7
6	Cerebrovascular diseases (I60-I69)	318	4.8	19.3
7	Human immunodeficiency virus (HIV) disease (B20-B24)	238	3.6	14.4
8	Viral hepatitis (B15-B19)	142	2.2	8.6
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	114	1.7	6.9
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	104	1.6	6.3
11	Influenza and pneumonia (J10-J18)	88	1.3	5.3
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	88	1.3	5.3
13	Septicemia (A40-A41)	74	1.1	4.5
14	Chronic lower respiratory diseases (J40-J47)	73	1.1	4.4
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	38	0.6	2.3
...	All other causes (Residual)	1,031	15.6	62.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total	
			deaths	Rate
Hispanic, both sexes, 55-59 years				
...	All causes	7,177	100.0	590.7
1	Malignant neoplasms (C00-C97)	2,152	30.0	177.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,541	21.5	126.8
3	Diabetes mellitus (E10-E14)	492	6.9	40.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	433	6.0	35.6
5	Cerebrovascular diseases (I60-I69)	353	4.9	29.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	338	4.7	27.8
7	Human immunodeficiency virus (HIV) disease (B20-B24)	150	2.1	12.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	146	2.0	12.0
9	Chronic lower respiratory diseases (J40-J47)	114	1.6	9.4
10	Influenza and pneumonia (J10-J18)	96	1.3	7.9
11	Viral hepatitis (B15-B19)	93	1.3	7.7
12	Septicemia (A40-A41)	90	1.3	7.4
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	88	1.2	7.2
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	52	0.7	4.3
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	37	0.5	3.0
...	All other causes (Residual)	1,002	14.0	82.5
Hispanic, both sexes, 60-64 years				
...	All causes	7,736	100.0	886.5
1	Malignant neoplasms (C00-C97)	2,330	30.1	267.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,913	24.7	219.2
3	Diabetes mellitus (E10-E14)	629	8.1	72.1
4	Cerebrovascular diseases (I60-I69)	383	5.0	43.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	369	4.8	42.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	301	3.9	34.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	172	2.2	19.7
8	Chronic lower respiratory diseases (J40-J47)	152	2.0	17.4
9	Influenza and pneumonia (J10-J18)	128	1.7	14.7
10	Septicemia (A40-A41)	100	1.3	11.5
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	69	0.9	7.9
12	Viral hepatitis (B15-B19)	58	0.7	6.6
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	57	0.7	6.5
14	Human immunodeficiency virus (HIV) disease (B20-B24)	54	0.7	6.2
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	38	0.5	4.4
...	All other causes (Residual)	983	12.7	112.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total	
			deaths	Rate
Hispanic, both sexes, 65-69 years				
...	All causes	9,230	100.0	1,352.1
1	Malignant neoplasms (C00-C97)	2,789	30.2	408.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,410	26.1	353.0
3	Diabetes mellitus (E10-E14)	719	7.8	105.3
4	Cerebrovascular diseases (I60-I69)	529	5.7	77.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	322	3.5	47.2
6	Chronic lower respiratory diseases (J40-J47)	252	2.7	36.9
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	239	2.6	35.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	225	2.4	33.0
9	Influenza and pneumonia (J10-J18)	168	1.8	24.6
10	Septicemia (A40-A41)	130	1.4	19.0
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	86	0.9	12.6
12	Viral hepatitis (B15-B19)	59	0.6	8.6
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	49	0.5	7.2
14	Alzheimer's disease (G30)	44	0.5	6.4
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	44	0.5	6.4
...	All other causes (Residual)	1,165	12.6	170.7
Hispanic, both sexes, 70-74 years				
...	All causes	11,366	100.0	2,107.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,225	28.4	598.1
2	Malignant neoplasms (C00-C97)	3,140	27.6	582.3
3	Diabetes mellitus (E10-E14)	892	7.8	165.4
4	Cerebrovascular diseases (I60-I69)	673	5.9	124.8
5	Chronic lower respiratory diseases (J40-J47)	399	3.5	74.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	302	2.7	56.0
7	Influenza and pneumonia (J10-J18)	260	2.3	48.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	259	2.3	48.0
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	227	2.0	42.1
10	Septicemia (A40-A41)	140	1.2	26.0
11	Alzheimer's disease (G30)	135	1.2	25.0
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	118	1.0	21.9
13	Parkinson's disease (G20-G21)	71	0.6	13.2
14	Aortic aneurysm and dissection (I71)	63	0.6	11.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	57	0.5	10.6
...	All other causes (Residual)	1,405	12.4	260.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 75-79 years				
...	All causes	13,406	100.0	3,380.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,966	29.6	1,000.2
2	Malignant neoplasms (C00-C97)	3,248	24.2	819.1
3	Cerebrovascular diseases (I60-I69)	1,001	7.5	252.4
4	Diabetes mellitus (E10-E14)	945	7.0	238.3
5	Chronic lower respiratory diseases (J40-J47)	515	3.8	129.9
6	Influenza and pneumonia (J10-J18)	371	2.8	93.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	320	2.4	80.7
8	Alzheimer's disease (G30)	268	2.0	67.6
9	Chronic liver disease and cirrhosis (K70,K73-K74)	265	2.0	66.8
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	234	1.7	59.0
11	Septicemia (A40-A41)	199	1.5	50.2
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	136	1.0	34.3
13	Parkinson's disease (G20-G21)	112	0.8	28.2
14	Aortic aneurysm and dissection (I71)	99	0.7	25.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	88	0.7	22.2
...	All other causes (Residual)	1,639	12.2	413.3
Hispanic, both sexes, 80-84 years				
...	All causes	12,813	100.0	5,251.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,111	32.1	1,685.0
2	Malignant neoplasms (C00-C97)	2,453	19.1	1,005.4
3	Cerebrovascular diseases (I60-I69)	1,028	8.0	421.4
4	Diabetes mellitus (E10-E14)	798	6.2	327.1
5	Chronic lower respiratory diseases (J40-J47)	580	4.5	237.7
6	Influenza and pneumonia (J10-J18)	489	3.8	200.4
7	Alzheimer's disease (G30)	396	3.1	162.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	258	2.0	105.7
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	220	1.7	90.2
10	Septicemia (A40-A41)	198	1.5	81.2
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	149	1.2	61.1
12	Parkinson's disease (G20-G21)	146	1.1	59.8
13	Chronic liver disease and cirrhosis (K70,K73-K74)	132	1.0	54.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	97	0.8	39.8
15	Pneumonitis due to solids and liquids (J69)	83	0.6	34.0
...	All other causes (Residual)	1,675	13.1	686.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 85-89 years				
...	All causes	9,960	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,522	35.4	...
2	Malignant neoplasms (C00-C97)	1,450	14.6	...
3	Cerebrovascular diseases (I60-I69)	863	8.7	...
4	Chronic lower respiratory diseases (J40-J47)	491	4.9	...
5	Diabetes mellitus (E10-E14)	471	4.7	...
6	Influenza and pneumonia (J10-J18)	415	4.2	...
7	Alzheimer's disease (G30)	402	4.0	...
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	248	2.5	...
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	157	1.6	...
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	153	1.5	...
11	Septicemia (A40-A41)	143	1.4	...
12	Parkinson's disease (G20-G21)	92	0.9	...
13	Pneumonitis due to solids and liquids (J69)	89	0.9	...
14	Atherosclerosis (I70)	64	0.6	...
15	Aortic aneurysm and dissection (I71)	59	0.6	...
...	All other causes (Residual)	1,341	13.5	...
Hispanic, both sexes, 90-94 years				
...	All causes	6,835	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,686	39.3	...
2	Malignant neoplasms (C00-C97)	728	10.7	...
3	Cerebrovascular diseases (I60-I69)	581	8.5	...
4	Alzheimer's disease (G30)	364	5.3	...
5	Influenza and pneumonia (J10-J18)	344	5.0	...
6	Chronic lower respiratory diseases (J40-J47)	277	4.1	...
7	Diabetes mellitus (E10-E14)	267	3.9	...
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	136	2.0	...
9	Septicemia (A40-A41)	108	1.6	...
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	107	1.6	...
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	92	1.3	...
12	Pneumonitis due to solids and liquids (J69)	73	1.1	...
13	Atherosclerosis (I70)	64	0.9	...
14	Parkinson's disease (G20-G21)	44	0.6	...
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	30	0.4	...
...	All other causes (Residual)	934	13.7	...

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Hispanic, both sexes, 95-99 years			
...	All causes	2,703	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,067	39.5	...
2	Cerebrovascular diseases (I60-I69)	222	8.2	...
3	Influenza and pneumonia (J10-J18)	188	7.0	...
4	Malignant neoplasms (C00-C97)	181	6.7	...
5	Alzheimer's disease (G30)	155	5.7	...
6	Chronic lower respiratory diseases (J40-J47)	103	3.8	...
7	Diabetes mellitus (E10-E14)	83	3.1	...
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	56	2.1	...
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	52	1.9	...
10	Pneumonitis due to solids and liquids (J69)	46	1.7	...
11	Septicemia (A40-A41)	38	1.4	...
12	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	37	1.4	...
13	Atherosclerosis (I70)	29	1.1	...
14	Parkinson's disease (G20-G21)	14	0.5	...
15	Nutritional deficiencies (E40-E64)	13	0.5	...
...	All other causes (Residual)	419	15.5	...
	Hispanic, both sexes, 100 years and over			
...	All causes	631	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	314	49.8	...
2	Influenza and pneumonia (J10-J18)	49	7.8	...
3	Cerebrovascular diseases (I60-I69)	33	5.2	...
4	Alzheimer's disease (G30)	31	4.9	...
5	Malignant neoplasms (C00-C97)	24	3.8	...
6	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	18	2.9	...
7	Chronic lower respiratory diseases (J40-J47)	17	2.7	...
8	Pneumonitis due to solids and liquids (J69)	10	1.6	...
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	1.6	...
10	Atherosclerosis (I70)	9	1.4	...
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7	1.1	...
12	Diabetes mellitus (E10-E14)	5	0.8	...
13	Septicemia (A40-A41)	4	0.6	...
14	Nutritional deficiencies (E40-E64)	3	0.5	...
14	Aortic aneurysm and dissection (I71)	3	0.5	...
...	All other causes (Residual)	94	14.9	...

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total	
			deaths	Rate
Hispanic, male, all ages				
...	All causes	68,119	100.0	330.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,867	21.8	72.2
2	Malignant neoplasms (C00-C97)	12,671	18.6	61.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,899	11.6	38.3
4	Cerebrovascular diseases (I60-I69)	3,070	4.5	14.9
5	Diabetes mellitus (E10-E14)	2,976	4.4	14.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,809	4.1	13.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,387	3.5	11.6
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,711	2.5	8.3
9	Chronic lower respiratory diseases (J40-J47)	1,697	2.5	8.2
10	Human immunodeficiency virus (HIV) disease (B20-B24)	1,465	2.2	7.1
11	Certain conditions originating in the perinatal period (P00-P96)	1,459	2.1	7.1
12	Influenza and pneumonia (J10-J18)	1,429	2.1	6.9
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,069	1.6	5.2
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	937	1.4	4.5
15	Septicemia (A40-A41)	721	1.1	3.5
...	All other causes (Residual)	10,952	16.1	53.2
Hispanic, male, 1-4 years				
...	All causes	569	100.0	33.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	200	35.1	11.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	62	10.9	3.7
3	Malignant neoplasms (C00-C97)	60	10.5	3.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	36	6.3	2.1
5	Influenza and pneumonia (J10-J18)	24	4.2	1.4
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	19	3.3	*
7	Chronic lower respiratory diseases (J40-J47)	8	1.4	*
7	Certain conditions originating in the perinatal period (P00-P96)	8	1.4	*
9	Cerebrovascular diseases (I60-I69)	7	1.2	*
10	Septicemia (A40-A41)	6	1.1	*
11	Acute bronchitis and bronchiolitis (J20-J21)	4	0.7	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.7	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.5	*
14	Anemias (D50-D64)	2	0.4	*
14	Nutritional deficiencies (E40-E64)	2	0.4	*
14	Meningitis (G00,G03)	2	0.4	*
14	Pneumonitis due to solids and liquids (J69)	2	0.4	*
14	Hernia (K40-K46)	2	0.4	*
...	All other causes (Residual)	118	20.7	7.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 5-9 years				
...	All causes	314	100.0	16.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	138	43.9	7.0
2	Malignant neoplasms (C00-C97)	70	22.3	3.6
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	15	4.8	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7	2.2	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	1.6	*
5	Influenza and pneumonia (J10-J18)	5	1.6	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	1.3	*
8	Septicemia (A40-A41)	3	1.0	*
8	Cerebrovascular diseases (I60-I69)	3	1.0	*
10	Chronic lower respiratory diseases (J40-J47)	2	0.6	*
10	Certain conditions originating in the perinatal period (P00-P96)	2	0.6	*
12	Anemias (D50-D64)	1	0.3	*
12	Meningitis (G00,G03)	1	0.3	*
12	Pneumonitis due to solids and liquids (J69)	1	0.3	*
12	Diseases of appendix (K35-K38)	1	0.3	*
...	All other causes (Residual)	56	17.8	2.9
Hispanic, male, 10-14 years				
...	All causes	392	100.0	20.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	151	38.5	8.1
2	Malignant neoplasms (C00-C97)	53	13.5	2.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	31	7.9	1.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	29	7.4	1.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	14	3.6	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	13	3.3	*
7	Chronic lower respiratory diseases (J40-J47)	8	2.0	*
8	Influenza and pneumonia (J10-J18)	7	1.8	*
9	Cerebrovascular diseases (I60-I69)	5	1.3	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.8	*
11	Meningitis (G00,G03)	2	0.5	*
11	Certain conditions originating in the perinatal period (P00-P96)	2	0.5	*
13	Septicemia (A40-A41)	1	0.3	*
13	Anemias (D50-D64)	1	0.3	*
13	Diabetes mellitus (E10-E14)	1	0.3	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
...	All other causes (Residual)	70	17.9	3.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 15-19 years				
...	All causes	1,659	100.0	98.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	756	45.6	44.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	423	25.5	25.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	155	9.3	9.2
4	Malignant neoplasms (C00-C97)	91	5.5	5.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	30	1.8	1.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	17	1.0	*
7	Cerebrovascular diseases (I60-I69)	7	0.4	*
7	Chronic lower respiratory diseases (J40-J47)	7	0.4	*
9	Influenza and pneumonia (J10-J18)	6	0.4	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.4	*
9	Legal intervention (Y35,Y89.0)	6	0.4	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.3	*
13	Septicemia (A40-A41)	4	0.2	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.2	*
13	Diabetes mellitus (E10-E14)	4	0.2	*
...	All other causes (Residual)	138	8.3	8.2
Hispanic, male, 20-24 years				
...	All causes	2,585	100.0	124.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,149	44.4	55.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	716	27.7	34.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	266	10.3	12.8
4	Malignant neoplasms (C00-C97)	95	3.7	4.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	71	2.7	3.4
6	Legal intervention (Y35,Y89.0)	22	0.9	1.1
7	Human immunodeficiency virus (HIV) disease (B20-B24)	16	0.6	*
8	Influenza and pneumonia (J10-J18)	14	0.5	*
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	13	0.5	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.3	*
10	Cerebrovascular diseases (I60-I69)	8	0.3	*
12	Diabetes mellitus (E10-E14)	7	0.3	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.3	*
14	Chronic lower respiratory diseases (J40-J47)	5	0.2	*
15	Septicemia (A40-A41)	4	0.2	*
15	Meningitis (G00,G03)	4	0.2	*
15	Pneumonitis due to solids and liquids (J69)	4	0.2	*
...	All other causes (Residual)	176	6.8	8.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 25-29 years				
...	All causes	2,421	100.0	115.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,022	42.2	48.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	527	21.8	25.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	220	9.1	10.5
4	Malignant neoplasms (C00-C97)	116	4.8	5.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	93	3.8	4.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	55	2.3	2.6
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	22	0.9	1.1
8	Cerebrovascular diseases (I60-I69)	18	0.7	*
9	Influenza and pneumonia (J10-J18)	16	0.7	*
10	Diabetes mellitus (E10-E14)	14	0.6	*
10	Legal intervention (Y35,Y89.0)	14	0.6	*
12	Chronic liver disease and cirrhosis (K70,K73-K74)	12	0.5	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	0.4	*
14	Chronic lower respiratory diseases (J40-J47)	9	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.3	*
...	All other causes (Residual)	265	10.9	12.7
Hispanic, male, 30-34 years				
...	All causes	2,321	100.0	120.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	805	34.7	41.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	325	14.0	16.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	214	9.2	11.1
4	Malignant neoplasms (C00-C97)	158	6.8	8.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	151	6.5	7.9
6	Human immunodeficiency virus (HIV) disease (B20-B24)	129	5.6	6.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	67	2.9	3.5
8	Cerebrovascular diseases (I60-I69)	33	1.4	1.7
9	Diabetes mellitus (E10-E14)	19	0.8	*
9	Influenza and pneumonia (J10-J18)	19	0.8	*
11	Legal intervention (Y35,Y89.0)	18	0.8	*
12	Septicemia (A40-A41)	17	0.7	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	15	0.6	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.5	*
15	Aortic aneurysm and dissection (I71)	9	0.4	*
...	All other causes (Residual)	331	14.3	17.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 35-39 years				
...	All causes	2,587	100.0	153.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	763	29.5	45.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	245	9.5	14.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	244	9.4	14.5
4	Human immunodeficiency virus (HIV) disease (B20-B24)	239	9.2	14.2
5	Malignant neoplasms (C00-C97)	214	8.3	12.7
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	173	6.7	10.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	133	5.1	7.9
8	Cerebrovascular diseases (I60-I69)	48	1.9	2.9
9	Diabetes mellitus (E10-E14)	37	1.4	2.2
10	Viral hepatitis (B15-B19)	20	0.8	1.2
11	Septicemia (A40-A41)	18	0.7	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	0.5	*
13	Influenza and pneumonia (J10-J18)	12	0.5	*
13	Chronic lower respiratory diseases (J40-J47)	12	0.5	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	10	0.4	*
...	All other causes (Residual)	405	15.7	24.1
Hispanic, male, 40-44 years				
...	All causes	3,300	100.0	232.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	725	22.0	51.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	410	12.4	28.9
3	Malignant neoplasms (C00-C97)	352	10.7	24.8
4	Human immunodeficiency virus (HIV) disease (B20-B24)	311	9.4	21.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	247	7.5	17.4
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	168	5.1	11.8
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	156	4.7	11.0
8	Cerebrovascular diseases (I60-I69)	114	3.5	8.0
9	Diabetes mellitus (E10-E14)	84	2.5	5.9
10	Viral hepatitis (B15-B19)	60	1.8	4.2
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	35	1.1	2.5
12	Septicemia (A40-A41)	30	0.9	2.1
13	Influenza and pneumonia (J10-J18)	29	0.9	2.0
14	Chronic lower respiratory diseases (J40-J47)	18	0.5	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	15	0.5	*
...	All other causes (Residual)	546	16.5	38.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total	
			deaths	Rate
Hispanic, male, 45-49 years				
...	All causes	4,010	100.0	365.3
1	Malignant neoplasms (C00-C97)	647	16.1	58.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	637	15.9	58.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	585	14.6	53.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	389	9.7	35.4
5	Human immunodeficiency virus (HIV) disease (B20-B24)	306	7.6	27.9
6	Cerebrovascular diseases (I60-I69)	156	3.9	14.2
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	130	3.2	11.8
8	Diabetes mellitus (E10-E14)	127	3.2	11.6
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	123	3.1	11.2
10	Viral hepatitis (B15-B19)	97	2.4	8.8
11	Septicemia (A40-A41)	48	1.2	4.4
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	39	1.0	3.6
13	Influenza and pneumonia (J10-J18)	38	0.9	3.5
14	Chronic lower respiratory diseases (J40-J47)	31	0.8	2.8
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	26	0.6	2.4
...	All other causes (Residual)	631	15.7	57.5
Hispanic, male, 50-54 years				
...	All causes	4,326	100.0	532.3
1	Malignant neoplasms (C00-C97)	903	20.9	111.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	883	20.4	108.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	381	8.8	46.9
4	Chronic liver disease and cirrhosis (K70,K73-K74)	379	8.8	46.6
5	Diabetes mellitus (E10-E14)	218	5.0	26.8
6	Human immunodeficiency virus (HIV) disease (B20-B24)	193	4.5	23.7
7	Cerebrovascular diseases (I60-I69)	189	4.4	23.3
8	Viral hepatitis (B15-B19)	110	2.5	13.5
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	90	2.1	11.1
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	71	1.6	8.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	68	1.6	8.4
12	Influenza and pneumonia (J10-J18)	62	1.4	7.6
13	Septicemia (A40-A41)	46	1.1	5.7
14	Chronic lower respiratory diseases (J40-J47)	30	0.7	3.7
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	27	0.6	3.3
...	All other causes (Residual)	676	15.6	83.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total	
			deaths	Rate
Hispanic, male, 55-59 years				
...	All causes	4,463	100.0	763.9
1	Malignant neoplasms (C00-C97)	1,134	25.4	194.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,091	24.4	186.7
3	Chronic liver disease and cirrhosis (K70,K73-K74)	331	7.4	56.7
4	Diabetes mellitus (E10-E14)	264	5.9	45.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	246	5.5	42.1
6	Cerebrovascular diseases (I60-I69)	203	4.5	34.7
7	Human immunodeficiency virus (HIV) disease (B20-B24)	118	2.6	20.2
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	78	1.7	13.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	75	1.7	12.8
10	Chronic lower respiratory diseases (J40-J47)	69	1.5	11.8
11	Viral hepatitis (B15-B19)	67	1.5	11.5
12	Septicemia (A40-A41)	53	1.2	9.1
12	Influenza and pneumonia (J10-J18)	53	1.2	9.1
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	26	0.6	4.5
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	25	0.6	4.3
...	All other causes (Residual)	630	14.1	107.8
Hispanic, male, 60-64 years				
...	All causes	4,659	100.0	1,146.5
1	Malignant neoplasms (C00-C97)	1,308	28.1	321.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,243	26.7	305.9
3	Diabetes mellitus (E10-E14)	354	7.6	87.1
4	Chronic liver disease and cirrhosis (K70,K73-K74)	285	6.1	70.1
5	Cerebrovascular diseases (I60-I69)	228	4.9	56.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	203	4.4	50.0
7	Chronic lower respiratory diseases (J40-J47)	95	2.0	23.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	85	1.8	20.9
9	Influenza and pneumonia (J10-J18)	62	1.3	15.3
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	51	1.1	12.6
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	47	1.0	11.6
12	Septicemia (A40-A41)	42	0.9	10.3
13	Viral hepatitis (B15-B19)	36	0.8	8.9
14	Human immunodeficiency virus (HIV) disease (B20-B24)	35	0.8	8.6
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	34	0.7	8.4
...	All other causes (Residual)	551	11.8	135.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 65-69 years				
...	All causes	5,235	100.0	1,701.1
1	Malignant neoplasms (C00-C97)	1,568	30.0	509.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,501	28.7	487.8
3	Diabetes mellitus (E10-E14)	359	6.9	116.7
4	Cerebrovascular diseases (I60-I69)	280	5.3	91.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	184	3.5	59.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	153	2.9	49.7
7	Chronic lower respiratory diseases (J40-J47)	126	2.4	40.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	111	2.1	36.1
9	Influenza and pneumonia (J10-J18)	95	1.8	30.9
10	Septicemia (A40-A41)	70	1.3	22.7
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	38	0.7	12.3
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	34	0.6	11.0
13	Human immunodeficiency virus (HIV) disease (B20-B24)	31	0.6	10.1
14	Viral hepatitis (B15-B19)	28	0.5	9.1
15	Aortic aneurysm and dissection (I71)	26	0.5	8.4
...	All other causes (Residual)	631	12.1	205.0
Hispanic, male, 70-74 years				
...	All causes	6,230	100.0	2,660.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,886	30.3	805.4
2	Malignant neoplasms (C00-C97)	1,779	28.6	759.7
3	Diabetes mellitus (E10-E14)	434	7.0	185.3
4	Cerebrovascular diseases (I60-I69)	337	5.4	143.9
5	Chronic lower respiratory diseases (J40-J47)	243	3.9	103.8
6	Chronic liver disease and cirrhosis (K70,K73-K74)	148	2.4	63.2
7	Influenza and pneumonia (J10-J18)	140	2.2	59.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	136	2.2	58.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	119	1.9	50.8
10	Septicemia (A40-A41)	67	1.1	28.6
11	Alzheimer's disease (G30)	57	0.9	24.3
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	50	0.8	21.4
13	Parkinson's disease (G20-G21)	47	0.8	20.1
14	Aortic aneurysm and dissection (I71)	42	0.7	17.9
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	33	0.5	14.1
...	All other causes (Residual)	712	11.4	304.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total	
			deaths	Rate
Hispanic, male, 75-79 years				
...	All causes	6,788	100.0	4,124.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,074	30.6	1,260.2
2	Malignant neoplasms (C00-C97)	1,751	25.8	1,064.0
3	Cerebrovascular diseases (I60-I69)	454	6.7	275.9
4	Diabetes mellitus (E10-E14)	419	6.2	254.6
5	Chronic lower respiratory diseases (J40-J47)	281	4.1	170.7
6	Influenza and pneumonia (J10-J18)	184	2.7	111.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	151	2.2	91.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	146	2.2	88.7
9	Chronic liver disease and cirrhosis (K70,K73-K74)	131	1.9	79.6
10	Alzheimer's disease (G30)	114	1.7	69.3
11	Septicemia (A40-A41)	93	1.4	56.5
12	Parkinson's disease (G20-G21)	64	0.9	38.9
13	Aortic aneurysm and dissection (I71)	61	0.9	37.1
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	55	0.8	33.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	43	0.6	26.1
...	All other causes (Residual)	767	11.3	466.1
Hispanic, male, 80-84 years				
...	All causes	6,116	100.0	6,357.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,951	31.9	2,028.2
2	Malignant neoplasms (C00-C97)	1,284	21.0	1,334.8
3	Cerebrovascular diseases (I60-I69)	425	6.9	441.8
4	Diabetes mellitus (E10-E14)	356	5.8	370.1
5	Chronic lower respiratory diseases (J40-J47)	322	5.3	334.7
6	Influenza and pneumonia (J10-J18)	248	4.1	257.8
7	Alzheimer's disease (G30)	150	2.5	155.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	129	2.1	134.1
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	121	2.0	125.8
10	Parkinson's disease (G20-G21)	88	1.4	91.5
11	Septicemia (A40-A41)	82	1.3	85.2
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	61	1.0	63.4
13	Chronic liver disease and cirrhosis (K70,K73-K74)	55	0.9	57.2
14	Pneumonitis due to solids and liquids (J69)	50	0.8	52.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	45	0.7	46.8
...	All other causes (Residual)	749	12.2	778.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 85-89 years				
...	All causes	3,926	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,331	33.9	...
2	Malignant neoplasms (C00-C97)	656	16.7	...
3	Cerebrovascular diseases (I60-I69)	319	8.1	...
4	Chronic lower respiratory diseases (J40-J47)	235	6.0	...
5	Influenza and pneumonia (J10-J18)	171	4.4	...
6	Diabetes mellitus (E10-E14)	167	4.3	...
7	Alzheimer's disease (G30)	123	3.1	...
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	116	3.0	...
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	77	2.0	...
10	Septicemia (A40-A41)	60	1.5	...
11	Parkinson's disease (G20-G21)	58	1.5	...
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	44	1.1	...
13	Aortic aneurysm and dissection (I71)	37	0.9	...
14	Pneumonitis due to solids and liquids (J69)	35	0.9	...
15	Atherosclerosis (I70)	23	0.6	...
...	All other causes (Residual)	474	12.1	...
Hispanic, male, 90-94 years				
...	All causes	2,276	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	832	36.6	...
2	Malignant neoplasms (C00-C97)	342	15.0	...
3	Cerebrovascular diseases (I60-I69)	155	6.8	...
4	Influenza and pneumonia (J10-J18)	128	5.6	...
4	Chronic lower respiratory diseases (J40-J47)	128	5.6	...
6	Alzheimer's disease (G30)	102	4.5	...
7	Diabetes mellitus (E10-E14)	84	3.7	...
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	49	2.2	...
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	37	1.6	...
10	Septicemia (A40-A41)	32	1.4	...
11	Pneumonitis due to solids and liquids (J69)	27	1.2	...
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	25	1.1	...
13	Atherosclerosis (I70)	23	1.0	...
14	Parkinson's disease (G20-G21)	20	0.9	...
15	Aortic aneurysm and dissection (I71)	11	0.5	...
...	All other causes (Residual)	281	12.3	...

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Hispanic, male, 95-99 years			
...	All causes	804	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	281	35.0	...
2	Malignant neoplasms (C00-C97)	75	9.3	...
3	Cerebrovascular diseases (I60-I69)	65	8.1	...
4	Influenza and pneumonia (J10-J18)	57	7.1	...
5	Alzheimer's disease (G30)	48	6.0	...
6	Chronic lower respiratory diseases (J40-J47)	46	5.7	...
7	Diabetes mellitus (E10-E14)	24	3.0	...
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	2.9	...
9	Pneumonitis due to solids and liquids (J69)	16	2.0	...
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	14	1.7	...
11	Atherosclerosis (I70)	13	1.6	...
12	Septicemia (A40-A41)	12	1.5	...
13	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11	1.4	...
14	Parkinson's disease (G20-G21)	6	0.7	...
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.5	...
15	Aortic aneurysm and dissection (I71)	4	0.5	...
...	All other causes (Residual)	105	13.1	...
	Hispanic, male, 100 years and over			
...	All causes	159	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	73	45.9	...
2	Influenza and pneumonia (J10-J18)	19	11.9	...
3	Malignant neoplasms (C00-C97)	11	6.9	...
3	Chronic lower respiratory diseases (J40-J47)	11	6.9	...
5	Cerebrovascular diseases (I60-I69)	5	3.1	...
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	3.1	...
7	Alzheimer's disease (G30)	4	2.5	...
7	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	4	2.5	...
9	Atherosclerosis (I70)	3	1.9	...
10	Diabetes mellitus (E10-E14)	2	1.3	...
10	Pneumonitis due to solids and liquids (J69)	2	1.3	...
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2	1.3	...
13	Anemias (D50-D64)	1	0.6	...
13	Nutritional deficiencies (E40-E64)	1	0.6	...
13	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.6	...
...	All other causes (Residual)	15	9.4	...

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total	
			deaths	Rate
Hispanic, female, all ages				
...	All causes	53,907	100.0	279.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,431	24.9	69.6
2	Malignant neoplasms (C00-C97)	11,399	21.1	59.1
3	Cerebrovascular diseases (I60-I69)	3,588	6.7	18.6
4	Diabetes mellitus (E10-E14)	3,203	5.9	16.6
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,519	4.7	13.1
6	Influenza and pneumonia (J10-J18)	1,519	2.8	7.9
7	Chronic lower respiratory diseases (J40-J47)	1,477	2.7	7.7
8	Alzheimer's disease (G30)	1,202	2.2	6.2
9	Certain conditions originating in the perinatal period (P00-P96)	1,169	2.2	6.1
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,101	2.0	5.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	995	1.8	5.2
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	850	1.6	4.4
13	Septicemia (A40-A41)	784	1.5	4.1
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	616	1.1	3.2
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	546	1.0	2.8
...	All other causes (Residual)	9,508	17.6	49.3
Hispanic, female, 1-4 years				
...	All causes	424	100.0	26.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	125	29.5	7.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	60	14.2	3.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	46	10.8	2.9
4	Malignant neoplasms (C00-C97)	38	9.0	2.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	16	3.8	*
6	Influenza and pneumonia (J10-J18)	13	3.1	*
7	Certain conditions originating in the perinatal period (P00-P96)	7	1.7	*
8	Septicemia (A40-A41)	4	0.9	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.9	*
10	Anemias (D50-D64)	3	0.7	*
10	Chronic lower respiratory diseases (J40-J47)	3	0.7	*
12	Meningitis (G00,G03)	2	0.5	*
12	Acute bronchitis and bronchiolitis (J20-J21)	2	0.5	*
14	Meningococcal infection (A39)	1	0.2	*
14	Cerebrovascular diseases (I60-I69)	1	0.2	*
14	Pneumonitis due to solids and liquids (J69)	1	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
...	All other causes (Residual)	97	22.9	6.0

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 5-9 years				
...	All causes	224	100.0	12.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	80	35.7	4.3
2	Malignant neoplasms (C00-C97)	41	18.3	2.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	16	7.1	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	9	4.0	*
5	Influenza and pneumonia (J10-J18)	8	3.6	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	1.8	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	1.8	*
8	Diseases of appendix (K35-K38)	3	1.3	*
9	Septicemia (A40-A41)	2	0.9	*
9	Cerebrovascular diseases (I60-I69)	2	0.9	*
9	Chronic lower respiratory diseases (J40-J47)	2	0.9	*
9	Certain conditions originating in the perinatal period (P00-P96)	2	0.9	*
13	Salmonella infections (A01-A02)	1	0.4	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.4	*
13	Anemias (D50-D64)	1	0.4	*
13	Diabetes mellitus (E10-E14)	1	0.4	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.4	*
...	All other causes (Residual)	46	20.5	2.5
Hispanic, female, 10-14 years				
...	All causes	243	100.0	13.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	62	25.5	3.5
2	Malignant neoplasms (C00-C97)	53	21.8	3.0
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	21	8.6	1.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	12	4.9	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	8	3.3	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8	3.3	*
7	Cerebrovascular diseases (I60-I69)	4	1.6	*
8	Chronic lower respiratory diseases (J40-J47)	3	1.2	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.8	*
9	Influenza and pneumonia (J10-J18)	2	0.8	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.8	*
9	Certain conditions originating in the perinatal period (P00-P96)	2	0.8	*
13	Meningococcal infection (A39)	1	0.4	*
13	Septicemia (A40-A41)	1	0.4	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.4	*
13	Anemias (D50-D64)	1	0.4	*
13	Diabetes mellitus (E10-E14)	1	0.4	*
13	Pneumonitis due to solids and liquids (J69)	1	0.4	*
13	Diseases of appendix (K35-K38)	1	0.4	*
13	Legal intervention (Y35,Y89.0)	1	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.4	*
...	All other causes (Residual)	55	22.6	3.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 15-19 years				
...	All causes	524	100.0	33.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	249	47.5	15.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	48	9.2	3.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	37	7.1	2.4
4	Malignant neoplasms (C00-C97)	33	6.3	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	16	3.1	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	10	1.9	*
7	Pregnancy, childbirth and the puerperium (O00-099)	8	1.5	*
8	Influenza and pneumonia (J10-J18)	5	1.0	*
8	Chronic lower respiratory diseases (J40-J47)	5	1.0	*
10	Septicemia (A40-A41)	3	0.6	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	3	0.6	*
12	Diabetes mellitus (E10-E14)	2	0.4	*
12	Cerebrovascular diseases (I60-I69)	2	0.4	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
15	Tuberculosis (A16-A19)	1	0.2	*
15	Meningococcal infection (A39)	1	0.2	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.2	*
15	Anemias (D50-D64)	1	0.2	*
15	Meningitis (G00,G03)	1	0.2	*
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1	0.2	*
15	Aortic aneurysm and dissection (I71)	1	0.2	*
15	Certain conditions originating in the perinatal period (P00-P96)	1	0.2	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
...	All other causes (Residual)	92	17.6	5.9
Hispanic, female, 20-24 years				
...	All causes	626	100.0	37.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	212	33.9	12.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	99	15.8	5.9
3	Malignant neoplasms (C00-C97)	69	11.0	4.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	33	5.3	2.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	18	2.9	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	17	2.7	*
7	Pregnancy, childbirth and the puerperium (O00-099)	16	2.6	*
8	Cerebrovascular diseases (I60-I69)	11	1.8	*
8	Influenza and pneumonia (J10-J18)	11	1.8	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	8	1.3	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	1.1	*
12	Septicemia (A40-A41)	6	1.0	*
12	Diabetes mellitus (E10-E14)	6	1.0	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.6	*
15	Chronic lower respiratory diseases (J40-J47)	3	0.5	*
...	All other causes (Residual)	106	16.9	6.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total	
			deaths	Rate
Hispanic, female, 25-29 years				
...	All causes	667	100.0	39.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	186	27.9	10.9
2	Malignant neoplasms (C00-C97)	92	13.8	5.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	65	9.7	3.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	41	6.1	2.4
5	Pregnancy, childbirth and the puerperium (O00-099)	31	4.6	1.8
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	31	4.6	1.8
7	Human immunodeficiency virus (HIV) disease (B20-B24)	20	3.0	1.2
8	Cerebrovascular diseases (I60-I69)	17	2.5	*
9	Diabetes mellitus (E10-E14)	14	2.1	*
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	13	1.9	*
11	Septicemia (A40-A41)	9	1.3	*
12	Influenza and pneumonia (J10-J18)	8	1.2	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	1.0	*
14	Chronic lower respiratory diseases (J40-J47)	5	0.7	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.6	*
...	All other causes (Residual)	124	18.6	7.3
Hispanic, female, 30-34 years				
...	All causes	780	100.0	47.1
1	Malignant neoplasms (C00-C97)	156	20.0	9.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	154	19.7	9.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	62	7.9	3.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	54	6.9	3.3
5	Human immunodeficiency virus (HIV) disease (B20-B24)	40	5.1	2.4
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	34	4.4	2.1
7	Cerebrovascular diseases (I60-I69)	21	2.7	1.3
8	Chronic liver disease and cirrhosis (K70,K73-K74)	20	2.6	1.2
9	Diabetes mellitus (E10-E14)	17	2.2	*
10	Pregnancy, childbirth and the puerperium (O00-099)	15	1.9	*
11	Influenza and pneumonia (J10-J18)	11	1.4	*
12	Chronic lower respiratory diseases (J40-J47)	9	1.2	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	1.0	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	1.0	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.9	*
...	All other causes (Residual)	164	21.0	9.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 35-39 years				
...	All causes	1,096	100.0	73.4
1	Malignant neoplasms (C00-C97)	299	27.3	20.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	185	16.9	12.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	83	7.6	5.6
4	Human immunodeficiency virus (HIV) disease (B20-B24)	72	6.6	4.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	51	4.7	3.4
6	Cerebrovascular diseases (I60-I69)	32	2.9	2.1
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	30	2.7	2.0
8	Chronic liver disease and cirrhosis (K70,K73-K74)	23	2.1	1.5
9	Diabetes mellitus (E10-E14)	22	2.0	1.5
10	Pregnancy, childbirth and the puerperium (O00-099)	17	1.6	*
11	Influenza and pneumonia (J10-J18)	15	1.4	*
12	Septicemia (A40-A41)	13	1.2	*
12	Chronic lower respiratory diseases (J40-J47)	13	1.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	1.0	*
15	Viral hepatitis (B15-B19)	9	0.8	*
...	All other causes (Residual)	221	20.2	14.8
Hispanic, female, 40-44 years				
...	All causes	1,593	100.0	120.5
1	Malignant neoplasms (C00-C97)	490	30.8	37.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	204	12.8	15.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	142	8.9	10.7
4	Human immunodeficiency virus (HIV) disease (B20-B24)	105	6.6	7.9
5	Cerebrovascular diseases (I60-I69)	73	4.6	5.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	66	4.1	5.0
7	Diabetes mellitus (E10-E14)	54	3.4	4.1
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	50	3.1	3.8
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	31	1.9	2.3
10	Septicemia (A40-A41)	25	1.6	1.9
11	Viral hepatitis (B15-B19)	18	1.1	*
11	Influenza and pneumonia (J10-J18)	18	1.1	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	1.0	*
14	Chronic lower respiratory diseases (J40-J47)	15	0.9	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	13	0.8	*
...	All other causes (Residual)	273	17.1	20.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 45-49 years				
...	All causes	1,907	100.0	177.9
1	Malignant neoplasms (C00-C97)	669	35.1	62.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	228	12.0	21.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	150	7.9	14.0
4	Cerebrovascular diseases (I60-I69)	95	5.0	8.9
5	Diabetes mellitus (E10-E14)	92	4.8	8.6
6	Chronic liver disease and cirrhosis (K70,K73-K74)	75	3.9	7.0
7	Human immunodeficiency virus (HIV) disease (B20-B24)	62	3.3	5.8
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	35	1.8	3.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	31	1.6	2.9
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	31	1.6	2.9
11	Chronic lower respiratory diseases (J40-J47)	30	1.6	2.8
12	Viral hepatitis (B15-B19)	27	1.4	2.5
13	Septicemia (A40-A41)	26	1.4	2.4
14	Influenza and pneumonia (J10-J18)	25	1.3	2.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	0.7	*
...	All other causes (Residual)	317	16.6	29.6
Hispanic, female, 50-54 years				
...	All causes	2,273	100.0	271.9
1	Malignant neoplasms (C00-C97)	862	37.9	103.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	287	12.6	34.3
3	Diabetes mellitus (E10-E14)	156	6.9	18.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	138	6.1	16.5
5	Cerebrovascular diseases (I60-I69)	129	5.7	15.4
6	Chronic liver disease and cirrhosis (K70,K73-K74)	84	3.7	10.0
7	Human immunodeficiency virus (HIV) disease (B20-B24)	45	2.0	5.4
8	Chronic lower respiratory diseases (J40-J47)	43	1.9	5.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	36	1.6	4.3
10	Viral hepatitis (B15-B19)	32	1.4	3.8
11	Septicemia (A40-A41)	28	1.2	3.3
12	Influenza and pneumonia (J10-J18)	26	1.1	3.1
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	24	1.1	2.9
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	17	0.7	*
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	11	0.5	*
...	All other causes (Residual)	355	15.6	42.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Hispanic, female, 55-59 years			
...	All causes	2,714	100.0	430.3
1	Malignant neoplasms (C00-C97)	1,018	37.5	161.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	450	16.6	71.3
3	Diabetes mellitus (E10-E14)	228	8.4	36.1
4	Cerebrovascular diseases (I60-I69)	150	5.5	23.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	102	3.8	16.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	92	3.4	14.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	71	2.6	11.3
8	Chronic lower respiratory diseases (J40-J47)	45	1.7	7.1
9	Influenza and pneumonia (J10-J18)	43	1.6	6.8
10	Septicemia (A40-A41)	37	1.4	5.9
11	Human immunodeficiency virus (HIV) disease (B20-B24)	32	1.2	5.1
12	Viral hepatitis (B15-B19)	26	1.0	4.1
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	26	1.0	4.1
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	12	0.4	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	0.4	*
15	Pneumonitis due to solids and liquids (J69)	10	0.4	*
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	10	0.4	*
...	All other causes (Residual)	352	13.0	55.8
	Hispanic, female, 60-64 years			
...	All causes	3,077	100.0	659.9
1	Malignant neoplasms (C00-C97)	1,022	33.2	219.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	670	21.8	143.7
3	Diabetes mellitus (E10-E14)	275	8.9	59.0
4	Cerebrovascular diseases (I60-I69)	155	5.0	33.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	98	3.2	21.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	87	2.8	18.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	84	2.7	18.0
8	Influenza and pneumonia (J10-J18)	66	2.1	14.2
9	Septicemia (A40-A41)	58	1.9	12.4
10	Chronic lower respiratory diseases (J40-J47)	57	1.9	12.2
11	Viral hepatitis (B15-B19)	22	0.7	4.7
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	22	0.7	4.7
13	Human immunodeficiency virus (HIV) disease (B20-B24)	19	0.6	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	15	0.5	*
15	Alzheimer's disease (G30)	14	0.5	*
...	All other causes (Residual)	413	13.4	88.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total	
			deaths	Rate
Hispanic, female, 65-69 years				
...	All causes	3,995	100.0	1,065.6
1	Malignant neoplasms (C00-C97)	1,221	30.6	325.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	909	22.8	242.5
3	Diabetes mellitus (E10-E14)	360	9.0	96.0
4	Cerebrovascular diseases (I60-I69)	249	6.2	66.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	138	3.5	36.8
6	Chronic lower respiratory diseases (J40-J47)	126	3.2	33.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	114	2.9	30.4
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	86	2.2	22.9
9	Influenza and pneumonia (J10-J18)	73	1.8	19.5
10	Septicemia (A40-A41)	60	1.5	16.0
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	52	1.3	13.9
12	Viral hepatitis (B15-B19)	31	0.8	8.3
13	Alzheimer's disease (G30)	26	0.7	6.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	24	0.6	6.4
15	Aortic aneurysm and dissection (I71)	13	0.3	*
...	All other causes (Residual)	513	12.8	136.8
Hispanic, female, 70-74 years				
...	All causes	5,136	100.0	1,683.6
1	Malignant neoplasms (C00-C97)	1,361	26.5	446.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,339	26.1	438.9
3	Diabetes mellitus (E10-E14)	458	8.9	150.1
4	Cerebrovascular diseases (I60-I69)	336	6.5	110.1
5	Chronic lower respiratory diseases (J40-J47)	156	3.0	51.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	154	3.0	50.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	140	2.7	45.9
8	Influenza and pneumonia (J10-J18)	120	2.3	39.3
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	91	1.8	29.8
10	Alzheimer's disease (G30)	78	1.5	25.6
11	Septicemia (A40-A41)	73	1.4	23.9
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	68	1.3	22.3
13	Viral hepatitis (B15-B19)	33	0.6	10.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	28	0.5	9.2
15	Parkinson's disease (G20-G21)	24	0.5	7.9
...	All other causes (Residual)	677	13.2	221.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 75-79 years				
...	All causes	6,618	100.0	2,853.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,892	28.6	815.6
2	Malignant neoplasms (C00-C97)	1,497	22.6	645.4
3	Cerebrovascular diseases (I60-I69)	547	8.3	235.8
4	Diabetes mellitus (E10-E14)	526	7.9	226.8
5	Chronic lower respiratory diseases (J40-J47)	234	3.5	100.9
6	Influenza and pneumonia (J10-J18)	187	2.8	80.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	169	2.6	72.9
8	Alzheimer's disease (G30)	154	2.3	66.4
9	Chronic liver disease and cirrhosis (K70,K73-K74)	134	2.0	57.8
10	Septicemia (A40-A41)	106	1.6	45.7
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	88	1.3	37.9
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	81	1.2	34.9
13	Parkinson's disease (G20-G21)	48	0.7	20.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	45	0.7	19.4
15	Aortic aneurysm and dissection (I71)	38	0.6	16.4
...	All other causes (Residual)	872	13.2	375.9
Hispanic, female, 80-84 years				
...	All causes	6,697	100.0	4,531.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,160	32.3	1,461.7
2	Malignant neoplasms (C00-C97)	1,169	17.5	791.1
3	Cerebrovascular diseases (I60-I69)	603	9.0	408.1
4	Diabetes mellitus (E10-E14)	442	6.6	299.1
5	Chronic lower respiratory diseases (J40-J47)	258	3.9	174.6
6	Alzheimer's disease (G30)	246	3.7	166.5
7	Influenza and pneumonia (J10-J18)	241	3.6	163.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	129	1.9	87.3
9	Septicemia (A40-A41)	116	1.7	78.5
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	99	1.5	67.0
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	88	1.3	59.5
12	Chronic liver disease and cirrhosis (K70,K73-K74)	77	1.1	52.1
13	Parkinson's disease (G20-G21)	58	0.9	39.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	52	0.8	35.2
15	Atherosclerosis (I70)	42	0.6	28.4
...	All other causes (Residual)	917	13.7	620.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Hispanic, female, 85-89 years			
...	All causes	6,034	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,191	36.3	...
2	Malignant neoplasms (C00-C97)	794	13.2	...
3	Cerebrovascular diseases (I60-I69)	544	9.0	...
4	Diabetes mellitus (E10-E14)	304	5.0	...
5	Alzheimer's disease (G30)	279	4.6	...
6	Chronic lower respiratory diseases (J40-J47)	256	4.2	...
7	Influenza and pneumonia (J10-J18)	244	4.0	...
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	132	2.2	...
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	109	1.8	...
10	Septicemia (A40-A41)	83	1.4	...
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	80	1.3	...
12	Pneumonitis due to solids and liquids (J69)	54	0.9	...
13	Atherosclerosis (I70)	41	0.7	...
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	35	0.6	...
15	Parkinson's disease (G20-G21)	34	0.6	...
...	All other causes (Residual)	854	14.2	...
	Hispanic, female, 90-94 years			
...	All causes	4,559	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,854	40.7	...
2	Cerebrovascular diseases (I60-I69)	426	9.3	...
3	Malignant neoplasms (C00-C97)	386	8.5	...
4	Alzheimer's disease (G30)	262	5.7	...
5	Influenza and pneumonia (J10-J18)	216	4.7	...
6	Diabetes mellitus (E10-E14)	183	4.0	...
7	Chronic lower respiratory diseases (J40-J47)	149	3.3	...
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	87	1.9	...
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	82	1.8	...
10	Septicemia (A40-A41)	76	1.7	...
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	55	1.2	...
12	Pneumonitis due to solids and liquids (J69)	46	1.0	...
13	Atherosclerosis (I70)	41	0.9	...
14	Parkinson's disease (G20-G21)	24	0.5	...
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	0.4	...
...	All other causes (Residual)	652	14.3	...

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 95-99 years				
...	All causes	1,899	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	786	41.4	...
2	Cerebrovascular diseases (I60-I69)	157	8.3	...
3	Influenza and pneumonia (J10-J18)	131	6.9	...
4	Alzheimer's disease (G30)	107	5.6	...
5	Malignant neoplasms (C00-C97)	106	5.6	...
6	Diabetes mellitus (E10-E14)	59	3.1	...
7	Chronic lower respiratory diseases (J40-J47)	57	3.0	...
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	38	2.0	...
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	33	1.7	...
10	Pneumonitis due to solids and liquids (J69)	30	1.6	...
11	Septicemia (A40-A41)	26	1.4	...
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	26	1.4	...
13	Atherosclerosis (I70)	16	0.8	...
14	Nutritional deficiencies (E40-E64)	10	0.5	...
15	Parkinson's disease (G20-G21)	8	0.4	...
...	All other causes (Residual)	309	16.3	...
Hispanic, female, 100 years and over				
...	All causes	472	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	241	51.1	...
2	Influenza and pneumonia (J10-J18)	30	6.4	...
3	Cerebrovascular diseases (I60-I69)	28	5.9	...
4	Alzheimer's disease (G30)	27	5.7	...
5	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	14	3.0	...
6	Malignant neoplasms (C00-C97)	13	2.8	...
7	Pneumonitis due to solids and liquids (J69)	8	1.7	...
8	Atherosclerosis (I70)	6	1.3	...
8	Chronic lower respiratory diseases (J40-J47)	6	1.3	...
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	1.1	...
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5	1.1	...
12	Septicemia (A40-A41)	4	0.8	...
13	Diabetes mellitus (E10-E14)	3	0.6	...
13	Aortic aneurysm and dissection (I71)	3	0.6	...
15	Nutritional deficiencies (E40-E64)	2	0.4	...
...	All other causes (Residual)	77	16.3	...

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total	
			deaths	Rate
White, non-Hispanic, both sexes, all ages				
...	All causes	1,979,465	100.0	993.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	565,808	28.6	284.0
2	Malignant neoplasms (C00-C97)	457,127	23.1	229.5
3	Cerebrovascular diseases (I60-I69)	127,969	6.5	64.2
4	Chronic lower respiratory diseases (J40-J47)	113,541	5.7	57.0
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	82,815	4.2	41.6
6	Alzheimer's disease (G30)	57,268	2.9	28.7
7	Influenza and pneumonia (J10-J18)	54,617	2.8	27.4
8	Diabetes mellitus (E10-E14)	52,894	2.7	26.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	31,497	1.6	15.8
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	26,434	1.3	13.3
11	Septicemia (A40-A41)	25,664	1.3	12.9
12	Chronic liver disease and cirrhosis (K70,K73-K74)	20,602	1.0	10.3
13	Parkinson's disease (G20-G21)	16,499	0.8	8.3
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	15,878	0.8	8.0
15	Pneumonitis due to solids and liquids (J69)	14,845	0.7	7.5
...	All other causes (Residual)	316,007	16.0	158.6
White, non-Hispanic, both sexes, 1-4 years				
...	All causes	2,539	100.0	27.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	985	38.8	10.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	272	10.7	3.0
3	Malignant neoplasms (C00-C97)	221	8.7	2.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	123	4.8	1.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	83	3.3	0.9
6	Influenza and pneumonia (J10-J18)	80	3.2	0.9
7	Septicemia (A40-A41)	43	1.7	0.5
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	32	1.3	0.3
9	Certain conditions originating in the perinatal period (P00-P96)	31	1.2	0.3
10	Cerebrovascular diseases (I60-I69)	23	0.9	0.2
11	Chronic lower respiratory diseases (J40-J47)	15	0.6	*
12	Meningitis (G00,G03)	14	0.6	*
13	Meningococcal infection (A39)	10	0.4	*
13	Acute bronchitis and bronchiolitis (J20-J21)	10	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	9	0.4	*
...	All other causes (Residual)	579	22.8	6.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 5-9 years				
...	All causes	1,579	100.0	13.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	592	37.5	5.0
2	Malignant neoplasms (C00-C97)	315	19.9	2.7
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	98	6.2	0.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	56	3.5	0.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	45	2.8	0.4
6	Influenza and pneumonia (J10-J18)	37	2.3	0.3
7	Septicemia (A40-A41)	21	1.3	0.2
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	1.2	*
9	Cerebrovascular diseases (I60-I69)	16	1.0	*
10	Chronic lower respiratory diseases (J40-J47)	10	0.6	*
11	Diabetes mellitus (E10-E14)	8	0.5	*
12	Meningitis (G00,G03)	7	0.4	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.3	*
13	Certain conditions originating in the perinatal period (P00-P96)	5	0.3	*
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5	0.3	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.3	*
...	All other causes (Residual)	335	21.2	2.8
White, non-Hispanic, both sexes, 10-14 years				
...	All causes	2,305	100.0	17.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	943	40.9	7.2
2	Malignant neoplasms (C00-C97)	322	14.0	2.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	146	6.3	1.1
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	124	5.4	0.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	90	3.9	0.7
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	64	2.8	0.5
7	Influenza and pneumonia (J10-J18)	49	2.1	0.4
8	Chronic lower respiratory diseases (J40-J47)	29	1.3	0.2
9	Septicemia (A40-A41)	26	1.1	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	24	1.0	0.2
11	Cerebrovascular diseases (I60-I69)	21	0.9	0.2
12	Anemias (D50-D64)	15	0.7	*
13	Diabetes mellitus (E10-E14)	11	0.5	*
14	Pneumonitis due to solids and liquids (J69)	9	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.3	*
...	All other causes (Residual)	426	18.5	3.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	White, non-Hispanic, both sexes, 15-19 years			
...	All causes	8,230	100.0	62.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,774	58.0	36.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,088	13.2	8.3
3	Malignant neoplasms (C00-C97)	442	5.4	3.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	331	4.0	2.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	218	2.6	1.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	143	1.7	1.1
7	Cerebrovascular diseases (I60-I69)	54	0.7	0.4
8	Influenza and pneumonia (J10-J18)	46	0.6	0.4
9	Septicemia (A40-A41)	30	0.4	0.2
10	Chronic lower respiratory diseases (J40-J47)	27	0.3	0.2
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	26	0.3	0.2
12	Diabetes mellitus (E10-E14)	22	0.3	0.2
13	Complications of medical and surgical care (Y40-Y84,Y88)	14	0.2	*
14	Meningococcal infection (A39)	13	0.2	*
14	Anemias (D50-D64)	13	0.2	*
...	All other causes (Residual)	989	12.0	7.6
	White, non-Hispanic, both sexes, 20-24 years			
...	All causes	11,410	100.0	88.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,864	51.4	45.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,744	15.3	13.6
3	Malignant neoplasms (C00-C97)	597	5.2	4.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	576	5.0	4.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	387	3.4	3.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	142	1.2	1.1
7	Influenza and pneumonia (J10-J18)	91	0.8	0.7
8	Cerebrovascular diseases (I60-I69)	70	0.6	0.5
9	Diabetes mellitus (E10-E14)	68	0.6	0.5
10	Septicemia (A40-A41)	56	0.5	0.4
11	Chronic lower respiratory diseases (J40-J47)	55	0.5	0.4
12	Pregnancy, childbirth and the puerperium (O00-099)	34	0.3	0.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	32	0.3	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	28	0.2	0.2
15	Human immunodeficiency virus (HIV) disease (B20-B24)	23	0.2	0.2
...	All other causes (Residual)	1,643	14.4	12.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 25-29 years				
...	All causes	9,937	100.0	86.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,020	40.5	34.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,647	16.6	14.3
3	Malignant neoplasms (C00-C97)	808	8.1	7.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	549	5.5	4.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	459	4.6	4.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	123	1.2	1.1
7	Diabetes mellitus (E10-E14)	107	1.1	0.9
8	Cerebrovascular diseases (I60-I69)	90	0.9	0.8
9	Influenza and pneumonia (J10-J18)	78	0.8	0.7
10	Human immunodeficiency virus (HIV) disease (B20-B24)	63	0.6	0.5
11	Septicemia (A40-A41)	62	0.6	0.5
12	Chronic lower respiratory diseases (J40-J47)	56	0.6	0.5
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	48	0.5	0.4
14	Pregnancy, childbirth and the puerperium (O00-099)	43	0.4	0.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	42	0.4	0.4
...	All other causes (Residual)	1,742	17.5	15.1
White, non-Hispanic, both sexes, 30-34 years				
...	All causes	13,620	100.0	104.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,218	31.0	32.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,164	15.9	16.6
3	Malignant neoplasms (C00-C97)	1,546	11.4	11.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,239	9.1	9.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	494	3.6	3.8
6	Human immunodeficiency virus (HIV) disease (B20-B24)	312	2.3	2.4
7	Diabetes mellitus (E10-E14)	259	1.9	2.0
8	Cerebrovascular diseases (I60-I69)	208	1.5	1.6
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	162	1.2	1.2
10	Chronic liver disease and cirrhosis (K70,K73-K74)	146	1.1	1.1
11	Influenza and pneumonia (J10-J18)	128	0.9	1.0
12	Septicemia (A40-A41)	93	0.7	0.7
13	Chronic lower respiratory diseases (J40-J47)	84	0.6	0.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	80	0.6	0.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	71	0.5	0.5
...	All other causes (Residual)	2,416	17.7	18.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 35-39 years				
...	All causes	21,279	100.0	149.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,259	24.7	37.0
2	Malignant neoplasms (C00-C97)	3,308	15.5	23.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,722	12.8	19.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,416	11.4	17.0
5	Human immunodeficiency virus (HIV) disease (B20-B24)	732	3.4	5.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	573	2.7	4.0
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	555	2.6	3.9
8	Diabetes mellitus (E10-E14)	428	2.0	3.0
9	Cerebrovascular diseases (I60-I69)	375	1.8	2.6
10	Influenza and pneumonia (J10-J18)	245	1.2	1.7
11	Chronic lower respiratory diseases (J40-J47)	186	0.9	1.3
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	183	0.9	1.3
13	Septicemia (A40-A41)	171	0.8	1.2
14	Viral hepatitis (B15-B19)	117	0.5	0.8
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	113	0.5	0.8
...	All other causes (Residual)	3,896	18.3	27.4
White, non-Hispanic, both sexes, 40-44 years				
...	All causes	36,816	100.0	227.3
1	Malignant neoplasms (C00-C97)	7,415	20.1	45.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,847	18.6	42.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,172	16.8	38.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,147	8.5	19.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,495	4.1	9.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	971	2.6	6.0
7	Cerebrovascular diseases (I60-I69)	864	2.3	5.3
8	Diabetes mellitus (E10-E14)	812	2.2	5.0
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	592	1.6	3.7
10	Chronic lower respiratory diseases (J40-J47)	431	1.2	2.7
11	Influenza and pneumonia (J10-J18)	392	1.1	2.4
12	Septicemia (A40-A41)	342	0.9	2.1
13	Viral hepatitis (B15-B19)	308	0.8	1.9
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	247	0.7	1.5
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	227	0.6	1.4
...	All other causes (Residual)	6,554	17.8	40.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total	
			deaths	Rate
White, non-Hispanic, both sexes, 45-49 years				
...	All causes	53,153	100.0	333.6
1	Malignant neoplasms (C00-C97)	13,917	26.2	87.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,546	19.8	66.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,593	12.4	41.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,082	5.8	19.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,444	4.6	15.3
6	Diabetes mellitus (E10-E14)	1,368	2.6	8.6
7	Cerebrovascular diseases (I60-I69)	1,364	2.6	8.6
8	Chronic lower respiratory diseases (J40-J47)	952	1.8	6.0
9	Human immunodeficiency virus (HIV) disease (B20-B24)	828	1.6	5.2
10	Viral hepatitis (B15-B19)	753	1.4	4.7
11	Influenza and pneumonia (J10-J18)	601	1.1	3.8
12	Septicemia (A40-A41)	544	1.0	3.4
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	543	1.0	3.4
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	402	0.8	2.5
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	245	0.5	1.5
...	All other causes (Residual)	8,971	16.9	56.3
White, non-Hispanic, both sexes, 50-54 years				
...	All causes	68,679	100.0	478.6
1	Malignant neoplasms (C00-C97)	22,251	32.4	155.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,436	22.5	107.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,128	7.5	35.7
4	Chronic liver disease and cirrhosis (K70,K73-K74)	2,866	4.2	20.0
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,692	3.9	18.8
6	Diabetes mellitus (E10-E14)	2,044	3.0	14.2
7	Cerebrovascular diseases (I60-I69)	1,930	2.8	13.4
8	Chronic lower respiratory diseases (J40-J47)	1,773	2.6	12.4
9	Influenza and pneumonia (J10-J18)	823	1.2	5.7
10	Viral hepatitis (B15-B19)	813	1.2	5.7
11	Septicemia (A40-A41)	751	1.1	5.2
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	575	0.8	4.0
13	Human immunodeficiency virus (HIV) disease (B20-B24)	532	0.8	3.7
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	404	0.6	2.8
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	320	0.5	2.2
...	All other causes (Residual)	10,341	15.1	72.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 55-59 years				
...	All causes	88,382	100.0	716.6
1	Malignant neoplasms (C00-C97)	32,981	37.3	267.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	21,427	24.2	173.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,978	4.5	32.3
4	Chronic lower respiratory diseases (J40-J47)	3,650	4.1	29.6
5	Diabetes mellitus (E10-E14)	3,064	3.5	24.8
6	Cerebrovascular diseases (I60-I69)	2,693	3.0	21.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,524	2.9	20.5
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,085	2.4	16.9
9	Septicemia (A40-A41)	1,088	1.2	8.8
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	907	1.0	7.4
11	Influenza and pneumonia (J10-J18)	894	1.0	7.2
12	Aortic aneurysm and dissection (I71)	452	0.5	3.7
13	Viral hepatitis (B15-B19)	423	0.5	3.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	397	0.4	3.2
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	387	0.4	3.1
...	All other causes (Residual)	11,432	12.9	92.7
White, non-Hispanic, both sexes, 60-64 years				
...	All causes	109,216	100.0	1,142.1
1	Malignant neoplasms (C00-C97)	42,752	39.1	447.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	26,664	24.4	278.8
3	Chronic lower respiratory diseases (J40-J47)	6,815	6.2	71.3
4	Diabetes mellitus (E10-E14)	3,809	3.5	39.8
5	Cerebrovascular diseases (I60-I69)	3,600	3.3	37.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,065	2.8	32.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,182	2.0	22.8
8	Septicemia (A40-A41)	1,383	1.3	14.5
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,377	1.3	14.4
10	Influenza and pneumonia (J10-J18)	1,316	1.2	13.8
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,290	1.2	13.5
12	Aortic aneurysm and dissection (I71)	726	0.7	7.6
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	598	0.5	6.3
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	531	0.5	5.6
15	Alzheimer's disease (G30)	353	0.3	3.7
...	All other causes (Residual)	12,755	11.7	133.4

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total	
			deaths	Rate
White, non-Hispanic, both sexes, 65-69 years				
...	All causes	136,254	100.0	1,762.4
1	Malignant neoplasms (C00-C97)	51,177	37.6	661.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	33,729	24.8	436.3
3	Chronic lower respiratory diseases (J40-J47)	10,564	7.8	136.6
4	Cerebrovascular diseases (I60-I69)	5,407	4.0	69.9
5	Diabetes mellitus (E10-E14)	4,843	3.6	62.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,874	2.1	37.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,099	1.5	27.1
8	Influenza and pneumonia (J10-J18)	1,929	1.4	25.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,886	1.4	24.4
10	Septicemia (A40-A41)	1,688	1.2	21.8
11	Aortic aneurysm and dissection (I71)	1,137	0.8	14.7
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,043	0.8	13.5
13	Alzheimer's disease (G30)	815	0.6	10.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	770	0.6	10.0
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	734	0.5	9.5
...	All other causes (Residual)	15,559	11.4	201.2
White, non-Hispanic, both sexes, 70-74 years				
...	All causes	193,316	100.0	2,761.7
1	Malignant neoplasms (C00-C97)	65,021	33.6	928.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	49,750	25.7	710.7
3	Chronic lower respiratory diseases (J40-J47)	16,338	8.5	233.4
4	Cerebrovascular diseases (I60-I69)	9,852	5.1	140.7
5	Diabetes mellitus (E10-E14)	6,473	3.3	92.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,592	1.9	51.3
7	Influenza and pneumonia (J10-J18)	3,333	1.7	47.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,005	1.6	42.9
9	Septicemia (A40-A41)	2,627	1.4	37.5
10	Alzheimer's disease (G30)	2,454	1.3	35.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	2,157	1.1	30.8
12	Aortic aneurysm and dissection (I71)	1,707	0.9	24.4
13	Parkinson's disease (G20-G21)	1,521	0.8	21.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,200	0.6	17.1
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,128	0.6	16.1
...	All other causes (Residual)	23,158	12.0	330.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total	
			deaths	Rate
White, non-Hispanic, both sexes, 75-79 years				
...	All causes	273,142	100.0	4,371.2
1	Malignant neoplasms (C00-C97)	75,855	27.8	1,213.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	75,414	27.6	1,206.9
3	Chronic lower respiratory diseases (J40-J47)	22,460	8.2	359.4
4	Cerebrovascular diseases (I60-I69)	17,759	6.5	284.2
5	Diabetes mellitus (E10-E14)	8,535	3.1	136.6
6	Alzheimer's disease (G30)	6,469	2.4	103.5
7	Influenza and pneumonia (J10-J18)	6,406	2.3	102.5
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,065	1.9	81.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,813	1.8	77.0
10	Septicemia (A40-A41)	4,019	1.5	64.3
11	Parkinson's disease (G20-G21)	3,352	1.2	53.6
12	Aortic aneurysm and dissection (I71)	2,417	0.9	38.7
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,053	0.8	32.9
14	Chronic liver disease and cirrhosis (K70,K73-K74)	1,918	0.7	30.7
15	Pneumonitis due to solids and liquids (J69)	1,900	0.7	30.4
...	All other causes (Residual)	34,707	12.7	555.4
White, non-Hispanic, both sexes, 80-84 years				
...	All causes	327,421	100.0	7,056.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	100,621	30.7	2,168.5
2	Malignant neoplasms (C00-C97)	68,157	20.8	1,468.8
3	Cerebrovascular diseases (I60-I69)	26,518	8.1	571.5
4	Chronic lower respiratory diseases (J40-J47)	22,717	6.9	489.6
5	Alzheimer's disease (G30)	12,468	3.8	268.7
6	Influenza and pneumonia (J10-J18)	10,139	3.1	218.5
7	Diabetes mellitus (E10-E14)	9,066	2.8	195.4
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,422	2.0	138.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,366	1.9	137.2
10	Septicemia (A40-A41)	4,784	1.5	103.1
11	Parkinson's disease (G20-G21)	4,668	1.4	100.6
12	Pneumonitis due to solids and liquids (J69)	3,078	0.9	66.3
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,973	0.9	64.1
14	Aortic aneurysm and dissection (I71)	2,466	0.8	53.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,277	0.7	49.1
...	All other causes (Residual)	44,701	13.7	963.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total	
			deaths	Rate
White, non-Hispanic, both sexes, 85-89 years				
...	All causes	307,581	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	104,708	34.0	...
2	Malignant neoplasms (C00-C97)	44,609	14.5	...
3	Cerebrovascular diseases (I60-I69)	28,371	9.2	...
4	Chronic lower respiratory diseases (J40-J47)	16,739	5.4	...
5	Alzheimer's disease (G30)	15,885	5.2	...
6	Influenza and pneumonia (J10-J18)	12,184	4.0	...
7	Diabetes mellitus (E10-E14)	7,035	2.3	...
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,282	2.0	...
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,147	2.0	...
10	Septicemia (A40-A41)	4,302	1.4	...
11	Parkinson's disease (G20-G21)	3,869	1.3	...
12	Pneumonitis due to solids and liquids (J69)	3,496	1.1	...
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	3,266	1.1	...
14	Atherosclerosis (I70)	2,684	0.9	...
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,006	0.7	...
...	All other causes (Residual)	45,998	15.0	...
White, non-Hispanic, both sexes, 90-94 years				
...	All causes	205,431	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	77,242	37.6	...
2	Cerebrovascular diseases (I60-I69)	19,865	9.7	...
3	Malignant neoplasms (C00-C97)	19,783	9.6	...
4	Alzheimer's disease (G30)	12,589	6.1	...
5	Influenza and pneumonia (J10-J18)	9,962	4.8	...
6	Chronic lower respiratory diseases (J40-J47)	8,091	3.9	...
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,991	1.9	...
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,881	1.9	...
9	Diabetes mellitus (E10-E14)	3,654	1.8	...
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,694	1.3	...
11	Pneumonitis due to solids and liquids (J69)	2,616	1.3	...
12	Septicemia (A40-A41)	2,516	1.2	...
13	Atherosclerosis (I70)	2,334	1.1	...
14	Parkinson's disease (G20-G21)	1,775	0.9	...
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,098	0.5	...
...	All other causes (Residual)	33,340	16.2	...

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 95-99 years				
...	All causes	79,709	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	31,761	39.8	...
2	Cerebrovascular diseases (I60-I69)	7,521	9.4	...
3	Alzheimer's disease (G30)	5,184	6.5	...
4	Malignant neoplasms (C00-C97)	4,977	6.2	...
5	Influenza and pneumonia (J10-J18)	4,577	5.7	...
6	Chronic lower respiratory diseases (J40-J47)	2,236	2.8	...
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,505	1.9	...
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,372	1.7	...
9	Atherosclerosis (I70)	1,256	1.6	...
10	Diabetes mellitus (E10-E14)	1,124	1.4	...
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,119	1.4	...
12	Pneumonitis due to solids and liquids (J69)	1,055	1.3	...
13	Septicemia (A40-A41)	890	1.1	...
14	Parkinson's disease (G20-G21)	391	0.5	...
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	303	0.4	...
...	All other causes (Residual)	14,438	18.1	...
White, non-Hispanic, both sexes, 100 years and over				
...	All causes	16,122	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,778	42.0	...
2	Cerebrovascular diseases (I60-I69)	1,321	8.2	...
3	Influenza and pneumonia (J10-J18)	1,163	7.2	...
4	Alzheimer's disease (G30)	843	5.2	...
5	Malignant neoplasms (C00-C97)	612	3.8	...
6	Atherosclerosis (I70)	332	2.1	...
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	300	1.9	...
8	Chronic lower respiratory diseases (J40-J47)	295	1.8	...
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	293	1.8	...
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	247	1.5	...
11	Pneumonitis due to solids and liquids (J69)	207	1.3	...
12	Diabetes mellitus (E10-E14)	155	1.0	...
13	Septicemia (A40-A41)	132	0.8	...
14	Anemias (D50-D64)	63	0.4	...
15	Nutritional deficiencies (E40-E64)	52	0.3	...
...	All other causes (Residual)	3,329	20.6	...

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, all ages				
...	All causes	956,194	100.0	979.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	276,241	28.9	282.9
2	Malignant neoplasms (C00-C97)	236,150	24.7	241.8
3	Chronic lower respiratory diseases (J40-J47)	53,593	5.6	54.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	51,888	5.4	53.1
5	Cerebrovascular diseases (I60-I69)	48,549	5.1	49.7
6	Diabetes mellitus (E10-E14)	25,940	2.7	26.6
7	Influenza and pneumonia (J10-J18)	23,541	2.5	24.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	21,085	2.2	21.6
9	Alzheimer's disease (G30)	16,427	1.7	16.8
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15,323	1.6	15.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	13,254	1.4	13.6
12	Septicemia (A40-A41)	11,181	1.2	11.4
13	Parkinson's disease (G20-G21)	9,303	1.0	9.5
14	Aortic aneurysm and dissection (I71)	7,731	0.8	7.9
15	Pneumonitis due to solids and liquids (J69)	7,454	0.8	7.6
...	All other causes (Residual)	138,534	14.5	141.9
White, non-Hispanic, male, 1-4 years				
...	All causes	1,423	100.0	30.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	610	42.9	12.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	136	9.6	2.9
3	Malignant neoplasms (C00-C97)	123	8.6	2.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	61	4.3	1.3
5	Influenza and pneumonia (J10-J18)	44	3.1	0.9
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	34	2.4	0.7
7	Septicemia (A40-A41)	29	2.0	0.6
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	1.3	*
9	Certain conditions originating in the perinatal period (P00-P96)	17	1.2	*
10	Cerebrovascular diseases (I60-I69)	8	0.6	*
11	Meningitis (G00,G03)	7	0.5	*
11	Chronic lower respiratory diseases (J40-J47)	7	0.5	*
13	Meningococcal infection (A39)	6	0.4	*
14	Acute bronchitis and bronchiolitis (J20-J21)	5	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.4	*
...	All other causes (Residual)	307	21.6	6.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 5-9 years				
...	All causes	893	100.0	14.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	338	37.8	5.6
2	Malignant neoplasms (C00-C97)	177	19.8	2.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	50	5.6	0.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	29	3.2	0.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	27	3.0	0.4
6	Influenza and pneumonia (J10-J18)	18	2.0	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	1.1	*
7	Cerebrovascular diseases (I60-I69)	10	1.1	*
9	Septicemia (A40-A41)	8	0.9	*
10	Diabetes mellitus (E10-E14)	5	0.6	*
10	Meningitis (G00,G03)	5	0.6	*
10	Chronic lower respiratory diseases (J40-J47)	5	0.6	*
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5	0.6	*
14	Certain conditions originating in the perinatal period (P00-P96)	3	0.3	*
15	Anemias (D50-D64)	2	0.2	*
15	Nutritional deficiencies (E40-E64)	2	0.2	*
15	Acute bronchitis and bronchiolitis (J20-J21)	2	0.2	*
15	Pneumonitis due to solids and liquids (J69)	2	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.2	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.2	*
...	All other causes (Residual)	191	21.4	3.1
White, non-Hispanic, male, 10-14 years				
...	All causes	1,426	100.0	21.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	616	43.2	9.2
2	Malignant neoplasms (C00-C97)	171	12.0	2.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	117	8.2	1.7
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	67	4.7	1.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	59	4.1	0.9
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	45	3.2	0.7
7	Influenza and pneumonia (J10-J18)	21	1.5	0.3
8	Chronic lower respiratory diseases (J40-J47)	16	1.1	*
9	Septicemia (A40-A41)	14	1.0	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	1.0	*
11	Cerebrovascular diseases (I60-I69)	13	0.9	*
12	Anemias (D50-D64)	7	0.5	*
12	Diabetes mellitus (E10-E14)	7	0.5	*
14	Pneumonitis due to solids and liquids (J69)	4	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.3	*
...	All other causes (Residual)	251	17.6	3.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 15-19 years				
...	All causes	5,661	100.0	84.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,251	57.4	48.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	896	15.8	13.3
3	Malignant neoplasms (C00-C97)	269	4.8	4.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	242	4.3	3.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	137	2.4	2.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	92	1.6	1.4
7	Cerebrovascular diseases (I60-I69)	29	0.5	0.4
7	Influenza and pneumonia (J10-J18)	29	0.5	0.4
9	Septicemia (A40-A41)	21	0.4	0.3
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	15	0.3	*
11	Chronic lower respiratory diseases (J40-J47)	14	0.2	*
12	Diabetes mellitus (E10-E14)	13	0.2	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	12	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.1	*
15	Aortic aneurysm and dissection (I71)	7	0.1	*
...	All other causes (Residual)	626	11.1	9.3
White, non-Hispanic, male, 20-24 years				
...	All causes	8,361	100.0	128.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,447	53.2	68.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,516	18.1	23.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	423	5.1	6.5
4	Malignant neoplasms (C00-C97)	338	4.0	5.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	231	2.8	3.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	83	1.0	1.3
7	Influenza and pneumonia (J10-J18)	46	0.6	0.7
8	Cerebrovascular diseases (I60-I69)	43	0.5	0.7
9	Diabetes mellitus (E10-E14)	41	0.5	0.6
10	Chronic lower respiratory diseases (J40-J47)	34	0.4	0.5
11	Septicemia (A40-A41)	27	0.3	0.4
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	0.2	*
12	Legal intervention (Y35,Y89.0)	19	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	0.2	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	15	0.2	*
...	All other causes (Residual)	1,063	12.7	16.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 25-29 years				
...	All causes	6,942	100.0	119.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,026	43.6	52.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,363	19.6	23.4
3	Malignant neoplasms (C00-C97)	426	6.1	7.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	353	5.1	6.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	312	4.5	5.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	73	1.1	1.3
7	Influenza and pneumonia (J10-J18)	52	0.7	0.9
8	Human immunodeficiency virus (HIV) disease (B20-B24)	50	0.7	0.9
9	Diabetes mellitus (E10-E14)	49	0.7	0.8
10	Cerebrovascular diseases (I60-I69)	48	0.7	0.8
11	Septicemia (A40-A41)	28	0.4	0.5
12	Chronic liver disease and cirrhosis (K70,K73-K74)	26	0.4	0.4
13	Chronic lower respiratory diseases (J40-J47)	25	0.4	0.4
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	25	0.4	0.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	22	0.3	0.4
...	All other causes (Residual)	1,064	15.3	18.3
White, non-Hispanic, male, 30-34 years				
...	All causes	9,155	100.0	139.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,112	34.0	47.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,736	19.0	26.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	859	9.4	13.1
4	Malignant neoplasms (C00-C97)	733	8.0	11.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	336	3.7	5.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	244	2.7	3.7
7	Diabetes mellitus (E10-E14)	163	1.8	2.5
8	Cerebrovascular diseases (I60-I69)	111	1.2	1.7
9	Chronic liver disease and cirrhosis (K70,K73-K74)	100	1.1	1.5
10	Influenza and pneumonia (J10-J18)	79	0.9	1.2
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	79	0.9	1.2
12	Septicemia (A40-A41)	46	0.5	0.7
13	Chronic lower respiratory diseases (J40-J47)	39	0.4	0.6
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	39	0.4	0.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	35	0.4	0.5
...	All other causes (Residual)	1,444	15.8	22.0

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 35-39 years				
...	All causes	13,718	100.0	192.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,703	27.0	51.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,943	14.2	27.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,898	13.8	26.6
4	Malignant neoplasms (C00-C97)	1,467	10.7	20.6
5	Human immunodeficiency virus (HIV) disease (B20-B24)	603	4.4	8.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	355	2.6	5.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	354	2.6	5.0
8	Diabetes mellitus (E10-E14)	258	1.9	3.6
9	Cerebrovascular diseases (I60-I69)	195	1.4	2.7
10	Influenza and pneumonia (J10-J18)	141	1.0	2.0
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	98	0.7	1.4
12	Septicemia (A40-A41)	94	0.7	1.3
13	Chronic lower respiratory diseases (J40-J47)	88	0.6	1.2
14	Viral hepatitis (B15-B19)	84	0.6	1.2
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	60	0.4	0.8
...	All other causes (Residual)	2,377	17.3	33.3
White, non-Hispanic, male, 40-44 years				
...	All causes	23,405	100.0	289.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,871	20.8	60.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,486	19.2	55.5
3	Malignant neoplasms (C00-C97)	3,395	14.5	42.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,398	10.2	29.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,012	4.3	12.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	826	3.5	10.2
7	Diabetes mellitus (E10-E14)	536	2.3	6.6
8	Cerebrovascular diseases (I60-I69)	442	1.9	5.5
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	392	1.7	4.8
10	Viral hepatitis (B15-B19)	216	0.9	2.7
11	Influenza and pneumonia (J10-J18)	215	0.9	2.7
12	Chronic lower respiratory diseases (J40-J47)	187	0.8	2.3
13	Septicemia (A40-A41)	161	0.7	2.0
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	145	0.6	1.8
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	135	0.6	1.7
...	All other causes (Residual)	3,988	17.0	49.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 45-49 years				
...	All causes	33,819	100.0	426.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,883	23.3	99.5
2	Malignant neoplasms (C00-C97)	6,970	20.6	87.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,643	13.7	58.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,309	6.8	29.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,756	5.2	22.2
6	Diabetes mellitus (E10-E14)	868	2.6	11.0
7	Cerebrovascular diseases (I60-I69)	747	2.2	9.4
8	Human immunodeficiency virus (HIV) disease (B20-B24)	702	2.1	8.9
9	Viral hepatitis (B15-B19)	549	1.6	6.9
10	Chronic lower respiratory diseases (J40-J47)	518	1.5	6.5
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	362	1.1	4.6
12	Influenza and pneumonia (J10-J18)	343	1.0	4.3
13	Septicemia (A40-A41)	298	0.9	3.8
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	241	0.7	3.0
15	Aortic aneurysm and dissection (I71)	176	0.5	2.2
...	All other causes (Residual)	5,454	16.1	68.8
White, non-Hispanic, male, 50-54 years				
...	All causes	43,287	100.0	609.7
1	Malignant neoplasms (C00-C97)	11,631	26.9	163.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,621	26.8	163.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,576	8.3	50.4
4	Chronic liver disease and cirrhosis (K70,K73-K74)	2,138	4.9	30.1
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,048	4.7	28.8
6	Diabetes mellitus (E10-E14)	1,254	2.9	17.7
7	Cerebrovascular diseases (I60-I69)	1,066	2.5	15.0
8	Chronic lower respiratory diseases (J40-J47)	908	2.1	12.8
9	Viral hepatitis (B15-B19)	613	1.4	8.6
10	Influenza and pneumonia (J10-J18)	501	1.2	7.1
11	Human immunodeficiency virus (HIV) disease (B20-B24)	470	1.1	6.6
12	Septicemia (A40-A41)	400	0.9	5.6
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	336	0.8	4.7
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	283	0.7	4.0
15	Aortic aneurysm and dissection (I71)	252	0.6	3.5
...	All other causes (Residual)	6,190	14.3	87.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total	
			deaths	Rate
White, non-Hispanic, male, 55-59 years				
...	All causes	53,763	100.0	888.4
1	Malignant neoplasms (C00-C97)	17,674	32.9	292.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,485	28.8	255.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,752	5.1	45.5
4	Chronic lower respiratory diseases (J40-J47)	1,867	3.5	30.9
5	Diabetes mellitus (E10-E14)	1,822	3.4	30.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1,788	3.3	29.5
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,588	3.0	26.2
8	Cerebrovascular diseases (I60-I69)	1,433	2.7	23.7
9	Septicemia (A40-A41)	562	1.0	9.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	528	1.0	8.7
11	Influenza and pneumonia (J10-J18)	503	0.9	8.3
12	Aortic aneurysm and dissection (I71)	355	0.7	5.9
13	Viral hepatitis (B15-B19)	310	0.6	5.1
14	Human immunodeficiency virus (HIV) disease (B20-B24)	280	0.5	4.6
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	245	0.5	4.0
...	All other causes (Residual)	6,571	12.2	108.6
White, non-Hispanic, male, 60-64 years				
...	All causes	64,620	100.0	1,402.1
1	Malignant neoplasms (C00-C97)	23,352	36.1	506.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	18,386	28.5	398.9
3	Chronic lower respiratory diseases (J40-J47)	3,460	5.4	75.1
4	Diabetes mellitus (E10-E14)	2,219	3.4	48.1
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,089	3.2	45.3
6	Cerebrovascular diseases (I60-I69)	1,994	3.1	43.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,497	2.3	32.5
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,090	1.7	23.7
9	Septicemia (A40-A41)	718	1.1	15.6
10	Influenza and pneumonia (J10-J18)	716	1.1	15.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	688	1.1	14.9
12	Aortic aneurysm and dissection (I71)	542	0.8	11.8
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	331	0.5	7.2
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	282	0.4	6.1
15	Pneumonitis due to solids and liquids (J69)	200	0.3	4.3
...	All other causes (Residual)	7,056	10.9	153.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total	
			deaths	Rate
White, non-Hispanic, male, 65-69 years				
...	All causes	78,944	100.0	2,169.8
1	Malignant neoplasms (C00-C97)	28,381	36.0	780.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	21,997	27.9	604.6
3	Chronic lower respiratory diseases (J40-J47)	5,330	6.8	146.5
4	Cerebrovascular diseases (I60-I69)	2,936	3.7	80.7
5	Diabetes mellitus (E10-E14)	2,741	3.5	75.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,833	2.3	50.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,364	1.7	37.5
8	Influenza and pneumonia (J10-J18)	1,109	1.4	30.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,027	1.3	28.2
10	Septicemia (A40-A41)	895	1.1	24.6
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	860	1.1	23.6
12	Aortic aneurysm and dissection (I71)	801	1.0	22.0
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	438	0.6	12.0
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	421	0.5	11.6
15	Alzheimer's disease (G30)	385	0.5	10.6
...	All other causes (Residual)	8,426	10.7	231.6
White, non-Hispanic, male, 70-74 years				
...	All causes	107,168	100.0	3,393.7
1	Malignant neoplasms (C00-C97)	35,638	33.3	1,128.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	30,151	28.1	954.8
3	Chronic lower respiratory diseases (J40-J47)	8,274	7.7	262.0
4	Cerebrovascular diseases (I60-I69)	5,011	4.7	158.7
5	Diabetes mellitus (E10-E14)	3,465	3.2	109.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,119	2.0	67.1
7	Influenza and pneumonia (J10-J18)	1,857	1.7	58.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,587	1.5	50.3
9	Septicemia (A40-A41)	1,312	1.2	41.5
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,263	1.2	40.0
11	Aortic aneurysm and dissection (I71)	1,125	1.0	35.6
12	Alzheimer's disease (G30)	1,058	1.0	33.5
13	Parkinson's disease (G20-G21)	996	0.9	31.5
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	909	0.8	28.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	704	0.7	22.3
...	All other causes (Residual)	11,699	10.9	370.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Percent of total deaths		
		Number	Rate	Rate
White, non-Hispanic, male, 75-79 years				
...	All causes	141,742	100.0	5,403.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	41,666	29.4	1,588.2
2	Malignant neoplasms (C00-C97)	40,432	28.5	1,541.2
3	Chronic lower respiratory diseases (J40-J47)	11,014	7.8	419.8
4	Cerebrovascular diseases (I60-I69)	8,161	5.8	311.1
5	Diabetes mellitus (E10-E14)	4,303	3.0	164.0
6	Influenza and pneumonia (J10-J18)	3,348	2.4	127.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,784	2.0	106.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,540	1.8	96.8
9	Alzheimer's disease (G30)	2,523	1.8	96.2
10	Parkinson's disease (G20-G21)	2,110	1.5	80.4
11	Septicemia (A40-A41)	1,886	1.3	71.9
12	Aortic aneurysm and dissection (I71)	1,497	1.1	57.1
13	Pneumonitis due to solids and liquids (J69)	1,092	0.8	41.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	994	0.7	37.9
15	Chronic liver disease and cirrhosis (K70,K73-K74)	986	0.7	37.6
...	All other causes (Residual)	16,406	11.6	625.4
White, non-Hispanic, male, 80-84 years				
...	All causes	152,032	100.0	8,607.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	48,319	31.8	2,735.5
2	Malignant neoplasms (C00-C97)	34,536	22.7	1,955.2
3	Chronic lower respiratory diseases (J40-J47)	10,539	6.9	596.6
4	Cerebrovascular diseases (I60-I69)	10,538	6.9	596.6
5	Influenza and pneumonia (J10-J18)	4,793	3.2	271.3
6	Alzheimer's disease (G30)	4,273	2.8	241.9
7	Diabetes mellitus (E10-E14)	4,114	2.7	232.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,210	2.1	181.7
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,173	2.1	179.6
10	Parkinson's disease (G20-G21)	2,674	1.8	151.4
11	Septicemia (A40-A41)	2,086	1.4	118.1
12	Pneumonitis due to solids and liquids (J69)	1,649	1.1	93.4
13	Aortic aneurysm and dissection (I71)	1,380	0.9	78.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,131	0.7	64.0
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,059	0.7	60.0
...	All other causes (Residual)	18,558	12.2	1,050.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 85-89 years				
...	All causes	120,507	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	41,591	34.5	...
2	Malignant neoplasms (C00-C97)	20,704	17.2	...
3	Cerebrovascular diseases (I60-I69)	9,261	7.7	...
4	Chronic lower respiratory diseases (J40-J47)	7,311	6.1	...
5	Influenza and pneumonia (J10-J18)	4,993	4.1	...
6	Alzheimer's disease (G30)	4,510	3.7	...
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,812	2.3	...
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,735	2.3	...
9	Diabetes mellitus (E10-E14)	2,720	2.3	...
10	Parkinson's disease (G20-G21)	2,038	1.7	...
11	Pneumonitis due to solids and liquids (J69)	1,683	1.4	...
12	Septicemia (A40-A41)	1,599	1.3	...
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	954	0.8	...
14	Aortic aneurysm and dissection (I71)	940	0.8	...
15	Atherosclerosis (I70)	917	0.8	...
...	All other causes (Residual)	15,739	13.1	...
White, non-Hispanic, male, 90-94 years				
...	All causes	62,124	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	23,226	37.4	...
2	Malignant neoplasms (C00-C97)	7,872	12.7	...
3	Cerebrovascular diseases (I60-I69)	4,988	8.0	...
4	Influenza and pneumonia (J10-J18)	3,257	5.2	...
5	Chronic lower respiratory diseases (J40-J47)	3,170	5.1	...
6	Alzheimer's disease (G30)	2,651	4.3	...
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,537	2.5	...
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,445	2.3	...
9	Pneumonitis due to solids and liquids (J69)	1,097	1.8	...
10	Diabetes mellitus (E10-E14)	1,071	1.7	...
11	Parkinson's disease (G20-G21)	785	1.3	...
12	Septicemia (A40-A41)	735	1.2	...
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	639	1.0	...
14	Atherosclerosis (I70)	602	1.0	...
15	Aortic aneurysm and dissection (I71)	376	0.6	...
...	All other causes (Residual)	8,673	14.0	...

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	White, non-Hispanic, male, 95-99 years			
...	All causes	17,126	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,715	39.2	...
2	Malignant neoplasms (C00-C97)	1,660	9.7	...
3	Cerebrovascular diseases (I60-I69)	1,339	7.8	...
4	Influenza and pneumonia (J10-J18)	1,168	6.8	...
5	Alzheimer's disease (G30)	708	4.1	...
6	Chronic lower respiratory diseases (J40-J47)	695	4.1	...
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	405	2.4	...
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	399	2.3	...
9	Pneumonitis due to solids and liquids (J69)	354	2.1	...
10	Diabetes mellitus (E10-E14)	252	1.5	...
11	Atherosclerosis (I70)	244	1.4	...
12	Septicemia (A40-A41)	195	1.1	...
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	189	1.1	...
14	Parkinson's disease (G20-G21)	127	0.7	...
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	82	0.5	...
...	All other causes (Residual)	2,594	15.1	...
	White, non-Hispanic, male, 100 years and over			
...	All causes	2,391	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	950	39.7	...
2	Influenza and pneumonia (J10-J18)	221	9.2	...
3	Malignant neoplasms (C00-C97)	168	7.0	...
4	Cerebrovascular diseases (I60-I69)	153	6.4	...
5	Alzheimer's disease (G30)	77	3.2	...
6	Chronic lower respiratory diseases (J40-J47)	75	3.1	...
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	62	2.6	...
8	Pneumonitis due to solids and liquids (J69)	60	2.5	...
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	58	2.4	...
10	Atherosclerosis (I70)	45	1.9	...
11	Diabetes mellitus (E10-E14)	33	1.4	...
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	29	1.2	...
13	Septicemia (A40-A41)	21	0.9	...
14	Parkinson's disease (G20-G21)	12	0.5	...
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.3	...
...	All other causes (Residual)	419	17.5	...

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total	
			deaths	Rate
White, non-Hispanic, female, all ages				
...	All causes	1,023,271	100.0	1,007.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	289,567	28.3	285.1
2	Malignant neoplasms (C00-C97)	220,977	21.6	217.6
3	Cerebrovascular diseases (I60-I69)	79,420	7.8	78.2
4	Chronic lower respiratory diseases (J40-J47)	59,948	5.9	59.0
5	Alzheimer's disease (G30)	40,841	4.0	40.2
6	Influenza and pneumonia (J10-J18)	31,076	3.0	30.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	30,927	3.0	30.5
8	Diabetes mellitus (E10-E14)	26,954	2.6	26.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16,174	1.6	15.9
10	Septicemia (A40-A41)	14,483	1.4	14.3
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	10,205	1.0	10.0
12	Pneumonitis due to solids and liquids (J69)	7,391	0.7	7.3
13	Chronic liver disease and cirrhosis (K70,K73-K74)	7,348	0.7	7.2
14	Atherosclerosis (I70)	7,294	0.7	7.2
15	Parkinson's disease (G20-G21)	7,196	0.7	7.1
...	All other causes (Residual)	173,470	17.0	170.8
White, non-Hispanic, female, 1-4 years				
...	All causes	1,116	100.0	24.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	375	33.6	8.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	136	12.2	3.0
3	Malignant neoplasms (C00-C97)	98	8.8	2.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	62	5.6	1.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	49	4.4	1.1
6	Influenza and pneumonia (J10-J18)	36	3.2	0.8
7	Cerebrovascular diseases (I60-I69)	15	1.3	*
8	Septicemia (A40-A41)	14	1.3	*
8	Certain conditions originating in the perinatal period (P00-P96)	14	1.3	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	1.2	*
11	Chronic lower respiratory diseases (J40-J47)	8	0.7	*
12	Meningitis (G00,G03)	7	0.6	*
13	Acute bronchitis and bronchiolitis (J20-J21)	5	0.4	*
14	Meningococcal infection (A39)	4	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.4	*
...	All other causes (Residual)	272	24.4	6.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 5-9 years				
...	All causes	686	100.0	11.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	254	37.0	4.4
2	Malignant neoplasms (C00-C97)	138	20.1	2.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	48	7.0	0.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	27	3.9	0.5
5	Influenza and pneumonia (J10-J18)	19	2.8	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	18	2.6	*
7	Septicemia (A40-A41)	13	1.9	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	1.3	*
9	Cerebrovascular diseases (I60-I69)	6	0.9	*
10	Chronic lower respiratory diseases (J40-J47)	5	0.7	*
11	Diabetes mellitus (E10-E14)	3	0.4	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.4	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.4	*
14	Meningitis (G00,G03)	2	0.3	*
14	Certain conditions originating in the perinatal period (P00-P96)	2	0.3	*
...	All other causes (Residual)	136	19.8	2.4
White, non-Hispanic, female, 10-14 years				
...	All causes	879	100.0	13.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	327	37.2	5.1
2	Malignant neoplasms (C00-C97)	151	17.2	2.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	57	6.5	0.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	31	3.5	0.5
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	29	3.3	0.5
6	Influenza and pneumonia (J10-J18)	28	3.2	0.4
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	19	2.2	*
8	Chronic lower respiratory diseases (J40-J47)	13	1.5	*
9	Septicemia (A40-A41)	12	1.4	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	1.1	*
11	Anemias (D50-D64)	8	0.9	*
11	Cerebrovascular diseases (I60-I69)	8	0.9	*
13	Pneumonitis due to solids and liquids (J69)	5	0.6	*
14	Diabetes mellitus (E10-E14)	4	0.5	*
15	Meningococcal infection (A39)	3	0.3	*
...	All other causes (Residual)	174	19.8	2.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	White, non-Hispanic, female, 15-19 years			
...	All causes	2,569	100.0	40.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,523	59.3	24.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	192	7.5	3.0
3	Malignant neoplasms (C00-C97)	173	6.7	2.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	89	3.5	1.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	81	3.2	1.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	51	2.0	0.8
7	Cerebrovascular diseases (I60-I69)	25	1.0	0.4
8	Influenza and pneumonia (J10-J18)	17	0.7	*
9	Chronic lower respiratory diseases (J40-J47)	13	0.5	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.4	*
11	Septicemia (A40-A41)	9	0.4	*
11	Diabetes mellitus (E10-E14)	9	0.4	*
13	Anemias (D50-D64)	8	0.3	*
14	Meningococcal infection (A39)	7	0.3	*
14	Pregnancy, childbirth and the puerperium (O00-099)	7	0.3	*
...	All other causes (Residual)	354	13.8	5.6
	White, non-Hispanic, female, 20-24 years			
...	All causes	3,049	100.0	48.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,417	46.5	22.4
2	Malignant neoplasms (C00-C97)	259	8.5	4.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	228	7.5	3.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	156	5.1	2.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	153	5.0	2.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	59	1.9	0.9
7	Influenza and pneumonia (J10-J18)	45	1.5	0.7
8	Pregnancy, childbirth and the puerperium (O00-099)	34	1.1	0.5
9	Septicemia (A40-A41)	29	1.0	0.5
10	Diabetes mellitus (E10-E14)	27	0.9	0.4
10	Cerebrovascular diseases (I60-I69)	27	0.9	0.4
12	Chronic lower respiratory diseases (J40-J47)	21	0.7	0.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	11	0.4	*
...	All other causes (Residual)	558	18.3	8.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	White, non-Hispanic, female, 25-29 years			
...	All causes	2,995	100.0	52.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	994	33.2	17.4
2	Malignant neoplasms (C00-C97)	382	12.8	6.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	284	9.5	5.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	196	6.5	3.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	147	4.9	2.6
6	Diabetes mellitus (E10-E14)	58	1.9	1.0
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	50	1.7	0.9
8	Pregnancy, childbirth and the puerperium (O00-099)	43	1.4	0.8
9	Cerebrovascular diseases (I60-I69)	42	1.4	0.7
10	Septicemia (A40-A41)	34	1.1	0.6
11	Chronic lower respiratory diseases (J40-J47)	31	1.0	0.5
12	Influenza and pneumonia (J10-J18)	26	0.9	0.5
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	0.8	0.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	0.7	0.4
15	Chronic liver disease and cirrhosis (K70,K73-K74)	14	0.5	*
...	All other causes (Residual)	651	21.7	11.4
	White, non-Hispanic, female, 30-34 years			
...	All causes	4,465	100.0	68.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,106	24.8	17.1
2	Malignant neoplasms (C00-C97)	813	18.2	12.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	428	9.6	6.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	380	8.5	5.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	158	3.5	2.4
6	Cerebrovascular diseases (I60-I69)	97	2.2	1.5
7	Diabetes mellitus (E10-E14)	96	2.2	1.5
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	83	1.9	1.3
9	Human immunodeficiency virus (HIV) disease (B20-B24)	68	1.5	1.0
10	Influenza and pneumonia (J10-J18)	49	1.1	0.8
11	Septicemia (A40-A41)	47	1.1	0.7
12	Chronic liver disease and cirrhosis (K70,K73-K74)	46	1.0	0.7
13	Chronic lower respiratory diseases (J40-J47)	45	1.0	0.7
14	Pregnancy, childbirth and the puerperium (O00-099)	44	1.0	0.7
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	41	0.9	0.6
...	All other causes (Residual)	964	21.6	14.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 35-39 years				
...	All causes	7,561	100.0	106.6
1	Malignant neoplasms (C00-C97)	1,841	24.3	26.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,556	20.6	21.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	779	10.3	11.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	518	6.9	7.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	219	2.9	3.1
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	200	2.6	2.8
7	Cerebrovascular diseases (I60-I69)	180	2.4	2.5
8	Diabetes mellitus (E10-E14)	170	2.2	2.4
9	Human immunodeficiency virus (HIV) disease (B20-B24)	129	1.7	1.8
10	Influenza and pneumonia (J10-J18)	104	1.4	1.5
11	Chronic lower respiratory diseases (J40-J47)	98	1.3	1.4
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	85	1.1	1.2
13	Septicemia (A40-A41)	77	1.0	1.1
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	53	0.7	0.7
15	Pregnancy, childbirth and the puerperium (O00-099)	39	0.5	0.5
...	All other causes (Residual)	1,513	20.0	21.3
White, non-Hispanic, female, 40-44 years				
...	All causes	13,411	100.0	165.4
1	Malignant neoplasms (C00-C97)	4,020	30.0	49.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,976	14.7	24.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,686	12.6	20.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	749	5.6	9.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	483	3.6	6.0
6	Cerebrovascular diseases (I60-I69)	422	3.1	5.2
7	Diabetes mellitus (E10-E14)	276	2.1	3.4
8	Chronic lower respiratory diseases (J40-J47)	244	1.8	3.0
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	200	1.5	2.5
10	Septicemia (A40-A41)	181	1.3	2.2
11	Influenza and pneumonia (J10-J18)	177	1.3	2.2
12	Human immunodeficiency virus (HIV) disease (B20-B24)	145	1.1	1.8
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	112	0.8	1.4
14	Viral hepatitis (B15-B19)	92	0.7	1.1
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	82	0.6	1.0
...	All other causes (Residual)	2,566	19.1	31.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total	
			deaths	Rate
White, non-Hispanic, female, 45-49 years				
...	All causes	19,334	100.0	241.4
1	Malignant neoplasms (C00-C97)	6,947	35.9	86.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,663	13.8	33.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,950	10.1	24.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	773	4.0	9.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	688	3.6	8.6
6	Cerebrovascular diseases (I60-I69)	617	3.2	7.7
7	Diabetes mellitus (E10-E14)	500	2.6	6.2
8	Chronic lower respiratory diseases (J40-J47)	434	2.2	5.4
9	Influenza and pneumonia (J10-J18)	258	1.3	3.2
10	Septicemia (A40-A41)	246	1.3	3.1
11	Viral hepatitis (B15-B19)	204	1.1	2.5
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	181	0.9	2.3
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	161	0.8	2.0
14	Human immunodeficiency virus (HIV) disease (B20-B24)	126	0.7	1.6
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	118	0.6	1.5
...	All other causes (Residual)	3,468	17.9	43.3
White, non-Hispanic, female, 50-54 years				
...	All causes	25,392	100.0	350.2
1	Malignant neoplasms (C00-C97)	10,620	41.8	146.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,815	15.0	52.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,552	6.1	21.4
4	Chronic lower respiratory diseases (J40-J47)	865	3.4	11.9
5	Cerebrovascular diseases (I60-I69)	864	3.4	11.9
6	Diabetes mellitus (E10-E14)	790	3.1	10.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	728	2.9	10.0
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	644	2.5	8.9
9	Septicemia (A40-A41)	351	1.4	4.8
10	Influenza and pneumonia (J10-J18)	322	1.3	4.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	239	0.9	3.3
12	Viral hepatitis (B15-B19)	200	0.8	2.8
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	141	0.6	1.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	130	0.5	1.8
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	121	0.5	1.7
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	121	0.5	1.7
...	All other causes (Residual)	3,889	15.3	53.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	White, non-Hispanic, female, 55-59 years			
...	All causes	34,619	100.0	551.1
1	Malignant neoplasms (C00-C97)	15,307	44.2	243.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,942	17.2	94.6
3	Chronic lower respiratory diseases (J40-J47)	1,783	5.2	28.4
4	Cerebrovascular diseases (I60-I69)	1,260	3.6	20.1
5	Diabetes mellitus (E10-E14)	1,242	3.6	19.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,226	3.5	19.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	736	2.1	11.7
8	Septicemia (A40-A41)	526	1.5	8.4
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	497	1.4	7.9
10	Influenza and pneumonia (J10-J18)	391	1.1	6.2
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	379	1.1	6.0
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	184	0.5	2.9
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	159	0.5	2.5
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	142	0.4	2.3
15	Viral hepatitis (B15-B19)	113	0.3	1.8
...	All other causes (Residual)	4,732	13.7	75.3
	White, non-Hispanic, female, 60-64 years			
...	All causes	44,596	100.0	900.2
1	Malignant neoplasms (C00-C97)	19,400	43.5	391.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,278	18.6	167.1
3	Chronic lower respiratory diseases (J40-J47)	3,355	7.5	67.7
4	Cerebrovascular diseases (I60-I69)	1,606	3.6	32.4
5	Diabetes mellitus (E10-E14)	1,590	3.6	32.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	976	2.2	19.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	685	1.5	13.8
8	Septicemia (A40-A41)	665	1.5	13.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	602	1.3	12.2
10	Influenza and pneumonia (J10-J18)	600	1.3	12.1
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	287	0.6	5.8
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	267	0.6	5.4
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	249	0.6	5.0
14	Alzheimer's disease (G30)	209	0.5	4.2
15	Aortic aneurysm and dissection (I71)	184	0.4	3.7
...	All other causes (Residual)	5,643	12.7	113.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	White, non-Hispanic, female, 65-69 years			
...	All causes	57,310	100.0	1,400.2
1	Malignant neoplasms (C00-C97)	22,796	39.8	556.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,732	20.5	286.6
3	Chronic lower respiratory diseases (J40-J47)	5,234	9.1	127.9
4	Cerebrovascular diseases (I60-I69)	2,471	4.3	60.4
5	Diabetes mellitus (E10-E14)	2,102	3.7	51.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,041	1.8	25.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	859	1.5	21.0
8	Influenza and pneumonia (J10-J18)	820	1.4	20.0
9	Septicemia (A40-A41)	793	1.4	19.4
10	Chronic liver disease and cirrhosis (K70,K73-K74)	735	1.3	18.0
11	Alzheimer's disease (G30)	430	0.8	10.5
12	Aortic aneurysm and dissection (I71)	336	0.6	8.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	332	0.6	8.1
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	313	0.5	7.6
15	Parkinson's disease (G20-G21)	193	0.3	4.7
...	All other causes (Residual)	7,123	12.4	174.0
	White, non-Hispanic, female, 70-74 years			
...	All causes	86,148	100.0	2,242.3
1	Malignant neoplasms (C00-C97)	29,383	34.1	764.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	19,599	22.8	510.1
3	Chronic lower respiratory diseases (J40-J47)	8,064	9.4	209.9
4	Cerebrovascular diseases (I60-I69)	4,841	5.6	126.0
5	Diabetes mellitus (E10-E14)	3,008	3.5	78.3
6	Influenza and pneumonia (J10-J18)	1,476	1.7	38.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,473	1.7	38.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,418	1.6	36.9
9	Alzheimer's disease (G30)	1,396	1.6	36.3
10	Septicemia (A40-A41)	1,315	1.5	34.2
11	Chronic liver disease and cirrhosis (K70,K73-K74)	894	1.0	23.3
12	Aortic aneurysm and dissection (I71)	582	0.7	15.1
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	573	0.7	14.9
14	Parkinson's disease (G20-G21)	525	0.6	13.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	496	0.6	12.9
...	All other causes (Residual)	11,105	12.9	289.0

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total	
			deaths	Rate
White, non-Hispanic, female, 75-79 years				
...	All causes	131,400	100.0	3,624.5
1	Malignant neoplasms (C00-C97)	35,423	27.0	977.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	33,748	25.7	930.9
3	Chronic lower respiratory diseases (J40-J47)	11,446	8.7	315.7
4	Cerebrovascular diseases (I60-I69)	9,598	7.3	264.8
5	Diabetes mellitus (E10-E14)	4,232	3.2	116.7
6	Alzheimer's disease (G30)	3,946	3.0	108.8
7	Influenza and pneumonia (J10-J18)	3,058	2.3	84.4
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,281	1.7	62.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,273	1.7	62.7
10	Septicemia (A40-A41)	2,133	1.6	58.8
11	Parkinson's disease (G20-G21)	1,242	0.9	34.3
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,196	0.9	33.0
13	Chronic liver disease and cirrhosis (K70,K73-K74)	932	0.7	25.7
14	Aortic aneurysm and dissection (I71)	920	0.7	25.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	876	0.7	24.2
...	All other causes (Residual)	18,096	13.8	499.2
White, non-Hispanic, female, 80-84 years				
...	All causes	175,389	100.0	6,103.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	52,302	29.8	1,820.0
2	Malignant neoplasms (C00-C97)	33,621	19.2	1,169.9
3	Cerebrovascular diseases (I60-I69)	15,980	9.1	556.1
4	Chronic lower respiratory diseases (J40-J47)	12,178	6.9	423.8
5	Alzheimer's disease (G30)	8,195	4.7	285.2
6	Influenza and pneumonia (J10-J18)	5,346	3.0	186.0
7	Diabetes mellitus (E10-E14)	4,952	2.8	172.3
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,249	1.9	113.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,156	1.8	109.8
10	Septicemia (A40-A41)	2,698	1.5	93.9
11	Parkinson's disease (G20-G21)	1,994	1.1	69.4
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,914	1.1	66.6
13	Pneumonitis due to solids and liquids (J69)	1,429	0.8	49.7
14	Atherosclerosis (I70)	1,198	0.7	41.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,146	0.7	39.9
...	All other causes (Residual)	26,031	14.8	905.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	White, non-Hispanic, female, 85-89 years			
...	All causes	187,074	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	63,117	33.7	...
2	Malignant neoplasms (C00-C97)	23,905	12.8	...
3	Cerebrovascular diseases (I60-I69)	19,110	10.2	...
4	Alzheimer's disease (G30)	11,375	6.1	...
5	Chronic lower respiratory diseases (J40-J47)	9,428	5.0	...
6	Influenza and pneumonia (J10-J18)	7,191	3.8	...
7	Diabetes mellitus (E10-E14)	4,315	2.3	...
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,547	1.9	...
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,335	1.8	...
10	Septicemia (A40-A41)	2,703	1.4	...
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,312	1.2	...
12	Parkinson's disease (G20-G21)	1,831	1.0	...
13	Pneumonitis due to solids and liquids (J69)	1,813	1.0	...
14	Atherosclerosis (I70)	1,767	0.9	...
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,123	0.6	...
...	All other causes (Residual)	30,202	16.1	...
	White, non-Hispanic, female, 90-94 years			
...	All causes	143,307	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	54,016	37.7	...
2	Cerebrovascular diseases (I60-I69)	14,877	10.4	...
3	Malignant neoplasms (C00-C97)	11,911	8.3	...
4	Alzheimer's disease (G30)	9,938	6.9	...
5	Influenza and pneumonia (J10-J18)	6,705	4.7	...
6	Chronic lower respiratory diseases (J40-J47)	4,921	3.4	...
7	Diabetes mellitus (E10-E14)	2,583	1.8	...
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,546	1.8	...
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,344	1.6	...
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,055	1.4	...
11	Septicemia (A40-A41)	1,781	1.2	...
12	Atherosclerosis (I70)	1,732	1.2	...
13	Pneumonitis due to solids and liquids (J69)	1,519	1.1	...
14	Parkinson's disease (G20-G21)	990	0.7	...
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	728	0.5	...
...	All other causes (Residual)	24,661	17.2	...

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 95-99 years				
...	All causes	62,583	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	25,046	40.0	...
2	Cerebrovascular diseases (I60-I69)	6,182	9.9	...
3	Alzheimer's disease (G30)	4,476	7.2	...
4	Influenza and pneumonia (J10-J18)	3,409	5.4	...
5	Malignant neoplasms (C00-C97)	3,317	5.3	...
6	Chronic lower respiratory diseases (J40-J47)	1,541	2.5	...
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,100	1.8	...
8	Atherosclerosis (I70)	1,012	1.6	...
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	973	1.6	...
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	930	1.5	...
11	Diabetes mellitus (E10-E14)	872	1.4	...
12	Pneumonitis due to solids and liquids (J69)	701	1.1	...
13	Septicemia (A40-A41)	695	1.1	...
14	Parkinson's disease (G20-G21)	264	0.4	...
15	Anemias (D50-D64)	236	0.4	...
...	All other causes (Residual)	11,829	18.9	...
White, non-Hispanic, female, 100 years and over				
...	All causes	13,731	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,828	42.4	...
2	Cerebrovascular diseases (I60-I69)	1,168	8.5	...
3	Influenza and pneumonia (J10-J18)	942	6.9	...
4	Alzheimer's disease (G30)	766	5.6	...
5	Malignant neoplasms (C00-C97)	444	3.2	...
6	Atherosclerosis (I70)	287	2.1	...
7	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	264	1.9	...
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	242	1.8	...
9	Chronic lower respiratory diseases (J40-J47)	220	1.6	...
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	185	1.3	...
11	Pneumonitis due to solids and liquids (J69)	147	1.1	...
12	Diabetes mellitus (E10-E14)	122	0.9	...
13	Septicemia (A40-A41)	111	0.8	...
14	Anemias (D50-D64)	56	0.4	...
15	Nutritional deficiencies (E40-E64)	46	0.3	...
...	All other causes (Residual)	2,903	21.1	...

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, all ages				
...	All causes	287,968	100.0	788.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	76,452	26.5	209.4
2	Malignant neoplasms (C00-C97)	62,054	21.5	170.0
3	Cerebrovascular diseases (I60-I69)	18,649	6.5	51.1
4	Diabetes mellitus (E10-E14)	12,787	4.4	35.0
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,154	4.2	33.3
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8,276	2.9	22.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,804	2.7	21.4
8	Chronic lower respiratory diseases (J40-J47)	7,632	2.7	20.9
9	Human immunodeficiency virus (HIV) disease (B20-B24)	7,351	2.6	20.1
10	Septicemia (A40-A41)	6,141	2.1	16.8
11	Influenza and pneumonia (J10-J18)	5,798	2.0	15.9
12	Certain conditions originating in the perinatal period (P00-P96)	4,716	1.6	12.9
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	4,445	1.5	12.2
14	Alzheimer's disease (G30)	3,693	1.3	10.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,582	0.9	7.1
...	All other causes (Residual)	47,434	16.5	129.9
Black, non-Hispanic, both sexes, 1-4 years				
...	All causes	1,169	100.0	48.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	333	28.5	13.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	155	13.3	6.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	118	10.1	4.8
4	Malignant neoplasms (C00-C97)	55	4.7	2.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	55	4.7	2.3
6	Certain conditions originating in the perinatal period (P00-P96)	29	2.5	1.2
7	Influenza and pneumonia (J10-J18)	27	2.3	1.1
8	Septicemia (A40-A41)	24	2.1	1.0
8	Chronic lower respiratory diseases (J40-J47)	24	2.1	1.0
10	Cerebrovascular diseases (I60-I69)	14	1.2	*
11	Anemias (D50-D64)	12	1.0	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.8	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	9	0.8	*
14	Acute bronchitis and bronchiolitis (J20-J21)	7	0.6	*
15	Meningococcal infection (A39)	6	0.5	*
...	All other causes (Residual)	292	25.0	12.0

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 5-9 years				
...	All causes	628	100.0	20.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	233	37.1	7.7
2	Malignant neoplasms (C00-C97)	73	11.6	2.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	54	8.6	1.8
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	36	5.7	1.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	33	5.3	1.1
6	Chronic lower respiratory diseases (J40-J47)	22	3.5	0.7
7	Influenza and pneumonia (J10-J18)	19	3.0	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	13	2.1	*
9	Anemias (D50-D64)	12	1.9	*
10	Septicemia (A40-A41)	10	1.6	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	1.4	*
12	Cerebrovascular diseases (I60-I69)	6	1.0	*
13	Meningitis (G00,G03)	2	0.3	*
13	Diseases of appendix (K35-K38)	2	0.3	*
13	Hernia (K40-K46)	2	0.3	*
...	All other causes (Residual)	102	16.2	3.3
Black, non-Hispanic, both sexes, 10-14 years				
...	All causes	902	100.0	26.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	284	31.5	8.4
2	Malignant neoplasms (C00-C97)	100	11.1	3.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	93	10.3	2.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	43	4.8	1.3
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	42	4.7	1.2
6	Chronic lower respiratory diseases (J40-J47)	40	4.4	1.2
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	37	4.1	1.1
8	Human immunodeficiency virus (HIV) disease (B20-B24)	18	2.0	*
9	Anemias (D50-D64)	13	1.4	*
10	Cerebrovascular diseases (I60-I69)	10	1.1	*
11	Influenza and pneumonia (J10-J18)	9	1.0	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.9	*
13	Diabetes mellitus (E10-E14)	6	0.7	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.7	*
15	Septicemia (A40-A41)	5	0.6	*
...	All other causes (Residual)	188	20.8	5.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 15-19 years				
...	All causes	2,530	100.0	82.3
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,052	41.6	34.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	656	25.9	21.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	119	4.7	3.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	108	4.3	3.5
5	Malignant neoplasms (C00-C97)	93	3.7	3.0
6	Chronic lower respiratory diseases (J40-J47)	40	1.6	1.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	40	1.6	1.3
8	Cerebrovascular diseases (I60-I69)	21	0.8	0.7
9	Human immunodeficiency virus (HIV) disease (B20-B24)	20	0.8	0.7
10	Anemias (D50-D64)	19	0.8	*
11	Septicemia (A40-A41)	18	0.7	*
12	Influenza and pneumonia (J10-J18)	17	0.7	*
13	Diabetes mellitus (E10-E14)	16	0.6	*
14	Pregnancy, childbirth and the puerperium (O00-099)	12	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	0.4	*
...	All other causes (Residual)	288	11.4	9.4
Black, non-Hispanic, both sexes, 20-24 years				
...	All causes	4,491	100.0	152.2
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,903	42.4	64.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	934	20.8	31.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	319	7.1	10.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	238	5.3	8.1
5	Malignant neoplasms (C00-C97)	152	3.4	5.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	97	2.2	3.3
7	Anemias (D50-D64)	59	1.3	2.0
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	52	1.2	1.8
9	Chronic lower respiratory diseases (J40-J47)	41	0.9	1.4
10	Cerebrovascular diseases (I60-I69)	37	0.8	1.3
11	Pregnancy, childbirth and the puerperium (O00-099)	30	0.7	1.0
12	Diabetes mellitus (E10-E14)	29	0.6	1.0
13	Legal intervention (Y35,Y89.0)	27	0.6	0.9
14	Influenza and pneumonia (J10-J18)	25	0.6	0.8
15	Septicemia (A40-A41)	23	0.5	0.8
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	0.5	0.8
...	All other causes (Residual)	502	11.2	17.0

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 25-29 years				
...	All causes	4,447	100.0	175.4
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,443	32.4	56.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	777	17.5	30.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	345	7.8	13.6
4	Human immunodeficiency virus (HIV) disease (B20-B24)	295	6.6	11.6
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	249	5.6	9.8
6	Malignant neoplasms (C00-C97)	226	5.1	8.9
7	Diabetes mellitus (E10-E14)	85	1.9	3.4
8	Cerebrovascular diseases (I60-I69)	53	1.2	2.1
9	Anemias (D50-D64)	52	1.2	2.1
10	Chronic lower respiratory diseases (J40-J47)	45	1.0	1.8
11	Influenza and pneumonia (J10-J18)	43	1.0	1.7
12	Septicemia (A40-A41)	42	0.9	1.7
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	37	0.8	1.5
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	33	0.7	1.3
15	Pregnancy, childbirth and the puerperium (O00-099)	32	0.7	1.3
...	All other causes (Residual)	690	15.5	27.2
Black, non-Hispanic, both sexes, 30-34 years				
...	All causes	5,348	100.0	201.8
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,000	18.7	37.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	825	15.4	31.1
3	Human immunodeficiency virus (HIV) disease (B20-B24)	643	12.0	24.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	629	11.8	23.7
5	Malignant neoplasms (C00-C97)	430	8.0	16.2
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	249	4.7	9.4
7	Diabetes mellitus (E10-E14)	123	2.3	4.6
8	Cerebrovascular diseases (I60-I69)	114	2.1	4.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	81	1.5	3.1
10	Anemias (D50-D64)	62	1.2	2.3
11	Septicemia (A40-A41)	60	1.1	2.3
12	Chronic lower respiratory diseases (J40-J47)	59	1.1	2.2
13	Influenza and pneumonia (J10-J18)	58	1.1	2.2
14	Pregnancy, childbirth and the puerperium (O00-099)	50	0.9	1.9
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	42	0.8	1.6
...	All other causes (Residual)	923	17.3	34.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 35-39 years				
...	All causes	7,746	100.0	285.1
1	Human immunodeficiency virus (HIV) disease (B20-B24)	1,291	16.7	47.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,118	14.4	41.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	997	12.9	36.7
4	Malignant neoplasms (C00-C97)	873	11.3	32.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	715	9.2	26.3
6	Cerebrovascular diseases (I60-I69)	248	3.2	9.1
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	209	2.7	7.7
8	Diabetes mellitus (E10-E14)	208	2.7	7.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	124	1.6	4.6
10	Septicemia (A40-A41)	101	1.3	3.7
10	Chronic liver disease and cirrhosis (K70,K73-K74)	101	1.3	3.7
12	Influenza and pneumonia (J10-J18)	93	1.2	3.4
13	Chronic lower respiratory diseases (J40-J47)	81	1.0	3.0
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	63	0.8	2.3
15	Anemias (D50-D64)	61	0.8	2.2
...	All other causes (Residual)	1,463	18.9	53.9
Black, non-Hispanic, both sexes, 40-44 years				
...	All causes	11,815	100.0	421.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,274	19.2	81.1
2	Malignant neoplasms (C00-C97)	1,914	16.2	68.2
3	Human immunodeficiency virus (HIV) disease (B20-B24)	1,495	12.7	53.3
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,198	10.1	42.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	620	5.2	22.1
6	Cerebrovascular diseases (I60-I69)	538	4.6	19.2
7	Diabetes mellitus (E10-E14)	339	2.9	12.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	231	2.0	8.2
9	Chronic liver disease and cirrhosis (K70,K73-K74)	222	1.9	7.9
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	200	1.7	7.1
11	Septicemia (A40-A41)	182	1.5	6.5
12	Chronic lower respiratory diseases (J40-J47)	164	1.4	5.8
13	Influenza and pneumonia (J10-J18)	158	1.3	5.6
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	111	0.9	4.0
15	Viral hepatitis (B15-B19)	77	0.7	2.7
...	All other causes (Residual)	2,092	17.7	74.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 45-49 years				
...	All causes	16,498	100.0	647.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,642	22.1	143.0
2	Malignant neoplasms (C00-C97)	3,604	21.8	141.5
3	Human immunodeficiency virus (HIV) disease (B20-B24)	1,388	8.4	54.5
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,247	7.6	49.0
5	Cerebrovascular diseases (I60-I69)	820	5.0	32.2
6	Diabetes mellitus (E10-E14)	578	3.5	22.7
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	460	2.8	18.1
8	Chronic liver disease and cirrhosis (K70,K73-K74)	390	2.4	15.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	332	2.0	13.0
10	Septicemia (A40-A41)	274	1.7	10.8
11	Chronic lower respiratory diseases (J40-J47)	239	1.4	9.4
12	Influenza and pneumonia (J10-J18)	221	1.3	8.7
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	183	1.1	7.2
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	140	0.8	5.5
15	Viral hepatitis (B15-B19)	136	0.8	5.3
...	All other causes (Residual)	2,844	17.2	111.7
Black, non-Hispanic, both sexes, 50-54 years				
...	All causes	20,086	100.0	959.3
1	Malignant neoplasms (C00-C97)	5,198	25.9	248.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,983	24.8	238.0
3	Cerebrovascular diseases (I60-I69)	1,129	5.6	53.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,042	5.2	49.8
5	Human immunodeficiency virus (HIV) disease (B20-B24)	994	4.9	47.5
6	Diabetes mellitus (E10-E14)	853	4.2	40.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	546	2.7	26.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	448	2.2	21.4
9	Septicemia (A40-A41)	376	1.9	18.0
10	Chronic lower respiratory diseases (J40-J47)	358	1.8	17.1
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	271	1.3	12.9
12	Influenza and pneumonia (J10-J18)	266	1.3	12.7
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	258	1.3	12.3
14	Viral hepatitis (B15-B19)	216	1.1	10.3
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	126	0.6	6.0
...	All other causes (Residual)	3,022	15.0	144.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 55-59 years				
...	All causes	20,251	100.0	1,321.4
1	Malignant neoplasms (C00-C97)	5,990	29.6	390.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,440	26.9	355.0
3	Cerebrovascular diseases (I60-I69)	1,132	5.6	73.9
4	Diabetes mellitus (E10-E14)	1,092	5.4	71.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	683	3.4	44.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	538	2.7	35.1
7	Human immunodeficiency virus (HIV) disease (B20-B24)	527	2.6	34.4
8	Chronic lower respiratory diseases (J40-J47)	491	2.4	32.0
9	Chronic liver disease and cirrhosis (K70,K73-K74)	409	2.0	26.7
10	Septicemia (A40-A41)	396	2.0	25.8
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	323	1.6	21.1
12	Influenza and pneumonia (J10-J18)	247	1.2	16.1
13	Viral hepatitis (B15-B19)	147	0.7	9.6
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	139	0.7	9.1
15	Aortic aneurysm and dissection (I71)	90	0.4	5.9
...	All other causes (Residual)	2,607	12.9	170.1
Black, non-Hispanic, both sexes, 60-64 years				
...	All causes	22,099	100.0	1,920.6
1	Malignant neoplasms (C00-C97)	6,913	31.3	600.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,224	28.2	540.9
3	Cerebrovascular diseases (I60-I69)	1,317	6.0	114.5
4	Diabetes mellitus (E10-E14)	1,255	5.7	109.1
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	634	2.9	55.1
6	Chronic lower respiratory diseases (J40-J47)	618	2.8	53.7
7	Septicemia (A40-A41)	485	2.2	42.2
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	469	2.1	40.8
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	341	1.5	29.6
10	Influenza and pneumonia (J10-J18)	318	1.4	27.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	291	1.3	25.3
12	Human immunodeficiency virus (HIV) disease (B20-B24)	275	1.2	23.9
13	Aortic aneurysm and dissection (I71)	106	0.5	9.2
14	Pneumonitis due to solids and liquids (J69)	99	0.4	8.6
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	79	0.4	6.9
...	All other causes (Residual)	2,675	12.1	232.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total	
			deaths	Rate
Black, non-Hispanic, both sexes, 65-69 years				
...	All causes	24,501	100.0	2,640.7
1	Malignant neoplasms (C00-C97)	7,577	30.9	816.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,093	28.9	764.5
3	Diabetes mellitus (E10-E14)	1,484	6.1	159.9
4	Cerebrovascular diseases (I60-I69)	1,483	6.1	159.8
5	Chronic lower respiratory diseases (J40-J47)	820	3.3	88.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	795	3.2	85.7
7	Septicemia (A40-A41)	575	2.3	62.0
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	451	1.8	48.6
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	405	1.7	43.7
10	Influenza and pneumonia (J10-J18)	391	1.6	42.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	239	1.0	25.8
12	Human immunodeficiency virus (HIV) disease (B20-B24)	159	0.6	17.1
13	Aortic aneurysm and dissection (I71)	111	0.5	12.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	107	0.4	11.5
15	Pneumonitis due to solids and liquids (J69)	103	0.4	11.1
...	All other causes (Residual)	2,708	11.1	291.9
Black, non-Hispanic, both sexes, 70-74 years				
...	All causes	28,082	100.0	3,810.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,316	29.6	1,128.5
2	Malignant neoplasms (C00-C97)	8,089	28.8	1,097.7
3	Cerebrovascular diseases (I60-I69)	1,928	6.9	261.6
4	Diabetes mellitus (E10-E14)	1,661	5.9	225.4
5	Chronic lower respiratory diseases (J40-J47)	1,100	3.9	149.3
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	963	3.4	130.7
7	Septicemia (A40-A41)	657	2.3	89.2
8	Influenza and pneumonia (J10-J18)	524	1.9	71.1
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	513	1.8	69.6
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	399	1.4	54.1
11	Alzheimer's disease (G30)	226	0.8	30.7
12	Chronic liver disease and cirrhosis (K70,K73-K74)	162	0.6	22.0
13	Aortic aneurysm and dissection (I71)	131	0.5	17.8
14	Pneumonitis due to solids and liquids (J69)	128	0.5	17.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	115	0.4	15.6
...	All other causes (Residual)	3,170	11.3	430.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Percent of total deaths		
		Number	Rate	Rate
Black, non-Hispanic, both sexes, 75-79 years				
...	All causes	31,540	100.0	5,490.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,512	30.2	1,655.8
2	Malignant neoplasms (C00-C97)	8,071	25.6	1,405.0
3	Cerebrovascular diseases (I60-I69)	2,530	8.0	440.4
4	Diabetes mellitus (E10-E14)	1,743	5.5	303.4
5	Chronic lower respiratory diseases (J40-J47)	1,233	3.9	214.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	998	3.2	173.7
7	Septicemia (A40-A41)	778	2.5	135.4
8	Influenza and pneumonia (J10-J18)	712	2.3	123.9
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	594	1.9	103.4
10	Alzheimer's disease (G30)	535	1.7	93.1
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	393	1.2	68.4
12	Pneumonitis due to solids and liquids (J69)	234	0.7	40.7
13	Aortic aneurysm and dissection (I71)	151	0.5	26.3
14	Parkinson's disease (G20-G21)	134	0.4	23.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	129	0.4	22.5
...	All other causes (Residual)	3,793	12.0	660.3
Black, non-Hispanic, both sexes, 80-84 years				
...	All causes	30,230	100.0	7,836.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,534	31.5	2,471.6
2	Malignant neoplasms (C00-C97)	6,411	21.2	1,662.0
3	Cerebrovascular diseases (I60-I69)	2,646	8.8	686.0
4	Diabetes mellitus (E10-E14)	1,522	5.0	394.6
5	Chronic lower respiratory diseases (J40-J47)	1,057	3.5	274.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,017	3.4	263.6
7	Alzheimer's disease (G30)	812	2.7	210.5
8	Influenza and pneumonia (J10-J18)	802	2.7	207.9
9	Septicemia (A40-A41)	758	2.5	196.5
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	568	1.9	147.2
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	405	1.3	105.0
12	Pneumonitis due to solids and liquids (J69)	299	1.0	77.5
13	Atherosclerosis (I70)	176	0.6	45.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	155	0.5	40.2
15	Parkinson's disease (G20-G21)	148	0.5	38.4
...	All other causes (Residual)	3,920	13.0	1,016.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 85-89 years				
...	All causes	23,345	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,888	33.8	...
2	Malignant neoplasms (C00-C97)	3,814	16.3	...
3	Cerebrovascular diseases (I60-I69)	2,273	9.7	...
4	Diabetes mellitus (E10-E14)	1,024	4.4	...
5	Alzheimer's disease (G30)	900	3.9	...
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	769	3.3	...
7	Influenza and pneumonia (J10-J18)	750	3.2	...
8	Chronic lower respiratory diseases (J40-J47)	670	2.9	...
9	Septicemia (A40-A41)	654	2.8	...
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	446	1.9	...
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	276	1.2	...
12	Pneumonitis due to solids and liquids (J69)	252	1.1	...
13	Atherosclerosis (I70)	153	0.7	...
14	Parkinson's disease (G20-G21)	112	0.5	...
15	Aortic aneurysm and dissection (I71)	99	0.4	...
...	All other causes (Residual)	3,265	14.0	...
Black, non-Hispanic, both sexes, 90-94 years				
...	All causes	15,434	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,490	35.6	...
2	Malignant neoplasms (C00-C97)	1,789	11.6	...
3	Cerebrovascular diseases (I60-I69)	1,525	9.9	...
4	Alzheimer's disease (G30)	694	4.5	...
5	Influenza and pneumonia (J10-J18)	635	4.1	...
6	Diabetes mellitus (E10-E14)	537	3.5	...
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	494	3.2	...
8	Septicemia (A40-A41)	384	2.5	...
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	366	2.4	...
10	Chronic lower respiratory diseases (J40-J47)	360	2.3	...
11	Pneumonitis due to solids and liquids (J69)	204	1.3	...
12	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	189	1.2	...
13	Atherosclerosis (I70)	148	1.0	...
14	Anemias (D50-D64)	75	0.5	...
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	70	0.5	...
...	All other causes (Residual)	2,474	16.0	...

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 95-99 years				
...	All causes	6,458	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,466	38.2	...
2	Cerebrovascular diseases (I60-I69)	598	9.3	...
3	Malignant neoplasms (C00-C97)	560	8.7	...
4	Alzheimer's disease (G30)	304	4.7	...
5	Influenza and pneumonia (J10-J18)	278	4.3	...
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	190	2.9	...
7	Diabetes mellitus (E10-E14)	183	2.8	...
8	Septicemia (A40-A41)	170	2.6	...
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	154	2.4	...
10	Chronic lower respiratory diseases (J40-J47)	128	2.0	...
11	Atherosclerosis (I70)	78	1.2	...
12	Pneumonitis due to solids and liquids (J69)	77	1.2	...
13	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	59	0.9	...
14	Nutritional deficiencies (E40-E64)	35	0.5	...
15	Anemias (D50-D64)	30	0.5	...
...	All other causes (Residual)	1,148	17.8	...
Black, non-Hispanic, both sexes, 100 years and over				
...	All causes	2,189	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	890	40.7	...
2	Cerebrovascular diseases (I60-I69)	196	9.0	...
3	Malignant neoplasms (C00-C97)	115	5.3	...
4	Influenza and pneumonia (J10-J18)	113	5.2	...
5	Alzheimer's disease (G30)	89	4.1	...
6	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	57	2.6	...
7	Septicemia (A40-A41)	49	2.2	...
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	48	2.2	...
9	Diabetes mellitus (E10-E14)	45	2.1	...
10	Chronic lower respiratory diseases (J40-J47)	35	1.6	...
11	Atherosclerosis (I70)	34	1.6	...
12	Pneumonitis due to solids and liquids (J69)	24	1.1	...
13	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	20	0.9	...
14	Nutritional deficiencies (E40-E64)	15	0.7	...
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.4	...
...	All other causes (Residual)	451	20.6	...

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, all ages				
...	All causes	146,136	100.0	840.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	36,962	25.3	212.6
2	Malignant neoplasms (C00-C97)	32,131	22.0	184.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,237	5.6	47.4
4	Cerebrovascular diseases (I60-I69)	7,798	5.3	44.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6,980	4.8	40.1
6	Diabetes mellitus (E10-E14)	5,380	3.7	30.9
7	Human immunodeficiency virus (HIV) disease (B20-B24)	4,947	3.4	28.5
8	Chronic lower respiratory diseases (J40-J47)	4,202	2.9	24.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,614	2.5	20.8
10	Influenza and pneumonia (J10-J18)	2,861	2.0	16.5
11	Septicemia (A40-A41)	2,748	1.9	15.8
12	Certain conditions originating in the perinatal period (P00-P96)	2,687	1.8	15.5
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,875	1.3	10.8
14	Chronic liver disease and cirrhosis (K70,K73-K74)	1,696	1.2	9.8
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,566	1.1	9.0
...	All other causes (Residual)	22,452	15.4	129.1
Black, non-Hispanic, male, 1-4 years				
...	All causes	681	100.0	55.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	197	28.9	15.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	96	14.1	7.8
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	62	9.1	5.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	31	4.6	2.5
5	Malignant neoplasms (C00-C97)	30	4.4	2.4
6	Chronic lower respiratory diseases (J40-J47)	19	2.8	*
7	Certain conditions originating in the perinatal period (P00-P96)	16	2.3	*
8	Septicemia (A40-A41)	14	2.1	*
9	Influenza and pneumonia (J10-J18)	13	1.9	*
10	Cerebrovascular diseases (I60-I69)	9	1.3	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	1.2	*
12	Anemias (D50-D64)	7	1.0	*
13	Acute bronchitis and bronchiolitis (J20-J21)	6	0.9	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.7	*
15	Meningococcal infection (A39)	3	0.4	*
...	All other causes (Residual)	165	24.2	13.4

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 5-9 years				
...	All causes	352	100.0	22.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	141	40.1	9.1
2	Malignant neoplasms (C00-C97)	40	11.4	2.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	33	9.4	2.1
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	20	5.7	1.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	18	5.1	*
6	Chronic lower respiratory diseases (J40-J47)	16	4.5	*
7	Influenza and pneumonia (J10-J18)	12	3.4	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	7	2.0	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	1.4	*
9	Anemias (D50-D64)	5	1.4	*
9	Cerebrovascular diseases (I60-I69)	5	1.4	*
12	Septicemia (A40-A41)	4	1.1	*
13	Meningococcal infection (A39)	1	0.3	*
13	Diabetes mellitus (E10-E14)	1	0.3	*
13	Meningitis (G00,G03)	1	0.3	*
13	Pneumonitis due to solids and liquids (J69)	1	0.3	*
13	Diseases of appendix (K35-K38)	1	0.3	*
13	Hernia (K40-K46)	1	0.3	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1	0.3	*
...	All other causes (Residual)	38	10.8	2.5
Black, non-Hispanic, male, 10-14 years				
...	All causes	559	100.0	32.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	189	33.8	11.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	65	11.6	3.8
3	Malignant neoplasms (C00-C97)	56	10.0	3.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	33	5.9	1.9
5	Chronic lower respiratory diseases (J40-J47)	27	4.8	1.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	24	4.3	1.4
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	18	3.2	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	10	1.8	*
9	Anemias (D50-D64)	7	1.3	*
10	Cerebrovascular diseases (I60-I69)	4	0.7	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.5	*
11	Influenza and pneumonia (J10-J18)	3	0.5	*
11	Diseases of appendix (K35-K38)	3	0.5	*
14	Septicemia (A40-A41)	2	0.4	*
14	Diabetes mellitus (E10-E14)	2	0.4	*
...	All other causes (Residual)	113	20.2	6.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 15-19 years				
...	All causes	1,938	100.0	124.3
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	946	48.8	60.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	482	24.9	30.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	105	5.4	6.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	72	3.7	4.6
5	Malignant neoplasms (C00-C97)	56	2.9	3.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	27	1.4	1.7
7	Chronic lower respiratory diseases (J40-J47)	25	1.3	1.6
8	Cerebrovascular diseases (I60-I69)	13	0.7	*
9	Anemias (D50-D64)	12	0.6	*
10	Diabetes mellitus (E10-E14)	10	0.5	*
10	Influenza and pneumonia (J10-J18)	10	0.5	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	9	0.5	*
13	Septicemia (A40-A41)	8	0.4	*
14	Legal intervention (Y35,Y89.0)	6	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.3	*
...	All other causes (Residual)	152	7.8	9.7
Black, non-Hispanic, male, 20-24 years				
...	All causes	3,405	100.0	232.0
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,696	49.8	115.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	723	21.2	49.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	272	8.0	18.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	153	4.5	10.4
5	Malignant neoplasms (C00-C97)	85	2.5	5.8
6	Human immunodeficiency virus (HIV) disease (B20-B24)	48	1.4	3.3
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	28	0.8	1.9
8	Anemias (D50-D64)	27	0.8	1.8
9	Legal intervention (Y35,Y89.0)	25	0.7	1.7
10	Chronic lower respiratory diseases (J40-J47)	23	0.7	1.6
11	Diabetes mellitus (E10-E14)	16	0.5	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	0.4	*
13	Influenza and pneumonia (J10-J18)	14	0.4	*
14	Cerebrovascular diseases (I60-I69)	12	0.4	*
15	Septicemia (A40-A41)	11	0.3	*
...	All other causes (Residual)	257	7.5	17.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 25-29 years				
...	All causes	3,161	100.0	259.9
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,292	40.9	106.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	583	18.4	47.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	220	7.0	18.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	202	6.4	16.6
5	Human immunodeficiency virus (HIV) disease (B20-B24)	160	5.1	13.2
6	Malignant neoplasms (C00-C97)	89	2.8	7.3
7	Diabetes mellitus (E10-E14)	53	1.7	4.4
8	Cerebrovascular diseases (I60-I69)	32	1.0	2.6
9	Chronic lower respiratory diseases (J40-J47)	31	1.0	2.5
10	Anemias (D50-D64)	28	0.9	2.3
11	Legal intervention (Y35,Y89.0)	27	0.9	2.2
12	Influenza and pneumonia (J10-J18)	21	0.7	1.7
13	Septicemia (A40-A41)	20	0.6	1.6
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	19	0.6	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18	0.6	*
...	All other causes (Residual)	366	11.6	30.1
Black, non-Hispanic, male, 30-34 years				
...	All causes	3,431	100.0	273.1
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	836	24.4	66.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	614	17.9	48.9
3	Human immunodeficiency virus (HIV) disease (B20-B24)	377	11.0	30.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	370	10.8	29.5
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	204	5.9	16.2
6	Malignant neoplasms (C00-C97)	174	5.1	13.9
7	Diabetes mellitus (E10-E14)	69	2.0	5.5
8	Cerebrovascular diseases (I60-I69)	54	1.6	4.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	46	1.3	3.7
10	Chronic lower respiratory diseases (J40-J47)	41	1.2	3.3
11	Influenza and pneumonia (J10-J18)	37	1.1	2.9
12	Septicemia (A40-A41)	36	1.0	2.9
13	Anemias (D50-D64)	30	0.9	2.4
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	28	0.8	2.2
15	Legal intervention (Y35,Y89.0)	24	0.7	1.9
...	All other causes (Residual)	491	14.3	39.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 35-39 years				
...	All causes	4,571	100.0	357.4
1	Human immunodeficiency virus (HIV) disease (B20-B24)	817	17.9	63.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	725	15.9	56.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	682	14.9	53.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	580	12.7	45.4
5	Malignant neoplasms (C00-C97)	288	6.3	22.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	171	3.7	13.4
7	Cerebrovascular diseases (I60-I69)	129	2.8	10.1
8	Diabetes mellitus (E10-E14)	113	2.5	8.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	71	1.6	5.6
10	Chronic liver disease and cirrhosis (K70,K73-K74)	58	1.3	4.5
11	Septicemia (A40-A41)	48	1.1	3.8
12	Influenza and pneumonia (J10-J18)	47	1.0	3.7
13	Chronic lower respiratory diseases (J40-J47)	38	0.8	3.0
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	31	0.7	2.4
15	Anemias (D50-D64)	28	0.6	2.2
...	All other causes (Residual)	745	16.3	58.3
Black, non-Hispanic, male, 40-44 years				
...	All causes	6,834	100.0	520.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,384	20.3	105.4
2	Human immunodeficiency virus (HIV) disease (B20-B24)	997	14.6	75.9
3	Malignant neoplasms (C00-C97)	824	12.1	62.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	814	11.9	62.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	498	7.3	37.9
6	Cerebrovascular diseases (I60-I69)	256	3.7	19.5
7	Diabetes mellitus (E10-E14)	196	2.9	14.9
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	153	2.2	11.6
9	Chronic liver disease and cirrhosis (K70,K73-K74)	130	1.9	9.9
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	120	1.8	9.1
11	Influenza and pneumonia (J10-J18)	95	1.4	7.2
12	Septicemia (A40-A41)	86	1.3	6.5
13	Chronic lower respiratory diseases (J40-J47)	68	1.0	5.2
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	61	0.9	4.6
15	Viral hepatitis (B15-B19)	48	0.7	3.7
...	All other causes (Residual)	1,104	16.2	84.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 45-49 years				
...	All causes	9,713	100.0	819.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,280	23.5	192.4
2	Malignant neoplasms (C00-C97)	1,767	18.2	149.1
3	Human immunodeficiency virus (HIV) disease (B20-B24)	962	9.9	81.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	896	9.2	75.6
5	Cerebrovascular diseases (I60-I69)	415	4.3	35.0
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	363	3.7	30.6
7	Diabetes mellitus (E10-E14)	310	3.2	26.2
8	Chronic liver disease and cirrhosis (K70,K73-K74)	261	2.7	22.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	181	1.9	15.3
10	Septicemia (A40-A41)	143	1.5	12.1
11	Influenza and pneumonia (J10-J18)	124	1.3	10.5
12	Chronic lower respiratory diseases (J40-J47)	119	1.2	10.0
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	117	1.2	9.9
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	108	1.1	9.1
15	Viral hepatitis (B15-B19)	92	0.9	7.8
...	All other causes (Residual)	1,575	16.2	132.9
Black, non-Hispanic, male, 50-54 years				
...	All causes	11,995	100.0	1,251.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,163	26.4	329.9
2	Malignant neoplasms (C00-C97)	2,766	23.1	288.5
3	Human immunodeficiency virus (HIV) disease (B20-B24)	731	6.1	76.3
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	729	6.1	76.0
5	Cerebrovascular diseases (I60-I69)	616	5.1	64.3
6	Diabetes mellitus (E10-E14)	431	3.6	45.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	403	3.4	42.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	286	2.4	29.8
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	213	1.8	22.2
10	Septicemia (A40-A41)	195	1.6	20.3
11	Chronic lower respiratory diseases (J40-J47)	174	1.5	18.2
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	161	1.3	16.8
12	Influenza and pneumonia (J10-J18)	161	1.3	16.8
14	Viral hepatitis (B15-B19)	158	1.3	16.5
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	91	0.8	9.5
...	All other causes (Residual)	1,717	14.3	179.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 55-59 years				
...	All causes	11,905	100.0	1,722.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,382	28.4	489.3
2	Malignant neoplasms (C00-C97)	3,287	27.6	475.6
3	Cerebrovascular diseases (I60-I69)	639	5.4	92.5
4	Diabetes mellitus (E10-E14)	545	4.6	78.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	493	4.1	71.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	404	3.4	58.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	306	2.6	44.3
8	Chronic liver disease and cirrhosis (K70,K73-K74)	294	2.5	42.5
9	Chronic lower respiratory diseases (J40-J47)	269	2.3	38.9
10	Septicemia (A40-A41)	226	1.9	32.7
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	173	1.5	25.0
12	Influenza and pneumonia (J10-J18)	152	1.3	22.0
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	120	1.0	17.4
14	Viral hepatitis (B15-B19)	106	0.9	15.3
15	Aortic aneurysm and dissection (I71)	57	0.5	8.2
...	All other causes (Residual)	1,452	12.2	210.1
Black, non-Hispanic, male, 60-64 years				
...	All causes	12,563	100.0	2,493.7
1	Malignant neoplasms (C00-C97)	3,848	30.6	763.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,658	29.1	726.1
3	Cerebrovascular diseases (I60-I69)	724	5.8	143.7
4	Diabetes mellitus (E10-E14)	624	5.0	123.9
5	Chronic lower respiratory diseases (J40-J47)	346	2.8	68.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	331	2.6	65.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	321	2.6	63.7
8	Septicemia (A40-A41)	236	1.9	46.8
9	Human immunodeficiency virus (HIV) disease (B20-B24)	210	1.7	41.7
10	Influenza and pneumonia (J10-J18)	204	1.6	40.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	198	1.6	39.3
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	186	1.5	36.9
13	Aortic aneurysm and dissection (I71)	75	0.6	14.9
14	Pneumonitis due to solids and liquids (J69)	59	0.5	11.7
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	58	0.5	11.5
...	All other causes (Residual)	1,485	11.8	294.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 65-69 years				
...	All causes	13,395	100.0	3,390.4
1	Malignant neoplasms (C00-C97)	4,208	31.4	1,065.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,999	29.9	1,012.2
3	Cerebrovascular diseases (I60-I69)	739	5.5	187.0
4	Diabetes mellitus (E10-E14)	691	5.2	174.9
5	Chronic lower respiratory diseases (J40-J47)	452	3.4	114.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	381	2.8	96.4
7	Septicemia (A40-A41)	301	2.2	76.2
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	298	2.2	75.4
9	Influenza and pneumonia (J10-J18)	245	1.8	62.0
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	200	1.5	50.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	147	1.1	37.2
12	Human immunodeficiency virus (HIV) disease (B20-B24)	117	0.9	29.6
13	Aortic aneurysm and dissection (I71)	65	0.5	16.5
14	Pneumonitis due to solids and liquids (J69)	58	0.4	14.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	50	0.4	12.7
...	All other causes (Residual)	1,444	10.8	365.5
Black, non-Hispanic, male, 70-74 years				
...	All causes	14,562	100.0	4,907.6
1	Malignant neoplasms (C00-C97)	4,385	30.1	1,477.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,378	30.1	1,475.4
3	Cerebrovascular diseases (I60-I69)	934	6.4	314.8
4	Diabetes mellitus (E10-E14)	701	4.8	236.2
5	Chronic lower respiratory diseases (J40-J47)	641	4.4	216.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	472	3.2	159.1
7	Influenza and pneumonia (J10-J18)	302	2.1	101.8
8	Septicemia (A40-A41)	300	2.1	101.1
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	224	1.5	75.5
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	220	1.5	74.1
11	Alzheimer's disease (G30)	97	0.7	32.7
12	Chronic liver disease and cirrhosis (K70,K73-K74)	86	0.6	29.0
13	Pneumonitis due to solids and liquids (J69)	79	0.5	26.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	64	0.4	21.6
15	Aortic aneurysm and dissection (I71)	59	0.4	19.9
...	All other causes (Residual)	1,620	11.1	546.0

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total	
			deaths	Rate
Black, non-Hispanic, male, 75-79 years				
...	All causes	15,124	100.0	7,066.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,503	29.8	2,104.0
2	Malignant neoplasms (C00-C97)	4,289	28.4	2,004.0
3	Cerebrovascular diseases (I60-I69)	1,094	7.2	511.2
4	Chronic lower respiratory diseases (J40-J47)	710	4.7	331.7
5	Diabetes mellitus (E10-E14)	667	4.4	311.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	417	2.8	194.8
7	Influenza and pneumonia (J10-J18)	384	2.5	179.4
8	Septicemia (A40-A41)	348	2.3	162.6
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	227	1.5	106.1
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	221	1.5	103.3
11	Alzheimer's disease (G30)	203	1.3	94.8
12	Pneumonitis due to solids and liquids (J69)	130	0.9	60.7
13	Parkinson's disease (G20-G21)	81	0.5	37.8
14	Aortic aneurysm and dissection (I71)	71	0.5	33.2
15	Atherosclerosis (I70)	62	0.4	29.0
...	All other causes (Residual)	1,717	11.4	802.2
Black, non-Hispanic, male, 80-84 years				
...	All causes	13,173	100.0	9,941.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,985	30.3	3,007.3
2	Malignant neoplasms (C00-C97)	3,251	24.7	2,453.4
3	Cerebrovascular diseases (I60-I69)	977	7.4	737.3
4	Chronic lower respiratory diseases (J40-J47)	630	4.8	475.4
5	Diabetes mellitus (E10-E14)	521	4.0	393.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	445	3.4	335.8
7	Influenza and pneumonia (J10-J18)	381	2.9	287.5
8	Septicemia (A40-A41)	315	2.4	237.7
9	Alzheimer's disease (G30)	278	2.1	209.8
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	202	1.5	152.4
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	198	1.5	149.4
12	Pneumonitis due to solids and liquids (J69)	146	1.1	110.2
13	Atherosclerosis (I70)	78	0.6	58.9
14	Parkinson's disease (G20-G21)	77	0.6	58.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	70	0.5	52.8
...	All other causes (Residual)	1,619	12.3	1,221.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 85-89 years				
...	All causes	8,119	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,587	31.9	...
2	Malignant neoplasms (C00-C97)	1,710	21.1	...
3	Cerebrovascular diseases (I60-I69)	656	8.1	...
4	Chronic lower respiratory diseases (J40-J47)	346	4.3	...
5	Diabetes mellitus (E10-E14)	285	3.5	...
6	Influenza and pneumonia (J10-J18)	283	3.5	...
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	252	3.1	...
8	Alzheimer's disease (G30)	233	2.9	...
9	Septicemia (A40-A41)	229	2.8	...
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	136	1.7	...
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	121	1.5	...
12	Pneumonitis due to solids and liquids (J69)	102	1.3	...
13	Parkinson's disease (G20-G21)	62	0.8	...
14	Atherosclerosis (I70)	50	0.6	...
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	34	0.4	...
...	All other causes (Residual)	1,033	12.7	...
Black, non-Hispanic, male, 90-94 years				
...	All causes	4,264	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,374	32.2	...
2	Malignant neoplasms (C00-C97)	731	17.1	...
3	Cerebrovascular diseases (I60-I69)	343	8.0	...
4	Influenza and pneumonia (J10-J18)	214	5.0	...
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	171	4.0	...
6	Chronic lower respiratory diseases (J40-J47)	168	3.9	...
7	Alzheimer's disease (G30)	147	3.4	...
8	Diabetes mellitus (E10-E14)	119	2.8	...
9	Septicemia (A40-A41)	112	2.6	...
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	87	2.0	...
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	71	1.7	...
12	Pneumonitis due to solids and liquids (J69)	70	1.6	...
13	Atherosclerosis (I70)	33	0.8	...
14	Parkinson's disease (G20-G21)	25	0.6	...
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	0.5	...
...	All other causes (Residual)	576	13.5	...

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 95-99 years				
...	All causes	1,433	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	497	34.7	...
2	Malignant neoplasms (C00-C97)	216	15.1	...
3	Cerebrovascular diseases (I60-I69)	113	7.9	...
4	Influenza and pneumonia (J10-J18)	76	5.3	...
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	58	4.0	...
6	Alzheimer's disease (G30)	49	3.4	...
7	Chronic lower respiratory diseases (J40-J47)	45	3.1	...
8	Septicemia (A40-A41)	36	2.5	...
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	32	2.2	...
10	Pneumonitis due to solids and liquids (J69)	26	1.8	...
11	Diabetes mellitus (E10-E14)	22	1.5	...
12	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17	1.2	...
13	Atherosclerosis (I70)	12	0.8	...
14	Anemias (D50-D64)	9	0.6	...
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.4	...
...	All other causes (Residual)	219	15.3	...
Black, non-Hispanic, male, 100 years and over				
...	All causes	340	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	139	40.9	...
2	Influenza and pneumonia (J10-J18)	30	8.8	...
3	Malignant neoplasms (C00-C97)	27	7.9	...
4	Cerebrovascular diseases (I60-I69)	15	4.4	...
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	3.8	...
6	Chronic lower respiratory diseases (J40-J47)	12	3.5	...
7	Alzheimer's disease (G30)	8	2.4	...
7	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	8	2.4	...
9	Septicemia (A40-A41)	7	2.1	...
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5	1.5	...
11	Atherosclerosis (I70)	4	1.2	...
11	Pneumonitis due to solids and liquids (J69)	4	1.2	...
13	Diabetes mellitus (E10-E14)	3	0.9	...
13	Nutritional deficiencies (E40-E64)	3	0.9	...
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.3	...
15	Parkinson's disease (G20-G21)	1	0.3	...
15	Hernia (K40-K46)	1	0.3	...
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.3	...
...	All other causes (Residual)	58	17.1	...

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, all ages				
...	All causes	141,832	100.0	741.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	39,490	27.8	206.5
2	Malignant neoplasms (C00-C97)	29,923	21.1	156.5
3	Cerebrovascular diseases (I60-I69)	10,851	7.7	56.7
4	Diabetes mellitus (E10-E14)	7,407	5.2	38.7
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,190	3.0	21.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,917	2.8	20.5
7	Chronic lower respiratory diseases (J40-J47)	3,430	2.4	17.9
8	Septicemia (A40-A41)	3,393	2.4	17.7
9	Influenza and pneumonia (J10-J18)	2,937	2.1	15.4
10	Alzheimer's disease (G30)	2,614	1.8	13.7
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,570	1.8	13.4
12	Human immunodeficiency virus (HIV) disease (B20-B24)	2,404	1.7	12.6
13	Certain conditions originating in the perinatal period (P00-P96)	2,029	1.4	10.6
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,296	0.9	6.8
15	Chronic liver disease and cirrhosis (K70,K73-K74)	886	0.6	4.6
...	All other causes (Residual)	24,495	17.3	128.1
Black, non-Hispanic, female, 1-4 years				
...	All causes	488	100.0	40.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	136	27.9	11.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	59	12.1	4.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	56	11.5	4.7
4	Malignant neoplasms (C00-C97)	25	5.1	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	24	4.9	2.0
6	Influenza and pneumonia (J10-J18)	14	2.9	*
7	Certain conditions originating in the perinatal period (P00-P96)	13	2.7	*
8	Septicemia (A40-A41)	10	2.0	*
9	Anemias (D50-D64)	5	1.0	*
9	Cerebrovascular diseases (I60-I69)	5	1.0	*
9	Chronic lower respiratory diseases (J40-J47)	5	1.0	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.8	*
13	Meningococcal infection (A39)	3	0.6	*
13	Meningitis (G00,G03)	3	0.6	*
15	Diseases of appendix (K35-K38)	2	0.4	*
15	Hernia (K40-K46)	2	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
...	All other causes (Residual)	120	24.6	10.0

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Black, non-Hispanic, female, 5-9 years			
...	All causes	276	100.0	18.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	92	33.3	6.1
2	Malignant neoplasms (C00-C97)	33	12.0	2.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	21	7.6	1.4
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	16	5.8	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	15	5.4	*
6	Anemias (D50-D64)	7	2.5	*
6	Influenza and pneumonia (J10-J18)	7	2.5	*
8	Septicemia (A40-A41)	6	2.2	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	6	2.2	*
8	Chronic lower respiratory diseases (J40-J47)	6	2.2	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	1.4	*
12	Meningitis (G00,G03)	1	0.4	*
12	Parkinson's disease (G20-G21)	1	0.4	*
12	Cerebrovascular diseases (I60-I69)	1	0.4	*
12	Diseases of appendix (K35-K38)	1	0.4	*
12	Hernia (K40-K46)	1	0.4	*
...	All other causes (Residual)	58	21.0	3.9
	Black, non-Hispanic, female, 10-14 years			
...	All causes	343	100.0	20.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	95	27.7	5.7
2	Malignant neoplasms (C00-C97)	44	12.8	2.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	28	8.2	1.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	19	5.5	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	19	5.5	*
6	Chronic lower respiratory diseases (J40-J47)	13	3.8	*
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	9	2.6	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	8	2.3	*
9	Anemias (D50-D64)	6	1.7	*
9	Cerebrovascular diseases (I60-I69)	6	1.7	*
9	Influenza and pneumonia (J10-J18)	6	1.7	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	1.5	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	1.5	*
14	Diabetes mellitus (E10-E14)	4	1.2	*
15	Meningococcal infection (A39)	3	0.9	*
15	Septicemia (A40-A41)	3	0.9	*
...	All other causes (Residual)	70	20.4	4.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 15-19 years				
...	All causes	592	100.0	39.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	174	29.4	11.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	106	17.9	7.0
3	Malignant neoplasms (C00-C97)	37	6.3	2.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	36	6.1	2.4
5	Chronic lower respiratory diseases (J40-J47)	15	2.5	*
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	14	2.4	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	13	2.2	*
8	Pregnancy, childbirth and the puerperium (O00-099)	12	2.0	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	11	1.9	*
10	Septicemia (A40-A41)	10	1.7	*
11	Cerebrovascular diseases (I60-I69)	8	1.4	*
12	Anemias (D50-D64)	7	1.2	*
12	Influenza and pneumonia (J10-J18)	7	1.2	*
14	Diabetes mellitus (E10-E14)	6	1.0	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	1.0	*
...	All other causes (Residual)	130	22.0	8.6
Black, non-Hispanic, female, 20-24 years				
...	All causes	1,086	100.0	73.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	211	19.4	14.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	207	19.1	14.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	85	7.8	5.7
4	Malignant neoplasms (C00-C97)	67	6.2	4.5
5	Human immunodeficiency virus (HIV) disease (B20-B24)	49	4.5	3.3
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	47	4.3	3.2
7	Anemias (D50-D64)	32	2.9	2.2
8	Pregnancy, childbirth and the puerperium (O00-099)	30	2.8	2.0
9	Cerebrovascular diseases (I60-I69)	25	2.3	1.7
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	24	2.2	1.6
11	Chronic lower respiratory diseases (J40-J47)	18	1.7	*
12	Diabetes mellitus (E10-E14)	13	1.2	*
13	Septicemia (A40-A41)	12	1.1	*
14	Influenza and pneumonia (J10-J18)	11	1.0	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.7	*
...	All other causes (Residual)	247	22.7	16.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 25-29 years				
...	All causes	1,286	100.0	97.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	194	15.1	14.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	151	11.7	11.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	143	11.1	10.8
4	Malignant neoplasms (C00-C97)	137	10.7	10.4
5	Human immunodeficiency virus (HIV) disease (B20-B24)	135	10.5	10.2
6	Diabetes mellitus (E10-E14)	32	2.5	2.4
6	Pregnancy, childbirth and the puerperium (O00-099)	32	2.5	2.4
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	29	2.3	2.2
9	Anemias (D50-D64)	24	1.9	1.8
10	Septicemia (A40-A41)	22	1.7	1.7
10	Influenza and pneumonia (J10-J18)	22	1.7	1.7
12	Cerebrovascular diseases (I60-I69)	21	1.6	1.6
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19	1.5	*
14	Chronic lower respiratory diseases (J40-J47)	14	1.1	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	14	1.1	*
...	All other causes (Residual)	297	23.1	22.5
Black, non-Hispanic, female, 30-34 years				
...	All causes	1,917	100.0	137.5
1	Human immunodeficiency virus (HIV) disease (B20-B24)	266	13.9	19.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	259	13.5	18.6
3	Malignant neoplasms (C00-C97)	256	13.4	18.4
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	211	11.0	15.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	164	8.6	11.8
6	Cerebrovascular diseases (I60-I69)	60	3.1	4.3
7	Diabetes mellitus (E10-E14)	54	2.8	3.9
8	Pregnancy, childbirth and the puerperium (O00-099)	50	2.6	3.6
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	45	2.3	3.2
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	35	1.8	2.5
11	Anemias (D50-D64)	32	1.7	2.3
12	Septicemia (A40-A41)	24	1.3	1.7
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	23	1.2	1.6
14	Influenza and pneumonia (J10-J18)	21	1.1	1.5
15	Chronic lower respiratory diseases (J40-J47)	18	0.9	*
...	All other causes (Residual)	399	20.8	28.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 35-39 years				
...	All causes	3,175	100.0	220.8
1	Malignant neoplasms (C00-C97)	585	18.4	40.7
2	Human immunodeficiency virus (HIV) disease (B20-B24)	474	14.9	33.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	436	13.7	30.3
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	272	8.6	18.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	135	4.3	9.4
6	Cerebrovascular diseases (I60-I69)	119	3.7	8.3
7	Diabetes mellitus (E10-E14)	95	3.0	6.6
8	Septicemia (A40-A41)	53	1.7	3.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	53	1.7	3.7
10	Influenza and pneumonia (J10-J18)	46	1.4	3.2
11	Chronic lower respiratory diseases (J40-J47)	43	1.4	3.0
11	Chronic liver disease and cirrhosis (K70,K73-K74)	43	1.4	3.0
13	Pregnancy, childbirth and the puerperium (O00-099)	40	1.3	2.8
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	38	1.2	2.6
15	Anemias (D50-D64)	33	1.0	2.3
...	All other causes (Residual)	710	22.4	49.4
Black, non-Hispanic, female, 40-44 years				
...	All causes	4,981	100.0	333.8
1	Malignant neoplasms (C00-C97)	1,090	21.9	73.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	890	17.9	59.6
3	Human immunodeficiency virus (HIV) disease (B20-B24)	498	10.0	33.4
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	384	7.7	25.7
5	Cerebrovascular diseases (I60-I69)	282	5.7	18.9
6	Diabetes mellitus (E10-E14)	143	2.9	9.6
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	122	2.4	8.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	111	2.2	7.4
9	Septicemia (A40-A41)	96	1.9	6.4
9	Chronic lower respiratory diseases (J40-J47)	96	1.9	6.4
11	Chronic liver disease and cirrhosis (K70,K73-K74)	92	1.8	6.2
12	Influenza and pneumonia (J10-J18)	63	1.3	4.2
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	50	1.0	3.4
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	47	0.9	3.1
15	Anemias (D50-D64)	32	0.6	2.1
...	All other causes (Residual)	985	19.8	66.0

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 45-49 years				
...	All causes	6,785	100.0	498.1
1	Malignant neoplasms (C00-C97)	1,837	27.1	134.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,362	20.1	100.0
3	Human immunodeficiency virus (HIV) disease (B20-B24)	426	6.3	31.3
4	Cerebrovascular diseases (I60-I69)	405	6.0	29.7
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	351	5.2	25.8
6	Diabetes mellitus (E10-E14)	268	3.9	19.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	151	2.2	11.1
8	Septicemia (A40-A41)	131	1.9	9.6
9	Chronic liver disease and cirrhosis (K70,K73-K74)	129	1.9	9.5
10	Chronic lower respiratory diseases (J40-J47)	120	1.8	8.8
11	Influenza and pneumonia (J10-J18)	97	1.4	7.1
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	97	1.4	7.1
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	66	1.0	4.8
14	Viral hepatitis (B15-B19)	44	0.6	3.2
15	Anemias (D50-D64)	38	0.6	2.8
...	All other causes (Residual)	1,263	18.6	92.7
Black, non-Hispanic, female, 50-54 years				
...	All causes	8,091	100.0	712.7
1	Malignant neoplasms (C00-C97)	2,432	30.1	214.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,820	22.5	160.3
3	Cerebrovascular diseases (I60-I69)	513	6.3	45.2
4	Diabetes mellitus (E10-E14)	422	5.2	37.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	313	3.9	27.6
6	Human immunodeficiency virus (HIV) disease (B20-B24)	263	3.3	23.2
7	Chronic lower respiratory diseases (J40-J47)	184	2.3	16.2
8	Septicemia (A40-A41)	181	2.2	15.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	162	2.0	14.3
10	Chronic liver disease and cirrhosis (K70,K73-K74)	143	1.8	12.6
11	Influenza and pneumonia (J10-J18)	105	1.3	9.2
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	97	1.2	8.5
13	Viral hepatitis (B15-B19)	58	0.7	5.1
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	58	0.7	5.1
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	35	0.4	3.1
...	All other causes (Residual)	1,305	16.1	115.0

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total	
			deaths	Rate
Black, non-Hispanic, female, 55-59 years				
...	All causes	8,346	100.0	992.0
1	Malignant neoplasms (C00-C97)	2,703	32.4	321.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,058	24.7	244.6
3	Diabetes mellitus (E10-E14)	547	6.6	65.0
4	Cerebrovascular diseases (I60-I69)	493	5.9	58.6
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	232	2.8	27.6
6	Chronic lower respiratory diseases (J40-J47)	222	2.7	26.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	190	2.3	22.6
8	Septicemia (A40-A41)	170	2.0	20.2
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	150	1.8	17.8
10	Human immunodeficiency virus (HIV) disease (B20-B24)	123	1.5	14.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	115	1.4	13.7
12	Influenza and pneumonia (J10-J18)	95	1.1	11.3
13	Viral hepatitis (B15-B19)	41	0.5	4.9
14	Aortic aneurysm and dissection (I71)	33	0.4	3.9
15	Anemias (D50-D64)	29	0.3	3.4
...	All other causes (Residual)	1,145	13.7	136.1
Black, non-Hispanic, female, 60-64 years				
...	All causes	9,536	100.0	1,474.3
1	Malignant neoplasms (C00-C97)	3,065	32.1	473.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,566	26.9	396.7
3	Diabetes mellitus (E10-E14)	631	6.6	97.6
4	Cerebrovascular diseases (I60-I69)	593	6.2	91.7
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	313	3.3	48.4
6	Chronic lower respiratory diseases (J40-J47)	272	2.9	42.1
7	Septicemia (A40-A41)	249	2.6	38.5
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	155	1.6	24.0
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	138	1.4	21.3
10	Influenza and pneumonia (J10-J18)	114	1.2	17.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	93	1.0	14.4
12	Human immunodeficiency virus (HIV) disease (B20-B24)	65	0.7	10.0
13	Pneumonitis due to solids and liquids (J69)	40	0.4	6.2
14	Viral hepatitis (B15-B19)	35	0.4	5.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	34	0.4	5.3
...	All other causes (Residual)	1,173	12.3	181.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Black, non-Hispanic, female, 65-69 years			
...	All causes	11,106	100.0	2,084.8
1	Malignant neoplasms (C00-C97)	3,369	30.3	632.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,094	27.9	580.8
3	Diabetes mellitus (E10-E14)	793	7.1	148.9
4	Cerebrovascular diseases (I60-I69)	744	6.7	139.7
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	414	3.7	77.7
6	Chronic lower respiratory diseases (J40-J47)	368	3.3	69.1
7	Septicemia (A40-A41)	274	2.5	51.4
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	205	1.8	38.5
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	153	1.4	28.7
10	Influenza and pneumonia (J10-J18)	146	1.3	27.4
11	Chronic liver disease and cirrhosis (K70,K73-K74)	92	0.8	17.3
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	57	0.5	10.7
13	Alzheimer's disease (G30)	55	0.5	10.3
14	Aortic aneurysm and dissection (I71)	46	0.4	8.6
15	Pneumonitis due to solids and liquids (J69)	45	0.4	8.4
...	All other causes (Residual)	1,251	11.3	234.8
	Black, non-Hispanic, female, 70-74 years			
...	All causes	13,520	100.0	3,071.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,938	29.1	894.7
2	Malignant neoplasms (C00-C97)	3,704	27.4	841.5
3	Cerebrovascular diseases (I60-I69)	994	7.4	225.8
4	Diabetes mellitus (E10-E14)	960	7.1	218.1
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	491	3.6	111.6
6	Chronic lower respiratory diseases (J40-J47)	459	3.4	104.3
7	Septicemia (A40-A41)	357	2.6	81.1
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	289	2.1	65.7
9	Influenza and pneumonia (J10-J18)	222	1.6	50.4
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	179	1.3	40.7
11	Alzheimer's disease (G30)	129	1.0	29.3
12	Chronic liver disease and cirrhosis (K70,K73-K74)	76	0.6	17.3
13	Aortic aneurysm and dissection (I71)	72	0.5	16.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	51	0.4	11.6
15	Pneumonitis due to solids and liquids (J69)	49	0.4	11.1
...	All other causes (Residual)	1,550	11.5	352.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total	
			deaths	Rate
Black, non-Hispanic, female, 75-79 years				
...	All causes	16,416	100.0	4,554.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,009	30.5	1,389.7
2	Malignant neoplasms (C00-C97)	3,782	23.0	1,049.3
3	Cerebrovascular diseases (I60-I69)	1,436	8.7	398.4
4	Diabetes mellitus (E10-E14)	1,076	6.6	298.5
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	581	3.5	161.2
6	Chronic lower respiratory diseases (J40-J47)	523	3.2	145.1
7	Septicemia (A40-A41)	430	2.6	119.3
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	373	2.3	103.5
9	Alzheimer's disease (G30)	332	2.0	92.1
10	Influenza and pneumonia (J10-J18)	328	2.0	91.0
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	166	1.0	46.1
12	Pneumonitis due to solids and liquids (J69)	104	0.6	28.9
13	Aortic aneurysm and dissection (I71)	80	0.5	22.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	72	0.4	20.0
15	Atherosclerosis (I70)	58	0.4	16.1
...	All other causes (Residual)	2,066	12.6	573.2
Black, non-Hispanic, female, 80-84 years				
...	All causes	17,057	100.0	6,735.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,549	32.5	2,191.3
2	Malignant neoplasms (C00-C97)	3,160	18.5	1,247.9
3	Cerebrovascular diseases (I60-I69)	1,669	9.8	659.1
4	Diabetes mellitus (E10-E14)	1,001	5.9	395.3
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	572	3.4	225.9
6	Alzheimer's disease (G30)	534	3.1	210.9
7	Septicemia (A40-A41)	443	2.6	174.9
8	Chronic lower respiratory diseases (J40-J47)	427	2.5	168.6
9	Influenza and pneumonia (J10-J18)	421	2.5	166.3
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	370	2.2	146.1
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	203	1.2	80.2
12	Pneumonitis due to solids and liquids (J69)	153	0.9	60.4
13	Atherosclerosis (I70)	98	0.6	38.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	85	0.5	33.6
15	Aortic aneurysm and dissection (I71)	74	0.4	29.2
...	All other causes (Residual)	2,298	13.5	907.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 85-89 years				
...	All causes	15,226	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,301	34.8	...
2	Malignant neoplasms (C00-C97)	2,104	13.8	...
3	Cerebrovascular diseases (I60-I69)	1,617	10.6	...
4	Diabetes mellitus (E10-E14)	739	4.9	...
5	Alzheimer's disease (G30)	667	4.4	...
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	517	3.4	...
7	Influenza and pneumonia (J10-J18)	467	3.1	...
8	Septicemia (A40-A41)	425	2.8	...
9	Chronic lower respiratory diseases (J40-J47)	324	2.1	...
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	310	2.0	...
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	155	1.0	...
12	Pneumonitis due to solids and liquids (J69)	150	1.0	...
13	Atherosclerosis (I70)	103	0.7	...
14	Aortic aneurysm and dissection (I71)	66	0.4	...
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	64	0.4	...
...	All other causes (Residual)	2,217	14.6	...
Black, non-Hispanic, female, 90-94 years				
...	All causes	11,170	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,116	36.8	...
2	Cerebrovascular diseases (I60-I69)	1,182	10.6	...
3	Malignant neoplasms (C00-C97)	1,058	9.5	...
4	Alzheimer's disease (G30)	547	4.9	...
5	Influenza and pneumonia (J10-J18)	421	3.8	...
6	Diabetes mellitus (E10-E14)	418	3.7	...
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	323	2.9	...
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	279	2.5	...
9	Septicemia (A40-A41)	272	2.4	...
10	Chronic lower respiratory diseases (J40-J47)	192	1.7	...
11	Pneumonitis due to solids and liquids (J69)	134	1.2	...
12	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	118	1.1	...
13	Atherosclerosis (I70)	115	1.0	...
14	Anemias (D50-D64)	56	0.5	...
15	Nutritional deficiencies (E40-E64)	50	0.4	...
...	All other causes (Residual)	1,889	16.9	...

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Black, non-Hispanic, female, 95-99 years			
...	All causes	5,025	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,969	39.2	...
2	Cerebrovascular diseases (I60-I69)	485	9.7	...
3	Malignant neoplasms (C00-C97)	344	6.8	...
4	Alzheimer's disease (G30)	255	5.1	...
5	Influenza and pneumonia (J10-J18)	202	4.0	...
6	Diabetes mellitus (E10-E14)	161	3.2	...
7	Septicemia (A40-A41)	134	2.7	...
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	132	2.6	...
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	122	2.4	...
10	Chronic lower respiratory diseases (J40-J47)	83	1.7	...
11	Atherosclerosis (I70)	66	1.3	...
12	Pneumonitis due to solids and liquids (J69)	51	1.0	...
13	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	42	0.8	...
14	Nutritional deficiencies (E40-E64)	30	0.6	...
15	Anemias (D50-D64)	21	0.4	...
...	All other causes (Residual)	928	18.5	...
	Black, non-Hispanic, female, 100 years and over			
...	All causes	1,849	100.0	...
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	751	40.6	...
2	Cerebrovascular diseases (I60-I69)	181	9.8	...
3	Malignant neoplasms (C00-C97)	88	4.8	...
4	Influenza and pneumonia (J10-J18)	83	4.5	...
5	Alzheimer's disease (G30)	81	4.4	...
6	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	49	2.7	...
7	Septicemia (A40-A41)	42	2.3	...
7	Diabetes mellitus (E10-E14)	42	2.3	...
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	35	1.9	...
10	Atherosclerosis (I70)	30	1.6	...
11	Chronic lower respiratory diseases (J40-J47)	23	1.2	...
12	Pneumonitis due to solids and liquids (J69)	20	1.1	...
13	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15	0.8	...
14	Nutritional deficiencies (E40-E64)	12	0.6	...
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.4	...
...	All other causes (Residual)	390	21.1	...

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available