

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, all ages				
...	All causes	131,161	100.0	307.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	29,555	22.5	69.2
2	Malignant neoplasms (C00-C97)	26,156	19.9	61.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,464	8.7	26.9
4	Cerebrovascular diseases (I60-I69)	6,830	5.2	16.0
5	Diabetes mellitus (E10-E14)	6,665	5.1	15.6
6	Chronic liver disease and cirrhosis (K70,K73-K74)	3,555	2.7	8.3
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,520	2.7	8.2
8	Chronic lower respiratory diseases (J40-J47)	3,457	2.6	8.1
9	Influenza and pneumonia (J10-J18)	3,085	2.4	7.2
10	Certain conditions originating in the perinatal period (P00-P96)	2,816	2.1	6.6
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,329	1.8	5.5
12	Alzheimer's disease (G30)	2,201	1.7	5.2
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,188	1.7	5.1
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,896	1.4	4.4
15	Septicemia (A40-A41)	1,704	1.3	4.0
...	All other causes (Residual)	23,740	18.1	55.6
Hispanic, both sexes, 1-4 years				
...	All causes	1,041	100.0	28.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	380	36.5	10.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	119	11.4	3.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	82	7.9	2.3
4	Malignant neoplasms (C00-C97)	80	7.7	2.2
5	Influenza and pneumonia (J10-J18)	32	3.1	0.9
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	23	2.2	0.6
7	Cerebrovascular diseases (I60-I69)	15	1.4	*
8	Septicemia (A40-A41)	14	1.3	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	1.2	*
10	Meningitis (G00,G03)	8	0.8	*
11	Certain conditions originating in the perinatal period (P00-P96)	6	0.6	*
12	Chronic lower respiratory diseases (J40-J47)	5	0.5	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.5	*
14	Anemias (D50-D64)	4	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.4	*
...	All other causes (Residual)	252	24.2	7.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 5-9 years				
...	All causes	474	100.0	12.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	182	38.4	4.6
2	Malignant neoplasms (C00-C97)	94	19.8	2.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	43	9.1	1.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	19	4.0	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	11	2.3	*
5	Cerebrovascular diseases (I60-I69)	11	2.3	*
7	Septicemia (A40-A41)	6	1.3	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.3	*
9	Chronic lower respiratory diseases (J40-J47)	5	1.1	*
10	Influenza and pneumonia (J10-J18)	4	0.8	*
11	Pneumonitis due to solids and liquids (J69)	2	0.4	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
11	Certain conditions originating in the perinatal period (P00-P96)	2	0.4	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
15	Shigellosis and amebiasis (A03,A06)	1	0.2	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.2	*
15	Diabetes mellitus (E10-E14)	1	0.2	*
15	Meningitis (G00,G03)	1	0.2	*
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1	0.2	*
15	Diseases of appendix (K35-K38)	1	0.2	*
...	All other causes (Residual)	79	16.7	2.0
Hispanic, both sexes, 10-14 years				
...	All causes	594	100.0	15.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	188	31.6	4.9
2	Malignant neoplasms (C00-C97)	100	16.8	2.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	43	7.2	1.1
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	42	7.1	1.1
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	39	6.6	1.0
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	23	3.9	0.6
7	Chronic lower respiratory diseases (J40-J47)	10	1.7	*
8	Cerebrovascular diseases (I60-I69)	8	1.3	*
9	Septicemia (A40-A41)	6	1.0	*
10	Influenza and pneumonia (J10-J18)	5	0.8	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.7	*
11	Anemias (D50-D64)	4	0.7	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	3	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.5	*
15	Diabetes mellitus (E10-E14)	2	0.3	*
...	All other causes (Residual)	114	19.2	3.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 15-19 years				
...	All causes	2,323	100.0	67.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,053	45.3	30.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	544	23.4	15.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	210	9.0	6.1
4	Malignant neoplasms (C00-C97)	140	6.0	4.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	46	2.0	1.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	36	1.5	1.0
7	Cerebrovascular diseases (I60-I69)	16	0.7	*
8	Pregnancy, childbirth and the puerperium (O00-099)	12	0.5	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.3	*
10	Chronic lower respiratory diseases (J40-J47)	6	0.3	*
11	Septicemia (A40-A41)	5	0.2	*
11	Aortic aneurysm and dissection (I71)	5	0.2	*
11	Legal intervention (Y35,Y89.0)	5	0.2	*
14	Tuberculosis (A16-A19)	4	0.2	*
14	Diabetes mellitus (E10-E14)	4	0.2	*
14	Influenza and pneumonia (J10-J18)	4	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.2	*
...	All other causes (Residual)	221	9.5	6.4
Hispanic, both sexes, 20-24 years				
...	All causes	3,514	100.0	94.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,617	46.0	43.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	763	21.7	20.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	343	9.8	9.2
4	Malignant neoplasms (C00-C97)	183	5.2	4.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	91	2.6	2.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	38	1.1	1.0
7	Human immunodeficiency virus (HIV) disease (B20-B24)	28	0.8	0.8
7	Cerebrovascular diseases (I60-I69)	28	0.8	0.8
9	Legal intervention (Y35,Y89.0)	27	0.8	0.7
10	Pregnancy, childbirth and the puerperium (O00-099)	23	0.7	0.6
11	Septicemia (A40-A41)	13	0.4	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	0.3	*
13	Diabetes mellitus (E10-E14)	10	0.3	*
14	Influenza and pneumonia (J10-J18)	7	0.2	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.1	*
15	Anemias (D50-D64)	5	0.1	*
...	All other causes (Residual)	322	9.2	8.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 25-29 years				
...	All causes	3,257	100.0	80.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,252	38.4	31.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	635	19.5	15.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	284	8.7	7.0
4	Malignant neoplasms (C00-C97)	220	6.8	5.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	163	5.0	4.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	62	1.9	1.5
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	35	1.1	0.9
8	Diabetes mellitus (E10-E14)	29	0.9	0.7
9	Cerebrovascular diseases (I60-I69)	27	0.8	0.7
10	Pregnancy, childbirth and the puerperium (O00-099)	23	0.7	0.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	22	0.7	0.5
12	Influenza and pneumonia (J10-J18)	19	0.6	*
13	Septicemia (A40-A41)	14	0.4	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	0.4	*
15	Legal intervention (Y35,Y89.0)	11	0.3	*
...	All other causes (Residual)	449	13.8	11.1
Hispanic, both sexes, 30-34 years				
...	All causes	3,154	100.0	83.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,050	33.3	27.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	450	14.3	11.8
3	Malignant neoplasms (C00-C97)	310	9.8	8.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	250	7.9	6.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	191	6.1	5.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	130	4.1	3.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	56	1.8	1.5
8	Cerebrovascular diseases (I60-I69)	55	1.7	1.4
9	Diabetes mellitus (E10-E14)	40	1.3	1.1
10	Pregnancy, childbirth and the puerperium (O00-099)	37	1.2	1.0
11	Septicemia (A40-A41)	22	0.7	0.6
12	Influenza and pneumonia (J10-J18)	21	0.7	0.6
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21	0.7	0.6
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	17	0.5	*
15	Chronic lower respiratory diseases (J40-J47)	15	0.5	*
...	All other causes (Residual)	489	15.5	12.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 35-39 years				
...	All causes	3,722	100.0	109.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	961	25.8	28.4
2	Malignant neoplasms (C00-C97)	498	13.4	14.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	330	8.9	9.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	309	8.3	9.1
5	Human immunodeficiency virus (HIV) disease (B20-B24)	244	6.6	7.2
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	244	6.6	7.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	149	4.0	4.4
8	Cerebrovascular diseases (I60-I69)	112	3.0	3.3
9	Diabetes mellitus (E10-E14)	67	1.8	2.0
10	Influenza and pneumonia (J10-J18)	37	1.0	1.1
11	Viral hepatitis (B15-B19)	34	0.9	1.0
12	Septicemia (A40-A41)	28	0.8	0.8
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	26	0.7	0.8
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	23	0.6	0.7
15	Pregnancy, childbirth and the puerperium (O00-099)	22	0.6	0.6
...	All other causes (Residual)	638	17.1	18.8
Hispanic, both sexes, 40-44 years				
...	All causes	5,154	100.0	171.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	959	18.6	31.9
2	Malignant neoplasms (C00-C97)	917	17.8	30.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	627	12.2	20.9
4	Human immunodeficiency virus (HIV) disease (B20-B24)	372	7.2	12.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	318	6.2	10.6
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	228	4.4	7.6
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	211	4.1	7.0
8	Cerebrovascular diseases (I60-I69)	184	3.6	6.1
9	Diabetes mellitus (E10-E14)	149	2.9	5.0
10	Viral hepatitis (B15-B19)	80	1.6	2.7
11	Influenza and pneumonia (J10-J18)	56	1.1	1.9
12	Septicemia (A40-A41)	46	0.9	1.5
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	42	0.8	1.4
14	Chronic lower respiratory diseases (J40-J47)	36	0.7	1.2
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	25	0.5	0.8
...	All other causes (Residual)	904	17.5	30.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 45-49 years				
...	All causes	6,206	100.0	256.1
1	Malignant neoplasms (C00-C97)	1,365	22.0	56.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	902	14.5	37.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	818	13.2	33.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	473	7.6	19.5
5	Human immunodeficiency virus (HIV) disease (B20-B24)	335	5.4	13.8
6	Cerebrovascular diseases (I60-I69)	303	4.9	12.5
7	Diabetes mellitus (E10-E14)	262	4.2	10.8
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	175	2.8	7.2
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	152	2.4	6.3
10	Viral hepatitis (B15-B19)	116	1.9	4.8
11	Influenza and pneumonia (J10-J18)	82	1.3	3.4
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	73	1.2	3.0
13	Septicemia (A40-A41)	68	1.1	2.8
14	Chronic lower respiratory diseases (J40-J47)	58	0.9	2.4
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	32	0.5	1.3
...	All other causes (Residual)	992	16.0	40.9
Hispanic, both sexes, 50-54 years				
...	All causes	7,363	100.0	398.8
1	Malignant neoplasms (C00-C97)	1,959	26.6	106.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,331	18.1	72.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	637	8.7	34.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	530	7.2	28.7
5	Diabetes mellitus (E10-E14)	370	5.0	20.0
6	Cerebrovascular diseases (I60-I69)	336	4.6	18.2
7	Human immunodeficiency virus (HIV) disease (B20-B24)	222	3.0	12.0
8	Viral hepatitis (B15-B19)	177	2.4	9.6
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	112	1.5	6.1
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	108	1.5	5.8
11	Chronic lower respiratory diseases (J40-J47)	93	1.3	5.0
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	92	1.2	5.0
13	Influenza and pneumonia (J10-J18)	90	1.2	4.9
14	Septicemia (A40-A41)	89	1.2	4.8
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	50	0.7	2.7
...	All other causes (Residual)	1,167	15.8	63.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 55-59 years				
...	All causes	7,709	100.0	549.3
1	Malignant neoplasms (C00-C97)	2,249	29.2	160.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,656	21.5	118.0
3	Diabetes mellitus (E10-E14)	514	6.7	36.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	511	6.6	36.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	414	5.4	29.5
6	Cerebrovascular diseases (I60-I69)	371	4.8	26.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	151	2.0	10.8
8	Viral hepatitis (B15-B19)	130	1.7	9.3
9	Human immunodeficiency virus (HIV) disease (B20-B24)	120	1.6	8.6
9	Chronic lower respiratory diseases (J40-J47)	120	1.6	8.6
11	Influenza and pneumonia (J10-J18)	114	1.5	8.1
12	Septicemia (A40-A41)	108	1.4	7.7
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	85	1.1	6.1
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	64	0.8	4.6
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	45	0.6	3.2
...	All other causes (Residual)	1,057	13.7	75.3
Hispanic, both sexes, 60-64 years				
...	All causes	8,380	100.0	859.4
1	Malignant neoplasms (C00-C97)	2,587	30.9	265.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,986	23.7	203.7
3	Diabetes mellitus (E10-E14)	654	7.8	67.1
4	Cerebrovascular diseases (I60-I69)	424	5.1	43.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	366	4.4	37.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	306	3.7	31.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	200	2.4	20.5
8	Chronic lower respiratory diseases (J40-J47)	178	2.1	18.3
9	Influenza and pneumonia (J10-J18)	133	1.6	13.6
10	Septicemia (A40-A41)	122	1.5	12.5
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	86	1.0	8.8
12	Viral hepatitis (B15-B19)	72	0.9	7.4
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	57	0.7	5.8
14	Human immunodeficiency virus (HIV) disease (B20-B24)	51	0.6	5.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	41	0.5	4.2
...	All other causes (Residual)	1,117	13.3	114.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 65-69 years				
...	All causes	9,513	100.0	1,276.5
1	Malignant neoplasms (C00-C97)	2,984	31.4	400.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,338	24.6	313.7
3	Diabetes mellitus (E10-E14)	755	7.9	101.3
4	Cerebrovascular diseases (I60-I69)	481	5.1	64.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	350	3.7	47.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	262	2.8	35.2
7	Chronic lower respiratory diseases (J40-J47)	248	2.6	33.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	219	2.3	29.4
9	Influenza and pneumonia (J10-J18)	182	1.9	24.4
10	Septicemia (A40-A41)	153	1.6	20.5
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	109	1.1	14.6
12	Viral hepatitis (B15-B19)	60	0.6	8.1
13	Alzheimer's disease (G30)	51	0.5	6.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	45	0.5	6.0
15	Aortic aneurysm and dissection (I71)	44	0.5	5.9
...	All other causes (Residual)	1,232	13.0	165.3
Hispanic, both sexes, 70-74 years				
...	All causes	11,909	100.0	2,067.5
1	Malignant neoplasms (C00-C97)	3,375	28.3	585.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,213	27.0	557.8
3	Diabetes mellitus (E10-E14)	915	7.7	158.9
4	Cerebrovascular diseases (I60-I69)	700	5.9	121.5
5	Chronic lower respiratory diseases (J40-J47)	420	3.5	72.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	295	2.5	51.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	287	2.4	49.8
8	Influenza and pneumonia (J10-J18)	273	2.3	47.4
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	258	2.2	44.8
10	Septicemia (A40-A41)	184	1.5	31.9
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	137	1.2	23.8
12	Alzheimer's disease (G30)	121	1.0	21.0
13	Parkinson's disease (G20-G21)	82	0.7	14.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	62	0.5	10.8
15	Viral hepatitis (B15-B19)	53	0.4	9.2
...	All other causes (Residual)	1,534	12.9	266.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 75-79 years				
...	All causes	14,156	100.0	3,196.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,996	28.2	902.2
2	Malignant neoplasms (C00-C97)	3,525	24.9	795.9
3	Diabetes mellitus (E10-E14)	1,064	7.5	240.2
4	Cerebrovascular diseases (I60-I69)	910	6.4	205.5
5	Chronic lower respiratory diseases (J40-J47)	605	4.3	136.6
6	Influenza and pneumonia (J10-J18)	394	2.8	89.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	364	2.6	82.2
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	281	2.0	63.4
9	Alzheimer's disease (G30)	278	2.0	62.8
10	Chronic liver disease and cirrhosis (K70,K73-K74)	257	1.8	58.0
11	Septicemia (A40-A41)	206	1.5	46.5
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	173	1.2	39.1
13	Parkinson's disease (G20-G21)	162	1.1	36.6
14	Aortic aneurysm and dissection (I71)	78	0.6	17.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	76	0.5	17.2
...	All other causes (Residual)	1,787	12.6	403.5
Hispanic, both sexes, 80-84 years				
...	All causes	14,854	100.0	5,186.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,587	30.9	1,601.6
2	Malignant neoplasms (C00-C97)	2,913	19.6	1,017.1
3	Cerebrovascular diseases (I60-I69)	1,099	7.4	383.7
4	Diabetes mellitus (E10-E14)	883	5.9	308.3
5	Chronic lower respiratory diseases (J40-J47)	649	4.4	226.6
6	Alzheimer's disease (G30)	523	3.5	182.6
7	Influenza and pneumonia (J10-J18)	498	3.4	173.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	335	2.3	117.0
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	300	2.0	104.7
10	Septicemia (A40-A41)	239	1.6	83.4
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	213	1.4	74.4
12	Parkinson's disease (G20-G21)	192	1.3	67.0
13	Chronic liver disease and cirrhosis (K70,K73-K74)	137	0.9	47.8
14	Pneumonitis due to solids and liquids (J69)	113	0.8	39.5
15	Atherosclerosis (I70)	78	0.5	27.2
...	All other causes (Residual)	2,095	14.1	731.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 85-89 years				
...	All causes	11,101	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,756	33.8	---
2	Malignant neoplasms (C00-C97)	1,640	14.8	---
3	Cerebrovascular diseases (I60-I69)	918	8.3	---
4	Diabetes mellitus (E10-E14)	569	5.1	---
5	Alzheimer's disease (G30)	549	4.9	---
6	Chronic lower respiratory diseases (J40-J47)	522	4.7	---
7	Influenza and pneumonia (J10-J18)	467	4.2	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	237	2.1	---
9	Septicemia (A40-A41)	180	1.6	---
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	170	1.5	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	162	1.5	---
12	Parkinson's disease (G20-G21)	152	1.4	---
13	Pneumonitis due to solids and liquids (J69)	121	1.1	---
14	Atherosclerosis (I70)	74	0.7	---
15	Chronic liver disease and cirrhosis (K70,K73-K74)	59	0.5	---
...	All other causes (Residual)	1,525	13.7	---
Hispanic, both sexes, 90-94 years				
...	All causes	7,355	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,786	37.9	---
2	Malignant neoplasms (C00-C97)	760	10.3	---
3	Cerebrovascular diseases (I60-I69)	566	7.7	---
4	Alzheimer's disease (G30)	394	5.4	---
5	Influenza and pneumonia (J10-J18)	389	5.3	---
6	Chronic lower respiratory diseases (J40-J47)	294	4.0	---
7	Diabetes mellitus (E10-E14)	278	3.8	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	151	2.1	---
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	140	1.9	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	139	1.9	---
11	Septicemia (A40-A41)	87	1.2	---
12	Pneumonitis due to solids and liquids (J69)	81	1.1	---
13	Atherosclerosis (I70)	60	0.8	---
14	Parkinson's disease (G20-G21)	50	0.7	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	35	0.5	---
...	All other causes (Residual)	1,145	15.6	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Hispanic, both sexes, 95-99 years			---
...	All causes	2,890	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,111	38.4	---
2	Malignant neoplasms (C00-C97)	216	7.5	---
3	Alzheimer's disease (G30)	212	7.3	---
4	Cerebrovascular diseases (I60-I69)	201	7.0	---
5	Influenza and pneumonia (J10-J18)	162	5.6	---
6	Chronic lower respiratory diseases (J40-J47)	132	4.6	---
7	Diabetes mellitus (E10-E14)	84	2.9	---
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	65	2.2	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	56	1.9	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	48	1.7	---
11	Pneumonitis due to solids and liquids (J69)	45	1.6	---
12	Septicemia (A40-A41)	40	1.4	---
13	Atherosclerosis (I70)	36	1.2	---
14	Parkinson's disease (G20-G21)	19	0.7	---
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	11	0.4	---
...	All other causes (Residual)	452	15.6	---
	Hispanic, both sexes, 100 years and over			
...	All causes	712	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	312	43.8	---
2	Influenza and pneumonia (J10-J18)	50	7.0	---
3	Cerebrovascular diseases (I60-I69)	41	5.8	---
4	Alzheimer's disease (G30)	32	4.5	---
5	Malignant neoplasms (C00-C97)	21	2.9	---
5	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	21	2.9	---
5	Chronic lower respiratory diseases (J40-J47)	21	2.9	---
8	Atherosclerosis (I70)	16	2.2	---
9	Diabetes mellitus (E10-E14)	15	2.1	---
10	Pneumonitis due to solids and liquids (J69)	10	1.4	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10	1.4	---
12	Septicemia (A40-A41)	8	1.1	---
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	1.1	---
14	Nutritional deficiencies (E40-E64)	6	0.8	---
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	4	0.6	---
...	All other causes (Residual)	137	19.2	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, all ages				
...	All causes	73,788	100.0	334.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,900	21.5	72.1
2	Malignant neoplasms (C00-C97)	13,896	18.8	63.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,612	11.7	39.0
4	Diabetes mellitus (E10-E14)	3,296	4.5	14.9
5	Cerebrovascular diseases (I60-I69)	3,188	4.3	14.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,008	4.1	13.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,561	3.5	11.6
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,841	2.5	8.3
9	Chronic lower respiratory diseases (J40-J47)	1,839	2.5	8.3
10	Certain conditions originating in the perinatal period (P00-P96)	1,612	2.2	7.3
11	Influenza and pneumonia (J10-J18)	1,506	2.0	6.8
12	Human immunodeficiency virus (HIV) disease (B20-B24)	1,309	1.8	5.9
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,186	1.6	5.4
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	996	1.3	4.5
	Septicemia (A40-A41)	838	1.1	3.8
...	All other causes (Residual)	12,200	16.5	55.3
Hispanic, male, 1-4 years				
...	All causes	610	100.0	33.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	244	40.0	13.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	65	10.7	3.5
3	Malignant neoplasms (C00-C97)	52	8.5	2.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	46	7.5	2.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	17	2.8	*
6	Influenza and pneumonia (J10-J18)	15	2.5	*
7	Cerebrovascular diseases (I60-I69)	8	1.3	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	1.1	*
9	Septicemia (A40-A41)	6	1.0	*
10	Meningococcal infection (A39)	3	0.5	*
10	Chronic lower respiratory diseases (J40-J47)	3	0.5	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.5	*
13	Tuberculosis (A16-A19)	2	0.3	*
13	Meningitis (G00,G03)	2	0.3	*
13	Acute bronchitis and bronchiolitis (J20-J21)	2	0.3	*
13	Certain conditions originating in the perinatal period (P00-P96)	2	0.3	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.3	*
...	All other causes (Residual)	131	21.5	7.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 5-9 years				
...	All causes	258	100.0	12.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	105	40.7	5.2
2	Malignant neoplasms (C00-C97)	47	18.2	2.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	24	9.3	1.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	9	3.5	*
5	Cerebrovascular diseases (I60-I69)	6	2.3	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	1.9	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	1.6	*
8	Septicemia (A40-A41)	3	1.2	*
9	Influenza and pneumonia (J10-J18)	2	0.8	*
9	Chronic lower respiratory diseases (J40-J47)	2	0.8	*
11	Diabetes mellitus (E10-E14)	1	0.4	*
11	Pneumonitis due to solids and liquids (J69)	1	0.4	*
11	Diseases of appendix (K35-K38)	1	0.4	*
11	Certain conditions originating in the perinatal period (P00-P96)	1	0.4	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.4	*
...	All other causes (Residual)	46	17.8	2.3
Hispanic, male, 10-14 years				
...	All causes	354	100.0	17.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	117	33.1	5.9
2	Malignant neoplasms (C00-C97)	52	14.7	2.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	31	8.8	1.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	27	7.6	1.4
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	23	6.5	1.2
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	16	4.5	*
7	Chronic lower respiratory diseases (J40-J47)	7	2.0	*
8	Influenza and pneumonia (J10-J18)	5	1.4	*
9	Anemias (D50-D64)	4	1.1	*
10	Septicemia (A40-A41)	3	0.8	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.8	*
10	Cerebrovascular diseases (I60-I69)	3	0.8	*
13	Whooping cough (A37)	1	0.3	*
13	Meningococcal infection (A39)	1	0.3	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.3	*
13	Diabetes mellitus (E10-E14)	1	0.3	*
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1	0.3	*
13	Pneumonitis due to solids and liquids (J69)	1	0.3	*
13	Diseases of appendix (K35-K38)	1	0.3	*
13	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.3	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
13	Certain conditions originating in the perinatal period (P00-P96)	1	0.3	*
...	All other causes (Residual)	53	15.0	2.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 15-19 years				
...	All causes	1,759	100.0	98.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	775	44.1	43.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	496	28.2	27.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	170	9.7	9.5
4	Malignant neoplasms (C00-C97)	89	5.1	5.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	35	2.0	2.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	21	1.2	1.2
7	Cerebrovascular diseases (I60-I69)	7	0.4	*
8	Chronic lower respiratory diseases (J40-J47)	5	0.3	*
8	Legal intervention (Y35,Y89.0)	5	0.3	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.2	*
10	Influenza and pneumonia (J10-J18)	4	0.2	*
12	Septicemia (A40-A41)	3	0.2	*
12	Diabetes mellitus (E10-E14)	3	0.2	*
12	Aortic aneurysm and dissection (I71)	3	0.2	*
12	Certain conditions originating in the perinatal period (P00-P96)	3	0.2	*
...	All other causes (Residual)	136	7.7	7.6
Hispanic, male, 20-24 years				
...	All causes	2,845	100.0	139.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,358	47.7	66.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	691	24.3	33.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	292	10.3	14.3
4	Malignant neoplasms (C00-C97)	113	4.0	5.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	56	2.0	2.7
6	Legal intervention (Y35,Y89.0)	27	0.9	1.3
7	Human immunodeficiency virus (HIV) disease (B20-B24)	22	0.8	1.1
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	22	0.8	1.1
9	Cerebrovascular diseases (I60-I69)	18	0.6	*
10	Septicemia (A40-A41)	7	0.2	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.2	*
12	Diabetes mellitus (E10-E14)	6	0.2	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.1	*
13	Influenza and pneumonia (J10-J18)	4	0.1	*
15	Anemias (D50-D64)	3	0.1	*
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	3	0.1	*
15	Aortic aneurysm and dissection (I71)	3	0.1	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	3	0.1	*
...	All other causes (Residual)	206	7.2	10.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 25-29 years				
...	All causes	2,597	100.0	115.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,061	40.9	47.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	568	21.9	25.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	254	9.8	11.3
4	Malignant neoplasms (C00-C97)	126	4.9	5.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	124	4.8	5.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	46	1.8	2.1
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	20	0.8	0.9
8	Diabetes mellitus (E10-E14)	17	0.7	*
8	Cerebrovascular diseases (I60-I69)	17	0.7	*
10	Chronic liver disease and cirrhosis (K70,K73-K74)	16	0.6	*
11	Septicemia (A40-A41)	11	0.4	*
11	Influenza and pneumonia (J10-J18)	11	0.4	*
11	Legal intervention (Y35,Y89.0)	11	0.4	*
14	Chronic lower respiratory diseases (J40-J47)	10	0.4	*
15	Viral hepatitis (B15-B19)	6	0.2	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.2	*
...	All other causes (Residual)	293	11.3	13.1
Hispanic, male, 30-34 years				
...	All causes	2,363	100.0	115.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	869	36.8	42.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	388	16.4	18.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	207	8.8	10.1
4	Malignant neoplasms (C00-C97)	154	6.5	7.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	148	6.3	7.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	96	4.1	4.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	46	1.9	2.2
8	Cerebrovascular diseases (I60-I69)	33	1.4	1.6
9	Diabetes mellitus (E10-E14)	23	1.0	1.1
10	Influenza and pneumonia (J10-J18)	15	0.6	*
11	Legal intervention (Y35,Y89.0)	13	0.6	*
12	Septicemia (A40-A41)	12	0.5	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	0.5	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	10	0.4	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.3	*
...	All other causes (Residual)	329	13.9	16.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 35-39 years				
...	All causes	2,624	100.0	145.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	766	29.2	42.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	254	9.7	14.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	251	9.6	13.9
4	Malignant neoplasms (C00-C97)	204	7.8	11.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	204	7.8	11.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	198	7.5	10.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	120	4.6	6.6
8	Cerebrovascular diseases (I60-I69)	71	2.7	3.9
9	Diabetes mellitus (E10-E14)	50	1.9	2.8
10	Viral hepatitis (B15-B19)	27	1.0	1.5
11	Influenza and pneumonia (J10-J18)	26	1.0	1.4
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	0.6	*
12	Legal intervention (Y35,Y89.0)	15	0.6	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	12	0.5	*
15	Septicemia (A40-A41)	11	0.4	*
15	Chronic lower respiratory diseases (J40-J47)	11	0.4	*
...	All other causes (Residual)	389	14.8	21.5
Hispanic, male, 40-44 years				
...	All causes	3,521	100.0	224.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	747	21.2	47.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	479	13.6	30.6
3	Malignant neoplasms (C00-C97)	397	11.3	25.3
4	Human immunodeficiency virus (HIV) disease (B20-B24)	298	8.5	19.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	249	7.1	15.9
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	195	5.5	12.4
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	177	5.0	11.3
8	Cerebrovascular diseases (I60-I69)	106	3.0	6.8
9	Diabetes mellitus (E10-E14)	103	2.9	6.6
10	Viral hepatitis (B15-B19)	58	1.6	3.7
11	Influenza and pneumonia (J10-J18)	33	0.9	2.1
12	Septicemia (A40-A41)	24	0.7	1.5
13	Chronic lower respiratory diseases (J40-J47)	23	0.7	1.5
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22	0.6	1.4
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	20	0.6	1.3
15	Aortic aneurysm and dissection (I71)	20	0.6	1.3
...	All other causes (Residual)	570	16.2	36.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 45-49 years				
...	All causes	4,199	100.0	339.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	674	16.1	54.5
2	Malignant neoplasms (C00-C97)	651	15.5	52.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	647	15.4	52.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	392	9.3	31.7
5	Human immunodeficiency virus (HIV) disease (B20-B24)	278	6.6	22.5
6	Cerebrovascular diseases (I60-I69)	177	4.2	14.3
7	Diabetes mellitus (E10-E14)	163	3.9	13.2
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	147	3.5	11.9
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	121	2.9	9.8
10	Viral hepatitis (B15-B19)	93	2.2	7.5
11	Influenza and pneumonia (J10-J18)	52	1.2	4.2
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	51	1.2	4.1
13	Septicemia (A40-A41)	41	1.0	3.3
14	Chronic lower respiratory diseases (J40-J47)	24	0.6	1.9
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	20	0.5	1.6
...	All other causes (Residual)	668	15.9	54.0
Hispanic, male, 50-54 years				
...	All causes	4,795	100.0	522.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,005	21.0	109.4
2	Malignant neoplasms (C00-C97)	984	20.5	107.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	494	10.3	53.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	422	8.8	46.0
5	Diabetes mellitus (E10-E14)	215	4.5	23.4
6	Cerebrovascular diseases (I60-I69)	207	4.3	22.5
7	Human immunodeficiency virus (HIV) disease (B20-B24)	170	3.5	18.5
8	Viral hepatitis (B15-B19)	129	2.7	14.0
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	97	2.0	10.6
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	90	1.9	9.8
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	61	1.3	6.6
12	Influenza and pneumonia (J10-J18)	58	1.2	6.3
13	Chronic lower respiratory diseases (J40-J47)	53	1.1	5.8
14	Septicemia (A40-A41)	52	1.1	5.7
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	26	0.5	2.8
...	All other causes (Residual)	732	15.3	79.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 55-59 years				
...	All causes	4,842	100.0	712.8
1	Malignant neoplasms (C00-C97)	1,240	25.6	182.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,167	24.1	171.8
3	Chronic liver disease and cirrhosis (K70,K73-K74)	384	7.9	56.5
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	309	6.4	45.5
5	Diabetes mellitus (E10-E14)	291	6.0	42.8
6	Cerebrovascular diseases (I60-I69)	210	4.3	30.9
7	Human immunodeficiency virus (HIV) disease (B20-B24)	97	2.0	14.3
8	Viral hepatitis (B15-B19)	87	1.8	12.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	87	1.8	12.8
10	Influenza and pneumonia (J10-J18)	77	1.6	11.3
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	70	1.4	10.3
12	Chronic lower respiratory diseases (J40-J47)	64	1.3	9.4
13	Septicemia (A40-A41)	63	1.3	9.3
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	32	0.7	4.7
		29	0.6	4.3
...	All other causes (Residual)	635	13.1	93.5
Hispanic, male, 60-64 years				
...	All causes	5,112	100.0	1,118.0
1	Malignant neoplasms (C00-C97)	1,453	28.4	317.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,325	25.9	289.8
3	Diabetes mellitus (E10-E14)	371	7.3	81.1
4	Chronic liver disease and cirrhosis (K70,K73-K74)	280	5.5	61.2
5	Cerebrovascular diseases (I60-I69)	248	4.9	54.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	221	4.3	48.3
7	Chronic lower respiratory diseases (J40-J47)	116	2.3	25.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	114	2.2	24.9
9	Influenza and pneumonia (J10-J18)	70	1.4	15.3
10	Septicemia (A40-A41)	58	1.1	12.7
11	Human immunodeficiency virus (HIV) disease (B20-B24)	46	0.9	10.1
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	46	0.9	10.1
13	Viral hepatitis (B15-B19)	43	0.8	9.4
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	40	0.8	8.7
15	Aortic aneurysm and dissection (I71)	30	0.6	6.6
...	All other causes (Residual)	651	12.7	142.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 65-69 years				
...	All causes	5,410	100.0	1,596.8
1	Malignant neoplasms (C00-C97)	1,677	31.0	495.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,449	26.8	427.7
3	Diabetes mellitus (E10-E14)	387	7.2	114.2
4	Cerebrovascular diseases (I60-I69)	254	4.7	75.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	224	4.1	66.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	178	3.3	52.5
7	Chronic lower respiratory diseases (J40-J47)	143	2.6	42.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	100	1.8	29.5
9	Influenza and pneumonia (J10-J18)	97	1.8	28.6
10	Septicemia (A40-A41)	75	1.4	22.1
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	62	1.1	18.3
12	Viral hepatitis (B15-B19)	34	0.6	10.0
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	30	0.6	8.9
13	Aortic aneurysm and dissection (I71)	30	0.6	8.9
15	Human immunodeficiency virus (HIV) disease (B20-B24)	28	0.5	8.3
...	All other causes (Residual)	642	11.9	189.5
Hispanic, male, 70-74 years				
...	All causes	6,574	100.0	2,611.5
1	Malignant neoplasms (C00-C97)	1,887	28.7	749.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,867	28.4	741.7
3	Diabetes mellitus (E10-E14)	463	7.0	183.9
4	Cerebrovascular diseases (I60-I69)	373	5.7	148.2
5	Chronic lower respiratory diseases (J40-J47)	238	3.6	94.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	184	2.8	73.1
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	167	2.5	66.3
8	Influenza and pneumonia (J10-J18)	153	2.3	60.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	144	2.2	57.2
10	Septicemia (A40-A41)	100	1.5	39.7
11	Alzheimer's disease (G30)	62	0.9	24.6
12	Parkinson's disease (G20-G21)	59	0.9	23.4
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	59	0.9	23.4
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	38	0.6	15.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	33	0.5	13.1
15	Aortic aneurysm and dissection (I71)	33	0.5	13.1
...	All other causes (Residual)	714	10.9	283.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 75-79 years				
...	All causes	7,383	100.0	3,987.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,183	29.6	1,179.1
2	Malignant neoplasms (C00-C97)	1,905	25.8	1,028.9
3	Diabetes mellitus (E10-E14)	493	6.7	266.3
4	Cerebrovascular diseases (I60-I69)	423	5.7	228.5
5	Chronic lower respiratory diseases (J40-J47)	341	4.6	184.2
6	Influenza and pneumonia (J10-J18)	201	2.7	108.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	183	2.5	98.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	161	2.2	87.0
9	Chronic liver disease and cirrhosis (K70,K73-K74)	129	1.7	69.7
10	Alzheimer's disease (G30)	128	1.7	69.1
11	Septicemia (A40-A41)	104	1.4	56.2
12	Parkinson's disease (G20-G21)	98	1.3	52.9
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	71	1.0	38.3
14	Aortic aneurysm and dissection (I71)	57	0.8	30.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	38	0.5	20.5
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	38	0.5	20.5
...	All other causes (Residual)	830	11.2	448.3
Hispanic, male, 80-84 years				
...	All causes	7,156	100.0	6,265.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,215	31.0	1,939.4
2	Malignant neoplasms (C00-C97)	1,573	22.0	1,377.3
3	Cerebrovascular diseases (I60-I69)	457	6.4	400.1
4	Diabetes mellitus (E10-E14)	384	5.4	336.2
5	Chronic lower respiratory diseases (J40-J47)	360	5.0	315.2
6	Influenza and pneumonia (J10-J18)	247	3.5	216.3
7	Alzheimer's disease (G30)	209	2.9	183.0
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	157	2.2	137.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	155	2.2	135.7
10	Septicemia (A40-A41)	118	1.6	103.3
11	Parkinson's disease (G20-G21)	115	1.6	100.7
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	91	1.3	79.7
13	Chronic liver disease and cirrhosis (K70,K73-K74)	77	1.1	67.4
14	Pneumonitis due to solids and liquids (J69)	63	0.9	55.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	44	0.6	38.5
...	All other causes (Residual)	891	12.5	780.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 85-89 years				
...	All causes	4,618	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,545	33.5	---
2	Malignant neoplasms (C00-C97)	821	17.8	---
3	Cerebrovascular diseases (I60-I69)	334	7.2	---
4	Chronic lower respiratory diseases (J40-J47)	248	5.4	---
5	Diabetes mellitus (E10-E14)	221	4.8	---
6	Influenza and pneumonia (J10-J18)	191	4.1	---
7	Alzheimer's disease (G30)	186	4.0	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	117	2.5	---
9	Parkinson's disease (G20-G21)	87	1.9	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	69	1.5	---
11	Septicemia (A40-A41)	66	1.4	---
12	Pneumonitis due to solids and liquids (J69)	58	1.3	---
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	55	1.2	---
14	Atherosclerosis (I70)	36	0.8	---
15	Chronic liver disease and cirrhosis (K70,K73-K74)	23	0.5	---
...	All other causes (Residual)	561	12.1	---
Hispanic, male, 90-94 years				
...	All causes	2,505	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	927	37.0	---
2	Malignant neoplasms (C00-C97)	360	14.4	---
3	Cerebrovascular diseases (I60-I69)	163	6.5	---
4	Influenza and pneumonia (J10-J18)	142	5.7	---
5	Chronic lower respiratory diseases (J40-J47)	116	4.6	---
6	Alzheimer's disease (G30)	90	3.6	---
7	Diabetes mellitus (E10-E14)	82	3.3	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	64	2.6	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	48	1.9	---
10	Pneumonitis due to solids and liquids (J69)	41	1.6	---
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	39	1.6	---
12	Septicemia (A40-A41)	31	1.2	---
13	Atherosclerosis (I70)	20	0.8	---
14	Parkinson's disease (G20-G21)	19	0.8	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	0.5	---
...	All other causes (Residual)	351	14.0	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 95-99 years				
...	All causes	853	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	297	34.8	---
2	Malignant neoplasms (C00-C97)	97	11.4	---
3	Influenza and pneumonia (J10-J18)	55	6.4	---
4	Chronic lower respiratory diseases (J40-J47)	53	6.2	---
5	Cerebrovascular diseases (I60-I69)	51	6.0	---
6	Alzheimer's disease (G30)	49	5.7	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22	2.6	---
8	Diabetes mellitus (E10-E14)	21	2.5	---
8	Pneumonitis due to solids and liquids (J69)	21	2.5	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	16	1.9	---
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	14	1.6	---
12	Septicemia (A40-A41)	11	1.3	---
13	Atherosclerosis (I70)	9	1.1	---
14	Nutritional deficiencies (E40-E64)	7	0.8	---
15	Parkinson's disease (G20-G21)	5	0.6	---
15	Aortic aneurysm and dissection (I71)	5	0.6	---
...	All other causes (Residual)	120	14.1	---
Hispanic, male, 100 years and over				
...	All causes	167	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	69	41.3	---
2	Influenza and pneumonia (J10-J18)	11	6.6	---
3	Malignant neoplasms (C00-C97)	8	4.8	---
3	Cerebrovascular diseases (I60-I69)	8	4.8	---
5	Alzheimer's disease (G30)	6	3.6	---
5	Chronic lower respiratory diseases (J40-J47)	6	3.6	---
7	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	5	3.0	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	3.0	---
9	Pneumonitis due to solids and liquids (J69)	4	2.4	---
10	Septicemia (A40-A41)	2	1.2	---
10	Atherosclerosis (I70)	2	1.2	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2	1.2	---
13	Diabetes mellitus (E10-E14)	1	0.6	---
13	Acute bronchitis and bronchiolitis (J20-J21)	1	0.6	---
13	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.6	---
13	Hyperplasia of prostate (N40)	1	0.6	---
...	All other causes (Residual)	35	21.0	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, all ages				
...	All causes	57,373	100.0	278.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,655	23.8	66.2
2	Malignant neoplasms (C00-C97)	12,260	21.4	59.5
3	Cerebrovascular diseases (I60-I69)	3,642	6.3	17.7
4	Diabetes mellitus (E10-E14)	3,369	5.9	16.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,852	5.0	13.8
6	Chronic lower respiratory diseases (J40-J47)	1,618	2.8	7.8
7	Influenza and pneumonia (J10-J18)	1,579	2.8	7.7
8	Alzheimer's disease (G30)	1,437	2.5	7.0
9	Certain conditions originating in the perinatal period (P00-P96)	1,204	2.1	5.8
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,143	2.0	5.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	994	1.7	4.8
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	900	1.6	4.4
13	Septicemia (A40-A41)	866	1.5	4.2
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	767	1.3	3.7
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	512	0.9	2.5
...	All other causes (Residual)	10,575	18.4	51.3
Hispanic, female, 1-4 years				
...	All causes	431	100.0	24.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	136	31.6	7.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	54	12.5	3.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	36	8.4	2.0
4	Malignant neoplasms (C00-C97)	28	6.5	1.6
5	Influenza and pneumonia (J10-J18)	17	3.9	*
6	Septicemia (A40-A41)	8	1.9	*
7	Cerebrovascular diseases (I60-I69)	7	1.6	*
8	Meningitis (G00,G03)	6	1.4	*
8	Diseases of heart (I00-I09,I11,I13,I20-I51)	6	1.4	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	1.2	*
11	Certain conditions originating in the perinatal period (P00-P96)	4	0.9	*
12	Anemias (D50-D64)	3	0.7	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.7	*
14	Chronic lower respiratory diseases (J40-J47)	2	0.5	*
15	Shigellosis and amebiasis (A03,A06)	1	0.2	*
15	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
15	Pneumoconioses and chemical effects (J60-J66,J68)	1	0.2	*
15	Pneumonitis due to solids and liquids (J69)	1	0.2	*
15	Diseases of appendix (K35-K38)	1	0.2	*
15	Hernia (K40-K46)	1	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
15	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.2	*
...	All other causes (Residual)	108	25.1	6.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 5-9 years				
...	All causes	216	100.0	11.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	77	35.6	4.0
2	Malignant neoplasms (C00-C97)	47	21.8	2.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	19	8.8	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	10	4.6	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	6	2.8	*
6	Cerebrovascular diseases (I60-I69)	5	2.3	*
7	Septicemia (A40-A41)	3	1.4	*
7	Chronic lower respiratory diseases (J40-J47)	3	1.4	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.9	*
9	Influenza and pneumonia (J10-J18)	2	0.9	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.9	*
12	Shigellosis and amebiasis (A03,A06)	1	0.5	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.5	*
12	Meningitis (G00,G03)	1	0.5	*
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1	0.5	*
12	Pneumonitis due to solids and liquids (J69)	1	0.5	*
12	Certain conditions originating in the perinatal period (P00-P96)	1	0.5	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.5	*
...	All other causes (Residual)	33	15.3	1.7
Hispanic, female, 10-14 years				
...	All causes	240	100.0	12.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	71	29.6	3.8
2	Malignant neoplasms (C00-C97)	48	20.0	2.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	19	7.9	*
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	12	5.0	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	12	5.0	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	7	2.9	*
7	Cerebrovascular diseases (I60-I69)	5	2.1	*
8	Septicemia (A40-A41)	3	1.3	*
8	Chronic lower respiratory diseases (J40-J47)	3	1.3	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.8	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.8	*
12	Tuberculosis (A16-A19)	1	0.4	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.4	*
12	Diabetes mellitus (E10-E14)	1	0.4	*
12	Meningitis (G00,G03)	1	0.4	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.4	*
...	All other causes (Residual)	51	21.3	2.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 15-19 years				
...	All causes	564	100.0	33.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	278	49.3	16.6
2	Malignant neoplasms (C00-C97)	51	9.0	3.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	48	8.5	2.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	40	7.1	2.4
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	15	2.7	*
6	Pregnancy, childbirth and the puerperium (O00-099)	12	2.1	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	11	2.0	*
8	Cerebrovascular diseases (I60-I69)	9	1.6	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.7	*
10	Tuberculosis (A16-A19)	3	0.5	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.5	*
12	Septicemia (A40-A41)	2	0.4	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.4	*
12	Aortic aneurysm and dissection (I71)	2	0.4	*
15	Diabetes mellitus (E10-E14)	1	0.2	*
15	Chronic lower respiratory diseases (J40-J47)	1	0.2	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
...	All other causes (Residual)	81	14.4	4.8
Hispanic, female, 20-24 years				
...	All causes	669	100.0	39.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	259	38.7	15.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	72	10.8	4.3
3	Malignant neoplasms (C00-C97)	70	10.5	4.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	51	7.6	3.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	35	5.2	2.1
6	Pregnancy, childbirth and the puerperium (O00-099)	23	3.4	1.4
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	16	2.4	*
8	Cerebrovascular diseases (I60-I69)	10	1.5	*
9	Septicemia (A40-A41)	6	0.9	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	6	0.9	*
11	Diabetes mellitus (E10-E14)	4	0.6	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.6	*
13	Influenza and pneumonia (J10-J18)	3	0.4	*
14	Anemias (D50-D64)	2	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.3	*
...	All other causes (Residual)	106	15.8	6.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 25-29 years				
...	All causes	660	100.0	36.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	191	28.9	10.7
2	Malignant neoplasms (C00-C97)	94	14.2	5.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	67	10.2	3.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	39	5.9	2.2
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	30	4.5	1.7
6	Pregnancy, childbirth and the puerperium (O00-099)	23	3.5	1.3
7	Human immunodeficiency virus (HIV) disease (B20-B24)	16	2.4	*
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	15	2.3	*
9	Diabetes mellitus (E10-E14)	12	1.8	*
10	Cerebrovascular diseases (I60-I69)	10	1.5	*
11	Influenza and pneumonia (J10-J18)	8	1.2	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.9	*
12	Chronic liver disease and cirrhosis (K70,K73-K74)	6	0.9	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.8	*
15	Septicemia (A40-A41)	3	0.5	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.5	*
...	All other causes (Residual)	132	20.0	7.4
Hispanic, female, 30-34 years				
...	All causes	791	100.0	45.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	181	22.9	10.4
2	Malignant neoplasms (C00-C97)	156	19.7	8.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	62	7.8	3.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	43	5.4	2.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	43	5.4	2.5
6	Pregnancy, childbirth and the puerperium (O00-099)	37	4.7	2.1
7	Human immunodeficiency virus (HIV) disease (B20-B24)	34	4.3	1.9
8	Cerebrovascular diseases (I60-I69)	22	2.8	1.3
9	Diabetes mellitus (E10-E14)	17	2.1	*
10	Septicemia (A40-A41)	10	1.3	*
10	Chronic liver disease and cirrhosis (K70,K73-K74)	10	1.3	*
12	Chronic lower respiratory diseases (J40-J47)	9	1.1	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	1.1	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	0.9	*
15	Influenza and pneumonia (J10-J18)	6	0.8	*
...	All other causes (Residual)	145	18.3	8.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 35-39 years				
...	All causes	1,098	100.0	69.5
1	Malignant neoplasms (C00-C97)	294	26.8	18.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	195	17.8	12.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	76	6.9	4.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	58	5.3	3.7
5	Human immunodeficiency virus (HIV) disease (B20-B24)	46	4.2	2.9
6	Cerebrovascular diseases (I60-I69)	41	3.7	2.6
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	40	3.6	2.5
8	Chronic liver disease and cirrhosis (K70,K73-K74)	29	2.6	1.8
9	Pregnancy, childbirth and the puerperium (O00-099)	22	2.0	1.4
10	Septicemia (A40-A41)	17	1.5	*
10	Diabetes mellitus (E10-E14)	17	1.5	*
12	Influenza and pneumonia (J10-J18)	11	1.0	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	1.0	*
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	11	1.0	*
15	Viral hepatitis (B15-B19)	7	0.6	*
...	All other causes (Residual)	223	20.3	14.1
Hispanic, female, 40-44 years				
...	All causes	1,633	100.0	113.8
1	Malignant neoplasms (C00-C97)	520	31.8	36.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	212	13.0	14.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	148	9.1	10.3
4	Cerebrovascular diseases (I60-I69)	78	4.8	5.4
5	Human immunodeficiency virus (HIV) disease (B20-B24)	74	4.5	5.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	69	4.2	4.8
7	Diabetes mellitus (E10-E14)	46	2.8	3.2
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	34	2.1	2.4
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	33	2.0	2.3
10	Influenza and pneumonia (J10-J18)	23	1.4	1.6
11	Septicemia (A40-A41)	22	1.3	1.5
11	Viral hepatitis (B15-B19)	22	1.3	1.5
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20	1.2	1.4
14	Chronic lower respiratory diseases (J40-J47)	13	0.8	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.7	*
...	All other causes (Residual)	308	18.9	21.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 45-49 years				
...	All causes	2,007	100.0	169.1
1	Malignant neoplasms (C00-C97)	714	35.6	60.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	228	11.4	19.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	171	8.5	14.4
4	Cerebrovascular diseases (I60-I69)	126	6.3	10.6
5	Diabetes mellitus (E10-E14)	99	4.9	8.3
6	Chronic liver disease and cirrhosis (K70,K73-K74)	81	4.0	6.8
7	Human immunodeficiency virus (HIV) disease (B20-B24)	57	2.8	4.8
8	Chronic lower respiratory diseases (J40-J47)	34	1.7	2.9
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	31	1.5	2.6
10	Influenza and pneumonia (J10-J18)	30	1.5	2.5
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	28	1.4	2.4
12	Septicemia (A40-A41)	27	1.3	2.3
13	Viral hepatitis (B15-B19)	23	1.1	1.9
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22	1.1	1.9
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	16	0.8	*
...	All other causes (Residual)	320	15.9	27.0
Hispanic, female, 50-54 years				
...	All causes	2,568	100.0	276.8
1	Malignant neoplasms (C00-C97)	975	38.0	105.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	326	12.7	35.1
3	Diabetes mellitus (E10-E14)	155	6.0	16.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	143	5.6	15.4
5	Cerebrovascular diseases (I60-I69)	129	5.0	13.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	108	4.2	11.6
7	Human immunodeficiency virus (HIV) disease (B20-B24)	52	2.0	5.6
8	Viral hepatitis (B15-B19)	48	1.9	5.2
9	Chronic lower respiratory diseases (J40-J47)	40	1.6	4.3
10	Septicemia (A40-A41)	37	1.4	4.0
11	Influenza and pneumonia (J10-J18)	32	1.2	3.4
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	31	1.2	3.3
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	24	0.9	2.6
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	18	0.7	*
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	15	0.6	*
...	All other causes (Residual)	435	16.9	46.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 55-59 years				
...	All causes	2,867	100.0	395.9
1	Malignant neoplasms (C00-C97)	1,009	35.2	139.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	489	17.1	67.5
3	Diabetes mellitus (E10-E14)	223	7.8	30.8
4	Cerebrovascular diseases (I60-I69)	161	5.6	22.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	127	4.4	17.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	105	3.7	14.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	64	2.2	8.8
8	Chronic lower respiratory diseases (J40-J47)	56	2.0	7.7
9	Septicemia (A40-A41)	45	1.6	6.2
10	Viral hepatitis (B15-B19)	43	1.5	5.9
11	Influenza and pneumonia (J10-J18)	37	1.3	5.1
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	35	1.2	4.8
13	Human immunodeficiency virus (HIV) disease (B20-B24)	23	0.8	3.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	0.6	*
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	15	0.5	*
...	All other causes (Residual)	419	14.6	57.9
Hispanic, female, 60-64 years				
...	All causes	3,268	100.0	631.0
1	Malignant neoplasms (C00-C97)	1,134	34.7	219.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	661	20.2	127.6
3	Diabetes mellitus (E10-E14)	283	8.7	54.6
4	Cerebrovascular diseases (I60-I69)	176	5.4	34.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	86	2.6	16.6
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	86	2.6	16.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	85	2.6	16.4
8	Septicemia (A40-A41)	64	2.0	12.4
9	Influenza and pneumonia (J10-J18)	63	1.9	12.2
10	Chronic lower respiratory diseases (J40-J47)	62	1.9	12.0
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	46	1.4	8.9
12	Viral hepatitis (B15-B19)	29	0.9	5.6
13	Alzheimer's disease (G30)	16	0.5	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	15	0.5	*
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11	0.3	*
...	All other causes (Residual)	451	13.8	87.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 65-69 years				
...	All causes	4,103	100.0	1,009.5
1	Malignant neoplasms (C00-C97)	1,307	31.9	321.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	889	21.7	218.7
3	Diabetes mellitus (E10-E14)	368	9.0	90.5
4	Cerebrovascular diseases (I60-I69)	227	5.5	55.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	126	3.1	31.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	119	2.9	29.3
7	Chronic lower respiratory diseases (J40-J47)	105	2.6	25.8
8	Influenza and pneumonia (J10-J18)	85	2.1	20.9
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	84	2.0	20.7
10	Septicemia (A40-A41)	78	1.9	19.2
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	47	1.1	11.6
12	Alzheimer's disease (G30)	31	0.8	7.6
13	Viral hepatitis (B15-B19)	26	0.6	6.4
14	Parkinson's disease (G20-G21)	16	0.4	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	15	0.4	*
...	All other causes (Residual)	580	14.1	142.7
Hispanic, female, 70-74 years				
...	All causes	5,335	100.0	1,645.2
1	Malignant neoplasms (C00-C97)	1,488	27.9	458.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,346	25.2	415.1
3	Diabetes mellitus (E10-E14)	452	8.5	139.4
4	Cerebrovascular diseases (I60-I69)	327	6.1	100.8
5	Chronic lower respiratory diseases (J40-J47)	182	3.4	56.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	143	2.7	44.1
7	Influenza and pneumonia (J10-J18)	120	2.2	37.0
8	Chronic liver disease and cirrhosis (K70,K73-K74)	111	2.1	34.2
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	91	1.7	28.1
10	Septicemia (A40-A41)	84	1.6	25.9
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	78	1.5	24.1
12	Alzheimer's disease (G30)	59	1.1	18.2
13	Viral hepatitis (B15-B19)	34	0.6	10.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	29	0.5	8.9
15	Pneumonitis due to solids and liquids (J69)	26	0.5	8.0
...	All other causes (Residual)	765	14.3	235.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 75-79 years				
...	All causes	6,773	100.0	2,627.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,813	26.8	703.4
2	Malignant neoplasms (C00-C97)	1,620	23.9	628.5
3	Diabetes mellitus (E10-E14)	571	8.4	221.5
4	Cerebrovascular diseases (I60-I69)	487	7.2	188.9
5	Chronic lower respiratory diseases (J40-J47)	264	3.9	102.4
6	Influenza and pneumonia (J10-J18)	193	2.8	74.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	181	2.7	70.2
8	Alzheimer's disease (G30)	150	2.2	58.2
9	Chronic liver disease and cirrhosis (K70,K73-K74)	128	1.9	49.7
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	120	1.8	46.6
11	Septicemia (A40-A41)	102	1.5	39.6
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	102	1.5	39.6
13	Parkinson's disease (G20-G21)	64	0.9	24.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	38	0.6	14.7
15	Pneumonitis due to solids and liquids (J69)	31	0.5	12.0
...	All other causes (Residual)	909	13.4	352.7
Hispanic, female, 80-84 years				
...	All causes	7,698	100.0	4,470.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,372	30.8	1,377.5
2	Malignant neoplasms (C00-C97)	1,340	17.4	778.2
3	Cerebrovascular diseases (I60-I69)	642	8.3	372.8
4	Diabetes mellitus (E10-E14)	499	6.5	289.8
5	Alzheimer's disease (G30)	314	4.1	182.4
6	Chronic lower respiratory diseases (J40-J47)	289	3.8	167.8
7	Influenza and pneumonia (J10-J18)	251	3.3	145.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	180	2.3	104.5
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	143	1.9	83.0
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	122	1.6	70.9
11	Septicemia (A40-A41)	121	1.6	70.3
12	Parkinson's disease (G20-G21)	77	1.0	44.7
13	Chronic liver disease and cirrhosis (K70,K73-K74)	60	0.8	34.8
14	Pneumonitis due to solids and liquids (J69)	50	0.6	29.0
15	Atherosclerosis (I70)	44	0.6	25.6
...	All other causes (Residual)	1,194	15.5	693.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 85-89 years				
...	All causes	6,483	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,211	34.1	---
2	Malignant neoplasms (C00-C97)	819	12.6	---
3	Cerebrovascular diseases (I60-I69)	584	9.0	---
4	Alzheimer's disease (G30)	363	5.6	---
5	Diabetes mellitus (E10-E14)	348	5.4	---
6	Influenza and pneumonia (J10-J18)	276	4.3	---
7	Chronic lower respiratory diseases (J40-J47)	274	4.2	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	120	1.9	---
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	115	1.8	---
10	Septicemia (A40-A41)	114	1.8	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	93	1.4	---
12	Parkinson's disease (G20-G21)	65	1.0	---
13	Pneumonitis due to solids and liquids (J69)	63	1.0	---
14	Atherosclerosis (I70)	38	0.6	---
15	Chronic liver disease and cirrhosis (K70,K73-K74)	36	0.6	---
...	All other causes (Residual)	964	14.9	---
Hispanic, female, 90-94 years				
...	All causes	4,850	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,859	38.3	---
2	Cerebrovascular diseases (I60-I69)	403	8.3	---
3	Malignant neoplasms (C00-C97)	400	8.2	---
4	Alzheimer's disease (G30)	304	6.3	---
5	Influenza and pneumonia (J10-J18)	247	5.1	---
6	Diabetes mellitus (E10-E14)	196	4.0	---
7	Chronic lower respiratory diseases (J40-J47)	178	3.7	---
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	101	2.1	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	91	1.9	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	87	1.8	---
11	Septicemia (A40-A41)	56	1.2	---
12	Atherosclerosis (I70)	40	0.8	---
12	Pneumonitis due to solids and liquids (J69)	40	0.8	---
14	Parkinson's disease (G20-G21)	31	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	0.5	---
...	All other causes (Residual)	794	16.4	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 95-99 years				
...	All causes	2,037	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	814	40.0	---
2	Alzheimer's disease (G30)	163	8.0	---
3	Cerebrovascular diseases (I60-I69)	150	7.4	---
4	Malignant neoplasms (C00-C97)	119	5.8	---
5	Influenza and pneumonia (J10-J18)	107	5.3	---
6	Chronic lower respiratory diseases (J40-J47)	79	3.9	---
7	Diabetes mellitus (E10-E14)	63	3.1	---
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	51	2.5	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	34	1.7	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	32	1.6	---
11	Septicemia (A40-A41)	29	1.4	---
12	Atherosclerosis (I70)	27	1.3	---
13	Pneumonitis due to solids and liquids (J69)	24	1.2	---
14	Parkinson's disease (G20-G21)	14	0.7	---
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	9	0.4	---
...	All other causes (Residual)	322	15.8	---
Hispanic, female, 100 years and over				
...	All causes	545	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	243	44.6	---
2	Influenza and pneumonia (J10-J18)	39	7.2	---
3	Cerebrovascular diseases (I60-I69)	33	6.1	---
4	Alzheimer's disease (G30)	26	4.8	---
5	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	16	2.9	---
6	Chronic lower respiratory diseases (J40-J47)	15	2.8	---
7	Diabetes mellitus (E10-E14)	14	2.6	---
7	Atherosclerosis (I70)	14	2.6	---
9	Malignant neoplasms (C00-C97)	13	2.4	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8	1.5	---
11	Septicemia (A40-A41)	6	1.1	---
11	Nutritional deficiencies (E40-E64)	6	1.1	---
11	Pneumonitis due to solids and liquids (J69)	6	1.1	---
14	Cholelithiasis and other disorders of gallbladder (K80-K82)	3	0.6	---
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.6	---
...	All other causes (Residual)	100	18.3	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, all ages				
...	All causes	1,967,142	100.0	981.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	535,101	27.2	267.1
2	Malignant neoplasms (C00-C97)	456,133	23.2	227.7
3	Chronic lower respiratory diseases (J40-J47)	117,337	6.0	58.6
4	Cerebrovascular diseases (I60-I69)	115,074	5.8	57.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	88,970	4.5	44.4
6	Alzheimer's disease (G30)	63,958	3.3	31.9
7	Diabetes mellitus (E10-E14)	53,159	2.7	26.5
8	Influenza and pneumonia (J10-J18)	52,431	2.7	26.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	32,492	1.7	16.2
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	27,305	1.4	13.6
11	Septicemia (A40-A41)	25,496	1.3	12.7
12	Chronic liver disease and cirrhosis (K70,K73-K74)	20,428	1.0	10.2
		17,949	0.9	9.0
14	Parkinson's disease (G20-G21)	17,767	0.9	8.9
15	Pneumonitis due to solids and liquids (J69)	14,739	0.7	7.4
...	All other causes (Residual)	328,803	16.7	164.1
White, non-Hispanic, both sexes, 1-4 years				
...	All causes	2,404	100.0	26.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	898	37.4	9.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	270	11.2	2.9
3	Malignant neoplasms (C00-C97)	215	8.9	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	138	5.7	1.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	70	2.9	0.8
6	Influenza and pneumonia (J10-J18)	46	1.9	0.5
7	Septicemia (A40-A41)	44	1.8	0.5
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	31	1.3	0.3
9	Cerebrovascular diseases (I60-I69)	30	1.2	0.3
10	Certain conditions originating in the perinatal period (P00-P96)	29	1.2	0.3
11	Chronic lower respiratory diseases (J40-J47)	20	0.8	0.2
12	Meningitis (G00,G03)	12	0.5	*
13	Acute bronchitis and bronchiolitis (J20-J21)	10	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	9	0.4	*
...	All other causes (Residual)	573	23.8	6.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 5-9 years				
...	All causes	1,561	100.0	13.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	596	38.2	5.1
2	Malignant neoplasms (C00-C97)	288	18.4	2.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	101	6.5	0.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	53	3.4	0.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	49	3.1	0.4
6	Influenza and pneumonia (J10-J18)	32	2.0	0.3
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	24	1.5	0.2
7	Cerebrovascular diseases (I60-I69)	24	1.5	0.2
9	Chronic lower respiratory diseases (J40-J47)	21	1.3	0.2
10	Septicemia (A40-A41)	18	1.2	*
11	Certain conditions originating in the perinatal period (P00-P96)	10	0.6	*
12	Meningitis (G00,G03)	9	0.6	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.3	*
15	Anemias (D50-D64)	3	0.2	*
15	Diabetes mellitus (E10-E14)	3	0.2	*
15	Acute bronchitis and bronchiolitis (J20-J21)	3	0.2	*
15	Pneumonitis due to solids and liquids (J69)	3	0.2	*
15	Diseases of appendix (K35-K38)	3	0.2	*
...	All other causes (Residual)	309	19.8	2.7
White, non-Hispanic, both sexes, 10-14 years				
...	All causes	2,120	100.0	16.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	827	39.0	6.6
2	Malignant neoplasms (C00-C97)	309	14.6	2.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	169	8.0	1.3
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	106	5.0	0.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	74	3.5	0.6
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	62	2.9	0.5
7	Influenza and pneumonia (J10-J18)	35	1.7	0.3
8	Septicemia (A40-A41)	29	1.4	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	1.1	0.2
10	Cerebrovascular diseases (I60-I69)	21	1.0	0.2
11	Diabetes mellitus (E10-E14)	19	0.9	*
12	Chronic lower respiratory diseases (J40-J47)	10	0.5	*
13	Certain conditions originating in the perinatal period (P00-P96)	9	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.4	*
15	Meningitis (G00,G03)	7	0.3	*
...	All other causes (Residual)	412	19.4	3.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 15-19 years				
...	All causes	7,989	100.0	60.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,536	56.8	34.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,160	14.5	8.8
3	Malignant neoplasms (C00-C97)	452	5.7	3.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	335	4.2	2.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	216	2.7	1.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	152	1.9	1.1
7	Influenza and pneumonia (J10-J18)	47	0.6	0.4
8	Diabetes mellitus (E10-E14)	37	0.5	0.3
9	Cerebrovascular diseases (I60-I69)	35	0.4	0.3
10	Septicemia (A40-A41)	30	0.4	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	30	0.4	0.2
12	Chronic lower respiratory diseases (J40-J47)	28	0.4	0.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	0.2	*
14	Pneumonitis due to solids and liquids (J69)	15	0.2	*
15	Meningococcal infection (A39)	14	0.2	*
...	All other causes (Residual)	885	11.1	6.7
White, non-Hispanic, both sexes, 20-24 years				
...	All causes	11,694	100.0	89.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,175	52.8	47.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,832	15.7	14.0
3	Malignant neoplasms (C00-C97)	596	5.1	4.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	560	4.8	4.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	388	3.3	3.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	161	1.4	1.2
7	Influenza and pneumonia (J10-J18)	76	0.6	0.6
8	Diabetes mellitus (E10-E14)	73	0.6	0.6
9	Pregnancy, childbirth and the puerperium (O00-099)	66	0.6	0.5
10	Cerebrovascular diseases (I60-I69)	57	0.5	0.4
11	Septicemia (A40-A41)	47	0.4	0.4
12	Chronic lower respiratory diseases (J40-J47)	36	0.3	0.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	33	0.3	0.3
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	28	0.2	0.2
15	Legal intervention (Y35,Y89.0)	21	0.2	0.2
...	All other causes (Residual)	1,545	13.2	11.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 25-29 years				
...	All causes	10,883	100.0	90.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,825	44.3	40.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,698	15.6	14.1
3	Malignant neoplasms (C00-C97)	776	7.1	6.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	600	5.5	5.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	516	4.7	4.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	145	1.3	1.2
7	Diabetes mellitus (E10-E14)	94	0.9	0.8
8	Influenza and pneumonia (J10-J18)	93	0.9	0.8
9	Cerebrovascular diseases (I60-I69)	91	0.8	0.8
10	Septicemia (A40-A41)	66	0.6	0.5
11	Human immunodeficiency virus (HIV) disease (B20-B24)	65	0.6	0.5
12	Pregnancy, childbirth and the puerperium (O00-099)	56	0.5	0.5
13	Chronic lower respiratory diseases (J40-J47)	54	0.5	0.4
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	51	0.5	0.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	41	0.4	0.3
...	All other causes (Residual)	1,712	15.7	14.3
White, non-Hispanic, both sexes, 30-34 years				
...	All causes	12,943	100.0	106.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,522	34.9	37.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,010	15.5	16.5
3	Malignant neoplasms (C00-C97)	1,399	10.8	11.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,142	8.8	9.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	457	3.5	3.7
6	Human immunodeficiency virus (HIV) disease (B20-B24)	212	1.6	1.7
6	Diabetes mellitus (E10-E14)	212	1.6	1.7
8	Cerebrovascular diseases (I60-I69)	183	1.4	1.5
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	142	1.1	1.2
10	Influenza and pneumonia (J10-J18)	131	1.0	1.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	117	0.9	1.0
12	Septicemia (A40-A41)	91	0.7	0.7
13	Chronic lower respiratory diseases (J40-J47)	75	0.6	0.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	72	0.6	0.6
15	Pregnancy, childbirth and the puerperium (O00-099)	65	0.5	0.5
...	All other causes (Residual)	2,113	16.3	17.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 35-39 years				
...	All causes	19,504	100.0	143.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,211	26.7	38.3
2	Malignant neoplasms (C00-C97)	2,890	14.8	21.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,470	12.7	18.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,338	12.0	17.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	490	2.5	3.6
6	Human immunodeficiency virus (HIV) disease (B20-B24)	469	2.4	3.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	459	2.4	3.4
8	Diabetes mellitus (E10-E14)	445	2.3	3.3
9	Cerebrovascular diseases (I60-I69)	350	1.8	2.6
10	Influenza and pneumonia (J10-J18)	196	1.0	1.4
11	Septicemia (A40-A41)	172	0.9	1.3
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	169	0.9	1.2
13	Chronic lower respiratory diseases (J40-J47)	164	0.8	1.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	121	0.6	0.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	95	0.5	0.7
...	All other causes (Residual)	3,465	17.8	25.5
White, non-Hispanic, both sexes, 40-44 years				
...	All causes	35,282	100.0	224.0
1	Malignant neoplasms (C00-C97)	7,086	20.1	45.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,041	20.0	44.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,683	16.1	36.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,096	8.8	19.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,355	3.8	8.6
6	Human immunodeficiency virus (HIV) disease (B20-B24)	883	2.5	5.6
7	Cerebrovascular diseases (I60-I69)	808	2.3	5.1
8	Diabetes mellitus (E10-E14)	770	2.2	4.9
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	609	1.7	3.9
10	Chronic lower respiratory diseases (J40-J47)	429	1.2	2.7
11	Influenza and pneumonia (J10-J18)	397	1.1	2.5
12	Septicemia (A40-A41)	308	0.9	2.0
13	Viral hepatitis (B15-B19)	238	0.7	1.5
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	212	0.6	1.3
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	212	0.6	1.3
...	All other causes (Residual)	6,155	17.4	39.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 45-49 years				
...	All causes	54,109	100.0	333.9
1	Malignant neoplasms (C00-C97)	13,796	25.5	85.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,430	19.3	64.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,545	13.9	46.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,313	6.1	20.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,368	4.4	14.6
6	Cerebrovascular diseases (I60-I69)	1,383	2.6	8.5
7	Diabetes mellitus (E10-E14)	1,331	2.5	8.2
8	Chronic lower respiratory diseases (J40-J47)	1,084	2.0	6.7
9	Human immunodeficiency virus (HIV) disease (B20-B24)	774	1.4	4.8
10	Viral hepatitis (B15-B19)	669	1.2	4.1
11	Influenza and pneumonia (J10-J18)	626	1.2	3.9
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	549	1.0	3.4
13	Septicemia (A40-A41)	533	1.0	3.3
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	428	0.8	2.6
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	250	0.5	1.5
...	All other causes (Residual)	9,030	16.7	55.7
White, non-Hispanic, both sexes, 50-54 years				
...	All causes	72,560	100.0	486.6
1	Malignant neoplasms (C00-C97)	22,637	31.2	151.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,705	21.6	105.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,112	8.4	41.0
4	Chronic liver disease and cirrhosis (K70,K73-K74)	2,955	4.1	19.8
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,895	4.0	19.4
6	Diabetes mellitus (E10-E14)	2,123	2.9	14.2
7	Cerebrovascular diseases (I60-I69)	2,078	2.9	13.9
8	Chronic lower respiratory diseases (J40-J47)	1,970	2.7	13.2
9	Viral hepatitis (B15-B19)	933	1.3	6.3
10	Influenza and pneumonia (J10-J18)	823	1.1	5.5
11	Septicemia (A40-A41)	809	1.1	5.4
12	Human immunodeficiency virus (HIV) disease (B20-B24)	634	0.9	4.3
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	631	0.9	4.2
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	371	0.5	2.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	312	0.4	2.1
...	All other causes (Residual)	11,572	15.9	77.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 55-59 years				
...	All causes	93,451	100.0	699.3
1	Malignant neoplasms (C00-C97)	34,030	36.4	254.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	21,306	22.8	159.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,680	5.0	35.0
4	Chronic lower respiratory diseases (J40-J47)	3,931	4.2	29.4
5	Diabetes mellitus (E10-E14)	3,278	3.5	24.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	2,936	3.1	22.0
7	Cerebrovascular diseases (I60-I69)	2,733	2.9	20.5
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,241	2.4	16.8
9	Septicemia (A40-A41)	1,204	1.3	9.0
10	Influenza and pneumonia (J10-J18)	1,060	1.1	7.9
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	978	1.0	7.3
12	Viral hepatitis (B15-B19)	596	0.6	4.5
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	457	0.5	3.4
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	454	0.5	3.4
15	Aortic aneurysm and dissection (I71)	445	0.5	3.3
...	All other causes (Residual)	13,122	14.0	98.2
White, non-Hispanic, both sexes, 60-64 years				
...	All causes	112,627	100.0	1,101.0
1	Malignant neoplasms (C00-C97)	43,587	38.7	426.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	26,438	23.5	258.4
3	Chronic lower respiratory diseases (J40-J47)	7,127	6.3	69.7
4	Diabetes mellitus (E10-E14)	4,015	3.6	39.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,623	3.2	35.4
6	Cerebrovascular diseases (I60-I69)	3,618	3.2	35.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,322	2.1	22.7
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,581	1.4	15.5
9	Septicemia (A40-A41)	1,437	1.3	14.0
10	Influenza and pneumonia (J10-J18)	1,397	1.2	13.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,339	1.2	13.1
12	Aortic aneurysm and dissection (I71)	703	0.6	6.9
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	605	0.5	5.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	577	0.5	5.6
15	Alzheimer's disease (G30)	386	0.3	3.8
...	All other causes (Residual)	13,872	12.3	135.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 65-69 years				
...	All causes	133,683	100.0	1,675.1
1	Malignant neoplasms (C00-C97)	50,729	37.9	635.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	31,085	23.3	389.5
3	Chronic lower respiratory diseases (J40-J47)	10,694	8.0	134.0
4	Cerebrovascular diseases (I60-I69)	5,228	3.9	65.5
5	Diabetes mellitus (E10-E14)	4,747	3.6	59.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,187	2.4	39.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,950	1.5	24.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,856	1.4	23.3
9	Influenza and pneumonia (J10-J18)	1,836	1.4	23.0
10	Septicemia (A40-A41)	1,830	1.4	22.9
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,094	0.8	13.7
12	Aortic aneurysm and dissection (I71)	973	0.7	12.2
13	Alzheimer's disease (G30)	816	0.6	10.2
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	812	0.6	10.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	796	0.6	10.0
...	All other causes (Residual)	16,050	12.0	201.1
White, non-Hispanic, both sexes, 70-74 years				
...	All causes	181,275	100.0	2,652.5
1	Malignant neoplasms (C00-C97)	62,211	34.3	910.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	43,631	24.1	638.4
3	Chronic lower respiratory diseases (J40-J47)	16,130	8.9	236.0
4	Cerebrovascular diseases (I60-I69)	8,446	4.7	123.6
5	Diabetes mellitus (E10-E14)	6,079	3.4	89.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,703	2.0	54.2
7	Influenza and pneumonia (J10-J18)	3,186	1.8	46.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,990	1.6	43.8
9	Septicemia (A40-A41)	2,583	1.4	37.8
10	Alzheimer's disease (G30)	2,392	1.3	35.0
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1,979	1.1	29.0
12	Parkinson's disease (G20-G21)	1,523	0.8	22.3
13	Aortic aneurysm and dissection (I71)	1,470	0.8	21.5
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,166	0.6	17.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,162	0.6	17.0
...	All other causes (Residual)	22,624	12.5	331.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 75-79 years				
...	All causes	256,164	100.0	4,179.8
1	Malignant neoplasms (C00-C97)	72,762	28.4	1,187.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	66,255	25.9	1,081.1
3	Chronic lower respiratory diseases (J40-J47)	22,260	8.7	363.2
4	Cerebrovascular diseases (I60-I69)	15,084	5.9	246.1
5	Diabetes mellitus (E10-E14)	8,114	3.2	132.4
6	Alzheimer's disease (G30)	6,618	2.6	108.0
7	Influenza and pneumonia (J10-J18)	5,831	2.3	95.1
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,165	2.0	84.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,787	1.9	78.1
10	Septicemia (A40-A41)	3,643	1.4	59.4
11	Parkinson's disease (G20-G21)	3,357	1.3	54.8
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,117	0.8	34.5
13	Aortic aneurysm and dissection (I71)	2,048	0.8	33.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,800	0.7	29.4
15	Pneumonitis due to solids and liquids (J69)	1,763	0.7	28.8
...	All other causes (Residual)	34,560	13.5	563.9
White, non-Hispanic, both sexes, 80-84 years				
...	All causes	325,548	100.0	6,803.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	93,829	28.8	1,960.7
2	Malignant neoplasms (C00-C97)	69,552	21.4	1,453.4
3	Cerebrovascular diseases (I60-I69)	23,746	7.3	496.2
4	Chronic lower respiratory diseases (J40-J47)	23,681	7.3	494.9
5	Alzheimer's disease (G30)	13,851	4.3	289.4
6	Influenza and pneumonia (J10-J18)	9,807	3.0	204.9
7	Diabetes mellitus (E10-E14)	9,179	2.8	191.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,777	2.1	141.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,511	2.0	136.1
10	Parkinson's disease (G20-G21)	5,088	1.6	106.3
11	Septicemia (A40-A41)	4,738	1.5	99.0
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	3,298	1.0	68.9
13	Pneumonitis due to solids and liquids (J69)	2,987	0.9	62.4
14	Aortic aneurysm and dissection (I71)	2,342	0.7	48.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,262	0.7	47.3
...	All other causes (Residual)	47,900	14.7	1,001.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 85-89 years				
...	All causes	310,106	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	100,682	32.5	---
2	Malignant neoplasms (C00-C97)	46,235	14.9	---
3	Cerebrovascular diseases (I60-I69)	25,299	8.2	---
4	Alzheimer's disease (G30)	17,994	5.8	---
5	Chronic lower respiratory diseases (J40-J47)	17,744	5.7	---
6	Influenza and pneumonia (J10-J18)	11,540	3.7	---
7	Diabetes mellitus (E10-E14)	7,434	2.4	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,669	2.2	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,462	2.1	---
10	Parkinson's disease (G20-G21)	4,279	1.4	---
11	Septicemia (A40-A41)	4,226	1.4	---
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	3,759	1.2	---
13	Pneumonitis due to solids and liquids (J69)	3,530	1.1	---
14	Atherosclerosis (I70)	2,467	0.8	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,044	0.7	---
...	All other causes (Residual)	49,742	16.0	---
White, non-Hispanic, both sexes, 90-94 years				
...	All causes	212,418	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	76,861	36.2	---
2	Malignant neoplasms (C00-C97)	20,733	9.8	---
3	Cerebrovascular diseases (I60-I69)	18,067	8.5	---
4	Alzheimer's disease (G30)	14,574	6.9	---
5	Influenza and pneumonia (J10-J18)	9,738	4.6	---
6	Chronic lower respiratory diseases (J40-J47)	8,947	4.2	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,354	2.0	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,185	2.0	---
9	Diabetes mellitus (E10-E14)	3,884	1.8	---
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	3,223	1.5	---
11	Pneumonitis due to solids and liquids (J69)	2,665	1.3	---
12	Septicemia (A40-A41)	2,596	1.2	---
13	Atherosclerosis (I70)	2,277	1.1	---
14	Parkinson's disease (G20-G21)	2,024	1.0	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,148	0.5	---
...	All other causes (Residual)	37,142	17.5	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 95-99 years				
...	All causes	81,590	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	31,364	38.4	---
2	Cerebrovascular diseases (I60-I69)	6,548	8.0	---
3	Alzheimer's disease (G30)	6,047	7.4	---
4	Malignant neoplasms (C00-C97)	5,179	6.3	---
5	Influenza and pneumonia (J10-J18)	4,383	5.4	---
6	Chronic lower respiratory diseases (J40-J47)	2,581	3.2	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,606	2.0	---
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,507	1.8	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,478	1.8	---
10	Diabetes mellitus (E10-E14)	1,162	1.4	---
11	Atherosclerosis (I70)	1,133	1.4	---
12	Pneumonitis due to solids and liquids (J69)	995	1.2	---
13	Septicemia (A40-A41)	867	1.1	---
14	Parkinson's disease (G20-G21)	440	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	325	0.4	---
...	All other causes (Residual)	15,975	19.6	---
White, non-Hispanic, both sexes, 100 years and over				
...	All causes	16,132	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,638	41.1	---
2	Cerebrovascular diseases (I60-I69)	1,189	7.4	---
3	Influenza and pneumonia (J10-J18)	1,056	6.5	---
4	Alzheimer's disease (G30)	1,031	6.4	---
5	Malignant neoplasms (C00-C97)	623	3.9	---
6	Chronic lower respiratory diseases (J40-J47)	339	2.1	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	333	2.1	---
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	299	1.9	---
9	Atherosclerosis (I70)	285	1.8	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	262	1.6	---
11	Pneumonitis due to solids and liquids (J69)	180	1.1	---
12	Diabetes mellitus (E10-E14)	158	1.0	---
13	Septicemia (A40-A41)	125	0.8	---
14	Anemias (D50-D64)	77	0.5	---
15	Nutritional deficiencies (E40-E64)	52	0.3	---
...	All other causes (Residual)	3,485	21.6	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, all ages				
...	All causes	954,402	100.0	970.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	263,311	27.6	267.8
2	Malignant neoplasms (C00-C97)	236,659	24.8	240.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	56,021	5.9	57.0
4	Chronic lower respiratory diseases (J40-J47)	55,015	5.8	56.0
5	Cerebrovascular diseases (I60-I69)	44,038	4.6	44.8
6	Diabetes mellitus (E10-E14)	26,365	2.8	26.8
7	Influenza and pneumonia (J10-J18)	22,903	2.4	23.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	21,605	2.3	22.0
9	Alzheimer's disease (G30)	18,208	1.9	18.5
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15,953	1.7	16.2
11	Chronic liver disease and cirrhosis (K70,K73-K74)	13,088	1.4	13.3
12	Septicemia (A40-A41)	11,337	1.2	11.5
13	Parkinson's disease (G20-G21)	10,232	1.1	10.4
14	Pneumonitis due to solids and liquids (J69)	7,430	0.8	7.6
15	Aortic aneurysm and dissection (I71)	6,877	0.7	7.0
...	All other causes (Residual)	145,360	15.2	147.8
White, non-Hispanic, male, 1-4 years				
...	All causes	1,408	100.0	29.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	557	39.6	11.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	139	9.9	3.0
3	Malignant neoplasms (C00-C97)	118	8.4	2.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	74	5.3	1.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	38	2.7	0.8
6	Septicemia (A40-A41)	26	1.8	0.6
7	Influenza and pneumonia (J10-J18)	23	1.6	0.5
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	1.4	0.4
8	Cerebrovascular diseases (I60-I69)	20	1.4	0.4
10	Certain conditions originating in the perinatal period (P00-P96)	18	1.3	*
11	Meningitis (G00,G03)	11	0.8	*
12	Chronic lower respiratory diseases (J40-J47)	9	0.6	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	8	0.6	*
14	Acute bronchitis and bronchiolitis (J20-J21)	6	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.4	*
...	All other causes (Residual)	335	23.8	7.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 5-9 years				
...	All causes	860	100.0	14.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	354	41.2	6.0
2	Malignant neoplasms (C00-C97)	155	18.0	2.6
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	41	4.8	0.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	33	3.8	0.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	28	3.3	0.5
6	Influenza and pneumonia (J10-J18)	14	1.6	*
7	Chronic lower respiratory diseases (J40-J47)	13	1.5	*
8	Septicemia (A40-A41)	11	1.3	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	1.3	*
8	Cerebrovascular diseases (I60-I69)	11	1.3	*
11	Certain conditions originating in the perinatal period (P00-P96)	6	0.7	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.6	*
13	Meningitis (G00,G03)	3	0.3	*
13	Acute bronchitis and bronchiolitis (J20-J21)	3	0.3	*
13	Pneumonitis due to solids and liquids (J69)	3	0.3	*
13	Diseases of appendix (K35-K38)	3	0.3	*
...	All other causes (Residual)	166	19.3	2.8
White, non-Hispanic, male, 10-14 years				
...	All causes	1,303	100.0	20.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	532	40.8	8.2
2	Malignant neoplasms (C00-C97)	178	13.7	2.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	132	10.1	2.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	60	4.6	0.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	42	3.2	0.6
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	28	2.1	0.4
7	Septicemia (A40-A41)	18	1.4	*
8	Cerebrovascular diseases (I60-I69)	15	1.2	*
8	Influenza and pneumonia (J10-J18)	15	1.2	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.8	*
11	Diabetes mellitus (E10-E14)	8	0.6	*
12	Meningitis (G00,G03)	6	0.5	*
13	Chronic lower respiratory diseases (J40-J47)	4	0.3	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.3	*
15	Meningococcal infection (A39)	3	0.2	*
15	Aortic aneurysm and dissection (I71)	3	0.2	*
15	Certain conditions originating in the perinatal period (P00-P96)	3	0.2	*
...	All other causes (Residual)	241	18.5	3.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 15-19 years				
...	All causes	5,568	100.0	82.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,134	56.3	46.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	949	17.0	14.0
3	Malignant neoplasms (C00-C97)	270	4.8	4.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	239	4.3	3.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	141	2.5	2.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	108	1.9	1.6
7	Influenza and pneumonia (J10-J18)	33	0.6	0.5
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	0.4	0.3
9	Diabetes mellitus (E10-E14)	19	0.3	*
10	Chronic lower respiratory diseases (J40-J47)	17	0.3	*
11	Septicemia (A40-A41)	16	0.3	*
12	Cerebrovascular diseases (I60-I69)	14	0.3	*
13	Pneumonitis due to solids and liquids (J69)	8	0.1	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.1	*
15	Meningococcal infection (A39)	7	0.1	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.1	*
...	All other causes (Residual)	578	10.4	8.5
White, non-Hispanic, male, 20-24 years				
...	All causes	8,699	100.0	130.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,808	55.3	71.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,538	17.7	23.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	400	4.6	6.0
4	Malignant neoplasms (C00-C97)	358	4.1	5.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	281	3.2	4.2
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	102	1.2	1.5
7	Influenza and pneumonia (J10-J18)	49	0.6	0.7
8	Diabetes mellitus (E10-E14)	45	0.5	0.7
9	Septicemia (A40-A41)	30	0.3	0.4
10	Cerebrovascular diseases (I60-I69)	26	0.3	0.4
11	Chronic lower respiratory diseases (J40-J47)	23	0.3	0.3
12	Legal intervention (Y35,Y89.0)	20	0.2	0.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	0.2	*
15	Aortic aneurysm and dissection (I71)	11	0.1	*
...	All other causes (Residual)	973	11.2	14.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 25-29 years				
...	All causes	7,765	100.0	128.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,687	47.5	60.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,403	18.1	23.2
3	Malignant neoplasms (C00-C97)	450	5.8	7.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	431	5.6	7.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	355	4.6	5.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	81	1.0	1.3
7	Diabetes mellitus (E10-E14)	54	0.7	0.9
8	Cerebrovascular diseases (I60-I69)	51	0.7	0.8
9	Influenza and pneumonia (J10-J18)	47	0.6	0.8
10	Human immunodeficiency virus (HIV) disease (B20-B24)	44	0.6	0.7
11	Septicemia (A40-A41)	35	0.5	0.6
12	Chronic lower respiratory diseases (J40-J47)	26	0.3	0.4
12	Chronic liver disease and cirrhosis (K70,K73-K74)	26	0.3	0.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	0.3	0.4
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	0.3	0.4
...	All other causes (Residual)	1,029	13.3	17.0
White, non-Hispanic, male, 30-34 years				
...	All causes	8,601	100.0	140.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,279	38.1	53.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,592	18.5	25.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	775	9.0	12.6
4	Malignant neoplasms (C00-C97)	653	7.6	10.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	309	3.6	5.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	151	1.8	2.5
7	Diabetes mellitus (E10-E14)	135	1.6	2.2
8	Cerebrovascular diseases (I60-I69)	96	1.1	1.6
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	82	1.0	1.3
10	Influenza and pneumonia (J10-J18)	70	0.8	1.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	66	0.8	1.1
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	44	0.5	0.7
13	Septicemia (A40-A41)	39	0.5	0.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	32	0.4	0.5
15	Aortic aneurysm and dissection (I71)	31	0.4	0.5
15	Chronic lower respiratory diseases (J40-J47)	31	0.4	0.5
...	All other causes (Residual)	1,216	14.1	19.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 35-39 years				
...	All causes	12,443	100.0	182.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,687	29.6	53.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,816	14.6	26.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,750	14.1	25.6
4	Malignant neoplasms (C00-C97)	1,256	10.1	18.4
5	Human immunodeficiency virus (HIV) disease (B20-B24)	385	3.1	5.6
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	308	2.5	4.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	290	2.3	4.2
8	Diabetes mellitus (E10-E14)	251	2.0	3.7
9	Cerebrovascular diseases (I60-I69)	171	1.4	2.5
10	Influenza and pneumonia (J10-J18)	113	0.9	1.7
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	90	0.7	1.3
12	Septicemia (A40-A41)	79	0.6	1.2
13	Chronic lower respiratory diseases (J40-J47)	74	0.6	1.1
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	68	0.5	1.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	52	0.4	0.8
...	All other causes (Residual)	2,053	16.5	30.0
White, non-Hispanic, male, 40-44 years				
...	All causes	22,276	100.0	283.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,898	22.0	62.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,170	18.7	53.0
3	Malignant neoplasms (C00-C97)	3,224	14.5	41.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,348	10.5	29.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	877	3.9	11.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	737	3.3	9.4
7	Diabetes mellitus (E10-E14)	509	2.3	6.5
8	Cerebrovascular diseases (I60-I69)	410	1.8	5.2
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	402	1.8	5.1
10	Influenza and pneumonia (J10-J18)	221	1.0	2.8
11	Chronic lower respiratory diseases (J40-J47)	200	0.9	2.5
12	Septicemia (A40-A41)	162	0.7	2.1
12	Viral hepatitis (B15-B19)	162	0.7	2.1
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	133	0.6	1.7
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	105	0.5	1.3
...	All other causes (Residual)	3,718	16.7	47.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 45-49 years				
...	All causes	33,926	100.0	420.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,719	22.8	95.7
2	Malignant neoplasms (C00-C97)	6,662	19.6	82.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,238	15.4	65.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,471	7.3	30.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,607	4.7	19.9
6	Diabetes mellitus (E10-E14)	879	2.6	10.9
7	Cerebrovascular diseases (I60-I69)	711	2.1	8.8
8	Human immunodeficiency virus (HIV) disease (B20-B24)	650	1.9	8.1
9	Chronic lower respiratory diseases (J40-J47)	532	1.6	6.6
10	Viral hepatitis (B15-B19)	494	1.5	6.1
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	392	1.2	4.9
12	Influenza and pneumonia (J10-J18)	363	1.1	4.5
13	Septicemia (A40-A41)	291	0.9	3.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	265	0.8	3.3
15	Aortic aneurysm and dissection (I71)	163	0.5	2.0
...	All other causes (Residual)	5,489	16.2	68.1
White, non-Hispanic, male, 50-54 years				
...	All causes	45,934	100.0	622.6
1	Malignant neoplasms (C00-C97)	11,912	25.9	161.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,766	25.6	159.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,324	9.4	58.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	2,192	4.8	29.7
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,185	4.8	29.6
6	Diabetes mellitus (E10-E14)	1,300	2.8	17.6
7	Cerebrovascular diseases (I60-I69)	1,157	2.5	15.7
8	Chronic lower respiratory diseases (J40-J47)	1,030	2.2	14.0
9	Viral hepatitis (B15-B19)	693	1.5	9.4
10	Human immunodeficiency virus (HIV) disease (B20-B24)	527	1.1	7.1
11	Influenza and pneumonia (J10-J18)	484	1.1	6.6
12	Septicemia (A40-A41)	417	0.9	5.7
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	391	0.9	5.3
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	272	0.6	3.7
15	Aortic aneurysm and dissection (I71)	207	0.5	2.8
...	All other causes (Residual)	7,077	15.4	95.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 55-59 years				
...	All causes	57,313	100.0	872.9
1	Malignant neoplasms (C00-C97)	18,330	32.0	279.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,485	27.0	235.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,234	5.6	49.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	2,131	3.7	32.5
5	Diabetes mellitus (E10-E14)	2,011	3.5	30.6
6	Chronic lower respiratory diseases (J40-J47)	1,981	3.5	30.2
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,706	3.0	26.0
8	Cerebrovascular diseases (I60-I69)	1,534	2.7	23.4
9	Septicemia (A40-A41)	630	1.1	9.6
10	Influenza and pneumonia (J10-J18)	600	1.0	9.1
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	554	1.0	8.4
12	Viral hepatitis (B15-B19)	457	0.8	7.0
13	Aortic aneurysm and dissection (I71)	344	0.6	5.2
14	Human immunodeficiency virus (HIV) disease (B20-B24)	307	0.5	4.7
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	298	0.5	4.5
...	All other causes (Residual)	7,711	13.5	117.4
White, non-Hispanic, male, 60-64 years				
...	All causes	66,916	100.0	1,354.3
1	Malignant neoplasms (C00-C97)	23,798	35.6	481.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	18,226	27.2	368.9
3	Chronic lower respiratory diseases (J40-J47)	3,648	5.5	73.8
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,415	3.6	48.9
5	Diabetes mellitus (E10-E14)	2,373	3.5	48.0
6	Cerebrovascular diseases (I60-I69)	2,000	3.0	40.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,607	2.4	32.5
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,247	1.9	25.2
9	Influenza and pneumonia (J10-J18)	799	1.2	16.2
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	791	1.2	16.0
11	Septicemia (A40-A41)	786	1.2	15.9
12	Aortic aneurysm and dissection (I71)	505	0.8	10.2
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	360	0.5	7.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	354	0.5	7.2
15	Pneumonitis due to solids and liquids (J69)	232	0.3	4.7
...	All other causes (Residual)	7,775	11.6	157.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 65-69 years				
...	All causes	76,971	100.0	2,042.8
1	Malignant neoplasms (C00-C97)	28,240	36.7	749.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,076	26.1	532.8
3	Chronic lower respiratory diseases (J40-J47)	5,512	7.2	146.3
4	Cerebrovascular diseases (I60-I69)	2,793	3.6	74.1
5	Diabetes mellitus (E10-E14)	2,677	3.5	71.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,008	2.6	53.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,206	1.6	32.0
8	Influenza and pneumonia (J10-J18)	1,029	1.3	27.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,007	1.3	26.7
10	Septicemia (A40-A41)	953	1.2	25.3
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	901	1.2	23.9
12	Aortic aneurysm and dissection (I71)	669	0.9	17.8
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	475	0.6	12.6
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	435	0.6	11.5
15	Parkinson's disease (G20-G21)	386	0.5	10.2
...	All other causes (Residual)	8,604	11.2	228.3
White, non-Hispanic, male, 70-74 years				
...	All causes	100,546	100.0	3,242.9
1	Malignant neoplasms (C00-C97)	34,288	34.1	1,105.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	26,462	26.3	853.5
3	Chronic lower respiratory diseases (J40-J47)	8,138	8.1	262.5
4	Cerebrovascular diseases (I60-I69)	4,239	4.2	136.7
5	Diabetes mellitus (E10-E14)	3,264	3.2	105.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,224	2.2	71.7
7	Influenza and pneumonia (J10-J18)	1,724	1.7	55.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,618	1.6	52.2
9	Septicemia (A40-A41)	1,296	1.3	41.8
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,142	1.1	36.8
11	Parkinson's disease (G20-G21)	1,006	1.0	32.4
12	Alzheimer's disease (G30)	1,003	1.0	32.3
13	Aortic aneurysm and dissection (I71)	990	1.0	31.9
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	884	0.9	28.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	668	0.7	21.5
...	All other causes (Residual)	11,600	11.5	374.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 75-79 years				
...	All causes	133,639	100.0	5,122.3
1	Malignant neoplasms (C00-C97)	39,102	29.3	1,498.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	36,844	27.6	1,412.2
3	Chronic lower respiratory diseases (J40-J47)	10,975	8.2	420.7
4	Cerebrovascular diseases (I60-I69)	6,994	5.2	268.1
5	Diabetes mellitus (E10-E14)	4,092	3.1	156.8
6	Influenza and pneumonia (J10-J18)	3,074	2.3	117.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,905	2.2	111.3
8	Alzheimer's disease (G30)	2,587	1.9	99.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,486	1.9	95.3
10	Parkinson's disease (G20-G21)	2,118	1.6	81.2
11	Septicemia (A40-A41)	1,749	1.3	67.0
12	Aortic aneurysm and dissection (I71)	1,197	0.9	45.9
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,006	0.8	38.6
14	Pneumonitis due to solids and liquids (J69)	1,000	0.7	38.3
15	Chronic liver disease and cirrhosis (K70,K73-K74)	931	0.7	35.7
...	All other causes (Residual)	16,579	12.4	635.5
White, non-Hispanic, male, 80-84 years				
...	All causes	151,969	100.0	8,261.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	45,420	29.9	2,469.1
2	Malignant neoplasms (C00-C97)	35,354	23.3	1,921.9
3	Chronic lower respiratory diseases (J40-J47)	10,998	7.2	597.9
4	Cerebrovascular diseases (I60-I69)	9,385	6.2	510.2
5	Alzheimer's disease (G30)	4,739	3.1	257.6
6	Influenza and pneumonia (J10-J18)	4,693	3.1	255.1
7	Diabetes mellitus (E10-E14)	4,210	2.8	228.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,406	2.2	185.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,316	2.2	180.3
10	Parkinson's disease (G20-G21)	3,046	2.0	165.6
11	Septicemia (A40-A41)	2,086	1.4	113.4
12	Pneumonitis due to solids and liquids (J69)	1,611	1.1	87.6
13	Aortic aneurysm and dissection (I71)	1,275	0.8	69.3
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,250	0.8	68.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,159	0.8	63.0
...	All other causes (Residual)	20,021	13.2	1,088.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 85-89 years				
...	All causes	123,778	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	41,168	33.3	---
2	Malignant neoplasms (C00-C97)	21,769	17.6	---
3	Cerebrovascular diseases (I60-I69)	8,457	6.8	---
4	Chronic lower respiratory diseases (J40-J47)	7,519	6.1	---
5	Alzheimer's disease (G30)	5,209	4.2	---
6	Influenza and pneumonia (J10-J18)	4,777	3.9	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,035	2.5	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,025	2.4	---
9	Diabetes mellitus (E10-E14)	2,979	2.4	---
10	Parkinson's disease (G20-G21)	2,279	1.8	---
11	Pneumonitis due to solids and liquids (J69)	1,770	1.4	---
12	Septicemia (A40-A41)	1,644	1.3	---
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,212	1.0	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	899	0.7	---
15	Atherosclerosis (I70)	866	0.7	---
...	All other causes (Residual)	17,170	13.9	---
White, non-Hispanic, male, 90-94 years				
...	All causes	66,543	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	24,350	36.6	---
2	Malignant neoplasms (C00-C97)	8,619	13.0	---
3	Cerebrovascular diseases (I60-I69)	4,619	6.9	---
4	Chronic lower respiratory diseases (J40-J47)	3,408	5.1	---
5	Influenza and pneumonia (J10-J18)	3,401	5.1	---
6	Alzheimer's disease (G30)	3,102	4.7	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,631	2.5	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,497	2.2	---
9	Diabetes mellitus (E10-E14)	1,272	1.9	---
10	Pneumonitis due to solids and liquids (J69)	1,132	1.7	---
11	Parkinson's disease (G20-G21)	939	1.4	---
12	Septicemia (A40-A41)	783	1.2	---
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	730	1.1	---
14	Atherosclerosis (I70)	653	1.0	---
15	Aortic aneurysm and dissection (I71)	387	0.6	---
...	All other causes (Residual)	10,020	15.1	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 95-99 years				
...	All causes	18,092	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,049	39.0	---
2	Malignant neoplasms (C00-C97)	1,747	9.7	---
3	Cerebrovascular diseases (I60-I69)	1,169	6.5	---
4	Influenza and pneumonia (J10-J18)	1,125	6.2	---
5	Alzheimer's disease (G30)	828	4.6	---
6	Chronic lower respiratory diseases (J40-J47)	792	4.4	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	459	2.5	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	438	2.4	---
9	Pneumonitis due to solids and liquids (J69)	304	1.7	---
10	Diabetes mellitus (E10-E14)	266	1.5	---
11	Atherosclerosis (I70)	226	1.2	---
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	220	1.2	---
13	Septicemia (A40-A41)	203	1.1	---
14	Parkinson's disease (G20-G21)	166	0.9	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	76	0.4	---
...	All other causes (Residual)	3,024	16.7	---
White, non-Hispanic, male, 100 years and over				
...	All causes	2,382	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	988	41.5	---
2	Influenza and pneumonia (J10-J18)	188	7.9	---
3	Malignant neoplasms (C00-C97)	143	6.0	---
4	Cerebrovascular diseases (I60-I69)	133	5.6	---
5	Alzheimer's disease (G30)	80	3.4	---
6	Chronic lower respiratory diseases (J40-J47)	78	3.3	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	64	2.7	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	54	2.3	---
9	Pneumonitis due to solids and liquids (J69)	49	2.1	---
10	Atherosclerosis (I70)	42	1.8	---
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	39	1.6	---
12	Septicemia (A40-A41)	20	0.8	---
12	Diabetes mellitus (E10-E14)	20	0.8	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	0.4	---
15	Nutritional deficiencies (E40-E64)	9	0.4	---
15	Parkinson's disease (G20-G21)	9	0.4	---
...	All other causes (Residual)	456	19.1	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, all ages				
...	All causes	1,012,740	100.0	992.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	271,790	26.8	266.4
2	Malignant neoplasms (C00-C97)	219,474	21.7	215.1
3	Cerebrovascular diseases (I60-I69)	71,036	7.0	69.6
4	Chronic lower respiratory diseases (J40-J47)	62,322	6.2	61.1
5	Alzheimer's disease (G30)	45,750	4.5	44.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	32,949	3.3	32.3
7	Influenza and pneumonia (J10-J18)	29,528	2.9	28.9
8	Diabetes mellitus (E10-E14)	26,794	2.6	26.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16,539	1.6	16.2
10	Septicemia (A40-A41)	14,159	1.4	13.9
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	11,475	1.1	11.2
12	Parkinson's disease (G20-G21)	7,535	0.7	7.4
13	Chronic liver disease and cirrhosis (K70,K73-K74)	7,340	0.7	7.2
14	Pneumonitis due to solids and liquids (J69)	7,309	0.7	7.2
15	Atherosclerosis (I70)	6,523	0.6	6.4
...	All other causes (Residual)	182,217	18.0	178.6
White, non-Hispanic, female, 1-4 years				
...	All causes	996	100.0	22.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	341	34.2	7.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	131	13.2	2.9
3	Malignant neoplasms (C00-C97)	97	9.7	2.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	64	6.4	1.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	32	3.2	0.7
6	Influenza and pneumonia (J10-J18)	23	2.3	0.5
7	Septicemia (A40-A41)	18	1.8	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	1.1	*
8	Chronic lower respiratory diseases (J40-J47)	11	1.1	*
8	Certain conditions originating in the perinatal period (P00-P96)	11	1.1	*
11	Cerebrovascular diseases (I60-I69)	10	1.0	*
12	Meningococcal infection (A39)	5	0.5	*
13	Anemias (D50-D64)	4	0.4	*
13	Acute bronchitis and bronchiolitis (J20-J21)	4	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.3	*
...	All other causes (Residual)	231	23.2	5.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 5-9 years				
...	All causes	701	100.0	12.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	242	34.5	4.3
2	Malignant neoplasms (C00-C97)	133	19.0	2.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	60	8.6	1.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	25	3.6	0.4
5	Influenza and pneumonia (J10-J18)	18	2.6	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	16	2.3	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	1.9	*
7	Cerebrovascular diseases (I60-I69)	13	1.9	*
9	Chronic lower respiratory diseases (J40-J47)	8	1.1	*
10	Septicemia (A40-A41)	7	1.0	*
11	Meningitis (G00,G03)	6	0.9	*
12	Certain conditions originating in the perinatal period (P00-P96)	4	0.6	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.4	*
14	Anemias (D50-D64)	2	0.3	*
14	Diabetes mellitus (E10-E14)	2	0.3	*
14	Nutritional deficiencies (E40-E64)	2	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.3	*
...	All other causes (Residual)	145	20.7	2.6
White, non-Hispanic, female, 10-14 years				
...	All causes	817	100.0	13.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	295	36.1	4.8
2	Malignant neoplasms (C00-C97)	131	16.0	2.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	46	5.6	0.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	37	4.5	0.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	34	4.2	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	32	3.9	0.5
7	Influenza and pneumonia (J10-J18)	20	2.4	0.3
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	1.5	*
9	Septicemia (A40-A41)	11	1.3	*
9	Diabetes mellitus (E10-E14)	11	1.3	*
11	Cerebrovascular diseases (I60-I69)	6	0.7	*
11	Chronic lower respiratory diseases (J40-J47)	6	0.7	*
11	Certain conditions originating in the perinatal period (P00-P96)	6	0.7	*
14	Anemias (D50-D64)	4	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.5	*
...	All other causes (Residual)	162	19.8	2.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 15-19 years				
...	All causes	2,421	100.0	37.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,402	57.9	21.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	211	8.7	3.3
3	Malignant neoplasms (C00-C97)	182	7.5	2.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	96	4.0	1.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	75	3.1	1.2
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	44	1.8	0.7
7	Cerebrovascular diseases (I60-I69)	21	0.9	0.3
8	Diabetes mellitus (E10-E14)	18	0.7	*
9	Septicemia (A40-A41)	14	0.6	*
9	Influenza and pneumonia (J10-J18)	14	0.6	*
11	Chronic lower respiratory diseases (J40-J47)	11	0.5	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	0.4	*
12	Pregnancy, childbirth and the puerperium (O00-099)	10	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.4	*
15	Meningococcal infection (A39)	7	0.3	*
15	Pneumonitis due to solids and liquids (J69)	7	0.3	*
...	All other causes (Residual)	290	12.0	4.5
White, non-Hispanic, female, 20-24 years				
...	All causes	2,995	100.0	46.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,367	45.6	21.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	294	9.8	4.6
3	Malignant neoplasms (C00-C97)	238	7.9	3.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	160	5.3	2.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	107	3.6	1.7
6	Pregnancy, childbirth and the puerperium (O00-099)	66	2.2	1.0
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	59	2.0	0.9
8	Cerebrovascular diseases (I60-I69)	31	1.0	0.5
9	Diabetes mellitus (E10-E14)	28	0.9	0.4
10	Influenza and pneumonia (J10-J18)	27	0.9	0.4
11	Septicemia (A40-A41)	17	0.6	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	0.5	*
13	Chronic lower respiratory diseases (J40-J47)	13	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	0.4	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	8	0.3	*
15	Pneumonitis due to solids and liquids (J69)	8	0.3	*
...	All other causes (Residual)	546	18.2	8.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 25-29 years				
...	All causes	3,118	100.0	52.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,138	36.5	19.1
2	Malignant neoplasms (C00-C97)	326	10.5	5.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	295	9.5	5.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	169	5.4	2.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	161	5.2	2.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	64	2.1	1.1
7	Pregnancy, childbirth and the puerperium (O00-099)	56	1.8	0.9
8	Influenza and pneumonia (J10-J18)	46	1.5	0.8
9	Diabetes mellitus (E10-E14)	40	1.3	0.7
9	Cerebrovascular diseases (I60-I69)	40	1.3	0.7
11	Septicemia (A40-A41)	31	1.0	0.5
12	Chronic lower respiratory diseases (J40-J47)	28	0.9	0.5
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	28	0.9	0.5
14	Human immunodeficiency virus (HIV) disease (B20-B24)	21	0.7	0.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	0.6	*
...	All other causes (Residual)	657	21.1	11.0
White, non-Hispanic, female, 30-34 years				
...	All causes	4,342	100.0	71.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,243	28.6	20.5
2	Malignant neoplasms (C00-C97)	746	17.2	12.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	418	9.6	6.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	367	8.5	6.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	148	3.4	2.4
6	Cerebrovascular diseases (I60-I69)	87	2.0	1.4
7	Diabetes mellitus (E10-E14)	77	1.8	1.3
8	Pregnancy, childbirth and the puerperium (O00-099)	65	1.5	1.1
9	Human immunodeficiency virus (HIV) disease (B20-B24)	61	1.4	1.0
9	Influenza and pneumonia (J10-J18)	61	1.4	1.0
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	60	1.4	1.0
12	Septicemia (A40-A41)	52	1.2	0.9
13	Chronic liver disease and cirrhosis (K70,K73-K74)	51	1.2	0.8
14	Chronic lower respiratory diseases (J40-J47)	44	1.0	0.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	32	0.7	0.5
...	All other causes (Residual)	830	19.1	13.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 35-39 years				
...	All causes	7,061	100.0	104.4
1	Malignant neoplasms (C00-C97)	1,634	23.1	24.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,524	21.6	22.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	720	10.2	10.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	522	7.4	7.7
5	Diabetes mellitus (E10-E14)	194	2.7	2.9
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	182	2.6	2.7
7	Cerebrovascular diseases (I60-I69)	179	2.5	2.6
8	Chronic liver disease and cirrhosis (K70,K73-K74)	169	2.4	2.5
9	Septicemia (A40-A41)	93	1.3	1.4
10	Chronic lower respiratory diseases (J40-J47)	90	1.3	1.3
11	Human immunodeficiency virus (HIV) disease (B20-B24)	84	1.2	1.2
12	Influenza and pneumonia (J10-J18)	83	1.2	1.2
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	79	1.1	1.2
14	Pregnancy, childbirth and the puerperium (O00-099)	58	0.8	0.9
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	53	0.8	0.8
...	All other causes (Residual)	1,397	19.8	20.7
White, non-Hispanic, female, 40-44 years				
...	All causes	13,006	100.0	165.0
1	Malignant neoplasms (C00-C97)	3,862	29.7	49.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,143	16.5	27.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,513	11.6	19.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	748	5.8	9.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	478	3.7	6.1
6	Cerebrovascular diseases (I60-I69)	398	3.1	5.0
7	Diabetes mellitus (E10-E14)	261	2.0	3.3
8	Chronic lower respiratory diseases (J40-J47)	229	1.8	2.9
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	207	1.6	2.6
10	Influenza and pneumonia (J10-J18)	176	1.4	2.2
11	Septicemia (A40-A41)	146	1.1	1.9
11	Human immunodeficiency virus (HIV) disease (B20-B24)	146	1.1	1.9
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	107	0.8	1.4
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	79	0.6	1.0
15	Viral hepatitis (B15-B19)	76	0.6	1.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	76	0.6	1.0
...	All other causes (Residual)	2,361	18.2	29.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 45-49 years				
...	All causes	20,183	100.0	248.0
1	Malignant neoplasms (C00-C97)	7,134	35.3	87.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,711	13.4	33.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,307	11.4	28.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	842	4.2	10.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	761	3.8	9.3
6	Cerebrovascular diseases (I60-I69)	672	3.3	8.3
7	Chronic lower respiratory diseases (J40-J47)	552	2.7	6.8
8	Diabetes mellitus (E10-E14)	452	2.2	5.6
9	Influenza and pneumonia (J10-J18)	263	1.3	3.2
10	Septicemia (A40-A41)	242	1.2	3.0
11	Viral hepatitis (B15-B19)	175	0.9	2.1
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	163	0.8	2.0
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	157	0.8	1.9
14	Human immunodeficiency virus (HIV) disease (B20-B24)	124	0.6	1.5
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	116	0.6	1.4
...	All other causes (Residual)	3,512	17.4	43.1
White, non-Hispanic, female, 50-54 years				
...	All causes	26,626	100.0	353.5
1	Malignant neoplasms (C00-C97)	10,725	40.3	142.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,939	14.8	52.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,788	6.7	23.7
4	Chronic lower respiratory diseases (J40-J47)	940	3.5	12.5
5	Cerebrovascular diseases (I60-I69)	921	3.5	12.2
6	Diabetes mellitus (E10-E14)	823	3.1	10.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	763	2.9	10.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	710	2.7	9.4
9	Septicemia (A40-A41)	392	1.5	5.2
10	Influenza and pneumonia (J10-J18)	339	1.3	4.5
11	Viral hepatitis (B15-B19)	240	0.9	3.2
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	240	0.9	3.2
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	150	0.6	2.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	134	0.5	1.8
15	Human immunodeficiency virus (HIV) disease (B20-B24)	107	0.4	1.4
...	All other causes (Residual)	4,415	16.6	58.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 55-59 years				
...	All causes	36,138	100.0	531.6
1	Malignant neoplasms (C00-C97)	15,700	43.4	231.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,821	16.1	85.6
3	Chronic lower respiratory diseases (J40-J47)	1,950	5.4	28.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,446	4.0	21.3
5	Diabetes mellitus (E10-E14)	1,267	3.5	18.6
6	Cerebrovascular diseases (I60-I69)	1,199	3.3	17.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	805	2.2	11.8
8	Septicemia (A40-A41)	574	1.6	8.4
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	535	1.5	7.9
10	Influenza and pneumonia (J10-J18)	460	1.3	6.8
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	424	1.2	6.2
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	212	0.6	3.1
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	158	0.4	2.3
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	156	0.4	2.3
15	Viral hepatitis (B15-B19)	139	0.4	2.0
...	All other causes (Residual)	5,292	14.6	77.9
White, non-Hispanic, female, 60-64 years				
...	All causes	45,711	100.0	864.3
1	Malignant neoplasms (C00-C97)	19,789	43.3	374.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,212	18.0	155.3
3	Chronic lower respiratory diseases (J40-J47)	3,479	7.6	65.8
4	Diabetes mellitus (E10-E14)	1,642	3.6	31.0
5	Cerebrovascular diseases (I60-I69)	1,618	3.5	30.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,208	2.6	22.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	715	1.6	13.5
8	Septicemia (A40-A41)	651	1.4	12.3
9	Influenza and pneumonia (J10-J18)	598	1.3	11.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	548	1.2	10.4
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	334	0.7	6.3
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	245	0.5	4.6
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	223	0.5	4.2
14	Alzheimer's disease (G30)	208	0.5	3.9
15	Aortic aneurysm and dissection (I71)	198	0.4	3.7
...	All other causes (Residual)	6,043	13.2	114.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 65-69 years				
...	All causes	56,712	100.0	1,346.2
1	Malignant neoplasms (C00-C97)	22,489	39.7	533.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,009	19.4	261.3
3	Chronic lower respiratory diseases (J40-J47)	5,182	9.1	123.0
4	Cerebrovascular diseases (I60-I69)	2,435	4.3	57.8
5	Diabetes mellitus (E10-E14)	2,070	3.7	49.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,179	2.1	28.0
7	Septicemia (A40-A41)	877	1.5	20.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	849	1.5	20.2
9	Influenza and pneumonia (J10-J18)	807	1.4	19.2
10	Chronic liver disease and cirrhosis (K70,K73-K74)	744	1.3	17.7
11	Alzheimer's disease (G30)	439	0.8	10.4
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	377	0.7	8.9
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	321	0.6	7.6
14	Aortic aneurysm and dissection (I71)	304	0.5	7.2
15	Pneumonitis due to solids and liquids (J69)	198	0.3	4.7
...	All other causes (Residual)	7,432	13.1	176.4
White, non-Hispanic, female, 70-74 years				
...	All causes	80,729	100.0	2,162.2
1	Malignant neoplasms (C00-C97)	27,923	34.6	747.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	17,169	21.3	459.9
3	Chronic lower respiratory diseases (J40-J47)	7,992	9.9	214.1
4	Cerebrovascular diseases (I60-I69)	4,207	5.2	112.7
5	Diabetes mellitus (E10-E14)	2,815	3.5	75.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,479	1.8	39.6
7	Influenza and pneumonia (J10-J18)	1,462	1.8	39.2
8	Alzheimer's disease (G30)	1,389	1.7	37.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,372	1.7	36.7
10	Septicemia (A40-A41)	1,287	1.6	34.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	837	1.0	22.4
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	621	0.8	16.6
13	Parkinson's disease (G20-G21)	517	0.6	13.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	494	0.6	13.2
15	Aortic aneurysm and dissection (I71)	480	0.6	12.9
...	All other causes (Residual)	10,685	13.2	286.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 75-79 years				
...	All causes	122,525	100.0	3,481.1
1	Malignant neoplasms (C00-C97)	33,660	27.5	956.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	29,411	24.0	835.6
3	Chronic lower respiratory diseases (J40-J47)	11,285	9.2	320.6
4	Cerebrovascular diseases (I60-I69)	8,090	6.6	229.8
5	Alzheimer's disease (G30)	4,031	3.3	114.5
6	Diabetes mellitus (E10-E14)	4,022	3.3	114.3
7	Influenza and pneumonia (J10-J18)	2,757	2.3	78.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,301	1.9	65.4
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,260	1.8	64.2
10	Septicemia (A40-A41)	1,894	1.5	53.8
11	Parkinson's disease (G20-G21)	1,239	1.0	35.2
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,204	1.0	34.2
13	Aortic aneurysm and dissection (I71)	851	0.7	24.2
14	Chronic liver disease and cirrhosis (K70,K73-K74)	831	0.7	23.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	794	0.6	22.6
...	All other causes (Residual)	17,895	14.6	508.4
White, non-Hispanic, female, 80-84 years				
...	All causes	173,579	100.0	5,892.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	48,409	27.9	1,643.3
2	Malignant neoplasms (C00-C97)	34,198	19.7	1,160.9
3	Cerebrovascular diseases (I60-I69)	14,361	8.3	487.5
4	Chronic lower respiratory diseases (J40-J47)	12,683	7.3	430.5
5	Alzheimer's disease (G30)	9,112	5.2	309.3
6	Influenza and pneumonia (J10-J18)	5,114	2.9	173.6
7	Diabetes mellitus (E10-E14)	4,969	2.9	168.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,371	1.9	114.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,195	1.8	108.5
10	Septicemia (A40-A41)	2,652	1.5	90.0
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,048	1.2	69.5
12	Parkinson's disease (G20-G21)	2,042	1.2	69.3
13	Pneumonitis due to solids and liquids (J69)	1,376	0.8	46.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,103	0.6	37.4
15	Atherosclerosis (I70)	1,083	0.6	36.8
...	All other causes (Residual)	27,863	16.1	945.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 85-89 years				
...	All causes	186,328	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	59,514	31.9	---
2	Malignant neoplasms (C00-C97)	24,466	13.1	---
3	Cerebrovascular diseases (I60-I69)	16,842	9.0	---
4	Alzheimer's disease (G30)	12,785	6.9	---
5	Chronic lower respiratory diseases (J40-J47)	10,225	5.5	---
6	Influenza and pneumonia (J10-J18)	6,763	3.6	---
7	Diabetes mellitus (E10-E14)	4,455	2.4	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,644	2.0	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,427	1.8	---
10	Septicemia (A40-A41)	2,582	1.4	---
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,547	1.4	---
12	Parkinson's disease (G20-G21)	2,000	1.1	---
13	Pneumonitis due to solids and liquids (J69)	1,760	0.9	---
14	Atherosclerosis (I70)	1,601	0.9	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,145	0.6	---
...	All other causes (Residual)	32,572	17.5	---
White, non-Hispanic, female, 90-94 years				
...	All causes	145,875	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	52,511	36.0	---
2	Cerebrovascular diseases (I60-I69)	13,448	9.2	---
3	Malignant neoplasms (C00-C97)	12,114	8.3	---
4	Alzheimer's disease (G30)	11,472	7.9	---
5	Influenza and pneumonia (J10-J18)	6,337	4.3	---
6	Chronic lower respiratory diseases (J40-J47)	5,539	3.8	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,857	2.0	---
8	Diabetes mellitus (E10-E14)	2,612	1.8	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,554	1.8	---
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,493	1.7	---
11	Septicemia (A40-A41)	1,813	1.2	---
12	Atherosclerosis (I70)	1,624	1.1	---
13	Pneumonitis due to solids and liquids (J69)	1,533	1.1	---
14	Parkinson's disease (G20-G21)	1,085	0.7	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	765	0.5	---
...	All other causes (Residual)	27,118	18.6	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 95-99 years				
...	All causes	63,498	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	24,315	38.3	---
2	Cerebrovascular diseases (I60-I69)	5,379	8.5	---
3	Alzheimer's disease (G30)	5,219	8.2	---
4	Malignant neoplasms (C00-C97)	3,432	5.4	---
5	Influenza and pneumonia (J10-J18)	3,258	5.1	---
6	Chronic lower respiratory diseases (J40-J47)	1,789	2.8	---
7	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,287	2.0	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,168	1.8	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,019	1.6	---
10	Atherosclerosis (I70)	907	1.4	---
11	Diabetes mellitus (E10-E14)	896	1.4	---
12	Pneumonitis due to solids and liquids (J69)	691	1.1	---
13	Septicemia (A40-A41)	664	1.0	---
14	Parkinson's disease (G20-G21)	274	0.4	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	249	0.4	---
...	All other causes (Residual)	12,951	20.4	---
White, non-Hispanic, female, 100 years and over				
...	All causes	13,750	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,650	41.1	---
2	Cerebrovascular diseases (I60-I69)	1,056	7.7	---
3	Alzheimer's disease (G30)	951	6.9	---
4	Influenza and pneumonia (J10-J18)	868	6.3	---
5	Malignant neoplasms (C00-C97)	480	3.5	---
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	279	2.0	---
7	Chronic lower respiratory diseases (J40-J47)	261	1.9	---
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	260	1.9	---
9	Atherosclerosis (I70)	243	1.8	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	198	1.4	---
11	Diabetes mellitus (E10-E14)	138	1.0	---
12	Pneumonitis due to solids and liquids (J69)	131	1.0	---
13	Septicemia (A40-A41)	105	0.8	---
14	Anemias (D50-D64)	70	0.5	---
15	Nutritional deficiencies (E40-E64)	43	0.3	---
...	All other causes (Residual)	3,017	21.9	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, all ages				
...	All causes	289,163	100.0	774.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	73,302	25.3	196.3
2	Malignant neoplasms (C00-C97)	62,510	21.6	167.4
3	Cerebrovascular diseases (I60-I69)	17,378	6.0	46.5
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,393	4.6	35.9
5	Diabetes mellitus (E10-E14)	12,835	4.4	34.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8,522	2.9	22.8
7	Chronic lower respiratory diseases (J40-J47)	8,136	2.8	21.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,017	2.8	21.5
9	Human immunodeficiency virus (HIV) disease (B20-B24)	6,908	2.4	18.5
10	Septicemia (A40-A41)	6,159	2.1	16.5
11	Influenza and pneumonia (J10-J18)	5,716	2.0	15.3
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	4,906	1.7	13.1
13	Certain conditions originating in the perinatal period (P00-P96)	4,837	1.7	13.0
14	Alzheimer's disease (G30)	4,579	1.6	12.3
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,495	0.9	6.7
...	All other causes (Residual)	49,470	17.1	132.5
Black, non-Hispanic, both sexes, 1-4 years				
...	All causes	1,065	100.0	42.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	316	29.7	12.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	138	13.0	5.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	104	9.8	4.1
4	Malignant neoplasms (C00-C97)	62	5.8	2.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	45	4.2	1.8
6	Chronic lower respiratory diseases (J40-J47)	30	2.8	1.2
7	Influenza and pneumonia (J10-J18)	28	2.6	1.1
8	Certain conditions originating in the perinatal period (P00-P96)	23	2.2	0.9
9	Septicemia (A40-A41)	20	1.9	0.8
10	Cerebrovascular diseases (I60-I69)	14	1.3	*
11	Anemias (D50-D64)	11	1.0	*
12	Meningitis (G00,G03)	6	0.6	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.5	*
13	Pneumonitis due to solids and liquids (J69)	5	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.3	*
...	All other causes (Residual)	255	23.9	10.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 5-9 years				
...	All causes	645	100.0	21.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	235	36.4	8.0
2	Malignant neoplasms (C00-C97)	79	12.2	2.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	44	6.8	1.5
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	36	5.6	1.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	34	5.3	1.2
6	Chronic lower respiratory diseases (J40-J47)	22	3.4	0.7
7	Cerebrovascular diseases (I60-I69)	13	2.0	*
8	Septicemia (A40-A41)	11	1.7	*
8	Influenza and pneumonia (J10-J18)	11	1.7	*
10	Anemias (D50-D64)	8	1.2	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	1.1	*
12	Meningitis (G00,G03)	3	0.5	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.3	*
13	Pneumonitis due to solids and liquids (J69)	2	0.3	*
13	Certain conditions originating in the perinatal period (P00-P96)	2	0.3	*
...	All other causes (Residual)	136	21.1	4.6
Black, non-Hispanic, both sexes, 10-14 years				
...	All causes	855	100.0	25.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	262	30.6	7.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	101	11.8	3.1
3	Malignant neoplasms (C00-C97)	86	10.1	2.6
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	44	5.1	1.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	41	4.8	1.2
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	40	4.7	1.2
7	Chronic lower respiratory diseases (J40-J47)	29	3.4	0.9
8	Diabetes mellitus (E10-E14)	12	1.4	*
9	Influenza and pneumonia (J10-J18)	11	1.3	*
10	Cerebrovascular diseases (I60-I69)	10	1.2	*
11	Septicemia (A40-A41)	9	1.1	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.8	*
12	Anemias (D50-D64)	7	0.8	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	6	0.7	*
15	Meningitis (G00,G03)	4	0.5	*
15	Diseases of appendix (K35-K38)	4	0.5	*
...	All other causes (Residual)	182	21.3	5.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 15-19 years				
...	All causes	2,793	100.0	86.2
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,120	40.1	34.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	754	27.0	23.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	145	5.2	4.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	116	4.2	3.6
5	Malignant neoplasms (C00-C97)	101	3.6	3.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	55	2.0	1.7
7	Anemias (D50-D64)	30	1.1	0.9
8	Chronic lower respiratory diseases (J40-J47)	25	0.9	0.8
9	Diabetes mellitus (E10-E14)	24	0.9	0.7
10	Human immunodeficiency virus (HIV) disease (B20-B24)	22	0.8	0.7
11	Septicemia (A40-A41)	20	0.7	0.6
11	Pregnancy, childbirth and the puerperium (O00-099)	20	0.7	0.6
13	Cerebrovascular diseases (I60-I69)	19	0.7	*
14	Influenza and pneumonia (J10-J18)	13	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	0.4	*
...	All other causes (Residual)	317	11.3	9.8
Black, non-Hispanic, both sexes, 20-24 years				
...	All causes	4,451	100.0	147.6
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,910	42.9	63.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	980	22.0	32.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	283	6.4	9.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	217	4.9	7.2
5	Malignant neoplasms (C00-C97)	158	3.5	5.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	86	1.9	2.9
7	Diabetes mellitus (E10-E14)	49	1.1	1.6
7	Chronic lower respiratory diseases (J40-J47)	49	1.1	1.6
7	Pregnancy, childbirth and the puerperium (O00-099)	49	1.1	1.6
10	Anemias (D50-D64)	47	1.1	1.6
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	44	1.0	1.5
12	Cerebrovascular diseases (I60-I69)	32	0.7	1.1
13	Septicemia (A40-A41)	24	0.5	0.8
14	Influenza and pneumonia (J10-J18)	18	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	0.4	*
...	All other causes (Residual)	488	11.0	16.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 25-29 years				
...	All causes	4,559	100.0	168.3
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,492	32.7	55.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	826	18.1	30.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	370	8.1	13.7
4	Human immunodeficiency virus (HIV) disease (B20-B24)	274	6.0	10.1
5	Malignant neoplasms (C00-C97)	256	5.6	9.4
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	249	5.5	9.2
7	Diabetes mellitus (E10-E14)	80	1.8	3.0
8	Anemias (D50-D64)	65	1.4	2.4
8	Pregnancy, childbirth and the puerperium (O00-099)	65	1.4	2.4
10	Cerebrovascular diseases (I60-I69)	64	1.4	2.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	46	1.0	1.7
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	46	1.0	1.7
13	Septicemia (A40-A41)	44	1.0	1.6
14	Chronic lower respiratory diseases (J40-J47)	34	0.7	1.3
15	Influenza and pneumonia (J10-J18)	27	0.6	1.0
...	All other causes (Residual)	621	13.6	22.9
Black, non-Hispanic, both sexes, 30-34 years				
...	All causes	5,291	100.0	204.6
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,046	19.8	40.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	928	17.5	35.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	638	12.1	24.7
4	Human immunodeficiency virus (HIV) disease (B20-B24)	550	10.4	21.3
5	Malignant neoplasms (C00-C97)	418	7.9	16.2
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	225	4.3	8.7
7	Diabetes mellitus (E10-E14)	138	2.6	5.3
8	Cerebrovascular diseases (I60-I69)	99	1.9	3.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	74	1.4	2.9
10	Anemias (D50-D64)	66	1.2	2.6
11	Septicemia (A40-A41)	62	1.2	2.4
11	Chronic lower respiratory diseases (J40-J47)	62	1.2	2.4
13	Influenza and pneumonia (J10-J18)	50	0.9	1.9
13	Pregnancy, childbirth and the puerperium (O00-099)	50	0.9	1.9
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	36	0.7	1.4
...	All other causes (Residual)	849	16.0	32.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 35-39 years				
...	All causes	6,948	100.0	263.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,142	16.4	43.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	977	14.1	37.0
3	Human immunodeficiency virus (HIV) disease (B20-B24)	943	13.6	35.7
4	Malignant neoplasms (C00-C97)	765	11.0	29.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	725	10.4	27.5
6	Diabetes mellitus (E10-E14)	221	3.2	8.4
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	197	2.8	7.5
8	Cerebrovascular diseases (I60-I69)	187	2.7	7.1
9	Septicemia (A40-A41)	99	1.4	3.8
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	98	1.4	3.7
11	Influenza and pneumonia (J10-J18)	70	1.0	2.7
11	Chronic lower respiratory diseases (J40-J47)	70	1.0	2.7
13	Anemias (D50-D64)	61	0.9	2.3
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	60	0.9	2.3
15	Chronic liver disease and cirrhosis (K70,K73-K74)	56	0.8	2.1
...	All other causes (Residual)	1,277	18.4	48.4
Black, non-Hispanic, both sexes, 40-44 years				
...	All causes	10,982	100.0	389.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,023	18.4	71.8
2	Malignant neoplasms (C00-C97)	1,756	16.0	62.3
3	Human immunodeficiency virus (HIV) disease (B20-B24)	1,355	12.3	48.1
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,184	10.8	42.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	626	5.7	22.2
6	Cerebrovascular diseases (I60-I69)	496	4.5	17.6
7	Diabetes mellitus (E10-E14)	313	2.9	11.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	218	2.0	7.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	214	1.9	7.6
10	Chronic liver disease and cirrhosis (K70,K73-K74)	178	1.6	6.3
11	Septicemia (A40-A41)	159	1.4	5.6
12	Chronic lower respiratory diseases (J40-J47)	147	1.3	5.2
13	Influenza and pneumonia (J10-J18)	144	1.3	5.1
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	134	1.2	4.8
15	Anemias (D50-D64)	83	0.8	2.9
...	All other causes (Residual)	1,952	17.8	69.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 45-49 years				
...	All causes	16,504	100.0	616.8
1	Malignant neoplasms (C00-C97)	3,540	21.4	132.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,511	21.3	131.2
3	Human immunodeficiency virus (HIV) disease (B20-B24)	1,398	8.5	52.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,349	8.2	50.4
5	Cerebrovascular diseases (I60-I69)	838	5.1	31.3
6	Diabetes mellitus (E10-E14)	567	3.4	21.2
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	471	2.9	17.6
8	Chronic liver disease and cirrhosis (K70,K73-K74)	392	2.4	14.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	333	2.0	12.4
10	Chronic lower respiratory diseases (J40-J47)	289	1.8	10.8
11	Septicemia (A40-A41)	275	1.7	10.3
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	236	1.4	8.8
13	Influenza and pneumonia (J10-J18)	216	1.3	8.1
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	167	1.0	6.2
15	Viral hepatitis (B15-B19)	130	0.8	4.9
...	All other causes (Residual)	2,792	16.9	104.3
Black, non-Hispanic, both sexes, 50-54 years				
...	All causes	20,794	100.0	935.4
1	Malignant neoplasms (C00-C97)	5,265	25.3	236.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,102	24.5	229.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,298	6.2	58.4
4	Cerebrovascular diseases (I60-I69)	1,148	5.5	51.6
5	Human immunodeficiency virus (HIV) disease (B20-B24)	1,077	5.2	48.5
6	Diabetes mellitus (E10-E14)	829	4.0	37.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	492	2.4	22.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	412	2.0	18.5
9	Chronic lower respiratory diseases (J40-J47)	408	2.0	18.4
10	Septicemia (A40-A41)	379	1.8	17.0
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	314	1.5	14.1
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	291	1.4	13.1
13	Influenza and pneumonia (J10-J18)	278	1.3	12.5
14	Viral hepatitis (B15-B19)	219	1.1	9.9
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	142	0.7	6.4
...	All other causes (Residual)	3,140	15.1	141.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 55-59 years				
...	All causes	22,417	100.0	1,281.3
1	Malignant neoplasms (C00-C97)	6,700	29.9	383.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,832	26.0	333.3
3	Cerebrovascular diseases (I60-I69)	1,178	5.3	67.3
4	Diabetes mellitus (E10-E14)	1,141	5.1	65.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	877	3.9	50.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	644	2.9	36.8
7	Human immunodeficiency virus (HIV) disease (B20-B24)	605	2.7	34.6
8	Chronic lower respiratory diseases (J40-J47)	545	2.4	31.2
9	Septicemia (A40-A41)	455	2.0	26.0
10	Chronic liver disease and cirrhosis (K70,K73-K74)	436	1.9	24.9
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	328	1.5	18.7
12	Influenza and pneumonia (J10-J18)	301	1.3	17.2
13	Viral hepatitis (B15-B19)	163	0.7	9.3
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	147	0.7	8.4
15	Aortic aneurysm and dissection (I71)	98	0.4	5.6
...	All other causes (Residual)	2,967	13.2	169.6
Black, non-Hispanic, both sexes, 60-64 years				
...	All causes	22,520	100.0	1,846.9
1	Malignant neoplasms (C00-C97)	7,135	31.7	585.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,145	27.3	504.0
3	Diabetes mellitus (E10-E14)	1,274	5.7	104.5
4	Cerebrovascular diseases (I60-I69)	1,242	5.5	101.9
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	696	3.1	57.1
6	Chronic lower respiratory diseases (J40-J47)	650	2.9	53.3
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	564	2.5	46.3
8	Septicemia (A40-A41)	479	2.1	39.3
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	356	1.6	29.2
10	Chronic liver disease and cirrhosis (K70,K73-K74)	315	1.4	25.8
11	Influenza and pneumonia (J10-J18)	302	1.3	24.8
12	Human immunodeficiency virus (HIV) disease (B20-B24)	297	1.3	24.4
13	Aortic aneurysm and dissection (I71)	109	0.5	8.9
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	97	0.4	8.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	94	0.4	7.7
...	All other causes (Residual)	2,765	12.3	226.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 65-69 years				
...	All causes	24,098	100.0	2,511.2
1	Malignant neoplasms (C00-C97)	7,656	31.8	797.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,460	26.8	673.2
3	Diabetes mellitus (E10-E14)	1,473	6.1	153.5
4	Cerebrovascular diseases (I60-I69)	1,412	5.9	147.1
5	Chronic lower respiratory diseases (J40-J47)	898	3.7	93.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	782	3.2	81.5
7	Septicemia (A40-A41)	545	2.3	56.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	465	1.9	48.5
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	435	1.8	45.3
10	Influenza and pneumonia (J10-J18)	384	1.6	40.0
11	Chronic liver disease and cirrhosis (K70,K73-K74)	195	0.8	20.3
12	Human immunodeficiency virus (HIV) disease (B20-B24)	161	0.7	16.8
13	Aortic aneurysm and dissection (I71)	131	0.5	13.7
14	Alzheimer's disease (G30)	102	0.4	10.6
15	Pneumonitis due to solids and liquids (J69)	101	0.4	10.5
...	All other causes (Residual)	2,898	12.0	302.0
Black, non-Hispanic, both sexes, 70-74 years				
...	All causes	27,068	100.0	3,572.9
1	Malignant neoplasms (C00-C97)	7,904	29.2	1,043.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,380	27.3	974.1
3	Cerebrovascular diseases (I60-I69)	1,782	6.6	235.2
4	Diabetes mellitus (E10-E14)	1,628	6.0	214.9
5	Chronic lower respiratory diseases (J40-J47)	1,100	4.1	145.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	962	3.6	127.0
7	Septicemia (A40-A41)	626	2.3	82.6
8	Influenza and pneumonia (J10-J18)	530	2.0	70.0
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	513	1.9	67.7
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	456	1.7	60.2
11	Alzheimer's disease (G30)	288	1.1	38.0
12	Chronic liver disease and cirrhosis (K70,K73-K74)	157	0.6	20.7
13	Pneumonitis due to solids and liquids (J69)	131	0.5	17.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	127	0.5	16.8
15	Aortic aneurysm and dissection (I71)	123	0.5	16.2
...	All other causes (Residual)	3,361	12.4	443.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 75-79 years				
...	All causes	30,563	100.0	5,236.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,844	28.9	1,515.4
2	Malignant neoplasms (C00-C97)	7,688	25.2	1,317.3
3	Cerebrovascular diseases (I60-I69)	2,201	7.2	377.1
4	Diabetes mellitus (E10-E14)	1,673	5.5	286.7
5	Chronic lower respiratory diseases (J40-J47)	1,305	4.3	223.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,050	3.4	179.9
7	Septicemia (A40-A41)	747	2.4	128.0
8	Influenza and pneumonia (J10-J18)	684	2.2	117.2
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	600	2.0	102.8
10	Alzheimer's disease (G30)	593	1.9	101.6
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	511	1.7	87.6
12	Pneumonitis due to solids and liquids (J69)	244	0.8	41.8
13	Aortic aneurysm and dissection (I71)	152	0.5	26.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	142	0.5	24.3
15	Parkinson's disease (G20-G21)	134	0.4	23.0
...	All other causes (Residual)	3,995	13.1	684.5
Black, non-Hispanic, both sexes, 80-84 years				
...	All causes	30,756	100.0	7,641.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,274	30.2	2,304.1
2	Malignant neoplasms (C00-C97)	6,468	21.0	1,607.0
3	Cerebrovascular diseases (I60-I69)	2,490	8.1	618.6
4	Diabetes mellitus (E10-E14)	1,532	5.0	380.6
5	Chronic lower respiratory diseases (J40-J47)	1,109	3.6	275.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,040	3.4	258.4
7	Alzheimer's disease (G30)	986	3.2	245.0
8	Septicemia (A40-A41)	830	2.7	206.2
9	Influenza and pneumonia (J10-J18)	813	2.6	202.0
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	626	2.0	155.5
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	453	1.5	112.5
12	Pneumonitis due to solids and liquids (J69)	281	0.9	69.8
13	Parkinson's disease (G20-G21)	205	0.7	50.9
14	Aortic aneurysm and dissection (I71)	155	0.5	38.5
15	Atherosclerosis (I70)	139	0.5	34.5
...	All other causes (Residual)	4,355	14.2	1,082.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 85-89 years				
...	All causes	23,772	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,484	31.5	---
2	Malignant neoplasms (C00-C97)	3,949	16.6	---
3	Cerebrovascular diseases (I60-I69)	2,038	8.6	---
4	Alzheimer's disease (G30)	1,094	4.6	---
5	Diabetes mellitus (E10-E14)	1,042	4.4	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	826	3.5	---
7	Influenza and pneumonia (J10-J18)	774	3.3	---
8	Chronic lower respiratory diseases (J40-J47)	755	3.2	---
9	Septicemia (A40-A41)	624	2.6	---
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	565	2.4	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	312	1.3	---
12	Pneumonitis due to solids and liquids (J69)	263	1.1	---
13	Atherosclerosis (I70)	147	0.6	---
14	Parkinson's disease (G20-G21)	128	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	116	0.5	---
...	All other causes (Residual)	3,655	15.4	---
Black, non-Hispanic, both sexes, 90-94 years				
...	All causes	15,851	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,308	33.5	---
2	Malignant neoplasms (C00-C97)	1,835	11.6	---
3	Cerebrovascular diseases (I60-I69)	1,400	8.8	---
4	Alzheimer's disease (G30)	915	5.8	---
5	Diabetes mellitus (E10-E14)	606	3.8	---
6	Influenza and pneumonia (J10-J18)	586	3.7	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	500	3.2	---
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	431	2.7	---
9	Chronic lower respiratory diseases (J40-J47)	406	2.6	---
10	Septicemia (A40-A41)	405	2.6	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	217	1.4	---
12	Pneumonitis due to solids and liquids (J69)	191	1.2	---
13	Atherosclerosis (I70)	122	0.8	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	58	0.4	---
14	Aortic aneurysm and dissection (I71)	58	0.4	---
...	All other causes (Residual)	2,813	17.7	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 95-99 years				
...	All causes	6,736	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,417	35.9	---
2	Malignant neoplasms (C00-C97)	578	8.6	---
3	Cerebrovascular diseases (I60-I69)	508	7.5	---
4	Alzheimer's disease (G30)	415	6.2	---
5	Influenza and pneumonia (J10-J18)	270	4.0	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	201	3.0	---
7	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	197	2.9	---
8	Diabetes mellitus (E10-E14)	185	2.7	---
9	Septicemia (A40-A41)	172	2.6	---
10	Chronic lower respiratory diseases (J40-J47)	157	2.3	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	95	1.4	---
12	Pneumonitis due to solids and liquids (J69)	91	1.4	---
13	Atherosclerosis (I70)	63	0.9	---
14	Nutritional deficiencies (E40-E64)	36	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	26	0.4	---
...	All other causes (Residual)	1,325	19.7	---
Black, non-Hispanic, both sexes, 100 years and over				
...	All causes	2,122	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	807	38.0	---
2	Cerebrovascular diseases (I60-I69)	164	7.7	---
3	Alzheimer's disease (G30)	135	6.4	---
4	Influenza and pneumonia (J10-J18)	111	5.2	---
5	Malignant neoplasms (C00-C97)	99	4.7	---
6	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	71	3.3	---
7	Septicemia (A40-A41)	47	2.2	---
7	Diabetes mellitus (E10-E14)	47	2.2	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	47	2.2	---
10	Chronic lower respiratory diseases (J40-J47)	34	1.6	---
11	Atherosclerosis (I70)	33	1.6	---
12	Pneumonitis due to solids and liquids (J69)	24	1.1	---
13	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	23	1.1	---
14	Nutritional deficiencies (E40-E64)	11	0.5	---
15	Parkinson's disease (G20-G21)	5	0.2	---
...	All other causes (Residual)	464	21.9	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, all ages				
...	All causes	147,010	100.0	825.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	35,874	24.4	201.5
2	Malignant neoplasms (C00-C97)	32,363	22.0	181.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,130	6.2	51.3
4	Cerebrovascular diseases (I60-I69)	7,448	5.1	41.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7,288	5.0	40.9
6	Diabetes mellitus (E10-E14)	5,666	3.9	31.8
7	Human immunodeficiency virus (HIV) disease (B20-B24)	4,600	3.1	25.8
8	Chronic lower respiratory diseases (J40-J47)	4,406	3.0	24.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,612	2.5	20.3
10	Septicemia (A40-A41)	2,755	1.9	15.5
11	Certain conditions originating in the perinatal period (P00-P96)	2,712	1.8	15.2
12	Influenza and pneumonia (J10-J18)	2,697	1.8	15.1
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,108	1.4	11.8
14	Chronic liver disease and cirrhosis (K70,K73-K74)	1,634	1.1	9.2
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,592	1.1	8.9
...	All other causes (Residual)	23,125	15.7	129.9
Black, non-Hispanic, male, 1-4 years				
...	All causes	606	100.0	47.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	179	29.5	14.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	88	14.5	6.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	46	7.6	3.6
4	Malignant neoplasms (C00-C97)	36	5.9	2.8
5	Chronic lower respiratory diseases (J40-J47)	21	3.5	1.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	19	3.1	*
7	Influenza and pneumonia (J10-J18)	16	2.6	*
7	Certain conditions originating in the perinatal period (P00-P96)	16	2.6	*
9	Septicemia (A40-A41)	13	2.1	*
10	Cerebrovascular diseases (I60-I69)	10	1.7	*
11	Anemias (D50-D64)	5	0.8	*
12	Meningitis (G00,G03)	3	0.5	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.5	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.3	*
14	Pneumonitis due to solids and liquids (J69)	2	0.3	*
...	All other causes (Residual)	147	24.3	11.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 5-9 years				
...	All causes	354	100.0	23.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	132	37.3	8.8
2	Malignant neoplasms (C00-C97)	46	13.0	3.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	24	6.8	1.6
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	16	4.5	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	14	4.0	*
6	Chronic lower respiratory diseases (J40-J47)	12	3.4	*
7	Septicemia (A40-A41)	9	2.5	*
8	Cerebrovascular diseases (I60-I69)	8	2.3	*
9	Influenza and pneumonia (J10-J18)	7	2.0	*
10	Anemias (D50-D64)	5	1.4	*
11	Meningitis (G00,G03)	3	0.8	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.6	*
13	Nutritional deficiencies (E40-E64)	1	0.3	*
13	Acute bronchitis and bronchiolitis (J20-J21)	1	0.3	*
13	Hernia (K40-K46)	1	0.3	*
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1	0.3	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.3	*
...	All other causes (Residual)	71	20.1	4.7
Black, non-Hispanic, male, 10-14 years				
...	All causes	531	100.0	31.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	180	33.9	10.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	69	13.0	4.1
3	Malignant neoplasms (C00-C97)	52	9.8	3.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	26	4.9	1.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	21	4.0	1.3
5	Chronic lower respiratory diseases (J40-J47)	21	4.0	1.3
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	21	4.0	1.3
8	Cerebrovascular diseases (I60-I69)	6	1.1	*
8	Influenza and pneumonia (J10-J18)	6	1.1	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	5	0.9	*
11	Septicemia (A40-A41)	4	0.8	*
11	Anemias (D50-D64)	4	0.8	*
13	Diabetes mellitus (E10-E14)	3	0.6	*
13	Meningitis (G00,G03)	3	0.6	*
13	Diseases of appendix (K35-K38)	3	0.6	*
...	All other causes (Residual)	107	20.2	6.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 15-19 years				
...	All causes	2,155	100.0	131.3
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,018	47.2	62.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	546	25.3	33.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	122	5.7	7.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	85	3.9	5.2
5	Malignant neoplasms (C00-C97)	56	2.6	3.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	36	1.7	2.2
7	Chronic lower respiratory diseases (J40-J47)	20	0.9	1.2
8	Anemias (D50-D64)	18	0.8	*
9	Diabetes mellitus (E10-E14)	17	0.8	*
10	Septicemia (A40-A41)	13	0.6	*
11	Legal intervention (Y35,Y89.0)	11	0.5	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	10	0.5	*
13	Influenza and pneumonia (J10-J18)	9	0.4	*
14	Cerebrovascular diseases (I60-I69)	8	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.2	*
...	All other causes (Residual)	182	8.4	11.1
Black, non-Hispanic, male, 20-24 years				
...	All causes	3,460	100.0	227.8
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,732	50.1	114.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	755	21.8	49.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	253	7.3	16.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	143	4.1	9.4
5	Malignant neoplasms (C00-C97)	95	2.7	6.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	47	1.4	3.1
7	Anemias (D50-D64)	27	0.8	1.8
7	Chronic lower respiratory diseases (J40-J47)	27	0.8	1.8
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	27	0.8	1.8
10	Diabetes mellitus (E10-E14)	22	0.6	1.4
11	Cerebrovascular diseases (I60-I69)	16	0.5	*
12	Influenza and pneumonia (J10-J18)	14	0.4	*
13	Legal intervention (Y35,Y89.0)	13	0.4	*
14	Septicemia (A40-A41)	11	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.2	*
...	All other causes (Residual)	271	7.8	17.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 25-29 years				
...	All causes	3,268	100.0	248.7
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,349	41.3	102.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	642	19.6	48.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	227	6.9	17.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	207	6.3	15.8
5	Human immunodeficiency virus (HIV) disease (B20-B24)	152	4.7	11.6
6	Malignant neoplasms (C00-C97)	124	3.8	9.4
7	Diabetes mellitus (E10-E14)	41	1.3	3.1
8	Cerebrovascular diseases (I60-I69)	36	1.1	2.7
9	Anemias (D50-D64)	34	1.0	2.6
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	29	0.9	2.2
11	Septicemia (A40-A41)	23	0.7	1.8
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	0.7	1.8
13	Chronic lower respiratory diseases (J40-J47)	21	0.6	1.6
14	Influenza and pneumonia (J10-J18)	15	0.5	*
15	Legal intervention (Y35,Y89.0)	12	0.4	*
...	All other causes (Residual)	333	10.2	25.3
Black, non-Hispanic, male, 30-34 years				
...	All causes	3,438	100.0	280.3
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	921	26.8	75.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	690	20.1	56.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	397	11.5	32.4
4	Human immunodeficiency virus (HIV) disease (B20-B24)	303	8.8	24.7
5	Malignant neoplasms (C00-C97)	189	5.5	15.4
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	187	5.4	15.2
7	Diabetes mellitus (E10-E14)	86	2.5	7.0
8	Cerebrovascular diseases (I60-I69)	53	1.5	4.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	37	1.1	3.0
10	Anemias (D50-D64)	35	1.0	2.9
11	Chronic lower respiratory diseases (J40-J47)	31	0.9	2.5
12	Influenza and pneumonia (J10-J18)	29	0.8	2.4
13	Legal intervention (Y35,Y89.0)	27	0.8	2.2
14	Septicemia (A40-A41)	26	0.8	2.1
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	21	0.6	1.7
...	All other causes (Residual)	406	11.8	33.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 35-39 years				
...	All causes	4,172	100.0	334.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	726	17.4	58.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	709	17.0	56.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	577	13.8	46.3
4	Human immunodeficiency virus (HIV) disease (B20-B24)	576	13.8	46.2
5	Malignant neoplasms (C00-C97)	280	6.7	22.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	156	3.7	12.5
7	Diabetes mellitus (E10-E14)	140	3.4	11.2
8	Cerebrovascular diseases (I60-I69)	92	2.2	7.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	48	1.2	3.9
10	Septicemia (A40-A41)	46	1.1	3.7
11	Influenza and pneumonia (J10-J18)	43	1.0	3.5
12	Aortic aneurysm and dissection (I71)	38	0.9	3.0
13	Anemias (D50-D64)	35	0.8	2.8
13	Chronic lower respiratory diseases (J40-J47)	35	0.8	2.8
15	Chronic liver disease and cirrhosis (K70,K73-K74)	33	0.8	2.6
...	All other causes (Residual)	638	15.3	51.2
Black, non-Hispanic, male, 40-44 years				
...	All causes	6,279	100.0	476.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,243	19.8	94.3
2	Human immunodeficiency virus (HIV) disease (B20-B24)	878	14.0	66.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	812	12.9	61.6
4	Malignant neoplasms (C00-C97)	680	10.8	51.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	498	7.9	37.8
6	Cerebrovascular diseases (I60-I69)	237	3.8	18.0
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	177	2.8	13.4
8	Diabetes mellitus (E10-E14)	167	2.7	12.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	116	1.8	8.8
10	Chronic liver disease and cirrhosis (K70,K73-K74)	108	1.7	8.2
11	Influenza and pneumonia (J10-J18)	84	1.3	6.4
12	Septicemia (A40-A41)	75	1.2	5.7
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	72	1.1	5.5
14	Chronic lower respiratory diseases (J40-J47)	68	1.1	5.2
15	Aortic aneurysm and dissection (I71)	43	0.7	3.3
...	All other causes (Residual)	1,021	16.3	77.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 45-49 years				
...	All causes	9,622	100.0	772.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,237	23.2	179.6
2	Malignant neoplasms (C00-C97)	1,619	16.8	130.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	954	9.9	76.6
4	Human immunodeficiency virus (HIV) disease (B20-B24)	951	9.9	76.4
5	Cerebrovascular diseases (I60-I69)	408	4.2	32.8
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	375	3.9	30.1
7	Diabetes mellitus (E10-E14)	329	3.4	26.4
8	Chronic liver disease and cirrhosis (K70,K73-K74)	268	2.8	21.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	203	2.1	16.3
10	Chronic lower respiratory diseases (J40-J47)	141	1.5	11.3
11	Septicemia (A40-A41)	139	1.4	11.2
12	Influenza and pneumonia (J10-J18)	132	1.4	10.6
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	130	1.4	10.4
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	127	1.3	10.2
15	Viral hepatitis (B15-B19)	86	0.9	6.9
...	All other causes (Residual)	1,523	15.8	122.3
Black, non-Hispanic, male, 50-54 years				
...	All causes	12,324	100.0	1,211.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,260	26.5	320.4
2	Malignant neoplasms (C00-C97)	2,690	21.8	264.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	943	7.7	92.7
4	Human immunodeficiency virus (HIV) disease (B20-B24)	777	6.3	76.4
5	Cerebrovascular diseases (I60-I69)	631	5.1	62.0
6	Diabetes mellitus (E10-E14)	443	3.6	43.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	331	2.7	32.5
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	249	2.0	24.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	235	1.9	23.1
10	Chronic lower respiratory diseases (J40-J47)	210	1.7	20.6
11	Septicemia (A40-A41)	205	1.7	20.2
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	180	1.5	17.7
13	Influenza and pneumonia (J10-J18)	167	1.4	16.4
14	Viral hepatitis (B15-B19)	148	1.2	14.5
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	108	0.9	10.6
...	All other causes (Residual)	1,747	14.2	171.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 55-59 years				
...	All causes	13,447	100.0	1,701.2
1	Malignant neoplasms (C00-C97)	3,720	27.7	470.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,708	27.6	469.1
3	Cerebrovascular diseases (I60-I69)	681	5.1	86.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	642	4.8	81.2
5	Diabetes mellitus (E10-E14)	632	4.7	80.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	467	3.5	59.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	373	2.8	47.2
8	Chronic liver disease and cirrhosis (K70,K73-K74)	308	2.3	39.0
9	Chronic lower respiratory diseases (J40-J47)	303	2.3	38.3
10	Septicemia (A40-A41)	248	1.8	31.4
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	197	1.5	24.9
12	Influenza and pneumonia (J10-J18)	193	1.4	24.4
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	122	0.9	15.4
14	Viral hepatitis (B15-B19)	114	0.8	14.4
15	Aortic aneurysm and dissection (I71)	69	0.5	8.7
...	All other causes (Residual)	1,670	12.4	211.3
Black, non-Hispanic, male, 60-64 years				
...	All causes	12,949	100.0	2,427.3
1	Malignant neoplasms (C00-C97)	3,975	30.7	745.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,696	28.5	692.8
3	Cerebrovascular diseases (I60-I69)	730	5.6	136.8
4	Diabetes mellitus (E10-E14)	642	5.0	120.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	429	3.3	80.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	373	2.9	69.9
7	Chronic lower respiratory diseases (J40-J47)	357	2.8	66.9
8	Septicemia (A40-A41)	237	1.8	44.4
9	Human immunodeficiency virus (HIV) disease (B20-B24)	222	1.7	41.6
10	Chronic liver disease and cirrhosis (K70,K73-K74)	220	1.7	41.2
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	198	1.5	37.1
12	Influenza and pneumonia (J10-J18)	179	1.4	33.6
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	71	0.5	13.3
14	Aortic aneurysm and dissection (I71)	65	0.5	12.2
15	Viral hepatitis (B15-B19)	57	0.4	10.7
...	All other causes (Residual)	1,498	11.6	280.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 65-69 years				
...	All causes	13,198	100.0	3,232.7
1	Malignant neoplasms (C00-C97)	4,265	32.3	1,044.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,646	27.6	893.1
3	Cerebrovascular diseases (I60-I69)	768	5.8	188.1
4	Diabetes mellitus (E10-E14)	673	5.1	164.8
5	Chronic lower respiratory diseases (J40-J47)	525	4.0	128.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	332	2.5	81.3
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	305	2.3	74.7
8	Septicemia (A40-A41)	282	2.1	69.1
9	Influenza and pneumonia (J10-J18)	224	1.7	54.9
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	218	1.7	53.4
11	Chronic liver disease and cirrhosis (K70,K73-K74)	127	1.0	31.1
12	Human immunodeficiency virus (HIV) disease (B20-B24)	123	0.9	30.1
13	Aortic aneurysm and dissection (I71)	79	0.6	19.4
14	Pneumonitis due to solids and liquids (J69)	63	0.5	15.4
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	53	0.4	13.0
...	All other causes (Residual)	1,515	11.5	371.1
Black, non-Hispanic, male, 70-74 years				
...	All causes	14,105	100.0	4,585.9
1	Malignant neoplasms (C00-C97)	4,365	30.9	1,419.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,940	27.9	1,281.0
3	Cerebrovascular diseases (I60-I69)	868	6.2	282.2
4	Diabetes mellitus (E10-E14)	740	5.2	240.6
5	Chronic lower respiratory diseases (J40-J47)	621	4.4	201.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	418	3.0	135.9
7	Septicemia (A40-A41)	304	2.2	98.8
8	Influenza and pneumonia (J10-J18)	286	2.0	93.0
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	284	2.0	92.3
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	247	1.8	80.3
11	Alzheimer's disease (G30)	114	0.8	37.1
12	Chronic liver disease and cirrhosis (K70,K73-K74)	98	0.7	31.9
13	Pneumonitis due to solids and liquids (J69)	80	0.6	26.0
14	Parkinson's disease (G20-G21)	59	0.4	19.2
15	Human immunodeficiency virus (HIV) disease (B20-B24)	56	0.4	18.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	56	0.4	18.2
...	All other causes (Residual)	1,569	11.1	510.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 75-79 years				
...	All causes	14,530	100.0	6,672.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,149	28.6	1,905.3
2	Malignant neoplasms (C00-C97)	3,989	27.5	1,831.8
3	Cerebrovascular diseases (I60-I69)	899	6.2	412.8
4	Chronic lower respiratory diseases (J40-J47)	761	5.2	349.5
5	Diabetes mellitus (E10-E14)	687	4.7	315.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	464	3.2	213.1
7	Septicemia (A40-A41)	319	2.2	146.5
8	Influenza and pneumonia (J10-J18)	318	2.2	146.0
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	283	1.9	130.0
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	277	1.9	127.2
11	Alzheimer's disease (G30)	228	1.6	104.7
12	Pneumonitis due to solids and liquids (J69)	139	1.0	63.8
13	Parkinson's disease (G20-G21)	80	0.6	36.7
14	Aortic aneurysm and dissection (I71)	75	0.5	34.4
15	Chronic liver disease and cirrhosis (K70,K73-K74)	72	0.5	33.1
...	All other causes (Residual)	1,790	12.3	822.0
Black, non-Hispanic, male, 80-84 years				
...	All causes	13,193	100.0	9,492.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,803	28.8	2,736.4
2	Malignant neoplasms (C00-C97)	3,315	25.1	2,385.2
3	Cerebrovascular diseases (I60-I69)	922	7.0	663.4
4	Chronic lower respiratory diseases (J40-J47)	599	4.5	431.0
5	Diabetes mellitus (E10-E14)	549	4.2	395.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	427	3.2	307.2
7	Influenza and pneumonia (J10-J18)	365	2.8	262.6
8	Septicemia (A40-A41)	335	2.5	241.0
9	Alzheimer's disease (G30)	317	2.4	228.1
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	235	1.8	169.1
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	219	1.7	157.6
12	Pneumonitis due to solids and liquids (J69)	150	1.1	107.9
13	Parkinson's disease (G20-G21)	116	0.9	83.5
14	Aortic aneurysm and dissection (I71)	65	0.5	46.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	61	0.5	43.9
...	All other causes (Residual)	1,715	13.0	1,234.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 85-89 years				
...	All causes	8,526	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,517	29.5	---
2	Malignant neoplasms (C00-C97)	1,850	21.7	---
3	Cerebrovascular diseases (I60-I69)	628	7.4	---
4	Chronic lower respiratory diseases (J40-J47)	382	4.5	---
5	Diabetes mellitus (E10-E14)	309	3.6	---
6	Alzheimer's disease (G30)	305	3.6	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	292	3.4	---
8	Influenza and pneumonia (J10-J18)	289	3.4	---
9	Septicemia (A40-A41)	249	2.9	---
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	168	2.0	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	133	1.6	---
12	Pneumonitis due to solids and liquids (J69)	117	1.4	---
13	Parkinson's disease (G20-G21)	68	0.8	---
14	Aortic aneurysm and dissection (I71)	39	0.5	---
15	Atherosclerosis (I70)	38	0.4	---
...	All other causes (Residual)	1,142	13.4	---
Black, non-Hispanic, male, 90-94 years				
...	All causes	4,367	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,388	31.8	---
2	Malignant neoplasms (C00-C97)	759	17.4	---
3	Cerebrovascular diseases (I60-I69)	318	7.3	---
4	Alzheimer's disease (G30)	187	4.3	---
5	Chronic lower respiratory diseases (J40-J47)	184	4.2	---
6	Influenza and pneumonia (J10-J18)	176	4.0	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	154	3.5	---
8	Diabetes mellitus (E10-E14)	142	3.3	---
9	Septicemia (A40-A41)	107	2.5	---
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	91	2.1	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	72	1.6	---
12	Pneumonitis due to solids and liquids (J69)	66	1.5	---
13	Atherosclerosis (I70)	29	0.7	---
14	Parkinson's disease (G20-G21)	26	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	0.3	---
15	Aortic aneurysm and dissection (I71)	14	0.3	---
...	All other causes (Residual)	640	14.7	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 95-99 years				
...	All causes	1,466	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	467	31.9	---
2	Malignant neoplasms (C00-C97)	219	14.9	---
3	Cerebrovascular diseases (I60-I69)	86	5.9	---
4	Influenza and pneumonia (J10-J18)	74	5.0	---
5	Chronic lower respiratory diseases (J40-J47)	55	3.8	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	53	3.6	---
7	Alzheimer's disease (G30)	49	3.3	---
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	41	2.8	---
9	Diabetes mellitus (E10-E14)	39	2.7	---
10	Septicemia (A40-A41)	36	2.5	---
11	Pneumonitis due to solids and liquids (J69)	29	2.0	---
12	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	28	1.9	---
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.6	---
13	Atherosclerosis (I70)	9	0.6	---
15	Parkinson's disease (G20-G21)	8	0.5	---
...	All other causes (Residual)	264	18.0	---
Black, non-Hispanic, male, 100 years and over				
...	All causes	331	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	123	37.2	---
2	Malignant neoplasms (C00-C97)	31	9.4	---
3	Influenza and pneumonia (J10-J18)	26	7.9	---
4	Cerebrovascular diseases (I60-I69)	17	5.1	---
5	Alzheimer's disease (G30)	13	3.9	---
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	3.9	---
7	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	9	2.7	---
8	Septicemia (A40-A41)	7	2.1	---
9	Chronic lower respiratory diseases (J40-J47)	6	1.8	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6	1.8	---
11	Diabetes mellitus (E10-E14)	5	1.5	---
11	Atherosclerosis (I70)	5	1.5	---
13	Pneumonitis due to solids and liquids (J69)	4	1.2	---
14	Hernia (K40-K46)	3	0.9	---
15	Nutritional deficiencies (E40-E64)	2	0.6	---
15	Parkinson's disease (G20-G21)	2	0.6	---
...	All other causes (Residual)	59	17.8	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, all ages				
...	All causes	142,153	100.0	727.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	37,428	26.3	191.6
2	Malignant neoplasms (C00-C97)	30,147	21.2	154.3
3	Cerebrovascular diseases (I60-I69)	9,930	7.0	50.8
4	Diabetes mellitus (E10-E14)	7,169	5.0	36.7
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,405	3.1	22.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,263	3.0	21.8
7	Chronic lower respiratory diseases (J40-J47)	3,730	2.6	19.1
8	Septicemia (A40-A41)	3,404	2.4	17.4
9	Alzheimer's disease (G30)	3,296	2.3	16.9
10	Influenza and pneumonia (J10-J18)	3,019	2.1	15.5
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,798	2.0	14.3
12	Human immunodeficiency virus (HIV) disease (B20-B24)	2,308	1.6	11.8
13	Certain conditions originating in the perinatal period (P00-P96)	2,125	1.5	10.9
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,234	0.9	6.3
15	Chronic liver disease and cirrhosis (K70,K73-K74)	861	0.6	4.4
...	All other causes (Residual)	26,036	18.3	133.3
Black, non-Hispanic, female, 1-4 years				
...	All causes	459	100.0	37.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	137	29.8	11.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	58	12.6	4.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	50	10.9	4.1
4	Malignant neoplasms (C00-C97)	26	5.7	2.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	26	5.7	2.1
6	Influenza and pneumonia (J10-J18)	12	2.6	*
7	Chronic lower respiratory diseases (J40-J47)	9	2.0	*
8	Septicemia (A40-A41)	7	1.5	*
8	Certain conditions originating in the perinatal period (P00-P96)	7	1.5	*
10	Anemias (D50-D64)	6	1.3	*
11	Cerebrovascular diseases (I60-I69)	4	0.9	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.7	*
12	Meningitis (G00,G03)	3	0.7	*
12	Pneumonitis due to solids and liquids (J69)	3	0.7	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
...	All other causes (Residual)	106	23.1	8.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 5-9 years				
...	All causes	291	100.0	20.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	103	35.4	7.1
2	Malignant neoplasms (C00-C97)	33	11.3	2.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	20	6.9	1.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	20	6.9	1.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	20	6.9	1.4
6	Chronic lower respiratory diseases (J40-J47)	10	3.4	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	1.7	*
7	Cerebrovascular diseases (I60-I69)	5	1.7	*
9	Influenza and pneumonia (J10-J18)	4	1.4	*
10	Anemias (D50-D64)	3	1.0	*
11	Septicemia (A40-A41)	2	0.7	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.7	*
11	Pneumonitis due to solids and liquids (J69)	2	0.7	*
11	Certain conditions originating in the perinatal period (P00-P96)	2	0.7	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
...	All other causes (Residual)	59	20.3	4.1
Black, non-Hispanic, female, 10-14 years				
...	All causes	324	100.0	19.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	82	25.3	5.0
2	Malignant neoplasms (C00-C97)	34	10.5	2.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	32	9.9	2.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	23	7.1	1.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	20	6.2	1.2
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	14	4.3	*
7	Diabetes mellitus (E10-E14)	9	2.8	*
8	Chronic lower respiratory diseases (J40-J47)	8	2.5	*
9	Septicemia (A40-A41)	5	1.5	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	1.5	*
9	Influenza and pneumonia (J10-J18)	5	1.5	*
12	Cerebrovascular diseases (I60-I69)	4	1.2	*
13	Anemias (D50-D64)	3	0.9	*
14	Syphilis (A50-A53)	1	0.3	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.3	*
14	Meningitis (G00,G03)	1	0.3	*
14	Diseases of appendix (K35-K38)	1	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
14	Certain conditions originating in the perinatal period (P00-P96)	1	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.3	*
...	All other causes (Residual)	73	22.5	4.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 15-19 years				
...	All causes	638	100.0	39.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	208	32.6	13.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	102	16.0	6.4
3	Malignant neoplasms (C00-C97)	45	7.1	2.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	31	4.9	1.9
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	23	3.6	1.4
6	Pregnancy, childbirth and the puerperium (000-099)	20	3.1	1.3
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	19	3.0	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	12	1.9	*
8	Anemias (D50-D64)	12	1.9	*
10	Cerebrovascular diseases (I60-I69)	11	1.7	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	1.3	*
12	Septicemia (A40-A41)	7	1.1	*
12	Diabetes mellitus (E10-E14)	7	1.1	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.8	*
14	Chronic lower respiratory diseases (J40-J47)	5	0.8	*
...	All other causes (Residual)	123	19.3	7.7
Black, non-Hispanic, female, 20-24 years				
...	All causes	991	100.0	66.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	225	22.7	15.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	178	18.0	11.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	74	7.5	4.9
4	Malignant neoplasms (C00-C97)	63	6.4	4.2
5	Pregnancy, childbirth and the puerperium (000-099)	49	4.9	3.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	39	3.9	2.6
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	30	3.0	2.0
8	Diabetes mellitus (E10-E14)	27	2.7	1.8
9	Chronic lower respiratory diseases (J40-J47)	22	2.2	1.5
10	Anemias (D50-D64)	20	2.0	1.3
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	17	1.7	*
12	Cerebrovascular diseases (I60-I69)	16	1.6	*
13	Septicemia (A40-A41)	13	1.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	1.0	*
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	4	0.4	*
15	Influenza and pneumonia (J10-J18)	4	0.4	*
...	All other causes (Residual)	200	20.2	13.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 25-29 years				
...	All causes	1,291	100.0	92.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	184	14.3	13.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	143	11.1	10.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	143	11.1	10.2
4	Malignant neoplasms (C00-C97)	132	10.2	9.5
5	Human immunodeficiency virus (HIV) disease (B20-B24)	122	9.5	8.7
6	Pregnancy, childbirth and the puerperium (O00-099)	65	5.0	4.7
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	42	3.3	3.0
8	Diabetes mellitus (E10-E14)	39	3.0	2.8
9	Anemias (D50-D64)	31	2.4	2.2
10	Cerebrovascular diseases (I60-I69)	28	2.2	2.0
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	1.8	1.6
12	Septicemia (A40-A41)	21	1.6	1.5
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	17	1.3	*
14	Chronic lower respiratory diseases (J40-J47)	13	1.0	*
15	Influenza and pneumonia (J10-J18)	12	0.9	*
...	All other causes (Residual)	276	21.4	19.8
Black, non-Hispanic, female, 30-34 years				
...	All causes	1,853	100.0	136.3
1	Human immunodeficiency virus (HIV) disease (B20-B24)	247	13.3	18.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	241	13.0	17.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	238	12.8	17.5
4	Malignant neoplasms (C00-C97)	229	12.4	16.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	125	6.7	9.2
6	Diabetes mellitus (E10-E14)	52	2.8	3.8
7	Pregnancy, childbirth and the puerperium (O00-099)	50	2.7	3.7
8	Cerebrovascular diseases (I60-I69)	46	2.5	3.4
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	38	2.1	2.8
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	37	2.0	2.7
11	Septicemia (A40-A41)	36	1.9	2.6
12	Anemias (D50-D64)	31	1.7	2.3
12	Chronic lower respiratory diseases (J40-J47)	31	1.7	2.3
14	Influenza and pneumonia (J10-J18)	21	1.1	1.5
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	21	1.1	1.5
...	All other causes (Residual)	410	22.1	30.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 35-39 years				
...	All causes	2,776	100.0	199.2
1	Malignant neoplasms (C00-C97)	485	17.5	34.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	416	15.0	29.9
3	Human immunodeficiency virus (HIV) disease (B20-B24)	367	13.2	26.3
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	268	9.7	19.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	148	5.3	10.6
6	Cerebrovascular diseases (I60-I69)	95	3.4	6.8
7	Diabetes mellitus (E10-E14)	81	2.9	5.8
8	Septicemia (A40-A41)	53	1.9	3.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	50	1.8	3.6
10	Pregnancy, childbirth and the puerperium (O00-099)	41	1.5	2.9
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	41	1.5	2.9
12	Chronic lower respiratory diseases (J40-J47)	35	1.3	2.5
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	29	1.0	2.1
14	Influenza and pneumonia (J10-J18)	27	1.0	1.9
15	Anemias (D50-D64)	26	0.9	1.9
...	All other causes (Residual)	614	22.1	44.1
Black, non-Hispanic, female, 40-44 years				
...	All causes	4,703	100.0	313.6
1	Malignant neoplasms (C00-C97)	1,076	22.9	71.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	780	16.6	52.0
3	Human immunodeficiency virus (HIV) disease (B20-B24)	477	10.1	31.8
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	372	7.9	24.8
5	Cerebrovascular diseases (I60-I69)	259	5.5	17.3
6	Diabetes mellitus (E10-E14)	146	3.1	9.7
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	128	2.7	8.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	98	2.1	6.5
9	Septicemia (A40-A41)	84	1.8	5.6
10	Chronic lower respiratory diseases (J40-J47)	79	1.7	5.3
11	Chronic liver disease and cirrhosis (K70,K73-K74)	70	1.5	4.7
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	62	1.3	4.1
13	Influenza and pneumonia (J10-J18)	60	1.3	4.0
14	Anemias (D50-D64)	41	0.9	2.7
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	41	0.9	2.7
...	All other causes (Residual)	930	19.8	62.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 45-49 years				
...	All causes	6,882	100.0	481.1
1	Malignant neoplasms (C00-C97)	1,921	27.9	134.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,274	18.5	89.1
3	Human immunodeficiency virus (HIV) disease (B20-B24)	447	6.5	31.2
4	Cerebrovascular diseases (I60-I69)	430	6.2	30.1
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	395	5.7	27.6
6	Diabetes mellitus (E10-E14)	238	3.5	16.6
7	Chronic lower respiratory diseases (J40-J47)	148	2.2	10.3
8	Septicemia (A40-A41)	136	2.0	9.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	130	1.9	9.1
10	Chronic liver disease and cirrhosis (K70,K73-K74)	124	1.8	8.7
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	109	1.6	7.6
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	96	1.4	6.7
13	Influenza and pneumonia (J10-J18)	84	1.2	5.9
14	Viral hepatitis (B15-B19)	44	0.6	3.1
15	Anemias (D50-D64)	38	0.6	2.7
...	All other causes (Residual)	1,268	18.4	88.6
Black, non-Hispanic, female, 50-54 years				
...	All causes	8,470	100.0	702.6
1	Malignant neoplasms (C00-C97)	2,575	30.4	213.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,842	21.7	152.8
3	Cerebrovascular diseases (I60-I69)	517	6.1	42.9
4	Diabetes mellitus (E10-E14)	386	4.6	32.0
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	355	4.2	29.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	300	3.5	24.9
7	Chronic lower respiratory diseases (J40-J47)	198	2.3	16.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	177	2.1	14.7
9	Septicemia (A40-A41)	174	2.1	14.4
10	Chronic liver disease and cirrhosis (K70,K73-K74)	161	1.9	13.4
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	111	1.3	9.2
11	Influenza and pneumonia (J10-J18)	111	1.3	9.2
13	Viral hepatitis (B15-B19)	71	0.8	5.9
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	65	0.8	5.4
15	Anemias (D50-D64)	39	0.5	3.2
...	All other causes (Residual)	1,388	16.4	115.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 55-59 years				
...	All causes	8,970	100.0	935.2
1	Malignant neoplasms (C00-C97)	2,980	33.2	310.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,124	23.7	221.5
3	Diabetes mellitus (E10-E14)	509	5.7	53.1
4	Cerebrovascular diseases (I60-I69)	497	5.5	51.8
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	271	3.0	28.3
6	Chronic lower respiratory diseases (J40-J47)	242	2.7	25.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	235	2.6	24.5
8	Septicemia (A40-A41)	207	2.3	21.6
9	Human immunodeficiency virus (HIV) disease (B20-B24)	138	1.5	14.4
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	131	1.5	13.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	128	1.4	13.3
12	Influenza and pneumonia (J10-J18)	108	1.2	11.3
13	Viral hepatitis (B15-B19)	49	0.5	5.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	36	0.4	3.8
15	Anemias (D50-D64)	29	0.3	3.0
15	Aortic aneurysm and dissection (I71)	29	0.3	3.0
...	All other causes (Residual)	1,257	14.0	131.1
Black, non-Hispanic, female, 60-64 years				
...	All causes	9,571	100.0	1,395.5
1	Malignant neoplasms (C00-C97)	3,160	33.0	460.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,449	25.6	357.1
3	Diabetes mellitus (E10-E14)	632	6.6	92.1
4	Cerebrovascular diseases (I60-I69)	512	5.3	74.7
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	323	3.4	47.1
6	Chronic lower respiratory diseases (J40-J47)	293	3.1	42.7
7	Septicemia (A40-A41)	242	2.5	35.3
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	158	1.7	23.0
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	135	1.4	19.7
10	Influenza and pneumonia (J10-J18)	123	1.3	17.9
11	Chronic liver disease and cirrhosis (K70,K73-K74)	95	1.0	13.9
12	Human immunodeficiency virus (HIV) disease (B20-B24)	75	0.8	10.9
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	44	0.5	6.4
13	Aortic aneurysm and dissection (I71)	44	0.5	6.4
15	Viral hepatitis (B15-B19)	33	0.3	4.8
...	All other causes (Residual)	1,253	13.1	182.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 65-69 years				
...	All causes	10,900	100.0	1,977.0
1	Malignant neoplasms (C00-C97)	3,391	31.1	615.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,814	25.8	510.4
3	Diabetes mellitus (E10-E14)	800	7.3	145.1
4	Cerebrovascular diseases (I60-I69)	644	5.9	116.8
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	450	4.1	81.6
6	Chronic lower respiratory diseases (J40-J47)	373	3.4	67.7
7	Septicemia (A40-A41)	263	2.4	47.7
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	217	2.0	39.4
9	Influenza and pneumonia (J10-J18)	160	1.5	29.0
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	160	1.5	29.0
11	Chronic liver disease and cirrhosis (K70,K73-K74)	68	0.6	12.3
12	Alzheimer's disease (G30)	53	0.5	9.6
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	52	0.5	9.4
13	Aortic aneurysm and dissection (I71)	52	0.5	9.4
15	Human immunodeficiency virus (HIV) disease (B20-B24)	38	0.3	6.9
15	Pneumonitis due to solids and liquids (J69)	38	0.3	6.9
...	All other causes (Residual)	1,327	12.2	240.7
Black, non-Hispanic, female, 70-74 years				
...	All causes	12,963	100.0	2,880.6
1	Malignant neoplasms (C00-C97)	3,539	27.3	786.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,440	26.5	764.4
3	Cerebrovascular diseases (I60-I69)	914	7.1	203.1
4	Diabetes mellitus (E10-E14)	888	6.9	197.3
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	544	4.2	120.9
6	Chronic lower respiratory diseases (J40-J47)	479	3.7	106.4
7	Septicemia (A40-A41)	322	2.5	71.6
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	266	2.1	59.1
9	Influenza and pneumonia (J10-J18)	244	1.9	54.2
10	Alzheimer's disease (G30)	174	1.3	38.7
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	172	1.3	38.2
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	71	0.5	15.8
13	Aortic aneurysm and dissection (I71)	70	0.5	15.6
14	Chronic liver disease and cirrhosis (K70,K73-K74)	59	0.5	13.1
15	Pneumonitis due to solids and liquids (J69)	51	0.4	11.3
...	All other causes (Residual)	1,730	13.3	384.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 75-79 years				
...	All causes	16,033	100.0	4,382.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,695	29.3	1,283.3
2	Malignant neoplasms (C00-C97)	3,699	23.1	1,011.1
3	Cerebrovascular diseases (I60-I69)	1,302	8.1	355.9
4	Diabetes mellitus (E10-E14)	986	6.1	269.5
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	586	3.7	160.2
6	Chronic lower respiratory diseases (J40-J47)	544	3.4	148.7
7	Septicemia (A40-A41)	428	2.7	117.0
8	Influenza and pneumonia (J10-J18)	366	2.3	100.0
9	Alzheimer's disease (G30)	365	2.3	99.8
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	323	2.0	88.3
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	228	1.4	62.3
12	Pneumonitis due to solids and liquids (J69)	105	0.7	28.7
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	80	0.5	21.9
14	Aortic aneurysm and dissection (I71)	77	0.5	21.0
15	Atherosclerosis (I70)	60	0.4	16.4
...	All other causes (Residual)	2,189	13.7	598.3
Black, non-Hispanic, female, 80-84 years				
...	All causes	17,563	100.0	6,664.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,471	31.2	2,076.1
2	Malignant neoplasms (C00-C97)	3,153	18.0	1,196.5
3	Cerebrovascular diseases (I60-I69)	1,568	8.9	595.0
4	Diabetes mellitus (E10-E14)	983	5.6	373.0
5	Alzheimer's disease (G30)	669	3.8	253.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	613	3.5	232.6
7	Chronic lower respiratory diseases (J40-J47)	510	2.9	193.5
8	Septicemia (A40-A41)	495	2.8	187.8
9	Influenza and pneumonia (J10-J18)	448	2.6	170.0
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	407	2.3	154.4
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	218	1.2	82.7
12	Pneumonitis due to solids and liquids (J69)	131	0.7	49.7
13	Aortic aneurysm and dissection (I71)	90	0.5	34.2
14	Parkinson's disease (G20-G21)	89	0.5	33.8
15	Atherosclerosis (I70)	82	0.5	31.1
...	All other causes (Residual)	2,636	15.0	1,000.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 85-89 years				
...	All causes	15,246	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,967	32.6	---
2	Malignant neoplasms (C00-C97)	2,099	13.8	---
3	Cerebrovascular diseases (I60-I69)	1,410	9.2	---
4	Alzheimer's disease (G30)	789	5.2	---
5	Diabetes mellitus (E10-E14)	733	4.8	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	534	3.5	---
7	Influenza and pneumonia (J10-J18)	485	3.2	---
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	397	2.6	---
9	Septicemia (A40-A41)	375	2.5	---
10	Chronic lower respiratory diseases (J40-J47)	373	2.4	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	179	1.2	---
12	Pneumonitis due to solids and liquids (J69)	146	1.0	---
13	Atherosclerosis (I70)	109	0.7	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	80	0.5	---
15	Aortic aneurysm and dissection (I71)	71	0.5	---
...	All other causes (Residual)	2,499	16.4	---
Black, non-Hispanic, female, 90-94 years				
...	All causes	11,484	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,920	34.1	---
2	Cerebrovascular diseases (I60-I69)	1,082	9.4	---
3	Malignant neoplasms (C00-C97)	1,076	9.4	---
4	Alzheimer's disease (G30)	728	6.3	---
5	Diabetes mellitus (E10-E14)	464	4.0	---
6	Influenza and pneumonia (J10-J18)	410	3.6	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	346	3.0	---
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	340	3.0	---
9	Septicemia (A40-A41)	298	2.6	---
10	Chronic lower respiratory diseases (J40-J47)	222	1.9	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	145	1.3	---
12	Pneumonitis due to solids and liquids (J69)	125	1.1	---
13	Atherosclerosis (I70)	93	0.8	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	44	0.4	---
14	Aortic aneurysm and dissection (I71)	44	0.4	---
...	All other causes (Residual)	2,147	18.7	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 95-99 years				
...	All causes	5,270	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,950	37.0	---
2	Cerebrovascular diseases (I60-I69)	422	8.0	---
3	Alzheimer's disease (G30)	366	6.9	---
4	Malignant neoplasms (C00-C97)	359	6.8	---
5	Influenza and pneumonia (J10-J18)	196	3.7	---
6	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	156	3.0	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	148	2.8	---
8	Diabetes mellitus (E10-E14)	146	2.8	---
9	Septicemia (A40-A41)	136	2.6	---
10	Chronic lower respiratory diseases (J40-J47)	102	1.9	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	67	1.3	---
12	Pneumonitis due to solids and liquids (J69)	62	1.2	---
13	Atherosclerosis (I70)	54	1.0	---
14	Nutritional deficiencies (E40-E64)	29	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	0.3	---
15	Anemias (D50-D64)	17	0.3	---
15	Aortic aneurysm and dissection (I71)	17	0.3	---
...	All other causes (Residual)	1,026	19.5	---
Black, non-Hispanic, female, 100 years and over				
...	All causes	1,791	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	684	38.2	---
2	Cerebrovascular diseases (I60-I69)	147	8.2	---
3	Alzheimer's disease (G30)	122	6.8	---
4	Influenza and pneumonia (J10-J18)	85	4.7	---
5	Malignant neoplasms (C00-C97)	68	3.8	---
6	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	62	3.5	---
7	Diabetes mellitus (E10-E14)	42	2.3	---
8	Septicemia (A40-A41)	40	2.2	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	34	1.9	---
10	Atherosclerosis (I70)	28	1.6	---
10	Chronic lower respiratory diseases (J40-J47)	28	1.6	---
12	Pneumonitis due to solids and liquids (J69)	20	1.1	---
13	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17	0.9	---
14	Nutritional deficiencies (E40-E64)	9	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.2	---
15	Anemias (D50-D64)	4	0.2	---
...	All other causes (Residual)	397	22.2	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available