

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, all ages				
...	All causes	144,490	100.0	286.2
1	Malignant neoplasms (C00-C97)	31,119	21.5	61.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	30,006	20.8	59.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,476	7.3	20.8
4	Cerebrovascular diseases (I60-I69)	7,274	5.0	14.4
5	Diabetes mellitus (E10-E14)	6,556	4.5	13.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	4,348	3.0	8.6
7	Chronic lower respiratory diseases (J40-J47)	4,172	2.9	8.3
8	Alzheimer's disease (G30)	3,427	2.4	6.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,252	2.3	6.4
10	Influenza and pneumonia (J09-J18)	3,025	2.1	6.0
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,890	2.0	5.7
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,661	1.8	5.3
13	Certain conditions originating in the perinatal period (P00-P96)	2,529	1.8	5.0
14	Septicemia (A40-A41)	2,035	1.4	4.0
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,786	1.2	3.5
...	All other causes (Residual)	28,934	20.0	57.3
Hispanic, both sexes, 1-4 years				
...	All causes	930	100.0	22.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	330	35.5	8.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	122	13.1	3.0
3	Malignant neoplasms (C00-C97)	75	8.1	1.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	57	6.1	1.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	37	4.0	0.9
6	Influenza and pneumonia (J09-J18)	21	2.3	0.5
7	Septicemia (A40-A41)	18	1.9	*
8	Chronic lower respiratory diseases (J40-J47)	10	1.1	*
9	Cerebrovascular diseases (I60-I69)	8	0.9	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.8	*
10	Certain conditions originating in the perinatal period (P00-P96)	7	0.8	*
12	Meningococcal infection (A39)	5	0.5	*
12	Pneumonitis due to solids and liquids (J69)	5	0.5	*
14	Acute bronchitis and bronchiolitis (J20-J21)	4	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.4	*
...	All other causes (Residual)	216	23.2	5.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 5-9 years				
...	All causes	423	100.0	8.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	117	27.7	2.4
2	Malignant neoplasms (C00-C97)	100	23.6	2.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	36	8.5	0.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	20	4.7	0.4
5	Cerebrovascular diseases (I60-I69)	12	2.8	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	11	2.6	*
7	Chronic lower respiratory diseases (J40-J47)	9	2.1	*
8	Influenza and pneumonia (J09-J18)	7	1.7	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.9	*
10	Septicemia (A40-A41)	3	0.7	*
11	Anemias (D50-D64)	2	0.5	*
11	Meningitis (G00,G03)	2	0.5	*
11	Pneumonitis due to solids and liquids (J69)	2	0.5	*
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2	0.5	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.5	*
...	All other causes (Residual)	94	22.2	2.0
Hispanic, both sexes, 10-14 years				
...	All causes	528	100.0	11.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	135	25.6	3.0
2	Malignant neoplasms (C00-C97)	112	21.2	2.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	42	8.0	0.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	34	6.4	0.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	15	2.8	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	14	2.7	*
7	Septicemia (A40-A41)	11	2.1	*
7	Cerebrovascular diseases (I60-I69)	11	2.1	*
7	Chronic lower respiratory diseases (J40-J47)	11	2.1	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	1.5	*
11	Influenza and pneumonia (J09-J18)	6	1.1	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.6	*
13	Anemias (D50-D64)	1	0.2	*
13	Diabetes mellitus (E10-E14)	1	0.2	*
13	Meningitis (G00,G03)	1	0.2	*
13	Aortic aneurysm and dissection (I71)	1	0.2	*
13	Certain conditions originating in the perinatal period (P00-P96)	1	0.2	*
...	All other causes (Residual)	121	22.9	2.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 15-19 years				
...	All causes	1,883	100.0	41.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	696	37.0	15.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	465	24.7	10.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	253	13.4	5.6
4	Malignant neoplasms (C00-C97)	133	7.1	2.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	46	2.4	1.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	26	1.4	0.6
7	Cerebrovascular diseases (I60-I69)	10	0.5	*
8	Septicemia (A40-A41)	8	0.4	*
8	Influenza and pneumonia (J09-J18)	8	0.4	*
10	Diabetes mellitus (E10-E14)	7	0.4	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.4	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.3	*
13	Aortic aneurysm and dissection (I71)	5	0.3	*
14	Pregnancy, childbirth and the puerperium (O00-O99)	4	0.2	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	3	0.2	*
15	Anemias (D50-D64)	3	0.2	*
15	Chronic lower respiratory diseases (J40-J47)	3	0.2	*
15	Pneumonitis due to solids and liquids (J69)	3	0.2	*
15	Legal intervention (Y35,Y89.0)	3	0.2	*
...	All other causes (Residual)	194	10.3	4.3
Hispanic, both sexes, 20-24 years				
...	All causes	2,912	100.0	67.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,195	41.0	27.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	557	19.1	12.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	375	12.9	8.7
4	Malignant neoplasms (C00-C97)	211	7.2	4.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	80	2.7	1.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	33	1.1	0.8
7	Pregnancy, childbirth and the puerperium (O00-O99)	28	1.0	0.6
8	Human immunodeficiency virus (HIV) disease (B20-B24)	22	0.8	0.5
9	Cerebrovascular diseases (I60-I69)	19	0.7	*
10	Influenza and pneumonia (J09-J18)	16	0.5	*
11	Septicemia (A40-A41)	13	0.4	*
11	Legal intervention (Y35,Y89.0)	13	0.4	*
13	Chronic lower respiratory diseases (J40-J47)	12	0.4	*
14	Diabetes mellitus (E10-E14)	11	0.4	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.3	*
...	All other causes (Residual)	319	11.0	7.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 25-29 years				
...	All causes	2,940	100.0	68.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,048	35.6	24.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	486	16.5	11.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	323	11.0	7.5
4	Malignant neoplasms (C00-C97)	254	8.6	5.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	144	4.9	3.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	42	1.4	1.0
7	Pregnancy, childbirth and the puerperium (O00-O99)	41	1.4	1.0
8	Influenza and pneumonia (J09-J18)	33	1.1	0.8
9	Cerebrovascular diseases (I60-I69)	31	1.1	0.7
10	Diabetes mellitus (E10-E14)	23	0.8	0.5
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	23	0.8	0.5
12	Septicemia (A40-A41)	20	0.7	0.5
12	Chronic liver disease and cirrhosis (K70,K73-K74)	20	0.7	0.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	0.6	*
15	Legal intervention (Y35,Y89.0)	13	0.4	*
...	All other causes (Residual)	421	14.3	9.8
Hispanic, both sexes, 30-34 years				
...	All causes	3,082	100.0	74.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	921	29.9	22.3
2	Malignant neoplasms (C00-C97)	398	12.9	9.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	360	11.7	8.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	265	8.6	6.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	228	7.4	5.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	92	3.0	2.2
7	Human immunodeficiency virus (HIV) disease (B20-B24)	90	2.9	2.2
8	Cerebrovascular diseases (I60-I69)	60	1.9	1.5
9	Diabetes mellitus (E10-E14)	40	1.3	1.0
10	Pregnancy, childbirth and the puerperium (O00-O99)	30	1.0	0.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	29	0.9	0.7
12	Influenza and pneumonia (J09-J18)	26	0.8	0.6
13	Septicemia (A40-A41)	24	0.8	0.6
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	22	0.7	0.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	0.6	*
15	Chronic lower respiratory diseases (J40-J47)	17	0.6	*
...	All other causes (Residual)	463	15.0	11.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 35-39 years				
...	All causes	3,497	100.0	90.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	848	24.2	22.0
2	Malignant neoplasms (C00-C97)	520	14.9	13.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	351	10.0	9.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	270	7.7	7.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	232	6.6	6.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	155	4.4	4.0
7	Human immunodeficiency virus (HIV) disease (B20-B24)	111	3.2	2.9
8	Cerebrovascular diseases (I60-I69)	95	2.7	2.5
9	Diabetes mellitus (E10-E14)	92	2.6	2.4
10	Influenza and pneumonia (J09-J18)	40	1.1	1.0
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	37	1.1	1.0
12	Septicemia (A40-A41)	27	0.8	0.7
12	Pregnancy, childbirth and the puerperium (O00-O99)	27	0.8	0.7
14	Chronic lower respiratory diseases (J40-J47)	25	0.7	0.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	21	0.6	0.5
...	All other causes (Residual)	646	18.5	16.8
Hispanic, both sexes, 40-44 years				
...	All causes	4,645	100.0	134.9
1	Malignant neoplasms (C00-C97)	934	20.1	27.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	871	18.8	25.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	568	12.2	16.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	303	6.5	8.8
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	235	5.1	6.8
6	Cerebrovascular diseases (I60-I69)	209	4.5	6.1
7	Human immunodeficiency virus (HIV) disease (B20-B24)	185	4.0	5.4
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	183	3.9	5.3
9	Diabetes mellitus (E10-E14)	150	3.2	4.4
10	Viral hepatitis (B15-B19)	70	1.5	2.0
11	Influenza and pneumonia (J09-J18)	57	1.2	1.7
12	Septicemia (A40-A41)	45	1.0	1.3
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	42	0.9	1.2
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	32	0.7	0.9
15	Chronic lower respiratory diseases (J40-J47)	30	0.6	0.9
...	All other causes (Residual)	731	15.7	21.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 45-49 years				
...	All causes	6,640	100.0	219.7
1	Malignant neoplasms (C00-C97)	1,611	24.3	53.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,000	15.1	33.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	809	12.2	26.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	537	8.1	17.8
5	Cerebrovascular diseases (I60-I69)	282	4.2	9.3
6	Diabetes mellitus (E10-E14)	265	4.0	8.8
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	239	3.6	7.9
8	Human immunodeficiency virus (HIV) disease (B20-B24)	221	3.3	7.3
9	Viral hepatitis (B15-B19)	160	2.4	5.3
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	144	2.2	4.8
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	96	1.4	3.2
12	Septicemia (A40-A41)	81	1.2	2.7
13	Chronic lower respiratory diseases (J40-J47)	72	1.1	2.4
14	Influenza and pneumonia (J09-J18)	71	1.1	2.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	37	0.6	1.2
...	All other causes (Residual)	1,015	15.3	33.6
Hispanic, both sexes, 50-54 years				
...	All causes	8,275	100.0	338.9
1	Malignant neoplasms (C00-C97)	2,313	28.0	94.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,478	17.9	60.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	703	8.5	28.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	701	8.5	28.7
5	Cerebrovascular diseases (I60-I69)	359	4.3	14.7
6	Diabetes mellitus (E10-E14)	330	4.0	13.5
7	Viral hepatitis (B15-B19)	207	2.5	8.5
8	Human immunodeficiency virus (HIV) disease (B20-B24)	190	2.3	7.8
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	183	2.2	7.5
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	130	1.6	5.3
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	127	1.5	5.2
12	Septicemia (A40-A41)	119	1.4	4.9
13	Chronic lower respiratory diseases (J40-J47)	101	1.2	4.1
14	Influenza and pneumonia (J09-J18)	87	1.1	3.6
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	65	0.8	2.7
...	All other causes (Residual)	1,182	14.3	48.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 55-59 years				
...	All causes	9,504	100.0	516.1
1	Malignant neoplasms (C00-C97)	2,886	30.4	156.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,786	18.8	97.0
3	Chronic liver disease and cirrhosis (K70,K73-K74)	664	7.0	36.1
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	553	5.8	30.0
5	Diabetes mellitus (E10-E14)	549	5.8	29.8
6	Cerebrovascular diseases (I60-I69)	394	4.1	21.4
7	Viral hepatitis (B15-B19)	268	2.8	14.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	218	2.3	11.8
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	167	1.8	9.1
10	Chronic lower respiratory diseases (J40-J47)	158	1.7	8.6
11	Septicemia (A40-A41)	154	1.6	8.4
12	Influenza and pneumonia (J09-J18)	122	1.3	6.6
13	Human immunodeficiency virus (HIV) disease (B20-B24)	120	1.3	6.5
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	102	1.1	5.5
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	54	0.6	2.9
...	All other causes (Residual)	1,309	13.8	71.1
Hispanic, both sexes, 60-64 years				
...	All causes	10,562	100.0	769.6
1	Malignant neoplasms (C00-C97)	3,360	31.8	244.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,281	21.6	166.2
3	Diabetes mellitus (E10-E14)	690	6.5	50.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	531	5.0	38.7
5	Cerebrovascular diseases (I60-I69)	467	4.4	34.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	399	3.8	29.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	267	2.5	19.5
8	Chronic lower respiratory diseases (J40-J47)	235	2.2	17.1
9	Influenza and pneumonia (J09-J18)	165	1.6	12.0
10	Septicemia (A40-A41)	164	1.6	11.9
11	Viral hepatitis (B15-B19)	158	1.5	11.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	121	1.1	8.8
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	86	0.8	6.3
14	Human immunodeficiency virus (HIV) disease (B20-B24)	68	0.6	5.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	45	0.4	3.3
...	All other causes (Residual)	1,525	14.4	111.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 65-69 years				
...	All causes	10,765	100.0	1,134.9
1	Malignant neoplasms (C00-C97)	3,494	32.5	368.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,299	21.4	242.4
3	Diabetes mellitus (E10-E14)	736	6.8	77.6
4	Cerebrovascular diseases (I60-I69)	522	4.8	55.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	393	3.7	41.4
6	Chronic lower respiratory diseases (J40-J47)	319	3.0	33.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	312	2.9	32.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	290	2.7	30.6
9	Influenza and pneumonia (J09-J18)	211	2.0	22.2
10	Septicemia (A40-A41)	176	1.6	18.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	129	1.2	13.6
12	Viral hepatitis (B15-B19)	83	0.8	8.7
13	Alzheimer's disease (G30)	80	0.7	8.4
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	73	0.7	7.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	54	0.5	5.7
...	All other causes (Residual)	1,594	14.8	168.0
Hispanic, both sexes, 70-74 years				
...	All causes	12,197	100.0	1,742.1
1	Malignant neoplasms (C00-C97)	3,808	31.2	543.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,760	22.6	394.2
3	Diabetes mellitus (E10-E14)	794	6.5	113.4
4	Cerebrovascular diseases (I60-I69)	652	5.3	93.1
5	Chronic lower respiratory diseases (J40-J47)	443	3.6	63.3
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	388	3.2	55.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	337	2.8	48.1
8	Influenza and pneumonia (J09-J18)	289	2.4	41.3
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	260	2.1	37.1
10	Septicemia (A40-A41)	205	1.7	29.3
11	Alzheimer's disease (G30)	157	1.3	22.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	150	1.2	21.4
13	Parkinson's disease (G20-G21)	87	0.7	12.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	70	0.6	10.0
15	Viral hepatitis (B15-B19)	63	0.5	9.0
15	Pneumonitis due to solids and liquids (J69)	63	0.5	9.0
...	All other causes (Residual)	1,671	13.7	238.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 75-79 years				
...	All causes	14,654	100.0	2,868.8
1	Malignant neoplasms (C00-C97)	3,782	25.8	740.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,534	24.1	691.8
3	Cerebrovascular diseases (I60-I69)	946	6.5	185.2
4	Diabetes mellitus (E10-E14)	884	6.0	173.1
5	Chronic lower respiratory diseases (J40-J47)	660	4.5	129.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	404	2.8	79.1
7	Alzheimer's disease (G30)	375	2.6	73.4
8	Influenza and pneumonia (J09-J18)	338	2.3	66.2
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	313	2.1	61.3
10	Chronic liver disease and cirrhosis (K70,K73-K74)	273	1.9	53.4
11	Septicemia (A40-A41)	239	1.6	46.8
12	Parkinson's disease (G20-G21)	215	1.5	42.1
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	193	1.3	37.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	105	0.7	20.6
15	Pneumonitis due to solids and liquids (J69)	92	0.6	18.0
...	All other causes (Residual)	2,301	15.7	450.5
Hispanic, both sexes, 80-84 years				
...	All causes	16,710	100.0	4,754.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,412	26.4	1,255.2
2	Malignant neoplasms (C00-C97)	3,513	21.0	999.5
3	Cerebrovascular diseases (I60-I69)	1,123	6.7	319.5
4	Diabetes mellitus (E10-E14)	863	5.2	245.5
5	Chronic lower respiratory diseases (J40-J47)	734	4.4	208.8
6	Alzheimer's disease (G30)	713	4.3	202.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	505	3.0	143.7
8	Influenza and pneumonia (J09-J18)	498	3.0	141.7
9	Septicemia (A40-A41)	299	1.8	85.1
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	294	1.8	83.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	274	1.6	78.0
12	Parkinson's disease (G20-G21)	253	1.5	72.0
13	Chronic liver disease and cirrhosis (K70,K73-K74)	193	1.2	54.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	117	0.7	33.3
15	Pneumonitis due to solids and liquids (J69)	109	0.7	31.0
...	All other causes (Residual)	2,810	16.8	799.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 85-89 years				
...	All causes	15,687	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,561	29.1	---
2	Malignant neoplasms (C00-C97)	2,373	15.1	---
3	Cerebrovascular diseases (I60-I69)	1,145	7.3	---
4	Alzheimer's disease (G30)	974	6.2	---
5	Chronic lower respiratory diseases (J40-J47)	724	4.6	---
6	Diabetes mellitus (E10-E14)	666	4.2	---
7	Influenza and pneumonia (J09-J18)	462	2.9	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	434	2.8	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	314	2.0	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	299	1.9	---
11	Parkinson's disease (G20-G21)	225	1.4	---
12	Septicemia (A40-A41)	224	1.4	---
13	Pneumonitis due to solids and liquids (J69)	139	0.9	---
14	Chronic liver disease and cirrhosis (K70,K73-K74)	105	0.7	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	95	0.6	---
...	All other causes (Residual)	2,947	18.8	---
Hispanic, both sexes, 90-94 years				
...	All causes	8,898	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,820	31.7	---
2	Malignant neoplasms (C00-C97)	915	10.3	---
3	Alzheimer's disease (G30)	671	7.5	---
4	Cerebrovascular diseases (I60-I69)	613	6.9	---
5	Chronic lower respiratory diseases (J40-J47)	379	4.3	---
6	Influenza and pneumonia (J09-J18)	346	3.9	---
7	Diabetes mellitus (E10-E14)	302	3.4	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	237	2.7	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	191	2.1	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	149	1.7	---
11	Septicemia (A40-A41)	117	1.3	---
12	Parkinson's disease (G20-G21)	105	1.2	---
13	Pneumonitis due to solids and liquids (J69)	87	1.0	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	51	0.6	---
15	Atherosclerosis (I70)	47	0.5	---
...	All other causes (Residual)	1,868	21.0	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 95-99 years				
...	All causes	3,700	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,237	33.4	---
2	Alzheimer's disease (G30)	324	8.8	---
3	Malignant neoplasms (C00-C97)	273	7.4	---
4	Cerebrovascular diseases (I60-I69)	240	6.5	---
5	Chronic lower respiratory diseases (J40-J47)	166	4.5	---
6	Influenza and pneumonia (J09-J18)	135	3.6	---
7	Diabetes mellitus (E10-E14)	131	3.5	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	77	2.1	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	66	1.8	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	66	1.8	---
11	Pneumonitis due to solids and liquids (J69)	32	0.9	---
12	Septicemia (A40-A41)	29	0.8	---
13	Atherosclerosis (I70)	28	0.8	---
14	Parkinson's disease (G20-G21)	22	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	0.4	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	14	0.4	---
...	All other causes (Residual)	846	22.9	---
Hispanic, both sexes, 100 years and over				
...	All causes	881	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	295	33.5	---
2	Alzheimer's disease (G30)	76	8.6	---
3	Influenza and pneumonia (J09-J18)	53	6.0	---
4	Cerebrovascular diseases (I60-I69)	51	5.8	---
5	Chronic lower respiratory diseases (J40-J47)	42	4.8	---
6	Malignant neoplasms (C00-C97)	38	4.3	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21	2.4	---
8	Diabetes mellitus (E10-E14)	20	2.3	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	18	2.0	---
10	Septicemia (A40-A41)	15	1.7	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11	1.2	---
12	Pneumonitis due to solids and liquids (J69)	10	1.1	---
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.6	---
13	Parkinson's disease (G20-G21)	5	0.6	---
15	Atherosclerosis (I70)	4	0.5	---
...	All other causes (Residual)	217	24.6	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, all ages				
...	All causes	79,622	100.0	310.8
1	Malignant neoplasms (C00-C97)	16,450	20.7	64.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	16,421	20.6	64.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,594	9.5	29.6
4	Cerebrovascular diseases (I60-I69)	3,382	4.2	13.2
5	Diabetes mellitus (E10-E14)	3,372	4.2	13.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	3,067	3.9	12.0
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,435	3.1	9.5
8	Chronic lower respiratory diseases (J40-J47)	2,174	2.7	8.5
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,168	2.7	8.5
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,670	2.1	6.5
11	Influenza and pneumonia (J09-J18)	1,565	2.0	6.1
12	Certain conditions originating in the perinatal period (P00-P96)	1,456	1.8	5.7
13	Alzheimer's disease (G30)	1,171	1.5	4.6
14	Septicemia (A40-A41)	1,012	1.3	4.0
15	Human immunodeficiency virus (HIV) disease (B20-B24)	910	1.1	3.6
...	All other causes (Residual)	14,775	18.6	57.7
Hispanic, male, 1-4 years				
...	All causes	524	100.0	25.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	209	39.9	10.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	59	11.3	2.8
3	Malignant neoplasms (C00-C97)	42	8.0	2.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	42	8.0	2.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	16	3.1	*
6	Influenza and pneumonia (J09-J18)	13	2.5	*
7	Septicemia (A40-A41)	12	2.3	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.1	*
9	Chronic lower respiratory diseases (J40-J47)	5	1.0	*
10	Pneumonitis due to solids and liquids (J69)	3	0.6	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.6	*
12	Meningitis (G00,G03)	2	0.4	*
12	Acute bronchitis and bronchiolitis (J20-J21)	2	0.4	*
12	Certain conditions originating in the perinatal period (P00-P96)	2	0.4	*
15	Meningococcal infection (A39)	1	0.2	*
15	Cerebrovascular diseases (I60-I69)	1	0.2	*
15	Diseases of appendix (K35-K38)	1	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
...	All other causes (Residual)	104	19.8	5.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 5-9 years				
...	All causes	245	100.0	10.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	70	28.6	2.9
2	Malignant neoplasms (C00-C97)	58	23.7	2.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	21	8.6	0.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	10	4.1	*
5	Chronic lower respiratory diseases (J40-J47)	9	3.7	*
6	Cerebrovascular diseases (I60-I69)	7	2.9	*
7	Influenza and pneumonia (J09-J18)	6	2.4	*
8	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	2.0	*
9	Septicemia (A40-A41)	2	0.8	*
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2	0.8	*
9	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.8	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.4	*
12	Anemias (D50-D64)	1	0.4	*
12	Meningitis (G00,G03)	1	0.4	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.4	*
...	All other causes (Residual)	49	20.0	2.0
Hispanic, male, 10-14 years				
...	All causes	298	100.0	12.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	80	26.8	3.5
2	Malignant neoplasms (C00-C97)	54	18.1	2.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	25	8.4	1.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	20	6.7	0.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	9	3.0	*
5	Chronic lower respiratory diseases (J40-J47)	9	3.0	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	3.0	*
8	Septicemia (A40-A41)	7	2.3	*
9	Cerebrovascular diseases (I60-I69)	6	2.0	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	1.7	*
11	Influenza and pneumonia (J09-J18)	3	1.0	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	1.0	*
13	Anemias (D50-D64)	1	0.3	*
13	Aortic aneurysm and dissection (I71)	1	0.3	*
13	Certain conditions originating in the perinatal period (P00-P96)	1	0.3	*
...	All other causes (Residual)	65	21.8	2.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 15-19 years				
...	All causes	1,436	100.0	61.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	508	35.4	21.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	420	29.2	17.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	190	13.2	8.1
4	Malignant neoplasms (C00-C97)	92	6.4	3.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	37	2.6	1.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	14	1.0	*
7	Influenza and pneumonia (J09-J18)	8	0.6	*
8	Cerebrovascular diseases (I60-I69)	7	0.5	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.3	*
10	Septicemia (A40-A41)	4	0.3	*
10	Diabetes mellitus (E10-E14)	4	0.3	*
10	Aortic aneurysm and dissection (I71)	4	0.3	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	3	0.2	*
13	Anemias (D50-D64)	3	0.2	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.2	*
13	Legal intervention (Y35,Y89.0)	3	0.2	*
...	All other causes (Residual)	131	9.1	5.6
Hispanic, male, 20-24 years				
...	All causes	2,253	100.0	97.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	970	43.1	42.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	494	21.9	21.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	307	13.6	13.3
4	Malignant neoplasms (C00-C97)	133	5.9	5.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	57	2.5	2.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	20	0.9	0.9
7	Human immunodeficiency virus (HIV) disease (B20-B24)	17	0.8	*
8	Legal intervention (Y35,Y89.0)	13	0.6	*
9	Cerebrovascular diseases (I60-I69)	12	0.5	*
10	Influenza and pneumonia (J09-J18)	9	0.4	*
11	Chronic lower respiratory diseases (J40-J47)	6	0.3	*
12	Septicemia (A40-A41)	5	0.2	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.2	*
12	Diabetes mellitus (E10-E14)	5	0.2	*
12	Aortic aneurysm and dissection (I71)	5	0.2	*
...	All other causes (Residual)	195	8.7	8.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 25-29 years				
...	All causes	2,217	100.0	97.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	846	38.2	37.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	426	19.2	18.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	272	12.3	12.0
4	Malignant neoplasms (C00-C97)	137	6.2	6.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	107	4.8	4.7
6	Human immunodeficiency virus (HIV) disease (B20-B24)	35	1.6	1.5
7	Influenza and pneumonia (J09-J18)	23	1.0	1.0
8	Cerebrovascular diseases (I60-I69)	20	0.9	0.9
9	Diabetes mellitus (E10-E14)	16	0.7	*
9	Chronic liver disease and cirrhosis (K70,K73-K74)	16	0.7	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	14	0.6	*
12	Legal intervention (Y35,Y89.0)	13	0.6	*
13	Aortic aneurysm and dissection (I71)	12	0.5	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.5	*
15	Septicemia (A40-A41)	9	0.4	*
...	All other causes (Residual)	260	11.7	11.4
Hispanic, male, 30-34 years				
...	All causes	2,244	100.0	104.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	742	33.1	34.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	315	14.0	14.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	232	10.3	10.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	189	8.4	8.8
5	Malignant neoplasms (C00-C97)	179	8.0	8.4
6	Chronic liver disease and cirrhosis (K70,K73-K74)	77	3.4	3.6
7	Human immunodeficiency virus (HIV) disease (B20-B24)	72	3.2	3.4
8	Cerebrovascular diseases (I60-I69)	39	1.7	1.8
9	Diabetes mellitus (E10-E14)	25	1.1	1.2
10	Septicemia (A40-A41)	14	0.6	*
10	Legal intervention (Y35,Y89.0)	14	0.6	*
12	Influenza and pneumonia (J09-J18)	13	0.6	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	0.6	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	0.5	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	12	0.5	*
...	All other causes (Residual)	296	13.2	13.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 35-39 years				
...	All causes	2,399	100.0	121.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	665	27.7	33.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	266	11.1	13.5
3	Malignant neoplasms (C00-C97)	242	10.1	12.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	211	8.8	10.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	192	8.0	9.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	128	5.3	6.5
7	Human immunodeficiency virus (HIV) disease (B20-B24)	85	3.5	4.3
8	Cerebrovascular diseases (I60-I69)	62	2.6	3.1
9	Diabetes mellitus (E10-E14)	53	2.2	2.7
10	Influenza and pneumonia (J09-J18)	29	1.2	1.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21	0.9	1.1
12	Aortic aneurysm and dissection (I71)	15	0.6	*
13	Chronic lower respiratory diseases (J40-J47)	14	0.6	*
14	Septicemia (A40-A41)	13	0.5	*
14	Legal intervention (Y35,Y89.0)	13	0.5	*
...	All other causes (Residual)	390	16.3	19.8
Hispanic, male, 40-44 years				
...	All causes	3,061	100.0	173.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	683	22.3	38.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	430	14.0	24.4
3	Malignant neoplasms (C00-C97)	376	12.3	21.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	238	7.8	13.5
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	196	6.4	11.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	145	4.7	8.2
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	141	4.6	8.0
8	Cerebrovascular diseases (I60-I69)	129	4.2	7.3
9	Diabetes mellitus (E10-E14)	92	3.0	5.2
10	Viral hepatitis (B15-B19)	54	1.8	3.1
11	Influenza and pneumonia (J09-J18)	38	1.2	2.2
12	Septicemia (A40-A41)	31	1.0	1.8
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	26	0.8	1.5
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	22	0.7	1.2
15	Chronic lower respiratory diseases (J40-J47)	18	0.6	*
...	All other causes (Residual)	442	14.4	25.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 45-49 years				
...	All causes	4,310	100.0	282.6
1	Malignant neoplasms (C00-C97)	744	17.3	48.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	720	16.7	47.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	610	14.2	40.0
4	Chronic liver disease and cirrhosis (K70,K73-K74)	408	9.5	26.8
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	197	4.6	12.9
6	Cerebrovascular diseases (I60-I69)	181	4.2	11.9
7	Human immunodeficiency virus (HIV) disease (B20-B24)	175	4.1	11.5
8	Diabetes mellitus (E10-E14)	171	4.0	11.2
9	Viral hepatitis (B15-B19)	127	2.9	8.3
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	114	2.6	7.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	59	1.4	3.9
12	Septicemia (A40-A41)	53	1.2	3.5
13	Influenza and pneumonia (J09-J18)	45	1.0	3.0
14	Chronic lower respiratory diseases (J40-J47)	34	0.8	2.2
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	24	0.6	1.6
...	All other causes (Residual)	648	15.0	42.5
Hispanic, male, 50-54 years				
...	All causes	5,317	100.0	439.1
1	Malignant neoplasms (C00-C97)	1,162	21.9	96.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,089	20.5	89.9
3	Chronic liver disease and cirrhosis (K70,K73-K74)	537	10.1	44.3
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	514	9.7	42.4
5	Cerebrovascular diseases (I60-I69)	210	3.9	17.3
6	Diabetes mellitus (E10-E14)	205	3.9	16.9
7	Human immunodeficiency virus (HIV) disease (B20-B24)	158	3.0	13.0
8	Viral hepatitis (B15-B19)	157	3.0	13.0
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	150	2.8	12.4
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	105	2.0	8.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	75	1.4	6.2
12	Septicemia (A40-A41)	68	1.3	5.6
13	Chronic lower respiratory diseases (J40-J47)	50	0.9	4.1
14	Influenza and pneumonia (J09-J18)	47	0.9	3.9
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	38	0.7	3.1
...	All other causes (Residual)	752	14.1	62.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 55-59 years				
...	All causes	6,100	100.0	686.1
1	Malignant neoplasms (C00-C97)	1,589	26.0	178.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,298	21.3	146.0
3	Chronic liver disease and cirrhosis (K70,K73-K74)	514	8.4	57.8
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	395	6.5	44.4
5	Diabetes mellitus (E10-E14)	363	6.0	40.8
6	Cerebrovascular diseases (I60-I69)	231	3.8	26.0
7	Viral hepatitis (B15-B19)	194	3.2	21.8
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	138	2.3	15.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	125	2.0	14.1
10	Human immunodeficiency virus (HIV) disease (B20-B24)	98	1.6	11.0
11	Chronic lower respiratory diseases (J40-J47)	96	1.6	10.8
12	Septicemia (A40-A41)	94	1.5	10.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	72	1.2	8.1
14	Influenza and pneumonia (J09-J18)	67	1.1	7.5
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	48	0.8	5.4
...	All other causes (Residual)	778	12.8	87.5
Hispanic, male, 60-64 years				
...	All causes	6,409	100.0	992.8
1	Malignant neoplasms (C00-C97)	1,871	29.2	289.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,555	24.3	240.9
3	Diabetes mellitus (E10-E14)	398	6.2	61.7
4	Chronic liver disease and cirrhosis (K70,K73-K74)	394	6.1	61.0
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	278	4.3	43.1
6	Cerebrovascular diseases (I60-I69)	259	4.0	40.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	158	2.5	24.5
8	Chronic lower respiratory diseases (J40-J47)	129	2.0	20.0
9	Viral hepatitis (B15-B19)	96	1.5	14.9
10	Influenza and pneumonia (J09-J18)	92	1.4	14.3
11	Septicemia (A40-A41)	86	1.3	13.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	70	1.1	10.8
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	67	1.0	10.4
14	Human immunodeficiency virus (HIV) disease (B20-B24)	58	0.9	9.0
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	28	0.4	4.3
...	All other causes (Residual)	870	13.6	134.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 65-69 years				
...	All causes	6,223	100.0	1,450.9
1	Malignant neoplasms (C00-C97)	1,953	31.4	455.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,464	23.5	341.3
3	Diabetes mellitus (E10-E14)	409	6.6	95.4
4	Cerebrovascular diseases (I60-I69)	287	4.6	66.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	258	4.1	60.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	190	3.1	44.3
7	Chronic lower respiratory diseases (J40-J47)	167	2.7	38.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	157	2.5	36.6
9	Influenza and pneumonia (J09-J18)	120	1.9	28.0
10	Septicemia (A40-A41)	81	1.3	18.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	66	1.1	15.4
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	57	0.9	13.3
13	Viral hepatitis (B15-B19)	56	0.9	13.1
14	Alzheimer's disease (G30)	35	0.6	8.2
15	Human immunodeficiency virus (HIV) disease (B20-B24)	33	0.5	7.7
...	All other causes (Residual)	890	14.3	207.5
Hispanic, male, 70-74 years				
...	All causes	6,817	100.0	2,229.5
1	Malignant neoplasms (C00-C97)	2,102	30.8	687.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,655	24.3	541.3
3	Diabetes mellitus (E10-E14)	416	6.1	136.1
4	Cerebrovascular diseases (I60-I69)	333	4.9	108.9
5	Chronic lower respiratory diseases (J40-J47)	257	3.8	84.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	200	2.9	65.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	195	2.9	63.8
8	Influenza and pneumonia (J09-J18)	168	2.5	54.9
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	166	2.4	54.3
10	Septicemia (A40-A41)	114	1.7	37.3
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	69	1.0	22.6
12	Alzheimer's disease (G30)	64	0.9	20.9
13	Parkinson's disease (G20-G21)	57	0.8	18.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	45	0.7	14.7
15	Pneumonitis due to solids and liquids (J69)	44	0.6	14.4
...	All other causes (Residual)	932	13.7	304.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 75-79 years				
...	All causes	7,664	100.0	3,592.4
1	Malignant neoplasms (C00-C97)	2,093	27.3	981.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,941	25.3	909.8
3	Cerebrovascular diseases (I60-I69)	449	5.9	210.5
4	Diabetes mellitus (E10-E14)	430	5.6	201.6
5	Chronic lower respiratory diseases (J40-J47)	358	4.7	167.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	212	2.8	99.4
7	Influenza and pneumonia (J09-J18)	191	2.5	89.5
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	185	2.4	86.7
9	Alzheimer's disease (G30)	171	2.2	80.2
10	Chronic liver disease and cirrhosis (K70,K73-K74)	146	1.9	68.4
11	Parkinson's disease (G20-G21)	133	1.7	62.3
12	Septicemia (A40-A41)	111	1.4	52.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	68	0.9	31.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	54	0.7	25.3
15	Pneumonitis due to solids and liquids (J69)	52	0.7	24.4
...	All other causes (Residual)	1,070	14.0	501.5
Hispanic, male, 80-84 years				
...	All causes	8,059	100.0	5,795.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,150	26.7	1,546.3
2	Malignant neoplasms (C00-C97)	1,849	22.9	1,329.8
3	Cerebrovascular diseases (I60-I69)	490	6.1	352.4
4	Diabetes mellitus (E10-E14)	396	4.9	284.8
5	Chronic lower respiratory diseases (J40-J47)	392	4.9	281.9
6	Alzheimer's disease (G30)	260	3.2	187.0
7	Influenza and pneumonia (J09-J18)	259	3.2	186.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	249	3.1	179.1
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	160	2.0	115.1
10	Septicemia (A40-A41)	140	1.7	100.7
11	Parkinson's disease (G20-G21)	131	1.6	94.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	131	1.6	94.2
13	Chronic liver disease and cirrhosis (K70,K73-K74)	94	1.2	67.6
14	Pneumonitis due to solids and liquids (J69)	62	0.8	44.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	54	0.7	38.8
...	All other causes (Residual)	1,242	15.4	893.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 85-89 years				
...	All causes	6,737	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,969	29.2	---
2	Malignant neoplasms (C00-C97)	1,216	18.0	---
3	Cerebrovascular diseases (I60-I69)	407	6.0	---
4	Chronic lower respiratory diseases (J40-J47)	360	5.3	---
5	Alzheimer's disease (G30)	333	4.9	---
6	Diabetes mellitus (E10-E14)	255	3.8	---
7	Influenza and pneumonia (J09-J18)	223	3.3	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	201	3.0	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	134	2.0	---
10	Parkinson's disease (G20-G21)	133	2.0	---
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	110	1.6	---
12	Septicemia (A40-A41)	86	1.3	---
13	Pneumonitis due to solids and liquids (J69)	83	1.2	---
14	Chronic liver disease and cirrhosis (K70,K73-K74)	51	0.8	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	47	0.7	---
...	All other causes (Residual)	1,129	16.8	---
Hispanic, male, 90-94 years				
...	All causes	3,159	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,016	32.2	---
2	Malignant neoplasms (C00-C97)	416	13.2	---
3	Alzheimer's disease (G30)	206	6.5	---
4	Cerebrovascular diseases (I60-I69)	171	5.4	---
5	Chronic lower respiratory diseases (J40-J47)	159	5.0	---
6	Influenza and pneumonia (J09-J18)	133	4.2	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	104	3.3	---
8	Diabetes mellitus (E10-E14)	94	3.0	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	74	2.3	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	59	1.9	---
11	Septicemia (A40-A41)	43	1.4	---
11	Parkinson's disease (G20-G21)	43	1.4	---
13	Pneumonitis due to solids and liquids (J69)	35	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	26	0.8	---
15	Atherosclerosis (I70)	21	0.7	---
...	All other causes (Residual)	559	17.7	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 95-99 years				
...	All causes	1,075	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	339	31.5	---
2	Malignant neoplasms (C00-C97)	119	11.1	---
3	Alzheimer's disease (G30)	70	6.5	---
4	Chronic lower respiratory diseases (J40-J47)	67	6.2	---
5	Cerebrovascular diseases (I60-I69)	62	5.8	---
6	Influenza and pneumonia (J09-J18)	45	4.2	---
7	Diabetes mellitus (E10-E14)	37	3.4	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	32	3.0	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	23	2.1	---
10	Pneumonitis due to solids and liquids (J69)	16	1.5	---
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	14	1.3	---
12	Septicemia (A40-A41)	10	0.9	---
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.8	---
14	Parkinson's disease (G20-G21)	8	0.7	---
15	Atherosclerosis (I70)	6	0.6	---
...	All other causes (Residual)	218	20.3	---
Hispanic, male, 100 years and over				
...	All causes	200	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	69	34.5	---
2	Chronic lower respiratory diseases (J40-J47)	16	8.0	---
3	Malignant neoplasms (C00-C97)	14	7.0	---
3	Influenza and pneumonia (J09-J18)	14	7.0	---
5	Alzheimer's disease (G30)	9	4.5	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	3.5	---
7	Septicemia (A40-A41)	5	2.5	---
8	Cerebrovascular diseases (I60-I69)	4	2.0	---
9	Diabetes mellitus (E10-E14)	3	1.5	---
9	Atherosclerosis (I70)	3	1.5	---
9	Pneumonitis due to solids and liquids (J69)	3	1.5	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3	1.5	---
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2	1.0	---
14	Anemias (D50-D64)	1	0.5	---
...	All other causes (Residual)	47	23.5	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, all ages				
...	All causes	64,868	100.0	260.9
1	Malignant neoplasms (C00-C97)	14,669	22.6	59.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,585	20.9	54.6
3	Cerebrovascular diseases (I60-I69)	3,892	6.0	15.7
4	Diabetes mellitus (E10-E14)	3,184	4.9	12.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,882	4.4	11.6
6	Alzheimer's disease (G30)	2,256	3.5	9.1
7	Chronic lower respiratory diseases (J40-J47)	1,998	3.1	8.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,582	2.4	6.4
9	Influenza and pneumonia (J09-J18)	1,460	2.3	5.9
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,281	2.0	5.2
11	Certain conditions originating in the perinatal period (P00-P96)	1,073	1.7	4.3
12	Septicemia (A40-A41)	1,023	1.6	4.1
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	948	1.5	3.8
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	883	1.4	3.6
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	493	0.8	2.0
...	All other causes (Residual)	13,659	21.1	54.9
Hispanic, female, 1-4 years				
...	All causes	406	100.0	20.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	121	29.8	6.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	63	15.5	3.1
3	Malignant neoplasms (C00-C97)	33	8.1	1.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	21	5.2	1.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	15	3.7	*
6	Influenza and pneumonia (J09-J18)	8	2.0	*
7	Cerebrovascular diseases (I60-I69)	7	1.7	*
8	Septicemia (A40-A41)	6	1.5	*
9	Chronic lower respiratory diseases (J40-J47)	5	1.2	*
9	Certain conditions originating in the perinatal period (P00-P96)	5	1.2	*
11	Meningococcal infection (A39)	4	1.0	*
12	Anemias (D50-D64)	3	0.7	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.7	*
14	Acute bronchitis and bronchiolitis (J20-J21)	2	0.5	*
14	Pneumonitis due to solids and liquids (J69)	2	0.5	*
...	All other causes (Residual)	108	26.6	5.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 5-9 years				
...	All causes	178	100.0	7.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	47	26.4	2.0
2	Malignant neoplasms (C00-C97)	42	23.6	1.8
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	15	8.4	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	10	5.6	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	6	3.4	*
6	Cerebrovascular diseases (I60-I69)	5	2.8	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	1.7	*
8	Pneumonitis due to solids and liquids (J69)	2	1.1	*
9	Tuberculosis (A16-A19)	1	0.6	*
9	Meningococcal infection (A39)	1	0.6	*
9	Septicemia (A40-A41)	1	0.6	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.6	*
9	Anemias (D50-D64)	1	0.6	*
9	Diabetes mellitus (E10-E14)	1	0.6	*
9	Meningitis (G00,G03)	1	0.6	*
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.6	*
9	Influenza and pneumonia (J09-J18)	1	0.6	*
9	Acute bronchitis and bronchiolitis (J20-J21)	1	0.6	*
9	Diseases of appendix (K35-K38)	1	0.6	*
...	All other causes (Residual)	37	20.8	1.6
Hispanic, female, 10-14 years				
...	All causes	230	100.0	10.4
1	Malignant neoplasms (C00-C97)	58	25.2	2.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	55	23.9	2.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	22	9.6	1.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	9	3.9	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	6	2.6	*
6	Cerebrovascular diseases (I60-I69)	5	2.2	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	2.2	*
8	Septicemia (A40-A41)	4	1.7	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	1.3	*
9	Influenza and pneumonia (J09-J18)	3	1.3	*
11	Chronic lower respiratory diseases (J40-J47)	2	0.9	*
12	Diabetes mellitus (E10-E14)	1	0.4	*
12	Meningitis (G00,G03)	1	0.4	*
...	All other causes (Residual)	56	24.3	2.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 15-19 years				
...	All causes	447	100.0	20.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	188	42.1	8.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	63	14.1	2.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	45	10.1	2.1
4	Malignant neoplasms (C00-C97)	41	9.2	1.9
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	12	2.7	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	9	2.0	*
7	Septicemia (A40-A41)	4	0.9	*
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.9	*
7	Pregnancy, childbirth and the puerperium (O00-O99)	4	0.9	*
10	Diabetes mellitus (E10-E14)	3	0.7	*
10	Cerebrovascular diseases (I60-I69)	3	0.7	*
12	Tuberculosis (A16-A19)	1	0.2	*
12	Meningococcal infection (A39)	1	0.2	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.2	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.2	*
12	Aortic aneurysm and dissection (I71)	1	0.2	*
12	Chronic lower respiratory diseases (J40-J47)	1	0.2	*
12	Pneumonitis due to solids and liquids (J69)	1	0.2	*
12	Certain conditions originating in the perinatal period (P00-P96)	1	0.2	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
12	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.2	*
...	All other causes (Residual)	61	13.6	2.8
Hispanic, female, 20-24 years				
...	All causes	659	100.0	32.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	225	34.1	11.1
2	Malignant neoplasms (C00-C97)	78	11.8	3.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	68	10.3	3.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	63	9.6	3.1
5	Pregnancy, childbirth and the puerperium (O00-O99)	28	4.2	1.4
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	23	3.5	1.1
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	13	2.0	*
8	Septicemia (A40-A41)	8	1.2	*
9	Cerebrovascular diseases (I60-I69)	7	1.1	*
9	Influenza and pneumonia (J09-J18)	7	1.1	*
11	Diabetes mellitus (E10-E14)	6	0.9	*
11	Chronic lower respiratory diseases (J40-J47)	6	0.9	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	5	0.8	*
14	Chronic liver disease and cirrhosis (K70,K73-K74)	4	0.6	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.5	*
...	All other causes (Residual)	112	17.0	5.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 25-29 years				
...	All causes	723	100.0	35.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	202	27.9	9.9
2	Malignant neoplasms (C00-C97)	117	16.2	5.8
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	60	8.3	2.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	51	7.1	2.5
5	Pregnancy, childbirth and the puerperium (O00-O99)	41	5.7	2.0
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	37	5.1	1.8
7	Septicemia (A40-A41)	11	1.5	*
7	Cerebrovascular diseases (I60-I69)	11	1.5	*
9	Influenza and pneumonia (J09-J18)	10	1.4	*
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	1.2	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	7	1.0	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	1.0	*
11	Diabetes mellitus (E10-E14)	7	1.0	*
14	Chronic lower respiratory diseases (J40-J47)	4	0.6	*
14	Chronic liver disease and cirrhosis (K70,K73-K74)	4	0.6	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.6	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.6	*
...	All other causes (Residual)	137	18.9	6.7
Hispanic, female, 30-34 years				
...	All causes	838	100.0	42.3
1	Malignant neoplasms (C00-C97)	219	26.1	11.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	179	21.4	9.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	45	5.4	2.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	39	4.7	2.0
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	33	3.9	1.7
6	Pregnancy, childbirth and the puerperium (O00-O99)	30	3.6	1.5
7	Cerebrovascular diseases (I60-I69)	21	2.5	1.1
8	Human immunodeficiency virus (HIV) disease (B20-B24)	18	2.1	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	1.9	*
10	Diabetes mellitus (E10-E14)	15	1.8	*
10	Chronic liver disease and cirrhosis (K70,K73-K74)	15	1.8	*
12	Influenza and pneumonia (J09-J18)	13	1.6	*
13	Septicemia (A40-A41)	10	1.2	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	10	1.2	*
15	Chronic lower respiratory diseases (J40-J47)	6	0.7	*
...	All other causes (Residual)	169	20.2	8.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 35-39 years				
...	All causes	1,098	100.0	58.2
1	Malignant neoplasms (C00-C97)	278	25.3	14.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	183	16.7	9.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	85	7.7	4.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	59	5.4	3.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	40	3.6	2.1
6	Diabetes mellitus (E10-E14)	39	3.6	2.1
7	Cerebrovascular diseases (I60-I69)	33	3.0	1.8
8	Chronic liver disease and cirrhosis (K70,K73-K74)	27	2.5	1.4
8	Pregnancy, childbirth and the puerperium (O00-O99)	27	2.5	1.4
10	Human immunodeficiency virus (HIV) disease (B20-B24)	26	2.4	1.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	1.5	*
12	Septicemia (A40-A41)	14	1.3	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	1.0	*
13	Influenza and pneumonia (J09-J18)	11	1.0	*
13	Chronic lower respiratory diseases (J40-J47)	11	1.0	*
...	All other causes (Residual)	238	21.7	12.6
Hispanic, female, 40-44 years				
...	All causes	1,584	100.0	94.4
1	Malignant neoplasms (C00-C97)	558	35.2	33.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	188	11.9	11.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	138	8.7	8.2
4	Cerebrovascular diseases (I60-I69)	80	5.1	4.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	65	4.1	3.9
6	Diabetes mellitus (E10-E14)	58	3.7	3.5
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	42	2.7	2.5
8	Human immunodeficiency virus (HIV) disease (B20-B24)	40	2.5	2.4
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	39	2.5	2.3
10	Influenza and pneumonia (J09-J18)	19	1.2	*
11	Viral hepatitis (B15-B19)	16	1.0	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	1.0	*
13	Septicemia (A40-A41)	14	0.9	*
14	Chronic lower respiratory diseases (J40-J47)	12	0.8	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10	0.6	*
15	Pregnancy, childbirth and the puerperium (O00-O99)	10	0.6	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	10	0.6	*
...	All other causes (Residual)	269	17.0	16.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 45-49 years				
...	All causes	2,330	100.0	155.6
1	Malignant neoplasms (C00-C97)	867	37.2	57.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	280	12.0	18.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	199	8.5	13.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	129	5.5	8.6
5	Cerebrovascular diseases (I60-I69)	101	4.3	6.7
6	Diabetes mellitus (E10-E14)	94	4.0	6.3
7	Human immunodeficiency virus (HIV) disease (B20-B24)	46	2.0	3.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	42	1.8	2.8
9	Chronic lower respiratory diseases (J40-J47)	38	1.6	2.5
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	37	1.6	2.5
11	Viral hepatitis (B15-B19)	33	1.4	2.2
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	30	1.3	2.0
13	Septicemia (A40-A41)	28	1.2	1.9
14	Influenza and pneumonia (J09-J18)	26	1.1	1.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	24	1.0	1.6
...	All other causes (Residual)	356	15.3	23.8
Hispanic, female, 50-54 years				
...	All causes	2,958	100.0	240.4
1	Malignant neoplasms (C00-C97)	1,151	38.9	93.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	389	13.2	31.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	189	6.4	15.4
4	Chronic liver disease and cirrhosis (K70,K73-K74)	164	5.5	13.3
5	Cerebrovascular diseases (I60-I69)	149	5.0	12.1
6	Diabetes mellitus (E10-E14)	125	4.2	10.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	52	1.8	4.2
8	Septicemia (A40-A41)	51	1.7	4.1
8	Chronic lower respiratory diseases (J40-J47)	51	1.7	4.1
10	Viral hepatitis (B15-B19)	50	1.7	4.1
11	Influenza and pneumonia (J09-J18)	40	1.4	3.3
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	33	1.1	2.7
13	Human immunodeficiency virus (HIV) disease (B20-B24)	32	1.1	2.6
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	27	0.9	2.2
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	25	0.8	2.0
...	All other causes (Residual)	430	14.5	34.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 55-59 years				
...	All causes	3,404	100.0	357.5
1	Malignant neoplasms (C00-C97)	1,297	38.1	136.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	488	14.3	51.2
3	Diabetes mellitus (E10-E14)	186	5.5	19.5
4	Cerebrovascular diseases (I60-I69)	163	4.8	17.1
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	158	4.6	16.6
6	Chronic liver disease and cirrhosis (K70,K73-K74)	150	4.4	15.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	93	2.7	9.8
8	Viral hepatitis (B15-B19)	74	2.2	7.8
9	Chronic lower respiratory diseases (J40-J47)	62	1.8	6.5
10	Septicemia (A40-A41)	60	1.8	6.3
11	Influenza and pneumonia (J09-J18)	55	1.6	5.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	30	0.9	3.2
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	29	0.9	3.0
14	Human immunodeficiency virus (HIV) disease (B20-B24)	22	0.6	2.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	0.5	*
...	All other causes (Residual)	519	15.2	54.5
Hispanic, female, 60-64 years				
...	All causes	4,153	100.0	571.4
1	Malignant neoplasms (C00-C97)	1,489	35.9	204.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	726	17.5	99.9
3	Diabetes mellitus (E10-E14)	292	7.0	40.2
4	Cerebrovascular diseases (I60-I69)	208	5.0	28.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	137	3.3	18.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	121	2.9	16.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	109	2.6	15.0
8	Chronic lower respiratory diseases (J40-J47)	106	2.6	14.6
9	Septicemia (A40-A41)	78	1.9	10.7
10	Influenza and pneumonia (J09-J18)	73	1.8	10.0
11	Viral hepatitis (B15-B19)	62	1.5	8.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	51	1.2	7.0
13	Alzheimer's disease (G30)	23	0.6	3.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	22	0.5	3.0
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	19	0.5	*
...	All other causes (Residual)	637	15.3	87.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 65-69 years				
...	All causes	4,542	100.0	874.0
1	Malignant neoplasms (C00-C97)	1,541	33.9	296.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	835	18.4	160.7
3	Diabetes mellitus (E10-E14)	327	7.2	62.9
4	Cerebrovascular diseases (I60-I69)	235	5.2	45.2
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	155	3.4	29.8
6	Chronic lower respiratory diseases (J40-J47)	152	3.3	29.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	135	3.0	26.0
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	100	2.2	19.2
9	Septicemia (A40-A41)	95	2.1	18.3
10	Influenza and pneumonia (J09-J18)	91	2.0	17.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	63	1.4	12.1
12	Alzheimer's disease (G30)	45	1.0	8.7
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	29	0.6	5.6
14	Viral hepatitis (B15-B19)	27	0.6	5.2
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	16	0.4	*
...	All other causes (Residual)	696	15.3	133.9
Hispanic, female, 70-74 years				
...	All causes	5,380	100.0	1,364.2
1	Malignant neoplasms (C00-C97)	1,706	31.7	432.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,105	20.5	280.2
3	Diabetes mellitus (E10-E14)	378	7.0	95.8
4	Cerebrovascular diseases (I60-I69)	319	5.9	80.9
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	188	3.5	47.7
6	Chronic lower respiratory diseases (J40-J47)	186	3.5	47.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	142	2.6	36.0
8	Influenza and pneumonia (J09-J18)	121	2.2	30.7
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	94	1.7	23.8
10	Alzheimer's disease (G30)	93	1.7	23.6
11	Septicemia (A40-A41)	91	1.7	23.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	81	1.5	20.5
13	Viral hepatitis (B15-B19)	33	0.6	8.4
14	Parkinson's disease (G20-G21)	30	0.6	7.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	25	0.5	6.3
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	25	0.5	6.3
...	All other causes (Residual)	763	14.2	193.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 75-79 years				
...	All causes	6,990	100.0	2,349.8
1	Malignant neoplasms (C00-C97)	1,689	24.2	567.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,593	22.8	535.5
3	Cerebrovascular diseases (I60-I69)	497	7.1	167.1
4	Diabetes mellitus (E10-E14)	454	6.5	152.6
5	Chronic lower respiratory diseases (J40-J47)	302	4.3	101.5
6	Alzheimer's disease (G30)	204	2.9	68.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	192	2.7	64.5
8	Influenza and pneumonia (J09-J18)	147	2.1	49.4
9	Septicemia (A40-A41)	128	1.8	43.0
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	128	1.8	43.0
11	Chronic liver disease and cirrhosis (K70,K73-K74)	127	1.8	42.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	125	1.8	42.0
13	Parkinson's disease (G20-G21)	82	1.2	27.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	51	0.7	17.1
15	Pneumonitis due to solids and liquids (J69)	40	0.6	13.4
...	All other causes (Residual)	1,231	17.6	413.8
Hispanic, female, 80-84 years				
...	All causes	8,651	100.0	4,072.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,262	26.1	1,064.8
2	Malignant neoplasms (C00-C97)	1,664	19.2	783.3
3	Cerebrovascular diseases (I60-I69)	633	7.3	298.0
4	Diabetes mellitus (E10-E14)	467	5.4	219.8
5	Alzheimer's disease (G30)	453	5.2	213.2
6	Chronic lower respiratory diseases (J40-J47)	342	4.0	161.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	256	3.0	120.5
8	Influenza and pneumonia (J09-J18)	239	2.8	112.5
9	Septicemia (A40-A41)	159	1.8	74.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	143	1.7	67.3
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	134	1.5	63.1
12	Parkinson's disease (G20-G21)	122	1.4	57.4
13	Chronic liver disease and cirrhosis (K70,K73-K74)	99	1.1	46.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	63	0.7	29.7
15	Pneumonitis due to solids and liquids (J69)	47	0.5	22.1
...	All other causes (Residual)	1,568	18.1	738.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 85-89 years				
...	All causes	8,950	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,592	29.0	---
2	Malignant neoplasms (C00-C97)	1,157	12.9	---
3	Cerebrovascular diseases (I60-I69)	738	8.2	---
4	Alzheimer's disease (G30)	641	7.2	---
5	Diabetes mellitus (E10-E14)	411	4.6	---
6	Chronic lower respiratory diseases (J40-J47)	364	4.1	---
7	Influenza and pneumonia (J09-J18)	239	2.7	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	233	2.6	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	189	2.1	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	180	2.0	---
11	Septicemia (A40-A41)	138	1.5	---
12	Parkinson's disease (G20-G21)	92	1.0	---
13	Pneumonitis due to solids and liquids (J69)	56	0.6	---
14	Chronic liver disease and cirrhosis (K70,K73-K74)	54	0.6	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	50	0.6	---
...	All other causes (Residual)	1,816	20.3	---
Hispanic, female, 90-94 years				
...	All causes	5,739	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,804	31.4	---
2	Malignant neoplasms (C00-C97)	499	8.7	---
3	Alzheimer's disease (G30)	465	8.1	---
4	Cerebrovascular diseases (I60-I69)	442	7.7	---
5	Chronic lower respiratory diseases (J40-J47)	220	3.8	---
6	Influenza and pneumonia (J09-J18)	213	3.7	---
7	Diabetes mellitus (E10-E14)	208	3.6	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	133	2.3	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	132	2.3	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	75	1.3	---
11	Septicemia (A40-A41)	74	1.3	---
12	Parkinson's disease (G20-G21)	62	1.1	---
13	Pneumonitis due to solids and liquids (J69)	52	0.9	---
14	Atherosclerosis (I70)	26	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	25	0.4	---
15	Chronic liver disease and cirrhosis (K70,K73-K74)	25	0.4	---
...	All other causes (Residual)	1,284	22.4	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 95-99 years				
...	All causes	2,625	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	898	34.2	---
2	Alzheimer's disease (G30)	254	9.7	---
3	Cerebrovascular diseases (I60-I69)	178	6.8	---
4	Malignant neoplasms (C00-C97)	154	5.9	---
5	Chronic lower respiratory diseases (J40-J47)	99	3.8	---
6	Diabetes mellitus (E10-E14)	94	3.6	---
7	Influenza and pneumonia (J09-J18)	90	3.4	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	52	2.0	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	45	1.7	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	43	1.6	---
11	Atherosclerosis (I70)	22	0.8	---
12	Septicemia (A40-A41)	19	0.7	---
13	Pneumonitis due to solids and liquids (J69)	16	0.6	---
14	Parkinson's disease (G20-G21)	14	0.5	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	12	0.5	---
...	All other causes (Residual)	635	24.2	---
Hispanic, female, 100 years and over				
...	All causes	681	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	226	33.2	---
2	Alzheimer's disease (G30)	67	9.8	---
3	Cerebrovascular diseases (I60-I69)	47	6.9	---
4	Influenza and pneumonia (J09-J18)	39	5.7	---
5	Chronic lower respiratory diseases (J40-J47)	26	3.8	---
6	Malignant neoplasms (C00-C97)	24	3.5	---
7	Diabetes mellitus (E10-E14)	17	2.5	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	16	2.3	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	2.1	---
10	Septicemia (A40-A41)	10	1.5	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8	1.2	---
12	Pneumonitis due to solids and liquids (J69)	7	1.0	---
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.7	---
13	Parkinson's disease (G20-G21)	5	0.7	---
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	3	0.4	---
...	All other causes (Residual)	167	24.5	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic white, both sexes, all ages				
...	All causes	1,969,916	100.0	984.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	483,973	24.6	241.8
2	Malignant neoplasms (C00-C97)	460,567	23.4	230.1
3	Chronic lower respiratory diseases (J40-J47)	122,887	6.2	61.4
4	Cerebrovascular diseases (I60-I69)	101,849	5.2	50.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	94,420	4.8	47.2
6	Alzheimer's disease (G30)	73,502	3.7	36.7
7	Diabetes mellitus (E10-E14)	47,746	2.4	23.9
8	Influenza and pneumonia (J09-J18)	40,244	2.0	20.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	36,963	1.9	18.5
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	32,010	1.6	16.0
11	Septicemia (A40-A41)	25,951	1.3	13.0
12	Chronic liver disease and cirrhosis (K70,K73-K74)	23,655	1.2	11.8
13	Parkinson's disease (G20-G21)	19,818	1.0	9.9
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	18,878	1.0	9.4
15	Pneumonitis due to solids and liquids (J69)	14,539	0.7	7.3
...	All other causes (Residual)	372,914	18.9	186.3
Non-Hispanic white, both sexes, 1-4 years				
...	All causes	2,139	100.0	24.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	744	34.8	8.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	229	10.7	2.6
3	Malignant neoplasms (C00-C97)	196	9.2	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	159	7.4	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	60	2.8	0.7
6	Influenza and pneumonia (J09-J18)	38	1.8	0.4
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	37	1.7	0.4
8	Septicemia (A40-A41)	31	1.4	0.4
9	Certain conditions originating in the perinatal period (P00-P96)	27	1.3	0.3
10	Cerebrovascular diseases (I60-I69)	23	1.1	0.3
11	Chronic lower respiratory diseases (J40-J47)	17	0.8	*
12	Meningitis (G00,G03)	13	0.6	*
13	Anemias (D50-D64)	11	0.5	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	9	0.4	*
15	Acute bronchitis and bronchiolitis (J20-J21)	6	0.3	*
...	All other causes (Residual)	539	25.2	6.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic white, both sexes, 5-9 years				
...	All causes	1,283	100.0	11.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	458	35.7	4.1
2	Malignant neoplasms (C00-C97)	243	18.9	2.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	89	6.9	0.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	50	3.9	0.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	30	2.3	0.3
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	22	1.7	0.2
6	Influenza and pneumonia (J09-J18)	22	1.7	0.2
8	Chronic lower respiratory diseases (J40-J47)	17	1.3	*
9	Septicemia (A40-A41)	16	1.2	*
9	Cerebrovascular diseases (I60-I69)	16	1.2	*
11	Certain conditions originating in the perinatal period (P00-P96)	10	0.8	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.5	*
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5	0.4	*
14	Meningitis (G00,G03)	4	0.3	*
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	4	0.3	*
...	All other causes (Residual)	290	22.6	2.6
Non-Hispanic white, both sexes, 10-14 years				
...	All causes	1,627	100.0	13.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	542	33.3	4.6
2	Malignant neoplasms (C00-C97)	257	15.8	2.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	161	9.9	1.4
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	75	4.6	0.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	60	3.7	0.5
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	44	2.7	0.4
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	25	1.5	0.2
8	Cerebrovascular diseases (I60-I69)	23	1.4	0.2
9	Influenza and pneumonia (J09-J18)	20	1.2	0.2
10	Diabetes mellitus (E10-E14)	17	1.0	*
10	Chronic lower respiratory diseases (J40-J47)	17	1.0	*
12	Septicemia (A40-A41)	15	0.9	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.6	*
14	Certain conditions originating in the perinatal period (P00-P96)	6	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.3	*
...	All other causes (Residual)	351	21.6	3.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic white, both sexes, 15-19 years				
...	All causes	6,013	100.0	47.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,063	50.9	24.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,138	18.9	9.0
3	Malignant neoplasms (C00-C97)	333	5.5	2.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	229	3.8	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	176	2.9	1.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	116	1.9	0.9
7	Cerebrovascular diseases (I60-I69)	47	0.8	0.4
8	Diabetes mellitus (E10-E14)	29	0.5	0.2
9	Chronic lower respiratory diseases (J40-J47)	28	0.5	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	26	0.4	0.2
11	Influenza and pneumonia (J09-J18)	25	0.4	0.2
11	Pregnancy, childbirth and the puerperium (O00-O99)	25	0.4	0.2
13	Septicemia (A40-A41)	24	0.4	0.2
14	Meningitis (G00,G03)	10	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	0.2	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	10	0.2	*
...	All other causes (Residual)	724	12.0	5.7
Non-Hispanic white, both sexes, 20-24 years				
...	All causes	10,834	100.0	85.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,444	50.2	42.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,052	18.9	16.2
3	Malignant neoplasms (C00-C97)	568	5.2	4.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	519	4.8	4.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	344	3.2	2.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	129	1.2	1.0
7	Influenza and pneumonia (J09-J18)	85	0.8	0.7
8	Diabetes mellitus (E10-E14)	66	0.6	0.5
9	Cerebrovascular diseases (I60-I69)	53	0.5	0.4
10	Septicemia (A40-A41)	52	0.5	0.4
11	Pregnancy, childbirth and the puerperium (O00-O99)	48	0.4	0.4
12	Chronic lower respiratory diseases (J40-J47)	40	0.4	0.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	29	0.3	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21	0.2	0.2
15	Pneumonitis due to solids and liquids (J69)	20	0.2	0.2
15	Legal intervention (Y35,Y89.0)	20	0.2	0.2
...	All other causes (Residual)	1,344	12.4	10.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic white, both sexes, 25-29 years				
...	All causes	12,207	100.0	98.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,435	44.5	43.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,232	18.3	17.9
3	Malignant neoplasms (C00-C97)	855	7.0	6.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	621	5.1	5.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	497	4.1	4.0
6	Diabetes mellitus (E10-E14)	128	1.0	1.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	128	1.0	1.0
8	Influenza and pneumonia (J09-J18)	97	0.8	0.8
9	Cerebrovascular diseases (I60-I69)	85	0.7	0.7
10	Pregnancy, childbirth and the puerperium (O00-O99)	80	0.7	0.6
11	Septicemia (A40-A41)	74	0.6	0.6
12	Chronic liver disease and cirrhosis (K70,K73-K74)	69	0.6	0.6
13	Chronic lower respiratory diseases (J40-J47)	58	0.5	0.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	41	0.3	0.3
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	41	0.3	0.3
...	All other causes (Residual)	1,766	14.5	14.2
Non-Hispanic white, both sexes, 30-34 years				
...	All causes	13,279	100.0	113.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,990	37.6	42.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,103	15.8	18.0
3	Malignant neoplasms (C00-C97)	1,299	9.8	11.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,183	8.9	10.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	391	2.9	3.3
6	Chronic liver disease and cirrhosis (K70,K73-K74)	213	1.6	1.8
7	Diabetes mellitus (E10-E14)	197	1.5	1.7
8	Cerebrovascular diseases (I60-I69)	175	1.3	1.5
9	Influenza and pneumonia (J09-J18)	129	1.0	1.1
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	126	0.9	1.1
11	Septicemia (A40-A41)	93	0.7	0.8
12	Human immunodeficiency virus (HIV) disease (B20-B24)	86	0.6	0.7
13	Chronic lower respiratory diseases (J40-J47)	76	0.6	0.7
14	Pregnancy, childbirth and the puerperium (O00-O99)	69	0.5	0.6
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	64	0.5	0.5
...	All other causes (Residual)	2,085	15.7	17.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic white, both sexes, 35-39 years				
...	All causes	17,508	100.0	144.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,094	29.1	41.9
2	Malignant neoplasms (C00-C97)	2,464	14.1	20.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,454	14.0	20.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,130	12.2	17.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	463	2.6	3.8
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	393	2.2	3.2
7	Diabetes mellitus (E10-E14)	333	1.9	2.7
8	Cerebrovascular diseases (I60-I69)	287	1.6	2.4
9	Influenza and pneumonia (J09-J18)	186	1.1	1.5
10	Septicemia (A40-A41)	175	1.0	1.4
11	Human immunodeficiency virus (HIV) disease (B20-B24)	173	1.0	1.4
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	162	0.9	1.3
13	Chronic lower respiratory diseases (J40-J47)	136	0.8	1.1
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	130	0.7	1.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	64	0.4	0.5
...	All other causes (Residual)	2,864	16.4	23.6
Non-Hispanic white, both sexes, 40-44 years				
...	All causes	27,491	100.0	205.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,886	21.4	44.0
2	Malignant neoplasms (C00-C97)	5,173	18.8	38.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,436	16.1	33.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,905	10.6	21.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,111	4.0	8.3
6	Diabetes mellitus (E10-E14)	633	2.3	4.7
7	Cerebrovascular diseases (I60-I69)	628	2.3	4.7
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	478	1.7	3.6
9	Human immunodeficiency virus (HIV) disease (B20-B24)	331	1.2	2.5
10	Chronic lower respiratory diseases (J40-J47)	319	1.2	2.4
11	Influenza and pneumonia (J09-J18)	306	1.1	2.3
12	Septicemia (A40-A41)	281	1.0	2.1
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	204	0.7	1.5
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	172	0.6	1.3
15	Viral hepatitis (B15-B19)	156	0.6	1.2
...	All other causes (Residual)	4,472	16.3	33.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic white, both sexes, 45-49 years				
...	All causes	50,736	100.0	327.0
1	Malignant neoplasms (C00-C97)	12,255	24.2	79.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,496	18.7	61.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,743	15.3	49.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,806	7.5	24.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,598	5.1	16.7
6	Diabetes mellitus (E10-E14)	1,284	2.5	8.3
7	Cerebrovascular diseases (I60-I69)	1,231	2.4	7.9
8	Chronic lower respiratory diseases (J40-J47)	1,054	2.1	6.8
9	Septicemia (A40-A41)	557	1.1	3.6
10	Human immunodeficiency virus (HIV) disease (B20-B24)	551	1.1	3.6
11	Viral hepatitis (B15-B19)	519	1.0	3.3
12	Influenza and pneumonia (J09-J18)	517	1.0	3.3
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	490	1.0	3.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	399	0.8	2.6
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	250	0.5	1.6
...	All other causes (Residual)	7,986	15.7	51.5
Non-Hispanic white, both sexes, 50-54 years				
...	All causes	77,298	100.0	485.2
1	Malignant neoplasms (C00-C97)	23,534	30.4	147.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,586	20.2	97.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,516	9.7	47.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,949	5.1	24.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	3,631	4.7	22.8
6	Chronic lower respiratory diseases (J40-J47)	2,501	3.2	15.7
7	Diabetes mellitus (E10-E14)	2,099	2.7	13.2
8	Cerebrovascular diseases (I60-I69)	1,998	2.6	12.5
9	Viral hepatitis (B15-B19)	1,097	1.4	6.9
10	Septicemia (A40-A41)	902	1.2	5.7
11	Influenza and pneumonia (J09-J18)	790	1.0	5.0
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	718	0.9	4.5
13	Human immunodeficiency virus (HIV) disease (B20-B24)	474	0.6	3.0
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	462	0.6	2.9
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	459	0.6	2.9
...	All other causes (Residual)	11,582	15.0	72.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic white, both sexes, 55-59 years				
...	All causes	100,742	100.0	691.4
1	Malignant neoplasms (C00-C97)	34,673	34.4	238.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	21,483	21.3	147.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,089	6.0	41.8
4	Chronic lower respiratory diseases (J40-J47)	4,413	4.4	30.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	3,944	3.9	27.1
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,370	3.3	23.1
7	Diabetes mellitus (E10-E14)	3,167	3.1	21.7
8	Cerebrovascular diseases (I60-I69)	2,679	2.7	18.4
9	Viral hepatitis (B15-B19)	1,389	1.4	9.5
10	Septicemia (A40-A41)	1,336	1.3	9.2
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,127	1.1	7.7
12	Influenza and pneumonia (J09-J18)	1,118	1.1	7.7
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	624	0.6	4.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	412	0.4	2.8
15	Human immunodeficiency virus (HIV) disease (B20-B24)	361	0.4	2.5
...	All other causes (Residual)	14,557	14.4	99.9
Non-Hispanic white, both sexes, 60-64 years				
...	All causes	128,523	100.0	995.3
1	Malignant neoplasms (C00-C97)	48,578	37.8	376.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	27,793	21.6	215.2
3	Chronic lower respiratory diseases (J40-J47)	7,729	6.0	59.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,742	3.7	36.7
5	Diabetes mellitus (E10-E14)	4,297	3.3	33.3
6	Cerebrovascular diseases (I60-I69)	3,985	3.1	30.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	3,301	2.6	25.6
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,364	1.8	18.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,764	1.4	13.7
10	Septicemia (A40-A41)	1,729	1.3	13.4
11	Influenza and pneumonia (J09-J18)	1,487	1.2	11.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	861	0.7	6.7
13	Viral hepatitis (B15-B19)	760	0.6	5.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	702	0.5	5.4
15	Aortic aneurysm and dissection (I71)	592	0.5	4.6
...	All other causes (Residual)	17,839	13.9	138.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic white, both sexes, 65-69 years				
...	All causes	148,016	100.0	1,518.9
1	Malignant neoplasms (C00-C97)	56,517	38.2	580.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	31,287	21.1	321.1
3	Chronic lower respiratory diseases (J40-J47)	12,157	8.2	124.8
4	Cerebrovascular diseases (I60-I69)	5,227	3.5	53.6
5	Diabetes mellitus (E10-E14)	4,688	3.2	48.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,745	2.5	38.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,491	1.7	25.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,380	1.6	24.4
9	Septicemia (A40-A41)	2,222	1.5	22.8
10	Influenza and pneumonia (J09-J18)	1,877	1.3	19.3
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,532	1.0	15.7
12	Alzheimer's disease (G30)	1,018	0.7	10.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,018	0.7	10.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	898	0.6	9.2
15	Aortic aneurysm and dissection (I71)	790	0.5	8.1
...	All other causes (Residual)	20,169	13.6	207.0
Non-Hispanic white, both sexes, 70-74 years				
...	All causes	171,789	100.0	2,353.0
1	Malignant neoplasms (C00-C97)	59,963	34.9	821.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	36,802	21.4	504.1
3	Chronic lower respiratory diseases (J40-J47)	16,201	9.4	221.9
4	Cerebrovascular diseases (I60-I69)	7,446	4.3	102.0
5	Diabetes mellitus (E10-E14)	5,103	3.0	69.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,836	2.2	52.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,225	1.9	44.2
8	Influenza and pneumonia (J09-J18)	2,584	1.5	35.4
9	Alzheimer's disease (G30)	2,544	1.5	34.8
10	Septicemia (A40-A41)	2,521	1.5	34.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1,938	1.1	26.5
12	Parkinson's disease (G20-G21)	1,570	0.9	21.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,256	0.7	17.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,243	0.7	17.0
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,151	0.7	15.8
...	All other causes (Residual)	24,406	14.2	334.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic white, both sexes, 75-79 years				
...	All causes	223,128	100.0	3,786.8
1	Malignant neoplasms (C00-C97)	64,712	29.0	1,098.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	50,854	22.8	863.1
3	Chronic lower respiratory diseases (J40-J47)	20,493	9.2	347.8
4	Cerebrovascular diseases (I60-I69)	11,797	5.3	200.2
5	Alzheimer's disease (G30)	6,758	3.0	114.7
6	Diabetes mellitus (E10-E14)	6,152	2.8	104.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,976	2.2	84.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,666	2.1	79.2
9	Influenza and pneumonia (J09-J18)	4,076	1.8	69.2
10	Parkinson's disease (G20-G21)	3,430	1.5	58.2
11	Septicemia (A40-A41)	3,353	1.5	56.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,907	0.9	32.4
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,722	0.8	29.2
14	Chronic liver disease and cirrhosis (K70,K73-K74)	1,597	0.7	27.1
15	Pneumonitis due to solids and liquids (J69)	1,496	0.7	25.4
...	All other causes (Residual)	35,139	15.7	596.4
Non-Hispanic white, both sexes, 80-84 years				
...	All causes	297,855	100.0	6,246.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	75,072	25.2	1,574.3
2	Malignant neoplasms (C00-C97)	66,773	22.4	1,400.2
3	Chronic lower respiratory diseases (J40-J47)	23,334	7.8	489.3
4	Cerebrovascular diseases (I60-I69)	18,654	6.3	391.2
5	Alzheimer's disease (G30)	14,176	4.8	297.3
6	Diabetes mellitus (E10-E14)	7,391	2.5	155.0
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,052	2.4	147.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,891	2.3	144.5
9	Influenza and pneumonia (J09-J18)	6,699	2.2	140.5
10	Parkinson's disease (G20-G21)	5,364	1.8	112.5
11	Septicemia (A40-A41)	4,402	1.5	92.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,985	1.0	62.6
13	Pneumonitis due to solids and liquids (J69)	2,772	0.9	58.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,354	0.8	49.4
15	Aortic aneurysm and dissection (I71)	1,672	0.6	35.1
...	All other causes (Residual)	52,264	17.5	1,096.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic white, both sexes, 85-89 years				
...	All causes	329,386	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	94,251	28.6	---
2	Malignant neoplasms (C00-C97)	51,881	15.8	---
3	Cerebrovascular diseases (I60-I69)	23,165	7.0	---
4	Alzheimer's disease (G30)	21,496	6.5	---
5	Chronic lower respiratory diseases (J40-J47)	20,324	6.2	---
6	Influenza and pneumonia (J09-J18)	8,799	2.7	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,152	2.5	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,800	2.4	---
9	Diabetes mellitus (E10-E14)	6,904	2.1	---
10	Parkinson's disease (G20-G21)	5,217	1.6	---
11	Septicemia (A40-A41)	4,467	1.4	---
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,903	1.2	---
13	Pneumonitis due to solids and liquids (J69)	3,464	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,341	0.7	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1,582	0.5	---
...	All other causes (Residual)	65,640	19.9	---
Non-Hispanic white, both sexes, 90-94 years				
...	All causes	229,162	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	73,683	32.2	---
2	Malignant neoplasms (C00-C97)	23,397	10.2	---
3	Alzheimer's disease (G30)	17,818	7.8	---
4	Cerebrovascular diseases (I60-I69)	16,749	7.3	---
5	Chronic lower respiratory diseases (J40-J47)	10,403	4.5	---
6	Influenza and pneumonia (J09-J18)	7,212	3.1	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,662	2.5	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,228	2.3	---
9	Diabetes mellitus (E10-E14)	3,902	1.7	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,311	1.4	---
11	Pneumonitis due to solids and liquids (J69)	2,742	1.2	---
12	Septicemia (A40-A41)	2,587	1.1	---
13	Parkinson's disease (G20-G21)	2,482	1.1	---
14	Atherosclerosis (I70)	1,390	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,357	0.6	---
...	All other causes (Residual)	51,239	22.4	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic white, both sexes, 95-99 years				
...	All causes	91,531	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	31,763	34.7	---
2	Alzheimer's disease (G30)	7,553	8.3	---
3	Cerebrovascular diseases (I60-I69)	6,421	7.0	---
4	Malignant neoplasms (C00-C97)	6,124	6.7	---
5	Influenza and pneumonia (J09-J18)	3,278	3.6	---
6	Chronic lower respiratory diseases (J40-J47)	3,114	3.4	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,267	2.5	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,902	2.1	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,677	1.8	---
10	Diabetes mellitus (E10-E14)	1,198	1.3	---
11	Pneumonitis due to solids and liquids (J69)	1,053	1.2	---
12	Septicemia (A40-A41)	904	1.0	---
13	Atherosclerosis (I70)	679	0.7	---
14	Parkinson's disease (G20-G21)	568	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	427	0.5	---
...	All other causes (Residual)	22,603	24.7	---
Non-Hispanic white, both sexes, 100 years and over				
...	All causes	18,282	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,692	36.6	---
2	Alzheimer's disease (G30)	1,394	7.6	---
3	Cerebrovascular diseases (I60-I69)	1,106	6.0	---
4	Influenza and pneumonia (J09-J18)	828	4.5	---
5	Malignant neoplasms (C00-C97)	734	4.0	---
6	Chronic lower respiratory diseases (J40-J47)	440	2.4	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	424	2.3	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	411	2.2	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	342	1.9	---
10	Pneumonitis due to solids and liquids (J69)	205	1.1	---
11	Atherosclerosis (I70)	191	1.0	---
12	Diabetes mellitus (E10-E14)	156	0.9	---
13	Septicemia (A40-A41)	136	0.7	---
14	Anemias (D50-D64)	81	0.4	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	68	0.4	---
...	All other causes (Residual)	5,074	27.8	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic white, male, all ages				
...	All causes	971,604	100.0	987.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	247,765	25.5	251.8
2	Malignant neoplasms (C00-C97)	241,816	24.9	245.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	57,747	5.9	58.7
4	Chronic lower respiratory diseases (J40-J47)	57,390	5.9	58.3
5	Cerebrovascular diseases (I60-I69)	40,040	4.1	40.7
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	25,238	2.6	25.7
7	Diabetes mellitus (E10-E14)	25,134	2.6	25.5
8	Alzheimer's disease (G30)	22,278	2.3	22.6
9	Influenza and pneumonia (J09-J18)	18,657	1.9	19.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18,503	1.9	18.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	15,270	1.6	15.5
12	Septicemia (A40-A41)	11,963	1.2	12.2
13	Parkinson's disease (G20-G21)	11,627	1.2	11.8
14	Pneumonitis due to solids and liquids (J69)	7,862	0.8	8.0
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7,476	0.8	7.6
...	All other causes (Residual)	162,838	16.8	165.5
Non-Hispanic white, male, 1-4 years				
...	All causes	1,219	100.0	27.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	456	37.4	10.3
2	Malignant neoplasms (C00-C97)	109	8.9	2.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	101	8.3	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	90	7.4	2.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	37	3.0	0.8
6	Influenza and pneumonia (J09-J18)	20	1.6	0.5
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	1.4	*
7	Cerebrovascular diseases (I60-I69)	17	1.4	*
7	Certain conditions originating in the perinatal period (P00-P96)	17	1.4	*
10	Septicemia (A40-A41)	13	1.1	*
11	Chronic lower respiratory diseases (J40-J47)	10	0.8	*
12	Meningitis (G00,G03)	9	0.7	*
13	Anemias (D50-D64)	8	0.7	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.5	*
15	Acute bronchitis and bronchiolitis (J20-J21)	5	0.4	*
...	All other causes (Residual)	304	24.9	6.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic white, male, 5-9 years				
...	All causes	722	100.0	12.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	293	40.6	5.1
2	Malignant neoplasms (C00-C97)	137	19.0	2.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	38	5.3	0.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	27	3.7	0.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	12	1.7	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	1.4	*
7	Septicemia (A40-A41)	9	1.2	*
7	Certain conditions originating in the perinatal period (P00-P96)	9	1.2	*
9	Cerebrovascular diseases (I60-I69)	8	1.1	*
10	Chronic lower respiratory diseases (J40-J47)	6	0.8	*
11	Influenza and pneumonia (J09-J18)	5	0.7	*
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5	0.7	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.4	*
14	Meningitis (G00,G03)	2	0.3	*
14	Pneumonitis due to solids and liquids (J69)	2	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.3	*
...	All other causes (Residual)	154	21.3	2.7
Non-Hispanic white, male, 10-14 years				
...	All causes	969	100.0	16.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	346	35.7	5.7
2	Malignant neoplasms (C00-C97)	140	14.4	2.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	123	12.7	2.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	36	3.7	0.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	29	3.0	0.5
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	28	2.9	0.5
7	Cerebrovascular diseases (I60-I69)	15	1.5	*
8	Diabetes mellitus (E10-E14)	11	1.1	*
9	Influenza and pneumonia (J09-J18)	10	1.0	*
10	Septicemia (A40-A41)	9	0.9	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.9	*
12	Chronic lower respiratory diseases (J40-J47)	8	0.8	*
13	Certain conditions originating in the perinatal period (P00-P96)	3	0.3	*
14	Chronic liver disease and cirrhosis (K70,K73-K74)	2	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.2	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.2	*
...	All other causes (Residual)	196	20.2	3.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic white, male, 15-19 years				
...	All causes	4,152	100.0	63.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,117	51.0	32.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	921	22.2	14.2
3	Malignant neoplasms (C00-C97)	195	4.7	3.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	153	3.7	2.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	122	2.9	1.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	70	1.7	1.1
7	Cerebrovascular diseases (I60-I69)	30	0.7	0.5
8	Chronic lower respiratory diseases (J40-J47)	16	0.4	*
9	Diabetes mellitus (E10-E14)	15	0.4	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	0.3	*
10	Influenza and pneumonia (J09-J18)	14	0.3	*
12	Septicemia (A40-A41)	11	0.3	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.1	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.1	*
15	Meningitis (G00,G03)	5	0.1	*
15	Aortic aneurysm and dissection (I71)	5	0.1	*
...	All other causes (Residual)	452	10.9	7.0
Non-Hispanic white, male, 20-24 years				
...	All causes	7,927	100.0	123.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,055	51.2	63.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,717	21.7	26.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	375	4.7	5.8
4	Malignant neoplasms (C00-C97)	370	4.7	5.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	260	3.3	4.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	80	1.0	1.2
7	Influenza and pneumonia (J09-J18)	45	0.6	0.7
8	Diabetes mellitus (E10-E14)	39	0.5	0.6
9	Cerebrovascular diseases (I60-I69)	31	0.4	0.5
10	Septicemia (A40-A41)	27	0.3	0.4
11	Chronic lower respiratory diseases (J40-J47)	26	0.3	0.4
12	Legal intervention (Y35,Y89.0)	20	0.3	0.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	0.2	*
14	Aortic aneurysm and dissection (I71)	13	0.2	*
14	Pneumonitis due to solids and liquids (J69)	13	0.2	*
...	All other causes (Residual)	838	10.6	13.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic white, male, 25-29 years				
...	All causes	8,637	100.0	137.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,060	47.0	64.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,822	21.1	29.0
3	Malignant neoplasms (C00-C97)	479	5.5	7.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	438	5.1	7.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	340	3.9	5.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	74	0.9	1.2
7	Diabetes mellitus (E10-E14)	71	0.8	1.1
8	Influenza and pneumonia (J09-J18)	55	0.6	0.9
9	Chronic liver disease and cirrhosis (K70,K73-K74)	40	0.5	0.6
10	Cerebrovascular diseases (I60-I69)	38	0.4	0.6
11	Septicemia (A40-A41)	34	0.4	0.5
12	Chronic lower respiratory diseases (J40-J47)	33	0.4	0.5
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	30	0.3	0.5
14	Human immunodeficiency virus (HIV) disease (B20-B24)	26	0.3	0.4
15	Legal intervention (Y35,Y89.0)	22	0.3	0.4
...	All other causes (Residual)	1,075	12.4	17.1
Non-Hispanic white, male, 30-34 years				
...	All causes	8,846	100.0	150.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,578	40.4	60.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,658	18.7	28.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	812	9.2	13.8
4	Malignant neoplasms (C00-C97)	627	7.1	10.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	265	3.0	4.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	132	1.5	2.2
7	Diabetes mellitus (E10-E14)	115	1.3	2.0
8	Cerebrovascular diseases (I60-I69)	78	0.9	1.3
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	72	0.8	1.2
10	Influenza and pneumonia (J09-J18)	69	0.8	1.2
11	Human immunodeficiency virus (HIV) disease (B20-B24)	62	0.7	1.1
12	Septicemia (A40-A41)	49	0.6	0.8
13	Chronic lower respiratory diseases (J40-J47)	42	0.5	0.7
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	37	0.4	0.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	33	0.4	0.6
...	All other causes (Residual)	1,217	13.8	20.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic white, male, 35-39 years				
...	All causes	11,045	100.0	180.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,454	31.3	56.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,887	17.1	30.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,494	13.5	24.5
4	Malignant neoplasms (C00-C97)	1,135	10.3	18.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	289	2.6	4.7
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	252	2.3	4.1
7	Diabetes mellitus (E10-E14)	197	1.8	3.2
8	Cerebrovascular diseases (I60-I69)	159	1.4	2.6
9	Human immunodeficiency virus (HIV) disease (B20-B24)	128	1.2	2.1
10	Influenza and pneumonia (J09-J18)	98	0.9	1.6
11	Septicemia (A40-A41)	97	0.9	1.6
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	90	0.8	1.5
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	68	0.6	1.1
14	Chronic lower respiratory diseases (J40-J47)	60	0.5	1.0
15	Aortic aneurysm and dissection (I71)	45	0.4	0.7
...	All other causes (Residual)	1,592	14.4	26.1
Non-Hispanic white, male, 40-44 years				
...	All causes	17,025	100.0	253.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,943	23.2	58.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,193	18.8	47.6
3	Malignant neoplasms (C00-C97)	2,327	13.7	34.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,213	13.0	33.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	723	4.2	10.8
6	Diabetes mellitus (E10-E14)	376	2.2	5.6
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	318	1.9	4.7
8	Cerebrovascular diseases (I60-I69)	315	1.9	4.7
9	Human immunodeficiency virus (HIV) disease (B20-B24)	265	1.6	4.0
10	Influenza and pneumonia (J09-J18)	161	0.9	2.4
11	Chronic lower respiratory diseases (J40-J47)	142	0.8	2.1
12	Septicemia (A40-A41)	138	0.8	2.1
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	110	0.6	1.6
14	Viral hepatitis (B15-B19)	102	0.6	1.5
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	101	0.6	1.5
...	All other causes (Residual)	2,598	15.3	38.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic white, male, 45-49 years				
...	All causes	31,406	100.0	406.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,873	21.9	89.1
2	Malignant neoplasms (C00-C97)	5,986	19.1	77.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,124	16.3	66.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,928	9.3	37.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,699	5.4	22.0
6	Diabetes mellitus (E10-E14)	850	2.7	11.0
7	Cerebrovascular diseases (I60-I69)	680	2.2	8.8
8	Human immunodeficiency virus (HIV) disease (B20-B24)	479	1.5	6.2
9	Chronic lower respiratory diseases (J40-J47)	452	1.4	5.9
10	Viral hepatitis (B15-B19)	345	1.1	4.5
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	316	1.0	4.1
12	Septicemia (A40-A41)	287	0.9	3.7
13	Influenza and pneumonia (J09-J18)	278	0.9	3.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	243	0.8	3.1
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	154	0.5	2.0
...	All other causes (Residual)	4,712	15.0	61.1
Non-Hispanic white, male, 50-54 years				
...	All causes	47,889	100.0	607.1
1	Malignant neoplasms (C00-C97)	12,299	25.7	155.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,420	23.8	144.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,972	10.4	63.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,996	6.3	38.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,513	5.2	31.9
6	Diabetes mellitus (E10-E14)	1,356	2.8	17.2
7	Chronic lower respiratory diseases (J40-J47)	1,196	2.5	15.2
8	Cerebrovascular diseases (I60-I69)	1,131	2.4	14.3
9	Viral hepatitis (B15-B19)	763	1.6	9.7
10	Septicemia (A40-A41)	503	1.1	6.4
11	Influenza and pneumonia (J09-J18)	435	0.9	5.5
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	425	0.9	5.4
13	Human immunodeficiency virus (HIV) disease (B20-B24)	398	0.8	5.0
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	329	0.7	4.2
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	317	0.7	4.0
...	All other causes (Residual)	6,836	14.3	86.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic white, male, 55-59 years				
...	All causes	62,997	100.0	881.6
1	Malignant neoplasms (C00-C97)	19,160	30.4	268.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,748	25.0	220.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,133	6.6	57.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	2,852	4.5	39.9
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,549	4.0	35.7
6	Chronic lower respiratory diseases (J40-J47)	2,283	3.6	31.9
7	Diabetes mellitus (E10-E14)	2,024	3.2	28.3
8	Cerebrovascular diseases (I60-I69)	1,571	2.5	22.0
9	Viral hepatitis (B15-B19)	1,010	1.6	14.1
10	Septicemia (A40-A41)	709	1.1	9.9
11	Influenza and pneumonia (J09-J18)	666	1.1	9.3
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	645	1.0	9.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	414	0.7	5.8
14	Human immunodeficiency virus (HIV) disease (B20-B24)	308	0.5	4.3
15	Aortic aneurysm and dissection (I71)	273	0.4	3.8
...	All other causes (Residual)	8,652	13.7	121.1
Non-Hispanic white, male, 60-64 years				
...	All causes	77,555	100.0	1,233.3
1	Malignant neoplasms (C00-C97)	27,080	34.9	430.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	19,443	25.1	309.2
3	Chronic lower respiratory diseases (J40-J47)	3,969	5.1	63.1
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,227	4.2	51.3
5	Diabetes mellitus (E10-E14)	2,670	3.4	42.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	2,260	2.9	35.9
7	Cerebrovascular diseases (I60-I69)	2,243	2.9	35.7
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,806	2.3	28.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,014	1.3	16.1
10	Septicemia (A40-A41)	916	1.2	14.6
11	Influenza and pneumonia (J09-J18)	862	1.1	13.7
12	Viral hepatitis (B15-B19)	560	0.7	8.9
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	557	0.7	8.9
14	Aortic aneurysm and dissection (I71)	435	0.6	6.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	421	0.5	6.7
...	All other causes (Residual)	10,092	13.0	160.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic white, male, 65-69 years				
...	All causes	85,858	100.0	1,844.8
1	Malignant neoplasms (C00-C97)	31,310	36.5	672.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,695	24.1	444.7
3	Chronic lower respiratory diseases (J40-J47)	6,180	7.2	132.8
4	Cerebrovascular diseases (I60-I69)	2,867	3.3	61.6
5	Diabetes mellitus (E10-E14)	2,806	3.3	60.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,415	2.8	51.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,612	1.9	34.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,354	1.6	29.1
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,210	1.4	26.0
10	Septicemia (A40-A41)	1,167	1.4	25.1
11	Influenza and pneumonia (J09-J18)	1,077	1.3	23.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	583	0.7	12.5
13	Aortic aneurysm and dissection (I71)	560	0.7	12.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	529	0.6	11.4
15	Alzheimer's disease (G30)	454	0.5	9.8
...	All other causes (Residual)	11,039	12.9	237.2
Non-Hispanic white, male, 70-74 years				
...	All causes	95,709	100.0	2,822.4
1	Malignant neoplasms (C00-C97)	32,928	34.4	971.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	22,923	24.0	676.0
3	Chronic lower respiratory diseases (J40-J47)	8,106	8.5	239.0
4	Cerebrovascular diseases (I60-I69)	3,781	4.0	111.5
5	Diabetes mellitus (E10-E14)	2,883	3.0	85.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,353	2.5	69.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,754	1.8	51.7
8	Influenza and pneumonia (J09-J18)	1,425	1.5	42.0
9	Septicemia (A40-A41)	1,268	1.3	37.4
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,139	1.2	33.6
11	Alzheimer's disease (G30)	1,119	1.2	33.0
12	Parkinson's disease (G20-G21)	1,041	1.1	30.7
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	972	1.0	28.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	718	0.8	21.2
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	653	0.7	19.3
...	All other causes (Residual)	12,646	13.2	372.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic white, male, 75-79 years				
...	All causes	117,699	100.0	4,518.2
1	Malignant neoplasms (C00-C97)	34,950	29.7	1,341.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	29,062	24.7	1,115.6
3	Chronic lower respiratory diseases (J40-J47)	9,911	8.4	380.5
4	Cerebrovascular diseases (I60-I69)	5,473	4.6	210.1
5	Diabetes mellitus (E10-E14)	3,358	2.9	128.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,817	2.4	108.1
7	Alzheimer's disease (G30)	2,702	2.3	103.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,541	2.2	97.5
9	Parkinson's disease (G20-G21)	2,218	1.9	85.1
10	Influenza and pneumonia (J09-J18)	2,182	1.9	83.8
11	Septicemia (A40-A41)	1,625	1.4	62.4
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	978	0.8	37.5
13	Chronic liver disease and cirrhosis (K70,K73-K74)	892	0.8	34.2
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	878	0.7	33.7
15	Pneumonitis due to solids and liquids (J69)	876	0.7	33.6
...	All other causes (Residual)	17,236	14.6	661.7
Non-Hispanic white, male, 80-84 years				
...	All causes	144,024	100.0	7,459.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	38,584	26.8	1,998.5
2	Malignant neoplasms (C00-C97)	34,647	24.1	1,794.6
3	Chronic lower respiratory diseases (J40-J47)	10,951	7.6	567.2
4	Cerebrovascular diseases (I60-I69)	7,618	5.3	394.6
5	Alzheimer's disease (G30)	5,185	3.6	268.6
6	Diabetes mellitus (E10-E14)	3,605	2.5	186.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,594	2.5	186.2
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,546	2.5	183.7
9	Parkinson's disease (G20-G21)	3,338	2.3	172.9
10	Influenza and pneumonia (J09-J18)	3,324	2.3	172.2
11	Septicemia (A40-A41)	2,051	1.4	106.2
12	Pneumonitis due to solids and liquids (J69)	1,604	1.1	83.1
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,263	0.9	65.4
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,180	0.8	61.1
15	Aortic aneurysm and dissection (I71)	855	0.6	44.3
...	All other causes (Residual)	22,679	15.7	1,174.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic white, male, 85-89 years				
...	All causes	137,518	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	40,963	29.8	---
2	Malignant neoplasms (C00-C97)	25,149	18.3	---
3	Chronic lower respiratory diseases (J40-J47)	8,934	6.5	---
4	Cerebrovascular diseases (I60-I69)	7,924	5.8	---
5	Alzheimer's disease (G30)	6,658	4.8	---
6	Influenza and pneumonia (J09-J18)	3,940	2.9	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,695	2.7	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,689	2.7	---
9	Diabetes mellitus (E10-E14)	2,985	2.2	---
10	Parkinson's disease (G20-G21)	2,863	2.1	---
11	Pneumonitis due to solids and liquids (J69)	1,857	1.4	---
12	Septicemia (A40-A41)	1,820	1.3	---
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,342	1.0	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,143	0.8	---
15	Aortic aneurysm and dissection (I71)	672	0.5	---
...	All other causes (Residual)	23,884	17.4	---
Non-Hispanic white, male, 90-94 years				
...	All causes	78,623	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	26,226	33.4	---
2	Malignant neoplasms (C00-C97)	10,322	13.1	---
3	Cerebrovascular diseases (I60-I69)	4,662	5.9	---
4	Alzheimer's disease (G30)	4,454	5.7	---
5	Chronic lower respiratory diseases (J40-J47)	3,992	5.1	---
6	Influenza and pneumonia (J09-J18)	2,813	3.6	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,213	2.8	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,114	2.7	---
9	Diabetes mellitus (E10-E14)	1,415	1.8	---
10	Pneumonitis due to solids and liquids (J69)	1,352	1.7	---
11	Parkinson's disease (G20-G21)	1,176	1.5	---
12	Septicemia (A40-A41)	914	1.2	---
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	896	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	552	0.7	---
15	Atherosclerosis (I70)	447	0.6	---
...	All other causes (Residual)	15,075	19.2	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic white, male, 95-99 years				
...	All causes	22,681	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,241	36.3	---
2	Malignant neoplasms (C00-C97)	2,247	9.9	---
3	Alzheimer's disease (G30)	1,267	5.6	---
4	Cerebrovascular diseases (I60-I69)	1,228	5.4	---
5	Influenza and pneumonia (J09-J18)	954	4.2	---
6	Chronic lower respiratory diseases (J40-J47)	953	4.2	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	663	2.9	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	646	2.8	---
9	Pneumonitis due to solids and liquids (J69)	414	1.8	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	339	1.5	---
11	Diabetes mellitus (E10-E14)	335	1.5	---
12	Septicemia (A40-A41)	245	1.1	---
13	Parkinson's disease (G20-G21)	219	1.0	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	144	0.6	---
15	Atherosclerosis (I70)	143	0.6	---
...	All other causes (Residual)	4,643	20.5	---
Non-Hispanic white, male, 100 years and over				
...	All causes	2,919	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,091	37.4	---
2	Malignant neoplasms (C00-C97)	198	6.8	---
3	Influenza and pneumonia (J09-J18)	184	6.3	---
4	Cerebrovascular diseases (I60-I69)	144	4.9	---
5	Alzheimer's disease (G30)	124	4.2	---
6	Chronic lower respiratory diseases (J40-J47)	112	3.8	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	92	3.2	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	80	2.7	---
9	Pneumonitis due to solids and liquids (J69)	60	2.1	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	52	1.8	---
11	Atherosclerosis (I70)	27	0.9	---
12	Septicemia (A40-A41)	24	0.8	---
13	Diabetes mellitus (E10-E14)	22	0.8	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	0.7	---
15	Anemias (D50-D64)	13	0.4	---
...	All other causes (Residual)	677	23.2	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic white, female, all ages				
...	All causes	998,312	100.0	981.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	236,208	23.7	232.2
2	Malignant neoplasms (C00-C97)	218,751	21.9	215.0
3	Chronic lower respiratory diseases (J40-J47)	65,497	6.6	64.4
4	Cerebrovascular diseases (I60-I69)	61,809	6.2	60.8
5	Alzheimer's disease (G30)	51,224	5.1	50.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	36,673	3.7	36.0
7	Diabetes mellitus (E10-E14)	22,612	2.3	22.2
8	Influenza and pneumonia (J09-J18)	21,587	2.2	21.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18,460	1.8	18.1
10	Septicemia (A40-A41)	13,988	1.4	13.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	11,402	1.1	11.2
12	Chronic liver disease and cirrhosis (K70,K73-K74)	8,385	0.8	8.2
13	Parkinson's disease (G20-G21)	8,191	0.8	8.1
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,772	0.7	6.7
15	Pneumonitis due to solids and liquids (J69)	6,677	0.7	6.6
...	All other causes (Residual)	210,076	21.0	206.5
Non-Hispanic white, female, 1-4 years				
...	All causes	920	100.0	21.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	288	31.3	6.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	128	13.9	3.0
3	Malignant neoplasms (C00-C97)	87	9.5	2.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	69	7.5	1.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	23	2.5	0.5
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	2.2	0.5
7	Septicemia (A40-A41)	18	2.0	*
7	Influenza and pneumonia (J09-J18)	18	2.0	*
9	Certain conditions originating in the perinatal period (P00-P96)	10	1.1	*
10	Chronic lower respiratory diseases (J40-J47)	7	0.8	*
11	Cerebrovascular diseases (I60-I69)	6	0.7	*
12	Meningitis (G00,G03)	4	0.4	*
13	Meningococcal infection (A39)	3	0.3	*
13	Anemias (D50-D64)	3	0.3	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.3	*
...	All other causes (Residual)	233	25.3	5.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic white, female, 5-9 years				
...	All causes	561	100.0	10.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	165	29.4	3.0
2	Malignant neoplasms (C00-C97)	106	18.9	1.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	51	9.1	0.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	23	4.1	0.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	18	3.2	*
6	Influenza and pneumonia (J09-J18)	17	3.0	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	2.1	*
8	Chronic lower respiratory diseases (J40-J47)	11	2.0	*
9	Cerebrovascular diseases (I60-I69)	8	1.4	*
10	Septicemia (A40-A41)	7	1.2	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.7	*
11	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	4	0.7	*
13	Anemias (D50-D64)	2	0.4	*
13	Meningitis (G00,G03)	2	0.4	*
15	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
15	Pneumonitis due to solids and liquids (J69)	1	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
15	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.2	*
15	Certain conditions originating in the perinatal period (P00-P96)	1	0.2	*
...	All other causes (Residual)	126	22.5	2.3
Non-Hispanic white, female, 10-14 years				
...	All causes	658	100.0	11.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	196	29.8	3.4
2	Malignant neoplasms (C00-C97)	117	17.8	2.0
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	39	5.9	0.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	38	5.8	0.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	31	4.7	0.5
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	2.4	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	16	2.4	*
8	Influenza and pneumonia (J09-J18)	10	1.5	*
9	Chronic lower respiratory diseases (J40-J47)	9	1.4	*
10	Cerebrovascular diseases (I60-I69)	8	1.2	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	1.1	*
12	Septicemia (A40-A41)	6	0.9	*
12	Diabetes mellitus (E10-E14)	6	0.9	*
14	Anemias (D50-D64)	3	0.5	*
14	Pneumonitis due to solids and liquids (J69)	3	0.5	*
14	Diseases of appendix (K35-K38)	3	0.5	*
14	Certain conditions originating in the perinatal period (P00-P96)	3	0.5	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.5	*
...	All other causes (Residual)	144	21.9	2.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic white, female, 15-19 years				
...	All causes	1,861	100.0	30.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	946	50.8	15.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	217	11.7	3.5
3	Malignant neoplasms (C00-C97)	138	7.4	2.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	76	4.1	1.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	54	2.9	0.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	46	2.5	0.7
7	Pregnancy, childbirth and the puerperium (O00-O99)	25	1.3	0.4
8	Cerebrovascular diseases (I60-I69)	17	0.9	*
9	Diabetes mellitus (E10-E14)	14	0.8	*
10	Septicemia (A40-A41)	13	0.7	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	0.6	*
11	Chronic lower respiratory diseases (J40-J47)	12	0.6	*
13	Influenza and pneumonia (J09-J18)	11	0.6	*
14	Meningitis (G00,G03)	5	0.3	*
15	Anemias (D50-D64)	4	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.2	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.2	*
...	All other causes (Residual)	263	14.1	4.3
Non-Hispanic white, female, 20-24 years				
...	All causes	2,907	100.0	46.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,389	47.8	22.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	335	11.5	5.4
3	Malignant neoplasms (C00-C97)	198	6.8	3.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	144	5.0	2.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	84	2.9	1.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	49	1.7	0.8
7	Pregnancy, childbirth and the puerperium (O00-O99)	48	1.7	0.8
8	Influenza and pneumonia (J09-J18)	40	1.4	0.6
9	Diabetes mellitus (E10-E14)	27	0.9	0.4
10	Septicemia (A40-A41)	25	0.9	0.4
11	Cerebrovascular diseases (I60-I69)	22	0.8	0.4
12	Chronic lower respiratory diseases (J40-J47)	14	0.5	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	0.5	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.4	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	9	0.3	*
...	All other causes (Residual)	498	17.1	8.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic white, female, 25-29 years				
...	All causes	3,570	100.0	57.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,375	38.5	22.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	410	11.5	6.6
3	Malignant neoplasms (C00-C97)	376	10.5	6.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	183	5.1	3.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	157	4.4	2.5
6	Pregnancy, childbirth and the puerperium (O00-O99)	80	2.2	1.3
7	Diabetes mellitus (E10-E14)	57	1.6	0.9
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	54	1.5	0.9
9	Cerebrovascular diseases (I60-I69)	47	1.3	0.8
10	Influenza and pneumonia (J09-J18)	42	1.2	0.7
11	Septicemia (A40-A41)	40	1.1	0.6
12	Chronic liver disease and cirrhosis (K70,K73-K74)	29	0.8	0.5
13	Chronic lower respiratory diseases (J40-J47)	25	0.7	0.4
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	25	0.7	0.4
15	Human immunodeficiency virus (HIV) disease (B20-B24)	13	0.4	*
...	All other causes (Residual)	657	18.4	10.6
Non-Hispanic white, female, 30-34 years				
...	All causes	4,433	100.0	76.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,412	31.9	24.4
2	Malignant neoplasms (C00-C97)	672	15.2	11.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	445	10.0	7.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	371	8.4	6.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	126	2.8	2.2
6	Cerebrovascular diseases (I60-I69)	97	2.2	1.7
7	Diabetes mellitus (E10-E14)	82	1.8	1.4
8	Chronic liver disease and cirrhosis (K70,K73-K74)	81	1.8	1.4
9	Pregnancy, childbirth and the puerperium (O00-O99)	69	1.6	1.2
10	Influenza and pneumonia (J09-J18)	60	1.4	1.0
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	54	1.2	0.9
12	Septicemia (A40-A41)	44	1.0	0.8
13	Chronic lower respiratory diseases (J40-J47)	34	0.8	0.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	27	0.6	0.5
15	Human immunodeficiency virus (HIV) disease (B20-B24)	24	0.5	0.4
...	All other causes (Residual)	835	18.8	14.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic white, female, 35-39 years				
...	All causes	6,463	100.0	106.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,640	25.4	27.1
2	Malignant neoplasms (C00-C97)	1,329	20.6	22.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	636	9.8	10.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	567	8.8	9.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	174	2.7	2.9
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	141	2.2	2.3
7	Diabetes mellitus (E10-E14)	136	2.1	2.2
8	Cerebrovascular diseases (I60-I69)	128	2.0	2.1
9	Influenza and pneumonia (J09-J18)	88	1.4	1.5
10	Septicemia (A40-A41)	78	1.2	1.3
11	Chronic lower respiratory diseases (J40-J47)	76	1.2	1.3
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	72	1.1	1.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	62	1.0	1.0
14	Pregnancy, childbirth and the puerperium (O00-O99)	56	0.9	0.9
15	Human immunodeficiency virus (HIV) disease (B20-B24)	45	0.7	0.7
...	All other causes (Residual)	1,235	19.1	20.4
Non-Hispanic white, female, 40-44 years				
...	All causes	10,466	100.0	156.9
1	Malignant neoplasms (C00-C97)	2,846	27.2	42.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,943	18.6	29.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,243	11.9	18.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	692	6.6	10.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	388	3.7	5.8
6	Cerebrovascular diseases (I60-I69)	313	3.0	4.7
7	Diabetes mellitus (E10-E14)	257	2.5	3.9
8	Chronic lower respiratory diseases (J40-J47)	177	1.7	2.7
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	160	1.5	2.4
10	Influenza and pneumonia (J09-J18)	145	1.4	2.2
11	Septicemia (A40-A41)	143	1.4	2.1
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	94	0.9	1.4
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	71	0.7	1.1
14	Human immunodeficiency virus (HIV) disease (B20-B24)	66	0.6	1.0
15	Viral hepatitis (B15-B19)	54	0.5	0.8
...	All other causes (Residual)	1,874	17.9	28.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic white, female, 45-49 years				
...	All causes	19,330	100.0	247.9
1	Malignant neoplasms (C00-C97)	6,269	32.4	80.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,623	13.6	33.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,619	13.5	33.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	899	4.7	11.5
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	878	4.5	11.3
6	Chronic lower respiratory diseases (J40-J47)	602	3.1	7.7
7	Cerebrovascular diseases (I60-I69)	551	2.9	7.1
8	Diabetes mellitus (E10-E14)	434	2.2	5.6
9	Septicemia (A40-A41)	270	1.4	3.5
10	Influenza and pneumonia (J09-J18)	239	1.2	3.1
11	Viral hepatitis (B15-B19)	174	0.9	2.2
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	174	0.9	2.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	156	0.8	2.0
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	106	0.5	1.4
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	96	0.5	1.2
...	All other causes (Residual)	3,240	16.8	41.6
Non-Hispanic white, female, 50-54 years				
...	All causes	29,409	100.0	365.7
1	Malignant neoplasms (C00-C97)	11,235	38.2	139.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,166	14.2	51.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,544	8.7	31.6
4	Chronic lower respiratory diseases (J40-J47)	1,305	4.4	16.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,118	3.8	13.9
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	953	3.2	11.8
7	Cerebrovascular diseases (I60-I69)	867	2.9	10.8
8	Diabetes mellitus (E10-E14)	743	2.5	9.2
9	Septicemia (A40-A41)	399	1.4	5.0
10	Influenza and pneumonia (J09-J18)	355	1.2	4.4
11	Viral hepatitis (B15-B19)	334	1.1	4.2
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	293	1.0	3.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	145	0.5	1.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	143	0.5	1.8
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	136	0.5	1.7
...	All other causes (Residual)	4,673	15.9	58.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic white, female, 55-59 years				
...	All causes	37,745	100.0	508.4
1	Malignant neoplasms (C00-C97)	15,513	41.1	208.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,735	15.2	77.2
3	Chronic lower respiratory diseases (J40-J47)	2,130	5.6	28.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,956	5.2	26.3
5	Diabetes mellitus (E10-E14)	1,143	3.0	15.4
6	Cerebrovascular diseases (I60-I69)	1,108	2.9	14.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,092	2.9	14.7
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	821	2.2	11.1
9	Septicemia (A40-A41)	627	1.7	8.4
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	482	1.3	6.5
11	Influenza and pneumonia (J09-J18)	452	1.2	6.1
12	Viral hepatitis (B15-B19)	379	1.0	5.1
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	210	0.6	2.8
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	169	0.4	2.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	168	0.4	2.3
...	All other causes (Residual)	5,760	15.3	77.6
Non-Hispanic white, female, 60-64 years				
...	All causes	50,968	100.0	769.4
1	Malignant neoplasms (C00-C97)	21,498	42.2	324.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,350	16.4	126.1
3	Chronic lower respiratory diseases (J40-J47)	3,760	7.4	56.8
4	Cerebrovascular diseases (I60-I69)	1,742	3.4	26.3
5	Diabetes mellitus (E10-E14)	1,627	3.2	24.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,515	3.0	22.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,041	2.0	15.7
8	Septicemia (A40-A41)	813	1.6	12.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	750	1.5	11.3
10	Influenza and pneumonia (J09-J18)	625	1.2	9.4
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	558	1.1	8.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	304	0.6	4.6
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	281	0.6	4.2
14	Alzheimer's disease (G30)	261	0.5	3.9
15	Viral hepatitis (B15-B19)	200	0.4	3.0
...	All other causes (Residual)	7,643	15.0	115.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic white, female, 65-69 years				
...	All causes	62,158	100.0	1,221.1
1	Malignant neoplasms (C00-C97)	25,207	40.6	495.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,592	17.0	208.1
3	Chronic lower respiratory diseases (J40-J47)	5,977	9.6	117.4
4	Cerebrovascular diseases (I60-I69)	2,360	3.8	46.4
5	Diabetes mellitus (E10-E14)	1,882	3.0	37.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,330	2.1	26.1
7	Septicemia (A40-A41)	1,055	1.7	20.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,026	1.7	20.2
9	Chronic liver disease and cirrhosis (K70,K73-K74)	879	1.4	17.3
10	Influenza and pneumonia (J09-J18)	800	1.3	15.7
11	Alzheimer's disease (G30)	564	0.9	11.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	435	0.7	8.5
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	369	0.6	7.2
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	322	0.5	6.3
15	Pneumonitis due to solids and liquids (J69)	251	0.4	4.9
...	All other causes (Residual)	9,109	14.7	178.9
Non-Hispanic white, female, 70-74 years				
...	All causes	76,080	100.0	1,945.9
1	Malignant neoplasms (C00-C97)	27,035	35.5	691.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,879	18.2	355.0
3	Chronic lower respiratory diseases (J40-J47)	8,095	10.6	207.0
4	Cerebrovascular diseases (I60-I69)	3,665	4.8	93.7
5	Diabetes mellitus (E10-E14)	2,220	2.9	56.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,483	1.9	37.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,471	1.9	37.6
8	Alzheimer's disease (G30)	1,425	1.9	36.4
9	Septicemia (A40-A41)	1,253	1.6	32.0
10	Influenza and pneumonia (J09-J18)	1,159	1.5	29.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	799	1.1	20.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	603	0.8	15.4
13	Parkinson's disease (G20-G21)	529	0.7	13.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	525	0.7	13.4
15	Pneumonitis due to solids and liquids (J69)	364	0.5	9.3
...	All other causes (Residual)	11,575	15.2	296.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic white, female, 75-79 years				
...	All causes	105,429	100.0	3,207.2
1	Malignant neoplasms (C00-C97)	29,762	28.2	905.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	21,792	20.7	662.9
3	Chronic lower respiratory diseases (J40-J47)	10,582	10.0	321.9
4	Cerebrovascular diseases (I60-I69)	6,324	6.0	192.4
5	Alzheimer's disease (G30)	4,056	3.8	123.4
6	Diabetes mellitus (E10-E14)	2,794	2.7	85.0
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,159	2.0	65.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,125	2.0	64.6
9	Influenza and pneumonia (J09-J18)	1,894	1.8	57.6
10	Septicemia (A40-A41)	1,728	1.6	52.6
11	Parkinson's disease (G20-G21)	1,212	1.1	36.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,052	1.0	32.0
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	744	0.7	22.6
14	Chronic liver disease and cirrhosis (K70,K73-K74)	705	0.7	21.4
15	Pneumonitis due to solids and liquids (J69)	620	0.6	18.9
...	All other causes (Residual)	17,880	17.0	543.9
Non-Hispanic white, female, 80-84 years				
...	All causes	153,831	100.0	5,420.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	36,488	23.7	1,285.7
2	Malignant neoplasms (C00-C97)	32,126	20.9	1,132.0
3	Chronic lower respiratory diseases (J40-J47)	12,383	8.0	436.3
4	Cerebrovascular diseases (I60-I69)	11,036	7.2	388.9
5	Alzheimer's disease (G30)	8,991	5.8	316.8
6	Diabetes mellitus (E10-E14)	3,786	2.5	133.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,506	2.3	123.5
8	Influenza and pneumonia (J09-J18)	3,375	2.2	118.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,297	2.1	116.2
10	Septicemia (A40-A41)	2,351	1.5	82.8
11	Parkinson's disease (G20-G21)	2,026	1.3	71.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,805	1.2	63.6
13	Pneumonitis due to solids and liquids (J69)	1,168	0.8	41.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,091	0.7	38.4
15	Aortic aneurysm and dissection (I71)	817	0.5	28.8
...	All other causes (Residual)	29,585	19.2	1,042.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic white, female, 85-89 years				
...	All causes	191,868	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	53,288	27.8	---
2	Malignant neoplasms (C00-C97)	26,732	13.9	---
3	Cerebrovascular diseases (I60-I69)	15,241	7.9	---
4	Alzheimer's disease (G30)	14,838	7.7	---
5	Chronic lower respiratory diseases (J40-J47)	11,390	5.9	---
6	Influenza and pneumonia (J09-J18)	4,859	2.5	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,463	2.3	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,105	2.1	---
9	Diabetes mellitus (E10-E14)	3,919	2.0	---
10	Septicemia (A40-A41)	2,647	1.4	---
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,561	1.3	---
12	Parkinson's disease (G20-G21)	2,354	1.2	---
13	Pneumonitis due to solids and liquids (J69)	1,607	0.8	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,198	0.6	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	988	0.5	---
...	All other causes (Residual)	41,678	21.7	---
Non-Hispanic white, female, 90-94 years				
...	All causes	150,539	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	47,457	31.5	---
2	Alzheimer's disease (G30)	13,364	8.9	---
3	Malignant neoplasms (C00-C97)	13,075	8.7	---
4	Cerebrovascular diseases (I60-I69)	12,087	8.0	---
5	Chronic lower respiratory diseases (J40-J47)	6,411	4.3	---
6	Influenza and pneumonia (J09-J18)	4,399	2.9	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,548	2.4	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,015	2.0	---
9	Diabetes mellitus (E10-E14)	2,487	1.7	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,415	1.6	---
11	Septicemia (A40-A41)	1,673	1.1	---
12	Pneumonitis due to solids and liquids (J69)	1,390	0.9	---
13	Parkinson's disease (G20-G21)	1,306	0.9	---
14	Atherosclerosis (I70)	943	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	805	0.5	---
...	All other causes (Residual)	36,164	24.0	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic white, female, 95-99 years				
...	All causes	68,850	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	23,522	34.2	---
2	Alzheimer's disease (G30)	6,286	9.1	---
3	Cerebrovascular diseases (I60-I69)	5,193	7.5	---
4	Malignant neoplasms (C00-C97)	3,877	5.6	---
5	Influenza and pneumonia (J09-J18)	2,324	3.4	---
6	Chronic lower respiratory diseases (J40-J47)	2,161	3.1	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,621	2.4	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,338	1.9	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,239	1.8	---
10	Diabetes mellitus (E10-E14)	863	1.3	---
11	Septicemia (A40-A41)	659	1.0	---
12	Pneumonitis due to solids and liquids (J69)	639	0.9	---
13	Atherosclerosis (I70)	536	0.8	---
14	Parkinson's disease (G20-G21)	349	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	283	0.4	---
...	All other causes (Residual)	17,960	26.1	---
Non-Hispanic white, female, 100 years and over				
...	All causes	15,363	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,601	36.5	---
2	Alzheimer's disease (G30)	1,270	8.3	---
3	Cerebrovascular diseases (I60-I69)	962	6.3	---
4	Influenza and pneumonia (J09-J18)	644	4.2	---
5	Malignant neoplasms (C00-C97)	536	3.5	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	372	2.4	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	331	2.2	---
8	Chronic lower respiratory diseases (J40-J47)	328	2.1	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	250	1.6	---
10	Atherosclerosis (I70)	164	1.1	---
11	Pneumonitis due to solids and liquids (J69)	145	0.9	---
12	Diabetes mellitus (E10-E14)	134	0.9	---
13	Septicemia (A40-A41)	112	0.7	---
14	Anemias (D50-D64)	68	0.4	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	49	0.3	---
...	All other causes (Residual)	4,397	28.6	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic black, both sexes, all ages				
...	All causes	283,438	100.0	718.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	68,215	24.1	173.0
2	Malignant neoplasms (C00-C97)	65,276	23.0	165.5
3	Cerebrovascular diseases (I60-I69)	15,833	5.6	40.1
4	Diabetes mellitus (E10-E14)	11,996	4.2	30.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,853	4.2	30.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,769	3.1	22.2
7	Chronic lower respiratory diseases (J40-J47)	8,614	3.0	21.8
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7,679	2.7	19.5
9	Septicemia (A40-A41)	5,958	2.1	15.1
10	Alzheimer's disease (G30)	5,169	1.8	13.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,059	1.8	12.8
12	Influenza and pneumonia (J09-J18)	4,854	1.7	12.3
13	Human immunodeficiency virus (HIV) disease (B20-B24)	4,598	1.6	11.7
14	Certain conditions originating in the perinatal period (P00-P96)	3,846	1.4	9.8
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,601	0.9	6.6
...	All other causes (Residual)	53,118	18.7	134.7
Non-Hispanic black, both sexes, 1-4 years				
...	All causes	993	100.0	40.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	249	25.1	10.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	153	15.4	6.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	124	12.5	5.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	50	5.0	2.0
5	Malignant neoplasms (C00-C97)	47	4.7	1.9
6	Influenza and pneumonia (J09-J18)	25	2.5	1.0
7	Chronic lower respiratory diseases (J40-J47)	19	1.9	*
8	Certain conditions originating in the perinatal period (P00-P96)	15	1.5	*
9	Anemias (D50-D64)	14	1.4	*
10	Cerebrovascular diseases (I60-I69)	13	1.3	*
11	Septicemia (A40-A41)	11	1.1	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	1.0	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.5	*
14	Meningitis (G00,G03)	4	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.4	*
...	All other causes (Residual)	250	25.2	10.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic black, both sexes, 5-9 years				
...	All causes	481	100.0	15.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	150	31.2	5.0
2	Malignant neoplasms (C00-C97)	64	13.3	2.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	35	7.3	1.2
4	Chronic lower respiratory diseases (J40-J47)	29	6.0	1.0
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	24	5.0	0.8
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	21	4.4	0.7
7	Cerebrovascular diseases (I60-I69)	14	2.9	*
8	Septicemia (A40-A41)	8	1.7	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	1.5	*
9	Anemias (D50-D64)	7	1.5	*
11	Influenza and pneumonia (J09-J18)	4	0.8	*
12	Certain conditions originating in the perinatal period (P00-P96)	3	0.6	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.6	*
14	Tuberculosis (A16-A19)	1	0.2	*
14	Diabetes mellitus (E10-E14)	1	0.2	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.2	*
14	Diseases of appendix (K35-K38)	1	0.2	*
14	Hernia (K40-K46)	1	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
...	All other causes (Residual)	106	22.0	3.5
Non-Hispanic black, both sexes, 10-14 years				
...	All causes	637	100.0	20.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	162	25.4	5.1
2	Malignant neoplasms (C00-C97)	78	12.2	2.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	64	10.0	2.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	43	6.8	1.4
5	Chronic lower respiratory diseases (J40-J47)	42	6.6	1.3
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	39	6.1	1.2
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	38	6.0	1.2
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	1.6	*
9	Anemias (D50-D64)	7	1.1	*
9	Cerebrovascular diseases (I60-I69)	7	1.1	*
9	Influenza and pneumonia (J09-J18)	7	1.1	*
12	Septicemia (A40-A41)	6	0.9	*
13	Diabetes mellitus (E10-E14)	5	0.8	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.6	*
15	Certain conditions originating in the perinatal period (P00-P96)	3	0.5	*
...	All other causes (Residual)	122	19.2	3.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic black, both sexes, 15-19 years				
...	All causes	2,482	100.0	70.6
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,089	43.9	31.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	573	23.1	16.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	146	5.9	4.2
4	Malignant neoplasms (C00-C97)	104	4.2	3.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	104	4.2	3.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	48	1.9	1.4
7	Anemias (D50-D64)	28	1.1	0.8
7	Chronic lower respiratory diseases (J40-J47)	28	1.1	0.8
9	Cerebrovascular diseases (I60-I69)	24	1.0	0.7
10	Human immunodeficiency virus (HIV) disease (B20-B24)	17	0.7	*
11	Septicemia (A40-A41)	11	0.4	*
11	Influenza and pneumonia (J09-J18)	11	0.4	*
11	Pregnancy, childbirth and the puerperium (O00-O99)	11	0.4	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.4	*
14	Diabetes mellitus (E10-E14)	9	0.4	*
...	All other causes (Residual)	270	10.9	7.7
Non-Hispanic black, both sexes, 20-24 years				
...	All causes	4,064	100.0	129.5
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,673	41.2	53.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	853	21.0	27.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	308	7.6	9.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	223	5.5	7.1
5	Malignant neoplasms (C00-C97)	160	3.9	5.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	92	2.3	2.9
7	Anemias (D50-D64)	51	1.3	1.6
8	Diabetes mellitus (E10-E14)	42	1.0	1.3
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	41	1.0	1.3
10	Pregnancy, childbirth and the puerperium (O00-O99)	39	1.0	1.2
11	Chronic lower respiratory diseases (J40-J47)	32	0.8	1.0
12	Septicemia (A40-A41)	30	0.7	1.0
13	Influenza and pneumonia (J09-J18)	28	0.7	0.9
14	Cerebrovascular diseases (I60-I69)	25	0.6	0.8
15	Legal intervention (Y35,Y89.0)	20	0.5	0.6
...	All other causes (Residual)	447	11.0	14.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic black, both sexes, 25-29 years				
...	All causes	4,172	100.0	149.2
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,347	32.3	48.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	796	19.1	28.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	344	8.2	12.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	282	6.8	10.1
5	Malignant neoplasms (C00-C97)	211	5.1	7.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	164	3.9	5.9
7	Diabetes mellitus (E10-E14)	76	1.8	2.7
8	Pregnancy, childbirth and the puerperium (O00-O99)	58	1.4	2.1
9	Anemias (D50-D64)	53	1.3	1.9
10	Cerebrovascular diseases (I60-I69)	47	1.1	1.7
11	Chronic lower respiratory diseases (J40-J47)	40	1.0	1.4
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	40	1.0	1.4
13	Influenza and pneumonia (J09-J18)	38	0.9	1.4
14	Septicemia (A40-A41)	34	0.8	1.2
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	31	0.7	1.1
...	All other causes (Residual)	611	14.6	21.8
Non-Hispanic black, both sexes, 30-34 years				
...	All causes	4,614	100.0	175.4
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,019	22.1	38.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	784	17.0	29.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	561	12.2	21.3
4	Malignant neoplasms (C00-C97)	364	7.9	13.8
5	Human immunodeficiency virus (HIV) disease (B20-B24)	307	6.7	11.7
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	212	4.6	8.1
7	Diabetes mellitus (E10-E14)	130	2.8	4.9
8	Cerebrovascular diseases (I60-I69)	90	2.0	3.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	62	1.3	2.4
10	Pregnancy, childbirth and the puerperium (O00-O99)	60	1.3	2.3
11	Chronic lower respiratory diseases (J40-J47)	56	1.2	2.1
12	Anemias (D50-D64)	54	1.2	2.1
13	Septicemia (A40-A41)	51	1.1	1.9
14	Influenza and pneumonia (J09-J18)	44	1.0	1.7
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	35	0.8	1.3
...	All other causes (Residual)	785	17.0	29.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic black, both sexes, 35-39 years				
...	All causes	5,710	100.0	218.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	993	17.4	38.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	747	13.1	28.6
3	Malignant neoplasms (C00-C97)	736	12.9	28.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	653	11.4	25.0
5	Human immunodeficiency virus (HIV) disease (B20-B24)	409	7.2	15.7
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	227	4.0	8.7
7	Cerebrovascular diseases (I60-I69)	196	3.4	7.5
8	Diabetes mellitus (E10-E14)	181	3.2	6.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	110	1.9	4.2
10	Septicemia (A40-A41)	86	1.5	3.3
11	Chronic lower respiratory diseases (J40-J47)	85	1.5	3.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	70	1.2	2.7
13	Anemias (D50-D64)	69	1.2	2.6
14	Chronic liver disease and cirrhosis (K70,K73-K74)	66	1.2	2.5
15	Influenza and pneumonia (J09-J18)	46	0.8	1.8
...	All other causes (Residual)	1,036	18.1	39.7
Non-Hispanic black, both sexes, 40-44 years				
...	All causes	8,097	100.0	304.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,695	20.9	63.7
2	Malignant neoplasms (C00-C97)	1,349	16.7	50.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	821	10.1	30.8
4	Human immunodeficiency virus (HIV) disease (B20-B24)	656	8.1	24.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	412	5.1	15.5
6	Cerebrovascular diseases (I60-I69)	357	4.4	13.4
7	Diabetes mellitus (E10-E14)	326	4.0	12.2
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	208	2.6	7.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	169	2.1	6.3
10	Septicemia (A40-A41)	124	1.5	4.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	112	1.4	4.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	102	1.3	3.8
13	Influenza and pneumonia (J09-J18)	97	1.2	3.6
14	Chronic lower respiratory diseases (J40-J47)	94	1.2	3.5
15	Anemias (D50-D64)	61	0.8	2.3
...	All other causes (Residual)	1,514	18.7	56.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic black, both sexes, 45-49 years				
...	All causes	13,423	100.0	475.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,164	23.6	112.1
2	Malignant neoplasms (C00-C97)	2,955	22.0	104.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,069	8.0	37.9
4	Human immunodeficiency virus (HIV) disease (B20-B24)	830	6.2	29.4
5	Cerebrovascular diseases (I60-I69)	654	4.9	23.2
6	Diabetes mellitus (E10-E14)	558	4.2	19.8
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	384	2.9	13.6
8	Chronic liver disease and cirrhosis (K70,K73-K74)	280	2.1	9.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	278	2.1	9.8
10	Septicemia (A40-A41)	227	1.7	8.0
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	217	1.6	7.7
12	Chronic lower respiratory diseases (J40-J47)	210	1.6	7.4
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	176	1.3	6.2
14	Influenza and pneumonia (J09-J18)	166	1.2	5.9
15	Viral hepatitis (B15-B19)	83	0.6	2.9
...	All other causes (Residual)	2,172	16.2	76.9
Non-Hispanic black, both sexes, 50-54 years				
...	All causes	20,275	100.0	753.0
1	Malignant neoplasms (C00-C97)	5,512	27.2	204.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,862	24.0	180.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,205	5.9	44.8
4	Cerebrovascular diseases (I60-I69)	1,056	5.2	39.2
5	Diabetes mellitus (E10-E14)	827	4.1	30.7
6	Human immunodeficiency virus (HIV) disease (B20-B24)	780	3.8	29.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	524	2.6	19.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	514	2.5	19.1
9	Chronic lower respiratory diseases (J40-J47)	432	2.1	16.0
10	Septicemia (A40-A41)	378	1.9	14.0
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	311	1.5	11.6
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	298	1.5	11.1
13	Viral hepatitis (B15-B19)	227	1.1	8.4
14	Influenza and pneumonia (J09-J18)	222	1.1	8.2
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	156	0.8	5.8
...	All other causes (Residual)	2,971	14.7	110.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic black, both sexes, 55-59 years				
...	All causes	24,964	100.0	1,132.8
1	Malignant neoplasms (C00-C97)	7,578	30.4	343.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,082	24.4	276.0
3	Cerebrovascular diseases (I60-I69)	1,308	5.2	59.4
4	Diabetes mellitus (E10-E14)	1,159	4.6	52.6
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,032	4.1	46.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	693	2.8	31.4
7	Human immunodeficiency virus (HIV) disease (B20-B24)	636	2.5	28.9
8	Chronic lower respiratory diseases (J40-J47)	621	2.5	28.2
9	Chronic liver disease and cirrhosis (K70,K73-K74)	572	2.3	26.0
10	Septicemia (A40-A41)	515	2.1	23.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	397	1.6	18.0
12	Viral hepatitis (B15-B19)	290	1.2	13.2
13	Influenza and pneumonia (J09-J18)	277	1.1	12.6
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	209	0.8	9.5
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	123	0.5	5.6
...	All other causes (Residual)	3,472	13.9	157.6
Non-Hispanic black, both sexes, 60-64 years				
...	All causes	26,324	100.0	1,561.9
1	Malignant neoplasms (C00-C97)	8,643	32.8	512.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,604	25.1	391.8
3	Cerebrovascular diseases (I60-I69)	1,329	5.0	78.9
4	Diabetes mellitus (E10-E14)	1,305	5.0	77.4
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	834	3.2	49.5
6	Chronic lower respiratory diseases (J40-J47)	828	3.1	49.1
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	721	2.7	42.8
8	Septicemia (A40-A41)	584	2.2	34.7
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	475	1.8	28.2
10	Chronic liver disease and cirrhosis (K70,K73-K74)	441	1.7	26.2
11	Human immunodeficiency virus (HIV) disease (B20-B24)	378	1.4	22.4
12	Influenza and pneumonia (J09-J18)	311	1.2	18.5
13	Viral hepatitis (B15-B19)	205	0.8	12.2
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	108	0.4	6.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	107	0.4	6.3
...	All other causes (Residual)	3,451	13.1	204.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic black, both sexes, 65-69 years				
...	All causes	25,491	100.0	2,195.3
1	Malignant neoplasms (C00-C97)	8,355	32.8	719.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,380	25.0	549.5
3	Cerebrovascular diseases (I60-I69)	1,459	5.7	125.7
4	Diabetes mellitus (E10-E14)	1,358	5.3	117.0
5	Chronic lower respiratory diseases (J40-J47)	987	3.9	85.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	881	3.5	75.9
7	Septicemia (A40-A41)	625	2.5	53.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	483	1.9	41.6
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	473	1.9	40.7
10	Influenza and pneumonia (J09-J18)	392	1.5	33.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	236	0.9	20.3
12	Human immunodeficiency virus (HIV) disease (B20-B24)	156	0.6	13.4
13	Alzheimer's disease (G30)	106	0.4	9.1
14	Pneumonitis due to solids and liquids (J69)	103	0.4	8.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	94	0.4	8.1
...	All other causes (Residual)	3,403	13.3	293.1
Non-Hispanic black, both sexes, 70-74 years				
...	All causes	26,439	100.0	3,110.7
1	Malignant neoplasms (C00-C97)	8,112	30.7	954.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,641	25.1	781.3
3	Cerebrovascular diseases (I60-I69)	1,637	6.2	192.6
4	Diabetes mellitus (E10-E14)	1,387	5.2	163.2
5	Chronic lower respiratory diseases (J40-J47)	1,173	4.4	138.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,065	4.0	125.3
7	Septicemia (A40-A41)	633	2.4	74.5
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	467	1.8	54.9
9	Influenza and pneumonia (J09-J18)	466	1.8	54.8
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	417	1.6	49.1
11	Alzheimer's disease (G30)	296	1.1	34.8
12	Chronic liver disease and cirrhosis (K70,K73-K74)	146	0.6	17.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	136	0.5	16.0
14	Pneumonitis due to solids and liquids (J69)	128	0.5	15.1
15	Human immunodeficiency virus (HIV) disease (B20-B24)	98	0.4	11.5
15	Aortic aneurysm and dissection (I71)	98	0.4	11.5
...	All other causes (Residual)	3,539	13.4	416.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic black, both sexes, 75-79 years				
...	All causes	27,956	100.0	4,542.9
1	Malignant neoplasms (C00-C97)	7,490	26.8	1,217.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,368	26.4	1,197.3
3	Cerebrovascular diseases (I60-I69)	1,839	6.6	298.8
4	Diabetes mellitus (E10-E14)	1,493	5.3	242.6
5	Chronic lower respiratory diseases (J40-J47)	1,126	4.0	183.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,073	3.8	174.4
7	Septicemia (A40-A41)	691	2.5	112.3
8	Alzheimer's disease (G30)	602	2.2	97.8
9	Influenza and pneumonia (J09-J18)	519	1.9	84.3
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	504	1.8	81.9
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	422	1.5	68.6
12	Pneumonitis due to solids and liquids (J69)	163	0.6	26.5
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	147	0.5	23.9
14	Parkinson's disease (G20-G21)	142	0.5	23.1
15	Aortic aneurysm and dissection (I71)	100	0.4	16.3
...	All other causes (Residual)	4,277	15.3	695.0
Non-Hispanic black, both sexes, 80-84 years				
...	All causes	28,986	100.0	6,834.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,865	27.1	1,854.4
2	Malignant neoplasms (C00-C97)	6,418	22.1	1,513.2
3	Cerebrovascular diseases (I60-I69)	2,053	7.1	484.1
4	Diabetes mellitus (E10-E14)	1,303	4.5	307.2
5	Chronic lower respiratory diseases (J40-J47)	1,186	4.1	279.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,135	3.9	267.6
7	Alzheimer's disease (G30)	1,048	3.6	247.1
8	Septicemia (A40-A41)	724	2.5	170.7
9	Influenza and pneumonia (J09-J18)	693	2.4	163.4
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	611	2.1	144.1
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	391	1.3	92.2
12	Pneumonitis due to solids and liquids (J69)	237	0.8	55.9
13	Parkinson's disease (G20-G21)	178	0.6	42.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	141	0.5	33.2
15	Aortic aneurysm and dissection (I71)	102	0.4	24.0
...	All other causes (Residual)	4,901	16.9	1,155.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic black, both sexes, 85-89 years				
...	All causes	25,797	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,320	28.4	---
2	Malignant neoplasms (C00-C97)	4,405	17.1	---
3	Cerebrovascular diseases (I60-I69)	1,849	7.2	---
4	Alzheimer's disease (G30)	1,328	5.1	---
5	Diabetes mellitus (E10-E14)	1,039	4.0	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	961	3.7	---
7	Chronic lower respiratory diseases (J40-J47)	954	3.7	---
8	Influenza and pneumonia (J09-J18)	664	2.6	---
9	Septicemia (A40-A41)	637	2.5	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	611	2.4	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	321	1.2	---
12	Pneumonitis due to solids and liquids (J69)	227	0.9	---
13	Parkinson's disease (G20-G21)	189	0.7	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	120	0.5	---
15	Atherosclerosis (I70)	91	0.4	---
...	All other causes (Residual)	5,081	19.7	---
Non-Hispanic black, both sexes, 90-94 years				
...	All causes	15,990	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,788	29.9	---
2	Malignant neoplasms (C00-C97)	1,959	12.3	---
3	Cerebrovascular diseases (I60-I69)	1,175	7.3	---
4	Alzheimer's disease (G30)	1,056	6.6	---
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	595	3.7	---
6	Diabetes mellitus (E10-E14)	533	3.3	---
7	Influenza and pneumonia (J09-J18)	466	2.9	---
8	Chronic lower respiratory diseases (J40-J47)	460	2.9	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	421	2.6	---
10	Septicemia (A40-A41)	329	2.1	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	209	1.3	---
12	Pneumonitis due to solids and liquids (J69)	133	0.8	---
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	92	0.6	---
14	Atherosclerosis (I70)	69	0.4	---
15	Parkinson's disease (G20-G21)	66	0.4	---
...	All other causes (Residual)	3,639	22.8	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic black, both sexes, 95-99 years				
...	All causes	7,289	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,285	31.3	---
2	Malignant neoplasms (C00-C97)	622	8.5	---
3	Cerebrovascular diseases (I60-I69)	526	7.2	---
4	Alzheimer's disease (G30)	519	7.1	---
5	Essential hypertension and hypertensive renal disease (I10,I12,I15)	247	3.4	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	246	3.4	---
7	Influenza and pneumonia (J09-J18)	222	3.0	---
8	Diabetes mellitus (E10-E14)	212	2.9	---
9	Chronic lower respiratory diseases (J40-J47)	163	2.2	---
10	Septicemia (A40-A41)	133	1.8	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	94	1.3	---
12	Pneumonitis due to solids and liquids (J69)	74	1.0	---
13	Atherosclerosis (I70)	48	0.7	---
14	Anemias (D50-D64)	31	0.4	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	30	0.4	---
15	Nutritional deficiencies (E40-E64)	30	0.4	---
...	All other causes (Residual)	1,807	24.8	---
Non-Hispanic black, both sexes, 100 years and over				
...	All causes	2,166	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	729	33.7	---
2	Alzheimer's disease (G30)	154	7.1	---
3	Cerebrovascular diseases (I60-I69)	130	6.0	---
4	Malignant neoplasms (C00-C97)	105	4.8	---
5	Essential hypertension and hypertensive renal disease (I10,I12,I15)	85	3.9	---
5	Influenza and pneumonia (J09-J18)	85	3.9	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	58	2.7	---
8	Diabetes mellitus (E10-E14)	48	2.2	---
9	Chronic lower respiratory diseases (J40-J47)	40	1.8	---
10	Septicemia (A40-A41)	34	1.6	---
11	Atherosclerosis (I70)	21	1.0	---
12	Pneumonitis due to solids and liquids (J69)	19	0.9	---
13	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14	0.6	---
14	Nutritional deficiencies (E40-E64)	10	0.5	---
15	Anemias (D50-D64)	7	0.3	---
...	All other causes (Residual)	627	28.9	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic black, male, all ages				
...	All causes	143,824	100.0	764.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	34,597	24.1	183.9
2	Malignant neoplasms (C00-C97)	33,607	23.4	178.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,920	5.5	42.1
4	Cerebrovascular diseases (I60-I69)	6,890	4.8	36.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6,580	4.6	35.0
6	Diabetes mellitus (E10-E14)	5,572	3.9	29.6
7	Chronic lower respiratory diseases (J40-J47)	4,480	3.1	23.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,984	2.8	21.2
9	Human immunodeficiency virus (HIV) disease (B20-B24)	3,000	2.1	15.9
10	Septicemia (A40-A41)	2,673	1.9	14.2
11	Influenza and pneumonia (J09-J18)	2,340	1.6	12.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,196	1.5	11.7
13	Certain conditions originating in the perinatal period (P00-P96)	2,116	1.5	11.2
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,712	1.2	9.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,694	1.2	9.0
...	All other causes (Residual)	24,463	17.0	130.0
Non-Hispanic black, male, 1-4 years				
...	All causes	570	100.0	45.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	159	27.9	12.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	91	16.0	7.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	57	10.0	4.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	28	4.9	2.2
5	Malignant neoplasms (C00-C97)	26	4.6	2.1
6	Influenza and pneumonia (J09-J18)	15	2.6	*
7	Chronic lower respiratory diseases (J40-J47)	10	1.8	*
8	Anemias (D50-D64)	9	1.6	*
9	Certain conditions originating in the perinatal period (P00-P96)	8	1.4	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.1	*
11	Cerebrovascular diseases (I60-I69)	5	0.9	*
12	Septicemia (A40-A41)	4	0.7	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.5	*
14	Meningococcal infection (A39)	2	0.4	*
14	Meningitis (G00,G03)	2	0.4	*
14	Pneumonitis due to solids and liquids (J69)	2	0.4	*
14	Hernia (K40-K46)	2	0.4	*
...	All other causes (Residual)	141	24.7	11.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic black, male, 5-9 years				
...	All causes	275	100.0	17.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	95	34.5	6.2
2	Malignant neoplasms (C00-C97)	24	8.7	1.6
2	Chronic lower respiratory diseases (J40-J47)	24	8.7	1.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	21	7.6	1.4
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	16	5.8	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	15	5.5	*
7	Cerebrovascular diseases (I60-I69)	9	3.3	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	1.8	*
9	Anemias (D50-D64)	4	1.5	*
10	Septicemia (A40-A41)	3	1.1	*
11	Influenza and pneumonia (J09-J18)	2	0.7	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.4	*
12	Hernia (K40-K46)	1	0.4	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.4	*
12	Certain conditions originating in the perinatal period (P00-P96)	1	0.4	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.4	*
...	All other causes (Residual)	52	18.9	3.4
Non-Hispanic black, male, 10-14 years				
...	All causes	374	100.0	23.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	112	29.9	7.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	46	12.3	2.9
3	Malignant neoplasms (C00-C97)	38	10.2	2.4
4	Chronic lower respiratory diseases (J40-J47)	27	7.2	1.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	23	6.1	1.4
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	23	6.1	1.4
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	18	4.8	*
8	Cerebrovascular diseases (I60-I69)	5	1.3	*
8	Influenza and pneumonia (J09-J18)	5	1.3	*
10	Septicemia (A40-A41)	4	1.1	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	1.1	*
12	Diabetes mellitus (E10-E14)	3	0.8	*
13	Anemias (D50-D64)	2	0.5	*
13	Certain conditions originating in the perinatal period (P00-P96)	2	0.5	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.3	*
15	Meningitis (G00,G03)	1	0.3	*
15	Pneumonitis due to solids and liquids (J69)	1	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
...	All other causes (Residual)	58	15.5	3.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic black, male, 15-19 years				
...	All causes	1,929	100.0	108.0
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	965	50.0	54.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	435	22.6	24.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	126	6.5	7.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	70	3.6	3.9
5	Malignant neoplasms (C00-C97)	56	2.9	3.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	33	1.7	1.8
7	Anemias (D50-D64)	21	1.1	1.2
8	Chronic lower respiratory diseases (J40-J47)	17	0.9	*
9	Cerebrovascular diseases (I60-I69)	13	0.7	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	10	0.5	*
11	Legal intervention (Y35,Y89.0)	8	0.4	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.4	*
13	Septicemia (A40-A41)	6	0.3	*
14	Diabetes mellitus (E10-E14)	5	0.3	*
14	Influenza and pneumonia (J09-J18)	5	0.3	*
...	All other causes (Residual)	152	7.9	8.5
Non-Hispanic black, male, 20-24 years				
...	All causes	3,104	100.0	200.2
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,534	49.4	99.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	618	19.9	39.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	259	8.3	16.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	134	4.3	8.6
5	Malignant neoplasms (C00-C97)	92	3.0	5.9
6	Human immunodeficiency virus (HIV) disease (B20-B24)	60	1.9	3.9
7	Anemias (D50-D64)	34	1.1	2.2
8	Chronic lower respiratory diseases (J40-J47)	24	0.8	1.5
9	Diabetes mellitus (E10-E14)	21	0.7	1.4
10	Septicemia (A40-A41)	20	0.6	1.3
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	20	0.6	1.3
12	Influenza and pneumonia (J09-J18)	19	0.6	*
12	Legal intervention (Y35,Y89.0)	19	0.6	*
14	Cerebrovascular diseases (I60-I69)	8	0.3	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.2	*
15	Pneumonitis due to solids and liquids (J69)	5	0.2	*
...	All other causes (Residual)	232	7.5	15.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic black, male, 25-29 years				
...	All causes	2,940	100.0	219.0
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,185	40.3	88.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	586	19.9	43.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	239	8.1	17.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	214	7.3	15.9
5	Human immunodeficiency virus (HIV) disease (B20-B24)	107	3.6	8.0
6	Malignant neoplasms (C00-C97)	97	3.3	7.2
7	Diabetes mellitus (E10-E14)	35	1.2	2.6
8	Chronic lower respiratory diseases (J40-J47)	29	1.0	2.2
9	Anemias (D50-D64)	27	0.9	2.0
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	27	0.9	2.0
11	Cerebrovascular diseases (I60-I69)	25	0.9	1.9
12	Influenza and pneumonia (J09-J18)	19	0.6	*
13	Septicemia (A40-A41)	15	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	0.5	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	14	0.5	*
15	Legal intervention (Y35,Y89.0)	14	0.5	*
...	All other causes (Residual)	292	9.9	21.8
Non-Hispanic black, male, 30-34 years				
...	All causes	3,031	100.0	243.4
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	914	30.2	73.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	571	18.8	45.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	355	11.7	28.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	186	6.1	14.9
5	Human immunodeficiency virus (HIV) disease (B20-B24)	167	5.5	13.4
6	Malignant neoplasms (C00-C97)	153	5.0	12.3
7	Diabetes mellitus (E10-E14)	77	2.5	6.2
8	Cerebrovascular diseases (I60-I69)	50	1.6	4.0
9	Chronic lower respiratory diseases (J40-J47)	39	1.3	3.1
10	Influenza and pneumonia (J09-J18)	29	1.0	2.3
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	26	0.9	2.1
12	Anemias (D50-D64)	24	0.8	1.9
13	Septicemia (A40-A41)	20	0.7	1.6
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	15	0.5	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	15	0.5	*
...	All other causes (Residual)	390	12.9	31.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic black, male, 35-39 years				
...	All causes	3,368	100.0	274.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	624	18.5	50.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	556	16.5	45.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	520	15.4	42.4
4	Malignant neoplasms (C00-C97)	247	7.3	20.1
5	Human immunodeficiency virus (HIV) disease (B20-B24)	226	6.7	18.4
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	180	5.3	14.7
7	Cerebrovascular diseases (I60-I69)	101	3.0	8.2
8	Diabetes mellitus (E10-E14)	97	2.9	7.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	62	1.8	5.1
10	Septicemia (A40-A41)	49	1.5	4.0
11	Chronic lower respiratory diseases (J40-J47)	42	1.2	3.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	41	1.2	3.3
13	Anemias (D50-D64)	38	1.1	3.1
13	Chronic liver disease and cirrhosis (K70,K73-K74)	38	1.1	3.1
15	Aortic aneurysm and dissection (I71)	37	1.1	3.0
...	All other causes (Residual)	510	15.1	41.6
Non-Hispanic black, male, 40-44 years				
...	All causes	4,607	100.0	366.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,024	22.2	81.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	559	12.1	44.5
3	Malignant neoplasms (C00-C97)	537	11.7	42.7
4	Human immunodeficiency virus (HIV) disease (B20-B24)	421	9.1	33.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	334	7.2	26.6
6	Diabetes mellitus (E10-E14)	203	4.4	16.1
7	Cerebrovascular diseases (I60-I69)	186	4.0	14.8
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	160	3.5	12.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	92	2.0	7.3
10	Chronic liver disease and cirrhosis (K70,K73-K74)	66	1.4	5.2
11	Influenza and pneumonia (J09-J18)	58	1.3	4.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	56	1.2	4.5
13	Septicemia (A40-A41)	55	1.2	4.4
14	Chronic lower respiratory diseases (J40-J47)	47	1.0	3.7
15	Aortic aneurysm and dissection (I71)	44	1.0	3.5
...	All other causes (Residual)	765	16.6	60.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic black, male, 45-49 years				
...	All causes	7,568	100.0	569.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,004	26.5	150.7
2	Malignant neoplasms (C00-C97)	1,273	16.8	95.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	721	9.5	54.2
4	Human immunodeficiency virus (HIV) disease (B20-B24)	545	7.2	41.0
5	Cerebrovascular diseases (I60-I69)	339	4.5	25.5
6	Diabetes mellitus (E10-E14)	321	4.2	24.1
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	300	4.0	22.6
8	Chronic liver disease and cirrhosis (K70,K73-K74)	172	2.3	12.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	142	1.9	10.7
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	140	1.8	10.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	127	1.7	9.6
12	Septicemia (A40-A41)	117	1.5	8.8
13	Influenza and pneumonia (J09-J18)	93	1.2	7.0
14	Chronic lower respiratory diseases (J40-J47)	90	1.2	6.8
15	Viral hepatitis (B15-B19)	55	0.7	4.1
...	All other causes (Residual)	1,129	14.9	84.9
Non-Hispanic black, male, 50-54 years				
...	All causes	11,578	100.0	918.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,102	26.8	246.1
2	Malignant neoplasms (C00-C97)	2,761	23.8	219.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	816	7.0	64.7
4	Cerebrovascular diseases (I60-I69)	565	4.9	44.8
5	Human immunodeficiency virus (HIV) disease (B20-B24)	526	4.5	41.7
6	Diabetes mellitus (E10-E14)	465	4.0	36.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	341	2.9	27.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	253	2.2	20.1
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	244	2.1	19.4
10	Septicemia (A40-A41)	197	1.7	15.6
11	Chronic lower respiratory diseases (J40-J47)	193	1.7	15.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	160	1.4	12.7
13	Viral hepatitis (B15-B19)	153	1.3	12.1
14	Influenza and pneumonia (J09-J18)	133	1.1	10.6
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	119	1.0	9.4
...	All other causes (Residual)	1,550	13.4	123.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic black, male, 55-59 years				
...	All causes	14,768	100.0	1,462.7
1	Malignant neoplasms (C00-C97)	4,180	28.3	414.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,847	26.0	381.0
3	Cerebrovascular diseases (I60-I69)	756	5.1	74.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	720	4.9	71.3
5	Diabetes mellitus (E10-E14)	648	4.4	64.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	439	3.0	43.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	411	2.8	40.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	389	2.6	38.5
9	Chronic lower respiratory diseases (J40-J47)	322	2.2	31.9
10	Septicemia (A40-A41)	271	1.8	26.8
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	240	1.6	23.8
12	Viral hepatitis (B15-B19)	196	1.3	19.4
13	Influenza and pneumonia (J09-J18)	164	1.1	16.2
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	159	1.1	15.7
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	97	0.7	9.6
...	All other causes (Residual)	1,929	13.1	191.1
Non-Hispanic black, male, 60-64 years				
...	All causes	15,370	100.0	2,027.9
1	Malignant neoplasms (C00-C97)	4,839	31.5	638.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,082	26.6	538.6
3	Cerebrovascular diseases (I60-I69)	763	5.0	100.7
4	Diabetes mellitus (E10-E14)	691	4.5	91.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	535	3.5	70.6
6	Chronic lower respiratory diseases (J40-J47)	465	3.0	61.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	453	2.9	59.8
8	Chronic liver disease and cirrhosis (K70,K73-K74)	309	2.0	40.8
9	Septicemia (A40-A41)	305	2.0	40.2
10	Human immunodeficiency virus (HIV) disease (B20-B24)	267	1.7	35.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	254	1.7	33.5
12	Influenza and pneumonia (J09-J18)	172	1.1	22.7
13	Viral hepatitis (B15-B19)	150	1.0	19.8
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	80	0.5	10.6
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	70	0.5	9.2
...	All other causes (Residual)	1,935	12.6	255.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic black, male, 65-69 years				
...	All causes	14,113	100.0	2,802.4
1	Malignant neoplasms (C00-C97)	4,612	32.7	915.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,762	26.7	747.0
3	Cerebrovascular diseases (I60-I69)	806	5.7	160.0
4	Diabetes mellitus (E10-E14)	682	4.8	135.4
5	Chronic lower respiratory diseases (J40-J47)	548	3.9	108.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	443	3.1	88.0
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	317	2.2	62.9
8	Septicemia (A40-A41)	300	2.1	59.6
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	242	1.7	48.1
10	Influenza and pneumonia (J09-J18)	222	1.6	44.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	163	1.2	32.4
12	Human immunodeficiency virus (HIV) disease (B20-B24)	111	0.8	22.0
13	Pneumonitis due to solids and liquids (J69)	68	0.5	13.5
14	Viral hepatitis (B15-B19)	55	0.4	10.9
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	55	0.4	10.9
...	All other causes (Residual)	1,727	12.2	342.9
Non-Hispanic black, male, 70-74 years				
...	All causes	13,947	100.0	3,947.8
1	Malignant neoplasms (C00-C97)	4,392	31.5	1,243.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,644	26.1	1,031.5
3	Cerebrovascular diseases (I60-I69)	801	5.7	226.7
4	Chronic lower respiratory diseases (J40-J47)	656	4.7	185.7
5	Diabetes mellitus (E10-E14)	646	4.6	182.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	522	3.7	147.8
7	Septicemia (A40-A41)	287	2.1	81.2
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	272	2.0	77.0
9	Influenza and pneumonia (J09-J18)	262	1.9	74.2
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	237	1.7	67.1
11	Alzheimer's disease (G30)	128	0.9	36.2
12	Chronic liver disease and cirrhosis (K70,K73-K74)	85	0.6	24.1
13	Pneumonitis due to solids and liquids (J69)	74	0.5	20.9
14	Human immunodeficiency virus (HIV) disease (B20-B24)	71	0.5	20.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	67	0.5	19.0
...	All other causes (Residual)	1,803	12.9	510.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic black, male, 75-79 years				
...	All causes	13,738	100.0	5,789.4
1	Malignant neoplasms (C00-C97)	3,953	28.8	1,665.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,691	26.9	1,555.4
3	Cerebrovascular diseases (I60-I69)	804	5.9	338.8
4	Diabetes mellitus (E10-E14)	669	4.9	281.9
5	Chronic lower respiratory diseases (J40-J47)	606	4.4	255.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	484	3.5	204.0
7	Septicemia (A40-A41)	327	2.4	137.8
8	Influenza and pneumonia (J09-J18)	262	1.9	110.4
9	Alzheimer's disease (G30)	246	1.8	103.7
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	236	1.7	99.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	219	1.6	92.3
12	Pneumonitis due to solids and liquids (J69)	94	0.7	39.6
13	Parkinson's disease (G20-G21)	91	0.7	38.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	68	0.5	28.7
15	Chronic liver disease and cirrhosis (K70,K73-K74)	58	0.4	24.4
...	All other causes (Residual)	1,930	14.0	813.3
Non-Hispanic black, male, 80-84 years				
...	All causes	12,543	100.0	8,566.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,342	26.6	2,282.5
2	Malignant neoplasms (C00-C97)	3,206	25.6	2,189.6
3	Cerebrovascular diseases (I60-I69)	757	6.0	517.0
4	Chronic lower respiratory diseases (J40-J47)	609	4.9	415.9
5	Diabetes mellitus (E10-E14)	492	3.9	336.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	469	3.7	320.3
7	Alzheimer's disease (G30)	360	2.9	245.9
8	Influenza and pneumonia (J09-J18)	320	2.6	218.5
9	Septicemia (A40-A41)	291	2.3	198.7
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	227	1.8	155.0
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	202	1.6	138.0
12	Pneumonitis due to solids and liquids (J69)	136	1.1	92.9
13	Parkinson's disease (G20-G21)	98	0.8	66.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	54	0.4	36.9
15	Atherosclerosis (I70)	39	0.3	26.6
...	All other causes (Residual)	1,941	15.5	1,325.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic black, male, 85-89 years				
...	All causes	9,570	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,668	27.9	---
2	Malignant neoplasms (C00-C97)	2,069	21.6	---
3	Cerebrovascular diseases (I60-I69)	514	5.4	---
4	Chronic lower respiratory diseases (J40-J47)	478	5.0	---
5	Alzheimer's disease (G30)	372	3.9	---
6	Diabetes mellitus (E10-E14)	340	3.6	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	337	3.5	---
8	Influenza and pneumonia (J09-J18)	251	2.6	---
9	Septicemia (A40-A41)	228	2.4	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	181	1.9	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	146	1.5	---
12	Parkinson's disease (G20-G21)	92	1.0	---
13	Pneumonitis due to solids and liquids (J69)	89	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	49	0.5	---
15	Atherosclerosis (I70)	28	0.3	---
...	All other causes (Residual)	1,728	18.1	---
Non-Hispanic black, male, 90-94 years				
...	All causes	4,645	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,344	28.9	---
2	Malignant neoplasms (C00-C97)	813	17.5	---
3	Cerebrovascular diseases (I60-I69)	262	5.6	---
4	Alzheimer's disease (G30)	203	4.4	---
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	198	4.3	---
6	Chronic lower respiratory diseases (J40-J47)	186	4.0	---
7	Influenza and pneumonia (J09-J18)	161	3.5	---
8	Diabetes mellitus (E10-E14)	128	2.8	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	119	2.6	---
10	Septicemia (A40-A41)	95	2.0	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	77	1.7	---
12	Pneumonitis due to solids and liquids (J69)	54	1.2	---
13	Parkinson's disease (G20-G21)	33	0.7	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	25	0.5	---
15	Atherosclerosis (I70)	16	0.3	---
...	All other causes (Residual)	931	20.0	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic black, male, 95-99 years				
...	All causes	1,510	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	448	29.7	---
2	Malignant neoplasms (C00-C97)	207	13.7	---
3	Cerebrovascular diseases (I60-I69)	88	5.8	---
4	Alzheimer's disease (G30)	80	5.3	---
5	Influenza and pneumonia (J09-J18)	66	4.4	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	63	4.2	---
7	Chronic lower respiratory diseases (J40-J47)	47	3.1	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	43	2.8	---
9	Diabetes mellitus (E10-E14)	38	2.5	---
10	Septicemia (A40-A41)	28	1.9	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	25	1.7	---
12	Pneumonitis due to solids and liquids (J69)	21	1.4	---
13	Atherosclerosis (I70)	11	0.7	---
14	Nutritional deficiencies (E40-E64)	10	0.7	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.5	---
...	All other causes (Residual)	328	21.7	---
Non-Hispanic black, male, 100 years and over				
...	All causes	330	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	112	33.9	---
2	Malignant neoplasms (C00-C97)	28	8.5	---
3	Alzheimer's disease (G30)	15	4.5	---
3	Cerebrovascular diseases (I60-I69)	15	4.5	---
5	Influenza and pneumonia (J09-J18)	14	4.2	---
5	Chronic lower respiratory diseases (J40-J47)	14	4.2	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	12	3.6	---
8	Diabetes mellitus (E10-E14)	8	2.4	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	2.4	---
10	Septicemia (A40-A41)	6	1.8	---
11	Atherosclerosis (I70)	5	1.5	---
12	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3	0.9	---
13	Parkinson's disease (G20-G21)	2	0.6	---
13	Pneumonitis due to solids and liquids (J69)	2	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.3	---
15	Nutritional deficiencies (E40-E64)	1	0.3	---
15	Pneumoconioses and chemical effects (J60-J66,J68)	1	0.3	---
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.3	---
15	Hyperplasia of prostate (N40)	1	0.3	---
...	All other causes (Residual)	81	24.5	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic black, female, all ages				
...	All causes	139,614	100.0	676.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	33,618	24.1	163.0
2	Malignant neoplasms (C00-C97)	31,669	22.7	153.5
3	Cerebrovascular diseases (I60-I69)	8,943	6.4	43.4
4	Diabetes mellitus (E10-E14)	6,424	4.6	31.1
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,785	3.4	23.2
6	Chronic lower respiratory diseases (J40-J47)	4,134	3.0	20.0
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,933	2.8	19.1
8	Alzheimer's disease (G30)	3,698	2.6	17.9
9	Septicemia (A40-A41)	3,285	2.4	15.9
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,863	2.1	13.9
11	Influenza and pneumonia (J09-J18)	2,514	1.8	12.2
12	Certain conditions originating in the perinatal period (P00-P96)	1,730	1.2	8.4
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,598	1.1	7.7
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,099	0.8	5.3
15	Chronic liver disease and cirrhosis (K70,K73-K74)	907	0.6	4.4
...	All other causes (Residual)	28,414	20.4	137.8
Non-Hispanic black, female, 1-4 years				
...	All causes	423	100.0	34.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	90	21.3	7.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	67	15.8	5.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	62	14.7	5.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	22	5.2	1.8
5	Malignant neoplasms (C00-C97)	21	5.0	1.7
6	Influenza and pneumonia (J09-J18)	10	2.4	*
7	Chronic lower respiratory diseases (J40-J47)	9	2.1	*
8	Cerebrovascular diseases (I60-I69)	8	1.9	*
9	Septicemia (A40-A41)	7	1.7	*
9	Certain conditions originating in the perinatal period (P00-P96)	7	1.7	*
11	Anemias (D50-D64)	5	1.2	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.9	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.7	*
14	Meningitis (G00,G03)	2	0.5	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.5	*
...	All other causes (Residual)	104	24.6	8.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic black, female, 5-9 years				
...	All causes	206	100.0	13.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	55	26.7	3.7
2	Malignant neoplasms (C00-C97)	40	19.4	2.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	14	6.8	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	3.9	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	6	2.9	*
6	Septicemia (A40-A41)	5	2.4	*
6	Cerebrovascular diseases (I60-I69)	5	2.4	*
6	Chronic lower respiratory diseases (J40-J47)	5	2.4	*
9	Anemias (D50-D64)	3	1.5	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	1.0	*
10	Influenza and pneumonia (J09-J18)	2	1.0	*
10	Certain conditions originating in the perinatal period (P00-P96)	2	1.0	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	2	1.0	*
14	Tuberculosis (A16-A19)	1	0.5	*
14	Diabetes mellitus (E10-E14)	1	0.5	*
14	Diseases of appendix (K35-K38)	1	0.5	*
...	All other causes (Residual)	54	26.2	3.6
Non-Hispanic black, female, 10-14 years				
...	All causes	263	100.0	17.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	50	19.0	3.2
2	Malignant neoplasms (C00-C97)	40	15.2	2.6
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	20	7.6	1.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	20	7.6	1.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	18	6.8	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	16	6.1	*
7	Chronic lower respiratory diseases (J40-J47)	15	5.7	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	2.3	*
9	Anemias (D50-D64)	5	1.9	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	1.1	*
11	Septicemia (A40-A41)	2	0.8	*
11	Diabetes mellitus (E10-E14)	2	0.8	*
11	Cerebrovascular diseases (I60-I69)	2	0.8	*
11	Influenza and pneumonia (J09-J18)	2	0.8	*
15	Viral hepatitis (B15-B19)	1	0.4	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.4	*
15	Diseases of appendix (K35-K38)	1	0.4	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.4	*
15	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.4	*
15	Certain conditions originating in the perinatal period (P00-P96)	1	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.4	*
...	All other causes (Residual)	55	20.9	3.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic black, female, 15-19 years				
...	All causes	553	100.0	31.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	138	25.0	8.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	124	22.4	7.2
3	Malignant neoplasms (C00-C97)	48	8.7	2.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	34	6.1	2.0
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	20	3.6	1.2
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	15	2.7	*
7	Cerebrovascular diseases (I60-I69)	11	2.0	*
7	Chronic lower respiratory diseases (J40-J47)	11	2.0	*
7	Pregnancy, childbirth and the puerperium (O00-O99)	11	2.0	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	7	1.3	*
10	Anemias (D50-D64)	7	1.3	*
12	Influenza and pneumonia (J09-J18)	6	1.1	*
13	Septicemia (A40-A41)	5	0.9	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.9	*
15	Diabetes mellitus (E10-E14)	4	0.7	*
...	All other causes (Residual)	107	19.3	6.2
Non-Hispanic black, female, 20-24 years				
...	All causes	960	100.0	60.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	235	24.5	14.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	139	14.5	8.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	89	9.3	5.6
4	Malignant neoplasms (C00-C97)	68	7.1	4.3
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	49	5.1	3.1
6	Pregnancy, childbirth and the puerperium (O00-O99)	39	4.1	2.5
7	Human immunodeficiency virus (HIV) disease (B20-B24)	32	3.3	2.0
8	Diabetes mellitus (E10-E14)	21	2.2	1.3
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	21	2.2	1.3
10	Anemias (D50-D64)	17	1.8	*
10	Cerebrovascular diseases (I60-I69)	17	1.8	*
12	Septicemia (A40-A41)	10	1.0	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	1.0	*
14	Influenza and pneumonia (J09-J18)	9	0.9	*
15	Chronic lower respiratory diseases (J40-J47)	8	0.8	*
...	All other causes (Residual)	196	20.4	12.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic black, female, 25-29 years				
...	All causes	1,232	100.0	84.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	210	17.0	14.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	162	13.1	11.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	130	10.6	8.9
4	Malignant neoplasms (C00-C97)	114	9.3	7.8
5	Pregnancy, childbirth and the puerperium (O00-O99)	58	4.7	4.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	57	4.6	3.9
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	43	3.5	3.0
8	Diabetes mellitus (E10-E14)	41	3.3	2.8
9	Anemias (D50-D64)	26	2.1	1.8
10	Cerebrovascular diseases (I60-I69)	22	1.8	1.5
11	Septicemia (A40-A41)	19	1.5	*
11	Influenza and pneumonia (J09-J18)	19	1.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	1.3	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	13	1.1	*
15	Chronic lower respiratory diseases (J40-J47)	11	0.9	*
...	All other causes (Residual)	291	23.6	20.0
Non-Hispanic black, female, 30-34 years				
...	All causes	1,583	100.0	114.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	213	13.5	15.4
2	Malignant neoplasms (C00-C97)	211	13.3	15.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	206	13.0	14.9
4	Human immunodeficiency virus (HIV) disease (B20-B24)	140	8.8	10.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	105	6.6	7.6
6	Pregnancy, childbirth and the puerperium (O00-O99)	60	3.8	4.3
7	Diabetes mellitus (E10-E14)	53	3.3	3.8
8	Cerebrovascular diseases (I60-I69)	40	2.5	2.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	36	2.3	2.6
10	Septicemia (A40-A41)	31	2.0	2.2
11	Anemias (D50-D64)	30	1.9	2.2
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	26	1.6	1.9
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	20	1.3	1.4
14	Chronic lower respiratory diseases (J40-J47)	17	1.1	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	16	1.0	*
...	All other causes (Residual)	379	23.9	27.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic black, female, 35-39 years				
...	All causes	2,342	100.0	169.2
1	Malignant neoplasms (C00-C97)	489	20.9	35.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	369	15.8	26.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	227	9.7	16.4
4	Human immunodeficiency virus (HIV) disease (B20-B24)	183	7.8	13.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	97	4.1	7.0
6	Cerebrovascular diseases (I60-I69)	95	4.1	6.9
7	Diabetes mellitus (E10-E14)	84	3.6	6.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	48	2.0	3.5
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	47	2.0	3.4
10	Pregnancy, childbirth and the puerperium (O00-O99)	45	1.9	3.3
11	Chronic lower respiratory diseases (J40-J47)	43	1.8	3.1
12	Septicemia (A40-A41)	37	1.6	2.7
13	Anemias (D50-D64)	31	1.3	2.2
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	29	1.2	2.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	28	1.2	2.0
...	All other causes (Residual)	490	20.9	35.4
Non-Hispanic black, female, 40-44 years				
...	All causes	3,490	100.0	248.5
1	Malignant neoplasms (C00-C97)	812	23.3	57.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	671	19.2	47.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	262	7.5	18.7
4	Human immunodeficiency virus (HIV) disease (B20-B24)	235	6.7	16.7
5	Cerebrovascular diseases (I60-I69)	171	4.9	12.2
6	Diabetes mellitus (E10-E14)	123	3.5	8.8
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	78	2.2	5.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	77	2.2	5.5
9	Septicemia (A40-A41)	69	2.0	4.9
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	48	1.4	3.4
11	Chronic lower respiratory diseases (J40-J47)	47	1.3	3.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	46	1.3	3.3
12	Chronic liver disease and cirrhosis (K70,K73-K74)	46	1.3	3.3
14	Influenza and pneumonia (J09-J18)	39	1.1	2.8
15	Anemias (D50-D64)	33	0.9	2.3
15	Pregnancy, childbirth and the puerperium (O00-O99)	33	0.9	2.3
...	All other causes (Residual)	700	20.1	49.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic black, female, 45-49 years				
...	All causes	5,855	100.0	392.0
1	Malignant neoplasms (C00-C97)	1,682	28.7	112.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,160	19.8	77.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	348	5.9	23.3
4	Cerebrovascular diseases (I60-I69)	315	5.4	21.1
5	Human immunodeficiency virus (HIV) disease (B20-B24)	285	4.9	19.1
6	Diabetes mellitus (E10-E14)	237	4.0	15.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	136	2.3	9.1
8	Chronic lower respiratory diseases (J40-J47)	120	2.0	8.0
9	Septicemia (A40-A41)	110	1.9	7.4
10	Chronic liver disease and cirrhosis (K70,K73-K74)	108	1.8	7.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	90	1.5	6.0
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	84	1.4	5.6
13	Influenza and pneumonia (J09-J18)	73	1.2	4.9
14	Anemias (D50-D64)	39	0.7	2.6
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	36	0.6	2.4
...	All other causes (Residual)	1,032	17.6	69.1
Non-Hispanic black, female, 50-54 years				
...	All causes	8,697	100.0	607.4
1	Malignant neoplasms (C00-C97)	2,751	31.6	192.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,760	20.2	122.9
3	Cerebrovascular diseases (I60-I69)	491	5.6	34.3
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	389	4.5	27.2
5	Diabetes mellitus (E10-E14)	362	4.2	25.3
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	261	3.0	18.2
7	Human immunodeficiency virus (HIV) disease (B20-B24)	254	2.9	17.7
8	Chronic lower respiratory diseases (J40-J47)	239	2.7	16.7
9	Chronic liver disease and cirrhosis (K70,K73-K74)	183	2.1	12.8
10	Septicemia (A40-A41)	181	2.1	12.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	151	1.7	10.5
12	Influenza and pneumonia (J09-J18)	89	1.0	6.2
13	Viral hepatitis (B15-B19)	74	0.9	5.2
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	54	0.6	3.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	41	0.5	2.9
...	All other causes (Residual)	1,417	16.3	99.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic black, female, 55-59 years				
...	All causes	10,196	100.0	853.9
1	Malignant neoplasms (C00-C97)	3,398	33.3	284.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,235	21.9	187.2
3	Cerebrovascular diseases (I60-I69)	552	5.4	46.2
4	Diabetes mellitus (E10-E14)	511	5.0	42.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	312	3.1	26.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	304	3.0	25.5
7	Chronic lower respiratory diseases (J40-J47)	299	2.9	25.0
8	Septicemia (A40-A41)	244	2.4	20.4
9	Human immunodeficiency virus (HIV) disease (B20-B24)	197	1.9	16.5
10	Chronic liver disease and cirrhosis (K70,K73-K74)	161	1.6	13.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	157	1.5	13.1
12	Influenza and pneumonia (J09-J18)	113	1.1	9.5
13	Viral hepatitis (B15-B19)	94	0.9	7.9
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	50	0.5	4.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	39	0.4	3.3
...	All other causes (Residual)	1,530	15.0	128.1
Non-Hispanic black, female, 60-64 years				
...	All causes	10,954	100.0	1,181.1
1	Malignant neoplasms (C00-C97)	3,804	34.7	410.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,522	23.0	271.9
3	Diabetes mellitus (E10-E14)	614	5.6	66.2
4	Cerebrovascular diseases (I60-I69)	566	5.2	61.0
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	381	3.5	41.1
6	Chronic lower respiratory diseases (J40-J47)	363	3.3	39.1
7	Septicemia (A40-A41)	279	2.5	30.1
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	221	2.0	23.8
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	186	1.7	20.1
10	Influenza and pneumonia (J09-J18)	139	1.3	15.0
11	Chronic liver disease and cirrhosis (K70,K73-K74)	132	1.2	14.2
12	Human immunodeficiency virus (HIV) disease (B20-B24)	111	1.0	12.0
13	Viral hepatitis (B15-B19)	55	0.5	5.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	52	0.5	5.6
15	Anemias (D50-D64)	37	0.3	4.0
...	All other causes (Residual)	1,492	13.6	160.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic black, female, 65-69 years				
...	All causes	11,378	100.0	1,730.3
1	Malignant neoplasms (C00-C97)	3,743	32.9	569.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,618	23.0	398.1
3	Diabetes mellitus (E10-E14)	676	5.9	102.8
4	Cerebrovascular diseases (I60-I69)	653	5.7	99.3
5	Chronic lower respiratory diseases (J40-J47)	439	3.9	66.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	438	3.8	66.6
7	Septicemia (A40-A41)	325	2.9	49.4
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	231	2.0	35.1
9	Influenza and pneumonia (J09-J18)	170	1.5	25.9
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	166	1.5	25.2
11	Chronic liver disease and cirrhosis (K70,K73-K74)	73	0.6	11.1
12	Alzheimer's disease (G30)	66	0.6	10.0
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	51	0.4	7.8
14	Human immunodeficiency virus (HIV) disease (B20-B24)	45	0.4	6.8
15	Viral hepatitis (B15-B19)	36	0.3	5.5
...	All other causes (Residual)	1,648	14.5	250.6
Non-Hispanic black, female, 70-74 years				
...	All causes	12,492	100.0	2,515.2
1	Malignant neoplasms (C00-C97)	3,720	29.8	749.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,997	24.0	603.4
3	Cerebrovascular diseases (I60-I69)	836	6.7	168.3
4	Diabetes mellitus (E10-E14)	741	5.9	149.2
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	543	4.3	109.3
6	Chronic lower respiratory diseases (J40-J47)	517	4.1	104.1
7	Septicemia (A40-A41)	346	2.8	69.7
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	230	1.8	46.3
9	Influenza and pneumonia (J09-J18)	204	1.6	41.1
10	Alzheimer's disease (G30)	168	1.3	33.8
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	145	1.2	29.2
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	69	0.6	13.9
13	Chronic liver disease and cirrhosis (K70,K73-K74)	61	0.5	12.3
14	Pneumonitis due to solids and liquids (J69)	54	0.4	10.9
15	Aortic aneurysm and dissection (I71)	41	0.3	8.3
...	All other causes (Residual)	1,820	14.6	366.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic black, female, 75-79 years				
...	All causes	14,218	100.0	3,760.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,677	25.9	972.6
2	Malignant neoplasms (C00-C97)	3,537	24.9	935.5
3	Cerebrovascular diseases (I60-I69)	1,035	7.3	273.8
4	Diabetes mellitus (E10-E14)	824	5.8	217.9
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	589	4.1	155.8
6	Chronic lower respiratory diseases (J40-J47)	520	3.7	137.5
7	Septicemia (A40-A41)	364	2.6	96.3
8	Alzheimer's disease (G30)	356	2.5	94.2
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	285	2.0	75.4
10	Influenza and pneumonia (J09-J18)	257	1.8	68.0
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	186	1.3	49.2
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	79	0.6	20.9
13	Pneumonitis due to solids and liquids (J69)	69	0.5	18.3
14	Aortic aneurysm and dissection (I71)	53	0.4	14.0
15	Parkinson's disease (G20-G21)	51	0.4	13.5
...	All other causes (Residual)	2,336	16.4	617.9
Non-Hispanic black, female, 80-84 years				
...	All causes	16,443	100.0	5,921.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,523	27.5	1,628.7
2	Malignant neoplasms (C00-C97)	3,212	19.5	1,156.6
3	Cerebrovascular diseases (I60-I69)	1,296	7.9	466.7
4	Diabetes mellitus (E10-E14)	811	4.9	292.0
5	Alzheimer's disease (G30)	688	4.2	247.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	666	4.1	239.8
7	Chronic lower respiratory diseases (J40-J47)	577	3.5	207.8
8	Septicemia (A40-A41)	433	2.6	155.9
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	384	2.3	138.3
10	Influenza and pneumonia (J09-J18)	373	2.3	134.3
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	189	1.1	68.1
12	Pneumonitis due to solids and liquids (J69)	101	0.6	36.4
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	87	0.5	31.3
14	Parkinson's disease (G20-G21)	80	0.5	28.8
15	Aortic aneurysm and dissection (I71)	67	0.4	24.1
...	All other causes (Residual)	2,956	18.0	1,064.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic black, female, 85-89 years				
...	All causes	16,227	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,652	28.7	---
2	Malignant neoplasms (C00-C97)	2,336	14.4	---
3	Cerebrovascular diseases (I60-I69)	1,335	8.2	---
4	Alzheimer's disease (G30)	956	5.9	---
5	Diabetes mellitus (E10-E14)	699	4.3	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	624	3.8	---
7	Chronic lower respiratory diseases (J40-J47)	476	2.9	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	430	2.6	---
9	Influenza and pneumonia (J09-J18)	413	2.5	---
10	Septicemia (A40-A41)	409	2.5	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	175	1.1	---
12	Pneumonitis due to solids and liquids (J69)	138	0.9	---
13	Parkinson's disease (G20-G21)	97	0.6	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	71	0.4	---
15	Atherosclerosis (I70)	63	0.4	---
...	All other causes (Residual)	3,353	20.7	---
Non-Hispanic black, female, 90-94 years				
...	All causes	11,345	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,444	30.4	---
2	Malignant neoplasms (C00-C97)	1,146	10.1	---
3	Cerebrovascular diseases (I60-I69)	913	8.0	---
4	Alzheimer's disease (G30)	853	7.5	---
5	Diabetes mellitus (E10-E14)	405	3.6	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	397	3.5	---
7	Influenza and pneumonia (J09-J18)	305	2.7	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	302	2.7	---
9	Chronic lower respiratory diseases (J40-J47)	274	2.4	---
10	Septicemia (A40-A41)	234	2.1	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	132	1.2	---
12	Pneumonitis due to solids and liquids (J69)	79	0.7	---
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	67	0.6	---
14	Atherosclerosis (I70)	53	0.5	---
15	Anemias (D50-D64)	45	0.4	---
...	All other causes (Residual)	2,696	23.8	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic black, female, 95-99 years				
...	All causes	5,779	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,837	31.8	---
2	Alzheimer's disease (G30)	439	7.6	---
3	Cerebrovascular diseases (I60-I69)	438	7.6	---
4	Malignant neoplasms (C00-C97)	415	7.2	---
5	Essential hypertension and hypertensive renal disease (I10,I12,I15)	204	3.5	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	183	3.2	---
7	Diabetes mellitus (E10-E14)	174	3.0	---
8	Influenza and pneumonia (J09-J18)	156	2.7	---
9	Chronic lower respiratory diseases (J40-J47)	116	2.0	---
10	Septicemia (A40-A41)	105	1.8	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	69	1.2	---
12	Pneumonitis due to solids and liquids (J69)	53	0.9	---
13	Atherosclerosis (I70)	37	0.6	---
14	Anemias (D50-D64)	25	0.4	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	0.4	---
...	All other causes (Residual)	1,505	26.0	---
Non-Hispanic black, female, 100 years and over				
...	All causes	1,836	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	617	33.6	---
2	Alzheimer's disease (G30)	139	7.6	---
3	Cerebrovascular diseases (I60-I69)	115	6.3	---
4	Malignant neoplasms (C00-C97)	77	4.2	---
5	Essential hypertension and hypertensive renal disease (I10,I12,I15)	73	4.0	---
6	Influenza and pneumonia (J09-J18)	71	3.9	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	50	2.7	---
8	Diabetes mellitus (E10-E14)	40	2.2	---
9	Septicemia (A40-A41)	28	1.5	---
10	Chronic lower respiratory diseases (J40-J47)	26	1.4	---
11	Pneumonitis due to solids and liquids (J69)	17	0.9	---
12	Atherosclerosis (I70)	16	0.9	---
13	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11	0.6	---
14	Nutritional deficiencies (E40-E64)	9	0.5	---
15	Anemias (D50-D64)	7	0.4	---
...	All other causes (Residual)	540	29.4	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available