

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Hispanic, both sexes, all ages				
...	All causes	122,416	100.0	296.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	27,788	22.7	67.2
2	Malignant neoplasms (C00-C97)	24,522	20.0	59.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,408	8.5	25.2
4	Cerebrovascular diseases (I60-I69)	6,781	5.5	16.4
5	Diabetes mellitus (E10-E14)	5,999	4.9	14.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	3,383	2.8	8.2
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,271	2.7	7.9
8	Chronic lower respiratory diseases (J40-J47)	3,107	2.5	7.5
9	Influenza and pneumonia (J10-J18)	2,912	2.4	7.0
10	Certain conditions originating in the perinatal period (P00-P96)	2,681	2.2	6.5
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,207	1.8	5.3
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,196	1.8	5.3
13	Alzheimer's disease (G30)	1,897	1.5	4.6
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,765	1.4	4.3
15	Human immunodeficiency virus (HIV) disease (B20-B24)	1,758	1.4	4.3
...	All other causes (Residual)	21,741	17.8	52.6
Hispanic, both sexes, 1-4 years				
...	All causes	947	100.0	27.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	313	33.1	9.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	138	14.6	4.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	80	8.4	2.3
4	Malignant neoplasms (C00-C97)	70	7.4	2.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	39	4.1	1.1
6	Septicemia (A40-A41)	24	2.5	0.7
6	Influenza and pneumonia (J10-J18)	24	2.5	0.7
8	Chronic lower respiratory diseases (J40-J47)	11	1.2	*
9	Certain conditions originating in the perinatal period (P00-P96)	10	1.1	*
10	Meningococcal infection (A39)	5	0.5	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.5	*
10	Anemias (D50-D64)	5	0.5	*
10	Acute bronchitis and bronchiolitis (J20-J21)	5	0.5	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.5	*
15	Meningitis (G00,G03)	3	0.3	*
15	Cerebrovascular diseases (I60-I69)	3	0.3	*
15	Pneumonitis due to solids and liquids (J69)	3	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.3	*
...	All other causes (Residual)	201	21.2	5.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Hispanic, both sexes, 5-9 years				
...	All causes	487	100.0	12.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	178	36.6	4.6
2	Malignant neoplasms (C00-C97)	99	20.3	2.6
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	36	7.4	0.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	27	5.5	0.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	16	3.3	*
6	Septicemia (A40-A41)	8	1.6	*
7	Chronic lower respiratory diseases (J40-J47)	6	1.2	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	1.0	*
8	Influenza and pneumonia (J10-J18)	5	1.0	*
8	Certain conditions originating in the perinatal period (P00-P96)	5	1.0	*
11	Cerebrovascular diseases (I60-I69)	4	0.8	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.8	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.6	*
14	Anemias (D50-D64)	2	0.4	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.2	*
15	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
15	Pneumonitis due to solids and liquids (J69)	1	0.2	*
15	Diseases of appendix (K35-K38)	1	0.2	*
...	All other causes (Residual)	85	17.5	2.2
Hispanic, both sexes, 10-14 years				
...	All causes	618	100.0	16.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	213	34.5	5.6
2	Malignant neoplasms (C00-C97)	112	18.1	3.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	40	6.5	1.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	39	6.3	1.0
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	34	5.5	0.9
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	18	2.9	*
7	Influenza and pneumonia (J10-J18)	8	1.3	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	1.1	*
9	Cerebrovascular diseases (I60-I69)	6	1.0	*
9	Chronic lower respiratory diseases (J40-J47)	6	1.0	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	3	0.5	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.5	*
11	Certain conditions originating in the perinatal period (P00-P96)	3	0.5	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.5	*
15	Meningitis (G00,G03)	2	0.3	*
...	All other causes (Residual)	121	19.6	3.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Hispanic, both sexes, 15-19 years				
...	All causes	2,141	100.0	64.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	956	44.7	28.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	492	23.0	14.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	214	10.0	6.4
4	Malignant neoplasms (C00-C97)	137	6.4	4.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	43	2.0	1.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	29	1.4	0.9
7	Cerebrovascular diseases (I60-I69)	11	0.5	*
7	Influenza and pneumonia (J10-J18)	11	0.5	*
7	Pregnancy, childbirth and the puerperium (O00-099)	11	0.5	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.4	*
10	Legal intervention (Y35,Y89.0)	8	0.4	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	6	0.3	*
12	Chronic lower respiratory diseases (J40-J47)	6	0.3	*
14	Anemias (D50-D64)	5	0.2	*
15	Septicemia (A40-A41)	4	0.2	*
15	Diabetes mellitus (E10-E14)	4	0.2	*
...	All other causes (Residual)	196	9.2	5.9
Hispanic, both sexes, 20-24 years				
...	All causes	3,287	100.0	87.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,416	43.1	37.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	743	22.6	19.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	357	10.9	9.5
4	Malignant neoplasms (C00-C97)	168	5.1	4.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	86	2.6	2.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	32	1.0	0.8
7	Cerebrovascular diseases (I60-I69)	25	0.8	0.7
8	Legal intervention (Y35,Y89.0)	22	0.7	0.6
9	Influenza and pneumonia (J10-J18)	21	0.6	0.6
10	Human immunodeficiency virus (HIV) disease (B20-B24)	19	0.6	*
11	Pregnancy, childbirth and the puerperium (O00-099)	18	0.5	*
12	Chronic lower respiratory diseases (J40-J47)	11	0.3	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	0.3	*
14	Septicemia (A40-A41)	9	0.3	*
14	Diabetes mellitus (E10-E14)	9	0.3	*
...	All other causes (Residual)	341	10.4	9.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
Hispanic, both sexes, 25-29 years				
...	All causes	3,019	100.0	77.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,164	38.6	29.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	594	19.7	15.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	297	9.8	7.6
4	Malignant neoplasms (C00-C97)	197	6.5	5.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	111	3.7	2.8
6	Human immunodeficiency virus (HIV) disease (B20-B24)	65	2.2	1.7
7	Cerebrovascular diseases (I60-I69)	37	1.2	0.9
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	36	1.2	0.9
9	Pregnancy, childbirth and the puerperium (O00-099)	27	0.9	0.7
10	Diabetes mellitus (E10-E14)	21	0.7	0.5
11	Septicemia (A40-A41)	19	0.6	*
12	Influenza and pneumonia (J10-J18)	18	0.6	*
13	Chronic lower respiratory diseases (J40-J47)	15	0.5	*
14	Legal intervention (Y35,Y89.0)	13	0.4	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	0.4	*
...	All other causes (Residual)	393	13.0	10.0
Hispanic, both sexes, 30-34 years				
...	All causes	2,940	100.0	79.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	922	31.4	24.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	368	12.5	9.9
3	Malignant neoplasms (C00-C97)	329	11.2	8.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	248	8.4	6.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	191	6.5	5.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	146	5.0	3.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	62	2.1	1.7
8	Cerebrovascular diseases (I60-I69)	56	1.9	1.5
9	Diabetes mellitus (E10-E14)	43	1.5	1.2
10	Influenza and pneumonia (J10-J18)	30	1.0	0.8
11	Pregnancy, childbirth and the puerperium (O00-099)	22	0.7	0.6
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	0.6	*
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	17	0.6	*
12	Legal intervention (Y35,Y89.0)	17	0.6	*
15	Septicemia (A40-A41)	16	0.5	*
...	All other causes (Residual)	456	15.5	12.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Hispanic, both sexes, 35-39 years				
...	All causes	3,686	100.0	112.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	940	25.5	28.7
2	Malignant neoplasms (C00-C97)	520	14.1	15.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	325	8.8	9.9
4	Human immunodeficiency virus (HIV) disease (B20-B24)	319	8.7	9.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	269	7.3	8.2
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	229	6.2	7.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	138	3.7	4.2
8	Cerebrovascular diseases (I60-I69)	106	2.9	3.2
9	Diabetes mellitus (E10-E14)	66	1.8	2.0
10	Septicemia (A40-A41)	41	1.1	1.3
11	Influenza and pneumonia (J10-J18)	37	1.0	1.1
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	27	0.7	0.8
13	Viral hepatitis (B15-B19)	24	0.7	0.7
14	Pregnancy, childbirth and the puerperium (O00-099)	16	0.4	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	16	0.4	*
...	All other causes (Residual)	613	16.6	18.7
Hispanic, both sexes, 40-44 years				
...	All causes	4,860	100.0	168.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	883	18.2	30.6
2	Malignant neoplasms (C00-C97)	825	17.0	28.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	609	12.5	21.1
4	Human immunodeficiency virus (HIV) disease (B20-B24)	357	7.3	12.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	330	6.8	11.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	208	4.3	7.2
7	Cerebrovascular diseases (I60-I69)	198	4.1	6.9
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	192	4.0	6.6
9	Diabetes mellitus (E10-E14)	121	2.5	4.2
10	Viral hepatitis (B15-B19)	86	1.8	3.0
11	Chronic lower respiratory diseases (J40-J47)	51	1.0	1.8
12	Septicemia (A40-A41)	47	1.0	1.6
13	Influenza and pneumonia (J10-J18)	45	0.9	1.6
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	45	0.9	1.6
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	24	0.5	0.8
...	All other causes (Residual)	839	17.3	29.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Hispanic, both sexes, 45-49 years				
...	All causes	5,903	100.0	258.4
1	Malignant neoplasms (C00-C97)	1,253	21.2	54.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	890	15.1	39.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	729	12.3	31.9
4	Chronic liver disease and cirrhosis (K70,K73-K74)	463	7.8	20.3
5	Human immunodeficiency virus (HIV) disease (B20-B24)	327	5.5	14.3
6	Cerebrovascular diseases (I60-I69)	260	4.4	11.4
7	Diabetes mellitus (E10-E14)	235	4.0	10.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	176	3.0	7.7
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	139	2.4	6.1
10	Viral hepatitis (B15-B19)	132	2.2	5.8
11	Septicemia (A40-A41)	79	1.3	3.5
12	Influenza and pneumonia (J10-J18)	65	1.1	2.8
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	59	1.0	2.6
14	Chronic lower respiratory diseases (J40-J47)	47	0.8	2.1
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	32	0.5	1.4
...	All other causes (Residual)	1,017	17.2	44.5
Hispanic, both sexes, 50-54 years				
...	All causes	6,725	100.0	385.6
1	Malignant neoplasms (C00-C97)	1,787	26.6	102.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,220	18.1	70.0
3	Chronic liver disease and cirrhosis (K70,K73-K74)	535	8.0	30.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	520	7.7	29.8
5	Cerebrovascular diseases (I60-I69)	330	4.9	18.9
6	Diabetes mellitus (E10-E14)	300	4.5	17.2
7	Human immunodeficiency virus (HIV) disease (B20-B24)	242	3.6	13.9
8	Viral hepatitis (B15-B19)	143	2.1	8.2
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	128	1.9	7.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	123	1.8	7.1
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	99	1.5	5.7
12	Influenza and pneumonia (J10-J18)	88	1.3	5.0
13	Septicemia (A40-A41)	83	1.2	4.8
14	Chronic lower respiratory diseases (J40-J47)	77	1.1	4.4
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	57	0.8	3.3
...	All other causes (Residual)	993	14.8	56.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
Hispanic, both sexes, 55-59 years				
...	All causes	7,248	100.0	557.4
1	Malignant neoplasms (C00-C97)	2,168	29.9	166.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,519	21.0	116.8
3	Diabetes mellitus (E10-E14)	477	6.6	36.7
4	Chronic liver disease and cirrhosis (K70,K73-K74)	412	5.7	31.7
5	Cerebrovascular diseases (I60-I69)	375	5.2	28.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	374	5.2	28.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	145	2.0	11.2
8	Human immunodeficiency virus (HIV) disease (B20-B24)	125	1.7	9.6
9	Chronic lower respiratory diseases (J40-J47)	122	1.7	9.4
10	Septicemia (A40-A41)	102	1.4	7.8
11	Influenza and pneumonia (J10-J18)	99	1.4	7.6
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	98	1.4	7.5
13	Viral hepatitis (B15-B19)	96	1.3	7.4
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	56	0.8	4.3
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	51	0.7	3.9
...	All other causes (Residual)	1,029	14.2	79.1
Hispanic, both sexes, 60-64 years				
...	All causes	7,950	100.0	862.6
1	Malignant neoplasms (C00-C97)	2,491	31.3	270.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,941	24.4	210.6
3	Diabetes mellitus (E10-E14)	596	7.5	64.7
4	Cerebrovascular diseases (I60-I69)	401	5.0	43.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	357	4.5	38.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	304	3.8	33.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	174	2.2	18.9
8	Chronic lower respiratory diseases (J40-J47)	150	1.9	16.3
9	Influenza and pneumonia (J10-J18)	115	1.4	12.5
10	Septicemia (A40-A41)	114	1.4	12.4
11	Viral hepatitis (B15-B19)	75	0.9	8.1
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	65	0.8	7.1
13	Human immunodeficiency virus (HIV) disease (B20-B24)	61	0.8	6.6
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	58	0.7	6.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	53	0.7	5.8
...	All other causes (Residual)	995	12.5	108.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Hispanic, both sexes, 65-69 years				
...	All causes	9,231	100.0	1,289.9
1	Malignant neoplasms (C00-C97)	2,847	30.8	397.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,362	25.6	330.1
3	Diabetes mellitus (E10-E14)	734	8.0	102.6
4	Cerebrovascular diseases (I60-I69)	524	5.7	73.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	312	3.4	43.6
6	Chronic lower respiratory diseases (J40-J47)	256	2.8	35.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	247	2.7	34.5
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	238	2.6	33.3
9	Influenza and pneumonia (J10-J18)	163	1.8	22.8
10	Septicemia (A40-A41)	137	1.5	19.1
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	88	1.0	12.3
12	Viral hepatitis (B15-B19)	52	0.6	7.3
13	Human immunodeficiency virus (HIV) disease (B20-B24)	46	0.5	6.4
14	Alzheimer's disease (G30)	43	0.5	6.0
14	Aortic aneurysm and dissection (I71)	43	0.5	6.0
...	All other causes (Residual)	1,139	12.3	159.2
Hispanic, both sexes, 70-74 years				
...	All causes	11,189	100.0	2,017.4
1	Malignant neoplasms (C00-C97)	3,149	28.1	567.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,027	27.1	545.8
3	Diabetes mellitus (E10-E14)	853	7.6	153.8
4	Cerebrovascular diseases (I60-I69)	708	6.3	127.7
5	Chronic lower respiratory diseases (J40-J47)	401	3.6	72.3
6	Chronic liver disease and cirrhosis (K70,K73-K74)	289	2.6	52.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	276	2.5	49.8
8	Influenza and pneumonia (J10-J18)	259	2.3	46.7
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	235	2.1	42.4
10	Septicemia (A40-A41)	182	1.6	32.8
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	117	1.0	21.1
12	Alzheimer's disease (G30)	110	1.0	19.8
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	74	0.7	13.3
14	Parkinson's disease (G20-G21)	62	0.6	11.2
15	Aortic aneurysm and dissection (I71)	57	0.5	10.3
...	All other causes (Residual)	1,390	12.4	250.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
Hispanic, both sexes, 75-79 years				
...	All causes	13,454	100.0	3,218.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,839	28.5	918.3
2	Malignant neoplasms (C00-C97)	3,281	24.4	784.8
3	Cerebrovascular diseases (I60-I69)	961	7.1	229.9
4	Diabetes mellitus (E10-E14)	960	7.1	229.6
5	Chronic lower respiratory diseases (J40-J47)	520	3.9	124.4
6	Influenza and pneumonia (J10-J18)	402	3.0	96.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	310	2.3	74.2
8	Chronic liver disease and cirrhosis (K70,K73-K74)	262	1.9	62.7
9	Alzheimer's disease (G30)	259	1.9	62.0
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	249	1.9	59.6
11	Septicemia (A40-A41)	214	1.6	51.2
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	146	1.1	34.9
13	Parkinson's disease (G20-G21)	132	1.0	31.6
14	Pneumonitis due to solids and liquids (J69)	76	0.6	18.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	75	0.6	17.9
...	All other causes (Residual)	1,768	13.1	422.9
Hispanic, both sexes, 80-84 years				
...	All causes	13,425	100.0	5,038.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,222	31.4	1,584.5
2	Malignant neoplasms (C00-C97)	2,689	20.0	1,009.2
3	Cerebrovascular diseases (I60-I69)	1,045	7.8	392.2
4	Diabetes mellitus (E10-E14)	809	6.0	303.6
5	Chronic lower respiratory diseases (J40-J47)	566	4.2	212.4
6	Influenza and pneumonia (J10-J18)	455	3.4	170.8
7	Alzheimer's disease (G30)	422	3.1	158.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	319	2.4	119.7
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	259	1.9	97.2
10	Septicemia (A40-A41)	223	1.7	83.7
11	Parkinson's disease (G20-G21)	170	1.3	63.8
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	157	1.2	58.9
13	Chronic liver disease and cirrhosis (K70,K73-K74)	150	1.1	56.3
14	Pneumonitis due to solids and liquids (J69)	104	0.8	39.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	85	0.6	31.9
...	All other causes (Residual)	1,750	13.0	656.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
Hispanic, both sexes, 85-89 years				
...	All causes	9,835	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,344	34.0	---
2	Malignant neoplasms (C00-C97)	1,454	14.8	---
3	Cerebrovascular diseases (I60-I69)	881	9.0	---
4	Influenza and pneumonia (J10-J18)	453	4.6	---
5	Alzheimer's disease (G30)	443	4.5	---
6	Diabetes mellitus (E10-E14)	442	4.5	---
7	Chronic lower respiratory diseases (J40-J47)	433	4.4	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	201	2.0	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	167	1.7	---
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	158	1.6	---
11	Septicemia (A40-A41)	145	1.5	---
12	Parkinson's disease (G20-G21)	113	1.1	---
13	Pneumonitis due to solids and liquids (J69)	103	1.0	---
14	Atherosclerosis (I70)	62	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	47	0.5	---
...	All other causes (Residual)	1,389	14.1	---
Hispanic, both sexes, 90-94 years				
...	All causes	6,766	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,536	37.5	---
2	Malignant neoplasms (C00-C97)	700	10.3	---
3	Cerebrovascular diseases (I60-I69)	569	8.4	---
4	Alzheimer's disease (G30)	377	5.6	---
5	Influenza and pneumonia (J10-J18)	336	5.0	---
6	Chronic lower respiratory diseases (J40-J47)	278	4.1	---
7	Diabetes mellitus (E10-E14)	248	3.7	---
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	143	2.1	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	131	1.9	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	120	1.8	---
11	Septicemia (A40-A41)	100	1.5	---
12	Pneumonitis due to solids and liquids (J69)	73	1.1	---
13	Atherosclerosis (I70)	57	0.8	---
14	Parkinson's disease (G20-G21)	53	0.8	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	34	0.5	---
...	All other causes (Residual)	1,011	14.9	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
Hispanic, both sexes, 95-99 years				
...	All causes	2,703	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,097	40.6	---
2	Cerebrovascular diseases (I60-I69)	221	8.2	---
3	Malignant neoplasms (C00-C97)	201	7.4	---
4	Alzheimer's disease (G30)	174	6.4	---
5	Influenza and pneumonia (J10-J18)	158	5.8	---
6	Chronic lower respiratory diseases (J40-J47)	95	3.5	---
7	Diabetes mellitus (E10-E14)	67	2.5	---
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	49	1.8	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	46	1.7	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	44	1.6	---
11	Septicemia (A40-A41)	39	1.4	---
12	Atherosclerosis (I70)	29	1.1	---
12	Pneumonitis due to solids and liquids (J69)	29	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	0.7	---
15	Parkinson's disease (G20-G21)	13	0.5	---
...	All other causes (Residual)	421	15.6	---
Hispanic, both sexes, 100 years and over				
...	All causes	633	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	269	42.5	---
2	Influenza and pneumonia (J10-J18)	67	10.6	---
3	Alzheimer's disease (G30)	38	6.0	---
3	Cerebrovascular diseases (I60-I69)	38	6.0	---
5	Malignant neoplasms (C00-C97)	23	3.6	---
6	Chronic lower respiratory diseases (J40-J47)	17	2.7	---
7	Diabetes mellitus (E10-E14)	12	1.9	---
7	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	12	1.9	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	1.6	---
10	Atherosclerosis (I70)	9	1.4	---
11	Pneumonitis due to solids and liquids (J69)	8	1.3	---
12	Septicemia (A40-A41)	6	0.9	---
13	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5	0.8	---
14	Anemias (D50-D64)	4	0.6	---
14	Nutritional deficiencies (E40-E64)	4	0.6	---
...	All other causes (Residual)	111	17.5	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
Hispanic, male, all ages				
...	All causes	68,544	100.0	321.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,983	21.9	70.2
2	Malignant neoplasms (C00-C97)	13,002	19.0	60.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,796	11.4	36.5
4	Cerebrovascular diseases (I60-I69)	3,207	4.7	15.0
5	Diabetes mellitus (E10-E14)	2,855	4.2	13.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,782	4.1	13.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,376	3.5	11.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,840	2.7	8.6
9	Chronic lower respiratory diseases (J40-J47)	1,631	2.4	7.6
10	Certain conditions originating in the perinatal period (P00-P96)	1,538	2.2	7.2
11	Influenza and pneumonia (J10-J18)	1,403	2.0	6.6
12	Human immunodeficiency virus (HIV) disease (B20-B24)	1,371	2.0	6.4
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,101	1.6	5.2
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	902	1.3	4.2
15	Septicemia (A40-A41)	801	1.2	3.8
...	All other causes (Residual)	10,956	16.0	51.3
Hispanic, male, 1-4 years				
...	All causes	535	100.0	30.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	186	34.8	10.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	82	15.3	4.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	45	8.4	2.5
4	Malignant neoplasms (C00-C97)	29	5.4	1.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	24	4.5	1.4
6	Septicemia (A40-A41)	14	2.6	*
7	Influenza and pneumonia (J10-J18)	9	1.7	*
8	Chronic lower respiratory diseases (J40-J47)	7	1.3	*
9	Certain conditions originating in the perinatal period (P00-P96)	6	1.1	*
10	Anemias (D50-D64)	3	0.6	*
10	Cerebrovascular diseases (I60-I69)	3	0.6	*
12	Meningococcal infection (A39)	2	0.4	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.4	*
12	Diabetes mellitus (E10-E14)	2	0.4	*
12	Meningitis (G00,G03)	2	0.4	*
12	Acute bronchitis and bronchiolitis (J20-J21)	2	0.4	*
12	Pneumonitis due to solids and liquids (J69)	2	0.4	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
...	All other causes (Residual)	111	20.7	6.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Hispanic, male, 5-9 years				
...	All causes	273	100.0	13.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	102	37.4	5.1
2	Malignant neoplasms (C00-C97)	58	21.2	2.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	24	8.8	1.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	12	4.4	*
5	Septicemia (A40-A41)	6	2.2	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	1.8	*
6	Influenza and pneumonia (J10-J18)	5	1.8	*
8	Chronic lower respiratory diseases (J40-J47)	3	1.1	*
8	Certain conditions originating in the perinatal period (P00-P96)	3	1.1	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.7	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.7	*
12	Anemias (D50-D64)	1	0.4	*
12	Cerebrovascular diseases (I60-I69)	1	0.4	*
12	Acute bronchitis and bronchiolitis (J20-J21)	1	0.4	*
12	Pneumonitis due to solids and liquids (J69)	1	0.4	*
12	Diseases of appendix (K35-K38)	1	0.4	*
...	All other causes (Residual)	46	16.8	2.3
Hispanic, male, 10-14 years				
...	All causes	383	100.0	19.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	139	36.3	7.2
2	Malignant neoplasms (C00-C97)	60	15.7	3.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	28	7.3	1.4
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	25	6.5	1.3
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	24	6.3	1.2
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	10	2.6	*
7	Influenza and pneumonia (J10-J18)	5	1.3	*
8	Cerebrovascular diseases (I60-I69)	4	1.0	*
8	Chronic lower respiratory diseases (J40-J47)	4	1.0	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	1.0	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.5	*
11	Meningitis (G00,G03)	2	0.5	*
11	Certain conditions originating in the perinatal period (P00-P96)	2	0.5	*
14	Shigellosis and amebiasis (A03,A06)	1	0.3	*
14	Tuberculosis (A16-A19)	1	0.3	*
14	Whooping cough (A37)	1	0.3	*
14	Meningococcal infection (A39)	1	0.3	*
14	Viral hepatitis (B15-B19)	1	0.3	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.3	*
14	Anemias (D50-D64)	1	0.3	*
14	Nutritional deficiencies (E40-E64)	1	0.3	*
14	Diseases of appendix (K35-K38)	1	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.3	*
...	All other causes (Residual)	64	16.7	3.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Hispanic, male, 15-19 years				
...	All causes	1,662	100.0	96.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	740	44.5	43.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	445	26.8	25.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	170	10.2	9.9
4	Malignant neoplasms (C00-C97)	94	5.7	5.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	33	2.0	1.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	18	1.1	*
7	Influenza and pneumonia (J10-J18)	8	0.5	*
7	Legal intervention (Y35,Y89.0)	8	0.5	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.4	*
9	Cerebrovascular diseases (I60-I69)	6	0.4	*
11	Diabetes mellitus (E10-E14)	4	0.2	*
11	Chronic lower respiratory diseases (J40-J47)	4	0.2	*
13	Septicemia (A40-A41)	2	0.1	*
13	Aortic aneurysm and dissection (I71)	2	0.1	*
13	Pneumonitis due to solids and liquids (J69)	2	0.1	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.1	*
...	All other causes (Residual)	118	7.1	6.9
Hispanic, male, 20-24 years				
...	All causes	2,714	100.0	130.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,186	43.7	57.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	680	25.1	32.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	317	11.7	15.3
4	Malignant neoplasms (C00-C97)	113	4.2	5.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	68	2.5	3.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	22	0.8	1.1
6	Legal intervention (Y35,Y89.0)	22	0.8	1.1
8	Human immunodeficiency virus (HIV) disease (B20-B24)	18	0.7	*
9	Influenza and pneumonia (J10-J18)	16	0.6	*
10	Cerebrovascular diseases (I60-I69)	14	0.5	*
11	Chronic lower respiratory diseases (J40-J47)	7	0.3	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.3	*
13	Septicemia (A40-A41)	5	0.2	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.2	*
13	Aortic aneurysm and dissection (I71)	5	0.2	*
...	All other causes (Residual)	229	8.4	11.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Hispanic, male, 25-29 years				
...	All causes	2,361	100.0	108.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	979	41.5	45.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	532	22.5	24.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	258	10.9	11.9
4	Malignant neoplasms (C00-C97)	104	4.4	4.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	90	3.8	4.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	52	2.2	2.4
7	Cerebrovascular diseases (I60-I69)	27	1.1	1.2
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	17	0.7	*
9	Legal intervention (Y35,Y89.0)	13	0.6	*
10	Diabetes mellitus (E10-E14)	11	0.5	*
11	Chronic lower respiratory diseases (J40-J47)	10	0.4	*
12	Influenza and pneumonia (J10-J18)	9	0.4	*
13	Chronic liver disease and cirrhosis (K70,K73-K74)	7	0.3	*
14	Septicemia (A40-A41)	6	0.3	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.3	*
...	All other causes (Residual)	240	10.2	11.1
Hispanic, male, 30-34 years				
...	All causes	2,186	100.0	109.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	754	34.5	37.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	312	14.3	15.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	209	9.6	10.5
4	Malignant neoplasms (C00-C97)	163	7.5	8.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	154	7.0	7.7
6	Human immunodeficiency virus (HIV) disease (B20-B24)	117	5.4	5.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	54	2.5	2.7
8	Cerebrovascular diseases (I60-I69)	34	1.6	1.7
9	Diabetes mellitus (E10-E14)	26	1.2	1.3
10	Influenza and pneumonia (J10-J18)	22	1.0	1.1
11	Legal intervention (Y35,Y89.0)	17	0.8	*
12	Septicemia (A40-A41)	10	0.5	*
12	Chronic lower respiratory diseases (J40-J47)	10	0.5	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.4	*
15	Aortic aneurysm and dissection (I71)	7	0.3	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	0.3	*
...	All other causes (Residual)	281	12.9	14.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Hispanic, male, 35-39 years				
...	All causes	2,636	100.0	151.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	763	28.9	43.8
2	Human immunodeficiency virus (HIV) disease (B20-B24)	250	9.5	14.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	245	9.3	14.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	228	8.6	13.1
5	Malignant neoplasms (C00-C97)	227	8.6	13.0
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	193	7.3	11.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	109	4.1	6.3
8	Cerebrovascular diseases (I60-I69)	75	2.8	4.3
9	Diabetes mellitus (E10-E14)	38	1.4	2.2
10	Septicemia (A40-A41)	23	0.9	1.3
11	Influenza and pneumonia (J10-J18)	20	0.8	1.1
12	Viral hepatitis (B15-B19)	16	0.6	*
13	Aortic aneurysm and dissection (I71)	13	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	0.5	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	12	0.5	*
...	All other causes (Residual)	412	15.6	23.7
Hispanic, male, 40-44 years				
...	All causes	3,337	100.0	222.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	691	20.7	46.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	470	14.1	31.3
3	Malignant neoplasms (C00-C97)	377	11.3	25.1
4	Human immunodeficiency virus (HIV) disease (B20-B24)	272	8.2	18.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	268	8.0	17.8
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	158	4.7	10.5
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	156	4.7	10.4
8	Cerebrovascular diseases (I60-I69)	115	3.4	7.7
9	Diabetes mellitus (E10-E14)	73	2.2	4.9
10	Viral hepatitis (B15-B19)	69	2.1	4.6
11	Chronic lower respiratory diseases (J40-J47)	36	1.1	2.4
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	33	1.0	2.2
13	Septicemia (A40-A41)	30	0.9	2.0
14	Influenza and pneumonia (J10-J18)	28	0.8	1.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	0.4	*
...	All other causes (Residual)	549	16.5	36.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Hispanic, male, 45-49 years				
...	All causes	3,984	100.0	343.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	680	17.1	58.6
2	Malignant neoplasms (C00-C97)	607	15.2	52.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	573	14.4	49.4
4	Chronic liver disease and cirrhosis (K70,K73-K74)	379	9.5	32.7
5	Human immunodeficiency virus (HIV) disease (B20-B24)	256	6.4	22.1
6	Cerebrovascular diseases (I60-I69)	159	4.0	13.7
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	139	3.5	12.0
8	Diabetes mellitus (E10-E14)	132	3.3	11.4
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	112	2.8	9.7
10	Viral hepatitis (B15-B19)	100	2.5	8.6
11	Septicemia (A40-A41)	52	1.3	4.5
12	Influenza and pneumonia (J10-J18)	43	1.1	3.7
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	37	0.9	3.2
14	Chronic lower respiratory diseases (J40-J47)	20	0.5	1.7
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	18	0.5	*
...	All other causes (Residual)	677	17.0	58.3
Hispanic, male, 50-54 years				
...	All causes	4,467	100.0	518.0
1	Malignant neoplasms (C00-C97)	959	21.5	111.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	889	19.9	103.1
3	Chronic liver disease and cirrhosis (K70,K73-K74)	446	10.0	51.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	395	8.8	45.8
5	Cerebrovascular diseases (I60-I69)	195	4.4	22.6
6	Diabetes mellitus (E10-E14)	191	4.3	22.1
7	Human immunodeficiency virus (HIV) disease (B20-B24)	189	4.2	21.9
8	Viral hepatitis (B15-B19)	106	2.4	12.3
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	103	2.3	11.9
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	81	1.8	9.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	70	1.6	8.1
12	Influenza and pneumonia (J10-J18)	57	1.3	6.6
13	Septicemia (A40-A41)	54	1.2	6.3
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	40	0.9	4.6
15	Chronic lower respiratory diseases (J40-J47)	36	0.8	4.2
...	All other causes (Residual)	656	14.7	76.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
Hispanic, male, 55-59 years				
...	All causes	4,438	100.0	706.7
1	Malignant neoplasms (C00-C97)	1,140	25.7	181.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,072	24.2	170.7
3	Chronic liver disease and cirrhosis (K70,K73-K74)	310	7.0	49.4
4	Diabetes mellitus (E10-E14)	261	5.9	41.6
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	260	5.9	41.4
6	Cerebrovascular diseases (I60-I69)	219	4.9	34.9
7	Human immunodeficiency virus (HIV) disease (B20-B24)	98	2.2	15.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	85	1.9	13.5
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	73	1.6	11.6
10	Viral hepatitis (B15-B19)	71	1.6	11.3
11	Chronic lower respiratory diseases (J40-J47)	60	1.4	9.6
12	Influenza and pneumonia (J10-J18)	57	1.3	9.1
13	Septicemia (A40-A41)	51	1.1	8.1
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	41	0.9	6.5
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	37	0.8	5.9
...	All other causes (Residual)	603	13.6	96.0
Hispanic, male, 60-64 years				
...	All causes	4,808	100.0	1,118.2
1	Malignant neoplasms (C00-C97)	1,388	28.9	322.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,304	27.1	303.3
3	Diabetes mellitus (E10-E14)	341	7.1	79.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	244	5.1	56.7
5	Cerebrovascular diseases (I60-I69)	235	4.9	54.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	214	4.5	49.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	95	2.0	22.1
8	Chronic lower respiratory diseases (J40-J47)	85	1.8	19.8
9	Influenza and pneumonia (J10-J18)	65	1.4	15.1
10	Septicemia (A40-A41)	58	1.2	13.5
11	Human immunodeficiency virus (HIV) disease (B20-B24)	54	1.1	12.6
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	47	1.0	10.9
13	Viral hepatitis (B15-B19)	39	0.8	9.1
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	31	0.6	7.2
14	Aortic aneurysm and dissection (I71)	31	0.6	7.2
...	All other causes (Residual)	577	12.0	134.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Hispanic, male, 65-69 years				
...	All causes	5,206	100.0	1,606.8
1	Malignant neoplasms (C00-C97)	1,559	29.9	481.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,467	28.2	452.8
3	Diabetes mellitus (E10-E14)	357	6.9	110.2
4	Cerebrovascular diseases (I60-I69)	304	5.8	93.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	178	3.4	54.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	158	3.0	48.8
7	Chronic lower respiratory diseases (J40-J47)	147	2.8	45.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	112	2.2	34.6
9	Influenza and pneumonia (J10-J18)	87	1.7	26.9
10	Septicemia (A40-A41)	74	1.4	22.8
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	52	1.0	16.0
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	37	0.7	11.4
13	Viral hepatitis (B15-B19)	31	0.6	9.6
13	Human immunodeficiency virus (HIV) disease (B20-B24)	31	0.6	9.6
15	Aortic aneurysm and dissection (I71)	26	0.5	8.0
...	All other causes (Residual)	586	11.3	180.9
Hispanic, male, 70-74 years				
...	All causes	6,073	100.0	2,514.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,769	29.1	732.3
2	Malignant neoplasms (C00-C97)	1,754	28.9	726.1
3	Diabetes mellitus (E10-E14)	402	6.6	166.4
4	Cerebrovascular diseases (I60-I69)	343	5.6	142.0
5	Chronic lower respiratory diseases (J40-J47)	240	4.0	99.4
6	Chronic liver disease and cirrhosis (K70,K73-K74)	158	2.6	65.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	139	2.3	57.5
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	136	2.2	56.3
9	Influenza and pneumonia (J10-J18)	133	2.2	55.1
10	Septicemia (A40-A41)	80	1.3	33.1
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	52	0.9	21.5
12	Alzheimer's disease (G30)	50	0.8	20.7
13	Aortic aneurysm and dissection (I71)	45	0.7	18.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	40	0.7	16.6
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	40	0.7	16.6
...	All other causes (Residual)	692	11.4	286.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
Hispanic, male, 75-79 years				
...	All causes	6,928	100.0	3,990.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,083	30.1	1,199.7
2	Malignant neoplasms (C00-C97)	1,828	26.4	1,052.8
3	Cerebrovascular diseases (I60-I69)	437	6.3	251.7
4	Diabetes mellitus (E10-E14)	433	6.3	249.4
5	Chronic lower respiratory diseases (J40-J47)	290	4.2	167.0
6	Influenza and pneumonia (J10-J18)	208	3.0	119.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	144	2.1	82.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	143	2.1	82.4
9	Chronic liver disease and cirrhosis (K70,K73-K74)	132	1.9	76.0
10	Alzheimer's disease (G30)	100	1.4	57.6
11	Septicemia (A40-A41)	98	1.4	56.4
12	Parkinson's disease (G20-G21)	85	1.2	49.0
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	66	1.0	38.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	38	0.5	21.9
14	Pneumonitis due to solids and liquids (J69)	38	0.5	21.9
...	All other causes (Residual)	805	11.6	463.6
Hispanic, male, 80-84 years				
...	All causes	6,469	100.0	6,105.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,080	32.2	1,963.0
2	Malignant neoplasms (C00-C97)	1,455	22.5	1,373.1
3	Cerebrovascular diseases (I60-I69)	467	7.2	440.7
4	Diabetes mellitus (E10-E14)	348	5.4	328.4
5	Chronic lower respiratory diseases (J40-J47)	287	4.4	270.9
6	Influenza and pneumonia (J10-J18)	217	3.4	204.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	162	2.5	152.9
8	Alzheimer's disease (G30)	160	2.5	151.0
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	135	2.1	127.4
10	Parkinson's disease (G20-G21)	101	1.6	95.3
11	Septicemia (A40-A41)	97	1.5	91.5
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	70	1.1	66.1
13	Chronic liver disease and cirrhosis (K70,K73-K74)	66	1.0	62.3
14	Pneumonitis due to solids and liquids (J69)	51	0.8	48.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	40	0.6	37.7
...	All other causes (Residual)	733	11.3	691.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
Hispanic, male, 85-89 years				
...	All causes	3,988	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,321	33.1	---
2	Malignant neoplasms (C00-C97)	689	17.3	---
3	Cerebrovascular diseases (I60-I69)	352	8.8	---
4	Chronic lower respiratory diseases (J40-J47)	206	5.2	---
5	Influenza and pneumonia (J10-J18)	193	4.8	---
6	Diabetes mellitus (E10-E14)	142	3.6	---
7	Alzheimer's disease (G30)	135	3.4	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	91	2.3	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	76	1.9	---
10	Septicemia (A40-A41)	62	1.6	---
10	Parkinson's disease (G20-G21)	62	1.6	---
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	52	1.3	---
13	Pneumonitis due to solids and liquids (J69)	42	1.1	---
14	Atherosclerosis (I70)	29	0.7	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	0.6	---
...	All other causes (Residual)	513	12.9	---
Hispanic, male, 90-94 years				
...	All causes	2,177	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	813	37.3	---
2	Malignant neoplasms (C00-C97)	289	13.3	---
3	Cerebrovascular diseases (I60-I69)	145	6.7	---
4	Chronic lower respiratory diseases (J40-J47)	119	5.5	---
5	Influenza and pneumonia (J10-J18)	117	5.4	---
6	Alzheimer's disease (G30)	78	3.6	---
7	Diabetes mellitus (E10-E14)	71	3.3	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	56	2.6	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	42	1.9	---
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	39	1.8	---
11	Pneumonitis due to solids and liquids (J69)	30	1.4	---
12	Septicemia (A40-A41)	29	1.3	---
13	Atherosclerosis (I70)	23	1.1	---
14	Parkinson's disease (G20-G21)	22	1.0	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	0.6	---
...	All other causes (Residual)	292	13.4	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
Hispanic, male, 95-99 years				
...	All causes	749	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	294	39.3	---
2	Malignant neoplasms (C00-C97)	90	12.0	---
3	Cerebrovascular diseases (I60-I69)	51	6.8	---
4	Influenza and pneumonia (J10-J18)	46	6.1	---
5	Alzheimer's disease (G30)	41	5.5	---
6	Chronic lower respiratory diseases (J40-J47)	37	4.9	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17	2.3	---
8	Diabetes mellitus (E10-E14)	16	2.1	---
9	Septicemia (A40-A41)	14	1.9	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	1.7	---
11	Atherosclerosis (I70)	10	1.3	---
11	Pneumonitis due to solids and liquids (J69)	10	1.3	---
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	9	1.2	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.8	---
14	Parkinson's disease (G20-G21)	6	0.8	---
...	All other causes (Residual)	89	11.9	---
Hispanic, male, 100 years and over				
...	All causes	176	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	68	38.6	---
2	Influenza and pneumonia (J10-J18)	21	11.9	---
3	Alzheimer's disease (G30)	13	7.4	---
4	Malignant neoplasms (C00-C97)	9	5.1	---
5	Cerebrovascular diseases (I60-I69)	8	4.5	---
5	Chronic lower respiratory diseases (J40-J47)	8	4.5	---
7	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	5	2.8	---
7	Pneumonitis due to solids and liquids (J69)	5	2.8	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	2.8	---
10	Diabetes mellitus (E10-E14)	3	1.7	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3	1.7	---
12	Anemias (D50-D64)	1	0.6	---
12	Nutritional deficiencies (E40-E64)	1	0.6	---
12	Atherosclerosis (I70)	1	0.6	---
12	Aortic aneurysm and dissection (I71)	1	0.6	---
12	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.6	---
...	All other causes (Residual)	23	13.1	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
Hispanic, female, all ages				
...	All causes	53,872	100.0	269.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,805	23.8	64.1
2	Malignant neoplasms (C00-C97)	11,520	21.4	57.7
3	Cerebrovascular diseases (I60-I69)	3,574	6.6	17.9
4	Diabetes mellitus (E10-E14)	3,144	5.8	15.7
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,612	4.8	13.1
6	Influenza and pneumonia (J10-J18)	1,509	2.8	7.6
7	Chronic lower respiratory diseases (J40-J47)	1,476	2.7	7.4
8	Alzheimer's disease (G30)	1,291	2.4	6.5
9	Certain conditions originating in the perinatal period (P00-P96)	1,143	2.1	5.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,095	2.0	5.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1,007	1.9	5.0
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	863	1.6	4.3
13	Septicemia (A40-A41)	847	1.6	4.2
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	624	1.2	3.1
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	489	0.9	2.4
...	All other causes (Residual)	9,873	18.3	49.4
Hispanic, female, 1-4 years				
...	All causes	412	100.0	24.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	127	30.8	7.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	56	13.6	3.3
3	Malignant neoplasms (C00-C97)	41	10.0	2.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	35	8.5	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	15	3.6	*
5	Influenza and pneumonia (J10-J18)	15	3.6	*
7	Septicemia (A40-A41)	10	2.4	*
8	Chronic lower respiratory diseases (J40-J47)	4	1.0	*
8	Certain conditions originating in the perinatal period (P00-P96)	4	1.0	*
10	Meningococcal infection (A39)	3	0.7	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.7	*
10	Acute bronchitis and bronchiolitis (J20-J21)	3	0.7	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.7	*
14	Anemias (D50-D64)	2	0.5	*
15	Meningitis (G00,G03)	1	0.2	*
15	Pneumonitis due to solids and liquids (J69)	1	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
...	All other causes (Residual)	88	21.4	5.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Hispanic, female, 5-9 years				
...	All causes	214	100.0	11.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	76	35.5	4.0
2	Malignant neoplasms (C00-C97)	41	19.2	2.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	15	7.0	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	12	5.6	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	11	5.1	*
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	1.9	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	1.4	*
7	Cerebrovascular diseases (I60-I69)	3	1.4	*
7	Chronic lower respiratory diseases (J40-J47)	3	1.4	*
10	Septicemia (A40-A41)	2	0.9	*
10	Certain conditions originating in the perinatal period (P00-P96)	2	0.9	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.5	*
12	Anemias (D50-D64)	1	0.5	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.5	*
...	All other causes (Residual)	39	18.2	2.1
Hispanic, female, 10-14 years				
...	All causes	235	100.0	12.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	74	31.5	4.0
2	Malignant neoplasms (C00-C97)	52	22.1	2.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	15	6.4	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	12	5.1	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	3.8	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	8	3.4	*
7	Influenza and pneumonia (J10-J18)	3	1.3	*
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	1.3	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.9	*
9	Cerebrovascular diseases (I60-I69)	2	0.9	*
9	Chronic lower respiratory diseases (J40-J47)	2	0.9	*
9	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.9	*
13	Septicemia (A40-A41)	1	0.4	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.4	*
13	Aortic aneurysm and dissection (I71)	1	0.4	*
13	Hernia (K40-K46)	1	0.4	*
13	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.4	*
13	Certain conditions originating in the perinatal period (P00-P96)	1	0.4	*
...	All other causes (Residual)	45	19.1	2.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Hispanic, female, 15-19 years				
...	All causes	479	100.0	29.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	216	45.1	13.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	47	9.8	2.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	44	9.2	2.7
4	Malignant neoplasms (C00-C97)	43	9.0	2.7
5	Pregnancy, childbirth and the puerperium (O00-099)	11	2.3	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	11	2.3	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	10	2.1	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	5	1.0	*
8	Anemias (D50-D64)	5	1.0	*
8	Cerebrovascular diseases (I60-I69)	5	1.0	*
11	Influenza and pneumonia (J10-J18)	3	0.6	*
12	Meningococcal infection (A39)	2	0.4	*
12	Septicemia (A40-A41)	2	0.4	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.4	*
12	Chronic lower respiratory diseases (J40-J47)	2	0.4	*
...	All other causes (Residual)	71	14.8	4.4
Hispanic, female, 20-24 years				
...	All causes	573	100.0	33.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	230	40.1	13.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	63	11.0	3.7
2	Malignant neoplasms (C00-C97)	55	9.6	3.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	40	7.0	2.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	18	3.1	*
5	Pregnancy, childbirth and the puerperium (O00-099)	18	3.1	*
7	Cerebrovascular diseases (I60-I69)	11	1.9	*
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	10	1.7	*
9	Diabetes mellitus (E10-E14)	5	0.9	*
9	Influenza and pneumonia (J10-J18)	5	0.9	*
11	Septicemia (A40-A41)	4	0.7	*
11	Chronic lower respiratory diseases (J40-J47)	4	0.7	*
13	Anemias (D50-D64)	3	0.5	*
13	Meningitis (G00,G03)	3	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.5	*
...	All other causes (Residual)	101	17.6	6.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Hispanic, female, 25-29 years				
...	All causes	658	100.0	37.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	185	28.1	10.6
2	Malignant neoplasms (C00-C97)	93	14.1	5.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	62	9.4	3.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	39	5.9	2.2
5	Pregnancy, childbirth and the puerperium (O00-099)	27	4.1	1.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	21	3.2	1.2
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	19	2.9	*
8	Septicemia (A40-A41)	13	2.0	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	13	2.0	*
10	Diabetes mellitus (E10-E14)	10	1.5	*
10	Cerebrovascular diseases (I60-I69)	10	1.5	*
12	Influenza and pneumonia (J10-J18)	9	1.4	*
13	Anemias (D50-D64)	7	1.1	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.9	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.9	*
...	All other causes (Residual)	138	21.0	7.9
Hispanic, female, 30-34 years				
...	All causes	754	100.0	44.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	168	22.3	9.8
2	Malignant neoplasms (C00-C97)	166	22.0	9.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	56	7.4	3.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	39	5.2	2.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	37	4.9	2.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	29	3.8	1.7
7	Cerebrovascular diseases (I60-I69)	22	2.9	1.3
7	Pregnancy, childbirth and the puerperium (O00-099)	22	2.9	1.3
9	Diabetes mellitus (E10-E14)	17	2.3	*
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	10	1.3	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	1.1	*
11	Influenza and pneumonia (J10-J18)	8	1.1	*
11	Chronic liver disease and cirrhosis (K70,K73-K74)	8	1.1	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	1.1	*
15	Septicemia (A40-A41)	6	0.8	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.8	*
...	All other causes (Residual)	144	19.1	8.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Hispanic, female, 35-39 years				
...	All causes	1,050	100.0	68.6
1	Malignant neoplasms (C00-C97)	293	27.9	19.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	177	16.9	11.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	80	7.6	5.2
4	Human immunodeficiency virus (HIV) disease (B20-B24)	69	6.6	4.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	41	3.9	2.7
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	36	3.4	2.4
7	Cerebrovascular diseases (I60-I69)	31	3.0	2.0
8	Chronic liver disease and cirrhosis (K70,K73-K74)	29	2.8	1.9
9	Diabetes mellitus (E10-E14)	28	2.7	1.8
10	Septicemia (A40-A41)	18	1.7	*
11	Influenza and pneumonia (J10-J18)	17	1.6	*
12	Pregnancy, childbirth and the puerperium (O00-099)	16	1.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	1.4	*
14	Viral hepatitis (B15-B19)	8	0.8	*
15	Chronic lower respiratory diseases (J40-J47)	6	0.6	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.6	*
...	All other causes (Residual)	180	17.1	11.8
Hispanic, female, 40-44 years				
...	All causes	1,523	100.0	109.8
1	Malignant neoplasms (C00-C97)	448	29.4	32.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	192	12.6	13.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	139	9.1	10.0
4	Human immunodeficiency virus (HIV) disease (B20-B24)	85	5.6	6.1
5	Cerebrovascular diseases (I60-I69)	83	5.4	6.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	62	4.1	4.5
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	52	3.4	3.7
8	Diabetes mellitus (E10-E14)	48	3.2	3.5
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	34	2.2	2.5
10	Septicemia (A40-A41)	17	1.1	*
10	Viral hepatitis (B15-B19)	17	1.1	*
10	Influenza and pneumonia (J10-J18)	17	1.1	*
13	Chronic lower respiratory diseases (J40-J47)	15	1.0	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	14	0.9	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	0.8	*
...	All other causes (Residual)	288	18.9	20.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
	Hispanic, female, 45-49 years			
...	All causes	1,919	100.0	170.7
1	Malignant neoplasms (C00-C97)	646	33.7	57.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	210	10.9	18.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	156	8.1	13.9
4	Diabetes mellitus (E10-E14)	103	5.4	9.2
5	Cerebrovascular diseases (I60-I69)	101	5.3	9.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	84	4.4	7.5
7	Human immunodeficiency virus (HIV) disease (B20-B24)	71	3.7	6.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	37	1.9	3.3
9	Viral hepatitis (B15-B19)	32	1.7	2.8
10	Septicemia (A40-A41)	27	1.4	2.4
10	Chronic lower respiratory diseases (J40-J47)	27	1.4	2.4
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	27	1.4	2.4
13	Influenza and pneumonia (J10-J18)	22	1.1	2.0
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22	1.1	2.0
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	14	0.7	*
...	All other causes (Residual)	340	17.7	30.2
	Hispanic, female, 50-54 years			
...	All causes	2,258	100.0	256.1
1	Malignant neoplasms (C00-C97)	828	36.7	93.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	331	14.7	37.5
3	Cerebrovascular diseases (I60-I69)	135	6.0	15.3
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	125	5.5	14.2
5	Diabetes mellitus (E10-E14)	109	4.8	12.4
6	Chronic liver disease and cirrhosis (K70,K73-K74)	89	3.9	10.1
7	Human immunodeficiency virus (HIV) disease (B20-B24)	53	2.3	6.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	53	2.3	6.0
9	Chronic lower respiratory diseases (J40-J47)	41	1.8	4.7
10	Viral hepatitis (B15-B19)	37	1.6	4.2
11	Influenza and pneumonia (J10-J18)	31	1.4	3.5
12	Septicemia (A40-A41)	29	1.3	3.3
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	25	1.1	2.8
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	18	0.8	*
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	17	0.8	*
...	All other causes (Residual)	337	14.9	38.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
Hispanic, female, 55-59 years				
...	All causes	2,810	100.0	417.9
1	Malignant neoplasms (C00-C97)	1,028	36.6	152.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	447	15.9	66.5
3	Diabetes mellitus (E10-E14)	216	7.7	32.1
4	Cerebrovascular diseases (I60-I69)	156	5.6	23.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	114	4.1	17.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	102	3.6	15.2
7	Chronic lower respiratory diseases (J40-J47)	62	2.2	9.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	60	2.1	8.9
9	Septicemia (A40-A41)	51	1.8	7.6
10	Influenza and pneumonia (J10-J18)	42	1.5	6.2
11	Human immunodeficiency virus (HIV) disease (B20-B24)	27	1.0	4.0
12	Viral hepatitis (B15-B19)	25	0.9	3.7
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	25	0.9	3.7
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	19	0.7	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	15	0.5	*
...	All other causes (Residual)	421	15.0	62.6
Hispanic, female, 60-64 years				
...	All causes	3,142	100.0	639.0
1	Malignant neoplasms (C00-C97)	1,103	35.1	224.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	637	20.3	129.5
3	Diabetes mellitus (E10-E14)	255	8.1	51.9
4	Cerebrovascular diseases (I60-I69)	166	5.3	33.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	113	3.6	23.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	90	2.9	18.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	79	2.5	16.1
8	Chronic lower respiratory diseases (J40-J47)	65	2.1	13.2
9	Septicemia (A40-A41)	56	1.8	11.4
10	Influenza and pneumonia (J10-J18)	50	1.6	10.2
11	Viral hepatitis (B15-B19)	36	1.1	7.3
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	34	1.1	6.9
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	26	0.8	5.3
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11	0.4	*
15	Atherosclerosis (I70)	9	0.3	*
...	All other causes (Residual)	412	13.1	83.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Hispanic, female, 65-69 years				
...	All causes	4,025	100.0	1,027.7
1	Malignant neoplasms (C00-C97)	1,288	32.0	328.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	895	22.2	228.5
3	Diabetes mellitus (E10-E14)	377	9.4	96.3
4	Cerebrovascular diseases (I60-I69)	220	5.5	56.2
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	135	3.4	34.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	134	3.3	34.2
7	Chronic lower respiratory diseases (J40-J47)	109	2.7	27.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	80	2.0	20.4
9	Influenza and pneumonia (J10-J18)	76	1.9	19.4
10	Septicemia (A40-A41)	63	1.6	16.1
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	36	0.9	9.2
12	Alzheimer's disease (G30)	28	0.7	7.1
13	Viral hepatitis (B15-B19)	21	0.5	5.4
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	21	0.5	5.4
15	Aortic aneurysm and dissection (I71)	17	0.4	*
...	All other causes (Residual)	525	13.0	134.1
Hispanic, female, 70-74 years				
...	All causes	5,116	100.0	1,634.2
1	Malignant neoplasms (C00-C97)	1,395	27.3	445.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,258	24.6	401.8
3	Diabetes mellitus (E10-E14)	451	8.8	144.1
4	Cerebrovascular diseases (I60-I69)	365	7.1	116.6
5	Chronic lower respiratory diseases (J40-J47)	161	3.1	51.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	137	2.7	43.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	131	2.6	41.8
8	Influenza and pneumonia (J10-J18)	126	2.5	40.2
9	Septicemia (A40-A41)	102	2.0	32.6
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	99	1.9	31.6
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	65	1.3	20.8
12	Alzheimer's disease (G30)	60	1.2	19.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	34	0.7	10.9
14	Parkinson's disease (G20-G21)	24	0.5	7.7
15	Viral hepatitis (B15-B19)	23	0.4	7.3
...	All other causes (Residual)	685	13.4	218.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
Hispanic, female, 75-79 years				
...	All causes	6,526	100.0	2,669.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,756	26.9	718.4
2	Malignant neoplasms (C00-C97)	1,453	22.3	594.4
3	Diabetes mellitus (E10-E14)	527	8.1	215.6
4	Cerebrovascular diseases (I60-I69)	524	8.0	214.4
5	Chronic lower respiratory diseases (J40-J47)	230	3.5	94.1
6	Influenza and pneumonia (J10-J18)	194	3.0	79.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	166	2.5	67.9
8	Alzheimer's disease (G30)	159	2.4	65.0
9	Chronic liver disease and cirrhosis (K70,K73-K74)	130	2.0	53.2
10	Septicemia (A40-A41)	116	1.8	47.5
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	106	1.6	43.4
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	80	1.2	32.7
13	Parkinson's disease (G20-G21)	47	0.7	19.2
14	Pneumonitis due to solids and liquids (J69)	38	0.6	15.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	37	0.6	15.1
...	All other causes (Residual)	963	14.8	394.0
Hispanic, female, 80-84 years				
...	All causes	6,956	100.0	4,334.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,142	30.8	1,334.7
2	Malignant neoplasms (C00-C97)	1,234	17.7	768.9
3	Cerebrovascular diseases (I60-I69)	578	8.3	360.1
4	Diabetes mellitus (E10-E14)	461	6.6	287.2
5	Chronic lower respiratory diseases (J40-J47)	279	4.0	173.8
6	Alzheimer's disease (G30)	262	3.8	163.3
7	Influenza and pneumonia (J10-J18)	238	3.4	148.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	157	2.3	97.8
9	Septicemia (A40-A41)	126	1.8	78.5
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	124	1.8	77.3
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	87	1.3	54.2
12	Chronic liver disease and cirrhosis (K70,K73-K74)	84	1.2	52.3
13	Parkinson's disease (G20-G21)	69	1.0	43.0
14	Pneumonitis due to solids and liquids (J69)	53	0.8	33.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	45	0.6	28.0
...	All other causes (Residual)	1,017	14.6	633.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Hispanic, female, 85-89 years				
...	All causes	5,847	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,023	34.6	---
2	Malignant neoplasms (C00-C97)	765	13.1	---
3	Cerebrovascular diseases (I60-I69)	529	9.0	---
4	Alzheimer's disease (G30)	308	5.3	---
5	Diabetes mellitus (E10-E14)	300	5.1	---
6	Influenza and pneumonia (J10-J18)	260	4.4	---
7	Chronic lower respiratory diseases (J40-J47)	227	3.9	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	110	1.9	---
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	106	1.8	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	91	1.6	---
11	Septicemia (A40-A41)	83	1.4	---
12	Pneumonitis due to solids and liquids (J69)	61	1.0	---
13	Parkinson's disease (G20-G21)	51	0.9	---
14	Atherosclerosis (I70)	33	0.6	---
15	Chronic liver disease and cirrhosis (K70,K73-K74)	28	0.5	---
...	All other causes (Residual)	872	14.9	---
Hispanic, female, 90-94 years				
...	All causes	4,589	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,723	37.5	---
2	Cerebrovascular diseases (I60-I69)	424	9.2	---
3	Malignant neoplasms (C00-C97)	411	9.0	---
4	Alzheimer's disease (G30)	299	6.5	---
5	Influenza and pneumonia (J10-J18)	219	4.8	---
6	Diabetes mellitus (E10-E14)	177	3.9	---
7	Chronic lower respiratory diseases (J40-J47)	159	3.5	---
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	104	2.3	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	78	1.7	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	75	1.6	---
11	Septicemia (A40-A41)	71	1.5	---
12	Pneumonitis due to solids and liquids (J69)	43	0.9	---
13	Atherosclerosis (I70)	34	0.7	---
14	Parkinson's disease (G20-G21)	31	0.7	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	22	0.5	---
...	All other causes (Residual)	719	15.7	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
Hispanic, female, 95-99 years				
...	All causes	1,954	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	803	41.1	---
2	Cerebrovascular diseases (I60-I69)	170	8.7	---
3	Alzheimer's disease (G30)	133	6.8	---
4	Influenza and pneumonia (J10-J18)	112	5.7	---
5	Malignant neoplasms (C00-C97)	111	5.7	---
6	Chronic lower respiratory diseases (J40-J47)	58	3.0	---
7	Diabetes mellitus (E10-E14)	51	2.6	---
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	40	2.0	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	31	1.6	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	29	1.5	---
11	Septicemia (A40-A41)	25	1.3	---
12	Atherosclerosis (I70)	19	1.0	---
12	Pneumonitis due to solids and liquids (J69)	19	1.0	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	0.7	---
15	Nutritional deficiencies (E40-E64)	10	0.5	---
...	All other causes (Residual)	329	16.8	---
Hispanic, female, 100 years and over				
...	All causes	457	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	201	44.0	---
2	Influenza and pneumonia (J10-J18)	46	10.1	---
3	Cerebrovascular diseases (I60-I69)	30	6.6	---
4	Alzheimer's disease (G30)	25	5.5	---
5	Malignant neoplasms (C00-C97)	14	3.1	---
6	Diabetes mellitus (E10-E14)	9	2.0	---
6	Chronic lower respiratory diseases (J40-J47)	9	2.0	---
8	Atherosclerosis (I70)	8	1.8	---
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	7	1.5	---
10	Septicemia (A40-A41)	6	1.3	---
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	1.1	---
12	Anemias (D50-D64)	3	0.7	---
12	Nutritional deficiencies (E40-E64)	3	0.7	---
12	Pneumonitis due to solids and liquids (J69)	3	0.7	---
15	Aortic aneurysm and dissection (I71)	2	0.4	---
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	2	0.4	---
15	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2	0.4	---
...	All other causes (Residual)	82	17.9	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
White, non-Hispanic, both sexes, all ages				
...	All causes	1,933,382	100.0	967.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	537,512	27.8	269.1
2	Malignant neoplasms (C00-C97)	453,534	23.5	227.0
3	Cerebrovascular diseases (I60-I69)	121,065	6.3	60.6
4	Chronic lower respiratory diseases (J40-J47)	109,670	5.7	54.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	85,466	4.4	42.8
6	Alzheimer's disease (G30)	59,153	3.1	29.6
7	Diabetes mellitus (E10-E14)	52,101	2.7	26.1
8	Influenza and pneumonia (J10-J18)	49,456	2.6	24.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	31,508	1.6	15.8
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	27,001	1.4	13.5
11	Septicemia (A40-A41)	25,041	1.3	12.5
12	Chronic liver disease and cirrhosis (K70,K73-K74)	20,144	1.0	10.1
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	16,569	0.9	8.3
14	Parkinson's disease (G20-G21)	16,428	0.8	8.2
15	Pneumonitis due to solids and liquids (J69)	14,411	0.7	7.2
...	All other causes (Residual)	314,323	16.3	157.3
White, non-Hispanic, both sexes, 1-4 years				
...	All causes	2,458	100.0	26.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	933	38.0	10.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	270	11.0	2.9
3	Malignant neoplasms (C00-C97)	244	9.9	2.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	131	5.3	1.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	80	3.3	0.9
6	Influenza and pneumonia (J10-J18)	55	2.2	0.6
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	39	1.6	0.4
8	Septicemia (A40-A41)	36	1.5	0.4
9	Cerebrovascular diseases (I60-I69)	25	1.0	0.3
10	Certain conditions originating in the perinatal period (P00-P96)	24	1.0	0.3
11	Meningitis (G00,G03)	16	0.7	*
12	Chronic lower respiratory diseases (J40-J47)	12	0.5	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	11	0.4	*
14	Meningococcal infection (A39)	10	0.4	*
15	Anemias (D50-D64)	8	0.3	*
...	All other causes (Residual)	564	22.9	6.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
White, non-Hispanic, both sexes, 5-9 years				
...	All causes	1,573	100.0	13.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	620	39.4	5.3
2	Malignant neoplasms (C00-C97)	322	20.5	2.8
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	111	7.1	0.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	48	3.1	0.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	38	2.4	0.3
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	25	1.6	0.2
7	Septicemia (A40-A41)	23	1.5	0.2
8	Cerebrovascular diseases (I60-I69)	22	1.4	0.2
9	Influenza and pneumonia (J10-J18)	14	0.9	*
10	Certain conditions originating in the perinatal period (P00-P96)	13	0.8	*
11	Chronic lower respiratory diseases (J40-J47)	11	0.7	*
12	Meningitis (G00,G03)	10	0.6	*
13	Anemias (D50-D64)	8	0.5	*
14	Meningococcal infection (A39)	7	0.4	*
15	Diabetes mellitus (E10-E14)	4	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.3	*
...	All other causes (Residual)	293	18.6	2.5
White, non-Hispanic, both sexes, 10-14 years				
...	All causes	2,247	100.0	17.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	953	42.4	7.4
2	Malignant neoplasms (C00-C97)	265	11.8	2.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	179	8.0	1.4
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	106	4.7	0.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	87	3.9	0.7
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	71	3.2	0.6
7	Chronic lower respiratory diseases (J40-J47)	30	1.3	0.2
8	Influenza and pneumonia (J10-J18)	29	1.3	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	27	1.2	0.2
10	Cerebrovascular diseases (I60-I69)	26	1.2	0.2
11	Septicemia (A40-A41)	18	0.8	*
12	Diabetes mellitus (E10-E14)	15	0.7	*
13	Anemias (D50-D64)	9	0.4	*
14	Meningitis (G00,G03)	7	0.3	*
14	Pneumonitis due to solids and liquids (J69)	7	0.3	*
...	All other causes (Residual)	418	18.6	3.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
White, non-Hispanic, both sexes, 15-19 years				
...	All causes	8,346	100.0	63.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,911	58.8	37.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,211	14.5	9.2
3	Malignant neoplasms (C00-C97)	442	5.3	3.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	341	4.1	2.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	191	2.3	1.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	155	1.9	1.2
7	Influenza and pneumonia (J10-J18)	41	0.5	0.3
8	Cerebrovascular diseases (I60-I69)	39	0.5	0.3
9	Chronic lower respiratory diseases (J40-J47)	33	0.4	0.3
10	Septicemia (A40-A41)	30	0.4	0.2
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	28	0.3	0.2
12	Diabetes mellitus (E10-E14)	25	0.3	0.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	0.2	*
14	Anemias (D50-D64)	16	0.2	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	16	0.2	*
...	All other causes (Residual)	850	10.2	6.5
White, non-Hispanic, both sexes, 20-24 years				
...	All causes	11,338	100.0	87.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,936	52.4	45.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,834	16.2	14.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	576	5.1	4.4
4	Malignant neoplasms (C00-C97)	574	5.1	4.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	362	3.2	2.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	135	1.2	1.0
7	Cerebrovascular diseases (I60-I69)	78	0.7	0.6
8	Influenza and pneumonia (J10-J18)	67	0.6	0.5
9	Diabetes mellitus (E10-E14)	62	0.5	0.5
10	Pregnancy, childbirth and the puerperium (O00-099)	47	0.4	0.4
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	44	0.4	0.3
12	Chronic lower respiratory diseases (J40-J47)	42	0.4	0.3
13	Septicemia (A40-A41)	40	0.4	0.3
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	24	0.2	0.2
15	Human immunodeficiency virus (HIV) disease (B20-B24)	22	0.2	0.2
...	All other causes (Residual)	1,495	13.2	11.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
White, non-Hispanic, both sexes, 25-29 years				
...	All causes	10,452	100.0	89.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,461	42.7	38.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,707	16.3	14.6
3	Malignant neoplasms (C00-C97)	795	7.6	6.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	580	5.5	4.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	497	4.8	4.2
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	139	1.3	1.2
7	Cerebrovascular diseases (I60-I69)	102	1.0	0.9
8	Diabetes mellitus (E10-E14)	95	0.9	0.8
9	Influenza and pneumonia (J10-J18)	64	0.6	0.5
10	Human immunodeficiency virus (HIV) disease (B20-B24)	62	0.6	0.5
11	Septicemia (A40-A41)	58	0.6	0.5
12	Pregnancy, childbirth and the puerperium (O00-099)	52	0.5	0.4
13	Chronic lower respiratory diseases (J40-J47)	48	0.5	0.4
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	43	0.4	0.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	37	0.4	0.3
...	All other causes (Residual)	1,712	16.4	14.6
White, non-Hispanic, both sexes, 30-34 years				
...	All causes	12,945	100.0	102.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,251	32.8	33.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,040	15.8	16.1
3	Malignant neoplasms (C00-C97)	1,475	11.4	11.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,165	9.0	9.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	479	3.7	3.8
6	Diabetes mellitus (E10-E14)	238	1.8	1.9
7	Human immunodeficiency virus (HIV) disease (B20-B24)	234	1.8	1.8
8	Cerebrovascular diseases (I60-I69)	178	1.4	1.4
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	156	1.2	1.2
10	Chronic liver disease and cirrhosis (K70,K73-K74)	149	1.2	1.2
11	Influenza and pneumonia (J10-J18)	119	0.9	0.9
12	Septicemia (A40-A41)	114	0.9	0.9
13	Chronic lower respiratory diseases (J40-J47)	72	0.6	0.6
14	Pregnancy, childbirth and the puerperium (O00-099)	70	0.5	0.6
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	65	0.5	0.5
...	All other causes (Residual)	2,140	16.5	16.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
White, non-Hispanic, both sexes, 35-39 years				
...	All causes	20,033	100.0	145.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,159	25.8	37.4
2	Malignant neoplasms (C00-C97)	2,994	14.9	21.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,567	12.8	18.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,404	12.0	17.4
5	Human immunodeficiency virus (HIV) disease (B20-B24)	603	3.0	4.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	513	2.6	3.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	503	2.5	3.6
8	Diabetes mellitus (E10-E14)	419	2.1	3.0
9	Cerebrovascular diseases (I60-I69)	388	1.9	2.8
10	Septicemia (A40-A41)	178	0.9	1.3
10	Influenza and pneumonia (J10-J18)	178	0.9	1.3
12	Chronic lower respiratory diseases (J40-J47)	175	0.9	1.3
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	150	0.7	1.1
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	144	0.7	1.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	90	0.4	0.7
...	All other causes (Residual)	3,568	17.8	25.9
White, non-Hispanic, both sexes, 40-44 years				
...	All causes	35,510	100.0	220.9
1	Malignant neoplasms (C00-C97)	7,068	19.9	44.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,866	19.3	42.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,844	16.5	36.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,194	9.0	19.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,360	3.8	8.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	926	2.6	5.8
7	Cerebrovascular diseases (I60-I69)	868	2.4	5.4
8	Diabetes mellitus (E10-E14)	819	2.3	5.1
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	611	1.7	3.8
10	Chronic lower respiratory diseases (J40-J47)	408	1.1	2.5
11	Influenza and pneumonia (J10-J18)	374	1.1	2.3
12	Viral hepatitis (B15-B19)	305	0.9	1.9
13	Septicemia (A40-A41)	296	0.8	1.8
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	250	0.7	1.6
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	219	0.6	1.4
...	All other causes (Residual)	6,102	17.2	38.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
White, non-Hispanic, both sexes, 45-49 years				
...	All causes	53,263	100.0	331.1
1	Malignant neoplasms (C00-C97)	13,736	25.8	85.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,463	19.6	65.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,124	13.4	44.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,269	6.1	20.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,368	4.4	14.7
6	Cerebrovascular diseases (I60-I69)	1,400	2.6	8.7
7	Diabetes mellitus (E10-E14)	1,389	2.6	8.6
8	Chronic lower respiratory diseases (J40-J47)	892	1.7	5.5
9	Human immunodeficiency virus (HIV) disease (B20-B24)	789	1.5	4.9
10	Viral hepatitis (B15-B19)	658	1.2	4.1
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	551	1.0	3.4
12	Influenza and pneumonia (J10-J18)	521	1.0	3.2
13	Septicemia (A40-A41)	510	1.0	3.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	400	0.8	2.5
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	258	0.5	1.6
...	All other causes (Residual)	8,935	16.8	55.5
White, non-Hispanic, both sexes, 50-54 years				
...	All causes	69,774	100.0	477.6
1	Malignant neoplasms (C00-C97)	22,280	31.9	152.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,395	22.1	105.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,620	8.1	38.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	2,962	4.2	20.3
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,832	4.1	19.4
6	Diabetes mellitus (E10-E14)	2,073	3.0	14.2
7	Cerebrovascular diseases (I60-I69)	1,955	2.8	13.4
8	Chronic lower respiratory diseases (J40-J47)	1,803	2.6	12.3
9	Viral hepatitis (B15-B19)	859	1.2	5.9
10	Septicemia (A40-A41)	849	1.2	5.8
11	Influenza and pneumonia (J10-J18)	721	1.0	4.9
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	605	0.9	4.1
13	Human immunodeficiency virus (HIV) disease (B20-B24)	540	0.8	3.7
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	382	0.5	2.6
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	306	0.4	2.1
...	All other causes (Residual)	10,592	15.2	72.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
White, non-Hispanic, both sexes, 55-59 years				
...	All causes	89,100	100.0	696.6
1	Malignant neoplasms (C00-C97)	33,363	37.4	260.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,700	23.2	161.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,106	4.6	32.1
4	Chronic lower respiratory diseases (J40-J47)	3,531	4.0	27.6
5	Diabetes mellitus (E10-E14)	3,089	3.5	24.1
6	Cerebrovascular diseases (I60-I69)	2,670	3.0	20.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,609	2.9	20.4
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,088	2.3	16.3
9	Septicemia (A40-A41)	1,134	1.3	8.9
10	Influenza and pneumonia (J10-J18)	973	1.1	7.6
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	946	1.1	7.4
12	Viral hepatitis (B15-B19)	500	0.6	3.9
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	440	0.5	3.4
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	399	0.4	3.1
15	Aortic aneurysm and dissection (I71)	357	0.4	2.8
...	All other causes (Residual)	12,195	13.7	95.3
White, non-Hispanic, both sexes, 60-64 years				
...	All causes	109,742	100.0	1,105.0
1	Malignant neoplasms (C00-C97)	43,037	39.2	433.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	26,234	23.9	264.2
3	Chronic lower respiratory diseases (J40-J47)	6,659	6.1	67.1
4	Diabetes mellitus (E10-E14)	3,876	3.5	39.0
5	Cerebrovascular diseases (I60-I69)	3,651	3.3	36.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,284	3.0	33.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,229	2.0	22.4
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,496	1.4	15.1
9	Septicemia (A40-A41)	1,379	1.3	13.9
10	Influenza and pneumonia (J10-J18)	1,322	1.2	13.3
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,292	1.2	13.0
12	Aortic aneurysm and dissection (I71)	628	0.6	6.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	568	0.5	5.7
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	561	0.5	5.6
15	Alzheimer's disease (G30)	338	0.3	3.4
...	All other causes (Residual)	13,188	12.0	132.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
White, non-Hispanic, both sexes, 65-69 years				
...	All causes	133,916	100.0	1,700.5
1	Malignant neoplasms (C00-C97)	51,307	38.3	651.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	31,971	23.9	406.0
3	Chronic lower respiratory diseases (J40-J47)	10,125	7.6	128.6
4	Cerebrovascular diseases (I60-I69)	5,461	4.1	69.3
5	Diabetes mellitus (E10-E14)	4,704	3.5	59.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,027	2.3	38.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,994	1.5	25.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,838	1.4	23.3
9	Influenza and pneumonia (J10-J18)	1,735	1.3	22.0
10	Septicemia (A40-A41)	1,729	1.3	22.0
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,022	0.8	13.0
12	Aortic aneurysm and dissection (I71)	959	0.7	12.2
13	Alzheimer's disease (G30)	851	0.6	10.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	761	0.6	9.7
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	756	0.6	9.6
...	All other causes (Residual)	15,676	11.7	199.1
White, non-Hispanic, both sexes, 70-74 years				
...	All causes	183,690	100.0	2,669.8
1	Malignant neoplasms (C00-C97)	62,841	34.2	913.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	45,681	24.9	663.9
3	Chronic lower respiratory diseases (J40-J47)	15,440	8.4	224.4
4	Cerebrovascular diseases (I60-I69)	9,102	5.0	132.3
5	Diabetes mellitus (E10-E14)	6,171	3.4	89.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,505	1.9	50.9
7	Influenza and pneumonia (J10-J18)	3,159	1.7	45.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,907	1.6	42.3
9	Septicemia (A40-A41)	2,599	1.4	37.8
10	Alzheimer's disease (G30)	2,256	1.2	32.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	2,009	1.1	29.2
12	Aortic aneurysm and dissection (I71)	1,574	0.9	22.9
13	Parkinson's disease (G20-G21)	1,431	0.8	20.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,188	0.6	17.3
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,184	0.6	17.2
...	All other causes (Residual)	22,643	12.3	329.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
White, non-Hispanic, both sexes, 75-79 years				
...	All causes	259,718	100.0	4,207.6
1	Malignant neoplasms (C00-C97)	73,822	28.4	1,196.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	69,171	26.6	1,120.6
3	Chronic lower respiratory diseases (J40-J47)	21,351	8.2	345.9
4	Cerebrovascular diseases (I60-I69)	16,445	6.3	266.4
5	Diabetes mellitus (E10-E14)	8,066	3.1	130.7
6	Alzheimer's disease (G30)	6,458	2.5	104.6
7	Influenza and pneumonia (J10-J18)	5,705	2.2	92.4
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,044	1.9	81.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,821	1.9	78.1
10	Septicemia (A40-A41)	3,777	1.5	61.2
11	Parkinson's disease (G20-G21)	3,187	1.2	51.6
12	Aortic aneurysm and dissection (I71)	2,155	0.8	34.9
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,016	0.8	32.7
14	Chronic liver disease and cirrhosis (K70,K73-K74)	1,831	0.7	29.7
15	Pneumonitis due to solids and liquids (J69)	1,777	0.7	28.8
...	All other causes (Residual)	34,092	13.1	552.3
White, non-Hispanic, both sexes, 80-84 years				
...	All causes	322,736	100.0	6,807.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	96,088	29.8	2,026.9
2	Malignant neoplasms (C00-C97)	68,639	21.3	1,447.9
3	Cerebrovascular diseases (I60-I69)	25,225	7.8	532.1
4	Chronic lower respiratory diseases (J40-J47)	22,262	6.9	469.6
5	Alzheimer's disease (G30)	12,980	4.0	273.8
6	Influenza and pneumonia (J10-J18)	9,438	2.9	199.1
7	Diabetes mellitus (E10-E14)	9,154	2.8	193.1
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,730	2.1	142.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,285	1.9	132.6
10	Parkinson's disease (G20-G21)	4,812	1.5	101.5
11	Septicemia (A40-A41)	4,649	1.4	98.1
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	3,082	1.0	65.0
13	Pneumonitis due to solids and liquids (J69)	3,034	0.9	64.0
14	Aortic aneurysm and dissection (I71)	2,418	0.7	51.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,337	0.7	49.3
...	All other causes (Residual)	45,603	14.1	961.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
White, non-Hispanic, both sexes, 85-89 years				
...	All causes	299,301	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	99,626	33.3	---
2	Malignant neoplasms (C00-C97)	44,719	14.9	---
3	Cerebrovascular diseases (I60-I69)	26,554	8.9	---
4	Alzheimer's disease (G30)	16,698	5.6	---
5	Chronic lower respiratory diseases (J40-J47)	16,437	5.5	---
6	Influenza and pneumonia (J10-J18)	10,788	3.6	---
7	Diabetes mellitus (E10-E14)	6,979	2.3	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,221	2.1	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,080	2.0	---
10	Septicemia (A40-A41)	4,096	1.4	---
11	Parkinson's disease (G20-G21)	3,860	1.3	---
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	3,506	1.2	---
13	Pneumonitis due to solids and liquids (J69)	3,439	1.1	---
14	Atherosclerosis (I70)	2,447	0.8	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,978	0.7	---
...	All other causes (Residual)	45,873	15.3	---
White, non-Hispanic, both sexes, 90-94 years				
...	All causes	201,488	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	74,261	36.9	---
2	Malignant neoplasms (C00-C97)	19,877	9.9	---
3	Cerebrovascular diseases (I60-I69)	18,757	9.3	---
4	Alzheimer's disease (G30)	13,147	6.5	---
5	Influenza and pneumonia (J10-J18)	8,997	4.5	---
6	Chronic lower respiratory diseases (J40-J47)	7,843	3.9	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,274	2.1	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,009	2.0	---
9	Diabetes mellitus (E10-E14)	3,714	1.8	---
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,781	1.4	---
11	Pneumonitis due to solids and liquids (J69)	2,501	1.2	---
12	Septicemia (A40-A41)	2,436	1.2	---
13	Atherosclerosis (I70)	2,177	1.1	---
14	Parkinson's disease (G20-G21)	1,783	0.9	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,070	0.5	---
...	All other causes (Residual)	33,861	16.8	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
White, non-Hispanic, both sexes, 95-99 years				
...	All causes	77,094	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	30,248	39.2	---
2	Cerebrovascular diseases (I60-I69)	6,838	8.9	---
3	Alzheimer's disease (G30)	5,272	6.8	---
4	Malignant neoplasms (C00-C97)	5,042	6.5	---
5	Influenza and pneumonia (J10-J18)	4,034	5.2	---
6	Chronic lower respiratory diseases (J40-J47)	2,201	2.9	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,585	2.1	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,456	1.9	---
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,284	1.7	---
10	Atherosclerosis (I70)	1,135	1.5	---
11	Diabetes mellitus (E10-E14)	1,055	1.4	---
12	Pneumonitis due to solids and liquids (J69)	981	1.3	---
13	Septicemia (A40-A41)	856	1.1	---
14	Parkinson's disease (G20-G21)	403	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	327	0.4	---
...	All other causes (Residual)	14,377	18.6	---
White, non-Hispanic, both sexes, 100 years and over				
...	All causes	15,499	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,534	42.2	---
2	Cerebrovascular diseases (I60-I69)	1,222	7.9	---
3	Influenza and pneumonia (J10-J18)	1,010	6.5	---
4	Alzheimer's disease (G30)	923	6.0	---
5	Malignant neoplasms (C00-C97)	645	4.2	---
6	Atherosclerosis (I70)	305	2.0	---
7	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	304	2.0	---
8	Chronic lower respiratory diseases (J40-J47)	280	1.8	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	264	1.7	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	251	1.6	---
11	Pneumonitis due to solids and liquids (J69)	203	1.3	---
12	Diabetes mellitus (E10-E14)	148	1.0	---
13	Septicemia (A40-A41)	121	0.8	---
14	Anemias (D50-D64)	63	0.4	---
15	Nutritional deficiencies (E40-E64)	48	0.3	---
...	All other causes (Residual)	3,178	20.5	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
White, non-Hispanic, male, all ages				
...	All causes	938,143	100.0	957.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	263,638	28.1	269.1
2	Malignant neoplasms (C00-C97)	234,392	25.0	239.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	53,465	5.7	54.6
4	Chronic lower respiratory diseases (J40-J47)	51,909	5.5	53.0
5	Cerebrovascular diseases (I60-I69)	46,087	4.9	47.0
6	Diabetes mellitus (E10-E14)	25,773	2.7	26.3
7	Influenza and pneumonia (J10-J18)	21,905	2.3	22.4
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	21,214	2.3	21.6
9	Alzheimer's disease (G30)	16,989	1.8	17.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15,247	1.6	15.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	12,931	1.4	13.2
12	Septicemia (A40-A41)	11,188	1.2	11.4
13	Parkinson's disease (G20-G21)	9,469	1.0	9.7
14	Pneumonitis due to solids and liquids (J69)	7,367	0.8	7.5
15	Aortic aneurysm and dissection (I71)	6,906	0.7	7.0
...	All other causes (Residual)	139,663	14.9	142.5
White, non-Hispanic, male, 1-4 years				
...	All causes	1,365	100.0	29.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	553	40.5	11.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	145	10.6	3.1
3	Malignant neoplasms (C00-C97)	134	9.8	2.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	73	5.3	1.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	37	2.7	0.8
6	Influenza and pneumonia (J10-J18)	23	1.7	0.5
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	21	1.5	0.4
8	Certain conditions originating in the perinatal period (P00-P96)	17	1.2	*
9	Septicemia (A40-A41)	15	1.1	*
9	Cerebrovascular diseases (I60-I69)	15	1.1	*
11	Meningitis (G00,G03)	10	0.7	*
12	Meningococcal infection (A39)	8	0.6	*
12	Chronic lower respiratory diseases (J40-J47)	8	0.6	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.3	*
...	All other causes (Residual)	296	21.7	6.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
White, non-Hispanic, male, 5-9 years				
...	All causes	914	100.0	15.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	377	41.2	6.3
2	Malignant neoplasms (C00-C97)	188	20.6	3.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	59	6.5	1.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	25	2.7	0.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	17	1.9	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	1.5	*
7	Cerebrovascular diseases (I60-I69)	13	1.4	*
8	Septicemia (A40-A41)	11	1.2	*
9	Certain conditions originating in the perinatal period (P00-P96)	8	0.9	*
10	Meningitis (G00,G03)	6	0.7	*
10	Influenza and pneumonia (J10-J18)	6	0.7	*
12	Chronic lower respiratory diseases (J40-J47)	5	0.5	*
13	Meningococcal infection (A39)	4	0.4	*
13	Anemias (D50-D64)	4	0.4	*
15	Diseases of appendix (K35-K38)	3	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.3	*
...	All other causes (Residual)	171	18.7	2.8
White, non-Hispanic, male, 10-14 years				
...	All causes	1,350	100.0	20.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	614	45.5	9.3
2	Malignant neoplasms (C00-C97)	151	11.2	2.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	118	8.7	1.8
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	52	3.9	0.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	50	3.7	0.8
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	45	3.3	0.7
7	Chronic lower respiratory diseases (J40-J47)	18	1.3	*
8	Influenza and pneumonia (J10-J18)	16	1.2	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	15	1.1	*
10	Cerebrovascular diseases (I60-I69)	11	0.8	*
11	Septicemia (A40-A41)	9	0.7	*
12	Diabetes mellitus (E10-E14)	7	0.5	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.3	*
14	Meningococcal infection (A39)	3	0.2	*
14	Meningitis (G00,G03)	3	0.2	*
14	Aortic aneurysm and dissection (I71)	3	0.2	*
14	Pneumonitis due to solids and liquids (J69)	3	0.2	*
14	Diseases of appendix (K35-K38)	3	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.2	*
14	Certain conditions originating in the perinatal period (P00-P96)	3	0.2	*
...	All other causes (Residual)	219	16.2	3.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
White, non-Hispanic, male, 15-19 years				
...	All causes	5,636	100.0	83.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,322	58.9	49.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	961	17.1	14.2
3	Malignant neoplasms (C00-C97)	254	4.5	3.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	223	4.0	3.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	121	2.1	1.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	86	1.5	1.3
7	Influenza and pneumonia (J10-J18)	26	0.5	0.4
8	Cerebrovascular diseases (I60-I69)	21	0.4	0.3
8	Chronic lower respiratory diseases (J40-J47)	21	0.4	0.3
10	Septicemia (A40-A41)	17	0.3	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	0.2	*
12	Legal intervention (Y35,Y89.0)	13	0.2	*
13	Diabetes mellitus (E10-E14)	12	0.2	*
14	Anemias (D50-D64)	11	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.1	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	8	0.1	*
...	All other causes (Residual)	518	9.2	7.7
White, non-Hispanic, male, 20-24 years				
...	All causes	8,381	100.0	126.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,563	54.4	68.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,574	18.8	23.8
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	401	4.8	6.1
4	Malignant neoplasms (C00-C97)	339	4.0	5.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	242	2.9	3.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	70	0.8	1.1
7	Influenza and pneumonia (J10-J18)	47	0.6	0.7
8	Cerebrovascular diseases (I60-I69)	42	0.5	0.6
9	Diabetes mellitus (E10-E14)	33	0.4	0.5
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	27	0.3	0.4
11	Chronic lower respiratory diseases (J40-J47)	24	0.3	0.4
12	Septicemia (A40-A41)	22	0.3	0.3
13	Aortic aneurysm and dissection (I71)	16	0.2	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	14	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	0.2	*
15	Legal intervention (Y35,Y89.0)	13	0.2	*
...	All other causes (Residual)	941	11.2	14.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
White, non-Hispanic, male, 25-29 years				
...	All causes	7,473	100.0	126.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,374	45.1	57.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,430	19.1	24.1
3	Malignant neoplasms (C00-C97)	443	5.9	7.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	386	5.2	6.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	344	4.6	5.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	86	1.2	1.5
7	Cerebrovascular diseases (I60-I69)	58	0.8	1.0
8	Diabetes mellitus (E10-E14)	54	0.7	0.9
9	Influenza and pneumonia (J10-J18)	44	0.6	0.7
10	Human immunodeficiency virus (HIV) disease (B20-B24)	43	0.6	0.7
11	Septicemia (A40-A41)	28	0.4	0.5
12	Aortic aneurysm and dissection (I71)	26	0.3	0.4
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	24	0.3	0.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	22	0.3	0.4
15	Chronic lower respiratory diseases (J40-J47)	20	0.3	0.3
...	All other causes (Residual)	1,091	14.6	18.4
White, non-Hispanic, male, 30-34 years				
...	All causes	8,625	100.0	135.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,084	35.8	48.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,634	18.9	25.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	808	9.4	12.7
4	Malignant neoplasms (C00-C97)	701	8.1	11.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	328	3.8	5.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	184	2.1	2.9
7	Diabetes mellitus (E10-E14)	137	1.6	2.1
8	Chronic liver disease and cirrhosis (K70,K73-K74)	104	1.2	1.6
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	82	1.0	1.3
10	Cerebrovascular diseases (I60-I69)	79	0.9	1.2
11	Influenza and pneumonia (J10-J18)	71	0.8	1.1
12	Septicemia (A40-A41)	57	0.7	0.9
13	Chronic lower respiratory diseases (J40-J47)	43	0.5	0.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	37	0.4	0.6
15	Aortic aneurysm and dissection (I71)	26	0.3	0.4
...	All other causes (Residual)	1,250	14.5	19.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
White, non-Hispanic, male, 35-39 years				
...	All causes	12,860	100.0	185.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,645	28.3	52.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,864	14.5	26.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,789	13.9	25.8
4	Malignant neoplasms (C00-C97)	1,317	10.2	19.0
5	Human immunodeficiency virus (HIV) disease (B20-B24)	475	3.7	6.9
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	357	2.8	5.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	320	2.5	4.6
8	Diabetes mellitus (E10-E14)	270	2.1	3.9
9	Cerebrovascular diseases (I60-I69)	190	1.5	2.7
10	Influenza and pneumonia (J10-J18)	103	0.8	1.5
11	Septicemia (A40-A41)	99	0.8	1.4
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	86	0.7	1.2
13	Chronic lower respiratory diseases (J40-J47)	78	0.6	1.1
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	72	0.6	1.0
15	Aortic aneurysm and dissection (I71)	55	0.4	0.8
...	All other causes (Residual)	2,140	16.6	30.9
White, non-Hispanic, male, 40-44 years				
...	All causes	22,393	100.0	279.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,767	21.3	59.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,291	19.2	53.5
3	Malignant neoplasms (C00-C97)	3,150	14.1	39.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,385	10.7	29.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	920	4.1	11.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	784	3.5	9.8
7	Diabetes mellitus (E10-E14)	525	2.3	6.5
8	Cerebrovascular diseases (I60-I69)	463	2.1	5.8
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	394	1.8	4.9
10	Viral hepatitis (B15-B19)	200	0.9	2.5
11	Influenza and pneumonia (J10-J18)	197	0.9	2.5
12	Chronic lower respiratory diseases (J40-J47)	192	0.9	2.4
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	156	0.7	1.9
14	Septicemia (A40-A41)	144	0.6	1.8
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	119	0.5	1.5
...	All other causes (Residual)	3,706	16.5	46.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
	White, non-Hispanic, male, 45-49 years			
...	All causes	33,465	100.0	418.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,802	23.3	97.5
2	Malignant neoplasms (C00-C97)	6,763	20.2	84.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,953	14.8	61.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,382	7.1	29.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,612	4.8	20.1
6	Diabetes mellitus (E10-E14)	875	2.6	10.9
7	Cerebrovascular diseases (I60-I69)	726	2.2	9.1
8	Human immunodeficiency virus (HIV) disease (B20-B24)	665	2.0	8.3
9	Viral hepatitis (B15-B19)	482	1.4	6.0
10	Chronic lower respiratory diseases (J40-J47)	453	1.4	5.7
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	365	1.1	4.6
12	Influenza and pneumonia (J10-J18)	298	0.9	3.7
13	Septicemia (A40-A41)	267	0.8	3.3
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	230	0.7	2.9
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	152	0.5	1.9
...	All other causes (Residual)	5,440	16.3	68.0
	White, non-Hispanic, male, 50-54 years			
...	All causes	44,276	100.0	612.7
1	Malignant neoplasms (C00-C97)	11,793	26.6	163.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,527	26.0	159.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,982	9.0	55.1
4	Chronic liver disease and cirrhosis (K70,K73-K74)	2,153	4.9	29.8
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,083	4.7	28.8
6	Diabetes mellitus (E10-E14)	1,309	3.0	18.1
7	Cerebrovascular diseases (I60-I69)	1,115	2.5	15.4
8	Chronic lower respiratory diseases (J40-J47)	945	2.1	13.1
9	Viral hepatitis (B15-B19)	650	1.5	9.0
10	Human immunodeficiency virus (HIV) disease (B20-B24)	465	1.1	6.4
11	Septicemia (A40-A41)	457	1.0	6.3
12	Influenza and pneumonia (J10-J18)	443	1.0	6.1
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	356	0.8	4.9
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	284	0.6	3.9
15	Aortic aneurysm and dissection (I71)	220	0.5	3.0
...	All other causes (Residual)	6,494	14.7	89.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
White, non-Hispanic, male, 55-59 years				
...	All causes	54,247	100.0	863.8
1	Malignant neoplasms (C00-C97)	17,937	33.1	285.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,021	27.7	239.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,815	5.2	44.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	1,883	3.5	30.0
5	Diabetes mellitus (E10-E14)	1,815	3.3	28.9
6	Chronic lower respiratory diseases (J40-J47)	1,766	3.3	28.1
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,575	2.9	25.1
8	Cerebrovascular diseases (I60-I69)	1,435	2.6	22.8
9	Septicemia (A40-A41)	610	1.1	9.7
10	Influenza and pneumonia (J10-J18)	566	1.0	9.0
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	538	1.0	8.6
12	Viral hepatitis (B15-B19)	368	0.7	5.9
13	Human immunodeficiency virus (HIV) disease (B20-B24)	308	0.6	4.9
14	Aortic aneurysm and dissection (I71)	280	0.5	4.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	248	0.5	3.9
...	All other causes (Residual)	7,082	13.1	112.8
White, non-Hispanic, male, 60-64 years				
...	All causes	64,924	100.0	1,355.2
1	Malignant neoplasms (C00-C97)	23,531	36.2	491.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	18,057	27.8	376.9
3	Chronic lower respiratory diseases (J40-J47)	3,360	5.2	70.1
4	Diabetes mellitus (E10-E14)	2,313	3.6	48.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,140	3.3	44.7
6	Cerebrovascular diseases (I60-I69)	2,012	3.1	42.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,504	2.3	31.4
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,179	1.8	24.6
9	Influenza and pneumonia (J10-J18)	762	1.2	15.9
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	696	1.1	14.5
11	Septicemia (A40-A41)	689	1.1	14.4
12	Aortic aneurysm and dissection (I71)	450	0.7	9.4
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	345	0.5	7.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	337	0.5	7.0
15	Pneumonitis due to solids and liquids (J69)	182	0.3	3.8
...	All other causes (Residual)	7,367	11.3	153.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
White, non-Hispanic, male, 65-69 years				
...	All causes	77,307	100.0	2,082.3
1	Malignant neoplasms (C00-C97)	28,376	36.7	764.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,843	27.0	561.4
3	Chronic lower respiratory diseases (J40-J47)	5,157	6.7	138.9
4	Cerebrovascular diseases (I60-I69)	2,940	3.8	79.2
5	Diabetes mellitus (E10-E14)	2,659	3.4	71.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,944	2.5	52.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,279	1.7	34.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	996	1.3	26.8
9	Influenza and pneumonia (J10-J18)	992	1.3	26.7
10	Septicemia (A40-A41)	875	1.1	23.6
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	840	1.1	22.6
12	Aortic aneurysm and dissection (I71)	671	0.9	18.1
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	466	0.6	12.6
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	414	0.5	11.2
15	Alzheimer's disease (G30)	376	0.5	10.1
...	All other causes (Residual)	8,479	11.0	228.4
White, non-Hispanic, male, 70-74 years				
...	All causes	101,335	100.0	3,257.1
1	Malignant neoplasms (C00-C97)	34,234	33.8	1,100.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	27,574	27.2	886.3
3	Chronic lower respiratory diseases (J40-J47)	7,708	7.6	247.7
4	Cerebrovascular diseases (I60-I69)	4,565	4.5	146.7
5	Diabetes mellitus (E10-E14)	3,325	3.3	106.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,083	2.1	67.0
7	Influenza and pneumonia (J10-J18)	1,758	1.7	56.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,544	1.5	49.6
9	Septicemia (A40-A41)	1,338	1.3	43.0
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,184	1.2	38.1
11	Aortic aneurysm and dissection (I71)	1,024	1.0	32.9
12	Alzheimer's disease (G30)	959	0.9	30.8
13	Parkinson's disease (G20-G21)	943	0.9	30.3
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	861	0.8	27.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	662	0.7	21.3
...	All other causes (Residual)	11,573	11.4	372.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
White, non-Hispanic, male, 75-79 years				
...	All causes	135,156	100.0	5,178.6
1	Malignant neoplasms (C00-C97)	39,534	29.3	1,514.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	38,364	28.4	1,469.9
3	Chronic lower respiratory diseases (J40-J47)	10,600	7.8	406.1
4	Cerebrovascular diseases (I60-I69)	7,558	5.6	289.6
5	Diabetes mellitus (E10-E14)	4,059	3.0	155.5
6	Influenza and pneumonia (J10-J18)	2,970	2.2	113.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,761	2.0	105.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,509	1.9	96.1
9	Alzheimer's disease (G30)	2,439	1.8	93.5
10	Parkinson's disease (G20-G21)	1,984	1.5	76.0
11	Septicemia (A40-A41)	1,837	1.4	70.4
12	Aortic aneurysm and dissection (I71)	1,240	0.9	47.5
13	Pneumonitis due to solids and liquids (J69)	1,025	0.8	39.3
14	Chronic liver disease and cirrhosis (K70,K73-K74)	995	0.7	38.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	942	0.7	36.1
...	All other causes (Residual)	16,339	12.1	626.0
White, non-Hispanic, male, 80-84 years				
...	All causes	150,596	100.0	8,308.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	46,330	30.8	2,556.1
2	Malignant neoplasms (C00-C97)	34,644	23.0	1,911.4
3	Chronic lower respiratory diseases (J40-J47)	10,476	7.0	578.0
4	Cerebrovascular diseases (I60-I69)	9,995	6.6	551.4
5	Influenza and pneumonia (J10-J18)	4,579	3.0	252.6
6	Alzheimer's disease (G30)	4,506	3.0	248.6
7	Diabetes mellitus (E10-E14)	4,217	2.8	232.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,379	2.2	186.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,212	2.1	177.2
10	Parkinson's disease (G20-G21)	2,880	1.9	158.9
11	Septicemia (A40-A41)	2,077	1.4	114.6
12	Pneumonitis due to solids and liquids (J69)	1,639	1.1	90.4
13	Aortic aneurysm and dissection (I71)	1,339	0.9	73.9
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,203	0.8	66.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,169	0.8	64.5
...	All other causes (Residual)	18,951	12.6	1,045.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
White, non-Hispanic, male, 85-89 years				
...	All causes	118,869	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	39,954	33.6	---
2	Malignant neoplasms (C00-C97)	20,976	17.6	---
3	Cerebrovascular diseases (I60-I69)	8,699	7.3	---
4	Chronic lower respiratory diseases (J40-J47)	7,233	6.1	---
5	Alzheimer's disease (G30)	4,821	4.1	---
6	Influenza and pneumonia (J10-J18)	4,557	3.8	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,763	2.3	---
8	Diabetes mellitus (E10-E14)	2,751	2.3	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,704	2.3	---
10	Parkinson's disease (G20-G21)	2,086	1.8	---
11	Pneumonitis due to solids and liquids (J69)	1,708	1.4	---
12	Septicemia (A40-A41)	1,611	1.4	---
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,136	1.0	---
14	Atherosclerosis (I70)	918	0.8	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	868	0.7	---
...	All other causes (Residual)	16,084	13.5	---
White, non-Hispanic, male, 90-94 years				
...	All causes	61,968	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	22,721	36.7	---
2	Malignant neoplasms (C00-C97)	8,017	12.9	---
3	Cerebrovascular diseases (I60-I69)	4,714	7.6	---
4	Influenza and pneumonia (J10-J18)	3,065	4.9	---
5	Chronic lower respiratory diseases (J40-J47)	3,003	4.8	---
6	Alzheimer's disease (G30)	2,788	4.5	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,566	2.5	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,546	2.5	---
9	Diabetes mellitus (E10-E14)	1,147	1.9	---
10	Pneumonitis due to solids and liquids (J69)	1,086	1.8	---
11	Parkinson's disease (G20-G21)	822	1.3	---
12	Septicemia (A40-A41)	715	1.2	---
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	669	1.1	---
14	Atherosclerosis (I70)	592	1.0	---
15	Aortic aneurysm and dissection (I71)	399	0.6	---
...	All other causes (Residual)	9,118	14.7	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
White, non-Hispanic, male, 95-99 years				
...	All causes	17,161	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,611	38.5	---
2	Malignant neoplasms (C00-C97)	1,713	10.0	---
3	Cerebrovascular diseases (I60-I69)	1,250	7.3	---
4	Influenza and pneumonia (J10-J18)	1,111	6.5	---
5	Alzheimer's disease (G30)	780	4.5	---
6	Chronic lower respiratory diseases (J40-J47)	723	4.2	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	436	2.5	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	432	2.5	---
9	Pneumonitis due to solids and liquids (J69)	342	2.0	---
10	Diabetes mellitus (E10-E14)	230	1.3	---
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	228	1.3	---
12	Atherosclerosis (I70)	218	1.3	---
13	Septicemia (A40-A41)	217	1.3	---
14	Parkinson's disease (G20-G21)	140	0.8	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	98	0.6	---
...	All other causes (Residual)	2,632	15.3	---
White, non-Hispanic, male, 100 years and over				
...	All causes	2,359	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	964	40.9	---
2	Influenza and pneumonia (J10-J18)	206	8.7	---
3	Malignant neoplasms (C00-C97)	172	7.3	---
4	Cerebrovascular diseases (I60-I69)	152	6.4	---
5	Chronic lower respiratory diseases (J40-J47)	68	2.9	---
6	Pneumonitis due to solids and liquids (J69)	64	2.7	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	64	2.7	---
8	Alzheimer's disease (G30)	62	2.6	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	59	2.5	---
10	Atherosclerosis (I70)	41	1.7	---
11	Diabetes mellitus (E10-E14)	33	1.4	---
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	31	1.3	---
13	Septicemia (A40-A41)	25	1.1	---
14	Anemias (D50-D64)	13	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.4	---
...	All other causes (Residual)	396	16.8	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
White, non-Hispanic, female, all ages				
...	All causes	995,239	100.0	977.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	273,874	27.5	269.1
2	Malignant neoplasms (C00-C97)	219,142	22.0	215.3
3	Cerebrovascular diseases (I60-I69)	74,978	7.5	73.7
4	Chronic lower respiratory diseases (J40-J47)	57,761	5.8	56.7
5	Alzheimer's disease (G30)	42,164	4.2	41.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	32,001	3.2	31.4
7	Influenza and pneumonia (J10-J18)	27,551	2.8	27.1
8	Diabetes mellitus (E10-E14)	26,328	2.6	25.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16,261	1.6	16.0
10	Septicemia (A40-A41)	13,853	1.4	13.6
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	10,422	1.0	10.2
12	Chronic liver disease and cirrhosis (K70,K73-K74)	7,213	0.7	7.1
13	Pneumonitis due to solids and liquids (J69)	7,044	0.7	6.9
14	Parkinson's disease (G20-G21)	6,959	0.7	6.8
15	Atherosclerosis (I70)	6,534	0.7	6.4
...	All other causes (Residual)	173,154	17.4	170.1
White, non-Hispanic, female, 1-4 years				
...	All causes	1,093	100.0	24.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	380	34.8	8.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	125	11.4	2.8
3	Malignant neoplasms (C00-C97)	110	10.1	2.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	58	5.3	1.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	43	3.9	1.0
6	Influenza and pneumonia (J10-J18)	32	2.9	0.7
7	Septicemia (A40-A41)	21	1.9	0.5
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	1.6	*
9	Cerebrovascular diseases (I60-I69)	10	0.9	*
10	Certain conditions originating in the perinatal period (P00-P96)	7	0.6	*
11	Meningitis (G00,G03)	6	0.5	*
12	Anemias (D50-D64)	5	0.5	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.5	*
14	Chronic lower respiratory diseases (J40-J47)	4	0.4	*
15	Infections of kidney (N10-N12,N13.6,N15.1)	3	0.3	*
...	All other causes (Residual)	266	24.3	5.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
White, non-Hispanic, female, 5-9 years				
...	All causes	659	100.0	11.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	243	36.9	4.3
2	Malignant neoplasms (C00-C97)	134	20.3	2.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	52	7.9	0.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	23	3.5	0.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	21	3.2	0.4
6	Septicemia (A40-A41)	12	1.8	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	1.7	*
8	Cerebrovascular diseases (I60-I69)	9	1.4	*
9	Influenza and pneumonia (J10-J18)	8	1.2	*
10	Chronic lower respiratory diseases (J40-J47)	6	0.9	*
11	Certain conditions originating in the perinatal period (P00-P96)	5	0.8	*
12	Anemias (D50-D64)	4	0.6	*
12	Diabetes mellitus (E10-E14)	4	0.6	*
12	Meningitis (G00,G03)	4	0.6	*
15	Meningococcal infection (A39)	3	0.5	*
...	All other causes (Residual)	120	18.2	2.1
White, non-Hispanic, female, 10-14 years				
...	All causes	897	100.0	14.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	339	37.8	5.4
2	Malignant neoplasms (C00-C97)	114	12.7	1.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	61	6.8	1.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	54	6.0	0.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	37	4.1	0.6
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	26	2.9	0.4
7	Cerebrovascular diseases (I60-I69)	15	1.7	*
8	Influenza and pneumonia (J10-J18)	13	1.4	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	1.3	*
9	Chronic lower respiratory diseases (J40-J47)	12	1.3	*
11	Septicemia (A40-A41)	9	1.0	*
12	Diabetes mellitus (E10-E14)	8	0.9	*
13	Anemias (D50-D64)	7	0.8	*
14	Meningitis (G00,G03)	4	0.4	*
14	Pneumonitis due to solids and liquids (J69)	4	0.4	*
...	All other causes (Residual)	182	20.3	2.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
White, non-Hispanic, female, 15-19 years				
...	All causes	2,710	100.0	42.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,589	58.6	24.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	250	9.2	3.9
3	Malignant neoplasms (C00-C97)	188	6.9	2.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	118	4.4	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	70	2.6	1.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	69	2.5	1.1
7	Cerebrovascular diseases (I60-I69)	18	0.7	*
8	Influenza and pneumonia (J10-J18)	15	0.6	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	0.5	*
10	Septicemia (A40-A41)	13	0.5	*
10	Diabetes mellitus (E10-E14)	13	0.5	*
10	Pregnancy, childbirth and the puerperium (O00-099)	13	0.5	*
13	Chronic lower respiratory diseases (J40-J47)	12	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	8	0.3	*
...	All other causes (Residual)	311	11.5	4.9
White, non-Hispanic, female, 20-24 years				
...	All causes	2,957	100.0	46.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,373	46.4	21.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	260	8.8	4.1
3	Malignant neoplasms (C00-C97)	235	7.9	3.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	175	5.9	2.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	120	4.1	1.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	65	2.2	1.0
7	Pregnancy, childbirth and the puerperium (O00-099)	47	1.6	0.7
8	Cerebrovascular diseases (I60-I69)	36	1.2	0.6
9	Diabetes mellitus (E10-E14)	29	1.0	0.5
10	Influenza and pneumonia (J10-J18)	20	0.7	0.3
11	Septicemia (A40-A41)	18	0.6	*
11	Chronic lower respiratory diseases (J40-J47)	18	0.6	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	0.6	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	0.4	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	8	0.3	*
...	All other causes (Residual)	525	17.8	8.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
White, non-Hispanic, female, 25-29 years				
...	All causes	2,979	100.0	51.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,087	36.5	18.7
2	Malignant neoplasms (C00-C97)	352	11.8	6.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	277	9.3	4.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	194	6.5	3.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	153	5.1	2.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	53	1.8	0.9
7	Pregnancy, childbirth and the puerperium (O00-099)	52	1.7	0.9
8	Cerebrovascular diseases (I60-I69)	44	1.5	0.8
9	Diabetes mellitus (E10-E14)	41	1.4	0.7
10	Septicemia (A40-A41)	30	1.0	0.5
11	Chronic lower respiratory diseases (J40-J47)	28	0.9	0.5
12	Influenza and pneumonia (J10-J18)	20	0.7	0.3
13	Human immunodeficiency virus (HIV) disease (B20-B24)	19	0.6	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19	0.6	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	15	0.5	*
...	All other causes (Residual)	595	20.0	10.3
White, non-Hispanic, female, 30-34 years				
...	All causes	4,320	100.0	68.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,167	27.0	18.6
2	Malignant neoplasms (C00-C97)	774	17.9	12.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	406	9.4	6.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	357	8.3	5.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	151	3.5	2.4
6	Diabetes mellitus (E10-E14)	101	2.3	1.6
7	Cerebrovascular diseases (I60-I69)	99	2.3	1.6
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	74	1.7	1.2
9	Pregnancy, childbirth and the puerperium (O00-099)	70	1.6	1.1
10	Septicemia (A40-A41)	57	1.3	0.9
11	Human immunodeficiency virus (HIV) disease (B20-B24)	50	1.2	0.8
12	Influenza and pneumonia (J10-J18)	48	1.1	0.8
13	Chronic liver disease and cirrhosis (K70,K73-K74)	45	1.0	0.7
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	42	1.0	0.7
15	Chronic lower respiratory diseases (J40-J47)	29	0.7	0.5
...	All other causes (Residual)	850	19.7	13.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
White, non-Hispanic, female, 35-39 years				
...	All causes	7,173	100.0	104.4
1	Malignant neoplasms (C00-C97)	1,677	23.4	24.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,514	21.1	22.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	778	10.8	11.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	540	7.5	7.9
5	Cerebrovascular diseases (I60-I69)	198	2.8	2.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	183	2.6	2.7
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	156	2.2	2.3
8	Diabetes mellitus (E10-E14)	149	2.1	2.2
9	Human immunodeficiency virus (HIV) disease (B20-B24)	128	1.8	1.9
10	Chronic lower respiratory diseases (J40-J47)	97	1.4	1.4
11	Septicemia (A40-A41)	79	1.1	1.2
12	Influenza and pneumonia (J10-J18)	75	1.0	1.1
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	72	1.0	1.0
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	64	0.9	0.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	46	0.6	0.7
...	All other causes (Residual)	1,417	19.8	20.6
White, non-Hispanic, female, 40-44 years				
...	All causes	13,117	100.0	163.0
1	Malignant neoplasms (C00-C97)	3,918	29.9	48.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,099	16.0	26.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,553	11.8	19.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	809	6.2	10.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	440	3.4	5.5
6	Cerebrovascular diseases (I60-I69)	405	3.1	5.0
7	Diabetes mellitus (E10-E14)	294	2.2	3.7
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	217	1.7	2.7
9	Chronic lower respiratory diseases (J40-J47)	216	1.6	2.7
10	Influenza and pneumonia (J10-J18)	177	1.3	2.2
11	Septicemia (A40-A41)	152	1.2	1.9
12	Human immunodeficiency virus (HIV) disease (B20-B24)	142	1.1	1.8
13	Viral hepatitis (B15-B19)	105	0.8	1.3
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	100	0.8	1.2
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	94	0.7	1.2
...	All other causes (Residual)	2,396	18.3	29.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
White, non-Hispanic, female, 45-49 years				
...	All causes	19,798	100.0	244.9
1	Malignant neoplasms (C00-C97)	6,973	35.2	86.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,661	13.4	32.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,171	11.0	26.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	887	4.5	11.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	756	3.8	9.4
6	Cerebrovascular diseases (I60-I69)	674	3.4	8.3
7	Diabetes mellitus (E10-E14)	514	2.6	6.4
8	Chronic lower respiratory diseases (J40-J47)	439	2.2	5.4
9	Septicemia (A40-A41)	243	1.2	3.0
10	Influenza and pneumonia (J10-J18)	223	1.1	2.8
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	186	0.9	2.3
12	Viral hepatitis (B15-B19)	176	0.9	2.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	170	0.9	2.1
14	Human immunodeficiency virus (HIV) disease (B20-B24)	124	0.6	1.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	107	0.5	1.3
...	All other causes (Residual)	3,494	17.6	43.2
White, non-Hispanic, female, 50-54 years				
...	All causes	25,498	100.0	345.4
1	Malignant neoplasms (C00-C97)	10,487	41.1	142.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,868	15.2	52.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,638	6.4	22.2
4	Chronic lower respiratory diseases (J40-J47)	858	3.4	11.6
5	Cerebrovascular diseases (I60-I69)	840	3.3	11.4
6	Chronic liver disease and cirrhosis (K70,K73-K74)	809	3.2	11.0
7	Diabetes mellitus (E10-E14)	764	3.0	10.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	749	2.9	10.1
9	Septicemia (A40-A41)	392	1.5	5.3
10	Influenza and pneumonia (J10-J18)	278	1.1	3.8
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	249	1.0	3.4
12	Viral hepatitis (B15-B19)	209	0.8	2.8
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	146	0.6	2.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	117	0.5	1.6
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	103	0.4	1.4
...	All other causes (Residual)	3,991	15.7	54.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
	White, non-Hispanic, female, 55-59 years			
...	All causes	34,853	100.0	535.3
1	Malignant neoplasms (C00-C97)	15,426	44.3	236.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,679	16.3	87.2
3	Chronic lower respiratory diseases (J40-J47)	1,765	5.1	27.1
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,291	3.7	19.8
5	Diabetes mellitus (E10-E14)	1,274	3.7	19.6
6	Cerebrovascular diseases (I60-I69)	1,235	3.5	19.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	726	2.1	11.2
8	Septicemia (A40-A41)	524	1.5	8.0
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	513	1.5	7.9
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	408	1.2	6.3
11	Influenza and pneumonia (J10-J18)	407	1.2	6.3
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	192	0.6	2.9
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	172	0.5	2.6
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	147	0.4	2.3
15	Viral hepatitis (B15-B19)	132	0.4	2.0
...	All other causes (Residual)	4,962	14.2	76.2
	White, non-Hispanic, female, 60-64 years			
...	All causes	44,818	100.0	871.8
1	Malignant neoplasms (C00-C97)	19,506	43.5	379.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,177	18.2	159.1
3	Chronic lower respiratory diseases (J40-J47)	3,299	7.4	64.2
4	Cerebrovascular diseases (I60-I69)	1,639	3.7	31.9
5	Diabetes mellitus (E10-E14)	1,563	3.5	30.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,144	2.6	22.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	725	1.6	14.1
8	Septicemia (A40-A41)	690	1.5	13.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	596	1.3	11.6
10	Influenza and pneumonia (J10-J18)	560	1.2	10.9
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	317	0.7	6.2
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	231	0.5	4.5
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	216	0.5	4.2
14	Aortic aneurysm and dissection (I71)	178	0.4	3.5
15	Alzheimer's disease (G30)	177	0.4	3.4
...	All other causes (Residual)	5,800	12.9	112.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
	White, non-Hispanic, female, 65-69 years			
...	All causes	56,609	100.0	1,360.0
1	Malignant neoplasms (C00-C97)	22,931	40.5	550.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,128	19.7	267.3
3	Chronic lower respiratory diseases (J40-J47)	4,968	8.8	119.4
4	Cerebrovascular diseases (I60-I69)	2,521	4.5	60.6
5	Diabetes mellitus (E10-E14)	2,045	3.6	49.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,083	1.9	26.0
7	Septicemia (A40-A41)	854	1.5	20.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	842	1.5	20.2
9	Influenza and pneumonia (J10-J18)	743	1.3	17.8
10	Chronic liver disease and cirrhosis (K70,K73-K74)	715	1.3	17.2
11	Alzheimer's disease (G30)	475	0.8	11.4
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	342	0.6	8.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	295	0.5	7.1
14	Aortic aneurysm and dissection (I71)	288	0.5	6.9
15	Parkinson's disease (G20-G21)	187	0.3	4.5
...	All other causes (Residual)	7,192	12.7	172.8
	White, non-Hispanic, female, 70-74 years			
...	All causes	82,355	100.0	2,185.0
1	Malignant neoplasms (C00-C97)	28,607	34.7	759.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	18,107	22.0	480.4
3	Chronic lower respiratory diseases (J40-J47)	7,732	9.4	205.1
4	Cerebrovascular diseases (I60-I69)	4,537	5.5	120.4
5	Diabetes mellitus (E10-E14)	2,846	3.5	75.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,422	1.7	37.7
7	Influenza and pneumonia (J10-J18)	1,401	1.7	37.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,363	1.7	36.2
9	Alzheimer's disease (G30)	1,297	1.6	34.4
10	Septicemia (A40-A41)	1,261	1.5	33.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	825	1.0	21.9
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	630	0.8	16.7
13	Aortic aneurysm and dissection (I71)	550	0.7	14.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	526	0.6	14.0
15	Parkinson's disease (G20-G21)	488	0.6	12.9
...	All other causes (Residual)	10,763	13.1	285.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
	White, non-Hispanic, female, 75-79 years			
...	All causes	124,562	100.0	3,496.3
1	Malignant neoplasms (C00-C97)	34,288	27.5	962.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	30,807	24.7	864.7
3	Chronic lower respiratory diseases (J40-J47)	10,751	8.6	301.8
4	Cerebrovascular diseases (I60-I69)	8,887	7.1	249.4
5	Alzheimer's disease (G30)	4,019	3.2	112.8
6	Diabetes mellitus (E10-E14)	4,007	3.2	112.5
7	Influenza and pneumonia (J10-J18)	2,735	2.2	76.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,312	1.9	64.9
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,283	1.8	64.1
10	Septicemia (A40-A41)	1,940	1.6	54.5
11	Parkinson's disease (G20-G21)	1,203	1.0	33.8
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,144	0.9	32.1
13	Aortic aneurysm and dissection (I71)	915	0.7	25.7
14	Chronic liver disease and cirrhosis (K70,K73-K74)	836	0.7	23.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	802	0.6	22.5
...	All other causes (Residual)	17,633	14.2	494.9
	White, non-Hispanic, female, 80-84 years			
...	All causes	172,140	100.0	5,878.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	49,758	28.9	1,699.3
2	Malignant neoplasms (C00-C97)	33,995	19.7	1,161.0
3	Cerebrovascular diseases (I60-I69)	15,230	8.8	520.1
4	Chronic lower respiratory diseases (J40-J47)	11,786	6.8	402.5
5	Alzheimer's disease (G30)	8,474	4.9	289.4
6	Diabetes mellitus (E10-E14)	4,937	2.9	168.6
7	Influenza and pneumonia (J10-J18)	4,859	2.8	165.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,351	1.9	114.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,073	1.8	104.9
10	Septicemia (A40-A41)	2,572	1.5	87.8
11	Parkinson's disease (G20-G21)	1,932	1.1	66.0
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,879	1.1	64.2
13	Pneumonitis due to solids and liquids (J69)	1,395	0.8	47.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,168	0.7	39.9
15	Aortic aneurysm and dissection (I71)	1,079	0.6	36.8
...	All other causes (Residual)	26,652	15.5	910.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
	White, non-Hispanic, female, 85-89 years			
...	All causes	180,432	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	59,672	33.1	---
2	Malignant neoplasms (C00-C97)	23,743	13.2	---
3	Cerebrovascular diseases (I60-I69)	17,855	9.9	---
4	Alzheimer's disease (G30)	11,877	6.6	---
5	Chronic lower respiratory diseases (J40-J47)	9,204	5.1	---
6	Influenza and pneumonia (J10-J18)	6,231	3.5	---
7	Diabetes mellitus (E10-E14)	4,228	2.3	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,517	1.9	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,317	1.8	---
10	Septicemia (A40-A41)	2,485	1.4	---
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,370	1.3	---
12	Parkinson's disease (G20-G21)	1,774	1.0	---
13	Pneumonitis due to solids and liquids (J69)	1,731	1.0	---
14	Atherosclerosis (I70)	1,529	0.8	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,110	0.6	---
...	All other causes (Residual)	29,789	16.5	---
	White, non-Hispanic, female, 90-94 years			
...	All causes	139,520	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	51,540	36.9	---
2	Cerebrovascular diseases (I60-I69)	14,043	10.1	---
3	Malignant neoplasms (C00-C97)	11,860	8.5	---
4	Alzheimer's disease (G30)	10,359	7.4	---
5	Influenza and pneumonia (J10-J18)	5,932	4.3	---
6	Chronic lower respiratory diseases (J40-J47)	4,840	3.5	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,728	2.0	---
8	Diabetes mellitus (E10-E14)	2,567	1.8	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,443	1.8	---
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,112	1.5	---
11	Septicemia (A40-A41)	1,721	1.2	---
12	Atherosclerosis (I70)	1,585	1.1	---
13	Pneumonitis due to solids and liquids (J69)	1,415	1.0	---
14	Parkinson's disease (G20-G21)	961	0.7	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	679	0.5	---
...	All other causes (Residual)	24,735	17.7	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
White, non-Hispanic, female, 95-99 years				
...	All causes	59,933	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	23,637	39.4	---
2	Cerebrovascular diseases (I60-I69)	5,588	9.3	---
3	Alzheimer's disease (G30)	4,492	7.5	---
4	Malignant neoplasms (C00-C97)	3,329	5.6	---
5	Influenza and pneumonia (J10-J18)	2,923	4.9	---
6	Chronic lower respiratory diseases (J40-J47)	1,478	2.5	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,149	1.9	---
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,056	1.8	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,024	1.7	---
10	Atherosclerosis (I70)	917	1.5	---
11	Diabetes mellitus (E10-E14)	825	1.4	---
12	Septicemia (A40-A41)	639	1.1	---
12	Pneumonitis due to solids and liquids (J69)	639	1.1	---
14	Parkinson's disease (G20-G21)	263	0.4	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	229	0.4	---
...	All other causes (Residual)	11,745	19.6	---
White, non-Hispanic, female, 100 years and over				
...	All causes	13,140	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,570	42.4	---
2	Cerebrovascular diseases (I60-I69)	1,070	8.1	---
3	Alzheimer's disease (G30)	861	6.6	---
4	Influenza and pneumonia (J10-J18)	804	6.1	---
5	Malignant neoplasms (C00-C97)	473	3.6	---
6	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	273	2.1	---
7	Atherosclerosis (I70)	264	2.0	---
8	Chronic lower respiratory diseases (J40-J47)	212	1.6	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	205	1.6	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	187	1.4	---
11	Pneumonitis due to solids and liquids (J69)	139	1.1	---
12	Diabetes mellitus (E10-E14)	115	0.9	---
13	Septicemia (A40-A41)	96	0.7	---
14	Anemias (D50-D64)	50	0.4	---
15	Nutritional deficiencies (E40-E64)	44	0.3	---
...	All other causes (Residual)	2,777	21.1	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
Black, non-Hispanic, both sexes, all ages				
...	All causes	283,859	100.0	768.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	73,373	25.8	198.7
2	Malignant neoplasms (C00-C97)	61,842	21.8	167.5
3	Cerebrovascular diseases (I60-I69)	17,980	6.3	48.7
4	Diabetes mellitus (E10-E14)	12,685	4.5	34.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,446	4.4	33.7
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8,000	2.8	21.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,759	2.7	21.0
8	Chronic lower respiratory diseases (J40-J47)	7,331	2.6	19.9
9	Human immunodeficiency virus (HIV) disease (B20-B24)	7,163	2.5	19.4
10	Septicemia (A40-A41)	5,945	2.1	16.1
11	Influenza and pneumonia (J10-J18)	5,620	2.0	15.2
12	Certain conditions originating in the perinatal period (P00-P96)	4,740	1.7	12.8
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	4,709	1.7	12.8
14	Alzheimer's disease (G30)	4,196	1.5	11.4
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,506	0.9	6.8
...	All other causes (Residual)	47,564	16.8	128.8
Black, non-Hispanic, both sexes, 1-4 years				
...	All causes	1,128	100.0	45.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	317	28.1	12.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	154	13.7	6.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	127	11.3	5.1
4	Malignant neoplasms (C00-C97)	62	5.5	2.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	61	5.4	2.5
6	Influenza and pneumonia (J10-J18)	34	3.0	1.4
7	Certain conditions originating in the perinatal period (P00-P96)	23	2.0	0.9
8	Chronic lower respiratory diseases (J40-J47)	21	1.9	0.8
9	Septicemia (A40-A41)	19	1.7	*
9	Anemias (D50-D64)	19	1.7	*
11	Cerebrovascular diseases (I60-I69)	12	1.1	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.6	*
13	Acute bronchitis and bronchiolitis (J20-J21)	6	0.5	*
14	Meningitis (G00,G03)	5	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.4	*
...	All other causes (Residual)	257	22.8	10.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Black, non-Hispanic, both sexes, 5-9 years				
...	All causes	665	100.0	22.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	262	39.4	8.8
2	Malignant neoplasms (C00-C97)	75	11.3	2.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	45	6.8	1.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	43	6.5	1.4
5	Chronic lower respiratory diseases (J40-J47)	27	4.1	0.9
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	26	3.9	0.9
7	Anemias (D50-D64)	12	1.8	*
7	Influenza and pneumonia (J10-J18)	12	1.8	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	1.4	*
10	Cerebrovascular diseases (I60-I69)	8	1.2	*
11	Certain conditions originating in the perinatal period (P00-P96)	6	0.9	*
12	Septicemia (A40-A41)	5	0.8	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.6	*
14	Diabetes mellitus (E10-E14)	2	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.3	*
...	All other causes (Residual)	127	19.1	4.3
Black, non-Hispanic, both sexes, 10-14 years				
...	All causes	886	100.0	26.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	295	33.3	8.7
2	Malignant neoplasms (C00-C97)	90	10.2	2.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	87	9.8	2.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	52	5.9	1.5
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	48	5.4	1.4
6	Chronic lower respiratory diseases (J40-J47)	37	4.2	1.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	37	4.2	1.1
8	Diabetes mellitus (E10-E14)	14	1.6	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	13	1.5	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	1.4	*
11	Anemias (D50-D64)	11	1.2	*
12	Cerebrovascular diseases (I60-I69)	9	1.0	*
13	Influenza and pneumonia (J10-J18)	8	0.9	*
14	Septicemia (A40-A41)	7	0.8	*
15	Meningitis (G00,G03)	5	0.6	*
...	All other causes (Residual)	161	18.2	4.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Black, non-Hispanic, both sexes, 15-19 years				
...	All causes	2,633	100.0	83.7
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,030	39.1	32.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	698	26.5	22.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	150	5.7	4.8
4	Malignant neoplasms (C00-C97)	123	4.7	3.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	115	4.4	3.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	59	2.2	1.9
7	Chronic lower respiratory diseases (J40-J47)	42	1.6	1.3
8	Anemias (D50-D64)	29	1.1	0.9
9	Human immunodeficiency virus (HIV) disease (B20-B24)	24	0.9	0.8
10	Diabetes mellitus (E10-E14)	19	0.7	*
11	Cerebrovascular diseases (I60-I69)	18	0.7	*
12	Influenza and pneumonia (J10-J18)	14	0.5	*
13	Septicemia (A40-A41)	13	0.5	*
14	Pregnancy, childbirth and the puerperium (O00-099)	12	0.5	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.4	*
...	All other causes (Residual)	276	10.5	8.8
Black, non-Hispanic, both sexes, 20-24 years				
...	All causes	4,291	100.0	143.3
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,741	40.6	58.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	916	21.3	30.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	300	7.0	10.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	195	4.5	6.5
5	Malignant neoplasms (C00-C97)	184	4.3	6.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	117	2.7	3.9
7	Anemias (D50-D64)	61	1.4	2.0
7	Pregnancy, childbirth and the puerperium (O00-099)	61	1.4	2.0
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	52	1.2	1.7
10	Diabetes mellitus (E10-E14)	41	1.0	1.4
11	Chronic lower respiratory diseases (J40-J47)	37	0.9	1.2
12	Septicemia (A40-A41)	32	0.7	1.1
13	Cerebrovascular diseases (I60-I69)	30	0.7	1.0
14	Influenza and pneumonia (J10-J18)	26	0.6	0.9
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	24	0.6	0.8
...	All other causes (Residual)	474	11.0	15.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Black, non-Hispanic, both sexes, 25-29 years				
...	All causes	4,541	100.0	174.2
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,437	31.6	55.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	871	19.2	33.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	363	8.0	13.9
4	Human immunodeficiency virus (HIV) disease (B20-B24)	297	6.5	11.4
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	276	6.1	10.6
6	Malignant neoplasms (C00-C97)	241	5.3	9.2
7	Anemias (D50-D64)	60	1.3	2.3
7	Cerebrovascular diseases (I60-I69)	60	1.3	2.3
9	Pregnancy, childbirth and the puerperium (O00-099)	59	1.3	2.3
10	Diabetes mellitus (E10-E14)	56	1.2	2.1
11	Chronic lower respiratory diseases (J40-J47)	47	1.0	1.8
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	36	0.8	1.4
13	Septicemia (A40-A41)	34	0.7	1.3
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	30	0.7	1.2
15	Influenza and pneumonia (J10-J18)	22	0.5	0.8
...	All other causes (Residual)	652	14.4	25.0
Black, non-Hispanic, both sexes, 30-34 years				
...	All causes	5,227	100.0	198.6
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	968	18.5	36.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	833	15.9	31.7
3	Human immunodeficiency virus (HIV) disease (B20-B24)	639	12.2	24.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	608	11.6	23.1
5	Malignant neoplasms (C00-C97)	410	7.8	15.6
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	234	4.5	8.9
7	Diabetes mellitus (E10-E14)	124	2.4	4.7
8	Cerebrovascular diseases (I60-I69)	113	2.2	4.3
9	Septicemia (A40-A41)	68	1.3	2.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	67	1.3	2.5
11	Anemias (D50-D64)	66	1.3	2.5
12	Chronic lower respiratory diseases (J40-J47)	54	1.0	2.1
13	Pregnancy, childbirth and the puerperium (O00-099)	47	0.9	1.8
14	Influenza and pneumonia (J10-J18)	41	0.8	1.6
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	34	0.7	1.3
...	All other causes (Residual)	921	17.6	35.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
Black, non-Hispanic, both sexes, 35-39 years				
...	All causes	7,003	100.0	263.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,053	15.0	39.6
2	Human immunodeficiency virus (HIV) disease (B20-B24)	1,030	14.7	38.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	958	13.7	36.0
4	Malignant neoplasms (C00-C97)	781	11.2	29.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	694	9.9	26.1
6	Diabetes mellitus (E10-E14)	206	2.9	7.7
7	Cerebrovascular diseases (I60-I69)	195	2.8	7.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	184	2.6	6.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	114	1.6	4.3
10	Chronic lower respiratory diseases (J40-J47)	82	1.2	3.1
10	Chronic liver disease and cirrhosis (K70,K73-K74)	82	1.2	3.1
12	Influenza and pneumonia (J10-J18)	81	1.2	3.0
13	Septicemia (A40-A41)	73	1.0	2.7
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	72	1.0	2.7
15	Anemias (D50-D64)	67	1.0	2.5
...	All other causes (Residual)	1,331	19.0	50.0
Black, non-Hispanic, both sexes, 40-44 years				
...	All causes	11,234	100.0	397.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,149	19.1	76.0
2	Malignant neoplasms (C00-C97)	1,868	16.6	66.0
3	Human immunodeficiency virus (HIV) disease (B20-B24)	1,492	13.3	52.8
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,114	9.9	39.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	582	5.2	20.6
6	Cerebrovascular diseases (I60-I69)	503	4.5	17.8
7	Diabetes mellitus (E10-E14)	334	3.0	11.8
8	Chronic liver disease and cirrhosis (K70,K73-K74)	225	2.0	8.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	202	1.8	7.1
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	185	1.6	6.5
11	Septicemia (A40-A41)	182	1.6	6.4
12	Influenza and pneumonia (J10-J18)	142	1.3	5.0
13	Chronic lower respiratory diseases (J40-J47)	138	1.2	4.9
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	105	0.9	3.7
15	Anemias (D50-D64)	72	0.6	2.5
...	All other causes (Residual)	1,941	17.3	68.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
Black, non-Hispanic, both sexes, 45-49 years				
...	All causes	16,211	100.0	621.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,600	22.2	138.1
2	Malignant neoplasms (C00-C97)	3,466	21.4	132.9
3	Human immunodeficiency virus (HIV) disease (B20-B24)	1,383	8.5	53.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,305	8.1	50.1
5	Cerebrovascular diseases (I60-I69)	845	5.2	32.4
6	Diabetes mellitus (E10-E14)	515	3.2	19.8
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	438	2.7	16.8
8	Chronic liver disease and cirrhosis (K70,K73-K74)	382	2.4	14.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	337	2.1	12.9
10	Septicemia (A40-A41)	273	1.7	10.5
11	Chronic lower respiratory diseases (J40-J47)	258	1.6	9.9
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	225	1.4	8.6
13	Influenza and pneumonia (J10-J18)	189	1.2	7.2
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	167	1.0	6.4
15	Viral hepatitis (B15-B19)	136	0.8	5.2
...	All other causes (Residual)	2,692	16.6	103.2
Black, non-Hispanic, both sexes, 50-54 years				
...	All causes	20,036	100.0	928.8
1	Malignant neoplasms (C00-C97)	5,195	25.9	240.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,972	24.8	230.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,113	5.6	51.6
4	Cerebrovascular diseases (I60-I69)	1,060	5.3	49.1
5	Human immunodeficiency virus (HIV) disease (B20-B24)	1,040	5.2	48.2
6	Diabetes mellitus (E10-E14)	856	4.3	39.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	508	2.5	23.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	476	2.4	22.1
9	Septicemia (A40-A41)	377	1.9	17.5
10	Chronic lower respiratory diseases (J40-J47)	363	1.8	16.8
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	294	1.5	13.6
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	272	1.4	12.6
13	Influenza and pneumonia (J10-J18)	254	1.3	11.8
14	Viral hepatitis (B15-B19)	226	1.1	10.5
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	112	0.6	5.2
...	All other causes (Residual)	2,918	14.6	135.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Black, non-Hispanic, both sexes, 55-59 years				
...	All causes	21,096	100.0	1,296.4
1	Malignant neoplasms (C00-C97)	6,267	29.7	385.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,601	26.6	344.2
3	Cerebrovascular diseases (I60-I69)	1,108	5.3	68.1
4	Diabetes mellitus (E10-E14)	1,075	5.1	66.1
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	747	3.5	45.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	593	2.8	36.4
7	Human immunodeficiency virus (HIV) disease (B20-B24)	547	2.6	33.6
8	Chronic lower respiratory diseases (J40-J47)	465	2.2	28.6
9	Chronic liver disease and cirrhosis (K70,K73-K74)	428	2.0	26.3
10	Septicemia (A40-A41)	423	2.0	26.0
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	326	1.5	20.0
12	Influenza and pneumonia (J10-J18)	253	1.2	15.5
13	Viral hepatitis (B15-B19)	169	0.8	10.4
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	153	0.7	9.4
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	106	0.5	6.5
...	All other causes (Residual)	2,835	13.4	174.2
Black, non-Hispanic, both sexes, 60-64 years				
...	All causes	21,902	100.0	1,845.3
1	Malignant neoplasms (C00-C97)	6,882	31.4	579.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,940	27.1	500.5
3	Cerebrovascular diseases (I60-I69)	1,312	6.0	110.5
4	Diabetes mellitus (E10-E14)	1,229	5.6	103.5
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	690	3.2	58.1
6	Chronic lower respiratory diseases (J40-J47)	636	2.9	53.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	504	2.3	42.5
8	Septicemia (A40-A41)	483	2.2	40.7
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	359	1.6	30.2
10	Chronic liver disease and cirrhosis (K70,K73-K74)	300	1.4	25.3
11	Human immunodeficiency virus (HIV) disease (B20-B24)	294	1.3	24.8
12	Influenza and pneumonia (J10-J18)	284	1.3	23.9
13	Aortic aneurysm and dissection (I71)	120	0.5	10.1
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	94	0.4	7.9
15	Pneumonitis due to solids and liquids (J69)	89	0.4	7.5
...	All other causes (Residual)	2,686	12.3	226.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
Black, non-Hispanic, both sexes, 65-69 years				
...	All causes	24,048	100.0	2,554.0
1	Malignant neoplasms (C00-C97)	7,698	32.0	817.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,658	27.7	707.1
3	Cerebrovascular diseases (I60-I69)	1,489	6.2	158.1
4	Diabetes mellitus (E10-E14)	1,423	5.9	151.1
5	Chronic lower respiratory diseases (J40-J47)	782	3.3	83.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	761	3.2	80.8
7	Septicemia (A40-A41)	556	2.3	59.0
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	418	1.7	44.4
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	399	1.7	42.4
10	Influenza and pneumonia (J10-J18)	382	1.6	40.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	219	0.9	23.3
12	Human immunodeficiency virus (HIV) disease (B20-B24)	165	0.7	17.5
13	Aortic aneurysm and dissection (I71)	111	0.5	11.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	98	0.4	10.4
14	Alzheimer's disease (G30)	98	0.4	10.4
...	All other causes (Residual)	2,791	11.6	296.4
Black, non-Hispanic, both sexes, 70-74 years				
...	All causes	27,078	100.0	3,637.4
1	Malignant neoplasms (C00-C97)	8,014	29.6	1,076.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,714	28.5	1,036.2
3	Cerebrovascular diseases (I60-I69)	1,873	6.9	251.6
4	Diabetes mellitus (E10-E14)	1,655	6.1	222.3
5	Chronic lower respiratory diseases (J40-J47)	969	3.6	130.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	868	3.2	116.6
7	Septicemia (A40-A41)	647	2.4	86.9
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	491	1.8	66.0
9	Influenza and pneumonia (J10-J18)	468	1.7	62.9
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	400	1.5	53.7
11	Alzheimer's disease (G30)	229	0.8	30.8
12	Pneumonitis due to solids and liquids (J69)	154	0.6	20.7
13	Chronic liver disease and cirrhosis (K70,K73-K74)	152	0.6	20.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	122	0.5	16.4
15	Aortic aneurysm and dissection (I71)	113	0.4	15.2
...	All other causes (Residual)	3,209	11.9	431.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
Black, non-Hispanic, both sexes, 75-79 years				
...	All causes	30,776	100.0	5,336.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,959	29.1	1,553.5
2	Malignant neoplasms (C00-C97)	7,921	25.7	1,373.5
3	Cerebrovascular diseases (I60-I69)	2,452	8.0	425.2
4	Diabetes mellitus (E10-E14)	1,733	5.6	300.5
5	Chronic lower respiratory diseases (J40-J47)	1,202	3.9	208.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,028	3.3	178.3
7	Septicemia (A40-A41)	746	2.4	129.4
8	Influenza and pneumonia (J10-J18)	684	2.2	118.6
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	611	2.0	105.9
10	Alzheimer's disease (G30)	563	1.8	97.6
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	453	1.5	78.6
12	Pneumonitis due to solids and liquids (J69)	202	0.7	35.0
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	153	0.5	26.5
14	Aortic aneurysm and dissection (I71)	152	0.5	26.4
15	Parkinson's disease (G20-G21)	132	0.4	22.9
...	All other causes (Residual)	3,785	12.3	656.3
Black, non-Hispanic, both sexes, 80-84 years				
...	All causes	29,972	100.0	7,587.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,180	30.6	2,324.0
2	Malignant neoplasms (C00-C97)	6,323	21.1	1,600.7
3	Cerebrovascular diseases (I60-I69)	2,592	8.6	656.2
4	Diabetes mellitus (E10-E14)	1,536	5.1	388.9
5	Chronic lower respiratory diseases (J40-J47)	993	3.3	251.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	965	3.2	244.3
7	Alzheimer's disease (G30)	947	3.2	239.7
8	Influenza and pneumonia (J10-J18)	855	2.9	216.5
9	Septicemia (A40-A41)	788	2.6	199.5
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	603	2.0	152.7
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	404	1.3	102.3
12	Pneumonitis due to solids and liquids (J69)	261	0.9	66.1
13	Aortic aneurysm and dissection (I71)	163	0.5	41.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	158	0.5	40.0
14	Parkinson's disease (G20-G21)	158	0.5	40.0
...	All other causes (Residual)	4,046	13.5	1,024.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
Black, non-Hispanic, both sexes, 85-89 years				
...	All causes	22,966	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,400	32.2	---
2	Malignant neoplasms (C00-C97)	3,783	16.5	---
3	Cerebrovascular diseases (I60-I69)	2,101	9.1	---
4	Diabetes mellitus (E10-E14)	1,062	4.6	---
5	Alzheimer's disease (G30)	1,013	4.4	---
6	Influenza and pneumonia (J10-J18)	795	3.5	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	749	3.3	---
8	Chronic lower respiratory diseases (J40-J47)	690	3.0	---
9	Septicemia (A40-A41)	588	2.6	---
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	512	2.2	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	256	1.1	---
12	Pneumonitis due to solids and liquids (J69)	234	1.0	---
13	Atherosclerosis (I70)	136	0.6	---
14	Parkinson's disease (G20-G21)	126	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	110	0.5	---
...	All other causes (Residual)	3,411	14.9	---
Black, non-Hispanic, both sexes, 90-94 years				
...	All causes	15,615	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,443	34.9	---
2	Malignant neoplasms (C00-C97)	1,831	11.7	---
3	Cerebrovascular diseases (I60-I69)	1,452	9.3	---
4	Alzheimer's disease (G30)	837	5.4	---
5	Influenza and pneumonia (J10-J18)	598	3.8	---
6	Diabetes mellitus (E10-E14)	566	3.6	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	526	3.4	---
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	436	2.8	---
9	Septicemia (A40-A41)	363	2.3	---
10	Chronic lower respiratory diseases (J40-J47)	337	2.2	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	203	1.3	---
12	Pneumonitis due to solids and liquids (J69)	171	1.1	---
13	Atherosclerosis (I70)	124	0.8	---
14	Parkinson's disease (G20-G21)	63	0.4	---
15	Anemias (D50-D64)	59	0.4	---
...	All other causes (Residual)	2,606	16.7	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
Black, non-Hispanic, both sexes, 95-99 years				
...	All causes	6,267	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,372	37.8	---
2	Cerebrovascular diseases (I60-I69)	554	8.8	---
3	Malignant neoplasms (C00-C97)	505	8.1	---
4	Alzheimer's disease (G30)	361	5.8	---
5	Influenza and pneumonia (J10-J18)	284	4.5	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	190	3.0	---
7	Diabetes mellitus (E10-E14)	176	2.8	---
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	172	2.7	---
9	Septicemia (A40-A41)	133	2.1	---
10	Chronic lower respiratory diseases (J40-J47)	111	1.8	---
11	Pneumonitis due to solids and liquids (J69)	85	1.4	---
12	Atherosclerosis (I70)	68	1.1	---
13	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	59	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	30	0.5	---
15	Nutritional deficiencies (E40-E64)	23	0.4	---
...	All other causes (Residual)	1,144	18.3	---
Black, non-Hispanic, both sexes, 100 years and over				
...	All causes	2,029	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	803	39.6	---
2	Cerebrovascular diseases (I60-I69)	153	7.5	---
3	Malignant neoplasms (C00-C97)	115	5.7	---
4	Alzheimer's disease (G30)	106	5.2	---
5	Influenza and pneumonia (J10-J18)	101	5.0	---
6	Diabetes mellitus (E10-E14)	60	3.0	---
6	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	60	3.0	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	54	2.7	---
9	Septicemia (A40-A41)	44	2.2	---
10	Atherosclerosis (I70)	32	1.6	---
11	Chronic lower respiratory diseases (J40-J47)	29	1.4	---
12	Pneumonitis due to solids and liquids (J69)	22	1.1	---
13	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	21	1.0	---
14	Anemias (D50-D64)	11	0.5	---
14	Nutritional deficiencies (E40-E64)	11	0.5	---
...	All other causes (Residual)	407	20.1	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
Black, non-Hispanic, male, all ages				
...	All causes	144,022	100.0	818.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	35,749	24.8	203.2
2	Malignant neoplasms (C00-C97)	32,089	22.3	182.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,430	5.9	47.9
4	Cerebrovascular diseases (I60-I69)	7,586	5.3	43.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6,730	4.7	38.3
6	Diabetes mellitus (E10-E14)	5,498	3.8	31.3
7	Human immunodeficiency virus (HIV) disease (B20-B24)	4,693	3.3	26.7
8	Chronic lower respiratory diseases (J40-J47)	4,010	2.8	22.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,505	2.4	19.9
10	Influenza and pneumonia (J10-J18)	2,729	1.9	15.5
11	Septicemia (A40-A41)	2,660	1.8	15.1
12	Certain conditions originating in the perinatal period (P00-P96)	2,653	1.8	15.1
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,945	1.4	11.1
14	Chronic liver disease and cirrhosis (K70,K73-K74)	1,650	1.1	9.4
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,614	1.1	9.2
...	All other causes (Residual)	22,481	15.6	127.8
Black, non-Hispanic, male, 1-4 years				
...	All causes	623	100.0	49.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	182	29.2	14.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	79	12.7	6.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	74	11.9	5.9
4	Malignant neoplasms (C00-C97)	35	5.6	2.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	32	5.1	2.5
6	Chronic lower respiratory diseases (J40-J47)	14	2.2	*
7	Certain conditions originating in the perinatal period (P00-P96)	13	2.1	*
8	Anemias (D50-D64)	12	1.9	*
8	Influenza and pneumonia (J10-J18)	12	1.9	*
10	Septicemia (A40-A41)	9	1.4	*
11	Cerebrovascular diseases (I60-I69)	5	0.8	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.6	*
12	Acute bronchitis and bronchiolitis (J20-J21)	4	0.6	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.5	*
15	Meningitis (G00,G03)	2	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.3	*
...	All other causes (Residual)	141	22.6	11.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Black, non-Hispanic, male, 5-9 years				
...	All causes	361	100.0	23.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	163	45.2	10.8
2	Malignant neoplasms (C00-C97)	38	10.5	2.5
3	Chronic lower respiratory diseases (J40-J47)	21	5.8	1.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	21	5.8	1.4
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	19	5.3	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	10	2.8	*
7	Influenza and pneumonia (J10-J18)	8	2.2	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	1.1	*
8	Anemias (D50-D64)	4	1.1	*
8	Cerebrovascular diseases (I60-I69)	4	1.1	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	3	0.8	*
11	Certain conditions originating in the perinatal period (P00-P96)	3	0.8	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.6	*
14	Meningococcal infection (A39)	1	0.3	*
14	Septicemia (A40-A41)	1	0.3	*
14	Diabetes mellitus (E10-E14)	1	0.3	*
14	Nutritional deficiencies (E40-E64)	1	0.3	*
14	Diseases of appendix (K35-K38)	1	0.3	*
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1	0.3	*
...	All other causes (Residual)	55	15.2	3.6
Black, non-Hispanic, male, 10-14 years				
...	All causes	509	100.0	29.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	180	35.4	10.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	60	11.8	3.5
3	Malignant neoplasms (C00-C97)	44	8.6	2.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	29	5.7	1.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	25	4.9	1.5
6	Chronic lower respiratory diseases (J40-J47)	23	4.5	1.3
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	19	3.7	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	1.6	*
8	Anemias (D50-D64)	8	1.6	*
10	Diabetes mellitus (E10-E14)	7	1.4	*
11	Cerebrovascular diseases (I60-I69)	6	1.2	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	5	1.0	*
13	Influenza and pneumonia (J10-J18)	4	0.8	*
14	Septicemia (A40-A41)	3	0.6	*
15	Meningitis (G00,G03)	2	0.4	*
15	Certain conditions originating in the perinatal period (P00-P96)	2	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
...	All other causes (Residual)	82	16.1	4.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Black, non-Hispanic, male, 15-19 years				
...	All causes	1,986	100.0	124.6
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	907	45.7	56.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	509	25.6	31.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	120	6.0	7.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	77	3.9	4.8
5	Malignant neoplasms (C00-C97)	68	3.4	4.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	33	1.7	2.1
7	Chronic lower respiratory diseases (J40-J47)	27	1.4	1.7
8	Anemias (D50-D64)	20	1.0	1.3
9	Diabetes mellitus (E10-E14)	14	0.7	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	13	0.7	*
10	Cerebrovascular diseases (I60-I69)	13	0.7	*
12	Influenza and pneumonia (J10-J18)	8	0.4	*
13	Septicemia (A40-A41)	6	0.3	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.2	*
...	All other causes (Residual)	163	8.2	10.2
Black, non-Hispanic, male, 20-24 years				
...	All causes	3,253	100.0	216.9
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,588	48.8	105.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	682	21.0	45.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	262	8.1	17.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	125	3.8	8.3
5	Malignant neoplasms (C00-C97)	106	3.3	7.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	60	1.8	4.0
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	40	1.2	2.7
8	Anemias (D50-D64)	30	0.9	2.0
9	Chronic lower respiratory diseases (J40-J47)	21	0.6	1.4
10	Cerebrovascular diseases (I60-I69)	17	0.5	*
10	Legal intervention (Y35,Y89.0)	17	0.5	*
12	Septicemia (A40-A41)	16	0.5	*
12	Diabetes mellitus (E10-E14)	16	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	0.5	*
15	Influenza and pneumonia (J10-J18)	11	0.3	*
...	All other causes (Residual)	247	7.6	16.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Black, non-Hispanic, male, 25-29 years				
...	All causes	3,209	100.0	255.3
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,294	40.3	102.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	672	20.9	53.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	236	7.4	18.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	219	6.8	17.4
5	Human immunodeficiency virus (HIV) disease (B20-B24)	158	4.9	12.6
6	Malignant neoplasms (C00-C97)	102	3.2	8.1
7	Diabetes mellitus (E10-E14)	33	1.0	2.6
8	Chronic lower respiratory diseases (J40-J47)	32	1.0	2.5
9	Anemias (D50-D64)	31	1.0	2.5
10	Cerebrovascular diseases (I60-I69)	23	0.7	1.8
11	Legal intervention (Y35,Y89.0)	21	0.7	1.7
12	Influenza and pneumonia (J10-J18)	16	0.5	*
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	16	0.5	*
14	Septicemia (A40-A41)	15	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	0.4	*
...	All other causes (Residual)	327	10.2	26.0
Black, non-Hispanic, male, 30-34 years				
...	All causes	3,355	100.0	269.0
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	836	24.9	67.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	631	18.8	50.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	369	11.0	29.6
4	Human immunodeficiency virus (HIV) disease (B20-B24)	343	10.2	27.5
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	184	5.5	14.8
6	Malignant neoplasms (C00-C97)	152	4.5	12.2
7	Diabetes mellitus (E10-E14)	68	2.0	5.5
8	Cerebrovascular diseases (I60-I69)	54	1.6	4.3
9	Anemias (D50-D64)	39	1.2	3.1
10	Septicemia (A40-A41)	37	1.1	3.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	37	1.1	3.0
12	Chronic lower respiratory diseases (J40-J47)	34	1.0	2.7
13	Influenza and pneumonia (J10-J18)	25	0.7	2.0
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	20	0.6	1.6
15	Aortic aneurysm and dissection (I71)	17	0.5	*
...	All other causes (Residual)	509	15.2	40.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Black, non-Hispanic, male, 35-39 years				
...	All causes	4,199	100.0	334.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	666	15.9	53.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	663	15.8	52.9
3	Human immunodeficiency virus (HIV) disease (B20-B24)	651	15.5	51.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	560	13.3	44.6
5	Malignant neoplasms (C00-C97)	258	6.1	20.6
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	145	3.5	11.6
7	Diabetes mellitus (E10-E14)	124	3.0	9.9
8	Cerebrovascular diseases (I60-I69)	112	2.7	8.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	68	1.6	5.4
10	Chronic liver disease and cirrhosis (K70,K73-K74)	53	1.3	4.2
11	Influenza and pneumonia (J10-J18)	48	1.1	3.8
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	45	1.1	3.6
13	Chronic lower respiratory diseases (J40-J47)	41	1.0	3.3
14	Anemias (D50-D64)	40	1.0	3.2
15	Septicemia (A40-A41)	32	0.8	2.6
...	All other causes (Residual)	693	16.5	55.2
Black, non-Hispanic, male, 40-44 years				
...	All causes	6,353	100.0	480.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,273	20.0	96.2
2	Human immunodeficiency virus (HIV) disease (B20-B24)	950	15.0	71.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	768	12.1	58.0
4	Malignant neoplasms (C00-C97)	767	12.1	58.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	445	7.0	33.6
6	Cerebrovascular diseases (I60-I69)	232	3.7	17.5
7	Diabetes mellitus (E10-E14)	190	3.0	14.4
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	148	2.3	11.2
9	Chronic liver disease and cirrhosis (K70,K73-K74)	144	2.3	10.9
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	98	1.5	7.4
11	Septicemia (A40-A41)	93	1.5	7.0
12	Influenza and pneumonia (J10-J18)	78	1.2	5.9
13	Chronic lower respiratory diseases (J40-J47)	66	1.0	5.0
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	58	0.9	4.4
15	Aortic aneurysm and dissection (I71)	39	0.6	2.9
...	All other causes (Residual)	1,004	15.8	75.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
Black, non-Hispanic, male, 45-49 years				
...	All causes	9,495	100.0	782.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,269	23.9	187.0
2	Malignant neoplasms (C00-C97)	1,669	17.6	137.6
3	Human immunodeficiency virus (HIV) disease (B20-B24)	950	10.0	78.3
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	890	9.4	73.4
5	Cerebrovascular diseases (I60-I69)	442	4.7	36.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	335	3.5	27.6
7	Diabetes mellitus (E10-E14)	302	3.2	24.9
8	Chronic liver disease and cirrhosis (K70,K73-K74)	262	2.8	21.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	192	2.0	15.8
10	Septicemia (A40-A41)	146	1.5	12.0
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	128	1.3	10.6
12	Chronic lower respiratory diseases (J40-J47)	124	1.3	10.2
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	123	1.3	10.1
14	Influenza and pneumonia (J10-J18)	107	1.1	8.8
15	Viral hepatitis (B15-B19)	100	1.1	8.2
...	All other causes (Residual)	1,456	15.3	120.0
Black, non-Hispanic, male, 50-54 years				
...	All causes	11,954	100.0	1,211.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,124	26.1	316.5
2	Malignant neoplasms (C00-C97)	2,770	23.2	280.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	824	6.9	83.5
4	Human immunodeficiency virus (HIV) disease (B20-B24)	729	6.1	73.9
5	Cerebrovascular diseases (I60-I69)	570	4.8	57.8
6	Diabetes mellitus (E10-E14)	446	3.7	45.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	346	2.9	35.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	282	2.4	28.6
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	239	2.0	24.2
10	Chronic lower respiratory diseases (J40-J47)	193	1.6	19.6
11	Septicemia (A40-A41)	188	1.6	19.0
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	168	1.4	17.0
13	Viral hepatitis (B15-B19)	162	1.4	16.4
14	Influenza and pneumonia (J10-J18)	159	1.3	16.1
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	86	0.7	8.7
...	All other causes (Residual)	1,668	14.0	169.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
Black, non-Hispanic, male, 55-59 years				
...	All causes	12,498	100.0	1,702.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,529	28.2	480.8
2	Malignant neoplasms (C00-C97)	3,422	27.4	466.2
3	Cerebrovascular diseases (I60-I69)	639	5.1	87.1
4	Diabetes mellitus (E10-E14)	561	4.5	76.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	547	4.4	74.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	417	3.3	56.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	307	2.5	41.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	298	2.4	40.6
9	Chronic lower respiratory diseases (J40-J47)	283	2.3	38.6
10	Septicemia (A40-A41)	229	1.8	31.2
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	168	1.3	22.9
12	Influenza and pneumonia (J10-J18)	140	1.1	19.1
13	Viral hepatitis (B15-B19)	125	1.0	17.0
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	122	1.0	16.6
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	93	0.7	12.7
...	All other causes (Residual)	1,618	12.9	220.4
Black, non-Hispanic, male, 60-64 years				
...	All causes	12,552	100.0	2,416.8
1	Malignant neoplasms (C00-C97)	3,854	30.7	742.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,620	28.8	697.0
3	Cerebrovascular diseases (I60-I69)	742	5.9	142.9
4	Diabetes mellitus (E10-E14)	610	4.9	117.4
5	Chronic lower respiratory diseases (J40-J47)	361	2.9	69.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	354	2.8	68.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	342	2.7	65.8
8	Septicemia (A40-A41)	236	1.9	45.4
9	Human immunodeficiency virus (HIV) disease (B20-B24)	220	1.8	42.4
10	Chronic liver disease and cirrhosis (K70,K73-K74)	194	1.5	37.4
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	193	1.5	37.2
12	Influenza and pneumonia (J10-J18)	171	1.4	32.9
13	Aortic aneurysm and dissection (I71)	83	0.7	16.0
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	68	0.5	13.1
15	Pneumonitis due to solids and liquids (J69)	59	0.5	11.4
...	All other causes (Residual)	1,445	11.5	278.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
Black, non-Hispanic, male, 65-69 years				
...	All causes	13,228	100.0	3,298.9
1	Malignant neoplasms (C00-C97)	4,230	32.0	1,054.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,792	28.7	945.7
3	Cerebrovascular diseases (I60-I69)	770	5.8	192.0
4	Diabetes mellitus (E10-E14)	692	5.2	172.6
5	Chronic lower respiratory diseases (J40-J47)	444	3.4	110.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	379	2.9	94.5
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	273	2.1	68.1
8	Septicemia (A40-A41)	266	2.0	66.3
9	Influenza and pneumonia (J10-J18)	249	1.9	62.1
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	201	1.5	50.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	145	1.1	36.2
12	Human immunodeficiency virus (HIV) disease (B20-B24)	118	0.9	29.4
13	Aortic aneurysm and dissection (I71)	74	0.6	18.5
14	Pneumonitis due to solids and liquids (J69)	56	0.4	14.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	51	0.4	12.7
...	All other causes (Residual)	1,488	11.2	371.1
Black, non-Hispanic, male, 70-74 years				
...	All causes	14,000	100.0	4,655.2
1	Malignant neoplasms (C00-C97)	4,412	31.5	1,467.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,006	28.6	1,332.0
3	Cerebrovascular diseases (I60-I69)	915	6.5	304.2
4	Diabetes mellitus (E10-E14)	725	5.2	241.1
5	Chronic lower respiratory diseases (J40-J47)	556	4.0	184.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	392	2.8	130.3
7	Septicemia (A40-A41)	316	2.3	105.1
8	Influenza and pneumonia (J10-J18)	265	1.9	88.1
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	247	1.8	82.1
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	203	1.5	67.5
11	Alzheimer's disease (G30)	98	0.7	32.6
12	Chronic liver disease and cirrhosis (K70,K73-K74)	97	0.7	32.3
13	Pneumonitis due to solids and liquids (J69)	95	0.7	31.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	59	0.4	19.6
15	Aortic aneurysm and dissection (I71)	56	0.4	18.6
...	All other causes (Residual)	1,558	11.1	518.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
Black, non-Hispanic, male, 75-79 years				
...	All causes	14,666	100.0	6,839.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,211	28.7	1,963.8
2	Malignant neoplasms (C00-C97)	4,186	28.5	1,952.1
3	Cerebrovascular diseases (I60-I69)	1,026	7.0	478.5
4	Diabetes mellitus (E10-E14)	671	4.6	312.9
5	Chronic lower respiratory diseases (J40-J47)	658	4.5	306.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	435	3.0	202.9
7	Influenza and pneumonia (J10-J18)	369	2.5	172.1
8	Septicemia (A40-A41)	336	2.3	156.7
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	269	1.8	125.4
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	249	1.7	116.1
11	Alzheimer's disease (G30)	196	1.3	91.4
12	Pneumonitis due to solids and liquids (J69)	112	0.8	52.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	75	0.5	35.0
13	Parkinson's disease (G20-G21)	75	0.5	35.0
15	Aortic aneurysm and dissection (I71)	64	0.4	29.8
...	All other causes (Residual)	1,734	11.8	808.6
Black, non-Hispanic, male, 80-84 years				
...	All causes	12,728	100.0	9,346.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,720	29.2	2,731.7
2	Malignant neoplasms (C00-C97)	3,236	25.4	2,376.3
3	Cerebrovascular diseases (I60-I69)	933	7.3	685.1
4	Chronic lower respiratory diseases (J40-J47)	542	4.3	398.0
5	Diabetes mellitus (E10-E14)	536	4.2	393.6
6	Influenza and pneumonia (J10-J18)	406	3.2	298.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	392	3.1	287.9
8	Septicemia (A40-A41)	339	2.7	248.9
9	Alzheimer's disease (G30)	290	2.3	213.0
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	219	1.7	160.8
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	218	1.7	160.1
12	Pneumonitis due to solids and liquids (J69)	129	1.0	94.7
13	Parkinson's disease (G20-G21)	81	0.6	59.5
14	Aortic aneurysm and dissection (I71)	67	0.5	49.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	65	0.5	47.7
...	All other causes (Residual)	1,555	12.2	1,141.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
Black, non-Hispanic, male, 85-89 years				
...	All causes	8,181	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,516	30.8	---
2	Malignant neoplasms (C00-C97)	1,719	21.0	---
3	Cerebrovascular diseases (I60-I69)	592	7.2	---
4	Chronic lower respiratory diseases (J40-J47)	358	4.4	---
5	Diabetes mellitus (E10-E14)	318	3.9	---
6	Influenza and pneumonia (J10-J18)	313	3.8	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	285	3.5	---
8	Alzheimer's disease (G30)	268	3.3	---
9	Septicemia (A40-A41)	198	2.4	---
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	143	1.7	---
11	Pneumonitis due to solids and liquids (J69)	107	1.3	---
12	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	103	1.3	---
13	Parkinson's disease (G20-G21)	69	0.8	---
14	Aortic aneurysm and dissection (I71)	44	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	39	0.5	---
...	All other causes (Residual)	1,109	13.6	---
Black, non-Hispanic, male, 90-94 years				
...	All causes	4,480	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,483	33.1	---
2	Malignant neoplasms (C00-C97)	792	17.7	---
3	Cerebrovascular diseases (I60-I69)	345	7.7	---
4	Influenza and pneumonia (J10-J18)	184	4.1	---
5	Alzheimer's disease (G30)	170	3.8	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	157	3.5	---
7	Chronic lower respiratory diseases (J40-J47)	151	3.4	---
8	Diabetes mellitus (E10-E14)	148	3.3	---
9	Septicemia (A40-A41)	104	2.3	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	90	2.0	---
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	77	1.7	---
12	Pneumonitis due to solids and liquids (J69)	62	1.4	---
13	Atherosclerosis (I70)	29	0.6	---
14	Parkinson's disease (G20-G21)	24	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	0.4	---
...	All other causes (Residual)	647	14.4	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
Black, non-Hispanic, male, 95-99 years				
...	All causes	1,395	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	480	34.4	---
2	Malignant neoplasms (C00-C97)	191	13.7	---
3	Cerebrovascular diseases (I60-I69)	99	7.1	---
4	Influenza and pneumonia (J10-J18)	71	5.1	---
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	58	4.2	---
6	Alzheimer's disease (G30)	48	3.4	---
7	Chronic lower respiratory diseases (J40-J47)	42	3.0	---
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	39	2.8	---
9	Pneumonitis due to solids and liquids (J69)	32	2.3	---
10	Septicemia (A40-A41)	30	2.2	---
11	Diabetes mellitus (E10-E14)	28	2.0	---
12	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17	1.2	---
13	Parkinson's disease (G20-G21)	10	0.7	---
13	Atherosclerosis (I70)	10	0.7	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.4	---
15	Nutritional deficiencies (E40-E64)	5	0.4	---
...	All other causes (Residual)	230	16.5	---
Black, non-Hispanic, male, 100 years and over				
...	All causes	359	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	142	39.6	---
2	Malignant neoplasms (C00-C97)	35	9.7	---
3	Influenza and pneumonia (J10-J18)	30	8.4	---
4	Cerebrovascular diseases (I60-I69)	24	6.7	---
5	Alzheimer's disease (G30)	13	3.6	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	3.3	---
7	Septicemia (A40-A41)	9	2.5	---
7	Chronic lower respiratory diseases (J40-J47)	9	2.5	---
9	Diabetes mellitus (E10-E14)	6	1.7	---
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	5	1.4	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5	1.4	---
12	Atherosclerosis (I70)	3	0.8	---
12	Pneumonitis due to solids and liquids (J69)	3	0.8	---
12	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.8	---
15	Parkinson's disease (G20-G21)	2	0.6	---
...	All other causes (Residual)	58	16.2	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
Black, non-Hispanic, female, all ages				
...	All causes	139,837	100.0	723.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	37,624	26.9	194.6
2	Malignant neoplasms (C00-C97)	29,753	21.3	153.9
3	Cerebrovascular diseases (I60-I69)	10,394	7.4	53.8
4	Diabetes mellitus (E10-E14)	7,187	5.1	37.2
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,254	3.0	22.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,016	2.9	20.8
7	Chronic lower respiratory diseases (J40-J47)	3,321	2.4	17.2
8	Septicemia (A40-A41)	3,285	2.3	17.0
9	Alzheimer's disease (G30)	3,058	2.2	15.8
10	Influenza and pneumonia (J10-J18)	2,891	2.1	15.0
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,764	2.0	14.3
12	Human immunodeficiency virus (HIV) disease (B20-B24)	2,470	1.8	12.8
13	Certain conditions originating in the perinatal period (P00-P96)	2,087	1.5	10.8
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,270	0.9	6.6
15	Chronic liver disease and cirrhosis (K70,K73-K74)	856	0.6	4.4
...	All other causes (Residual)	24,607	17.6	127.3
Black, non-Hispanic, female, 1-4 years				
...	All causes	505	100.0	41.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	135	26.7	11.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	75	14.9	6.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	53	10.5	4.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	29	5.7	2.4
5	Malignant neoplasms (C00-C97)	27	5.3	2.2
6	Influenza and pneumonia (J10-J18)	22	4.4	1.8
7	Septicemia (A40-A41)	10	2.0	*
7	Certain conditions originating in the perinatal period (P00-P96)	10	2.0	*
9	Anemias (D50-D64)	7	1.4	*
9	Cerebrovascular diseases (I60-I69)	7	1.4	*
9	Chronic lower respiratory diseases (J40-J47)	7	1.4	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.6	*
12	Meningitis (G00,G03)	3	0.6	*
14	Meningococcal infection (A39)	2	0.4	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.4	*
14	Acute bronchitis and bronchiolitis (J20-J21)	2	0.4	*
...	All other causes (Residual)	111	22.0	9.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
Black, non-Hispanic, female, 5-9 years				
...	All causes	304	100.0	20.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	99	32.6	6.8
2	Malignant neoplasms (C00-C97)	37	12.2	2.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	26	8.6	1.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	22	7.2	1.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	16	5.3	*
6	Anemias (D50-D64)	8	2.6	*
7	Chronic lower respiratory diseases (J40-J47)	6	2.0	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	1.6	*
9	Septicemia (A40-A41)	4	1.3	*
9	Cerebrovascular diseases (I60-I69)	4	1.3	*
9	Influenza and pneumonia (J10-J18)	4	1.3	*
12	Certain conditions originating in the perinatal period (P00-P96)	3	1.0	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.3	*
13	Diabetes mellitus (E10-E14)	1	0.3	*
13	Meningitis (G00,G03)	1	0.3	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.3	*
...	All other causes (Residual)	66	21.7	4.5
Black, non-Hispanic, female, 10-14 years				
...	All causes	377	100.0	22.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	115	30.5	6.9
2	Malignant neoplasms (C00-C97)	46	12.2	2.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	27	7.2	1.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	27	7.2	1.6
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	19	5.0	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	18	4.8	*
7	Chronic lower respiratory diseases (J40-J47)	14	3.7	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	8	2.1	*
9	Diabetes mellitus (E10-E14)	7	1.9	*
10	Septicemia (A40-A41)	4	1.1	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	1.1	*
10	Influenza and pneumonia (J10-J18)	4	1.1	*
13	Anemias (D50-D64)	3	0.8	*
13	Meningitis (G00,G03)	3	0.8	*
13	Cerebrovascular diseases (I60-I69)	3	0.8	*
...	All other causes (Residual)	75	19.9	4.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Black, non-Hispanic, female, 15-19 years				
...	All causes	647	100.0	41.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	189	29.2	12.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	123	19.0	7.9
3	Malignant neoplasms (C00-C97)	55	8.5	3.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	38	5.9	2.5
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	30	4.6	1.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	26	4.0	1.7
7	Chronic lower respiratory diseases (J40-J47)	15	2.3	*
8	Pregnancy, childbirth and the puerperium (O00-099)	12	1.9	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	11	1.7	*
10	Anemias (D50-D64)	9	1.4	*
11	Septicemia (A40-A41)	7	1.1	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	1.1	*
13	Influenza and pneumonia (J10-J18)	6	0.9	*
14	Diabetes mellitus (E10-E14)	5	0.8	*
14	Cerebrovascular diseases (I60-I69)	5	0.8	*
...	All other causes (Residual)	109	16.8	7.0
Black, non-Hispanic, female, 20-24 years				
...	All causes	1,038	100.0	69.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	234	22.5	15.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	153	14.7	10.2
3	Malignant neoplasms (C00-C97)	78	7.5	5.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	70	6.7	4.7
5	Pregnancy, childbirth and the puerperium (O00-099)	61	5.9	4.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	57	5.5	3.8
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	38	3.7	2.5
8	Anemias (D50-D64)	31	3.0	2.1
9	Diabetes mellitus (E10-E14)	25	2.4	1.7
10	Septicemia (A40-A41)	16	1.5	*
10	Chronic lower respiratory diseases (J40-J47)	16	1.5	*
12	Influenza and pneumonia (J10-J18)	15	1.4	*
13	Cerebrovascular diseases (I60-I69)	13	1.3	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	12	1.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.9	*
...	All other causes (Residual)	210	20.2	14.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Black, non-Hispanic, female, 25-29 years				
...	All causes	1,332	100.0	98.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	199	14.9	14.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	144	10.8	10.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	143	10.7	10.6
4	Human immunodeficiency virus (HIV) disease (B20-B24)	139	10.4	10.3
4	Malignant neoplasms (C00-C97)	139	10.4	10.3
6	Pregnancy, childbirth and the puerperium (O00-099)	59	4.4	4.4
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	40	3.0	3.0
8	Cerebrovascular diseases (I60-I69)	37	2.8	2.7
9	Anemias (D50-D64)	29	2.2	2.1
10	Diabetes mellitus (E10-E14)	23	1.7	1.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22	1.7	1.6
12	Septicemia (A40-A41)	19	1.4	*
13	Chronic lower respiratory diseases (J40-J47)	15	1.1	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	14	1.1	*
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	8	0.6	*
...	All other causes (Residual)	302	22.7	22.4
Black, non-Hispanic, female, 30-34 years				
...	All causes	1,872	100.0	135.2
1	Human immunodeficiency virus (HIV) disease (B20-B24)	296	15.8	21.4
2	Malignant neoplasms (C00-C97)	258	13.8	18.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	239	12.8	17.3
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	202	10.8	14.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	132	7.1	9.5
6	Cerebrovascular diseases (I60-I69)	59	3.2	4.3
7	Diabetes mellitus (E10-E14)	56	3.0	4.0
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	50	2.7	3.6
9	Pregnancy, childbirth and the puerperium (O00-099)	47	2.5	3.4
10	Septicemia (A40-A41)	31	1.7	2.2
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	30	1.6	2.2
12	Anemias (D50-D64)	27	1.4	2.0
13	Chronic lower respiratory diseases (J40-J47)	20	1.1	1.4
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	18	1.0	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	0.9	*
15	Influenza and pneumonia (J10-J18)	16	0.9	*
...	All other causes (Residual)	375	20.0	27.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Black, non-Hispanic, female, 35-39 years				
...	All causes	2,804	100.0	199.5
1	Malignant neoplasms (C00-C97)	523	18.7	37.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	390	13.9	27.8
3	Human immunodeficiency virus (HIV) disease (B20-B24)	379	13.5	27.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	292	10.4	20.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	134	4.8	9.5
6	Cerebrovascular diseases (I60-I69)	83	3.0	5.9
7	Diabetes mellitus (E10-E14)	82	2.9	5.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	46	1.6	3.3
9	Septicemia (A40-A41)	41	1.5	2.9
9	Chronic lower respiratory diseases (J40-J47)	41	1.5	2.9
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	39	1.4	2.8
12	Influenza and pneumonia (J10-J18)	33	1.2	2.3
13	Chronic liver disease and cirrhosis (K70,K73-K74)	29	1.0	2.1
13	Pregnancy, childbirth and the puerperium (O00-099)	29	1.0	2.1
15	Anemias (D50-D64)	27	1.0	1.9
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	27	1.0	1.9
...	All other causes (Residual)	609	21.7	43.3
Black, non-Hispanic, female, 40-44 years				
...	All causes	4,881	100.0	324.4
1	Malignant neoplasms (C00-C97)	1,101	22.6	73.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	876	17.9	58.2
3	Human immunodeficiency virus (HIV) disease (B20-B24)	542	11.1	36.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	346	7.1	23.0
5	Cerebrovascular diseases (I60-I69)	271	5.6	18.0
6	Diabetes mellitus (E10-E14)	144	3.0	9.6
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	137	2.8	9.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	104	2.1	6.9
9	Septicemia (A40-A41)	89	1.8	5.9
10	Chronic liver disease and cirrhosis (K70,K73-K74)	81	1.7	5.4
11	Chronic lower respiratory diseases (J40-J47)	72	1.5	4.8
12	Influenza and pneumonia (J10-J18)	64	1.3	4.3
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	47	1.0	3.1
14	Anemias (D50-D64)	38	0.8	2.5
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	37	0.8	2.5
...	All other causes (Residual)	932	19.1	61.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
Black, non-Hispanic, female, 45-49 years				
...	All causes	6,716	100.0	481.7
1	Malignant neoplasms (C00-C97)	1,797	26.8	128.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,331	19.8	95.5
3	Human immunodeficiency virus (HIV) disease (B20-B24)	433	6.4	31.1
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	415	6.2	29.8
5	Cerebrovascular diseases (I60-I69)	403	6.0	28.9
6	Diabetes mellitus (E10-E14)	213	3.2	15.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	145	2.2	10.4
8	Chronic lower respiratory diseases (J40-J47)	134	2.0	9.6
9	Septicemia (A40-A41)	127	1.9	9.1
10	Chronic liver disease and cirrhosis (K70,K73-K74)	120	1.8	8.6
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	103	1.5	7.4
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	102	1.5	7.3
13	Influenza and pneumonia (J10-J18)	82	1.2	5.9
14	Anemias (D50-D64)	40	0.6	2.9
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	39	0.6	2.8
...	All other causes (Residual)	1,232	18.3	88.4
Black, non-Hispanic, female, 50-54 years				
...	All causes	8,082	100.0	690.6
1	Malignant neoplasms (C00-C97)	2,425	30.0	207.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,848	22.9	157.9
3	Cerebrovascular diseases (I60-I69)	490	6.1	41.9
4	Diabetes mellitus (E10-E14)	410	5.1	35.0
5	Human immunodeficiency virus (HIV) disease (B20-B24)	311	3.8	26.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	289	3.6	24.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	194	2.4	16.6
8	Septicemia (A40-A41)	189	2.3	16.1
9	Chronic lower respiratory diseases (J40-J47)	170	2.1	14.5
10	Chronic liver disease and cirrhosis (K70,K73-K74)	162	2.0	13.8
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	104	1.3	8.9
12	Influenza and pneumonia (J10-J18)	95	1.2	8.1
13	Viral hepatitis (B15-B19)	64	0.8	5.5
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	55	0.7	4.7
15	Anemias (D50-D64)	37	0.5	3.2
...	All other causes (Residual)	1,239	15.3	105.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
Black, non-Hispanic, female, 55-59 years				
...	All causes	8,598	100.0	962.5
1	Malignant neoplasms (C00-C97)	2,845	33.1	318.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,072	24.1	232.0
3	Diabetes mellitus (E10-E14)	514	6.0	57.5
4	Cerebrovascular diseases (I60-I69)	469	5.5	52.5
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	295	3.4	33.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	200	2.3	22.4
7	Septicemia (A40-A41)	194	2.3	21.7
8	Chronic lower respiratory diseases (J40-J47)	182	2.1	20.4
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	158	1.8	17.7
10	Human immunodeficiency virus (HIV) disease (B20-B24)	130	1.5	14.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	121	1.4	13.5
12	Influenza and pneumonia (J10-J18)	113	1.3	12.6
13	Viral hepatitis (B15-B19)	44	0.5	4.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	42	0.5	4.7
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	31	0.4	3.5
...	All other causes (Residual)	1,188	13.8	133.0
Black, non-Hispanic, female, 60-64 years				
...	All causes	9,350	100.0	1,400.6
1	Malignant neoplasms (C00-C97)	3,028	32.4	453.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,320	24.8	347.5
3	Diabetes mellitus (E10-E14)	619	6.6	92.7
4	Cerebrovascular diseases (I60-I69)	570	6.1	85.4
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	348	3.7	52.1
6	Chronic lower respiratory diseases (J40-J47)	275	2.9	41.2
7	Septicemia (A40-A41)	247	2.6	37.0
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	166	1.8	24.9
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	150	1.6	22.5
10	Influenza and pneumonia (J10-J18)	113	1.2	16.9
11	Chronic liver disease and cirrhosis (K70,K73-K74)	106	1.1	15.9
12	Human immunodeficiency virus (HIV) disease (B20-B24)	74	0.8	11.1
13	Aortic aneurysm and dissection (I71)	37	0.4	5.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	35	0.4	5.2
15	Pneumonitis due to solids and liquids (J69)	30	0.3	4.5
...	All other causes (Residual)	1,232	13.2	184.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
Black, non-Hispanic, female, 65-69 years				
...	All causes	10,820	100.0	2,001.4
1	Malignant neoplasms (C00-C97)	3,468	32.1	641.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,866	26.5	530.1
3	Diabetes mellitus (E10-E14)	731	6.8	135.2
4	Cerebrovascular diseases (I60-I69)	719	6.6	133.0
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	382	3.5	70.7
6	Chronic lower respiratory diseases (J40-J47)	338	3.1	62.5
7	Septicemia (A40-A41)	290	2.7	53.6
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	198	1.8	36.6
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	145	1.3	26.8
10	Influenza and pneumonia (J10-J18)	133	1.2	24.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	74	0.7	13.7
12	Alzheimer's disease (G30)	60	0.6	11.1
13	Human immunodeficiency virus (HIV) disease (B20-B24)	47	0.4	8.7
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	47	0.4	8.7
15	Aortic aneurysm and dissection (I71)	37	0.3	6.8
...	All other causes (Residual)	1,285	11.9	237.7
Black, non-Hispanic, female, 70-74 years				
...	All causes	13,078	100.0	2,947.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,708	28.4	835.7
2	Malignant neoplasms (C00-C97)	3,602	27.5	811.8
3	Cerebrovascular diseases (I60-I69)	958	7.3	215.9
4	Diabetes mellitus (E10-E14)	930	7.1	209.6
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	476	3.6	107.3
6	Chronic lower respiratory diseases (J40-J47)	413	3.2	93.1
7	Septicemia (A40-A41)	331	2.5	74.6
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	288	2.2	64.9
9	Influenza and pneumonia (J10-J18)	203	1.6	45.8
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	153	1.2	34.5
11	Alzheimer's disease (G30)	131	1.0	29.5
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	63	0.5	14.2
13	Pneumonitis due to solids and liquids (J69)	59	0.5	13.3
14	Aortic aneurysm and dissection (I71)	57	0.4	12.8
15	Chronic liver disease and cirrhosis (K70,K73-K74)	55	0.4	12.4
...	All other causes (Residual)	1,651	12.6	372.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
Black, non-Hispanic, female, 75-79 years				
...	All causes	16,110	100.0	4,447.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,748	29.5	1,310.7
2	Malignant neoplasms (C00-C97)	3,735	23.2	1,031.0
3	Cerebrovascular diseases (I60-I69)	1,426	8.9	393.6
4	Diabetes mellitus (E10-E14)	1,062	6.6	293.2
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	593	3.7	163.7
6	Chronic lower respiratory diseases (J40-J47)	544	3.4	150.2
7	Septicemia (A40-A41)	410	2.5	113.2
8	Alzheimer's disease (G30)	367	2.3	101.3
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	342	2.1	94.4
10	Influenza and pneumonia (J10-J18)	315	2.0	87.0
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	204	1.3	56.3
12	Pneumonitis due to solids and liquids (J69)	90	0.6	24.8
13	Aortic aneurysm and dissection (I71)	88	0.5	24.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	78	0.5	21.5
15	Atherosclerosis (I70)	58	0.4	16.0
...	All other causes (Residual)	2,050	12.7	565.9
Black, non-Hispanic, female, 80-84 years				
...	All causes	17,244	100.0	6,662.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,460	31.7	2,109.5
2	Malignant neoplasms (C00-C97)	3,087	17.9	1,192.7
3	Cerebrovascular diseases (I60-I69)	1,659	9.6	641.0
4	Diabetes mellitus (E10-E14)	1,000	5.8	386.4
5	Alzheimer's disease (G30)	657	3.8	253.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	573	3.3	221.4
7	Chronic lower respiratory diseases (J40-J47)	451	2.6	174.2
8	Septicemia (A40-A41)	449	2.6	173.5
8	Influenza and pneumonia (J10-J18)	449	2.6	173.5
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	384	2.2	148.4
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	186	1.1	71.9
12	Pneumonitis due to solids and liquids (J69)	132	0.8	51.0
13	Aortic aneurysm and dissection (I71)	96	0.6	37.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	93	0.5	35.9
15	Parkinson's disease (G20-G21)	77	0.4	29.7
...	All other causes (Residual)	2,491	14.4	962.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
Black, non-Hispanic, female, 85-89 years				
...	All causes	14,785	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,884	33.0	---
2	Malignant neoplasms (C00-C97)	2,064	14.0	---
3	Cerebrovascular diseases (I60-I69)	1,509	10.2	---
4	Alzheimer's disease (G30)	745	5.0	---
5	Diabetes mellitus (E10-E14)	744	5.0	---
6	Influenza and pneumonia (J10-J18)	482	3.3	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	464	3.1	---
8	Septicemia (A40-A41)	390	2.6	---
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	369	2.5	---
10	Chronic lower respiratory diseases (J40-J47)	332	2.2	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	153	1.0	---
12	Pneumonitis due to solids and liquids (J69)	127	0.9	---
13	Atherosclerosis (I70)	101	0.7	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	71	0.5	---
15	Aortic aneurysm and dissection (I71)	61	0.4	---
...	All other causes (Residual)	2,289	15.5	---
Black, non-Hispanic, female, 90-94 years				
...	All causes	11,135	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,960	35.6	---
2	Cerebrovascular diseases (I60-I69)	1,107	9.9	---
3	Malignant neoplasms (C00-C97)	1,039	9.3	---
4	Alzheimer's disease (G30)	667	6.0	---
5	Diabetes mellitus (E10-E14)	418	3.8	---
6	Influenza and pneumonia (J10-J18)	414	3.7	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	369	3.3	---
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	359	3.2	---
9	Septicemia (A40-A41)	259	2.3	---
10	Chronic lower respiratory diseases (J40-J47)	186	1.7	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	113	1.0	---
12	Pneumonitis due to solids and liquids (J69)	109	1.0	---
13	Atherosclerosis (I70)	95	0.9	---
14	Anemias (D50-D64)	50	0.4	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	39	0.4	---
15	Parkinson's disease (G20-G21)	39	0.4	---
...	All other causes (Residual)	1,912	17.2	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
Black, non-Hispanic, female, 95-99 years				
...	All causes	4,872	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,892	38.8	---
2	Cerebrovascular diseases (I60-I69)	455	9.3	---
3	Malignant neoplasms (C00-C97)	314	6.4	---
4	Alzheimer's disease (G30)	313	6.4	---
5	Influenza and pneumonia (J10-J18)	213	4.4	---
6	Diabetes mellitus (E10-E14)	148	3.0	---
7	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	133	2.7	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	132	2.7	---
9	Septicemia (A40-A41)	103	2.1	---
10	Chronic lower respiratory diseases (J40-J47)	69	1.4	---
11	Atherosclerosis (I70)	58	1.2	---
12	Pneumonitis due to solids and liquids (J69)	53	1.1	---
13	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	42	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	25	0.5	---
15	Anemias (D50-D64)	19	0.4	---
...	All other causes (Residual)	903	18.5	---
Black, non-Hispanic, female, 100 years and over				
...	All causes	1,670	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	661	39.6	---
2	Cerebrovascular diseases (I60-I69)	129	7.7	---
3	Alzheimer's disease (G30)	93	5.6	---
4	Malignant neoplasms (C00-C97)	80	4.8	---
5	Influenza and pneumonia (J10-J18)	71	4.3	---
6	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	55	3.3	---
7	Diabetes mellitus (E10-E14)	54	3.2	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	42	2.5	---
9	Septicemia (A40-A41)	35	2.1	---
10	Atherosclerosis (I70)	29	1.7	---
11	Chronic lower respiratory diseases (J40-J47)	20	1.2	---
12	Pneumonitis due to solids and liquids (J69)	19	1.1	---
13	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	16	1.0	---
14	Anemias (D50-D64)	10	0.6	---
14	Nutritional deficiencies (E40-E64)	10	0.6	---
...	All other causes (Residual)	346	20.7	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available