

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Hispanic, both sexes, all ages				
...	All causes	107,254	100.0	330.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	25,819	24.1	79.5
2	Malignant neoplasms (C00-C97)	21,160	19.7	65.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,830	8.2	27.2
4	Cerebrovascular diseases (I60-I69)	6,187	5.8	19.1
5	Diabetes mellitus (E10-E14)	5,328	5.0	16.4
6	Chronic liver disease and cirrhosis (K70,K73-K74)	3,187	3.0	9.8
7	Assault (homicide) (X85-Y09,Y87.1)	2,917	2.7	9.0
8	Chronic lower respiratory diseases (J40-J47)	2,689	2.5	8.3
9	Influenza and pneumonia (J10-J18)	2,625	2.4	8.1
10	Certain conditions originating in the perinatal period (P00-P96)	2,145	2.0	6.6
11	Human immunodeficiency virus (HIV) disease (B20-B24)	1,908	1.8	5.9
12	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,787	1.7	5.5
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,646	1.5	5.1
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,612	1.5	5.0
15	Septicemia (A40-A41)	1,226	1.1	3.8
...	All other causes (Residual)	18,188	17.0	56.0
Hispanic, both sexes, 1-4 years				
...	All causes	872	100.0	31.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	306	35.1	10.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	111	12.7	3.9
3	Malignant neoplasms (C00-C97)	75	8.6	2.7
4	Assault (homicide) (X85-Y09,Y87.1)	57	6.5	2.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	29	3.3	1.0
6	Influenza and pneumonia (J10-J18)	17	1.9	*
7	Septicemia (A40-A41)	15	1.7	*
8	Certain conditions originating in the perinatal period (P00-P96)	12	1.4	*
9	Chronic lower respiratory diseases (J40-J47)	9	1.0	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.8	*
11	Cerebrovascular diseases (I60-I69)	6	0.7	*
12	Meningitis (G00,G03)	4	0.5	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.5	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.5	*
15	Meningococcal infection (A39)	3	0.3	*
...	All other causes (Residual)	213	24.4	7.6

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Hispanic, both sexes, 5-14 years				
...	All causes	1,065	100.0	17.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	383	36.0	6.1
2	Malignant neoplasms (C00-C97)	198	18.6	3.2
3	Assault (homicide) (X85-Y09,Y87.1)	76	7.1	1.2
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	52	4.9	0.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	38	3.6	0.6
6	Intentional self-harm (suicide) (X60-X84,Y87.0)	29	2.7	0.5
7	Influenza and pneumonia (J10-J18)	21	2.0	0.3
8	Chronic lower respiratory diseases (J40-J47)	15	1.4	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	1.2	*
10	Certain conditions originating in the perinatal period (P00-P96)	10	0.9	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	9	0.8	*
12	Cerebrovascular diseases (I60-I69)	8	0.8	*
13	Septicemia (A40-A41)	7	0.7	*
13	Anemias (D50-D64)	7	0.7	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.6	*
...	All other causes (Residual)	193	18.1	3.1
Hispanic, both sexes, 15-24 years				
...	All causes	4,793	100.0	85.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,095	43.7	37.2
2	Assault (homicide) (X85-Y09,Y87.1)	1,126	23.5	20.0
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	447	9.3	7.9
4	Malignant neoplasms (C00-C97)	275	5.7	4.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	125	2.6	2.2
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	54	1.1	1.0
7	Cerebrovascular diseases (I60-I69)	31	0.6	0.5
8	Pregnancy, childbirth and the puerperium (O00-099)	24	0.5	0.4
9	Human immunodeficiency virus (HIV) disease (B20-B24)	23	0.5	0.4
10	Influenza and pneumonia (J10-J18)	21	0.4	0.4
11	Legal intervention (Y35,Y89.0)	18	0.4	*
12	Chronic lower respiratory diseases (J40-J47)	16	0.3	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	0.3	*
14	Diabetes mellitus (E10-E14)	13	0.3	*
15	Septicemia (A40-A41)	12	0.3	*
...	All other causes (Residual)	499	10.4	8.9

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Hispanic, both sexes, 25-34 years				
...	All causes	5,509	100.0	104.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,776	32.2	33.6
2	Assault (homicide) (X85-Y09,Y87.1)	810	14.7	15.3
3	Malignant neoplasms (C00-C97)	476	8.6	9.0
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	420	7.6	7.9
5	Human immunodeficiency virus (HIV) disease (B20-B24)	365	6.6	6.9
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	280	5.1	5.3
7	Cerebrovascular diseases (I60-I69)	108	2.0	2.0
8	Chronic liver disease and cirrhosis (K70,K73-K74)	92	1.7	1.7
9	Diabetes mellitus (E10-E14)	64	1.2	1.2
10	Influenza and pneumonia (J10-J18)	48	0.9	0.9
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	46	0.8	0.9
12	Pregnancy, childbirth and the puerperium (O00-099)	36	0.7	0.7
13	Septicemia (A40-A41)	31	0.6	0.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	26	0.5	0.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	25	0.5	0.5
...	All other causes (Residual)	906	16.4	17.1
Hispanic, both sexes, 35-44 years				
...	All causes	8,085	100.0	163.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,618	20.0	32.8
2	Malignant neoplasms (C00-C97)	1,295	16.0	26.2
3	Human immunodeficiency virus (HIV) disease (B20-B24)	762	9.4	15.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	760	9.4	15.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	501	6.2	10.1
6	Assault (homicide) (X85-Y09,Y87.1)	447	5.5	9.1
7	Intentional self-harm (suicide) (X60-X84,Y87.0)	385	4.8	7.8
8	Cerebrovascular diseases (I60-I69)	267	3.3	5.4
9	Diabetes mellitus (E10-E14)	154	1.9	3.1
10	Viral hepatitis (B15-B19)	126	1.6	2.6
11	Influenza and pneumonia (J10-J18)	82	1.0	1.7
12	Septicemia (A40-A41)	79	1.0	1.6
13	Chronic lower respiratory diseases (J40-J47)	55	0.7	1.1
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	53	0.7	1.1
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	43	0.5	0.9
...	All other causes (Residual)	1,458	18.0	29.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Hispanic, both sexes, 45-54 years				
...	All causes	10,356	100.0	333.1
1	Malignant neoplasms (C00-C97)	2,575	24.9	82.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,754	16.9	56.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	953	9.2	30.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	854	8.2	27.5
5	Human immunodeficiency virus (HIV) disease (B20-B24)	531	5.1	17.1
6	Cerebrovascular diseases (I60-I69)	477	4.6	15.3
7	Diabetes mellitus (E10-E14)	439	4.2	14.1
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	244	2.4	7.8
9	Viral hepatitis (B15-B19)	218	2.1	7.0
10	Assault (homicide) (X85-Y09,Y87.1)	211	2.0	6.8
11	Septicemia (A40-A41)	122	1.2	3.9
12	Influenza and pneumonia (J10-J18)	120	1.2	3.9
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	109	1.1	3.5
14	Chronic lower respiratory diseases (J40-J47)	105	1.0	3.4
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	55	0.5	1.8
...	All other causes (Residual)	1,589	15.3	51.1
Hispanic, both sexes, 55-64 years				
...	All causes	12,733	100.0	721.3
1	Malignant neoplasms (C00-C97)	3,780	29.7	214.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,136	24.6	177.6
3	Diabetes mellitus (E10-E14)	950	7.5	53.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	708	5.6	40.1
5	Cerebrovascular diseases (I60-I69)	681	5.3	38.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	492	3.9	27.9
7	Chronic lower respiratory diseases (J40-J47)	244	1.9	13.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	232	1.8	13.1
9	Influenza and pneumonia (J10-J18)	191	1.5	10.8
10	Human immunodeficiency virus (HIV) disease (B20-B24)	163	1.3	9.2
11	Viral hepatitis (B15-B19)	147	1.2	8.3
12	Septicemia (A40-A41)	145	1.1	8.2
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	109	0.9	6.2
14	Intentional self-harm (suicide) (X60-X84,Y87.0)	102	0.8	5.8
15	Assault (homicide) (X85-Y09,Y87.1)	77	0.6	4.4
...	All other causes (Residual)	1,576	12.4	89.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
	Hispanic, both sexes, 65-74 years			
...	All causes	19,420	100.0	1,683.4
1	Malignant neoplasms (C00-C97)	5,557	28.6	481.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,453	28.1	472.7
3	Diabetes mellitus (E10-E14)	1,592	8.2	138.0
4	Cerebrovascular diseases (I60-I69)	1,219	6.3	105.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	628	3.2	54.4
6	Chronic lower respiratory diseases (J40-J47)	572	2.9	49.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	418	2.2	36.2
8	Influenza and pneumonia (J10-J18)	405	2.1	35.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	404	2.1	35.0
10	Septicemia (A40-A41)	235	1.2	20.4
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	162	0.8	14.0
12	Alzheimer's disease (G30)	128	0.7	11.1
13	Aortic aneurysm and dissection (I71)	121	0.6	10.5
14	Viral hepatitis (B15-B19)	120	0.6	10.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	98	0.5	8.5
...	All other causes (Residual)	2,308	11.9	200.1
	Hispanic, both sexes, 75-84 years			
...	All causes	21,951	100.0	3,704.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,067	32.2	1,192.5
2	Malignant neoplasms (C00-C97)	4,827	22.0	814.5
3	Cerebrovascular diseases (I60-I69)	1,742	7.9	294.0
4	Diabetes mellitus (E10-E14)	1,410	6.4	237.9
5	Chronic lower respiratory diseases (J40-J47)	921	4.2	155.4
6	Influenza and pneumonia (J10-J18)	690	3.1	116.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	429	2.0	72.4
8	Alzheimer's disease (G30)	401	1.8	67.7
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	365	1.7	61.6
10	Chronic liver disease and cirrhosis (K70,K73-K74)	323	1.5	54.5
11	Septicemia (A40-A41)	300	1.4	50.6
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	232	1.1	39.1
13	Parkinson's disease (G20-G21)	191	0.9	32.2
14	Pneumonitis due to solids and liquids (J69)	141	0.6	23.8
15	Aortic aneurysm and dissection (I71)	137	0.6	23.1
...	All other causes (Residual)	2,775	12.6	468.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
	Hispanic, both sexes, 85 years and over			
...	All causes	17,818	100.0	8,715.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,086	39.8	3,465.8
2	Malignant neoplasms (C00-C97)	2,085	11.7	1,019.8
3	Cerebrovascular diseases (I60-I69)	1,632	9.2	798.2
4	Influenza and pneumonia (J10-J18)	970	5.4	474.4
5	Chronic lower respiratory diseases (J40-J47)	725	4.1	354.6
6	Diabetes mellitus (E10-E14)	701	3.9	342.9
7	Alzheimer's disease (G30)	573	3.2	280.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	346	1.9	169.2
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	278	1.6	136.0
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	223	1.3	109.1
11	Septicemia (A40-A41)	218	1.2	106.6
12	Pneumonitis due to solids and liquids (J69)	207	1.2	101.2
13	Atherosclerosis (I70)	200	1.1	97.8
14	Parkinson's disease (G20-G21)	130	0.7	63.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	106	0.6	51.8
...	All other causes (Residual)	2,338	13.1	1,143.5
	Hispanic, male, all ages			
...	All causes	60,172	100.0	368.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,566	22.5	83.2
2	Malignant neoplasms (C00-C97)	11,138	18.5	68.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,696	11.1	41.1
4	Cerebrovascular diseases (I60-I69)	2,865	4.8	17.6
5	Diabetes mellitus (E10-E14)	2,507	4.2	15.4
6	Assault (homicide) (X85-Y09,Y87.1)	2,431	4.0	14.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,312	3.8	14.2
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,525	2.5	9.3
9	Human immunodeficiency virus (HIV) disease (B20-B24)	1,493	2.5	9.2
10	Chronic lower respiratory diseases (J40-J47)	1,451	2.4	8.9
11	Influenza and pneumonia (J10-J18)	1,303	2.2	8.0
12	Certain conditions originating in the perinatal period (P00-P96)	1,194	2.0	7.3
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	857	1.4	5.3
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	805	1.3	4.9
15	Septicemia (A40-A41)	625	1.0	3.8
...	All other causes (Residual)	9,404	15.6	57.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Hispanic, male, 1-4 years				
...	All causes	475	100.0	33.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	174	36.6	12.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	60	12.6	4.2
3	Malignant neoplasms (C00-C97)	43	9.1	3.0
4	Assault (homicide) (X85-Y09,Y87.1)	30	6.3	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	21	4.4	1.5
6	Septicemia (A40-A41)	9	1.9	*
7	Certain conditions originating in the perinatal period (P00-P96)	7	1.5	*
8	Influenza and pneumonia (J10-J18)	5	1.1	*
8	Chronic lower respiratory diseases (J40-J47)	5	1.1	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.6	*
11	Meningococcal infection (A39)	2	0.4	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.4	*
11	Meningitis (G00,G03)	2	0.4	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
15	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
15	Diseases of appendix (K35-K38)	1	0.2	*
...	All other causes (Residual)	108	22.7	7.5
Hispanic, male, 5-14 years				
...	All causes	622	100.0	19.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	210	33.8	6.6
2	Malignant neoplasms (C00-C97)	122	19.6	3.8
3	Assault (homicide) (X85-Y09,Y87.1)	56	9.0	1.8
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	28	4.5	0.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	26	4.2	0.8
6	Intentional self-harm (suicide) (X60-X84,Y87.0)	22	3.5	0.7
7	Influenza and pneumonia (J10-J18)	10	1.6	*
8	Chronic lower respiratory diseases (J40-J47)	9	1.4	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	1.3	*
10	Certain conditions originating in the perinatal period (P00-P96)	6	1.0	*
11	Septicemia (A40-A41)	5	0.8	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	5	0.8	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.6	*
14	Anemias (D50-D64)	3	0.5	*
14	Cerebrovascular diseases (I60-I69)	3	0.5	*
14	Diseases of appendix (K35-K38)	3	0.5	*
...	All other causes (Residual)	102	16.4	3.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total deaths		
		Number	Rate	Rate
	Hispanic, male, 15-24 years			
...	All causes	3,837	100.0	131.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,717	44.7	58.7
2	Assault (homicide) (X85-Y09,Y87.1)	1,014	26.4	34.7
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	387	10.1	13.2
4	Malignant neoplasms (C00-C97)	162	4.2	5.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	87	2.3	3.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	39	1.0	1.3
7	Human immunodeficiency virus (HIV) disease (B20-B24)	18	0.5	*
8	Cerebrovascular diseases (I60-I69)	17	0.4	*
8	Legal intervention (Y35,Y89.0)	17	0.4	*
10	Chronic lower respiratory diseases (J40-J47)	13	0.3	*
11	Influenza and pneumonia (J10-J18)	10	0.3	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.2	*
13	Diabetes mellitus (E10-E14)	7	0.2	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.2	*
15	Meningococcal infection (A39)	6	0.2	*
15	Septicemia (A40-A41)	6	0.2	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.2	*
...	All other causes (Residual)	316	8.2	10.8
	Hispanic, male, 25-34 years			
...	All causes	4,200	100.0	155.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,499	35.7	55.4
2	Assault (homicide) (X85-Y09,Y87.1)	697	16.6	25.7
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	370	8.8	13.7
4	Human immunodeficiency virus (HIV) disease (B20-B24)	274	6.5	10.1
5	Malignant neoplasms (C00-C97)	241	5.7	8.9
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	199	4.7	7.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	82	2.0	3.0
8	Cerebrovascular diseases (I60-I69)	65	1.5	2.4
9	Diabetes mellitus (E10-E14)	48	1.1	1.8
10	Influenza and pneumonia (J10-J18)	27	0.6	1.0
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	26	0.6	1.0
12	Septicemia (A40-A41)	21	0.5	0.8
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	15	0.4	*
13	Legal intervention (Y35,Y89.0)	15	0.4	*
15	Chronic lower respiratory diseases (J40-J47)	12	0.3	*
...	All other causes (Residual)	609	14.5	22.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Hispanic, male, 35-44 years				
...	All causes	5,597	100.0	221.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,317	23.5	52.2
2	Human immunodeficiency virus (HIV) disease (B20-B24)	603	10.8	23.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	550	9.8	21.8
4	Malignant neoplasms (C00-C97)	534	9.5	21.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	399	7.1	15.8
6	Assault (homicide) (X85-Y09,Y87.1)	359	6.4	14.2
7	Intentional self-harm (suicide) (X60-X84,Y87.0)	320	5.7	12.7
8	Cerebrovascular diseases (I60-I69)	145	2.6	5.7
9	Viral hepatitis (B15-B19)	100	1.8	4.0
10	Diabetes mellitus (E10-E14)	95	1.7	3.8
11	Influenza and pneumonia (J10-J18)	56	1.0	2.2
12	Septicemia (A40-A41)	46	0.8	1.8
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	36	0.6	1.4
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	27	0.5	1.1
15	Chronic lower respiratory diseases (J40-J47)	24	0.4	1.0
...	All other causes (Residual)	986	17.6	39.1
Hispanic, male, 45-54 years				
...	All causes	6,808	100.0	447.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,307	19.2	85.9
2	Malignant neoplasms (C00-C97)	1,232	18.1	80.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	750	11.0	49.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	707	10.4	46.5
5	Human immunodeficiency virus (HIV) disease (B20-B24)	426	6.3	28.0
6	Cerebrovascular diseases (I60-I69)	280	4.1	18.4
7	Diabetes mellitus (E10-E14)	240	3.5	15.8
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	194	2.8	12.7
9	Viral hepatitis (B15-B19)	165	2.4	10.8
10	Assault (homicide) (X85-Y09,Y87.1)	158	2.3	10.4
11	Influenza and pneumonia (J10-J18)	82	1.2	5.4
12	Septicemia (A40-A41)	78	1.1	5.1
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	69	1.0	4.5
14	Chronic lower respiratory diseases (J40-J47)	53	0.8	3.5
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	44	0.6	2.9
...	All other causes (Residual)	1,023	15.0	67.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Hispanic, male, 55-64 years				
...	All causes	7,760	100.0	948.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,128	27.4	260.1
2	Malignant neoplasms (C00-C97)	2,034	26.2	248.7
3	Chronic liver disease and cirrhosis (K70,K73-K74)	531	6.8	64.9
4	Diabetes mellitus (E10-E14)	520	6.7	63.6
5	Cerebrovascular diseases (I60-I69)	392	5.1	47.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	347	4.5	42.4
7	Chronic lower respiratory diseases (J40-J47)	146	1.9	17.8
8	Human immunodeficiency virus (HIV) disease (B20-B24)	123	1.6	15.0
9	Influenza and pneumonia (J10-J18)	119	1.5	14.5
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	115	1.5	14.1
11	Viral hepatitis (B15-B19)	94	1.2	11.5
12	Intentional self-harm (suicide) (X60-X84,Y87.0)	89	1.1	10.9
13	Septicemia (A40-A41)	80	1.0	9.8
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	65	0.8	7.9
15	Assault (homicide) (X85-Y09,Y87.1)	57	0.7	7.0
...	All other causes (Residual)	920	11.9	112.5
Hispanic, male, 65-74 years				
...	All causes	10,842	100.0	2,127.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,245	29.9	636.8
2	Malignant neoplasms (C00-C97)	3,086	28.5	605.6
3	Diabetes mellitus (E10-E14)	762	7.0	149.5
4	Cerebrovascular diseases (I60-I69)	645	5.9	126.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	395	3.6	77.5
6	Chronic lower respiratory diseases (J40-J47)	343	3.2	67.3
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	271	2.5	53.2
8	Influenza and pneumonia (J10-J18)	218	2.0	42.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	194	1.8	38.1
10	Septicemia (A40-A41)	126	1.2	24.7
11	Aortic aneurysm and dissection (I71)	76	0.7	14.9
12	Intentional self-harm (suicide) (X60-X84,Y87.0)	74	0.7	14.5
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	72	0.7	14.1
14	Alzheimer's disease (G30)	63	0.6	12.4
15	Viral hepatitis (B15-B19)	52	0.5	10.2
...	All other causes (Residual)	1,220	11.3	239.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
	Hispanic, male, 75-84 years			
...	All causes	10,973	100.0	4,470.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,525	32.1	1,436.0
2	Malignant neoplasms (C00-C97)	2,657	24.2	1,082.4
3	Cerebrovascular diseases (I60-I69)	799	7.3	325.5
4	Diabetes mellitus (E10-E14)	609	5.5	248.1
5	Chronic lower respiratory diseases (J40-J47)	509	4.6	207.4
6	Influenza and pneumonia (J10-J18)	367	3.3	149.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	206	1.9	83.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	191	1.7	77.8
9	Alzheimer's disease (G30)	166	1.5	67.6
10	Chronic liver disease and cirrhosis (K70,K73-K74)	163	1.5	66.4
11	Septicemia (A40-A41)	143	1.3	58.3
12	Parkinson's disease (G20-G21)	112	1.0	45.6
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	107	1.0	43.6
14	Pneumonitis due to solids and liquids (J69)	87	0.8	35.4
15	Aortic aneurysm and dissection (I71)	86	0.8	35.0
...	All other causes (Residual)	1,246	11.4	507.6
	Hispanic, male, 85 years and over			
...	All causes	6,493	100.0	9,385.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,430	37.4	3,512.4
2	Malignant neoplasms (C00-C97)	1,017	15.7	1,470.0
3	Cerebrovascular diseases (I60-I69)	511	7.9	738.6
4	Influenza and pneumonia (J10-J18)	379	5.8	547.8
5	Chronic lower respiratory diseases (J40-J47)	333	5.1	481.3
6	Diabetes mellitus (E10-E14)	223	3.4	322.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	145	2.2	209.6
8	Alzheimer's disease (G30)	141	2.2	203.8
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	125	1.9	180.7
10	Pneumonitis due to solids and liquids (J69)	84	1.3	121.4
11	Septicemia (A40-A41)	78	1.2	112.7
12	Parkinson's disease (G20-G21)	71	1.1	102.6
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	70	1.1	101.2
14	Atherosclerosis (I70)	52	0.8	75.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	43	0.7	62.2
...	All other causes (Residual)	791	12.2	1,143.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Hispanic, female, all ages				
...	All causes	47,082	100.0	291.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,253	26.0	75.9
2	Malignant neoplasms (C00-C97)	10,022	21.3	62.0
3	Cerebrovascular diseases (I60-I69)	3,322	7.1	20.6
4	Diabetes mellitus (E10-E14)	2,821	6.0	17.5
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,134	4.5	13.2
6	Influenza and pneumonia (J10-J18)	1,322	2.8	8.2
7	Chronic lower respiratory diseases (J40-J47)	1,238	2.6	7.7
8	Certain conditions originating in the perinatal period (P00-P96)	951	2.0	5.9
9	Chronic liver disease and cirrhosis (K70,K73-K74)	875	1.9	5.4
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	841	1.8	5.2
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	755	1.6	4.7
12	Alzheimer's disease (G30)	744	1.6	4.6
13	Septicemia (A40-A41)	601	1.3	3.7
14	Assault (homicide) (X85-Y09,Y87.1)	486	1.0	3.0
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	435	0.9	2.7
...	All other causes (Residual)	8,282	17.6	51.3
Hispanic, female, 1-4 years				
...	All causes	397	100.0	28.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	132	33.2	9.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	51	12.8	3.7
3	Malignant neoplasms (C00-C97)	32	8.1	2.3
4	Assault (homicide) (X85-Y09,Y87.1)	27	6.8	2.0
5	Influenza and pneumonia (J10-J18)	12	3.0	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	8	2.0	*
7	Septicemia (A40-A41)	6	1.5	*
7	Cerebrovascular diseases (I60-I69)	6	1.5	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	1.3	*
9	Certain conditions originating in the perinatal period (P00-P96)	5	1.3	*
11	Chronic lower respiratory diseases (J40-J47)	4	1.0	*
12	Tuberculosis (A16-A19)	2	0.5	*
12	Meningitis (G00,G03)	2	0.5	*
12	Pneumonitis due to solids and liquids (J69)	2	0.5	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.5	*
...	All other causes (Residual)	101	25.4	7.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Hispanic, female, 5-14 years				
...	All causes	443	100.0	14.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	173	39.1	5.7
2	Malignant neoplasms (C00-C97)	76	17.2	2.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	24	5.4	0.8
4	Assault (homicide) (X85-Y09,Y87.1)	20	4.5	0.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	12	2.7	*
6	Influenza and pneumonia (J10-J18)	11	2.5	*
7	Intentional self-harm (suicide) (X60-X84,Y87.0)	7	1.6	*
8	Chronic lower respiratory diseases (J40-J47)	6	1.4	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	1.1	*
9	Cerebrovascular diseases (I60-I69)	5	1.1	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.9	*
11	Anemias (D50-D64)	4	0.9	*
11	Certain conditions originating in the perinatal period (P00-P96)	4	0.9	*
14	Meningococcal infection (A39)	2	0.5	*
14	Septicemia (A40-A41)	2	0.5	*
14	Diabetes mellitus (E10-E14)	2	0.5	*
14	Meningitis (G00,G03)	2	0.5	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.5	*
...	All other causes (Residual)	82	18.5	2.7
Hispanic, female, 15-24 years				
...	All causes	956	100.0	35.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	378	39.5	13.9
2	Malignant neoplasms (C00-C97)	113	11.8	4.2
3	Assault (homicide) (X85-Y09,Y87.1)	112	11.7	4.1
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	60	6.3	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	38	4.0	1.4
6	Pregnancy, childbirth and the puerperium (O00-099)	24	2.5	0.9
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	15	1.6	*
8	Cerebrovascular diseases (I60-I69)	14	1.5	*
9	Influenza and pneumonia (J10-J18)	11	1.2	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.7	*
11	Septicemia (A40-A41)	6	0.6	*
11	Diabetes mellitus (E10-E14)	6	0.6	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	5	0.5	*
14	Anemias (D50-D64)	4	0.4	*
14	Aortic aneurysm and dissection (I71)	4	0.4	*
...	All other causes (Residual)	159	16.6	5.9

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Hispanic, female, 25-34 years				
...	All causes	1,309	100.0	50.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	277	21.2	10.8
2	Malignant neoplasms (C00-C97)	235	18.0	9.1
3	Assault (homicide) (X85-Y09,Y87.1)	113	8.6	4.4
4	Human immunodeficiency virus (HIV) disease (B20-B24)	91	7.0	3.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	81	6.2	3.1
6	Intentional self-harm (suicide) (X60-X84,Y87.0)	50	3.8	1.9
7	Cerebrovascular diseases (I60-I69)	43	3.3	1.7
8	Pregnancy, childbirth and the puerperium (O00-099)	36	2.8	1.4
9	Influenza and pneumonia (J10-J18)	21	1.6	0.8
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	20	1.5	0.8
11	Diabetes mellitus (E10-E14)	16	1.2	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	1.1	*
13	Septicemia (A40-A41)	10	0.8	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	0.8	*
13	Chronic liver disease and cirrhosis (K70,K73-K74)	10	0.8	*
...	All other causes (Residual)	281	21.5	10.9
Hispanic, female, 35-44 years				
...	All causes	2,488	100.0	103.0
1	Malignant neoplasms (C00-C97)	761	30.6	31.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	301	12.1	12.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	210	8.4	8.7
4	Human immunodeficiency virus (HIV) disease (B20-B24)	159	6.4	6.6
5	Cerebrovascular diseases (I60-I69)	122	4.9	5.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	102	4.1	4.2
7	Assault (homicide) (X85-Y09,Y87.1)	88	3.5	3.6
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	65	2.6	2.7
9	Diabetes mellitus (E10-E14)	59	2.4	2.4
10	Septicemia (A40-A41)	33	1.3	1.4
11	Chronic lower respiratory diseases (J40-J47)	31	1.2	1.3
12	Viral hepatitis (B15-B19)	26	1.0	1.1
12	Influenza and pneumonia (J10-J18)	26	1.0	1.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	0.9	1.0
14	Pregnancy, childbirth and the puerperium (O00-099)	23	0.9	1.0
...	All other causes (Residual)	459	18.4	19.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total deaths		
		Number	Rate	Rate
	Hispanic, female, 45-54 years			
...	All causes	3,548	100.0	223.5
1	Malignant neoplasms (C00-C97)	1,343	37.9	84.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	447	12.6	28.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	203	5.7	12.8
4	Diabetes mellitus (E10-E14)	199	5.6	12.5
5	Cerebrovascular diseases (I60-I69)	197	5.6	12.4
6	Chronic liver disease and cirrhosis (K70,K73-K74)	147	4.1	9.3
7	Human immunodeficiency virus (HIV) disease (B20-B24)	105	3.0	6.6
8	Viral hepatitis (B15-B19)	53	1.5	3.3
8	Assault (homicide) (X85-Y09,Y87.1)	53	1.5	3.3
10	Chronic lower respiratory diseases (J40-J47)	52	1.5	3.3
11	Intentional self-harm (suicide) (X60-X84,Y87.0)	50	1.4	3.1
12	Septicemia (A40-A41)	44	1.2	2.8
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	40	1.1	2.5
14	Influenza and pneumonia (J10-J18)	38	1.1	2.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	25	0.7	1.6
...	All other causes (Residual)	552	15.6	34.8
	Hispanic, female, 55-64 years			
...	All causes	4,973	100.0	525.0
1	Malignant neoplasms (C00-C97)	1,746	35.1	184.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,008	20.3	106.4
3	Diabetes mellitus (E10-E14)	430	8.6	45.4
4	Cerebrovascular diseases (I60-I69)	289	5.8	30.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	177	3.6	18.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	145	2.9	15.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	117	2.4	12.4
8	Chronic lower respiratory diseases (J40-J47)	98	2.0	10.3
9	Influenza and pneumonia (J10-J18)	72	1.4	7.6
10	Septicemia (A40-A41)	65	1.3	6.9
11	Viral hepatitis (B15-B19)	53	1.1	5.6
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	44	0.9	4.6
13	Human immunodeficiency virus (HIV) disease (B20-B24)	40	0.8	4.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	28	0.6	3.0
15	Assault (homicide) (X85-Y09,Y87.1)	20	0.4	2.1
...	All other causes (Residual)	641	12.9	67.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Hispanic, female, 65-74 years				
...	All causes	8,578	100.0	1,331.9
1	Malignant neoplasms (C00-C97)	2,471	28.8	383.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,208	25.7	342.8
3	Diabetes mellitus (E10-E14)	830	9.7	128.9
4	Cerebrovascular diseases (I60-I69)	574	6.7	89.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	233	2.7	36.2
6	Chronic lower respiratory diseases (J40-J47)	229	2.7	35.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	210	2.4	32.6
8	Influenza and pneumonia (J10-J18)	187	2.2	29.0
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	147	1.7	22.8
10	Septicemia (A40-A41)	109	1.3	16.9
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	90	1.0	14.0
12	Viral hepatitis (B15-B19)	68	0.8	10.6
13	Alzheimer's disease (G30)	65	0.8	10.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	52	0.6	8.1
15	Aortic aneurysm and dissection (I71)	45	0.5	7.0
...	All other causes (Residual)	1,060	12.4	164.6
Hispanic, female, 75-84 years				
...	All causes	10,978	100.0	3,162.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,542	32.3	1,020.3
2	Malignant neoplasms (C00-C97)	2,170	19.8	625.1
3	Cerebrovascular diseases (I60-I69)	943	8.6	271.6
4	Diabetes mellitus (E10-E14)	801	7.3	230.7
5	Chronic lower respiratory diseases (J40-J47)	412	3.8	118.7
6	Influenza and pneumonia (J10-J18)	323	2.9	93.0
7	Alzheimer's disease (G30)	235	2.1	67.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	223	2.0	64.2
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	174	1.6	50.1
10	Chronic liver disease and cirrhosis (K70,K73-K74)	160	1.5	46.1
11	Septicemia (A40-A41)	157	1.4	45.2
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	125	1.1	36.0
13	Parkinson's disease (G20-G21)	79	0.7	22.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	69	0.6	19.9
15	Atherosclerosis (I70)	59	0.5	17.0
...	All other causes (Residual)	1,506	13.7	433.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
	Hispanic, female, 85 years and over			
...	All causes	11,325	100.0	8,372.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,656	41.1	3,442.0
2	Cerebrovascular diseases (I60-I69)	1,121	9.9	828.7
3	Malignant neoplasms (C00-C97)	1,068	9.4	789.5
4	Influenza and pneumonia (J10-J18)	591	5.2	436.9
5	Diabetes mellitus (E10-E14)	478	4.2	353.4
6	Alzheimer's disease (G30)	432	3.8	319.4
7	Chronic lower respiratory diseases (J40-J47)	392	3.5	289.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	201	1.8	148.6
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	153	1.4	113.1
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	153	1.4	113.1
11	Atherosclerosis (I70)	148	1.3	109.4
12	Septicemia (A40-A41)	140	1.2	103.5
13	Pneumonitis due to solids and liquids (J69)	123	1.1	90.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	63	0.6	46.6
15	Parkinson's disease (G20-G21)	59	0.5	43.6
...	All other causes (Residual)	1,547	13.7	1,143.6
	White, non-Hispanic, both sexes, all ages			
...	All causes	1,959,919	100.0	996.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	594,465	30.3	302.3
2	Malignant neoplasms (C00-C97)	458,200	23.4	233.0
3	Cerebrovascular diseases (I60-I69)	138,210	7.1	70.3
4	Chronic lower respiratory diseases (J40-J47)	109,901	5.6	55.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	73,536	3.8	37.4
6	Influenza and pneumonia (J10-J18)	55,135	2.8	28.0
7	Diabetes mellitus (E10-E14)	50,163	2.6	25.5
8	Alzheimer's disease (G30)	45,246	2.3	23.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	27,924	1.4	14.2
10	Intentional self-harm (suicide) (X60-X84,Y87.0)	24,616	1.3	12.5
11	Septicemia (A40-A41)	23,517	1.2	12.0
12	Chronic liver disease and cirrhosis (K70,K73-K74)	19,739	1.0	10.0
13	Parkinson's disease (G20-G21)	14,497	0.7	7.4
14	Pneumonitis due to solids and liquids (J69)	14,343	0.7	7.3
15	Aortic aneurysm and dissection (I71)	13,722	0.7	7.0
...	All other causes (Residual)	296,705	15.1	150.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
White, non-Hispanic, both sexes, 1-4 years				
...	All causes	2,641	100.0	27.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,033	39.1	10.9
2	Malignant neoplasms (C00-C97)	256	9.7	2.7
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	240	9.1	2.5
4	Assault (homicide) (X85-Y09,Y87.1)	144	5.5	1.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	98	3.7	1.0
6	Septicemia (A40-A41)	51	1.9	0.5
6	Influenza and pneumonia (J10-J18)	51	1.9	0.5
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	37	1.4	0.4
9	Certain conditions originating in the perinatal period (P00-P96)	34	1.3	0.4
10	Cerebrovascular diseases (I60-I69)	25	0.9	0.3
11	Meningitis (G00,G03)	19	0.7	*
11	Chronic lower respiratory diseases (J40-J47)	19	0.7	*
13	Anemias (D50-D64)	14	0.5	*
14	Meningococcal infection (A39)	11	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	8	0.3	*
...	All other causes (Residual)	601	22.8	6.4
White, non-Hispanic, both sexes, 5-14 years				
...	All causes	4,397	100.0	17.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,843	41.9	7.2
2	Malignant neoplasms (C00-C97)	627	14.3	2.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	240	5.5	0.9
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	225	5.1	0.9
5	Assault (homicide) (X85-Y09,Y87.1)	164	3.7	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	140	3.2	0.5
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	63	1.4	0.2
8	Influenza and pneumonia (J10-J18)	49	1.1	0.2
9	Chronic lower respiratory diseases (J40-J47)	47	1.1	0.2
10	Cerebrovascular diseases (I60-I69)	46	1.0	0.2
11	Septicemia (A40-A41)	42	1.0	0.2
12	Certain conditions originating in the perinatal period (P00-P96)	20	0.5	0.1
13	Meningitis (G00,G03)	18	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	17	0.4	*
...	All other causes (Residual)	839	19.1	3.3

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total deaths		
		Number	Rate	Rate
White, non-Hispanic, both sexes, 15-24 years				
...	All causes	18,261	100.0	72.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,735	53.3	38.4
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	2,868	15.7	11.3
3	Malignant neoplasms (C00-C97)	1,059	5.8	4.2
4	Assault (homicide) (X85-Y09,Y87.1)	874	4.8	3.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	531	2.9	2.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	279	1.5	1.1
7	Cerebrovascular diseases (I60-I69)	118	0.6	0.5
8	Influenza and pneumonia (J10-J18)	116	0.6	0.5
9	Diabetes mellitus (E10-E14)	92	0.5	0.4
10	Chronic lower respiratory diseases (J40-J47)	87	0.5	0.3
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	63	0.3	0.2
12	Septicemia (A40-A41)	48	0.3	0.2
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	48	0.3	0.2
14	Meningococcal infection (A39)	37	0.2	0.1
15	Pregnancy, childbirth and the puerperium (O00-099)	33	0.2	0.1
...	All other causes (Residual)	2,273	12.4	9.0
White, non-Hispanic, both sexes, 25-34 years				
...	All causes	23,191	100.0	92.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,730	33.3	30.9
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	3,658	15.8	14.6
3	Malignant neoplasms (C00-C97)	2,599	11.2	10.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,653	7.1	6.6
5	Assault (homicide) (X85-Y09,Y87.1)	1,023	4.4	4.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	619	2.7	2.5
7	Diabetes mellitus (E10-E14)	357	1.5	1.4
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	344	1.5	1.4
9	Cerebrovascular diseases (I60-I69)	313	1.3	1.2
10	Chronic liver disease and cirrhosis (K70,K73-K74)	217	0.9	0.9
11	Influenza and pneumonia (J10-J18)	213	0.9	0.9
12	Chronic lower respiratory diseases (J40-J47)	144	0.6	0.6
13	Septicemia (A40-A41)	134	0.6	0.5
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	110	0.5	0.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	105	0.5	0.4
...	All other causes (Residual)	3,972	17.1	15.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
White, non-Hispanic, both sexes, 35-44 years				
...	All causes	58,231	100.0	179.7
1	Malignant neoplasms (C00-C97)	11,458	19.7	35.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,008	18.9	34.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,743	15.0	27.0
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	5,540	9.5	17.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,240	3.8	6.9
6	Human immunodeficiency virus (HIV) disease (B20-B24)	2,000	3.4	6.2
7	Cerebrovascular diseases (I60-I69)	1,378	2.4	4.3
8	Assault (homicide) (X85-Y09,Y87.1)	1,273	2.2	3.9
9	Diabetes mellitus (E10-E14)	1,260	2.2	3.9
10	Influenza and pneumonia (J10-J18)	678	1.2	2.1
11	Chronic lower respiratory diseases (J40-J47)	570	1.0	1.8
12	Viral hepatitis (B15-B19)	501	0.9	1.5
13	Septicemia (A40-A41)	498	0.9	1.5
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	419	0.7	1.3
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	359	0.6	1.1
...	All other causes (Residual)	10,306	17.7	31.8
White, non-Hispanic, both sexes, 45-54 years				
...	All causes	111,964	100.0	393.1
1	Malignant neoplasms (C00-C97)	35,861	32.0	125.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	25,153	22.5	88.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,882	7.9	31.2
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	4,774	4.3	16.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	4,637	4.1	16.3
6	Cerebrovascular diseases (I60-I69)	3,433	3.1	12.1
7	Diabetes mellitus (E10-E14)	3,078	2.7	10.8
8	Chronic lower respiratory diseases (J40-J47)	2,463	2.2	8.6
9	Human immunodeficiency virus (HIV) disease (B20-B24)	1,325	1.2	4.7
10	Viral hepatitis (B15-B19)	1,243	1.1	4.4
11	Influenza and pneumonia (J10-J18)	1,165	1.0	4.1
12	Septicemia (A40-A41)	1,092	1.0	3.8
13	Assault (homicide) (X85-Y09,Y87.1)	834	0.7	2.9
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	829	0.7	2.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	516	0.5	1.8
...	All other causes (Residual)	16,679	14.9	58.6

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
White, non-Hispanic, both sexes, 55-64 years				
...	All causes	182,348	100.0	960.0
1	Malignant neoplasms (C00-C97)	70,865	38.9	373.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	47,292	25.9	249.0
3	Chronic lower respiratory diseases (J40-J47)	9,253	5.1	48.7
4	Cerebrovascular diseases (I60-I69)	6,555	3.6	34.5
5	Diabetes mellitus (E10-E14)	5,923	3.2	31.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,753	3.2	30.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	4,235	2.3	22.3
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	2,659	1.5	14.0
9	Influenza and pneumonia (J10-J18)	2,061	1.1	10.9
10	Septicemia (A40-A41)	1,919	1.1	10.1
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,776	1.0	9.4
12	Aortic aneurysm and dissection (I71)	1,201	0.7	6.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,004	0.6	5.3
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	736	0.4	3.9
15	Viral hepatitis (B15-B19)	520	0.3	2.7
...	All other causes (Residual)	20,596	11.3	108.4
White, non-Hispanic, both sexes, 65-74 years				
...	All causes	356,869	100.0	2,409.4
1	Malignant neoplasms (C00-C97)	124,842	35.0	842.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	97,748	27.4	659.9
3	Chronic lower respiratory diseases (J40-J47)	28,122	7.9	189.9
4	Cerebrovascular diseases (I60-I69)	17,989	5.0	121.5
5	Diabetes mellitus (E10-E14)	11,475	3.2	77.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,181	1.7	41.7
7	Influenza and pneumonia (J10-J18)	5,546	1.6	37.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,741	1.3	32.0
9	Chronic liver disease and cirrhosis (K70,K73-K74)	4,285	1.2	28.9
10	Septicemia (A40-A41)	4,173	1.2	28.2
11	Aortic aneurysm and dissection (I71)	3,526	1.0	23.8
12	Alzheimer's disease (G30)	3,004	0.8	20.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,144	0.6	14.5
14	Intentional self-harm (suicide) (X60-X84,Y87.0)	2,075	0.6	14.0
15	Parkinson's disease (G20-G21)	1,895	0.5	12.8
...	All other causes (Residual)	39,123	11.0	264.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
White, non-Hispanic, both sexes, 75-84 years				
...	All causes	604,658	100.0	5,728.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	189,366	31.3	1,794.1
2	Malignant neoplasms (C00-C97)	143,079	23.7	1,355.6
3	Cerebrovascular diseases (I60-I69)	48,556	8.0	460.0
4	Chronic lower respiratory diseases (J40-J47)	44,024	7.3	417.1
5	Influenza and pneumonia (J10-J18)	17,068	2.8	161.7
6	Diabetes mellitus (E10-E14)	17,040	2.8	161.4
7	Alzheimer's disease (G30)	15,704	2.6	148.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,356	1.7	98.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10,026	1.7	95.0
10	Septicemia (A40-A41)	7,901	1.3	74.9
11	Parkinson's disease (G20-G21)	7,106	1.2	67.3
12	Aortic aneurysm and dissection (I71)	5,250	0.9	49.7
13	Pneumonitis due to solids and liquids (J69)	4,793	0.8	45.4
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	4,252	0.7	40.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,171	0.7	39.5
...	All other causes (Residual)	75,966	12.6	719.7
White, non-Hispanic, both sexes, 85 years and over				
...	All causes	583,757	100.0	15,826.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	223,454	38.3	6,058.1
2	Malignant neoplasms (C00-C97)	67,487	11.6	1,829.7
3	Cerebrovascular diseases (I60-I69)	59,738	10.2	1,619.6
4	Influenza and pneumonia (J10-J18)	28,073	4.8	761.1
5	Alzheimer's disease (G30)	26,052	4.5	706.3
6	Chronic lower respiratory diseases (J40-J47)	25,155	4.3	682.0
7	Diabetes mellitus (E10-E14)	10,917	1.9	296.0
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,552	1.8	286.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9,953	1.7	269.8
10	Septicemia (A40-A41)	7,550	1.3	204.7
11	Atherosclerosis (I70)	7,270	1.2	197.1
12	Pneumonitis due to solids and liquids (J69)	7,111	1.2	192.8
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	5,641	1.0	152.9
14	Parkinson's disease (G20-G21)	5,216	0.9	141.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3,237	0.6	87.8
...	All other causes (Residual)	86,351	14.8	2,341.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
White, non-Hispanic, male, all ages				
...	All causes	944,781	100.0	980.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	287,210	30.4	298.2
2	Malignant neoplasms (C00-C97)	235,932	25.0	245.0
3	Chronic lower respiratory diseases (J40-J47)	53,231	5.6	55.3
4	Cerebrovascular diseases (I60-I69)	52,000	5.5	54.0
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	46,470	4.9	48.2
6	Influenza and pneumonia (J10-J18)	23,609	2.5	24.5
7	Diabetes mellitus (E10-E14)	23,465	2.5	24.4
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	19,711	2.1	20.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13,567	1.4	14.1
10	Alzheimer's disease (G30)	13,123	1.4	13.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	12,640	1.3	13.1
12	Septicemia (A40-A41)	10,056	1.1	10.4
13	Aortic aneurysm and dissection (I71)	8,488	0.9	8.8
14	Parkinson's disease (G20-G21)	8,116	0.9	8.4
15	Pneumonitis due to solids and liquids (J69)	7,214	0.8	7.5
...	All other causes (Residual)	129,949	13.8	134.9
White, non-Hispanic, male, 1-4 years				
...	All causes	1,541	100.0	31.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	638	41.4	13.2
2	Malignant neoplasms (C00-C97)	146	9.5	3.0
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	124	8.0	2.6
4	Assault (homicide) (X85-Y09,Y87.1)	86	5.6	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	59	3.8	1.2
6	Septicemia (A40-A41)	26	1.7	0.5
7	Influenza and pneumonia (J10-J18)	24	1.6	0.5
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	1.5	0.5
8	Certain conditions originating in the perinatal period (P00-P96)	23	1.5	0.5
10	Cerebrovascular diseases (I60-I69)	14	0.9	*
11	Meningitis (G00,G03)	12	0.8	*
12	Anemias (D50-D64)	8	0.5	*
12	Chronic lower respiratory diseases (J40-J47)	8	0.5	*
14	Meningococcal infection (A39)	6	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.4	*
...	All other causes (Residual)	338	21.9	7.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
White, non-Hispanic, male, 5-14 years				
...	All causes	2,652	100.0	20.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,176	44.3	9.0
2	Malignant neoplasms (C00-C97)	338	12.7	2.6
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	179	6.7	1.4
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	119	4.5	0.9
5	Assault (homicide) (X85-Y09,Y87.1)	93	3.5	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	87	3.3	0.7
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	31	1.2	0.2
8	Influenza and pneumonia (J10-J18)	29	1.1	0.2
9	Septicemia (A40-A41)	25	0.9	0.2
10	Chronic lower respiratory diseases (J40-J47)	24	0.9	0.2
11	Cerebrovascular diseases (I60-I69)	23	0.9	0.2
12	Certain conditions originating in the perinatal period (P00-P96)	14	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	0.5	*
14	Anemias (D50-D64)	9	0.3	*
14	Meningitis (G00,G03)	9	0.3	*
...	All other causes (Residual)	484	18.3	3.7
White, non-Hispanic, male, 15-24 years				
...	All causes	13,066	100.0	100.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,065	54.1	54.4
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	2,467	18.9	19.0
3	Malignant neoplasms (C00-C97)	636	4.9	4.9
4	Assault (homicide) (X85-Y09,Y87.1)	588	4.5	4.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	326	2.5	2.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	163	1.2	1.3
7	Influenza and pneumonia (J10-J18)	68	0.5	0.5
8	Cerebrovascular diseases (I60-I69)	67	0.5	0.5
9	Diabetes mellitus (E10-E14)	57	0.4	0.4
10	Chronic lower respiratory diseases (J40-J47)	52	0.4	0.4
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	40	0.3	0.3
12	Septicemia (A40-A41)	33	0.3	0.3
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	25	0.2	0.2
13	Legal intervention (Y35,Y89.0)	25	0.2	0.2
15	Meningococcal infection (A39)	22	0.2	0.2
...	All other causes (Residual)	1,432	11.0	11.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
White, non-Hispanic, male, 25-34 years				
...	All causes	15,934	100.0	127.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,861	36.8	46.8
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	2,989	18.8	23.9
3	Malignant neoplasms (C00-C97)	1,259	7.9	10.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,115	7.0	8.9
5	Assault (homicide) (X85-Y09,Y87.1)	677	4.2	5.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	475	3.0	3.8
7	Diabetes mellitus (E10-E14)	220	1.4	1.8
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	203	1.3	1.6
9	Cerebrovascular diseases (I60-I69)	161	1.0	1.3
10	Chronic liver disease and cirrhosis (K70,K73-K74)	140	0.9	1.1
11	Influenza and pneumonia (J10-J18)	111	0.7	0.9
12	Septicemia (A40-A41)	74	0.5	0.6
13	Chronic lower respiratory diseases (J40-J47)	71	0.4	0.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	62	0.4	0.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	55	0.3	0.4
15	Legal intervention (Y35,Y89.0)	55	0.3	0.4
...	All other causes (Residual)	2,406	15.1	19.2
White, non-Hispanic, male, 35-44 years				
...	All causes	37,622	100.0	231.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,113	21.6	50.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,446	17.1	39.7
3	Malignant neoplasms (C00-C97)	5,201	13.8	32.1
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	4,287	11.4	26.4
5	Human immunodeficiency virus (HIV) disease (B20-B24)	1,678	4.5	10.3
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1,518	4.0	9.4
7	Assault (homicide) (X85-Y09,Y87.1)	836	2.2	5.2
8	Diabetes mellitus (E10-E14)	766	2.0	4.7
9	Cerebrovascular diseases (I60-I69)	694	1.8	4.3
10	Influenza and pneumonia (J10-J18)	394	1.0	2.4
11	Viral hepatitis (B15-B19)	342	0.9	2.1
12	Chronic lower respiratory diseases (J40-J47)	267	0.7	1.6
13	Septicemia (A40-A41)	262	0.7	1.6
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	233	0.6	1.4
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	206	0.5	1.3
...	All other causes (Residual)	6,379	17.0	39.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
White, non-Hispanic, male, 45-54 years				
...	All causes	70,788	100.0	501.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	19,101	27.0	135.4
2	Malignant neoplasms (C00-C97)	18,092	25.6	128.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,521	9.2	46.2
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	3,627	5.1	25.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	3,477	4.9	24.6
6	Cerebrovascular diseases (I60-I69)	1,856	2.6	13.2
7	Diabetes mellitus (E10-E14)	1,842	2.6	13.1
8	Chronic lower respiratory diseases (J40-J47)	1,259	1.8	8.9
9	Human immunodeficiency virus (HIV) disease (B20-B24)	1,161	1.6	8.2
10	Viral hepatitis (B15-B19)	958	1.4	6.8
11	Influenza and pneumonia (J10-J18)	667	0.9	4.7
12	Septicemia (A40-A41)	613	0.9	4.3
13	Assault (homicide) (X85-Y09,Y87.1)	560	0.8	4.0
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	491	0.7	3.5
15	Aortic aneurysm and dissection (I71)	355	0.5	2.5
...	All other causes (Residual)	10,208	14.4	72.4
White, non-Hispanic, male, 55-64 years				
...	All causes	109,031	100.0	1,185.7
1	Malignant neoplasms (C00-C97)	38,368	35.2	417.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	33,179	30.4	360.8
3	Chronic lower respiratory diseases (J40-J47)	4,728	4.3	51.4
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,871	3.6	42.1
5	Cerebrovascular diseases (I60-I69)	3,605	3.3	39.2
6	Diabetes mellitus (E10-E14)	3,298	3.0	35.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,935	2.7	31.9
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	2,031	1.9	22.1
9	Influenza and pneumonia (J10-J18)	1,176	1.1	12.8
10	Septicemia (A40-A41)	1,015	0.9	11.0
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	990	0.9	10.8
12	Aortic aneurysm and dissection (I71)	914	0.8	9.9
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	539	0.5	5.9
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	434	0.4	4.7
15	Human immunodeficiency virus (HIV) disease (B20-B24)	369	0.3	4.0
...	All other causes (Residual)	11,579	10.6	125.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
	White, non-Hispanic, male, 65-74 years			
...	All causes	202,038	100.0	2,986.8
1	Malignant neoplasms (C00-C97)	69,085	34.2	1,021.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	61,024	30.2	902.1
3	Chronic lower respiratory diseases (J40-J47)	14,398	7.1	212.8
4	Cerebrovascular diseases (I60-I69)	9,176	4.5	135.7
5	Diabetes mellitus (E10-E14)	5,967	3.0	88.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,692	1.8	54.6
7	Influenza and pneumonia (J10-J18)	3,084	1.5	45.6
8	Chronic liver disease and cirrhosis (K70,K73-K74)	2,602	1.3	38.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,574	1.3	38.1
10	Aortic aneurysm and dissection (I71)	2,453	1.2	36.3
11	Septicemia (A40-A41)	2,131	1.1	31.5
12	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,701	0.8	25.1
13	Alzheimer's disease (G30)	1,367	0.7	20.2
14	Parkinson's disease (G20-G21)	1,214	0.6	17.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,202	0.6	17.8
...	All other causes (Residual)	20,368	10.1	301.1
	White, non-Hispanic, male, 75-84 years			
...	All causes	294,842	100.0	6,913.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	95,576	32.4	2,241.0
2	Malignant neoplasms (C00-C97)	73,952	25.1	1,734.0
3	Chronic lower respiratory diseases (J40-J47)	21,703	7.4	508.9
4	Cerebrovascular diseases (I60-I69)	20,379	6.9	477.8
5	Influenza and pneumonia (J10-J18)	8,346	2.8	195.7
6	Diabetes mellitus (E10-E14)	7,796	2.6	182.8
7	Alzheimer's disease (G30)	5,686	1.9	133.3
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,292	1.8	124.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,141	1.7	120.5
10	Parkinson's disease (G20-G21)	4,191	1.4	98.3
11	Septicemia (A40-A41)	3,579	1.2	83.9
12	Aortic aneurysm and dissection (I71)	3,136	1.1	73.5
13	Pneumonitis due to solids and liquids (J69)	2,648	0.9	62.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,108	0.7	49.4
15	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,761	0.6	41.3
...	All other causes (Residual)	33,548	11.4	786.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
White, non-Hispanic, male, 85 years and over				
...	All causes	189,537	100.0	17,269.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	70,138	37.0	6,390.4
2	Malignant neoplasms (C00-C97)	28,814	15.2	2,625.3
3	Cerebrovascular diseases (I60-I69)	15,987	8.4	1,456.6
4	Chronic lower respiratory diseases (J40-J47)	10,708	5.6	975.6
5	Influenza and pneumonia (J10-J18)	9,641	5.1	878.4
6	Alzheimer's disease (G30)	5,857	3.1	533.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,023	2.1	366.5
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,968	2.1	361.5
9	Diabetes mellitus (E10-E14)	3,508	1.9	319.6
10	Pneumonitis due to solids and liquids (J69)	3,059	1.6	278.7
11	Parkinson's disease (G20-G21)	2,533	1.3	230.8
12	Septicemia (A40-A41)	2,242	1.2	204.3
13	Atherosclerosis (I70)	1,949	1.0	177.6
14	Aortic aneurysm and dissection (I71)	1,436	0.8	130.8
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,399	0.7	127.5
...	All other causes (Residual)	24,275	12.8	2,211.7
White, non-Hispanic, female, all ages				
...	All causes	1,015,138	100.0	1,011.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	307,255	30.3	306.2
2	Malignant neoplasms (C00-C97)	222,268	21.9	221.5
3	Cerebrovascular diseases (I60-I69)	86,210	8.5	85.9
4	Chronic lower respiratory diseases (J40-J47)	56,670	5.6	56.5
5	Alzheimer's disease (G30)	32,123	3.2	32.0
6	Influenza and pneumonia (J10-J18)	31,526	3.1	31.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	27,066	2.7	27.0
8	Diabetes mellitus (E10-E14)	26,698	2.6	26.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14,357	1.4	14.3
10	Septicemia (A40-A41)	13,461	1.3	13.4
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	8,196	0.8	8.2
12	Atherosclerosis (I70)	8,052	0.8	8.0
13	Pneumonitis due to solids and liquids (J69)	7,129	0.7	7.1
14	Chronic liver disease and cirrhosis (K70,K73-K74)	7,099	0.7	7.1
15	Parkinson's disease (G20-G21)	6,381	0.6	6.4
...	All other causes (Residual)	160,647	15.8	160.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
White, non-Hispanic, female, 1-4 years				
...	All causes	1,100	100.0	23.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	395	35.9	8.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	116	10.5	2.5
3	Malignant neoplasms (C00-C97)	110	10.0	2.4
4	Assault (homicide) (X85-Y09,Y87.1)	58	5.3	1.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	39	3.5	0.8
6	Influenza and pneumonia (J10-J18)	27	2.5	0.6
7	Septicemia (A40-A41)	25	2.3	0.5
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	1.3	*
9	Cerebrovascular diseases (I60-I69)	11	1.0	*
9	Chronic lower respiratory diseases (J40-J47)	11	1.0	*
9	Certain conditions originating in the perinatal period (P00-P96)	11	1.0	*
12	Meningitis (G00,G03)	7	0.6	*
13	Anemias (D50-D64)	6	0.5	*
14	Meningococcal infection (A39)	5	0.5	*
15	Diabetes mellitus (E10-E14)	4	0.4	*
...	All other causes (Residual)	261	23.7	5.7
White, non-Hispanic, female, 5-14 years				
...	All causes	1,745	100.0	14.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	667	38.2	5.4
2	Malignant neoplasms (C00-C97)	289	16.6	2.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	121	6.9	1.0
4	Assault (homicide) (X85-Y09,Y87.1)	71	4.1	0.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	53	3.0	0.4
6	Intentional self-harm (suicide) (X60-X84,Y87.0)	46	2.6	0.4
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	32	1.8	0.3
8	Cerebrovascular diseases (I60-I69)	23	1.3	0.2
8	Chronic lower respiratory diseases (J40-J47)	23	1.3	0.2
10	Influenza and pneumonia (J10-J18)	20	1.1	0.2
11	Septicemia (A40-A41)	17	1.0	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	10	0.6	*
13	Meningitis (G00,G03)	9	0.5	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	8	0.5	*
14	Diseases of appendix (K35-K38)	8	0.5	*
...	All other causes (Residual)	348	19.9	2.8

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
White, non-Hispanic, female, 15-24 years				
...	All causes	5,195	100.0	42.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,670	51.4	21.7
2	Malignant neoplasms (C00-C97)	423	8.1	3.4
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	401	7.7	3.3
4	Assault (homicide) (X85-Y09,Y87.1)	286	5.5	2.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	205	3.9	1.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	116	2.2	0.9
7	Cerebrovascular diseases (I60-I69)	51	1.0	0.4
8	Influenza and pneumonia (J10-J18)	48	0.9	0.4
9	Diabetes mellitus (E10-E14)	35	0.7	0.3
9	Chronic lower respiratory diseases (J40-J47)	35	0.7	0.3
11	Pregnancy, childbirth and the puerperium (O00-099)	33	0.6	0.3
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	0.4	0.2
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	0.4	0.2
14	Meningococcal infection (A39)	15	0.3	*
14	Septicemia (A40-A41)	15	0.3	*
...	All other causes (Residual)	816	15.7	6.6
White, non-Hispanic, female, 25-34 years				
...	All causes	7,257	100.0	57.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,869	25.8	14.9
2	Malignant neoplasms (C00-C97)	1,340	18.5	10.7
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	669	9.2	5.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	538	7.4	4.3
5	Assault (homicide) (X85-Y09,Y87.1)	346	4.8	2.8
6	Cerebrovascular diseases (I60-I69)	152	2.1	1.2
7	Human immunodeficiency virus (HIV) disease (B20-B24)	144	2.0	1.1
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	141	1.9	1.1
9	Diabetes mellitus (E10-E14)	137	1.9	1.1
10	Influenza and pneumonia (J10-J18)	102	1.4	0.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	77	1.1	0.6
12	Chronic lower respiratory diseases (J40-J47)	73	1.0	0.6
13	Pregnancy, childbirth and the puerperium (O00-099)	72	1.0	0.6
14	Septicemia (A40-A41)	60	0.8	0.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	50	0.7	0.4
...	All other causes (Residual)	1,487	20.5	11.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
White, non-Hispanic, female, 35-44 years				
...	All causes	20,609	100.0	127.4
1	Malignant neoplasms (C00-C97)	6,257	30.4	38.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,895	14.0	17.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,297	11.1	14.2
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,253	6.1	7.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	722	3.5	4.5
6	Cerebrovascular diseases (I60-I69)	684	3.3	4.2
7	Diabetes mellitus (E10-E14)	494	2.4	3.1
8	Assault (homicide) (X85-Y09,Y87.1)	437	2.1	2.7
9	Human immunodeficiency virus (HIV) disease (B20-B24)	322	1.6	2.0
10	Chronic lower respiratory diseases (J40-J47)	303	1.5	1.9
11	Influenza and pneumonia (J10-J18)	284	1.4	1.8
12	Septicemia (A40-A41)	236	1.1	1.5
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	186	0.9	1.1
14	Viral hepatitis (B15-B19)	159	0.8	1.0
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	153	0.7	0.9
...	All other causes (Residual)	3,927	19.1	24.3
White, non-Hispanic, female, 45-54 years				
...	All causes	41,176	100.0	286.4
1	Malignant neoplasms (C00-C97)	17,769	43.2	123.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,052	14.7	42.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,361	5.7	16.4
4	Cerebrovascular diseases (I60-I69)	1,577	3.8	11.0
5	Diabetes mellitus (E10-E14)	1,236	3.0	8.6
6	Chronic lower respiratory diseases (J40-J47)	1,204	2.9	8.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,160	2.8	8.1
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,147	2.8	8.0
9	Influenza and pneumonia (J10-J18)	498	1.2	3.5
10	Septicemia (A40-A41)	479	1.2	3.3
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	338	0.8	2.4
12	Viral hepatitis (B15-B19)	285	0.7	2.0
13	Assault (homicide) (X85-Y09,Y87.1)	274	0.7	1.9
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	225	0.5	1.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	215	0.5	1.5
...	All other causes (Residual)	6,356	15.4	44.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
White, non-Hispanic, female, 55-64 years				
...	All causes	73,317	100.0	748.2
1	Malignant neoplasms (C00-C97)	32,497	44.3	331.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,113	19.2	144.0
3	Chronic lower respiratory diseases (J40-J47)	4,525	6.2	46.2
4	Cerebrovascular diseases (I60-I69)	2,950	4.0	30.1
5	Diabetes mellitus (E10-E14)	2,625	3.6	26.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,882	2.6	19.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,300	1.8	13.3
8	Septicemia (A40-A41)	904	1.2	9.2
9	Influenza and pneumonia (J10-J18)	885	1.2	9.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	786	1.1	8.0
11	Intentional self-harm (suicide) (X60-X84,Y87.0)	628	0.9	6.4
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	465	0.6	4.7
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	302	0.4	3.1
14	Aortic aneurysm and dissection (I71)	287	0.4	2.9
15	Alzheimer's disease (G30)	245	0.3	2.5
...	All other causes (Residual)	8,923	12.2	91.1
White, non-Hispanic, female, 65-74 years				
...	All causes	154,831	100.0	1,924.0
1	Malignant neoplasms (C00-C97)	55,757	36.0	692.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	36,724	23.7	456.4
3	Chronic lower respiratory diseases (J40-J47)	13,724	8.9	170.5
4	Cerebrovascular diseases (I60-I69)	8,813	5.7	109.5
5	Diabetes mellitus (E10-E14)	5,508	3.6	68.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,489	1.6	30.9
7	Influenza and pneumonia (J10-J18)	2,462	1.6	30.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,167	1.4	26.9
9	Septicemia (A40-A41)	2,042	1.3	25.4
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,683	1.1	20.9
11	Alzheimer's disease (G30)	1,637	1.1	20.3
12	Aortic aneurysm and dissection (I71)	1,073	0.7	13.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	942	0.6	11.7
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	892	0.6	11.1
15	Parkinson's disease (G20-G21)	681	0.4	8.5
...	All other causes (Residual)	18,237	11.8	226.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
	White, non-Hispanic, female, 75-84 years			
...	All causes	309,816	100.0	4,925.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	93,790	30.3	1,491.1
2	Malignant neoplasms (C00-C97)	69,127	22.3	1,099.0
3	Cerebrovascular diseases (I60-I69)	28,177	9.1	448.0
4	Chronic lower respiratory diseases (J40-J47)	22,321	7.2	354.9
5	Alzheimer's disease (G30)	10,018	3.2	159.3
6	Diabetes mellitus (E10-E14)	9,244	3.0	147.0
7	Influenza and pneumonia (J10-J18)	8,722	2.8	138.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,064	1.6	80.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,885	1.6	77.7
10	Septicemia (A40-A41)	4,322	1.4	68.7
11	Parkinson's disease (G20-G21)	2,915	0.9	46.3
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,581	0.8	41.0
13	Pneumonitis due to solids and liquids (J69)	2,145	0.7	34.1
14	Aortic aneurysm and dissection (I71)	2,114	0.7	33.6
15	Atherosclerosis (I70)	2,066	0.7	32.8
...	All other causes (Residual)	42,325	13.7	672.9
	White, non-Hispanic, female, 85 years and over			
...	All causes	394,220	100.0	15,215.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	153,316	38.9	5,917.4
2	Cerebrovascular diseases (I60-I69)	43,751	11.1	1,688.6
3	Malignant neoplasms (C00-C97)	38,673	9.8	1,492.6
4	Alzheimer's disease (G30)	20,195	5.1	779.4
5	Influenza and pneumonia (J10-J18)	18,432	4.7	711.4
6	Chronic lower respiratory diseases (J40-J47)	14,447	3.7	557.6
7	Diabetes mellitus (E10-E14)	7,409	1.9	286.0
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,584	1.7	254.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,930	1.5	228.9
10	Atherosclerosis (I70)	5,321	1.3	205.4
11	Septicemia (A40-A41)	5,308	1.3	204.9
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	4,242	1.1	163.7
13	Pneumonitis due to solids and liquids (J69)	4,052	1.0	156.4
14	Parkinson's disease (G20-G21)	2,683	0.7	103.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,036	0.5	78.6
...	All other causes (Residual)	61,841	15.7	2,386.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Black, non-Hispanic, both sexes, all ages				
...	All causes	282,676	100.0	844.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	76,706	27.1	229.1
2	Malignant neoplasms (C00-C97)	61,390	21.7	183.4
3	Cerebrovascular diseases (I60-I69)	19,053	6.7	56.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,087	4.3	36.1
5	Diabetes mellitus (E10-E14)	11,918	4.2	35.6
6	Assault (homicide) (X85-Y09,Y87.1)	7,748	2.7	23.1
7	Human immunodeficiency virus (HIV) disease (B20-B24)	7,708	2.7	23.0
8	Chronic lower respiratory diseases (J40-J47)	7,528	2.7	22.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,853	2.4	20.5
10	Influenza and pneumonia (J10-J18)	5,920	2.1	17.7
11	Septicemia (A40-A41)	5,863	2.1	17.5
12	Certain conditions originating in the perinatal period (P00-P96)	4,878	1.7	14.6
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	3,862	1.4	11.5
14	Alzheimer's disease (G30)	2,719	1.0	8.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,705	1.0	8.1
...	All other causes (Residual)	45,738	16.2	136.6
Black, non-Hispanic, both sexes, 1-4 years				
...	All causes	1,217	100.0	58.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	396	32.5	19.1
2	Assault (homicide) (X85-Y09,Y87.1)	143	11.8	6.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	123	10.1	5.9
4	Malignant neoplasms (C00-C97)	66	5.4	3.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	47	3.9	2.3
6	Influenza and pneumonia (J10-J18)	29	2.4	1.4
7	Certain conditions originating in the perinatal period (P00-P96)	27	2.2	1.3
8	Septicemia (A40-A41)	25	2.1	1.2
9	Chronic lower respiratory diseases (J40-J47)	21	1.7	1.0
10	Anemias (D50-D64)	19	1.6	*
11	Meningitis (G00,G03)	13	1.1	*
11	Cerebrovascular diseases (I60-I69)	13	1.1	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	12	1.0	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.7	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	8	0.7	*
...	All other causes (Residual)	266	21.9	12.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total		
		Number	deaths	Rate
Black, non-Hispanic, both sexes, 5-14 years				
...	All causes	1,620	100.0	27.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	619	38.2	10.5
2	Malignant neoplasms (C00-C97)	147	9.1	2.5
3	Assault (homicide) (X85-Y09,Y87.1)	122	7.5	2.1
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	89	5.5	1.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	75	4.6	1.3
6	Chronic lower respiratory diseases (J40-J47)	70	4.3	1.2
7	Intentional self-harm (suicide) (X60-X84,Y87.0)	39	2.4	0.7
8	Anemias (D50-D64)	30	1.9	0.5
9	Human immunodeficiency virus (HIV) disease (B20-B24)	23	1.4	0.4
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	1.1	*
11	Cerebrovascular diseases (I60-I69)	15	0.9	*
12	Septicemia (A40-A41)	11	0.7	*
12	Influenza and pneumonia (J10-J18)	11	0.7	*
14	Certain conditions originating in the perinatal period (P00-P96)	10	0.6	*
15	Diabetes mellitus (E10-E14)	9	0.6	*
15	Meningitis (G00,G03)	9	0.6	*
...	All other causes (Residual)	323	19.9	5.5
Black, non-Hispanic, both sexes, 15-24 years				
...	All causes	6,926	100.0	125.2
1	Assault (homicide) (X85-Y09,Y87.1)	2,771	40.0	50.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,662	24.0	30.0
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	471	6.8	8.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	331	4.8	6.0
5	Malignant neoplasms (C00-C97)	297	4.3	5.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	125	1.8	2.3
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	91	1.3	1.6
8	Chronic lower respiratory diseases (J40-J47)	80	1.2	1.4
9	Anemias (D50-D64)	61	0.9	1.1
10	Diabetes mellitus (E10-E14)	55	0.8	1.0
11	Influenza and pneumonia (J10-J18)	44	0.6	0.8
12	Cerebrovascular diseases (I60-I69)	42	0.6	0.8
13	Pregnancy, childbirth and the puerperium (O00-099)	34	0.5	0.6
14	Septicemia (A40-A41)	33	0.5	0.6
15	Legal intervention (Y35,Y89.0)	27	0.4	0.5
...	All other causes (Residual)	802	11.6	14.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Black, non-Hispanic, both sexes, 25-34 years				
...	All causes	10,103	100.0	204.2
1	Assault (homicide) (X85-Y09,Y87.1)	2,193	21.7	44.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,740	17.2	35.2
3	Human immunodeficiency virus (HIV) disease (B20-B24)	1,414	14.0	28.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	929	9.2	18.8
5	Malignant neoplasms (C00-C97)	641	6.3	13.0
6	Intentional self-harm (suicide) (X60-X84,Y87.0)	469	4.6	9.5
7	Diabetes mellitus (E10-E14)	187	1.9	3.8
8	Cerebrovascular diseases (I60-I69)	158	1.6	3.2
9	Anemias (D50-D64)	124	1.2	2.5
10	Chronic lower respiratory diseases (J40-J47)	114	1.1	2.3
11	Septicemia (A40-A41)	107	1.1	2.2
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	99	1.0	2.0
13	Influenza and pneumonia (J10-J18)	92	0.9	1.9
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	74	0.7	1.5
15	Pregnancy, childbirth and the puerperium (O00-099)	65	0.6	1.3
...	All other causes (Residual)	1,697	16.8	34.3
Black, non-Hispanic, both sexes, 35-44 years				
...	All causes	20,279	100.0	374.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,259	16.1	60.2
2	Malignant neoplasms (C00-C97)	3,072	15.1	56.8
3	Human immunodeficiency virus (HIV) disease (B20-B24)	3,028	14.9	56.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,253	11.1	41.6
5	Assault (homicide) (X85-Y09,Y87.1)	1,359	6.7	25.1
6	Cerebrovascular diseases (I60-I69)	829	4.1	15.3
7	Diabetes mellitus (E10-E14)	458	2.3	8.5
8	Chronic liver disease and cirrhosis (K70,K73-K74)	446	2.2	8.2
9	Intentional self-harm (suicide) (X60-X84,Y87.0)	427	2.1	7.9
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	305	1.5	5.6
11	Chronic lower respiratory diseases (J40-J47)	272	1.3	5.0
12	Septicemia (A40-A41)	269	1.3	5.0
13	Influenza and pneumonia (J10-J18)	254	1.3	4.7
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	176	0.9	3.3
15	Anemias (D50-D64)	139	0.7	2.6
...	All other causes (Residual)	3,733	18.4	69.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Black, non-Hispanic, both sexes, 45-54 years				
...	All causes	32,911	100.0	834.2
1	Malignant neoplasms (C00-C97)	8,068	24.5	204.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,606	23.1	192.8
3	Human immunodeficiency virus (HIV) disease (B20-B24)	2,211	6.7	56.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,961	6.0	49.7
5	Cerebrovascular diseases (I60-I69)	1,819	5.5	46.1
6	Diabetes mellitus (E10-E14)	1,270	3.9	32.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	929	2.8	23.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	664	2.0	16.8
9	Assault (homicide) (X85-Y09,Y87.1)	615	1.9	15.6
10	Chronic lower respiratory diseases (J40-J47)	593	1.8	15.0
11	Septicemia (A40-A41)	572	1.7	14.5
12	Influenza and pneumonia (J10-J18)	423	1.3	10.7
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	405	1.2	10.3
14	Viral hepatitis (B15-B19)	329	1.0	8.3
15	Intentional self-harm (suicide) (X60-X84,Y87.0)	266	0.8	6.7
...	All other causes (Residual)	5,180	15.7	131.3
Black, non-Hispanic, both sexes, 55-64 years				
...	All causes	38,867	100.0	1,681.8
1	Malignant neoplasms (C00-C97)	12,019	30.9	520.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,257	29.0	487.1
3	Cerebrovascular diseases (I60-I69)	2,265	5.8	98.0
4	Diabetes mellitus (E10-E14)	1,995	5.1	86.3
5	Chronic lower respiratory diseases (J40-J47)	1,063	2.7	46.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	983	2.5	42.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	980	2.5	42.4
8	Septicemia (A40-A41)	756	1.9	32.7
9	Human immunodeficiency virus (HIV) disease (B20-B24)	625	1.6	27.0
10	Chronic liver disease and cirrhosis (K70,K73-K74)	622	1.6	26.9
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	555	1.4	24.0
12	Influenza and pneumonia (J10-J18)	531	1.4	23.0
13	Assault (homicide) (X85-Y09,Y87.1)	201	0.5	8.7
14	Aortic aneurysm and dissection (I71)	196	0.5	8.5
15	Viral hepatitis (B15-B19)	138	0.4	6.0
...	All other causes (Residual)	4,681	12.0	202.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
	Black, non-Hispanic, both sexes, 65-74 years			
...	All causes	54,645	100.0	3,345.3
1	Malignant neoplasms (C00-C97)	16,485	30.2	1,009.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	16,413	30.0	1,004.8
3	Cerebrovascular diseases (I60-I69)	3,660	6.7	224.1
4	Diabetes mellitus (E10-E14)	3,083	5.6	188.7
5	Chronic lower respiratory diseases (J40-J47)	1,993	3.6	122.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,637	3.0	100.2
7	Septicemia (A40-A41)	1,173	2.1	71.8
8	Influenza and pneumonia (J10-J18)	984	1.8	60.2
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	857	1.6	52.5
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	801	1.5	49.0
11	Chronic liver disease and cirrhosis (K70,K73-K74)	429	0.8	26.3
12	Pneumonitis due to solids and liquids (J69)	289	0.5	17.7
13	Aortic aneurysm and dissection (I71)	279	0.5	17.1
14	Alzheimer's disease (G30)	261	0.5	16.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	245	0.4	15.0
...	All other causes (Residual)	6,056	11.1	370.7
	Black, non-Hispanic, both sexes, 75-84 years			
...	All causes	60,874	100.0	6,983.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	19,600	32.2	2,248.6
2	Malignant neoplasms (C00-C97)	14,244	23.4	1,634.2
3	Cerebrovascular diseases (I60-I69)	5,344	8.8	613.1
4	Diabetes mellitus (E10-E14)	3,193	5.2	366.3
5	Chronic lower respiratory diseases (J40-J47)	2,128	3.5	244.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,776	2.9	203.8
7	Septicemia (A40-A41)	1,576	2.6	180.8
8	Influenza and pneumonia (J10-J18)	1,550	2.5	177.8
9	Alzheimer's disease (G30)	1,004	1.6	115.2
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	980	1.6	112.4
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	768	1.3	88.1
12	Pneumonitis due to solids and liquids (J69)	468	0.8	53.7
13	Aortic aneurysm and dissection (I71)	331	0.5	38.0
14	Atherosclerosis (I70)	320	0.5	36.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	282	0.5	32.4
...	All other causes (Residual)	7,310	12.0	838.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
	Black, non-Hispanic, both sexes, 85 years and over			
...	All causes	46,660	100.0	15,010.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	17,059	36.6	5,488.0
2	Malignant neoplasms (C00-C97)	6,337	13.6	2,038.6
3	Cerebrovascular diseases (I60-I69)	4,861	10.4	1,563.8
4	Influenza and pneumonia (J10-J18)	1,901	4.1	611.6
5	Diabetes mellitus (E10-E14)	1,667	3.6	536.3
6	Alzheimer's disease (G30)	1,418	3.0	456.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,290	2.8	415.0
8	Septicemia (A40-A41)	1,245	2.7	400.5
9	Chronic lower respiratory diseases (J40-J47)	1,183	2.5	380.6
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	899	1.9	289.2
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	576	1.2	185.3
12	Pneumonitis due to solids and liquids (J69)	505	1.1	162.5
13	Atherosclerosis (I70)	441	0.9	141.9
14	Nutritional deficiencies (E40-E64)	212	0.5	68.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	202	0.4	65.0
...	All other causes (Residual)	6,864	14.7	2,208.2
	Black, non-Hispanic, male, all ages			
...	All causes	143,297	100.0	903.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	36,282	25.3	228.7
2	Malignant neoplasms (C00-C97)	32,493	22.7	204.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,380	5.8	52.8
4	Cerebrovascular diseases (I60-I69)	7,945	5.5	50.1
5	Assault (homicide) (X85-Y09,Y87.1)	6,382	4.5	40.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	5,291	3.7	33.4
7	Diabetes mellitus (E10-E14)	4,720	3.3	29.8
8	Chronic lower respiratory diseases (J40-J47)	4,184	2.9	26.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,045	2.1	19.2
10	Influenza and pneumonia (J10-J18)	2,880	2.0	18.2
11	Certain conditions originating in the perinatal period (P00-P96)	2,760	1.9	17.4
12	Septicemia (A40-A41)	2,559	1.8	16.1
13	Chronic liver disease and cirrhosis (K70,K73-K74)	1,709	1.2	10.8
14	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,601	1.1	10.1
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,576	1.1	9.9
...	All other causes (Residual)	21,490	15.0	135.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Black, non-Hispanic, male, 1-4 years				
...	All causes	672	100.0	63.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	232	34.5	22.1
2	Assault (homicide) (X85-Y09,Y87.1)	73	10.9	6.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	67	10.0	6.4
4	Malignant neoplasms (C00-C97)	31	4.6	2.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	24	3.6	2.3
6	Influenza and pneumonia (J10-J18)	20	3.0	1.9
7	Certain conditions originating in the perinatal period (P00-P96)	18	2.7	*
8	Chronic lower respiratory diseases (J40-J47)	14	2.1	*
9	Septicemia (A40-A41)	13	1.9	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	1.2	*
10	Anemias (D50-D64)	8	1.2	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	6	0.9	*
12	Meningitis (G00,G03)	6	0.9	*
14	Meningococcal infection (A39)	4	0.6	*
15	Cerebrovascular diseases (I60-I69)	3	0.4	*
...	All other causes (Residual)	145	21.6	13.8
Black, non-Hispanic, male, 5-14 years				
...	All causes	959	100.0	32.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	406	42.3	13.6
2	Malignant neoplasms (C00-C97)	86	9.0	2.9
3	Assault (homicide) (X85-Y09,Y87.1)	70	7.3	2.3
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	50	5.2	1.7
5	Chronic lower respiratory diseases (J40-J47)	43	4.5	1.4
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	41	4.3	1.4
7	Intentional self-harm (suicide) (X60-X84,Y87.0)	34	3.5	1.1
8	Anemias (D50-D64)	17	1.8	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	12	1.3	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.8	*
11	Influenza and pneumonia (J10-J18)	7	0.7	*
12	Septicemia (A40-A41)	6	0.6	*
12	Meningitis (G00,G03)	6	0.6	*
14	Cerebrovascular diseases (I60-I69)	5	0.5	*
15	Diabetes mellitus (E10-E14)	4	0.4	*
...	All other causes (Residual)	164	17.1	5.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Black, non-Hispanic, male, 15-24 years				
...	All causes	5,224	100.0	188.2
1	Assault (homicide) (X85-Y09,Y87.1)	2,459	47.1	88.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,233	23.6	44.4
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	407	7.8	14.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	192	3.7	6.9
5	Malignant neoplasms (C00-C97)	178	3.4	6.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	60	1.1	2.2
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	58	1.1	2.1
8	Chronic lower respiratory diseases (J40-J47)	51	1.0	1.8
9	Anemias (D50-D64)	29	0.6	1.0
10	Diabetes mellitus (E10-E14)	28	0.5	1.0
11	Legal intervention (Y35,Y89.0)	27	0.5	1.0
12	Influenza and pneumonia (J10-J18)	21	0.4	0.8
13	Cerebrovascular diseases (I60-I69)	19	0.4	*
14	Septicemia (A40-A41)	18	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	0.2	*
...	All other causes (Residual)	434	8.3	15.6
Black, non-Hispanic, male, 25-34 years				
...	All causes	6,649	100.0	284.0
1	Assault (homicide) (X85-Y09,Y87.1)	1,883	28.3	80.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,316	19.8	56.2
3	Human immunodeficiency virus (HIV) disease (B20-B24)	822	12.4	35.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	543	8.2	23.2
5	Intentional self-harm (suicide) (X60-X84,Y87.0)	397	6.0	17.0
6	Malignant neoplasms (C00-C97)	259	3.9	11.1
7	Diabetes mellitus (E10-E14)	98	1.5	4.2
8	Cerebrovascular diseases (I60-I69)	70	1.1	3.0
8	Chronic lower respiratory diseases (J40-J47)	70	1.1	3.0
10	Septicemia (A40-A41)	59	0.9	2.5
11	Influenza and pneumonia (J10-J18)	58	0.9	2.5
12	Anemias (D50-D64)	55	0.8	2.3
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	50	0.8	2.1
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	43	0.6	1.8
15	Legal intervention (Y35,Y89.0)	33	0.5	1.4
...	All other causes (Residual)	893	13.4	38.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Black, non-Hispanic, male, 35-44 years				
...	All causes	12,061	100.0	476.4
1	Human immunodeficiency virus (HIV) disease (B20-B24)	2,060	17.1	81.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,972	16.4	77.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,649	13.7	65.1
4	Malignant neoplasms (C00-C97)	1,292	10.7	51.0
5	Assault (homicide) (X85-Y09,Y87.1)	1,027	8.5	40.6
6	Cerebrovascular diseases (I60-I69)	394	3.3	15.6
7	Intentional self-harm (suicide) (X60-X84,Y87.0)	346	2.9	13.7
8	Chronic liver disease and cirrhosis (K70,K73-K74)	279	2.3	11.0
9	Diabetes mellitus (E10-E14)	252	2.1	10.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	175	1.5	6.9
11	Influenza and pneumonia (J10-J18)	151	1.3	6.0
12	Chronic lower respiratory diseases (J40-J47)	129	1.1	5.1
13	Septicemia (A40-A41)	125	1.0	4.9
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	98	0.8	3.9
15	Viral hepatitis (B15-B19)	77	0.6	3.0
...	All other causes (Residual)	2,035	16.9	80.4
Black, non-Hispanic, male, 45-54 years				
...	All causes	19,647	100.0	1,099.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,787	24.4	267.8
2	Malignant neoplasms (C00-C97)	4,152	21.1	232.3
3	Human immunodeficiency virus (HIV) disease (B20-B24)	1,642	8.4	91.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,454	7.4	81.3
5	Cerebrovascular diseases (I60-I69)	955	4.9	53.4
6	Diabetes mellitus (E10-E14)	648	3.3	36.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	633	3.2	35.4
8	Assault (homicide) (X85-Y09,Y87.1)	498	2.5	27.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	394	2.0	22.0
10	Septicemia (A40-A41)	319	1.6	17.8
11	Chronic lower respiratory diseases (J40-J47)	289	1.5	16.2
12	Influenza and pneumonia (J10-J18)	271	1.4	15.2
13	Viral hepatitis (B15-B19)	234	1.2	13.1
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	229	1.2	12.8
15	Intentional self-harm (suicide) (X60-X84,Y87.0)	210	1.1	11.7
...	All other causes (Residual)	2,932	14.9	164.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Black, non-Hispanic, male, 55-64 years				
...	All causes	22,402	100.0	2,245.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,786	30.3	680.1
2	Malignant neoplasms (C00-C97)	6,763	30.2	677.8
3	Cerebrovascular diseases (I60-I69)	1,240	5.5	124.3
4	Diabetes mellitus (E10-E14)	952	4.2	95.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	736	3.3	73.8
6	Chronic lower respiratory diseases (J40-J47)	587	2.6	58.8
7	Human immunodeficiency virus (HIV) disease (B20-B24)	492	2.2	49.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	484	2.2	48.5
9	Chronic liver disease and cirrhosis (K70,K73-K74)	408	1.8	40.9
10	Septicemia (A40-A41)	381	1.7	38.2
11	Influenza and pneumonia (J10-J18)	322	1.4	32.3
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	301	1.3	30.2
13	Assault (homicide) (X85-Y09,Y87.1)	159	0.7	15.9
14	Aortic aneurysm and dissection (I71)	123	0.5	12.3
15	Viral hepatitis (B15-B19)	86	0.4	8.6
...	All other causes (Residual)	2,582	11.5	258.8
Black, non-Hispanic, male, 65-74 years				
...	All causes	28,745	100.0	4,182.2
1	Malignant neoplasms (C00-C97)	9,235	32.1	1,343.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,560	29.8	1,245.4
3	Cerebrovascular diseases (I60-I69)	1,817	6.3	264.4
4	Diabetes mellitus (E10-E14)	1,248	4.3	181.6
5	Chronic lower respiratory diseases (J40-J47)	1,195	4.2	173.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	706	2.5	102.7
7	Influenza and pneumonia (J10-J18)	562	2.0	81.8
8	Septicemia (A40-A41)	545	1.9	79.3
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	537	1.9	78.1
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	344	1.2	50.0
11	Chronic liver disease and cirrhosis (K70,K73-K74)	255	0.9	37.1
12	Pneumonitis due to solids and liquids (J69)	190	0.7	27.6
13	Human immunodeficiency virus (HIV) disease (B20-B24)	159	0.6	23.1
14	Aortic aneurysm and dissection (I71)	151	0.5	22.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	130	0.5	18.9
...	All other causes (Residual)	3,111	10.8	452.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Black, non-Hispanic, male, 75-84 years				
...	All causes	27,715	100.0	8,426.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,476	30.6	2,577.2
2	Malignant neoplasms (C00-C97)	7,689	27.7	2,337.9
3	Cerebrovascular diseases (I60-I69)	2,155	7.8	655.2
4	Chronic lower respiratory diseases (J40-J47)	1,207	4.4	367.0
5	Diabetes mellitus (E10-E14)	1,100	4.0	334.5
6	Influenza and pneumonia (J10-J18)	763	2.8	232.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	759	2.7	230.8
8	Septicemia (A40-A41)	677	2.4	205.8
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	432	1.6	131.4
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	358	1.3	108.9
11	Alzheimer's disease (G30)	320	1.2	97.3
12	Pneumonitis due to solids and liquids (J69)	243	0.9	73.9
13	Parkinson's disease (G20-G21)	154	0.6	46.8
14	Aortic aneurysm and dissection (I71)	153	0.6	46.5
15	Atherosclerosis (I70)	131	0.5	39.8
...	All other causes (Residual)	3,098	11.2	942.0
Black, non-Hispanic, male, 85 years and over				
...	All causes	14,431	100.0	15,798.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,835	33.5	5,293.1
2	Malignant neoplasms (C00-C97)	2,802	19.4	3,067.5
3	Cerebrovascular diseases (I60-I69)	1,260	8.7	1,379.4
4	Influenza and pneumonia (J10-J18)	650	4.5	711.6
5	Chronic lower respiratory diseases (J40-J47)	593	4.1	649.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	423	2.9	463.1
7	Diabetes mellitus (E10-E14)	389	2.7	425.9
8	Septicemia (A40-A41)	368	2.6	402.9
9	Alzheimer's disease (G30)	322	2.2	352.5
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	236	1.6	258.4
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	229	1.6	250.7
12	Pneumonitis due to solids and liquids (J69)	179	1.2	196.0
13	Atherosclerosis (I70)	128	0.9	140.1
14	Parkinson's disease (G20-G21)	75	0.5	82.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	65	0.5	71.2
...	All other causes (Residual)	1,877	13.0	2,054.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Black, non-Hispanic, female, all ages				
...	All causes	139,379	100.0	791.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	40,424	29.0	229.5
2	Malignant neoplasms (C00-C97)	28,897	20.7	164.1
3	Cerebrovascular diseases (I60-I69)	11,108	8.0	63.1
4	Diabetes mellitus (E10-E14)	7,198	5.2	40.9
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,808	2.7	21.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,707	2.7	21.0
7	Chronic lower respiratory diseases (J40-J47)	3,344	2.4	19.0
8	Septicemia (A40-A41)	3,304	2.4	18.8
9	Influenza and pneumonia (J10-J18)	3,040	2.2	17.3
10	Human immunodeficiency virus (HIV) disease (B20-B24)	2,417	1.7	13.7
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,286	1.6	13.0
12	Certain conditions originating in the perinatal period (P00-P96)	2,118	1.5	12.0
13	Alzheimer's disease (G30)	1,945	1.4	11.0
14	Assault (homicide) (X85-Y09,Y87.1)	1,366	1.0	7.8
15	Chronic liver disease and cirrhosis (K70,K73-K74)	996	0.7	5.7
...	All other causes (Residual)	23,421	16.8	133.0
Black, non-Hispanic, female, 1-4 years				
...	All causes	545	100.0	53.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	164	30.1	16.1
2	Assault (homicide) (X85-Y09,Y87.1)	70	12.8	6.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	56	10.3	5.5
4	Malignant neoplasms (C00-C97)	35	6.4	3.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	23	4.2	2.3
6	Septicemia (A40-A41)	12	2.2	*
7	Anemias (D50-D64)	11	2.0	*
8	Cerebrovascular diseases (I60-I69)	10	1.8	*
9	Influenza and pneumonia (J10-J18)	9	1.7	*
9	Certain conditions originating in the perinatal period (P00-P96)	9	1.7	*
11	Meningitis (G00,G03)	7	1.3	*
11	Chronic lower respiratory diseases (J40-J47)	7	1.3	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	6	1.1	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	6	1.1	*
15	Acute bronchitis and bronchiolitis (J20-J21)	3	0.6	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.6	*
...	All other causes (Residual)	114	20.9	11.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of total deaths		
		Number	Rate	Rate
Black, non-Hispanic, female, 5-14 years				
...	All causes	661	100.0	22.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	213	32.2	7.3
2	Malignant neoplasms (C00-C97)	61	9.2	2.1
3	Assault (homicide) (X85-Y09,Y87.1)	52	7.9	1.8
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	39	5.9	1.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	34	5.1	1.2
6	Chronic lower respiratory diseases (J40-J47)	27	4.1	0.9
7	Anemias (D50-D64)	13	2.0	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	11	1.7	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	1.5	*
9	Cerebrovascular diseases (I60-I69)	10	1.5	*
11	Certain conditions originating in the perinatal period (P00-P96)	7	1.1	*
12	Meningococcal infection (A39)	5	0.8	*
12	Septicemia (A40-A41)	5	0.8	*
12	Diabetes mellitus (E10-E14)	5	0.8	*
12	Intentional self-harm (suicide) (X60-X84,Y87.0)	5	0.8	*
...	All other causes (Residual)	164	24.8	5.6
Black, non-Hispanic, female, 15-24 years				
...	All causes	1,702	100.0	61.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	429	25.2	15.6
2	Assault (homicide) (X85-Y09,Y87.1)	312	18.3	11.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	139	8.2	5.0
4	Malignant neoplasms (C00-C97)	119	7.0	4.3
5	Human immunodeficiency virus (HIV) disease (B20-B24)	65	3.8	2.4
6	Intentional self-harm (suicide) (X60-X84,Y87.0)	64	3.8	2.3
7	Pregnancy, childbirth and the puerperium (O00-099)	34	2.0	1.2
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	33	1.9	1.2
9	Anemias (D50-D64)	32	1.9	1.2
10	Chronic lower respiratory diseases (J40-J47)	29	1.7	1.1
11	Diabetes mellitus (E10-E14)	27	1.6	1.0
12	Cerebrovascular diseases (I60-I69)	23	1.4	0.8
12	Influenza and pneumonia (J10-J18)	23	1.4	0.8
14	Septicemia (A40-A41)	15	0.9	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	0.8	*
...	All other causes (Residual)	344	20.2	12.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
	Black, non-Hispanic, female, 25-34 years			
...	All causes	3,454	100.0	132.5
1	Human immunodeficiency virus (HIV) disease (B20-B24)	592	17.1	22.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	424	12.3	16.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	386	11.2	14.8
4	Malignant neoplasms (C00-C97)	382	11.1	14.7
5	Assault (homicide) (X85-Y09,Y87.1)	310	9.0	11.9
6	Diabetes mellitus (E10-E14)	89	2.6	3.4
7	Cerebrovascular diseases (I60-I69)	88	2.5	3.4
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	72	2.1	2.8
9	Anemias (D50-D64)	69	2.0	2.6
10	Pregnancy, childbirth and the puerperium (O00-099)	65	1.9	2.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	49	1.4	1.9
12	Septicemia (A40-A41)	48	1.4	1.8
13	Chronic lower respiratory diseases (J40-J47)	44	1.3	1.7
14	Influenza and pneumonia (J10-J18)	34	1.0	1.3
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	31	0.9	1.2
...	All other causes (Residual)	771	22.3	29.6
	Black, non-Hispanic, female, 35-44 years			
...	All causes	8,218	100.0	285.5
1	Malignant neoplasms (C00-C97)	1,780	21.7	61.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,287	15.7	44.7
3	Human immunodeficiency virus (HIV) disease (B20-B24)	968	11.8	33.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	604	7.3	21.0
5	Cerebrovascular diseases (I60-I69)	435	5.3	15.1
6	Assault (homicide) (X85-Y09,Y87.1)	332	4.0	11.5
7	Diabetes mellitus (E10-E14)	206	2.5	7.2
8	Chronic liver disease and cirrhosis (K70,K73-K74)	167	2.0	5.8
9	Septicemia (A40-A41)	144	1.8	5.0
10	Chronic lower respiratory diseases (J40-J47)	143	1.7	5.0
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	130	1.6	4.5
12	Influenza and pneumonia (J10-J18)	103	1.3	3.6
13	Anemias (D50-D64)	84	1.0	2.9
14	Intentional self-harm (suicide) (X60-X84,Y87.0)	81	1.0	2.8
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	78	0.9	2.7
...	All other causes (Residual)	1,676	20.4	58.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
	Black, non-Hispanic, female, 45-54 years			
...	All causes	13,264	100.0	614.7
1	Malignant neoplasms (C00-C97)	3,916	29.5	181.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,819	21.3	130.6
3	Cerebrovascular diseases (I60-I69)	864	6.5	40.0
4	Diabetes mellitus (E10-E14)	622	4.7	28.8
5	Human immunodeficiency virus (HIV) disease (B20-B24)	569	4.3	26.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	507	3.8	23.5
7	Chronic lower respiratory diseases (J40-J47)	304	2.3	14.1
8	Chronic liver disease and cirrhosis (K70,K73-K74)	296	2.2	13.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	270	2.0	12.5
10	Septicemia (A40-A41)	253	1.9	11.7
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	176	1.3	8.2
12	Influenza and pneumonia (J10-J18)	152	1.1	7.0
13	Assault (homicide) (X85-Y09,Y87.1)	117	0.9	5.4
14	Viral hepatitis (B15-B19)	95	0.7	4.4
15	Anemias (D50-D64)	64	0.5	3.0
...	All other causes (Residual)	2,240	16.9	103.8
	Black, non-Hispanic, female, 55-64 years			
...	All causes	16,465	100.0	1,253.7
1	Malignant neoplasms (C00-C97)	5,256	31.9	400.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,471	27.2	340.4
3	Diabetes mellitus (E10-E14)	1,043	6.3	79.4
4	Cerebrovascular diseases (I60-I69)	1,025	6.2	78.0
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	496	3.0	37.8
6	Chronic lower respiratory diseases (J40-J47)	476	2.9	36.2
7	Septicemia (A40-A41)	375	2.3	28.6
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	254	1.5	19.3
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	247	1.5	18.8
10	Chronic liver disease and cirrhosis (K70,K73-K74)	214	1.3	16.3
11	Influenza and pneumonia (J10-J18)	209	1.3	15.9
12	Human immunodeficiency virus (HIV) disease (B20-B24)	133	0.8	10.1
13	Aortic aneurysm and dissection (I71)	73	0.4	5.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	59	0.4	4.5
15	Viral hepatitis (B15-B19)	52	0.3	4.0
15	Complications of medical and surgical care (Y40-Y84,Y88)	52	0.3	4.0
...	All other causes (Residual)	2,030	12.3	154.6

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
	Black, non-Hispanic, female, 65-74 years			
...	All causes	25,900	100.0	2,737.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,853	30.3	830.0
2	Malignant neoplasms (C00-C97)	7,250	28.0	766.3
3	Cerebrovascular diseases (I60-I69)	1,843	7.1	194.8
4	Diabetes mellitus (E10-E14)	1,835	7.1	193.9
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	931	3.6	98.4
6	Chronic lower respiratory diseases (J40-J47)	798	3.1	84.3
7	Septicemia (A40-A41)	628	2.4	66.4
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	457	1.8	48.3
9	Influenza and pneumonia (J10-J18)	422	1.6	44.6
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	320	1.2	33.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	174	0.7	18.4
12	Alzheimer's disease (G30)	148	0.6	15.6
13	Aortic aneurysm and dissection (I71)	128	0.5	13.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	115	0.4	12.2
15	Pneumonitis due to solids and liquids (J69)	99	0.4	10.5
...	All other causes (Residual)	2,899	11.2	306.4
	Black, non-Hispanic, female, 75-84 years			
...	All causes	33,159	100.0	6,109.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,124	33.5	2,049.6
2	Malignant neoplasms (C00-C97)	6,555	19.8	1,207.7
3	Cerebrovascular diseases (I60-I69)	3,189	9.6	587.6
4	Diabetes mellitus (E10-E14)	2,093	6.3	385.6
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,017	3.1	187.4
6	Chronic lower respiratory diseases (J40-J47)	921	2.8	169.7
7	Septicemia (A40-A41)	899	2.7	165.6
8	Influenza and pneumonia (J10-J18)	787	2.4	145.0
9	Alzheimer's disease (G30)	684	2.1	126.0
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	622	1.9	114.6
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	336	1.0	61.9
12	Pneumonitis due to solids and liquids (J69)	225	0.7	41.5
13	Atherosclerosis (I70)	189	0.6	34.8
14	Aortic aneurysm and dissection (I71)	178	0.5	32.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	164	0.5	30.2
...	All other causes (Residual)	4,176	12.6	769.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups,  
by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
	Black, non-Hispanic, female, 85 years and over			
...	All causes	32,229	100.0	14,683.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,224	37.9	5,569.1
2	Cerebrovascular diseases (I60-I69)	3,601	11.2	1,640.6
3	Malignant neoplasms (C00-C97)	3,535	11.0	1,610.5
4	Diabetes mellitus (E10-E14)	1,278	4.0	582.2
5	Influenza and pneumonia (J10-J18)	1,251	3.9	569.9
6	Alzheimer's disease (G30)	1,096	3.4	499.3
7	Septicemia (A40-A41)	877	2.7	399.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	867	2.7	395.0
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	670	2.1	305.2
10	Chronic lower respiratory diseases (J40-J47)	590	1.8	268.8
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	340	1.1	154.9
12	Pneumonitis due to solids and liquids (J69)	326	1.0	148.5
13	Atherosclerosis (I70)	313	1.0	142.6
14	Nutritional deficiencies (E40-E64)	157	0.5	71.5
15	Aortic aneurysm and dissection (I71)	138	0.4	62.9
...	All other causes (Residual)	4,966	15.4	2,262.4
12	Atherosclerosis (I70)	342	1.1	158.3
13	Pneumonitis due to solids and liquids (J69)	328	1.0	151.8
14	Nutritional deficiencies (E40-E64)	155	0.5	71.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	140	0.4	64.8
...	All other causes (Residual)	4,579	14.6	2,119.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable