

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, all ages				
...	All causes	113,413	100.0	306.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	27,090	23.9	73.3
2	Malignant neoplasms (C00-C97)	22,371	19.7	60.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,523	8.4	25.8
4	Cerebrovascular diseases (I60-I69)	6,416	5.7	17.4
5	Diabetes mellitus (E10-E14)	5,663	5.0	15.3
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,331	2.9	9.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	3,301	2.9	8.9
8	Chronic lower respiratory diseases (J40-J47)	2,832	2.5	7.7
9	Influenza and pneumonia (J10-J18)	2,722	2.4	7.4
10	Certain conditions originating in the perinatal period (P00-P96)	2,227	2.0	6.0
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,857	1.6	5.0
12	Human immunodeficiency virus (HIV) disease (B20-B24)	1,853	1.6	5.0
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,850	1.6	5.0
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,632	1.4	4.4
15	Septicemia (A40-A41)	1,386	1.2	3.7
...	All other causes (Residual)	19,359	17.1	52.4
Hispanic, both sexes, 1-4 years				
...	All causes	917	100.0	30.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	310	33.8	10.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	103	11.2	3.4
3	Malignant neoplasms (C00-C97)	84	9.2	2.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	69	7.5	2.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	33	3.6	1.1
6	Influenza and pneumonia (J10-J18)	32	3.5	1.1
7	Septicemia (A40-A41)	18	2.0	*
8	Certain conditions originating in the perinatal period (P00-P96)	14	1.5	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	1.4	*
10	Cerebrovascular diseases (I60-I69)	10	1.1	*
11	Chronic lower respiratory diseases (J40-J47)	8	0.9	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.7	*
13	Meningococcal infection (A39)	5	0.5	*
14	Meningitis (G00,G03)	4	0.4	*
14	Diseases of appendix (K35-K38)	4	0.4	*
...	All other causes (Residual)	204	22.2	6.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 5-14 years				
...	All causes	1,038	100.0	14.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	379	36.5	5.4
2	Malignant neoplasms (C00-C97)	177	17.1	2.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	58	5.6	0.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	56	5.4	0.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	38	3.7	0.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	37	3.6	0.5
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	21	2.0	0.3
8	Septicemia (A40-A41)	18	1.7	*
9	Cerebrovascular diseases (I60-I69)	16	1.5	*
10	Influenza and pneumonia (J10-J18)	10	1.0	*
11	Chronic lower respiratory diseases (J40-J47)	9	0.9	*
12	Anemias (D50-D64)	5	0.5	*
12	Pneumonitis due to solids and liquids (J69)	5	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.4	*
...	All other causes (Residual)	201	19.4	2.8
Hispanic, both sexes, 15-24 years				
...	All causes	5,058	100.0	75.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,182	43.1	32.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,222	24.2	18.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	414	8.2	6.2
4	Malignant neoplasms (C00-C97)	289	5.7	4.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	112	2.2	1.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	73	1.4	1.1
7	Cerebrovascular diseases (I60-I69)	40	0.8	0.6
8	Human immunodeficiency virus (HIV) disease (B20-B24)	32	0.6	0.5
9	Pregnancy, childbirth and the puerperium (O00-099)	26	0.5	0.4
10	Legal intervention (Y35,Y89.0)	23	0.5	0.3
11	Diabetes mellitus (E10-E14)	19	0.4	*
12	Influenza and pneumonia (J10-J18)	18	0.4	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	0.3	*
14	Septicemia (A40-A41)	13	0.3	*
15	Anemias (D50-D64)	11	0.2	*
...	All other causes (Residual)	568	11.2	8.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 25-34 years				
...	All causes	5,797	100.0	84.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,938	33.4	28.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	960	16.6	14.0
3	Malignant neoplasms (C00-C97)	497	8.6	7.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	490	8.5	7.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	314	5.4	4.6
6	Human immunodeficiency virus (HIV) disease (B20-B24)	300	5.2	4.4
7	Cerebrovascular diseases (I60-I69)	99	1.7	1.4
8	Chronic liver disease and cirrhosis (K70,K73-K74)	81	1.4	1.2
9	Diabetes mellitus (E10-E14)	49	0.8	0.7
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	45	0.8	0.7
11	Pregnancy, childbirth and the puerperium (O00-099)	38	0.7	0.6
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	25	0.4	0.4
13	Legal intervention (Y35,Y89.0)	24	0.4	0.3
14	Septicemia (A40-A41)	23	0.4	0.3
14	Influenza and pneumonia (J10-J18)	23	0.4	0.3
...	All other causes (Residual)	891	15.4	13.0
Hispanic, both sexes, 35-44 years				
...	All causes	8,441	100.0	154.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,760	20.9	32.3
2	Malignant neoplasms (C00-C97)	1,396	16.5	25.6
3	Human immunodeficiency virus (HIV) disease (B20-B24)	759	9.0	13.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	750	8.9	13.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	567	6.7	10.4
6	Chronic liver disease and cirrhosis (K70,K73-K74)	530	6.3	9.7
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	397	4.7	7.3
8	Cerebrovascular diseases (I60-I69)	275	3.3	5.0
9	Diabetes mellitus (E10-E14)	167	2.0	3.1
10	Viral hepatitis (B15-B19)	126	1.5	2.3
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	74	0.9	1.4
12	Influenza and pneumonia (J10-J18)	73	0.9	1.3
13	Septicemia (A40-A41)	72	0.9	1.3
14	Chronic lower respiratory diseases (J40-J47)	52	0.6	1.0
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	42	0.5	0.8
...	All other causes (Residual)	1,401	16.6	25.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 45-54 years				
...	All causes	11,362	100.0	334.4
1	Malignant neoplasms (C00-C97)	2,815	24.8	82.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,875	16.5	55.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,078	9.5	31.7
4	Chronic liver disease and cirrhosis (K70,K73-K74)	951	8.4	28.0
5	Cerebrovascular diseases (I60-I69)	539	4.7	15.9
6	Human immunodeficiency virus (HIV) disease (B20-B24)	526	4.6	15.5
7	Diabetes mellitus (E10-E14)	507	4.5	14.9
8	Viral hepatitis (B15-B19)	266	2.3	7.8
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	248	2.2	7.3
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	239	2.1	7.0
11	Septicemia (A40-A41)	133	1.2	3.9
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	132	1.2	3.9
13	Chronic lower respiratory diseases (J40-J47)	124	1.1	3.6
14	Influenza and pneumonia (J10-J18)	121	1.1	3.6
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	60	0.5	1.8
...	All other causes (Residual)	1,748	15.4	51.4
Hispanic, both sexes, 55-64 years				
...	All causes	13,337	100.0	722.2
1	Malignant neoplasms (C00-C97)	4,126	30.9	223.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,152	23.6	170.7
3	Diabetes mellitus (E10-E14)	982	7.4	53.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	721	5.4	39.0
5	Cerebrovascular diseases (I60-I69)	654	4.9	35.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	564	4.2	30.5
7	Chronic lower respiratory diseases (J40-J47)	258	1.9	14.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	237	1.8	12.8
9	Viral hepatitis (B15-B19)	192	1.4	10.4
10	Human immunodeficiency virus (HIV) disease (B20-B24)	169	1.3	9.2
11	Influenza and pneumonia (J10-J18)	168	1.3	9.1
12	Septicemia (A40-A41)	151	1.1	8.2
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	114	0.9	6.2
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	83	0.6	4.5
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	82	0.6	4.4
...	All other causes (Residual)	1,684	12.6	91.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 65-74 years				
...	All causes	20,172	100.0	1,776.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,730	28.4	504.5
2	Malignant neoplasms (C00-C97)	5,724	28.4	504.0
3	Diabetes mellitus (E10-E14)	1,604	8.0	141.2
4	Cerebrovascular diseases (I60-I69)	1,206	6.0	106.2
5	Chronic lower respiratory diseases (J40-J47)	634	3.1	55.8
6	Chronic liver disease and cirrhosis (K70,K73-K74)	618	3.1	54.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	470	2.3	41.4
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	454	2.3	40.0
9	Influenza and pneumonia (J10-J18)	402	2.0	35.4
10	Septicemia (A40-A41)	288	1.4	25.4
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	187	0.9	16.5
12	Viral hepatitis (B15-B19)	113	0.6	9.9
13	Alzheimer's disease (G30)	112	0.6	9.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	100	0.5	8.8
15	Parkinson's disease (G20-G21)	98	0.5	8.6
...	All other causes (Residual)	2,432	12.1	214.1
Hispanic, both sexes, 75-84 years				
...	All causes	23,637	100.0	4,262.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,632	32.3	1,376.3
2	Malignant neoplasms (C00-C97)	5,068	21.4	913.9
3	Cerebrovascular diseases (I60-I69)	1,892	8.0	341.2
4	Diabetes mellitus (E10-E14)	1,548	6.5	279.2
5	Chronic lower respiratory diseases (J40-J47)	940	4.0	169.5
6	Influenza and pneumonia (J10-J18)	761	3.2	137.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	524	2.2	94.5
8	Alzheimer's disease (G30)	465	2.0	83.9
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	409	1.7	73.8
10	Septicemia (A40-A41)	346	1.5	62.4
11	Chronic liver disease and cirrhosis (K70,K73-K74)	335	1.4	60.4
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	271	1.1	48.9
13	Parkinson's disease (G20-G21)	230	1.0	41.5
14	Pneumonitis due to solids and liquids (J69)	152	0.6	27.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	128	0.5	23.1
15	Atherosclerosis (I70)	128	0.5	23.1
...	All other causes (Residual)	2,808	11.9	506.4

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Hispanic, both sexes, 85 years and over			
...	All causes	18,907	100.0	11,821.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,369	39.0	4,607.5
2	Malignant neoplasms (C00-C97)	2,181	11.5	1,363.7
3	Cerebrovascular diseases (I60-I69)	1,668	8.8	1,042.9
4	Influenza and pneumonia (J10-J18)	1,056	5.6	660.3
5	Diabetes mellitus (E10-E14)	783	4.1	489.6
6	Chronic lower respiratory diseases (J40-J47)	768	4.1	480.2
7	Alzheimer's disease (G30)	639	3.4	399.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	355	1.9	222.0
9	Septicemia (A40-A41)	274	1.4	171.3
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	266	1.4	166.3
11	Pneumonitis due to solids and liquids (J69)	258	1.4	161.3
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	256	1.4	160.1
13	Atherosclerosis (I70)	209	1.1	130.7
14	Parkinson's disease (G20-G21)	138	0.7	86.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	92	0.5	57.5
...	All other causes (Residual)	2,595	13.7	1,622.5
	Hispanic, male, all ages			
...	All causes	63,317	100.0	332.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,195	22.4	74.6
2	Malignant neoplasms (C00-C97)	11,825	18.7	62.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,157	11.3	37.6
4	Cerebrovascular diseases (I60-I69)	2,982	4.7	15.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,756	4.4	14.5
6	Diabetes mellitus (E10-E14)	2,590	4.1	13.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,410	3.8	12.7
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,576	2.5	8.3
9	Chronic lower respiratory diseases (J40-J47)	1,482	2.3	7.8
10	Human immunodeficiency virus (HIV) disease (B20-B24)	1,437	2.3	7.6
11	Influenza and pneumonia (J10-J18)	1,309	2.1	6.9
12	Certain conditions originating in the perinatal period (P00-P96)	1,268	2.0	6.7
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	892	1.4	4.7
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	892	1.4	4.7
15	Septicemia (A40-A41)	684	1.1	3.6
...	All other causes (Residual)	9,862	15.6	51.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 1-4 years				
...	All causes	518	100.0	33.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	195	37.6	12.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	54	10.4	3.5
3	Malignant neoplasms (C00-C97)	44	8.5	2.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	38	7.3	2.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	16	3.1	*
5	Influenza and pneumonia (J10-J18)	16	3.1	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	1.4	*
8	Certain conditions originating in the perinatal period (P00-P96)	6	1.2	*
9	Septicemia (A40-A41)	5	1.0	*
10	Cerebrovascular diseases (I60-I69)	4	0.8	*
10	Chronic lower respiratory diseases (J40-J47)	4	0.8	*
12	Meningococcal infection (A39)	3	0.6	*
12	Meningitis (G00,G03)	3	0.6	*
12	Diseases of appendix (K35-K38)	3	0.6	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.6	*
...	All other causes (Residual)	117	22.6	7.6
Hispanic, male, 5-14 years				
...	All causes	600	100.0	16.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	224	37.3	6.2
2	Malignant neoplasms (C00-C97)	90	15.0	2.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	38	6.3	1.1
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	36	6.0	1.0
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	26	4.3	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	19	3.2	*
7	Septicemia (A40-A41)	14	2.3	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	2.0	*
9	Cerebrovascular diseases (I60-I69)	10	1.7	*
10	Influenza and pneumonia (J10-J18)	6	1.0	*
10	Chronic lower respiratory diseases (J40-J47)	6	1.0	*
12	Anemias (D50-D64)	5	0.8	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.5	*
14	Meningococcal infection (A39)	2	0.3	*
14	Viral hepatitis (B15-B19)	2	0.3	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.3	*
14	Diabetes mellitus (E10-E14)	2	0.3	*
14	Meningitis (G00,G03)	2	0.3	*
14	Pneumonitis due to solids and liquids (J69)	2	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.3	*
14	Certain conditions originating in the perinatal period (P00-P96)	2	0.3	*
...	All other causes (Residual)	95	15.8	2.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 15-24 years				
...	All causes	4,019	100.0	111.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,773	44.1	49.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,099	27.3	30.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	343	8.5	9.5
4	Malignant neoplasms (C00-C97)	199	5.0	5.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	76	1.9	2.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	39	1.0	1.1
7	Cerebrovascular diseases (I60-I69)	26	0.6	0.7
8	Legal intervention (Y35,Y89.0)	23	0.6	0.6
9	Human immunodeficiency virus (HIV) disease (B20-B24)	21	0.5	0.6
10	Influenza and pneumonia (J10-J18)	15	0.4	*
11	Diabetes mellitus (E10-E14)	11	0.3	*
12	Septicemia (A40-A41)	9	0.2	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.2	*
14	Tuberculosis (A16-A19)	6	0.1	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.1	*
14	Anemias (D50-D64)	6	0.1	*
14	Chronic lower respiratory diseases (J40-J47)	6	0.1	*
...	All other causes (Residual)	352	8.8	9.8
Hispanic, male, 25-34 years				
...	All causes	4,366	100.0	118.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,619	37.1	43.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	800	18.3	21.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	419	9.6	11.3
4	Malignant neoplasms (C00-C97)	233	5.3	6.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	226	5.2	6.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	222	5.1	6.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	67	1.5	1.8
8	Cerebrovascular diseases (I60-I69)	60	1.4	1.6
9	Diabetes mellitus (E10-E14)	28	0.6	0.8
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	26	0.6	0.7
11	Legal intervention (Y35,Y89.0)	24	0.5	0.6
12	Septicemia (A40-A41)	15	0.3	*
13	Aortic aneurysm and dissection (I71)	14	0.3	*
13	Influenza and pneumonia (J10-J18)	14	0.3	*
15	Chronic lower respiratory diseases (J40-J47)	13	0.3	*
...	All other causes (Residual)	586	13.4	15.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 35-44 years				
...	All causes	5,895	100.0	208.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,418	24.1	50.1
2	Malignant neoplasms (C00-C97)	615	10.4	21.7
3	Human immunodeficiency virus (HIV) disease (B20-B24)	589	10.0	20.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	565	9.6	20.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	451	7.7	15.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	425	7.2	15.0
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	350	5.9	12.4
8	Cerebrovascular diseases (I60-I69)	151	2.6	5.3
9	Diabetes mellitus (E10-E14)	109	1.8	3.9
10	Viral hepatitis (B15-B19)	89	1.5	3.1
11	Influenza and pneumonia (J10-J18)	48	0.8	1.7
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	46	0.8	1.6
13	Septicemia (A40-A41)	43	0.7	1.5
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	27	0.5	1.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	24	0.4	0.8
...	All other causes (Residual)	945	16.0	33.4
Hispanic, male, 45-54 years				
...	All causes	7,477	100.0	443.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,397	18.7	82.9
2	Malignant neoplasms (C00-C97)	1,372	18.3	81.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	822	11.0	48.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	782	10.5	46.4
5	Human immunodeficiency virus (HIV) disease (B20-B24)	420	5.6	24.9
6	Cerebrovascular diseases (I60-I69)	315	4.2	18.7
7	Diabetes mellitus (E10-E14)	311	4.2	18.5
8	Viral hepatitis (B15-B19)	206	2.8	12.2
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	192	2.6	11.4
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	190	2.5	11.3
11	Influenza and pneumonia (J10-J18)	85	1.1	5.0
12	Septicemia (A40-A41)	80	1.1	4.7
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	64	0.9	3.8
14	Chronic lower respiratory diseases (J40-J47)	54	0.7	3.2
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	39	0.5	2.3
...	All other causes (Residual)	1,148	15.4	68.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Hispanic, male, 55-64 years			
...	All causes	8,026	100.0	923.9
1	Malignant neoplasms (C00-C97)	2,202	27.4	253.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,104	26.2	242.2
3	Chronic liver disease and cirrhosis (K70,K73-K74)	555	6.9	63.9
4	Diabetes mellitus (E10-E14)	510	6.4	58.7
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	391	4.9	45.0
6	Cerebrovascular diseases (I60-I69)	378	4.7	43.5
7	Chronic lower respiratory diseases (J40-J47)	140	1.7	16.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	132	1.6	15.2
9	Human immunodeficiency virus (HIV) disease (B20-B24)	129	1.6	14.8
10	Viral hepatitis (B15-B19)	123	1.5	14.2
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	100	1.2	11.5
12	Influenza and pneumonia (J10-J18)	99	1.2	11.4
13	Septicemia (A40-A41)	86	1.1	9.9
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	59	0.7	6.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	35	0.4	4.0
...	All other causes (Residual)	983	12.2	113.2
	Hispanic, male, 65-74 years			
...	All causes	11,231	100.0	2,242.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,424	30.5	683.7
2	Malignant neoplasms (C00-C97)	3,219	28.7	642.8
3	Diabetes mellitus (E10-E14)	758	6.7	151.4
4	Cerebrovascular diseases (I60-I69)	637	5.7	127.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	386	3.4	77.1
6	Chronic lower respiratory diseases (J40-J47)	378	3.4	75.5
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	275	2.4	54.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	217	1.9	43.3
9	Influenza and pneumonia (J10-J18)	204	1.8	40.7
10	Septicemia (A40-A41)	149	1.3	29.8
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	84	0.7	16.8
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	77	0.7	15.4
13	Aortic aneurysm and dissection (I71)	71	0.6	14.2
14	Parkinson's disease (G20-G21)	56	0.5	11.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	50	0.4	10.0
...	All other causes (Residual)	1,246	11.1	248.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 75-84 years				
...	All causes	11,759	100.0	5,258.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,808	32.4	1,702.7
2	Malignant neoplasms (C00-C97)	2,814	23.9	1,258.3
3	Cerebrovascular diseases (I60-I69)	864	7.3	386.3
4	Diabetes mellitus (E10-E14)	638	5.4	285.3
5	Chronic lower respiratory diseases (J40-J47)	501	4.3	224.0
6	Influenza and pneumonia (J10-J18)	388	3.3	173.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	251	2.1	112.2
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	235	2.0	105.1
9	Alzheimer's disease (G30)	186	1.6	83.2
10	Septicemia (A40-A41)	170	1.4	76.0
10	Chronic liver disease and cirrhosis (K70,K73-K74)	170	1.4	76.0
12	Parkinson's disease (G20-G21)	141	1.2	63.0
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	114	1.0	51.0
14	Pneumonitis due to solids and liquids (J69)	81	0.7	36.2
15	Aortic aneurysm and dissection (I71)	78	0.7	34.9
...	All other causes (Residual)	1,320	11.2	590.2
Hispanic, male, 85 years and over				
...	All causes	6,767	100.0	12,888.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,512	37.1	4,784.3
2	Malignant neoplasms (C00-C97)	1,033	15.3	1,967.4
3	Cerebrovascular diseases (I60-I69)	528	7.8	1,005.6
4	Influenza and pneumonia (J10-J18)	402	5.9	765.6
5	Chronic lower respiratory diseases (J40-J47)	355	5.2	676.1
6	Diabetes mellitus (E10-E14)	222	3.3	422.8
7	Alzheimer's disease (G30)	161	2.4	306.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	148	2.2	281.9
9	Pneumonitis due to solids and liquids (J69)	116	1.7	220.9
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	104	1.5	198.1
11	Septicemia (A40-A41)	84	1.2	160.0
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	68	1.0	129.5
13	Parkinson's disease (G20-G21)	67	1.0	127.6
13	Atherosclerosis (I70)	67	1.0	127.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	37	0.5	70.5
...	All other causes (Residual)	863	12.8	1,643.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, all ages				
...	All causes	50,096	100.0	279.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,895	25.7	71.8
2	Malignant neoplasms (C00-C97)	10,546	21.1	58.7
3	Cerebrovascular diseases (I60-I69)	3,434	6.9	19.1
4	Diabetes mellitus (E10-E14)	3,073	6.1	17.1
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,366	4.7	13.2
6	Influenza and pneumonia (J10-J18)	1,413	2.8	7.9
7	Chronic lower respiratory diseases (J40-J47)	1,350	2.7	7.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	965	1.9	5.4
9	Certain conditions originating in the perinatal period (P00-P96)	959	1.9	5.3
10	Chronic liver disease and cirrhosis (K70,K73-K74)	891	1.8	5.0
11	Alzheimer's disease (G30)	838	1.7	4.7
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	740	1.5	4.1
13	Septicemia (A40-A41)	702	1.4	3.9
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	575	1.1	3.2
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	524	1.0	2.9
...	All other causes (Residual)	8,825	17.6	49.2
Hispanic, female, 1-4 years				
...	All causes	399	100.0	27.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	115	28.8	7.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	49	12.3	3.3
3	Malignant neoplasms (C00-C97)	40	10.0	2.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	31	7.8	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	17	4.3	*
6	Influenza and pneumonia (J10-J18)	16	4.0	*
7	Septicemia (A40-A41)	13	3.3	*
8	Certain conditions originating in the perinatal period (P00-P96)	8	2.0	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.5	*
9	Cerebrovascular diseases (I60-I69)	6	1.5	*
11	Chronic lower respiratory diseases (J40-J47)	4	1.0	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.8	*
13	Meningococcal infection (A39)	2	0.5	*
13	Pneumonitis due to solids and liquids (J69)	2	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.5	*
...	All other causes (Residual)	85	21.3	5.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 5-14 years				
...	All causes	438	100.0	12.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	155	35.4	4.5
2	Malignant neoplasms (C00-C97)	87	19.9	2.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	22	5.0	0.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	19	4.3	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	18	4.1	*
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11	2.5	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	2.1	*
8	Cerebrovascular diseases (I60-I69)	6	1.4	*
9	Septicemia (A40-A41)	4	0.9	*
9	Influenza and pneumonia (J10-J18)	4	0.9	*
11	Chronic lower respiratory diseases (J40-J47)	3	0.7	*
11	Pneumonitis due to solids and liquids (J69)	3	0.7	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.5	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.2	*
14	Meningitis (G00,G03)	1	0.2	*
14	Pregnancy, childbirth and the puerperium (O00-099)	1	0.2	*
14	Certain conditions originating in the perinatal period (P00-P96)	1	0.2	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
...	All other causes (Residual)	90	20.5	2.6
Hispanic, female, 15-24 years				
...	All causes	1,039	100.0	33.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	409	39.4	13.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	123	11.8	4.0
3	Malignant neoplasms (C00-C97)	90	8.7	2.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	71	6.8	2.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	36	3.5	1.2
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	34	3.3	1.1
7	Pregnancy, childbirth and the puerperium (O00-099)	26	2.5	0.8
8	Cerebrovascular diseases (I60-I69)	14	1.3	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	11	1.1	*
10	Diabetes mellitus (E10-E14)	8	0.8	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.7	*
12	Anemias (D50-D64)	5	0.5	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.5	*
14	Septicemia (A40-A41)	4	0.4	*
14	Viral hepatitis (B15-B19)	4	0.4	*
14	Chronic lower respiratory diseases (J40-J47)	4	0.4	*
...	All other causes (Residual)	188	18.1	6.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 25-34 years				
...	All causes	1,431	100.0	45.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	319	22.3	10.1
2	Malignant neoplasms (C00-C97)	264	18.4	8.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	160	11.2	5.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	88	6.1	2.8
5	Human immunodeficiency virus (HIV) disease (B20-B24)	78	5.5	2.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	71	5.0	2.2
7	Cerebrovascular diseases (I60-I69)	39	2.7	1.2
8	Pregnancy, childbirth and the puerperium (O00-099)	38	2.7	1.2
9	Diabetes mellitus (E10-E14)	21	1.5	0.7
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	19	1.3	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	1.0	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	1.0	*
12	Chronic liver disease and cirrhosis (K70,K73-K74)	14	1.0	*
14	Influenza and pneumonia (J10-J18)	9	0.6	*
14	Chronic lower respiratory diseases (J40-J47)	9	0.6	*
...	All other causes (Residual)	273	19.1	8.6
Hispanic, female, 35-44 years				
...	All causes	2,546	100.0	97.0
1	Malignant neoplasms (C00-C97)	781	30.7	29.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	342	13.4	13.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	185	7.3	7.1
4	Human immunodeficiency virus (HIV) disease (B20-B24)	170	6.7	6.5
5	Cerebrovascular diseases (I60-I69)	124	4.9	4.7
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	116	4.6	4.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	105	4.1	4.0
8	Diabetes mellitus (E10-E14)	58	2.3	2.2
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	47	1.8	1.8
10	Viral hepatitis (B15-B19)	37	1.5	1.4
11	Chronic lower respiratory diseases (J40-J47)	31	1.2	1.2
12	Septicemia (A40-A41)	29	1.1	1.1
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	28	1.1	1.1
14	Influenza and pneumonia (J10-J18)	25	1.0	1.0
15	Pregnancy, childbirth and the puerperium (O00-099)	17	0.7	*
...	All other causes (Residual)	451	17.7	17.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Hispanic, female, 45-54 years			
...	All causes	3,885	100.0	226.7
1	Malignant neoplasms (C00-C97)	1,443	37.1	84.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	478	12.3	27.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	256	6.6	14.9
4	Cerebrovascular diseases (I60-I69)	224	5.8	13.1
5	Diabetes mellitus (E10-E14)	196	5.0	11.4
6	Chronic liver disease and cirrhosis (K70,K73-K74)	169	4.4	9.9
7	Human immunodeficiency virus (HIV) disease (B20-B24)	106	2.7	6.2
8	Chronic lower respiratory diseases (J40-J47)	70	1.8	4.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	68	1.8	4.0
10	Viral hepatitis (B15-B19)	60	1.5	3.5
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	58	1.5	3.4
12	Septicemia (A40-A41)	53	1.4	3.1
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	47	1.2	2.7
14	Influenza and pneumonia (J10-J18)	36	0.9	2.1
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	21	0.5	1.2
...	All other causes (Residual)	600	15.4	35.0
	Hispanic, female, 55-64 years			
...	All causes	5,311	100.0	543.0
1	Malignant neoplasms (C00-C97)	1,924	36.2	196.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,048	19.7	107.2
3	Diabetes mellitus (E10-E14)	472	8.9	48.3
4	Cerebrovascular diseases (I60-I69)	276	5.2	28.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	173	3.3	17.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	166	3.1	17.0
7	Chronic lower respiratory diseases (J40-J47)	118	2.2	12.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	105	2.0	10.7
9	Viral hepatitis (B15-B19)	69	1.3	7.1
9	Influenza and pneumonia (J10-J18)	69	1.3	7.1
11	Septicemia (A40-A41)	65	1.2	6.6
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	49	0.9	5.0
13	Human immunodeficiency virus (HIV) disease (B20-B24)	40	0.8	4.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	32	0.6	3.3
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	23	0.4	2.4
...	All other causes (Residual)	682	12.8	69.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 65-74 years				
...	All causes	8,941	100.0	1,408.0
1	Malignant neoplasms (C00-C97)	2,505	28.0	394.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,306	25.8	363.1
3	Diabetes mellitus (E10-E14)	846	9.5	133.2
4	Cerebrovascular diseases (I60-I69)	569	6.4	89.6
5	Chronic lower respiratory diseases (J40-J47)	256	2.9	40.3
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	253	2.8	39.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	232	2.6	36.5
8	Influenza and pneumonia (J10-J18)	198	2.2	31.2
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	179	2.0	28.2
10	Septicemia (A40-A41)	139	1.6	21.9
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	103	1.2	16.2
12	Viral hepatitis (B15-B19)	78	0.9	12.3
13	Alzheimer's disease (G30)	68	0.8	10.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	50	0.6	7.9
15	Parkinson's disease (G20-G21)	42	0.5	6.6
...	All other causes (Residual)	1,117	12.5	175.9
Hispanic, female, 75-84 years				
...	All causes	11,878	100.0	3,589.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,824	32.2	1,155.7
2	Malignant neoplasms (C00-C97)	2,254	19.0	681.2
3	Cerebrovascular diseases (I60-I69)	1,028	8.7	310.7
4	Diabetes mellitus (E10-E14)	910	7.7	275.0
5	Chronic lower respiratory diseases (J40-J47)	439	3.7	132.7
6	Influenza and pneumonia (J10-J18)	373	3.1	112.7
7	Alzheimer's disease (G30)	279	2.3	84.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	273	2.3	82.5
9	Septicemia (A40-A41)	176	1.5	53.2
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	174	1.5	52.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	165	1.4	49.9
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	157	1.3	47.4
13	Parkinson's disease (G20-G21)	89	0.7	26.9
14	Pneumonitis due to solids and liquids (J69)	71	0.6	21.5
15	Atherosclerosis (I70)	67	0.6	20.2
...	All other causes (Residual)	1,599	13.5	483.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 85 years and over				
...	All causes	12,140	100.0	11,300.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,857	40.0	4,521.1
2	Malignant neoplasms (C00-C97)	1,148	9.5	1,068.6
3	Cerebrovascular diseases (I60-I69)	1,140	9.4	1,061.2
4	Influenza and pneumonia (J10-J18)	654	5.4	608.8
5	Diabetes mellitus (E10-E14)	561	4.6	522.2
6	Alzheimer's disease (G30)	478	3.9	444.9
7	Chronic lower respiratory diseases (J40-J47)	413	3.4	384.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	207	1.7	192.7
9	Septicemia (A40-A41)	190	1.6	176.9
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	188	1.5	175.0
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	162	1.3	150.8
12	Atherosclerosis (I70)	142	1.2	132.2
12	Pneumonitis due to solids and liquids (J69)	142	1.2	132.2
14	Parkinson's disease (G20-G21)	71	0.6	66.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	55	0.5	51.2
...	All other causes (Residual)	1,732	14.3	1,612.2
White, non-Hispanic, both sexes, all ages				
...	All causes	1,962,810	100.0	991.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	582,349	29.7	294.1
2	Malignant neoplasms (C00-C97)	456,709	23.3	230.6
3	Cerebrovascular diseases (I60-I69)	133,879	6.8	67.6
4	Chronic lower respiratory diseases (J40-J47)	110,753	5.6	55.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	76,262	3.9	38.5
6	Influenza and pneumonia (J10-J18)	51,952	2.6	26.2
7	Diabetes mellitus (E10-E14)	51,482	2.6	26.0
8	Alzheimer's disease (G30)	49,030	2.5	24.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	29,449	1.5	14.9
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	25,813	1.3	13.0
11	Septicemia (A40-A41)	24,389	1.2	12.3
12	Chronic liver disease and cirrhosis (K70,K73-K74)	20,065	1.0	10.1
13	Parkinson's disease (G20-G21)	15,239	0.8	7.7
14	Pneumonitis due to solids and liquids (J69)	14,843	0.8	7.5
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	13,837	0.7	7.0
...	All other causes (Residual)	306,759	15.6	154.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 1-4 years				
...	All causes	2,772	100.0	30.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,006	36.3	10.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	287	10.4	3.1
3	Malignant neoplasms (C00-C97)	268	9.7	2.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	160	5.8	1.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	108	3.9	1.2
6	Septicemia (A40-A41)	63	2.3	0.7
7	Influenza and pneumonia (J10-J18)	48	1.7	0.5
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	36	1.3	0.4
9	Certain conditions originating in the perinatal period (P00-P96)	34	1.2	0.4
10	Cerebrovascular diseases (I60-I69)	28	1.0	0.3
11	Meningitis (G00,G03)	15	0.5	*
12	Chronic lower respiratory diseases (J40-J47)	13	0.5	*
13	Acute bronchitis and bronchiolitis (J20-J21)	12	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	0.4	*
15	Meningococcal infection (A39)	10	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	10	0.4	*
...	All other causes (Residual)	663	23.9	7.2
White, non-Hispanic, both sexes, 5-14 years				
...	All causes	4,151	100.0	16.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,688	40.7	6.6
2	Malignant neoplasms (C00-C97)	607	14.6	2.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	226	5.4	0.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	183	4.4	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	164	4.0	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	152	3.7	0.6
7	Influenza and pneumonia (J10-J18)	57	1.4	0.2
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	56	1.3	0.2
9	Cerebrovascular diseases (I60-I69)	42	1.0	0.2
10	Chronic lower respiratory diseases (J40-J47)	39	0.9	0.2
11	Septicemia (A40-A41)	35	0.8	0.1
12	Anemias (D50-D64)	22	0.5	0.1
13	Diabetes mellitus (E10-E14)	19	0.5	*
14	Meningococcal infection (A39)	18	0.4	*
14	Certain conditions originating in the perinatal period (P00-P96)	18	0.4	*
...	All other causes (Residual)	825	19.9	3.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 15-24 years				
...	All causes	18,730	100.0	74.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,867	52.7	39.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,926	15.6	11.6
3	Malignant neoplasms (C00-C97)	1,076	5.7	4.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,058	5.6	4.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	522	2.8	2.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	302	1.6	1.2
7	Influenza and pneumonia (J10-J18)	108	0.6	0.4
8	Cerebrovascular diseases (I60-I69)	98	0.5	0.4
9	Chronic lower respiratory diseases (J40-J47)	75	0.4	0.3
10	Septicemia (A40-A41)	69	0.4	0.3
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	66	0.4	0.3
11	Diabetes mellitus (E10-E14)	66	0.4	0.3
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	45	0.2	0.2
14	Meningococcal infection (A39)	38	0.2	0.2
14	Pregnancy, childbirth and the puerperium (O00-099)	38	0.2	0.2
...	All other causes (Residual)	2,376	12.7	9.4
White, non-Hispanic, both sexes, 25-34 years				
...	All causes	24,191	100.0	96.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,798	32.2	31.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,871	16.0	15.4
3	Malignant neoplasms (C00-C97)	2,594	10.7	10.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,775	7.3	7.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,642	6.8	6.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	519	2.1	2.1
7	Diabetes mellitus (E10-E14)	335	1.4	1.3
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	330	1.4	1.3
9	Cerebrovascular diseases (I60-I69)	320	1.3	1.3
10	Chronic liver disease and cirrhosis (K70,K73-K74)	227	0.9	0.9
11	Influenza and pneumonia (J10-J18)	207	0.9	0.8
12	Septicemia (A40-A41)	145	0.6	0.6
13	Chronic lower respiratory diseases (J40-J47)	134	0.6	0.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	106	0.4	0.4
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	100	0.4	0.4
...	All other causes (Residual)	4,088	16.9	16.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	White, non-Hispanic, both sexes, 35-44 years			
...	All causes	59,546	100.0	188.3
1	Malignant neoplasms (C00-C97)	11,551	19.4	36.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,340	19.0	35.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,734	14.7	27.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,604	9.4	17.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,212	3.7	7.0
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,081	3.5	6.6
7	Human immunodeficiency virus (HIV) disease (B20-B24)	1,911	3.2	6.0
8	Cerebrovascular diseases (I60-I69)	1,333	2.2	4.2
9	Diabetes mellitus (E10-E14)	1,233	2.1	3.9
10	Chronic lower respiratory diseases (J40-J47)	613	1.0	1.9
11	Influenza and pneumonia (J10-J18)	605	1.0	1.9
12	Viral hepatitis (B15-B19)	522	0.9	1.7
13	Septicemia (A40-A41)	458	0.8	1.4
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	439	0.7	1.4
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	359	0.6	1.1
...	All other causes (Residual)	10,551	17.7	33.4
	White, non-Hispanic, both sexes, 45-54 years			
...	All causes	117,086	100.0	396.0
1	Malignant neoplasms (C00-C97)	36,836	31.5	124.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	25,544	21.8	86.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,704	8.3	32.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,292	4.5	17.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	5,035	4.3	17.0
6	Diabetes mellitus (E10-E14)	3,332	2.8	11.3
7	Cerebrovascular diseases (I60-I69)	3,200	2.7	10.8
8	Chronic lower respiratory diseases (J40-J47)	2,570	2.2	8.7
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,373	1.2	4.6
10	Viral hepatitis (B15-B19)	1,364	1.2	4.6
11	Human immunodeficiency virus (HIV) disease (B20-B24)	1,248	1.1	4.2
12	Influenza and pneumonia (J10-J18)	1,193	1.0	4.0
13	Septicemia (A40-A41)	1,181	1.0	4.0
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	904	0.8	3.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	561	0.5	1.9
...	All other causes (Residual)	17,749	15.2	60.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 55-64 years				
...	All causes	184,109	100.0	924.0
1	Malignant neoplasms (C00-C97)	71,389	38.8	358.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	46,433	25.2	233.0
3	Chronic lower respiratory diseases (J40-J47)	9,678	5.3	48.6
4	Cerebrovascular diseases (I60-I69)	6,208	3.4	31.2
5	Diabetes mellitus (E10-E14)	6,118	3.3	30.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,776	3.1	29.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	4,189	2.3	21.0
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,979	1.6	15.0
9	Septicemia (A40-A41)	2,081	1.1	10.4
10	Influenza and pneumonia (J10-J18)	1,963	1.1	9.9
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,861	1.0	9.3
12	Aortic aneurysm and dissection (I71)	1,117	0.6	5.6
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,006	0.5	5.0
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	776	0.4	3.9
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	652	0.4	3.3
...	All other causes (Residual)	21,883	11.9	109.8
White, non-Hispanic, both sexes, 65-74 years				
...	All causes	346,389	100.0	2,325.5
1	Malignant neoplasms (C00-C97)	121,529	35.1	815.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	91,855	26.5	616.7
3	Chronic lower respiratory diseases (J40-J47)	27,693	8.0	185.9
4	Cerebrovascular diseases (I60-I69)	16,838	4.9	113.0
5	Diabetes mellitus (E10-E14)	11,464	3.3	77.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,309	1.8	42.4
7	Influenza and pneumonia (J10-J18)	5,214	1.5	35.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,984	1.4	33.5
9	Septicemia (A40-A41)	4,385	1.3	29.4
10	Chronic liver disease and cirrhosis (K70,K73-K74)	4,307	1.2	28.9
11	Aortic aneurysm and dissection (I71)	3,079	0.9	20.7
12	Alzheimer's disease (G30)	3,020	0.9	20.3
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,187	0.6	14.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,185	0.6	14.7
15	Parkinson's disease (G20-G21)	1,911	0.6	12.8
...	All other causes (Residual)	39,429	11.4	264.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 75-84 years				
...	All causes	603,143	100.0	5,604.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	185,536	30.8	1,724.0
2	Malignant neoplasms (C00-C97)	142,760	23.7	1,326.5
3	Cerebrovascular diseases (I60-I69)	47,304	7.8	439.5
4	Chronic lower respiratory diseases (J40-J47)	43,972	7.3	408.6
5	Diabetes mellitus (E10-E14)	17,507	2.9	162.7
6	Alzheimer's disease (G30)	16,742	2.8	155.6
7	Influenza and pneumonia (J10-J18)	15,837	2.6	147.2
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,157	1.8	103.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10,398	1.7	96.6
10	Septicemia (A40-A41)	8,352	1.4	77.6
11	Parkinson's disease (G20-G21)	7,521	1.2	69.9
12	Aortic aneurysm and dissection (I71)	5,155	0.9	47.9
13	Pneumonitis due to solids and liquids (J69)	5,012	0.8	46.6
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	4,499	0.7	41.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,150	0.7	38.6
...	All other causes (Residual)	77,241	12.8	717.7
White, non-Hispanic, both sexes, 85 years and over				
...	All causes	589,269	100.0	15,371.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	221,430	37.6	5,776.0
2	Malignant neoplasms (C00-C97)	68,053	11.5	1,775.2
3	Cerebrovascular diseases (I60-I69)	58,461	9.9	1,525.0
4	Alzheimer's disease (G30)	28,720	4.9	749.2
5	Influenza and pneumonia (J10-J18)	26,602	4.5	693.9
6	Chronic lower respiratory diseases (J40-J47)	25,950	4.4	676.9
7	Diabetes mellitus (E10-E14)	11,402	1.9	297.4
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,092	1.9	289.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10,723	1.8	279.7
10	Septicemia (A40-A41)	7,498	1.3	195.6
11	Pneumonitis due to solids and liquids (J69)	7,329	1.2	191.2
12	Atherosclerosis (I70)	7,186	1.2	187.4
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	6,242	1.1	162.8
14	Parkinson's disease (G20-G21)	5,502	0.9	143.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3,251	0.6	84.8
...	All other causes (Residual)	89,828	15.2	2,343.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, all ages				
...	All causes	945,967	100.0	975.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	280,713	29.7	289.5
2	Malignant neoplasms (C00-C97)	236,004	24.9	243.4
3	Chronic lower respiratory diseases (J40-J47)	52,952	5.6	54.6
4	Cerebrovascular diseases (I60-I69)	50,382	5.3	52.0
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	48,180	5.1	49.7
6	Diabetes mellitus (E10-E14)	24,302	2.6	25.1
7	Influenza and pneumonia (J10-J18)	22,389	2.4	23.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	20,710	2.2	21.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14,331	1.5	14.8
10	Alzheimer's disease (G30)	14,295	1.5	14.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	12,609	1.3	13.0
12	Septicemia (A40-A41)	10,668	1.1	11.0
13	Parkinson's disease (G20-G21)	8,685	0.9	9.0
14	Aortic aneurysm and dissection (I71)	7,924	0.8	8.2
15	Pneumonitis due to solids and liquids (J69)	7,411	0.8	7.6
...	All other causes (Residual)	134,412	14.2	138.6
White, non-Hispanic, male, 1-4 years				
...	All causes	1,592	100.0	33.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	624	39.2	13.2
2	Malignant neoplasms (C00-C97)	144	9.0	3.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	144	9.0	3.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	95	6.0	2.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	59	3.7	1.3
6	Septicemia (A40-A41)	41	2.6	0.9
7	Influenza and pneumonia (J10-J18)	26	1.6	0.6
8	Certain conditions originating in the perinatal period (P00-P96)	22	1.4	0.5
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	1.1	*
10	Cerebrovascular diseases (I60-I69)	14	0.9	*
11	Meningitis (G00,G03)	10	0.6	*
12	Chronic lower respiratory diseases (J40-J47)	9	0.6	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.4	*
14	Meningococcal infection (A39)	5	0.3	*
14	Acute bronchitis and bronchiolitis (J20-J21)	5	0.3	*
...	All other causes (Residual)	369	23.2	7.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 5-14 years				
...	All causes	2,428	100.0	18.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,066	43.9	8.1
2	Malignant neoplasms (C00-C97)	310	12.8	2.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	144	5.9	1.1
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	135	5.6	1.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	83	3.4	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	80	3.3	0.6
7	Influenza and pneumonia (J10-J18)	28	1.2	0.2
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	25	1.0	0.2
9	Cerebrovascular diseases (I60-I69)	24	1.0	0.2
10	Chronic lower respiratory diseases (J40-J47)	22	0.9	0.2
11	Septicemia (A40-A41)	18	0.7	*
12	Certain conditions originating in the perinatal period (P00-P96)	14	0.6	*
13	Diabetes mellitus (E10-E14)	13	0.5	*
14	Anemias (D50-D64)	11	0.5	*
15	Meningococcal infection (A39)	10	0.4	*
...	All other causes (Residual)	445	18.3	3.4
White, non-Hispanic, male, 15-24 years				
...	All causes	13,542	100.0	105.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,297	53.9	56.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,522	18.6	19.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	729	5.4	5.7
4	Malignant neoplasms (C00-C97)	632	4.7	4.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	336	2.5	2.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	183	1.4	1.4
7	Influenza and pneumonia (J10-J18)	64	0.5	0.5
8	Cerebrovascular diseases (I60-I69)	55	0.4	0.4
9	Diabetes mellitus (E10-E14)	42	0.3	0.3
10	Chronic lower respiratory diseases (J40-J47)	39	0.3	0.3
11	Septicemia (A40-A41)	37	0.3	0.3
12	Legal intervention (Y35,Y89.0)	35	0.3	0.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	31	0.2	0.2
14	Aortic aneurysm and dissection (I71)	21	0.2	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21	0.2	0.2
...	All other causes (Residual)	1,498	11.1	11.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 25-34 years				
...	All causes	16,613	100.0	131.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,906	35.6	46.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,190	19.2	25.2
3	Malignant neoplasms (C00-C97)	1,260	7.6	10.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,184	7.1	9.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,177	7.1	9.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	410	2.5	3.2
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	186	1.1	1.5
8	Diabetes mellitus (E10-E14)	184	1.1	1.5
9	Cerebrovascular diseases (I60-I69)	161	1.0	1.3
10	Chronic liver disease and cirrhosis (K70,K73-K74)	145	0.9	1.1
11	Influenza and pneumonia (J10-J18)	112	0.7	0.9
12	Septicemia (A40-A41)	69	0.4	0.5
13	Chronic lower respiratory diseases (J40-J47)	62	0.4	0.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	58	0.3	0.5
14	Aortic aneurysm and dissection (I71)	58	0.3	0.5
...	All other causes (Residual)	2,451	14.8	19.4
White, non-Hispanic, male, 35-44 years				
...	All causes	38,213	100.0	241.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,188	21.4	51.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,395	16.7	40.4
3	Malignant neoplasms (C00-C97)	5,160	13.5	32.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,329	11.3	27.4
5	Human immunodeficiency virus (HIV) disease (B20-B24)	1,639	4.3	10.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,498	3.9	9.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,403	3.7	8.9
8	Diabetes mellitus (E10-E14)	767	2.0	4.8
9	Cerebrovascular diseases (I60-I69)	684	1.8	4.3
10	Viral hepatitis (B15-B19)	368	1.0	2.3
11	Influenza and pneumonia (J10-J18)	333	0.9	2.1
12	Chronic lower respiratory diseases (J40-J47)	293	0.8	1.9
13	Septicemia (A40-A41)	266	0.7	1.7
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	236	0.6	1.5
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	211	0.6	1.3
...	All other causes (Residual)	6,443	16.9	40.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 45-54 years				
...	All causes	73,731	100.0	502.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	19,178	26.0	130.7
2	Malignant neoplasms (C00-C97)	18,679	25.3	127.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,034	9.5	48.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,016	5.4	27.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	3,645	4.9	24.8
6	Diabetes mellitus (E10-E14)	2,014	2.7	13.7
7	Cerebrovascular diseases (I60-I69)	1,745	2.4	11.9
8	Chronic lower respiratory diseases (J40-J47)	1,331	1.8	9.1
9	Human immunodeficiency virus (HIV) disease (B20-B24)	1,106	1.5	7.5
10	Viral hepatitis (B15-B19)	1,062	1.4	7.2
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	990	1.3	6.7
12	Influenza and pneumonia (J10-J18)	707	1.0	4.8
13	Septicemia (A40-A41)	615	0.8	4.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	543	0.7	3.7
15	Aortic aneurysm and dissection (I71)	342	0.5	2.3
...	All other causes (Residual)	10,724	14.5	73.1
White, non-Hispanic, male, 55-64 years				
...	All causes	109,993	100.0	1,136.3
1	Malignant neoplasms (C00-C97)	38,884	35.4	401.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	32,504	29.6	335.8
3	Chronic lower respiratory diseases (J40-J47)	4,885	4.4	50.5
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,905	3.6	40.3
5	Diabetes mellitus (E10-E14)	3,524	3.2	36.4
6	Cerebrovascular diseases (I60-I69)	3,400	3.1	35.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,875	2.6	29.7
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,284	2.1	23.6
9	Influenza and pneumonia (J10-J18)	1,132	1.0	11.7
10	Septicemia (A40-A41)	1,071	1.0	11.1
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,012	0.9	10.5
12	Aortic aneurysm and dissection (I71)	841	0.8	8.7
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	547	0.5	5.7
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	454	0.4	4.7
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	439	0.4	4.5
...	All other causes (Residual)	12,236	11.1	126.4

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 65-74 years				
...	All causes	196,158	100.0	2,869.4
1	Malignant neoplasms (C00-C97)	67,554	34.4	988.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	57,059	29.1	834.7
3	Chronic lower respiratory diseases (J40-J47)	14,089	7.2	206.1
4	Cerebrovascular diseases (I60-I69)	8,753	4.5	128.0
5	Diabetes mellitus (E10-E14)	6,142	3.1	89.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,837	2.0	56.1
7	Influenza and pneumonia (J10-J18)	2,919	1.5	42.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,729	1.4	39.9
9	Chronic liver disease and cirrhosis (K70,K73-K74)	2,606	1.3	38.1
10	Septicemia (A40-A41)	2,223	1.1	32.5
11	Aortic aneurysm and dissection (I71)	2,109	1.1	30.9
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,836	0.9	26.9
13	Alzheimer's disease (G30)	1,305	0.7	19.1
14	Parkinson's disease (G20-G21)	1,252	0.6	18.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,250	0.6	18.3
...	All other causes (Residual)	20,495	10.4	299.8
White, non-Hispanic, male, 75-84 years				
...	All causes	294,013	100.0	6,851.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	93,996	32.0	2,190.4
2	Malignant neoplasms (C00-C97)	73,885	25.1	1,721.8
3	Chronic lower respiratory diseases (J40-J47)	21,496	7.3	500.9
4	Cerebrovascular diseases (I60-I69)	19,762	6.7	460.5
5	Diabetes mellitus (E10-E14)	7,874	2.7	183.5
6	Influenza and pneumonia (J10-J18)	7,830	2.7	182.5
7	Alzheimer's disease (G30)	6,137	2.1	143.0
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,800	2.0	135.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,310	1.8	123.7
10	Parkinson's disease (G20-G21)	4,486	1.5	104.5
11	Septicemia (A40-A41)	3,851	1.3	89.7
12	Aortic aneurysm and dissection (I71)	3,002	1.0	70.0
13	Pneumonitis due to solids and liquids (J69)	2,765	0.9	64.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,152	0.7	50.1
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,763	0.6	41.1
...	All other causes (Residual)	33,904	11.5	790.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 85 years and over				
...	All causes	192,084	100.0	17,055.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	69,775	36.3	6,195.4
2	Malignant neoplasms (C00-C97)	29,471	15.3	2,616.8
3	Cerebrovascular diseases (I60-I69)	15,756	8.2	1,399.0
4	Chronic lower respiratory diseases (J40-J47)	10,716	5.6	951.5
5	Influenza and pneumonia (J10-J18)	9,166	4.8	813.9
6	Alzheimer's disease (G30)	6,613	3.4	587.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,422	2.3	392.6
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,220	2.2	374.7
9	Diabetes mellitus (E10-E14)	3,740	1.9	332.1
10	Pneumonitis due to solids and liquids (J69)	3,119	1.6	276.9
11	Parkinson's disease (G20-G21)	2,722	1.4	241.7
12	Septicemia (A40-A41)	2,406	1.3	213.6
13	Atherosclerosis (I70)	1,973	1.0	175.2
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,600	0.8	142.1
15	Aortic aneurysm and dissection (I71)	1,384	0.7	122.9
...	All other causes (Residual)	25,001	13.0	2,219.9
White, non-Hispanic, female, all ages				
...	All causes	1,016,843	100.0	1,006.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	301,636	29.7	298.4
2	Malignant neoplasms (C00-C97)	220,705	21.7	218.4
3	Cerebrovascular diseases (I60-I69)	83,497	8.2	82.6
4	Chronic lower respiratory diseases (J40-J47)	57,801	5.7	57.2
5	Alzheimer's disease (G30)	34,735	3.4	34.4
6	Influenza and pneumonia (J10-J18)	29,563	2.9	29.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	28,082	2.8	27.8
8	Diabetes mellitus (E10-E14)	27,180	2.7	26.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15,118	1.5	15.0
10	Septicemia (A40-A41)	13,721	1.3	13.6
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	8,867	0.9	8.8
12	Atherosclerosis (I70)	8,006	0.8	7.9
13	Chronic liver disease and cirrhosis (K70,K73-K74)	7,456	0.7	7.4
14	Pneumonitis due to solids and liquids (J69)	7,432	0.7	7.4
15	Parkinson's disease (G20-G21)	6,554	0.6	6.5
...	All other causes (Residual)	166,490	16.4	164.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 1-4 years				
...	All causes	1,180	100.0	26.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	382	32.4	8.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	143	12.1	3.2
3	Malignant neoplasms (C00-C97)	124	10.5	2.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	65	5.5	1.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	49	4.2	1.1
6	Septicemia (A40-A41)	22	1.9	0.5
6	Influenza and pneumonia (J10-J18)	22	1.9	0.5
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	1.5	*
9	Cerebrovascular diseases (I60-I69)	14	1.2	*
10	Certain conditions originating in the perinatal period (P00-P96)	12	1.0	*
11	Acute bronchitis and bronchiolitis (J20-J21)	7	0.6	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.6	*
13	Pneumonitis due to solids and liquids (J69)	6	0.5	*
14	Meningococcal infection (A39)	5	0.4	*
14	Anemias (D50-D64)	5	0.4	*
14	Meningitis (G00,G03)	5	0.4	*
...	All other causes (Residual)	294	24.9	6.6
White, non-Hispanic, female, 5-14 years				
...	All causes	1,723	100.0	13.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	622	36.1	5.0
2	Malignant neoplasms (C00-C97)	297	17.2	2.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	91	5.3	0.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	81	4.7	0.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	72	4.2	0.6
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	39	2.3	0.3
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	31	1.8	0.3
8	Influenza and pneumonia (J10-J18)	29	1.7	0.2
9	Cerebrovascular diseases (I60-I69)	18	1.0	*
10	Septicemia (A40-A41)	17	1.0	*
10	Chronic lower respiratory diseases (J40-J47)	17	1.0	*
12	Anemias (D50-D64)	11	0.6	*
13	Meningococcal infection (A39)	8	0.5	*
14	Meningitis (G00,G03)	7	0.4	*
15	Diabetes mellitus (E10-E14)	6	0.3	*
...	All other causes (Residual)	377	21.9	3.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 15-24 years				
...	All causes	5,188	100.0	41.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,570	49.5	20.7
2	Malignant neoplasms (C00-C97)	444	8.6	3.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	404	7.8	3.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	329	6.3	2.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	186	3.6	1.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	119	2.3	1.0
7	Influenza and pneumonia (J10-J18)	44	0.8	0.4
8	Cerebrovascular diseases (I60-I69)	43	0.8	0.3
9	Pregnancy, childbirth and the puerperium (O00-099)	38	0.7	0.3
10	Chronic lower respiratory diseases (J40-J47)	36	0.7	0.3
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	35	0.7	0.3
12	Septicemia (A40-A41)	32	0.6	0.3
13	Diabetes mellitus (E10-E14)	24	0.5	0.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	24	0.5	0.2
15	Meningococcal infection (A39)	20	0.4	0.2
...	All other causes (Residual)	840	16.2	6.8
White, non-Hispanic, female, 25-34 years				
...	All causes	7,578	100.0	60.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,892	25.0	15.2
2	Malignant neoplasms (C00-C97)	1,334	17.6	10.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	681	9.0	5.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	591	7.8	4.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	465	6.1	3.7
6	Cerebrovascular diseases (I60-I69)	159	2.1	1.3
7	Diabetes mellitus (E10-E14)	151	2.0	1.2
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	144	1.9	1.2
9	Human immunodeficiency virus (HIV) disease (B20-B24)	109	1.4	0.9
10	Influenza and pneumonia (J10-J18)	95	1.3	0.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	82	1.1	0.7
12	Pregnancy, childbirth and the puerperium (O00-099)	77	1.0	0.6
13	Septicemia (A40-A41)	76	1.0	0.6
14	Chronic lower respiratory diseases (J40-J47)	72	1.0	0.6
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	55	0.7	0.4
...	All other causes (Residual)	1,595	21.0	12.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 35-44 years				
...	All causes	21,333	100.0	134.9
1	Malignant neoplasms (C00-C97)	6,391	30.0	40.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,152	14.8	19.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,339	11.0	14.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,275	6.0	8.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	809	3.8	5.1
6	Cerebrovascular diseases (I60-I69)	649	3.0	4.1
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	583	2.7	3.7
8	Diabetes mellitus (E10-E14)	466	2.2	2.9
9	Chronic lower respiratory diseases (J40-J47)	320	1.5	2.0
10	Human immunodeficiency virus (HIV) disease (B20-B24)	272	1.3	1.7
10	Influenza and pneumonia (J10-J18)	272	1.3	1.7
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	203	1.0	1.3
13	Septicemia (A40-A41)	192	0.9	1.2
14	Viral hepatitis (B15-B19)	154	0.7	1.0
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	148	0.7	0.9
...	All other causes (Residual)	4,108	19.3	26.0
White, non-Hispanic, female, 45-54 years				
...	All causes	43,355	100.0	291.0
1	Malignant neoplasms (C00-C97)	18,157	41.9	121.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,366	14.7	42.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,670	6.2	17.9
4	Cerebrovascular diseases (I60-I69)	1,455	3.4	9.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,390	3.2	9.3
6	Diabetes mellitus (E10-E14)	1,318	3.0	8.8
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,276	2.9	8.6
8	Chronic lower respiratory diseases (J40-J47)	1,239	2.9	8.3
9	Septicemia (A40-A41)	566	1.3	3.8
10	Influenza and pneumonia (J10-J18)	486	1.1	3.3
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	383	0.9	2.6
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	361	0.8	2.4
13	Viral hepatitis (B15-B19)	302	0.7	2.0
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	252	0.6	1.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	249	0.6	1.7
...	All other causes (Residual)	6,885	15.9	46.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 55-64 years				
...	All causes	74,116	100.0	723.5
1	Malignant neoplasms (C00-C97)	32,505	43.9	317.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,929	18.8	136.0
3	Chronic lower respiratory diseases (J40-J47)	4,793	6.5	46.8
4	Cerebrovascular diseases (I60-I69)	2,808	3.8	27.4
5	Diabetes mellitus (E10-E14)	2,594	3.5	25.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,871	2.5	18.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,314	1.8	12.8
8	Septicemia (A40-A41)	1,010	1.4	9.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	849	1.1	8.3
10	Influenza and pneumonia (J10-J18)	831	1.1	8.1
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	695	0.9	6.8
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	459	0.6	4.5
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	337	0.5	3.3
14	Aortic aneurysm and dissection (I71)	276	0.4	2.7
15	Alzheimer's disease (G30)	275	0.4	2.7
...	All other causes (Residual)	9,570	12.9	93.4
White, non-Hispanic, female, 65-74 years				
...	All causes	150,231	100.0	1,864.1
1	Malignant neoplasms (C00-C97)	53,975	35.9	669.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	34,796	23.2	431.8
3	Chronic lower respiratory diseases (J40-J47)	13,604	9.1	168.8
4	Cerebrovascular diseases (I60-I69)	8,085	5.4	100.3
5	Diabetes mellitus (E10-E14)	5,322	3.5	66.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,472	1.6	30.7
7	Influenza and pneumonia (J10-J18)	2,295	1.5	28.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,255	1.5	28.0
9	Septicemia (A40-A41)	2,162	1.4	26.8
10	Alzheimer's disease (G30)	1,715	1.1	21.3
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1,701	1.1	21.1
12	Aortic aneurysm and dissection (I71)	970	0.6	12.0
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	935	0.6	11.6
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	874	0.6	10.8
15	Parkinson's disease (G20-G21)	659	0.4	8.2
...	All other causes (Residual)	18,411	12.3	228.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	White, non-Hispanic, female, 75-84 years			
...	All causes	309,130	100.0	4,777.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	91,540	29.6	1,414.7
2	Malignant neoplasms (C00-C97)	68,875	22.3	1,064.4
3	Cerebrovascular diseases (I60-I69)	27,542	8.9	425.6
4	Chronic lower respiratory diseases (J40-J47)	22,476	7.3	347.3
5	Alzheimer's disease (G30)	10,605	3.4	163.9
6	Diabetes mellitus (E10-E14)	9,633	3.1	148.9
7	Influenza and pneumonia (J10-J18)	8,007	2.6	123.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,357	1.7	82.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,088	1.6	78.6
10	Septicemia (A40-A41)	4,501	1.5	69.6
11	Parkinson's disease (G20-G21)	3,035	1.0	46.9
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,798	0.9	43.2
13	Pneumonitis due to solids and liquids (J69)	2,247	0.7	34.7
14	Aortic aneurysm and dissection (I71)	2,153	0.7	33.3
15	Atherosclerosis (I70)	2,105	0.7	32.5
...	All other causes (Residual)	43,168	14.0	667.1
	White, non-Hispanic, female, 85 years and over			
...	All causes	397,185	100.0	14,670.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	151,655	38.2	5,601.6
2	Cerebrovascular diseases (I60-I69)	42,705	10.8	1,577.4
3	Malignant neoplasms (C00-C97)	38,582	9.7	1,425.1
4	Alzheimer's disease (G30)	22,107	5.6	816.6
5	Influenza and pneumonia (J10-J18)	17,436	4.4	644.0
6	Chronic lower respiratory diseases (J40-J47)	15,234	3.8	562.7
7	Diabetes mellitus (E10-E14)	7,662	1.9	283.0
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,872	1.7	253.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,301	1.6	232.7
10	Atherosclerosis (I70)	5,213	1.3	192.6
11	Septicemia (A40-A41)	5,092	1.3	188.1
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	4,642	1.2	171.5
13	Pneumonitis due to solids and liquids (J69)	4,210	1.1	155.5
14	Parkinson's disease (G20-G21)	2,780	0.7	102.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,996	0.5	73.7
...	All other causes (Residual)	64,698	16.3	2,389.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, all ages				
...	All causes	284,343	100.0	798.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	76,794	27.0	215.5
2	Malignant neoplasms (C00-C97)	61,580	21.7	172.8
3	Cerebrovascular diseases (I60-I69)	18,844	6.6	52.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,239	4.3	34.4
5	Diabetes mellitus (E10-E14)	12,167	4.3	34.1
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8,077	2.8	22.7
7	Human immunodeficiency virus (HIV) disease (B20-B24)	7,710	2.7	21.6
8	Chronic lower respiratory diseases (J40-J47)	7,504	2.6	21.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,214	2.5	20.2
10	Septicemia (A40-A41)	5,829	2.0	16.4
11	Influenza and pneumonia (J10-J18)	5,686	2.0	16.0
12	Certain conditions originating in the perinatal period (P00-P96)	4,713	1.7	13.2
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	4,012	1.4	11.3
14	Alzheimer's disease (G30)	3,104	1.1	8.7
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,733	1.0	7.7
...	All other causes (Residual)	46,137	16.2	129.5
Black, non-Hispanic, both sexes, 1-4 years				
...	All causes	1,156	100.0	49.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	331	28.6	14.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	166	14.4	7.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	127	11.0	5.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	73	6.3	3.1
5	Malignant neoplasms (C00-C97)	46	4.0	2.0
6	Influenza and pneumonia (J10-J18)	24	2.1	1.0
7	Chronic lower respiratory diseases (J40-J47)	22	1.9	0.9
7	Certain conditions originating in the perinatal period (P00-P96)	22	1.9	0.9
9	Septicemia (A40-A41)	21	1.8	0.9
10	Cerebrovascular diseases (I60-I69)	12	1.0	*
11	Anemias (D50-D64)	11	1.0	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	10	0.9	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.6	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.4	*
14	Meningitis (G00,G03)	5	0.4	*
...	All other causes (Residual)	274	23.7	11.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 5-14 years				
...	All causes	1,557	100.0	24.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	630	40.5	9.7
2	Malignant neoplasms (C00-C97)	172	11.0	2.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	92	5.9	1.4
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	73	4.7	1.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	70	4.5	1.1
6	Chronic lower respiratory diseases (J40-J47)	53	3.4	0.8
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	40	2.6	0.6
8	Human immunodeficiency virus (HIV) disease (B20-B24)	26	1.7	0.4
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	26	1.7	0.4
8	Anemias (D50-D64)	26	1.7	0.4
11	Influenza and pneumonia (J10-J18)	17	1.1	*
12	Diabetes mellitus (E10-E14)	15	1.0	*
13	Cerebrovascular diseases (I60-I69)	14	0.9	*
14	Certain conditions originating in the perinatal period (P00-P96)	11	0.7	*
15	Septicemia (A40-A41)	10	0.6	*
...	All other causes (Residual)	282	18.1	4.4
Black, non-Hispanic, both sexes, 15-24 years				
...	All causes	7,024	100.0	121.4
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,817	40.1	48.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,713	24.4	29.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	425	6.1	7.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	322	4.6	5.6
5	Malignant neoplasms (C00-C97)	257	3.7	4.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	161	2.3	2.8
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	113	1.6	2.0
8	Chronic lower respiratory diseases (J40-J47)	77	1.1	1.3
9	Anemias (D50-D64)	72	1.0	1.2
10	Diabetes mellitus (E10-E14)	56	0.8	1.0
11	Cerebrovascular diseases (I60-I69)	51	0.7	0.9
12	Influenza and pneumonia (J10-J18)	45	0.6	0.8
12	Pregnancy, childbirth and the puerperium (O00-099)	45	0.6	0.8
14	Septicemia (A40-A41)	41	0.6	0.7
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	27	0.4	0.5
...	All other causes (Residual)	802	11.4	13.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 25-34 years				
...	All causes	9,859	100.0	191.5
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,335	23.7	45.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,600	16.2	31.1
3	Human immunodeficiency virus (HIV) disease (B20-B24)	1,234	12.5	24.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	961	9.7	18.7
5	Malignant neoplasms (C00-C97)	691	7.0	13.4
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	472	4.8	9.2
7	Diabetes mellitus (E10-E14)	198	2.0	3.8
8	Cerebrovascular diseases (I60-I69)	153	1.6	3.0
9	Chronic lower respiratory diseases (J40-J47)	128	1.3	2.5
10	Anemias (D50-D64)	117	1.2	2.3
11	Septicemia (A40-A41)	98	1.0	1.9
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	93	0.9	1.8
13	Influenza and pneumonia (J10-J18)	87	0.9	1.7
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	67	0.7	1.3
15	Pregnancy, childbirth and the puerperium (O00-099)	65	0.7	1.3
...	All other causes (Residual)	1,560	15.8	30.3
Black, non-Hispanic, both sexes, 35-44 years				
...	All causes	20,436	100.0	365.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,424	16.8	61.3
2	Human immunodeficiency virus (HIV) disease (B20-B24)	3,072	15.0	55.0
3	Malignant neoplasms (C00-C97)	2,950	14.4	52.8
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,282	11.2	40.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,427	7.0	25.5
6	Cerebrovascular diseases (I60-I69)	758	3.7	13.6
7	Diabetes mellitus (E10-E14)	506	2.5	9.1
8	Chronic liver disease and cirrhosis (K70,K73-K74)	420	2.1	7.5
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	418	2.0	7.5
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	306	1.5	5.5
11	Influenza and pneumonia (J10-J18)	274	1.3	4.9
12	Chronic lower respiratory diseases (J40-J47)	273	1.3	4.9
13	Septicemia (A40-A41)	268	1.3	4.8
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	177	0.9	3.2
15	Anemias (D50-D64)	154	0.8	2.8
...	All other causes (Residual)	3,727	18.2	66.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 45-54 years				
...	All causes	34,098	100.0	786.9
1	Malignant neoplasms (C00-C97)	8,175	24.0	188.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,956	23.3	183.6
3	Human immunodeficiency virus (HIV) disease (B20-B24)	2,249	6.6	51.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,103	6.2	48.5
5	Cerebrovascular diseases (I60-I69)	1,886	5.5	43.5
6	Diabetes mellitus (E10-E14)	1,306	3.8	30.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	993	2.9	22.9
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	704	2.1	16.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	702	2.1	16.2
10	Septicemia (A40-A41)	595	1.7	13.7
11	Chronic lower respiratory diseases (J40-J47)	549	1.6	12.7
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	439	1.3	10.1
13	Influenza and pneumonia (J10-J18)	430	1.3	9.9
14	Viral hepatitis (B15-B19)	349	1.0	8.1
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	268	0.8	6.2
...	All other causes (Residual)	5,394	15.8	124.5
Black, non-Hispanic, both sexes, 55-64 years				
...	All causes	39,523	100.0	1,602.5
1	Malignant neoplasms (C00-C97)	12,196	30.9	494.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,209	28.4	454.5
3	Cerebrovascular diseases (I60-I69)	2,289	5.8	92.8
4	Diabetes mellitus (E10-E14)	2,133	5.4	86.5
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,048	2.7	42.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,045	2.6	42.4
7	Chronic lower respiratory diseases (J40-J47)	1,019	2.6	41.3
8	Septicemia (A40-A41)	804	2.0	32.6
9	Human immunodeficiency virus (HIV) disease (B20-B24)	690	1.7	28.0
10	Chronic liver disease and cirrhosis (K70,K73-K74)	649	1.6	26.3
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	553	1.4	22.4
12	Influenza and pneumonia (J10-J18)	471	1.2	19.1
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	219	0.6	8.9
14	Aortic aneurysm and dissection (I71)	189	0.5	7.7
15	Viral hepatitis (B15-B19)	167	0.4	6.8
...	All other causes (Residual)	4,842	12.3	196.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Black, non-Hispanic, both sexes, 65-74 years			
...	All causes	53,803	100.0	3,300.7
1	Malignant neoplasms (C00-C97)	16,392	30.5	1,005.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,940	29.6	977.9
3	Cerebrovascular diseases (I60-I69)	3,684	6.8	226.0
4	Diabetes mellitus (E10-E14)	3,110	5.8	190.8
5	Chronic lower respiratory diseases (J40-J47)	1,962	3.6	120.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,698	3.2	104.2
7	Septicemia (A40-A41)	1,186	2.2	72.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	836	1.6	51.3
9	Influenza and pneumonia (J10-J18)	796	1.5	48.8
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	789	1.5	48.4
11	Chronic liver disease and cirrhosis (K70,K73-K74)	412	0.8	25.3
12	Alzheimer's disease (G30)	253	0.5	15.5
13	Pneumonitis due to solids and liquids (J69)	251	0.5	15.4
14	Aortic aneurysm and dissection (I71)	240	0.4	14.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	236	0.4	14.5
...	All other causes (Residual)	6,018	11.2	369.2
	Black, non-Hispanic, both sexes, 75-84 years			
...	All causes	61,592	100.0	6,645.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	19,681	32.0	2,123.4
2	Malignant neoplasms (C00-C97)	14,454	23.5	1,559.5
3	Cerebrovascular diseases (I60-I69)	5,257	8.5	567.2
4	Diabetes mellitus (E10-E14)	3,132	5.1	337.9
5	Chronic lower respiratory diseases (J40-J47)	2,190	3.6	236.3
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,891	3.1	204.0
7	Influenza and pneumonia (J10-J18)	1,571	2.6	169.5
8	Septicemia (A40-A41)	1,501	2.4	161.9
9	Alzheimer's disease (G30)	1,176	1.9	126.9
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,056	1.7	113.9
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	830	1.3	89.6
12	Pneumonitis due to solids and liquids (J69)	497	0.8	53.6
13	Aortic aneurysm and dissection (I71)	352	0.6	38.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	296	0.5	31.9
15	Atherosclerosis (I70)	285	0.5	30.7
...	All other causes (Residual)	7,423	12.1	800.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 85 years and over				
...	All causes	46,968	100.0	14,660.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	17,004	36.2	5,307.7
2	Malignant neoplasms (C00-C97)	6,239	13.3	1,947.5
3	Cerebrovascular diseases (I60-I69)	4,695	10.0	1,465.5
4	Influenza and pneumonia (J10-J18)	1,868	4.0	583.1
5	Diabetes mellitus (E10-E14)	1,710	3.6	533.8
6	Alzheimer's disease (G30)	1,644	3.5	513.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,387	3.0	432.9
8	Chronic lower respiratory diseases (J40-J47)	1,215	2.6	379.3
9	Septicemia (A40-A41)	1,181	2.5	368.6
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	934	2.0	291.5
11	Pneumonitis due to solids and liquids (J69)	600	1.3	187.3
12	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	589	1.3	183.9
13	Atherosclerosis (I70)	404	0.9	126.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	233	0.5	72.7
15	Parkinson's disease (G20-G21)	189	0.4	59.0
...	All other causes (Residual)	7,076	15.1	2,208.7
Black, non-Hispanic, male, all ages				
...	All causes	143,971	100.0	849.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	36,541	25.4	215.7
2	Malignant neoplasms (C00-C97)	32,335	22.5	190.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,381	5.8	49.5
4	Cerebrovascular diseases (I60-I69)	7,828	5.4	46.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6,658	4.6	39.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	5,227	3.6	30.8
7	Diabetes mellitus (E10-E14)	4,983	3.5	29.4
8	Chronic lower respiratory diseases (J40-J47)	4,143	2.9	24.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,158	2.2	18.6
10	Influenza and pneumonia (J10-J18)	2,769	1.9	16.3
11	Certain conditions originating in the perinatal period (P00-P96)	2,674	1.9	15.8
12	Septicemia (A40-A41)	2,611	1.8	15.4
13	Chronic liver disease and cirrhosis (K70,K73-K74)	1,819	1.3	10.7
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,698	1.2	10.0
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,578	1.1	9.3
...	All other causes (Residual)	21,568	15.0	127.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 1-4 years				
...	All causes	655	100.0	55.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	188	28.7	15.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	100	15.3	8.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	65	9.9	5.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	40	6.1	3.4
5	Malignant neoplasms (C00-C97)	29	4.4	2.4
6	Chronic lower respiratory diseases (J40-J47)	16	2.4	*
7	Certain conditions originating in the perinatal period (P00-P96)	14	2.1	*
8	Influenza and pneumonia (J10-J18)	13	2.0	*
9	Septicemia (A40-A41)	12	1.8	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	8	1.2	*
11	Anemias (D50-D64)	7	1.1	*
12	Cerebrovascular diseases (I60-I69)	5	0.8	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.6	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.5	*
...	All other causes (Residual)	148	22.6	12.5
Black, non-Hispanic, male, 5-14 years				
...	All causes	941	100.0	28.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	432	45.9	13.1
2	Malignant neoplasms (C00-C97)	100	10.6	3.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	66	7.0	2.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	34	3.6	1.0
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	33	3.5	1.0
6	Chronic lower respiratory diseases (J40-J47)	32	3.4	1.0
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	31	3.3	0.9
8	Human immunodeficiency virus (HIV) disease (B20-B24)	15	1.6	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	1.4	*
10	Anemias (D50-D64)	10	1.1	*
11	Influenza and pneumonia (J10-J18)	9	1.0	*
12	Diabetes mellitus (E10-E14)	6	0.6	*
13	Cerebrovascular diseases (I60-I69)	5	0.5	*
13	Certain conditions originating in the perinatal period (P00-P96)	5	0.5	*
15	Meningococcal infection (A39)	3	0.3	*
15	Septicemia (A40-A41)	3	0.3	*
15	Meningitis (G00,G03)	3	0.3	*
...	All other causes (Residual)	141	15.0	4.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 15-24 years				
...	All causes	5,377	100.0	186.2
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,551	47.4	88.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,285	23.9	44.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	388	7.2	13.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	205	3.8	7.1
5	Malignant neoplasms (C00-C97)	148	2.8	5.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	72	1.3	2.5
7	Human immunodeficiency virus (HIV) disease (B20-B24)	68	1.3	2.4
8	Chronic lower respiratory diseases (J40-J47)	48	0.9	1.7
9	Anemias (D50-D64)	41	0.8	1.4
10	Diabetes mellitus (E10-E14)	27	0.5	0.9
11	Septicemia (A40-A41)	25	0.5	0.9
12	Legal intervention (Y35,Y89.0)	24	0.4	0.8
13	Influenza and pneumonia (J10-J18)	23	0.4	0.8
14	Cerebrovascular diseases (I60-I69)	20	0.4	0.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	0.2	*
...	All other causes (Residual)	439	8.2	15.2
Black, non-Hispanic, male, 25-34 years				
...	All causes	6,548	100.0	267.9
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,998	30.5	81.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,192	18.2	48.8
3	Human immunodeficiency virus (HIV) disease (B20-B24)	753	11.5	30.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	578	8.8	23.6
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	406	6.2	16.6
6	Malignant neoplasms (C00-C97)	267	4.1	10.9
7	Diabetes mellitus (E10-E14)	122	1.9	5.0
8	Cerebrovascular diseases (I60-I69)	75	1.1	3.1
9	Anemias (D50-D64)	73	1.1	3.0
10	Chronic lower respiratory diseases (J40-J47)	69	1.1	2.8
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	54	0.8	2.2
12	Influenza and pneumonia (J10-J18)	52	0.8	2.1
13	Septicemia (A40-A41)	46	0.7	1.9
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	30	0.5	1.2
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	29	0.4	1.2
...	All other causes (Residual)	804	12.3	32.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 35-44 years				
...	All causes	11,971	100.0	456.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,071	17.3	78.9
2	Human immunodeficiency virus (HIV) disease (B20-B24)	2,040	17.0	77.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,608	13.4	61.3
4	Malignant neoplasms (C00-C97)	1,201	10.0	45.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,036	8.7	39.5
6	Cerebrovascular diseases (I60-I69)	355	3.0	13.5
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	329	2.7	12.5
8	Diabetes mellitus (E10-E14)	275	2.3	10.5
9	Chronic liver disease and cirrhosis (K70,K73-K74)	273	2.3	10.4
10	Influenza and pneumonia (J10-J18)	155	1.3	5.9
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	139	1.2	5.3
12	Septicemia (A40-A41)	128	1.1	4.9
13	Chronic lower respiratory diseases (J40-J47)	124	1.0	4.7
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	102	0.9	3.9
15	Viral hepatitis (B15-B19)	79	0.7	3.0
...	All other causes (Residual)	2,056	17.2	78.4
Black, non-Hispanic, male, 45-54 years				
...	All causes	20,307	100.0	1,014.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,955	24.4	247.6
2	Malignant neoplasms (C00-C97)	4,196	20.7	209.6
3	Human immunodeficiency virus (HIV) disease (B20-B24)	1,629	8.0	81.4
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,574	7.8	78.6
5	Cerebrovascular diseases (I60-I69)	998	4.9	49.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	702	3.5	35.1
7	Diabetes mellitus (E10-E14)	701	3.5	35.0
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	544	2.7	27.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	399	2.0	19.9
10	Septicemia (A40-A41)	304	1.5	15.2
11	Chronic lower respiratory diseases (J40-J47)	271	1.3	13.5
12	Influenza and pneumonia (J10-J18)	270	1.3	13.5
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	263	1.3	13.1
14	Viral hepatitis (B15-B19)	240	1.2	12.0
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	196	1.0	9.8
...	All other causes (Residual)	3,065	15.1	153.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 55-64 years				
...	All causes	22,600	100.0	2,055.4
1	Malignant neoplasms (C00-C97)	6,752	29.9	614.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,713	29.7	610.5
3	Cerebrovascular diseases (I60-I69)	1,252	5.5	113.9
4	Diabetes mellitus (E10-E14)	1,031	4.6	93.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	750	3.3	68.2
6	Chronic lower respiratory diseases (J40-J47)	554	2.5	50.4
7	Human immunodeficiency virus (HIV) disease (B20-B24)	524	2.3	47.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	509	2.3	46.3
9	Chronic liver disease and cirrhosis (K70,K73-K74)	462	2.0	42.0
10	Septicemia (A40-A41)	411	1.8	37.4
11	Influenza and pneumonia (J10-J18)	293	1.3	26.6
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	284	1.3	25.8
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	165	0.7	15.0
14	Aortic aneurysm and dissection (I71)	124	0.5	11.3
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	102	0.5	9.3
...	All other causes (Residual)	2,674	11.8	243.2
Black, non-Hispanic, male, 65-74 years				
...	All causes	28,446	100.0	4,218.9
1	Malignant neoplasms (C00-C97)	9,134	32.1	1,354.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,491	29.8	1,259.3
3	Cerebrovascular diseases (I60-I69)	1,844	6.5	273.5
4	Diabetes mellitus (E10-E14)	1,278	4.5	189.5
5	Chronic lower respiratory diseases (J40-J47)	1,147	4.0	170.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	782	2.7	116.0
7	Septicemia (A40-A41)	586	2.1	86.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	505	1.8	74.9
9	Influenza and pneumonia (J10-J18)	472	1.7	70.0
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	374	1.3	55.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	268	0.9	39.7
12	Human immunodeficiency virus (HIV) disease (B20-B24)	144	0.5	21.4
13	Pneumonitis due to solids and liquids (J69)	140	0.5	20.8
14	Aortic aneurysm and dissection (I71)	134	0.5	19.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	119	0.4	17.6
...	All other causes (Residual)	3,028	10.6	449.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 75-84 years				
...	All causes	28,083	100.0	8,426.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,621	30.7	2,586.9
2	Malignant neoplasms (C00-C97)	7,700	27.4	2,310.5
3	Cerebrovascular diseases (I60-I69)	2,068	7.4	620.5
4	Chronic lower respiratory diseases (J40-J47)	1,285	4.6	385.6
5	Diabetes mellitus (E10-E14)	1,150	4.1	345.1
6	Influenza and pneumonia (J10-J18)	808	2.9	242.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	784	2.8	235.3
8	Septicemia (A40-A41)	656	2.3	196.8
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	461	1.6	138.3
10	Alzheimer's disease (G30)	412	1.5	123.6
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	399	1.4	119.7
12	Pneumonitis due to solids and liquids (J69)	252	0.9	75.6
13	Aortic aneurysm and dissection (I71)	158	0.6	47.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	141	0.5	42.3
15	Parkinson's disease (G20-G21)	127	0.5	38.1
...	All other causes (Residual)	3,061	10.9	918.5
Black, non-Hispanic, male, 85 years and over				
...	All causes	14,352	100.0	16,576.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,758	33.2	5,495.4
2	Malignant neoplasms (C00-C97)	2,804	19.5	3,238.5
3	Cerebrovascular diseases (I60-I69)	1,177	8.2	1,359.4
4	Influenza and pneumonia (J10-J18)	622	4.3	718.4
5	Chronic lower respiratory diseases (J40-J47)	591	4.1	682.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	448	3.1	517.4
7	Diabetes mellitus (E10-E14)	392	2.7	452.7
8	Septicemia (A40-A41)	373	2.6	430.8
9	Alzheimer's disease (G30)	364	2.5	420.4
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	240	1.7	277.2
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	222	1.5	256.4
12	Pneumonitis due to solids and liquids (J69)	211	1.5	243.7
13	Atherosclerosis (I70)	98	0.7	113.2
14	Parkinson's disease (G20-G21)	86	0.6	99.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	76	0.5	87.8
...	All other causes (Residual)	1,890	13.2	2,182.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, all ages				
...	All causes	140,372	100.0	751.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	40,253	28.7	215.4
2	Malignant neoplasms (C00-C97)	29,245	20.8	156.5
3	Cerebrovascular diseases (I60-I69)	11,016	7.8	59.0
4	Diabetes mellitus (E10-E14)	7,184	5.1	38.4
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,056	2.9	21.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,858	2.7	20.6
7	Chronic lower respiratory diseases (J40-J47)	3,361	2.4	18.0
8	Septicemia (A40-A41)	3,218	2.3	17.2
9	Influenza and pneumonia (J10-J18)	2,917	2.1	15.6
10	Human immunodeficiency virus (HIV) disease (B20-B24)	2,483	1.8	13.3
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,314	1.6	12.4
12	Alzheimer's disease (G30)	2,198	1.6	11.8
13	Certain conditions originating in the perinatal period (P00-P96)	2,039	1.5	10.9
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,419	1.0	7.6
15	Chronic liver disease and cirrhosis (K70,K73-K74)	914	0.7	4.9
...	All other causes (Residual)	23,897	17.0	127.9
Black, non-Hispanic, female, 1-4 years				
...	All causes	501	100.0	43.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	143	28.5	12.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	66	13.2	5.7
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	62	12.4	5.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	33	6.6	2.9
5	Malignant neoplasms (C00-C97)	17	3.4	*
6	Influenza and pneumonia (J10-J18)	11	2.2	*
7	Septicemia (A40-A41)	9	1.8	*
8	Certain conditions originating in the perinatal period (P00-P96)	8	1.6	*
9	Cerebrovascular diseases (I60-I69)	7	1.4	*
10	Chronic lower respiratory diseases (J40-J47)	6	1.2	*
11	Anemias (D50-D64)	4	0.8	*
12	Meningococcal infection (A39)	3	0.6	*
12	Meningitis (G00,G03)	3	0.6	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.6	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.4	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.4	*
...	All other causes (Residual)	122	24.4	10.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 5-14 years				
...	All causes	616	100.0	19.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	198	32.1	6.2
2	Malignant neoplasms (C00-C97)	72	11.7	2.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	40	6.5	1.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	36	5.8	1.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	26	4.2	0.8
6	Chronic lower respiratory diseases (J40-J47)	21	3.4	0.7
7	Anemias (D50-D64)	16	2.6	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	2.1	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	11	1.8	*
10	Diabetes mellitus (E10-E14)	9	1.5	*
10	Cerebrovascular diseases (I60-I69)	9	1.5	*
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	9	1.5	*
13	Influenza and pneumonia (J10-J18)	8	1.3	*
14	Septicemia (A40-A41)	7	1.1	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	1.1	*
...	All other causes (Residual)	134	21.8	4.2
Black, non-Hispanic, female, 15-24 years				
...	All causes	1,647	100.0	56.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	428	26.0	14.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	266	16.2	9.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	117	7.1	4.0
4	Malignant neoplasms (C00-C97)	109	6.6	3.8
5	Human immunodeficiency virus (HIV) disease (B20-B24)	93	5.6	3.2
6	Pregnancy, childbirth and the puerperium (O00-099)	45	2.7	1.6
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	41	2.5	1.4
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	37	2.2	1.3
9	Anemias (D50-D64)	31	1.9	1.1
9	Cerebrovascular diseases (I60-I69)	31	1.9	1.1
11	Diabetes mellitus (E10-E14)	29	1.8	1.0
11	Chronic lower respiratory diseases (J40-J47)	29	1.8	1.0
13	Influenza and pneumonia (J10-J18)	22	1.3	0.8
14	Septicemia (A40-A41)	16	1.0	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	0.9	*
...	All other causes (Residual)	338	20.5	11.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 25-34 years				
...	All causes	3,311	100.0	122.4
1	Human immunodeficiency virus (HIV) disease (B20-B24)	481	14.5	17.8
2	Malignant neoplasms (C00-C97)	424	12.8	15.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	408	12.3	15.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	383	11.6	14.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	337	10.2	12.5
6	Cerebrovascular diseases (I60-I69)	78	2.4	2.9
7	Diabetes mellitus (E10-E14)	76	2.3	2.8
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	66	2.0	2.4
9	Pregnancy, childbirth and the puerperium (O00-099)	65	2.0	2.4
10	Chronic lower respiratory diseases (J40-J47)	59	1.8	2.2
11	Septicemia (A40-A41)	52	1.6	1.9
12	Anemias (D50-D64)	44	1.3	1.6
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	39	1.2	1.4
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	37	1.1	1.4
15	Influenza and pneumonia (J10-J18)	35	1.1	1.3
...	All other causes (Residual)	727	22.0	26.9
Black, non-Hispanic, female, 35-44 years				
...	All causes	8,465	100.0	285.7
1	Malignant neoplasms (C00-C97)	1,749	20.7	59.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,353	16.0	45.7
3	Human immunodeficiency virus (HIV) disease (B20-B24)	1,032	12.2	34.8
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	674	8.0	22.7
5	Cerebrovascular diseases (I60-I69)	403	4.8	13.6
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	391	4.6	13.2
7	Diabetes mellitus (E10-E14)	231	2.7	7.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	167	2.0	5.6
9	Chronic lower respiratory diseases (J40-J47)	149	1.8	5.0
10	Chronic liver disease and cirrhosis (K70,K73-K74)	147	1.7	5.0
11	Septicemia (A40-A41)	140	1.7	4.7
12	Influenza and pneumonia (J10-J18)	119	1.4	4.0
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	89	1.1	3.0
14	Anemias (D50-D64)	88	1.0	3.0
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	75	0.9	2.5
...	All other causes (Residual)	1,658	19.6	56.0

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Black, non-Hispanic, female, 45-54 years			
...	All causes	13,791	100.0	591.5
1	Malignant neoplasms (C00-C97)	3,979	28.9	170.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,001	21.8	128.7
3	Cerebrovascular diseases (I60-I69)	888	6.4	38.1
4	Human immunodeficiency virus (HIV) disease (B20-B24)	620	4.5	26.6
5	Diabetes mellitus (E10-E14)	605	4.4	25.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	529	3.8	22.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	303	2.2	13.0
8	Septicemia (A40-A41)	291	2.1	12.5
8	Chronic liver disease and cirrhosis (K70,K73-K74)	291	2.1	12.5
10	Chronic lower respiratory diseases (J40-J47)	278	2.0	11.9
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	176	1.3	7.5
12	Influenza and pneumonia (J10-J18)	160	1.2	6.9
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	160	1.2	6.9
14	Viral hepatitis (B15-B19)	109	0.8	4.7
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	72	0.5	3.1
...	All other causes (Residual)	2,329	16.9	99.9
	Black, non-Hispanic, female, 55-64 years			
...	All causes	16,923	100.0	1,238.2
1	Malignant neoplasms (C00-C97)	5,444	32.2	398.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,496	26.6	328.9
3	Diabetes mellitus (E10-E14)	1,102	6.5	80.6
4	Cerebrovascular diseases (I60-I69)	1,037	6.1	75.9
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	539	3.2	39.4
6	Chronic lower respiratory diseases (J40-J47)	465	2.7	34.0
7	Septicemia (A40-A41)	393	2.3	28.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	295	1.7	21.6
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	269	1.6	19.7
10	Chronic liver disease and cirrhosis (K70,K73-K74)	187	1.1	13.7
11	Influenza and pneumonia (J10-J18)	178	1.1	13.0
12	Human immunodeficiency virus (HIV) disease (B20-B24)	166	1.0	12.1
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	74	0.4	5.4
14	Viral hepatitis (B15-B19)	68	0.4	5.0
15	Aortic aneurysm and dissection (I71)	65	0.4	4.8
...	All other causes (Residual)	2,145	12.7	156.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Black, non-Hispanic, female, 65-74 years			
...	All causes	25,357	100.0	2,653.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,449	29.4	779.4
2	Malignant neoplasms (C00-C97)	7,258	28.6	759.4
3	Cerebrovascular diseases (I60-I69)	1,840	7.3	192.5
4	Diabetes mellitus (E10-E14)	1,832	7.2	191.7
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	916	3.6	95.8
6	Chronic lower respiratory diseases (J40-J47)	815	3.2	85.3
7	Septicemia (A40-A41)	600	2.4	62.8
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	415	1.6	43.4
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	331	1.3	34.6
10	Influenza and pneumonia (J10-J18)	324	1.3	33.9
11	Chronic liver disease and cirrhosis (K70,K73-K74)	144	0.6	15.1
12	Alzheimer's disease (G30)	143	0.6	15.0
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	117	0.5	12.2
14	Pneumonitis due to solids and liquids (J69)	111	0.4	11.6
15	Aortic aneurysm and dissection (I71)	106	0.4	11.1
...	All other causes (Residual)	2,956	11.7	309.3
	Black, non-Hispanic, female, 75-84 years			
...	All causes	33,509	100.0	5,645.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,060	33.0	1,863.2
2	Malignant neoplasms (C00-C97)	6,754	20.2	1,137.8
3	Cerebrovascular diseases (I60-I69)	3,189	9.5	537.2
4	Diabetes mellitus (E10-E14)	1,982	5.9	333.9
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,107	3.3	186.5
6	Chronic lower respiratory diseases (J40-J47)	905	2.7	152.5
7	Septicemia (A40-A41)	845	2.5	142.4
8	Alzheimer's disease (G30)	764	2.3	128.7
9	Influenza and pneumonia (J10-J18)	763	2.3	128.5
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	657	2.0	110.7
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	369	1.1	62.2
12	Pneumonitis due to solids and liquids (J69)	245	0.7	41.3
13	Aortic aneurysm and dissection (I71)	194	0.6	32.7
14	Atherosclerosis (I70)	161	0.5	27.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	155	0.5	26.1
...	All other causes (Residual)	4,359	13.0	734.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Black, non-Hispanic, female, 85 years and over			
...	All causes	32,616	100.0	13,951.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,246	37.5	5,238.2
2	Cerebrovascular diseases (I60-I69)	3,518	10.8	1,504.8
3	Malignant neoplasms (C00-C97)	3,435	10.5	1,469.3
4	Diabetes mellitus (E10-E14)	1,318	4.0	563.8
5	Alzheimer's disease (G30)	1,280	3.9	547.5
6	Influenza and pneumonia (J10-J18)	1,246	3.8	533.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	939	2.9	401.7
8	Septicemia (A40-A41)	808	2.5	345.6
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	694	2.1	296.9
10	Chronic lower respiratory diseases (J40-J47)	624	1.9	266.9
11	Pneumonitis due to solids and liquids (J69)	389	1.2	166.4
12	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	367	1.1	157.0
13	Atherosclerosis (I70)	306	0.9	130.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	157	0.5	67.2
15	Aortic aneurysm and dissection (I71)	136	0.4	58.2
...	All other causes (Residual)	5,153	15.8	2,204.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable