

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, all ages				
...	All causes	117,135	100.0	302.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	27,887	23.8	71.9
2	Malignant neoplasms (C00-C97)	23,141	19.8	59.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,106	8.6	26.1
4	Cerebrovascular diseases (I60-I69)	6,451	5.5	16.6
5	Diabetes mellitus (E10-E14)	5,912	5.0	15.3
6	Chronic liver disease and cirrhosis (K70,K73-K74)	3,409	2.9	8.8
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,129	2.7	8.1
8	Chronic lower respiratory diseases (J40-J47)	3,058	2.6	7.9
9	Influenza and pneumonia (J10-J18)	2,824	2.4	7.3
10	Certain conditions originating in the perinatal period (P00-P96)	2,402	2.1	6.2
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,954	1.7	5.0
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,898	1.6	4.9
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,855	1.6	4.8
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,715	1.5	4.4
15	Septicemia (A40-A41)	1,521	1.3	3.9
...	All other causes (Residual)	19,873	17.0	51.3
Hispanic, both sexes, 1-4 years				
...	All causes	933	100.0	29.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	341	36.5	10.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	111	11.9	3.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	81	8.7	2.6
4	Malignant neoplasms (C00-C97)	72	7.7	2.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	24	2.6	0.8
6	Influenza and pneumonia (J10-J18)	17	1.8	*
7	Septicemia (A40-A41)	16	1.7	*
8	Certain conditions originating in the perinatal period (P00-P96)	14	1.5	*
9	Cerebrovascular diseases (I60-I69)	12	1.3	*
10	Chronic lower respiratory diseases (J40-J47)	9	1.0	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	7	0.8	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.8	*
13	Meningitis (G00,G03)	5	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.5	*
15	Anemias (D50-D64)	3	0.3	*
...	All other causes (Residual)	209	22.4	6.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 5-14 years				
...	All causes	1,124	100.0	15.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	389	34.6	5.4
2	Malignant neoplasms (C00-C97)	210	18.7	2.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	70	6.2	1.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	64	5.7	0.9
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	37	3.3	0.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	30	2.7	0.4
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	26	2.3	0.4
8	Cerebrovascular diseases (I60-I69)	20	1.8	0.3
9	Septicemia (A40-A41)	14	1.2	*
9	Influenza and pneumonia (J10-J18)	14	1.2	*
11	Chronic lower respiratory diseases (J40-J47)	13	1.2	*
12	Certain conditions originating in the perinatal period (P00-P96)	6	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.4	*
15	Anemias (D50-D64)	4	0.4	*
...	All other causes (Residual)	217	19.3	3.0
Hispanic, both sexes, 15-24 years				
...	All causes	5,255	100.0	77.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,413	45.9	35.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,200	22.8	17.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	455	8.7	6.7
4	Malignant neoplasms (C00-C97)	304	5.8	4.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	128	2.4	1.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	67	1.3	1.0
7	Human immunodeficiency virus (HIV) disease (B20-B24)	29	0.6	0.4
7	Cerebrovascular diseases (I60-I69)	29	0.6	0.4
9	Legal intervention (Y35,Y89.0)	22	0.4	0.3
10	Influenza and pneumonia (J10-J18)	19	0.4	*
10	Pregnancy, childbirth and the puerperium (O00-099)	19	0.4	*
12	Septicemia (A40-A41)	15	0.3	*
12	Diabetes mellitus (E10-E14)	15	0.3	*
12	Chronic lower respiratory diseases (J40-J47)	15	0.3	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	9	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.2	*
...	All other causes (Residual)	507	9.6	7.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 25-34 years				
...	All causes	5,816	100.0	79.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,027	34.9	27.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	903	15.5	12.3
3	Malignant neoplasms (C00-C97)	502	8.6	6.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	484	8.3	6.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	321	5.5	4.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	262	4.5	3.6
7	Cerebrovascular diseases (I60-I69)	93	1.6	1.3
8	Chronic liver disease and cirrhosis (K70,K73-K74)	81	1.4	1.1
9	Diabetes mellitus (E10-E14)	56	1.0	0.8
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	53	0.9	0.7
11	Influenza and pneumonia (J10-J18)	43	0.7	0.6
12	Septicemia (A40-A41)	38	0.7	0.5
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	33	0.6	0.5
14	Chronic lower respiratory diseases (J40-J47)	31	0.5	0.4
15	Pregnancy, childbirth and the puerperium (O00-099)	28	0.5	0.4
...	All other causes (Residual)	861	14.8	11.7
Hispanic, both sexes, 35-44 years				
...	All causes	8,467	100.0	145.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,841	21.7	31.7
2	Malignant neoplasms (C00-C97)	1,346	15.9	23.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	852	10.1	14.7
4	Human immunodeficiency virus (HIV) disease (B20-B24)	779	9.2	13.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	507	6.0	8.7
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	464	5.5	8.0
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	407	4.8	7.0
8	Cerebrovascular diseases (I60-I69)	256	3.0	4.4
9	Diabetes mellitus (E10-E14)	205	2.4	3.5
10	Viral hepatitis (B15-B19)	101	1.2	1.7
11	Influenza and pneumonia (J10-J18)	77	0.9	1.3
12	Chronic lower respiratory diseases (J40-J47)	70	0.8	1.2
13	Septicemia (A40-A41)	67	0.8	1.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	65	0.8	1.1
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	40	0.5	0.7
...	All other causes (Residual)	1,390	16.4	23.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 45-54 years				
...	All causes	11,745	100.0	321.3
1	Malignant neoplasms (C00-C97)	2,859	24.3	78.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,020	17.2	55.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,168	9.9	32.0
4	Chronic liver disease and cirrhosis (K70,K73-K74)	983	8.4	26.9
5	Cerebrovascular diseases (I60-I69)	560	4.8	15.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	537	4.6	14.7
7	Diabetes mellitus (E10-E14)	513	4.4	14.0
8	Viral hepatitis (B15-B19)	291	2.5	8.0
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	260	2.2	7.1
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	216	1.8	5.9
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	157	1.3	4.3
12	Septicemia (A40-A41)	136	1.2	3.7
13	Chronic lower respiratory diseases (J40-J47)	127	1.1	3.5
14	Influenza and pneumonia (J10-J18)	124	1.1	3.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	54	0.5	1.5
...	All other causes (Residual)	1,740	14.8	47.6
Hispanic, both sexes, 55-64 years				
...	All causes	14,215	100.0	719.8
1	Malignant neoplasms (C00-C97)	4,247	29.9	215.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,492	24.6	176.8
3	Diabetes mellitus (E10-E14)	1,029	7.2	52.1
4	Chronic liver disease and cirrhosis (K70,K73-K74)	758	5.3	38.4
5	Cerebrovascular diseases (I60-I69)	708	5.0	35.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	566	4.0	28.7
7	Chronic lower respiratory diseases (J40-J47)	274	1.9	13.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	267	1.9	13.5
9	Influenza and pneumonia (J10-J18)	207	1.5	10.5
10	Human immunodeficiency virus (HIV) disease (B20-B24)	188	1.3	9.5
11	Septicemia (A40-A41)	165	1.2	8.4
12	Viral hepatitis (B15-B19)	164	1.2	8.3
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	142	1.0	7.2
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	123	0.9	6.2
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	87	0.6	4.4
...	All other causes (Residual)	1,798	12.6	91.0

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 65-74 years				
...	All causes	20,476	100.0	1,734.1
1	Malignant neoplasms (C00-C97)	5,857	28.6	496.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,719	27.9	484.3
3	Diabetes mellitus (E10-E14)	1,591	7.8	134.7
4	Cerebrovascular diseases (I60-I69)	1,215	5.9	102.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	663	3.2	56.2
6	Chronic lower respiratory diseases (J40-J47)	635	3.1	53.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	466	2.3	39.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	427	2.1	36.2
9	Influenza and pneumonia (J10-J18)	399	1.9	33.8
10	Septicemia (A40-A41)	341	1.7	28.9
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	183	0.9	15.5
12	Alzheimer's disease (G30)	127	0.6	10.8
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	105	0.5	8.9
14	Viral hepatitis (B15-B19)	102	0.5	8.6
14	Parkinson's disease (G20-G21)	102	0.5	8.6
...	All other causes (Residual)	2,544	12.4	215.5
Hispanic, both sexes, 75-84 years				
...	All causes	24,838	100.0	4,143.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,779	31.3	1,297.6
2	Malignant neoplasms (C00-C97)	5,365	21.6	894.9
3	Cerebrovascular diseases (I60-I69)	1,918	7.7	319.9
4	Diabetes mellitus (E10-E14)	1,712	6.9	285.6
5	Chronic lower respiratory diseases (J40-J47)	1,055	4.2	176.0
6	Influenza and pneumonia (J10-J18)	855	3.4	142.6
7	Alzheimer's disease (G30)	535	2.2	89.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	532	2.1	88.7
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	464	1.9	77.4
10	Septicemia (A40-A41)	382	1.5	63.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	321	1.3	53.5
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	283	1.1	47.2
13	Parkinson's disease (G20-G21)	223	0.9	37.2
14	Pneumonitis due to solids and liquids (J69)	174	0.7	29.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	148	0.6	24.7
...	All other causes (Residual)	3,092	12.4	515.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 85 years and over				
...	All causes	19,265	100.0	10,679.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,436	38.6	4,121.9
2	Malignant neoplasms (C00-C97)	2,361	12.3	1,308.8
3	Cerebrovascular diseases (I60-I69)	1,620	8.4	898.0
4	Influenza and pneumonia (J10-J18)	1,021	5.3	566.0
5	Chronic lower respiratory diseases (J40-J47)	825	4.3	457.3
6	Alzheimer's disease (G30)	791	4.1	438.5
7	Diabetes mellitus (E10-E14)	790	4.1	437.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	363	1.9	201.2
9	Septicemia (A40-A41)	282	1.5	156.3
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	276	1.4	153.0
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	268	1.4	148.6
12	Pneumonitis due to solids and liquids (J69)	237	1.2	131.4
13	Atherosclerosis (I70)	196	1.0	108.6
14	Parkinson's disease (G20-G21)	138	0.7	76.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	88	0.5	48.8
...	All other causes (Residual)	2,573	13.4	1,426.3
Hispanic, male, all ages				
...	All causes	65,703	100.0	328.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,798	22.5	74.0
2	Malignant neoplasms (C00-C97)	12,235	18.6	61.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,698	11.7	38.5
4	Cerebrovascular diseases (I60-I69)	3,003	4.6	15.0
5	Diabetes mellitus (E10-E14)	2,779	4.2	13.9
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,635	4.0	13.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,437	3.7	12.2
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,651	2.5	8.3
9	Chronic lower respiratory diseases (J40-J47)	1,625	2.5	8.1
10	Human immunodeficiency virus (HIV) disease (B20-B24)	1,440	2.2	7.2
11	Influenza and pneumonia (J10-J18)	1,398	2.1	7.0
12	Certain conditions originating in the perinatal period (P00-P96)	1,352	2.1	6.8
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	937	1.4	4.7
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	903	1.4	4.5
15	Septicemia (A40-A41)	744	1.1	3.7
...	All other causes (Residual)	10,068	15.3	50.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 1-4 years				
...	All causes	546	100.0	34.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	215	39.4	13.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	60	11.0	3.8
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	49	9.0	3.1
4	Malignant neoplasms (C00-C97)	35	6.4	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	11	2.0	*
5	Influenza and pneumonia (J10-J18)	11	2.0	*
7	Septicemia (A40-A41)	9	1.6	*
7	Certain conditions originating in the perinatal period (P00-P96)	9	1.6	*
9	Cerebrovascular diseases (I60-I69)	5	0.9	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	4	0.7	*
10	Meningitis (G00,G03)	4	0.7	*
10	Chronic lower respiratory diseases (J40-J47)	4	0.7	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.7	*
14	Anemias (D50-D64)	1	0.2	*
14	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
...	All other causes (Residual)	123	22.5	7.7
Hispanic, male, 5-14 years				
...	All causes	647	100.0	17.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	244	37.7	6.6
2	Malignant neoplasms (C00-C97)	108	16.7	2.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	37	5.7	1.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	35	5.4	0.9
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	21	3.2	0.6
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	15	2.3	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	14	2.2	*
8	Septicemia (A40-A41)	8	1.2	*
8	Chronic lower respiratory diseases (J40-J47)	8	1.2	*
10	Cerebrovascular diseases (I60-I69)	7	1.1	*
11	Influenza and pneumonia (J10-J18)	6	0.9	*
12	Certain conditions originating in the perinatal period (P00-P96)	5	0.8	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.6	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	3	0.5	*
14	Anemias (D50-D64)	3	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.5	*
...	All other causes (Residual)	126	19.5	3.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 15-24 years				
...	All causes	4,182	100.0	114.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,958	46.8	53.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,081	25.8	29.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	388	9.3	10.6
4	Malignant neoplasms (C00-C97)	191	4.6	5.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	94	2.2	2.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	39	0.9	1.1
7	Legal intervention (Y35,Y89.0)	22	0.5	0.6
8	Human immunodeficiency virus (HIV) disease (B20-B24)	19	0.5	*
9	Cerebrovascular diseases (I60-I69)	17	0.4	*
10	Chronic lower respiratory diseases (J40-J47)	12	0.3	*
11	Influenza and pneumonia (J10-J18)	11	0.3	*
12	Diabetes mellitus (E10-E14)	10	0.2	*
13	Septicemia (A40-A41)	8	0.2	*
14	Viral hepatitis (B15-B19)	5	0.1	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	5	0.1	*
14	Anemias (D50-D64)	5	0.1	*
14	Meningitis (G00,G03)	5	0.1	*
14	Aortic aneurysm and dissection (I71)	5	0.1	*
14	Pneumonitis due to solids and liquids (J69)	5	0.1	*
14	Chronic liver disease and cirrhosis (K70,K73-K74)	5	0.1	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.1	*
...	All other causes (Residual)	292	7.0	8.0
Hispanic, male, 25-34 years				
...	All causes	4,474	100.0	112.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,709	38.2	43.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	789	17.6	19.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	426	9.5	10.7
4	Malignant neoplasms (C00-C97)	252	5.6	6.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	242	5.4	6.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	201	4.5	5.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	62	1.4	1.6
8	Cerebrovascular diseases (I60-I69)	61	1.4	1.5
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	36	0.8	0.9
10	Diabetes mellitus (E10-E14)	34	0.8	0.9
11	Influenza and pneumonia (J10-J18)	26	0.6	0.7
12	Septicemia (A40-A41)	25	0.6	0.6
13	Legal intervention (Y35,Y89.0)	21	0.5	0.5
14	Chronic lower respiratory diseases (J40-J47)	19	0.4	*
15	Viral hepatitis (B15-B19)	14	0.3	*
...	All other causes (Residual)	557	12.4	14.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 35-44 years				
...	All causes	5,828	100.0	192.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,468	25.2	48.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	625	10.7	20.6
3	Human immunodeficiency virus (HIV) disease (B20-B24)	607	10.4	20.1
4	Malignant neoplasms (C00-C97)	557	9.6	18.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	410	7.0	13.5
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	367	6.3	12.1
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	340	5.8	11.2
8	Cerebrovascular diseases (I60-I69)	154	2.6	5.1
9	Diabetes mellitus (E10-E14)	124	2.1	4.1
10	Viral hepatitis (B15-B19)	75	1.3	2.5
11	Influenza and pneumonia (J10-J18)	52	0.9	1.7
12	Septicemia (A40-A41)	47	0.8	1.6
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	38	0.7	1.3
14	Chronic lower respiratory diseases (J40-J47)	31	0.5	1.0
15	Aortic aneurysm and dissection (I71)	20	0.3	0.7
...	All other causes (Residual)	913	15.7	30.2
Hispanic, male, 45-54 years				
...	All causes	7,719	100.0	423.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,467	19.0	80.5
2	Malignant neoplasms (C00-C97)	1,430	18.5	78.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	888	11.5	48.7
4	Chronic liver disease and cirrhosis (K70,K73-K74)	805	10.4	44.2
5	Human immunodeficiency virus (HIV) disease (B20-B24)	421	5.5	23.1
6	Cerebrovascular diseases (I60-I69)	340	4.4	18.6
7	Diabetes mellitus (E10-E14)	305	4.0	16.7
8	Viral hepatitis (B15-B19)	218	2.8	12.0
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	206	2.7	11.3
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	166	2.2	9.1
11	Septicemia (A40-A41)	94	1.2	5.2
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	88	1.1	4.8
13	Influenza and pneumonia (J10-J18)	86	1.1	4.7
14	Chronic lower respiratory diseases (J40-J47)	70	0.9	3.8
15	Aortic aneurysm and dissection (I71)	29	0.4	1.6
...	All other causes (Residual)	1,106	14.3	60.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 55-64 years				
...	All causes	8,762	100.0	937.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,393	27.3	256.0
2	Malignant neoplasms (C00-C97)	2,377	27.1	254.3
3	Chronic liver disease and cirrhosis (K70,K73-K74)	561	6.4	60.0
4	Diabetes mellitus (E10-E14)	551	6.3	58.9
5	Cerebrovascular diseases (I60-I69)	421	4.8	45.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	407	4.6	43.5
7	Chronic lower respiratory diseases (J40-J47)	164	1.9	17.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	141	1.6	15.1
9	Human immunodeficiency virus (HIV) disease (B20-B24)	140	1.6	15.0
10	Influenza and pneumonia (J10-J18)	130	1.5	13.9
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	123	1.4	13.2
12	Viral hepatitis (B15-B19)	108	1.2	11.6
13	Septicemia (A40-A41)	84	1.0	9.0
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	72	0.8	7.7
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	66	0.8	7.1
...	All other causes (Residual)	1,024	11.7	109.6
Hispanic, male, 65-74 years				
...	All causes	11,476	100.0	2,193.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,441	30.0	657.7
2	Malignant neoplasms (C00-C97)	3,256	28.4	622.3
3	Diabetes mellitus (E10-E14)	752	6.6	143.7
4	Cerebrovascular diseases (I60-I69)	652	5.7	124.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	407	3.5	77.8
6	Chronic lower respiratory diseases (J40-J47)	362	3.2	69.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	307	2.7	58.7
8	Influenza and pneumonia (J10-J18)	229	2.0	43.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	214	1.9	40.9
10	Septicemia (A40-A41)	161	1.4	30.8
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	80	0.7	15.3
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	77	0.7	14.7
13	Parkinson's disease (G20-G21)	71	0.6	13.6
14	Aortic aneurysm and dissection (I71)	63	0.5	12.0
15	Alzheimer's disease (G30)	55	0.5	10.5
...	All other causes (Residual)	1,349	11.8	257.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 75-84 years				
...	All causes	12,291	100.0	5,043.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,898	31.7	1,599.5
2	Malignant neoplasms (C00-C97)	2,902	23.6	1,190.8
3	Cerebrovascular diseases (I60-I69)	825	6.7	338.5
4	Diabetes mellitus (E10-E14)	748	6.1	306.9
5	Chronic lower respiratory diseases (J40-J47)	593	4.8	243.3
6	Influenza and pneumonia (J10-J18)	422	3.4	173.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	273	2.2	112.0
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	258	2.1	105.9
9	Alzheimer's disease (G30)	196	1.6	80.4
10	Septicemia (A40-A41)	176	1.4	72.2
11	Chronic liver disease and cirrhosis (K70,K73-K74)	157	1.3	64.4
12	Parkinson's disease (G20-G21)	122	1.0	50.1
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	113	0.9	46.4
14	Pneumonitis due to solids and liquids (J69)	105	0.9	43.1
15	Aortic aneurysm and dissection (I71)	78	0.6	32.0
...	All other causes (Residual)	1,425	11.6	584.7
Hispanic, male, 85 years and over				
...	All causes	6,977	100.0	11,674.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,571	36.8	4,301.8
2	Malignant neoplasms (C00-C97)	1,117	16.0	1,869.0
3	Cerebrovascular diseases (I60-I69)	512	7.3	856.7
4	Influenza and pneumonia (J10-J18)	392	5.6	655.9
5	Chronic lower respiratory diseases (J40-J47)	360	5.2	602.4
6	Diabetes mellitus (E10-E14)	254	3.6	425.0
7	Alzheimer's disease (G30)	204	2.9	341.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	141	2.0	235.9
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	134	1.9	224.2
10	Pneumonitis due to solids and liquids (J69)	100	1.4	167.3
11	Septicemia (A40-A41)	98	1.4	164.0
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	79	1.1	132.2
13	Parkinson's disease (G20-G21)	68	1.0	113.8
14	Atherosclerosis (I70)	61	0.9	102.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	34	0.5	56.9
...	All other causes (Residual)	852	12.2	1,425.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, all ages				
...	All causes	51,432	100.0	274.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,089	25.4	69.7
2	Malignant neoplasms (C00-C97)	10,906	21.2	58.1
3	Cerebrovascular diseases (I60-I69)	3,448	6.7	18.4
4	Diabetes mellitus (E10-E14)	3,133	6.1	16.7
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,408	4.7	12.8
6	Chronic lower respiratory diseases (J40-J47)	1,433	2.8	7.6
7	Influenza and pneumonia (J10-J18)	1,426	2.8	7.6
8	Certain conditions originating in the perinatal period (P00-P96)	1,050	2.0	5.6
9	Alzheimer's disease (G30)	1,010	2.0	5.4
10	Chronic liver disease and cirrhosis (K70,K73-K74)	972	1.9	5.2
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	961	1.9	5.1
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	812	1.6	4.3
13	Septicemia (A40-A41)	777	1.5	4.1
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	557	1.1	3.0
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	494	1.0	2.6
...	All other causes (Residual)	8,956	17.4	47.7
Hispanic, female, 1-4 years				
...	All causes	387	100.0	25.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	126	32.6	8.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	51	13.2	3.3
3	Malignant neoplasms (C00-C97)	37	9.6	2.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	32	8.3	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	13	3.4	*
6	Septicemia (A40-A41)	7	1.8	*
6	Cerebrovascular diseases (I60-I69)	7	1.8	*
8	Influenza and pneumonia (J10-J18)	6	1.6	*
9	Chronic lower respiratory diseases (J40-J47)	5	1.3	*
9	Certain conditions originating in the perinatal period (P00-P96)	5	1.3	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	1.0	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	3	0.8	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.8	*
14	Anemias (D50-D64)	2	0.5	*
15	Meningitis (G00,G03)	1	0.3	*
15	Acute bronchitis and bronchiolitis (J20-J21)	1	0.3	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.3	*
...	All other causes (Residual)	83	21.4	5.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 5-14 years				
...	All causes	477	100.0	13.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	145	30.4	4.1
2	Malignant neoplasms (C00-C97)	102	21.4	2.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	35	7.3	1.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	27	5.7	0.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	16	3.4	*
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	16	3.4	*
7	Cerebrovascular diseases (I60-I69)	13	2.7	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	11	2.3	*
9	Influenza and pneumonia (J10-J18)	8	1.7	*
10	Septicemia (A40-A41)	6	1.3	*
11	Chronic lower respiratory diseases (J40-J47)	5	1.0	*
12	Pneumonitis due to solids and liquids (J69)	2	0.4	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
12	Infections of kidney (N10-N12,N13.6,N15.1)	2	0.4	*
15	Anemias (D50-D64)	1	0.2	*
15	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.2	*
15	Certain conditions originating in the perinatal period (P00-P96)	1	0.2	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
...	All other causes (Residual)	82	17.2	2.3
Hispanic, female, 15-24 years				
...	All causes	1,073	100.0	34.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	455	42.4	14.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	119	11.1	3.8
3	Malignant neoplasms (C00-C97)	113	10.5	3.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	67	6.2	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	34	3.2	1.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	28	2.6	0.9
7	Pregnancy, childbirth and the puerperium (O00-099)	19	1.8	*
8	Cerebrovascular diseases (I60-I69)	12	1.1	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	10	0.9	*
10	Influenza and pneumonia (J10-J18)	8	0.7	*
11	Septicemia (A40-A41)	7	0.7	*
12	Diabetes mellitus (E10-E14)	5	0.5	*
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	4	0.4	*
13	Chronic liver disease and cirrhosis (K70,K73-K74)	4	0.4	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.4	*
...	All other causes (Residual)	184	17.1	5.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 25-34 years				
...	All causes	1,342	100.0	40.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	318	23.7	9.5
2	Malignant neoplasms (C00-C97)	250	18.6	7.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	114	8.5	3.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	79	5.9	2.4
5	Human immunodeficiency virus (HIV) disease (B20-B24)	61	4.5	1.8
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	58	4.3	1.7
7	Cerebrovascular diseases (I60-I69)	32	2.4	1.0
8	Pregnancy, childbirth and the puerperium (O00-099)	28	2.1	0.8
9	Diabetes mellitus (E10-E14)	22	1.6	0.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21	1.6	0.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	19	1.4	*
12	Influenza and pneumonia (J10-J18)	17	1.3	*
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	17	1.3	*
14	Septicemia (A40-A41)	13	1.0	*
15	Chronic lower respiratory diseases (J40-J47)	12	0.9	*
...	All other causes (Residual)	281	20.9	8.4
Hispanic, female, 35-44 years				
...	All causes	2,639	100.0	94.9
1	Malignant neoplasms (C00-C97)	789	29.9	28.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	373	14.1	13.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	227	8.6	8.2
4	Human immunodeficiency virus (HIV) disease (B20-B24)	172	6.5	6.2
5	Cerebrovascular diseases (I60-I69)	102	3.9	3.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	97	3.7	3.5
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	97	3.7	3.5
8	Diabetes mellitus (E10-E14)	81	3.1	2.9
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	67	2.5	2.4
10	Chronic lower respiratory diseases (J40-J47)	39	1.5	1.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	27	1.0	1.0
12	Viral hepatitis (B15-B19)	26	1.0	0.9
13	Influenza and pneumonia (J10-J18)	25	0.9	0.9
14	Pregnancy, childbirth and the puerperium (O00-099)	21	0.8	0.8
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	21	0.8	0.8
...	All other causes (Residual)	475	18.0	17.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 45-54 years				
...	All causes	4,026	100.0	219.8
1	Malignant neoplasms (C00-C97)	1,429	35.5	78.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	553	13.7	30.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	280	7.0	15.3
4	Cerebrovascular diseases (I60-I69)	220	5.5	12.0
5	Diabetes mellitus (E10-E14)	208	5.2	11.4
6	Chronic liver disease and cirrhosis (K70,K73-K74)	178	4.4	9.7
7	Human immunodeficiency virus (HIV) disease (B20-B24)	116	2.9	6.3
8	Viral hepatitis (B15-B19)	73	1.8	4.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	69	1.7	3.8
10	Chronic lower respiratory diseases (J40-J47)	57	1.4	3.1
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	54	1.3	2.9
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	50	1.2	2.7
13	Septicemia (A40-A41)	42	1.0	2.3
14	Influenza and pneumonia (J10-J18)	38	0.9	2.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	28	0.7	1.5
...	All other causes (Residual)	631	15.7	34.5
Hispanic, female, 55-64 years				
...	All causes	5,453	100.0	524.3
1	Malignant neoplasms (C00-C97)	1,870	34.3	179.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,099	20.2	105.7
3	Diabetes mellitus (E10-E14)	478	8.8	46.0
4	Cerebrovascular diseases (I60-I69)	287	5.3	27.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	197	3.6	18.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	159	2.9	15.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	126	2.3	12.1
8	Chronic lower respiratory diseases (J40-J47)	110	2.0	10.6
9	Septicemia (A40-A41)	81	1.5	7.8
10	Influenza and pneumonia (J10-J18)	77	1.4	7.4
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	57	1.0	5.5
12	Viral hepatitis (B15-B19)	56	1.0	5.4
13	Human immunodeficiency virus (HIV) disease (B20-B24)	48	0.9	4.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	25	0.5	2.4
15	Aortic aneurysm and dissection (I71)	24	0.4	2.3
...	All other causes (Residual)	759	13.9	73.0

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 65-74 years				
...	All causes	9,000	100.0	1,368.7
1	Malignant neoplasms (C00-C97)	2,601	28.9	395.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,278	25.3	346.4
3	Diabetes mellitus (E10-E14)	839	9.3	127.6
4	Cerebrovascular diseases (I60-I69)	563	6.3	85.6
5	Chronic lower respiratory diseases (J40-J47)	273	3.0	41.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	256	2.8	38.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	213	2.4	32.4
8	Septicemia (A40-A41)	180	2.0	27.4
9	Influenza and pneumonia (J10-J18)	170	1.9	25.9
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	159	1.8	24.2
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	103	1.1	15.7
12	Alzheimer's disease (G30)	72	0.8	10.9
13	Viral hepatitis (B15-B19)	62	0.7	9.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	51	0.6	7.8
15	Parkinson's disease (G20-G21)	31	0.3	4.7
...	All other causes (Residual)	1,149	12.8	174.7
Hispanic, female, 75-84 years				
...	All causes	12,547	100.0	3,526.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,881	30.9	1,090.8
2	Malignant neoplasms (C00-C97)	2,463	19.6	692.2
3	Cerebrovascular diseases (I60-I69)	1,093	8.7	307.2
4	Diabetes mellitus (E10-E14)	964	7.7	270.9
5	Chronic lower respiratory diseases (J40-J47)	462	3.7	129.8
6	Influenza and pneumonia (J10-J18)	433	3.5	121.7
7	Alzheimer's disease (G30)	339	2.7	95.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	259	2.1	72.8
9	Septicemia (A40-A41)	206	1.6	57.9
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	206	1.6	57.9
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	170	1.4	47.8
12	Chronic liver disease and cirrhosis (K70,K73-K74)	164	1.3	46.1
13	Parkinson's disease (G20-G21)	101	0.8	28.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	71	0.6	20.0
14	Atherosclerosis (I70)	71	0.6	20.0
...	All other causes (Residual)	1,664	13.3	467.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 85 years and over				
...	All causes	12,288	100.0	10,186.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,865	39.6	4,032.8
2	Malignant neoplasms (C00-C97)	1,244	10.1	1,031.2
3	Cerebrovascular diseases (I60-I69)	1,108	9.0	918.5
4	Influenza and pneumonia (J10-J18)	629	5.1	521.4
5	Alzheimer's disease (G30)	587	4.8	486.6
6	Diabetes mellitus (E10-E14)	536	4.4	444.3
7	Chronic lower respiratory diseases (J40-J47)	465	3.8	385.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	222	1.8	184.0
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	189	1.5	156.7
10	Septicemia (A40-A41)	184	1.5	152.5
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	142	1.2	117.7
12	Pneumonitis due to solids and liquids (J69)	137	1.1	113.6
13	Atherosclerosis (I70)	135	1.1	111.9
14	Parkinson's disease (G20-G21)	70	0.6	58.0
15	Chronic liver disease and cirrhosis (K70,K73-K74)	55	0.4	45.6
...	All other causes (Residual)	1,720	14.0	1,425.8
White, non-Hispanic, both sexes, all ages				
...	All causes	1,981,973	100.0	997.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	577,761	29.2	290.8
2	Malignant neoplasms (C00-C97)	458,754	23.1	230.9
3	Cerebrovascular diseases (I60-I69)	133,118	6.7	67.0
4	Chronic lower respiratory diseases (J40-J47)	112,128	5.7	56.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	80,605	4.1	40.6
6	Influenza and pneumonia (J10-J18)	55,419	2.8	27.9
7	Alzheimer's disease (G30)	53,486	2.7	26.9
8	Diabetes mellitus (E10-E14)	52,463	2.6	26.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	30,669	1.5	15.4
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	26,691	1.3	13.4
11	Septicemia (A40-A41)	25,545	1.3	12.9
12	Chronic liver disease and cirrhosis (K70,K73-K74)	20,353	1.0	10.2
13	Parkinson's disease (G20-G21)	15,626	0.8	7.9
14	Pneumonitis due to solids and liquids (J69)	15,203	0.8	7.7
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	14,617	0.7	7.4
...	All other causes (Residual)	309,535	15.6	155.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 1-4 years				
...	All causes	2,502	100.0	27.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	865	34.6	9.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	281	11.2	3.0
3	Malignant neoplasms (C00-C97)	236	9.4	2.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	169	6.8	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	91	3.6	1.0
6	Influenza and pneumonia (J10-J18)	46	1.8	0.5
7	Septicemia (A40-A41)	42	1.7	0.5
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	36	1.4	0.4
9	Cerebrovascular diseases (I60-I69)	29	1.2	0.3
10	Chronic lower respiratory diseases (J40-J47)	23	0.9	0.2
11	Certain conditions originating in the perinatal period (P00-P96)	22	0.9	0.2
12	Acute bronchitis and bronchiolitis (J20-J21)	18	0.7	*
13	Anemias (D50-D64)	10	0.4	*
13	Meningitis (G00,G03)	10	0.4	*
15	Meningococcal infection (A39)	8	0.3	*
...	All other causes (Residual)	616	24.6	6.7
White, non-Hispanic, both sexes, 5-14 years				
...	All causes	4,031	100.0	16.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,581	39.2	6.3
2	Malignant neoplasms (C00-C97)	642	15.9	2.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	253	6.3	1.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	177	4.4	0.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	146	3.6	0.6
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	133	3.3	0.5
7	Septicemia (A40-A41)	55	1.4	0.2
7	Influenza and pneumonia (J10-J18)	55	1.4	0.2
9	Cerebrovascular diseases (I60-I69)	44	1.1	0.2
10	Chronic lower respiratory diseases (J40-J47)	42	1.0	0.2
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	41	1.0	0.2
12	Meningitis (G00,G03)	20	0.5	0.1
13	Anemias (D50-D64)	19	0.5	*
13	Diabetes mellitus (E10-E14)	19	0.5	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	16	0.4	*
...	All other causes (Residual)	788	19.5	3.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 15-24 years				
...	All causes	19,458	100.0	75.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,664	54.8	41.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,951	15.2	11.5
3	Malignant neoplasms (C00-C97)	1,097	5.6	4.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	958	4.9	3.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	547	2.8	2.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	315	1.6	1.2
7	Influenza and pneumonia (J10-J18)	103	0.5	0.4
8	Cerebrovascular diseases (I60-I69)	101	0.5	0.4
9	Diabetes mellitus (E10-E14)	95	0.5	0.4
10	Chronic lower respiratory diseases (J40-J47)	94	0.5	0.4
11	Septicemia (A40-A41)	72	0.4	0.3
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	59	0.3	0.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	33	0.2	0.1
14	Legal intervention (Y35,Y89.0)	28	0.1	0.1
15	Pneumonitis due to solids and liquids (J69)	27	0.1	0.1
...	All other causes (Residual)	2,314	11.9	9.0
White, non-Hispanic, both sexes, 25-34 years				
...	All causes	23,758	100.0	95.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,344	35.1	33.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,809	16.0	15.4
3	Malignant neoplasms (C00-C97)	2,483	10.5	10.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,718	7.2	6.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	977	4.1	3.9
6	Human immunodeficiency virus (HIV) disease (B20-B24)	429	1.8	1.7
7	Diabetes mellitus (E10-E14)	382	1.6	1.5
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	331	1.4	1.3
9	Cerebrovascular diseases (I60-I69)	285	1.2	1.2
10	Influenza and pneumonia (J10-J18)	207	0.9	0.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	204	0.9	0.8
12	Septicemia (A40-A41)	165	0.7	0.7
13	Chronic lower respiratory diseases (J40-J47)	157	0.7	0.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	119	0.5	0.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	105	0.4	0.4
...	All other causes (Residual)	4,043	17.0	16.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 35-44 years				
...	All causes	59,529	100.0	191.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,035	20.2	38.7
2	Malignant neoplasms (C00-C97)	11,234	18.9	36.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,114	15.3	29.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,795	9.7	18.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,148	3.6	6.9
6	Human immunodeficiency virus (HIV) disease (B20-B24)	1,790	3.0	5.8
7	Diabetes mellitus (E10-E14)	1,345	2.3	4.3
8	Cerebrovascular diseases (I60-I69)	1,288	2.2	4.1
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,268	2.1	4.1
10	Chronic lower respiratory diseases (J40-J47)	624	1.0	2.0
11	Influenza and pneumonia (J10-J18)	617	1.0	2.0
12	Septicemia (A40-A41)	510	0.9	1.6
13	Viral hepatitis (B15-B19)	496	0.8	1.6
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	423	0.7	1.4
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	353	0.6	1.1
...	All other causes (Residual)	10,489	17.6	33.8
White, non-Hispanic, both sexes, 45-54 years				
...	All causes	119,812	100.0	400.1
1	Malignant neoplasms (C00-C97)	36,609	30.6	122.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	26,206	21.9	87.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,859	9.1	36.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,603	4.7	18.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	5,023	4.2	16.8
6	Diabetes mellitus (E10-E14)	3,385	2.8	11.3
7	Cerebrovascular diseases (I60-I69)	3,320	2.8	11.1
8	Chronic lower respiratory diseases (J40-J47)	2,686	2.2	9.0
9	Viral hepatitis (B15-B19)	1,590	1.3	5.3
10	Human immunodeficiency virus (HIV) disease (B20-B24)	1,366	1.1	4.6
11	Influenza and pneumonia (J10-J18)	1,329	1.1	4.4
12	Septicemia (A40-A41)	1,274	1.1	4.3
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	922	0.8	3.1
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	898	0.7	3.0
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	562	0.5	1.9
...	All other causes (Residual)	18,180	15.2	60.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 55-64 years				
...	All causes	190,843	100.0	912.8
1	Malignant neoplasms (C00-C97)	73,809	38.7	353.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	47,494	24.9	227.2
3	Chronic lower respiratory diseases (J40-J47)	9,757	5.1	46.7
4	Cerebrovascular diseases (I60-I69)	6,432	3.4	30.8
5	Diabetes mellitus (E10-E14)	6,403	3.4	30.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,393	3.3	30.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	4,483	2.3	21.4
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,254	1.7	15.6
9	Septicemia (A40-A41)	2,241	1.2	10.7
10	Influenza and pneumonia (J10-J18)	2,152	1.1	10.3
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,978	1.0	9.5
12	Aortic aneurysm and dissection (I71)	1,156	0.6	5.5
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	962	0.5	4.6
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	805	0.4	3.9
15	Viral hepatitis (B15-B19)	635	0.3	3.0
...	All other causes (Residual)	22,889	12.0	109.5
White, non-Hispanic, both sexes, 65-74 years				
...	All causes	338,961	100.0	2,296.1
1	Malignant neoplasms (C00-C97)	119,464	35.2	809.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	88,528	26.1	599.7
3	Chronic lower respiratory diseases (J40-J47)	26,808	7.9	181.6
4	Cerebrovascular diseases (I60-I69)	16,355	4.8	110.8
5	Diabetes mellitus (E10-E14)	11,337	3.3	76.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,481	1.9	43.9
7	Influenza and pneumonia (J10-J18)	5,419	1.6	36.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,870	1.4	33.0
9	Septicemia (A40-A41)	4,583	1.4	31.0
10	Chronic liver disease and cirrhosis (K70,K73-K74)	4,205	1.2	28.5
11	Alzheimer's disease (G30)	3,149	0.9	21.3
12	Aortic aneurysm and dissection (I71)	3,057	0.9	20.7
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,228	0.7	15.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	1,985	0.6	13.4
15	Parkinson's disease (G20-G21)	1,959	0.6	13.3
...	All other causes (Residual)	38,533	11.4	261.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 75-84 years				
...	All causes	606,431	100.0	5,596.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	181,975	30.0	1,679.2
2	Malignant neoplasms (C00-C97)	143,957	23.7	1,328.4
3	Cerebrovascular diseases (I60-I69)	46,327	7.6	427.5
4	Chronic lower respiratory diseases (J40-J47)	45,117	7.4	416.3
5	Alzheimer's disease (G30)	18,227	3.0	168.2
6	Diabetes mellitus (E10-E14)	17,692	2.9	163.3
7	Influenza and pneumonia (J10-J18)	17,001	2.8	156.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,316	1.9	104.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11,110	1.8	102.5
10	Septicemia (A40-A41)	8,806	1.5	81.3
11	Parkinson's disease (G20-G21)	7,545	1.2	69.6
12	Pneumonitis due to solids and liquids (J69)	5,131	0.8	47.3
13	Aortic aneurysm and dissection (I71)	4,988	0.8	46.0
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	4,570	0.8	42.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	4,063	0.7	37.5
...	All other causes (Residual)	78,606	13.0	725.4
White, non-Hispanic, both sexes, 85 years and over				
...	All causes	603,063	100.0	15,149.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	221,670	36.8	5,568.4
2	Malignant neoplasms (C00-C97)	69,184	11.5	1,737.9
3	Cerebrovascular diseases (I60-I69)	58,884	9.8	1,479.2
4	Alzheimer's disease (G30)	31,605	5.2	793.9
5	Influenza and pneumonia (J10-J18)	28,373	4.7	712.7
6	Chronic lower respiratory diseases (J40-J47)	26,796	4.4	673.1
7	Diabetes mellitus (E10-E14)	11,801	2.0	296.4
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,579	1.9	290.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11,192	1.9	281.1
10	Septicemia (A40-A41)	7,666	1.3	192.6
11	Pneumonitis due to solids and liquids (J69)	7,616	1.3	191.3
12	Atherosclerosis (I70)	7,046	1.2	177.0
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	6,790	1.1	170.6
14	Parkinson's disease (G20-G21)	5,803	1.0	145.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	3,258	0.5	81.8
...	All other causes (Residual)	93,800	15.6	2,356.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, all ages				
...	All causes	957,645	100.0	983.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	281,442	29.4	289.2
2	Malignant neoplasms (C00-C97)	237,323	24.8	243.8
3	Chronic lower respiratory diseases (J40-J47)	53,662	5.6	55.1
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	50,652	5.3	52.0
5	Cerebrovascular diseases (I60-I69)	49,908	5.2	51.3
6	Diabetes mellitus (E10-E14)	25,273	2.6	26.0
7	Influenza and pneumonia (J10-J18)	23,934	2.5	24.6
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	21,323	2.2	21.9
9	Alzheimer's disease (G30)	15,399	1.6	15.8
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14,883	1.6	15.3
11	Chronic liver disease and cirrhosis (K70,K73-K74)	12,800	1.3	13.2
12	Septicemia (A40-A41)	11,148	1.2	11.5
13	Parkinson's disease (G20-G21)	8,838	0.9	9.1
14	Aortic aneurysm and dissection (I71)	7,834	0.8	8.0
15	Pneumonitis due to solids and liquids (J69)	7,654	0.8	7.9
...	All other causes (Residual)	135,572	14.2	139.3
White, non-Hispanic, male, 1-4 years				
...	All causes	1,431	100.0	30.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	527	36.8	11.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	151	10.6	3.2
3	Malignant neoplasms (C00-C97)	134	9.4	2.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	89	6.2	1.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	54	3.8	1.1
6	Septicemia (A40-A41)	25	1.7	0.5
7	Influenza and pneumonia (J10-J18)	24	1.7	0.5
8	Cerebrovascular diseases (I60-I69)	18	1.3	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	17	1.2	*
9	Chronic lower respiratory diseases (J40-J47)	17	1.2	*
11	Certain conditions originating in the perinatal period (P00-P96)	10	0.7	*
12	Meningococcal infection (A39)	6	0.4	*
12	Meningitis (G00,G03)	6	0.4	*
12	Acute bronchitis and bronchiolitis (J20-J21)	6	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.3	*
...	All other causes (Residual)	342	23.9	7.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 5-14 years				
...	All causes	2,365	100.0	18.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	997	42.2	7.7
2	Malignant neoplasms (C00-C97)	355	15.0	2.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	138	5.8	1.1
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	125	5.3	1.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	85	3.6	0.7
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	61	2.6	0.5
7	Septicemia (A40-A41)	34	1.4	0.3
8	Cerebrovascular diseases (I60-I69)	23	1.0	0.2
8	Chronic lower respiratory diseases (J40-J47)	23	1.0	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	22	0.9	0.2
11	Influenza and pneumonia (J10-J18)	20	0.8	0.2
12	Diabetes mellitus (E10-E14)	16	0.7	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	12	0.5	*
14	Meningitis (G00,G03)	11	0.5	*
15	Meningococcal infection (A39)	8	0.3	*
15	Anemias (D50-D64)	8	0.3	*
...	All other causes (Residual)	427	18.1	3.3
White, non-Hispanic, male, 15-24 years				
...	All causes	13,957	100.0	106.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,746	55.5	59.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,527	18.1	19.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	685	4.9	5.2
4	Malignant neoplasms (C00-C97)	639	4.6	4.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	370	2.7	2.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	178	1.3	1.4
7	Cerebrovascular diseases (I60-I69)	62	0.4	0.5
7	Influenza and pneumonia (J10-J18)	62	0.4	0.5
9	Diabetes mellitus (E10-E14)	51	0.4	0.4
10	Chronic lower respiratory diseases (J40-J47)	48	0.3	0.4
11	Septicemia (A40-A41)	37	0.3	0.3
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	32	0.2	0.2
13	Legal intervention (Y35,Y89.0)	25	0.2	0.2
14	Aortic aneurysm and dissection (I71)	19	0.1	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	0.1	*
...	All other causes (Residual)	1,459	10.5	11.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 25-34 years				
...	All causes	16,339	100.0	130.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,273	38.4	50.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,065	18.8	24.6
3	Malignant neoplasms (C00-C97)	1,218	7.5	9.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,177	7.2	9.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	660	4.0	5.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	335	2.1	2.7
7	Diabetes mellitus (E10-E14)	231	1.4	1.9
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	195	1.2	1.6
9	Cerebrovascular diseases (I60-I69)	136	0.8	1.1
10	Chronic liver disease and cirrhosis (K70,K73-K74)	130	0.8	1.0
11	Influenza and pneumonia (J10-J18)	111	0.7	0.9
12	Septicemia (A40-A41)	86	0.5	0.7
13	Chronic lower respiratory diseases (J40-J47)	81	0.5	0.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	64	0.4	0.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	62	0.4	0.5
...	All other causes (Residual)	2,515	15.4	20.2
White, non-Hispanic, male, 35-44 years				
...	All causes	38,049	100.0	244.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,550	22.5	55.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,657	17.5	42.9
3	Malignant neoplasms (C00-C97)	5,041	13.2	32.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,462	11.7	28.7
5	Human immunodeficiency virus (HIV) disease (B20-B24)	1,517	4.0	9.8
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1,371	3.6	8.8
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	874	2.3	5.6
8	Diabetes mellitus (E10-E14)	849	2.2	5.5
9	Cerebrovascular diseases (I60-I69)	617	1.6	4.0
10	Viral hepatitis (B15-B19)	351	0.9	2.3
11	Influenza and pneumonia (J10-J18)	340	0.9	2.2
12	Chronic lower respiratory diseases (J40-J47)	304	0.8	2.0
13	Septicemia (A40-A41)	249	0.7	1.6
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	223	0.6	1.4
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	191	0.5	1.2
...	All other causes (Residual)	6,453	17.0	41.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	White, non-Hispanic, male, 45-54 years			
...	All causes	75,724	100.0	509.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	19,774	26.1	133.1
2	Malignant neoplasms (C00-C97)	18,699	24.7	125.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,745	10.2	52.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,252	5.6	28.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	3,670	4.8	24.7
6	Diabetes mellitus (E10-E14)	2,054	2.7	13.8
7	Cerebrovascular diseases (I60-I69)	1,797	2.4	12.1
8	Chronic lower respiratory diseases (J40-J47)	1,383	1.8	9.3
9	Viral hepatitis (B15-B19)	1,212	1.6	8.2
10	Human immunodeficiency virus (HIV) disease (B20-B24)	1,183	1.6	8.0
11	Influenza and pneumonia (J10-J18)	768	1.0	5.2
12	Septicemia (A40-A41)	680	0.9	4.6
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	586	0.8	3.9
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	533	0.7	3.6
15	Aortic aneurysm and dissection (I71)	355	0.5	2.4
...	All other causes (Residual)	11,033	14.6	74.3
	White, non-Hispanic, male, 55-64 years			
...	All causes	114,535	100.0	1,126.5
1	Malignant neoplasms (C00-C97)	40,210	35.1	395.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	33,313	29.1	327.6
3	Chronic lower respiratory diseases (J40-J47)	4,909	4.3	48.3
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,374	3.8	43.0
5	Diabetes mellitus (E10-E14)	3,815	3.3	37.5
6	Cerebrovascular diseases (I60-I69)	3,511	3.1	34.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	3,097	2.7	30.5
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,546	2.2	25.0
9	Influenza and pneumonia (J10-J18)	1,228	1.1	12.1
10	Septicemia (A40-A41)	1,204	1.1	11.8
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,045	0.9	10.3
12	Aortic aneurysm and dissection (I71)	875	0.8	8.6
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	547	0.5	5.4
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	477	0.4	4.7
15	Viral hepatitis (B15-B19)	415	0.4	4.1
...	All other causes (Residual)	12,969	11.3	127.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 65-74 years				
...	All causes	191,844	100.0	2,824.1
1	Malignant neoplasms (C00-C97)	66,250	34.5	975.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	55,265	28.8	813.5
3	Chronic lower respiratory diseases (J40-J47)	13,615	7.1	200.4
4	Cerebrovascular diseases (I60-I69)	8,368	4.4	123.2
5	Diabetes mellitus (E10-E14)	6,041	3.1	88.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,991	2.1	58.8
7	Influenza and pneumonia (J10-J18)	3,050	1.6	44.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,694	1.4	39.7
9	Chronic liver disease and cirrhosis (K70,K73-K74)	2,489	1.3	36.6
10	Septicemia (A40-A41)	2,310	1.2	34.0
11	Aortic aneurysm and dissection (I71)	2,105	1.1	31.0
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,865	1.0	27.5
13	Alzheimer's disease (G30)	1,353	0.7	19.9
14	Parkinson's disease (G20-G21)	1,312	0.7	19.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	1,135	0.6	16.7
...	All other causes (Residual)	20,001	10.4	294.4
White, non-Hispanic, male, 75-84 years				
...	All causes	295,763	100.0	6,801.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	92,616	31.3	2,129.9
2	Malignant neoplasms (C00-C97)	74,639	25.2	1,716.5
3	Chronic lower respiratory diseases (J40-J47)	22,034	7.4	506.7
4	Cerebrovascular diseases (I60-I69)	19,181	6.5	441.1
5	Influenza and pneumonia (J10-J18)	8,402	2.8	193.2
6	Diabetes mellitus (E10-E14)	8,151	2.8	187.4
7	Alzheimer's disease (G30)	6,544	2.2	150.5
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,805	2.0	133.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,687	1.9	130.8
10	Parkinson's disease (G20-G21)	4,473	1.5	102.9
11	Septicemia (A40-A41)	4,006	1.4	92.1
12	Aortic aneurysm and dissection (I71)	2,917	1.0	67.1
13	Pneumonitis due to solids and liquids (J69)	2,822	1.0	64.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	2,069	0.7	47.6
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,812	0.6	41.7
...	All other causes (Residual)	34,605	11.7	795.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 85 years and over				
...	All causes	199,834	100.0	16,641.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	71,976	36.0	5,994.1
2	Malignant neoplasms (C00-C97)	30,112	15.1	2,507.7
3	Cerebrovascular diseases (I60-I69)	16,162	8.1	1,345.9
4	Chronic lower respiratory diseases (J40-J47)	11,232	5.6	935.4
5	Influenza and pneumonia (J10-J18)	9,854	4.9	820.6
6	Alzheimer's disease (G30)	7,270	3.6	605.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,594	2.3	382.6
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,342	2.2	361.6
9	Diabetes mellitus (E10-E14)	4,064	2.0	338.4
10	Pneumonitis due to solids and liquids (J69)	3,343	1.7	278.4
11	Parkinson's disease (G20-G21)	2,856	1.4	237.8
12	Septicemia (A40-A41)	2,448	1.2	203.9
13	Atherosclerosis (I70)	1,990	1.0	165.7
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,746	0.9	145.4
15	Aortic aneurysm and dissection (I71)	1,377	0.7	114.7
...	All other causes (Residual)	26,468	13.2	2,204.2
White, non-Hispanic, female, all ages				
...	All causes	1,024,328	100.0	1,010.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	296,319	28.9	292.3
2	Malignant neoplasms (C00-C97)	221,431	21.6	218.5
3	Cerebrovascular diseases (I60-I69)	83,210	8.1	82.1
4	Chronic lower respiratory diseases (J40-J47)	58,466	5.7	57.7
5	Alzheimer's disease (G30)	38,087	3.7	37.6
6	Influenza and pneumonia (J10-J18)	31,485	3.1	31.1
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	29,953	2.9	29.6
8	Diabetes mellitus (E10-E14)	27,190	2.7	26.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15,786	1.5	15.6
10	Septicemia (A40-A41)	14,397	1.4	14.2
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	9,292	0.9	9.2
12	Atherosclerosis (I70)	7,683	0.8	7.6
13	Chronic liver disease and cirrhosis (K70,K73-K74)	7,553	0.7	7.5
14	Pneumonitis due to solids and liquids (J69)	7,549	0.7	7.4
15	Parkinson's disease (G20-G21)	6,788	0.7	6.7
...	All other causes (Residual)	169,139	16.5	166.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 1-4 years				
...	All causes	1,071	100.0	23.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	338	31.6	7.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	130	12.1	2.9
3	Malignant neoplasms (C00-C97)	102	9.5	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	80	7.5	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	37	3.5	0.8
6	Influenza and pneumonia (J10-J18)	22	2.1	0.5
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	19	1.8	*
8	Septicemia (A40-A41)	17	1.6	*
9	Acute bronchitis and bronchiolitis (J20-J21)	12	1.1	*
9	Certain conditions originating in the perinatal period (P00-P96)	12	1.1	*
11	Cerebrovascular diseases (I60-I69)	11	1.0	*
12	Anemias (D50-D64)	6	0.6	*
12	Chronic lower respiratory diseases (J40-J47)	6	0.6	*
14	Meningitis (G00,G03)	4	0.4	*
15	Meningococcal infection (A39)	2	0.2	*
15	Diabetes mellitus (E10-E14)	2	0.2	*
15	Pneumonitis due to solids and liquids (J69)	2	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.2	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.2	*
...	All other causes (Residual)	265	24.7	5.9
White, non-Hispanic, female, 5-14 years				
...	All causes	1,666	100.0	13.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	584	35.1	4.8
2	Malignant neoplasms (C00-C97)	287	17.2	2.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	128	7.7	1.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	72	4.3	0.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	61	3.7	0.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	39	2.3	0.3
7	Influenza and pneumonia (J10-J18)	35	2.1	0.3
8	Septicemia (A40-A41)	21	1.3	0.2
8	Cerebrovascular diseases (I60-I69)	21	1.3	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	19	1.1	*
10	Chronic lower respiratory diseases (J40-J47)	19	1.1	*
12	Anemias (D50-D64)	11	0.7	*
13	Meningitis (G00,G03)	9	0.5	*
14	Certain conditions originating in the perinatal period (P00-P96)	8	0.5	*
15	Pneumonitis due to solids and liquids (J69)	4	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.2	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.2	*
...	All other causes (Residual)	340	20.4	2.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 15-24 years				
...	All causes	5,501	100.0	43.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,918	53.0	23.2
2	Malignant neoplasms (C00-C97)	458	8.3	3.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	424	7.7	3.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	273	5.0	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	177	3.2	1.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	137	2.5	1.1
7	Chronic lower respiratory diseases (J40-J47)	46	0.8	0.4
8	Diabetes mellitus (E10-E14)	44	0.8	0.4
9	Influenza and pneumonia (J10-J18)	41	0.7	0.3
10	Cerebrovascular diseases (I60-I69)	39	0.7	0.3
11	Septicemia (A40-A41)	35	0.6	0.3
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	27	0.5	0.2
13	Pregnancy, childbirth and the puerperium (O00-099)	21	0.4	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	0.3	*
15	Pneumonitis due to solids and liquids (J69)	12	0.2	*
...	All other causes (Residual)	833	15.1	6.6
White, non-Hispanic, female, 25-34 years				
...	All causes	7,419	100.0	60.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,071	27.9	16.8
2	Malignant neoplasms (C00-C97)	1,265	17.1	10.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	744	10.0	6.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	541	7.3	4.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	317	4.3	2.6
6	Diabetes mellitus (E10-E14)	151	2.0	1.2
7	Cerebrovascular diseases (I60-I69)	149	2.0	1.2
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	136	1.8	1.1
9	Influenza and pneumonia (J10-J18)	96	1.3	0.8
10	Human immunodeficiency virus (HIV) disease (B20-B24)	94	1.3	0.8
11	Septicemia (A40-A41)	79	1.1	0.6
12	Chronic lower respiratory diseases (J40-J47)	76	1.0	0.6
13	Chronic liver disease and cirrhosis (K70,K73-K74)	74	1.0	0.6
14	Pregnancy, childbirth and the puerperium (O00-099)	69	0.9	0.6
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	55	0.7	0.4
...	All other causes (Residual)	1,502	20.2	12.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	White, non-Hispanic, female, 35-44 years			
...	All causes	21,480	100.0	138.3
1	Malignant neoplasms (C00-C97)	6,193	28.8	39.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,485	16.2	22.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,457	11.4	15.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,333	6.2	8.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	777	3.6	5.0
6	Cerebrovascular diseases (I60-I69)	671	3.1	4.3
7	Diabetes mellitus (E10-E14)	496	2.3	3.2
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	394	1.8	2.5
9	Chronic lower respiratory diseases (J40-J47)	320	1.5	2.1
10	Influenza and pneumonia (J10-J18)	277	1.3	1.8
11	Human immunodeficiency virus (HIV) disease (B20-B24)	273	1.3	1.8
12	Septicemia (A40-A41)	261	1.2	1.7
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	200	0.9	1.3
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	162	0.8	1.0
15	Viral hepatitis (B15-B19)	145	0.7	0.9
...	All other causes (Residual)	4,036	18.8	26.0
	White, non-Hispanic, female, 45-54 years			
...	All causes	44,088	100.0	292.1
1	Malignant neoplasms (C00-C97)	17,910	40.6	118.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,432	14.6	42.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,114	7.1	20.6
4	Cerebrovascular diseases (I60-I69)	1,523	3.5	10.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,353	3.1	9.0
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,351	3.1	9.0
7	Diabetes mellitus (E10-E14)	1,331	3.0	8.8
8	Chronic lower respiratory diseases (J40-J47)	1,303	3.0	8.6
9	Septicemia (A40-A41)	594	1.3	3.9
10	Influenza and pneumonia (J10-J18)	561	1.3	3.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	389	0.9	2.6
12	Viral hepatitis (B15-B19)	378	0.9	2.5
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	312	0.7	2.1
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	267	0.6	1.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	247	0.6	1.6
...	All other causes (Residual)	7,023	15.9	46.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	White, non-Hispanic, female, 55-64 years			
...	All causes	76,308	100.0	710.5
1	Malignant neoplasms (C00-C97)	33,599	44.0	312.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,181	18.6	132.0
3	Chronic lower respiratory diseases (J40-J47)	4,848	6.4	45.1
4	Cerebrovascular diseases (I60-I69)	2,921	3.8	27.2
5	Diabetes mellitus (E10-E14)	2,588	3.4	24.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,019	2.6	18.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,386	1.8	12.9
8	Septicemia (A40-A41)	1,037	1.4	9.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	933	1.2	8.7
10	Influenza and pneumonia (J10-J18)	924	1.2	8.6
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	708	0.9	6.6
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	415	0.5	3.9
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	328	0.4	3.1
14	Aortic aneurysm and dissection (I71)	281	0.4	2.6
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	264	0.3	2.5
...	All other causes (Residual)	9,876	12.9	92.0
	White, non-Hispanic, female, 65-74 years			
...	All causes	147,117	100.0	1,846.0
1	Malignant neoplasms (C00-C97)	53,214	36.2	667.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	33,263	22.6	417.4
3	Chronic lower respiratory diseases (J40-J47)	13,193	9.0	165.5
4	Cerebrovascular diseases (I60-I69)	7,987	5.4	100.2
5	Diabetes mellitus (E10-E14)	5,296	3.6	66.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,490	1.7	31.2
7	Influenza and pneumonia (J10-J18)	2,369	1.6	29.7
8	Septicemia (A40-A41)	2,273	1.5	28.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,176	1.5	27.3
10	Alzheimer's disease (G30)	1,796	1.2	22.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1,716	1.2	21.5
12	Aortic aneurysm and dissection (I71)	952	0.6	11.9
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	918	0.6	11.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	850	0.6	10.7
15	Parkinson's disease (G20-G21)	647	0.4	8.1
...	All other causes (Residual)	17,977	12.2	225.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	White, non-Hispanic, female, 75-84 years			
...	All causes	310,668	100.0	4,787.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	89,359	28.8	1,377.2
2	Malignant neoplasms (C00-C97)	69,318	22.3	1,068.3
3	Cerebrovascular diseases (I60-I69)	27,146	8.7	418.4
4	Chronic lower respiratory diseases (J40-J47)	23,083	7.4	355.7
5	Alzheimer's disease (G30)	11,683	3.8	180.1
6	Diabetes mellitus (E10-E14)	9,541	3.1	147.0
7	Influenza and pneumonia (J10-J18)	8,599	2.8	132.5
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,511	1.8	84.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,423	1.7	83.6
10	Septicemia (A40-A41)	4,800	1.5	74.0
11	Parkinson's disease (G20-G21)	3,072	1.0	47.3
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,786	0.9	42.9
13	Pneumonitis due to solids and liquids (J69)	2,309	0.7	35.6
14	Aortic aneurysm and dissection (I71)	2,071	0.7	31.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	1,994	0.6	30.7
...	All other causes (Residual)	43,973	14.2	677.7
	White, non-Hispanic, female, 85 years and over			
...	All causes	403,229	100.0	14,504.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	149,694	37.1	5,384.5
2	Cerebrovascular diseases (I60-I69)	42,722	10.6	1,536.7
3	Malignant neoplasms (C00-C97)	39,072	9.7	1,405.4
4	Alzheimer's disease (G30)	24,335	6.0	875.3
5	Influenza and pneumonia (J10-J18)	18,519	4.6	666.1
6	Chronic lower respiratory diseases (J40-J47)	15,564	3.9	559.8
7	Diabetes mellitus (E10-E14)	7,737	1.9	278.3
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,237	1.8	260.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,598	1.6	237.3
10	Septicemia (A40-A41)	5,218	1.3	187.7
11	Atherosclerosis (I70)	5,056	1.3	181.9
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	5,044	1.3	181.4
13	Pneumonitis due to solids and liquids (J69)	4,273	1.1	153.7
14	Parkinson's disease (G20-G21)	2,947	0.7	106.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	1,985	0.5	71.4
...	All other causes (Residual)	67,228	16.7	2,418.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, all ages				
...	All causes	286,573	100.0	792.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	76,694	26.8	212.2
2	Malignant neoplasms (C00-C97)	61,996	21.6	171.5
3	Cerebrovascular diseases (I60-I69)	18,691	6.5	51.7
4	Diabetes mellitus (E10-E14)	12,583	4.4	34.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,285	4.3	34.0
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8,147	2.8	22.5
7	Chronic lower respiratory diseases (J40-J47)	7,730	2.7	21.4
8	Human immunodeficiency virus (HIV) disease (B20-B24)	7,714	2.7	21.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,410	2.6	20.5
10	Septicemia (A40-A41)	6,074	2.1	16.8
11	Influenza and pneumonia (J10-J18)	5,803	2.0	16.1
12	Certain conditions originating in the perinatal period (P00-P96)	4,777	1.7	13.2
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	4,165	1.5	11.5
14	Alzheimer's disease (G30)	3,303	1.2	9.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,545	0.9	7.0
...	All other causes (Residual)	46,656	16.3	129.1
Black, non-Hispanic, both sexes, 1-4 years				
...	All causes	1,163	100.0	48.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	350	30.1	14.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	156	13.4	6.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	113	9.7	4.7
4	Malignant neoplasms (C00-C97)	71	6.1	3.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	40	3.4	1.7
5	Influenza and pneumonia (J10-J18)	40	3.4	1.7
7	Chronic lower respiratory diseases (J40-J47)	29	2.5	1.2
8	Certain conditions originating in the perinatal period (P00-P96)	26	2.2	1.1
9	Septicemia (A40-A41)	17	1.5	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	14	1.2	*
11	Cerebrovascular diseases (I60-I69)	10	0.9	*
12	Anemias (D50-D64)	7	0.6	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	5	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.4	*
15	Meningitis (G00,G03)	4	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.3	*
...	All other causes (Residual)	272	23.4	11.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Black, non-Hispanic, both sexes, 5-14 years			
...	All causes	1,634	100.0	25.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	607	37.1	9.4
2	Malignant neoplasms (C00-C97)	168	10.3	2.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	132	8.1	2.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	81	5.0	1.3
5	Chronic lower respiratory diseases (J40-J47)	74	4.5	1.1
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	65	4.0	1.0
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	35	2.1	0.5
8	Anemias (D50-D64)	26	1.6	0.4
9	Cerebrovascular diseases (I60-I69)	23	1.4	0.4
10	Septicemia (A40-A41)	20	1.2	0.3
11	Human immunodeficiency virus (HIV) disease (B20-B24)	19	1.2	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	18	1.1	*
12	Influenza and pneumonia (J10-J18)	18	1.1	*
14	Diabetes mellitus (E10-E14)	14	0.9	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	14	0.9	*
...	All other causes (Residual)	320	19.6	5.0
	Black, non-Hispanic, both sexes, 15-24 years			
...	All causes	6,904	100.0	116.9
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,854	41.3	48.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,680	24.3	28.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	392	5.7	6.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	299	4.3	5.1
5	Malignant neoplasms (C00-C97)	242	3.5	4.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	124	1.8	2.1
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	98	1.4	1.7
8	Anemias (D50-D64)	78	1.1	1.3
8	Chronic lower respiratory diseases (J40-J47)	78	1.1	1.3
10	Diabetes mellitus (E10-E14)	57	0.8	1.0
11	Pregnancy, childbirth and the puerperium (O00-099)	45	0.7	0.8
12	Cerebrovascular diseases (I60-I69)	38	0.6	0.6
12	Influenza and pneumonia (J10-J18)	38	0.6	0.6
14	Legal intervention (Y35,Y89.0)	37	0.5	0.6
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	36	0.5	0.6
...	All other causes (Residual)	808	11.7	13.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Black, non-Hispanic, both sexes, 25-34 years			
...	All causes	10,009	100.0	193.9
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,425	24.2	47.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,676	16.7	32.5
3	Human immunodeficiency virus (HIV) disease (B20-B24)	1,110	11.1	21.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	993	9.9	19.2
5	Malignant neoplasms (C00-C97)	663	6.6	12.8
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	497	5.0	9.6
7	Diabetes mellitus (E10-E14)	194	1.9	3.8
8	Cerebrovascular diseases (I60-I69)	166	1.7	3.2
9	Anemias (D50-D64)	127	1.3	2.5
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	108	1.1	2.1
11	Septicemia (A40-A41)	102	1.0	2.0
11	Chronic lower respiratory diseases (J40-J47)	102	1.0	2.0
13	Influenza and pneumonia (J10-J18)	80	0.8	1.6
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	75	0.7	1.5
15	Pregnancy, childbirth and the puerperium (O00-099)	64	0.6	1.2
...	All other causes (Residual)	1,627	16.3	31.5
	Black, non-Hispanic, both sexes, 35-44 years			
...	All causes	19,955	100.0	357.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,312	16.6	59.4
2	Human immunodeficiency virus (HIV) disease (B20-B24)	3,011	15.1	54.0
3	Malignant neoplasms (C00-C97)	2,870	14.4	51.5
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,227	11.2	39.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,357	6.8	24.3
6	Cerebrovascular diseases (I60-I69)	741	3.7	13.3
7	Diabetes mellitus (E10-E14)	554	2.8	9.9
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	432	2.2	7.7
9	Chronic liver disease and cirrhosis (K70,K73-K74)	313	1.6	5.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	306	1.5	5.5
11	Chronic lower respiratory diseases (J40-J47)	292	1.5	5.2
12	Septicemia (A40-A41)	254	1.3	4.6
13	Influenza and pneumonia (J10-J18)	250	1.3	4.5
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	189	0.9	3.4
15	Anemias (D50-D64)	142	0.7	2.5
...	All other causes (Residual)	3,705	18.6	66.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 45-54 years				
...	All causes	35,211	100.0	782.0
1	Malignant neoplasms (C00-C97)	8,376	23.8	186.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,318	23.6	184.7
3	Human immunodeficiency virus (HIV) disease (B20-B24)	2,469	7.0	54.8
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,145	6.1	47.6
5	Cerebrovascular diseases (I60-I69)	1,874	5.3	41.6
6	Diabetes mellitus (E10-E14)	1,393	4.0	30.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	956	2.7	21.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	745	2.1	16.5
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	710	2.0	15.8
10	Septicemia (A40-A41)	593	1.7	13.2
11	Chronic lower respiratory diseases (J40-J47)	589	1.7	13.1
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	413	1.2	9.2
13	Influenza and pneumonia (J10-J18)	399	1.1	8.9
14	Viral hepatitis (B15-B19)	368	1.0	8.2
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	265	0.8	5.9
...	All other causes (Residual)	5,598	15.9	124.3
Black, non-Hispanic, both sexes, 55-64 years				
...	All causes	40,878	100.0	1,592.9
1	Malignant neoplasms (C00-C97)	12,693	31.1	494.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,554	28.3	450.2
3	Cerebrovascular diseases (I60-I69)	2,303	5.6	89.7
4	Diabetes mellitus (E10-E14)	2,204	5.4	85.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,075	2.6	41.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,072	2.6	41.8
7	Chronic lower respiratory diseases (J40-J47)	1,067	2.6	41.6
8	Septicemia (A40-A41)	868	2.1	33.8
9	Human immunodeficiency virus (HIV) disease (B20-B24)	675	1.7	26.3
10	Chronic liver disease and cirrhosis (K70,K73-K74)	649	1.6	25.3
11	Influenza and pneumonia (J10-J18)	554	1.4	21.6
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	550	1.3	21.4
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	216	0.5	8.4
14	Aortic aneurysm and dissection (I71)	192	0.5	7.5
15	Viral hepatitis (B15-B19)	184	0.5	7.2
...	All other causes (Residual)	5,022	12.3	195.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 65-74 years				
...	All causes	52,748	100.0	3,204.8
1	Malignant neoplasms (C00-C97)	16,084	30.5	977.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,411	29.2	936.3
3	Cerebrovascular diseases (I60-I69)	3,600	6.8	218.7
4	Diabetes mellitus (E10-E14)	3,152	6.0	191.5
5	Chronic lower respiratory diseases (J40-J47)	1,892	3.6	115.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,670	3.2	101.5
7	Septicemia (A40-A41)	1,251	2.4	76.0
8	Influenza and pneumonia (J10-J18)	837	1.6	50.9
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	818	1.6	49.7
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	795	1.5	48.3
11	Chronic liver disease and cirrhosis (K70,K73-K74)	358	0.7	21.8
12	Alzheimer's disease (G30)	277	0.5	16.8
13	Pneumonitis due to solids and liquids (J69)	250	0.5	15.2
14	Aortic aneurysm and dissection (I71)	244	0.5	14.8
15	Human immunodeficiency virus (HIV) disease (B20-B24)	240	0.5	14.6
...	All other causes (Residual)	5,869	11.1	356.6
Black, non-Hispanic, both sexes, 75-84 years				
...	All causes	62,183	100.0	6,584.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	19,520	31.4	2,067.0
2	Malignant neoplasms (C00-C97)	14,518	23.3	1,537.4
3	Cerebrovascular diseases (I60-I69)	5,277	8.5	558.8
4	Diabetes mellitus (E10-E14)	3,213	5.2	340.2
5	Chronic lower respiratory diseases (J40-J47)	2,366	3.8	250.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,941	3.1	205.5
7	Septicemia (A40-A41)	1,633	2.6	172.9
8	Influenza and pneumonia (J10-J18)	1,584	2.5	167.7
9	Alzheimer's disease (G30)	1,173	1.9	124.2
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,133	1.8	120.0
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	799	1.3	84.6
12	Pneumonitis due to solids and liquids (J69)	499	0.8	52.8
13	Atherosclerosis (I70)	349	0.6	37.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	300	0.5	31.8
15	Aortic aneurysm and dissection (I71)	285	0.5	30.2
...	All other causes (Residual)	7,593	12.2	804.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 85 years and over				
...	All causes	47,565	100.0	14,386.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	17,030	35.8	5,151.0
2	Malignant neoplasms (C00-C97)	6,296	13.2	1,904.3
3	Cerebrovascular diseases (I60-I69)	4,622	9.7	1,398.0
4	Influenza and pneumonia (J10-J18)	1,917	4.0	579.8
5	Alzheimer's disease (G30)	1,814	3.8	548.7
6	Diabetes mellitus (E10-E14)	1,801	3.8	544.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,468	3.1	444.0
8	Septicemia (A40-A41)	1,226	2.6	370.8
8	Chronic lower respiratory diseases (J40-J47)	1,226	2.6	370.8
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,033	2.2	312.4
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	580	1.2	175.4
12	Pneumonitis due to solids and liquids (J69)	553	1.2	167.3
13	Atherosclerosis (I70)	390	0.8	118.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	213	0.4	64.4
15	Nutritional deficiencies (E40-E64)	199	0.4	60.2
...	All other causes (Residual)	7,197	15.1	2,176.8
Black, non-Hispanic, male, all ages				
...	All causes	144,802	100.0	842.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	36,596	25.3	212.9
2	Malignant neoplasms (C00-C97)	32,287	22.3	187.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,428	5.8	49.0
4	Cerebrovascular diseases (I60-I69)	7,754	5.4	45.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6,780	4.7	39.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	5,203	3.6	30.3
7	Diabetes mellitus (E10-E14)	5,158	3.6	30.0
8	Chronic lower respiratory diseases (J40-J47)	4,282	3.0	24.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,389	2.3	19.7
10	Influenza and pneumonia (J10-J18)	2,734	1.9	15.9
11	Septicemia (A40-A41)	2,668	1.8	15.5
12	Certain conditions originating in the perinatal period (P00-P96)	2,640	1.8	15.4
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,701	1.2	9.9
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,595	1.1	9.3
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,592	1.1	9.3
...	All other causes (Residual)	21,995	15.2	127.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 1-4 years				
...	All causes	681	100.0	55.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	231	33.9	18.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	89	13.1	7.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	58	8.5	4.8
4	Malignant neoplasms (C00-C97)	33	4.8	2.7
5	Influenza and pneumonia (J10-J18)	22	3.2	1.8
6	Chronic lower respiratory diseases (J40-J47)	20	2.9	1.6
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	18	2.6	*
7	Certain conditions originating in the perinatal period (P00-P96)	18	2.6	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	8	1.2	*
10	Cerebrovascular diseases (I60-I69)	6	0.9	*
11	Septicemia (A40-A41)	5	0.7	*
12	Anemias (D50-D64)	4	0.6	*
13	Meningitis (G00,G03)	3	0.4	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.4	*
15	Meningococcal infection (A39)	2	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.3	*
...	All other causes (Residual)	159	23.3	13.0
Black, non-Hispanic, male, 5-14 years				
...	All causes	977	100.0	29.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	391	40.0	11.9
2	Malignant neoplasms (C00-C97)	103	10.5	3.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	84	8.6	2.6
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	47	4.8	1.4
5	Chronic lower respiratory diseases (J40-J47)	39	4.0	1.2
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	33	3.4	1.0
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	29	3.0	0.9
8	Cerebrovascular diseases (I60-I69)	15	1.5	*
9	Anemias (D50-D64)	14	1.4	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	10	1.0	*
10	Influenza and pneumonia (J10-J18)	10	1.0	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	10	1.0	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	9	0.9	*
14	Septicemia (A40-A41)	7	0.7	*
15	Certain conditions originating in the perinatal period (P00-P96)	6	0.6	*
...	All other causes (Residual)	170	17.4	5.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 15-24 years				
...	All causes	5,248	100.0	177.4
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,540	48.4	85.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,260	24.0	42.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	342	6.5	11.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	185	3.5	6.3
5	Malignant neoplasms (C00-C97)	137	2.6	4.6
6	Human immunodeficiency virus (HIV) disease (B20-B24)	60	1.1	2.0
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	59	1.1	2.0
8	Chronic lower respiratory diseases (J40-J47)	58	1.1	2.0
9	Anemias (D50-D64)	49	0.9	1.7
10	Legal intervention (Y35,Y89.0)	36	0.7	1.2
11	Diabetes mellitus (E10-E14)	34	0.6	1.1
12	Cerebrovascular diseases (I60-I69)	22	0.4	0.7
13	Influenza and pneumonia (J10-J18)	20	0.4	0.7
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	0.3	*
15	Septicemia (A40-A41)	10	0.2	*
...	All other causes (Residual)	421	8.0	14.2
Black, non-Hispanic, male, 25-34 years				
...	All causes	6,723	100.0	274.3
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,092	31.1	85.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,279	19.0	52.2
3	Human immunodeficiency virus (HIV) disease (B20-B24)	624	9.3	25.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	612	9.1	25.0
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	441	6.6	18.0
6	Malignant neoplasms (C00-C97)	287	4.3	11.7
7	Diabetes mellitus (E10-E14)	112	1.7	4.6
8	Anemias (D50-D64)	72	1.1	2.9
8	Cerebrovascular diseases (I60-I69)	72	1.1	2.9
10	Chronic lower respiratory diseases (J40-J47)	56	0.8	2.3
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	49	0.7	2.0
12	Septicemia (A40-A41)	47	0.7	1.9
13	Influenza and pneumonia (J10-J18)	43	0.6	1.8
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	43	0.6	1.8
15	Legal intervention (Y35,Y89.0)	27	0.4	1.1
...	All other causes (Residual)	867	12.9	35.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 35-44 years				
...	All causes	11,661	100.0	445.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,031	17.4	77.6
2	Human immunodeficiency virus (HIV) disease (B20-B24)	1,978	17.0	75.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,562	13.4	59.7
4	Malignant neoplasms (C00-C97)	1,158	9.9	44.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,042	8.9	39.8
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	345	3.0	13.2
7	Cerebrovascular diseases (I60-I69)	331	2.8	12.7
8	Diabetes mellitus (E10-E14)	316	2.7	12.1
9	Chronic liver disease and cirrhosis (K70,K73-K74)	180	1.5	6.9
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	161	1.4	6.2
11	Chronic lower respiratory diseases (J40-J47)	150	1.3	5.7
12	Influenza and pneumonia (J10-J18)	136	1.2	5.2
13	Septicemia (A40-A41)	130	1.1	5.0
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	112	1.0	4.3
15	Viral hepatitis (B15-B19)	67	0.6	2.6
...	All other causes (Residual)	1,962	16.8	75.0
Black, non-Hispanic, male, 45-54 years				
...	All causes	20,839	100.0	1,002.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,214	25.0	250.9
2	Malignant neoplasms (C00-C97)	4,203	20.2	202.3
3	Human immunodeficiency virus (HIV) disease (B20-B24)	1,791	8.6	86.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,523	7.3	73.3
5	Cerebrovascular diseases (I60-I69)	986	4.7	47.5
6	Diabetes mellitus (E10-E14)	744	3.6	35.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	637	3.1	30.7
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	572	2.7	27.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	428	2.1	20.6
10	Septicemia (A40-A41)	312	1.5	15.0
11	Chronic lower respiratory diseases (J40-J47)	304	1.5	14.6
12	Viral hepatitis (B15-B19)	262	1.3	12.6
13	Influenza and pneumonia (J10-J18)	239	1.1	11.5
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	234	1.1	11.3
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	212	1.0	10.2
...	All other causes (Residual)	3,178	15.3	152.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Black, non-Hispanic, male, 55-64 years			
...	All causes	23,706	100.0	2,074.5
1	Malignant neoplasms (C00-C97)	7,106	30.0	621.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,028	29.6	615.0
3	Cerebrovascular diseases (I60-I69)	1,288	5.4	112.7
4	Diabetes mellitus (E10-E14)	1,104	4.7	96.6
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	791	3.3	69.2
6	Chronic lower respiratory diseases (J40-J47)	607	2.6	53.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	573	2.4	50.1
8	Human immunodeficiency virus (HIV) disease (B20-B24)	512	2.2	44.8
9	Chronic liver disease and cirrhosis (K70,K73-K74)	436	1.8	38.2
10	Septicemia (A40-A41)	418	1.8	36.6
11	Influenza and pneumonia (J10-J18)	338	1.4	29.6
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	274	1.2	24.0
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	180	0.8	15.8
14	Aortic aneurysm and dissection (I71)	138	0.6	12.1
15	Viral hepatitis (B15-B19)	122	0.5	10.7
...	All other causes (Residual)	2,791	11.8	244.2
	Black, non-Hispanic, male, 65-74 years			
...	All causes	27,859	100.0	4,078.9
1	Malignant neoplasms (C00-C97)	8,835	31.7	1,293.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,247	29.6	1,207.5
3	Cerebrovascular diseases (I60-I69)	1,826	6.6	267.3
4	Diabetes mellitus (E10-E14)	1,320	4.7	193.3
5	Chronic lower respiratory diseases (J40-J47)	1,093	3.9	160.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	794	2.9	116.3
7	Septicemia (A40-A41)	598	2.1	87.6
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	527	1.9	77.2
9	Influenza and pneumonia (J10-J18)	454	1.6	66.5
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	359	1.3	52.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	221	0.8	32.4
12	Human immunodeficiency virus (HIV) disease (B20-B24)	182	0.7	26.6
13	Pneumonitis due to solids and liquids (J69)	152	0.5	22.3
14	Aortic aneurysm and dissection (I71)	132	0.5	19.3
15	Alzheimer's disease (G30)	122	0.4	17.9
...	All other causes (Residual)	2,997	10.8	438.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 75-84 years				
...	All causes	28,137	100.0	8,246.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,428	30.0	2,470.0
2	Malignant neoplasms (C00-C97)	7,669	27.3	2,247.6
3	Cerebrovascular diseases (I60-I69)	2,031	7.2	595.2
4	Chronic lower respiratory diseases (J40-J47)	1,350	4.8	395.7
5	Diabetes mellitus (E10-E14)	1,135	4.0	332.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	809	2.9	237.1
7	Influenza and pneumonia (J10-J18)	789	2.8	231.2
8	Septicemia (A40-A41)	729	2.6	213.7
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	448	1.6	131.3
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	442	1.6	129.5
11	Alzheimer's disease (G30)	418	1.5	122.5
12	Pneumonitis due to solids and liquids (J69)	253	0.9	74.1
13	Atherosclerosis (I70)	153	0.5	44.8
14	Parkinson's disease (G20-G21)	138	0.5	40.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	134	0.5	39.3
...	All other causes (Residual)	3,211	11.4	941.1
Black, non-Hispanic, male, 85 years and over				
...	All causes	14,422	100.0	15,776.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,718	32.7	5,161.2
2	Malignant neoplasms (C00-C97)	2,750	19.1	3,008.4
3	Cerebrovascular diseases (I60-I69)	1,158	8.0	1,266.8
4	Influenza and pneumonia (J10-J18)	636	4.4	695.8
5	Chronic lower respiratory diseases (J40-J47)	597	4.1	653.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	520	3.6	568.9
7	Diabetes mellitus (E10-E14)	387	2.7	423.4
8	Alzheimer's disease (G30)	379	2.6	414.6
9	Septicemia (A40-A41)	365	2.5	399.3
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	253	1.8	276.8
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	240	1.7	262.5
12	Pneumonitis due to solids and liquids (J69)	217	1.5	237.4
13	Atherosclerosis (I70)	100	0.7	109.4
14	Parkinson's disease (G20-G21)	80	0.6	87.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	74	0.5	81.0
...	All other causes (Residual)	1,948	13.5	2,131.0

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, all ages				
...	All causes	141,771	100.0	748.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	40,098	28.3	211.6
2	Malignant neoplasms (C00-C97)	29,709	21.0	156.7
3	Cerebrovascular diseases (I60-I69)	10,937	7.7	57.7
4	Diabetes mellitus (E10-E14)	7,425	5.2	39.2
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,021	2.8	21.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,857	2.7	20.3
7	Chronic lower respiratory diseases (J40-J47)	3,448	2.4	18.2
8	Septicemia (A40-A41)	3,406	2.4	18.0
9	Influenza and pneumonia (J10-J18)	3,069	2.2	16.2
10	Human immunodeficiency virus (HIV) disease (B20-B24)	2,511	1.8	13.2
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,464	1.7	13.0
12	Alzheimer's disease (G30)	2,363	1.7	12.5
13	Certain conditions originating in the perinatal period (P00-P96)	2,137	1.5	11.3
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,367	1.0	7.2
15	Chronic liver disease and cirrhosis (K70,K73-K74)	953	0.7	5.0
...	All other causes (Residual)	24,006	16.9	126.7
Black, non-Hispanic, female, 1-4 years				
...	All causes	482	100.0	40.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	119	24.7	10.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	67	13.9	5.7
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	55	11.4	4.7
4	Malignant neoplasms (C00-C97)	38	7.9	3.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	22	4.6	1.9
6	Influenza and pneumonia (J10-J18)	18	3.7	*
7	Septicemia (A40-A41)	12	2.5	*
8	Chronic lower respiratory diseases (J40-J47)	9	1.9	*
9	Certain conditions originating in the perinatal period (P00-P96)	8	1.7	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	6	1.2	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.8	*
11	Cerebrovascular diseases (I60-I69)	4	0.8	*
13	Anemias (D50-D64)	3	0.6	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.6	*
15	Pneumonitis due to solids and liquids (J69)	2	0.4	*
...	All other causes (Residual)	112	23.2	9.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 5-14 years				
...	All causes	657	100.0	20.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	216	32.9	6.8
2	Malignant neoplasms (C00-C97)	65	9.9	2.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	48	7.3	1.5
4	Chronic lower respiratory diseases (J40-J47)	35	5.3	1.1
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	34	5.2	1.1
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	32	4.9	1.0
7	Septicemia (A40-A41)	13	2.0	*
8	Anemias (D50-D64)	12	1.8	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	9	1.4	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	9	1.4	*
9	Diabetes mellitus (E10-E14)	9	1.4	*
12	Cerebrovascular diseases (I60-I69)	8	1.2	*
12	Influenza and pneumonia (J10-J18)	8	1.2	*
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6	0.9	*
15	Meningitis (G00,G03)	5	0.8	*
...	All other causes (Residual)	148	22.5	4.7
Black, non-Hispanic, female, 15-24 years				
...	All causes	1,656	100.0	56.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	420	25.4	14.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	314	19.0	10.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	114	6.9	3.9
4	Malignant neoplasms (C00-C97)	105	6.3	3.6
5	Human immunodeficiency virus (HIV) disease (B20-B24)	64	3.9	2.2
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	50	3.0	1.7
7	Pregnancy, childbirth and the puerperium (O00-099)	45	2.7	1.5
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	39	2.4	1.3
9	Anemias (D50-D64)	29	1.8	1.0
10	Diabetes mellitus (E10-E14)	23	1.4	0.8
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21	1.3	0.7
12	Chronic lower respiratory diseases (J40-J47)	20	1.2	0.7
13	Influenza and pneumonia (J10-J18)	18	1.1	*
14	Cerebrovascular diseases (I60-I69)	16	1.0	*
15	Septicemia (A40-A41)	14	0.8	*
...	All other causes (Residual)	364	22.0	12.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Black, non-Hispanic, female, 25-34 years			
...	All causes	3,286	100.0	121.3
1	Human immunodeficiency virus (HIV) disease (B20-B24)	486	14.8	17.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	397	12.1	14.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	381	11.6	14.1
4	Malignant neoplasms (C00-C97)	376	11.4	13.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	333	10.1	12.3
6	Cerebrovascular diseases (I60-I69)	94	2.9	3.5
7	Diabetes mellitus (E10-E14)	82	2.5	3.0
8	Pregnancy, childbirth and the puerperium (O00-099)	64	1.9	2.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	59	1.8	2.2
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	56	1.7	2.1
11	Septicemia (A40-A41)	55	1.7	2.0
11	Anemias (D50-D64)	55	1.7	2.0
13	Chronic lower respiratory diseases (J40-J47)	46	1.4	1.7
14	Influenza and pneumonia (J10-J18)	37	1.1	1.4
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	32	1.0	1.2
...	All other causes (Residual)	733	22.3	27.1
	Black, non-Hispanic, female, 35-44 years			
...	All causes	8,294	100.0	280.1
1	Malignant neoplasms (C00-C97)	1,712	20.6	57.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,281	15.4	43.3
3	Human immunodeficiency virus (HIV) disease (B20-B24)	1,033	12.5	34.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	665	8.0	22.5
5	Cerebrovascular diseases (I60-I69)	410	4.9	13.8
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	315	3.8	10.6
7	Diabetes mellitus (E10-E14)	238	2.9	8.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	145	1.7	4.9
9	Chronic lower respiratory diseases (J40-J47)	142	1.7	4.8
10	Chronic liver disease and cirrhosis (K70,K73-K74)	133	1.6	4.5
11	Septicemia (A40-A41)	124	1.5	4.2
12	Influenza and pneumonia (J10-J18)	114	1.4	3.9
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	87	1.0	2.9
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	77	0.9	2.6
15	Anemias (D50-D64)	76	0.9	2.6
...	All other causes (Residual)	1,742	21.0	58.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Black, non-Hispanic, female, 45-54 years			
...	All causes	14,372	100.0	592.7
1	Malignant neoplasms (C00-C97)	4,173	29.0	172.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,104	21.6	128.0
3	Cerebrovascular diseases (I60-I69)	888	6.2	36.6
4	Human immunodeficiency virus (HIV) disease (B20-B24)	678	4.7	28.0
5	Diabetes mellitus (E10-E14)	649	4.5	26.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	622	4.3	25.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	319	2.2	13.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	317	2.2	13.1
9	Chronic lower respiratory diseases (J40-J47)	285	2.0	11.8
10	Septicemia (A40-A41)	281	2.0	11.6
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	179	1.2	7.4
12	Influenza and pneumonia (J10-J18)	160	1.1	6.6
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	138	1.0	5.7
14	Viral hepatitis (B15-B19)	106	0.7	4.4
15	Anemias (D50-D64)	86	0.6	3.5
...	All other causes (Residual)	2,387	16.6	98.4
	Black, non-Hispanic, female, 55-64 years			
...	All causes	17,172	100.0	1,206.3
1	Malignant neoplasms (C00-C97)	5,587	32.5	392.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,526	26.4	317.9
3	Diabetes mellitus (E10-E14)	1,100	6.4	77.3
4	Cerebrovascular diseases (I60-I69)	1,015	5.9	71.3
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	499	2.9	35.1
6	Chronic lower respiratory diseases (J40-J47)	460	2.7	32.3
7	Septicemia (A40-A41)	450	2.6	31.6
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	284	1.7	20.0
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	276	1.6	19.4
10	Influenza and pneumonia (J10-J18)	216	1.3	15.2
11	Chronic liver disease and cirrhosis (K70,K73-K74)	213	1.2	15.0
12	Human immunodeficiency virus (HIV) disease (B20-B24)	163	0.9	11.5
13	Viral hepatitis (B15-B19)	62	0.4	4.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	61	0.4	4.3
15	Aortic aneurysm and dissection (I71)	54	0.3	3.8
...	All other causes (Residual)	2,206	12.8	155.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Black, non-Hispanic, female, 65-74 years			
...	All causes	24,889	100.0	2,584.8
1	Malignant neoplasms (C00-C97)	7,249	29.1	752.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,164	28.8	744.0
3	Diabetes mellitus (E10-E14)	1,832	7.4	190.3
4	Cerebrovascular diseases (I60-I69)	1,774	7.1	184.2
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	876	3.5	91.0
6	Chronic lower respiratory diseases (J40-J47)	799	3.2	83.0
7	Septicemia (A40-A41)	653	2.6	67.8
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	436	1.8	45.3
9	Influenza and pneumonia (J10-J18)	383	1.5	39.8
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	291	1.2	30.2
11	Alzheimer's disease (G30)	155	0.6	16.1
12	Chronic liver disease and cirrhosis (K70,K73-K74)	137	0.6	14.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	113	0.5	11.7
14	Aortic aneurysm and dissection (I71)	112	0.4	11.6
15	Pneumonitis due to solids and liquids (J69)	98	0.4	10.2
...	All other causes (Residual)	2,817	11.3	292.6
	Black, non-Hispanic, female, 75-84 years			
...	All causes	34,046	100.0	5,644.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,092	32.6	1,839.1
2	Malignant neoplasms (C00-C97)	6,849	20.1	1,135.6
3	Cerebrovascular diseases (I60-I69)	3,246	9.5	538.2
4	Diabetes mellitus (E10-E14)	2,078	6.1	344.5
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,132	3.3	187.7
6	Chronic lower respiratory diseases (J40-J47)	1,016	3.0	168.5
7	Septicemia (A40-A41)	904	2.7	149.9
8	Influenza and pneumonia (J10-J18)	795	2.3	131.8
9	Alzheimer's disease (G30)	755	2.2	125.2
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	691	2.0	114.6
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	351	1.0	58.2
12	Pneumonitis due to solids and liquids (J69)	246	0.7	40.8
13	Atherosclerosis (I70)	196	0.6	32.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	166	0.5	27.5
14	Aortic aneurysm and dissection (I71)	166	0.5	27.5
...	All other causes (Residual)	4,363	12.8	723.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Black, non-Hispanic, female, 85 years and over			
...	All causes	33,143	100.0	13,855.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,312	37.1	5,147.0
2	Malignant neoplasms (C00-C97)	3,546	10.7	1,482.4
3	Cerebrovascular diseases (I60-I69)	3,464	10.5	1,448.1
4	Alzheimer's disease (G30)	1,435	4.3	599.9
5	Diabetes mellitus (E10-E14)	1,414	4.3	591.1
6	Influenza and pneumonia (J10-J18)	1,281	3.9	535.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	948	2.9	396.3
8	Septicemia (A40-A41)	861	2.6	359.9
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	780	2.4	326.1
10	Chronic lower respiratory diseases (J40-J47)	629	1.9	263.0
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	340	1.0	142.1
12	Pneumonitis due to solids and liquids (J69)	336	1.0	140.5
13	Atherosclerosis (I70)	290	0.9	121.2
14	Nutritional deficiencies (E40-E64)	157	0.5	65.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	139	0.4	58.1
...	All other causes (Residual)	5,211	15.7	2,178.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable