

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, all ages				
...	All causes	122,026	100.0	305.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	28,298	23.2	70.9
2	Malignant neoplasms (C00-C97)	24,070	19.7	60.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,418	8.5	26.1
4	Cerebrovascular diseases (I60-I69)	6,658	5.5	16.7
5	Diabetes mellitus (E10-E14)	6,179	5.1	15.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	3,382	2.8	8.5
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,355	2.7	8.4
8	Chronic lower respiratory diseases (J40-J47)	3,174	2.6	8.0
9	Influenza and pneumonia (J10-J18)	2,948	2.4	7.4
10	Certain conditions originating in the perinatal period (P00-P96)	2,628	2.2	6.6
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,170	1.8	5.4
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,007	1.6	5.0
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,890	1.5	4.7
14	Alzheimer's disease (G30)	1,835	1.5	4.6
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,787	1.5	4.5
...	All other causes (Residual)	21,227	17.4	53.2
Hispanic, both sexes, 1-4 years				
...	All causes	993	100.0	30.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	325	32.7	9.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	122	12.3	3.7
3	Malignant neoplasms (C00-C97)	98	9.9	3.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	82	8.3	2.5
5	Influenza and pneumonia (J10-J18)	37	3.7	1.1
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	35	3.5	1.1
7	Certain conditions originating in the perinatal period (P00-P96)	15	1.5	*
8	Chronic lower respiratory diseases (J40-J47)	11	1.1	*
9	Septicemia (A40-A41)	10	1.0	*
10	Cerebrovascular diseases (I60-I69)	8	0.8	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.7	*
12	Acute bronchitis and bronchiolitis (J20-J21)	6	0.6	*
13	Anemias (D50-D64)	5	0.5	*
14	Meningitis (G00,G03)	4	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.4	*
...	All other causes (Residual)	224	22.6	6.8

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 5-14 years				
...	All causes	1,173	100.0	15.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	431	36.7	5.8
2	Malignant neoplasms (C00-C97)	217	18.5	2.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	65	5.5	0.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	53	4.5	0.7
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	43	3.7	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	31	2.6	0.4
7	Influenza and pneumonia (J10-J18)	22	1.9	0.3
8	Chronic lower respiratory diseases (J40-J47)	15	1.3	*
9	Cerebrovascular diseases (I60-I69)	14	1.2	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	1.1	*
11	Certain conditions originating in the perinatal period (P00-P96)	8	0.7	*
12	Septicemia (A40-A41)	7	0.6	*
13	Diseases of appendix (K35-K38)	5	0.4	*
14	Anemias (D50-D64)	4	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.3	*
...	All other causes (Residual)	241	20.5	3.2
Hispanic, both sexes, 15-24 years				
...	All causes	5,394	100.0	77.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,366	43.9	33.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,286	23.8	18.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	491	9.1	7.0
4	Malignant neoplasms (C00-C97)	288	5.3	4.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	135	2.5	1.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	57	1.1	0.8
7	Influenza and pneumonia (J10-J18)	36	0.7	0.5
8	Human immunodeficiency virus (HIV) disease (B20-B24)	31	0.6	0.4
9	Cerebrovascular diseases (I60-I69)	28	0.5	0.4
9	Legal intervention (Y35,Y89.0)	28	0.5	0.4
11	Pregnancy, childbirth and the puerperium (O00-099)	24	0.4	0.3
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	21	0.4	0.3
13	Chronic lower respiratory diseases (J40-J47)	20	0.4	0.3
14	Diabetes mellitus (E10-E14)	19	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19	0.4	*
...	All other causes (Residual)	545	10.1	7.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 25-34 years				
...	All causes	6,189	100.0	83.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,167	35.0	29.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	971	15.7	13.2
3	Malignant neoplasms (C00-C97)	522	8.4	7.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	499	8.1	6.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	347	5.6	4.7
6	Human immunodeficiency virus (HIV) disease (B20-B24)	244	3.9	3.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	100	1.6	1.4
8	Cerebrovascular diseases (I60-I69)	89	1.4	1.2
9	Diabetes mellitus (E10-E14)	64	1.0	0.9
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	58	0.9	0.8
11	Influenza and pneumonia (J10-J18)	54	0.9	0.7
12	Pregnancy, childbirth and the puerperium (O00-099)	46	0.7	0.6
13	Septicemia (A40-A41)	35	0.6	0.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	32	0.5	0.4
14	Legal intervention (Y35,Y89.0)	32	0.5	0.4
...	All other causes (Residual)	929	15.0	12.6
Hispanic, both sexes, 35-44 years				
...	All causes	8,576	100.0	144.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,877	21.9	31.7
2	Malignant neoplasms (C00-C97)	1,355	15.8	22.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	879	10.2	14.9
4	Human immunodeficiency virus (HIV) disease (B20-B24)	727	8.5	12.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	502	5.9	8.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	469	5.5	7.9
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	402	4.7	6.8
8	Cerebrovascular diseases (I60-I69)	267	3.1	4.5
9	Diabetes mellitus (E10-E14)	197	2.3	3.3
10	Viral hepatitis (B15-B19)	107	1.2	1.8
11	Septicemia (A40-A41)	86	1.0	1.5
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	76	0.9	1.3
13	Influenza and pneumonia (J10-J18)	74	0.9	1.3
14	Chronic lower respiratory diseases (J40-J47)	58	0.7	1.0
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	44	0.5	0.7
...	All other causes (Residual)	1,456	17.0	24.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 45-54 years				
...	All causes	12,516	100.0	327.8
1	Malignant neoplasms (C00-C97)	3,081	24.6	80.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,035	16.3	53.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,254	10.0	32.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	927	7.4	24.3
5	Human immunodeficiency virus (HIV) disease (B20-B24)	606	4.8	15.9
6	Diabetes mellitus (E10-E14)	593	4.7	15.5
7	Cerebrovascular diseases (I60-I69)	569	4.5	14.9
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	275	2.2	7.2
9	Viral hepatitis (B15-B19)	266	2.1	7.0
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	246	2.0	6.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	174	1.4	4.6
12	Influenza and pneumonia (J10-J18)	151	1.2	4.0
13	Septicemia (A40-A41)	148	1.2	3.9
14	Chronic lower respiratory diseases (J40-J47)	134	1.1	3.5
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	75	0.6	2.0
...	All other causes (Residual)	1,982	15.8	51.9
Hispanic, both sexes, 55-64 years				
...	All causes	14,913	100.0	714.3
1	Malignant neoplasms (C00-C97)	4,482	30.1	214.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,454	23.2	165.4
3	Diabetes mellitus (E10-E14)	1,121	7.5	53.7
4	Chronic liver disease and cirrhosis (K70,K73-K74)	802	5.4	38.4
5	Cerebrovascular diseases (I60-I69)	736	4.9	35.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	639	4.3	30.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	318	2.1	15.2
8	Chronic lower respiratory diseases (J40-J47)	266	1.8	12.7
9	Influenza and pneumonia (J10-J18)	224	1.5	10.7
10	Human immunodeficiency virus (HIV) disease (B20-B24)	204	1.4	9.8
11	Septicemia (A40-A41)	190	1.3	9.1
12	Viral hepatitis (B15-B19)	151	1.0	7.2
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	145	1.0	6.9
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	121	0.8	5.8
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	75	0.5	3.6
...	All other causes (Residual)	1,985	13.3	95.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 65-74 years				
...	All causes	20,596	100.0	1,685.6
1	Malignant neoplasms (C00-C97)	5,929	28.8	485.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,635	27.4	461.2
3	Diabetes mellitus (E10-E14)	1,611	7.8	131.8
4	Cerebrovascular diseases (I60-I69)	1,202	5.8	98.4
5	Chronic lower respiratory diseases (J40-J47)	651	3.2	53.3
6	Chronic liver disease and cirrhosis (K70,K73-K74)	624	3.0	51.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	484	2.3	39.6
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	466	2.3	38.1
9	Influenza and pneumonia (J10-J18)	428	2.1	35.0
10	Septicemia (A40-A41)	270	1.3	22.1
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	204	1.0	16.7
12	Alzheimer's disease (G30)	179	0.9	14.6
13	Viral hepatitis (B15-B19)	106	0.5	8.7
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	106	0.5	8.7
15	Aortic aneurysm and dissection (I71)	102	0.5	8.3
...	All other causes (Residual)	2,599	12.6	212.7
Hispanic, both sexes, 75-84 years				
...	All causes	26,219	100.0	4,093.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,077	30.8	1,261.0
2	Malignant neoplasms (C00-C97)	5,701	21.7	890.1
3	Cerebrovascular diseases (I60-I69)	2,029	7.7	316.8
4	Diabetes mellitus (E10-E14)	1,743	6.6	272.1
5	Chronic lower respiratory diseases (J40-J47)	1,095	4.2	171.0
6	Influenza and pneumonia (J10-J18)	860	3.3	134.3
7	Alzheimer's disease (G30)	664	2.5	103.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	578	2.2	90.2
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	454	1.7	70.9
10	Septicemia (A40-A41)	397	1.5	62.0
10	Chronic liver disease and cirrhosis (K70,K73-K74)	397	1.5	62.0
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	285	1.1	44.5
13	Parkinson's disease (G20-G21)	258	1.0	40.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	185	0.7	28.9
15	Aortic aneurysm and dissection (I71)	171	0.7	26.7
...	All other causes (Residual)	3,325	12.7	519.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 85 years and over				
...	All causes	20,129	100.0	10,431.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,589	37.7	3,932.9
2	Malignant neoplasms (C00-C97)	2,383	11.8	1,234.9
3	Cerebrovascular diseases (I60-I69)	1,699	8.4	880.5
4	Influenza and pneumonia (J10-J18)	996	4.9	516.2
5	Alzheimer's disease (G30)	952	4.7	493.4
6	Chronic lower respiratory diseases (J40-J47)	888	4.4	460.2
7	Diabetes mellitus (E10-E14)	826	4.1	428.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	450	2.2	233.2
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	330	1.6	171.0
10	Septicemia (A40-A41)	293	1.5	151.8
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	293	1.5	151.8
12	Pneumonitis due to solids and liquids (J69)	218	1.1	113.0
13	Atherosclerosis (I70)	166	0.8	86.0
14	Parkinson's disease (G20-G21)	150	0.7	77.7
15	Aortic aneurysm and dissection (I71)	95	0.5	49.2
...	All other causes (Residual)	2,801	13.9	1,451.6
Hispanic, male, all ages				
...	All causes	68,119	100.0	330.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,867	21.8	72.2
2	Malignant neoplasms (C00-C97)	12,671	18.6	61.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,899	11.6	38.3
4	Cerebrovascular diseases (I60-I69)	3,070	4.5	14.9
5	Diabetes mellitus (E10-E14)	2,976	4.4	14.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,809	4.1	13.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,387	3.5	11.6
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,711	2.5	8.3
9	Chronic lower respiratory diseases (J40-J47)	1,697	2.5	8.2
10	Human immunodeficiency virus (HIV) disease (B20-B24)	1,465	2.2	7.1
11	Certain conditions originating in the perinatal period (P00-P96)	1,459	2.1	7.1
12	Influenza and pneumonia (J10-J18)	1,429	2.1	6.9
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,069	1.6	5.2
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	937	1.4	4.5
15	Septicemia (A40-A41)	721	1.1	3.5
...	All other causes (Residual)	10,952	16.1	53.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 1-4 years				
...	All causes	569	100.0	33.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	200	35.1	11.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	62	10.9	3.7
3	Malignant neoplasms (C00-C97)	60	10.5	3.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	36	6.3	2.1
5	Influenza and pneumonia (J10-J18)	24	4.2	1.4
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	19	3.3	*
7	Chronic lower respiratory diseases (J40-J47)	8	1.4	*
7	Certain conditions originating in the perinatal period (P00-P96)	8	1.4	*
9	Cerebrovascular diseases (I60-I69)	7	1.2	*
10	Septicemia (A40-A41)	6	1.1	*
11	Acute bronchitis and bronchiolitis (J20-J21)	4	0.7	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.7	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.5	*
14	Anemias (D50-D64)	2	0.4	*
14	Nutritional deficiencies (E40-E64)	2	0.4	*
14	Meningitis (G00,G03)	2	0.4	*
14	Pneumonitis due to solids and liquids (J69)	2	0.4	*
14	Hernia (K40-K46)	2	0.4	*
...	All other causes (Residual)	118	20.7	7.0
Hispanic, male, 5-14 years				
...	All causes	706	100.0	18.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	289	40.9	7.5
2	Malignant neoplasms (C00-C97)	123	17.4	3.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	36	5.1	0.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	31	4.4	0.8
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	28	4.0	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	19	2.7	*
7	Influenza and pneumonia (J10-J18)	12	1.7	*
8	Chronic lower respiratory diseases (J40-J47)	10	1.4	*
9	Cerebrovascular diseases (I60-I69)	8	1.1	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	1.0	*
11	Septicemia (A40-A41)	4	0.6	*
11	Certain conditions originating in the perinatal period (P00-P96)	4	0.6	*
13	Meningitis (G00,G03)	3	0.4	*
14	Anemias (D50-D64)	2	0.3	*
15	Diabetes mellitus (E10-E14)	1	0.1	*
15	Pneumonitis due to solids and liquids (J69)	1	0.1	*
15	Diseases of appendix (K35-K38)	1	0.1	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.1	*
...	All other causes (Residual)	126	17.8	3.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 15-24 years				
...	All causes	4,244	100.0	112.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,905	44.9	50.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,139	26.8	30.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	421	9.9	11.2
4	Malignant neoplasms (C00-C97)	186	4.4	4.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	101	2.4	2.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	30	0.7	0.8
7	Legal intervention (Y35,Y89.0)	28	0.7	0.7
8	Human immunodeficiency virus (HIV) disease (B20-B24)	20	0.5	0.5
8	Influenza and pneumonia (J10-J18)	20	0.5	0.5
10	Cerebrovascular diseases (I60-I69)	15	0.4	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	0.3	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	0.3	*
13	Chronic lower respiratory diseases (J40-J47)	12	0.3	*
14	Diabetes mellitus (E10-E14)	11	0.3	*
15	Septicemia (A40-A41)	8	0.2	*
...	All other causes (Residual)	322	7.6	8.6
Hispanic, male, 25-34 years				
...	All causes	4,742	100.0	118.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,827	38.5	45.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	852	18.0	21.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	434	9.2	10.8
4	Malignant neoplasms (C00-C97)	274	5.8	6.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	244	5.1	6.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	184	3.9	4.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	79	1.7	2.0
8	Cerebrovascular diseases (I60-I69)	51	1.1	1.3
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	37	0.8	0.9
10	Influenza and pneumonia (J10-J18)	35	0.7	0.9
11	Diabetes mellitus (E10-E14)	33	0.7	0.8
12	Legal intervention (Y35,Y89.0)	32	0.7	0.8
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	21	0.4	0.5
14	Septicemia (A40-A41)	20	0.4	0.5
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	0.3	*
...	All other causes (Residual)	603	12.7	15.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 35-44 years				
...	All causes	5,887	100.0	189.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,488	25.3	48.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	654	11.1	21.1
3	Malignant neoplasms (C00-C97)	566	9.6	18.2
4	Human immunodeficiency virus (HIV) disease (B20-B24)	550	9.3	17.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	401	6.8	12.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	380	6.5	12.3
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	341	5.8	11.0
8	Cerebrovascular diseases (I60-I69)	162	2.8	5.2
9	Diabetes mellitus (E10-E14)	121	2.1	3.9
10	Viral hepatitis (B15-B19)	80	1.4	2.6
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	49	0.8	1.6
12	Septicemia (A40-A41)	48	0.8	1.5
13	Influenza and pneumonia (J10-J18)	41	0.7	1.3
14	Chronic lower respiratory diseases (J40-J47)	30	0.5	1.0
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	25	0.4	0.8
...	All other causes (Residual)	951	16.2	30.7
Hispanic, male, 45-54 years				
...	All causes	8,336	100.0	436.4
1	Malignant neoplasms (C00-C97)	1,550	18.6	81.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,520	18.2	79.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	966	11.6	50.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	768	9.2	40.2
5	Human immunodeficiency virus (HIV) disease (B20-B24)	499	6.0	26.1
6	Diabetes mellitus (E10-E14)	345	4.1	18.1
6	Cerebrovascular diseases (I60-I69)	345	4.1	18.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	220	2.6	11.5
9	Viral hepatitis (B15-B19)	207	2.5	10.8
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	194	2.3	10.2
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	107	1.3	5.6
12	Influenza and pneumonia (J10-J18)	100	1.2	5.2
13	Septicemia (A40-A41)	94	1.1	4.9
14	Chronic lower respiratory diseases (J40-J47)	61	0.7	3.2
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	53	0.6	2.8
...	All other causes (Residual)	1,307	15.7	68.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 55-64 years				
...	All causes	9,122	100.0	920.8
1	Malignant neoplasms (C00-C97)	2,442	26.8	246.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,334	25.6	235.6
3	Diabetes mellitus (E10-E14)	618	6.8	62.4
4	Chronic liver disease and cirrhosis (K70,K73-K74)	616	6.8	62.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	449	4.9	45.3
6	Cerebrovascular diseases (I60-I69)	431	4.7	43.5
7	Chronic lower respiratory diseases (J40-J47)	164	1.8	16.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	160	1.8	16.2
9	Human immunodeficiency virus (HIV) disease (B20-B24)	153	1.7	15.4
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	129	1.4	13.0
11	Influenza and pneumonia (J10-J18)	115	1.3	11.6
12	Viral hepatitis (B15-B19)	103	1.1	10.4
13	Septicemia (A40-A41)	95	1.0	9.6
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	73	0.8	7.4
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	59	0.6	6.0
...	All other causes (Residual)	1,181	12.9	119.2
Hispanic, male, 65-74 years				
...	All causes	11,465	100.0	2,115.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,387	29.5	625.0
2	Malignant neoplasms (C00-C97)	3,347	29.2	617.6
3	Diabetes mellitus (E10-E14)	793	6.9	146.3
4	Cerebrovascular diseases (I60-I69)	617	5.4	113.9
5	Chronic lower respiratory diseases (J40-J47)	369	3.2	68.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	332	2.9	61.3
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	289	2.5	53.3
8	Influenza and pneumonia (J10-J18)	235	2.0	43.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	230	2.0	42.4
10	Septicemia (A40-A41)	137	1.2	25.3
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	84	0.7	15.5
12	Alzheimer's disease (G30)	75	0.7	13.8
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	71	0.6	13.1
14	Aortic aneurysm and dissection (I71)	68	0.6	12.5
15	Parkinson's disease (G20-G21)	67	0.6	12.4
...	All other causes (Residual)	1,364	11.9	251.7

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 75-84 years				
...	All causes	12,904	100.0	4,948.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,025	31.2	1,543.5
2	Malignant neoplasms (C00-C97)	3,035	23.5	1,163.9
3	Cerebrovascular diseases (I60-I69)	879	6.8	337.1
4	Diabetes mellitus (E10-E14)	775	6.0	297.2
5	Chronic lower respiratory diseases (J40-J47)	603	4.7	231.2
6	Influenza and pneumonia (J10-J18)	432	3.3	165.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	280	2.2	107.4
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	267	2.1	102.4
9	Alzheimer's disease (G30)	264	2.0	101.2
10	Chronic liver disease and cirrhosis (K70,K73-K74)	186	1.4	71.3
11	Septicemia (A40-A41)	175	1.4	67.1
12	Parkinson's disease (G20-G21)	152	1.2	58.3
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	116	0.9	44.5
14	Aortic aneurysm and dissection (I71)	100	0.8	38.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	88	0.7	33.7
...	All other causes (Residual)	1,527	11.8	585.6
Hispanic, male, 85 years and over				
...	All causes	7,165	100.0	11,029.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,517	35.1	3,874.5
2	Malignant neoplasms (C00-C97)	1,084	15.1	1,668.6
3	Cerebrovascular diseases (I60-I69)	544	7.6	837.4
4	Chronic lower respiratory diseases (J40-J47)	420	5.9	646.5
5	Influenza and pneumonia (J10-J18)	375	5.2	577.2
6	Diabetes mellitus (E10-E14)	277	3.9	426.4
6	Alzheimer's disease (G30)	277	3.9	426.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	193	2.7	297.1
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	127	1.8	195.5
10	Septicemia (A40-A41)	104	1.5	160.1
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	87	1.2	133.9
12	Parkinson's disease (G20-G21)	84	1.2	129.3
13	Pneumonitis due to solids and liquids (J69)	80	1.1	123.1
14	Atherosclerosis (I70)	62	0.9	95.4
15	Aortic aneurysm and dissection (I71)	52	0.7	80.0
...	All other causes (Residual)	882	12.3	1,357.7

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, all ages				
...	All causes	53,907	100.0	279.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,431	24.9	69.6
2	Malignant neoplasms (C00-C97)	11,399	21.1	59.1
3	Cerebrovascular diseases (I60-I69)	3,588	6.7	18.6
4	Diabetes mellitus (E10-E14)	3,203	5.9	16.6
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,519	4.7	13.1
6	Influenza and pneumonia (J10-J18)	1,519	2.8	7.9
7	Chronic lower respiratory diseases (J40-J47)	1,477	2.7	7.7
8	Alzheimer's disease (G30)	1,202	2.2	6.2
9	Certain conditions originating in the perinatal period (P00-P96)	1,169	2.2	6.1
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,101	2.0	5.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	995	1.8	5.2
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	850	1.6	4.4
13	Septicemia (A40-A41)	784	1.5	4.1
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	616	1.1	3.2
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	546	1.0	2.8
...	All other causes (Residual)	9,508	17.6	49.3
Hispanic, female, 1-4 years				
...	All causes	424	100.0	26.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	125	29.5	7.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	60	14.2	3.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	46	10.8	2.9
4	Malignant neoplasms (C00-C97)	38	9.0	2.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	16	3.8	*
6	Influenza and pneumonia (J10-J18)	13	3.1	*
7	Certain conditions originating in the perinatal period (P00-P96)	7	1.7	*
8	Septicemia (A40-A41)	4	0.9	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.9	*
10	Anemias (D50-D64)	3	0.7	*
10	Chronic lower respiratory diseases (J40-J47)	3	0.7	*
12	Meningitis (G00,G03)	2	0.5	*
12	Acute bronchitis and bronchiolitis (J20-J21)	2	0.5	*
14	Meningococcal infection (A39)	1	0.2	*
14	Cerebrovascular diseases (I60-I69)	1	0.2	*
14	Pneumonitis due to solids and liquids (J69)	1	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
...	All other causes (Residual)	97	22.9	6.0

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Hispanic, female, 5-14 years			
...	All causes	467	100.0	12.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	142	30.4	3.9
2	Malignant neoplasms (C00-C97)	94	20.1	2.6
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	37	7.9	1.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	17	3.6	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	12	2.6	*
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	12	2.6	*
7	Influenza and pneumonia (J10-J18)	10	2.1	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.3	*
8	Cerebrovascular diseases (I60-I69)	6	1.3	*
10	Chronic lower respiratory diseases (J40-J47)	5	1.1	*
11	Diseases of appendix (K35-K38)	4	0.9	*
11	Certain conditions originating in the perinatal period (P00-P96)	4	0.9	*
13	Septicemia (A40-A41)	3	0.6	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.6	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.4	*
15	Anemias (D50-D64)	2	0.4	*
15	Diabetes mellitus (E10-E14)	2	0.4	*
...	All other causes (Residual)	106	22.7	2.9
	Hispanic, female, 15-24 years			
...	All causes	1,150	100.0	35.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	461	40.1	14.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	147	12.8	4.5
3	Malignant neoplasms (C00-C97)	102	8.9	3.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	70	6.1	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	34	3.0	1.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	27	2.3	0.8
7	Pregnancy, childbirth and the puerperium (O00-099)	24	2.1	0.7
8	Influenza and pneumonia (J10-J18)	16	1.4	*
9	Cerebrovascular diseases (I60-I69)	13	1.1	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	11	1.0	*
11	Septicemia (A40-A41)	9	0.8	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.7	*
12	Diabetes mellitus (E10-E14)	8	0.7	*
12	Chronic lower respiratory diseases (J40-J47)	8	0.7	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.5	*
...	All other causes (Residual)	206	17.9	6.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 25-34 years				
...	All causes	1,447	100.0	43.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	340	23.5	10.1
2	Malignant neoplasms (C00-C97)	248	17.1	7.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	119	8.2	3.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	103	7.1	3.1
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	65	4.5	1.9
6	Human immunodeficiency virus (HIV) disease (B20-B24)	60	4.1	1.8
7	Pregnancy, childbirth and the puerperium (O00-099)	46	3.2	1.4
8	Cerebrovascular diseases (I60-I69)	38	2.6	1.1
9	Diabetes mellitus (E10-E14)	31	2.1	0.9
10	Chronic liver disease and cirrhosis (K70,K73-K74)	21	1.5	0.6
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	21	1.5	0.6
12	Influenza and pneumonia (J10-J18)	19	1.3	*
13	Septicemia (A40-A41)	15	1.0	*
14	Chronic lower respiratory diseases (J40-J47)	14	1.0	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	1.0	*
...	All other causes (Residual)	293	20.2	8.7
Hispanic, female, 35-44 years				
...	All causes	2,689	100.0	95.5
1	Malignant neoplasms (C00-C97)	789	29.3	28.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	389	14.5	13.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	225	8.4	8.0
4	Human immunodeficiency virus (HIV) disease (B20-B24)	177	6.6	6.3
5	Cerebrovascular diseases (I60-I69)	105	3.9	3.7
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	101	3.8	3.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	89	3.3	3.2
8	Diabetes mellitus (E10-E14)	76	2.8	2.7
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	61	2.3	2.2
10	Septicemia (A40-A41)	38	1.4	1.3
11	Influenza and pneumonia (J10-J18)	33	1.2	1.2
12	Chronic lower respiratory diseases (J40-J47)	28	1.0	1.0
12	Pregnancy, childbirth and the puerperium (O00-099)	28	1.0	1.0
14	Viral hepatitis (B15-B19)	27	1.0	1.0
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	27	1.0	1.0
...	All other causes (Residual)	496	18.4	17.6

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 45-54 years				
...	All causes	4,180	100.0	219.1
1	Malignant neoplasms (C00-C97)	1,531	36.6	80.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	515	12.3	27.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	288	6.9	15.1
4	Diabetes mellitus (E10-E14)	248	5.9	13.0
5	Cerebrovascular diseases (I60-I69)	224	5.4	11.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	159	3.8	8.3
7	Human immunodeficiency virus (HIV) disease (B20-B24)	107	2.6	5.6
8	Chronic lower respiratory diseases (J40-J47)	73	1.7	3.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	67	1.6	3.5
10	Viral hepatitis (B15-B19)	59	1.4	3.1
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	55	1.3	2.9
12	Septicemia (A40-A41)	54	1.3	2.8
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	52	1.2	2.7
14	Influenza and pneumonia (J10-J18)	51	1.2	2.7
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	22	0.5	1.2
...	All other causes (Residual)	675	16.1	35.4
Hispanic, female, 55-64 years				
...	All causes	5,791	100.0	527.9
1	Malignant neoplasms (C00-C97)	2,040	35.2	185.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,120	19.3	102.1
3	Diabetes mellitus (E10-E14)	503	8.7	45.8
4	Cerebrovascular diseases (I60-I69)	305	5.3	27.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	190	3.3	17.3
6	Chronic liver disease and cirrhosis (K70,K73-K74)	186	3.2	17.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	158	2.7	14.4
8	Influenza and pneumonia (J10-J18)	109	1.9	9.9
9	Chronic lower respiratory diseases (J40-J47)	102	1.8	9.3
10	Septicemia (A40-A41)	95	1.6	8.7
11	Human immunodeficiency virus (HIV) disease (B20-B24)	51	0.9	4.6
12	Viral hepatitis (B15-B19)	48	0.8	4.4
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	48	0.8	4.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	25	0.4	2.3
15	Alzheimer's disease (G30)	16	0.3	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	16	0.3	*
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	16	0.3	*
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	16	0.3	*
...	All other causes (Residual)	747	12.9	68.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Hispanic, female, 65-74 years			
...	All causes	9,131	100.0	1,342.9
1	Malignant neoplasms (C00-C97)	2,582	28.3	379.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,248	24.6	330.6
3	Diabetes mellitus (E10-E14)	818	9.0	120.3
4	Cerebrovascular diseases (I60-I69)	585	6.4	86.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	292	3.2	42.9
6	Chronic lower respiratory diseases (J40-J47)	282	3.1	41.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	254	2.8	37.4
8	Influenza and pneumonia (J10-J18)	193	2.1	28.4
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	177	1.9	26.0
10	Septicemia (A40-A41)	133	1.5	19.6
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	120	1.3	17.6
12	Alzheimer's disease (G30)	104	1.1	15.3
13	Viral hepatitis (B15-B19)	64	0.7	9.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	52	0.6	7.6
15	Aortic aneurysm and dissection (I71)	34	0.4	5.0
...	All other causes (Residual)	1,193	13.1	175.5
	Hispanic, female, 75-84 years			
...	All causes	13,315	100.0	3,506.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,052	30.4	1,067.0
2	Malignant neoplasms (C00-C97)	2,666	20.0	702.1
3	Cerebrovascular diseases (I60-I69)	1,150	8.6	302.8
4	Diabetes mellitus (E10-E14)	968	7.3	254.9
5	Chronic lower respiratory diseases (J40-J47)	492	3.7	129.6
6	Influenza and pneumonia (J10-J18)	428	3.2	112.7
7	Alzheimer's disease (G30)	400	3.0	105.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	298	2.2	78.5
9	Septicemia (A40-A41)	222	1.7	58.5
10	Chronic liver disease and cirrhosis (K70,K73-K74)	211	1.6	55.6
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	187	1.4	49.2
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	169	1.3	44.5
13	Parkinson's disease (G20-G21)	106	0.8	27.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	97	0.7	25.5
15	Aortic aneurysm and dissection (I71)	71	0.5	18.7
...	All other causes (Residual)	1,798	13.5	473.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 85 years and over				
...	All causes	12,964	100.0	10,128.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,072	39.1	3,962.5
2	Malignant neoplasms (C00-C97)	1,299	10.0	1,014.8
3	Cerebrovascular diseases (I60-I69)	1,155	8.9	902.3
4	Alzheimer's disease (G30)	675	5.2	527.3
5	Influenza and pneumonia (J10-J18)	621	4.8	485.2
6	Diabetes mellitus (E10-E14)	549	4.2	428.9
7	Chronic lower respiratory diseases (J40-J47)	468	3.6	365.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	257	2.0	200.8
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	243	1.9	189.8
10	Septicemia (A40-A41)	189	1.5	147.7
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	166	1.3	129.7
12	Pneumonitis due to solids and liquids (J69)	138	1.1	107.8
13	Atherosclerosis (I70)	104	0.8	81.3
14	Parkinson's disease (G20-G21)	66	0.5	51.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	59	0.5	46.1
...	All other causes (Residual)	1,903	14.7	1,486.7
White, non-Hispanic, both sexes, all ages				
...	All causes	1,979,465	100.0	993.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	565,808	28.6	284.0
2	Malignant neoplasms (C00-C97)	457,127	23.1	229.5
3	Cerebrovascular diseases (I60-I69)	127,969	6.5	64.2
4	Chronic lower respiratory diseases (J40-J47)	113,541	5.7	57.0
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	82,815	4.2	41.6
6	Alzheimer's disease (G30)	57,268	2.9	28.7
7	Influenza and pneumonia (J10-J18)	54,617	2.8	27.4
8	Diabetes mellitus (E10-E14)	52,894	2.7	26.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	31,497	1.6	15.8
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	26,434	1.3	13.3
11	Septicemia (A40-A41)	25,664	1.3	12.9
12	Chronic liver disease and cirrhosis (K70,K73-K74)	20,602	1.0	10.3
13	Parkinson's disease (G20-G21)	16,499	0.8	8.3
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	15,878	0.8	8.0
15	Pneumonitis due to solids and liquids (J69)	14,845	0.7	7.5
...	All other causes (Residual)	316,007	16.0	158.6

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 1-4 years				
...	All causes	2,539	100.0	27.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	985	38.8	10.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	272	10.7	3.0
3	Malignant neoplasms (C00-C97)	221	8.7	2.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	123	4.8	1.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	83	3.3	0.9
6	Influenza and pneumonia (J10-J18)	80	3.2	0.9
7	Septicemia (A40-A41)	43	1.7	0.5
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	32	1.3	0.3
9	Certain conditions originating in the perinatal period (P00-P96)	31	1.2	0.3
10	Cerebrovascular diseases (I60-I69)	23	0.9	0.2
11	Chronic lower respiratory diseases (J40-J47)	15	0.6	*
12	Meningitis (G00,G03)	14	0.6	*
13	Meningococcal infection (A39)	10	0.4	*
13	Acute bronchitis and bronchiolitis (J20-J21)	10	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	9	0.4	*
...	All other causes (Residual)	579	22.8	6.3
White, non-Hispanic, both sexes, 5-14 years				
...	All causes	3,884	100.0	15.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,535	39.5	6.2
2	Malignant neoplasms (C00-C97)	637	16.4	2.6
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	222	5.7	0.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	151	3.9	0.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	146	3.8	0.6
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	109	2.8	0.4
7	Influenza and pneumonia (J10-J18)	86	2.2	0.3
8	Septicemia (A40-A41)	47	1.2	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	43	1.1	0.2
10	Chronic lower respiratory diseases (J40-J47)	39	1.0	0.2
11	Cerebrovascular diseases (I60-I69)	37	1.0	0.1
12	Diabetes mellitus (E10-E14)	19	0.5	*
13	Anemias (D50-D64)	17	0.4	*
14	Pneumonitis due to solids and liquids (J69)	11	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	0.3	*
...	All other causes (Residual)	774	19.9	3.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 15-24 years				
...	All causes	19,640	100.0	75.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,638	54.2	41.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,832	14.4	10.9
3	Malignant neoplasms (C00-C97)	1,039	5.3	4.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	907	4.6	3.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	605	3.1	2.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	285	1.5	1.1
7	Influenza and pneumonia (J10-J18)	137	0.7	0.5
8	Cerebrovascular diseases (I60-I69)	124	0.6	0.5
9	Diabetes mellitus (E10-E14)	90	0.5	0.3
10	Septicemia (A40-A41)	86	0.4	0.3
11	Chronic lower respiratory diseases (J40-J47)	82	0.4	0.3
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	58	0.3	0.2
13	Pregnancy, childbirth and the puerperium (O00-099)	41	0.2	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	38	0.2	0.1
15	Complications of medical and surgical care (Y40-Y84,Y88)	31	0.2	0.1
...	All other causes (Residual)	2,647	13.5	10.2
White, non-Hispanic, both sexes, 25-34 years				
...	All causes	23,557	100.0	95.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,238	35.0	33.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,811	16.2	15.5
3	Malignant neoplasms (C00-C97)	2,354	10.0	9.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,788	7.6	7.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	953	4.0	3.9
6	Human immunodeficiency virus (HIV) disease (B20-B24)	375	1.6	1.5
7	Diabetes mellitus (E10-E14)	366	1.6	1.5
8	Cerebrovascular diseases (I60-I69)	298	1.3	1.2
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	285	1.2	1.2
10	Influenza and pneumonia (J10-J18)	206	0.9	0.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	186	0.8	0.8
12	Septicemia (A40-A41)	155	0.7	0.6
13	Chronic lower respiratory diseases (J40-J47)	140	0.6	0.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	128	0.5	0.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	113	0.5	0.5
...	All other causes (Residual)	4,161	17.7	16.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 35-44 years				
...	All causes	58,095	100.0	190.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,106	20.8	39.8
2	Malignant neoplasms (C00-C97)	10,723	18.5	35.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,894	15.3	29.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,563	9.6	18.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,068	3.6	6.8
6	Human immunodeficiency virus (HIV) disease (B20-B24)	1,703	2.9	5.6
7	Diabetes mellitus (E10-E14)	1,240	2.1	4.1
8	Cerebrovascular diseases (I60-I69)	1,239	2.1	4.1
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,147	2.0	3.8
10	Influenza and pneumonia (J10-J18)	637	1.1	2.1
11	Chronic lower respiratory diseases (J40-J47)	617	1.1	2.0
12	Septicemia (A40-A41)	513	0.9	1.7
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	430	0.7	1.4
14	Viral hepatitis (B15-B19)	425	0.7	1.4
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	340	0.6	1.1
...	All other causes (Residual)	10,450	18.0	34.3
White, non-Hispanic, both sexes, 45-54 years				
...	All causes	121,832	100.0	402.3
1	Malignant neoplasms (C00-C97)	36,168	29.7	119.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	25,982	21.3	85.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,721	9.6	38.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,774	4.7	19.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	5,310	4.4	17.5
6	Diabetes mellitus (E10-E14)	3,412	2.8	11.3
7	Cerebrovascular diseases (I60-I69)	3,294	2.7	10.9
8	Chronic lower respiratory diseases (J40-J47)	2,725	2.2	9.0
9	Viral hepatitis (B15-B19)	1,566	1.3	5.2
10	Influenza and pneumonia (J10-J18)	1,424	1.2	4.7
11	Human immunodeficiency virus (HIV) disease (B20-B24)	1,360	1.1	4.5
12	Septicemia (A40-A41)	1,295	1.1	4.3
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	977	0.8	3.2
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	947	0.8	3.1
15	Aortic aneurysm and dissection (I71)	548	0.4	1.8
...	All other causes (Residual)	19,329	15.9	63.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 55-64 years				
...	All causes	197,598	100.0	902.4
1	Malignant neoplasms (C00-C97)	75,733	38.3	345.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	48,091	24.3	219.6
3	Chronic lower respiratory diseases (J40-J47)	10,465	5.3	47.8
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,043	3.6	32.2
5	Diabetes mellitus (E10-E14)	6,873	3.5	31.4
6	Cerebrovascular diseases (I60-I69)	6,293	3.2	28.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	4,706	2.4	21.5
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,462	1.8	15.8
9	Septicemia (A40-A41)	2,471	1.3	11.3
10	Influenza and pneumonia (J10-J18)	2,210	1.1	10.1
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,197	1.1	10.0
12	Aortic aneurysm and dissection (I71)	1,178	0.6	5.4
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	995	0.5	4.5
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	918	0.5	4.2
15	Viral hepatitis (B15-B19)	642	0.3	2.9
...	All other causes (Residual)	24,321	12.3	111.1
White, non-Hispanic, both sexes, 65-74 years				
...	All causes	329,570	100.0	2,237.2
1	Malignant neoplasms (C00-C97)	116,198	35.3	788.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	83,479	25.3	566.7
3	Chronic lower respiratory diseases (J40-J47)	26,902	8.2	182.6
4	Cerebrovascular diseases (I60-I69)	15,259	4.6	103.6
5	Diabetes mellitus (E10-E14)	11,316	3.4	76.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,466	2.0	43.9
7	Influenza and pneumonia (J10-J18)	5,262	1.6	35.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,891	1.5	33.2
9	Septicemia (A40-A41)	4,315	1.3	29.3
10	Chronic liver disease and cirrhosis (K70,K73-K74)	4,256	1.3	28.9
11	Alzheimer's disease (G30)	3,269	1.0	22.2
12	Aortic aneurysm and dissection (I71)	2,844	0.9	19.3
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,110	0.6	14.3
14	Parkinson's disease (G20-G21)	2,061	0.6	14.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,970	0.6	13.4
...	All other causes (Residual)	38,972	11.8	264.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
White, non-Hispanic, both sexes, 75-84 years				
...	All causes	600,563	100.0	5,515.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	176,035	29.3	1,616.7
2	Malignant neoplasms (C00-C97)	144,012	24.0	1,322.6
3	Chronic lower respiratory diseases (J40-J47)	45,177	7.5	414.9
4	Cerebrovascular diseases (I60-I69)	44,277	7.4	406.6
5	Alzheimer's disease (G30)	18,937	3.2	173.9
6	Diabetes mellitus (E10-E14)	17,601	2.9	161.6
7	Influenza and pneumonia (J10-J18)	16,545	2.8	151.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,487	1.9	105.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11,179	1.9	102.7
10	Septicemia (A40-A41)	8,803	1.5	80.8
11	Parkinson's disease (G20-G21)	8,020	1.3	73.7
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	5,026	0.8	46.2
13	Pneumonitis due to solids and liquids (J69)	4,978	0.8	45.7
14	Aortic aneurysm and dissection (I71)	4,883	0.8	44.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,147	0.7	38.1
...	All other causes (Residual)	79,456	13.2	729.7
White, non-Hispanic, both sexes, 85 years and over				
...	All causes	608,843	100.0	14,944.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	220,489	36.2	5,412.2
2	Malignant neoplasms (C00-C97)	69,981	11.5	1,717.8
3	Cerebrovascular diseases (I60-I69)	57,078	9.4	1,401.1
4	Alzheimer's disease (G30)	34,501	5.7	846.9
5	Influenza and pneumonia (J10-J18)	27,886	4.6	684.5
6	Chronic lower respiratory diseases (J40-J47)	27,361	4.5	671.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,078	2.0	296.5
8	Diabetes mellitus (E10-E14)	11,968	2.0	293.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11,647	1.9	285.9
10	Septicemia (A40-A41)	7,840	1.3	192.4
11	Pneumonitis due to solids and liquids (J69)	7,374	1.2	181.0
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	7,372	1.2	181.0
13	Atherosclerosis (I70)	6,606	1.1	162.2
14	Parkinson's disease (G20-G21)	6,061	1.0	148.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3,437	0.6	84.4
...	All other causes (Residual)	97,164	16.0	2,385.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, all ages				
...	All causes	956,194	100.0	979.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	276,241	28.9	282.9
2	Malignant neoplasms (C00-C97)	236,150	24.7	241.8
3	Chronic lower respiratory diseases (J40-J47)	53,593	5.6	54.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	51,888	5.4	53.1
5	Cerebrovascular diseases (I60-I69)	48,549	5.1	49.7
6	Diabetes mellitus (E10-E14)	25,940	2.7	26.6
7	Influenza and pneumonia (J10-J18)	23,541	2.5	24.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	21,085	2.2	21.6
9	Alzheimer's disease (G30)	16,427	1.7	16.8
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15,323	1.6	15.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	13,254	1.4	13.6
12	Septicemia (A40-A41)	11,181	1.2	11.4
13	Parkinson's disease (G20-G21)	9,303	1.0	9.5
14	Aortic aneurysm and dissection (I71)	7,731	0.8	7.9
15	Pneumonitis due to solids and liquids (J69)	7,454	0.8	7.6
...	All other causes (Residual)	138,534	14.5	141.9
White, non-Hispanic, male, 1-4 years				
...	All causes	1,423	100.0	30.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	610	42.9	12.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	136	9.6	2.9
3	Malignant neoplasms (C00-C97)	123	8.6	2.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	61	4.3	1.3
5	Influenza and pneumonia (J10-J18)	44	3.1	0.9
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	34	2.4	0.7
7	Septicemia (A40-A41)	29	2.0	0.6
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	1.3	*
9	Certain conditions originating in the perinatal period (P00-P96)	17	1.2	*
10	Cerebrovascular diseases (I60-I69)	8	0.6	*
11	Meningitis (G00,G03)	7	0.5	*
11	Chronic lower respiratory diseases (J40-J47)	7	0.5	*
13	Meningococcal infection (A39)	6	0.4	*
14	Acute bronchitis and bronchiolitis (J20-J21)	5	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.4	*
...	All other causes (Residual)	307	21.6	6.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 5-14 years				
...	All causes	2,319	100.0	18.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	954	41.1	7.5
2	Malignant neoplasms (C00-C97)	348	15.0	2.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	122	5.3	1.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	117	5.0	0.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	88	3.8	0.7
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	72	3.1	0.6
7	Influenza and pneumonia (J10-J18)	39	1.7	0.3
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	24	1.0	0.2
9	Cerebrovascular diseases (I60-I69)	23	1.0	0.2
10	Septicemia (A40-A41)	22	0.9	0.2
11	Chronic lower respiratory diseases (J40-J47)	21	0.9	0.2
12	Diabetes mellitus (E10-E14)	12	0.5	*
13	Anemias (D50-D64)	9	0.4	*
14	Meningitis (G00,G03)	6	0.3	*
14	Pneumonitis due to solids and liquids (J69)	6	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.3	*
14	Certain conditions originating in the perinatal period (P00-P96)	6	0.3	*
...	All other causes (Residual)	444	19.1	3.5
White, non-Hispanic, male, 15-24 years				
...	All causes	14,022	100.0	105.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,698	54.9	58.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,412	17.2	18.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	665	4.7	5.0
4	Malignant neoplasms (C00-C97)	607	4.3	4.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	368	2.6	2.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	175	1.2	1.3
7	Influenza and pneumonia (J10-J18)	75	0.5	0.6
8	Cerebrovascular diseases (I60-I69)	72	0.5	0.5
9	Diabetes mellitus (E10-E14)	54	0.4	0.4
10	Septicemia (A40-A41)	48	0.3	0.4
10	Chronic lower respiratory diseases (J40-J47)	48	0.3	0.4
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	34	0.2	0.3
13	Legal intervention (Y35,Y89.0)	25	0.2	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	24	0.2	0.2
15	Aortic aneurysm and dissection (I71)	21	0.1	0.2
...	All other causes (Residual)	1,696	12.1	12.8

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 25-34 years				
...	All causes	16,097	100.0	129.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,138	38.1	49.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,099	19.3	25.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,212	7.5	9.8
4	Malignant neoplasms (C00-C97)	1,159	7.2	9.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	648	4.0	5.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	294	1.8	2.4
7	Diabetes mellitus (E10-E14)	212	1.3	1.7
8	Cerebrovascular diseases (I60-I69)	159	1.0	1.3
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	152	0.9	1.2
10	Influenza and pneumonia (J10-J18)	131	0.8	1.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	126	0.8	1.0
12	Septicemia (A40-A41)	74	0.5	0.6
13	Chronic lower respiratory diseases (J40-J47)	64	0.4	0.5
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	64	0.4	0.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	57	0.4	0.5
...	All other causes (Residual)	2,508	15.6	20.2
White, non-Hispanic, male, 35-44 years				
...	All causes	37,123	100.0	243.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,574	23.1	56.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,429	17.3	42.2
3	Malignant neoplasms (C00-C97)	4,862	13.1	31.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,296	11.6	28.2
5	Human immunodeficiency virus (HIV) disease (B20-B24)	1,429	3.8	9.4
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1,366	3.7	9.0
7	Diabetes mellitus (E10-E14)	794	2.1	5.2
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	747	2.0	4.9
9	Cerebrovascular diseases (I60-I69)	637	1.7	4.2
10	Influenza and pneumonia (J10-J18)	356	1.0	2.3
11	Viral hepatitis (B15-B19)	300	0.8	2.0
12	Chronic lower respiratory diseases (J40-J47)	275	0.7	1.8
13	Septicemia (A40-A41)	255	0.7	1.7
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	233	0.6	1.5
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	205	0.6	1.3
...	All other causes (Residual)	6,365	17.1	41.8

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 45-54 years				
...	All causes	77,106	100.0	513.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	19,504	25.3	129.8
2	Malignant neoplasms (C00-C97)	18,601	24.1	123.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,219	10.7	54.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,357	5.7	29.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	3,894	5.1	25.9
6	Diabetes mellitus (E10-E14)	2,122	2.8	14.1
7	Cerebrovascular diseases (I60-I69)	1,813	2.4	12.1
8	Chronic lower respiratory diseases (J40-J47)	1,426	1.8	9.5
9	Human immunodeficiency virus (HIV) disease (B20-B24)	1,172	1.5	7.8
10	Viral hepatitis (B15-B19)	1,162	1.5	7.7
11	Influenza and pneumonia (J10-J18)	844	1.1	5.6
12	Septicemia (A40-A41)	698	0.9	4.6
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	645	0.8	4.3
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	577	0.7	3.8
15	Aortic aneurysm and dissection (I71)	428	0.6	2.8
...	All other causes (Residual)	11,644	15.1	77.5
White, non-Hispanic, male, 55-64 years				
...	All causes	118,383	100.0	1,110.5
1	Malignant neoplasms (C00-C97)	41,026	34.7	384.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	33,871	28.6	317.7
3	Chronic lower respiratory diseases (J40-J47)	5,327	4.5	50.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,841	4.1	45.4
5	Diabetes mellitus (E10-E14)	4,041	3.4	37.9
6	Cerebrovascular diseases (I60-I69)	3,427	2.9	32.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	3,285	2.8	30.8
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,678	2.3	25.1
9	Septicemia (A40-A41)	1,280	1.1	12.0
10	Influenza and pneumonia (J10-J18)	1,219	1.0	11.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,216	1.0	11.4
12	Aortic aneurysm and dissection (I71)	897	0.8	8.4
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	544	0.5	5.1
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	527	0.4	4.9
15	Viral hepatitis (B15-B19)	445	0.4	4.2
...	All other causes (Residual)	13,759	11.6	129.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 65-74 years				
...	All causes	186,112	100.0	2,738.5
1	Malignant neoplasms (C00-C97)	64,019	34.4	942.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	52,148	28.0	767.3
3	Chronic lower respiratory diseases (J40-J47)	13,604	7.3	200.2
4	Cerebrovascular diseases (I60-I69)	7,947	4.3	116.9
5	Diabetes mellitus (E10-E14)	6,206	3.3	91.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,952	2.1	58.2
7	Influenza and pneumonia (J10-J18)	2,966	1.6	43.6
8	Chronic liver disease and cirrhosis (K70,K73-K74)	2,627	1.4	38.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,614	1.4	38.5
10	Septicemia (A40-A41)	2,207	1.2	32.5
11	Aortic aneurysm and dissection (I71)	1,926	1.0	28.3
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,769	1.0	26.0
13	Alzheimer's disease (G30)	1,443	0.8	21.2
14	Parkinson's disease (G20-G21)	1,343	0.7	19.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,142	0.6	16.8
...	All other causes (Residual)	20,199	10.9	297.2
White, non-Hispanic, male, 75-84 years				
...	All causes	293,774	100.0	6,692.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	89,985	30.6	2,049.9
2	Malignant neoplasms (C00-C97)	74,968	25.5	1,707.8
3	Chronic lower respiratory diseases (J40-J47)	21,553	7.3	491.0
4	Cerebrovascular diseases (I60-I69)	18,699	6.4	426.0
5	Diabetes mellitus (E10-E14)	8,417	2.9	191.7
6	Influenza and pneumonia (J10-J18)	8,141	2.8	185.5
7	Alzheimer's disease (G30)	6,796	2.3	154.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,957	2.0	135.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,750	2.0	131.0
10	Parkinson's disease (G20-G21)	4,784	1.6	109.0
11	Septicemia (A40-A41)	3,972	1.4	90.5
12	Aortic aneurysm and dissection (I71)	2,877	1.0	65.5
13	Pneumonitis due to solids and liquids (J69)	2,741	0.9	62.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,125	0.7	48.4
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,916	0.7	43.6
...	All other causes (Residual)	35,093	11.9	799.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 85 years and over				
...	All causes	202,148	100.0	16,234.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	72,482	35.9	5,821.0
2	Malignant neoplasms (C00-C97)	30,404	15.0	2,441.7
3	Cerebrovascular diseases (I60-I69)	15,741	7.8	1,264.2
4	Chronic lower respiratory diseases (J40-J47)	11,251	5.6	903.6
5	Influenza and pneumonia (J10-J18)	9,639	4.8	774.1
6	Alzheimer's disease (G30)	7,946	3.9	638.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,810	2.4	386.3
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,643	2.3	372.9
9	Diabetes mellitus (E10-E14)	4,076	2.0	327.3
10	Pneumonitis due to solids and liquids (J69)	3,194	1.6	256.5
11	Parkinson's disease (G20-G21)	2,962	1.5	237.9
12	Septicemia (A40-A41)	2,550	1.3	204.8
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,811	0.9	145.4
14	Atherosclerosis (I70)	1,808	0.9	145.2
15	Aortic aneurysm and dissection (I71)	1,385	0.7	111.2
...	All other causes (Residual)	27,446	13.6	2,204.2
White, non-Hispanic, female, all ages				
...	All causes	1,023,271	100.0	1,007.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	289,567	28.3	285.1
2	Malignant neoplasms (C00-C97)	220,977	21.6	217.6
3	Cerebrovascular diseases (I60-I69)	79,420	7.8	78.2
4	Chronic lower respiratory diseases (J40-J47)	59,948	5.9	59.0
5	Alzheimer's disease (G30)	40,841	4.0	40.2
6	Influenza and pneumonia (J10-J18)	31,076	3.0	30.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	30,927	3.0	30.5
8	Diabetes mellitus (E10-E14)	26,954	2.6	26.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16,174	1.6	15.9
10	Septicemia (A40-A41)	14,483	1.4	14.3
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	10,205	1.0	10.0
12	Pneumonitis due to solids and liquids (J69)	7,391	0.7	7.3
13	Chronic liver disease and cirrhosis (K70,K73-K74)	7,348	0.7	7.2
14	Atherosclerosis (I70)	7,294	0.7	7.2
15	Parkinson's disease (G20-G21)	7,196	0.7	7.1
...	All other causes (Residual)	173,470	17.0	170.8

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 1-4 years				
...	All causes	1,116	100.0	24.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	375	33.6	8.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	136	12.2	3.0
3	Malignant neoplasms (C00-C97)	98	8.8	2.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	62	5.6	1.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	49	4.4	1.1
6	Influenza and pneumonia (J10-J18)	36	3.2	0.8
7	Cerebrovascular diseases (I60-I69)	15	1.3	*
8	Septicemia (A40-A41)	14	1.3	*
8	Certain conditions originating in the perinatal period (P00-P96)	14	1.3	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	1.2	*
11	Chronic lower respiratory diseases (J40-J47)	8	0.7	*
12	Meningitis (G00,G03)	7	0.6	*
13	Acute bronchitis and bronchiolitis (J20-J21)	5	0.4	*
14	Meningococcal infection (A39)	4	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.4	*
...	All other causes (Residual)	272	24.4	6.1
White, non-Hispanic, female, 5-14 years				
...	All causes	1,565	100.0	12.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	581	37.1	4.8
2	Malignant neoplasms (C00-C97)	289	18.5	2.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	105	6.7	0.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	58	3.7	0.5
5	Influenza and pneumonia (J10-J18)	47	3.0	0.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	37	2.4	0.3
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	29	1.9	0.2
8	Septicemia (A40-A41)	25	1.6	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	1.2	*
10	Chronic lower respiratory diseases (J40-J47)	18	1.2	*
11	Cerebrovascular diseases (I60-I69)	14	0.9	*
12	Anemias (D50-D64)	8	0.5	*
13	Diabetes mellitus (E10-E14)	7	0.4	*
14	Pneumonitis due to solids and liquids (J69)	5	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.3	*
...	All other causes (Residual)	318	20.3	2.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 15-24 years				
...	All causes	5,618	100.0	44.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,940	52.3	23.2
2	Malignant neoplasms (C00-C97)	432	7.7	3.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	420	7.5	3.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	242	4.3	1.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	237	4.2	1.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	110	2.0	0.9
7	Influenza and pneumonia (J10-J18)	62	1.1	0.5
8	Cerebrovascular diseases (I60-I69)	52	0.9	0.4
9	Pregnancy, childbirth and the puerperium (O00-099)	41	0.7	0.3
10	Septicemia (A40-A41)	38	0.7	0.3
11	Diabetes mellitus (E10-E14)	36	0.6	0.3
12	Chronic lower respiratory diseases (J40-J47)	34	0.6	0.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	24	0.4	0.2
14	Meningococcal infection (A39)	15	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	0.2	*
...	All other causes (Residual)	921	16.4	7.3
White, non-Hispanic, female, 25-34 years				
...	All causes	7,460	100.0	61.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,100	28.2	17.2
2	Malignant neoplasms (C00-C97)	1,195	16.0	9.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	712	9.5	5.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	576	7.7	4.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	305	4.1	2.5
6	Diabetes mellitus (E10-E14)	154	2.1	1.3
7	Cerebrovascular diseases (I60-I69)	139	1.9	1.1
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	133	1.8	1.1
9	Pregnancy, childbirth and the puerperium (O00-099)	87	1.2	0.7
10	Septicemia (A40-A41)	81	1.1	0.7
10	Human immunodeficiency virus (HIV) disease (B20-B24)	81	1.1	0.7
12	Chronic lower respiratory diseases (J40-J47)	76	1.0	0.6
13	Influenza and pneumonia (J10-J18)	75	1.0	0.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	64	0.9	0.5
15	Chronic liver disease and cirrhosis (K70,K73-K74)	60	0.8	0.5
...	All other causes (Residual)	1,622	21.7	13.3

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	White, non-Hispanic, female, 35-44 years			
...	All causes	20,972	100.0	138.0
1	Malignant neoplasms (C00-C97)	5,861	27.9	38.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,532	16.8	23.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,465	11.8	16.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,267	6.0	8.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	702	3.3	4.6
6	Cerebrovascular diseases (I60-I69)	602	2.9	4.0
7	Diabetes mellitus (E10-E14)	446	2.1	2.9
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	400	1.9	2.6
9	Chronic lower respiratory diseases (J40-J47)	342	1.6	2.2
10	Influenza and pneumonia (J10-J18)	281	1.3	1.8
11	Human immunodeficiency virus (HIV) disease (B20-B24)	274	1.3	1.8
12	Septicemia (A40-A41)	258	1.2	1.7
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	197	0.9	1.3
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	135	0.6	0.9
15	Viral hepatitis (B15-B19)	125	0.6	0.8
...	All other causes (Residual)	4,085	19.5	26.9
	White, non-Hispanic, female, 45-54 years			
...	All causes	44,726	100.0	293.1
1	Malignant neoplasms (C00-C97)	17,567	39.3	115.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,478	14.5	42.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,502	7.8	22.9
4	Cerebrovascular diseases (I60-I69)	1,481	3.3	9.7
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,417	3.2	9.3
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1,416	3.2	9.3
7	Chronic lower respiratory diseases (J40-J47)	1,299	2.9	8.5
8	Diabetes mellitus (E10-E14)	1,290	2.9	8.5
9	Septicemia (A40-A41)	597	1.3	3.9
10	Influenza and pneumonia (J10-J18)	580	1.3	3.8
11	Viral hepatitis (B15-B19)	404	0.9	2.6
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	400	0.9	2.6
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	302	0.7	2.0
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	259	0.6	1.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	227	0.5	1.5
...	All other causes (Residual)	7,507	16.8	49.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 55-64 years				
...	All causes	79,215	100.0	705.0
1	Malignant neoplasms (C00-C97)	34,707	43.8	308.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,220	18.0	126.6
3	Chronic lower respiratory diseases (J40-J47)	5,138	6.5	45.7
4	Cerebrovascular diseases (I60-I69)	2,866	3.6	25.5
5	Diabetes mellitus (E10-E14)	2,832	3.6	25.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,202	2.8	19.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,421	1.8	12.6
8	Septicemia (A40-A41)	1,191	1.5	10.6
9	Influenza and pneumonia (J10-J18)	991	1.3	8.8
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	981	1.2	8.7
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	784	1.0	7.0
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	451	0.6	4.0
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	391	0.5	3.5
14	Aortic aneurysm and dissection (I71)	281	0.4	2.5
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	281	0.4	2.5
...	All other causes (Residual)	10,478	13.2	93.3
White, non-Hispanic, female, 65-74 years				
...	All causes	143,458	100.0	1,807.9
1	Malignant neoplasms (C00-C97)	52,179	36.4	657.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	31,331	21.8	394.8
3	Chronic lower respiratory diseases (J40-J47)	13,298	9.3	167.6
4	Cerebrovascular diseases (I60-I69)	7,312	5.1	92.1
5	Diabetes mellitus (E10-E14)	5,110	3.6	64.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,514	1.8	31.7
7	Influenza and pneumonia (J10-J18)	2,296	1.6	28.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,277	1.6	28.7
9	Septicemia (A40-A41)	2,108	1.5	26.6
10	Alzheimer's disease (G30)	1,826	1.3	23.0
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1,629	1.1	20.5
12	Aortic aneurysm and dissection (I71)	918	0.6	11.6
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	886	0.6	11.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	828	0.6	10.4
15	Parkinson's disease (G20-G21)	718	0.5	9.0
...	All other causes (Residual)	18,228	12.7	229.7

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	White, non-Hispanic, female, 75-84 years			
...	All causes	306,789	100.0	4,720.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	86,050	28.0	1,324.0
2	Malignant neoplasms (C00-C97)	69,044	22.5	1,062.4
3	Cerebrovascular diseases (I60-I69)	25,578	8.3	393.6
4	Chronic lower respiratory diseases (J40-J47)	23,624	7.7	363.5
5	Alzheimer's disease (G30)	12,141	4.0	186.8
6	Diabetes mellitus (E10-E14)	9,184	3.0	141.3
7	Influenza and pneumonia (J10-J18)	8,404	2.7	129.3
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,530	1.8	85.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,429	1.8	83.5
10	Septicemia (A40-A41)	4,831	1.6	74.3
11	Parkinson's disease (G20-G21)	3,236	1.1	49.8
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	3,110	1.0	47.9
13	Pneumonitis due to solids and liquids (J69)	2,237	0.7	34.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,022	0.7	31.1
15	Aortic aneurysm and dissection (I71)	2,006	0.7	30.9
...	All other causes (Residual)	44,363	14.5	682.6
	White, non-Hispanic, female, 85 years and over			
...	All causes	406,695	100.0	14,377.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	148,007	36.4	5,232.2
2	Cerebrovascular diseases (I60-I69)	41,337	10.2	1,461.3
3	Malignant neoplasms (C00-C97)	39,577	9.7	1,399.1
4	Alzheimer's disease (G30)	26,555	6.5	938.8
5	Influenza and pneumonia (J10-J18)	18,247	4.5	645.1
6	Chronic lower respiratory diseases (J40-J47)	16,110	4.0	569.5
7	Diabetes mellitus (E10-E14)	7,892	1.9	279.0
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,435	1.8	262.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,837	1.7	241.7
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	5,561	1.4	196.6
11	Septicemia (A40-A41)	5,290	1.3	187.0
12	Atherosclerosis (I70)	4,798	1.2	169.6
13	Pneumonitis due to solids and liquids (J69)	4,180	1.0	147.8
14	Parkinson's disease (G20-G21)	3,099	0.8	109.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,094	0.5	74.0
...	All other causes (Residual)	69,676	17.1	2,463.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, all ages				
...	All causes	287,968	100.0	788.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	76,452	26.5	209.4
2	Malignant neoplasms (C00-C97)	62,054	21.5	170.0
3	Cerebrovascular diseases (I60-I69)	18,649	6.5	51.1
4	Diabetes mellitus (E10-E14)	12,787	4.4	35.0
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,154	4.2	33.3
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8,276	2.9	22.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,804	2.7	21.4
8	Chronic lower respiratory diseases (J40-J47)	7,632	2.7	20.9
9	Human immunodeficiency virus (HIV) disease (B20-B24)	7,351	2.6	20.1
10	Septicemia (A40-A41)	6,141	2.1	16.8
11	Influenza and pneumonia (J10-J18)	5,798	2.0	15.9
12	Certain conditions originating in the perinatal period (P00-P96)	4,716	1.6	12.9
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	4,445	1.5	12.2
14	Alzheimer's disease (G30)	3,693	1.3	10.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,582	0.9	7.1
...	All other causes (Residual)	47,434	16.5	129.9
Black, non-Hispanic, both sexes, 1-4 years				
...	All causes	1,169	100.0	48.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	333	28.5	13.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	155	13.3	6.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	118	10.1	4.8
4	Malignant neoplasms (C00-C97)	55	4.7	2.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	55	4.7	2.3
6	Certain conditions originating in the perinatal period (P00-P96)	29	2.5	1.2
7	Influenza and pneumonia (J10-J18)	27	2.3	1.1
8	Septicemia (A40-A41)	24	2.1	1.0
8	Chronic lower respiratory diseases (J40-J47)	24	2.1	1.0
10	Cerebrovascular diseases (I60-I69)	14	1.2	*
11	Anemias (D50-D64)	12	1.0	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.8	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	9	0.8	*
14	Acute bronchitis and bronchiolitis (J20-J21)	7	0.6	*
15	Meningococcal infection (A39)	6	0.5	*
...	All other causes (Residual)	292	25.0	12.0

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 5-14 years				
...	All causes	1,530	100.0	23.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	517	33.8	8.0
2	Malignant neoplasms (C00-C97)	173	11.3	2.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	147	9.6	2.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	76	5.0	1.2
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	73	4.8	1.1
6	Chronic lower respiratory diseases (J40-J47)	62	4.1	1.0
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	43	2.8	0.7
8	Human immunodeficiency virus (HIV) disease (B20-B24)	31	2.0	0.5
9	Influenza and pneumonia (J10-J18)	28	1.8	0.4
10	Anemias (D50-D64)	25	1.6	0.4
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	1.1	*
12	Cerebrovascular diseases (I60-I69)	16	1.0	*
13	Septicemia (A40-A41)	15	1.0	*
14	Diabetes mellitus (E10-E14)	7	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.5	*
...	All other causes (Residual)	293	19.2	4.6
Black, non-Hispanic, both sexes, 15-24 years				
...	All causes	7,021	100.0	116.6
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,955	42.1	49.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,590	22.6	26.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	438	6.2	7.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	346	4.9	5.7
5	Malignant neoplasms (C00-C97)	245	3.5	4.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	117	1.7	1.9
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	92	1.3	1.5
8	Chronic lower respiratory diseases (J40-J47)	81	1.2	1.3
9	Anemias (D50-D64)	78	1.1	1.3
10	Cerebrovascular diseases (I60-I69)	58	0.8	1.0
11	Diabetes mellitus (E10-E14)	45	0.6	0.7
12	Influenza and pneumonia (J10-J18)	42	0.6	0.7
12	Pregnancy, childbirth and the puerperium (O00-099)	42	0.6	0.7
14	Septicemia (A40-A41)	41	0.6	0.7
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	34	0.5	0.6
...	All other causes (Residual)	817	11.6	13.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Black, non-Hispanic, both sexes, 25-34 years			
...	All causes	9,795	100.0	188.9
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,443	24.9	47.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,602	16.4	30.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	974	9.9	18.8
4	Human immunodeficiency virus (HIV) disease (B20-B24)	938	9.6	18.1
5	Malignant neoplasms (C00-C97)	656	6.7	12.6
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	498	5.1	9.6
7	Diabetes mellitus (E10-E14)	208	2.1	4.0
8	Cerebrovascular diseases (I60-I69)	167	1.7	3.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	118	1.2	2.3
10	Anemias (D50-D64)	114	1.2	2.2
11	Chronic lower respiratory diseases (J40-J47)	104	1.1	2.0
12	Septicemia (A40-A41)	102	1.0	2.0
13	Influenza and pneumonia (J10-J18)	101	1.0	1.9
14	Pregnancy, childbirth and the puerperium (O00-099)	82	0.8	1.6
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	75	0.8	1.4
...	All other causes (Residual)	1,613	16.5	31.1
	Black, non-Hispanic, both sexes, 35-44 years			
...	All causes	19,561	100.0	354.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,392	17.3	61.4
2	Malignant neoplasms (C00-C97)	2,787	14.2	50.5
3	Human immunodeficiency virus (HIV) disease (B20-B24)	2,786	14.2	50.5
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,195	11.2	39.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,335	6.8	24.2
6	Cerebrovascular diseases (I60-I69)	786	4.0	14.2
7	Diabetes mellitus (E10-E14)	547	2.8	9.9
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	409	2.1	7.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	355	1.8	6.4
10	Chronic liver disease and cirrhosis (K70,K73-K74)	323	1.7	5.8
11	Septicemia (A40-A41)	283	1.4	5.1
12	Influenza and pneumonia (J10-J18)	251	1.3	4.5
13	Chronic lower respiratory diseases (J40-J47)	245	1.3	4.4
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	174	0.9	3.2
15	Anemias (D50-D64)	133	0.7	2.4
...	All other causes (Residual)	3,560	18.2	64.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 45-54 years				
...	All causes	36,584	100.0	788.3
1	Malignant neoplasms (C00-C97)	8,802	24.1	189.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,625	23.6	185.8
3	Human immunodeficiency virus (HIV) disease (B20-B24)	2,382	6.5	51.3
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,289	6.3	49.3
5	Cerebrovascular diseases (I60-I69)	1,949	5.3	42.0
6	Diabetes mellitus (E10-E14)	1,431	3.9	30.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	936	2.6	20.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	780	2.1	16.8
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	731	2.0	15.8
10	Septicemia (A40-A41)	650	1.8	14.0
11	Chronic lower respiratory diseases (J40-J47)	597	1.6	12.9
12	Influenza and pneumonia (J10-J18)	487	1.3	10.5
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	441	1.2	9.5
14	Viral hepatitis (B15-B19)	352	1.0	7.6
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	266	0.7	5.7
...	All other causes (Residual)	5,866	16.0	126.4
Black, non-Hispanic, both sexes, 55-64 years				
...	All causes	42,350	100.0	1,578.4
1	Malignant neoplasms (C00-C97)	12,903	30.5	480.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,664	27.5	434.7
3	Cerebrovascular diseases (I60-I69)	2,449	5.8	91.3
4	Diabetes mellitus (E10-E14)	2,347	5.5	87.5
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,172	2.8	43.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,152	2.7	42.9
7	Chronic lower respiratory diseases (J40-J47)	1,109	2.6	41.3
8	Septicemia (A40-A41)	881	2.1	32.8
9	Human immunodeficiency virus (HIV) disease (B20-B24)	802	1.9	29.9
10	Chronic liver disease and cirrhosis (K70,K73-K74)	700	1.7	26.1
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	664	1.6	24.7
12	Influenza and pneumonia (J10-J18)	565	1.3	21.1
13	Viral hepatitis (B15-B19)	221	0.5	8.2
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	218	0.5	8.1
15	Aortic aneurysm and dissection (I71)	196	0.5	7.3
...	All other causes (Residual)	5,307	12.5	197.8

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 65-74 years				
...	All causes	52,583	100.0	3,158.7
1	Malignant neoplasms (C00-C97)	15,666	29.8	941.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,409	29.3	925.6
3	Cerebrovascular diseases (I60-I69)	3,411	6.5	204.9
4	Diabetes mellitus (E10-E14)	3,145	6.0	188.9
5	Chronic lower respiratory diseases (J40-J47)	1,920	3.7	115.3
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,758	3.3	105.6
7	Septicemia (A40-A41)	1,232	2.3	74.0
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	918	1.7	55.1
9	Influenza and pneumonia (J10-J18)	915	1.7	55.0
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	850	1.6	51.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	401	0.8	24.1
12	Alzheimer's disease (G30)	326	0.6	19.6
13	Aortic aneurysm and dissection (I71)	242	0.5	14.5
14	Human immunodeficiency virus (HIV) disease (B20-B24)	239	0.5	14.4
15	Pneumonitis due to solids and liquids (J69)	231	0.4	13.9
...	All other causes (Residual)	5,920	11.3	355.6
Black, non-Hispanic, both sexes, 75-84 years				
...	All causes	61,770	100.0	6,433.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	19,046	30.8	1,983.5
2	Malignant neoplasms (C00-C97)	14,482	23.4	1,508.2
3	Cerebrovascular diseases (I60-I69)	5,176	8.4	539.1
4	Diabetes mellitus (E10-E14)	3,265	5.3	340.0
5	Chronic lower respiratory diseases (J40-J47)	2,290	3.7	238.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,015	3.3	209.9
7	Septicemia (A40-A41)	1,536	2.5	160.0
8	Influenza and pneumonia (J10-J18)	1,514	2.5	157.7
9	Alzheimer's disease (G30)	1,347	2.2	140.3
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,162	1.9	121.0
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	798	1.3	83.1
12	Pneumonitis due to solids and liquids (J69)	533	0.9	55.5
13	Atherosclerosis (I70)	296	0.5	30.8
14	Aortic aneurysm and dissection (I71)	292	0.5	30.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	284	0.5	29.6
...	All other causes (Residual)	7,734	12.5	805.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 85 years and over				
...	All causes	47,426	100.0	14,107.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	16,734	35.3	4,977.8
2	Malignant neoplasms (C00-C97)	6,278	13.2	1,867.5
3	Cerebrovascular diseases (I60-I69)	4,592	9.7	1,366.0
4	Alzheimer's disease (G30)	1,987	4.2	591.1
5	Diabetes mellitus (E10-E14)	1,789	3.8	532.2
6	Influenza and pneumonia (J10-J18)	1,776	3.7	528.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,501	3.2	446.5
8	Septicemia (A40-A41)	1,257	2.7	373.9
9	Chronic lower respiratory diseases (J40-J47)	1,193	2.5	354.9
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,023	2.2	304.3
11	Pneumonitis due to solids and liquids (J69)	557	1.2	165.7
12	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	544	1.1	161.8
13	Atherosclerosis (I70)	413	0.9	122.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	198	0.4	58.9
15	Nutritional deficiencies (E40-E64)	188	0.4	55.9
...	All other causes (Residual)	7,396	15.6	2,200.1
Black, non-Hispanic, male, all ages				
...	All causes	146,136	100.0	840.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	36,962	25.3	212.6
2	Malignant neoplasms (C00-C97)	32,131	22.0	184.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,237	5.6	47.4
4	Cerebrovascular diseases (I60-I69)	7,798	5.3	44.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6,980	4.8	40.1
6	Diabetes mellitus (E10-E14)	5,380	3.7	30.9
7	Human immunodeficiency virus (HIV) disease (B20-B24)	4,947	3.4	28.5
8	Chronic lower respiratory diseases (J40-J47)	4,202	2.9	24.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,614	2.5	20.8
10	Influenza and pneumonia (J10-J18)	2,861	2.0	16.5
11	Septicemia (A40-A41)	2,748	1.9	15.8
12	Certain conditions originating in the perinatal period (P00-P96)	2,687	1.8	15.5
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,875	1.3	10.8
14	Chronic liver disease and cirrhosis (K70,K73-K74)	1,696	1.2	9.8
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,566	1.1	9.0
...	All other causes (Residual)	22,452	15.4	129.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 1-4 years				
...	All causes	681	100.0	55.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	197	28.9	15.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	96	14.1	7.8
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	62	9.1	5.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	31	4.6	2.5
5	Malignant neoplasms (C00-C97)	30	4.4	2.4
6	Chronic lower respiratory diseases (J40-J47)	19	2.8	*
7	Certain conditions originating in the perinatal period (P00-P96)	16	2.3	*
8	Septicemia (A40-A41)	14	2.1	*
9	Influenza and pneumonia (J10-J18)	13	1.9	*
10	Cerebrovascular diseases (I60-I69)	9	1.3	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	1.2	*
12	Anemias (D50-D64)	7	1.0	*
13	Acute bronchitis and bronchiolitis (J20-J21)	6	0.9	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.7	*
15	Meningococcal infection (A39)	3	0.4	*
...	All other causes (Residual)	165	24.2	13.4
Black, non-Hispanic, male, 5-14 years				
...	All causes	911	100.0	27.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	330	36.2	10.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	98	10.8	3.0
3	Malignant neoplasms (C00-C97)	96	10.5	2.9
4	Chronic lower respiratory diseases (J40-J47)	43	4.7	1.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	42	4.6	1.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	38	4.2	1.2
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	34	3.7	1.0
8	Human immunodeficiency virus (HIV) disease (B20-B24)	17	1.9	*
9	Influenza and pneumonia (J10-J18)	15	1.6	*
10	Anemias (D50-D64)	12	1.3	*
11	Cerebrovascular diseases (I60-I69)	9	1.0	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.9	*
13	Septicemia (A40-A41)	6	0.7	*
14	Diseases of appendix (K35-K38)	4	0.4	*
15	Diabetes mellitus (E10-E14)	3	0.3	*
...	All other causes (Residual)	156	17.1	4.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 15-24 years				
...	All causes	5,343	100.0	176.5
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,642	49.4	87.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,205	22.6	39.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	377	7.1	12.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	225	4.2	7.4
5	Malignant neoplasms (C00-C97)	141	2.6	4.7
6	Human immunodeficiency virus (HIV) disease (B20-B24)	57	1.1	1.9
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	55	1.0	1.8
8	Chronic lower respiratory diseases (J40-J47)	48	0.9	1.6
9	Anemias (D50-D64)	39	0.7	1.3
10	Legal intervention (Y35,Y89.0)	31	0.6	1.0
11	Diabetes mellitus (E10-E14)	26	0.5	0.9
12	Cerebrovascular diseases (I60-I69)	25	0.5	0.8
13	Influenza and pneumonia (J10-J18)	24	0.4	0.8
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20	0.4	0.7
15	Septicemia (A40-A41)	19	0.4	*
...	All other causes (Residual)	409	7.7	13.5
Black, non-Hispanic, male, 25-34 years				
...	All causes	6,592	100.0	266.6
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,128	32.3	86.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,197	18.2	48.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	572	8.7	23.1
4	Human immunodeficiency virus (HIV) disease (B20-B24)	537	8.1	21.7
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	424	6.4	17.1
6	Malignant neoplasms (C00-C97)	263	4.0	10.6
7	Diabetes mellitus (E10-E14)	122	1.9	4.9
8	Cerebrovascular diseases (I60-I69)	86	1.3	3.5
9	Chronic lower respiratory diseases (J40-J47)	72	1.1	2.9
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	64	1.0	2.6
11	Anemias (D50-D64)	58	0.9	2.3
11	Influenza and pneumonia (J10-J18)	58	0.9	2.3
13	Septicemia (A40-A41)	56	0.8	2.3
14	Legal intervention (Y35,Y89.0)	51	0.8	2.1
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	38	0.6	1.5
...	All other causes (Residual)	866	13.1	35.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 35-44 years				
...	All causes	11,405	100.0	440.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,066	18.1	79.7
2	Human immunodeficiency virus (HIV) disease (B20-B24)	1,814	15.9	70.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,539	13.5	59.4
4	Malignant neoplasms (C00-C97)	1,112	9.8	42.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,078	9.5	41.6
6	Cerebrovascular diseases (I60-I69)	385	3.4	14.9
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	324	2.8	12.5
8	Diabetes mellitus (E10-E14)	309	2.7	11.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	191	1.7	7.4
10	Chronic liver disease and cirrhosis (K70,K73-K74)	188	1.6	7.3
11	Influenza and pneumonia (J10-J18)	142	1.2	5.5
12	Septicemia (A40-A41)	134	1.2	5.2
13	Chronic lower respiratory diseases (J40-J47)	106	0.9	4.1
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	92	0.8	3.5
15	Anemias (D50-D64)	68	0.6	2.6
...	All other causes (Residual)	1,857	16.3	71.6
Black, non-Hispanic, male, 45-54 years				
...	All causes	21,708	100.0	1,012.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,443	25.1	253.9
2	Malignant neoplasms (C00-C97)	4,533	20.9	211.5
3	Human immunodeficiency virus (HIV) disease (B20-B24)	1,693	7.8	79.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,625	7.5	75.8
5	Cerebrovascular diseases (I60-I69)	1,031	4.7	48.1
6	Diabetes mellitus (E10-E14)	741	3.4	34.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	664	3.1	31.0
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	576	2.7	26.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	467	2.2	21.8
10	Septicemia (A40-A41)	338	1.6	15.8
11	Chronic lower respiratory diseases (J40-J47)	293	1.3	13.7
12	Influenza and pneumonia (J10-J18)	285	1.3	13.3
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	278	1.3	13.0
14	Viral hepatitis (B15-B19)	250	1.2	11.7
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	199	0.9	9.3
...	All other causes (Residual)	3,292	15.2	153.6

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 55-64 years				
...	All causes	24,468	100.0	2,047.7
1	Malignant neoplasms (C00-C97)	7,135	29.2	597.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,040	28.8	589.2
3	Cerebrovascular diseases (I60-I69)	1,363	5.6	114.1
4	Diabetes mellitus (E10-E14)	1,169	4.8	97.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	824	3.4	69.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	627	2.6	52.5
7	Chronic lower respiratory diseases (J40-J47)	615	2.5	51.5
8	Human immunodeficiency virus (HIV) disease (B20-B24)	614	2.5	51.4
9	Chronic liver disease and cirrhosis (K70,K73-K74)	492	2.0	41.2
10	Septicemia (A40-A41)	462	1.9	38.7
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	359	1.5	30.0
12	Influenza and pneumonia (J10-J18)	356	1.5	29.8
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	178	0.7	14.9
14	Viral hepatitis (B15-B19)	145	0.6	12.1
15	Aortic aneurysm and dissection (I71)	132	0.5	11.0
...	All other causes (Residual)	2,957	12.1	247.5
Black, non-Hispanic, male, 65-74 years				
...	All causes	27,957	100.0	4,041.1
1	Malignant neoplasms (C00-C97)	8,593	30.7	1,242.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,377	30.0	1,210.9
3	Cerebrovascular diseases (I60-I69)	1,673	6.0	241.8
4	Diabetes mellitus (E10-E14)	1,392	5.0	201.2
5	Chronic lower respiratory diseases (J40-J47)	1,093	3.9	158.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	853	3.1	123.3
7	Septicemia (A40-A41)	601	2.1	86.9
8	Influenza and pneumonia (J10-J18)	547	2.0	79.1
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	518	1.9	74.9
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	424	1.5	61.3
11	Chronic liver disease and cirrhosis (K70,K73-K74)	233	0.8	33.7
12	Human immunodeficiency virus (HIV) disease (B20-B24)	175	0.6	25.3
13	Alzheimer's disease (G30)	142	0.5	20.5
14	Pneumonitis due to solids and liquids (J69)	137	0.5	19.8
15	Aortic aneurysm and dissection (I71)	124	0.4	17.9
...	All other causes (Residual)	3,075	11.0	444.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 75-84 years				
...	All causes	28,297	100.0	8,165.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,488	30.0	2,449.4
2	Malignant neoplasms (C00-C97)	7,540	26.6	2,175.8
3	Cerebrovascular diseases (I60-I69)	2,071	7.3	597.6
4	Chronic lower respiratory diseases (J40-J47)	1,340	4.7	386.7
5	Diabetes mellitus (E10-E14)	1,188	4.2	342.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	862	3.0	248.7
7	Influenza and pneumonia (J10-J18)	765	2.7	220.8
8	Septicemia (A40-A41)	663	2.3	191.3
9	Alzheimer's disease (G30)	481	1.7	138.8
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	429	1.5	123.8
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	419	1.5	120.9
12	Pneumonitis due to solids and liquids (J69)	276	1.0	79.6
13	Parkinson's disease (G20-G21)	158	0.6	45.6
14	Atherosclerosis (I70)	140	0.5	40.4
15	Aortic aneurysm and dissection (I71)	138	0.5	39.8
...	All other causes (Residual)	3,339	11.8	963.5
Black, non-Hispanic, male, 85 years and over				
...	All causes	14,156	100.0	15,082.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,597	32.5	4,897.8
2	Malignant neoplasms (C00-C97)	2,684	19.0	2,859.6
3	Cerebrovascular diseases (I60-I69)	1,127	8.0	1,200.7
4	Influenza and pneumonia (J10-J18)	603	4.3	642.5
5	Chronic lower respiratory diseases (J40-J47)	571	4.0	608.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	494	3.5	526.3
7	Alzheimer's disease (G30)	437	3.1	465.6
8	Diabetes mellitus (E10-E14)	429	3.0	457.1
9	Septicemia (A40-A41)	384	2.7	409.1
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	263	1.9	280.2
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	214	1.5	228.0
12	Pneumonitis due to solids and liquids (J69)	202	1.4	215.2
13	Atherosclerosis (I70)	99	0.7	105.5
14	Parkinson's disease (G20-G21)	93	0.7	99.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	64	0.5	68.2
...	All other causes (Residual)	1,895	13.4	2,019.0

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, all ages				
...	All causes	141,832	100.0	741.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	39,490	27.8	206.5
2	Malignant neoplasms (C00-C97)	29,923	21.1	156.5
3	Cerebrovascular diseases (I60-I69)	10,851	7.7	56.7
4	Diabetes mellitus (E10-E14)	7,407	5.2	38.7
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,190	3.0	21.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,917	2.8	20.5
7	Chronic lower respiratory diseases (J40-J47)	3,430	2.4	17.9
8	Septicemia (A40-A41)	3,393	2.4	17.7
9	Influenza and pneumonia (J10-J18)	2,937	2.1	15.4
10	Alzheimer's disease (G30)	2,614	1.8	13.7
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,570	1.8	13.4
12	Human immunodeficiency virus (HIV) disease (B20-B24)	2,404	1.7	12.6
13	Certain conditions originating in the perinatal period (P00-P96)	2,029	1.4	10.6
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,296	0.9	6.8
15	Chronic liver disease and cirrhosis (K70,K73-K74)	886	0.6	4.6
...	All other causes (Residual)	24,495	17.3	128.1
Black, non-Hispanic, female, 1-4 years				
...	All causes	488	100.0	40.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	136	27.9	11.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	59	12.1	4.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	56	11.5	4.7
4	Malignant neoplasms (C00-C97)	25	5.1	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	24	4.9	2.0
6	Influenza and pneumonia (J10-J18)	14	2.9	*
7	Certain conditions originating in the perinatal period (P00-P96)	13	2.7	*
8	Septicemia (A40-A41)	10	2.0	*
9	Anemias (D50-D64)	5	1.0	*
9	Cerebrovascular diseases (I60-I69)	5	1.0	*
9	Chronic lower respiratory diseases (J40-J47)	5	1.0	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.8	*
13	Meningococcal infection (A39)	3	0.6	*
13	Meningitis (G00,G03)	3	0.6	*
15	Diseases of appendix (K35-K38)	2	0.4	*
15	Hernia (K40-K46)	2	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
...	All other causes (Residual)	120	24.6	10.0

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 5-14 years				
...	All causes	619	100.0	19.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	187	30.2	5.9
2	Malignant neoplasms (C00-C97)	77	12.4	2.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	49	7.9	1.5
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	35	5.7	1.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	34	5.5	1.1
6	Chronic lower respiratory diseases (J40-J47)	19	3.1	*
7	Human immunodeficiency virus (HIV) disease (B20-B24)	14	2.3	*
8	Anemias (D50-D64)	13	2.1	*
8	Influenza and pneumonia (J10-J18)	13	2.1	*
10	Septicemia (A40-A41)	9	1.5	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	1.5	*
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	9	1.5	*
13	Cerebrovascular diseases (I60-I69)	7	1.1	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.8	*
15	Diabetes mellitus (E10-E14)	4	0.6	*
...	All other causes (Residual)	135	21.8	4.3
Black, non-Hispanic, female, 15-24 years				
...	All causes	1,678	100.0	56.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	385	22.9	12.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	313	18.7	10.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	121	7.2	4.0
4	Malignant neoplasms (C00-C97)	104	6.2	3.5
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	61	3.6	2.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	60	3.6	2.0
7	Pregnancy, childbirth and the puerperium (O00-099)	42	2.5	1.4
8	Anemias (D50-D64)	39	2.3	1.3
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	37	2.2	1.2
10	Cerebrovascular diseases (I60-I69)	33	2.0	1.1
10	Chronic lower respiratory diseases (J40-J47)	33	2.0	1.1
12	Septicemia (A40-A41)	22	1.3	0.7
13	Diabetes mellitus (E10-E14)	19	1.1	*
14	Influenza and pneumonia (J10-J18)	18	1.1	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	0.8	*
...	All other causes (Residual)	377	22.5	12.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 25-34 years				
...	All causes	3,203	100.0	118.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	405	12.6	14.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	402	12.6	14.8
3	Human immunodeficiency virus (HIV) disease (B20-B24)	401	12.5	14.8
4	Malignant neoplasms (C00-C97)	393	12.3	14.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	315	9.8	11.6
6	Diabetes mellitus (E10-E14)	86	2.7	3.2
7	Pregnancy, childbirth and the puerperium (O00-099)	82	2.6	3.0
8	Cerebrovascular diseases (I60-I69)	81	2.5	3.0
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	74	2.3	2.7
10	Anemias (D50-D64)	56	1.7	2.1
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	54	1.7	2.0
12	Septicemia (A40-A41)	46	1.4	1.7
13	Influenza and pneumonia (J10-J18)	43	1.3	1.6
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	37	1.2	1.4
15	Chronic lower respiratory diseases (J40-J47)	32	1.0	1.2
...	All other causes (Residual)	696	21.7	25.6
Black, non-Hispanic, female, 35-44 years				
...	All causes	8,156	100.0	278.4
1	Malignant neoplasms (C00-C97)	1,675	20.5	57.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,326	16.3	45.3
3	Human immunodeficiency virus (HIV) disease (B20-B24)	972	11.9	33.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	656	8.0	22.4
5	Cerebrovascular diseases (I60-I69)	401	4.9	13.7
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	257	3.2	8.8
7	Diabetes mellitus (E10-E14)	238	2.9	8.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	164	2.0	5.6
9	Septicemia (A40-A41)	149	1.8	5.1
10	Chronic lower respiratory diseases (J40-J47)	139	1.7	4.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	135	1.7	4.6
12	Influenza and pneumonia (J10-J18)	109	1.3	3.7
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	85	1.0	2.9
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	82	1.0	2.8
15	Anemias (D50-D64)	65	0.8	2.2
...	All other causes (Residual)	1,703	20.9	58.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Black, non-Hispanic, female, 45-54 years			
...	All causes	14,876	100.0	595.7
1	Malignant neoplasms (C00-C97)	4,269	28.7	170.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,182	21.4	127.4
3	Cerebrovascular diseases (I60-I69)	918	6.2	36.8
4	Diabetes mellitus (E10-E14)	690	4.6	27.6
5	Human immunodeficiency virus (HIV) disease (B20-B24)	689	4.6	27.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	664	4.5	26.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	313	2.1	12.5
8	Septicemia (A40-A41)	312	2.1	12.5
9	Chronic lower respiratory diseases (J40-J47)	304	2.0	12.2
10	Chronic liver disease and cirrhosis (K70,K73-K74)	272	1.8	10.9
11	Influenza and pneumonia (J10-J18)	202	1.4	8.1
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	163	1.1	6.5
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	155	1.0	6.2
14	Viral hepatitis (B15-B19)	102	0.7	4.1
15	Anemias (D50-D64)	70	0.5	2.8
...	All other causes (Residual)	2,571	17.3	102.9
	Black, non-Hispanic, female, 55-64 years			
...	All causes	17,882	100.0	1,201.6
1	Malignant neoplasms (C00-C97)	5,768	32.3	387.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,624	25.9	310.7
3	Diabetes mellitus (E10-E14)	1,178	6.6	79.2
4	Cerebrovascular diseases (I60-I69)	1,086	6.1	73.0
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	545	3.0	36.6
6	Chronic lower respiratory diseases (J40-J47)	494	2.8	33.2
7	Septicemia (A40-A41)	419	2.3	28.2
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	328	1.8	22.0
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	305	1.7	20.5
10	Influenza and pneumonia (J10-J18)	209	1.2	14.0
11	Chronic liver disease and cirrhosis (K70,K73-K74)	208	1.2	14.0
12	Human immunodeficiency virus (HIV) disease (B20-B24)	188	1.1	12.6
13	Viral hepatitis (B15-B19)	76	0.4	5.1
14	Aortic aneurysm and dissection (I71)	64	0.4	4.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	61	0.3	4.1
...	All other causes (Residual)	2,329	13.0	156.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Black, non-Hispanic, female, 65-74 years			
...	All causes	24,626	100.0	2,531.2
1	Malignant neoplasms (C00-C97)	7,073	28.7	727.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,032	28.6	722.8
3	Diabetes mellitus (E10-E14)	1,753	7.1	180.2
4	Cerebrovascular diseases (I60-I69)	1,738	7.1	178.6
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	905	3.7	93.0
6	Chronic lower respiratory diseases (J40-J47)	827	3.4	85.0
7	Septicemia (A40-A41)	631	2.6	64.9
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	494	2.0	50.8
9	Influenza and pneumonia (J10-J18)	368	1.5	37.8
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	332	1.3	34.1
11	Alzheimer's disease (G30)	184	0.7	18.9
12	Chronic liver disease and cirrhosis (K70,K73-K74)	168	0.7	17.3
13	Aortic aneurysm and dissection (I71)	118	0.5	12.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	108	0.4	11.1
15	Pneumonitis due to solids and liquids (J69)	94	0.4	9.7
...	All other causes (Residual)	2,801	11.4	287.9
	Black, non-Hispanic, female, 75-84 years			
...	All causes	33,473	100.0	5,454.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,558	31.5	1,720.5
2	Malignant neoplasms (C00-C97)	6,942	20.7	1,131.2
3	Cerebrovascular diseases (I60-I69)	3,105	9.3	506.0
4	Diabetes mellitus (E10-E14)	2,077	6.2	338.5
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,153	3.4	187.9
6	Chronic lower respiratory diseases (J40-J47)	950	2.8	154.8
7	Septicemia (A40-A41)	873	2.6	142.3
8	Alzheimer's disease (G30)	866	2.6	141.1
9	Influenza and pneumonia (J10-J18)	749	2.2	122.1
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	743	2.2	121.1
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	369	1.1	60.1
12	Pneumonitis due to solids and liquids (J69)	257	0.8	41.9
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	157	0.5	25.6
14	Atherosclerosis (I70)	156	0.5	25.4
15	Aortic aneurysm and dissection (I71)	154	0.5	25.1
...	All other causes (Residual)	4,364	13.0	711.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2003

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Black, non-Hispanic, female, 85 years and over			
...	All causes	33,270	100.0	13,730.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,137	36.5	5,008.9
2	Malignant neoplasms (C00-C97)	3,594	10.8	1,483.2
3	Cerebrovascular diseases (I60-I69)	3,465	10.4	1,430.0
4	Alzheimer's disease (G30)	1,550	4.7	639.7
5	Diabetes mellitus (E10-E14)	1,360	4.1	561.3
6	Influenza and pneumonia (J10-J18)	1,173	3.5	484.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,007	3.0	415.6
8	Septicemia (A40-A41)	873	2.6	360.3
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	760	2.3	313.6
10	Chronic lower respiratory diseases (J40-J47)	622	1.9	256.7
11	Pneumonitis due to solids and liquids (J69)	355	1.1	146.5
12	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	330	1.0	136.2
13	Atherosclerosis (I70)	314	0.9	129.6
14	Nutritional deficiencies (E40-E64)	142	0.4	58.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	134	0.4	55.3
...	All other causes (Residual)	5,454	16.4	2,250.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable