

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, all ages				
...	All causes	122,416	100.0	296.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	27,788	22.7	67.2
2	Malignant neoplasms (C00-C97)	24,522	20.0	59.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,408	8.5	25.2
4	Cerebrovascular diseases (I60-I69)	6,781	5.5	16.4
5	Diabetes mellitus (E10-E14)	5,999	4.9	14.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	3,383	2.8	8.2
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,271	2.7	7.9
8	Chronic lower respiratory diseases (J40-J47)	3,107	2.5	7.5
9	Influenza and pneumonia (J10-J18)	2,912	2.4	7.0
10	Certain conditions originating in the perinatal period (P00-P96)	2,681	2.2	6.5
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,207	1.8	5.3
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,196	1.8	5.3
13	Alzheimer's disease (G30)	1,897	1.5	4.6
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,765	1.4	4.3
15	Human immunodeficiency virus (HIV) disease (B20-B24)	1,758	1.4	4.3
...	All other causes (Residual)	21,741	17.8	52.6
Hispanic, both sexes, 1-4 years				
...	All causes	947	100.0	27.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	313	33.1	9.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	138	14.6	4.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	80	8.4	2.3
4	Malignant neoplasms (C00-C97)	70	7.4	2.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	39	4.1	1.1
6	Septicemia (A40-A41)	24	2.5	0.7
6	Influenza and pneumonia (J10-J18)	24	2.5	0.7
8	Chronic lower respiratory diseases (J40-J47)	11	1.2	*
9	Certain conditions originating in the perinatal period (P00-P96)	10	1.1	*
10	Meningococcal infection (A39)	5	0.5	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.5	*
10	Anemias (D50-D64)	5	0.5	*
10	Acute bronchitis and bronchiolitis (J20-J21)	5	0.5	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.5	*
15	Meningitis (G00,G03)	3	0.3	*
15	Cerebrovascular diseases (I60-I69)	3	0.3	*
15	Pneumonitis due to solids and liquids (J69)	3	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.3	*
...	All other causes (Residual)	201	21.2	5.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 5-14 years				
...	All causes	1,105	100.0	14.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	391	35.4	5.1
2	Malignant neoplasms (C00-C97)	211	19.1	2.8
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	70	6.3	0.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	67	6.1	0.9
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	39	3.5	0.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	34	3.1	0.4
7	Influenza and pneumonia (J10-J18)	13	1.2	*
8	Chronic lower respiratory diseases (J40-J47)	12	1.1	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	1.0	*
10	Cerebrovascular diseases (I60-I69)	10	0.9	*
11	Septicemia (A40-A41)	9	0.8	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.7	*
12	Certain conditions originating in the perinatal period (P00-P96)	8	0.7	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.5	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.4	*
...	All other causes (Residual)	212	19.2	2.8
Hispanic, both sexes, 15-24 years				
...	All causes	5,428	100.0	76.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,372	43.7	33.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,235	22.8	17.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	571	10.5	8.0
4	Malignant neoplasms (C00-C97)	305	5.6	4.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	129	2.4	1.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	61	1.1	0.9
7	Cerebrovascular diseases (I60-I69)	36	0.7	0.5
8	Influenza and pneumonia (J10-J18)	32	0.6	0.5
9	Legal intervention (Y35,Y89.0)	30	0.6	0.4
10	Pregnancy, childbirth and the puerperium (O00-099)	29	0.5	0.4
11	Human immunodeficiency virus (HIV) disease (B20-B24)	25	0.5	0.4
12	Chronic lower respiratory diseases (J40-J47)	17	0.3	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	15	0.3	*
14	Septicemia (A40-A41)	13	0.2	*
14	Diabetes mellitus (E10-E14)	13	0.2	*
...	All other causes (Residual)	545	10.0	7.7

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 25-34 years				
...	All causes	5,959	100.0	78.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,086	35.0	27.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	962	16.1	12.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	545	9.1	7.2
4	Malignant neoplasms (C00-C97)	526	8.8	6.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	302	5.1	4.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	211	3.5	2.8
7	Cerebrovascular diseases (I60-I69)	93	1.6	1.2
8	Chronic liver disease and cirrhosis (K70,K73-K74)	73	1.2	1.0
9	Diabetes mellitus (E10-E14)	64	1.1	0.8
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	53	0.9	0.7
11	Pregnancy, childbirth and the puerperium (O00-099)	49	0.8	0.6
12	Influenza and pneumonia (J10-J18)	48	0.8	0.6
13	Septicemia (A40-A41)	35	0.6	0.5
14	Chronic lower respiratory diseases (J40-J47)	30	0.5	0.4
14	Legal intervention (Y35,Y89.0)	30	0.5	0.4
...	All other causes (Residual)	852	14.3	11.2
Hispanic, both sexes, 35-44 years				
...	All causes	8,546	100.0	138.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,823	21.3	29.6
2	Malignant neoplasms (C00-C97)	1,345	15.7	21.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	934	10.9	15.2
4	Human immunodeficiency virus (HIV) disease (B20-B24)	676	7.9	11.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	477	5.6	7.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	468	5.5	7.6
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	421	4.9	6.8
8	Cerebrovascular diseases (I60-I69)	304	3.6	4.9
9	Diabetes mellitus (E10-E14)	187	2.2	3.0
10	Viral hepatitis (B15-B19)	110	1.3	1.8
11	Septicemia (A40-A41)	88	1.0	1.4
12	Influenza and pneumonia (J10-J18)	82	1.0	1.3
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	72	0.8	1.2
14	Chronic lower respiratory diseases (J40-J47)	66	0.8	1.1
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	40	0.5	0.6
...	All other causes (Residual)	1,453	17.0	23.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 45-54 years				
...	All causes	12,628	100.0	313.5
1	Malignant neoplasms (C00-C97)	3,040	24.1	75.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,110	16.7	52.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,249	9.9	31.0
4	Chronic liver disease and cirrhosis (K70,K73-K74)	998	7.9	24.8
5	Cerebrovascular diseases (I60-I69)	590	4.7	14.6
6	Human immunodeficiency virus (HIV) disease (B20-B24)	569	4.5	14.1
7	Diabetes mellitus (E10-E14)	535	4.2	13.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	304	2.4	7.5
9	Viral hepatitis (B15-B19)	275	2.2	6.8
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	238	1.9	5.9
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	182	1.4	4.5
12	Septicemia (A40-A41)	162	1.3	4.0
13	Influenza and pneumonia (J10-J18)	153	1.2	3.8
14	Chronic lower respiratory diseases (J40-J47)	124	1.0	3.1
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	89	0.7	2.2
...	All other causes (Residual)	2,010	15.9	49.9
Hispanic, both sexes, 55-64 years				
...	All causes	15,198	100.0	684.0
1	Malignant neoplasms (C00-C97)	4,659	30.7	209.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,460	22.8	155.7
3	Diabetes mellitus (E10-E14)	1,073	7.1	48.3
4	Cerebrovascular diseases (I60-I69)	776	5.1	34.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	769	5.1	34.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	678	4.5	30.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	319	2.1	14.4
8	Chronic lower respiratory diseases (J40-J47)	272	1.8	12.2
9	Septicemia (A40-A41)	216	1.4	9.7
10	Influenza and pneumonia (J10-J18)	214	1.4	9.6
11	Human immunodeficiency virus (HIV) disease (B20-B24)	186	1.2	8.4
12	Viral hepatitis (B15-B19)	171	1.1	7.7
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	156	1.0	7.0
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	121	0.8	5.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	84	0.6	3.8
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	84	0.6	3.8
...	All other causes (Residual)	1,960	12.9	88.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 65-74 years				
...	All causes	20,420	100.0	1,607.5
1	Malignant neoplasms (C00-C97)	5,996	29.4	472.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,389	26.4	424.2
3	Diabetes mellitus (E10-E14)	1,587	7.8	124.9
4	Cerebrovascular diseases (I60-I69)	1,232	6.0	97.0
5	Chronic lower respiratory diseases (J40-J47)	657	3.2	51.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	601	2.9	47.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	523	2.6	41.2
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	473	2.3	37.2
9	Influenza and pneumonia (J10-J18)	422	2.1	33.2
10	Septicemia (A40-A41)	319	1.6	25.1
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	205	1.0	16.1
12	Alzheimer's disease (G30)	153	0.7	12.0
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	110	0.5	8.7
14	Aortic aneurysm and dissection (I71)	100	0.5	7.9
15	Viral hepatitis (B15-B19)	89	0.4	7.0
...	All other causes (Residual)	2,564	12.6	201.8
Hispanic, both sexes, 75-84 years				
...	All causes	26,879	100.0	3,926.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,061	30.0	1,177.6
2	Malignant neoplasms (C00-C97)	5,970	22.2	872.2
3	Cerebrovascular diseases (I60-I69)	2,006	7.5	293.1
4	Diabetes mellitus (E10-E14)	1,769	6.6	258.4
5	Chronic lower respiratory diseases (J40-J47)	1,086	4.0	158.7
6	Influenza and pneumonia (J10-J18)	857	3.2	125.2
7	Alzheimer's disease (G30)	681	2.5	99.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	629	2.3	91.9
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	508	1.9	74.2
10	Septicemia (A40-A41)	437	1.6	63.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	412	1.5	60.2
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	303	1.1	44.3
13	Parkinson's disease (G20-G21)	302	1.1	44.1
14	Pneumonitis due to solids and liquids (J69)	180	0.7	26.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	160	0.6	23.4
...	All other causes (Residual)	3,518	13.1	513.9

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 85 years and over				
...	All causes	19,937	100.0	9,483.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,246	36.3	3,446.8
2	Malignant neoplasms (C00-C97)	2,378	11.9	1,131.2
3	Cerebrovascular diseases (I60-I69)	1,709	8.6	813.0
4	Alzheimer's disease (G30)	1,032	5.2	490.9
5	Influenza and pneumonia (J10-J18)	1,014	5.1	482.3
6	Chronic lower respiratory diseases (J40-J47)	823	4.1	391.5
7	Diabetes mellitus (E10-E14)	769	3.9	365.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	386	1.9	183.6
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	362	1.8	172.2
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	338	1.7	160.8
11	Septicemia (A40-A41)	290	1.5	138.0
12	Pneumonitis due to solids and liquids (J69)	213	1.1	101.3
13	Parkinson's disease (G20-G21)	179	0.9	85.1
14	Atherosclerosis (I70)	157	0.8	74.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	102	0.5	48.5
...	All other causes (Residual)	2,939	14.7	1,398.1
Hispanic, male, all ages				
...	All causes	68,544	100.0	321.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,983	21.9	70.2
2	Malignant neoplasms (C00-C97)	13,002	19.0	60.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,796	11.4	36.5
4	Cerebrovascular diseases (I60-I69)	3,207	4.7	15.0
5	Diabetes mellitus (E10-E14)	2,855	4.2	13.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,782	4.1	13.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,376	3.5	11.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,840	2.7	8.6
9	Chronic lower respiratory diseases (J40-J47)	1,631	2.4	7.6
10	Certain conditions originating in the perinatal period (P00-P96)	1,538	2.2	7.2
11	Influenza and pneumonia (J10-J18)	1,403	2.0	6.6
12	Human immunodeficiency virus (HIV) disease (B20-B24)	1,371	2.0	6.4
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,101	1.6	5.2
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	902	1.3	4.2
15	Septicemia (A40-A41)	801	1.2	3.8
...	All other causes (Residual)	10,956	16.0	51.3

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 1-4 years				
...	All causes	535	100.0	30.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	186	34.8	10.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	82	15.3	4.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	45	8.4	2.5
4	Malignant neoplasms (C00-C97)	29	5.4	1.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	24	4.5	1.4
6	Septicemia (A40-A41)	14	2.6	*
7	Influenza and pneumonia (J10-J18)	9	1.7	*
8	Chronic lower respiratory diseases (J40-J47)	7	1.3	*
9	Certain conditions originating in the perinatal period (P00-P96)	6	1.1	*
10	Anemias (D50-D64)	3	0.6	*
10	Cerebrovascular diseases (I60-I69)	3	0.6	*
12	Meningococcal infection (A39)	2	0.4	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.4	*
12	Diabetes mellitus (E10-E14)	2	0.4	*
12	Meningitis (G00,G03)	2	0.4	*
12	Acute bronchitis and bronchiolitis (J20-J21)	2	0.4	*
12	Pneumonitis due to solids and liquids (J69)	2	0.4	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
...	All other causes (Residual)	111	20.7	6.3
Hispanic, male, 5-14 years				
...	All causes	656	100.0	16.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	241	36.7	6.2
2	Malignant neoplasms (C00-C97)	118	18.0	3.0
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	49	7.5	1.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	40	6.1	1.0
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	24	3.7	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	15	2.3	*
7	Influenza and pneumonia (J10-J18)	10	1.5	*
8	Chronic lower respiratory diseases (J40-J47)	7	1.1	*
9	Septicemia (A40-A41)	6	0.9	*
10	Cerebrovascular diseases (I60-I69)	5	0.8	*
10	Certain conditions originating in the perinatal period (P00-P96)	5	0.8	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.6	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.6	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.5	*
15	Anemias (D50-D64)	2	0.3	*
15	Meningitis (G00,G03)	2	0.3	*
15	Diseases of appendix (K35-K38)	2	0.3	*
...	All other causes (Residual)	119	18.1	3.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 15-24 years				
...	All causes	4,376	100.0	115.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,926	44.0	50.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,125	25.7	29.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	487	11.1	12.8
4	Malignant neoplasms (C00-C97)	207	4.7	5.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	101	2.3	2.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	40	0.9	1.1
7	Legal intervention (Y35,Y89.0)	30	0.7	0.8
8	Influenza and pneumonia (J10-J18)	24	0.5	0.6
9	Cerebrovascular diseases (I60-I69)	20	0.5	0.5
10	Human immunodeficiency virus (HIV) disease (B20-B24)	19	0.4	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.3	*
11	Chronic lower respiratory diseases (J40-J47)	11	0.3	*
13	Diabetes mellitus (E10-E14)	8	0.2	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.2	*
15	Septicemia (A40-A41)	7	0.2	*
15	Aortic aneurysm and dissection (I71)	7	0.2	*
...	All other causes (Residual)	345	7.9	9.1
Hispanic, male, 25-34 years				
...	All causes	4,547	100.0	109.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,733	38.1	41.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	844	18.6	20.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	467	10.3	11.2
4	Malignant neoplasms (C00-C97)	267	5.9	6.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	244	5.4	5.9
6	Human immunodeficiency virus (HIV) disease (B20-B24)	169	3.7	4.1
7	Cerebrovascular diseases (I60-I69)	61	1.3	1.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	61	1.3	1.5
9	Diabetes mellitus (E10-E14)	37	0.8	0.9
10	Influenza and pneumonia (J10-J18)	31	0.7	0.7
11	Legal intervention (Y35,Y89.0)	30	0.7	0.7
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	24	0.5	0.6
13	Chronic lower respiratory diseases (J40-J47)	20	0.4	0.5
14	Septicemia (A40-A41)	16	0.4	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	15	0.3	*
...	All other causes (Residual)	528	11.6	12.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 35-44 years				
...	All causes	5,973	100.0	184.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,454	24.3	44.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	715	12.0	22.1
3	Malignant neoplasms (C00-C97)	604	10.1	18.6
4	Human immunodeficiency virus (HIV) disease (B20-B24)	522	8.7	16.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	384	6.4	11.8
6	Chronic liver disease and cirrhosis (K70,K73-K74)	377	6.3	11.6
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	351	5.9	10.8
8	Cerebrovascular diseases (I60-I69)	190	3.2	5.9
9	Diabetes mellitus (E10-E14)	111	1.9	3.4
10	Viral hepatitis (B15-B19)	85	1.4	2.6
11	Septicemia (A40-A41)	53	0.9	1.6
12	Influenza and pneumonia (J10-J18)	48	0.8	1.5
13	Chronic lower respiratory diseases (J40-J47)	45	0.8	1.4
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	45	0.8	1.4
15	Aortic aneurysm and dissection (I71)	24	0.4	0.7
...	All other causes (Residual)	965	16.2	29.8
Hispanic, male, 45-54 years				
...	All causes	8,451	100.0	417.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,569	18.6	77.6
2	Malignant neoplasms (C00-C97)	1,566	18.5	77.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	968	11.5	47.9
4	Chronic liver disease and cirrhosis (K70,K73-K74)	825	9.8	40.8
5	Human immunodeficiency virus (HIV) disease (B20-B24)	445	5.3	22.0
6	Cerebrovascular diseases (I60-I69)	354	4.2	17.5
7	Diabetes mellitus (E10-E14)	323	3.8	16.0
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	242	2.9	12.0
9	Viral hepatitis (B15-B19)	206	2.4	10.2
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	193	2.3	9.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	107	1.3	5.3
12	Septicemia (A40-A41)	106	1.3	5.2
13	Influenza and pneumonia (J10-J18)	100	1.2	4.9
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	58	0.7	2.9
15	Chronic lower respiratory diseases (J40-J47)	56	0.7	2.8
...	All other causes (Residual)	1,333	15.8	65.9

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 55-64 years				
...	All causes	9,246	100.0	874.0
1	Malignant neoplasms (C00-C97)	2,528	27.3	239.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,376	25.7	224.6
3	Diabetes mellitus (E10-E14)	602	6.5	56.9
4	Chronic liver disease and cirrhosis (K70,K73-K74)	554	6.0	52.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	474	5.1	44.8
6	Cerebrovascular diseases (I60-I69)	454	4.9	42.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	180	1.9	17.0
8	Human immunodeficiency virus (HIV) disease (B20-B24)	152	1.6	14.4
9	Chronic lower respiratory diseases (J40-J47)	145	1.6	13.7
10	Influenza and pneumonia (J10-J18)	122	1.3	11.5
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	120	1.3	11.3
12	Viral hepatitis (B15-B19)	110	1.2	10.4
13	Septicemia (A40-A41)	109	1.2	10.3
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	69	0.7	6.5
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	68	0.7	6.4
...	All other causes (Residual)	1,183	12.8	111.8
Hispanic, male, 65-74 years				
...	All causes	11,279	100.0	1,994.3
1	Malignant neoplasms (C00-C97)	3,313	29.4	585.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,236	28.7	572.2
3	Diabetes mellitus (E10-E14)	759	6.7	134.2
4	Cerebrovascular diseases (I60-I69)	647	5.7	114.4
5	Chronic lower respiratory diseases (J40-J47)	387	3.4	68.4
6	Chronic liver disease and cirrhosis (K70,K73-K74)	336	3.0	59.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	294	2.6	52.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	251	2.2	44.4
9	Influenza and pneumonia (J10-J18)	220	2.0	38.9
10	Septicemia (A40-A41)	154	1.4	27.2
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	104	0.9	18.4
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	77	0.7	13.6
13	Aortic aneurysm and dissection (I71)	71	0.6	12.6
14	Alzheimer's disease (G30)	65	0.6	11.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	55	0.5	9.7
...	All other causes (Residual)	1,310	11.6	231.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 75-84 years				
...	All causes	13,397	100.0	4,791.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,163	31.1	1,489.0
2	Malignant neoplasms (C00-C97)	3,283	24.5	1,174.2
3	Cerebrovascular diseases (I60-I69)	904	6.7	323.3
4	Diabetes mellitus (E10-E14)	781	5.8	279.3
5	Chronic lower respiratory diseases (J40-J47)	577	4.3	206.4
6	Influenza and pneumonia (J10-J18)	425	3.2	152.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	306	2.3	109.4
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	278	2.1	99.4
9	Alzheimer's disease (G30)	260	1.9	93.0
10	Chronic liver disease and cirrhosis (K70,K73-K74)	198	1.5	70.8
11	Septicemia (A40-A41)	195	1.5	69.7
12	Parkinson's disease (G20-G21)	186	1.4	66.5
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	136	1.0	48.6
14	Pneumonitis due to solids and liquids (J69)	89	0.7	31.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	78	0.6	27.9
...	All other causes (Residual)	1,538	11.5	550.1
Hispanic, male, 85 years and over				
...	All causes	7,090	100.0	9,932.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,496	35.2	3,496.8
2	Malignant neoplasms (C00-C97)	1,077	15.2	1,508.8
3	Cerebrovascular diseases (I60-I69)	556	7.8	778.9
4	Influenza and pneumonia (J10-J18)	377	5.3	528.2
5	Chronic lower respiratory diseases (J40-J47)	370	5.2	518.4
6	Alzheimer's disease (G30)	267	3.8	374.1
7	Diabetes mellitus (E10-E14)	232	3.3	325.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	165	2.3	231.2
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	138	1.9	193.3
10	Septicemia (A40-A41)	105	1.5	147.1
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	105	1.5	147.1
12	Parkinson's disease (G20-G21)	90	1.3	126.1
13	Pneumonitis due to solids and liquids (J69)	87	1.2	121.9
14	Atherosclerosis (I70)	63	0.9	88.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	41	0.6	57.4
...	All other causes (Residual)	921	13.0	1,290.3

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, all ages				
...	All causes	53,872	100.0	269.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,805	23.8	64.1
2	Malignant neoplasms (C00-C97)	11,520	21.4	57.7
3	Cerebrovascular diseases (I60-I69)	3,574	6.6	17.9
4	Diabetes mellitus (E10-E14)	3,144	5.8	15.7
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,612	4.8	13.1
6	Influenza and pneumonia (J10-J18)	1,509	2.8	7.6
7	Chronic lower respiratory diseases (J40-J47)	1,476	2.7	7.4
8	Alzheimer's disease (G30)	1,291	2.4	6.5
9	Certain conditions originating in the perinatal period (P00-P96)	1,143	2.1	5.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,095	2.0	5.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1,007	1.9	5.0
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	863	1.6	4.3
13	Septicemia (A40-A41)	847	1.6	4.2
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	624	1.2	3.1
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	489	0.9	2.4
...	All other causes (Residual)	9,873	18.3	49.4
Hispanic, female, 1-4 years				
...	All causes	412	100.0	24.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	127	30.8	7.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	56	13.6	3.3
3	Malignant neoplasms (C00-C97)	41	10.0	2.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	35	8.5	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	15	3.6	*
5	Influenza and pneumonia (J10-J18)	15	3.6	*
7	Septicemia (A40-A41)	10	2.4	*
8	Chronic lower respiratory diseases (J40-J47)	4	1.0	*
8	Certain conditions originating in the perinatal period (P00-P96)	4	1.0	*
10	Meningococcal infection (A39)	3	0.7	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.7	*
10	Acute bronchitis and bronchiolitis (J20-J21)	3	0.7	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.7	*
14	Anemias (D50-D64)	2	0.5	*
15	Meningitis (G00,G03)	1	0.2	*
15	Pneumonitis due to solids and liquids (J69)	1	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
...	All other causes (Residual)	88	21.4	5.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 5-14 years				
...	All causes	449	100.0	12.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	150	33.4	4.0
2	Malignant neoplasms (C00-C97)	93	20.7	2.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	27	6.0	0.7
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	21	4.7	0.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	19	4.2	*
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	15	3.3	*
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	1.6	*
8	Cerebrovascular diseases (I60-I69)	5	1.1	*
8	Chronic lower respiratory diseases (J40-J47)	5	1.1	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.9	*
11	Septicemia (A40-A41)	3	0.7	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	3	0.7	*
11	Influenza and pneumonia (J10-J18)	3	0.7	*
11	Certain conditions originating in the perinatal period (P00-P96)	3	0.7	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.7	*
...	All other causes (Residual)	88	19.6	2.4
Hispanic, female, 15-24 years				
...	All causes	1,052	100.0	31.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	446	42.4	13.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	110	10.5	3.3
3	Malignant neoplasms (C00-C97)	98	9.3	3.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	84	8.0	2.5
5	Pregnancy, childbirth and the puerperium (O00-099)	29	2.8	0.9
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	28	2.7	0.8
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	21	2.0	0.6
8	Cerebrovascular diseases (I60-I69)	16	1.5	*
9	Anemias (D50-D64)	8	0.8	*
9	Influenza and pneumonia (J10-J18)	8	0.8	*
11	Septicemia (A40-A41)	6	0.6	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	6	0.6	*
11	Chronic lower respiratory diseases (J40-J47)	6	0.6	*
14	Diabetes mellitus (E10-E14)	5	0.5	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.4	*
...	All other causes (Residual)	173	16.4	5.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 25-34 years				
...	All causes	1,412	100.0	40.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	353	25.0	10.2
2	Malignant neoplasms (C00-C97)	259	18.3	7.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	118	8.4	3.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	78	5.5	2.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	58	4.1	1.7
6	Pregnancy, childbirth and the puerperium (O00-099)	49	3.5	1.4
7	Human immunodeficiency virus (HIV) disease (B20-B24)	42	3.0	1.2
8	Cerebrovascular diseases (I60-I69)	32	2.3	0.9
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	29	2.1	0.8
10	Diabetes mellitus (E10-E14)	27	1.9	0.8
11	Septicemia (A40-A41)	19	1.3	*
12	Influenza and pneumonia (J10-J18)	17	1.2	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	1.0	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	1.0	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	12	0.8	*
...	All other causes (Residual)	291	20.6	8.4
Hispanic, female, 35-44 years				
...	All causes	2,573	100.0	88.2
1	Malignant neoplasms (C00-C97)	741	28.8	25.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	369	14.3	12.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	219	8.5	7.5
4	Human immunodeficiency virus (HIV) disease (B20-B24)	154	6.0	5.3
5	Cerebrovascular diseases (I60-I69)	114	4.4	3.9
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	93	3.6	3.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	91	3.5	3.1
8	Diabetes mellitus (E10-E14)	76	3.0	2.6
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	70	2.7	2.4
10	Septicemia (A40-A41)	35	1.4	1.2
11	Influenza and pneumonia (J10-J18)	34	1.3	1.2
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	27	1.0	0.9
13	Pregnancy, childbirth and the puerperium (O00-099)	26	1.0	0.9
14	Viral hepatitis (B15-B19)	25	1.0	0.9
15	Chronic lower respiratory diseases (J40-J47)	21	0.8	0.7
...	All other causes (Residual)	478	18.6	16.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 45-54 years				
...	All causes	4,177	100.0	208.3
1	Malignant neoplasms (C00-C97)	1,474	35.3	73.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	541	13.0	27.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	281	6.7	14.0
4	Cerebrovascular diseases (I60-I69)	236	5.6	11.8
5	Diabetes mellitus (E10-E14)	212	5.1	10.6
6	Chronic liver disease and cirrhosis (K70,K73-K74)	173	4.1	8.6
7	Human immunodeficiency virus (HIV) disease (B20-B24)	124	3.0	6.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	75	1.8	3.7
9	Viral hepatitis (B15-B19)	69	1.7	3.4
10	Chronic lower respiratory diseases (J40-J47)	68	1.6	3.4
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	62	1.5	3.1
12	Septicemia (A40-A41)	56	1.3	2.8
13	Influenza and pneumonia (J10-J18)	53	1.3	2.6
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	45	1.1	2.2
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	31	0.7	1.5
...	All other causes (Residual)	677	16.2	33.8
Hispanic, female, 55-64 years				
...	All causes	5,952	100.0	511.3
1	Malignant neoplasms (C00-C97)	2,131	35.8	183.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,084	18.2	93.1
3	Diabetes mellitus (E10-E14)	471	7.9	40.5
4	Cerebrovascular diseases (I60-I69)	322	5.4	27.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	215	3.6	18.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	204	3.4	17.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	139	2.3	11.9
8	Chronic lower respiratory diseases (J40-J47)	127	2.1	10.9
9	Septicemia (A40-A41)	107	1.8	9.2
10	Influenza and pneumonia (J10-J18)	92	1.5	7.9
11	Viral hepatitis (B15-B19)	61	1.0	5.2
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	53	0.9	4.6
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	41	0.7	3.5
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	36	0.6	3.1
15	Human immunodeficiency virus (HIV) disease (B20-B24)	34	0.6	2.9
...	All other causes (Residual)	835	14.0	71.7

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 65-74 years				
...	All causes	9,141	100.0	1,297.2
1	Malignant neoplasms (C00-C97)	2,683	29.4	380.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,153	23.6	305.5
3	Diabetes mellitus (E10-E14)	828	9.1	117.5
4	Cerebrovascular diseases (I60-I69)	585	6.4	83.0
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	272	3.0	38.6
6	Chronic lower respiratory diseases (J40-J47)	270	3.0	38.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	265	2.9	37.6
8	Influenza and pneumonia (J10-J18)	202	2.2	28.7
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	179	2.0	25.4
10	Septicemia (A40-A41)	165	1.8	23.4
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	101	1.1	14.3
12	Alzheimer's disease (G30)	88	1.0	12.5
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	55	0.6	7.8
14	Viral hepatitis (B15-B19)	44	0.5	6.2
15	Parkinson's disease (G20-G21)	33	0.4	4.7
...	All other causes (Residual)	1,218	13.3	172.8
Hispanic, female, 75-84 years				
...	All causes	13,482	100.0	3,329.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,898	28.9	962.7
2	Malignant neoplasms (C00-C97)	2,687	19.9	663.6
3	Cerebrovascular diseases (I60-I69)	1,102	8.2	272.2
4	Diabetes mellitus (E10-E14)	988	7.3	244.0
5	Chronic lower respiratory diseases (J40-J47)	509	3.8	125.7
6	Influenza and pneumonia (J10-J18)	432	3.2	106.7
7	Alzheimer's disease (G30)	421	3.1	104.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	323	2.4	79.8
9	Septicemia (A40-A41)	242	1.8	59.8
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	230	1.7	56.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	214	1.6	52.9
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	167	1.2	41.2
13	Parkinson's disease (G20-G21)	116	0.9	28.6
14	Pneumonitis due to solids and liquids (J69)	91	0.7	22.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	82	0.6	20.3
...	All other causes (Residual)	1,980	14.7	489.0

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 85 years and over				
...	All causes	12,847	100.0	9,253.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,750	37.0	3,421.2
2	Malignant neoplasms (C00-C97)	1,301	10.1	937.0
3	Cerebrovascular diseases (I60-I69)	1,153	9.0	830.4
4	Alzheimer's disease (G30)	765	6.0	551.0
5	Influenza and pneumonia (J10-J18)	637	5.0	458.8
6	Diabetes mellitus (E10-E14)	537	4.2	386.8
7	Chronic lower respiratory diseases (J40-J47)	453	3.5	326.3
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	257	2.0	185.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	221	1.7	159.2
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	200	1.6	144.0
11	Septicemia (A40-A41)	185	1.4	133.2
12	Pneumonitis due to solids and liquids (J69)	126	1.0	90.8
13	Atherosclerosis (I70)	94	0.7	67.7
14	Parkinson's disease (G20-G21)	89	0.7	64.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	61	0.5	43.9
...	All other causes (Residual)	2,018	15.7	1,453.5
White, non-Hispanic, both sexes, all ages				
...	All causes	1,933,382	100.0	967.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	537,512	27.8	269.1
2	Malignant neoplasms (C00-C97)	453,534	23.5	227.0
3	Cerebrovascular diseases (I60-I69)	121,065	6.3	60.6
4	Chronic lower respiratory diseases (J40-J47)	109,670	5.7	54.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	85,466	4.4	42.8
6	Alzheimer's disease (G30)	59,153	3.1	29.6
7	Diabetes mellitus (E10-E14)	52,101	2.7	26.1
8	Influenza and pneumonia (J10-J18)	49,456	2.6	24.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	31,508	1.6	15.8
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	27,001	1.4	13.5
11	Septicemia (A40-A41)	25,041	1.3	12.5
12	Chronic liver disease and cirrhosis (K70,K73-K74)	20,144	1.0	10.1
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	16,569	0.9	8.3
14	Parkinson's disease (G20-G21)	16,428	0.8	8.2
15	Pneumonitis due to solids and liquids (J69)	14,411	0.7	7.2
...	All other causes (Residual)	314,323	16.3	157.3

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 1-4 years				
...	All causes	2,458	100.0	26.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	933	38.0	10.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	270	11.0	2.9
3	Malignant neoplasms (C00-C97)	244	9.9	2.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	131	5.3	1.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	80	3.3	0.9
6	Influenza and pneumonia (J10-J18)	55	2.2	0.6
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	39	1.6	0.4
8	Septicemia (A40-A41)	36	1.5	0.4
9	Cerebrovascular diseases (I60-I69)	25	1.0	0.3
10	Certain conditions originating in the perinatal period (P00-P96)	24	1.0	0.3
11	Meningitis (G00,G03)	16	0.7	*
12	Chronic lower respiratory diseases (J40-J47)	12	0.5	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	11	0.4	*
14	Meningococcal infection (A39)	10	0.4	*
15	Anemias (D50-D64)	8	0.3	*
...	All other causes (Residual)	564	22.9	6.1
White, non-Hispanic, both sexes, 5-14 years				
...	All causes	3,820	100.0	15.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,573	41.2	6.4
2	Malignant neoplasms (C00-C97)	587	15.4	2.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	217	5.7	0.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	180	4.7	0.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	125	3.3	0.5
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	119	3.1	0.5
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	52	1.4	0.2
8	Cerebrovascular diseases (I60-I69)	48	1.3	0.2
9	Influenza and pneumonia (J10-J18)	43	1.1	0.2
10	Septicemia (A40-A41)	41	1.1	0.2
10	Chronic lower respiratory diseases (J40-J47)	41	1.1	0.2
12	Diabetes mellitus (E10-E14)	19	0.5	*
13	Certain conditions originating in the perinatal period (P00-P96)	18	0.5	*
14	Anemias (D50-D64)	17	0.4	*
14	Meningitis (G00,G03)	17	0.4	*
...	All other causes (Residual)	723	18.9	2.9

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 15-24 years				
...	All causes	19,684	100.0	75.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,847	55.1	41.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,045	15.5	11.6
3	Malignant neoplasms (C00-C97)	1,016	5.2	3.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	917	4.7	3.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	553	2.8	2.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	290	1.5	1.1
7	Cerebrovascular diseases (I60-I69)	117	0.6	0.4
8	Influenza and pneumonia (J10-J18)	108	0.5	0.4
9	Diabetes mellitus (E10-E14)	87	0.4	0.3
10	Chronic lower respiratory diseases (J40-J47)	75	0.4	0.3
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	72	0.4	0.3
12	Septicemia (A40-A41)	70	0.4	0.3
13	Pregnancy, childbirth and the puerperium (O00-099)	60	0.3	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	41	0.2	0.2
15	Anemias (D50-D64)	29	0.1	0.1
...	All other causes (Residual)	2,357	12.0	9.0
White, non-Hispanic, both sexes, 25-34 years				
...	All causes	23,397	100.0	95.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,712	37.2	35.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,747	16.0	15.4
3	Malignant neoplasms (C00-C97)	2,270	9.7	9.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,745	7.5	7.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	976	4.2	4.0
6	Diabetes mellitus (E10-E14)	333	1.4	1.4
7	Human immunodeficiency virus (HIV) disease (B20-B24)	296	1.3	1.2
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	295	1.3	1.2
9	Cerebrovascular diseases (I60-I69)	280	1.2	1.1
10	Influenza and pneumonia (J10-J18)	183	0.8	0.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	178	0.8	0.7
12	Septicemia (A40-A41)	172	0.7	0.7
13	Pregnancy, childbirth and the puerperium (O00-099)	122	0.5	0.5
14	Chronic lower respiratory diseases (J40-J47)	120	0.5	0.5
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	108	0.5	0.4
...	All other causes (Residual)	3,860	16.5	15.8

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 35-44 years				
...	All causes	55,543	100.0	185.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,025	21.6	40.3
2	Malignant neoplasms (C00-C97)	10,062	18.1	33.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,411	15.1	28.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,598	10.1	18.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,863	3.4	6.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	1,529	2.8	5.1
7	Cerebrovascular diseases (I60-I69)	1,256	2.3	4.2
8	Diabetes mellitus (E10-E14)	1,238	2.2	4.1
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,124	2.0	3.8
10	Chronic lower respiratory diseases (J40-J47)	583	1.0	2.0
11	Influenza and pneumonia (J10-J18)	552	1.0	1.8
12	Septicemia (A40-A41)	474	0.9	1.6
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	394	0.7	1.3
14	Viral hepatitis (B15-B19)	390	0.7	1.3
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	369	0.7	1.2
...	All other causes (Residual)	9,675	17.4	32.4
White, non-Hispanic, both sexes, 45-54 years				
...	All causes	123,037	100.0	400.8
1	Malignant neoplasms (C00-C97)	36,016	29.3	117.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	25,858	21.0	84.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,744	10.4	41.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,101	5.0	19.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	5,330	4.3	17.4
6	Diabetes mellitus (E10-E14)	3,462	2.8	11.3
7	Cerebrovascular diseases (I60-I69)	3,355	2.7	10.9
8	Chronic lower respiratory diseases (J40-J47)	2,695	2.2	8.8
9	Viral hepatitis (B15-B19)	1,517	1.2	4.9
10	Septicemia (A40-A41)	1,359	1.1	4.4
11	Human immunodeficiency virus (HIV) disease (B20-B24)	1,329	1.1	4.3
12	Influenza and pneumonia (J10-J18)	1,242	1.0	4.0
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,005	0.8	3.3
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	933	0.8	3.0
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	555	0.5	1.8
...	All other causes (Residual)	19,536	15.9	63.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 55-64 years				
...	All causes	198,842	100.0	875.1
1	Malignant neoplasms (C00-C97)	76,400	38.4	336.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	46,934	23.6	206.6
3	Chronic lower respiratory diseases (J40-J47)	10,190	5.1	44.8
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,390	3.7	32.5
5	Diabetes mellitus (E10-E14)	6,965	3.5	30.7
6	Cerebrovascular diseases (I60-I69)	6,321	3.2	27.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	4,838	2.4	21.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,584	1.8	15.8
9	Septicemia (A40-A41)	2,513	1.3	11.1
10	Influenza and pneumonia (J10-J18)	2,295	1.2	10.1
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,238	1.1	9.8
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,008	0.5	4.4
13	Aortic aneurysm and dissection (I71)	985	0.5	4.3
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	960	0.5	4.2
15	Viral hepatitis (B15-B19)	766	0.4	3.4
...	All other causes (Residual)	25,455	12.8	112.0
White, non-Hispanic, both sexes, 65-74 years				
...	All causes	317,606	100.0	2,152.5
1	Malignant neoplasms (C00-C97)	114,148	35.9	773.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	77,652	24.4	526.3
3	Chronic lower respiratory diseases (J40-J47)	25,565	8.0	173.3
4	Cerebrovascular diseases (I60-I69)	14,563	4.6	98.7
5	Diabetes mellitus (E10-E14)	10,875	3.4	73.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,532	2.1	44.3
7	Influenza and pneumonia (J10-J18)	4,894	1.5	33.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,745	1.5	32.2
9	Septicemia (A40-A41)	4,328	1.4	29.3
10	Chronic liver disease and cirrhosis (K70,K73-K74)	4,003	1.3	27.1
11	Alzheimer's disease (G30)	3,107	1.0	21.1
12	Aortic aneurysm and dissection (I71)	2,533	0.8	17.2
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,043	0.6	13.8
14	Parkinson's disease (G20-G21)	1,988	0.6	13.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,949	0.6	13.2
...	All other causes (Residual)	38,681	12.2	262.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 75-84 years				
...	All causes	582,454	100.0	5,337.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	165,259	28.4	1,514.3
2	Malignant neoplasms (C00-C97)	142,461	24.5	1,305.4
3	Chronic lower respiratory diseases (J40-J47)	43,613	7.5	399.6
4	Cerebrovascular diseases (I60-I69)	41,670	7.2	381.8
5	Alzheimer's disease (G30)	19,438	3.3	178.1
6	Diabetes mellitus (E10-E14)	17,220	3.0	157.8
7	Influenza and pneumonia (J10-J18)	15,143	2.6	138.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,774	2.0	107.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11,106	1.9	101.8
10	Septicemia (A40-A41)	8,426	1.4	77.2
11	Parkinson's disease (G20-G21)	7,999	1.4	73.3
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	5,098	0.9	46.7
13	Pneumonitis due to solids and liquids (J69)	4,811	0.8	44.1
14	Aortic aneurysm and dissection (I71)	4,573	0.8	41.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,081	0.7	37.4
...	All other causes (Residual)	79,782	13.7	731.1
White, non-Hispanic, both sexes, 85 years and over				
...	All causes	593,382	100.0	4,191.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	210,669	35.5	5,038.3
2	Malignant neoplasms (C00-C97)	70,283	11.8	1,680.9
3	Cerebrovascular diseases (I60-I69)	53,371	9.0	1,276.4
4	Alzheimer's disease (G30)	36,040	6.1	861.9
5	Chronic lower respiratory diseases (J40-J47)	26,761	4.5	640.0
6	Influenza and pneumonia (J10-J18)	24,829	4.2	593.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,344	2.1	295.2
8	Diabetes mellitus (E10-E14)	11,896	2.0	284.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11,796	2.0	282.1
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	7,875	1.3	188.3
11	Septicemia (A40-A41)	7,509	1.3	179.6
12	Pneumonitis due to solids and liquids (J69)	7,124	1.2	170.4
13	Parkinson's disease (G20-G21)	6,083	1.0	145.5
14	Atherosclerosis (I70)	6,064	1.0	145.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3,413	0.6	81.6
...	All other causes (Residual)	97,325	16.4	2,327.6

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, all ages				
...	All causes	938,143	100.0	957.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	263,638	28.1	269.1
2	Malignant neoplasms (C00-C97)	234,392	25.0	239.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	53,465	5.7	54.6
4	Chronic lower respiratory diseases (J40-J47)	51,909	5.5	53.0
5	Cerebrovascular diseases (I60-I69)	46,087	4.9	47.0
6	Diabetes mellitus (E10-E14)	25,773	2.7	26.3
7	Influenza and pneumonia (J10-J18)	21,905	2.3	22.4
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	21,214	2.3	21.6
9	Alzheimer's disease (G30)	16,989	1.8	17.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15,247	1.6	15.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	12,931	1.4	13.2
12	Septicemia (A40-A41)	11,188	1.2	11.4
13	Parkinson's disease (G20-G21)	9,469	1.0	9.7
14	Pneumonitis due to solids and liquids (J69)	7,367	0.8	7.5
15	Aortic aneurysm and dissection (I71)	6,906	0.7	7.0
...	All other causes (Residual)	139,663	14.9	142.5
White, non-Hispanic, male, 1-4 years				
...	All causes	1,365	100.0	29.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	553	40.5	11.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	145	10.6	3.1
3	Malignant neoplasms (C00-C97)	134	9.8	2.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	73	5.3	1.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	37	2.7	0.8
6	Influenza and pneumonia (J10-J18)	23	1.7	0.5
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	21	1.5	0.4
8	Certain conditions originating in the perinatal period (P00-P96)	17	1.2	*
9	Septicemia (A40-A41)	15	1.1	*
9	Cerebrovascular diseases (I60-I69)	15	1.1	*
11	Meningitis (G00,G03)	10	0.7	*
12	Meningococcal infection (A39)	8	0.6	*
12	Chronic lower respiratory diseases (J40-J47)	8	0.6	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.3	*
...	All other causes (Residual)	296	21.7	6.3

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 5-14 years				
...	All causes	2,264	100.0	17.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	991	43.8	7.9
2	Malignant neoplasms (C00-C97)	339	15.0	2.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	119	5.3	0.9
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	111	4.9	0.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	70	3.1	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	67	3.0	0.5
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	29	1.3	0.2
8	Cerebrovascular diseases (I60-I69)	24	1.1	0.2
9	Chronic lower respiratory diseases (J40-J47)	23	1.0	0.2
10	Influenza and pneumonia (J10-J18)	22	1.0	0.2
11	Septicemia (A40-A41)	20	0.9	0.2
12	Certain conditions originating in the perinatal period (P00-P96)	11	0.5	*
13	Meningitis (G00,G03)	9	0.4	*
14	Meningococcal infection (A39)	7	0.3	*
14	Diabetes mellitus (E10-E14)	7	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.3	*
...	All other causes (Residual)	408	18.0	3.2
White, non-Hispanic, male, 15-24 years				
...	All causes	14,017	100.0	104.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,885	56.3	58.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,535	18.1	19.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	624	4.5	4.7
4	Malignant neoplasms (C00-C97)	593	4.2	4.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	363	2.6	2.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	156	1.1	1.2
7	Influenza and pneumonia (J10-J18)	73	0.5	0.5
8	Cerebrovascular diseases (I60-I69)	63	0.4	0.5
9	Diabetes mellitus (E10-E14)	45	0.3	0.3
9	Chronic lower respiratory diseases (J40-J47)	45	0.3	0.3
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	41	0.3	0.3
12	Septicemia (A40-A41)	39	0.3	0.3
13	Legal intervention (Y35,Y89.0)	26	0.2	0.2
14	Aortic aneurysm and dissection (I71)	22	0.2	0.2
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21	0.1	0.2
...	All other causes (Residual)	1,486	10.6	11.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 25-34 years				
...	All causes	16,098	100.0	130.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,458	40.1	52.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,064	19.0	24.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,194	7.4	9.7
4	Malignant neoplasms (C00-C97)	1,144	7.1	9.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	672	4.2	5.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	227	1.4	1.8
7	Diabetes mellitus (E10-E14)	191	1.2	1.6
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	168	1.0	1.4
9	Cerebrovascular diseases (I60-I69)	137	0.9	1.1
10	Chronic liver disease and cirrhosis (K70,K73-K74)	121	0.8	1.0
11	Influenza and pneumonia (J10-J18)	115	0.7	0.9
12	Septicemia (A40-A41)	85	0.5	0.7
13	Chronic lower respiratory diseases (J40-J47)	63	0.4	0.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	59	0.4	0.5
15	Aortic aneurysm and dissection (I71)	52	0.3	0.4
...	All other causes (Residual)	2,348	14.6	19.1
White, non-Hispanic, male, 35-44 years				
...	All causes	35,253	100.0	235.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,412	23.9	56.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,080	17.2	40.6
3	Malignant neoplasms (C00-C97)	4,467	12.7	29.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,249	12.1	28.4
5	Human immunodeficiency virus (HIV) disease (B20-B24)	1,259	3.6	8.4
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1,240	3.5	8.3
7	Diabetes mellitus (E10-E14)	795	2.3	5.3
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	751	2.1	5.0
9	Cerebrovascular diseases (I60-I69)	653	1.9	4.4
10	Influenza and pneumonia (J10-J18)	300	0.9	2.0
11	Chronic lower respiratory diseases (J40-J47)	270	0.8	1.8
12	Viral hepatitis (B15-B19)	251	0.7	1.7
13	Septicemia (A40-A41)	243	0.7	1.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	228	0.6	1.5
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	205	0.6	1.4
...	All other causes (Residual)	5,850	16.6	39.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 45-54 years				
...	All causes	77,741	100.0	510.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	19,329	24.9	126.9
2	Malignant neoplasms (C00-C97)	18,556	23.9	121.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,935	11.5	58.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,465	5.7	29.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	3,765	4.8	24.7
6	Diabetes mellitus (E10-E14)	2,184	2.8	14.3
7	Cerebrovascular diseases (I60-I69)	1,841	2.4	12.1
8	Chronic lower respiratory diseases (J40-J47)	1,398	1.8	9.2
9	Viral hepatitis (B15-B19)	1,132	1.5	7.4
10	Human immunodeficiency virus (HIV) disease (B20-B24)	1,130	1.5	7.4
11	Influenza and pneumonia (J10-J18)	741	1.0	4.9
12	Septicemia (A40-A41)	724	0.9	4.8
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	649	0.8	4.3
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	586	0.8	3.8
15	Aortic aneurysm and dissection (I71)	367	0.5	2.4
...	All other causes (Residual)	11,939	15.4	78.4
White, non-Hispanic, male, 55-64 years				
...	All causes	119,171	100.0	1,076.4
1	Malignant neoplasms (C00-C97)	41,468	34.8	374.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	33,078	27.8	298.8
3	Chronic lower respiratory diseases (J40-J47)	5,126	4.3	46.3
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,955	4.2	44.8
5	Diabetes mellitus (E10-E14)	4,128	3.5	37.3
6	Cerebrovascular diseases (I60-I69)	3,447	2.9	31.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	3,387	2.8	30.6
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,754	2.3	24.9
9	Influenza and pneumonia (J10-J18)	1,328	1.1	12.0
10	Septicemia (A40-A41)	1,299	1.1	11.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,234	1.0	11.1
12	Aortic aneurysm and dissection (I71)	730	0.6	6.6
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	585	0.5	5.3
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	572	0.5	5.2
15	Viral hepatitis (B15-B19)	536	0.4	4.8
...	All other causes (Residual)	14,544	12.2	131.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 65-74 years				
...	All causes	178,642	100.0	2,617.9
1	Malignant neoplasms (C00-C97)	62,610	35.0	917.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	48,417	27.1	709.5
3	Chronic lower respiratory diseases (J40-J47)	12,865	7.2	188.5
4	Cerebrovascular diseases (I60-I69)	7,505	4.2	110.0
5	Diabetes mellitus (E10-E14)	5,984	3.3	87.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,027	2.3	59.0
7	Influenza and pneumonia (J10-J18)	2,750	1.5	40.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,540	1.4	37.2
9	Chronic liver disease and cirrhosis (K70,K73-K74)	2,463	1.4	36.1
10	Septicemia (A40-A41)	2,213	1.2	32.4
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,701	1.0	24.9
12	Aortic aneurysm and dissection (I71)	1,695	0.9	24.8
13	Alzheimer's disease (G30)	1,335	0.7	19.6
14	Parkinson's disease (G20-G21)	1,313	0.7	19.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,128	0.6	16.5
...	All other causes (Residual)	20,096	11.2	294.5
White, non-Hispanic, male, 75-84 years				
...	All causes	285,752	100.0	6,461.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	84,694	29.6	1,915.1
2	Malignant neoplasms (C00-C97)	74,178	26.0	1,677.3
3	Chronic lower respiratory diseases (J40-J47)	21,076	7.4	476.6
4	Cerebrovascular diseases (I60-I69)	17,553	6.1	396.9
5	Diabetes mellitus (E10-E14)	8,276	2.9	187.1
6	Influenza and pneumonia (J10-J18)	7,549	2.6	170.7
7	Alzheimer's disease (G30)	6,945	2.4	157.0
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,140	2.1	138.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,721	2.0	129.4
10	Parkinson's disease (G20-G21)	4,864	1.7	110.0
11	Septicemia (A40-A41)	3,914	1.4	88.5
12	Pneumonitis due to solids and liquids (J69)	2,664	0.9	60.2
13	Aortic aneurysm and dissection (I71)	2,579	0.9	58.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,111	0.7	47.7
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,075	0.7	46.9
...	All other causes (Residual)	35,413	12.4	800.8

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 85 years and over				
...	All causes	200,357	100.0	5,489.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	70,250	35.1	5,430.9
2	Malignant neoplasms (C00-C97)	30,878	15.4	2,387.1
3	Cerebrovascular diseases (I60-I69)	14,815	7.4	1,145.3
4	Chronic lower respiratory diseases (J40-J47)	11,027	5.5	852.5
5	Influenza and pneumonia (J10-J18)	8,939	4.5	691.1
6	Alzheimer's disease (G30)	8,451	4.2	653.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,825	2.4	373.0
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,745	2.4	366.8
9	Diabetes mellitus (E10-E14)	4,161	2.1	321.7
10	Pneumonitis due to solids and liquids (J69)	3,200	1.6	247.4
11	Parkinson's disease (G20-G21)	3,056	1.5	236.3
12	Septicemia (A40-A41)	2,568	1.3	198.5
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,064	1.0	159.6
14	Atherosclerosis (I70)	1,769	0.9	136.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,366	0.7	105.6
...	All other causes (Residual)	28,243	14.1	2,183.4
White, non-Hispanic, female, all ages				
...	All causes	995,239	100.0	977.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	273,874	27.5	269.1
2	Malignant neoplasms (C00-C97)	219,142	22.0	215.3
3	Cerebrovascular diseases (I60-I69)	74,978	7.5	73.7
4	Chronic lower respiratory diseases (J40-J47)	57,761	5.8	56.7
5	Alzheimer's disease (G30)	42,164	4.2	41.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	32,001	3.2	31.4
7	Influenza and pneumonia (J10-J18)	27,551	2.8	27.1
8	Diabetes mellitus (E10-E14)	26,328	2.6	25.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16,261	1.6	16.0
10	Septicemia (A40-A41)	13,853	1.4	13.6
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	10,422	1.0	10.2
12	Chronic liver disease and cirrhosis (K70,K73-K74)	7,213	0.7	7.1
13	Pneumonitis due to solids and liquids (J69)	7,044	0.7	6.9
14	Parkinson's disease (G20-G21)	6,959	0.7	6.8
15	Atherosclerosis (I70)	6,534	0.7	6.4
...	All other causes (Residual)	173,154	17.4	170.1

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	White, non-Hispanic, female, 1-4 years			
...	All causes	1,093	100.0	24.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	380	34.8	8.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	125	11.4	2.8
3	Malignant neoplasms (C00-C97)	110	10.1	2.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	58	5.3	1.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	43	3.9	1.0
6	Influenza and pneumonia (J10-J18)	32	2.9	0.7
7	Septicemia (A40-A41)	21	1.9	0.5
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	1.6	*
9	Cerebrovascular diseases (I60-I69)	10	0.9	*
10	Certain conditions originating in the perinatal period (P00-P96)	7	0.6	*
11	Meningitis (G00,G03)	6	0.5	*
12	Anemias (D50-D64)	5	0.5	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.5	*
14	Chronic lower respiratory diseases (J40-J47)	4	0.4	*
15	Infections of kidney (N10-N12,N13.6,N15.1)	3	0.3	*
...	All other causes (Residual)	266	24.3	5.9
	White, non-Hispanic, female, 5-14 years			
...	All causes	1,556	100.0	13.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	582	37.4	4.9
2	Malignant neoplasms (C00-C97)	248	15.9	2.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	106	6.8	0.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	61	3.9	0.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	58	3.7	0.5
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	49	3.1	0.4
7	Cerebrovascular diseases (I60-I69)	24	1.5	0.2
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	1.5	0.2
9	Septicemia (A40-A41)	21	1.3	0.2
9	Influenza and pneumonia (J10-J18)	21	1.3	0.2
11	Chronic lower respiratory diseases (J40-J47)	18	1.2	*
12	Diabetes mellitus (E10-E14)	12	0.8	*
13	Anemias (D50-D64)	11	0.7	*
14	Meningitis (G00,G03)	8	0.5	*
15	Certain conditions originating in the perinatal period (P00-P96)	7	0.4	*
...	All other causes (Residual)	307	19.7	2.6

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 15-24 years				
...	All causes	5,667	100.0	44.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,962	52.3	23.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	510	9.0	4.0
3	Malignant neoplasms (C00-C97)	423	7.5	3.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	293	5.2	2.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	190	3.4	1.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	134	2.4	1.0
7	Pregnancy, childbirth and the puerperium (O00-099)	60	1.1	0.5
8	Cerebrovascular diseases (I60-I69)	54	1.0	0.4
9	Diabetes mellitus (E10-E14)	42	0.7	0.3
10	Influenza and pneumonia (J10-J18)	35	0.6	0.3
11	Septicemia (A40-A41)	31	0.5	0.2
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	31	0.5	0.2
13	Chronic lower respiratory diseases (J40-J47)	30	0.5	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20	0.4	0.2
15	Complications of medical and surgical care (Y40-Y84,Y88)	15	0.3	*
...	All other causes (Residual)	837	14.8	6.5
White, non-Hispanic, female, 25-34 years				
...	All causes	7,299	100.0	60.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,254	30.9	18.6
2	Malignant neoplasms (C00-C97)	1,126	15.4	9.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	683	9.4	5.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	551	7.5	4.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	304	4.2	2.5
6	Cerebrovascular diseases (I60-I69)	143	2.0	1.2
7	Diabetes mellitus (E10-E14)	142	1.9	1.2
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	127	1.7	1.1
9	Pregnancy, childbirth and the puerperium (O00-099)	122	1.7	1.0
10	Septicemia (A40-A41)	87	1.2	0.7
11	Human immunodeficiency virus (HIV) disease (B20-B24)	69	0.9	0.6
12	Influenza and pneumonia (J10-J18)	68	0.9	0.6
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	61	0.8	0.5
14	Chronic lower respiratory diseases (J40-J47)	57	0.8	0.5
14	Chronic liver disease and cirrhosis (K70,K73-K74)	57	0.8	0.5
...	All other causes (Residual)	1,448	19.8	12.0

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 35-44 years				
...	All causes	20,290	100.0	136.0
1	Malignant neoplasms (C00-C97)	5,595	27.6	37.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,613	17.8	24.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,331	11.5	15.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,349	6.6	9.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	623	3.1	4.2
6	Cerebrovascular diseases (I60-I69)	603	3.0	4.0
7	Diabetes mellitus (E10-E14)	443	2.2	3.0
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	373	1.8	2.5
9	Chronic lower respiratory diseases (J40-J47)	313	1.5	2.1
10	Human immunodeficiency virus (HIV) disease (B20-B24)	270	1.3	1.8
11	Influenza and pneumonia (J10-J18)	252	1.2	1.7
12	Septicemia (A40-A41)	231	1.1	1.5
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	166	0.8	1.1
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	164	0.8	1.1
15	Viral hepatitis (B15-B19)	139	0.7	0.9
...	All other causes (Residual)	3,825	18.9	25.6
White, non-Hispanic, female, 45-54 years				
...	All causes	45,296	100.0	292.9
1	Malignant neoplasms (C00-C97)	17,460	38.5	112.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,529	14.4	42.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,809	8.4	24.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,636	3.6	10.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,565	3.5	10.1
6	Cerebrovascular diseases (I60-I69)	1,514	3.3	9.8
7	Chronic lower respiratory diseases (J40-J47)	1,297	2.9	8.4
8	Diabetes mellitus (E10-E14)	1,278	2.8	8.3
9	Septicemia (A40-A41)	635	1.4	4.1
10	Influenza and pneumonia (J10-J18)	501	1.1	3.2
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	419	0.9	2.7
12	Viral hepatitis (B15-B19)	385	0.8	2.5
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	284	0.6	1.8
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	252	0.6	1.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	224	0.5	1.4
...	All other causes (Residual)	7,508	16.6	48.5

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	White, non-Hispanic, female, 55-64 years			
...	All causes	79,671	100.0	683.8
1	Malignant neoplasms (C00-C97)	34,932	43.8	299.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,856	17.4	118.9
3	Chronic lower respiratory diseases (J40-J47)	5,064	6.4	43.5
4	Cerebrovascular diseases (I60-I69)	2,874	3.6	24.7
5	Diabetes mellitus (E10-E14)	2,837	3.6	24.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,435	3.1	20.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,451	1.8	12.5
8	Septicemia (A40-A41)	1,214	1.5	10.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,004	1.3	8.6
10	Influenza and pneumonia (J10-J18)	967	1.2	8.3
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	830	1.0	7.1
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	423	0.5	3.6
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	388	0.5	3.3
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	269	0.3	2.3
15	Aortic aneurysm and dissection (I71)	255	0.3	2.2
...	All other causes (Residual)	10,872	13.6	93.3
	White, non-Hispanic, female, 65-74 years			
...	All causes	138,964	100.0	1,752.0
1	Malignant neoplasms (C00-C97)	51,538	37.1	649.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	29,235	21.0	368.6
3	Chronic lower respiratory diseases (J40-J47)	12,700	9.1	160.1
4	Cerebrovascular diseases (I60-I69)	7,058	5.1	89.0
5	Diabetes mellitus (E10-E14)	4,891	3.5	61.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,505	1.8	31.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,205	1.6	27.8
8	Influenza and pneumonia (J10-J18)	2,144	1.5	27.0
9	Septicemia (A40-A41)	2,115	1.5	26.7
10	Alzheimer's disease (G30)	1,772	1.3	22.3
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1,540	1.1	19.4
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	972	0.7	12.3
13	Aortic aneurysm and dissection (I71)	838	0.6	10.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	821	0.6	10.4
15	Parkinson's disease (G20-G21)	675	0.5	8.5
...	All other causes (Residual)	17,955	12.9	226.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	White, non-Hispanic, female, 75-84 years			
...	All causes	296,702	100.0	4,571.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	80,565	27.2	1,241.2
2	Malignant neoplasms (C00-C97)	68,283	23.0	1,052.0
3	Cerebrovascular diseases (I60-I69)	24,117	8.1	371.6
4	Chronic lower respiratory diseases (J40-J47)	22,537	7.6	347.2
5	Alzheimer's disease (G30)	12,493	4.2	192.5
6	Diabetes mellitus (E10-E14)	8,944	3.0	137.8
7	Influenza and pneumonia (J10-J18)	7,594	2.6	117.0
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,634	1.9	86.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,385	1.8	83.0
10	Septicemia (A40-A41)	4,512	1.5	69.5
11	Parkinson's disease (G20-G21)	3,135	1.1	48.3
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	3,023	1.0	46.6
13	Pneumonitis due to solids and liquids (J69)	2,147	0.7	33.1
14	Aortic aneurysm and dissection (I71)	1,994	0.7	30.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,970	0.7	30.4
...	All other causes (Residual)	44,369	15.0	683.6
	White, non-Hispanic, female, 85 years and over			
...	All causes	393,025	100.0	3,609.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	140,419	35.7	4,862.4
2	Malignant neoplasms (C00-C97)	39,405	10.0	1,364.5
3	Cerebrovascular diseases (I60-I69)	38,556	9.8	1,335.1
4	Alzheimer's disease (G30)	27,589	7.0	955.3
5	Influenza and pneumonia (J10-J18)	15,890	4.0	550.2
6	Chronic lower respiratory diseases (J40-J47)	15,734	4.0	544.8
7	Diabetes mellitus (E10-E14)	7,735	2.0	267.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,599	1.9	263.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,971	1.8	241.4
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	5,811	1.5	201.2
11	Septicemia (A40-A41)	4,941	1.3	171.1
12	Atherosclerosis (I70)	4,295	1.1	148.7
13	Pneumonitis due to solids and liquids (J69)	3,924	1.0	135.9
14	Parkinson's disease (G20-G21)	3,027	0.8	104.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,047	0.5	70.9
...	All other causes (Residual)	69,082	17.6	2,392.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, all ages				
...	All causes	283,859	100.0	768.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	73,373	25.8	198.7
2	Malignant neoplasms (C00-C97)	61,842	21.8	167.5
3	Cerebrovascular diseases (I60-I69)	17,980	6.3	48.7
4	Diabetes mellitus (E10-E14)	12,685	4.5	34.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,446	4.4	33.7
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8,000	2.8	21.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,759	2.7	21.0
8	Chronic lower respiratory diseases (J40-J47)	7,331	2.6	19.9
9	Human immunodeficiency virus (HIV) disease (B20-B24)	7,163	2.5	19.4
10	Septicemia (A40-A41)	5,945	2.1	16.1
11	Influenza and pneumonia (J10-J18)	5,620	2.0	15.2
12	Certain conditions originating in the perinatal period (P00-P96)	4,740	1.7	12.8
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	4,709	1.7	12.8
14	Alzheimer's disease (G30)	4,196	1.5	11.4
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,506	0.9	6.8
...	All other causes (Residual)	47,564	16.8	128.8
Black, non-Hispanic, both sexes, 1-4 years				
...	All causes	1,128	100.0	45.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	317	28.1	12.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	154	13.7	6.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	127	11.3	5.1
4	Malignant neoplasms (C00-C97)	62	5.5	2.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	61	5.4	2.5
6	Influenza and pneumonia (J10-J18)	34	3.0	1.4
7	Certain conditions originating in the perinatal period (P00-P96)	23	2.0	0.9
8	Chronic lower respiratory diseases (J40-J47)	21	1.9	0.8
9	Septicemia (A40-A41)	19	1.7	*
9	Anemias (D50-D64)	19	1.7	*
11	Cerebrovascular diseases (I60-I69)	12	1.1	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.6	*
13	Acute bronchitis and bronchiolitis (J20-J21)	6	0.5	*
14	Meningitis (G00,G03)	5	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.4	*
...	All other causes (Residual)	257	22.8	10.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 5-14 years				
...	All causes	1,551	100.0	24.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	557	35.9	8.8
2	Malignant neoplasms (C00-C97)	165	10.6	2.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	130	8.4	2.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	82	5.3	1.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	78	5.0	1.2
6	Chronic lower respiratory diseases (J40-J47)	64	4.1	1.0
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	49	3.2	0.8
8	Anemias (D50-D64)	23	1.5	0.4
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	21	1.4	0.3
10	Influenza and pneumonia (J10-J18)	20	1.3	0.3
11	Human immunodeficiency virus (HIV) disease (B20-B24)	17	1.1	*
11	Cerebrovascular diseases (I60-I69)	17	1.1	*
13	Diabetes mellitus (E10-E14)	16	1.0	*
14	Septicemia (A40-A41)	12	0.8	*
15	Certain conditions originating in the perinatal period (P00-P96)	9	0.6	*
...	All other causes (Residual)	291	18.8	4.6
Black, non-Hispanic, both sexes, 15-24 years				
...	All causes	6,924	100.0	112.8
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,771	40.0	45.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,614	23.3	26.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	450	6.5	7.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	310	4.5	5.0
5	Malignant neoplasms (C00-C97)	307	4.4	5.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	141	2.0	2.3
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	111	1.6	1.8
8	Anemias (D50-D64)	90	1.3	1.5
9	Chronic lower respiratory diseases (J40-J47)	79	1.1	1.3
10	Pregnancy, childbirth and the puerperium (O00-099)	73	1.1	1.2
11	Diabetes mellitus (E10-E14)	60	0.9	1.0
12	Cerebrovascular diseases (I60-I69)	48	0.7	0.8
13	Septicemia (A40-A41)	45	0.6	0.7
14	Influenza and pneumonia (J10-J18)	40	0.6	0.7
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	31	0.4	0.5
...	All other causes (Residual)	754	10.9	12.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 25-34 years				
...	All causes	9,768	100.0	186.5
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,405	24.6	45.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,704	17.4	32.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	971	9.9	18.5
4	Human immunodeficiency virus (HIV) disease (B20-B24)	936	9.6	17.9
5	Malignant neoplasms (C00-C97)	651	6.7	12.4
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	510	5.2	9.7
7	Diabetes mellitus (E10-E14)	180	1.8	3.4
8	Cerebrovascular diseases (I60-I69)	173	1.8	3.3
9	Anemias (D50-D64)	126	1.3	2.4
10	Pregnancy, childbirth and the puerperium (O00-099)	106	1.1	2.0
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	103	1.1	2.0
12	Septicemia (A40-A41)	102	1.0	1.9
13	Chronic lower respiratory diseases (J40-J47)	101	1.0	1.9
14	Influenza and pneumonia (J10-J18)	63	0.6	1.2
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	59	0.6	1.1
...	All other causes (Residual)	1,578	16.2	30.1
Black, non-Hispanic, both sexes, 35-44 years				
...	All causes	18,237	100.0	332.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,202	17.6	58.3
2	Malignant neoplasms (C00-C97)	2,649	14.5	48.3
3	Human immunodeficiency virus (HIV) disease (B20-B24)	2,522	13.8	46.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,072	11.4	37.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,276	7.0	23.3
6	Cerebrovascular diseases (I60-I69)	698	3.8	12.7
7	Diabetes mellitus (E10-E14)	540	3.0	9.8
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	369	2.0	6.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	316	1.7	5.8
10	Chronic liver disease and cirrhosis (K70,K73-K74)	307	1.7	5.6
11	Septicemia (A40-A41)	255	1.4	4.6
12	Influenza and pneumonia (J10-J18)	223	1.2	4.1
13	Chronic lower respiratory diseases (J40-J47)	220	1.2	4.0
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	177	1.0	3.2
15	Anemias (D50-D64)	139	0.8	2.5
...	All other causes (Residual)	3,272	17.9	59.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 45-54 years				
...	All causes	36,247	100.0	760.8
1	Malignant neoplasms (C00-C97)	8,661	23.9	181.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,572	23.6	179.9
3	Human immunodeficiency virus (HIV) disease (B20-B24)	2,423	6.7	50.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,418	6.7	50.7
5	Cerebrovascular diseases (I60-I69)	1,905	5.3	40.0
6	Diabetes mellitus (E10-E14)	1,371	3.8	28.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	890	2.5	18.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	813	2.2	17.1
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	732	2.0	15.4
10	Septicemia (A40-A41)	650	1.8	13.6
11	Chronic lower respiratory diseases (J40-J47)	621	1.7	13.0
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	497	1.4	10.4
13	Influenza and pneumonia (J10-J18)	443	1.2	9.3
14	Viral hepatitis (B15-B19)	362	1.0	7.6
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	279	0.8	5.9
...	All other causes (Residual)	5,610	15.5	117.7
Black, non-Hispanic, both sexes, 55-64 years				
...	All causes	42,998	100.0	1,527.9
1	Malignant neoplasms (C00-C97)	13,149	30.6	467.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,541	26.8	410.1
3	Cerebrovascular diseases (I60-I69)	2,420	5.6	86.0
4	Diabetes mellitus (E10-E14)	2,304	5.4	81.9
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,283	3.0	45.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,251	2.9	44.5
7	Chronic lower respiratory diseases (J40-J47)	1,101	2.6	39.1
8	Septicemia (A40-A41)	906	2.1	32.2
9	Human immunodeficiency virus (HIV) disease (B20-B24)	841	2.0	29.9
10	Chronic liver disease and cirrhosis (K70,K73-K74)	728	1.7	25.9
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	685	1.6	24.3
12	Influenza and pneumonia (J10-J18)	537	1.2	19.1
13	Viral hepatitis (B15-B19)	247	0.6	8.8
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	247	0.6	8.8
15	Aortic aneurysm and dissection (I71)	215	0.5	7.6
...	All other causes (Residual)	5,543	12.9	197.0

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 65-74 years				
...	All causes	51,126	100.0	3,032.3
1	Malignant neoplasms (C00-C97)	15,712	30.7	931.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,372	28.1	852.4
3	Cerebrovascular diseases (I60-I69)	3,362	6.6	199.4
4	Diabetes mellitus (E10-E14)	3,078	6.0	182.6
5	Chronic lower respiratory diseases (J40-J47)	1,751	3.4	103.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,629	3.2	96.6
7	Septicemia (A40-A41)	1,203	2.4	71.4
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	890	1.7	52.8
9	Influenza and pneumonia (J10-J18)	850	1.7	50.4
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	818	1.6	48.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	371	0.7	22.0
12	Alzheimer's disease (G30)	327	0.6	19.4
13	Pneumonitis due to solids and liquids (J69)	239	0.5	14.2
14	Human immunodeficiency virus (HIV) disease (B20-B24)	229	0.4	13.6
15	Aortic aneurysm and dissection (I71)	224	0.4	13.3
...	All other causes (Residual)	6,071	11.9	360.1
Black, non-Hispanic, both sexes, 75-84 years				
...	All causes	60,748	100.0	6,251.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	18,139	29.9	1,866.7
2	Malignant neoplasms (C00-C97)	14,244	23.4	1,465.9
3	Cerebrovascular diseases (I60-I69)	5,044	8.3	519.1
4	Diabetes mellitus (E10-E14)	3,269	5.4	336.4
5	Chronic lower respiratory diseases (J40-J47)	2,195	3.6	225.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,993	3.3	205.1
7	Influenza and pneumonia (J10-J18)	1,539	2.5	158.4
8	Septicemia (A40-A41)	1,534	2.5	157.9
9	Alzheimer's disease (G30)	1,510	2.5	155.4
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,214	2.0	124.9
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	857	1.4	88.2
12	Pneumonitis due to solids and liquids (J69)	463	0.8	47.6
13	Aortic aneurysm and dissection (I71)	315	0.5	32.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	311	0.5	32.0
15	Parkinson's disease (G20-G21)	290	0.5	29.8
...	All other causes (Residual)	7,831	12.9	805.9

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 85 years and over				
...	All causes	46,877	100.0	3,492.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	16,018	34.2	4,610.3
2	Malignant neoplasms (C00-C97)	6,234	13.3	1,794.3
3	Cerebrovascular diseases (I60-I69)	4,260	9.1	1,226.1
4	Alzheimer's disease (G30)	2,317	4.9	666.9
5	Diabetes mellitus (E10-E14)	1,864	4.0	536.5
6	Influenza and pneumonia (J10-J18)	1,778	3.8	511.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,519	3.2	437.2
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,180	2.5	339.6
9	Chronic lower respiratory diseases (J40-J47)	1,167	2.5	335.9
10	Septicemia (A40-A41)	1,128	2.4	324.7
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	539	1.1	155.1
12	Pneumonitis due to solids and liquids (J69)	512	1.1	147.4
13	Atherosclerosis (I70)	360	0.8	103.6
14	Parkinson's disease (G20-G21)	214	0.5	61.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	201	0.4	57.9
...	All other causes (Residual)	7,586	16.2	2,183.4
Black, non-Hispanic, male, all ages				
...	All causes	144,022	100.0	818.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	35,749	24.8	203.2
2	Malignant neoplasms (C00-C97)	32,089	22.3	182.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,430	5.9	47.9
4	Cerebrovascular diseases (I60-I69)	7,586	5.3	43.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6,730	4.7	38.3
6	Diabetes mellitus (E10-E14)	5,498	3.8	31.3
7	Human immunodeficiency virus (HIV) disease (B20-B24)	4,693	3.3	26.7
8	Chronic lower respiratory diseases (J40-J47)	4,010	2.8	22.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,505	2.4	19.9
10	Influenza and pneumonia (J10-J18)	2,729	1.9	15.5
11	Septicemia (A40-A41)	2,660	1.8	15.1
12	Certain conditions originating in the perinatal period (P00-P96)	2,653	1.8	15.1
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,945	1.4	11.1
14	Chronic liver disease and cirrhosis (K70,K73-K74)	1,650	1.1	9.4
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,614	1.1	9.2
...	All other causes (Residual)	22,481	15.6	127.8

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 1-4 years				
...	All causes	623	100.0	49.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	182	29.2	14.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	79	12.7	6.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	74	11.9	5.9
4	Malignant neoplasms (C00-C97)	35	5.6	2.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	32	5.1	2.5
6	Chronic lower respiratory diseases (J40-J47)	14	2.2	*
7	Certain conditions originating in the perinatal period (P00-P96)	13	2.1	*
8	Anemias (D50-D64)	12	1.9	*
8	Influenza and pneumonia (J10-J18)	12	1.9	*
10	Septicemia (A40-A41)	9	1.4	*
11	Cerebrovascular diseases (I60-I69)	5	0.8	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.6	*
12	Acute bronchitis and bronchiolitis (J20-J21)	4	0.6	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.5	*
15	Meningitis (G00,G03)	2	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.3	*
...	All other causes (Residual)	141	22.6	11.2
Black, non-Hispanic, male, 5-14 years				
...	All causes	870	100.0	26.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	343	39.4	10.6
2	Malignant neoplasms (C00-C97)	82	9.4	2.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	81	9.3	2.5
4	Chronic lower respiratory diseases (J40-J47)	44	5.1	1.4
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	38	4.4	1.2
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	35	4.0	1.1
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	30	3.4	0.9
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	1.4	*
8	Anemias (D50-D64)	12	1.4	*
8	Influenza and pneumonia (J10-J18)	12	1.4	*
11	Cerebrovascular diseases (I60-I69)	10	1.1	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	8	0.9	*
12	Diabetes mellitus (E10-E14)	8	0.9	*
14	Certain conditions originating in the perinatal period (P00-P96)	5	0.6	*
15	Septicemia (A40-A41)	4	0.5	*
...	All other causes (Residual)	146	16.8	4.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 15-24 years				
...	All causes	5,239	100.0	169.4
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,495	47.6	80.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,191	22.7	38.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	382	7.3	12.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	202	3.9	6.5
5	Malignant neoplasms (C00-C97)	174	3.3	5.6
6	Human immunodeficiency virus (HIV) disease (B20-B24)	73	1.4	2.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	73	1.4	2.4
8	Anemias (D50-D64)	50	1.0	1.6
9	Chronic lower respiratory diseases (J40-J47)	48	0.9	1.6
10	Diabetes mellitus (E10-E14)	30	0.6	1.0
10	Cerebrovascular diseases (I60-I69)	30	0.6	1.0
12	Septicemia (A40-A41)	22	0.4	0.7
13	Influenza and pneumonia (J10-J18)	19	0.4	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19	0.4	*
13	Legal intervention (Y35,Y89.0)	19	0.4	*
...	All other causes (Residual)	412	7.9	13.3
Black, non-Hispanic, male, 25-34 years				
...	All causes	6,564	100.0	262.1
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,130	32.4	85.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,303	19.9	52.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	588	9.0	23.5
4	Human immunodeficiency virus (HIV) disease (B20-B24)	501	7.6	20.0
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	420	6.4	16.8
6	Malignant neoplasms (C00-C97)	254	3.9	10.1
7	Diabetes mellitus (E10-E14)	101	1.5	4.0
8	Cerebrovascular diseases (I60-I69)	77	1.2	3.1
9	Anemias (D50-D64)	70	1.1	2.8
10	Chronic lower respiratory diseases (J40-J47)	66	1.0	2.6
11	Septicemia (A40-A41)	52	0.8	2.1
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	51	0.8	2.0
13	Influenza and pneumonia (J10-J18)	41	0.6	1.6
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	36	0.5	1.4
15	Legal intervention (Y35,Y89.0)	33	0.5	1.3
...	All other causes (Residual)	841	12.8	33.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 35-44 years				
...	All causes	10,552	100.0	409.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,936	18.3	75.1
2	Human immunodeficiency virus (HIV) disease (B20-B24)	1,601	15.2	62.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,434	13.6	55.6
4	Malignant neoplasms (C00-C97)	1,025	9.7	39.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,005	9.5	39.0
6	Cerebrovascular diseases (I60-I69)	344	3.3	13.3
7	Diabetes mellitus (E10-E14)	314	3.0	12.2
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	293	2.8	11.4
9	Chronic liver disease and cirrhosis (K70,K73-K74)	197	1.9	7.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	166	1.6	6.4
11	Influenza and pneumonia (J10-J18)	126	1.2	4.9
12	Septicemia (A40-A41)	125	1.2	4.8
13	Chronic lower respiratory diseases (J40-J47)	107	1.0	4.2
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	103	1.0	4.0
15	Anemias (D50-D64)	74	0.7	2.9
...	All other causes (Residual)	1,702	16.1	66.0
Black, non-Hispanic, male, 45-54 years				
...	All causes	21,449	100.0	974.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,393	25.1	245.1
2	Malignant neoplasms (C00-C97)	4,439	20.7	201.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,714	8.0	77.9
4	Human immunodeficiency virus (HIV) disease (B20-B24)	1,679	7.8	76.3
5	Cerebrovascular diseases (I60-I69)	1,012	4.7	46.0
6	Diabetes mellitus (E10-E14)	748	3.5	34.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	608	2.8	27.6
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	574	2.7	26.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	474	2.2	21.5
10	Septicemia (A40-A41)	334	1.6	15.2
11	Chronic lower respiratory diseases (J40-J47)	317	1.5	14.4
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	291	1.4	13.2
13	Influenza and pneumonia (J10-J18)	266	1.2	12.1
14	Viral hepatitis (B15-B19)	262	1.2	11.9
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	214	1.0	9.7
...	All other causes (Residual)	3,124	14.6	142.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 55-64 years				
...	All causes	25,050	100.0	1,998.6
1	Malignant neoplasms (C00-C97)	7,276	29.0	580.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,149	28.5	570.4
3	Cerebrovascular diseases (I60-I69)	1,381	5.5	110.2
4	Diabetes mellitus (E10-E14)	1,171	4.7	93.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	901	3.6	71.9
6	Chronic lower respiratory diseases (J40-J47)	644	2.6	51.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	640	2.6	51.1
8	Human immunodeficiency virus (HIV) disease (B20-B24)	637	2.5	50.8
9	Chronic liver disease and cirrhosis (K70,K73-K74)	501	2.0	40.0
10	Septicemia (A40-A41)	465	1.9	37.1
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	361	1.4	28.8
12	Influenza and pneumonia (J10-J18)	311	1.2	24.8
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	190	0.8	15.2
14	Viral hepatitis (B15-B19)	177	0.7	14.1
15	Aortic aneurysm and dissection (I71)	148	0.6	11.8
...	All other causes (Residual)	3,098	12.4	247.2
Black, non-Hispanic, male, 65-74 years				
...	All causes	27,228	100.0	3,880.2
1	Malignant neoplasms (C00-C97)	8,642	31.7	1,231.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,798	28.6	1,111.3
3	Cerebrovascular diseases (I60-I69)	1,685	6.2	240.1
4	Diabetes mellitus (E10-E14)	1,417	5.2	201.9
5	Chronic lower respiratory diseases (J40-J47)	1,000	3.7	142.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	771	2.8	109.9
7	Septicemia (A40-A41)	582	2.1	82.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	520	1.9	74.1
9	Influenza and pneumonia (J10-J18)	514	1.9	73.2
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	404	1.5	57.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	242	0.9	34.5
12	Human immunodeficiency virus (HIV) disease (B20-B24)	164	0.6	23.4
13	Pneumonitis due to solids and liquids (J69)	151	0.6	21.5
14	Alzheimer's disease (G30)	136	0.5	19.4
15	Aortic aneurysm and dissection (I71)	130	0.5	18.5
...	All other causes (Residual)	3,072	11.3	437.8

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 75-84 years				
...	All causes	27,394	100.0	7,813.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,931	29.0	2,262.0
2	Malignant neoplasms (C00-C97)	7,422	27.1	2,116.9
3	Cerebrovascular diseases (I60-I69)	1,959	7.2	558.7
4	Diabetes mellitus (E10-E14)	1,207	4.4	344.3
5	Chronic lower respiratory diseases (J40-J47)	1,200	4.4	342.3
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	827	3.0	235.9
7	Influenza and pneumonia (J10-J18)	775	2.8	221.0
8	Septicemia (A40-A41)	675	2.5	192.5
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	488	1.8	139.2
10	Alzheimer's disease (G30)	486	1.8	138.6
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	467	1.7	133.2
12	Pneumonitis due to solids and liquids (J69)	241	0.9	68.7
13	Parkinson's disease (G20-G21)	156	0.6	44.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	140	0.5	39.9
15	Aortic aneurysm and dissection (I71)	131	0.5	37.4
...	All other causes (Residual)	3,289	12.0	938.1
Black, non-Hispanic, male, 85 years and over				
...	All causes	14,415	100.0	4,647.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,621	32.1	4,695.5
2	Malignant neoplasms (C00-C97)	2,737	19.0	2,781.1
3	Cerebrovascular diseases (I60-I69)	1,060	7.4	1,077.1
4	Influenza and pneumonia (J10-J18)	598	4.1	607.6
5	Chronic lower respiratory diseases (J40-J47)	560	3.9	569.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	512	3.6	520.3
7	Diabetes mellitus (E10-E14)	500	3.5	508.1
8	Alzheimer's disease (G30)	499	3.5	507.0
9	Septicemia (A40-A41)	341	2.4	346.5
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	264	1.8	268.3
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	215	1.5	218.5
12	Pneumonitis due to solids and liquids (J69)	204	1.4	207.3
13	Parkinson's disease (G20-G21)	105	0.7	106.7
14	Atherosclerosis (I70)	77	0.5	78.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	61	0.4	62.0
...	All other causes (Residual)	2,061	14.3	2,094.2

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, all ages				
...	All causes	139,837	100.0	723.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	37,624	26.9	194.6
2	Malignant neoplasms (C00-C97)	29,753	21.3	153.9
3	Cerebrovascular diseases (I60-I69)	10,394	7.4	53.8
4	Diabetes mellitus (E10-E14)	7,187	5.1	37.2
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,254	3.0	22.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,016	2.9	20.8
7	Chronic lower respiratory diseases (J40-J47)	3,321	2.4	17.2
8	Septicemia (A40-A41)	3,285	2.3	17.0
9	Alzheimer's disease (G30)	3,058	2.2	15.8
10	Influenza and pneumonia (J10-J18)	2,891	2.1	15.0
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,764	2.0	14.3
12	Human immunodeficiency virus (HIV) disease (B20-B24)	2,470	1.8	12.8
13	Certain conditions originating in the perinatal period (P00-P96)	2,087	1.5	10.8
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,270	0.9	6.6
15	Chronic liver disease and cirrhosis (K70,K73-K74)	856	0.6	4.4
...	All other causes (Residual)	24,607	17.6	127.3
Black, non-Hispanic, female, 1-4 years				
...	All causes	505	100.0	41.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	135	26.7	11.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	75	14.9	6.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	53	10.5	4.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	29	5.7	2.4
5	Malignant neoplasms (C00-C97)	27	5.3	2.2
6	Influenza and pneumonia (J10-J18)	22	4.4	1.8
7	Septicemia (A40-A41)	10	2.0	*
7	Certain conditions originating in the perinatal period (P00-P96)	10	2.0	*
9	Anemias (D50-D64)	7	1.4	*
9	Cerebrovascular diseases (I60-I69)	7	1.4	*
9	Chronic lower respiratory diseases (J40-J47)	7	1.4	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.6	*
12	Meningitis (G00,G03)	3	0.6	*
14	Meningococcal infection (A39)	2	0.4	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.4	*
14	Acute bronchitis and bronchiolitis (J20-J21)	2	0.4	*
...	All other causes (Residual)	111	22.0	9.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 5-14 years				
...	All causes	681	100.0	21.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	214	31.4	6.8
2	Malignant neoplasms (C00-C97)	83	12.2	2.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	49	7.2	1.6
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	44	6.5	1.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	43	6.3	1.4
6	Chronic lower respiratory diseases (J40-J47)	20	2.9	0.6
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	19	2.8	*
8	Anemias (D50-D64)	11	1.6	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	9	1.3	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	1.3	*
11	Septicemia (A40-A41)	8	1.2	*
11	Diabetes mellitus (E10-E14)	8	1.2	*
11	Influenza and pneumonia (J10-J18)	8	1.2	*
14	Cerebrovascular diseases (I60-I69)	7	1.0	*
15	Meningitis (G00,G03)	4	0.6	*
15	Certain conditions originating in the perinatal period (P00-P96)	4	0.6	*
...	All other causes (Residual)	141	20.7	4.5
Black, non-Hispanic, female, 15-24 years				
...	All causes	1,685	100.0	55.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	423	25.1	13.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	276	16.4	9.1
3	Malignant neoplasms (C00-C97)	133	7.9	4.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	108	6.4	3.5
5	Pregnancy, childbirth and the puerperium (O00-099)	73	4.3	2.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	68	4.0	2.2
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	68	4.0	2.2
8	Anemias (D50-D64)	40	2.4	1.3
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	38	2.3	1.2
10	Chronic lower respiratory diseases (J40-J47)	31	1.8	1.0
11	Diabetes mellitus (E10-E14)	30	1.8	1.0
12	Septicemia (A40-A41)	23	1.4	0.8
13	Influenza and pneumonia (J10-J18)	21	1.2	0.7
14	Cerebrovascular diseases (I60-I69)	18	1.1	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	0.7	*
...	All other causes (Residual)	323	19.2	10.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 25-34 years				
...	All causes	3,204	100.0	117.2
1	Human immunodeficiency virus (HIV) disease (B20-B24)	435	13.6	15.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	401	12.5	14.7
3	Malignant neoplasms (C00-C97)	397	12.4	14.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	383	12.0	14.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	275	8.6	10.1
6	Pregnancy, childbirth and the puerperium (O00-099)	106	3.3	3.9
7	Cerebrovascular diseases (I60-I69)	96	3.0	3.5
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	90	2.8	3.3
9	Diabetes mellitus (E10-E14)	79	2.5	2.9
10	Anemias (D50-D64)	56	1.7	2.0
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	52	1.6	1.9
12	Septicemia (A40-A41)	50	1.6	1.8
13	Chronic lower respiratory diseases (J40-J47)	35	1.1	1.3
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	26	0.8	1.0
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	23	0.7	0.8
...	All other causes (Residual)	700	21.8	25.6
Black, non-Hispanic, female, 35-44 years				
...	All causes	7,685	100.0	264.1
1	Malignant neoplasms (C00-C97)	1,624	21.1	55.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,266	16.5	43.5
3	Human immunodeficiency virus (HIV) disease (B20-B24)	921	12.0	31.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	638	8.3	21.9
5	Cerebrovascular diseases (I60-I69)	354	4.6	12.2
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	271	3.5	9.3
7	Diabetes mellitus (E10-E14)	226	2.9	7.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	150	2.0	5.2
9	Septicemia (A40-A41)	130	1.7	4.5
10	Chronic lower respiratory diseases (J40-J47)	113	1.5	3.9
11	Chronic liver disease and cirrhosis (K70,K73-K74)	110	1.4	3.8
12	Influenza and pneumonia (J10-J18)	97	1.3	3.3
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	76	1.0	2.6
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	74	1.0	2.5
15	Anemias (D50-D64)	65	0.8	2.2
...	All other causes (Residual)	1,570	20.4	53.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Black, non-Hispanic, female, 45-54 years			
...	All causes	14,798	100.0	577.0
1	Malignant neoplasms (C00-C97)	4,222	28.5	164.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,179	21.5	124.0
3	Cerebrovascular diseases (I60-I69)	893	6.0	34.8
4	Human immunodeficiency virus (HIV) disease (B20-B24)	744	5.0	29.0
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	704	4.8	27.5
6	Diabetes mellitus (E10-E14)	623	4.2	24.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	339	2.3	13.2
8	Septicemia (A40-A41)	316	2.1	12.3
9	Chronic lower respiratory diseases (J40-J47)	304	2.1	11.9
10	Chronic liver disease and cirrhosis (K70,K73-K74)	282	1.9	11.0
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	206	1.4	8.0
12	Influenza and pneumonia (J10-J18)	177	1.2	6.9
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	158	1.1	6.2
14	Viral hepatitis (B15-B19)	100	0.7	3.9
15	Anemias (D50-D64)	77	0.5	3.0
...	All other causes (Residual)	2,474	16.7	96.5
	Black, non-Hispanic, female, 55-64 years			
...	All causes	17,948	100.0	1,149.9
1	Malignant neoplasms (C00-C97)	5,873	32.7	376.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,392	24.5	281.4
3	Diabetes mellitus (E10-E14)	1,133	6.3	72.6
4	Cerebrovascular diseases (I60-I69)	1,039	5.8	66.6
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	643	3.6	41.2
6	Chronic lower respiratory diseases (J40-J47)	457	2.5	29.3
7	Septicemia (A40-A41)	441	2.5	28.3
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	350	2.0	22.4
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	324	1.8	20.8
10	Chronic liver disease and cirrhosis (K70,K73-K74)	227	1.3	14.5
11	Influenza and pneumonia (J10-J18)	226	1.3	14.5
12	Human immunodeficiency virus (HIV) disease (B20-B24)	204	1.1	13.1
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	77	0.4	4.9
14	Viral hepatitis (B15-B19)	70	0.4	4.5
15	Aortic aneurysm and dissection (I71)	67	0.4	4.3
...	All other causes (Residual)	2,425	13.5	155.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Black, non-Hispanic, female, 65-74 years			
...	All causes	23,898	100.0	2,427.9
1	Malignant neoplasms (C00-C97)	7,070	29.6	718.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,574	27.5	667.9
3	Cerebrovascular diseases (I60-I69)	1,677	7.0	170.4
4	Diabetes mellitus (E10-E14)	1,661	7.0	168.7
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	858	3.6	87.2
6	Chronic lower respiratory diseases (J40-J47)	751	3.1	76.3
7	Septicemia (A40-A41)	621	2.6	63.1
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	486	2.0	49.4
9	Influenza and pneumonia (J10-J18)	336	1.4	34.1
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	298	1.2	30.3
11	Alzheimer's disease (G30)	191	0.8	19.4
12	Chronic liver disease and cirrhosis (K70,K73-K74)	129	0.5	13.1
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	110	0.5	11.2
14	Aortic aneurysm and dissection (I71)	94	0.4	9.5
15	Pneumonitis due to solids and liquids (J69)	88	0.4	8.9
...	All other causes (Residual)	2,954	12.4	300.1
	Black, non-Hispanic, female, 75-84 years			
...	All causes	33,354	100.0	5,370.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,208	30.6	1,643.6
2	Malignant neoplasms (C00-C97)	6,822	20.5	1,098.4
3	Cerebrovascular diseases (I60-I69)	3,085	9.2	496.7
4	Diabetes mellitus (E10-E14)	2,062	6.2	332.0
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,166	3.5	187.7
6	Alzheimer's disease (G30)	1,024	3.1	164.9
7	Chronic lower respiratory diseases (J40-J47)	995	3.0	160.2
8	Septicemia (A40-A41)	859	2.6	138.3
9	Influenza and pneumonia (J10-J18)	764	2.3	123.0
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	726	2.2	116.9
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	390	1.2	62.8
12	Pneumonitis due to solids and liquids (J69)	222	0.7	35.7
13	Aortic aneurysm and dissection (I71)	184	0.6	29.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	171	0.5	27.5
15	Parkinson's disease (G20-G21)	134	0.4	21.6
...	All other causes (Residual)	4,542	13.6	731.3

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2004

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Percent		
		Number	of total deaths	Rate
	Black, non-Hispanic, female, 85 years and over			
...	All causes	32,462	100.0	3,035.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,397	35.1	4,576.6
2	Malignant neoplasms (C00-C97)	3,497	10.8	1,404.3
3	Cerebrovascular diseases (I60-I69)	3,200	9.9	1,285.0
4	Alzheimer's disease (G30)	1,818	5.6	730.0
5	Diabetes mellitus (E10-E14)	1,364	4.2	547.7
6	Influenza and pneumonia (J10-J18)	1,180	3.6	473.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,007	3.1	404.4
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	916	2.8	367.8
9	Septicemia (A40-A41)	787	2.4	316.0
10	Chronic lower respiratory diseases (J40-J47)	607	1.9	243.8
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	324	1.0	130.1
12	Pneumonitis due to solids and liquids (J69)	308	0.9	123.7
13	Atherosclerosis (I70)	283	0.9	113.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	140	0.4	56.2
15	Anemias (D50-D64)	127	0.4	51.0
...	All other causes (Residual)	5,507	17.0	2,211.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable