

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|---------------------------------|---|---------|-------------------------|-------|
| Hispanic, both sexes, all ages | | | | |
| ... | All causes | 107,254 | 100.0 | 330.4 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 25,819 | 24.1 | 79.5 |
| 2 | Malignant neoplasms (C00-C97) | 21,160 | 19.7 | 65.2 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 8,830 | 8.2 | 27.2 |
| 4 | Cerebrovascular diseases (I60-I69) | 6,187 | 5.8 | 19.1 |
| 5 | Diabetes mellitus (E10-E14) | 5,328 | 5.0 | 16.4 |
| 6 | Chronic liver disease and cirrhosis (K70,K73-K74) | 3,187 | 3.0 | 9.8 |
| 7 | Assault (homicide) (X85-Y09,Y87.1) | 2,917 | 2.7 | 9.0 |
| 8 | Chronic lower respiratory diseases (J40-J47) | 2,689 | 2.5 | 8.3 |
| 9 | Influenza and pneumonia (J10-J18) | 2,625 | 2.4 | 8.1 |
| 10 | Certain conditions originating in the perinatal period (P00-P96) | 2,145 | 2.0 | 6.6 |
| 11 | Human immunodeficiency virus (HIV) disease (B20-B24) | 1,908 | 1.8 | 5.9 |
| 12 | Intentional self-harm (suicide) (X60-X84,Y87.0) | 1,787 | 1.7 | 5.5 |
| 13 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1,646 | 1.5 | 5.1 |
| 14 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 1,612 | 1.5 | 5.0 |
| 15 | Septicemia (A40-A41) | 1,226 | 1.1 | 3.8 |
| ... | All other causes (Residual) | 18,188 | 17.0 | 56.0 |
| Hispanic, both sexes, 1-4 years | | | | |
| ... | All causes | 872 | 100.0 | 31.0 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 306 | 35.1 | 10.9 |
| 2 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 111 | 12.7 | 3.9 |
| 3 | Malignant neoplasms (C00-C97) | 75 | 8.6 | 2.7 |
| 4 | Assault (homicide) (X85-Y09,Y87.1) | 57 | 6.5 | 2.0 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 29 | 3.3 | 1.0 |
| 6 | Influenza and pneumonia (J10-J18) | 17 | 1.9 | * |
| 7 | Septicemia (A40-A41) | 15 | 1.7 | * |
| 8 | Certain conditions originating in the perinatal period (P00-P96) | 12 | 1.4 | * |
| 9 | Chronic lower respiratory diseases (J40-J47) | 9 | 1.0 | * |
| 10 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 7 | 0.8 | * |
| 11 | Cerebrovascular diseases (I60-I69) | 6 | 0.7 | * |
| 12 | Meningitis (G00,G03) | 4 | 0.5 | * |
| 12 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 4 | 0.5 | * |
| 12 | Complications of medical and surgical care (Y40-Y84,Y88) | 4 | 0.5 | * |
| 15 | Meningococcal infection (A39) | 3 | 0.3 | * |
| ... | All other causes (Residual) | 213 | 24.4 | 7.6 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|-----------------------------------|---|--------|-------------------------|------|
| Hispanic, both sexes, 5-14 years | | | | |
| ... | All causes | 1,065 | 100.0 | 17.1 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 383 | 36.0 | 6.1 |
| 2 | Malignant neoplasms (C00-C97) | 198 | 18.6 | 3.2 |
| 3 | Assault (homicide) (X85-Y09,Y87.1) | 76 | 7.1 | 1.2 |
| 4 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 52 | 4.9 | 0.8 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 38 | 3.6 | 0.6 |
| 6 | Intentional self-harm (suicide) (X60-X84,Y87.0) | 29 | 2.7 | 0.5 |
| 7 | Influenza and pneumonia (J10-J18) | 21 | 2.0 | 0.3 |
| 8 | Chronic lower respiratory diseases (J40-J47) | 15 | 1.4 | * |
| 9 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 13 | 1.2 | * |
| 10 | Certain conditions originating in the perinatal period (P00-P96) | 10 | 0.9 | * |
| 11 | Human immunodeficiency virus (HIV) disease (B20-B24) | 9 | 0.8 | * |
| 12 | Cerebrovascular diseases (I60-I69) | 8 | 0.8 | * |
| 13 | Septicemia (A40-A41) | 7 | 0.7 | * |
| 13 | Anemias (D50-D64) | 7 | 0.7 | * |
| 15 | Complications of medical and surgical care (Y40-Y84,Y88) | 6 | 0.6 | * |
| ... | All other causes (Residual) | 193 | 18.1 | 3.1 |
| Hispanic, both sexes, 15-24 years | | | | |
| ... | All causes | 4,793 | 100.0 | 85.0 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 2,095 | 43.7 | 37.2 |
| 2 | Assault (homicide) (X85-Y09,Y87.1) | 1,126 | 23.5 | 20.0 |
| 3 | Intentional self-harm (suicide) (X60-X84,Y87.0) | 447 | 9.3 | 7.9 |
| 4 | Malignant neoplasms (C00-C97) | 275 | 5.7 | 4.9 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 125 | 2.6 | 2.2 |
| 6 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 54 | 1.1 | 1.0 |
| 7 | Cerebrovascular diseases (I60-I69) | 31 | 0.6 | 0.5 |
| 8 | Pregnancy, childbirth and the puerperium (O00-099) | 24 | 0.5 | 0.4 |
| 9 | Human immunodeficiency virus (HIV) disease (B20-B24) | 23 | 0.5 | 0.4 |
| 10 | Influenza and pneumonia (J10-J18) | 21 | 0.4 | 0.4 |
| 11 | Legal intervention (Y35,Y89.0) | 18 | 0.4 | * |
| 12 | Chronic lower respiratory diseases (J40-J47) | 16 | 0.3 | * |
| 13 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 14 | 0.3 | * |
| 14 | Diabetes mellitus (E10-E14) | 13 | 0.3 | * |
| 15 | Septicemia (A40-A41) | 12 | 0.3 | * |
| ... | All other causes (Residual) | 499 | 10.4 | 8.9 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|-----------------------------------|---|--------|-------------------------|-------|
| Hispanic, both sexes, 25-44 years | | | | |
| ... | All causes | 13,594 | 100.0 | 133.0 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 3,394 | 25.0 | 33.2 |
| 2 | Malignant neoplasms (C00-C97) | 1,771 | 13.0 | 17.3 |
| 3 | Assault (homicide) (X85-Y09,Y87.1) | 1,257 | 9.2 | 12.3 |
| 4 | Human immunodeficiency virus (HIV) disease (B20-B24) | 1,127 | 8.3 | 11.0 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 1,040 | 7.7 | 10.2 |
| 6 | Intentional self-harm (suicide) (X60-X84,Y87.0) | 805 | 5.9 | 7.9 |
| 7 | Chronic liver disease and cirrhosis (K70,K73-K74) | 593 | 4.4 | 5.8 |
| 8 | Cerebrovascular diseases (I60-I69) | 375 | 2.8 | 3.7 |
| 9 | Diabetes mellitus (E10-E14) | 218 | 1.6 | 2.1 |
| 10 | Viral hepatitis (B15-B19) | 140 | 1.0 | 1.4 |
| 11 | Influenza and pneumonia (J10-J18) | 130 | 1.0 | 1.3 |
| 12 | Septicemia (A40-A41) | 110 | 0.8 | 1.1 |
| 13 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 89 | 0.7 | 0.9 |
| 14 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 79 | 0.6 | 0.8 |
| 15 | Chronic lower respiratory diseases (J40-J47) | 75 | 0.6 | 0.7 |
| ... | All other causes (Residual) | 2,391 | 17.6 | 23.4 |
| Hispanic, both sexes, 45-64 years | | | | |
| ... | All causes | 23,089 | 100.0 | 473.7 |
| 1 | Malignant neoplasms (C00-C97) | 6,355 | 27.5 | 130.4 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 4,890 | 21.2 | 100.3 |
| 3 | Chronic liver disease and cirrhosis (K70,K73-K74) | 1,562 | 6.8 | 32.0 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,445 | 6.3 | 29.6 |
| 5 | Diabetes mellitus (E10-E14) | 1,389 | 6.0 | 28.5 |
| 6 | Cerebrovascular diseases (I60-I69) | 1,158 | 5.0 | 23.8 |
| 7 | Human immunodeficiency virus (HIV) disease (B20-B24) | 694 | 3.0 | 14.2 |
| 8 | Viral hepatitis (B15-B19) | 365 | 1.6 | 7.5 |
| 9 | Chronic lower respiratory diseases (J40-J47) | 349 | 1.5 | 7.2 |
| 10 | Intentional self-harm (suicide) (X60-X84,Y87.0) | 346 | 1.5 | 7.1 |
| 11 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 341 | 1.5 | 7.0 |
| 12 | Influenza and pneumonia (J10-J18) | 311 | 1.3 | 6.4 |
| 13 | Assault (homicide) (X85-Y09,Y87.1) | 288 | 1.2 | 5.9 |
| 14 | Septicemia (A40-A41) | 267 | 1.2 | 5.5 |
| 15 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 164 | 0.7 | 3.4 |
| ... | All other causes (Residual) | 3,165 | 13.7 | 64.9 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|-------------------|---|--------|-------------------------|---------|
| | Hispanic, both sexes, 65 years and over | | | |
| ... | All causes | 59,189 | 100.0 | 3,034.3 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 19,606 | 33.1 | 1,005.1 |
| 2 | Malignant neoplasms (C00-C97) | 12,469 | 21.1 | 639.2 |
| 3 | Cerebrovascular diseases (I60-I69) | 4,593 | 7.8 | 235.5 |
| 4 | Diabetes mellitus (E10-E14) | 3,703 | 6.3 | 189.8 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 2,218 | 3.7 | 113.7 |
| 6 | Influenza and pneumonia (J10-J18) | 2,065 | 3.5 | 105.9 |
| 7 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1,179 | 2.0 | 60.4 |
| 8 | Alzheimer's disease (G30) | 1,102 | 1.9 | 56.5 |
| 9 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,061 | 1.8 | 54.4 |
| 10 | Chronic liver disease and cirrhosis (K70,K73-K74) | 1,024 | 1.7 | 52.5 |
| 11 | Septicemia (A40-A41) | 753 | 1.3 | 38.6 |
| 12 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 617 | 1.0 | 31.6 |
| 13 | Pneumonitis due to solids and liquids (J69) | 412 | 0.7 | 21.1 |
| 14 | Parkinson's disease (G20-G21) | 405 | 0.7 | 20.8 |
| 15 | Atherosclerosis (I70) | 387 | 0.7 | 19.8 |
| ... | All other causes (Residual) | 7,595 | 12.8 | 389.4 |
| | Hispanic, male, all ages | | | |
| ... | All causes | 60,172 | 100.0 | 368.9 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 13,566 | 22.5 | 83.2 |
| 2 | Malignant neoplasms (C00-C97) | 11,138 | 18.5 | 68.3 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 6,696 | 11.1 | 41.1 |
| 4 | Cerebrovascular diseases (I60-I69) | 2,865 | 4.8 | 17.6 |
| 5 | Diabetes mellitus (E10-E14) | 2,507 | 4.2 | 15.4 |
| 6 | Assault (homicide) (X85-Y09,Y87.1) | 2,431 | 4.0 | 14.9 |
| 7 | Chronic liver disease and cirrhosis (K70,K73-K74) | 2,312 | 3.8 | 14.2 |
| 8 | Intentional self-harm (suicide) (X60-X84,Y87.0) | 1,525 | 2.5 | 9.3 |
| 9 | Human immunodeficiency virus (HIV) disease (B20-B24) | 1,493 | 2.5 | 9.2 |
| 10 | Chronic lower respiratory diseases (J40-J47) | 1,451 | 2.4 | 8.9 |
| 11 | Influenza and pneumonia (J10-J18) | 1,303 | 2.2 | 8.0 |
| 12 | Certain conditions originating in the perinatal period (P00-P96) | 1,194 | 2.0 | 7.3 |
| 13 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 857 | 1.4 | 5.3 |
| 14 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 805 | 1.3 | 4.9 |
| 15 | Septicemia (A40-A41) | 625 | 1.0 | 3.8 |
| ... | All other causes (Residual) | 9,404 | 15.6 | 57.7 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|----------------------------|---|--------|-------------------------|------|
| Hispanic, male, 1-4 years | | | | |
| ... | All causes | 475 | 100.0 | 33.1 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 174 | 36.6 | 12.1 |
| 2 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 60 | 12.6 | 4.2 |
| 3 | Malignant neoplasms (C00-C97) | 43 | 9.1 | 3.0 |
| 4 | Assault (homicide) (X85-Y09,Y87.1) | 30 | 6.3 | 2.1 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 21 | 4.4 | 1.5 |
| 6 | Septicemia (A40-A41) | 9 | 1.9 | * |
| 7 | Certain conditions originating in the perinatal period (P00-P96) | 7 | 1.5 | * |
| 8 | Influenza and pneumonia (J10-J18) | 5 | 1.1 | * |
| 8 | Chronic lower respiratory diseases (J40-J47) | 5 | 1.1 | * |
| 10 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 3 | 0.6 | * |
| 11 | Meningococcal infection (A39) | 2 | 0.4 | * |
| 11 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 2 | 0.4 | * |
| 11 | Meningitis (G00,G03) | 2 | 0.4 | * |
| 11 | Complications of medical and surgical care (Y40-Y84,Y88) | 2 | 0.4 | * |
| 15 | Acute bronchitis and bronchiolitis (J20-J21) | 1 | 0.2 | * |
| 15 | Diseases of appendix (K35-K38) | 1 | 0.2 | * |
| ... | All other causes (Residual) | 108 | 22.7 | 7.5 |
| Hispanic, male, 5-14 years | | | | |
| ... | All causes | 622 | 100.0 | 19.6 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 210 | 33.8 | 6.6 |
| 2 | Malignant neoplasms (C00-C97) | 122 | 19.6 | 3.8 |
| 3 | Assault (homicide) (X85-Y09,Y87.1) | 56 | 9.0 | 1.8 |
| 4 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 28 | 4.5 | 0.9 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 26 | 4.2 | 0.8 |
| 6 | Intentional self-harm (suicide) (X60-X84,Y87.0) | 22 | 3.5 | 0.7 |
| 7 | Influenza and pneumonia (J10-J18) | 10 | 1.6 | * |
| 8 | Chronic lower respiratory diseases (J40-J47) | 9 | 1.4 | * |
| 9 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 8 | 1.3 | * |
| 10 | Certain conditions originating in the perinatal period (P00-P96) | 6 | 1.0 | * |
| 11 | Septicemia (A40-A41) | 5 | 0.8 | * |
| 11 | Human immunodeficiency virus (HIV) disease (B20-B24) | 5 | 0.8 | * |
| 13 | Complications of medical and surgical care (Y40-Y84,Y88) | 4 | 0.6 | * |
| 14 | Anemias (D50-D64) | 3 | 0.5 | * |
| 14 | Cerebrovascular diseases (I60-I69) | 3 | 0.5 | * |
| 14 | Diseases of appendix (K35-K38) | 3 | 0.5 | * |
| ... | All other causes (Residual) | 102 | 16.4 | 3.2 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|-------------------|---|--------|-------------------------|-------|
| | Hispanic, male, 15-24 years | | | |
| ... | All causes | 3,837 | 100.0 | 131.2 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,717 | 44.7 | 58.7 |
| 2 | Assault (homicide) (X85-Y09,Y87.1) | 1,014 | 26.4 | 34.7 |
| 3 | Intentional self-harm (suicide) (X60-X84,Y87.0) | 387 | 10.1 | 13.2 |
| 4 | Malignant neoplasms (C00-C97) | 162 | 4.2 | 5.5 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 87 | 2.3 | 3.0 |
| 6 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 39 | 1.0 | 1.3 |
| 7 | Human immunodeficiency virus (HIV) disease (B20-B24) | 18 | 0.5 | * |
| 8 | Cerebrovascular diseases (I60-I69) | 17 | 0.4 | * |
| 8 | Legal intervention (Y35,Y89.0) | 17 | 0.4 | * |
| 10 | Chronic lower respiratory diseases (J40-J47) | 13 | 0.3 | * |
| 11 | Influenza and pneumonia (J10-J18) | 10 | 0.3 | * |
| 12 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 8 | 0.2 | * |
| 13 | Diabetes mellitus (E10-E14) | 7 | 0.2 | * |
| 13 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 7 | 0.2 | * |
| 15 | Meningococcal infection (A39) | 6 | 0.2 | * |
| 15 | Septicemia (A40-A41) | 6 | 0.2 | * |
| 15 | Complications of medical and surgical care (Y40-Y84,Y88) | 6 | 0.2 | * |
| ... | All other causes (Residual) | 316 | 8.2 | 10.8 |
| | Hispanic, male, 25-44 years | | | |
| ... | All causes | 9,797 | 100.0 | 187.3 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 2,816 | 28.7 | 53.8 |
| 2 | Assault (homicide) (X85-Y09,Y87.1) | 1,056 | 10.8 | 20.2 |
| 3 | Human immunodeficiency virus (HIV) disease (B20-B24) | 877 | 9.0 | 16.8 |
| 4 | Malignant neoplasms (C00-C97) | 775 | 7.9 | 14.8 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 749 | 7.6 | 14.3 |
| 6 | Intentional self-harm (suicide) (X60-X84,Y87.0) | 690 | 7.0 | 13.2 |
| 7 | Chronic liver disease and cirrhosis (K70,K73-K74) | 481 | 4.9 | 9.2 |
| 8 | Cerebrovascular diseases (I60-I69) | 210 | 2.1 | 4.0 |
| 9 | Diabetes mellitus (E10-E14) | 143 | 1.5 | 2.7 |
| 10 | Viral hepatitis (B15-B19) | 111 | 1.1 | 2.1 |
| 11 | Influenza and pneumonia (J10-J18) | 83 | 0.8 | 1.6 |
| 12 | Septicemia (A40-A41) | 67 | 0.7 | 1.3 |
| 13 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 53 | 0.5 | 1.0 |
| 14 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 47 | 0.5 | 0.9 |
| 15 | Chronic lower respiratory diseases (J40-J47) | 36 | 0.4 | 0.7 |
| ... | All other causes (Residual) | 1,603 | 16.4 | 30.7 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|-------------------|---|--------|-------------------------|---------|
| | Hispanic, male, 45-64 years | | | |
| ... | All causes | 14,568 | 100.0 | 622.6 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 3,435 | 23.6 | 146.8 |
| 2 | Malignant neoplasms (C00-C97) | 3,266 | 22.4 | 139.6 |
| 3 | Chronic liver disease and cirrhosis (K70,K73-K74) | 1,238 | 8.5 | 52.9 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,097 | 7.5 | 46.9 |
| 5 | Diabetes mellitus (E10-E14) | 760 | 5.2 | 32.5 |
| 6 | Cerebrovascular diseases (I60-I69) | 672 | 4.6 | 28.7 |
| 7 | Human immunodeficiency virus (HIV) disease (B20-B24) | 549 | 3.8 | 23.5 |
| 8 | Intentional self-harm (suicide) (X60-X84,Y87.0) | 283 | 1.9 | 12.1 |
| 9 | Viral hepatitis (B15-B19) | 259 | 1.8 | 11.1 |
| 10 | Assault (homicide) (X85-Y09,Y87.1) | 215 | 1.5 | 9.2 |
| 11 | Influenza and pneumonia (J10-J18) | 201 | 1.4 | 8.6 |
| 12 | Chronic lower respiratory diseases (J40-J47) | 199 | 1.4 | 8.5 |
| 13 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 184 | 1.3 | 7.9 |
| 14 | Septicemia (A40-A41) | 158 | 1.1 | 6.8 |
| 15 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 109 | 0.7 | 4.7 |
| ... | All other causes (Residual) | 1,943 | 13.3 | 83.0 |
| | Hispanic, male, 65 years and over | | | |
| ... | All causes | 28,308 | 100.0 | 3,434.5 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 9,200 | 32.5 | 1,116.2 |
| 2 | Malignant neoplasms (C00-C97) | 6,760 | 23.9 | 820.2 |
| 3 | Cerebrovascular diseases (I60-I69) | 1,955 | 6.9 | 237.2 |
| 4 | Diabetes mellitus (E10-E14) | 1,594 | 5.6 | 193.4 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 1,185 | 4.2 | 143.8 |
| 6 | Influenza and pneumonia (J10-J18) | 964 | 3.4 | 117.0 |
| 7 | Chronic liver disease and cirrhosis (K70,K73-K74) | 588 | 2.1 | 71.3 |
| 8 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 587 | 2.1 | 71.2 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 545 | 1.9 | 66.1 |
| 10 | Alzheimer's disease (G30) | 370 | 1.3 | 44.9 |
| 11 | Septicemia (A40-A41) | 347 | 1.2 | 42.1 |
| 12 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 249 | 0.9 | 30.2 |
| 13 | Parkinson's disease (G20-G21) | 229 | 0.8 | 27.8 |
| 14 | Pneumonitis due to solids and liquids (J69) | 206 | 0.7 | 25.0 |
| 15 | Aortic aneurysm and dissection (I71) | 194 | 0.7 | 23.5 |
| ... | All other causes (Residual) | 3,335 | 11.8 | 404.6 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|-------------------|---|--------|-------------------------|-------|
| | Hispanic, female, all ages | | | |
| ... | All causes | 47,082 | 100.0 | 291.5 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 12,253 | 26.0 | 75.9 |
| 2 | Malignant neoplasms (C00-C97) | 10,022 | 21.3 | 62.0 |
| 3 | Cerebrovascular diseases (I60-I69) | 3,322 | 7.1 | 20.6 |
| 4 | Diabetes mellitus (E10-E14) | 2,821 | 6.0 | 17.5 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 2,134 | 4.5 | 13.2 |
| 6 | Influenza and pneumonia (J10-J18) | 1,322 | 2.8 | 8.2 |
| 7 | Chronic lower respiratory diseases (J40-J47) | 1,238 | 2.6 | 7.7 |
| 8 | Certain conditions originating in the perinatal period (P00-P96) | 951 | 2.0 | 5.9 |
| 9 | Chronic liver disease and cirrhosis (K70,K73-K74) | 875 | 1.9 | 5.4 |
| 10 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 841 | 1.8 | 5.2 |
| 11 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 755 | 1.6 | 4.7 |
| 12 | Alzheimer's disease (G30) | 744 | 1.6 | 4.6 |
| 13 | Septicemia (A40-A41) | 601 | 1.3 | 3.7 |
| 14 | Assault (homicide) (X85-Y09,Y87.1) | 486 | 1.0 | 3.0 |
| 15 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 435 | 0.9 | 2.7 |
| ... | All other causes (Residual) | 8,282 | 17.6 | 51.3 |
| | Hispanic, female, 1-4 years | | | |
| ... | All causes | 397 | 100.0 | 28.8 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 132 | 33.2 | 9.6 |
| 2 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 51 | 12.8 | 3.7 |
| 3 | Malignant neoplasms (C00-C97) | 32 | 8.1 | 2.3 |
| 4 | Assault (homicide) (X85-Y09,Y87.1) | 27 | 6.8 | 2.0 |
| 5 | Influenza and pneumonia (J10-J18) | 12 | 3.0 | * |
| 6 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 8 | 2.0 | * |
| 7 | Septicemia (A40-A41) | 6 | 1.5 | * |
| 7 | Cerebrovascular diseases (I60-I69) | 6 | 1.5 | * |
| 9 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 5 | 1.3 | * |
| 9 | Certain conditions originating in the perinatal period (P00-P96) | 5 | 1.3 | * |
| 11 | Chronic lower respiratory diseases (J40-J47) | 4 | 1.0 | * |
| 12 | Tuberculosis (A16-A19) | 2 | 0.5 | * |
| 12 | Meningitis (G00,G03) | 2 | 0.5 | * |
| 12 | Pneumonitis due to solids and liquids (J69) | 2 | 0.5 | * |
| 12 | Complications of medical and surgical care (Y40-Y84,Y88) | 2 | 0.5 | * |
| ... | All other causes (Residual) | 101 | 25.4 | 7.3 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|-------------------|---|--------|-------------------------|------|
| | Hispanic, female, 5-14 years | | | |
| ... | All causes | 443 | 100.0 | 14.5 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 173 | 39.1 | 5.7 |
| 2 | Malignant neoplasms (C00-C97) | 76 | 17.2 | 2.5 |
| 3 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 24 | 5.4 | 0.8 |
| 4 | Assault (homicide) (X85-Y09,Y87.1) | 20 | 4.5 | 0.7 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 12 | 2.7 | * |
| 6 | Influenza and pneumonia (J10-J18) | 11 | 2.5 | * |
| 7 | Intentional self-harm (suicide) (X60-X84,Y87.0) | 7 | 1.6 | * |
| 8 | Chronic lower respiratory diseases (J40-J47) | 6 | 1.4 | * |
| 9 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 5 | 1.1 | * |
| 9 | Cerebrovascular diseases (I60-I69) | 5 | 1.1 | * |
| 11 | Human immunodeficiency virus (HIV) disease (B20-B24) | 4 | 0.9 | * |
| 11 | Anemias (D50-D64) | 4 | 0.9 | * |
| 11 | Certain conditions originating in the perinatal period (P00-P96) | 4 | 0.9 | * |
| 14 | Meningococcal infection (A39) | 2 | 0.5 | * |
| 14 | Septicemia (A40-A41) | 2 | 0.5 | * |
| 14 | Diabetes mellitus (E10-E14) | 2 | 0.5 | * |
| 14 | Meningitis (G00,G03) | 2 | 0.5 | * |
| 14 | Complications of medical and surgical care (Y40-Y84,Y88) | 2 | 0.5 | * |
| ... | All other causes (Residual) | 82 | 18.5 | 2.7 |
| | Hispanic, female, 15-24 years | | | |
| ... | All causes | 956 | 100.0 | 35.2 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 378 | 39.5 | 13.9 |
| 2 | Malignant neoplasms (C00-C97) | 113 | 11.8 | 4.2 |
| 3 | Assault (homicide) (X85-Y09,Y87.1) | 112 | 11.7 | 4.1 |
| 4 | Intentional self-harm (suicide) (X60-X84,Y87.0) | 60 | 6.3 | 2.2 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 38 | 4.0 | 1.4 |
| 6 | Pregnancy, childbirth and the puerperium (O00-099) | 24 | 2.5 | 0.9 |
| 7 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 15 | 1.6 | * |
| 8 | Cerebrovascular diseases (I60-I69) | 14 | 1.5 | * |
| 9 | Influenza and pneumonia (J10-J18) | 11 | 1.2 | * |
| 10 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 7 | 0.7 | * |
| 11 | Septicemia (A40-A41) | 6 | 0.6 | * |
| 11 | Diabetes mellitus (E10-E14) | 6 | 0.6 | * |
| 13 | Human immunodeficiency virus (HIV) disease (B20-B24) | 5 | 0.5 | * |
| 14 | Anemias (D50-D64) | 4 | 0.4 | * |
| 14 | Aortic aneurysm and dissection (I71) | 4 | 0.4 | * |
| ... | All other causes (Residual) | 159 | 16.6 | 5.9 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|-------------------|---|--------|-------------------------|-------|
| | Hispanic, female, 25-44 years | | | |
| ... | All causes | 3,797 | 100.0 | 76.1 |
| 1 | Malignant neoplasms (C00-C97) | 996 | 26.2 | 20.0 |
| 2 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 578 | 15.2 | 11.6 |
| 3 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 291 | 7.7 | 5.8 |
| 4 | Human immunodeficiency virus (HIV) disease (B20-B24) | 250 | 6.6 | 5.0 |
| 5 | Assault (homicide) (X85-Y09,Y87.1) | 201 | 5.3 | 4.0 |
| 6 | Cerebrovascular diseases (I60-I69) | 165 | 4.3 | 3.3 |
| 7 | Intentional self-harm (suicide) (X60-X84,Y87.0) | 115 | 3.0 | 2.3 |
| 8 | Chronic liver disease and cirrhosis (K70,K73-K74) | 112 | 2.9 | 2.2 |
| 9 | Diabetes mellitus (E10-E14) | 75 | 2.0 | 1.5 |
| 10 | Pregnancy, childbirth and the puerperium (O00-099) | 59 | 1.6 | 1.2 |
| 11 | Influenza and pneumonia (J10-J18) | 47 | 1.2 | 0.9 |
| 12 | Septicemia (A40-A41) | 43 | 1.1 | 0.9 |
| 13 | Chronic lower respiratory diseases (J40-J47) | 39 | 1.0 | 0.8 |
| 14 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 36 | 0.9 | 0.7 |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 33 | 0.9 | 0.7 |
| ... | All other causes (Residual) | 757 | 19.9 | 15.2 |
| | Hispanic, female, 45-64 years | | | |
| ... | All causes | 8,521 | 100.0 | 336.2 |
| 1 | Malignant neoplasms (C00-C97) | 3,089 | 36.3 | 121.9 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 1,455 | 17.1 | 57.4 |
| 3 | Diabetes mellitus (E10-E14) | 629 | 7.4 | 24.8 |
| 4 | Cerebrovascular diseases (I60-I69) | 486 | 5.7 | 19.2 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 348 | 4.1 | 13.7 |
| 6 | Chronic liver disease and cirrhosis (K70,K73-K74) | 324 | 3.8 | 12.8 |
| 7 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 157 | 1.8 | 6.2 |
| 8 | Chronic lower respiratory diseases (J40-J47) | 150 | 1.8 | 5.9 |
| 9 | Human immunodeficiency virus (HIV) disease (B20-B24) | 145 | 1.7 | 5.7 |
| 10 | Influenza and pneumonia (J10-J18) | 110 | 1.3 | 4.3 |
| 11 | Septicemia (A40-A41) | 109 | 1.3 | 4.3 |
| 12 | Viral hepatitis (B15-B19) | 106 | 1.2 | 4.2 |
| 13 | Assault (homicide) (X85-Y09,Y87.1) | 73 | 0.9 | 2.9 |
| 14 | Intentional self-harm (suicide) (X60-X84,Y87.0) | 63 | 0.7 | 2.5 |
| 15 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 55 | 0.6 | 2.2 |
| ... | All other causes (Residual) | 1,222 | 14.3 | 48.2 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|-------------------|---|-----------|-------------------------|---------|
| | Hispanic, female, 65 years and over | | | |
| ... | All causes | 30,881 | 100.0 | 2,741.5 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 10,406 | 33.7 | 923.8 |
| 2 | Malignant neoplasms (C00-C97) | 5,709 | 18.5 | 506.8 |
| 3 | Cerebrovascular diseases (I60-I69) | 2,638 | 8.5 | 234.2 |
| 4 | Diabetes mellitus (E10-E14) | 2,109 | 6.8 | 187.2 |
| 5 | Influenza and pneumonia (J10-J18) | 1,101 | 3.6 | 97.7 |
| 6 | Chronic lower respiratory diseases (J40-J47) | 1,033 | 3.3 | 91.7 |
| 7 | Alzheimer's disease (G30) | 732 | 2.4 | 65.0 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 634 | 2.1 | 56.3 |
| 9 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 474 | 1.5 | 42.1 |
| 10 | Chronic liver disease and cirrhosis (K70,K73-K74) | 436 | 1.4 | 38.7 |
| 11 | Septicemia (A40-A41) | 406 | 1.3 | 36.0 |
| 12 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 368 | 1.2 | 32.7 |
| 13 | Atherosclerosis (I70) | 231 | 0.7 | 20.5 |
| 14 | Pneumonitis due to solids and liquids (J69) | 206 | 0.7 | 18.3 |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 184 | 0.6 | 16.3 |
| ... | All other causes (Residual) | 4,214 | 13.6 | 374.1 |
| | White, non-Hispanic, both sexes, all ages | | | |
| ... | All causes | 1,959,919 | 100.0 | 996.6 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 594,465 | 30.3 | 302.3 |
| 2 | Malignant neoplasms (C00-C97) | 458,200 | 23.4 | 233.0 |
| 3 | Cerebrovascular diseases (I60-I69) | 138,210 | 7.1 | 70.3 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 109,901 | 5.6 | 55.9 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 73,536 | 3.8 | 37.4 |
| 6 | Influenza and pneumonia (J10-J18) | 55,135 | 2.8 | 28.0 |
| 7 | Diabetes mellitus (E10-E14) | 50,163 | 2.6 | 25.5 |
| 8 | Alzheimer's disease (G30) | 45,246 | 2.3 | 23.0 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 27,924 | 1.4 | 14.2 |
| 10 | Intentional self-harm (suicide) (X60-X84,Y87.0) | 24,616 | 1.3 | 12.5 |
| 11 | Septicemia (A40-A41) | 23,517 | 1.2 | 12.0 |
| 12 | Chronic liver disease and cirrhosis (K70,K73-K74) | 19,739 | 1.0 | 10.0 |
| 13 | Parkinson's disease (G20-G21) | 14,497 | 0.7 | 7.4 |
| 14 | Pneumonitis due to solids and liquids (J69) | 14,343 | 0.7 | 7.3 |
| 15 | Aortic aneurysm and dissection (I71) | 13,722 | 0.7 | 7.0 |
| ... | All other causes (Residual) | 296,705 | 15.1 | 150.9 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|-------------------|---|--------|-------------------------|------|
| | White, non-Hispanic, both sexes, 1-4 years | | | |
| ... | All causes | 2,641 | 100.0 | 27.9 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,033 | 39.1 | 10.9 |
| 2 | Malignant neoplasms (C00-C97) | 256 | 9.7 | 2.7 |
| 3 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 240 | 9.1 | 2.5 |
| 4 | Assault (homicide) (X85-Y09,Y87.1) | 144 | 5.5 | 1.5 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 98 | 3.7 | 1.0 |
| 6 | Septicemia (A40-A41) | 51 | 1.9 | 0.5 |
| 6 | Influenza and pneumonia (J10-J18) | 51 | 1.9 | 0.5 |
| 8 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 37 | 1.4 | 0.4 |
| 9 | Certain conditions originating in the perinatal period (P00-P96) | 34 | 1.3 | 0.4 |
| 10 | Cerebrovascular diseases (I60-I69) | 25 | 0.9 | 0.3 |
| 11 | Meningitis (G00,G03) | 19 | 0.7 | * |
| 11 | Chronic lower respiratory diseases (J40-J47) | 19 | 0.7 | * |
| 13 | Anemias (D50-D64) | 14 | 0.5 | * |
| 14 | Meningococcal infection (A39) | 11 | 0.4 | * |
| 15 | Complications of medical and surgical care (Y40-Y84,Y88) | 8 | 0.3 | * |
| ... | All other causes (Residual) | 601 | 22.8 | 6.4 |
| | White, non-Hispanic, both sexes, 5-14 years | | | |
| ... | All causes | 4,397 | 100.0 | 17.2 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,843 | 41.9 | 7.2 |
| 2 | Malignant neoplasms (C00-C97) | 627 | 14.3 | 2.5 |
| 3 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 240 | 5.5 | 0.9 |
| 4 | Intentional self-harm (suicide) (X60-X84,Y87.0) | 225 | 5.1 | 0.9 |
| 5 | Assault (homicide) (X85-Y09,Y87.1) | 164 | 3.7 | 0.6 |
| 6 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 140 | 3.2 | 0.5 |
| 7 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 63 | 1.4 | 0.2 |
| 8 | Influenza and pneumonia (J10-J18) | 49 | 1.1 | 0.2 |
| 9 | Chronic lower respiratory diseases (J40-J47) | 47 | 1.1 | 0.2 |
| 10 | Cerebrovascular diseases (I60-I69) | 46 | 1.0 | 0.2 |
| 11 | Septicemia (A40-A41) | 42 | 1.0 | 0.2 |
| 12 | Certain conditions originating in the perinatal period (P00-P96) | 20 | 0.5 | 0.1 |
| 13 | Meningitis (G00,G03) | 18 | 0.4 | * |
| 14 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 17 | 0.4 | * |
| 14 | Complications of medical and surgical care (Y40-Y84,Y88) | 17 | 0.4 | * |
| ... | All other causes (Residual) | 839 | 19.1 | 3.3 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|-------|
| White, non-Hispanic, both sexes, 15-24 years | | | | |
| ... | All causes | 18,261 | 100.0 | 72.1 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 9,735 | 53.3 | 38.4 |
| 2 | Intentional self-harm (suicide) (X60-X84,Y87.0) | 2,868 | 15.7 | 11.3 |
| 3 | Malignant neoplasms (C00-C97) | 1,059 | 5.8 | 4.2 |
| 4 | Assault (homicide) (X85-Y09,Y87.1) | 874 | 4.8 | 3.5 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 531 | 2.9 | 2.1 |
| 6 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 279 | 1.5 | 1.1 |
| 7 | Cerebrovascular diseases (I60-I69) | 118 | 0.6 | 0.5 |
| 8 | Influenza and pneumonia (J10-J18) | 116 | 0.6 | 0.5 |
| 9 | Diabetes mellitus (E10-E14) | 92 | 0.5 | 0.4 |
| 10 | Chronic lower respiratory diseases (J40-J47) | 87 | 0.5 | 0.3 |
| 11 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 63 | 0.3 | 0.2 |
| 12 | Septicemia (A40-A41) | 48 | 0.3 | 0.2 |
| 12 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 48 | 0.3 | 0.2 |
| 14 | Meningococcal infection (A39) | 37 | 0.2 | 0.1 |
| 15 | Pregnancy, childbirth and the puerperium (O00-099) | 33 | 0.2 | 0.1 |
| ... | All other causes (Residual) | 2,273 | 12.4 | 9.0 |
| White, non-Hispanic, both sexes, 25-44 years | | | | |
| ... | All causes | 81,422 | 100.0 | 141.7 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 18,738 | 23.0 | 32.6 |
| 2 | Malignant neoplasms (C00-C97) | 14,057 | 17.3 | 24.5 |
| 3 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 10,396 | 12.8 | 18.1 |
| 4 | Intentional self-harm (suicide) (X60-X84,Y87.0) | 9,198 | 11.3 | 16.0 |
| 5 | Human immunodeficiency virus (HIV) disease (B20-B24) | 2,619 | 3.2 | 4.6 |
| 6 | Chronic liver disease and cirrhosis (K70,K73-K74) | 2,457 | 3.0 | 4.3 |
| 7 | Assault (homicide) (X85-Y09,Y87.1) | 2,296 | 2.8 | 4.0 |
| 8 | Cerebrovascular diseases (I60-I69) | 1,691 | 2.1 | 2.9 |
| 9 | Diabetes mellitus (E10-E14) | 1,617 | 2.0 | 2.8 |
| 10 | Influenza and pneumonia (J10-J18) | 891 | 1.1 | 1.6 |
| 11 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 763 | 0.9 | 1.3 |
| 12 | Chronic lower respiratory diseases (J40-J47) | 714 | 0.9 | 1.2 |
| 13 | Septicemia (A40-A41) | 632 | 0.8 | 1.1 |
| 14 | Viral hepatitis (B15-B19) | 538 | 0.7 | 0.9 |
| 15 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 469 | 0.6 | 0.8 |
| ... | All other causes (Residual) | 14,346 | 17.6 | 25.0 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|-------------------|---|-----------|-------------------------|---------|
| | White, non-Hispanic, both sexes, 45-64 years | | | |
| ... | All causes | 294,312 | 100.0 | 619.9 |
| 1 | Malignant neoplasms (C00-C97) | 106,726 | 36.3 | 224.8 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 72,445 | 24.6 | 152.6 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 14,635 | 5.0 | 30.8 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 11,716 | 4.0 | 24.7 |
| 5 | Cerebrovascular diseases (I60-I69) | 9,988 | 3.4 | 21.0 |
| 6 | Diabetes mellitus (E10-E14) | 9,001 | 3.1 | 19.0 |
| 7 | Chronic liver disease and cirrhosis (K70,K73-K74) | 8,872 | 3.0 | 18.7 |
| 8 | Intentional self-harm (suicide) (X60-X84,Y87.0) | 7,433 | 2.5 | 15.7 |
| 9 | Influenza and pneumonia (J10-J18) | 3,226 | 1.1 | 6.8 |
| 10 | Septicemia (A40-A41) | 3,011 | 1.0 | 6.3 |
| 11 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 2,605 | 0.9 | 5.5 |
| 12 | Viral hepatitis (B15-B19) | 1,763 | 0.6 | 3.7 |
| 13 | Human immunodeficiency virus (HIV) disease (B20-B24) | 1,745 | 0.6 | 3.7 |
| 14 | Aortic aneurysm and dissection (I71) | 1,655 | 0.6 | 3.5 |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 1,520 | 0.5 | 3.2 |
| ... | All other causes (Residual) | 37,971 | 12.9 | 80.0 |
| | White, non-Hispanic, both sexes, 65 years and over | | | |
| ... | All causes | 1,545,284 | 100.0 | 5,318.5 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 510,568 | 33.0 | 1,757.2 |
| 2 | Malignant neoplasms (C00-C97) | 335,408 | 21.7 | 1,154.4 |
| 3 | Cerebrovascular diseases (I60-I69) | 126,283 | 8.2 | 434.6 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 97,301 | 6.3 | 334.9 |
| 5 | Influenza and pneumonia (J10-J18) | 50,687 | 3.3 | 174.5 |
| 6 | Alzheimer's disease (G30) | 44,760 | 2.9 | 154.1 |
| 7 | Diabetes mellitus (E10-E14) | 39,432 | 2.6 | 135.7 |
| 8 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 27,089 | 1.8 | 93.2 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 24,720 | 1.6 | 85.1 |
| 10 | Septicemia (A40-A41) | 19,624 | 1.3 | 67.5 |
| 11 | Parkinson's disease (G20-G21) | 14,217 | 0.9 | 48.9 |
| 12 | Pneumonitis due to solids and liquids (J69) | 13,419 | 0.9 | 46.2 |
| 13 | Atherosclerosis (I70) | 12,233 | 0.8 | 42.1 |
| 14 | Aortic aneurysm and dissection (I71) | 11,805 | 0.8 | 40.6 |
| 15 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 11,637 | 0.8 | 40.1 |
| ... | All other causes (Residual) | 206,101 | 13.3 | 709.3 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|-------------------|---|---------|-------------------------|-------|
| | White, non-Hispanic, male, all ages | | | |
| ... | All causes | 944,781 | 100.0 | 980.9 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 287,210 | 30.4 | 298.2 |
| 2 | Malignant neoplasms (C00-C97) | 235,932 | 25.0 | 245.0 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 53,231 | 5.6 | 55.3 |
| 4 | Cerebrovascular diseases (I60-I69) | 52,000 | 5.5 | 54.0 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 46,470 | 4.9 | 48.2 |
| 6 | Influenza and pneumonia (J10-J18) | 23,609 | 2.5 | 24.5 |
| 7 | Diabetes mellitus (E10-E14) | 23,465 | 2.5 | 24.4 |
| 8 | Intentional self-harm (suicide) (X60-X84,Y87.0) | 19,711 | 2.1 | 20.5 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 13,567 | 1.4 | 14.1 |
| 10 | Alzheimer's disease (G30) | 13,123 | 1.4 | 13.6 |
| 11 | Chronic liver disease and cirrhosis (K70,K73-K74) | 12,640 | 1.3 | 13.1 |
| 12 | Septicemia (A40-A41) | 10,056 | 1.1 | 10.4 |
| 13 | Aortic aneurysm and dissection (I71) | 8,488 | 0.9 | 8.8 |
| 14 | Parkinson's disease (G20-G21) | 8,116 | 0.9 | 8.4 |
| 15 | Pneumonitis due to solids and liquids (J69) | 7,214 | 0.8 | 7.5 |
| ... | All other causes (Residual) | 129,949 | 13.8 | 134.9 |
| | White, non-Hispanic, male, 1-4 years | | | |
| ... | All causes | 1,541 | 100.0 | 31.8 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 638 | 41.4 | 13.2 |
| 2 | Malignant neoplasms (C00-C97) | 146 | 9.5 | 3.0 |
| 3 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 124 | 8.0 | 2.6 |
| 4 | Assault (homicide) (X85-Y09,Y87.1) | 86 | 5.6 | 1.8 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 59 | 3.8 | 1.2 |
| 6 | Septicemia (A40-A41) | 26 | 1.7 | 0.5 |
| 7 | Influenza and pneumonia (J10-J18) | 24 | 1.6 | 0.5 |
| 8 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 23 | 1.5 | 0.5 |
| 8 | Certain conditions originating in the perinatal period (P00-P96) | 23 | 1.5 | 0.5 |
| 10 | Cerebrovascular diseases (I60-I69) | 14 | 0.9 | * |
| 11 | Meningitis (G00,G03) | 12 | 0.8 | * |
| 12 | Anemias (D50-D64) | 8 | 0.5 | * |
| 12 | Chronic lower respiratory diseases (J40-J47) | 8 | 0.5 | * |
| 14 | Meningococcal infection (A39) | 6 | 0.4 | * |
| 14 | Complications of medical and surgical care (Y40-Y84,Y88) | 6 | 0.4 | * |
| ... | All other causes (Residual) | 338 | 21.9 | 7.0 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|-------|
| White, non-Hispanic, male, 5-14 years | | | | |
| ... | All causes | 2,652 | 100.0 | 20.2 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,176 | 44.3 | 9.0 |
| 2 | Malignant neoplasms (C00-C97) | 338 | 12.7 | 2.6 |
| 3 | Intentional self-harm (suicide) (X60-X84,Y87.0) | 179 | 6.7 | 1.4 |
| 4 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 119 | 4.5 | 0.9 |
| 5 | Assault (homicide) (X85-Y09,Y87.1) | 93 | 3.5 | 0.7 |
| 6 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 87 | 3.3 | 0.7 |
| 7 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 31 | 1.2 | 0.2 |
| 8 | Influenza and pneumonia (J10-J18) | 29 | 1.1 | 0.2 |
| 9 | Septicemia (A40-A41) | 25 | 0.9 | 0.2 |
| 10 | Chronic lower respiratory diseases (J40-J47) | 24 | 0.9 | 0.2 |
| 11 | Cerebrovascular diseases (I60-I69) | 23 | 0.9 | 0.2 |
| 12 | Certain conditions originating in the perinatal period (P00-P96) | 14 | 0.5 | * |
| 13 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 12 | 0.5 | * |
| 14 | Anemias (D50-D64) | 9 | 0.3 | * |
| 14 | Meningitis (G00,G03) | 9 | 0.3 | * |
| ... | All other causes (Residual) | 484 | 18.3 | 3.7 |
| White, non-Hispanic, male, 15-24 years | | | | |
| ... | All causes | 13,066 | 100.0 | 100.5 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 7,065 | 54.1 | 54.4 |
| 2 | Intentional self-harm (suicide) (X60-X84,Y87.0) | 2,467 | 18.9 | 19.0 |
| 3 | Malignant neoplasms (C00-C97) | 636 | 4.9 | 4.9 |
| 4 | Assault (homicide) (X85-Y09,Y87.1) | 588 | 4.5 | 4.5 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 326 | 2.5 | 2.5 |
| 6 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 163 | 1.2 | 1.3 |
| 7 | Influenza and pneumonia (J10-J18) | 68 | 0.5 | 0.5 |
| 8 | Cerebrovascular diseases (I60-I69) | 67 | 0.5 | 0.5 |
| 9 | Diabetes mellitus (E10-E14) | 57 | 0.4 | 0.4 |
| 10 | Chronic lower respiratory diseases (J40-J47) | 52 | 0.4 | 0.4 |
| 11 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 40 | 0.3 | 0.3 |
| 12 | Septicemia (A40-A41) | 33 | 0.3 | 0.3 |
| 13 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 25 | 0.2 | 0.2 |
| 13 | Legal intervention (Y35,Y89.0) | 25 | 0.2 | 0.2 |
| 15 | Meningococcal infection (A39) | 22 | 0.2 | 0.2 |
| ... | All other causes (Residual) | 1,432 | 11.0 | 11.0 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|-------------------|---|---------|-------------------------|-------|
| | White, non-Hispanic, male, 25-44 years | | | |
| ... | All causes | 53,556 | 100.0 | 186.4 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 13,974 | 26.1 | 48.6 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 7,561 | 14.1 | 26.3 |
| 3 | Intentional self-harm (suicide) (X60-X84,Y87.0) | 7,276 | 13.6 | 25.3 |
| 4 | Malignant neoplasms (C00-C97) | 6,460 | 12.1 | 22.5 |
| 5 | Human immunodeficiency virus (HIV) disease (B20-B24) | 2,153 | 4.0 | 7.5 |
| 6 | Chronic liver disease and cirrhosis (K70,K73-K74) | 1,658 | 3.1 | 5.8 |
| 7 | Assault (homicide) (X85-Y09,Y87.1) | 1,513 | 2.8 | 5.3 |
| 8 | Diabetes mellitus (E10-E14) | 986 | 1.8 | 3.4 |
| 9 | Cerebrovascular diseases (I60-I69) | 855 | 1.6 | 3.0 |
| 10 | Influenza and pneumonia (J10-J18) | 505 | 0.9 | 1.8 |
| 11 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 436 | 0.8 | 1.5 |
| 12 | Viral hepatitis (B15-B19) | 361 | 0.7 | 1.3 |
| 13 | Chronic lower respiratory diseases (J40-J47) | 338 | 0.6 | 1.2 |
| 14 | Septicemia (A40-A41) | 336 | 0.6 | 1.2 |
| 15 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 268 | 0.5 | 0.9 |
| ... | All other causes (Residual) | 8,876 | 16.6 | 30.9 |
| | White, non-Hispanic, male, 45-64 years | | | |
| ... | All causes | 179,819 | 100.0 | 771.7 |
| 1 | Malignant neoplasms (C00-C97) | 56,460 | 31.4 | 242.3 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 52,280 | 29.1 | 224.4 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 10,392 | 5.8 | 44.6 |
| 4 | Chronic liver disease and cirrhosis (K70,K73-K74) | 6,412 | 3.6 | 27.5 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 5,987 | 3.3 | 25.7 |
| 6 | Intentional self-harm (suicide) (X60-X84,Y87.0) | 5,658 | 3.1 | 24.3 |
| 7 | Cerebrovascular diseases (I60-I69) | 5,461 | 3.0 | 23.4 |
| 8 | Diabetes mellitus (E10-E14) | 5,140 | 2.9 | 22.1 |
| 9 | Influenza and pneumonia (J10-J18) | 1,843 | 1.0 | 7.9 |
| 10 | Septicemia (A40-A41) | 1,628 | 0.9 | 7.0 |
| 11 | Human immunodeficiency virus (HIV) disease (B20-B24) | 1,530 | 0.9 | 6.6 |
| 12 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1,481 | 0.8 | 6.4 |
| 13 | Viral hepatitis (B15-B19) | 1,297 | 0.7 | 5.6 |
| 14 | Aortic aneurysm and dissection (I71) | 1,269 | 0.7 | 5.4 |
| 15 | Assault (homicide) (X85-Y09,Y87.1) | 848 | 0.5 | 3.6 |
| ... | All other causes (Residual) | 22,133 | 12.3 | 95.0 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|-------------------|---|-----------|-------------------------|---------|
| | White, non-Hispanic, male, 65 years and over | | | |
| ... | All causes | 686,417 | 100.0 | 5,660.3 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 226,738 | 33.0 | 1,869.7 |
| 2 | Malignant neoplasms (C00-C97) | 171,851 | 25.0 | 1,417.1 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 46,809 | 6.8 | 386.0 |
| 4 | Cerebrovascular diseases (I60-I69) | 45,542 | 6.6 | 375.5 |
| 5 | Influenza and pneumonia (J10-J18) | 21,071 | 3.1 | 173.8 |
| 6 | Diabetes mellitus (E10-E14) | 17,271 | 2.5 | 142.4 |
| 7 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 12,952 | 1.9 | 106.8 |
| 8 | Alzheimer's disease (G30) | 12,910 | 1.9 | 106.5 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 11,738 | 1.7 | 96.8 |
| 10 | Septicemia (A40-A41) | 7,952 | 1.2 | 65.6 |
| 11 | Parkinson's disease (G20-G21) | 7,938 | 1.2 | 65.5 |
| 12 | Aortic aneurysm and dissection (I71) | 7,025 | 1.0 | 57.9 |
| 13 | Pneumonitis due to solids and liquids (J69) | 6,652 | 1.0 | 54.9 |
| 14 | Chronic liver disease and cirrhosis (K70,K73-K74) | 4,561 | 0.7 | 37.6 |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 4,511 | 0.7 | 37.2 |
| ... | All other causes (Residual) | 80,896 | 11.8 | 667.1 |
| | White, non-Hispanic, female, all ages | | | |
| ... | All causes | 1,015,138 | 100.0 | 1,011.7 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 307,255 | 30.3 | 306.2 |
| 2 | Malignant neoplasms (C00-C97) | 222,268 | 21.9 | 221.5 |
| 3 | Cerebrovascular diseases (I60-I69) | 86,210 | 8.5 | 85.9 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 56,670 | 5.6 | 56.5 |
| 5 | Alzheimer's disease (G30) | 32,123 | 3.2 | 32.0 |
| 6 | Influenza and pneumonia (J10-J18) | 31,526 | 3.1 | 31.4 |
| 7 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 27,066 | 2.7 | 27.0 |
| 8 | Diabetes mellitus (E10-E14) | 26,698 | 2.6 | 26.6 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 14,357 | 1.4 | 14.3 |
| 10 | Septicemia (A40-A41) | 13,461 | 1.3 | 13.4 |
| 11 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 8,196 | 0.8 | 8.2 |
| 12 | Atherosclerosis (I70) | 8,052 | 0.8 | 8.0 |
| 13 | Pneumonitis due to solids and liquids (J69) | 7,129 | 0.7 | 7.1 |
| 14 | Chronic liver disease and cirrhosis (K70,K73-K74) | 7,099 | 0.7 | 7.1 |
| 15 | Parkinson's disease (G20-G21) | 6,381 | 0.6 | 6.4 |
| ... | All other causes (Residual) | 160,647 | 15.8 | 160.1 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|---|---|--------|-------------------------|------|
| White, non-Hispanic, female, 1-4 years | | | | |
| ... | All causes | 1,100 | 100.0 | 23.9 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 395 | 35.9 | 8.6 |
| 2 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 116 | 10.5 | 2.5 |
| 3 | Malignant neoplasms (C00-C97) | 110 | 10.0 | 2.4 |
| 4 | Assault (homicide) (X85-Y09,Y87.1) | 58 | 5.3 | 1.3 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 39 | 3.5 | 0.8 |
| 6 | Influenza and pneumonia (J10-J18) | 27 | 2.5 | 0.6 |
| 7 | Septicemia (A40-A41) | 25 | 2.3 | 0.5 |
| 8 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 14 | 1.3 | * |
| 9 | Cerebrovascular diseases (I60-I69) | 11 | 1.0 | * |
| 9 | Chronic lower respiratory diseases (J40-J47) | 11 | 1.0 | * |
| 9 | Certain conditions originating in the perinatal period (P00-P96) | 11 | 1.0 | * |
| 12 | Meningitis (G00,G03) | 7 | 0.6 | * |
| 13 | Anemias (D50-D64) | 6 | 0.5 | * |
| 14 | Meningococcal infection (A39) | 5 | 0.5 | * |
| 15 | Diabetes mellitus (E10-E14) | 4 | 0.4 | * |
| ... | All other causes (Residual) | 261 | 23.7 | 5.7 |
| White, non-Hispanic, female, 5-14 years | | | | |
| ... | All causes | 1,745 | 100.0 | 14.0 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 667 | 38.2 | 5.4 |
| 2 | Malignant neoplasms (C00-C97) | 289 | 16.6 | 2.3 |
| 3 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 121 | 6.9 | 1.0 |
| 4 | Assault (homicide) (X85-Y09,Y87.1) | 71 | 4.1 | 0.6 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 53 | 3.0 | 0.4 |
| 6 | Intentional self-harm (suicide) (X60-X84,Y87.0) | 46 | 2.6 | 0.4 |
| 7 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 32 | 1.8 | 0.3 |
| 8 | Cerebrovascular diseases (I60-I69) | 23 | 1.3 | 0.2 |
| 8 | Chronic lower respiratory diseases (J40-J47) | 23 | 1.3 | 0.2 |
| 10 | Influenza and pneumonia (J10-J18) | 20 | 1.1 | 0.2 |
| 11 | Septicemia (A40-A41) | 17 | 1.0 | * |
| 12 | Complications of medical and surgical care (Y40-Y84,Y88) | 10 | 0.6 | * |
| 13 | Meningitis (G00,G03) | 9 | 0.5 | * |
| 14 | Human immunodeficiency virus (HIV) disease (B20-B24) | 8 | 0.5 | * |
| 14 | Diseases of appendix (K35-K38) | 8 | 0.5 | * |
| ... | All other causes (Residual) | 348 | 19.9 | 2.8 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|------|
| White, non-Hispanic, female, 15-24 years | | | | |
| ... | All causes | 5,195 | 100.0 | 42.2 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 2,670 | 51.4 | 21.7 |
| 2 | Malignant neoplasms (C00-C97) | 423 | 8.1 | 3.4 |
| 3 | Intentional self-harm (suicide) (X60-X84,Y87.0) | 401 | 7.7 | 3.3 |
| 4 | Assault (homicide) (X85-Y09,Y87.1) | 286 | 5.5 | 2.3 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 205 | 3.9 | 1.7 |
| 6 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 116 | 2.2 | 0.9 |
| 7 | Cerebrovascular diseases (I60-I69) | 51 | 1.0 | 0.4 |
| 8 | Influenza and pneumonia (J10-J18) | 48 | 0.9 | 0.4 |
| 9 | Diabetes mellitus (E10-E14) | 35 | 0.7 | 0.3 |
| 9 | Chronic lower respiratory diseases (J40-J47) | 35 | 0.7 | 0.3 |
| 11 | Pregnancy, childbirth and the puerperium (O00-099) | 33 | 0.6 | 0.3 |
| 12 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 23 | 0.4 | 0.2 |
| 12 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 23 | 0.4 | 0.2 |
| 14 | Meningococcal infection (A39) | 15 | 0.3 | * |
| 14 | Septicemia (A40-A41) | 15 | 0.3 | * |
| ... | All other causes (Residual) | 816 | 15.7 | 6.6 |
| White, non-Hispanic, female, 25-44 years | | | | |
| ... | All causes | 27,866 | 100.0 | 97.0 |
| 1 | Malignant neoplasms (C00-C97) | 7,597 | 27.3 | 26.5 |
| 2 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 4,764 | 17.1 | 16.6 |
| 3 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 2,835 | 10.2 | 9.9 |
| 4 | Intentional self-harm (suicide) (X60-X84,Y87.0) | 1,922 | 6.9 | 6.7 |
| 5 | Cerebrovascular diseases (I60-I69) | 836 | 3.0 | 2.9 |
| 6 | Chronic liver disease and cirrhosis (K70,K73-K74) | 799 | 2.9 | 2.8 |
| 7 | Assault (homicide) (X85-Y09,Y87.1) | 783 | 2.8 | 2.7 |
| 8 | Diabetes mellitus (E10-E14) | 631 | 2.3 | 2.2 |
| 9 | Human immunodeficiency virus (HIV) disease (B20-B24) | 466 | 1.7 | 1.6 |
| 10 | Influenza and pneumonia (J10-J18) | 386 | 1.4 | 1.3 |
| 11 | Chronic lower respiratory diseases (J40-J47) | 376 | 1.3 | 1.3 |
| 12 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 327 | 1.2 | 1.1 |
| 13 | Septicemia (A40-A41) | 296 | 1.1 | 1.0 |
| 14 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 201 | 0.7 | 0.7 |
| 15 | Viral hepatitis (B15-B19) | 177 | 0.6 | 0.6 |
| ... | All other causes (Residual) | 5,470 | 19.6 | 19.0 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|-------------------|---|---------|-------------------------|---------|
| | White, non-Hispanic, female, 45-64 years | | | |
| ... | All causes | 114,493 | 100.0 | 473.5 |
| 1 | Malignant neoplasms (C00-C97) | 50,266 | 43.9 | 207.9 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 20,165 | 17.6 | 83.4 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 5,729 | 5.0 | 23.7 |
| 4 | Cerebrovascular diseases (I60-I69) | 4,527 | 4.0 | 18.7 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 4,243 | 3.7 | 17.5 |
| 6 | Diabetes mellitus (E10-E14) | 3,861 | 3.4 | 16.0 |
| 7 | Chronic liver disease and cirrhosis (K70,K73-K74) | 2,460 | 2.1 | 10.2 |
| 8 | Intentional self-harm (suicide) (X60-X84,Y87.0) | 1,775 | 1.6 | 7.3 |
| 9 | Septicemia (A40-A41) | 1,383 | 1.2 | 5.7 |
| 9 | Influenza and pneumonia (J10-J18) | 1,383 | 1.2 | 5.7 |
| 11 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1,124 | 1.0 | 4.6 |
| 12 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 680 | 0.6 | 2.8 |
| 13 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 468 | 0.4 | 1.9 |
| 14 | Viral hepatitis (B15-B19) | 466 | 0.4 | 1.9 |
| 15 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 428 | 0.4 | 1.8 |
| ... | All other causes (Residual) | 15,535 | 13.6 | 64.3 |
| | White, non-Hispanic, female, 65 years and over | | | |
| ... | All causes | 858,867 | 100.0 | 5,073.6 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 283,830 | 33.0 | 1,676.7 |
| 2 | Malignant neoplasms (C00-C97) | 163,557 | 19.0 | 966.2 |
| 3 | Cerebrovascular diseases (I60-I69) | 80,741 | 9.4 | 477.0 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 50,492 | 5.9 | 298.3 |
| 5 | Alzheimer's disease (G30) | 31,850 | 3.7 | 188.1 |
| 6 | Influenza and pneumonia (J10-J18) | 29,616 | 3.4 | 175.0 |
| 7 | Diabetes mellitus (E10-E14) | 22,161 | 2.6 | 130.9 |
| 8 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 14,137 | 1.6 | 83.5 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 12,982 | 1.5 | 76.7 |
| 10 | Septicemia (A40-A41) | 11,672 | 1.4 | 69.0 |
| 11 | Atherosclerosis (I70) | 7,895 | 0.9 | 46.6 |
| 12 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 7,715 | 0.9 | 45.6 |
| 13 | Pneumonitis due to solids and liquids (J69) | 6,767 | 0.8 | 40.0 |
| 14 | Parkinson's disease (G20-G21) | 6,279 | 0.7 | 37.1 |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 5,041 | 0.6 | 29.8 |
| ... | All other causes (Residual) | 124,132 | 14.5 | 733.3 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|-------------------|---|---------|-------------------------|-------|
| | Black, non-Hispanic, both sexes, all ages | | | |
| ... | All causes | 282,676 | 100.0 | 844.4 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 76,706 | 27.1 | 229.1 |
| 2 | Malignant neoplasms (C00-C97) | 61,390 | 21.7 | 183.4 |
| 3 | Cerebrovascular diseases (I60-I69) | 19,053 | 6.7 | 56.9 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 12,087 | 4.3 | 36.1 |
| 5 | Diabetes mellitus (E10-E14) | 11,918 | 4.2 | 35.6 |
| 6 | Assault (homicide) (X85-Y09,Y87.1) | 7,748 | 2.7 | 23.1 |
| 7 | Human immunodeficiency virus (HIV) disease (B20-B24) | 7,708 | 2.7 | 23.0 |
| 8 | Chronic lower respiratory diseases (J40-J47) | 7,528 | 2.7 | 22.5 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 6,853 | 2.4 | 20.5 |
| 10 | Influenza and pneumonia (J10-J18) | 5,920 | 2.1 | 17.7 |
| 11 | Septicemia (A40-A41) | 5,863 | 2.1 | 17.5 |
| 12 | Certain conditions originating in the perinatal period (P00-P96) | 4,878 | 1.7 | 14.6 |
| 13 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 3,862 | 1.4 | 11.5 |
| 14 | Alzheimer's disease (G30) | 2,719 | 1.0 | 8.1 |
| 15 | Chronic liver disease and cirrhosis (K70,K73-K74) | 2,705 | 1.0 | 8.1 |
| ... | All other causes (Residual) | 45,738 | 16.2 | 136.6 |
| | Black, non-Hispanic, both sexes, 1-4 years | | | |
| ... | All causes | 1,217 | 100.0 | 58.8 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 396 | 32.5 | 19.1 |
| 2 | Assault (homicide) (X85-Y09,Y87.1) | 143 | 11.8 | 6.9 |
| 3 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 123 | 10.1 | 5.9 |
| 4 | Malignant neoplasms (C00-C97) | 66 | 5.4 | 3.2 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 47 | 3.9 | 2.3 |
| 6 | Influenza and pneumonia (J10-J18) | 29 | 2.4 | 1.4 |
| 7 | Certain conditions originating in the perinatal period (P00-P96) | 27 | 2.2 | 1.3 |
| 8 | Septicemia (A40-A41) | 25 | 2.1 | 1.2 |
| 9 | Chronic lower respiratory diseases (J40-J47) | 21 | 1.7 | 1.0 |
| 10 | Anemias (D50-D64) | 19 | 1.6 | * |
| 11 | Meningitis (G00,G03) | 13 | 1.1 | * |
| 11 | Cerebrovascular diseases (I60-I69) | 13 | 1.1 | * |
| 13 | Human immunodeficiency virus (HIV) disease (B20-B24) | 12 | 1.0 | * |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 9 | 0.7 | * |
| 15 | Complications of medical and surgical care (Y40-Y84,Y88) | 8 | 0.7 | * |
| ... | All other causes (Residual) | 266 | 21.9 | 12.8 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|-------------------|---|--------|-------------------------|-------|
| | Black, non-Hispanic, both sexes, 5-14 years | | | |
| ... | All causes | 1,620 | 100.0 | 27.5 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 619 | 38.2 | 10.5 |
| 2 | Malignant neoplasms (C00-C97) | 147 | 9.1 | 2.5 |
| 3 | Assault (homicide) (X85-Y09,Y87.1) | 122 | 7.5 | 2.1 |
| 4 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 89 | 5.5 | 1.5 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 75 | 4.6 | 1.3 |
| 6 | Chronic lower respiratory diseases (J40-J47) | 70 | 4.3 | 1.2 |
| 7 | Intentional self-harm (suicide) (X60-X84,Y87.0) | 39 | 2.4 | 0.7 |
| 8 | Anemias (D50-D64) | 30 | 1.9 | 0.5 |
| 9 | Human immunodeficiency virus (HIV) disease (B20-B24) | 23 | 1.4 | 0.4 |
| 10 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 18 | 1.1 | * |
| 11 | Cerebrovascular diseases (I60-I69) | 15 | 0.9 | * |
| 12 | Septicemia (A40-A41) | 11 | 0.7 | * |
| 12 | Influenza and pneumonia (J10-J18) | 11 | 0.7 | * |
| 14 | Certain conditions originating in the perinatal period (P00-P96) | 10 | 0.6 | * |
| 15 | Diabetes mellitus (E10-E14) | 9 | 0.6 | * |
| 15 | Meningitis (G00,G03) | 9 | 0.6 | * |
| ... | All other causes (Residual) | 323 | 19.9 | 5.5 |
| | Black, non-Hispanic, both sexes, 15-24 years | | | |
| ... | All causes | 6,926 | 100.0 | 125.2 |
| 1 | Assault (homicide) (X85-Y09,Y87.1) | 2,771 | 40.0 | 50.1 |
| 2 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,662 | 24.0 | 30.0 |
| 3 | Intentional self-harm (suicide) (X60-X84,Y87.0) | 471 | 6.8 | 8.5 |
| 4 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 331 | 4.8 | 6.0 |
| 5 | Malignant neoplasms (C00-C97) | 297 | 4.3 | 5.4 |
| 6 | Human immunodeficiency virus (HIV) disease (B20-B24) | 125 | 1.8 | 2.3 |
| 7 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 91 | 1.3 | 1.6 |
| 8 | Chronic lower respiratory diseases (J40-J47) | 80 | 1.2 | 1.4 |
| 9 | Anemias (D50-D64) | 61 | 0.9 | 1.1 |
| 10 | Diabetes mellitus (E10-E14) | 55 | 0.8 | 1.0 |
| 11 | Influenza and pneumonia (J10-J18) | 44 | 0.6 | 0.8 |
| 12 | Cerebrovascular diseases (I60-I69) | 42 | 0.6 | 0.8 |
| 13 | Pregnancy, childbirth and the puerperium (O00-099) | 34 | 0.5 | 0.6 |
| 14 | Septicemia (A40-A41) | 33 | 0.5 | 0.6 |
| 15 | Legal intervention (Y35,Y89.0) | 27 | 0.4 | 0.5 |
| ... | All other causes (Residual) | 802 | 11.6 | 14.5 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|-------------------|---|--------|-------------------------|---------|
| | Black, non-Hispanic, both sexes, 25-44 years | | | |
| ... | All causes | 30,382 | 100.0 | 293.3 |
| 1 | Human immunodeficiency virus (HIV) disease (B20-B24) | 4,442 | 14.6 | 42.9 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 4,188 | 13.8 | 40.4 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 3,993 | 13.1 | 38.5 |
| 4 | Malignant neoplasms (C00-C97) | 3,713 | 12.2 | 35.8 |
| 5 | Assault (homicide) (X85-Y09,Y87.1) | 3,552 | 11.7 | 34.3 |
| 6 | Cerebrovascular diseases (I60-I69) | 987 | 3.2 | 9.5 |
| 7 | Intentional self-harm (suicide) (X60-X84,Y87.0) | 896 | 2.9 | 8.6 |
| 8 | Diabetes mellitus (E10-E14) | 645 | 2.1 | 6.2 |
| 9 | Chronic liver disease and cirrhosis (K70,K73-K74) | 502 | 1.7 | 4.8 |
| 10 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 404 | 1.3 | 3.9 |
| 11 | Chronic lower respiratory diseases (J40-J47) | 386 | 1.3 | 3.7 |
| 12 | Septicemia (A40-A41) | 376 | 1.2 | 3.6 |
| 13 | Influenza and pneumonia (J10-J18) | 346 | 1.1 | 3.3 |
| 14 | Anemias (D50-D64) | 263 | 0.9 | 2.5 |
| 15 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 211 | 0.7 | 2.0 |
| ... | All other causes (Residual) | 5,478 | 18.0 | 52.9 |
| | Black, non-Hispanic, both sexes, 45-64 years | | | |
| ... | All causes | 71,778 | 100.0 | 1,147.3 |
| 1 | Malignant neoplasms (C00-C97) | 20,087 | 28.0 | 321.1 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 18,863 | 26.3 | 301.5 |
| 3 | Cerebrovascular diseases (I60-I69) | 4,084 | 5.7 | 65.3 |
| 4 | Diabetes mellitus (E10-E14) | 3,265 | 4.5 | 52.2 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 2,944 | 4.1 | 47.1 |
| 6 | Human immunodeficiency virus (HIV) disease (B20-B24) | 2,836 | 4.0 | 45.3 |
| 7 | Chronic lower respiratory diseases (J40-J47) | 1,656 | 2.3 | 26.5 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1,644 | 2.3 | 26.3 |
| 9 | Chronic liver disease and cirrhosis (K70,K73-K74) | 1,551 | 2.2 | 24.8 |
| 10 | Septicemia (A40-A41) | 1,328 | 1.9 | 21.2 |
| 11 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 960 | 1.3 | 15.3 |
| 12 | Influenza and pneumonia (J10-J18) | 954 | 1.3 | 15.2 |
| 13 | Assault (homicide) (X85-Y09,Y87.1) | 816 | 1.1 | 13.0 |
| 14 | Viral hepatitis (B15-B19) | 467 | 0.7 | 7.5 |
| 15 | Intentional self-harm (suicide) (X60-X84,Y87.0) | 367 | 0.5 | 5.9 |
| ... | All other causes (Residual) | 9,956 | 13.9 | 159.1 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|-------------------|---|---------|-------------------------|---------|
| | Black, non-Hispanic, both sexes, 65 years and over | | | |
| ... | All causes | 162,179 | 100.0 | 5,759.3 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 53,072 | 32.7 | 1,884.7 |
| 2 | Malignant neoplasms (C00-C97) | 37,066 | 22.9 | 1,316.3 |
| 3 | Cerebrovascular diseases (I60-I69) | 13,865 | 8.5 | 492.4 |
| 4 | Diabetes mellitus (E10-E14) | 7,943 | 4.9 | 282.1 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 5,304 | 3.3 | 188.4 |
| 6 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 4,703 | 2.9 | 167.0 |
| 7 | Influenza and pneumonia (J10-J18) | 4,435 | 2.7 | 157.5 |
| 8 | Septicemia (A40-A41) | 3,994 | 2.5 | 141.8 |
| 9 | Alzheimer's disease (G30) | 2,683 | 1.7 | 95.3 |
| 10 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 2,680 | 1.7 | 95.2 |
| 11 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 2,201 | 1.4 | 78.2 |
| 12 | Pneumonitis due to solids and liquids (J69) | 1,262 | 0.8 | 44.8 |
| 13 | Atherosclerosis (I70) | 908 | 0.6 | 32.2 |
| 14 | Aortic aneurysm and dissection (I71) | 808 | 0.5 | 28.7 |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 729 | 0.4 | 25.9 |
| ... | All other causes (Residual) | 20,526 | 12.7 | 728.9 |
| | Black, non-Hispanic, male, all ages | | | |
| ... | All causes | 143,297 | 100.0 | 903.3 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 36,282 | 25.3 | 228.7 |
| 2 | Malignant neoplasms (C00-C97) | 32,493 | 22.7 | 204.8 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 8,380 | 5.8 | 52.8 |
| 4 | Cerebrovascular diseases (I60-I69) | 7,945 | 5.5 | 50.1 |
| 5 | Assault (homicide) (X85-Y09,Y87.1) | 6,382 | 4.5 | 40.2 |
| 6 | Human immunodeficiency virus (HIV) disease (B20-B24) | 5,291 | 3.7 | 33.4 |
| 7 | Diabetes mellitus (E10-E14) | 4,720 | 3.3 | 29.8 |
| 8 | Chronic lower respiratory diseases (J40-J47) | 4,184 | 2.9 | 26.4 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 3,045 | 2.1 | 19.2 |
| 10 | Influenza and pneumonia (J10-J18) | 2,880 | 2.0 | 18.2 |
| 11 | Certain conditions originating in the perinatal period (P00-P96) | 2,760 | 1.9 | 17.4 |
| 12 | Septicemia (A40-A41) | 2,559 | 1.8 | 16.1 |
| 13 | Chronic liver disease and cirrhosis (K70,K73-K74) | 1,709 | 1.2 | 10.8 |
| 14 | Intentional self-harm (suicide) (X60-X84,Y87.0) | 1,601 | 1.1 | 10.1 |
| 15 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 1,576 | 1.1 | 9.9 |
| ... | All other causes (Residual) | 21,490 | 15.0 | 135.5 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|---------------------------------------|---|--------|-------------------------|------|
| Black, non-Hispanic, male, 1-4 years | | | | |
| ... | All causes | 672 | 100.0 | 63.9 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 232 | 34.5 | 22.1 |
| 2 | Assault (homicide) (X85-Y09,Y87.1) | 73 | 10.9 | 6.9 |
| 3 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 67 | 10.0 | 6.4 |
| 4 | Malignant neoplasms (C00-C97) | 31 | 4.6 | 2.9 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 24 | 3.6 | 2.3 |
| 6 | Influenza and pneumonia (J10-J18) | 20 | 3.0 | 1.9 |
| 7 | Certain conditions originating in the perinatal period (P00-P96) | 18 | 2.7 | * |
| 8 | Chronic lower respiratory diseases (J40-J47) | 14 | 2.1 | * |
| 9 | Septicemia (A40-A41) | 13 | 1.9 | * |
| 10 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 8 | 1.2 | * |
| 10 | Anemias (D50-D64) | 8 | 1.2 | * |
| 12 | Human immunodeficiency virus (HIV) disease (B20-B24) | 6 | 0.9 | * |
| 12 | Meningitis (G00,G03) | 6 | 0.9 | * |
| 14 | Meningococcal infection (A39) | 4 | 0.6 | * |
| 15 | Cerebrovascular diseases (I60-I69) | 3 | 0.4 | * |
| ... | All other causes (Residual) | 145 | 21.6 | 13.8 |
| Black, non-Hispanic, male, 5-14 years | | | | |
| ... | All causes | 959 | 100.0 | 32.0 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 406 | 42.3 | 13.6 |
| 2 | Malignant neoplasms (C00-C97) | 86 | 9.0 | 2.9 |
| 3 | Assault (homicide) (X85-Y09,Y87.1) | 70 | 7.3 | 2.3 |
| 4 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 50 | 5.2 | 1.7 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 43 | 4.5 | 1.4 |
| 6 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 41 | 4.3 | 1.4 |
| 7 | Intentional self-harm (suicide) (X60-X84,Y87.0) | 34 | 3.5 | 1.1 |
| 8 | Anemias (D50-D64) | 17 | 1.8 | * |
| 9 | Human immunodeficiency virus (HIV) disease (B20-B24) | 12 | 1.3 | * |
| 10 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 8 | 0.8 | * |
| 11 | Influenza and pneumonia (J10-J18) | 7 | 0.7 | * |
| 12 | Septicemia (A40-A41) | 6 | 0.6 | * |
| 12 | Meningitis (G00,G03) | 6 | 0.6 | * |
| 14 | Cerebrovascular diseases (I60-I69) | 5 | 0.5 | * |
| 15 | Diabetes mellitus (E10-E14) | 4 | 0.4 | * |
| ... | All other causes (Residual) | 164 | 17.1 | 5.5 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|-------|
| Black, non-Hispanic, male, 15-24 years | | | | |
| ... | All causes | 5,224 | 100.0 | 188.2 |
| 1 | Assault (homicide) (X85-Y09,Y87.1) | 2,459 | 47.1 | 88.6 |
| 2 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,233 | 23.6 | 44.4 |
| 3 | Intentional self-harm (suicide) (X60-X84,Y87.0) | 407 | 7.8 | 14.7 |
| 4 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 192 | 3.7 | 6.9 |
| 5 | Malignant neoplasms (C00-C97) | 178 | 3.4 | 6.4 |
| 6 | Human immunodeficiency virus (HIV) disease (B20-B24) | 60 | 1.1 | 2.2 |
| 7 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 58 | 1.1 | 2.1 |
| 8 | Chronic lower respiratory diseases (J40-J47) | 51 | 1.0 | 1.8 |
| 9 | Anemias (D50-D64) | 29 | 0.6 | 1.0 |
| 10 | Diabetes mellitus (E10-E14) | 28 | 0.5 | 1.0 |
| 11 | Legal intervention (Y35,Y89.0) | 27 | 0.5 | 1.0 |
| 12 | Influenza and pneumonia (J10-J18) | 21 | 0.4 | 0.8 |
| 13 | Cerebrovascular diseases (I60-I69) | 19 | 0.4 | * |
| 14 | Septicemia (A40-A41) | 18 | 0.3 | * |
| 15 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 10 | 0.2 | * |
| ... | All other causes (Residual) | 434 | 8.3 | 15.6 |
| Black, non-Hispanic, male, 25-44 years | | | | |
| ... | All causes | 18,710 | 100.0 | 384.0 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 2,965 | 15.8 | 60.9 |
| 2 | Assault (homicide) (X85-Y09,Y87.1) | 2,910 | 15.6 | 59.7 |
| 3 | Human immunodeficiency virus (HIV) disease (B20-B24) | 2,882 | 15.4 | 59.1 |
| 4 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 2,515 | 13.4 | 51.6 |
| 5 | Malignant neoplasms (C00-C97) | 1,551 | 8.3 | 31.8 |
| 6 | Intentional self-harm (suicide) (X60-X84,Y87.0) | 743 | 4.0 | 15.2 |
| 7 | Cerebrovascular diseases (I60-I69) | 464 | 2.5 | 9.5 |
| 8 | Diabetes mellitus (E10-E14) | 350 | 1.9 | 7.2 |
| 9 | Chronic liver disease and cirrhosis (K70,K73-K74) | 305 | 1.6 | 6.3 |
| 10 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 225 | 1.2 | 4.6 |
| 11 | Influenza and pneumonia (J10-J18) | 209 | 1.1 | 4.3 |
| 12 | Chronic lower respiratory diseases (J40-J47) | 199 | 1.1 | 4.1 |
| 13 | Septicemia (A40-A41) | 184 | 1.0 | 3.8 |
| 14 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 114 | 0.6 | 2.3 |
| 15 | Anemias (D50-D64) | 110 | 0.6 | 2.3 |
| ... | All other causes (Residual) | 2,984 | 15.9 | 61.2 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|-------------------|---|--------|-------------------------|---------|
| | Black, non-Hispanic, male, 45-64 years | | | |
| ... | All causes | 42,049 | 100.0 | 1,509.7 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 11,573 | 27.5 | 415.5 |
| 2 | Malignant neoplasms (C00-C97) | 10,915 | 26.0 | 391.9 |
| 3 | Cerebrovascular diseases (I60-I69) | 2,195 | 5.2 | 78.8 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 2,190 | 5.2 | 78.6 |
| 5 | Human immunodeficiency virus (HIV) disease (B20-B24) | 2,134 | 5.1 | 76.6 |
| 6 | Diabetes mellitus (E10-E14) | 1,600 | 3.8 | 57.4 |
| 7 | Chronic liver disease and cirrhosis (K70,K73-K74) | 1,041 | 2.5 | 37.4 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 878 | 2.1 | 31.5 |
| 9 | Chronic lower respiratory diseases (J40-J47) | 876 | 2.1 | 31.5 |
| 10 | Septicemia (A40-A41) | 700 | 1.7 | 25.1 |
| 11 | Assault (homicide) (X85-Y09,Y87.1) | 657 | 1.6 | 23.6 |
| 12 | Influenza and pneumonia (J10-J18) | 593 | 1.4 | 21.3 |
| 13 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 530 | 1.3 | 19.0 |
| 14 | Viral hepatitis (B15-B19) | 320 | 0.8 | 11.5 |
| 15 | Intentional self-harm (suicide) (X60-X84,Y87.0) | 290 | 0.7 | 10.4 |
| ... | All other causes (Residual) | 5,557 | 13.2 | 199.5 |
| | Black, non-Hispanic, male, 65 years and over | | | |
| ... | All causes | 70,891 | 100.0 | 6,400.7 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 21,871 | 30.9 | 1,974.7 |
| 2 | Malignant neoplasms (C00-C97) | 19,726 | 27.8 | 1,781.0 |
| 3 | Cerebrovascular diseases (I60-I69) | 5,232 | 7.4 | 472.4 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 2,995 | 4.2 | 270.4 |
| 5 | Diabetes mellitus (E10-E14) | 2,737 | 3.9 | 247.1 |
| 6 | Influenza and pneumonia (J10-J18) | 1,975 | 2.8 | 178.3 |
| 7 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1,888 | 2.7 | 170.5 |
| 8 | Septicemia (A40-A41) | 1,590 | 2.2 | 143.6 |
| 9 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,205 | 1.7 | 108.8 |
| 10 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 931 | 1.3 | 84.1 |
| 11 | Alzheimer's disease (G30) | 755 | 1.1 | 68.2 |
| 12 | Pneumonitis due to solids and liquids (J69) | 612 | 0.9 | 55.3 |
| 13 | Aortic aneurysm and dissection (I71) | 364 | 0.5 | 32.9 |
| 14 | Chronic liver disease and cirrhosis (K70,K73-K74) | 353 | 0.5 | 31.9 |
| 15 | Atherosclerosis (I70) | 333 | 0.5 | 30.1 |
| ... | All other causes (Residual) | 8,324 | 11.7 | 751.6 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|--|---|---------|-------------------------|-------|
| Black, non-Hispanic, female, all ages | | | | |
| ... | All causes | 139,379 | 100.0 | 791.4 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 40,424 | 29.0 | 229.5 |
| 2 | Malignant neoplasms (C00-C97) | 28,897 | 20.7 | 164.1 |
| 3 | Cerebrovascular diseases (I60-I69) | 11,108 | 8.0 | 63.1 |
| 4 | Diabetes mellitus (E10-E14) | 7,198 | 5.2 | 40.9 |
| 5 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 3,808 | 2.7 | 21.6 |
| 6 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 3,707 | 2.7 | 21.0 |
| 7 | Chronic lower respiratory diseases (J40-J47) | 3,344 | 2.4 | 19.0 |
| 8 | Septicemia (A40-A41) | 3,304 | 2.4 | 18.8 |
| 9 | Influenza and pneumonia (J10-J18) | 3,040 | 2.2 | 17.3 |
| 10 | Human immunodeficiency virus (HIV) disease (B20-B24) | 2,417 | 1.7 | 13.7 |
| 11 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 2,286 | 1.6 | 13.0 |
| 12 | Certain conditions originating in the perinatal period (P00-P96) | 2,118 | 1.5 | 12.0 |
| 13 | Alzheimer's disease (G30) | 1,945 | 1.4 | 11.0 |
| 14 | Assault (homicide) (X85-Y09,Y87.1) | 1,366 | 1.0 | 7.8 |
| 15 | Chronic liver disease and cirrhosis (K70,K73-K74) | 996 | 0.7 | 5.7 |
| ... | All other causes (Residual) | 23,421 | 16.8 | 133.0 |
| Black, non-Hispanic, female, 1-4 years | | | | |
| ... | All causes | 545 | 100.0 | 53.5 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 164 | 30.1 | 16.1 |
| 2 | Assault (homicide) (X85-Y09,Y87.1) | 70 | 12.8 | 6.9 |
| 3 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 56 | 10.3 | 5.5 |
| 4 | Malignant neoplasms (C00-C97) | 35 | 6.4 | 3.4 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 23 | 4.2 | 2.3 |
| 6 | Septicemia (A40-A41) | 12 | 2.2 | * |
| 7 | Anemias (D50-D64) | 11 | 2.0 | * |
| 8 | Cerebrovascular diseases (I60-I69) | 10 | 1.8 | * |
| 9 | Influenza and pneumonia (J10-J18) | 9 | 1.7 | * |
| 9 | Certain conditions originating in the perinatal period (P00-P96) | 9 | 1.7 | * |
| 11 | Meningitis (G00,G03) | 7 | 1.3 | * |
| 11 | Chronic lower respiratory diseases (J40-J47) | 7 | 1.3 | * |
| 13 | Human immunodeficiency virus (HIV) disease (B20-B24) | 6 | 1.1 | * |
| 13 | Complications of medical and surgical care (Y40-Y84,Y88) | 6 | 1.1 | * |
| 15 | Acute bronchitis and bronchiolitis (J20-J21) | 3 | 0.6 | * |
| 15 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 3 | 0.6 | * |
| ... | All other causes (Residual) | 114 | 20.9 | 11.2 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|-------------------|---|--------|-------------------------|------|
| | Black, non-Hispanic, female, 5-14 years | | | |
| ... | All causes | 661 | 100.0 | 22.8 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 213 | 32.2 | 7.3 |
| 2 | Malignant neoplasms (C00-C97) | 61 | 9.2 | 2.1 |
| 3 | Assault (homicide) (X85-Y09,Y87.1) | 52 | 7.9 | 1.8 |
| 4 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 39 | 5.9 | 1.3 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 34 | 5.1 | 1.2 |
| 6 | Chronic lower respiratory diseases (J40-J47) | 27 | 4.1 | 0.9 |
| 7 | Anemias (D50-D64) | 13 | 2.0 | * |
| 8 | Human immunodeficiency virus (HIV) disease (B20-B24) | 11 | 1.7 | * |
| 9 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 10 | 1.5 | * |
| 9 | Cerebrovascular diseases (I60-I69) | 10 | 1.5 | * |
| 11 | Certain conditions originating in the perinatal period (P00-P96) | 7 | 1.1 | * |
| 12 | Meningococcal infection (A39) | 5 | 0.8 | * |
| 12 | Septicemia (A40-A41) | 5 | 0.8 | * |
| 12 | Diabetes mellitus (E10-E14) | 5 | 0.8 | * |
| 12 | Intentional self-harm (suicide) (X60-X84,Y87.0) | 5 | 0.8 | * |
| ... | All other causes (Residual) | 164 | 24.8 | 5.6 |
| | Black, non-Hispanic, female, 15-24 years | | | |
| ... | All causes | 1,702 | 100.0 | 61.8 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 429 | 25.2 | 15.6 |
| 2 | Assault (homicide) (X85-Y09,Y87.1) | 312 | 18.3 | 11.3 |
| 3 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 139 | 8.2 | 5.0 |
| 4 | Malignant neoplasms (C00-C97) | 119 | 7.0 | 4.3 |
| 5 | Human immunodeficiency virus (HIV) disease (B20-B24) | 65 | 3.8 | 2.4 |
| 6 | Intentional self-harm (suicide) (X60-X84,Y87.0) | 64 | 3.8 | 2.3 |
| 7 | Pregnancy, childbirth and the puerperium (O00-099) | 34 | 2.0 | 1.2 |
| 8 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 33 | 1.9 | 1.2 |
| 9 | Anemias (D50-D64) | 32 | 1.9 | 1.2 |
| 10 | Chronic lower respiratory diseases (J40-J47) | 29 | 1.7 | 1.1 |
| 11 | Diabetes mellitus (E10-E14) | 27 | 1.6 | 1.0 |
| 12 | Cerebrovascular diseases (I60-I69) | 23 | 1.4 | 0.8 |
| 12 | Influenza and pneumonia (J10-J18) | 23 | 1.4 | 0.8 |
| 14 | Septicemia (A40-A41) | 15 | 0.9 | * |
| 15 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 14 | 0.8 | * |
| ... | All other causes (Residual) | 344 | 20.2 | 12.5 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|-------------------|---|--------|-------------------------|-------|
| | Black, non-Hispanic, female, 25-44 years | | | |
| ... | All causes | 11,672 | 100.0 | 212.8 |
| 1 | Malignant neoplasms (C00-C97) | 2,162 | 18.5 | 39.4 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 1,673 | 14.3 | 30.5 |
| 3 | Human immunodeficiency virus (HIV) disease (B20-B24) | 1,560 | 13.4 | 28.4 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,028 | 8.8 | 18.7 |
| 5 | Assault (homicide) (X85-Y09,Y87.1) | 642 | 5.5 | 11.7 |
| 6 | Cerebrovascular diseases (I60-I69) | 523 | 4.5 | 9.5 |
| 7 | Diabetes mellitus (E10-E14) | 295 | 2.5 | 5.4 |
| 8 | Chronic liver disease and cirrhosis (K70,K73-K74) | 197 | 1.7 | 3.6 |
| 9 | Septicemia (A40-A41) | 192 | 1.6 | 3.5 |
| 10 | Chronic lower respiratory diseases (J40-J47) | 187 | 1.6 | 3.4 |
| 11 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 179 | 1.5 | 3.3 |
| 12 | Anemias (D50-D64) | 153 | 1.3 | 2.8 |
| 12 | Intentional self-harm (suicide) (X60-X84,Y87.0) | 153 | 1.3 | 2.8 |
| 14 | Influenza and pneumonia (J10-J18) | 137 | 1.2 | 2.5 |
| 15 | Pregnancy, childbirth and the puerperium (O00-099) | 102 | 0.9 | 1.9 |
| ... | All other causes (Residual) | 2,489 | 21.3 | 45.4 |
| | Black, non-Hispanic, female, 45-64 years | | | |
| ... | All causes | 29,729 | 100.0 | 856.4 |
| 1 | Malignant neoplasms (C00-C97) | 9,172 | 30.9 | 264.2 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 7,290 | 24.5 | 210.0 |
| 3 | Cerebrovascular diseases (I60-I69) | 1,889 | 6.4 | 54.4 |
| 4 | Diabetes mellitus (E10-E14) | 1,665 | 5.6 | 48.0 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 780 | 2.6 | 22.5 |
| 6 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 766 | 2.6 | 22.1 |
| 7 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 754 | 2.5 | 21.7 |
| 8 | Human immunodeficiency virus (HIV) disease (B20-B24) | 702 | 2.4 | 20.2 |
| 9 | Septicemia (A40-A41) | 628 | 2.1 | 18.1 |
| 10 | Chronic liver disease and cirrhosis (K70,K73-K74) | 510 | 1.7 | 14.7 |
| 11 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 430 | 1.4 | 12.4 |
| 12 | Influenza and pneumonia (J10-J18) | 361 | 1.2 | 10.4 |
| 13 | Assault (homicide) (X85-Y09,Y87.1) | 159 | 0.5 | 4.6 |
| 14 | Viral hepatitis (B15-B19) | 147 | 0.5 | 4.2 |
| 15 | Aortic aneurysm and dissection (I71) | 119 | 0.4 | 3.4 |
| ... | All other causes (Residual) | 4,357 | 14.7 | 125.5 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2000

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Population denominators for 2000 death rates are based on the 1990 census population.]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age | Number | Percent of total deaths | Rate |
|-------------------|---|--------|-------------------------|---------|
| | Black, non-Hispanic, female, 65 years and over | | | |
| ... | All causes | 91,288 | 100.0 | 5,343.5 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 31,201 | 34.2 | 1,826.3 |
| 2 | Malignant neoplasms (C00-C97) | 17,340 | 19.0 | 1,015.0 |
| 3 | Cerebrovascular diseases (I60-I69) | 8,633 | 9.5 | 505.3 |
| 4 | Diabetes mellitus (E10-E14) | 5,206 | 5.7 | 304.7 |
| 5 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 2,815 | 3.1 | 164.8 |
| 6 | Influenza and pneumonia (J10-J18) | 2,460 | 2.7 | 144.0 |
| 7 | Septicemia (A40-A41) | 2,404 | 2.6 | 140.7 |
| 8 | Chronic lower respiratory diseases (J40-J47) | 2,309 | 2.5 | 135.2 |
| 9 | Alzheimer's disease (G30) | 1,928 | 2.1 | 112.9 |
| 10 | Essential (primary) hypertension and hypertensive renal disease (I10,I12) | 1,749 | 1.9 | 102.4 |
| 11 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 996 | 1.1 | 58.3 |
| 12 | Pneumonitis due to solids and liquids (J69) | 650 | 0.7 | 38.0 |
| 13 | Atherosclerosis (I70) | 575 | 0.6 | 33.7 |
| 14 | Aortic aneurysm and dissection (I71) | 444 | 0.5 | 26.0 |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 416 | 0.5 | 24.4 |
| ... | All other causes (Residual) | 12,162 | 13.3 | 711.9 |
| 11 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,167 | 1.3 | 69.0 |
| 12 | Pneumonitis due to solids and liquids (J69) | 632 | 0.7 | 37.3 |
| 13 | Atherosclerosis (I70) | 582 | 0.6 | 34.4 |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 439 | 0.5 | 25.9 |
| 15 | Aortic aneurysm and dissection (I71) | 427 | 0.5 | 25.2 |
| ... | All other causes (Residual) | 11,751 | 13.0 | 694.4 |

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable