

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, all ages				
...	All causes	113,413	100.0	306.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	27,090	23.9	73.3
2	Malignant neoplasms (C00-C97)	22,371	19.7	60.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,523	8.4	25.8
4	Cerebrovascular diseases (I60-I69)	6,416	5.7	17.4
5	Diabetes mellitus (E10-E14)	5,663	5.0	15.3
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,331	2.9	9.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	3,301	2.9	8.9
8	Chronic lower respiratory diseases (J40-J47)	2,832	2.5	7.7
9	Influenza and pneumonia (J10-J18)	2,722	2.4	7.4
10	Certain conditions originating in the perinatal period (P00-P96)	2,227	2.0	6.0
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,857	1.6	5.0
12	Human immunodeficiency virus (HIV) disease (B20-B24)	1,853	1.6	5.0
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,850	1.6	5.0
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,632	1.4	4.4
15	Septicemia (A40-A41)	1,386	1.2	3.7
...	All other causes (Residual)	19,359	17.1	52.4
Hispanic, both sexes, 1-4 years				
...	All causes	917	100.0	30.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	310	33.8	10.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	103	11.2	3.4
3	Malignant neoplasms (C00-C97)	84	9.2	2.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	69	7.5	2.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	33	3.6	1.1
6	Influenza and pneumonia (J10-J18)	32	3.5	1.1
7	Septicemia (A40-A41)	18	2.0	*
8	Certain conditions originating in the perinatal period (P00-P96)	14	1.5	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	1.4	*
10	Cerebrovascular diseases (I60-I69)	10	1.1	*
11	Chronic lower respiratory diseases (J40-J47)	8	0.9	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.7	*
13	Meningococcal infection (A39)	5	0.5	*
14	Meningitis (G00,G03)	4	0.4	*
14	Diseases of appendix (K35-K38)	4	0.4	*
...	All other causes (Residual)	204	22.2	6.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 5-14 years				
...	All causes	1,038	100.0	14.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	379	36.5	5.4
2	Malignant neoplasms (C00-C97)	177	17.1	2.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	58	5.6	0.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	56	5.4	0.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	38	3.7	0.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	37	3.6	0.5
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	21	2.0	0.3
8	Septicemia (A40-A41)	18	1.7	*
9	Cerebrovascular diseases (I60-I69)	16	1.5	*
10	Influenza and pneumonia (J10-J18)	10	1.0	*
11	Chronic lower respiratory diseases (J40-J47)	9	0.9	*
12	Anemias (D50-D64)	5	0.5	*
12	Pneumonitis due to solids and liquids (J69)	5	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.4	*
...	All other causes (Residual)	201	19.4	2.8
Hispanic, both sexes, 15-24 years				
...	All causes	5,058	100.0	75.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,182	43.1	32.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,222	24.2	18.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	414	8.2	6.2
4	Malignant neoplasms (C00-C97)	289	5.7	4.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	112	2.2	1.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	73	1.4	1.1
7	Cerebrovascular diseases (I60-I69)	40	0.8	0.6
8	Human immunodeficiency virus (HIV) disease (B20-B24)	32	0.6	0.5
9	Pregnancy, childbirth and the puerperium (O00-099)	26	0.5	0.4
10	Legal intervention (Y35,Y89.0)	23	0.5	0.3
11	Diabetes mellitus (E10-E14)	19	0.4	*
12	Influenza and pneumonia (J10-J18)	18	0.4	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	0.3	*
14	Septicemia (A40-A41)	13	0.3	*
15	Anemias (D50-D64)	11	0.2	*
...	All other causes (Residual)	568	11.2	8.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 25-44 years				
...	All causes	14,238	100.0	115.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,698	26.0	30.0
2	Malignant neoplasms (C00-C97)	1,893	13.3	15.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,527	10.7	12.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,064	7.5	8.6
5	Human immunodeficiency virus (HIV) disease (B20-B24)	1,059	7.4	8.6
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	887	6.2	7.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	611	4.3	5.0
8	Cerebrovascular diseases (I60-I69)	374	2.6	3.0
9	Diabetes mellitus (E10-E14)	216	1.5	1.8
10	Viral hepatitis (B15-B19)	136	1.0	1.1
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	99	0.7	0.8
12	Influenza and pneumonia (J10-J18)	96	0.7	0.8
13	Septicemia (A40-A41)	95	0.7	0.8
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	87	0.6	0.7
15	Chronic lower respiratory diseases (J40-J47)	74	0.5	0.6
...	All other causes (Residual)	2,322	16.3	18.9
Hispanic, both sexes, 45-64 years				
...	All causes	24,699	100.0	470.9
1	Malignant neoplasms (C00-C97)	6,941	28.1	132.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,027	20.4	95.9
3	Chronic liver disease and cirrhosis (K70,K73-K74)	1,672	6.8	31.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,642	6.6	31.3
5	Diabetes mellitus (E10-E14)	1,489	6.0	28.4
6	Cerebrovascular diseases (I60-I69)	1,193	4.8	22.7
7	Human immunodeficiency virus (HIV) disease (B20-B24)	695	2.8	13.3
8	Viral hepatitis (B15-B19)	458	1.9	8.7
9	Chronic lower respiratory diseases (J40-J47)	382	1.5	7.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	369	1.5	7.0
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	353	1.4	6.7
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	330	1.3	6.3
13	Influenza and pneumonia (J10-J18)	289	1.2	5.5
14	Septicemia (A40-A41)	284	1.1	5.4
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	143	0.6	2.7
...	All other causes (Residual)	3,432	13.9	65.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 65 years and over				
...	All causes	62,716	100.0	3,389.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,731	33.1	1,120.4
2	Malignant neoplasms (C00-C97)	12,973	20.7	701.1
3	Cerebrovascular diseases (I60-I69)	4,766	7.6	257.6
4	Diabetes mellitus (E10-E14)	3,935	6.3	212.7
5	Chronic lower respiratory diseases (J40-J47)	2,342	3.7	126.6
6	Influenza and pneumonia (J10-J18)	2,219	3.5	119.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,349	2.2	72.9
8	Alzheimer's disease (G30)	1,216	1.9	65.7
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,129	1.8	61.0
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,012	1.6	54.7
11	Septicemia (A40-A41)	908	1.4	49.1
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	714	1.1	38.6
13	Pneumonitis due to solids and liquids (J69)	479	0.8	25.9
14	Parkinson's disease (G20-G21)	466	0.7	25.2
15	Atherosclerosis (I70)	383	0.6	20.7
...	All other causes (Residual)	8,094	12.9	437.4
Hispanic, male, all ages				
...	All causes	63,317	100.0	332.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,195	22.4	74.6
2	Malignant neoplasms (C00-C97)	11,825	18.7	62.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,157	11.3	37.6
4	Cerebrovascular diseases (I60-I69)	2,982	4.7	15.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,756	4.4	14.5
6	Diabetes mellitus (E10-E14)	2,590	4.1	13.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,410	3.8	12.7
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,576	2.5	8.3
9	Chronic lower respiratory diseases (J40-J47)	1,482	2.3	7.8
10	Human immunodeficiency virus (HIV) disease (B20-B24)	1,437	2.3	7.6
11	Influenza and pneumonia (J10-J18)	1,309	2.1	6.9
12	Certain conditions originating in the perinatal period (P00-P96)	1,268	2.0	6.7
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	892	1.4	4.7
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	892	1.4	4.7
15	Septicemia (A40-A41)	684	1.1	3.6
...	All other causes (Residual)	9,862	15.6	51.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 1-4 years				
...	All causes	518	100.0	33.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	195	37.6	12.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	54	10.4	3.5
3	Malignant neoplasms (C00-C97)	44	8.5	2.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	38	7.3	2.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	16	3.1	*
5	Influenza and pneumonia (J10-J18)	16	3.1	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	1.4	*
8	Certain conditions originating in the perinatal period (P00-P96)	6	1.2	*
9	Septicemia (A40-A41)	5	1.0	*
10	Cerebrovascular diseases (I60-I69)	4	0.8	*
10	Chronic lower respiratory diseases (J40-J47)	4	0.8	*
12	Meningococcal infection (A39)	3	0.6	*
12	Meningitis (G00,G03)	3	0.6	*
12	Diseases of appendix (K35-K38)	3	0.6	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.6	*
...	All other causes (Residual)	117	22.6	7.6
Hispanic, male, 5-14 years				
...	All causes	600	100.0	16.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	224	37.3	6.2
2	Malignant neoplasms (C00-C97)	90	15.0	2.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	38	6.3	1.1
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	36	6.0	1.0
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	26	4.3	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	19	3.2	*
7	Septicemia (A40-A41)	14	2.3	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	2.0	*
9	Cerebrovascular diseases (I60-I69)	10	1.7	*
10	Influenza and pneumonia (J10-J18)	6	1.0	*
10	Chronic lower respiratory diseases (J40-J47)	6	1.0	*
12	Anemias (D50-D64)	5	0.8	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.5	*
14	Meningococcal infection (A39)	2	0.3	*
14	Viral hepatitis (B15-B19)	2	0.3	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.3	*
14	Diabetes mellitus (E10-E14)	2	0.3	*
14	Meningitis (G00,G03)	2	0.3	*
14	Pneumonitis due to solids and liquids (J69)	2	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.3	*
14	Certain conditions originating in the perinatal period (P00-P96)	2	0.3	*
...	All other causes (Residual)	95	15.8	2.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 15-24 years				
...	All causes	4,019	100.0	111.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,773	44.1	49.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,099	27.3	30.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	343	8.5	9.5
4	Malignant neoplasms (C00-C97)	199	5.0	5.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	76	1.9	2.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	39	1.0	1.1
7	Cerebrovascular diseases (I60-I69)	26	0.6	0.7
8	Legal intervention (Y35,Y89.0)	23	0.6	0.6
9	Human immunodeficiency virus (HIV) disease (B20-B24)	21	0.5	0.6
10	Influenza and pneumonia (J10-J18)	15	0.4	*
11	Diabetes mellitus (E10-E14)	11	0.3	*
12	Septicemia (A40-A41)	9	0.2	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.2	*
14	Tuberculosis (A16-A19)	6	0.1	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.1	*
14	Anemias (D50-D64)	6	0.1	*
14	Chronic lower respiratory diseases (J40-J47)	6	0.1	*
...	All other causes (Residual)	352	8.8	9.8
Hispanic, male, 25-44 years				
...	All causes	10,261	100.0	157.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,037	29.6	46.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,251	12.2	19.2
3	Malignant neoplasms (C00-C97)	848	8.3	13.0
4	Human immunodeficiency virus (HIV) disease (B20-B24)	811	7.9	12.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	791	7.7	12.1
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	769	7.5	11.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	492	4.8	7.5
8	Cerebrovascular diseases (I60-I69)	211	2.1	3.2
9	Diabetes mellitus (E10-E14)	137	1.3	2.1
10	Viral hepatitis (B15-B19)	92	0.9	1.4
11	Influenza and pneumonia (J10-J18)	62	0.6	0.9
12	Septicemia (A40-A41)	58	0.6	0.9
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	56	0.5	0.9
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	53	0.5	0.8
15	Legal intervention (Y35,Y89.0)	37	0.4	0.6
...	All other causes (Residual)	1,556	15.2	23.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 45-64 years				
...	All causes	15,503	100.0	607.2
1	Malignant neoplasms (C00-C97)	3,574	23.1	140.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,501	22.6	137.1
3	Chronic liver disease and cirrhosis (K70,K73-K74)	1,337	8.6	52.4
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,213	7.8	47.5
5	Diabetes mellitus (E10-E14)	821	5.3	32.2
6	Cerebrovascular diseases (I60-I69)	693	4.5	27.1
7	Human immunodeficiency virus (HIV) disease (B20-B24)	549	3.5	21.5
8	Viral hepatitis (B15-B19)	329	2.1	12.9
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	292	1.9	11.4
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	249	1.6	9.8
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	196	1.3	7.7
12	Chronic lower respiratory diseases (J40-J47)	194	1.3	7.6
13	Influenza and pneumonia (J10-J18)	184	1.2	7.2
14	Septicemia (A40-A41)	166	1.1	6.5
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	73	0.5	2.9
...	All other causes (Residual)	2,132	13.8	83.5
Hispanic, male, 65 years and over				
...	All causes	29,757	100.0	3,830.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,744	32.7	1,254.2
2	Malignant neoplasms (C00-C97)	7,066	23.7	909.5
3	Cerebrovascular diseases (I60-I69)	2,029	6.8	261.2
4	Diabetes mellitus (E10-E14)	1,618	5.4	208.3
5	Chronic lower respiratory diseases (J40-J47)	1,234	4.1	158.8
6	Influenza and pneumonia (J10-J18)	994	3.3	127.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	616	2.1	79.3
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	614	2.1	79.0
9	Chronic liver disease and cirrhosis (K70,K73-K74)	578	1.9	74.4
10	Septicemia (A40-A41)	403	1.4	51.9
11	Alzheimer's disease (G30)	391	1.3	50.3
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	266	0.9	34.2
13	Parkinson's disease (G20-G21)	264	0.9	34.0
14	Pneumonitis due to solids and liquids (J69)	239	0.8	30.8
15	Aortic aneurysm and dissection (I71)	177	0.6	22.8
...	All other causes (Residual)	3,524	11.8	453.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, all ages				
...	All causes	50,096	100.0	279.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,895	25.7	71.8
2	Malignant neoplasms (C00-C97)	10,546	21.1	58.7
3	Cerebrovascular diseases (I60-I69)	3,434	6.9	19.1
4	Diabetes mellitus (E10-E14)	3,073	6.1	17.1
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,366	4.7	13.2
6	Influenza and pneumonia (J10-J18)	1,413	2.8	7.9
7	Chronic lower respiratory diseases (J40-J47)	1,350	2.7	7.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	965	1.9	5.4
9	Certain conditions originating in the perinatal period (P00-P96)	959	1.9	5.3
10	Chronic liver disease and cirrhosis (K70,K73-K74)	891	1.8	5.0
11	Alzheimer's disease (G30)	838	1.7	4.7
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	740	1.5	4.1
13	Septicemia (A40-A41)	702	1.4	3.9
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	575	1.1	3.2
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	524	1.0	2.9
...	All other causes (Residual)	8,825	17.6	49.2
Hispanic, female, 1-4 years				
...	All causes	399	100.0	27.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	115	28.8	7.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	49	12.3	3.3
3	Malignant neoplasms (C00-C97)	40	10.0	2.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	31	7.8	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	17	4.3	*
6	Influenza and pneumonia (J10-J18)	16	4.0	*
7	Septicemia (A40-A41)	13	3.3	*
8	Certain conditions originating in the perinatal period (P00-P96)	8	2.0	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.5	*
9	Cerebrovascular diseases (I60-I69)	6	1.5	*
11	Chronic lower respiratory diseases (J40-J47)	4	1.0	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.8	*
13	Meningococcal infection (A39)	2	0.5	*
13	Pneumonitis due to solids and liquids (J69)	2	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.5	*
...	All other causes (Residual)	85	21.3	5.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 5-14 years				
...	All causes	438	100.0	12.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	155	35.4	4.5
2	Malignant neoplasms (C00-C97)	87	19.9	2.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	22	5.0	0.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	19	4.3	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	18	4.1	*
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11	2.5	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	2.1	*
8	Cerebrovascular diseases (I60-I69)	6	1.4	*
9	Septicemia (A40-A41)	4	0.9	*
9	Influenza and pneumonia (J10-J18)	4	0.9	*
11	Chronic lower respiratory diseases (J40-J47)	3	0.7	*
11	Pneumonitis due to solids and liquids (J69)	3	0.7	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.5	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.2	*
14	Meningitis (G00,G03)	1	0.2	*
14	Pregnancy, childbirth and the puerperium (O00-099)	1	0.2	*
14	Certain conditions originating in the perinatal period (P00-P96)	1	0.2	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
...	All other causes (Residual)	90	20.5	2.6
Hispanic, female, 15-24 years				
...	All causes	1,039	100.0	33.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	409	39.4	13.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	123	11.8	4.0
3	Malignant neoplasms (C00-C97)	90	8.7	2.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	71	6.8	2.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	36	3.5	1.2
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	34	3.3	1.1
7	Pregnancy, childbirth and the puerperium (O00-099)	26	2.5	0.8
8	Cerebrovascular diseases (I60-I69)	14	1.3	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	11	1.1	*
10	Diabetes mellitus (E10-E14)	8	0.8	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.7	*
12	Anemias (D50-D64)	5	0.5	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.5	*
14	Septicemia (A40-A41)	4	0.4	*
14	Viral hepatitis (B15-B19)	4	0.4	*
14	Chronic lower respiratory diseases (J40-J47)	4	0.4	*
...	All other causes (Residual)	188	18.1	6.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 25-44 years				
...	All causes	3,977	100.0	68.7
1	Malignant neoplasms (C00-C97)	1,045	26.3	18.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	661	16.6	11.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	276	6.9	4.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	273	6.9	4.7
5	Human immunodeficiency virus (HIV) disease (B20-B24)	248	6.2	4.3
6	Cerebrovascular diseases (I60-I69)	163	4.1	2.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	119	3.0	2.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	118	3.0	2.0
9	Diabetes mellitus (E10-E14)	79	2.0	1.4
10	Pregnancy, childbirth and the puerperium (O00-099)	55	1.4	1.0
11	Viral hepatitis (B15-B19)	44	1.1	0.8
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	43	1.1	0.7
13	Chronic lower respiratory diseases (J40-J47)	40	1.0	0.7
14	Septicemia (A40-A41)	37	0.9	0.6
15	Influenza and pneumonia (J10-J18)	34	0.9	0.6
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	34	0.9	0.6
...	All other causes (Residual)	708	17.8	12.2
Hispanic, female, 45-64 years				
...	All causes	9,196	100.0	341.7
1	Malignant neoplasms (C00-C97)	3,367	36.6	125.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,526	16.6	56.7
3	Diabetes mellitus (E10-E14)	668	7.3	24.8
4	Cerebrovascular diseases (I60-I69)	500	5.4	18.6
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	429	4.7	15.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	335	3.6	12.4
7	Chronic lower respiratory diseases (J40-J47)	188	2.0	7.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	173	1.9	6.4
9	Human immunodeficiency virus (HIV) disease (B20-B24)	146	1.6	5.4
10	Viral hepatitis (B15-B19)	129	1.4	4.8
11	Septicemia (A40-A41)	118	1.3	4.4
12	Influenza and pneumonia (J10-J18)	105	1.1	3.9
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	81	0.9	3.0
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	70	0.8	2.6
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	61	0.7	2.3
...	All other causes (Residual)	1,300	14.1	48.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 65 years and over				
...	All causes	32,959	100.0	3,070.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,987	33.3	1,023.6
2	Malignant neoplasms (C00-C97)	5,907	17.9	550.3
3	Cerebrovascular diseases (I60-I69)	2,737	8.3	255.0
4	Diabetes mellitus (E10-E14)	2,317	7.0	215.9
5	Influenza and pneumonia (J10-J18)	1,225	3.7	114.1
6	Chronic lower respiratory diseases (J40-J47)	1,108	3.4	103.2
7	Alzheimer's disease (G30)	825	2.5	76.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	733	2.2	68.3
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	515	1.6	48.0
10	Septicemia (A40-A41)	505	1.5	47.0
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	448	1.4	41.7
12	Chronic liver disease and cirrhosis (K70,K73-K74)	434	1.3	40.4
13	Pneumonitis due to solids and liquids (J69)	240	0.7	22.4
14	Atherosclerosis (I70)	231	0.7	21.5
15	Parkinson's disease (G20-G21)	202	0.6	18.8
...	All other causes (Residual)	4,545	13.8	423.4
White, non-Hispanic, both sexes, all ages				
...	All causes	1,962,810	100.0	991.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	582,349	29.7	294.1
2	Malignant neoplasms (C00-C97)	456,709	23.3	230.6
3	Cerebrovascular diseases (I60-I69)	133,879	6.8	67.6
4	Chronic lower respiratory diseases (J40-J47)	110,753	5.6	55.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	76,262	3.9	38.5
6	Influenza and pneumonia (J10-J18)	51,952	2.6	26.2
7	Diabetes mellitus (E10-E14)	51,482	2.6	26.0
8	Alzheimer's disease (G30)	49,030	2.5	24.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	29,449	1.5	14.9
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	25,813	1.3	13.0
11	Septicemia (A40-A41)	24,389	1.2	12.3
12	Chronic liver disease and cirrhosis (K70,K73-K74)	20,065	1.0	10.1
13	Parkinson's disease (G20-G21)	15,239	0.8	7.7
14	Pneumonitis due to solids and liquids (J69)	14,843	0.8	7.5
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	13,837	0.7	7.0
...	All other causes (Residual)	306,759	15.6	154.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 1-4 years				
...	All causes	2,772	100.0	30.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,006	36.3	10.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	287	10.4	3.1
3	Malignant neoplasms (C00-C97)	268	9.7	2.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	160	5.8	1.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	108	3.9	1.2
6	Septicemia (A40-A41)	63	2.3	0.7
7	Influenza and pneumonia (J10-J18)	48	1.7	0.5
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	36	1.3	0.4
9	Certain conditions originating in the perinatal period (P00-P96)	34	1.2	0.4
10	Cerebrovascular diseases (I60-I69)	28	1.0	0.3
11	Meningitis (G00,G03)	15	0.5	*
12	Chronic lower respiratory diseases (J40-J47)	13	0.5	*
13	Acute bronchitis and bronchiolitis (J20-J21)	12	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	0.4	*
15	Meningococcal infection (A39)	10	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	10	0.4	*
...	All other causes (Residual)	663	23.9	7.2
White, non-Hispanic, both sexes, 5-14 years				
...	All causes	4,151	100.0	16.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,688	40.7	6.6
2	Malignant neoplasms (C00-C97)	607	14.6	2.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	226	5.4	0.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	183	4.4	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	164	4.0	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	152	3.7	0.6
7	Influenza and pneumonia (J10-J18)	57	1.4	0.2
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	56	1.3	0.2
9	Cerebrovascular diseases (I60-I69)	42	1.0	0.2
10	Chronic lower respiratory diseases (J40-J47)	39	0.9	0.2
11	Septicemia (A40-A41)	35	0.8	0.1
12	Anemias (D50-D64)	22	0.5	0.1
13	Diabetes mellitus (E10-E14)	19	0.5	*
14	Meningococcal infection (A39)	18	0.4	*
14	Certain conditions originating in the perinatal period (P00-P96)	18	0.4	*
...	All other causes (Residual)	825	19.9	3.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 15-24 years				
...	All causes	18,730	100.0	74.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,867	52.7	39.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,926	15.6	11.6
3	Malignant neoplasms (C00-C97)	1,076	5.7	4.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,058	5.6	4.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	522	2.8	2.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	302	1.6	1.2
7	Influenza and pneumonia (J10-J18)	108	0.6	0.4
8	Cerebrovascular diseases (I60-I69)	98	0.5	0.4
9	Chronic lower respiratory diseases (J40-J47)	75	0.4	0.3
10	Septicemia (A40-A41)	69	0.4	0.3
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	66	0.4	0.3
11	Diabetes mellitus (E10-E14)	66	0.4	0.3
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	45	0.2	0.2
14	Meningococcal infection (A39)	38	0.2	0.2
14	Pregnancy, childbirth and the puerperium (O00-099)	38	0.2	0.2
...	All other causes (Residual)	2,376	12.7	9.4
White, non-Hispanic, both sexes, 25-44 years				
...	All causes	83,737	100.0	147.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19,138	22.9	33.7
2	Malignant neoplasms (C00-C97)	14,145	16.9	24.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,509	12.6	18.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	9,475	11.3	16.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,723	4.4	6.6
6	Chronic liver disease and cirrhosis (K70,K73-K74)	2,439	2.9	4.3
7	Human immunodeficiency virus (HIV) disease (B20-B24)	2,430	2.9	4.3
8	Cerebrovascular diseases (I60-I69)	1,653	2.0	2.9
9	Diabetes mellitus (E10-E14)	1,568	1.9	2.8
10	Influenza and pneumonia (J10-J18)	812	1.0	1.4
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	769	0.9	1.4
12	Chronic lower respiratory diseases (J40-J47)	747	0.9	1.3
13	Septicemia (A40-A41)	603	0.7	1.1
14	Viral hepatitis (B15-B19)	565	0.7	1.0
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	459	0.5	0.8
...	All other causes (Residual)	14,702	17.6	25.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 45-64 years				
...	All causes	301,195	100.0	608.6
1	Malignant neoplasms (C00-C97)	108,225	35.9	218.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	71,977	23.9	145.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15,480	5.1	31.3
4	Chronic lower respiratory diseases (J40-J47)	12,248	4.1	24.7
5	Diabetes mellitus (E10-E14)	9,450	3.1	19.1
6	Cerebrovascular diseases (I60-I69)	9,408	3.1	19.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	9,224	3.1	18.6
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8,271	2.7	16.7
9	Septicemia (A40-A41)	3,262	1.1	6.6
10	Influenza and pneumonia (J10-J18)	3,156	1.0	6.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,765	0.9	5.6
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,025	0.7	4.1
13	Viral hepatitis (B15-B19)	1,950	0.6	3.9
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1,671	0.6	3.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,567	0.5	3.2
...	All other causes (Residual)	40,516	13.5	81.9
White, non-Hispanic, both sexes, 65 years and over				
...	All causes	1,538,801	100.0	5,217.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	498,821	32.4	1,691.4
2	Malignant neoplasms (C00-C97)	332,342	21.6	1,126.9
3	Cerebrovascular diseases (I60-I69)	122,603	8.0	415.7
4	Chronic lower respiratory diseases (J40-J47)	97,615	6.3	331.0
5	Alzheimer's disease (G30)	48,482	3.2	164.4
6	Influenza and pneumonia (J10-J18)	47,653	3.1	161.6
7	Diabetes mellitus (E10-E14)	40,373	2.6	136.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	28,558	1.9	96.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	26,105	1.7	88.5
10	Septicemia (A40-A41)	20,235	1.3	68.6
11	Parkinson's disease (G20-G21)	14,934	1.0	50.6
12	Pneumonitis due to solids and liquids (J69)	13,810	0.9	46.8
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	12,488	0.8	42.3
14	Atherosclerosis (I70)	11,980	0.8	40.6
15	Aortic aneurysm and dissection (I71)	11,352	0.7	38.5
...	All other causes (Residual)	211,450	13.7	717.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, all ages				
...	All causes	945,967	100.0	975.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	280,713	29.7	289.5
2	Malignant neoplasms (C00-C97)	236,004	24.9	243.4
3	Chronic lower respiratory diseases (J40-J47)	52,952	5.6	54.6
4	Cerebrovascular diseases (I60-I69)	50,382	5.3	52.0
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	48,180	5.1	49.7
6	Diabetes mellitus (E10-E14)	24,302	2.6	25.1
7	Influenza and pneumonia (J10-J18)	22,389	2.4	23.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	20,710	2.2	21.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14,331	1.5	14.8
10	Alzheimer's disease (G30)	14,295	1.5	14.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	12,609	1.3	13.0
12	Septicemia (A40-A41)	10,668	1.1	11.0
13	Parkinson's disease (G20-G21)	8,685	0.9	9.0
14	Aortic aneurysm and dissection (I71)	7,924	0.8	8.2
15	Pneumonitis due to solids and liquids (J69)	7,411	0.8	7.6
...	All other causes (Residual)	134,412	14.2	138.6
White, non-Hispanic, male, 1-4 years				
...	All causes	1,592	100.0	33.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	624	39.2	13.2
2	Malignant neoplasms (C00-C97)	144	9.0	3.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	144	9.0	3.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	95	6.0	2.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	59	3.7	1.3
6	Septicemia (A40-A41)	41	2.6	0.9
7	Influenza and pneumonia (J10-J18)	26	1.6	0.6
8	Certain conditions originating in the perinatal period (P00-P96)	22	1.4	0.5
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	1.1	*
10	Cerebrovascular diseases (I60-I69)	14	0.9	*
11	Meningitis (G00,G03)	10	0.6	*
12	Chronic lower respiratory diseases (J40-J47)	9	0.6	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.4	*
14	Meningococcal infection (A39)	5	0.3	*
14	Acute bronchitis and bronchiolitis (J20-J21)	5	0.3	*
...	All other causes (Residual)	369	23.2	7.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 5-14 years				
...	All causes	2,428	100.0	18.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,066	43.9	8.1
2	Malignant neoplasms (C00-C97)	310	12.8	2.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	144	5.9	1.1
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	135	5.6	1.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	83	3.4	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	80	3.3	0.6
7	Influenza and pneumonia (J10-J18)	28	1.2	0.2
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	25	1.0	0.2
9	Cerebrovascular diseases (I60-I69)	24	1.0	0.2
10	Chronic lower respiratory diseases (J40-J47)	22	0.9	0.2
11	Septicemia (A40-A41)	18	0.7	*
12	Certain conditions originating in the perinatal period (P00-P96)	14	0.6	*
13	Diabetes mellitus (E10-E14)	13	0.5	*
14	Anemias (D50-D64)	11	0.5	*
15	Meningococcal infection (A39)	10	0.4	*
...	All other causes (Residual)	445	18.3	3.4
White, non-Hispanic, male, 15-24 years				
...	All causes	13,542	100.0	105.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,297	53.9	56.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,522	18.6	19.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	729	5.4	5.7
4	Malignant neoplasms (C00-C97)	632	4.7	4.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	336	2.5	2.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	183	1.4	1.4
7	Influenza and pneumonia (J10-J18)	64	0.5	0.5
8	Cerebrovascular diseases (I60-I69)	55	0.4	0.4
9	Diabetes mellitus (E10-E14)	42	0.3	0.3
10	Chronic lower respiratory diseases (J40-J47)	39	0.3	0.3
11	Septicemia (A40-A41)	37	0.3	0.3
12	Legal intervention (Y35,Y89.0)	35	0.3	0.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	31	0.2	0.2
14	Aortic aneurysm and dissection (I71)	21	0.2	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21	0.2	0.2
...	All other causes (Residual)	1,498	11.1	11.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 25-44 years				
...	All causes	54,826	100.0	192.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14,094	25.7	49.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,579	13.8	26.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7,519	13.7	26.4
4	Malignant neoplasms (C00-C97)	6,420	11.7	22.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,675	4.9	9.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	2,049	3.7	7.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,548	2.8	5.4
8	Diabetes mellitus (E10-E14)	951	1.7	3.3
9	Cerebrovascular diseases (I60-I69)	845	1.5	3.0
10	Influenza and pneumonia (J10-J18)	445	0.8	1.6
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	422	0.8	1.5
12	Viral hepatitis (B15-B19)	396	0.7	1.4
13	Chronic lower respiratory diseases (J40-J47)	355	0.6	1.2
14	Septicemia (A40-A41)	335	0.6	1.2
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	256	0.5	0.9
...	All other causes (Residual)	8,937	16.3	31.4
White, non-Hispanic, male, 45-64 years				
...	All causes	183,724	100.0	754.6
1	Malignant neoplasms (C00-C97)	57,563	31.3	236.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	51,682	28.1	212.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,939	6.0	44.9
4	Chronic liver disease and cirrhosis (K70,K73-K74)	6,520	3.5	26.8
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,300	3.4	25.9
6	Chronic lower respiratory diseases (J40-J47)	6,216	3.4	25.5
7	Diabetes mellitus (E10-E14)	5,538	3.0	22.7
8	Cerebrovascular diseases (I60-I69)	5,145	2.8	21.1
9	Influenza and pneumonia (J10-J18)	1,839	1.0	7.6
10	Septicemia (A40-A41)	1,686	0.9	6.9
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,555	0.8	6.4
12	Human immunodeficiency virus (HIV) disease (B20-B24)	1,476	0.8	6.1
13	Viral hepatitis (B15-B19)	1,446	0.8	5.9
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,444	0.8	5.9
15	Aortic aneurysm and dissection (I71)	1,183	0.6	4.9
...	All other causes (Residual)	23,192	12.6	95.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 65 years and over				
...	All causes	682,255	100.0	5,567.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	220,830	32.4	1,802.2
2	Malignant neoplasms (C00-C97)	170,910	25.1	1,394.8
3	Chronic lower respiratory diseases (J40-J47)	46,301	6.8	377.9
4	Cerebrovascular diseases (I60-I69)	44,271	6.5	361.3
5	Influenza and pneumonia (J10-J18)	19,915	2.9	162.5
6	Diabetes mellitus (E10-E14)	17,756	2.6	144.9
7	Alzheimer's disease (G30)	14,055	2.1	114.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,857	2.0	113.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12,461	1.8	101.7
10	Septicemia (A40-A41)	8,480	1.2	69.2
11	Parkinson's disease (G20-G21)	8,460	1.2	69.0
12	Pneumonitis due to solids and liquids (J69)	6,786	1.0	55.4
13	Aortic aneurysm and dissection (I71)	6,495	1.0	53.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,657	0.7	38.0
15	Chronic liver disease and cirrhosis (K70,K73-K74)	4,533	0.7	37.0
...	All other causes (Residual)	82,488	12.1	673.2
White, non-Hispanic, female, all ages				
...	All causes	1,016,843	100.0	1,006.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	301,636	29.7	298.4
2	Malignant neoplasms (C00-C97)	220,705	21.7	218.4
3	Cerebrovascular diseases (I60-I69)	83,497	8.2	82.6
4	Chronic lower respiratory diseases (J40-J47)	57,801	5.7	57.2
5	Alzheimer's disease (G30)	34,735	3.4	34.4
6	Influenza and pneumonia (J10-J18)	29,563	2.9	29.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	28,082	2.8	27.8
8	Diabetes mellitus (E10-E14)	27,180	2.7	26.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15,118	1.5	15.0
10	Septicemia (A40-A41)	13,721	1.3	13.6
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	8,867	0.9	8.8
12	Atherosclerosis (I70)	8,006	0.8	7.9
13	Chronic liver disease and cirrhosis (K70,K73-K74)	7,456	0.7	7.4
14	Pneumonitis due to solids and liquids (J69)	7,432	0.7	7.4
15	Parkinson's disease (G20-G21)	6,554	0.6	6.5
...	All other causes (Residual)	166,490	16.4	164.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 1-4 years				
...	All causes	1,180	100.0	26.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	382	32.4	8.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	143	12.1	3.2
3	Malignant neoplasms (C00-C97)	124	10.5	2.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	65	5.5	1.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	49	4.2	1.1
6	Septicemia (A40-A41)	22	1.9	0.5
6	Influenza and pneumonia (J10-J18)	22	1.9	0.5
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	1.5	*
9	Cerebrovascular diseases (I60-I69)	14	1.2	*
10	Certain conditions originating in the perinatal period (P00-P96)	12	1.0	*
11	Acute bronchitis and bronchiolitis (J20-J21)	7	0.6	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.6	*
13	Pneumonitis due to solids and liquids (J69)	6	0.5	*
14	Meningococcal infection (A39)	5	0.4	*
14	Anemias (D50-D64)	5	0.4	*
14	Meningitis (G00,G03)	5	0.4	*
...	All other causes (Residual)	294	24.9	6.6
White, non-Hispanic, female, 5-14 years				
...	All causes	1,723	100.0	13.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	622	36.1	5.0
2	Malignant neoplasms (C00-C97)	297	17.2	2.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	91	5.3	0.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	81	4.7	0.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	72	4.2	0.6
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	39	2.3	0.3
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	31	1.8	0.3
8	Influenza and pneumonia (J10-J18)	29	1.7	0.2
9	Cerebrovascular diseases (I60-I69)	18	1.0	*
10	Septicemia (A40-A41)	17	1.0	*
10	Chronic lower respiratory diseases (J40-J47)	17	1.0	*
12	Anemias (D50-D64)	11	0.6	*
13	Meningococcal infection (A39)	8	0.5	*
14	Meningitis (G00,G03)	7	0.4	*
15	Diabetes mellitus (E10-E14)	6	0.3	*
...	All other causes (Residual)	377	21.9	3.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 15-24 years				
...	All causes	5,188	100.0	41.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,570	49.5	20.7
2	Malignant neoplasms (C00-C97)	444	8.6	3.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	404	7.8	3.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	329	6.3	2.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	186	3.6	1.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	119	2.3	1.0
7	Influenza and pneumonia (J10-J18)	44	0.8	0.4
8	Cerebrovascular diseases (I60-I69)	43	0.8	0.3
9	Pregnancy, childbirth and the puerperium (O00-099)	38	0.7	0.3
10	Chronic lower respiratory diseases (J40-J47)	36	0.7	0.3
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	35	0.7	0.3
12	Septicemia (A40-A41)	32	0.6	0.3
13	Diabetes mellitus (E10-E14)	24	0.5	0.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	24	0.5	0.2
15	Meningococcal infection (A39)	20	0.4	0.2
...	All other causes (Residual)	840	16.2	6.8
White, non-Hispanic, female, 25-44 years				
...	All causes	28,911	100.0	102.3
1	Malignant neoplasms (C00-C97)	7,725	26.7	27.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,044	17.4	17.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,930	10.1	10.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,956	6.8	6.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,048	3.6	3.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	891	3.1	3.2
7	Cerebrovascular diseases (I60-I69)	808	2.8	2.9
8	Diabetes mellitus (E10-E14)	617	2.1	2.2
9	Chronic lower respiratory diseases (J40-J47)	392	1.4	1.4
10	Human immunodeficiency virus (HIV) disease (B20-B24)	381	1.3	1.3
11	Influenza and pneumonia (J10-J18)	367	1.3	1.3
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	347	1.2	1.2
13	Septicemia (A40-A41)	268	0.9	0.9
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	203	0.7	0.7
15	Viral hepatitis (B15-B19)	169	0.6	0.6
...	All other causes (Residual)	5,765	19.9	20.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 45-64 years				
...	All causes	117,471	100.0	467.2
1	Malignant neoplasms (C00-C97)	50,662	43.1	201.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,295	17.3	80.7
3	Chronic lower respiratory diseases (J40-J47)	6,032	5.1	24.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,541	3.9	18.1
5	Cerebrovascular diseases (I60-I69)	4,263	3.6	17.0
6	Diabetes mellitus (E10-E14)	3,912	3.3	15.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,704	2.3	10.8
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,971	1.7	7.8
9	Septicemia (A40-A41)	1,576	1.3	6.3
10	Influenza and pneumonia (J10-J18)	1,317	1.1	5.2
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,210	1.0	4.8
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	708	0.6	2.8
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	581	0.5	2.3
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	520	0.4	2.1
15	Viral hepatitis (B15-B19)	504	0.4	2.0
...	All other causes (Residual)	16,675	14.2	66.3
White, non-Hispanic, female, 65 years and over				
...	All causes	856,546	100.0	4,969.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	277,991	32.5	1,612.7
2	Malignant neoplasms (C00-C97)	161,432	18.8	936.5
3	Cerebrovascular diseases (I60-I69)	78,332	9.1	454.4
4	Chronic lower respiratory diseases (J40-J47)	51,314	6.0	297.7
5	Alzheimer's disease (G30)	34,427	4.0	199.7
6	Influenza and pneumonia (J10-J18)	27,738	3.2	160.9
7	Diabetes mellitus (E10-E14)	22,617	2.6	131.2
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14,701	1.7	85.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13,644	1.6	79.2
10	Septicemia (A40-A41)	11,755	1.4	68.2
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	8,314	1.0	48.2
12	Atherosclerosis (I70)	7,806	0.9	45.3
13	Pneumonitis due to solids and liquids (J69)	7,024	0.8	40.7
14	Parkinson's disease (G20-G21)	6,474	0.8	37.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,929	0.6	28.6
...	All other causes (Residual)	128,048	14.9	742.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, all ages				
...	All causes	284,343	100.0	798.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	76,794	27.0	215.5
2	Malignant neoplasms (C00-C97)	61,580	21.7	172.8
3	Cerebrovascular diseases (I60-I69)	18,844	6.6	52.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,239	4.3	34.4
5	Diabetes mellitus (E10-E14)	12,167	4.3	34.1
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8,077	2.8	22.7
7	Human immunodeficiency virus (HIV) disease (B20-B24)	7,710	2.7	21.6
8	Chronic lower respiratory diseases (J40-J47)	7,504	2.6	21.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,214	2.5	20.2
10	Septicemia (A40-A41)	5,829	2.0	16.4
11	Influenza and pneumonia (J10-J18)	5,686	2.0	16.0
12	Certain conditions originating in the perinatal period (P00-P96)	4,713	1.7	13.2
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	4,012	1.4	11.3
14	Alzheimer's disease (G30)	3,104	1.1	8.7
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,733	1.0	7.7
...	All other causes (Residual)	46,137	16.2	129.5
Black, non-Hispanic, both sexes, 1-4 years				
...	All causes	1,156	100.0	49.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	331	28.6	14.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	166	14.4	7.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	127	11.0	5.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	73	6.3	3.1
5	Malignant neoplasms (C00-C97)	46	4.0	2.0
6	Influenza and pneumonia (J10-J18)	24	2.1	1.0
7	Chronic lower respiratory diseases (J40-J47)	22	1.9	0.9
7	Certain conditions originating in the perinatal period (P00-P96)	22	1.9	0.9
9	Septicemia (A40-A41)	21	1.8	0.9
10	Cerebrovascular diseases (I60-I69)	12	1.0	*
11	Anemias (D50-D64)	11	1.0	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	10	0.9	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.6	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.4	*
14	Meningitis (G00,G03)	5	0.4	*
...	All other causes (Residual)	274	23.7	11.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 5-14 years				
...	All causes	1,557	100.0	24.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	630	40.5	9.7
2	Malignant neoplasms (C00-C97)	172	11.0	2.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	92	5.9	1.4
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	73	4.7	1.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	70	4.5	1.1
6	Chronic lower respiratory diseases (J40-J47)	53	3.4	0.8
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	40	2.6	0.6
8	Human immunodeficiency virus (HIV) disease (B20-B24)	26	1.7	0.4
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	26	1.7	0.4
8	Anemias (D50-D64)	26	1.7	0.4
11	Influenza and pneumonia (J10-J18)	17	1.1	*
12	Diabetes mellitus (E10-E14)	15	1.0	*
13	Cerebrovascular diseases (I60-I69)	14	0.9	*
14	Certain conditions originating in the perinatal period (P00-P96)	11	0.7	*
15	Septicemia (A40-A41)	10	0.6	*
...	All other causes (Residual)	282	18.1	4.4
Black, non-Hispanic, both sexes, 15-24 years				
...	All causes	7,024	100.0	121.4
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,817	40.1	48.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,713	24.4	29.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	425	6.1	7.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	322	4.6	5.6
5	Malignant neoplasms (C00-C97)	257	3.7	4.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	161	2.3	2.8
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	113	1.6	2.0
8	Chronic lower respiratory diseases (J40-J47)	77	1.1	1.3
9	Anemias (D50-D64)	72	1.0	1.2
10	Diabetes mellitus (E10-E14)	56	0.8	1.0
11	Cerebrovascular diseases (I60-I69)	51	0.7	0.9
12	Influenza and pneumonia (J10-J18)	45	0.6	0.8
12	Pregnancy, childbirth and the puerperium (O00-099)	45	0.6	0.8
14	Septicemia (A40-A41)	41	0.6	0.7
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	27	0.4	0.5
...	All other causes (Residual)	802	11.4	13.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 25-44 years				
...	All causes	30,295	100.0	282.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,385	14.5	40.8
2	Human immunodeficiency virus (HIV) disease (B20-B24)	4,306	14.2	40.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,882	12.8	36.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,762	12.4	35.0
5	Malignant neoplasms (C00-C97)	3,641	12.0	33.9
6	Cerebrovascular diseases (I60-I69)	911	3.0	8.5
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	890	2.9	8.3
8	Diabetes mellitus (E10-E14)	704	2.3	6.6
9	Chronic liver disease and cirrhosis (K70,K73-K74)	465	1.5	4.3
10	Chronic lower respiratory diseases (J40-J47)	401	1.3	3.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	399	1.3	3.7
12	Septicemia (A40-A41)	366	1.2	3.4
13	Influenza and pneumonia (J10-J18)	361	1.2	3.4
14	Anemias (D50-D64)	271	0.9	2.5
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	232	0.8	2.2
...	All other causes (Residual)	5,319	17.6	49.5
Black, non-Hispanic, both sexes, 45-64 years				
...	All causes	73,621	100.0	1,082.8
1	Malignant neoplasms (C00-C97)	20,371	27.7	299.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	19,165	26.0	281.9
3	Cerebrovascular diseases (I60-I69)	4,175	5.7	61.4
4	Diabetes mellitus (E10-E14)	3,439	4.7	50.6
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,148	4.3	46.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	2,939	4.0	43.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,750	2.4	25.7
8	Chronic liver disease and cirrhosis (K70,K73-K74)	1,642	2.2	24.1
9	Chronic lower respiratory diseases (J40-J47)	1,568	2.1	23.1
10	Septicemia (A40-A41)	1,399	1.9	20.6
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	992	1.3	14.6
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	923	1.3	13.6
13	Influenza and pneumonia (J10-J18)	901	1.2	13.3
14	Viral hepatitis (B15-B19)	516	0.7	7.6
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	394	0.5	5.8
...	All other causes (Residual)	10,299	14.0	151.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 65 years and over				
...	All causes	162,363	100.0	5,643.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	52,625	32.4	1,829.0
2	Malignant neoplasms (C00-C97)	37,085	22.8	1,288.9
3	Cerebrovascular diseases (I60-I69)	13,636	8.4	473.9
4	Diabetes mellitus (E10-E14)	7,952	4.9	276.4
5	Chronic lower respiratory diseases (J40-J47)	5,367	3.3	186.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,976	3.1	172.9
7	Influenza and pneumonia (J10-J18)	4,235	2.6	147.2
8	Septicemia (A40-A41)	3,868	2.4	134.4
9	Alzheimer's disease (G30)	3,073	1.9	106.8
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,779	1.7	96.6
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,255	1.4	78.4
12	Pneumonitis due to solids and liquids (J69)	1,348	0.8	46.9
13	Atherosclerosis (I70)	834	0.5	29.0
14	Aortic aneurysm and dissection (I71)	771	0.5	26.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	765	0.5	26.6
...	All other causes (Residual)	20,794	12.8	722.7
Black, non-Hispanic, male, all ages				
...	All causes	143,971	100.0	849.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	36,541	25.4	215.7
2	Malignant neoplasms (C00-C97)	32,335	22.5	190.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,381	5.8	49.5
4	Cerebrovascular diseases (I60-I69)	7,828	5.4	46.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6,658	4.6	39.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	5,227	3.6	30.8
7	Diabetes mellitus (E10-E14)	4,983	3.5	29.4
8	Chronic lower respiratory diseases (J40-J47)	4,143	2.9	24.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,158	2.2	18.6
10	Influenza and pneumonia (J10-J18)	2,769	1.9	16.3
11	Certain conditions originating in the perinatal period (P00-P96)	2,674	1.9	15.8
12	Septicemia (A40-A41)	2,611	1.8	15.4
13	Chronic liver disease and cirrhosis (K70,K73-K74)	1,819	1.3	10.7
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,698	1.2	10.0
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,578	1.1	9.3
...	All other causes (Residual)	21,568	15.0	127.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 1-4 years				
...	All causes	655	100.0	55.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	188	28.7	15.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	100	15.3	8.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	65	9.9	5.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	40	6.1	3.4
5	Malignant neoplasms (C00-C97)	29	4.4	2.4
6	Chronic lower respiratory diseases (J40-J47)	16	2.4	*
7	Certain conditions originating in the perinatal period (P00-P96)	14	2.1	*
8	Influenza and pneumonia (J10-J18)	13	2.0	*
9	Septicemia (A40-A41)	12	1.8	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	8	1.2	*
11	Anemias (D50-D64)	7	1.1	*
12	Cerebrovascular diseases (I60-I69)	5	0.8	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.6	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.5	*
...	All other causes (Residual)	148	22.6	12.5
Black, non-Hispanic, male, 5-14 years				
...	All causes	941	100.0	28.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	432	45.9	13.1
2	Malignant neoplasms (C00-C97)	100	10.6	3.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	66	7.0	2.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	34	3.6	1.0
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	33	3.5	1.0
6	Chronic lower respiratory diseases (J40-J47)	32	3.4	1.0
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	31	3.3	0.9
8	Human immunodeficiency virus (HIV) disease (B20-B24)	15	1.6	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	1.4	*
10	Anemias (D50-D64)	10	1.1	*
11	Influenza and pneumonia (J10-J18)	9	1.0	*
12	Diabetes mellitus (E10-E14)	6	0.6	*
13	Cerebrovascular diseases (I60-I69)	5	0.5	*
13	Certain conditions originating in the perinatal period (P00-P96)	5	0.5	*
15	Meningococcal infection (A39)	3	0.3	*
15	Septicemia (A40-A41)	3	0.3	*
15	Meningitis (G00,G03)	3	0.3	*
...	All other causes (Residual)	141	15.0	4.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 15-24 years				
...	All causes	5,377	100.0	186.2
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,551	47.4	88.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,285	23.9	44.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	388	7.2	13.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	205	3.8	7.1
5	Malignant neoplasms (C00-C97)	148	2.8	5.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	72	1.3	2.5
7	Human immunodeficiency virus (HIV) disease (B20-B24)	68	1.3	2.4
8	Chronic lower respiratory diseases (J40-J47)	48	0.9	1.7
9	Anemias (D50-D64)	41	0.8	1.4
10	Diabetes mellitus (E10-E14)	27	0.5	0.9
11	Septicemia (A40-A41)	25	0.5	0.9
12	Legal intervention (Y35,Y89.0)	24	0.4	0.8
13	Influenza and pneumonia (J10-J18)	23	0.4	0.8
14	Cerebrovascular diseases (I60-I69)	20	0.4	0.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	0.2	*
...	All other causes (Residual)	439	8.2	15.2
Black, non-Hispanic, male, 25-44 years				
...	All causes	18,519	100.0	365.4
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,034	16.4	59.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,800	15.1	55.3
3	Human immunodeficiency virus (HIV) disease (B20-B24)	2,793	15.1	55.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,649	14.3	52.3
5	Malignant neoplasms (C00-C97)	1,468	7.9	29.0
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	735	4.0	14.5
7	Cerebrovascular diseases (I60-I69)	430	2.3	8.5
8	Diabetes mellitus (E10-E14)	397	2.1	7.8
9	Chronic liver disease and cirrhosis (K70,K73-K74)	298	1.6	5.9
10	Influenza and pneumonia (J10-J18)	207	1.1	4.1
11	Chronic lower respiratory diseases (J40-J47)	193	1.0	3.8
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	193	1.0	3.8
13	Septicemia (A40-A41)	174	0.9	3.4
14	Anemias (D50-D64)	139	0.8	2.7
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	131	0.7	2.6
...	All other causes (Residual)	2,878	15.5	56.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 45-64 years				
...	All causes	42,907	100.0	1,383.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,668	27.2	376.3
2	Malignant neoplasms (C00-C97)	10,948	25.5	353.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,324	5.4	74.9
4	Cerebrovascular diseases (I60-I69)	2,250	5.2	72.6
5	Human immunodeficiency virus (HIV) disease (B20-B24)	2,153	5.0	69.4
6	Diabetes mellitus (E10-E14)	1,732	4.0	55.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,164	2.7	37.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	908	2.1	29.3
9	Chronic lower respiratory diseases (J40-J47)	825	1.9	26.6
10	Septicemia (A40-A41)	715	1.7	23.1
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	709	1.7	22.9
12	Influenza and pneumonia (J10-J18)	563	1.3	18.2
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	547	1.3	17.6
14	Viral hepatitis (B15-B19)	339	0.8	10.9
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	298	0.7	9.6
...	All other causes (Residual)	5,764	13.4	185.9
Black, non-Hispanic, male, 65 years and over				
...	All causes	70,881	100.0	6,478.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	21,870	30.9	1,998.9
2	Malignant neoplasms (C00-C97)	19,638	27.7	1,794.9
3	Cerebrovascular diseases (I60-I69)	5,089	7.2	465.1
4	Chronic lower respiratory diseases (J40-J47)	3,023	4.3	276.3
5	Diabetes mellitus (E10-E14)	2,820	4.0	257.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,014	2.8	184.1
7	Influenza and pneumonia (J10-J18)	1,902	2.7	173.8
8	Septicemia (A40-A41)	1,615	2.3	147.6
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,188	1.7	108.6
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,013	1.4	92.6
11	Alzheimer's disease (G30)	886	1.2	81.0
12	Pneumonitis due to solids and liquids (J69)	603	0.9	55.1
13	Chronic liver disease and cirrhosis (K70,K73-K74)	354	0.5	32.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	336	0.5	30.7
15	Aortic aneurysm and dissection (I71)	335	0.5	30.6
...	All other causes (Residual)	8,195	11.6	749.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, all ages				
...	All causes	140,372	100.0	751.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	40,253	28.7	215.4
2	Malignant neoplasms (C00-C97)	29,245	20.8	156.5
3	Cerebrovascular diseases (I60-I69)	11,016	7.8	59.0
4	Diabetes mellitus (E10-E14)	7,184	5.1	38.4
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,056	2.9	21.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,858	2.7	20.6
7	Chronic lower respiratory diseases (J40-J47)	3,361	2.4	18.0
8	Septicemia (A40-A41)	3,218	2.3	17.2
9	Influenza and pneumonia (J10-J18)	2,917	2.1	15.6
10	Human immunodeficiency virus (HIV) disease (B20-B24)	2,483	1.8	13.3
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,314	1.6	12.4
12	Alzheimer's disease (G30)	2,198	1.6	11.8
13	Certain conditions originating in the perinatal period (P00-P96)	2,039	1.5	10.9
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,419	1.0	7.6
15	Chronic liver disease and cirrhosis (K70,K73-K74)	914	0.7	4.9
...	All other causes (Residual)	23,897	17.0	127.9
Black, non-Hispanic, female, 1-4 years				
...	All causes	501	100.0	43.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	143	28.5	12.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	66	13.2	5.7
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	62	12.4	5.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	33	6.6	2.9
5	Malignant neoplasms (C00-C97)	17	3.4	*
6	Influenza and pneumonia (J10-J18)	11	2.2	*
7	Septicemia (A40-A41)	9	1.8	*
8	Certain conditions originating in the perinatal period (P00-P96)	8	1.6	*
9	Cerebrovascular diseases (I60-I69)	7	1.4	*
10	Chronic lower respiratory diseases (J40-J47)	6	1.2	*
11	Anemias (D50-D64)	4	0.8	*
12	Meningococcal infection (A39)	3	0.6	*
12	Meningitis (G00,G03)	3	0.6	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.6	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.4	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.4	*
...	All other causes (Residual)	122	24.4	10.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 5-14 years				
...	All causes	616	100.0	19.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	198	32.1	6.2
2	Malignant neoplasms (C00-C97)	72	11.7	2.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	40	6.5	1.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	36	5.8	1.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	26	4.2	0.8
6	Chronic lower respiratory diseases (J40-J47)	21	3.4	0.7
7	Anemias (D50-D64)	16	2.6	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	2.1	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	11	1.8	*
10	Diabetes mellitus (E10-E14)	9	1.5	*
10	Cerebrovascular diseases (I60-I69)	9	1.5	*
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	9	1.5	*
13	Influenza and pneumonia (J10-J18)	8	1.3	*
14	Septicemia (A40-A41)	7	1.1	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	1.1	*
...	All other causes (Residual)	134	21.8	4.2
Black, non-Hispanic, female, 15-24 years				
...	All causes	1,647	100.0	56.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	428	26.0	14.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	266	16.2	9.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	117	7.1	4.0
4	Malignant neoplasms (C00-C97)	109	6.6	3.8
5	Human immunodeficiency virus (HIV) disease (B20-B24)	93	5.6	3.2
6	Pregnancy, childbirth and the puerperium (O00-099)	45	2.7	1.6
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	41	2.5	1.4
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	37	2.2	1.3
9	Anemias (D50-D64)	31	1.9	1.1
9	Cerebrovascular diseases (I60-I69)	31	1.9	1.1
11	Diabetes mellitus (E10-E14)	29	1.8	1.0
11	Chronic lower respiratory diseases (J40-J47)	29	1.8	1.0
13	Influenza and pneumonia (J10-J18)	22	1.3	0.8
14	Septicemia (A40-A41)	16	1.0	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	0.9	*
...	All other causes (Residual)	338	20.5	11.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Black, non-Hispanic, female, 25-44 years			
...	All causes	11,776	100.0	207.8
1	Malignant neoplasms (C00-C97)	2,173	18.5	38.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,736	14.7	30.6
3	Human immunodeficiency virus (HIV) disease (B20-B24)	1,513	12.8	26.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,082	9.2	19.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	728	6.2	12.8
6	Cerebrovascular diseases (I60-I69)	481	4.1	8.5
7	Diabetes mellitus (E10-E14)	307	2.6	5.4
8	Chronic lower respiratory diseases (J40-J47)	208	1.8	3.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	206	1.7	3.6
10	Septicemia (A40-A41)	192	1.6	3.4
11	Chronic liver disease and cirrhosis (K70,K73-K74)	167	1.4	2.9
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	155	1.3	2.7
13	Influenza and pneumonia (J10-J18)	154	1.3	2.7
14	Anemias (D50-D64)	132	1.1	2.3
15	Pregnancy, childbirth and the puerperium (O00-099)	105	0.9	1.9
...	All other causes (Residual)	2,437	20.7	43.0
	Black, non-Hispanic, female, 45-64 years			
...	All causes	30,714	100.0	830.5
1	Malignant neoplasms (C00-C97)	9,423	30.7	254.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,497	24.4	202.7
3	Cerebrovascular diseases (I60-I69)	1,925	6.3	52.1
4	Diabetes mellitus (E10-E14)	1,707	5.6	46.2
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	842	2.7	22.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	824	2.7	22.3
7	Human immunodeficiency virus (HIV) disease (B20-B24)	786	2.6	21.3
8	Chronic lower respiratory diseases (J40-J47)	743	2.4	20.1
9	Septicemia (A40-A41)	684	2.2	18.5
10	Chronic liver disease and cirrhosis (K70,K73-K74)	478	1.6	12.9
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	445	1.4	12.0
12	Influenza and pneumonia (J10-J18)	338	1.1	9.1
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	214	0.7	5.8
14	Viral hepatitis (B15-B19)	177	0.6	4.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	125	0.4	3.4
...	All other causes (Residual)	4,506	14.7	121.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2001

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Black, non-Hispanic, female, 65 years and over			
...	All causes	91,482	100.0	5,130.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	30,755	33.6	1,724.7
2	Malignant neoplasms (C00-C97)	17,447	19.1	978.4
3	Cerebrovascular diseases (I60-I69)	8,547	9.3	479.3
4	Diabetes mellitus (E10-E14)	5,132	5.6	287.8
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,962	3.2	166.1
6	Chronic lower respiratory diseases (J40-J47)	2,344	2.6	131.5
7	Influenza and pneumonia (J10-J18)	2,333	2.6	130.8
8	Septicemia (A40-A41)	2,253	2.5	126.3
9	Alzheimer's disease (G30)	2,187	2.4	122.6
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,766	1.9	99.0
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,067	1.2	59.8
12	Pneumonitis due to solids and liquids (J69)	745	0.8	41.8
13	Atherosclerosis (I70)	538	0.6	30.2
14	Aortic aneurysm and dissection (I71)	436	0.5	24.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	429	0.5	24.1
...	All other causes (Residual)	12,541	13.7	703.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable