

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, all ages				
...	All causes	117,135	100.0	302.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	27,887	23.8	71.9
2	Malignant neoplasms (C00-C97)	23,141	19.8	59.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,106	8.6	26.1
4	Cerebrovascular diseases (I60-I69)	6,451	5.5	16.6
5	Diabetes mellitus (E10-E14)	5,912	5.0	15.3
6	Chronic liver disease and cirrhosis (K70,K73-K74)	3,409	2.9	8.8
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,129	2.7	8.1
8	Chronic lower respiratory diseases (J40-J47)	3,058	2.6	7.9
9	Influenza and pneumonia (J10-J18)	2,824	2.4	7.3
10	Certain conditions originating in the perinatal period (P00-P96)	2,402	2.1	6.2
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,954	1.7	5.0
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,898	1.6	4.9
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,855	1.6	4.8
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,715	1.5	4.4
15	Septicemia (A40-A41)	1,521	1.3	3.9
...	All other causes (Residual)	19,873	17.0	51.3
Hispanic, both sexes, 1-4 years				
...	All causes	933	100.0	29.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	341	36.5	10.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	111	11.9	3.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	81	8.7	2.6
4	Malignant neoplasms (C00-C97)	72	7.7	2.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	24	2.6	0.8
6	Influenza and pneumonia (J10-J18)	17	1.8	*
7	Septicemia (A40-A41)	16	1.7	*
8	Certain conditions originating in the perinatal period (P00-P96)	14	1.5	*
9	Cerebrovascular diseases (I60-I69)	12	1.3	*
10	Chronic lower respiratory diseases (J40-J47)	9	1.0	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	7	0.8	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.8	*
13	Meningitis (G00,G03)	5	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.5	*
15	Anemias (D50-D64)	3	0.3	*
...	All other causes (Residual)	209	22.4	6.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 5-14 years				
...	All causes	1,124	100.0	15.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	389	34.6	5.4
2	Malignant neoplasms (C00-C97)	210	18.7	2.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	70	6.2	1.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	64	5.7	0.9
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	37	3.3	0.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	30	2.7	0.4
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	26	2.3	0.4
8	Cerebrovascular diseases (I60-I69)	20	1.8	0.3
9	Septicemia (A40-A41)	14	1.2	*
9	Influenza and pneumonia (J10-J18)	14	1.2	*
11	Chronic lower respiratory diseases (J40-J47)	13	1.2	*
12	Certain conditions originating in the perinatal period (P00-P96)	6	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.4	*
15	Anemias (D50-D64)	4	0.4	*
...	All other causes (Residual)	217	19.3	3.0
Hispanic, both sexes, 15-24 years				
...	All causes	5,255	100.0	77.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,413	45.9	35.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,200	22.8	17.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	455	8.7	6.7
4	Malignant neoplasms (C00-C97)	304	5.8	4.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	128	2.4	1.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	67	1.3	1.0
7	Human immunodeficiency virus (HIV) disease (B20-B24)	29	0.6	0.4
7	Cerebrovascular diseases (I60-I69)	29	0.6	0.4
9	Legal intervention (Y35,Y89.0)	22	0.4	0.3
10	Influenza and pneumonia (J10-J18)	19	0.4	*
10	Pregnancy, childbirth and the puerperium (O00-099)	19	0.4	*
12	Septicemia (A40-A41)	15	0.3	*
12	Diabetes mellitus (E10-E14)	15	0.3	*
12	Chronic lower respiratory diseases (J40-J47)	15	0.3	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	9	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.2	*
...	All other causes (Residual)	507	9.6	7.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 25-44 years				
...	All causes	14,283	100.0	108.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,868	27.1	29.4
2	Malignant neoplasms (C00-C97)	1,848	12.9	14.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,367	9.6	10.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,173	8.2	8.9
5	Human immunodeficiency virus (HIV) disease (B20-B24)	1,041	7.3	7.9
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	891	6.2	6.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	588	4.1	4.5
8	Cerebrovascular diseases (I60-I69)	349	2.4	2.7
9	Diabetes mellitus (E10-E14)	261	1.8	2.0
10	Influenza and pneumonia (J10-J18)	120	0.8	0.9
11	Viral hepatitis (B15-B19)	118	0.8	0.9
12	Septicemia (A40-A41)	105	0.7	0.8
13	Chronic lower respiratory diseases (J40-J47)	101	0.7	0.8
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	98	0.7	0.7
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	93	0.7	0.7
...	All other causes (Residual)	2,262	15.8	17.2
Hispanic, both sexes, 45-64 years				
...	All causes	25,960	100.0	461.1
1	Malignant neoplasms (C00-C97)	7,106	27.4	126.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,512	21.2	97.9
3	Chronic liver disease and cirrhosis (K70,K73-K74)	1,741	6.7	30.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,734	6.7	30.8
5	Diabetes mellitus (E10-E14)	1,542	5.9	27.4
6	Cerebrovascular diseases (I60-I69)	1,268	4.9	22.5
7	Human immunodeficiency virus (HIV) disease (B20-B24)	725	2.8	12.9
8	Viral hepatitis (B15-B19)	455	1.8	8.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	424	1.6	7.5
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	402	1.5	7.1
11	Chronic lower respiratory diseases (J40-J47)	401	1.5	7.1
12	Influenza and pneumonia (J10-J18)	331	1.3	5.9
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	303	1.2	5.4
14	Septicemia (A40-A41)	301	1.2	5.3
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	162	0.6	2.9
...	All other causes (Residual)	3,553	13.7	63.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 65 years and over				
...	All causes	64,579	100.0	3,293.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,934	32.4	1,067.7
2	Malignant neoplasms (C00-C97)	13,583	21.0	692.8
3	Cerebrovascular diseases (I60-I69)	4,753	7.4	242.4
4	Diabetes mellitus (E10-E14)	4,093	6.3	208.8
5	Chronic lower respiratory diseases (J40-J47)	2,515	3.9	128.3
6	Influenza and pneumonia (J10-J18)	2,275	3.5	116.0
7	Alzheimer's disease (G30)	1,453	2.2	74.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,322	2.0	67.4
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,206	1.9	61.5
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,066	1.7	54.4
11	Septicemia (A40-A41)	1,005	1.6	51.3
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	734	1.1	37.4
13	Pneumonitis due to solids and liquids (J69)	473	0.7	24.1
14	Parkinson's disease (G20-G21)	463	0.7	23.6
15	Atherosclerosis (I70)	382	0.6	19.5
...	All other causes (Residual)	8,322	12.9	424.4
Hispanic, male, all ages				
...	All causes	65,703	100.0	328.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,798	22.5	74.0
2	Malignant neoplasms (C00-C97)	12,235	18.6	61.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,698	11.7	38.5
4	Cerebrovascular diseases (I60-I69)	3,003	4.6	15.0
5	Diabetes mellitus (E10-E14)	2,779	4.2	13.9
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,635	4.0	13.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,437	3.7	12.2
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,651	2.5	8.3
9	Chronic lower respiratory diseases (J40-J47)	1,625	2.5	8.1
10	Human immunodeficiency virus (HIV) disease (B20-B24)	1,440	2.2	7.2
11	Influenza and pneumonia (J10-J18)	1,398	2.1	7.0
12	Certain conditions originating in the perinatal period (P00-P96)	1,352	2.1	6.8
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	937	1.4	4.7
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	903	1.4	4.5
15	Septicemia (A40-A41)	744	1.1	3.7
...	All other causes (Residual)	10,068	15.3	50.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 1-4 years				
...	All causes	546	100.0	34.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	215	39.4	13.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	60	11.0	3.8
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	49	9.0	3.1
4	Malignant neoplasms (C00-C97)	35	6.4	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	11	2.0	*
5	Influenza and pneumonia (J10-J18)	11	2.0	*
7	Septicemia (A40-A41)	9	1.6	*
7	Certain conditions originating in the perinatal period (P00-P96)	9	1.6	*
9	Cerebrovascular diseases (I60-I69)	5	0.9	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	4	0.7	*
10	Meningitis (G00,G03)	4	0.7	*
10	Chronic lower respiratory diseases (J40-J47)	4	0.7	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.7	*
14	Anemias (D50-D64)	1	0.2	*
14	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
...	All other causes (Residual)	123	22.5	7.7
Hispanic, male, 5-14 years				
...	All causes	647	100.0	17.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	244	37.7	6.6
2	Malignant neoplasms (C00-C97)	108	16.7	2.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	37	5.7	1.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	35	5.4	0.9
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	21	3.2	0.6
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	15	2.3	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	14	2.2	*
8	Septicemia (A40-A41)	8	1.2	*
8	Chronic lower respiratory diseases (J40-J47)	8	1.2	*
10	Cerebrovascular diseases (I60-I69)	7	1.1	*
11	Influenza and pneumonia (J10-J18)	6	0.9	*
12	Certain conditions originating in the perinatal period (P00-P96)	5	0.8	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.6	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	3	0.5	*
14	Anemias (D50-D64)	3	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.5	*
...	All other causes (Residual)	126	19.5	3.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 15-24 years				
...	All causes	4,182	100.0	114.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,958	46.8	53.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,081	25.8	29.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	388	9.3	10.6
4	Malignant neoplasms (C00-C97)	191	4.6	5.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	94	2.2	2.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	39	0.9	1.1
7	Legal intervention (Y35,Y89.0)	22	0.5	0.6
8	Human immunodeficiency virus (HIV) disease (B20-B24)	19	0.5	*
9	Cerebrovascular diseases (I60-I69)	17	0.4	*
10	Chronic lower respiratory diseases (J40-J47)	12	0.3	*
11	Influenza and pneumonia (J10-J18)	11	0.3	*
12	Diabetes mellitus (E10-E14)	10	0.2	*
13	Septicemia (A40-A41)	8	0.2	*
14	Viral hepatitis (B15-B19)	5	0.1	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	5	0.1	*
14	Anemias (D50-D64)	5	0.1	*
14	Meningitis (G00,G03)	5	0.1	*
14	Aortic aneurysm and dissection (I71)	5	0.1	*
14	Pneumonitis due to solids and liquids (J69)	5	0.1	*
14	Chronic liver disease and cirrhosis (K70,K73-K74)	5	0.1	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.1	*
...	All other causes (Residual)	292	7.0	8.0
Hispanic, male, 25-44 years				
...	All causes	10,302	100.0	147.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,177	30.8	45.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,156	11.2	16.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	867	8.4	12.4
4	Malignant neoplasms (C00-C97)	809	7.9	11.5
5	Human immunodeficiency virus (HIV) disease (B20-B24)	808	7.8	11.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	766	7.4	10.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	472	4.6	6.7
8	Cerebrovascular diseases (I60-I69)	215	2.1	3.1
9	Diabetes mellitus (E10-E14)	158	1.5	2.3
10	Viral hepatitis (B15-B19)	89	0.9	1.3
11	Influenza and pneumonia (J10-J18)	78	0.8	1.1
12	Septicemia (A40-A41)	72	0.7	1.0
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	55	0.5	0.8
14	Chronic lower respiratory diseases (J40-J47)	50	0.5	0.7
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	50	0.5	0.7
...	All other causes (Residual)	1,480	14.4	21.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 45-64 years				
...	All causes	16,481	100.0	597.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,860	23.4	140.0
2	Malignant neoplasms (C00-C97)	3,807	23.1	138.0
3	Chronic liver disease and cirrhosis (K70,K73-K74)	1,366	8.3	49.5
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,295	7.9	47.0
5	Diabetes mellitus (E10-E14)	856	5.2	31.0
6	Cerebrovascular diseases (I60-I69)	761	4.6	27.6
7	Human immunodeficiency virus (HIV) disease (B20-B24)	561	3.4	20.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	329	2.0	11.9
9	Viral hepatitis (B15-B19)	326	2.0	11.8
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	238	1.4	8.6
11	Chronic lower respiratory diseases (J40-J47)	234	1.4	8.5
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	229	1.4	8.3
13	Influenza and pneumonia (J10-J18)	216	1.3	7.8
14	Septicemia (A40-A41)	178	1.1	6.5
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	88	0.5	3.2
...	All other causes (Residual)	2,137	13.0	77.5
Hispanic, male, 65 years and over				
...	All causes	30,744	100.0	3,719.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,910	32.2	1,198.8
2	Malignant neoplasms (C00-C97)	7,275	23.7	880.0
3	Cerebrovascular diseases (I60-I69)	1,989	6.5	240.6
4	Diabetes mellitus (E10-E14)	1,754	5.7	212.2
5	Chronic lower respiratory diseases (J40-J47)	1,315	4.3	159.1
6	Influenza and pneumonia (J10-J18)	1,043	3.4	126.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	699	2.3	84.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	628	2.0	76.0
9	Chronic liver disease and cirrhosis (K70,K73-K74)	591	1.9	71.5
10	Alzheimer's disease (G30)	455	1.5	55.0
11	Septicemia (A40-A41)	435	1.4	52.6
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	272	0.9	32.9
13	Parkinson's disease (G20-G21)	261	0.8	31.6
14	Pneumonitis due to solids and liquids (J69)	237	0.8	28.7
15	Aortic aneurysm and dissection (I71)	167	0.5	20.2
...	All other causes (Residual)	3,713	12.1	449.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, all ages				
...	All causes	51,432	100.0	274.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,089	25.4	69.7
2	Malignant neoplasms (C00-C97)	10,906	21.2	58.1
3	Cerebrovascular diseases (I60-I69)	3,448	6.7	18.4
4	Diabetes mellitus (E10-E14)	3,133	6.1	16.7
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,408	4.7	12.8
6	Chronic lower respiratory diseases (J40-J47)	1,433	2.8	7.6
7	Influenza and pneumonia (J10-J18)	1,426	2.8	7.6
8	Certain conditions originating in the perinatal period (P00-P96)	1,050	2.0	5.6
9	Alzheimer's disease (G30)	1,010	2.0	5.4
10	Chronic liver disease and cirrhosis (K70,K73-K74)	972	1.9	5.2
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	961	1.9	5.1
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	812	1.6	4.3
13	Septicemia (A40-A41)	777	1.5	4.1
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	557	1.1	3.0
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	494	1.0	2.6
...	All other causes (Residual)	8,956	17.4	47.7
Hispanic, female, 1-4 years				
...	All causes	387	100.0	25.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	126	32.6	8.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	51	13.2	3.3
3	Malignant neoplasms (C00-C97)	37	9.6	2.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	32	8.3	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	13	3.4	*
6	Septicemia (A40-A41)	7	1.8	*
6	Cerebrovascular diseases (I60-I69)	7	1.8	*
8	Influenza and pneumonia (J10-J18)	6	1.6	*
9	Chronic lower respiratory diseases (J40-J47)	5	1.3	*
9	Certain conditions originating in the perinatal period (P00-P96)	5	1.3	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	1.0	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	3	0.8	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.8	*
14	Anemias (D50-D64)	2	0.5	*
15	Meningitis (G00,G03)	1	0.3	*
15	Acute bronchitis and bronchiolitis (J20-J21)	1	0.3	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.3	*
...	All other causes (Residual)	83	21.4	5.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 5-14 years				
...	All causes	477	100.0	13.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	145	30.4	4.1
2	Malignant neoplasms (C00-C97)	102	21.4	2.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	35	7.3	1.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	27	5.7	0.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	16	3.4	*
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	16	3.4	*
7	Cerebrovascular diseases (I60-I69)	13	2.7	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	11	2.3	*
9	Influenza and pneumonia (J10-J18)	8	1.7	*
10	Septicemia (A40-A41)	6	1.3	*
11	Chronic lower respiratory diseases (J40-J47)	5	1.0	*
12	Pneumonitis due to solids and liquids (J69)	2	0.4	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
12	Infections of kidney (N10-N12,N13.6,N15.1)	2	0.4	*
15	Anemias (D50-D64)	1	0.2	*
15	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.2	*
15	Certain conditions originating in the perinatal period (P00-P96)	1	0.2	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
...	All other causes (Residual)	82	17.2	2.3
Hispanic, female, 15-24 years				
...	All causes	1,073	100.0	34.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	455	42.4	14.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	119	11.1	3.8
3	Malignant neoplasms (C00-C97)	113	10.5	3.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	67	6.2	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	34	3.2	1.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	28	2.6	0.9
7	Pregnancy, childbirth and the puerperium (O00-099)	19	1.8	*
8	Cerebrovascular diseases (I60-I69)	12	1.1	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	10	0.9	*
10	Influenza and pneumonia (J10-J18)	8	0.7	*
11	Septicemia (A40-A41)	7	0.7	*
12	Diabetes mellitus (E10-E14)	5	0.5	*
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	4	0.4	*
13	Chronic liver disease and cirrhosis (K70,K73-K74)	4	0.4	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.4	*
...	All other causes (Residual)	184	17.1	5.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 25-44 years				
...	All causes	3,981	100.0	64.9
1	Malignant neoplasms (C00-C97)	1,039	26.1	16.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	691	17.4	11.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	306	7.7	5.0
4	Human immunodeficiency virus (HIV) disease (B20-B24)	233	5.9	3.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	211	5.3	3.4
6	Cerebrovascular diseases (I60-I69)	134	3.4	2.2
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	125	3.1	2.0
8	Chronic liver disease and cirrhosis (K70,K73-K74)	116	2.9	1.9
9	Diabetes mellitus (E10-E14)	103	2.6	1.7
10	Chronic lower respiratory diseases (J40-J47)	51	1.3	0.8
11	Pregnancy, childbirth and the puerperium (O00-099)	49	1.2	0.8
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	48	1.2	0.8
13	Influenza and pneumonia (J10-J18)	42	1.1	0.7
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	38	1.0	0.6
15	Septicemia (A40-A41)	33	0.8	0.5
...	All other causes (Residual)	762	19.1	12.4
Hispanic, female, 45-64 years				
...	All causes	9,479	100.0	330.1
1	Malignant neoplasms (C00-C97)	3,299	34.8	114.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,652	17.4	57.5
3	Diabetes mellitus (E10-E14)	686	7.2	23.9
4	Cerebrovascular diseases (I60-I69)	507	5.3	17.7
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	439	4.6	15.3
6	Chronic liver disease and cirrhosis (K70,K73-K74)	375	4.0	13.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	195	2.1	6.8
8	Chronic lower respiratory diseases (J40-J47)	167	1.8	5.8
9	Human immunodeficiency virus (HIV) disease (B20-B24)	164	1.7	5.7
10	Viral hepatitis (B15-B19)	129	1.4	4.5
11	Septicemia (A40-A41)	123	1.3	4.3
12	Influenza and pneumonia (J10-J18)	115	1.2	4.0
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	74	0.8	2.6
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	73	0.8	2.5
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	65	0.7	2.3
...	All other causes (Residual)	1,416	14.9	49.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 65 years and over				
...	All causes	33,835	100.0	2,983.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,024	32.6	972.1
2	Malignant neoplasms (C00-C97)	6,308	18.6	556.3
3	Cerebrovascular diseases (I60-I69)	2,764	8.2	243.7
4	Diabetes mellitus (E10-E14)	2,339	6.9	206.3
5	Influenza and pneumonia (J10-J18)	1,232	3.6	108.6
6	Chronic lower respiratory diseases (J40-J47)	1,200	3.5	105.8
7	Alzheimer's disease (G30)	998	2.9	88.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	694	2.1	61.2
9	Septicemia (A40-A41)	570	1.7	50.3
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	507	1.5	44.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	475	1.4	41.9
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	462	1.4	40.7
13	Pneumonitis due to solids and liquids (J69)	236	0.7	20.8
14	Atherosclerosis (I70)	229	0.7	20.2
15	Parkinson's disease (G20-G21)	202	0.6	17.8
...	All other causes (Residual)	4,595	13.6	405.2
White, non-Hispanic, both sexes, all ages				
...	All causes	1,981,973	100.0	997.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	577,761	29.2	290.8
2	Malignant neoplasms (C00-C97)	458,754	23.1	230.9
3	Cerebrovascular diseases (I60-I69)	133,118	6.7	67.0
4	Chronic lower respiratory diseases (J40-J47)	112,128	5.7	56.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	80,605	4.1	40.6
6	Influenza and pneumonia (J10-J18)	55,419	2.8	27.9
7	Alzheimer's disease (G30)	53,486	2.7	26.9
8	Diabetes mellitus (E10-E14)	52,463	2.6	26.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	30,669	1.5	15.4
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	26,691	1.3	13.4
11	Septicemia (A40-A41)	25,545	1.3	12.9
12	Chronic liver disease and cirrhosis (K70,K73-K74)	20,353	1.0	10.2
13	Parkinson's disease (G20-G21)	15,626	0.8	7.9
14	Pneumonitis due to solids and liquids (J69)	15,203	0.8	7.7
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	14,617	0.7	7.4
...	All other causes (Residual)	309,535	15.6	155.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 1-4 years				
...	All causes	2,502	100.0	27.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	865	34.6	9.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	281	11.2	3.0
3	Malignant neoplasms (C00-C97)	236	9.4	2.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	169	6.8	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	91	3.6	1.0
6	Influenza and pneumonia (J10-J18)	46	1.8	0.5
7	Septicemia (A40-A41)	42	1.7	0.5
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	36	1.4	0.4
9	Cerebrovascular diseases (I60-I69)	29	1.2	0.3
10	Chronic lower respiratory diseases (J40-J47)	23	0.9	0.2
11	Certain conditions originating in the perinatal period (P00-P96)	22	0.9	0.2
12	Acute bronchitis and bronchiolitis (J20-J21)	18	0.7	*
13	Anemias (D50-D64)	10	0.4	*
13	Meningitis (G00,G03)	10	0.4	*
15	Meningococcal infection (A39)	8	0.3	*
...	All other causes (Residual)	616	24.6	6.7
White, non-Hispanic, both sexes, 5-14 years				
...	All causes	4,031	100.0	16.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,581	39.2	6.3
2	Malignant neoplasms (C00-C97)	642	15.9	2.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	253	6.3	1.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	177	4.4	0.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	146	3.6	0.6
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	133	3.3	0.5
7	Septicemia (A40-A41)	55	1.4	0.2
7	Influenza and pneumonia (J10-J18)	55	1.4	0.2
9	Cerebrovascular diseases (I60-I69)	44	1.1	0.2
10	Chronic lower respiratory diseases (J40-J47)	42	1.0	0.2
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	41	1.0	0.2
12	Meningitis (G00,G03)	20	0.5	0.1
13	Anemias (D50-D64)	19	0.5	*
13	Diabetes mellitus (E10-E14)	19	0.5	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	16	0.4	*
...	All other causes (Residual)	788	19.5	3.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 15-24 years				
...	All causes	19,458	100.0	75.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,664	54.8	41.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,951	15.2	11.5
3	Malignant neoplasms (C00-C97)	1,097	5.6	4.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	958	4.9	3.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	547	2.8	2.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	315	1.6	1.2
7	Influenza and pneumonia (J10-J18)	103	0.5	0.4
8	Cerebrovascular diseases (I60-I69)	101	0.5	0.4
9	Diabetes mellitus (E10-E14)	95	0.5	0.4
10	Chronic lower respiratory diseases (J40-J47)	94	0.5	0.4
11	Septicemia (A40-A41)	72	0.4	0.3
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	59	0.3	0.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	33	0.2	0.1
14	Legal intervention (Y35,Y89.0)	28	0.1	0.1
15	Pneumonitis due to solids and liquids (J69)	27	0.1	0.1
...	All other causes (Residual)	2,314	11.9	9.0
White, non-Hispanic, both sexes, 25-44 years				
...	All causes	83,287	100.0	149.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	20,379	24.5	36.5
2	Malignant neoplasms (C00-C97)	13,717	16.5	24.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,832	13.0	19.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	9,604	11.5	17.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,352	2.8	4.2
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,245	2.7	4.0
7	Human immunodeficiency virus (HIV) disease (B20-B24)	2,219	2.7	4.0
8	Diabetes mellitus (E10-E14)	1,727	2.1	3.1
9	Cerebrovascular diseases (I60-I69)	1,573	1.9	2.8
10	Influenza and pneumonia (J10-J18)	824	1.0	1.5
11	Chronic lower respiratory diseases (J40-J47)	781	0.9	1.4
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	754	0.9	1.4
13	Septicemia (A40-A41)	675	0.8	1.2
14	Viral hepatitis (B15-B19)	545	0.7	1.0
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	472	0.6	0.8
...	All other causes (Residual)	14,588	17.5	26.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 45-64 years				
...	All causes	310,655	100.0	610.9
1	Malignant neoplasms (C00-C97)	110,418	35.5	217.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	73,700	23.7	144.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17,252	5.6	33.9
4	Chronic lower respiratory diseases (J40-J47)	12,443	4.0	24.5
5	Diabetes mellitus (E10-E14)	9,788	3.2	19.2
6	Cerebrovascular diseases (I60-I69)	9,752	3.1	19.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	9,506	3.1	18.7
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8,857	2.9	17.4
9	Septicemia (A40-A41)	3,515	1.1	6.9
10	Influenza and pneumonia (J10-J18)	3,481	1.1	6.8
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,900	0.9	5.7
12	Viral hepatitis (B15-B19)	2,225	0.7	4.4
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,826	0.6	3.6
14	Aortic aneurysm and dissection (I71)	1,613	0.5	3.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	1,492	0.5	2.9
...	All other causes (Residual)	41,887	13.5	82.4
White, non-Hispanic, both sexes, 65 years and over				
...	All causes	1,548,455	100.0	5,234.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	492,173	31.8	1,663.8
2	Malignant neoplasms (C00-C97)	332,605	21.5	1,124.4
3	Cerebrovascular diseases (I60-I69)	121,566	7.9	411.0
4	Chronic lower respiratory diseases (J40-J47)	98,721	6.4	333.7
5	Alzheimer's disease (G30)	52,981	3.4	179.1
6	Influenza and pneumonia (J10-J18)	50,793	3.3	171.7
7	Diabetes mellitus (E10-E14)	40,830	2.6	138.0
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	29,376	1.9	99.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	27,172	1.8	91.9
10	Septicemia (A40-A41)	21,055	1.4	71.2
11	Parkinson's disease (G20-G21)	15,307	1.0	51.7
12	Pneumonitis due to solids and liquids (J69)	14,184	0.9	48.0
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	13,198	0.9	44.6
14	Atherosclerosis (I70)	11,646	0.8	39.4
15	Aortic aneurysm and dissection (I71)	10,942	0.7	37.0
...	All other causes (Residual)	215,906	13.9	729.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, all ages				
...	All causes	957,645	100.0	983.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	281,442	29.4	289.2
2	Malignant neoplasms (C00-C97)	237,323	24.8	243.8
3	Chronic lower respiratory diseases (J40-J47)	53,662	5.6	55.1
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	50,652	5.3	52.0
5	Cerebrovascular diseases (I60-I69)	49,908	5.2	51.3
6	Diabetes mellitus (E10-E14)	25,273	2.6	26.0
7	Influenza and pneumonia (J10-J18)	23,934	2.5	24.6
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	21,323	2.2	21.9
9	Alzheimer's disease (G30)	15,399	1.6	15.8
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14,883	1.6	15.3
11	Chronic liver disease and cirrhosis (K70,K73-K74)	12,800	1.3	13.2
12	Septicemia (A40-A41)	11,148	1.2	11.5
13	Parkinson's disease (G20-G21)	8,838	0.9	9.1
14	Aortic aneurysm and dissection (I71)	7,834	0.8	8.0
15	Pneumonitis due to solids and liquids (J69)	7,654	0.8	7.9
...	All other causes (Residual)	135,572	14.2	139.3
White, non-Hispanic, male, 1-4 years				
...	All causes	1,431	100.0	30.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	527	36.8	11.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	151	10.6	3.2
3	Malignant neoplasms (C00-C97)	134	9.4	2.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	89	6.2	1.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	54	3.8	1.1
6	Septicemia (A40-A41)	25	1.7	0.5
7	Influenza and pneumonia (J10-J18)	24	1.7	0.5
8	Cerebrovascular diseases (I60-I69)	18	1.3	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	17	1.2	*
9	Chronic lower respiratory diseases (J40-J47)	17	1.2	*
11	Certain conditions originating in the perinatal period (P00-P96)	10	0.7	*
12	Meningococcal infection (A39)	6	0.4	*
12	Meningitis (G00,G03)	6	0.4	*
12	Acute bronchitis and bronchiolitis (J20-J21)	6	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.3	*
...	All other causes (Residual)	342	23.9	7.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 5-14 years				
...	All causes	2,365	100.0	18.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	997	42.2	7.7
2	Malignant neoplasms (C00-C97)	355	15.0	2.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	138	5.8	1.1
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	125	5.3	1.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	85	3.6	0.7
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	61	2.6	0.5
7	Septicemia (A40-A41)	34	1.4	0.3
8	Cerebrovascular diseases (I60-I69)	23	1.0	0.2
8	Chronic lower respiratory diseases (J40-J47)	23	1.0	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	22	0.9	0.2
11	Influenza and pneumonia (J10-J18)	20	0.8	0.2
12	Diabetes mellitus (E10-E14)	16	0.7	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	12	0.5	*
14	Meningitis (G00,G03)	11	0.5	*
15	Meningococcal infection (A39)	8	0.3	*
15	Anemias (D50-D64)	8	0.3	*
...	All other causes (Residual)	427	18.1	3.3
White, non-Hispanic, male, 15-24 years				
...	All causes	13,957	100.0	106.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,746	55.5	59.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,527	18.1	19.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	685	4.9	5.2
4	Malignant neoplasms (C00-C97)	639	4.6	4.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	370	2.7	2.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	178	1.3	1.4
7	Cerebrovascular diseases (I60-I69)	62	0.4	0.5
7	Influenza and pneumonia (J10-J18)	62	0.4	0.5
9	Diabetes mellitus (E10-E14)	51	0.4	0.4
10	Chronic lower respiratory diseases (J40-J47)	48	0.3	0.4
11	Septicemia (A40-A41)	37	0.3	0.3
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	32	0.2	0.2
13	Legal intervention (Y35,Y89.0)	25	0.2	0.2
14	Aortic aneurysm and dissection (I71)	19	0.1	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	0.1	*
...	All other causes (Residual)	1,459	10.5	11.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 25-44 years				
...	All causes	54,388	100.0	194.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14,823	27.3	52.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,834	14.4	28.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7,527	13.8	26.9
4	Malignant neoplasms (C00-C97)	6,259	11.5	22.3
5	Human immunodeficiency virus (HIV) disease (B20-B24)	1,852	3.4	6.6
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,534	2.8	5.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,501	2.8	5.4
8	Diabetes mellitus (E10-E14)	1,080	2.0	3.9
9	Cerebrovascular diseases (I60-I69)	753	1.4	2.7
10	Influenza and pneumonia (J10-J18)	451	0.8	1.6
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	418	0.8	1.5
12	Chronic lower respiratory diseases (J40-J47)	385	0.7	1.4
13	Viral hepatitis (B15-B19)	384	0.7	1.4
14	Septicemia (A40-A41)	335	0.6	1.2
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	255	0.5	0.9
...	All other causes (Residual)	8,997	16.5	32.1
White, non-Hispanic, male, 45-64 years				
...	All causes	190,259	100.0	760.5
1	Malignant neoplasms (C00-C97)	58,909	31.0	235.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	53,087	27.9	212.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,119	6.4	48.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,798	3.6	27.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	6,767	3.6	27.0
6	Chronic lower respiratory diseases (J40-J47)	6,292	3.3	25.1
7	Diabetes mellitus (E10-E14)	5,869	3.1	23.5
8	Cerebrovascular diseases (I60-I69)	5,308	2.8	21.2
9	Influenza and pneumonia (J10-J18)	1,996	1.0	8.0
10	Septicemia (A40-A41)	1,884	1.0	7.5
11	Viral hepatitis (B15-B19)	1,627	0.9	6.5
12	Human immunodeficiency virus (HIV) disease (B20-B24)	1,592	0.8	6.4
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,578	0.8	6.3
14	Aortic aneurysm and dissection (I71)	1,230	0.6	4.9
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	918	0.5	3.7
...	All other causes (Residual)	24,285	12.8	97.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 65 years and over				
...	All causes	687,441	100.0	5,569.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	219,857	32.0	1,781.3
2	Malignant neoplasms (C00-C97)	171,001	24.9	1,385.5
3	Chronic lower respiratory diseases (J40-J47)	46,881	6.8	379.8
4	Cerebrovascular diseases (I60-I69)	43,711	6.4	354.2
5	Influenza and pneumonia (J10-J18)	21,306	3.1	172.6
6	Diabetes mellitus (E10-E14)	18,256	2.7	147.9
7	Alzheimer's disease (G30)	15,167	2.2	122.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14,138	2.1	114.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12,975	1.9	105.1
10	Septicemia (A40-A41)	8,764	1.3	71.0
11	Parkinson's disease (G20-G21)	8,641	1.3	70.0
12	Pneumonitis due to solids and liquids (J69)	7,039	1.0	57.0
13	Aortic aneurysm and dissection (I71)	6,399	0.9	51.8
14	Chronic liver disease and cirrhosis (K70,K73-K74)	4,515	0.7	36.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	4,477	0.7	36.3
...	All other causes (Residual)	84,314	12.3	683.1
White, non-Hispanic, female, all ages				
...	All causes	1,024,328	100.0	1,010.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	296,319	28.9	292.3
2	Malignant neoplasms (C00-C97)	221,431	21.6	218.5
3	Cerebrovascular diseases (I60-I69)	83,210	8.1	82.1
4	Chronic lower respiratory diseases (J40-J47)	58,466	5.7	57.7
5	Alzheimer's disease (G30)	38,087	3.7	37.6
6	Influenza and pneumonia (J10-J18)	31,485	3.1	31.1
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	29,953	2.9	29.6
8	Diabetes mellitus (E10-E14)	27,190	2.7	26.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15,786	1.5	15.6
10	Septicemia (A40-A41)	14,397	1.4	14.2
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	9,292	0.9	9.2
12	Atherosclerosis (I70)	7,683	0.8	7.6
13	Chronic liver disease and cirrhosis (K70,K73-K74)	7,553	0.7	7.5
14	Pneumonitis due to solids and liquids (J69)	7,549	0.7	7.4
15	Parkinson's disease (G20-G21)	6,788	0.7	6.7
...	All other causes (Residual)	169,139	16.5	166.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 1-4 years				
...	All causes	1,071	100.0	23.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	338	31.6	7.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	130	12.1	2.9
3	Malignant neoplasms (C00-C97)	102	9.5	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	80	7.5	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	37	3.5	0.8
6	Influenza and pneumonia (J10-J18)	22	2.1	0.5
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	19	1.8	*
8	Septicemia (A40-A41)	17	1.6	*
9	Acute bronchitis and bronchiolitis (J20-J21)	12	1.1	*
9	Certain conditions originating in the perinatal period (P00-P96)	12	1.1	*
11	Cerebrovascular diseases (I60-I69)	11	1.0	*
12	Anemias (D50-D64)	6	0.6	*
12	Chronic lower respiratory diseases (J40-J47)	6	0.6	*
14	Meningitis (G00,G03)	4	0.4	*
15	Meningococcal infection (A39)	2	0.2	*
15	Diabetes mellitus (E10-E14)	2	0.2	*
15	Pneumonitis due to solids and liquids (J69)	2	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.2	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.2	*
...	All other causes (Residual)	265	24.7	5.9
White, non-Hispanic, female, 5-14 years				
...	All causes	1,666	100.0	13.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	584	35.1	4.8
2	Malignant neoplasms (C00-C97)	287	17.2	2.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	128	7.7	1.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	72	4.3	0.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	61	3.7	0.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	39	2.3	0.3
7	Influenza and pneumonia (J10-J18)	35	2.1	0.3
8	Septicemia (A40-A41)	21	1.3	0.2
8	Cerebrovascular diseases (I60-I69)	21	1.3	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	19	1.1	*
10	Chronic lower respiratory diseases (J40-J47)	19	1.1	*
12	Anemias (D50-D64)	11	0.7	*
13	Meningitis (G00,G03)	9	0.5	*
14	Certain conditions originating in the perinatal period (P00-P96)	8	0.5	*
15	Pneumonitis due to solids and liquids (J69)	4	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.2	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.2	*
...	All other causes (Residual)	340	20.4	2.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 15-24 years				
...	All causes	5,501	100.0	43.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,918	53.0	23.2
2	Malignant neoplasms (C00-C97)	458	8.3	3.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	424	7.7	3.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	273	5.0	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	177	3.2	1.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	137	2.5	1.1
7	Chronic lower respiratory diseases (J40-J47)	46	0.8	0.4
8	Diabetes mellitus (E10-E14)	44	0.8	0.4
9	Influenza and pneumonia (J10-J18)	41	0.7	0.3
10	Cerebrovascular diseases (I60-I69)	39	0.7	0.3
11	Septicemia (A40-A41)	35	0.6	0.3
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	27	0.5	0.2
13	Pregnancy, childbirth and the puerperium (O00-099)	21	0.4	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	0.3	*
15	Pneumonitis due to solids and liquids (J69)	12	0.2	*
...	All other causes (Residual)	833	15.1	6.6
White, non-Hispanic, female, 25-44 years				
...	All causes	28,899	100.0	103.9
1	Malignant neoplasms (C00-C97)	7,458	25.8	26.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,556	19.2	20.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,998	10.4	10.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,077	7.2	7.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	851	2.9	3.1
6	Cerebrovascular diseases (I60-I69)	820	2.8	2.9
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	711	2.5	2.6
8	Diabetes mellitus (E10-E14)	647	2.2	2.3
9	Chronic lower respiratory diseases (J40-J47)	396	1.4	1.4
10	Influenza and pneumonia (J10-J18)	373	1.3	1.3
11	Human immunodeficiency virus (HIV) disease (B20-B24)	367	1.3	1.3
12	Septicemia (A40-A41)	340	1.2	1.2
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	336	1.2	1.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	217	0.8	0.8
15	Viral hepatitis (B15-B19)	161	0.6	0.6
...	All other causes (Residual)	5,591	19.3	20.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 45-64 years				
...	All causes	120,396	100.0	466.1
1	Malignant neoplasms (C00-C97)	51,509	42.8	199.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,613	17.1	79.8
3	Chronic lower respiratory diseases (J40-J47)	6,151	5.1	23.8
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,133	4.3	19.9
5	Cerebrovascular diseases (I60-I69)	4,444	3.7	17.2
6	Diabetes mellitus (E10-E14)	3,919	3.3	15.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,739	2.3	10.6
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,059	1.7	8.0
9	Septicemia (A40-A41)	1,631	1.4	6.3
10	Influenza and pneumonia (J10-J18)	1,485	1.2	5.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,322	1.1	5.1
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	662	0.5	2.6
13	Viral hepatitis (B15-B19)	598	0.5	2.3
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	531	0.4	2.1
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	487	0.4	1.9
...	All other causes (Residual)	17,113	14.2	66.2
White, non-Hispanic, female, 65 years and over				
...	All causes	861,014	100.0	4,994.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	272,316	31.6	1,579.7
2	Malignant neoplasms (C00-C97)	161,604	18.8	937.5
3	Cerebrovascular diseases (I60-I69)	77,855	9.0	451.6
4	Chronic lower respiratory diseases (J40-J47)	51,840	6.0	300.7
5	Alzheimer's disease (G30)	37,814	4.4	219.4
6	Influenza and pneumonia (J10-J18)	29,487	3.4	171.1
7	Diabetes mellitus (E10-E14)	22,574	2.6	131.0
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15,238	1.8	88.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14,197	1.6	82.4
10	Septicemia (A40-A41)	12,291	1.4	71.3
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	8,748	1.0	50.7
12	Atherosclerosis (I70)	7,482	0.9	43.4
13	Pneumonitis due to solids and liquids (J69)	7,145	0.8	41.4
14	Parkinson's disease (G20-G21)	6,666	0.8	38.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	4,829	0.6	28.0
...	All other causes (Residual)	130,928	15.2	759.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, all ages				
...	All causes	286,573	100.0	792.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	76,694	26.8	212.2
2	Malignant neoplasms (C00-C97)	61,996	21.6	171.5
3	Cerebrovascular diseases (I60-I69)	18,691	6.5	51.7
4	Diabetes mellitus (E10-E14)	12,583	4.4	34.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,285	4.3	34.0
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8,147	2.8	22.5
7	Chronic lower respiratory diseases (J40-J47)	7,730	2.7	21.4
8	Human immunodeficiency virus (HIV) disease (B20-B24)	7,714	2.7	21.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,410	2.6	20.5
10	Septicemia (A40-A41)	6,074	2.1	16.8
11	Influenza and pneumonia (J10-J18)	5,803	2.0	16.1
12	Certain conditions originating in the perinatal period (P00-P96)	4,777	1.7	13.2
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	4,165	1.5	11.5
14	Alzheimer's disease (G30)	3,303	1.2	9.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,545	0.9	7.0
...	All other causes (Residual)	46,656	16.3	129.1
Black, non-Hispanic, both sexes, 1-4 years				
...	All causes	1,163	100.0	48.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	350	30.1	14.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	156	13.4	6.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	113	9.7	4.7
4	Malignant neoplasms (C00-C97)	71	6.1	3.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	40	3.4	1.7
5	Influenza and pneumonia (J10-J18)	40	3.4	1.7
7	Chronic lower respiratory diseases (J40-J47)	29	2.5	1.2
8	Certain conditions originating in the perinatal period (P00-P96)	26	2.2	1.1
9	Septicemia (A40-A41)	17	1.5	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	14	1.2	*
11	Cerebrovascular diseases (I60-I69)	10	0.9	*
12	Anemias (D50-D64)	7	0.6	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	5	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.4	*
15	Meningitis (G00,G03)	4	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.3	*
...	All other causes (Residual)	272	23.4	11.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 5-14 years				
...	All causes	1,634	100.0	25.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	607	37.1	9.4
2	Malignant neoplasms (C00-C97)	168	10.3	2.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	132	8.1	2.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	81	5.0	1.3
5	Chronic lower respiratory diseases (J40-J47)	74	4.5	1.1
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	65	4.0	1.0
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	35	2.1	0.5
8	Anemias (D50-D64)	26	1.6	0.4
9	Cerebrovascular diseases (I60-I69)	23	1.4	0.4
10	Septicemia (A40-A41)	20	1.2	0.3
11	Human immunodeficiency virus (HIV) disease (B20-B24)	19	1.2	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	18	1.1	*
12	Influenza and pneumonia (J10-J18)	18	1.1	*
14	Diabetes mellitus (E10-E14)	14	0.9	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	14	0.9	*
...	All other causes (Residual)	320	19.6	5.0
Black, non-Hispanic, both sexes, 15-24 years				
...	All causes	6,904	100.0	116.9
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,854	41.3	48.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,680	24.3	28.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	392	5.7	6.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	299	4.3	5.1
5	Malignant neoplasms (C00-C97)	242	3.5	4.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	124	1.8	2.1
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	98	1.4	1.7
8	Anemias (D50-D64)	78	1.1	1.3
8	Chronic lower respiratory diseases (J40-J47)	78	1.1	1.3
10	Diabetes mellitus (E10-E14)	57	0.8	1.0
11	Pregnancy, childbirth and the puerperium (O00-099)	45	0.7	0.8
12	Cerebrovascular diseases (I60-I69)	38	0.6	0.6
12	Influenza and pneumonia (J10-J18)	38	0.6	0.6
14	Legal intervention (Y35,Y89.0)	37	0.5	0.6
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	36	0.5	0.6
...	All other causes (Residual)	808	11.7	13.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 25-44 years				
...	All causes	29,964	100.0	279.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,305	14.4	40.1
2	Human immunodeficiency virus (HIV) disease (B20-B24)	4,121	13.8	38.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,903	13.0	36.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,782	12.6	35.2
5	Malignant neoplasms (C00-C97)	3,533	11.8	32.9
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	929	3.1	8.7
7	Cerebrovascular diseases (I60-I69)	907	3.0	8.4
8	Diabetes mellitus (E10-E14)	748	2.5	7.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	414	1.4	3.9
10	Chronic lower respiratory diseases (J40-J47)	394	1.3	3.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	362	1.2	3.4
12	Septicemia (A40-A41)	356	1.2	3.3
13	Influenza and pneumonia (J10-J18)	330	1.1	3.1
14	Anemias (D50-D64)	269	0.9	2.5
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	233	0.8	2.2
...	All other causes (Residual)	5,378	17.9	50.1
Black, non-Hispanic, both sexes, 45-64 years				
...	All causes	76,089	100.0	1,076.4
1	Malignant neoplasms (C00-C97)	21,069	27.7	298.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	19,872	26.1	281.1
3	Cerebrovascular diseases (I60-I69)	4,177	5.5	59.1
4	Diabetes mellitus (E10-E14)	3,597	4.7	50.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,220	4.2	45.6
6	Human immunodeficiency virus (HIV) disease (B20-B24)	3,144	4.1	44.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,817	2.4	25.7
8	Chronic lower respiratory diseases (J40-J47)	1,656	2.2	23.4
9	Chronic liver disease and cirrhosis (K70,K73-K74)	1,605	2.1	22.7
10	Septicemia (A40-A41)	1,461	1.9	20.7
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	963	1.3	13.6
12	Influenza and pneumonia (J10-J18)	953	1.3	13.5
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	926	1.2	13.1
14	Viral hepatitis (B15-B19)	552	0.7	7.8
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	390	0.5	5.5
...	All other causes (Residual)	10,687	14.0	151.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 65 years and over				
...	All causes	162,496	100.0	5,563.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	51,961	32.0	1,779.0
2	Malignant neoplasms (C00-C97)	36,898	22.7	1,263.3
3	Cerebrovascular diseases (I60-I69)	13,499	8.3	462.2
4	Diabetes mellitus (E10-E14)	8,166	5.0	279.6
5	Chronic lower respiratory diseases (J40-J47)	5,484	3.4	187.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,079	3.1	173.9
7	Influenza and pneumonia (J10-J18)	4,338	2.7	148.5
8	Septicemia (A40-A41)	4,110	2.5	140.7
9	Alzheimer's disease (G30)	3,264	2.0	111.7
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,961	1.8	101.4
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,197	1.4	75.2
12	Pneumonitis due to solids and liquids (J69)	1,302	0.8	44.6
13	Atherosclerosis (I70)	873	0.5	29.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	724	0.4	24.8
15	Aortic aneurysm and dissection (I71)	696	0.4	23.8
...	All other causes (Residual)	20,944	12.9	717.0
Black, non-Hispanic, male, all ages				
...	All causes	144,802	100.0	842.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	36,596	25.3	212.9
2	Malignant neoplasms (C00-C97)	32,287	22.3	187.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,428	5.8	49.0
4	Cerebrovascular diseases (I60-I69)	7,754	5.4	45.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6,780	4.7	39.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	5,203	3.6	30.3
7	Diabetes mellitus (E10-E14)	5,158	3.6	30.0
8	Chronic lower respiratory diseases (J40-J47)	4,282	3.0	24.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,389	2.3	19.7
10	Influenza and pneumonia (J10-J18)	2,734	1.9	15.9
11	Septicemia (A40-A41)	2,668	1.8	15.5
12	Certain conditions originating in the perinatal period (P00-P96)	2,640	1.8	15.4
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,701	1.2	9.9
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,595	1.1	9.3
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,592	1.1	9.3
...	All other causes (Residual)	21,995	15.2	127.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 1-4 years				
...	All causes	681	100.0	55.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	231	33.9	18.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	89	13.1	7.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	58	8.5	4.8
4	Malignant neoplasms (C00-C97)	33	4.8	2.7
5	Influenza and pneumonia (J10-J18)	22	3.2	1.8
6	Chronic lower respiratory diseases (J40-J47)	20	2.9	1.6
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	18	2.6	*
7	Certain conditions originating in the perinatal period (P00-P96)	18	2.6	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	8	1.2	*
10	Cerebrovascular diseases (I60-I69)	6	0.9	*
11	Septicemia (A40-A41)	5	0.7	*
12	Anemias (D50-D64)	4	0.6	*
13	Meningitis (G00,G03)	3	0.4	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.4	*
15	Meningococcal infection (A39)	2	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.3	*
...	All other causes (Residual)	159	23.3	13.0
Black, non-Hispanic, male, 5-14 years				
...	All causes	977	100.0	29.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	391	40.0	11.9
2	Malignant neoplasms (C00-C97)	103	10.5	3.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	84	8.6	2.6
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	47	4.8	1.4
5	Chronic lower respiratory diseases (J40-J47)	39	4.0	1.2
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	33	3.4	1.0
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	29	3.0	0.9
8	Cerebrovascular diseases (I60-I69)	15	1.5	*
9	Anemias (D50-D64)	14	1.4	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	10	1.0	*
10	Influenza and pneumonia (J10-J18)	10	1.0	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	10	1.0	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	9	0.9	*
14	Septicemia (A40-A41)	7	0.7	*
15	Certain conditions originating in the perinatal period (P00-P96)	6	0.6	*
...	All other causes (Residual)	170	17.4	5.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 15-24 years				
...	All causes	5,248	100.0	177.4
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,540	48.4	85.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,260	24.0	42.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	342	6.5	11.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	185	3.5	6.3
5	Malignant neoplasms (C00-C97)	137	2.6	4.6
6	Human immunodeficiency virus (HIV) disease (B20-B24)	60	1.1	2.0
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	59	1.1	2.0
8	Chronic lower respiratory diseases (J40-J47)	58	1.1	2.0
9	Anemias (D50-D64)	49	0.9	1.7
10	Legal intervention (Y35,Y89.0)	36	0.7	1.2
11	Diabetes mellitus (E10-E14)	34	0.6	1.1
12	Cerebrovascular diseases (I60-I69)	22	0.4	0.7
13	Influenza and pneumonia (J10-J18)	20	0.4	0.7
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	0.3	*
15	Septicemia (A40-A41)	10	0.2	*
...	All other causes (Residual)	421	8.0	14.2
Black, non-Hispanic, male, 25-44 years				
...	All causes	18,384	100.0	362.8
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,134	17.0	61.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,841	15.5	56.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,643	14.4	52.2
4	Human immunodeficiency virus (HIV) disease (B20-B24)	2,602	14.2	51.3
5	Malignant neoplasms (C00-C97)	1,445	7.9	28.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	786	4.3	15.5
7	Diabetes mellitus (E10-E14)	428	2.3	8.4
8	Cerebrovascular diseases (I60-I69)	403	2.2	8.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	210	1.1	4.1
10	Chronic lower respiratory diseases (J40-J47)	206	1.1	4.1
10	Chronic liver disease and cirrhosis (K70,K73-K74)	206	1.1	4.1
12	Influenza and pneumonia (J10-J18)	179	1.0	3.5
13	Septicemia (A40-A41)	177	1.0	3.5
14	Anemias (D50-D64)	138	0.8	2.7
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	136	0.7	2.7
...	All other causes (Residual)	2,850	15.5	56.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 45-64 years				
...	All causes	44,545	100.0	1,383.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,242	27.5	380.1
2	Malignant neoplasms (C00-C97)	11,309	25.4	351.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,314	5.2	71.9
4	Human immunodeficiency virus (HIV) disease (B20-B24)	2,303	5.2	71.5
5	Cerebrovascular diseases (I60-I69)	2,274	5.1	70.6
6	Diabetes mellitus (E10-E14)	1,848	4.1	57.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,073	2.4	33.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,001	2.2	31.1
9	Chronic lower respiratory diseases (J40-J47)	911	2.0	28.3
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	752	1.7	23.3
11	Septicemia (A40-A41)	730	1.6	22.7
12	Influenza and pneumonia (J10-J18)	577	1.3	17.9
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	508	1.1	15.8
14	Viral hepatitis (B15-B19)	384	0.9	11.9
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	309	0.7	9.6
...	All other causes (Residual)	6,010	13.5	186.6
Black, non-Hispanic, male, 65 years and over				
...	All causes	70,418	100.0	6,312.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	21,393	30.4	1,917.6
2	Malignant neoplasms (C00-C97)	19,254	27.3	1,725.8
3	Cerebrovascular diseases (I60-I69)	5,015	7.1	449.5
4	Chronic lower respiratory diseases (J40-J47)	3,040	4.3	272.5
5	Diabetes mellitus (E10-E14)	2,842	4.0	254.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,123	3.0	190.3
7	Influenza and pneumonia (J10-J18)	1,879	2.7	168.4
8	Septicemia (A40-A41)	1,692	2.4	151.7
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,215	1.7	108.9
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,054	1.5	94.5
11	Alzheimer's disease (G30)	919	1.3	82.4
12	Pneumonitis due to solids and liquids (J69)	622	0.9	55.8
13	Atherosclerosis (I70)	323	0.5	29.0
14	Chronic liver disease and cirrhosis (K70,K73-K74)	311	0.4	27.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	306	0.4	27.4
...	All other causes (Residual)	8,430	12.0	755.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, all ages				
...	All causes	141,771	100.0	748.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	40,098	28.3	211.6
2	Malignant neoplasms (C00-C97)	29,709	21.0	156.7
3	Cerebrovascular diseases (I60-I69)	10,937	7.7	57.7
4	Diabetes mellitus (E10-E14)	7,425	5.2	39.2
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,021	2.8	21.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,857	2.7	20.3
7	Chronic lower respiratory diseases (J40-J47)	3,448	2.4	18.2
8	Septicemia (A40-A41)	3,406	2.4	18.0
9	Influenza and pneumonia (J10-J18)	3,069	2.2	16.2
10	Human immunodeficiency virus (HIV) disease (B20-B24)	2,511	1.8	13.2
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,464	1.7	13.0
12	Alzheimer's disease (G30)	2,363	1.7	12.5
13	Certain conditions originating in the perinatal period (P00-P96)	2,137	1.5	11.3
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,367	1.0	7.2
15	Chronic liver disease and cirrhosis (K70,K73-K74)	953	0.7	5.0
...	All other causes (Residual)	24,006	16.9	126.7
Black, non-Hispanic, female, 1-4 years				
...	All causes	482	100.0	40.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	119	24.7	10.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	67	13.9	5.7
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	55	11.4	4.7
4	Malignant neoplasms (C00-C97)	38	7.9	3.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	22	4.6	1.9
6	Influenza and pneumonia (J10-J18)	18	3.7	*
7	Septicemia (A40-A41)	12	2.5	*
8	Chronic lower respiratory diseases (J40-J47)	9	1.9	*
9	Certain conditions originating in the perinatal period (P00-P96)	8	1.7	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	6	1.2	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.8	*
11	Cerebrovascular diseases (I60-I69)	4	0.8	*
13	Anemias (D50-D64)	3	0.6	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.6	*
15	Pneumonitis due to solids and liquids (J69)	2	0.4	*
...	All other causes (Residual)	112	23.2	9.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 5-14 years				
...	All causes	657	100.0	20.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	216	32.9	6.8
2	Malignant neoplasms (C00-C97)	65	9.9	2.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	48	7.3	1.5
4	Chronic lower respiratory diseases (J40-J47)	35	5.3	1.1
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	34	5.2	1.1
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	32	4.9	1.0
7	Septicemia (A40-A41)	13	2.0	*
8	Anemias (D50-D64)	12	1.8	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	9	1.4	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	9	1.4	*
9	Diabetes mellitus (E10-E14)	9	1.4	*
12	Cerebrovascular diseases (I60-I69)	8	1.2	*
12	Influenza and pneumonia (J10-J18)	8	1.2	*
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6	0.9	*
15	Meningitis (G00,G03)	5	0.8	*
...	All other causes (Residual)	148	22.5	4.7
Black, non-Hispanic, female, 15-24 years				
...	All causes	1,656	100.0	56.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	420	25.4	14.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	314	19.0	10.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	114	6.9	3.9
4	Malignant neoplasms (C00-C97)	105	6.3	3.6
5	Human immunodeficiency virus (HIV) disease (B20-B24)	64	3.9	2.2
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	50	3.0	1.7
7	Pregnancy, childbirth and the puerperium (O00-099)	45	2.7	1.5
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	39	2.4	1.3
9	Anemias (D50-D64)	29	1.8	1.0
10	Diabetes mellitus (E10-E14)	23	1.4	0.8
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21	1.3	0.7
12	Chronic lower respiratory diseases (J40-J47)	20	1.2	0.7
13	Influenza and pneumonia (J10-J18)	18	1.1	*
14	Cerebrovascular diseases (I60-I69)	16	1.0	*
15	Septicemia (A40-A41)	14	0.8	*
...	All other causes (Residual)	364	22.0	12.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Black, non-Hispanic, female, 25-44 years			
...	All causes	11,580	100.0	204.2
1	Malignant neoplasms (C00-C97)	2,088	18.0	36.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,662	14.4	29.3
3	Human immunodeficiency virus (HIV) disease (B20-B24)	1,519	13.1	26.8
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,062	9.2	18.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	648	5.6	11.4
6	Cerebrovascular diseases (I60-I69)	504	4.4	8.9
7	Diabetes mellitus (E10-E14)	320	2.8	5.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	204	1.8	3.6
9	Chronic lower respiratory diseases (J40-J47)	188	1.6	3.3
10	Septicemia (A40-A41)	179	1.5	3.2
11	Chronic liver disease and cirrhosis (K70,K73-K74)	156	1.3	2.8
12	Influenza and pneumonia (J10-J18)	151	1.3	2.7
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	143	1.2	2.5
14	Anemias (D50-D64)	131	1.1	2.3
15	Pregnancy, childbirth and the puerperium (O00-099)	105	0.9	1.9
...	All other causes (Residual)	2,520	21.8	44.4
	Black, non-Hispanic, female, 45-64 years			
...	All causes	31,544	100.0	819.7
1	Malignant neoplasms (C00-C97)	9,760	30.9	253.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,630	24.2	198.3
3	Cerebrovascular diseases (I60-I69)	1,903	6.0	49.4
4	Diabetes mellitus (E10-E14)	1,749	5.5	45.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	906	2.9	23.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	841	2.7	21.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	816	2.6	21.2
8	Chronic lower respiratory diseases (J40-J47)	745	2.4	19.4
9	Septicemia (A40-A41)	731	2.3	19.0
10	Chronic liver disease and cirrhosis (K70,K73-K74)	532	1.7	13.8
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	455	1.4	11.8
12	Influenza and pneumonia (J10-J18)	376	1.2	9.8
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	174	0.6	4.5
14	Viral hepatitis (B15-B19)	168	0.5	4.4
15	Anemias (D50-D64)	134	0.4	3.5
...	All other causes (Residual)	4,624	14.7	120.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2002

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Black, non-Hispanic, female, 65 years and over			
...	All causes	92,078	100.0	5,100.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	30,568	33.2	1,693.3
2	Malignant neoplasms (C00-C97)	17,644	19.2	977.4
3	Cerebrovascular diseases (I60-I69)	8,484	9.2	470.0
4	Diabetes mellitus (E10-E14)	5,324	5.8	294.9
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,956	3.2	163.7
6	Influenza and pneumonia (J10-J18)	2,459	2.7	136.2
7	Chronic lower respiratory diseases (J40-J47)	2,444	2.7	135.4
8	Septicemia (A40-A41)	2,418	2.6	133.9
9	Alzheimer's disease (G30)	2,345	2.5	129.9
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,907	2.1	105.6
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	982	1.1	54.4
12	Pneumonitis due to solids and liquids (J69)	680	0.7	37.7
13	Atherosclerosis (I70)	550	0.6	30.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown beha	418	0.5	23.2
15	Aortic aneurysm and dissection (I71)	400	0.4	22.2
...	All other causes (Residual)	12,499	13.6	692.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable